

# Presentation

Carson James

May 13, 2022

## Definition

We define

$\Lambda_+^{n \times r} = \{\Sigma \in \mathbb{R}^{n \times r} : \Sigma \text{ is diagonal and positive semi-definite}\}$  and  
 $O_n = \{U \in \mathbb{R}^{n \times n} : U \text{ is orthogonal}\}.$

## Model

1. Fix  $M \in \mathbb{R}^{n \times n_M}$  columns of  $M$  are orthogonal and set  $P_M = M(M^T M)^{-1}M^T$ ?
2. Choose  $\Sigma_Z \in \Lambda_+^{n_M \times r}$ ,  $\Sigma_X \in \Lambda_+^{n \times p}$ ,  $U_Z \in O_r$  and  $U_X \in O_p$ .
3. Set  $V_Z^T = \Sigma_Z U_Z$  and  $V_X^T = \Sigma_X U_X$ .
4. Set  $J_Z = M V_Z^T$  and  $J_X = M V_X^T$ .
5. Choose  $I_Z \in \mathbb{R}^{n \times r}$ ,  $I_X \in \mathbb{R}^{n \times p}$  such that  $\mathcal{C}(I_Z) \cup \mathcal{C}(I_X) \subset \mathcal{C}(I - P_M)$ .
6. Choose  $E_X \in \mathbb{R}_{n \times p}$  with  $(E_X)_{i,j} \sim N(0, \sigma^2)$
7. Set  $Z = J_Z + I_Z$  and  $X = J_X + I_X + E_X$

Then  $\mathcal{C}(M) \perp \mathcal{C}(I_Z), \mathcal{C}(I_X)$ .

## Model

1. Fix  $M \in \mathbb{R}^{n \times n_M}$  columns of  $M$  are orthogonal and set  $P_M = M(M^T M)^{-1}M^T$ ?
2. Choose  $\Sigma_Z \in \Lambda_+^{n_M \times r}$ ,  $\Sigma_X \in \Lambda_+^{n \times p}$ ,  $U_Z \in O_r$  and  $U_X \in O_p$ .
3. Set  $V_Z^T = \Sigma_Z U_Z$  and  $V_X^T = \Sigma_X U_X$ .
4. Set  $J_Z = M V_Z^T$  and  $J_X = M V_X^T$ .
5. Choose  $I_Z \in \mathbb{R}^{n \times r}$ ,  $I_X \in \mathbb{R}^{n \times p}$  such that  $\mathcal{C}(I_Z) \cup \mathcal{C}(I_X) \subset \mathcal{C}(I - P_M)$ .
6. Choose  $E_X \in \mathbb{R}_{n \times p}$  with  $(E_X)_{i,j} \sim N(0, \sigma^2)$
7. Set  $Z = J_Z + I_Z$  and  $X = J_X + I_X + E_X$

Then  $\mathcal{C}(M) \perp \mathcal{C}(I_Z), \mathcal{C}(I_X)$ .

## Model

We consider a modification of the planted partition model which is a submodel of the stochastic block model with  $n$  nodes and  $r$  blocks (for now  $r = 2$ ).

1. Choose  $U \in \mathbb{R}^{n \times 2}$  such that for each  $i \in \{1, \dots, n\}$ ,

$$U_{i,j} = \begin{cases} 1 & \text{node } i \text{ is in block } j \\ 0 & \text{else} \end{cases}$$

as in the stochastic block model. We choose

$$U = \begin{pmatrix} 1 & 0 \\ \vdots & \vdots \\ 1 & 0 \\ 0 & 1 \\ \vdots & \vdots \\ 0 & 1 \end{pmatrix}$$

2. Choose  $W \in \mathbb{R}^{n \times 1}$ , with  $W = (1, -1, \dots, 1, -1)^T$  and

## Model

We consider another modification of the planted partition model which is a submodel of the stochastic block model with  $n$  nodes and  $r$  blocks (for now  $r = 2$ ).

1. Choose  $U \in \mathbb{R}^{n \times 2}$  such that for each  $i \in \{1, \dots, n\}$ ,

$$U_{i,j} = \begin{cases} 1 & \text{node } i \text{ is in block } j \\ 0 & \text{else} \end{cases}$$

as in the stochastic block model. We choose

$$U = \begin{pmatrix} 1 & 0 \\ \vdots & \vdots \\ 1 & 0 \\ 0 & 1 \\ \vdots & \vdots \\ 0 & 1 \end{pmatrix}$$

2. Choose  $W \in \mathbb{R}^{n \times 1}$ , with  $W = (1, -1, \dots, 1, -1)^T$  and