

# INTRODUCTION TO ANALYSIS

CARSON JAMES

## CONTENTS

Preface	2
1. Set Theory	3
1.1. Product Sets	3
1.2. Quotient Sets	4
2. Real and Complex Numbers	4
2.1. Real Numbers	4
3. Metric Spaces	5
3.1. Introduction	5
3.2. Product Spaces	9
4. Topology	10
4.1. Introduction	10
4.2. Continuous Maps	14
4.3. Nets	20
4.4. Product Topology	26
4.5. Subspace Topology	27
4.6. Quotient Topology	28
4.7. Separation and Countability	31
4.8. Compactness	33
4.9. Semi-continuity	34
5. Locally Convex Spaces	36
5.1. Topological Vector Spaces	36
5.2. Sublinear Functionals	40
5.3. Seminorms	43
5.4. Minkowski Functionals	48
5.5. Locally Convex Spaces	58
5.6. Quotient Spaces	62
5.7. Duality	64
6. Banach Spaces	66
6.1. Introduction	66
6.2. Bounded Operators	68
6.3. Direct Sums	74
6.4. Quotient Spaces	77
6.5. Applications of the Hahn-Banach Theorem	81
6.6. The Baire Category and Closed Graph Theorems	86
6.7. Duality	91
6.8. Compact Operators	97
6.9. Multilinear Maps	98

6.10. Banach Algebras	101
7. Hilbert Spaces	104
7.1. Introduction	104
7.2. Operators and Functionals	110
7.3. Tensor Products	113
8. Differentiation	118
8.1. The Gateaux Derivative	118
8.2. The Frechet Derivative	122
8.3. The Calc I Derivative	126
8.4. Mean Value Theorem	128
8.5. Taylor's Theorem	130
8.6. The Gradient	136
9. Convexity	137
9.1. Introduction	137
9.2. The Subdifferential	143
9.3. Conjugacy	150
10. Topological Groups	153
10.1. Topological Groups	153
10.2. Automorphism Groups of Metric Spaces	156
11. Group Actions on Metric Spaces	158
11.1. Introduction	158
11.2. Induced Metrics on Orbit Spaces	159
11.3. Fundamental Examples	163
12. Appendix	165
12.1. Summation	165
12.2. Asymptotic Notation	166
References	167

## PREFACE

[cc-by-nc-sa](#)

## 1. SET THEORY

## 1.1. Product Sets.

**Definition 1.1.1.** Let  $(X_\alpha)_{\alpha \in A}$  be a collection of sets. We define the **Cartesian product**, denoted  $\prod_{\alpha \in A} X_\alpha$ , by

$$\prod_{\alpha \in A} X_\alpha = \{f : A \rightarrow \bigcup_{\alpha \in A} X_\alpha : \text{for each } \alpha \in A, f(\alpha) \in X_\alpha\}$$

**Definition 1.1.2.** Let  $(X_\alpha)_{\alpha \in A}$  be a collection of sets. For  $\alpha \in A$ , we define the **projection map onto**  $X_\alpha$ , denoted  $\pi_\alpha : \prod_{\alpha \in A} X_\alpha \rightarrow X_\alpha$ , by

$$\pi_\alpha(f) = f(\alpha)$$

## 1.2. Quotient Sets.

**Definition 1.2.1.** Let  $X$  be a set and  $\sim$  an equivalence relation on  $X$ . We define the **quotient set** of  $X$  by  $\sim$ , denoted  $X/\sim$ , by

$$X/\sim = \{\bar{x} : x \in X\}$$

## 2. REAL AND COMPLEX NUMBERS

**Note 2.0.1.** As a starting point, we will take as fact the existence of the **natural numbers**

$$\mathbb{N} = \{1, 2, \dots\}$$

the **integers**

$$\mathbb{Z} = \{\dots, -2, -1, 0, 1, 2, \dots\}$$

and the **rational numbers**

$$\mathbb{Q} = \left\{ \frac{a}{b} : a \in \mathbb{Z}, b \in \mathbb{N} \right\}$$

## 2.1. Real Numbers.

**Definition 2.1.1.** Let  $X$  be a set and  $\leq$  a relation on  $X$ . Then  $\leq$  is said to be a **total order** if for each  $a, b, c \in X$ ,

- (1)  $a \leq a$
- (2)  $a \leq b$  and  $b \leq c$  implies that  $a \leq c$
- (3)  $a \leq b$  and  $b \leq a$  implies that  $a = b$
- (4)  $a \leq b$  or  $b \leq a$

**Exercise 2.1.2.** We define the relation  $\leq$  on  $\mathbb{Q}$  defined by

$$\frac{a}{b} \leq \frac{c}{d} \text{ iff } ad \leq bc$$

Then  $\leq$  is a total order of  $\mathbb{Q}$ .

*Proof.* Let  $\frac{a}{b}, \frac{c}{d}, \frac{e}{f} \in \mathbb{Q}$ . Then

- (1)  $\frac{a}{b} \leq \frac{a}{b}$  since  $ab \leq ab$ .
- (2) if  $\frac{a}{b} \leq \frac{c}{d}$  and  $\frac{c}{d} \leq \frac{e}{f}$ , then  $ad \leq bc$  and  $cf \leq de$ . Multiplying the first inequality by  $f$  and the second inequality by  $b$ , we obtain  $adf \leq bcf \leq bde$ . Dividing both sides by  $d$  yields  $af \leq be$ . Hence  $\frac{a}{b} \leq \frac{e}{f}$ .
- (3) if  $\frac{a}{b} \leq \frac{c}{d}$  and  $\frac{c}{d} \leq \frac{a}{b}$ , then  $ad \leq bc$  and  $bc \leq ad$ . This implies that  $ad = bc$ . Hence  $\frac{a}{b} = \frac{c}{d}$ .
- (4)

□

## 3. METRIC SPACES

## 3.1. Introduction.

**Definition 3.1.1.** Let  $M$  be a set and  $d : M \times M \rightarrow \mathbb{R}$ . Then  $d$  is said to be a **metric on**  $M$  if for each  $x, y, z \in M$ ,

- (1)  $d(x, y) = 0$  iff  $x = y$
- (2)  $d(x, y) \leq d(x, z) + d(z, y)$

**Exercise 3.1.2.** Let  $M$  be a set and  $d : M \times M \rightarrow \mathbb{R}$  a metric on  $M$ . Then for each  $x, y \in M$ ,  $d(x, y) \geq 0$ .

*Proof.* Let  $x, y, z \in M$ . Then  $d(x, z) \leq d(x, y) + d(y, z)$ . This implies that  $d(x, z) - d(x, y) \leq d(y, z)$ . Since  $z$  is arbitrary, taking  $z = x$ , we obtain

$$\begin{aligned} d(x, x) - d(x, y) &\leq d(y, x) \implies -d(x, y) \leq d(x, y) \\ &\implies 0 \leq 2d(x, y) \\ &\implies d(x, y) \geq 0 \end{aligned}$$

□

**Definition 3.1.3.** Let  $M$  be a set and  $d : M \times M \rightarrow [0, \infty)$  a metric. Then  $(M, d)$  is called a **metric space**.

**Note 3.1.4.** We usually suppress the metric and write  $M$  in place of  $(M, d)$ .

**Definition 3.1.5.** Let  $M$  be a metric space,  $x \in M$  and  $r > 0$ . We define the

- **open ball of radius  $r$  at  $x$** , denoted  $B(x, r)$ , by

$$B(x, r) = \{y \in M : d(x, y) < r\}$$

- **closed ball of radius  $r$  at  $x$** , denoted  $\bar{B}(x, r)$ , by

$$\bar{B}(x, r) = \{y \in M : d(x, y) \leq r\}$$

**Definition 3.1.6.** Let  $M$  be a metric space and  $A \subset M$ . Then  $A$  is said to be

- **open** if for each  $x \in A$ , there exists  $r > 0$  such that  $B(x, r) \subset A$
- **closed** if  $A^c$  is open

**Definition 3.1.7.** Let  $M$  be a metric space. Then  $M$  is said to be **separable** if there exists  $D \subset M$  such that  $D$  is countable and for each  $x \in M$  and  $\epsilon > 0$ , there exists  $y \in D$  such that  $d(x, y) < \epsilon$ .

**Exercise 3.1.8.** Let  $M$  be a metric space. If  $M$  is separable, then for each  $A \subset M$ , if  $A$  is open, then

- (1) there there exist  $(x_n)_{n \in \mathbb{N}} \subset X$  and  $(r_n)_{n \in \mathbb{N}} \subset (0, \infty)$  such that

$$A = \bigcup_{n \in \mathbb{N}} B(x_n, r_n)$$

i.e.  $A$  is a countable union of open balls

- (2) there there exist  $(x_n)_{n \in \mathbb{N}} \subset X$  and  $(r_n)_{n \in \mathbb{N}} \subset (0, \infty)$  such that

$$A = \bigcup_{n \in \mathbb{N}} \bar{B}(x_n, r_n)$$

i.e.  $A$  is a countable union of closed balls.

*Proof.* Suppose that  $M$  is separable. Then there exists  $(x_n)_{n \in \mathbb{N}} \subset M$  such that for each  $x \in M$  and  $\epsilon > 0$ , there exists  $N \in \mathbb{N}$  such that  $d(x, x_N) < \epsilon$ . Let  $A \subset X$ . Suppose that  $A$  is open.

(1) Set

$$\mathcal{B} = \{B(x_n, r) : r \in \mathbb{Q} \text{ and } B(x_n, r) \subset A\}$$

Note that  $\mathcal{B}$  is countable. Let  $x \in A$ . Since  $A$  is open, there exists  $s \in \mathbb{R}$  such that  $B(x, s) \subset A$ . Then there exists  $r \in \mathbb{Q} \cap (0, s)$ . Choose  $N \in \mathbb{N}$  such that  $d(x, x_N) < r/2$ . Let  $y \in B(x_N, r/2)$ , then

$$\begin{aligned} d(x, y) &\leq d(x, x_N) + d(x_N, y) \\ &< r/2 + r/2 \\ &= r \end{aligned}$$

Therefore

$$\begin{aligned} x &\in B(x_N, r/2) \\ &\subset B(x, r) \\ &\subset A \end{aligned}$$

Hence  $B(x_N, r/2) \in \mathcal{B}$  and  $x \in \bigcup_{B \in \mathcal{B}} B$ . Since  $x \in A$  is arbitrary,  $A \subset \bigcup_{B \in \mathcal{B}} B$ .

(2) Similar, but take  $r/4$  instead of  $r/2$ .

□

**Definition 3.1.9.** Let  $(M, d)$  be a metric space and  $A, B \subset M$ . We define the **distance between  $A$  and  $B$** , denoted  $d(A, B)$ , by

$$d(A, B) = \inf_{\substack{a \in A \\ b \in B}} d(a, b)$$

**Exercise 3.1.10.** Let  $(M, d)$  be a metric space. Then for each  $A, B \subset M$  and  $c \in M$ ,

$$d(A, B) \leq d(A, c) + d(c, B)$$

*Proof.* Let  $A, B \subset M$ ,  $c \in M$  and  $\epsilon > 0$ . Choose  $a \in A$  and  $b \in B$  such that  $d(a, c) < d(A, c) + \epsilon/2$  and  $d(c, b) < d(c, B) + \epsilon/2$ . Then

$$\begin{aligned} d(A, B) &\leq d(a, b) \\ &\leq d(a, c) + d(c, b) \\ &< d(A, c) + \frac{\epsilon}{2} + d(c, B) + \frac{\epsilon}{2} \\ &= d(A, c) + d(c, B) + \epsilon \end{aligned}$$

Since  $\epsilon > 0$  is arbitrary,  $d(A, B) \leq d(A, c) + d(c, B)$ . □

**Definition 3.1.11.** Let  $M$  be a set,  $d_1, d_2 : M \times M \rightarrow [0, \infty)$  metrics on  $M$ . Then  $d_1$  and  $d_2$  are said to be

- **topologically equivalent** if for each  $(x_n)_{n \in \mathbb{N}} \subset M$  and  $x \in M$ ,  $x_n \xrightarrow{d_1} x$  iff  $x_n \xrightarrow{d_2} x$
- **equivalent** if there exist  $A, B > 0$  such that

$$Ad_1 \leq d_2 \leq Bd_1$$

**Definition 3.1.12.** Let  $(X, d_X)$  and  $(Y, d_Y)$  be metric spaces and  $f : X \rightarrow Y$ . Then  $f$  is said to be **Lipchitz** if there exists  $K \geq 0$  such that for each  $a, b \in X$ ,

$$d_Y(f(a), f(b)) \leq K d_X(a, b)$$

**Exercise 3.1.13.** Let  $(X, d_X)$  and  $(Y, d_Y)$  be metric spaces and  $f : X \rightarrow Y$ . If  $f$  is Lipchitz, then  $f$  is uniformly continuous.

*Proof.* By definition, there exists  $K \geq 0$  such that for each  $a, b \in X$ ,

$$d_Y(f(a), f(b)) \leq K d_X(a, b)$$

Let  $\epsilon > 0$ . Choose  $\delta = \epsilon/(K + 1)$ . Let  $a, b \in X$ . Suppose that  $d_X(a, b) < \delta$ . Then

$$\begin{aligned} d_Y(f(a), f(b)) &\leq K d_X(a, b) \\ &< K \delta \\ &= K \frac{\epsilon}{K + 1} \\ &< \epsilon \end{aligned}$$

□

**Definition 3.1.14.** Let  $(X, d_X)$  and  $(Y, d_Y)$  be metric spaces and  $f : X \rightarrow Y$  and  $x_0 \in X$ . Then  $f$  is said to be **locally Lipchitz at  $x_0$**  if there exists  $U \in \mathcal{N}_{x_0}$  such that  $f$  is Lipschitz on  $U$ .

**Definition 3.1.15.** Let  $(X, d_X)$  and  $(Y, d_Y)$  be metric spaces and  $f : X \rightarrow Y$ . Then  $f$  is said to be **locally Lipchitz** if for each  $x_0 \in X$ ,  $f$  is locally Lipschitz at  $x_0$ .

**Definition 3.1.16.** Let  $X, Y$  be metric spaces and  $T : X \rightarrow Y$ . Then  $T$  is said to be an **isometry** if for each  $x_1, x_2 \in X$ ,  $d(Tx_1, Tx_2) = d(x_1, x_2)$ .

**Exercise 3.1.17.** Let  $X, Y$  be metric spaces and  $T : X \rightarrow Y$  and isometry. Then  $T$  is injective.

*Proof.* Let  $x_1, x_2 \in X$ . Suppose that  $Tx_1 = Tx_2$ . Then  $0 = d(Tx_1, Tx_2) = d(x_1, x_2)$ . So  $x_1 = x_2$ . Hence  $T$  is injective. □

**Note 3.1.18.** Let  $X, Y$  be metric spaces and  $T : X \rightarrow Y$  an isometry. Then  $T$  is clearly continuous. If  $T$  is surjective, then  $T^{-1}$  is an isometry and therefore continuous. Hence  $T$  is a homeomorphism.

**Definition 3.1.19.** Let  $(M, d)$  be a metric space. Then  $(M, d)$  is said to be a **Polish space** if  $(M, d)$  is complete and separable.

**Exercise 3.1.20.** Let  $(X, d)$  be a compact metric space,  $E \subset X$  closed,  $U \subset X$  open. Suppose that  $E \subset U$ . Then there exists  $\delta > 0$  such that for each  $x \in E$ ,  $B(x, \delta) \subset U$ .

*Proof.* Since  $X$  is compact,  $E$  and  $U^c$  are compact. Then there exist  $x_0 \in E$  and  $y_0 \in U^c$  such that  $d(E, U^c) = d(x_0, y_0)$ . Since  $E \cap U^c = \emptyset$ ,  $x_0 \neq y_0$  and  $d(E, U^c) > 0$ . Put  $\epsilon = d(E, U^c)$

and  $\delta = \frac{\epsilon}{2}$ . Let  $x \in E$ ,  $w \in B(x, \delta)$  and  $y \in U^c$ . Then

$$\begin{aligned} d(y, w) &\geq d(y, x) - d(x, w) \\ &> \epsilon - \delta \\ &= \epsilon - \frac{\epsilon}{2} \\ &= \frac{\epsilon}{2} \\ &> 0 \end{aligned}$$

So  $y \neq w$ . Since  $y \in U^c$  and  $w \in B(x, \delta)$  are arbitrary,  $B(x, \delta) \subset U$ .  $\square$

**Definition 3.1.21.** Let  $S$  be a set,  $(M, d)$  a metric space and  $B(S, M) = \{f : S \rightarrow M : f \text{ is bounded}\}$ . We define the **supremum metric**, denoted  $d_u : B(S, M) \times B(S, M) \rightarrow [0, \infty)$ , by

$$d_u(f, g) = \sup_{x \in X} d(f(x), g(x))$$

**Exercise 3.1.22.** Let  $X$  be a set,  $(Y, d_Y)$ ,  $(Z, d_Z)$  metric spaces,  $(f_n)_{n \in \mathbb{N}} \subset B(X, Y)$ ,  $f \in B(X, Y)$  and  $g \in C(Y, Z)$ . Suppose that  $g$  is uniformly continuous. If  $f_n \xrightarrow{u} f$ , then  $g \circ f_n \xrightarrow{u} g \circ f$ .

*Proof.* Suppose that  $f_n \xrightarrow{u} f$ . Let  $\epsilon > 0$ . Uniform continuity of  $g$  implies that there exists  $\delta > 0$  such that for each  $y_1, y_2 \in Y$ ,  $d_Y(y_1, y_2) < \delta$  implies that  $d_Z(g(y_1), g(y_2)) < \epsilon/2$ . Uniform convergence implies that there exists  $N \in \mathbb{N}$  such that for each  $n \in \mathbb{N}$ ,  $n \geq N$  implies that  $d_u(f_n, f) < \delta/2$ . Let  $n \in \mathbb{N}$ . Suppose that  $n \geq N$ . Let  $x \in X$ . Then  $d_Y(f_n(x), f(x)) < \delta$ . This implies that  $d_Z(g(f_n(x)), g(f(x))) < \epsilon/2$ . Hence  $\sup_{x \in X} d_Z(g \circ f_n(x), g \circ f(x)) \leq \epsilon/2$ . Thus

$d_u(g \circ f_n, g \circ f) < \epsilon$ . So  $g \circ f_n \xrightarrow{u} g \circ f$ .  $\square$

**Definition 3.1.23.** Let  $(X, d)$  be a metric space. Define

- (1)  $\text{Aut}(X) = \{\sigma : X \rightarrow X : \sigma \text{ is a homeomorphism}\}$
- (2)  $\text{Aut}(X, d) = \{\sigma : X \rightarrow X : \sigma \text{ is an isometric isomorphism}\}$

**Exercise 3.1.24.** Let  $(X, d)$  be a compact metric space,  $E \subset X$  closed,  $U \subset X$  open. Suppose that  $E \subset U$ . Let  $(f_n)_{n \in \mathbb{N}} \in \text{Aut}(X)$ ,  $f \in \text{Aut}(X)$ . Suppose that  $f_n \xrightarrow{u} f$ . Then there exists  $N \in \mathbb{N}$  such that for each  $n \geq N$ ,  $f(E) \subset f_n(U)$ .

*Proof.* Since  $f$  is a homeomorphism,  $E$  is closed and  $U$  is open,  $f(E)$  is compact and  $f(U)$  is open and  $f(E) \subset f(U)$ . Then  $d(f(E), f(U^c)) > 0$ . Put  $\epsilon = d(f(E), f(U^c))$ . Choose  $\delta = \epsilon/2$ . Then there exists  $N \in \mathbb{N}$  such that for each  $n \in \mathbb{N}$ ,  $n \geq N$  implies that  $\sup_{z \in X} d(f(z), f_n(z)) < \delta$ .

Let  $n \geq N$ ,  $x \in E$  and  $w \in B(f(x), \delta)$ . For the sake of contradiction, suppose that  $w \in f_n(U^c)$ . Then there exist  $p \in U^c$  such that  $w = f_n(p)$ . Put  $z = f(p) \in f(U^c)$ . Then

$$\begin{aligned} \epsilon &\leq d(f(x), z) \\ &\leq d(f(x), w) + d(w, z) \\ &= d(f(x), w) + d(f_n(p), f(p)) \\ &< \delta + \delta \\ &= \epsilon \end{aligned}$$

which is a contradiction. So  $w \in f_n(U)$ . Hence  $B(f(x), \delta) \subset f_n(U)$   $\square$



### 3.2. Product Spaces.

## 4. TOPOLOGY

## 4.1. Introduction.

**Definition 4.1.1.** Let  $X$  be a set and  $\mathcal{T} \subset \mathcal{P}(X)$ . Then  $\mathcal{T}$  is said to be a **topology on  $X$**  if

- (1)  $X, \emptyset \in \mathcal{T}$
- (2) for each  $(U_\alpha)_{\alpha \in A} \subset \mathcal{T}$ ,

$$\bigcup_{\alpha \in A} U_\alpha \in \mathcal{T}$$

- (3) for each  $(U_j)_{j=1}^n \subset \mathcal{T}$ ,

$$\bigcap_{j=1}^n U_j \in \mathcal{T}$$

**Exercise 4.1.2.** Let  $X$  be a set and  $(\mathcal{T}_i)_{i \in I}$  a collection of topologies on  $X$ . Then  $\bigcap_{i \in I} \mathcal{T}_i$  is a topology on  $X$ .

*Proof.*

- (1) Since for each  $i \in I$ ,  $X, \emptyset \in \mathcal{T}_i$ , we have that  $X, \emptyset \in \bigcap_{i \in I} \mathcal{T}_i$ .
- (2) Let  $(U_\alpha)_{\alpha \in A} \subset \bigcap_{i \in I} \mathcal{T}_i$ . Then for each  $i \in I$ ,  $(U_\alpha)_{\alpha \in A} \subset \mathcal{T}_i$ . So for each  $i \in I$ ,  $\bigcup_{\alpha \in A} U_\alpha \in \mathcal{T}_i$ . Thus  $\bigcup_{\alpha \in A} U_\alpha \in \bigcap_{i \in I} \mathcal{T}_i$ .
- (3) Let  $(U_j)_{j=1}^n \subset \bigcap_{i \in I} \mathcal{T}_i$ . Then for each  $i \in I$ ,  $(U_j)_{j=1}^n \subset \mathcal{T}_i$ . So for each  $i \in I$ ,  $\bigcap_{j=1}^n U_j \in \mathcal{T}_i$ . Thus  $\bigcap_{j=1}^n U_j \in \bigcap_{i \in I} \mathcal{T}_i$ .

So  $\bigcap_{i \in I} \mathcal{T}_i$  is a topology on  $X$ . □

**Definition 4.1.3.** Let  $X$  be a set and  $\mathcal{E} \subset \mathcal{P}(X)$ . Set

$$\mathcal{S} = \{\mathcal{T} \subset \mathcal{P}(X) : \mathcal{T} \text{ is a topology on } X \text{ and } \mathcal{E} \subset \mathcal{T}\}$$

We define the **topology generated by  $\mathcal{E}$  on  $X$** , denoted  $\tau(\mathcal{E})$ , by

$$\tau(\mathcal{E}) = \bigcap_{\mathcal{T} \in \mathcal{S}} \mathcal{T}$$

**Definition 4.1.4.** Let  $(X, d)$  be a metric space. We define the **metric topology on  $X$** , denoted  $\mathcal{T}_d$ , by

$$\mathcal{T}_d = \tau(\{B(x, \delta) : x \in X, \delta > 0\})$$

**Definition 4.1.5.** Let  $X$  be a set and  $\mathcal{T} \subset \mathcal{P}(X)$  a topology on  $X$ ,  $x \in X$  and  $\mathcal{B}_x \subset \mathcal{T}$ . Then  $\mathcal{B}_x$  is said to be a **local basis for  $\mathcal{T}$  at  $x$**  if

- (1) for each  $V \in \mathcal{B}_x$ ,  $x \in V$
- (2) for each  $U \in \mathcal{T}$ , if  $x \in U$ , then there exists  $V \in \mathcal{B}_x$  such that  $V \subset U$

**Exercise 4.1.6.** Let  $(X, d)$  be a metric space and  $x \in X$ . Set  $\mathcal{B}_x = \{B(x, \delta) : \delta > 0\}$ . Then  $\mathcal{B}_x$  is a local basis for  $\mathcal{T}_d$  at  $x$ .

**FINISH!!! right now not well defined.**

*Proof.* Clear. □

**Definition 4.1.7.** Let  $X$  be a set and  $\mathcal{T} \subset \mathcal{P}(X)$  a topology on  $X$  and  $\mathcal{B} \subset \mathcal{T}$ . Then  $\mathcal{B}$  is said to be a **basis for  $\mathcal{T}$**  if for each  $x \in X$ , there exists  $\mathcal{B}_x \subset \mathcal{B}$  such that  $\mathcal{B}_x$  is a local basis for  $\mathcal{T}$  at  $x$ . **FINISH!!! Needs fixing, this definition is non standard.**

**Exercise 4.1.8.** Let  $X$  be a set and  $\mathcal{T} \subset \mathcal{P}(X)$  a topology on  $X$  and  $\mathcal{B} \subset \mathcal{T}$ . Then  $\mathcal{B}$  is a basis for  $\mathcal{T}$  iff for each  $U \in \mathcal{T}$ ,  $U \neq \emptyset$  implies that there exists a collection  $(U_\alpha)_{\alpha \in A} \subset \mathcal{B}$  such that

$$U = \bigcup_{\alpha \in A} U_\alpha$$

*Proof.* Suppose that  $\mathcal{B}$  is a basis for  $\mathcal{T}$ . Let  $U \in \mathcal{T}$ . Since  $\mathcal{B}$  is a basis for  $\mathcal{T}$ , for each  $x \in U$ , there exists  $\mathcal{B}_x \subset \mathcal{B} \subset \mathcal{T}$  such that  $\mathcal{B}_x$  is a local basis for  $\mathcal{T}$  at  $x$ . This implies that for each  $x \in U$ , there exists  $V_x \in \mathcal{B}_x$  such that  $x \in V_x \subset U$ . Suppose that  $U \neq \emptyset$ . Then  $(V_x)_{x \in U} \subset \mathcal{B}$  satisfies  $U = \bigcup_{x \in U} V_x$ .

Conversely, suppose that for each  $U \in \mathcal{T}$ ,  $U \neq \emptyset$  implies that there exists a collection  $(U_\alpha)_{\alpha \in A} \subset \mathcal{B}$  such that  $U = \bigcup_{\alpha \in A} U_\alpha$ . Let  $x \in X$ . Set  $\mathcal{B}_x = \{V \in \mathcal{B} : x \in V\}$ . Let  $U \in \mathcal{T}$ .

Suppose that  $x \in U$ . Then  $U \neq \emptyset$ . By assumption, there exists a collection  $(U_\alpha)_{\alpha \in A} \subset \mathcal{B}$  such that  $U = \bigcup_{\alpha \in A} U_\alpha$ . Since  $x \in U$ , there exists  $\alpha^* \in A$  such that  $x \in U_{\alpha^*}$ . By definition,

$U_{\alpha^*} \in \mathcal{B}_x$  and  $U_{\alpha^*} \subset U$ . Hence  $\mathcal{B}_x$  is a local basis for  $\mathcal{T}$  at  $x$ . Therefore for each  $x \in X$ , there exists  $\mathcal{B}_x \subset \mathcal{B}$  such that  $\mathcal{B}_x$  is a local basis for  $\mathcal{T}$  at  $x$ . Thus  $\mathcal{B}$  is a basis for  $\mathcal{T}$ . □

**Exercise 4.1.9.** Let  $X$  be a set and  $\mathcal{T}_1, \mathcal{T}_2 \subset \mathcal{P}(X)$  topologies on  $X$  and  $\mathcal{B} \subset \mathcal{T}_1$ . Suppose that  $\mathcal{T}_1 \subset \mathcal{T}_2$ . If  $\mathcal{B}$  is a basis for  $\mathcal{T}_2$ , then  $\mathcal{B}$  is a basis for  $\mathcal{T}_1$ .

*Proof.* Suppose that  $\mathcal{B}$  is a basis for  $\mathcal{T}_2$ . Let  $U \in \mathcal{T}_1$ . Then  $U \in \mathcal{T}_2$ . Suppose that  $U \neq \emptyset$ . Since  $\mathcal{B}$  is a basis for  $\mathcal{T}_2$ , the previous exercise implies that there exists a collection  $(U_\alpha)_{\alpha \in A} \subset \mathcal{B}$  such that  $U = \bigcup_{\alpha \in A} U_\alpha$ . Thus the previous exercise implies that  $\mathcal{B}$  is a basis for  $\mathcal{T}_1$ . □

**Exercise 4.1.10.** Let  $X$  be a set and  $\mathcal{B} \subset \mathcal{P}(X)$ . Then there exists a topology  $\mathcal{T}$  on  $X$  such that  $\mathcal{B}$  is a basis for  $\mathcal{T}$  iff

- (1) for each  $x \in X$ , there exists  $V \in \mathcal{B}$  such that  $x \in V$
- (2) for each  $U, V \in \mathcal{B}$ , if  $x \in U \cap V$ , then there exists  $W \in \mathcal{B}$  such that  $x \in W \subset U \cap V$

*Proof.* Suppose that there exists a topology  $\mathcal{T}$  on  $X$  such that  $\mathcal{B}$  is a basis for  $\mathcal{T}$ . Then conditions (1) and (2) are clear by Definition 4.1.5 and Definition 4.1.7.

Conversely, suppose that (1) and (2) are satisfied. Define  $\mathcal{T} \subset \mathcal{P}(X)$  by

$$\mathcal{T} = \{U \subset X : \text{for each } x \in U, \text{ there exists } V \in \mathcal{B} \text{ such that } x \in V \subset U\}$$

Trivially  $\emptyset \in \mathcal{T}$ . By condition (1),  $X \in \mathcal{T}$ . Let  $(U_\alpha)_{\alpha \in A} \subset \mathcal{T}$  and  $x \in \bigcup_{\alpha \in A} U_\alpha$ . Then there exists  $\alpha \in A$  such that  $x \in U_\alpha$ . Hence there exists  $V \in \mathcal{B}$  such that

$$\begin{aligned} x &\in V \\ &\subset U_\alpha \\ &\subset \bigcup_{\alpha \in A} U_\alpha \end{aligned}$$

So  $\bigcup_{\alpha \in A} U_\alpha \in \mathcal{T}$ . Let  $(U_j)_{j=1}^n \subset \mathcal{T}$  and  $x \in \bigcap_{j=1}^n U_j$ . Then in particular,  $U_1, U_2 \in \mathcal{T}$  and  $x \in U_1 \cap U_2$ . Then for  $j \in \{1, 2\}$ , there exists  $V_j \in \mathcal{B}$  such that  $x \in V_j \subset U_j$ . This implies that  $x \in V_1 \cap V_2$  and by condition (2), there exists  $W \in \mathcal{B}$  such that

$$\begin{aligned} x &\in W \\ &\subset V_1 \cap V_2 \\ &\subset U_1 \cap U_2 \end{aligned}$$

Therefore  $U_1 \cap U_2 \in \mathcal{T}$ . Proceeding inductively, we obtain that  $\bigcap_{j=1}^n U_j \in \mathcal{T}$ . □

**Exercise 4.1.11.** Let  $X$  be a set and  $\mathcal{E} \subset \mathcal{P}(X)$ . Define  $\mathcal{B} \subset \mathcal{P}(X)$  by

$$\mathcal{B} = \{X, \emptyset\} \cup \left\{ \bigcap_{j=1}^n V_j : (V_j)_{j=1}^n \subset \mathcal{E} \right\}$$

Then

- $\mathcal{B}$  is a basis for  $\tau(\mathcal{E})$
- 

$$\tau(\mathcal{E}) = \left\{ \bigcup_{\alpha \in A} V_\alpha : (V_\alpha)_{\alpha \in A} \subset \mathcal{B} \right\}$$

That is, each element of  $\tau(\mathcal{E})$  is either  $X, \emptyset$  or a union of finite intersections of elements of  $\mathcal{E}$ .

*Proof.*

- Referring to Exercise 4.1.10, since  $X \in \mathcal{B}$ , condition (1) is satisfied and since for each  $U, V \in \mathcal{B}$ ,  $U \cap V \in \mathcal{B}$ , condition (2) is satisfied. Hence there exists a topology  $\mathcal{T}$  on  $X$  such that  $\mathcal{B}$  is a basis for  $\mathcal{T}$ . Since  $\mathcal{E} \subset \mathcal{B} \subset \mathcal{T}$ , we have that  $\tau(\mathcal{E}) = \tau(\mathcal{B}) \subset \mathcal{T}$ . Then Exercise 4.1.9 implies that  $\mathcal{B}$  is a basis for  $\tau(\mathcal{E})$ .
- Exercise 4.1.8 implies that

$$\tau(\mathcal{E}) = \left\{ \bigcup_{\alpha \in A} V_\alpha : (V_\alpha)_{\alpha \in A} \subset \mathcal{B} \right\}$$

□

**Definition 4.1.12.** Let  $X$  be a set and  $\mathcal{T}$  a topology on  $X$ . Then  $(X, \mathcal{T})$  is said to be a **topological space**. Let  $U \subset X$ . Then  $U$  is said to be **open** if  $U \in \mathcal{T}$  and  $U$  is said to be **closed** if  $U^c$  is open. We define  $\mathcal{F}_T = \{C \subset X : C^c \in \mathcal{T}\}$ .

**Definition 4.1.13.** Let  $X$  be a topological space and  $S, N \subset X$ . Then  $N$  is said to be a **neighborhood** of  $S$  if there exists  $U \subset X$  such that  $U$  is open and  $S \subset U \subset N$ . For  $S \in X$ , we denote the set of neighborhoods of  $S$  by  $\mathcal{N}_S$ .

**Definition 4.1.14.** Let  $X$  be a topological space and  $A \subset X$ . Set  $\mathcal{U}_A = \{U \subset X : U \subset A \text{ and } U \text{ is open}\}$  and  $\mathcal{C}_A = \{U \subset X : A \subset U \text{ and } U \text{ is closed}\}$ .

We define the **interior of A**, denoted  $A^\circ$ , by

$$A^\circ = \bigcup_{U \in \mathcal{U}_A} U$$

We define the **closure of  $A$** , denoted  $\overline{A}$ , by

$$\overline{A} = \bigcap_{U \in \mathcal{C}_A} U$$

**Definition 4.1.15.** Let  $X$  be a topological space and  $A \subset X$ . Then

- (1)  $A$  is open iff  $A = A^\circ$
- (2)  $A$  is closed iff  $A = \overline{A}$

*Proof.* Clear. □

**Exercise 4.1.16.** Let  $X$  be a topological space and  $A \subset X$ . Then  $(A^\circ)^c = \overline{A^c}$ .

*Proof.* □

**Exercise 4.1.17.** Let  $X$  be a topological space,  $A \subset X$  and  $x \in X$ . Then  $A \in \mathcal{N}_x$  iff  $x \in A^\circ$ .

*Proof.* Suppose that  $A \in \mathcal{N}_x$ . Then there exists  $U \subset X$  such that  $U$  is open and  $x \in U \subset A$ . By definition,  $U \subset A^\circ$ . Conversely, suppose that  $x \in A^\circ$ . Then by definition,  $A^\circ \in \mathcal{N}_x$ . □

**Exercise 4.1.18.** Let  $X$  be a topological space and  $A \subset X$ . Then  $A$  is open iff for each  $x \in A$ , there exists  $U \in \mathcal{N}_x$  such that  $U$  is open and  $U \subset A$ .

*Proof.* Suppose that  $A$  is open. Let  $x \in A$ . Then  $A \in \mathcal{N}_x$ ,  $A$  is open and  $A \subset A$ . Conversely, suppose that for each  $x \in A$ , there exists  $U_x \in \mathcal{N}_x$  such that  $U_x$  is open and  $U_x \subset A$ . Then

$$A = \bigcup_{x \in A} U_x$$

is open. □

**Definition 4.1.19.** Let  $X$  be a topological space,  $A \subset X$  and  $x \in X$ . Then  $x$  is said to be a **limit point of  $A$**  if for each  $U \in \mathcal{N}_x$ ,

$$A \cap (U \setminus \{x\}) \neq \emptyset$$

We define  $A' = \{x \in A : x \text{ is a limit point of } A\}$ .

**Exercise 4.1.20.** Let  $X$  be a topological space and  $A \subset X$ . Then  $\overline{A} = A \cup A'$ .

*Proof.* Let  $x \in A'$ . For the sake of contradiction, suppose that  $x \notin \overline{A}$ . Then there exists  $C \subset X$  such that  $C$  is closed,  $A \subset C$  and  $x \notin C$ . Hence  $x \in C^c \subset A^c$ . Since  $C^c$  is open,  $x \in (A^c)^\circ$ . Since  $x \in A'$  and  $(A^c)^\circ \in \mathcal{N}_x$ ,  $[(A^c)^\circ \setminus \{x\}] \cap A \neq \emptyset$ . This is a contradiction since  $(A^c)^\circ \setminus \{x\} \subset A^c$ . So  $x \notin \overline{A}$  and  $A' \subset \overline{A}$ . Since  $A \subset \overline{A}$ , we have that  $A \cup A' \subset \overline{A}$ .

Conversely, let  $x \in \overline{A}$ . For the sake of contradiction, suppose that  $x \notin A \cup A'$ . Then  $x \in A^c \cap (A')^c$ . Since  $x \in (A')^c$ , there exists  $U \in \mathcal{N}_x$  such that  $(U \setminus \{x\}) \cap A = \emptyset$ . Hence  $U \setminus \{x\} \subset A^c$ . Since  $x \in A^c$ ,

$$\begin{aligned} x &\in U^\circ \\ &\subset U \\ &= (U \setminus \{x\}) \cup \{x\} \\ &\subset A^c \end{aligned}$$

Since  $A \subset (U^\circ)^c$  which is closed,  $x \in \overline{A}$  implies that  $x \in (U^\circ)^c$  which is a contradiction. So  $x \in A \cup A'$  and  $\overline{A} \subset A \cup A'$ . Therefore  $\overline{A} = A \cup A'$ . □

## 4.2. Continuous Maps.

**Definition 4.2.1.** Let  $(X, \mathcal{A})$  and  $(Y, \mathcal{B})$  be topological spaces and  $f : X \rightarrow Y$ . Then  $f$  is said to be **continuous** if for each  $B \in \mathcal{B}$ ,  $f^{-1}(B) \in \mathcal{A}$ .

**Definition 4.2.2.** Let  $(X, \mathcal{A})$  and  $(Y, \mathcal{B})$  be topological spaces,  $f : X \rightarrow Y$  and  $x \in X$ . Then  $f$  is said to be **continuous at  $x$**  if for each  $V \in \mathcal{N}_{f(x)}$ , there exists  $U \in \mathcal{N}_x$  such that  $f(U) \subset V$ .

**Exercise 4.2.3.** Let  $(X, \mathcal{A})$  and  $(Y, \mathcal{B})$  be topological spaces,  $f : X \rightarrow Y$  and  $x \in X$ . Then  $f$  is continuous at  $x$  iff for each  $V \in \mathcal{N}_{f(x)}$ ,  $f^{-1}(V) \in \mathcal{N}_x$ .

**Hint:** for  $U \in \mathcal{N}_x$  and  $V \in \mathcal{N}_{f(x)}$ , consider  $f^{-1}(f(U))$  and  $f(f^{-1}(V))$

*Proof.* Suppose that  $f$  is continuous at  $x$ . Let  $V \in \mathcal{N}_{f(x)}$ . Then there exists  $U \in \mathcal{N}_x$  such that  $f(U) \subset V$ . Thus

$$\begin{aligned} x &\in U^\circ \\ &\subset U \\ &\subset f^{-1}(f(U)) \\ &\subset f^{-1}(V) \end{aligned}$$

So  $f^{-1}(V) \in \mathcal{N}_x$ .

Conversely, suppose that for each  $V \in \mathcal{N}_{f(x)}$ ,  $f^{-1}(V) \in \mathcal{N}_x$ . Let  $V \in \mathcal{N}_{f(x)}$ . Hence  $f^{-1}(V) \in \mathcal{N}_x$ . Set  $U = f^{-1}(V)$ . Then

$$\begin{aligned} f(U) &= f(f^{-1}(V)) \\ &\subset V \end{aligned}$$

Thus  $f$  is continuous at  $x$ . □

**Exercise 4.2.4.** Let  $(X, \mathcal{A})$  and  $(Y, \mathcal{B})$  be topological spaces and  $f : X \rightarrow Y$ . Then  $f$  is continuous iff for each  $x \in X$ ,  $f$  is continuous at  $x$ .

*Proof.* Suppose that  $f$  is continuous. Let  $x \in X$ . Let  $V \in \mathcal{N}_{f(x)}$ . Then  $V^\circ \in \mathcal{B}$  and  $f(x) \in V^\circ$ . Set  $U = f^{-1}(V^\circ)$ . By continuity,  $U \in \mathcal{A}$  and by construction,  $x \in U$ . Hence  $U \in \mathcal{N}_x$ . Then

$$\begin{aligned} f(U) &= f(f^{-1}(V^\circ)) \\ &\subset V^\circ \\ &\subset V \end{aligned}$$

So  $f$  is continuous at  $x$ .

Conversely, suppose that for each  $x \in X$ ,  $f$  is continuous at  $x$ . Let  $B \in \mathcal{B}$ . Let  $x \in f^{-1}(B)$ . Then  $B \in \mathcal{N}_{f(x)}$ . Continuity at  $x$  implies that  $f^{-1}(B) \in \mathcal{N}_x$ . Then  $x \in (f^{-1}(B))^\circ$ . Since  $x \in f^{-1}(B)$  is arbitrary,  $f^{-1}(B) \subset (f^{-1}(B))^\circ$ . Hence  $f^{-1}(B) = (f^{-1}(B))^\circ$  which implies that  $f^{-1}(B) \in \mathcal{A}$ . So  $f$  is continuous. □

**Definition 4.2.5.** Let  $(X, \mathcal{A})$  and  $(Y, \mathcal{B})$  be topological spaces and  $f : X \rightarrow Y$ . We define the

(1) **push-forward of  $\mathcal{A}$** , denoted  $f_*\mathcal{A}$ , by

$$f_*\mathcal{A} = \{B \subset Y : f^{-1}(B) \in \mathcal{A}\}$$

(2) **pull-back of  $\mathcal{B}$** , denoted  $f^*\mathcal{B}$ , by

$$f^*\mathcal{B} = \{f^{-1}(B) : B \in \mathcal{B}\}$$

**Exercise 4.2.6.** Let  $(X, \mathcal{A})$  and  $(Y, \mathcal{B})$  be topological spaces and  $f : X \rightarrow Y$ . Then

- (1)  $f_*\mathcal{A}$  is a topology on  $Y$
- (2)  $f^*\mathcal{B}$  is a topology on  $X$

*Proof.*

- (1) • Since  $f^{-1}(Y) = X \in \mathcal{A}$  and  $f^{-1}(\emptyset) = \emptyset \in \mathcal{A}$ ,  $Y, \emptyset \in f_*\mathcal{A}$ .
- Let  $(U_\alpha)_{\alpha \in A} \subset f_*\mathcal{A}$ . Then for each  $\alpha \in A$ ,  $f^{-1}(U_\alpha) \in \mathcal{A}$ . This implies that

$$f^{-1}\left(\bigcup_{\alpha \in A} U_\alpha\right) = \bigcup_{\alpha \in A} f^{-1}(U_\alpha) \in \mathcal{A}$$

Hence  $\bigcup_{\alpha \in A} U_\alpha \in f_*\mathcal{A}$ .

- Let  $(U_j)_{j=1}^n \subset f_*\mathcal{A}$ . Then for each  $j \in 1, \dots, n$ ,  $f^{-1}(U_j) \in \mathcal{A}$ . This implies that

$$f^{-1}\left(\bigcap_{j=1}^n U_j\right) = \bigcap_{j=1}^n f^{-1}(U_j) \in \mathcal{A}$$

Hence  $\bigcap_{j=1}^n U_j \in f_*\mathcal{A}$ .

So  $f_*\mathcal{A}$  is a topology on  $Y$ .

(2) Similar to (1).

□

**Exercise 4.2.7.** Let  $(X, \mathcal{A})$  and  $(Y, \mathcal{B})$  be topological spaces,  $f : X \rightarrow Y$  and  $\mathcal{E} \subset \mathcal{P}(Y)$ . Suppose that  $\mathcal{B} = \tau(\mathcal{E})$ . Then  $f$  is continuous iff for each  $B \in \mathcal{E}$ ,  $f^{-1}(B) \in \mathcal{A}$ .

*Proof.* Suppose that  $f$  is continuous. Since  $\mathcal{E} \subset \mathcal{B}$ , clearly for each  $B \in \mathcal{E}$ ,  $f^{-1}(B) \in \mathcal{A}$ . Conversely, suppose that for each  $B \in \mathcal{E}$ ,  $f^{-1}(B) \in \mathcal{A}$ . Then  $\mathcal{E} \subset f_*\mathcal{A}$ . Since  $f_*\mathcal{A}$  is a topology on  $Y$ , we have that  $\mathcal{B} = \tau(\mathcal{E}) \subset f_*\mathcal{A}$ . So  $f$  is continuous. □

**Definition 4.2.8.** Let  $X$  be a set,  $(Y_\alpha, \mathcal{B}_\alpha)_{\alpha \in A}$  a collection of topological spaces and  $\mathcal{F} \in \prod_{\alpha \in A} Y_\alpha^X$  (i.e.  $\mathcal{F} = (f_\alpha)_{\alpha \in A}$  where for each  $\alpha \in A$ ,  $f_\alpha : X \rightarrow Y_\alpha$ ). We define the **initial topology generated by  $\mathcal{F}$**  on  $X$ , denoted  $\tau_X(\mathcal{F})$ , by

$$\tau_X(\mathcal{F}) = \tau(\{f_\alpha^{-1}(B) : B \in \mathcal{B}_\alpha \text{ and } \alpha \in A\})$$

**Note 4.2.9.** The initial topology topology generated by  $\mathcal{F}$  is also called the **weak topology generated by  $\mathcal{F}$**  and if  $\mathcal{F} = \{f\}$ , then  $\tau_X(\mathcal{F}) = f^*\mathcal{B}$ .

**Note 4.2.10.** Essentially,  $\tau_X(\mathcal{F})$  is the smallest topology on  $X$  such that for each  $\alpha \in A$ ,  $f_\alpha : X \rightarrow Y_\alpha$  is continuous.

**Definition 4.2.11.** Let  $(X_\alpha, \mathcal{A}_\alpha)_{\alpha \in A}$  be a collection of topological spaces,  $Y$  a set and  $\mathcal{F} \in \prod_{\alpha \in A} Y^{X_\alpha}$  (i.e.  $\mathcal{F} = (f_\alpha)_{\alpha \in A}$  where for each  $\alpha \in A$ ,  $f_\alpha : X_\alpha \rightarrow Y$ ). We define the **final topology generated by  $\mathcal{F}$**  on  $X$ , denoted  $\tau_Y(\mathcal{F})$ , by

$$\tau_Y(\mathcal{F}) = \tau(\{V \subset Y : \text{for each } \alpha \in A, f_\alpha^{-1}(V) \in \mathcal{A}_\alpha\})$$

**Note 4.2.12.** If  $\mathcal{F} = \{f\}$ , then  $\tau_Y(\mathcal{F}) = f_*\mathcal{A}$ .

**Note 4.2.13.** Essentially,  $\tau_X(\mathcal{F})$  is the largest topology on  $X$  such that for each  $\alpha \in A$ ,  $f_\alpha : X_\alpha \rightarrow Y$  is continuous.

**Exercise 4.2.14.** Let  $(X_\alpha, \mathcal{A}_\alpha)_{\alpha \in A}$  be a collection of topological spaces,  $Y$  a set,  $(Z, \mathcal{C})$  a topological space,  $\mathcal{F} = (f_\alpha)_{\alpha \in A} \in \prod_{\alpha \in A} Y^{X_\alpha}$  and  $g : Y \rightarrow Z$ . Then  $g$  is  $\tau_Y(\mathcal{F})$ - $\mathcal{C}$  continuous iff for each  $\alpha \in A$ ,  $g \circ f_\alpha$  is  $X_\alpha$ - $\mathcal{C}$  continuous, i.e. for each  $\alpha \in A$ , the following diagram commutes in the category of topological spaces:

$$\begin{array}{ccc} X_\alpha & \xrightarrow{f_\alpha} & Y \\ & \searrow g \circ f_\alpha & \downarrow g \\ & & Z \end{array}$$

*Proof.* If  $g$  is  $\tau_Y(\mathcal{F})$ - $\mathcal{C}$  continuous, then clearly for each  $\alpha \in A$ ,  $g \circ f_\alpha$  is  $X_\alpha$ - $\mathcal{C}$  continuous. Conversely, suppose that for each  $\alpha \in A$ ,  $g \circ f_\alpha$  is  $X_\alpha$ - $\mathcal{C}$  continuous. Let  $V \in \mathcal{C}$ . Continuity implies that for each  $\alpha \in A$ ,  $f_\alpha^{-1}(g^{-1}(V)) \in \mathcal{A}_\alpha$ . By definition,  $g^{-1}(V) \in \tau_Y(\mathcal{F})$ . So  $g$  is  $\tau_Y(\mathcal{F})$ - $\mathcal{C}$  continuous.  $\square$

**Definition 4.2.15.** Let  $(X, \mathcal{A})$  and  $(Y, \mathcal{B})$  be topological spaces and  $f : X \rightarrow Y$ . Then

- (1)  $f$  is said to be **open** if for each  $A \in \mathcal{A}$ ,  $f(A) \in \mathcal{B}$ .
- (2)  $f$  is said to be **closed** if for each  $A \subset X$ , if  $A^c \in \mathcal{A}$ , then  $f(A)^c \in \mathcal{B}$ .

**Exercise 4.2.16.** Let  $(X, \mathcal{T}), (Y, \mathcal{S})$  be topological spaces,  $\mathcal{B} \subset \mathcal{T}$  a basis for  $\mathcal{T}$  and  $f : X \rightarrow Y$ . If for each  $U \in \mathcal{B}$ ,  $f(U) \in \mathcal{S}$ , then  $f$  is open.

**Hint:**  $f\left(\bigcup_{\alpha \in A} A_\alpha\right) = \bigcup_{\alpha \in A} f(A_\alpha)$ .

*Proof.* Suppose that for each  $U \in \mathcal{B}$ ,  $f(U) \in \mathcal{S}$ . Let  $U \in \mathcal{T}$ . Then there exists  $(U_\alpha)_{\alpha \in A} \subset \mathcal{B}$  such that  $U = \bigcup_{\alpha \in A} U_\alpha$ . Then

$$\begin{aligned} f(U) &= \bigcup_{\alpha \in A} f(U_\alpha) \\ &\in \mathcal{S} \end{aligned}$$

Since  $U \in \mathcal{T}$  is arbitrary,  $f$  is open.  $\square$

**Exercise 4.2.17. Doob-Dynkin Lemma:**

Let  $(X_1, \mathcal{T}_1), (X_2, \mathcal{T}_2)$  and  $(X_3, \mathcal{T}_3)$  be topological spaces and  $f : X_1 \rightarrow X_2$  and  $g : X_1 \rightarrow X_3$ . Suppose that  $f$  is surjective and  $\mathcal{T}_1$ - $\mathcal{T}_2$  continuous and  $g$  is  $\mathcal{T}_1$ - $\mathcal{T}_3$  continuous and  $(X_3, \mathcal{T}_3)$  is a  $\mathcal{T}_1$  space. Then  $g$  is  $f^*\mathcal{T}_2$ - $\mathcal{T}_3$  continuous iff there exists a unique  $\phi : X_2 \rightarrow X_3$  such that  $\phi$  is  $\mathcal{T}_2$ - $\mathcal{T}_3$  continuous and  $g = \phi \circ f$ .

**Hint:** For each  $t \in X_3$ , set  $A_t = g^{-1}(\{t\}) \in \mathcal{F}_{(f^*\mathcal{T}_2)}$  and choose  $B_t \in \mathcal{T}_2$  such that  $A_t = f^{-1}(B_t)$ . Set  $\phi(y) = t$  for  $y \in B_t \cap f(X_1)$  and  $t \in g(X_1)$ .



*Proof.* Suppose that there exists a unique  $\phi : X_2 \rightarrow X_3$  such that  $\phi$  is  $\mathcal{T}_2$  -  $\mathcal{T}_3$  measurable and  $g = \phi \circ f$ . Since  $f$  is  $f^*\mathcal{T}_2$  -  $\mathcal{T}_2$  continuous, we have that  $g = \phi \circ f$  is  $f^*\mathcal{T}_2$ - $\mathcal{T}_3$  continuous. Conversely, suppose that  $g$  is  $f^*\mathcal{T}_2$ - $\mathcal{T}_3$  continuous.

• **(Existence)**

For each  $t \in X_3$ , set  $A_t = g^{-1}(\{t\})$ . Since  $(X_3, \mathcal{T}_3)$  is a  $T_1$  space, for each  $t \in X_3$ ,  $A_t \in \mathcal{F}_{f^*\mathcal{T}_2}$  and thus, there exists  $B_t \in \mathcal{F}_{\mathcal{T}_2}$  such that  $A_t = f^{-1}(B_t)$ .

Note that

– for each  $t \in g(X_1)$ , there exists  $x \in A_t$  such that  $g(x) = t$ . Hence  $f(x) \in B_t$ .

– for  $t_1, t_2 \in g(X_1)$ ,  $t_1 \neq t_2$  implies that

$$\begin{aligned} f^{-1}(B_{t_1} \cap B_{t_2}) &= A_{t_1} \cap A_{t_2} \\ &= g^{-1}(\{t_1\} \cap \{t_2\}) \\ &= \emptyset \end{aligned}$$

and since  $f$  is surjective,

$$\begin{aligned} B_{t_1} \cap B_{t_2} &= f(f^{-1}(B_{t_1} \cap B_{t_2})) \\ &= f(\emptyset) \\ &= \emptyset \end{aligned}$$

– we have that

$$\begin{aligned} f^{-1}\left(\bigcup_{t \in g(X_1)} B_t\right) &= \bigcup_{t \in g(X_1)} A_t \\ &= \bigcup_{t \in g(X_1)} g^{-1}(\{t\}) \\ &= g^{-1}(g(X_1)) \\ &= X_1 \end{aligned}$$

Since  $f$  is surjective, we have that

$$\begin{aligned} X_2 &= f(X_1) \\ &= f\left(f^{-1}\left(\bigcup_{t \in g(X_1)} B_t\right)\right) \\ &= \bigcup_{t \in g(X_1)} B_t \end{aligned}$$

Therefore,

- for each  $t \in g(X_1)$ ,  $B_t \neq \emptyset$
- $(A_t)_{t \in g(X_1)}$  is a partition of  $X_1$
- $(B_t)_{t \in g(X_1)}$  is a partition of  $X_2$

Define  $\phi : X_2 \rightarrow X_3$  by  $\phi(y) = t$  for  $t \in g(X_1)$  and  $y \in B_t$ . Then the previous observations imply that  $\phi$  is well defined and  $\phi(X_2) = g(X_1)$ . Since for each  $t \in g(X_1)$  and  $x \in A_t$ ,  $f(x) \in B_t$  and  $g(x) = t$ , we have that  $\phi \circ f(x) = t = g(x)$ . So  $\phi \circ f = g$ .

To show that  $\phi$  is continuous, let  $C \in \mathcal{T}_3$ . Choose  $B \in \mathcal{T}_2$  such that  $g^{-1}(C) = f^{-1}(B)$ . Let  $y \in \phi^{-1}(C) \subset X_2$ . Set  $t = \phi(y) \in C$  and choose  $x \in X_1$  such that  $y = f(x)$ . Since

$$\begin{aligned} g(x) &= \phi \circ f(x) \\ &= \phi(y) \\ &= t \\ &\in C \end{aligned}$$

$x \in g^{-1}(C) = f^{-1}(B)$ . Therefore,  $y = f(x) \in B$ . So  $\phi^{-1}(C) \subset B$ . Let  $y \in B$ . Choose  $x \in X_1$  such that  $f(x) = y$ . Then  $x \in f^{-1}(B) = g^{-1}(C)$ . So

$$\begin{aligned} \phi(y) &= \phi \circ f(x) \\ &= g(x) \\ &\in C \end{aligned}$$

and  $y \in \phi^{-1}(C)$ . So  $B \subset \phi^{-1}(C)$ . Hence  $\phi^{-1}(C) = B \in \mathcal{T}_2$  and  $\phi$  is  $\mathcal{T}_2$ - $\mathcal{T}_3$  continuous.

• **(Uniqueness)**

Let  $\psi : X_2 \rightarrow X_3$ . Suppose that  $\psi$  is  $\mathcal{T}_2$ - $\mathcal{T}_3$  continuous and  $g = \psi \circ f$ . Let  $y \in X_2$ . Then there exists  $x \in X_1$  such that  $y = f(x)$ . Then

$$\begin{aligned} \psi(y) &= \psi \circ f(x) \\ &= g(x) \\ &= \phi \circ f(x) \\ &= \phi(y) \end{aligned}$$

So  $\psi = \phi$ .

□

**Exercise 4.2.18.** Let  $(X_1, \mathcal{T}_1)$ ,  $(X_2, \mathcal{T}_2)$  and  $(X_3, \mathcal{T}_3)$  be topological spaces and  $f : X_1 \rightarrow X_2$  and  $g : X_1 \rightarrow X_3$ . Suppose that  $f$  is  $\mathcal{T}_1$ - $\mathcal{T}_2$  continuous and  $g$  is  $\mathcal{T}_1$ - $\mathcal{T}_3$  continuous and  $(X_3, \mathcal{T}_3)$  is a  $T_1$  space. Then  $g$  is  $f^*\mathcal{T}_2$ - $\mathcal{T}_3$  continuous iff there exists a unique  $\phi : f(X_1) \rightarrow X_3$  such that  $\phi$  is  $\mathcal{T}_2 \cap f(X_1) - \mathcal{T}_3$  continuous and  $g = \phi \circ f$ .

*Proof.* A previous exercise implies that  $f : X_1 \rightarrow f(X_1)$  is  $\mathcal{T}_1 - \mathcal{T}_2 \cap f(X_1)$  continuous. Now apply the previous exercise. □

**Definition 4.2.19.** Let  $X$  be a topological space,  $x_0 \in X$  and  $f : X \rightarrow \mathbb{R}$ . We define the **limit inferior of  $f$  as  $x \rightarrow x_0$  (resp. limit inferior of  $f$  as  $x \rightarrow x_0$ )**, denoted  $\liminf_{x \rightarrow x_0} f(x)$  (resp.  $\liminf_{x \rightarrow x_0} f(x)$ ), by

$$\liminf_{x \rightarrow x_0} f(x) = \sup_{V \in \mathcal{N}_{x_0}} \inf_{x \in V \setminus \{x_0\}} f(x)$$

resp.

$$\limsup_{x \rightarrow x_0} f(x) = \inf_{V \in \mathcal{N}_{x_0}} \sup_{x \in V \setminus \{x_0\}} f(x)$$

**Exercise 4.2.20.** Let  $X$  be a topological space,  $x_0 \in X$  and  $f : X \rightarrow \mathbb{R}$ . Then  $f$  is continuous at  $x_0$  iff  $\liminf_{x \rightarrow x_0} f(x) = \limsup_{x \rightarrow x_0} f(x) = f(x_0)$

*Proof.* Suppose that

**FINISH!!!**

□

### 4.3. Nets.

**Definition 4.3.1.** Let  $A$  be a set and  $\leq$  a relation on  $A$ . Then  $(A, \leq)$  is said to be a **directed set** if ,

- (1) for each  $\alpha \in A$ ,  $\alpha \leq \alpha$
- (2) for each  $\alpha, \beta, \gamma \in A$ ,  $\alpha \leq \beta$  and  $\beta \leq \gamma$  implies that  $\alpha \leq \gamma$
- (3) for each  $\alpha, \beta \in A$ , there exists  $\gamma \in A$  such that  $\alpha, \beta \leq \gamma$
- (4)  $A \neq \emptyset$

**Definition 4.3.2.** Let  $X$  be a set. Define the **reverse inclusion ordering** on  $\mathcal{N}_x$ , denoted  $\leq$ , by  $U \leq V$  iff  $V \subset U$ .

**Exercise 4.3.3.** Let  $X$  be a topological space and  $x \in X$ . Then  $\mathcal{N}_x$  ordered by reverse inclusion is a directed set.

*Proof.*

- (1) Clearly, for each  $U \in \mathcal{N}_x$ ,  $U \leq U$ .
- (2) Let  $U, V, W \in \mathcal{N}_x$ . Suppose that  $U \leq V$  and  $V \leq W$ . Then  $W \subset V \subset U$  which implies that  $W \subset U$  and hence  $U \leq W$ .
- (3) Let  $U, V \in \mathcal{N}_x$ . Set  $W = U \cap V$ . Then  $W \in \mathcal{N}_x$  and  $U, V \leq W$ .

So  $\mathcal{N}_x$  is a directed set. □

**Definition 4.3.4.** Let  $X$  be a metric space and  $x_0 \in X$ . Define the **reverse distance from  $x_0$  ordering** on  $X \setminus \{x_0\}$ , denoted  $\leq_{x_0}$ , by  $x \leq_{x_0} y$  iff  $d(x, x_0) \geq d(y, x_0)$ .

**Exercise 4.3.5.** Let  $X$  be a metric space and  $x_0 \in X$ . Then  $(X \setminus \{x_0\}, \leq_{x_0})$  is a directed set.

*Proof.*

- (1) Let  $x \in X \setminus \{x_0\}$ . Since  $d(x, x_0) \geq d(x, x_0)$ ,  $x \leq_{x_0} x$ .
- (2) Let  $x, y, z \in X \setminus \{x_0\}$ . Suppose that  $x \leq_{x_0} y$  and  $y \leq_{x_0} z$ . Then  $d(x, x_0) \geq d(y, x_0)$  and  $d(y, x_0) \geq d(z, x_0)$ . Hence  $d(x, x_0) \geq d(z, x_0)$  so that  $x \leq_{x_0} z$ .
- (3) Let  $x, y \in X \setminus \{x_0\}$ . Set

$$\begin{aligned} z &= \arg \min_{a \in \{x, y\}} d(a, x_0) \\ &\in X \setminus \{x_0\} \end{aligned}$$

Then  $x, y \leq_{x_0} z$ . □

**Definition 4.3.6.** Let  $(A, \leq_A)$  and  $(B, \leq_B)$  be directed sets. We define the **product directed set of  $(A, \leq_A)$  and  $(B, \leq_B)$** , denoted  $(A \times B, \leq)$ , by

$$(a_1, b_1) \leq (a_2, b_2) \text{ iff } a_1 \leq_A a_2 \text{ and } b_1 \leq_B b_2$$

**Exercise 4.3.7.** Let  $(A, \leq_A)$  and  $(B, \leq_B)$  be directed sets. Then the product directed set of  $(A, \leq_A)$  and  $(B, \leq_B)$  is a directed set.

*Proof.*

- (1) Let  $(a, b) \in A \times B$ . Then  $a \leq_A a$  and  $b \leq_B b$ . So  $(a, b) \leq (a, b)$ .

- (2) Let  $(a_1, b_1), (a_2, b_2), (a_3, b_3) \in A \times B$ . Suppose that  $(a_1, b_1) \leq (a_2, b_2)$  and  $(a_2, b_2) \leq (a_3, b_3)$ . Then  $a_1 \leq_A a_2$ ,  $a_2 \leq_A a_3$ ,  $b_1 \leq_B b_2$  and  $b_2 \leq_B b_3$ . Therefore  $a_1 \leq_A a_3$  and  $b_1 \leq_B b_3$ . Hence  $(a_1, b_1) \leq (a_3, b_3)$ .
- (3) Let  $(a_1, b_1), (a_2, b_2) \in A \times B$ . Then there exist  $a \in A$  and  $b \in B$  such that  $a_1, a_2 \leq_A a$  and  $b_1, b_2 \leq_B b$ . Hence  $(a_1, b_1), (a_2, b_2) \leq (a, b)$ .

So  $(A \times B, \leq)$  is directed.  $\square$

**Definition 4.3.8.** Let  $X$  be a topological space,  $A$  a directed set and  $x : A \rightarrow Y$ . Then  $x$  is said to be a **net** in  $X$ . We typically write  $(x_\alpha)_{\alpha \in A}$ .

**Definition 4.3.9.** Let  $X$  be a topological space,  $(x_\alpha)_{\alpha \in A} \subset X$  a net and  $U \subset X$ . Then  $(x_\alpha)_{\alpha \in A}$  is said to be

- **eventually in  $U$**  if there exists  $\beta \in A$  such that for each  $\alpha \in A$   $\alpha \geq \beta$  implies that  $x_\alpha \in U$
- **frequently in  $U$**  if for each  $\alpha \in A$ , there exists  $\beta \in A$  such that  $\beta \geq \alpha$  and  $x_\beta \in U$

**Definition 4.3.10.** Let  $X$  be a topological space,  $(x_\alpha)_{\alpha \in A} \subset X$  a net and  $x \in X$ . Then  $(x_\alpha)_{\alpha \in A}$  is said to **converge to  $x$** , denoted  $x_\alpha \rightarrow x$ , if for each  $U \in \mathcal{N}_x$ ,  $(x_\alpha)_{\alpha \in A}$  is eventually in  $U$ .

**Definition 4.3.11.** Let  $X$  be a topological space and  $(x_\alpha)_{\alpha \in A} \subset X$  a net. Then  $(x_\alpha)_{\alpha \in A}$  is said to **converge** if there exists  $x \in X$  such that  $x_\alpha \rightarrow x$ .

**Exercise 4.3.12.** Let  $X$  be a metric space and  $x_0 \in X$ . Set  $A = X \setminus \{x_0\}$ . Order  $A$  by reverse distance from  $x_0$ . Define  $(x_\alpha)_{\alpha \in A} \subset X$  by  $x_\alpha = \alpha$ . Then  $x_\alpha \rightarrow x_0$ .

*Proof.* Let  $U \in \mathcal{N}_{x_0}$ . Since  $x_0 \in \text{Int } U$ , there exists  $\delta > 0$  such that  $B(x_0, \delta) \subset \text{Int } U$ . Choose  $\beta \in B^*(x_0, \delta)$ . Let  $\alpha \in A$ . Suppose that  $\alpha \geq \beta$ . Then  $d(\alpha, x_0) \leq d(\beta, x_0) < \delta$ . Hence

$$\begin{aligned} x_\alpha &= \alpha \\ &\in B^*(x_0, \delta) \\ &\subset U \end{aligned}$$

Since  $U \in \mathcal{N}_{x_0}$  is arbitrary,  $x_\alpha \rightarrow x_0$ .  $\square$

**Exercise 4.3.13.** Let  $X$  be a topological space,  $S \subset X$  and  $x \in X$ . Then  $x \in S'$  iff there exists a net  $(x_\alpha)_{\alpha \in A} \subset S \setminus \{x\}$  such that  $x_\alpha \rightarrow x$ .

*Proof.* Suppose that  $x \in S'$ . Set  $A = \mathcal{N}_x$ , ordered by reverse inclusion. Since  $x \in S'$ , for each  $\alpha \in A$ , there exists  $x_\alpha \in (\alpha \setminus \{x\}) \cap S$ . Then  $(x_\alpha)_{\alpha \in A} \subset S \setminus \{x\}$ . Let  $V \in \mathcal{N}_x$ . Choose  $\beta = V$ . Let  $\alpha \in \mathcal{N}_x$ . Suppose that  $\alpha \geq \beta$ . Then

$$\begin{aligned} x_\alpha &\in (\alpha \setminus \{x\}) \cap S \\ &\subset \alpha \\ &\subset \beta \\ &= V \end{aligned}$$

So  $(x_\alpha)_{\alpha \in \mathcal{N}_x}$  is eventually in  $V$ . Since  $V \in \mathcal{N}_x$  is arbitrary,  $x_\alpha \rightarrow x$ .

Conversely, suppose that there exists a net  $(x_\alpha)_{\alpha \in A} \subset S \setminus \{x\}$  such that  $x_\alpha \rightarrow x$ . Let  $U \in \mathcal{N}_x$ . Since  $(x_\alpha)_{\alpha \in A}$  is eventually in  $U$ , there exists  $\beta \in A$  such that  $x_\beta \in U$ . Then  $x_\beta \in (U \setminus \{x\}) \cap S$  and  $(U \setminus \{x\}) \cap S \neq \emptyset$ . Since  $U \in \mathcal{N}_x$  is arbitrary,  $x \in S'$ .  $\square$

**Exercise 4.3.14.** Let  $X$  be a topological space,  $S \subset X$  and  $x \in X$ . Then  $x \in \bar{S}$  iff there exists a net  $(x_\alpha)_{\alpha \in A} \subset S$  such that  $x_\alpha \rightarrow x$ .

*Proof.* Suppose that  $x \in \bar{S}$ . Since  $\bar{S} = S \cup S'$ ,  $x \in S$  or  $x \in S'$ . If  $x \in S$ , define  $(x_n)_{n \in \mathbb{N}} \subset S$  by  $x_n = x$ . Then  $x_n \rightarrow x$ . If  $x \in S'$ , the previous exercise implies that there exists a net  $(x_\alpha)_{\alpha \in A} \subset S \setminus \{x\} \subset S$  such that  $x_\alpha \rightarrow x$ .  $\square$

**Exercise 4.3.15.** Let  $X$  be a topological space and  $U \subset X$ . Then  $U$  is open iff for each net  $(x_\alpha)_{\alpha \in A} \subset X$  and  $x \in U$ ,  $x_\alpha \rightarrow x$  implies that  $(x_\alpha)_{\alpha \in A}$  is eventually in  $U$ .

*Proof.* Suppose that  $U$  is open. Let  $(x_\alpha)_{\alpha \in A} \subset X$  be a net and  $x \in U$ . Suppose that  $x_\alpha \rightarrow x$ . Since  $U \in \mathcal{N}_x$ ,  $(x_\alpha)_{\alpha \in A}$  is eventually in  $U$ .

Conversely, suppose that for each net  $(x_\alpha)_{\alpha \in A} \subset X$  and  $x \in U$ ,  $x_\alpha \rightarrow x$  implies that  $(x_\alpha)_{\alpha \in A}$  is eventually in  $U$ . For the sake of contradiction, suppose that  $U^c$  is not closed. Then there exist a net  $(x_\alpha)_{\alpha \in A} \subset U^c$  and  $x \in U$  such that  $x_\alpha \rightarrow x$ . By assumption,  $(x_\alpha)_{\alpha \in A}$  is eventually in  $U$ . This is a contradiction, so  $U^c$  is closed and hence  $U$  is open.  $\square$

**Exercise 4.3.16.** Let  $(X, \mathcal{A})$  and  $(Y, \mathcal{B})$  be topological spaces,  $f : X \rightarrow Y$  and  $x \in X$ . Then  $f$  is continuous at  $x$  iff for each net  $(x_\alpha)_{\alpha \in A} \subset X$ ,  $x_\alpha \rightarrow x$  implies that  $f(x_\alpha) \rightarrow f(x)$ .

*Proof.* Suppose that  $f$  is continuous at  $x$ . Let  $(x_\alpha)_{\alpha \in A} \subset X$  be a net. Suppose that  $x_\alpha \rightarrow x$ . Let  $V \in \mathcal{N}_{f(x)}$ . Continuity implies that  $f^{-1}(V) \in \mathcal{N}_x$ . Since  $x_\alpha \rightarrow x$ ,  $(x_\alpha)_{\alpha \in A}$  is eventually in  $f^{-1}(V)$ . So there exists  $\beta \in A$  such that for each  $\alpha \in A$ ,  $\alpha \geq \beta$  implies that  $x_\alpha \in f^{-1}(V)$ . Let  $\alpha \in A$ . Suppose that  $\alpha \geq \beta$ . Then  $f(x_\alpha) \in V$ . So  $(f(x_\alpha))_{\alpha \in A}$  is eventually in  $V$ . Since  $V \in \mathcal{N}_{f(x)}$  is arbitrary,  $f(x_\alpha) \rightarrow f(x)$ .

Conversely, suppose that  $f$  is not continuous at  $x$ . Then there exists  $V \in \mathcal{N}_{f(x)}$  such that  $f^{-1}(V) \notin \mathcal{N}_x$ . Then  $x \notin (f^{-1}(V))^\circ$ . So  $x \in ((f^{-1}(V))^\circ)^c = \overline{f^{-1}(V^c)}$ . This implies that there exists a net  $(x_\alpha)_{\alpha \in A} \subset f^{-1}(V^c)$  such that  $x_\alpha \rightarrow x$ . Since for each  $\alpha \in A$ ,  $f(x_\alpha) \in V^c$ ,  $f(x_\alpha)$  is not eventually in  $V$ . So  $f(x_\alpha) \not\rightarrow f(x)$ .  $\square$

**Exercise 4.3.17.** Let  $(Y_\alpha, \mathcal{B}_\alpha)_{\alpha \in A}$  be a collection of topological spaces,  $X$  a set and  $\mathcal{F} \in \prod_{\alpha \in A} Y_\alpha^X$  with  $\mathcal{F} = (f_\alpha)_{\alpha \in A}$ . Equip  $X$  with  $\tau_X(\mathcal{F})$ . Let  $(x_\gamma)_{\gamma \in \Gamma} \subset X$  be a net and  $x \in X$ . Then  $x_\gamma \rightarrow x$  iff for each  $\alpha \in A$ ,  $f_\alpha(x_\gamma) \rightarrow f_\alpha(x)$ .

*Proof.* Suppose that  $x_\gamma \rightarrow x$ . Let  $\alpha \in A$ . Since  $f_\alpha$  is continuous, the previous exercise implies that  $f_\alpha(x_\gamma) \rightarrow f_\alpha(x)$ .

Conversely, Suppose that for each  $\alpha \in A$ ,  $f_\alpha(x_\gamma) \rightarrow f_\alpha(x)$ . Let  $U \in \mathcal{N}_x$ . Since  $U^\circ \in \tau(\mathcal{F})$ , Exercise 4.1.11 implies there exist  $V_1 \in B_{\alpha_1}, \dots, V_n \in B_{\alpha_n}$  such that  $\bigcap_{j=1}^n f_{\alpha_j}^{-1}(V_j) \subset U^\circ$  and

$x \in \bigcap_{j=1}^n f_{\alpha_j}^{-1}(V_j)$ . Let  $j \in \{1, \dots, n\}$ . Since  $f_{\alpha_j}^{-1}(V_j) \in \mathcal{N}_x$ ,  $V_j \in \mathcal{N}_{f(x)}$ . By assumption,  $f_{\alpha_j}(x_\gamma)$  is eventually in  $V_j$ . Thus there exist there exist  $\gamma'_j \in \Gamma$  such that for each  $\gamma \geq \gamma'_j$ ,  $f_{\alpha_j}(x_\gamma) \in V_j$ , or equivalently,  $x_\gamma \in f_{\alpha_j}^{-1}(V_j)$ . Since  $\Gamma$  is directed, there exists  $\gamma' \in \Gamma$  such that for each  $j \in \{1, \dots, n\}$ ,  $\gamma' \geq \gamma'_j$ . Let  $\gamma \in \Gamma$ . Suppose that  $\gamma \geq \gamma'$ . Then

$$\begin{aligned} x_\gamma &\in \bigcap_{j=1}^n f_{\alpha_j}^{-1}(V_j) \\ &\subset U^\circ \\ &\subset U \end{aligned}$$

So  $(x_\gamma)_{\gamma \in \Gamma}$  is eventually in  $U$ . Since  $U \in \mathcal{N}_x$  is arbitrary,  $x_\gamma \rightarrow x$ .  $\square$

**Exercise 4.3.18.** Let  $X$  be a set and  $\mathcal{T}_1, \mathcal{T}_2$  topologies on  $X$ . Then the following are equivalent:

- (1)  $\mathcal{T}_1 = \mathcal{T}_2$
- (2) for each net  $(x_\alpha)_{\alpha \in A} \subset X$  and  $x \in X$ ,  $x_\alpha \rightarrow x$  in  $\mathcal{T}_1$  iff  $x_\alpha \rightarrow x$  in  $\mathcal{T}_2$ .

*Proof.*

- (1)  $\implies$  (2):  
Clear.

- (2)  $\implies$  (1):  
Let  $U \in \mathcal{T}_1$  and  $x \in U^c$ . Since  $U^c$  is closed in  $\mathcal{T}_1$ , there exists a net  $(x_\alpha)_{\alpha \in A} \subset U^c$  such that  $x_\alpha \rightarrow x$  in  $\mathcal{T}_1$ . By assumption,  $x_\alpha \rightarrow x$  in  $\mathcal{T}_2$ . So  $U^c$  is closed in  $\mathcal{T}_2$  and  $U \in \mathcal{T}_2$ . Hence  $\mathcal{T}_1 \subset \mathcal{T}_2$ .  
Similarly,  $\mathcal{T}_2 \subset \mathcal{T}_1$ .  $\square$

**Exercise 4.3.19.** Let  $X, Y$  be topological spaces and  $\phi : X \rightarrow Y$  a homeomorphism. Then for each  $E \subset X$ ,

- (1)  $\overline{\phi(E)} = \phi(\overline{E})$
- (2)  $\phi(E)^\circ = \phi(E^\circ)$

*Proof.*

- (1) Let  $E \subset X$ . Since  $E \subset \overline{E}$ , we have that  $\phi(E) \subset \phi(\overline{E})$ . Since  $\overline{E}$  is closed,  $\phi(\overline{E})$  is closed and thus  $\overline{\phi(E)} \subset \phi(\overline{E})$ . Conversely, let  $x \in \phi(\overline{E})$ . Then  $\phi^{-1}(x) \in \overline{E}$ . Then there exists a net  $(y_\alpha)_{\alpha \in A} \subset E$  such that  $y_\alpha \rightarrow \phi^{-1}(x)$ . Then  $(\phi(y_\alpha))_{\alpha \in A} \subset \phi(E)$  and  $\phi(y_\alpha) \rightarrow x$ . Thus  $x \in \overline{\phi(E)}$  and  $\phi(\overline{E}) \subset \overline{\phi(E)}$ .
- (2) Similar  $\square$

**Definition 4.3.20.** Let  $X$  be a topological space,  $(x_\alpha)_{\alpha \in A} \subset X$  a net and  $x \in X$ . Then  $x$  is said to be a **cluster point** of  $(x_\alpha)_{\alpha \in A}$  if for each  $U \in \mathcal{N}_x$ ,  $(x_\alpha)_{\alpha \in A}$  is frequently in  $U$ .

**Definition 4.3.21.** Let  $X$  be a topological space,  $(x_\alpha)_{\alpha \in A}, (y_\beta)_{\beta \in B} \subset X$  nets and  $\phi : B \rightarrow A$ . Then  $((y_\beta)_{\beta \in B}, \phi)$  is said to be a **subnet** of  $(x_\alpha)_{\alpha \in A}$  if

- (1) for each  $\beta \in B$ ,  $y_\beta = x_{\phi(\beta)}$
- (2) for each  $\alpha_0 \in A$ , there exists  $\beta_0 \in B$  such that for each  $\beta \in B$ ,  $\beta \geq \beta_0$  implies that  $\phi(\beta) \geq \alpha_0$

**Note 4.3.22.** We usually suppress  $\phi$  and write  $\alpha_\beta$  in place of  $\phi(\beta)$ .

**Exercise 4.3.23.** Let  $X$  be a topological space,  $(x_\alpha)_{\alpha \in A} \subset X$  a net and  $x \in X$ . Then  $x$  is a cluster point of  $(x_\alpha)_{\alpha \in A}$  iff there exists a subnet  $(x_{\alpha_\beta})_{\beta \in B}$  of  $(x_\alpha)_{\alpha \in A}$  such that  $x_{\alpha_\beta} \rightarrow x$ .

**Hint:** Order  $\mathcal{N}_x$  by reverse inclusion and consider the product directed set  $B = A \times \mathcal{N}_x$ . If  $x$  is a cluster point of  $(x_\alpha)_{\alpha \in A}$ , then for each  $\beta = (\gamma, U) \in B$ , there exists  $\alpha_\beta \in A$  such that  $\alpha_\beta \geq \gamma$  and  $\alpha_\beta \in U$ .

*Proof.* Suppose that  $x$  is a cluster point of  $(x_\alpha)_{\alpha \in A}$ . Set  $B = A \times \mathcal{N}_x$ . Since  $x$  is a cluster point of  $(x_\alpha)_{\alpha \in A}$ , for each  $(\gamma, U) \in B$ , there exists  $\alpha_{(\gamma, U)} \in A$  such that  $\alpha_{(\gamma, U)} \geq \gamma$  and

$x_{\alpha_{(\gamma,U)}} \in U$ . Let  $\alpha_0 \in A$ . Choose  $\beta_0 = (\alpha_0, X) \in B$ . Let  $\beta = (\gamma, U) \in B$ . Suppose that  $\beta \geq \beta_0$ . Then  $\gamma \geq \alpha_0$  and

$$\begin{aligned}\alpha_\beta &= \alpha_{(\gamma,U)} \\ &\geq \gamma \\ &\geq \alpha_0\end{aligned}$$

So that  $(x_{\alpha_\beta})_{\beta \in B}$  is a subnet of  $(x_\alpha)_{\alpha \in A}$ . Let  $U_0 \in \mathcal{N}_x$ . Choose  $\alpha_0 \in A$  and set  $\beta_0 = (\alpha_0, U_0)$ . Let  $\beta = (\gamma, U) \in B$ . Suppose that  $\beta \geq \beta_0$ . Then

$$\begin{aligned}x_{\alpha_\beta} &= x_{\alpha_{(\gamma,U)}} \\ &\in U \\ &\subset U_0\end{aligned}$$

Since  $U_0 \in \mathcal{N}_x$  is arbitrary,  $x_{\alpha_\beta} \rightarrow x$ .

Conversely, suppose that there exists a subnet  $(x_{\alpha_\beta})_{\beta \in B}$  of  $(x_\alpha)_{\alpha \in A}$  such that  $x_{\alpha_\beta} \rightarrow x$ . Let  $U \in \mathcal{N}_x$  and  $\alpha \in A$ . Since  $x_{\alpha_\beta} \rightarrow x$ , there exists  $\beta_1 \in B$  such that for each  $\beta \in B$ ,  $\beta \geq \beta_1$  implies that  $x_{\alpha_\beta} \in U$ . Since  $(x_{\alpha_\beta})_{\beta \in B}$  is a subnet of  $(x_\alpha)_{\alpha \in A}$ , there exists  $\beta_2 \in B$  such that for each  $\beta \in B$ ,  $\beta \geq \beta_2$  implies that  $\alpha_\beta \geq \alpha$ . Since  $B$  is directed, there exists  $\beta \in B$  such that  $\beta_1, \beta_2 \leq \beta$ . Then  $x_\beta \geq \alpha$  and  $x_\beta \in U$ . Since  $\alpha \in A$  is arbitrary,  $(x_{\alpha_\beta})_{\beta \in B}$  is frequently in  $U$ . Since  $U \in \mathcal{N}_x$  is arbitrary,  $x$  is a cluster point of  $(x_{\alpha_\beta})_{\beta \in B}$ .  $\square$

**Exercise 4.3.24.** Let  $X$  be a topological space,  $(x_\alpha)_{\alpha \in A} \subset X$  a net and  $x \in X$ . If  $x_\alpha \rightarrow x$ , then for each subnet  $(x_{\alpha_\beta})_{\beta \in B}$  of  $(x_\alpha)_{\alpha \in A}$ ,  $x_{\alpha_\beta} \rightarrow x$ .

*Proof.* Suppose that  $x_\alpha \rightarrow x$ . Let  $(x_{\alpha_\beta})_{\beta \in B}$  be a subnet of  $(x_\alpha)_{\alpha \in A}$  and  $U \in \mathcal{N}_x$ . Since  $x_\alpha \rightarrow x$ , there exists  $\alpha_0 \in A$  such that for each  $\alpha \geq \alpha_0$ ,  $x_\alpha \in U$ . Since  $(x_{\alpha_\beta})_{\beta \in B}$  is a subnet of  $(x_\alpha)_{\alpha \in A}$ , there exists  $\beta_0 \in B$  such that for each  $\beta \in B$ ,  $\beta \geq \beta_0$  implies that  $\alpha_{\beta_0} \geq \alpha_0$ . Then for each  $\beta \in B$ ,  $\beta \geq \beta_0$  implies that  $x_{\alpha_\beta} \in U$ . Since  $U \in \mathcal{N}_x$  is arbitrary,  $x_{\alpha_\beta} \rightarrow x$ .  $\square$

**Definition 4.3.25.** Let  $(x_\alpha)_{\alpha \in A} \subset \mathbb{R}$  a net. We define the **limit inferior** (resp. **limit superior**) of  $(x_\alpha)_{\alpha \in A}$ , denoted  $\liminf x_\alpha$  (resp.  $\limsup x_\alpha$ ), by

$$\liminf x_\alpha = \sup_{\beta \in A} \inf_{\alpha \geq \beta} x_\alpha$$

resp.

$$\limsup x_\alpha = \inf_{\beta \in A} \sup_{\alpha \geq \beta} x_\alpha$$

**Exercise 4.3.26.** Let  $(x_\alpha)_{\alpha \in A} \subset \mathbb{R}$  a net. Then

$$\liminf x_\alpha \leq \limsup x_\alpha$$

*Proof.* **FINISH!!!**  $\square$

**Exercise 4.3.27.** Let  $(x_\alpha)_{\alpha \in A} \subset \mathbb{R}$  a net and  $x \in \mathbb{R}$ . Then  $x_\alpha \rightarrow x$  iff

$$\liminf x_\alpha = \limsup x_\alpha = x$$



*Proof.* Suppose that  $x_\alpha \rightarrow x$ . Let  $\epsilon > 0$ . Then there exist  $\beta \in A$  such that for each  $\alpha \in A$ ,  $\alpha \geq \beta$  implies that  $x_\alpha \in B(x, \epsilon)$ . So  $\inf_{\alpha \geq \beta} x_\alpha \geq x - \epsilon$  and  $\sup_{\alpha \geq \beta} x_\alpha \leq x + \epsilon$ . Therefore

$$\begin{aligned} \liminf x_\alpha &= \sup_{\beta \in A} \inf_{\alpha \geq \beta} x_\alpha \\ &\geq x - \epsilon \end{aligned}$$

and

$$\begin{aligned} \limsup x_\alpha &= \inf_{\beta \in A} \sup_{\alpha \geq \beta} x_\alpha \\ &\leq x + \epsilon \end{aligned}$$

Since  $\epsilon > 0$  is arbitrary,

$$\limsup x_\alpha \leq x \leq \liminf x_\alpha$$

Since  $\liminf x_\alpha \leq \limsup x_\alpha$ , we have that  $\liminf x_\alpha = \limsup x_\alpha = x$ .  $\square$

**Exercise 4.3.28.** Let  $X$  be a topological space,  $f : X \rightarrow \mathbb{R}$ ,  $(x_\alpha)_{\alpha \in A} \subset X$  a net and  $x \in X$ . Suppose that  $x_\alpha \rightarrow x$  and for each  $\alpha \in A$ ,  $x_\alpha \neq x$ . Then

- (1)  $\liminf_{t \rightarrow x} f(t) \leq \liminf f(x_\alpha)$
- (2)  $\limsup_{t \rightarrow x} f(t) \geq \limsup f(x_\alpha)$

*Proof.*

- (1) Let  $V \in \mathcal{N}_x$ . Then there exists  $\beta \in A$  such that for each  $\alpha \geq \beta$ ,  $x_\alpha \in V \setminus \{x\}$ . Thus

$$\inf_{t \in V \setminus \{x\}} f(t) \leq \inf_{\alpha \geq \beta} f(x_\alpha)$$

which implies that

$$\inf_{t \in V \setminus \{x\}} f(t) \leq \sup_{\beta \in A} \inf_{\alpha \geq \beta} f(x_\alpha)$$

and since  $V \in \mathcal{N}_x$  is arbitrary, we have that

$$\begin{aligned} \liminf_{t \rightarrow x} f(t) &= \sup_{V \in \mathcal{N}_x} \inf_{t \in V \setminus \{x\}} f(t) \\ &\leq \sup_{\beta \in A} \inf_{\alpha \geq \beta} f(x_\alpha) \\ &= \liminf f(x_\alpha) \end{aligned}$$

- (2) Similar to (1).

$\square$

#### 4.4. Product Topology.

**Definition 4.4.1.** Let  $(X_\alpha)_{\alpha \in A}$  be a collection of topological spaces. We define the **product topology** on  $\prod_{\alpha \in A} X_\alpha$  to be the initial (weak) topology generated by  $(\pi_\alpha)_{\alpha \in A}$ .

**Exercise 4.4.2.** Let  $(X_\alpha, \mathcal{T}_\alpha)_{\alpha \in A}$  be a collection of topological spaces. Define

$$\mathcal{B} = \left\{ \prod_{\alpha \in A} B_\alpha : \text{for each } \alpha \in A, B_\alpha \in \mathcal{T}_\alpha \text{ and for finitely many } \alpha \in A, B_\alpha \neq X_\alpha \right\}$$

Then  $\mathcal{B}$  is a basis for the product topology on  $\prod_{\alpha \in A} X_\alpha$ .

*Proof.* Set  $X = \prod_{\alpha \in A} X_\alpha$ . Denote the product topology on  $X$  by  $\mathcal{T}_X$ . Set  $\mathcal{E} = \{\pi_\alpha^{-1}(B_\alpha) : \alpha \in A, B_\alpha \in \mathcal{T}_\alpha\}$ . By definition,  $\mathcal{T}_X = \tau_X(\mathcal{E})$ . Let  $\alpha \in A$  and  $B_\alpha \in \mathcal{T}_\alpha$ . For  $\beta \in A$ , set

$$C_\beta = \begin{cases} B_\beta & \beta = \alpha \\ X_\beta & \beta \neq \alpha \end{cases}$$

Then

$$\pi_\alpha^{-1}(B_\alpha) = \prod_{\beta \in A} C_\beta$$

Hence  $\mathcal{B} = \left\{ \bigcap_{j=1}^n V_j : (V_j)_{j=1}^n \subset \mathcal{E} \right\}$ . A previous exercise implies that  $\mathcal{B}$  is a basis for  $\mathcal{T}_X$ .  $\square$

**Definition 4.4.3.** Let  $X_1, X_2, Y_1, Y_2$  be sets and  $f_1 : X_1 \rightarrow Y_1, f_2 : X_2 \rightarrow Y_2$ . We define  $f_1 \times f_2 : X_1 \times X_2 \rightarrow Y_1 \times Y_2$  by

$$f_1 \times f_2(x_1, x_2) = (f_1(x_1), f_2(x_2))$$

**Exercise 4.4.4.** Let  $X_1, X_2, Y_1, Y_2$  be topological spaces and  $f_1 : X_1 \rightarrow Y_1, f_2 : X_2 \rightarrow Y_2$ . If  $f_1$  and  $f_2$  are continuous, then  $f_1 \times f_2$  is continuous.

*Proof.* Suppose that  $f_1$  and  $f_2$  are continuous. Let  $B_1 \subset Y_1, B_2 \subset Y_2$  be open. Then  $f_1 \times f_2^{-1}(B_1 \times B_2) = f_1^{-1}(B_1) \times f_2^{-1}(B_2)$  which is open in  $X_1 \times X_2$ . Since  $\mathcal{B} = \{B_1 \times B_2 : B_1 \subset Y_1 \text{ and } B_2 \subset Y_2 \text{ are open}\}$  generates the product topology on  $Y_1 \times Y_2$ , we have that  $f_1 \times f_2$  is continuous.  $\square$

**Exercise 4.4.5.** Let  $X_1, X_2, Y_1, Y_2$  be topological spaces and  $f_1 : X_1 \rightarrow Y_1, f_2 : X_2 \rightarrow Y_2$ . If  $f_1$  and  $f_2$  are open, then  $f_1 \times f_2$  is open.

*Proof.* Let  $A_1 \subset X_1, A_2 \subset X_2$  be open. Then  $f_1 \times f_2(A_1 \times A_2) = f_1(A_1) \times f_2(A_2)$  which is open in  $Y_1 \times Y_2$ . Since  $\mathcal{B} = \{A_1 \times A_2 : A_1 \subset X_1 \text{ and } A_2 \subset X_2 \text{ are open}\}$  is a basis for the product topology on  $X_1 \times X_2$ , an exercise in the section on continuous maps implies that  $f_1 \times f_2$  is open.  $\square$

#### 4.5. Subspace Topology.

**Definition 4.5.1.** Let  $X$  be a set and  $A \subset X$ . We define the **inclusion map** from  $A$  to  $X$ , denoted  $\iota : A \rightarrow X$ , by  $\iota(x) = x$ .

**Definition 4.5.2.** Let  $(X, \mathcal{T})$  be a topological space and  $A \subset X$ . We define the **subspace topology** on  $A$ , denoted  $\mathcal{T} \cap A$ , by

$$\mathcal{T} \cap A = \iota^*(\mathcal{T})$$

**Exercise 4.5.3.** Let  $(X, \mathcal{T})$  be a topological space and  $A \subset X$ . Then

$$\mathcal{T} \cap A = \{U \cap A : U \in \mathcal{T}\}$$

*Proof.* Clear. □

**Exercise 4.5.4.** universal property

*Proof.* **FINISH!!!** □

**Exercise 4.5.5.** Let  $(X, \mathcal{T})$  be a topological space,  $A \subset X$ ,  $(x_\gamma)_{\gamma \in \Gamma} \subset A$  a net and  $x \in A$ . Then  $x_\gamma \rightarrow x$  in  $(A, \mathcal{T} \cap A)$  iff  $x_\gamma \rightarrow x$  in  $(X, \mathcal{T})$ .

*Proof.* Suppose that  $x_\gamma \rightarrow x$  in  $(A, \mathcal{T} \cap A)$ . Since  $\iota : A \rightarrow X$  is continuous,

$$x_\gamma = \iota(x_\gamma) \rightarrow \iota(x) = x$$

So that  $x_\gamma \rightarrow x$  in  $(X, \mathcal{T})$ .

Conversely, suppose that  $x_\gamma \rightarrow x$  in  $(X, \mathcal{T})$ . Let  $V \in \mathcal{N}_x$  in  $(A, \mathcal{T} \cap A)$ . Then  $x \in V^\circ$  in  $(A, \mathcal{T} \cap A)$ . Hence there exists  $U \in \mathcal{T}$  such that  $V^\circ = U \cap A$ . Thus  $U \in \mathcal{N}_x$  in  $(X, \mathcal{T})$ . This implies that  $(x_\gamma)_{\gamma \in \Gamma}$  is eventually in  $U$ . Then  $(x_\gamma)_{\gamma \in \Gamma}$  is eventually in  $U \cap A = V^\circ \subset V$ . So  $x_\gamma \rightarrow x$  in  $(A, \mathcal{T} \cap A)$ . □

#### 4.6. Quotient Topology.

**Definition 4.6.1.** Let  $(X, \mathcal{A})$ ,  $(Y, \mathcal{B})$  be topological spaces and  $f : X \rightarrow Y$ . Then  $f$  is said to be a  **$\mathcal{A}$ - $\mathcal{B}$  quotient map** if

- (1)  $f$  is surjective
- (2)  $\mathcal{B}$  is the final topology on  $Y$  generated by  $f$ , i.e. for each  $V \subset Y$ ,  $V \in \mathcal{B}$  iff  $f^{-1}(V) \in \mathcal{A}$ .

**Note 4.6.2.** We typically avoid specifying the topologies when they are clear from the context.

**Exercise 4.6.3.** Let  $(X, \mathcal{A})$ ,  $(Y, \mathcal{B})$  be topological spaces and  $f : X \rightarrow Y$ . If  $f$  is a quotient map, then  $f$  is continuous.

*Proof.* Suppose that  $f$  is a quotient map. Let  $V \subset Y$ . Suppose that  $V$  is open. By definition,  $f^{-1}(V)$  is open. Hence  $f$  is continuous.  $\square$

**Exercise 4.6.4.** Let  $(X, \mathcal{A})$ ,  $(Y, \mathcal{B})$  be topological spaces and  $f : X \rightarrow Y$ . Suppose that  $f$  is continuous and surjective. Then  $f$  is a quotient map iff

for each  $C \subset Y$ ,  $C$  is closed iff  $f^{-1}(C)$  is closed

*Proof.*

- $(\implies)$

Suppose that  $f$  is a quotient map.

Let  $C \subset Y$ . If  $C$  is closed, then continuity implies that  $f^{-1}(C)$  is closed.

Conversely, suppose that  $f^{-1}(C)$  is closed. Then  $f^{-1}(C^c) = (f^{-1}(C))^c$  is open. Since  $f$  is a quotient map,  $f(f^{-1}(C^c))$  is open. Surjectivity implies that  $f(f^{-1}(C^c)) = C^c$ . So  $C$  is closed.

- $(\impliedby)$

Suppose that for each  $C \subset Y$ ,  $C$  is closed iff  $f^{-1}(C)$  is closed.

Let  $V \subset Y$ . If  $V$  is open. Continuity implies that  $f^{-1}(V)$  is open.

Conversely, suppose that  $f^{-1}(V)$  is open. Then  $f^{-1}(V^c) = (f^{-1}(V))^c$  is closed. Therefore,  $f(f^{-1}(V^c))$  is closed. Surjectivity implies that  $V^c = f(f^{-1}(V^c))$ . So  $V$  is open.

$\square$

**Exercise 4.6.5.** Let  $(X, \mathcal{A})$ ,  $(Y, \mathcal{B})$  be topological spaces and  $f : X \rightarrow Y$ . Suppose that  $f$  is continuous and surjective. If  $f$  is open or closed, then  $f$  is a quotient map.

*Proof.*

- Suppose that  $f$  is open. Let  $V \subset Y$ .

Suppose that  $V$  is open. Then continuity implies that  $f^{-1}(V)$  is open. Conversely, suppose that  $f^{-1}(V)$  is open. Since  $f$  is open  $f(f^{-1}(V))$  is open. Surjectivity implies that  $V = f(f^{-1}(V))$ . So  $V$  is open. By definition,  $f$  is a quotient map.

- Suppose that  $f$  is closed. Then similarly to above,  $f$  is a quotient map.

$\square$

**Exercise 4.6.6.** Let  $(X, \mathcal{A})$ ,  $(Y, \mathcal{B})$  be topological spaces and  $f : X \rightarrow Y$ . Suppose that  $f$  is a quotient map. Then  $f$  is open iff

for each  $U \subset X$ ,  $U$  is open implies that  $f^{-1}(f(U))$  is open

*Proof.*

- $(\implies)$

Suppose that  $f$  is open.

Let  $U \subset X$ . Suppose that  $U$  is open. Since  $f$  is open,  $f(U)$  is open. Continuity implies that  $f^{-1}(f(U))$  is open.

- $(\impliedby)$

Suppose that for each  $U \subset X$ ,  $U$  is open implies that  $f^{-1}(f(U))$  is open.

Since  $f$  is a quotient map,  $f(U)$  is open. So  $f$  is open.

□

**Exercise 4.6.7.** Let  $(X, \mathcal{A})$ ,  $(Y, \mathcal{B})$  be topological spaces, and  $f : X \rightarrow Y$ . Suppose that  $f$  is surjective and continuous. If  $f$  is open or closed, then  $f$  is a quotient map.

*Proof.* By continuity,  $\mathcal{B} \subset f_*\mathcal{A}$ .

- Suppose that  $f$  is open. Let  $V \in f_*\mathcal{A}$ . By definition,  $f^{-1}(V) \in \mathcal{A}$ . Since  $f$  is open,  $f(f^{-1}(V)) \in \mathcal{B}$ . Surjectivity implies that  $V = f(f^{-1}(V))$ . So  $f_*\mathcal{A} = \mathcal{B}$  and  $f$  is a  $\mathcal{A}$ - $\mathcal{B}$  quotient map.
- The case is similar if  $f$  is closed.

□

**Definition 4.6.8.** Let  $(X, \mathcal{T})$  be a topological space,  $Y$  a set and  $f : X \rightarrow Y$ . Suppose that  $f$  is surjective. We call  $f_*\mathcal{T}$  a **quotient topology** on  $Y$ .

**Exercise 4.6.9.** Let  $(X, \mathcal{T})$  be a topological space,  $Y$  a set and  $f : X \rightarrow Y$ . Suppose that  $f$  is surjective. Then  $f : X \rightarrow Y$  is a  $\mathcal{T}$ - $f_*\mathcal{T}$  quotient map.

*Proof.* Clear.

□

**Exercise 4.6.10.** Let  $(X, \mathcal{T})$  be a topological space,  $\sim$  an equivalence relation on  $X$  and  $\pi : X \rightarrow X/\sim$  the projection map given by  $x \mapsto \bar{x}$ . Then  $\pi$  is a  $\mathcal{T}$ - $\pi_*\mathcal{T}$  quotient map.

*Proof.* Since  $\pi$  is surjective, the previous exercise implies that  $\pi$  is a  $\mathcal{T}$ - $\pi_*\mathcal{T}$  quotient map. □

**Definition 4.6.11.** Let  $(X, \mathcal{T})$  be a topological space,  $\sim$  an equivalence relation on  $X$  and  $\pi : X \rightarrow X/\sim$  the projection map given by  $x \mapsto \bar{x}$ . We define the **quotient topology on  $X/\sim$**  on  $X/\sim$ , denoted  $\mathcal{T}_{X/\sim}$ , by

$$\mathcal{T}_{X/\sim} = \pi_*\mathcal{T}$$

**Definition 4.6.12.** Let  $X, Y$  be sets,  $\sim$  an equivalence relation on  $X$  and  $f : X \rightarrow Y$ . Then  $f$  is said to be  **$\sim$ -invariant** if for each  $a, b \in X$ ,  $\bar{a} = \bar{b}$  implies that  $f(a) = f(b)$ .

**Definition 4.6.13.** Let  $X, Y$  be sets,  $\sim$  an equivalence relation on  $X$  and  $f : X \rightarrow Y$ . Suppose that  $f$  is  $\sim$ -invariant

**Exercise 4.6.14.** Let  $(X, \mathcal{A})$ ,  $(Y, \mathcal{B})$  be topological spaces,  $\sim$  an equivalence relation on  $X$ ,  $\pi : X \rightarrow X/\sim$  the projection map and  $f : X \rightarrow Y$  continuous. If  $f$  is  $\sim$ -invariant, then there exists a unique  $\bar{f} : X/\sim \rightarrow Y$  such that

- (1)  $\bar{f} \circ \pi = f$
- (2)  $\bar{f}$  is  $\mathcal{A}$ - $\pi_*\mathcal{A}$  continuous

*Proof.* Suppose that  $f$  is  $\sim$ -invariant. Define  $\bar{f} : X/\sim \rightarrow Y$  by  $\bar{f}(\bar{x}) = f(x)$ . By assumption, for each  $a, b \in X$ ,  $\bar{a} = \bar{b}$  implies that  $f(a) = f(b)$ . Thus  $\bar{f}$  is well defined. By construction,  $f = \bar{f} \circ \pi$ . Let  $V \in \mathcal{B}$ . Continuity of  $f$  implies that  $f^{-1}(V) \in \mathcal{A}$ . Since

$$\begin{aligned} f^{-1}(V) &= \pi^{-1}(\bar{f}^{-1}(V)) \\ &\in \mathcal{A} \end{aligned}$$

by definition of the quotient topology,  $\bar{f}^{-1}(V) \in \pi_*\mathcal{A}$ . So  $\bar{f}$  is  $\mathcal{A}$ - $\pi_*\mathcal{A}$  continuous. □

**Exercise 4.6.15.** Let  $G$  be a group,  $X$  a topological space and  $\phi : G \times X \rightarrow X$  a group action. Suppose that for each  $g \in G$ , the map  $\phi_g \in \text{Sym}(X)$  defined by  $\phi_g(x) = g \cdot x$  is continuous. Then  $\pi : X \rightarrow X/G$  is open.

*Proof.* Suppose that for each  $g \in G$ ,  $\phi_g$  is continuous. Let  $g \in G$ . Since  $(\phi_g)^{-1} = \phi_{g^{-1}}$ ,  $\phi_g$  is a homeomorphism. Hence for each  $g \in G$  and  $U \subset X$ ,  $g \cdot U$  is open. Let  $U \subset X$ . Suppose that  $U$  is open. Then  $\pi^{-1}(\pi(U)) = \bigcup_{g \in G} g \cdot U$  is open. A previous exercise implies that  $\pi$  is open. □

#### 4.7. Separation and Countability.

**Definition 4.7.1.** Let  $X$  be a topological space. Then  $X$  is said to be  $\mathbf{T}_1$  if for each  $x, y \in X$ , if  $x \neq y$ , then there exists  $U \in \mathcal{N}_x$  such that  $U$  is open and  $y \notin U$ .

**Exercise 4.7.2.** Let  $X$  be a topological space. Then  $X$  is  $T_1$  iff for each  $x \in X$ ,  $\{x\}$  is closed.

*Proof.* Suppose that  $X$  is  $T_1$ . Let  $a \in \{x\}^c$ . Then there exists  $U_a \in \mathcal{N}_a$  such that  $U_a$  is open and  $U_a \subset \{x\}^c$ . Therefore

$$\{x\}^c = \bigcap_{a \in \{x\}^c} U_a$$

which is open. Hence  $\{x\}$  is closed.

Conversely, suppose that for each  $x \in X$ ,  $\{x\}$  is closed. Let  $x, y \in X$ . Suppose that  $x \neq y$ . Since  $\{y\}$  is closed,  $\{y\}^c$  is open and  $x \in \{y\}^c$ .  $\square$

**Definition 4.7.3.** Let  $X$  be a topological space. Then  $X$  is said to be  $\mathbf{T}_2$  or **Hausdorff** if for each  $x, y \in X$ , if  $x \neq y$ , then there exist  $U \in \mathcal{N}_x$  and  $V \in \mathcal{N}_y$  such that  $U$  and  $V$  are open and  $U \cap V = \emptyset$ .

**Exercise 4.7.4.** Let  $X$  be a topological space. If  $X$  is Hausdorff, then  $X$  is  $T_1$ .

*Proof.* Clear.  $\square$

**Exercise 4.7.5.** Let  $X$  be a topological space. Then the following are equivalent:

- (1)  $X$  is Hausdorff
- (2) for each net  $(x_\alpha)_{\alpha \in A} \subset X$  and  $x, y \in X$ , if  $x_\alpha \rightarrow x$  and  $x_\alpha \rightarrow y$ , then  $x = y$ .
- (3) The diagonal  $\Delta_X = \{(x, x) : x \in X\}$  is closed in  $X \times X$ .

*Proof.*

- (1)  $\implies$  (2):

Suppose that  $X$  is Hausdorff. Let  $(x_\alpha)_{\alpha \in A} \subset X$  be a net and  $x, y \in X$ . Suppose that  $x_\alpha \rightarrow x$  and  $x_\alpha \rightarrow y$ . For the sake of contradiction, suppose that  $x \neq y$ . Then there exist  $U \in \mathcal{N}_x$  and  $V \in \mathcal{N}_y$  such that  $U$  and  $V$  are open and  $U \cap V = \emptyset$ . Since  $x_\alpha \rightarrow x$ ,  $(x_\alpha)_{\alpha \in A}$  is eventually in  $U$  and there exists  $\beta_x \in A$  such that for each  $\alpha \in A$ ,  $\alpha \geq \beta_x$  implies that  $x_\alpha \in U$ . Since  $x_\alpha \rightarrow y$ ,  $(x_\alpha)_{\alpha \in A}$  is eventually in  $V$  and there exists  $\beta_y \in A$  such that for each  $\alpha \in A$ ,  $\alpha \geq \beta_y$  implies that  $x_\alpha \in V$ . Since  $A$  is directed, there exists  $\beta \in A$  such that  $\beta \geq \beta_x, \beta_y$ . Hence

$$\begin{aligned} x_\beta &\in U \cap V \\ &= \emptyset \end{aligned}$$

which is a contradiction. So  $x = y$ .

- (2)  $\implies$  (3):

Let  $(x_\alpha, y_\alpha)_{\alpha \in A} \subset \Delta_X$  be a net and  $(x, y) \in X \times X$ . Then for each  $\alpha \in A$ ,  $x_\alpha = y_\alpha$ . Suppose that  $(x_\alpha, y_\alpha) \rightarrow (x, y)$ . So  $x_\alpha \rightarrow x$  and  $x_\alpha \rightarrow y$ . Hence  $x = y$  and  $(x, y) \in \Delta_X$ . Thus  $\Delta_X$  is closed.

- (3)  $\implies$  (1):

Suppose that  $\Delta_X$  is closed. Let  $x, y \in X$ . Suppose that  $x \neq y$ . Then  $(x, y) \in \Delta_X^c$ . Recall that  $\mathcal{B} = \{A \times B : A, B \subset X \text{ and } A, B \text{ are open}\}$  is a basis for the product topology on  $X \times X$ . Since  $\Delta_X^c$  is open and  $(x, y) \in \Delta_X^c$ , there exist  $A \times B \in \mathcal{B}$  such

that  $(x, y) \in A \times B \subset \Delta_X^c$ . Suppose that  $A \cap B \neq \emptyset$ . Then there exists  $z \in A \cap B$ . Hence  $(z, z) \in A \times B$ . This is a contradiction since  $A \times B \subset \Delta_X^c$ . Thus  $x \in A$ ,  $y \in B$  and  $A \cap B = \emptyset$  and  $A, B$  are open. Since  $x, y \in X$  are arbitrary,  $X$  is Hausdorff.  $\square$

**Exercise 4.7.6.** Let  $X$  be a topological space and  $\sim$  an equivalence relation on  $X$ . If  $\pi : X \rightarrow X/\sim$  is open, then  $X/\sim$  is Hausdorff iff  $\sim$  is closed in  $X \times X$ .

*Proof.* Suppose that  $\pi : X \rightarrow X/\sim$  is open.

• ( $\implies$ ):

Suppose that  $X/\sim$  is Hausdorff. Let  $(x_\alpha, y_\alpha)_{\alpha \in A} \subset \sim$  be a net and  $(x, y) \in X \times X$ . Suppose that  $x_\alpha, y_\alpha \rightarrow (x, y)$ . Then  $x_\alpha \rightarrow x$  and  $y_\alpha \rightarrow y$ . Since  $\pi : X \rightarrow X/\sim$  is continuous,  $\pi(x_\alpha) \rightarrow \pi(x)$  and  $\pi(y_\alpha) \rightarrow \pi(y)$ . Since for each  $\alpha \in A$ ,  $x_\alpha \sim y_\alpha$ , we have that

$$\begin{aligned} \pi(x_\alpha) &= \pi(y_\alpha) \\ &\rightarrow \pi(y) \end{aligned}$$

Since  $X/\sim$  is Hausdorff,  $\pi(x) = \pi(y)$ . Hence  $x \sim y$  and  $(x, y) \in \sim$ . Thus  $\sim$  is closed in  $X \times X$ .

• ( $\impliedby$ ):

Conversely, suppose that  $\sim$  is closed in  $X \times X$  is closed. Let  $\bar{x}, \bar{y} \in X/\sim$ . Suppose that  $\bar{x} \neq \bar{y}$ . Then  $(x, y) \in \sim^c$ . Recall that  $\mathcal{B} = \{A \times B : A, B \subset X \text{ and } A, B \text{ are open}\}$  is a basis for  $X \times X$ . Since  $\sim^c$  is open and  $(x, y) \in \sim^c$ , there exist  $A, B \subset X$  such that  $A, B$  are open and  $(x, y) \in A \times B \subset \sim^c$ . Thus  $x \in A$  and  $y \in B$ . Since  $\pi$  is open,  $\pi(A) = \bar{A}$  and  $\pi(B) = \bar{B}$  are open. Suppose for the sake of contradiction that  $\pi(A) \cap \pi(B) \neq \emptyset$ . Then there exists  $z \in X$  such that  $\bar{z} \in \pi(A) \cap \pi(B)$ . Therefore there exist  $z_A \in A$  and  $z_B \in B$  such that  $z_A \sim z \sim z_B$ . Then  $(z_A, z_B) \in A \times B$  and  $(z_A, z_B) \in \sim$ . This is a contradiction since  $A \times B \subset \sim^c$ . So  $\pi(A) \cap \pi(B) = \emptyset$ . Thus  $\bar{x} \in \pi(A)$ ,  $\bar{y} \in \pi(B)$ ,  $\pi(A)$ ,  $\pi(B)$  are open and  $\pi(A) \cap \pi(B) = \emptyset$ . Since  $\bar{x}, \bar{y} \in X/\sim$  are arbitrary,  $X/\sim$  is Hausdorff.  $\square$



#### 4.8. Compactness.

**Definition 4.8.1.** Let  $X$  be a topological space and  $E \subset X$ . Then  $E$  is said to be **precompact** if  $\overline{E}$  is compact.

#### 4.9. Semi-continuity.

**Definition 4.9.1.** Let  $X$  be a topological space,  $f : X \rightarrow (\infty, \infty]$  and  $x_0 \in X$ . Then  $f$  is said to be **lower semicontinuous at**  $x_0$  if

$$\liminf_{x \rightarrow x_0} f(x) \geq f(x_0)$$

and  $f$  is said to be **lower semicontinuous** if for each  $x_0 \in X$ ,  $f$  is lower semicontinuous at  $x_0$ .

**Exercise 4.9.2.** Let  $X$  be a topological space and  $f : X \rightarrow (\infty, \infty]$ . Then  $f$  is lower semicontinuous iff for each  $\alpha \in \mathbb{R}$ ,  $f^{-1}((\alpha, \infty])$  is open.

*Proof.* Suppose that  $f$  is lower semicontinuous. Let  $\alpha \in \mathbb{R}$  and  $x_0 \in f^{-1}(\alpha, \infty]$ . Put  $\epsilon = f(x_0) - \alpha$ . By definition,

$$\sup_{V \in \mathcal{N}_{x_0}} \inf_{x \in V \setminus \{x_0\}} f(x) \geq f(x_0)$$

Choose  $V_\epsilon \in \mathcal{N}_{x_0}$  such that

$$\begin{aligned} \inf_{x \in V_\epsilon \setminus \{x_0\}} f(x) &> f(x_0) - \epsilon \\ &= \alpha \end{aligned}$$

Then  $V_\epsilon^o \in \mathcal{N}_{x_0}$  is open and

$$\begin{aligned} V_\epsilon^o &\subset V_\epsilon \\ &\subset f^{-1}((\alpha, \infty]) \end{aligned}$$

So  $f^{-1}((\alpha, \infty])$  is open.

Conversely, suppose that for each  $\alpha \in \mathbb{R}$ ,  $f^{-1}((\alpha, \infty])$  is open. Let  $x_0 \in X$ . Put  $\alpha = f(x_0)$ . For  $n \in \mathbb{N}$ , define  $V_n = f^{-1}((f(x_0) - 1/n, \infty])$ . Then for each  $n \in \mathbb{N}$ ,  $V_n \in \mathcal{N}_{x_0}$  and

$$\begin{aligned} \liminf_{x \rightarrow x_0} f(x) &= \sup_{V \in \mathcal{N}_{x_0}} \inf_{x \in V \setminus \{x_0\}} f(x) \\ &\geq \sup_{n \in \mathbb{N}} \inf_{x \in V_n \setminus \{x_0\}} f(x) \\ &\geq \sup_{n \in \mathbb{N}} f(x_0) - 1/n \\ &= f(x_0) \end{aligned}$$

So  $f$  is lower semicontinuous. □

**Definition 4.9.3.** Let  $X$  be a topological space and  $f : X \rightarrow \mathbb{R}$ . We define the **epigraph of**  $f$ , denoted  $\text{epi } f$ , by

$$\text{epi } f = \{(x, y) \in X \times \mathbb{R} : f(x) \leq y\}$$

**Exercise 4.9.4.** Let  $X$  be a topological space and  $f : X \rightarrow \mathbb{R}$ . Then  $f$  is lower semicontinuous iff  $\text{epi } f$  is closed.

*Proof.* Suppose that  $f$  is lower semicontinuous. Let  $(x_\alpha, y_\alpha)_{\alpha \in A} \subset \text{epi } f$  be a net and  $(x, y) \in X \times \mathbb{R}$ . Then for each  $\alpha \in A$ ,  $f(x_\alpha) \leq y_\alpha$ . Suppose that  $(x_\alpha, y_\alpha) \rightarrow (x, y)$ . Then  $x_\alpha \rightarrow x$  and  $y_\alpha \rightarrow y$ . Therefore

$$\begin{aligned} f(x) &\leq \liminf_{t \rightarrow x} f(t) \\ &\leq \liminf f(x_\alpha) \\ &\leq \liminf y_\alpha \\ &= y \end{aligned}$$

So  $(x, y) \in \text{epi } f$  and  $\text{epi } f$  is closed.

Conversely, suppose that  $\text{epi } f$  is closed. □

**Exercise 4.9.5.** Let  $X$  be a topological space and  $(f_\lambda)_{\lambda \in \Lambda} \subset (-\infty, \infty]^X$ . Suppose that for each  $\lambda \in \Lambda$ ,  $f_\lambda$  is lower semicontinuous. Set  $f = \sup_{\lambda \in \Lambda} f_\lambda$ . Then  $f$  is lower semicontinuous.

*Proof.* Let  $\alpha \in \mathbb{R}$  and  $x \in X$ . Then

$$\begin{aligned} x \in f^{-1}((\alpha, \infty]) &\iff \sup_{\lambda \in \Lambda} f_\lambda(x) > \alpha \\ &\iff \text{there exists } \lambda \in \Lambda \text{ such that } f_\lambda(x) > \alpha \\ &\iff \text{there exists } \lambda \in \Lambda \text{ such that } x \in f_\lambda^{-1}((\alpha, \infty]) \\ &\iff x \in \bigcup_{\lambda \in \Lambda} f_\lambda^{-1}((\alpha, \infty]) \end{aligned}$$

Since for each  $\lambda \in \Lambda$ ,  $f_\lambda^{-1}((\alpha, \infty])$  is open,  $f^{-1}((\alpha, \infty]) = \bigcup_{\lambda \in \Lambda} f_\lambda^{-1}((\alpha, \infty])$  is open. So  $f$  is lower semicontinuous. □

## 5. LOCALLY CONVEX SPACES

## 5.1. Topological Vector Spaces.

**Definition 5.1.1.** Let  $X$  be a vector space and  $\mathcal{T}$  a topology on  $X$ . Then  $X$  is said to be a **topological vector space** if

- (1) addition  $X \times X \rightarrow X$  is continuous
- (2) scalar multiplication  $\mathbb{C} \times X \rightarrow X$  is continuous

**Note 5.1.2.** We usually suppress the topology  $\mathcal{T}$ .

**Exercise 5.1.3.** Let  $X$  be a topological vector space,  $(\lambda_\alpha)_{\alpha \in A} \subset \mathbb{C}$ ,  $(x_\alpha)_{\alpha \in A}$ ,  $(y_\alpha)_{\alpha \in A} \subset X$  nets and  $\lambda \in \mathbb{C}$ ,  $x, y \in X$ . If  $\lambda_\alpha \rightarrow \lambda$ ,  $x_\alpha \rightarrow x$  and  $y_\alpha \rightarrow y$ , then  $x_\alpha + \lambda_\alpha y_\alpha \rightarrow x + \lambda y$ .

*Proof.* Clear since addition and scalar multiplication are continuous.  $\square$

**Exercise 5.1.4.** Let  $X$  be a topological vector space,  $a \in X$  and  $\lambda \in \mathbb{C}^\times$ . Define  $f, g : X \rightarrow X$  by  $f(x) = x + a$  and  $g(x) = \lambda x$ . Then  $f$  and  $g$  are homeomorphisms.

*Proof.* Since  $X$  is a topological vector space,  $f$  and  $g$  are continuous. Clearly  $f$  and  $g$  are bijections with  $f^{-1}(x) = x - a$  and  $g^{-1}(x) = \lambda^{-1}x$ . Again, since  $X$  is a topological vector space,  $f^{-1}$  and  $g^{-1}$  are continuous.  $\square$

**Exercise 5.1.5.** Let  $X$  be a topological vector space,  $x, y \in X$  and  $U \in \mathcal{N}_x$ . If  $U$  is open, then there exists  $r > 0$  such that for each  $t \in \mathbb{R}$ ,  $|t| \leq r$  implies that  $x + ty \in U$ .

*Proof.* Suppose that  $U$  is open. For the sake of contradiction, suppose that for each  $r > 0$ , there exists  $t \in \mathbb{R}$  such that  $t \leq r$  and  $x + ty \notin U$ . Then for each  $n \in \mathbb{N}$ , there exists  $t_n \in \mathbb{R}$  such that  $|t_n| \leq 1/n$  and  $x + t_n y \in U^c$ . Since  $t_n \rightarrow 0$ ,

$$\begin{aligned} x + t_n y &\rightarrow x + 0y \\ &= x \end{aligned}$$

Since  $U^c$  is closed,  $x \in U^c$ . This is a contradiction. Hence there exists  $r > 0$  such that for each  $t \in \mathbb{R}$ ,  $|t| \leq r$  implies that  $x + ty \in U$ .  $\square$

**Exercise 5.1.6.** Let  $X$  be a topological vector space and  $A, B \subset X$ . If  $A$  is open, then  $A + B$  is open.

*Proof.* Suppose that  $A$  is open. Then for each  $b \in B$ ,  $A + b$  is open. Since

$$A + B = \bigcup_{b \in B} A + b$$

we have that  $A + B$  is open.  $\square$

**Exercise 5.1.7.** Let  $X$  be a topological vector space and  $A, B \subset X$ . Suppose that  $A$  is compact,  $B$  is closed and  $A \cap B = \emptyset$ . Then there exists  $U \in \mathcal{N}_0$  such that  $U$  is open and  $(A + U) \cap B = \emptyset$ .

*Proof.* Set  $\Gamma = \{U \in \mathcal{N}_0 : U \text{ is open}\}$  and order  $\Gamma$  by reverse inclusion, so that  $\Gamma$  is a directed set. For the sake of contradiction, suppose that for each  $U \in \Gamma$ ,  $(A + U) \cap B \neq \emptyset$ . Then for each  $\gamma \in \Gamma$ , there exist  $a_\gamma \in A$  and  $u_\gamma \in \gamma$  such that  $a_\gamma + u_\gamma \in B$ . Let  $V \in \mathcal{N}_0$ . Since  $\text{Int } V \in \Gamma$

$$\begin{aligned} u_{\text{Int } V} &\in \text{Int } V \\ &\subset V \end{aligned}$$

Since  $V \in \mathcal{N}_0$  is arbitrary,  $u_\gamma \rightarrow 0$ . Since  $A$  is compact, there exists  $a \in A$  and a subnet  $(a_{\gamma_\zeta})_{\zeta \in Z}$  of  $(a_\gamma)_{\gamma \in \Gamma}$  such that  $a_{\gamma_\zeta} \rightarrow a$ . Then  $a_{\gamma_\zeta} + u_{\gamma_\zeta} \rightarrow a$ . Since  $(a_{\gamma_\zeta} + u_{\gamma_\zeta})_{\zeta \in Z} \subset B$  and  $B$  is closed, we have that  $a \in B$ . This is a contradiction since  $A \cap B = \emptyset$ . So there exists  $U \in \mathcal{N}_0$  such that  $U$  is open and  $(A + U) \cap B = \emptyset$ .  $\square$

**Exercise 5.1.8.** Let  $X$  be a topological vector space and  $U \in \mathcal{N}_0$ . If  $U$  is open, then there exists  $V \in \mathcal{N}_0$  such that  $V$  is open and  $V + V \subset U$ .

*Proof.* Suppose that  $U$  is open. Set  $\Gamma = \{V \in \mathcal{N}_0 : V \text{ is open}\}$  and order  $\Gamma$  by reverse inclusion, so that  $\Gamma$  is a directed set. For the sake of contradiction, suppose that for each  $V \in \mathcal{N}_0$ , if  $V$  is open, then  $V + V \not\subset U$ . Then for each  $\gamma \in \Gamma$ , there exists  $x_\gamma, y_\gamma \in \gamma$  such that  $x_\gamma + y_\gamma \in U^c$ . Let  $W \in \mathcal{N}_0$ . Set  $\beta = \text{Int } U$ . Then  $\beta \in \Gamma$ . Then for each  $\gamma \geq \beta$ ,

$$\begin{aligned} x_\gamma, y_\gamma &\in \gamma \\ &\subset \beta \\ &\subset W \end{aligned}$$

So that  $(x_\gamma)_{\gamma \in \Gamma}$  and  $(y_\gamma)_{\gamma \in \Gamma}$  are eventually in  $W$ . Since  $W \in \mathcal{N}_0$  is arbitrary,  $x_\gamma \rightarrow 0$  and  $y_\gamma \rightarrow 0$ . Therefore  $x_\gamma + y_\gamma \rightarrow 0$ . Since for each  $\gamma \in \Gamma$ ,  $x_\gamma + y_\gamma \in U^c$  and  $U^c$  is closed,  $0 \in U^c$ . This is a contradiction, so there exists  $V \in \mathcal{N}_0$  such that  $V$  is open and  $V + V \subset U$ .  $\square$

**Definition 5.1.9.** Let  $X$  be a vector space over  $\mathbb{C}$  and  $T : X \rightarrow \mathbb{C}$ . Then  $T$  is said to be a **linear functional on  $X$**  if  $T$  is linear. We define the **algebraic dual space of  $X$** , denoted  $X^*$ , by  $X^* = \{T : X \rightarrow \mathbb{C} : T \text{ is linear}\}$

**Note 5.1.10.** We define  $X^*$  similarly when  $X$  is a vector space over  $\mathbb{R}$ .

**Definition 5.1.11.** Let  $X$  be a topological vector space over  $\mathbb{C}$  and  $T : X \rightarrow \mathbb{C}$ . We define the **dual space of  $X$** , denoted  $X^*$ , by  $X^* = \{T : X \rightarrow \mathbb{C} : T \text{ is linear and continuous}\}$

**Note 5.1.12.** We define  $X^*$  similarly when  $X$  is a vector space over  $\mathbb{R}$ .

**Exercise 5.1.13.** Let  $X$  be a topological vector space. Then  $X^*$  is a vector space.

*Proof.* Clear.  $\square$

**Exercise 5.1.14.** Let  $X, Y$  be topological vector spaces and  $\phi : X \rightarrow Y$ . Suppose that  $\phi$  is linear. Then  $\phi$  is continuous iff  $\phi$  is continuous at 0.

*Proof.* If  $\phi$  is continuous, then  $\phi$  is continuous at 0.

Conversely, suppose that  $\phi$  is continuous at 0. Let  $(x_\alpha)_{\alpha \in A} \subset X$  be a net and  $x \in X$ . Suppose that  $x_\alpha \rightarrow x$ . Then  $x_\alpha - x \rightarrow 0$ . Hence

$$\begin{aligned} \phi(x_\alpha) - \phi(x) &= \phi(x_\alpha - x) \\ &\rightarrow \phi(0) \\ &= 0 \end{aligned}$$

Therefore  $\phi(x_\alpha) \rightarrow \phi(x)$  and  $\phi$  is continuous at  $x$ . Since  $x \in X$  is arbitrary,  $\phi$  is continuous.  $\square$

**Exercise 5.1.15.** Let  $X$  be a topological vector space and  $\phi : X \rightarrow \mathbb{C}$  linear. Then  $\phi \in X^*$  iff  $|\phi|$  is continuous.

*Proof.* Suppose that  $\phi$  is continuous. Since  $|\cdot| : \mathbb{C} \rightarrow [0, \infty)$  is continuous,  $|\phi|$  is continuous. Conversely, suppose that  $|\phi|$  is continuous. Let  $(x_\alpha)_{\alpha \in A} \subset X$  be a net and  $x \in X$ . Suppose that  $x_\alpha \rightarrow x$ . Then  $x_\alpha - x \rightarrow 0$ . Therefore

$$\begin{aligned} |\phi(x_\alpha) - \phi(x)| &= |\phi(x_\alpha - x)| \\ &\rightarrow |\phi(0)| \\ &= 0 \end{aligned}$$

So  $\phi(x_\alpha) \rightarrow \phi(x)$  and  $\phi$  is continuous.  $\square$

**Exercise 5.1.16.** Let  $X$  be a real topological vector space and  $\phi \in X^*$ . If  $\phi$  is not constant, then  $\phi$  is open.

**Hint:** There exists  $x_* \in X$  such that  $\phi(x_*) = 1$  and for each  $U \subset X$  open and  $x \in U$ , there exists  $r > 0$  such that for each  $t \in \mathbb{R}$ ,  $|t| \leq r$  implies that  $x + tx_* \in U$ .

*Proof.* Suppose that  $\phi$  is not constant. Then there exists  $x_0 \in X$  such that  $\phi(x_0) \neq 0$ . Set  $x_* = \phi(x_0)^{-1}x_0$ . Then

$$\begin{aligned} \phi(x_*) &= \phi(\phi(x_0)^{-1}x_0) \\ &= \phi(x_0)^{-1}\phi(x_0) \\ &= 1 \end{aligned}$$

Let  $U \subset X$  be open and  $y \in \phi(U)$ . Then there exists  $x \in U$  such that  $\phi(x) = y$ . Since  $U$  is open, a previous exercise implies that there exists  $r > 0$  such that for each  $t \in \mathbb{R}$ ,  $\|t\| \leq r$  implies that  $x + tx_* \in U$ . Let  $t \in (-r, r)$ . Then  $\phi(x + tx_*) \in \phi(U)$ . Since

$$\begin{aligned} \phi(x + tx_*) &= \phi(x) + t\phi(x_*) \\ &= y + t \end{aligned}$$

we have that  $(y - r, y + r) \subset \phi(U)$ . Since  $y \in U$  is arbitrary,  $\phi(U)$  is open thus  $\phi$  is open.  $\square$

**Definition 5.1.17.** Let  $X$  be a vector space and  $\phi : X \rightarrow \mathbb{C}$ . Then  $\phi$  is said to be **real-linear** if for each  $x, y \in X$  and  $\lambda \in \mathbb{R}$ ,  $\phi(x + \lambda y) = \phi(x) + \lambda\phi(y)$ .

**Exercise 5.1.18.** Let  $X$  be a topological vector space and  $\phi \in X^*$ . Then  $\text{Re } \phi$  is continuous and real-linear.

*Proof.* Clear.  $\square$

**Exercise 5.1.19.** Let  $X$  be a topological vector space and  $f : X \rightarrow \mathbb{R}$ . If  $f$  is continuous and real-linear, then there exists a unique  $\phi \in X^*$  such that  $\text{Re } \phi = f$ .

**Hint:** For each  $z \in \mathbb{C}$ ,  $z = \text{Re}(z) - i\text{Re}(iz)$

*Proof.* Suppose that  $f$  is continuous and real-linear. Define  $\phi : X \rightarrow \mathbb{C}$  by  $\phi(x) = f(x) - if(ix)$ . Then  $\phi$  is continuous. Let  $x, y \in X$  and  $\lambda \in \mathbb{C}$ . Write  $\lambda = a + bi$ . Then

$$\begin{aligned} \phi(x + \lambda y) &= f(x + \lambda y) - if(ix + \lambda y) \\ &= f(x + ay + iby) - if(ix + iay - by) \\ &= f(x) + af(y) + bf(iy) - if(ix) - iaf(iy) + ibf(y) \\ &= [f(x) - if(ix)] + a[f(y) - if(iy)] + ib[f(y) - if(iy)] \\ &= \phi(x) + a\phi(y) + ibf(y) \\ &= \phi(x) + \lambda\phi(y) \end{aligned}$$

So  $\phi$  is linear and  $\phi \in X^*$ . Let  $\psi \in X^*$ . Suppose that  $f = \operatorname{Re} \psi$ . Then for each  $x \in X$ ,

$$\begin{aligned}\phi(x) &= f(x) - if(ix) \\ &= \operatorname{Re} \psi(x) - i \operatorname{Re} \psi(ix) \\ &= \operatorname{Re} \psi(x) - \operatorname{Re} i\psi(x) \\ &= \operatorname{Re} \psi(x) + \operatorname{Im} \psi(x) \\ &= \psi(x)\end{aligned}$$

So  $\psi = \phi$  and  $\phi$  is unique.

□

## 5.2. Sublinear Functionals.

**Definition 5.2.1.** Let  $X$  be a real vector space and  $p : X \rightarrow \mathbb{R}$ . Then  $p$  is said to be a **sublinear functional** if for each  $x, y \in X$ ,  $\lambda \geq 0$ ,

- (1)  $p(x + y) \leq p(x) + p(y)$
- (2)  $p(\lambda x) = \lambda p(x)$

**Exercise 5.2.2.** Let  $X$  be a vector space and  $p : X \rightarrow \mathbb{R}$  be a sublinear functional. Then  $p(0) = 0$ .

*Proof.* Set  $\lambda = 0$ . Then

$$\begin{aligned} 0 &= \lambda p(0) \\ &= p(\lambda 0) \\ &= p(0) \end{aligned}$$

□

*Proof.* Clear

□

**Exercise 5.2.3.** Let  $X$  be a vector space and  $p : X \rightarrow \mathbb{R}$  a sublinear functional. Then for each  $x, y \in X$

- (1)  $-p(-x) \leq p(x)$
- (2)  $-p(y - x) \leq p(x) - p(y) \leq p(x - y)$

*Proof.* Let  $x, y \in X$ .

- (1) We have

$$\begin{aligned} 0 &= p(0) \\ &= p(x - x) \\ &\leq p(x) + p(-x) \end{aligned}$$

So  $-p(-x) \leq p(x)$ .

- (2) We have

$$\begin{aligned} p(x) &= p(x - y + y) \\ &\leq p(x - y) + p(y) \end{aligned}$$

So  $p(x) - p(y) \leq p(x - y)$ . Switching  $x$  and  $y$  gives us  $p(y) - p(x) \leq p(y - x)$  and multiplying both sides by  $-1$  yields  $-p(y - x) \leq p(x) - p(y)$

Putting these two together, we see that

$$-p(y - x) \leq p(x) - p(y) \leq p(x - y)$$

□

### Theorem 5.2.4. Hahn-Banach Theorem for Sublinear Functionals

Let  $X$  be a vector space,  $p : X \rightarrow \mathbb{R}$  a sublinear functional,  $M \subset X$  a subspace and  $f : M \rightarrow \mathbb{R}$  a linear functional. If for each  $x \in M$ ,  $f(x) \leq p(x)$ , then there exists a linear functional  $F : X \rightarrow \mathbb{R}$  such that for each  $x \in X$ ,  $F(x) \leq p(x)$  and  $F|_M = f$ .

**Exercise 5.2.5.** Let  $X$  be a vector space and  $p : X \rightarrow \mathbb{R}$  a sublinear functional. Then there exists a linear functional  $F : X \rightarrow \mathbb{R}$  such that for each  $x \in X$ ,  $F(x) \leq p(x)$ .



*Proof.* Take  $M = \{0\}$  and  $f \equiv 0$  and apply the Hahn-Banach theorem.  $\square$

**Exercise 5.2.6. Equivalency of linearity (General Case)** Let  $X$  be a vector space and  $p : X \rightarrow \mathbb{R}$  a sublinear functional. Then the following are equivalent:

- (1) there exists a unique  $F \in X^*$  such that  $F \leq p$
- (2) for each  $x \in X$ ,  $-p(-x) = p(x)$
- (3)  $p$  is linear

**Hint:** If there exists  $x \in X$  such that  $-p(-x) \neq p(x)$ , define  $f_1, f_2 : \text{span}(x) \rightarrow \mathbb{R}$  by  $f_1(tx) = tp(x)$  and  $f_2(tx) = -tp(-x)$

*Proof.*

- (1)  $\implies$  (2):

Suppose that there exists a unique  $F \in X^*$  such that  $F \leq p$ . For the sake of contradiction, suppose that there exists  $x \in X$  such that  $-p(-x) \neq p(x)$ . Define  $f_1, f_2 : \text{span}(x) \rightarrow \mathbb{R}$  by

$$f_1(tx) = tp(x)$$

and

$$f_2(tx) = -tp(-x)$$

Let  $y \in \text{span}(x)$ . Then there exists  $t \in \mathbb{R}$  such that  $y = tx$ . Then for each  $k \in \mathbb{R}$ ,

$$\begin{aligned} f_1(ky) &= f_1(ktx) \\ &= ktp(x) \\ &= kf_1(tx) \\ &= kf_1(y) \end{aligned}$$

Similarly,  $f_2(ky) = kf_2(y)$  and so  $f_1, f_2 \in \text{span}(x)^*$ . If  $t \geq 0$ , then

$$\begin{aligned} f_1(y) &= f_1(tx) \\ &= tp(x) \\ &= p(tx) \\ &= p(y) \end{aligned}$$

If  $t < 0$ , then

$$\begin{aligned} f_1(y) &= f_1(tx) \\ &= tp(x) \\ &= -|t|p(x) \\ &= -p(|t|x) \\ &= -p(-tx) \\ &\leq p(tx) \\ &= p(y) \end{aligned}$$

So  $f_1 \leq p$  on  $\text{span}(x)$ . Similarly,  $f_2 \leq p$  on  $\text{span}(x)$ . The Hahn-Banach theorem implies that there exist  $F_1, F_2 \in X^*$  such that  $F_1, F_2 \leq p$  and  $F_1 = f_1, F_2 = f_2$  on

$\text{span}(x)$ . By the assumption of uniqueness,  $F_1 = F_2$ . This is a contradiction since

$$\begin{aligned} F_1(x) &= p(x) \\ &\neq -p(-x) \\ &= F_2(x) \end{aligned}$$

So for each  $x \in X$ ,  $-p(-x) = p(x)$ .

- (2)  $\Rightarrow$  (3):

Suppose that for each  $x \in X$ ,  $-p(-x) = p(x)$ . The previous exercise implies that there exists  $F \in X^*$  such that  $F \leq p$ . Let  $x \in X$ . Then

$$\begin{aligned} -F(x) &= F(-x) \\ &\leq p(-x) \\ &= -p(x) \end{aligned}$$

So  $p(x) \leq F(x)$  and  $p \leq F$ . Therefore  $p = F$  and  $p$  is linear.

- (3)  $\Rightarrow$  (1):

Suppose that  $p$  is linear. Let  $F \in X^*$ . Suppose that  $F \leq p$ . Let  $x \in X$ . Then as in the case for (2)  $\Rightarrow$  (3), we have that

$$\begin{aligned} -F(x) &= F(-x) \\ &\leq p(-x) \\ &= -p(x) \end{aligned}$$

which implies that  $p = F$ . So  $p$  is the unique linear function  $F \in X^*$  such that  $F \leq p$ .

□

### 5.3. Seminorms.

**Definition 5.3.1.** Let  $X$  be a vector space and  $p : X \rightarrow \mathbb{R}$ . Then  $p$  is said to be a **seminorm** if for each  $x, y \in X$ ,  $\lambda \in \mathbb{R}$ ,

- (1)  $p(x + y) \leq p(x) + p(y)$
- (2)  $p(\lambda x) = |\lambda|p(x)$

**Exercise 5.3.2.** Let  $X$  be a vector space and  $p : X \rightarrow \mathbb{R}$  be a seminorm, then  $p$  is a sublinear functional.

*Proof.* Clear □

**Exercise 5.3.3.** Let  $X$  be a vector space and  $\phi \in X^*$ . Then  $|\phi|$  is a seminorm on  $X$ .

*Proof.* Clear. □

**Exercise 5.3.4.** Let  $X, Y$  be a vector spaces,  $T \in L(X, Y)$  and  $p$  a seminorm on  $Y$ . Then  $p \circ T$  is a seminorm on  $X$ .

*Proof.* Clear. □

**Exercise 5.3.5.** Let  $X$  be a vector space and  $p : X \rightarrow \mathbb{R}$  be a seminorm. Then  $p \geq 0$ .

*Proof.* Let  $x \in X$ . Then

$$\begin{aligned} 0 &= p(0) \\ &= p(x - x) \\ &\leq p(x) + p(-x) \\ &= p(x) + p(x) \\ &= 2p(x) \end{aligned}$$

So  $p(x) \geq 0$ . □

**Exercise 5.3.6. Reverse Triangle Inequality:**

Let  $X$  be a vector space and  $p : X \rightarrow [0, \infty)$  be a seminorm on  $X$ . Then for each  $x, y \in X$ ,  $|p(x) - p(y)| \leq p(x - y)$ .

*Proof.* Let  $x, y \in X$ . Then

$$\begin{aligned} p(x) &= p(x - y + y) \\ &\leq p(x - y) + p(y) \end{aligned}$$

So  $p(x) - p(y) \leq p(x - y)$ . Similarly,  $p(y) \leq p(y - x) + p(x)$  and so  $p(x) - p(y) \leq p(x - y)$ . Therefore  $|p(x) - p(y)| \leq p(x - y)$ . □

**Exercise 5.3.7.** Let  $X$  be a vector space,  $p : X \rightarrow [0, \infty)$  a seminorm and  $\phi \in X^*$ . Then  $\phi \leq p$  iff  $|\phi| \leq p$ .

*Proof.* Suppose that  $\phi \leq p$ . Let  $x \in X$ . Then

$$\begin{aligned} -\phi(x) &= \phi(-x) \\ &\leq p(-x) \\ &= p(x) \end{aligned}$$

So  $-p(x) \leq \phi(x)$ . Hence  $-p \leq \phi \leq p$ . Thus  $|\phi| \leq p$ .

Conversely, if  $|\phi| \leq p$ , then clearly  $\phi \leq p$ .  $\square$

**Definition 5.3.8.** Let  $X$  be a vector space and  $p : X \rightarrow [0, \infty)$  be a seminorm on  $X$ . We define the **kernel of  $p$** , denoted  $\ker p$ , by  $\ker p = p^{-1}(\{0\})$ .

**Exercise 5.3.9.** Let  $X$  be a vector space and  $p : X \rightarrow [0, \infty)$  a seminorm. Then  $\ker p$  is a subspace of  $X$ .

*Proof.* Let  $x, y \in \ker p$  and  $\lambda \in \mathbb{C}$ . Then  $p(x) = p(y) = 0$ . Thus

$$\begin{aligned} p(x + \lambda y) &\leq p(x) + p(\lambda y) \\ &= p(x) + |\lambda|p(y) \\ &= 0 \end{aligned}$$

So  $x + \lambda y \in \ker p$  and  $\ker p$  is a subspace.  $\square$

**Definition 5.3.10.** Let  $X$  be a vector space and  $p : X \rightarrow [0, \infty)$  a seminorm on  $X$ . We define the **norm induced by  $p$** , denoted  $\bar{p} : X/\ker p \rightarrow [0, \infty)$ , by

$$\bar{p}(\bar{x}) = p(x)$$

**Exercise 5.3.11.** Let  $X$  be a vector space and  $p : X \rightarrow [0, \infty)$  a seminorm on  $X$ . Then  $\bar{p} : X/\ker p \rightarrow [0, \infty)$  is well defined and a norm.

*Proof.* Let  $x, y \in X$ . Suppose that  $\bar{x} = \bar{y}$ . Then there exists  $n \in \ker p$  such that  $x = y + n$ . Therefore,

$$\begin{aligned} \bar{p}(\bar{x}) &= p(x) \\ &= p(y + n) \\ &\leq p(y) + p(n) \\ &= p(y) \\ &= \bar{p}(\bar{y}) \end{aligned}$$

and

$$\begin{aligned} \bar{p}(\bar{y}) &= p(y) \\ &= p(x - n) \\ &\leq p(x) + p(n) \\ &= p(x) \\ &= \bar{p}(\bar{x}) \end{aligned}$$

So  $\bar{p}(\bar{x}) = \bar{p}(\bar{y})$  and  $\bar{p} : X/\ker p \rightarrow [0, \infty)$  is well defined. Let  $x \in X$ . Suppose that  $\bar{x} = \bar{0}$ . Then there exists  $n \in \ker p$  such that  $x = n$ . Therefore

$$\begin{aligned} \bar{p}(\bar{x}) &= p(x) \\ &= p(n) \\ &= 0 \end{aligned}$$

So  $\bar{p}$  is a norm.  $\square$

**Definition 5.3.12.** Let  $X$  be a vector space,  $p : X \rightarrow [0, \infty)$  a seminorm on  $X$ ,  $x \in X$  and  $r > 0$ . We define the

- **open semiball of  $p$  at  $x$  of radius  $r$** , denoted  $B_p(x, r)$ , by

$$B_p(x, r) = \{y \in X : p(x - y) < r\}$$

- **closed semiball of  $p$  at  $x$  of radius  $r$** , denoted  $\bar{B}_p(x, r)$ , by

$$\bar{B}_p(x, r) = \{y \in X : p(x - y) \leq r\}$$

**Exercise 5.3.13.** Let  $X$  be a vector space,  $p : X \rightarrow [0, \infty)$  a seminorm on  $X$ ,  $x \in X$  and  $r > 0$ . Then  $B_p(x, r) = x + rB_p(0, 1)$ .

*Proof.* Let  $y \in B_p(x, r)$ . Then

$$\begin{aligned} p(r^{-1}(y - x)) &= r^{-1}p(y - x) \\ &< r^{-1}r \\ &= 1 \end{aligned}$$

So  $r^{-1}(y - x) \in B_p(0, 1)$ . By definition, there exists  $u \in B_p(0, 1)$  such that  $r^{-1}(y - x) = u$ , which implies that

$$\begin{aligned} y &= x + ru \\ &\in x + rB_p(0, 1) \end{aligned}$$

Conversely, let  $y \in x + rB_p(0, 1)$ . By definition, there exists  $u \in B_p(0, 1)$  such that  $y = x + ru$ . Then

$$\begin{aligned} p(y - x) &= p(ru) \\ &= rp(u) \\ &< r \end{aligned}$$

So  $y \in B_p(x, r)$  □

**Exercise 5.3.14.** Let  $X$  be a vector space and  $p, q : X \rightarrow [0, \infty)$  seminorms on  $X$ . Then  $p \leq q$  iff  $B_q(0, 1) \subset B_p(0, 1)$ .

*Proof.* Suppose that  $p \leq q$ . Let  $x \in B_q(0, 1)$ . Then

$$\begin{aligned} p(x) &\leq q(x) \\ &< 1 \end{aligned}$$

So  $x \in B_p(0, 1)$ .

Conversely, suppose that  $B_q(0, 1) \subset B_p(0, 1)$ . Let  $x \in X$ . If  $p(x) = 0$ , then  $p(x) \leq q(x)$ . Suppose that  $p(x) > 0$ . For the sake of contradiction, suppose that  $p(x) > q(x)$ . Then

$$\begin{aligned} q\left(\frac{x}{p(x)}\right) &= \frac{q(x)}{p(x)} \\ &< 1 \end{aligned}$$

Therefore,  $x/p(x) \in B_q(0, 1) \subset B_p(0, 1)$ . By definition,

$$\begin{aligned} \frac{p(x)}{p(x)} &= p\left(\frac{x}{p(x)}\right) \\ &< 1 \end{aligned}$$

which is a contradiction. So  $p(x) \leq q(x)$ . Since  $x \in X$  is arbitrary,  $p \leq q$ . □

**Exercise 5.3.15.** Let  $X$  be a topological vector space and  $p : X \rightarrow [0, \infty)$  a continuous seminorm. Then

- (1)  $B_p(0, 1)$  is open
- (2)  $\bar{B}_p(0, 1)$  is closed

*Proof.*

- (1) Let  $(x_\alpha)_{\alpha \in A}$  be a net in  $B_p(0, 1)^c$  and  $x \in X$ . Suppose that  $x_\alpha \rightarrow x$ . Then  $p(x_\alpha) \rightarrow p(x)$ . Since for each  $\alpha \in A$ ,  $p(x_\alpha) \geq 1$ ,  $p(x) \geq 1$ . Hence  $x \in B_p(0, 1)^c$ . So  $B_p(0, 1)^c$  is closed which implies that  $B_p(0, 1)$  is open.
- (2) Let  $(x_\alpha)_{\alpha \in A}$  be a net in  $\bar{B}_p(0, 1)$  and  $x \in X$ . Suppose that  $x_\alpha \rightarrow x$ . Then  $p(x_\alpha) \rightarrow p(x)$ . Since for each  $\alpha \in A$ ,  $p(x_\alpha) \leq 1$ ,  $p(x) \leq 1$ . Hence  $x \in \bar{B}_p(0, 1)$ . So  $\bar{B}_p(0, 1)$  is closed.

□

**Exercise 5.3.16.** Let  $X$  be a topological vector space and  $p : X \rightarrow [0, \infty)$  a seminorm. Then the following are equivalent:

- (1)  $p$  is continuous
- (2)  $B_p(0, 1)$  is open
- (3)  $\bar{B}_p(0, 1) \in \mathcal{N}_0$
- (4)  $p$  is continuous at 0.

*Proof.*

- (1)  $\implies$  (2):  
Clear from previous exercise.
- (2)  $\implies$  (3):  
Clear since  $B_p(0, 1) \subset \bar{B}_p(0, 1)$ .
- (3)  $\implies$  (4):  
Let  $(x_\alpha)_{\alpha \in A} \subset X$  be a net. Suppose that  $x_\alpha \rightarrow 0$ . Let  $U \subset \mathbb{R}$ . Suppose that  $U \in \mathcal{N}_0$ . Then there exists  $\epsilon > 0$  such that  $\bar{B}(0, \epsilon) \subset U$ . Since the map  $f_\epsilon : X \rightarrow X$  defined by  $f_\epsilon(x) = \epsilon x$  is a homeomorphism,  $\bar{B}_p(0, \epsilon) = \epsilon \bar{B}_p(0, 1) \in \mathcal{N}_0$ . Hence there exists  $\beta \in A$  such that for each  $\alpha \geq \beta$ ,  $x_\alpha \in \bar{B}_p(0, \epsilon)$ . Let  $\alpha \in A$ . Suppose that  $\alpha \geq \beta$ . By definition,  $p(x_\alpha) \leq \epsilon$ . So  $p(x_\alpha) \in \bar{B}(0, \epsilon) \subset U$ . Hence  $(p(x_\alpha))_{\alpha \in A}$  is eventually in  $U$ . Since  $U \in \mathcal{N}_0$  is arbitrary,  $p(x_\alpha) \rightarrow 0$ . So  $p$  is continuous at 0.
- (4)  $\implies$  (1):  
Let  $(x_\alpha)_{\alpha \in A} \subset X$  be a net and  $x \in X$ . Suppose that  $x_\alpha \rightarrow x$ . Then  $x_\alpha - x \rightarrow 0$ . Therefore  $p(x_\alpha - x) \rightarrow 0$ . The reverse triangle inequality implies that  $p(x_\alpha) \rightarrow p(x)$ . So  $p$  is continuous.

□

**Exercise 5.3.17.** Let  $X$  be a topological vector space and  $p : X \rightarrow [0, \infty)$  a seminorm. Then  $p$  is continuous iff there exists a continuous seminorm  $q : X \rightarrow [0, \infty)$  such that  $p \leq q$ .

*Proof.* Suppose that  $p$  is continuous. Set  $q = p$ . Then  $q$  is a continuous and  $p \leq q$ . Conversely, suppose that there exists a continuous seminorm  $q : X \rightarrow [0, \infty)$  such that  $p \leq q$ .

Then  $\bar{B}_q(0, 1) \subset \bar{B}_p(0, 1)$ . The previous exercise tells us that

$$\begin{aligned} q \text{ is continuous} &\iff \bar{B}_q(0, 1) \in \mathcal{N}_0 \\ &\implies \bar{B}_p(0, 1) \in \mathcal{N}_0 \\ &\iff p \text{ is continuous} \end{aligned}$$

□

**Theorem 5.3.18. Hahn-Banach Theorem for Seminorms**

Let  $X$  be a vector space,  $p : X \rightarrow \mathbb{R}$  a seminorm,  $M \subset X$  a subspace and  $f : M \rightarrow \mathbb{C}$  a linear functional. If for each  $x \in M$ ,  $|f(x)| \leq p(x)$ , then there exists a linear functional  $F : X \rightarrow \mathbb{C}$  such that for each  $x \in X$ ,  $|F(x)| \leq p(x)$  and  $F|_M = f$ .

#### 5.4. Minkowski Functionals.

**Definition 5.4.1.** Let  $X$  be a vector space and  $A \subset X$ . Then  $A$  is said to be **convex** if for each  $x, y \in A$ ,  $t \in [0, 1]$ ,  $tx + (1 - t)y \in A$ .

**Exercise 5.4.2.** Let  $X$  be a vector space and  $\mathcal{A} \subset \mathcal{P}(X)$ , Suppose that for each  $A \in \mathcal{A}$ ,  $A$  is convex. Then

$$\bigcap_{A \in \mathcal{A}} A$$

is convex.

*Proof.* Let  $x, y \in \bigcap_{A \in \mathcal{A}} A$  and  $t \in [0, 1]$ . Then for each  $A \in \mathcal{A}$ ,  $x, y \in A$ . Let  $A \in \mathcal{A}$ . Since  $A$  is convex,  $tx + (1 - t)y \in A$ . Since  $A \in \mathcal{A}$  is arbitrary,  $tx + (1 - t)y \in \bigcap_{A \in \mathcal{A}} A$ . So  $\bigcap_{A \in \mathcal{A}} A$  is convex.  $\square$

**Definition 5.4.3.** Let  $X$  be a vector space and  $A \subset X$ . Set

$$\mathcal{S} = \{S \subset X : S \text{ is convex and } A \subset S\}$$

We define the **convex hull of  $A$** , denoted  $\text{conv } A$ , by

$$\text{conv } A = \bigcap_{S \in \mathcal{S}} S$$

**Note 5.4.4.** We may think of  $\text{conv } A$  as the smallest convex set containing  $A$ .

**Definition 5.4.5.** Let  $X$  be a vector space,  $A \subset X$  and  $x \in X$ . Then  $x$  is said to be a **convex combinations of elements of  $A$**  if there exist  $(a_j)_{j=1}^n \subset A$  and  $(t_j)_{j=1}^n \subset [0, 1]$  such that  $x = \sum_{j=1}^n t_j a_j$  and  $\sum_{j=1}^n t_j = 1$ . We define  $C_A \subset X$  by

$$C_A = \{x \in X : x \text{ is a convex combination of elements of } A\}$$

**Exercise 5.4.6.** Let  $X$  be a vector space and  $A \subset X$ . Then

- (1)  $A \subset C_A$
- (2)  $C_A$  is convex

*Proof.*

- (1) Let  $x \in A$ , then

$$\begin{aligned} x &= 1x \\ &\in C_A \end{aligned}$$

So  $A \subset C_A$ .

- (2) Let  $x, y \in C_A$ . and  $\lambda \in [0, 1]$ . Then there exist  $(a_i)_{i=1}^n, (b_j)_{j=1}^m \subset A$  and  $(s_i)_{i=1}^n, (t_j)_{j=1}^m \subset [0, 1]$  such that  $x = \sum_{i=1}^n s_i a_i$  and  $y = \sum_{j=1}^m t_j b_j$ . Then

$$\begin{aligned} \lambda x + (1 - \lambda)y &= \lambda \left[ \sum_{i=1}^n s_i a_i \right] + (1 - \lambda) \left[ \sum_{j=1}^m t_j b_j \right] \\ &= \sum_{i=1}^n \lambda s_i a_i + \sum_{j=1}^m (1 - \lambda) t_j b_j \end{aligned}$$



Since

- (a) for each  $i \in \{1, \dots, n\}$  and  $j \in \{1, \dots, m\}$ , we have that  $\lambda s_i \in [0, 1]$  and  $(1 - \lambda)t_j \in [0, 1]$   
 (b)

$$\begin{aligned} \sum_{i=1}^n \lambda s_i + \sum_{j=1}^m (1 - \lambda)t_j &= \lambda \sum_{i=1}^n s_i + (1 - \lambda) \sum_{j=1}^m t_j \\ &= \lambda + (1 - \lambda) \\ &= 1 \end{aligned}$$

we have that  $\lambda x + (1 - \lambda)y \in C_A$ . So  $C_A$  is convex. □

**Exercise 5.4.7.** Let  $X$  be a vector space and  $A \subset X$ . Let  $(a_j)_{j=1}^n \subset A$  and  $(t_j)_{j=1}^n \subset [0, 1]$ .

Suppose that  $\sum_{j=1}^n t_j = 1$ . If  $A$  is convex, then  $\sum_{j=1}^n t_j a_j \in A$ .

**Hint:** proceed by induction on  $n$

*Proof.* Suppose that  $A$  is convex. If  $n = 2$ , then by definition,  $\sum_{j=1}^n t_j a_j \in A$ .

Suppose that the claim is true for  $n - 1$ . Since  $\sum_{j=1}^n t_j = 1$ , then there  $k \in \{1, \dots, n\}$  such that  $t_k > 0$ . Choose  $l \in \{1, \dots, n\}$  such that  $l \neq k$ . Set  $S = \{1, \dots, n\} \setminus \{l\}$ . Then  $1 - t_l > 0$  and

$$\begin{aligned} x &= \sum_{j=1}^n t_j a_j \\ &= t_l a_l + \sum_{j \in S} t_j a_j \\ &= t_l a_l + (1 - t_l) \sum_{j \in S} \frac{t_j}{1 - t_l} a_j \end{aligned}$$

Since

$$\begin{aligned} \sum_{j \in S} \frac{t_j}{1 - t_l} &= \frac{1 - t_l}{1 - t_l} \\ &= 1 \end{aligned}$$

our induction hypothesis implies that

$$\sum_{j \in S} \frac{t_j}{1 - t_l} a_j \in A$$

Since  $A$  is convex, by definition we have that

$$\begin{aligned} x &= t_l a_l + (1 - t_l) \left[ \sum_{j \in S} \frac{t_j}{1 - t_l} a_j \right] \\ &\in A \end{aligned}$$

□

**Exercise 5.4.8.** Let  $X$  be a vector space and  $A \subset X$ . Then

$$\text{conv } A = C_A$$

*Proof.* Since  $A \subset C_A$  and  $C_A$  is convex,  $\text{conv } A \subset C_A$ .

Conversely, Let  $x \in C_A$ . Then there exist  $(a_j)_{j=1}^n \subset A$  and  $(t_j)_{j=1}^n \subset [0, 1]$  such that  $x = \sum_{j=1}^n t_j a_j$  and  $\sum_{j=1}^n t_j = 1$ . Since  $A \subset \text{conv } A$  and  $\text{conv } A$  is convex, the previous exercise implies that  $x \in \text{conv } A$ . So  $C_A \subset \text{conv } A$ . Hence  $\text{conv } A = C_A$ .  $\square$

**Exercise 5.4.9.** Let  $X$  be a vector space and  $A, B \subset X$  convex and  $\lambda \in \mathbb{C}$ . Then

- (1)  $A + B$  is convex
- (2)  $\lambda A$  is convex

*Proof.*

- (1) Let  $x, y \in A + B$  and  $t \in [0, 1]$ . Then there exist  $a_x, a_y \in A$ ,  $b_x, b_y \in B$  such that  $x = a_x + b_x$  and  $y = a_y + b_y$ . Since  $A$  and  $B$  are convex,  $ta_x + (1-t)a_y \in A$  and  $tb_x + (1-t)b_y \in B$ . Hence

$$\begin{aligned} tx + (1-t)y &= ta_x + tb_x + (1-t)a_y + (1-t)b_y \\ &= [ta_x + (1-t)a_y] + [tb_x + (1-t)b_y] \\ &\in A + B \end{aligned}$$

So  $A + B$  is convex.

- (2) Let  $x, y \in \lambda A$  and  $t \in [0, 1]$ . Then there exist  $a_x, a_y \in A$  such that  $x = \lambda a_x$  and  $y = \lambda a_y$ . Since  $A$  is convex,  $ta_x + (1-t)a_y \in A$ . Therefore

$$\begin{aligned} tx + (1-t)y &= t\lambda a_x + (1-t)\lambda a_y \\ &= \lambda[ta_x + (1-t)a_y] \\ &\in \lambda A \end{aligned}$$

So  $\lambda A$  is convex.  $\square$

**Definition 5.4.10.** Let  $X$  be a vector space and  $A \subset X$ . Then  $A$  is said to be **balanced** if for each  $x \in A$ ,  $c \in \mathbb{C}$ ,  $|c| \leq 1$  implies that  $cx \in A$ .

**Exercise 5.4.11.** Let  $X$  be a vector space and  $\mathcal{A} \subset \mathcal{P}(X)$ , Suppose that for each  $A \in \mathcal{A}$ ,  $A$  is balanced. Then

$$\bigcup_{A \in \mathcal{A}} A$$

is balanced.

*Proof.* Let  $x \in \bigcap_{A \in \mathcal{A}} A$  and  $r \in \mathbb{C}$ . Suppose that  $|r| \leq 1$ . Then there exists  $B \in \mathcal{A}$  such that  $x \in B$ . Since  $B$  is balanced,

$$\begin{aligned} rx &\in B \\ &\subset \bigcap_{A \in \mathcal{A}} A \end{aligned}$$

So  $\bigcap_{A \in \mathcal{A}} A$  is balanced.  $\square$

**Definition 5.4.12.** Let  $X$  be a vector space and  $A \subset X$ . We define the **balanced hull** of  $A$ , denoted  $\text{bal } A$ , by

$$\text{bal } A = \bigcup_{\substack{r \in \mathbb{C} \\ |r| \leq 1}} rA$$

**Exercise 5.4.13.** Let  $X$  be a vector space and  $A \subset X$ . Then  $\text{bal } A$  is balanced.

*Proof.* Let  $x \in \text{bal } A$  and  $r \in \mathbb{C}$ . Suppose that  $|r| \leq 1$ . By definition, there exists  $s \in \mathbb{C}$  and  $a \in A$  such that  $|s| \leq 1$  and  $x = sa$ . Then

$$\begin{aligned} |rs| &= |r||s| \\ &\leq 1 \end{aligned}$$

which implies that

$$\begin{aligned} rx &= rsa \\ &\in rsA \\ &\subset \bigcup_{\substack{q \in \mathbb{C} \\ |q| \leq 1}} qA \\ &= \text{bal } A \end{aligned}$$

So  $\text{bal } A$  is balanced. □

**Note 5.4.14.** We may think of  $\text{bal } A$  as the smallest balanced set containing  $A$ .

**Exercise 5.4.15.** Let  $X$  be a vector space and  $A \subset X$ . Suppose that  $A \neq \emptyset$ . If  $A$  is balanced, then  $0 \in A$ .

*Proof.* Clear by definition. □

**Exercise 5.4.16.** Let  $X$  be a vector space,  $A \subset X$ ,  $x \in X$  and  $\lambda \in \mathbb{C}$ . Suppose that  $A$  is balanced. Then  $\lambda x \in A$  iff  $|\lambda|x \in A$ .

*Proof.* If  $\lambda = 0$ , then the claim is clearly true. Suppose that  $\lambda \neq 0$ . Set  $s = \text{sgn}(\lambda)$ . Suppose that  $\lambda x \in A$ . Since  $A$  is balanced and  $|s| = |s^{-1}| = 1$ ,

$$\begin{aligned} |\lambda|x &= s^{-1}\lambda x \\ &\in A \end{aligned}$$

Conversely, suppose that  $|\lambda|x \in A$ . Then

$$\begin{aligned} \lambda x &= s|\lambda|x \\ &\in A \end{aligned}$$

□

**Exercise 5.4.17.** Let  $X$  be a vector space and  $A \subset X$ . If  $A$  is balanced, then  $\text{conv } A$  is balanced.

*Proof.* Suppose that  $A$  is balanced. Let  $x \in \text{conv } A$  and  $r \in \mathbb{C}$ . Suppose that  $|r| \leq 1$ . Then there exist  $(a_j)_{j=1}^n \subset A$  and  $(t_j)_{j=1}^n \subset [0, 1]$  such that  $x = \sum_{j=1}^n t_j a_j$  and  $\sum_{j=1}^n t_j = 1$ . Since  $A$  is

balanced, for each  $j \in \{1, \dots, n\}$ ,

$$\begin{aligned} ra_j &\in A \\ &\subset \text{conv } A \end{aligned}$$

Since  $\text{conv } A$  is convex, we have that

$$\begin{aligned} rx &= r \sum_{j=1}^n t_j a_j \\ &= \sum_{j=1}^n t_j ra_j \\ &\in \text{conv } A \end{aligned}$$

Hence  $\text{conv } A$  is balanced.. □

**Definition 5.4.18.** Let  $X$  be a vector space and  $A \subset X$ . Then  $A$  is said to be **absorbing** if for each  $x \in X$ , there exists  $r > 0$  such that for each  $c \in \mathbb{R}$ ,  $|c| \geq r$  implies that  $x \in cA$ .

**Exercise 5.4.19.** Let  $X$  be a topological vector space and  $A \in \mathcal{N}_0$ . Then  $A$  is absorbing.

*Proof.* Let  $x \in A$ . For the sake of contradiction, suppose that for each  $r > 0$ , there exists  $c \in \mathbb{R}$  such that  $|c| \geq r$  and  $c^{-1}x \in A^c$ . Then there exists a sequence  $(c_n)_{n \in \mathbb{N}} \subset \mathbb{R}$  such that for each  $n \in \mathbb{N}$ ,  $c_n \geq n$  and  $c_n^{-1}x \in A^c$ . Since  $c_n^{-1} \rightarrow 0$ ,  $c_n^{-1}x \rightarrow 0$ . Since  $A \in \mathcal{N}_0$ ,  $(c_n^{-1}x)_{n \in \mathbb{N}}$  is eventually in  $A$ . This is a contradiction. So there exists  $r > 0$  such that for each  $c \in \mathbb{R}$ ,  $|c| \geq r$  implies that  $x \in cA$ . Hence  $A$  is absorbing. □

**Exercise 5.4.20.**

*Proof.* □

**Definition 5.4.21.** Let  $X$  be a vector space and  $A \subset X$ . For  $x \in X$ , set

$$T_x^A = \{t > 0 : x \in tA\}$$

We define the **Minkowski functional**, denoted  $p_A : X \rightarrow [0, \infty]$ , by

$$p_A(x) = \inf T_x^A$$

**Exercise 5.4.22.** Let  $X$  be a vector space and  $A \subset X$ . Suppose that  $A$  is convex, absorbing and  $0 \in A$ . Then

- (1)  $p_A : X \rightarrow [0, \infty)$
- (2)  $p(0) = 0$
- (3)  $p_A$  is a sublinear functional on  $X$

*Proof.*

- (1) Since  $A$  is absorbing, there exists  $r > 0$  such that for each  $c \in \mathbb{R}$ ,  $|c| \geq r$  implies that  $x \in cA$ . Therefore  $p_A(x) \leq |c|$  and  $p_A : X \rightarrow [0, \infty)$ .
- (2) Since  $0 \in A$ ,

$$\begin{aligned} p_A(0) &= \inf T_0^A \\ &= 0 \end{aligned}$$

- (3) • Let  $\epsilon > 0$ . Choose  $t_x \in T_x^A$  and  $t_y \in T_y^A$  such that  $t_x < p_A(x) + \epsilon/2$  and  $t_y < p_A(y) + \epsilon/2$ . By definition,  $t_x^{-1}x, t_y^{-1}y \in A$ . Set  $\theta = t_x(t_x + t_y)^{-1} \in (0, 1)$ . Since  $A$  is convex,

$$\begin{aligned} (t_x + t_y)^{-1}(x + y) &= (t_x + t_y)^{-1}x + (t_x + t_y)^{-1}y \\ &= \theta t_x^{-1}x + (1 - \theta)t_y^{-1}y \\ &\in A \end{aligned}$$

Therefore,  $t_x + t_y \in T_{x+y}^A$  and

$$\begin{aligned} p_A(x + y) &\leq t_x + t_y \\ &< p_A(x) + \frac{\epsilon}{2} + p_A(y) + \frac{\epsilon}{2} \\ &= p_A(x) + p_A(y) + \epsilon. \end{aligned}$$

Since  $\epsilon > 0$  is arbitrary,  $p_A(x + y) \leq p_A(x) + p_A(y)$ .

- If  $\lambda = 0$ , then

$$\begin{aligned} p_A(\lambda x) &= p_A(0) \\ &= 0 \\ &= |\lambda|p_A(x) \end{aligned}$$

Suppose that  $\lambda > 0$ . Let  $t > 0$ . Then

$$\begin{aligned} p_A(\lambda x) &= \inf\{t > 0 : \lambda x \in tA\} \\ &= \inf\{t > 0 : x \in \lambda^{-1}tA\} \\ &= \inf\{\lambda s > 0 : x \in sA\} \\ &= \lambda \inf\{s > 0 : x \in sA\} \\ &= \lambda p_A(x) \end{aligned}$$

So  $p$  is a sublinear functional on  $X$ . □

**Exercise 5.4.23.** Let  $X$  be a vector space and  $A \subset X$ . Suppose that  $A$  is convex, absorbing and  $0 \in A$ . Then  $p_A^{-1}[0, 1) \subset A$ .

*Proof.* Let  $x \in p_A^{-1}[0, 1)$ . Then  $p_A(x) < 1$ . By definition, there exists  $t \in (0, 1)$  such that  $x \in tA$ . Thus  $t^{-1}x \in A$ . Since  $0 \in A$  and  $A$  is convex, we have that

$$\begin{aligned} x &= t(t^{-1}x) + (1 - t)0 \\ &\in A \end{aligned}$$

Since  $x \in p_A^{-1}[0, 1)$  is arbitrary,  $p_A^{-1}[0, 1) \subset A$ . □

**Exercise 5.4.24.** Let  $X$  be a topological vector space and  $A \subset X$ . Suppose that  $A$  is open, convex, and  $0 \in A$ . Then  $p_A^{-1}[0, 1) = A$ .

**Hint:** for  $x \in A$ , consider the sequence  $(1 + 1/n)x$

*Proof.* Since  $A$  is open and  $0 \in A$ ,  $A \in \mathcal{N}_0$  which implies that  $A$  is absorbing. The previous exercise implies that  $p_A^{-1}[0, 1) \subset A$ .

Conversely, let  $x \in A$ . Since  $A$  is open,  $A \in \mathcal{N}_x$ . Since  $1 + 1/n \rightarrow 1$ ,  $(1 + 1/n)x \rightarrow x$ .

Therefore, there exists  $N \in \mathbb{N}$  such that for each  $n \in \mathbb{N}$ ,  $n \geq N$  implies that  $(1 + 1/n)x \in A$ . In particular,  $x \in (1 + 1/N)^{-1}A$ . Hence  $(1 + 1/N)^{-1} \in T_x^A$  and

$$\begin{aligned} p_A(x) &\leq (1 + 1/N)^{-1} \\ &< 1 \end{aligned}$$

So  $x \in p_A^{-1}[0, 1)$  and  $A \subset B_{p_A}(0, 1)$ . □

**Exercise 5.4.25.** Let  $X$  be a topological vector space,  $A \subset X$  and  $x_0 \in A^c$ . Suppose that  $A$  is convex,  $A \in \mathcal{N}_0$  and  $A$  is open. Then there exists  $F \in X^*$  such that  $\operatorname{Re} F(x_0) = 1$  and  $\operatorname{Re} F|_A < 1$ .

**Hint:** Assume  $X$  is real.

- (1) **Existence:** Consider a special  $f \in (\mathbb{R}x_0)^*$  and use  $p_A$  to apply the Hahn-Banach theorem.
- (2) **Continuity:** for  $\epsilon > 0$ , consider the neighborhood  $U_\epsilon = \epsilon A \cap -\epsilon A$

*Proof.* Assume that  $X$  is real.

- (1) Define  $f \in (\mathbb{R}x_0)^*$  by  $f(tx_0) = t$ . Then  $f(x_0) = 1$ . Since  $A \in \mathcal{N}_0$ ,  $0 \in A$  and a previous exercise implies that  $A$  is absorbing. Since  $A$  is convex, absorbing and  $0 \in A$ ,  $p_A : X \rightarrow [0, \infty)$  is a sublinear functional on  $X$ . Since  $x_0 \in A^c$ , the previous exercise implies that  $1 \leq p_A(x_0)$ . Let  $x \in \mathbb{R}x_0$ . Then there exists  $t \in \mathbb{R}$  such that  $x = tx_0$ .

- If  $t \geq 0$ , then

$$\begin{aligned} f(x) &= t \\ &\leq tp_A(x_0) \\ &= p_A(tx_0) \\ &= p_A(x) \end{aligned}$$

- If  $t < 0$ , then  $-t > 0$  and an exercise from the section on sublinear functionals implies that

$$\begin{aligned} f(x) &= t \\ &= -(-t) \\ &\leq p_A(x) \end{aligned}$$

So  $f \leq p_A$  on  $\mathbb{R}x_0$ . The Hahn-Banach theorem implies that there exists  $F : X \rightarrow \mathbb{R}$  such that  $F$  is linear,  $F|_{\mathbb{R}x_0} = f$  and  $F \leq p_A$ . The previous exercise implies that  $p_A|_A < 1$ . Hence  $F|_A < 1$ .

- (2) Let  $V \in \mathcal{N}_{0\mathbb{R}}$ . Choose  $\epsilon > 0$  such that  $B(0, \epsilon) \subset V$ . Set  $U_\epsilon = \epsilon A \cap -\epsilon A$ . Then  $U_\epsilon \in \mathcal{N}_0$ . Let  $u \in U_\epsilon$ . Then  $\epsilon^{-1}u, -\epsilon^{-1}u \in A$ . A previous exercise implies that  $p_A^{-1}([0, 1)) = A$ . Hence

$$\begin{aligned} \epsilon^{-1}F(u) &= F(\epsilon^{-1}u) \\ &\leq p_A(\epsilon^{-1}u) \\ &< 1 \end{aligned}$$

So  $F(u) < \epsilon$ . Similarly,  $F(-u) < \epsilon$ . So  $-\epsilon < F(u) < \epsilon$  and

$$\begin{aligned} F(U_\epsilon) &\subset B(0, \epsilon) \\ &\subset V \end{aligned}$$

Since  $V \in \mathcal{N}_{0_{\mathbb{R}}}$  is arbitrary,  $F$  is continuous at 0. Since  $F$  is linear and  $F$  is continuous at 0,  $F$  is continuous. Hence  $F \in X^*$ .

If  $X$  is complex, then the previous part implies that there exists  $G : X \rightarrow \mathbb{R}$  such that  $G$  is continuous, real-linear,  $G(x_0) = 1$  and  $G|_A < 1$ . A previous exercise implies that there exists a unique  $F \in X^*$  such that  $\operatorname{Re} F = G$ .  $\square$

**Exercise 5.4.26. Hahn-Banach Separation Theorem 1:**

Let  $X$  be a topological vector space and  $A, B \subset X$ . Suppose that  $A, B$  are nonempty, convex and disjoint. If  $A$  is open, then there exists  $\phi \in X^*$  and  $c \in \mathbb{R}$  such that for each  $x \in A, y \in B$ ,

$$\operatorname{Re} \phi(x) < c \leq \operatorname{Re} \phi(y)$$

**Hint:** Assume  $X$  is real.

- (1) Choose  $a_0 \in A$  and  $b_0 \in B$  and set  $x_0 = b_0 - a_0$  and  $C = A - B + x_0$ . Then there exists  $\phi \in X^*$  such that  $\phi(x_0) = 1$  and  $\phi|_C < 1$ .
- (2) For each  $a \in A, b \in B, \phi(a) < \phi(b)$ . Set  $c = \sup_{a \in A} \phi(a)$ . Since  $\phi$  is not constant,  $\phi$  is open.

*Proof.* Assume  $X$  is real.

- (1) Since  $A, B$  are nonempty, there exist  $a_0 \in A$  and  $b_0 \in B$ . Set  $x_0 = b_0 - a_0$ . Previous exercises imply that  $A - B$  is open and convex. Set  $C = A - B + x_0$ . Then  $C$  is open and convex. Since

$$\begin{aligned} 0 &= a_0 - b_0 + x_0 \\ &\in C \end{aligned}$$

$C \in \mathcal{N}_0$ . For the sake of contradiction, suppose that  $x_0 \in C$ . Then there exist  $a \in A, b \in B$  such that  $x_0 = a - b + x_0$ . This implies that  $a = b$ . This is a contradiction since  $A \cap B = \emptyset$ . Hence  $x_0 \notin C$ . The previous exercise implies that there exists a  $\phi \in X^*$  such that  $\phi(x_0) = 1$  and  $\phi|_C < 1$ .

- (2) Let  $x \in A$  and  $y \in B$ . Then

$$\begin{aligned} \phi(a) - \phi(b) + 1 &= \phi(a) - \phi(b) + \phi(x_0) \\ &= \phi(a - b + x_0) \\ &< 1 \end{aligned}$$

So  $\phi(a) < \phi(b)$ . Set  $c = \sup_{a \in A} \phi(a)$ . Since  $A$  is open and  $\phi \in X^*$  is open. Thus for each  $x \in A, y \in B$ ,

$$\phi(x) < c \leq \phi(y)$$

If  $X$  is complex, then the previous part implies that there exists  $f : X \rightarrow \mathbb{R}$  and  $c \in \mathbb{R}$  such that  $f$  is continuous, real-linear and for each  $x \in A$  and  $y \in B$ ,

$$f(x) < c \leq f(y)$$

A previous exercise implies that there exists a unique  $\phi \in X^*$  such that  $\operatorname{Re} \phi = f$ .  $\square$

**Definition 5.4.27.** Let  $X$  be a vector space and  $A \subset X$ . Then  $A$  is said to be an **absorbing disk** if  $A$  is convex, absorbing and balanced.

**Exercise 5.4.28.** Let  $X$  be a vector space,  $p : X \rightarrow [0, \infty)$  a seminorm on  $X$  and  $r > 0$ . Then  $B_p(0, r)$  is an absorbing disk.

*Proof.*

(1) Let  $a, b \in B_p(0, r)$  and  $t \in [0, 1]$ . Then  $p(a - x) < r$  and  $p(b) < r$ . So

$$\begin{aligned} p([ta + (1 - t)b]) &\leq p(ta + p((1 - t)b)) \\ &= tp(a) + (1 - t)p(b) \\ &< tr + (1 - t)r \\ &= r \end{aligned}$$

So  $ta + (1 - t)b \in B_p(0, r)$  and  $B_p(0, r)$  is convex.

(2) Let  $a \in X$ . Set  $s = (p(a) + 1)/r$ . Then for each  $t \geq s$ ,  $tr \geq p(a) + 1$  so that

$$\begin{aligned} a &\in B_p(0, p(a) + 1) \\ &\subset B_p(0, tr) \\ &= tB_p(0, r) \end{aligned}$$

So  $B_p(0, r)$  is absorbing.

(3) Let  $a \in B_p(0, r)$  and  $u \in \mathbb{C}$ . Suppose that  $|u| \leq 1$ . Then

$$\begin{aligned} p(ua) &= |u|p(a) \\ &< |u|r \\ &\leq r \end{aligned}$$

So  $ua \in B_p(0, r)$  and  $B_p(0, r)$  is balanced.

Since  $B_p(0, r)$  is convex, absorbing and balanced, it is an absorbing disk.  $\square$

**Exercise 5.4.29.** Let  $X$  be a vector space and  $A \subset X$ . Suppose that  $A$  is an absorbing disk. Then  $p_A : X \rightarrow [0, \infty)$  is a seminorm on  $X$ .

*Proof.* Since  $A$  is an absorbing disk,  $A$  is convex, absorbing and balanced. So  $0 \in A$  and the previous exercise tells us that  $p$  is a sublinear functional on  $X$ . Let  $x \in X$  and  $\lambda \in \mathbb{C}$ .

If  $\lambda = 0$ , then

$$\begin{aligned} p_A(\lambda x) &= p_A(0) \\ &= 0 \\ &= |\lambda|p_A(x) \end{aligned}$$

Suppose that  $\lambda \neq 0$ . Since  $A$  is balanced, for  $t > 0$ ,  $\lambda t^{-1}x \in A$  iff  $|\lambda|t^{-1}x \in A$ . So

$$\begin{aligned} p_A(\lambda x) &= \inf\{t > 0 : \lambda x \in tA\} \\ &= \inf\{t > 0 : x \in |\lambda|^{-1}tA\} \\ &= \inf\{|\lambda|s > 0 : x \in sA\} \\ &= |\lambda| \inf\{s > 0 : x \in sA\} \\ &= |\lambda|p_A(x) \end{aligned}$$

So  $p$  is a seminorm on  $X$ .  $\square$



**Exercise 5.4.30.** Let  $X$  be a topological vector space and  $A \subset X$ . Suppose that  $A$  is an absorbing disk and  $A$  is open. Then  $B_{p_A}(0, 1) = A$ .

*Proof.* Clear by previous exercise. □

**Exercise 5.4.31.** Let  $X$  be a topological vector space and  $A \subset X$ . Suppose that  $A$  is an absorbing disk. Then  $p_A : X \rightarrow [0, \infty)$  is continuous iff  $A$  is open.

*Proof.* If  $A$  is open, then

$$\begin{aligned} A &= B_{p_A}(0, 1) \\ &\subset \bar{B}_{p_A}(0, 1) \end{aligned}$$

which implies that  $\bar{B}_{p_A}(0, 1) \in \mathcal{N}_0$ . An exercise in the previous section implies that  $p_A$  is continuous.

Conversely, if  $p_A$  is continuous, then an exercise in the previous section implies that  $B_{p_A}(0, 1)$  is open. □

### 5.5. Locally Convex Spaces.

**Definition 5.5.1.** Let  $X$  be a vector space and  $p : X \rightarrow [0, \infty)$  a seminorm on  $X$ . We equip  $X/\ker p$  with the topology induced by the norm  $\bar{p} : X/\ker p \rightarrow [0, \infty)$ . We define the projection  $\pi_p : X \rightarrow X/\ker p$  by  $\pi_p(x) = \bar{x} = x + \ker p$ .

**Definition 5.5.2.** Let  $X$  be a vector space and  $\mathcal{P}$  a family of seminorms on  $X$ . Then  $\mathcal{P}$  is said to **separate points of  $X$**  if for each  $x \in X$ , if  $x \neq 0$ , then there exists  $p \in \mathcal{P}$  such that  $p(x) \neq 0$ .

**Definition 5.5.3.** Let  $X$  be a vector space,  $\mathcal{T}$  a topology on  $X$  and  $\mathcal{P}$  a family of seminorms. Then  $(X, \mathcal{T})$  is said to be a **locally convex space with associated family of seminorms  $\mathcal{P}$**  if

- $\mathcal{P}$  separates points of  $X$
- $\mathcal{T} = \tau_X(\pi_p : p \in \mathcal{P})$

**Note 5.5.4.** We will generally suppress the family  $\mathcal{P}$  of seminorms and the induced topology  $\mathcal{T}$ .

**Exercise 5.5.5.** Let  $X$  be a locally convex space and  $(x_\alpha)_{\alpha \in A} \subset X$  a net and  $x \in X$ . Then  $x_\alpha \rightarrow x$  iff for each  $p \in \mathcal{P}$ ,  $p(x_\alpha - x) \rightarrow 0$ .

*Proof.* Suppose that  $x_\alpha \rightarrow x$ . Let  $p \in \mathcal{P}$ . By assumption,

$$\begin{aligned} \bar{x}_\alpha &= \pi_p(x_\alpha) \\ &\rightarrow \pi_p(x) \\ &= \bar{x} \end{aligned}$$

So

$$\begin{aligned} p(x_\alpha - x) &= \bar{p}(\bar{x}_\alpha - \bar{x}) \\ &\rightarrow 0 \end{aligned}$$

Conversely, suppose that for each  $p \in \mathcal{P}$ ,  $p(x_\alpha - x) \rightarrow 0$ . Let  $p \in \mathcal{P}$ . Then

$$\begin{aligned} \bar{p}(\bar{x}_\alpha - \bar{x}) &= p(x_\alpha - x) \\ &\rightarrow 0 \end{aligned}$$

So  $\pi_p(x_\alpha) \rightarrow \pi_p(x)$ . Since  $p \in \mathcal{P}$  is arbitrary,  $x_\alpha \rightarrow x$ . □

**Exercise 5.5.6.** Let  $X$  be a locally convex space. Then for each  $p \in \mathcal{P}$ ,  $p$  is continuous.

*Proof.* Let  $(x_\alpha)_{\alpha \in A} \subset X$  be a net and  $x \in X$ . Suppose that  $x_\alpha \rightarrow x$ . Let  $p \in \mathcal{P}$ . Then  $p(x_\alpha - x) \rightarrow 0$ . The reverse triangle inequality implies that

$$\begin{aligned} |p(x_\alpha) - p(x)| &\leq p(x_\alpha - x) \\ &\rightarrow 0 \end{aligned}$$

So  $p(x_\alpha) \rightarrow p(x)$  and  $p$  is continuous. □

**Exercise 5.5.7.** Let  $X$  be a locally convex space. Then  $X$  is a Hausdorff topological vector space.

*Proof.*

- (1) Let  $(x_\alpha)_{\alpha \in A}$ ,  $(y_\alpha)_{\alpha \in A} \subset X$  and  $(\lambda_\alpha)_{\alpha \in A} \subset \mathbb{C}$  be nets and  $x, y \in X$ ,  $\lambda \in \mathbb{C}$ . Suppose that  $x_\alpha \rightarrow x$ ,  $y_\alpha \rightarrow y$  and  $\lambda_\alpha \rightarrow \lambda$ . Let  $P \in \mathcal{P}$ . Then

$$\begin{aligned} p([x_\alpha + y_\alpha] - [x + y]) &= p([x_\alpha - x] + [y_\alpha - y]) \\ &\leq p(x_\alpha - x) + p(y_\alpha - y) \\ &\rightarrow 0 \end{aligned}$$

Since  $p \in \mathcal{P}$  is arbitrary,  $x_\alpha + y_\alpha \rightarrow x + y$  and addition  $X \times X \rightarrow X$  is continuous.

- (2) Similarly,

$$\begin{aligned} p(\lambda_\alpha x_\alpha - \lambda x) &= p([\lambda_\alpha x_\alpha - \lambda x_\alpha] + [\lambda x_\alpha - \lambda x]) \\ &\leq p(\lambda_\alpha x_\alpha - \lambda x_\alpha) + p(\lambda x_\alpha - \lambda x) \\ &= p([\lambda_\alpha - \lambda]x_\alpha) + p(\lambda[x_\alpha - x]) \\ &= |\lambda_\alpha - \lambda|p(x_\alpha) + |\lambda|p(x_\alpha - x) \\ &\rightarrow 0 \end{aligned}$$

So scalar multiplication  $\mathbb{C} \times X \rightarrow X$  is continuous.

- (3) Let  $x, y \in X$ . Suppose that  $x \neq y$ . Since  $\mathcal{P}$  separates points of  $X$ , there exists  $p \in \mathcal{P}$  such that  $p(x - y) \neq 0$ . Thus  $\bar{p}(\bar{x} - \bar{y}) \neq 0$ . Thus  $\bar{x} \neq \bar{y}$ . Since  $X/\ker p$  is Hausdorff, there exists  $U' \in \mathcal{N}_{\bar{x}}$  and  $V' \in \mathcal{N}_{\bar{y}}$  such that  $U' \cap V' = \emptyset$ . Set  $U = \pi_p^{-1}(U')$  and  $V = \pi_p^{-1}(V')$ . Then  $U \in \mathcal{N}_x$ ,  $V \in \mathcal{N}_y$  and

$$\begin{aligned} U \cap V &= \pi_p^{-1}(U') \cap \pi_p^{-1}(V') \\ &= \pi_p^{-1}(U' \cap V') \\ &= \pi_p^{-1}(\emptyset) \\ &= \emptyset \end{aligned}$$

So  $X$  is Hausdorff. □

**Exercise 5.5.8.** Let  $X$  be a locally convex space and  $U \in \mathcal{N}_0$  open. Then there exist  $p \in \mathcal{P}$  and  $r > 0$  such that  $B_p(0, r) \subset U$ .

*Proof.* For the sake of contradiction, suppose that for each  $p \in \mathcal{P}$  and  $r > 0$ ,  $B_p(0, r) \not\subset U$ . Then there exists a sequence  $(x_n)_{n \in \mathbb{N}} \subset U^c$  such that for each  $p \in \mathcal{P}$  and  $n \in \mathbb{N}$ ,  $p(x_n) < 1/n$ . So  $x_n \rightarrow 0$ . Since  $U^c$  is closed,  $0 \in U^c$  which is a contradiction. Hence there exist  $p \in \mathcal{P}$  and  $r > 0$  such that  $B_p(0, r) \subset U$ . □

**Exercise 5.5.9.** Let  $X$  be a locally convex space. Then for each  $U \in \mathcal{N}_0$ , if  $U$  is open, then there exists  $V \subset U$  such that  $V$  is an open absorbing disk.

*Proof.* Let  $U \in \mathcal{N}_0$ . Suppose that  $U$  is open. The previous exercise implies that there exists  $p \in \mathcal{P}$  and  $r > 0$  such that  $B_p(0, 1) \subset U$ . A previous exercise tells us that  $B_p(0, 1)$  is an open absorbing disk. □

**Exercise 5.5.10.** Let  $(X, \mathcal{T})$  be a locally convex space with associated family of seminorms  $\mathcal{P}$  and  $M \subset X$  a subspace. Define  $\mathcal{P}_M = \{p|_M : p \in \mathcal{P}\}$ . Then  $(M, \mathcal{T} \cap M)$  is a locally convex space with associated family of seminorms  $\mathcal{P}_M$ .

*Proof.* Let  $(x_\alpha)_{\alpha \in A} \subset M$  be a net and  $x \in M$ . Suppose that  $x_\alpha \rightarrow x$  in  $\mathcal{T} \cap M$ . Then an exercise in the section on the subspace topology implies that  $x_\alpha \rightarrow x$  in  $\mathcal{T}$ . Let  $q \in \mathcal{P}_M$ . Then there exists  $p \in \mathcal{P}$  such that  $q = p|_M$ . Therefore

$$\begin{aligned} q(x_\alpha - x) &= p|_M(x_\alpha - x) \\ &= p(x_\alpha - x) \\ &\rightarrow 0 \end{aligned}$$

Hence  $x_\alpha \rightarrow x$  in  $\tau_X(\pi_q : q \in \mathcal{P}_M)$ .

Conversely, suppose that  $x_\alpha \rightarrow x$  in  $\tau_X(\pi_q : q \in \mathcal{P}_M)$ . Let  $p \in \mathcal{P}$ . Then

$$\begin{aligned} p(x_\alpha - x) &= p|_M(x_\alpha - x) \\ &\rightarrow 0 \end{aligned}$$

Hence  $x_\alpha \rightarrow x$  in  $\mathcal{T}$ . So  $x_\alpha \rightarrow x$  in  $\mathcal{T} \cap M$ . Therefore  $\mathcal{T} \cap M = \tau_X(\pi_q : q \in \mathcal{P}_M)$ .  $\square$

**Exercise 5.5.11.** Let  $X$  be a locally convex space,  $M \subset X$  a subspace and  $f \in M^*$ . Then there exists  $F \in X^*$  such that  $F|_M = f$ .

*Proof.* Set  $p_f = |f|$ . Since  $p_f$  is a continuous seminorm,  $B_{p_f}(0, 1)$  is open in  $M$ . Therefore, there exists  $U \subset X$  open such that  $B_{p_f}(0, 1) = U \cap M$ . A previous exercise implies that there exists  $p \in \mathcal{P}$  and  $r > 0$  such that  $B_p(0, r) \subset U$ . Set  $A = B_p(0, r)$ . Since  $A$  is open,  $p_A : X \rightarrow [0, \infty)$  is continuous and  $A = B_{p_A}(0, 1)$ . Hence

$$\begin{aligned} B_{p_A|_M}(0, 1) &= A \cap M \subset U \cap M \\ &= B_{p_f}(0, 1) \end{aligned}$$

Therefore  $p_f \leq p_A|_M$  and  $|f| \leq p_A$  on  $M$ . The Hahn-Banach theorem implies that there exists  $F : X \rightarrow \mathbb{C}$  such that  $F$  is linear,  $F|_M = f$  and  $|F| \leq p_A$ . Since  $p_A$  is continuous,  $|F|$  is continuous, which implies that  $F$  is continuous. So  $F \in X^*$ .  $\square$

**Exercise 5.5.12. Hahn-Banach Separation Theorem 2:**

Let  $X$  be a locally convex space and  $A, B \subset X$ . Suppose that  $A, B$  are nonempty, convex and disjoint. If  $A$  is compact and  $B$  is closed, then there exists  $\phi \in X^*$  and  $c_1, c_2 \in \mathbb{R}$  such that for each  $x \in A, y \in B$ ,

$$\operatorname{Re} \phi(x) < c_1 < c_2 \leq \operatorname{Re} \phi(y)$$

**Hint:** Assume  $X$  is real. Since  $X$  is locally convex, there exists  $V \subset U$  such that  $V$  is an open absorbing disk and  $(A + V) \cap B = \emptyset$ . Then apply the first Hahn-Banach separation theorem to  $A + V$  and  $B$ .

*Proof.* Assume  $X$  is real. Suppose that  $A$  is compact and  $B$  is closed. A previous exercise implies that there exists  $U \in \mathcal{N}_0$  such that  $U$  is open and  $(A + U) \cap B = \emptyset$ . Since  $X$  is locally convex, there exists  $V \subset U$  such that  $V$  is an open absorbing disk. Then  $(A + V)$  is open and convex. By the first Hahn-Banach separation theorem, there exist  $\phi \in X^*$  and  $c_2 \in \mathbb{R}$  such that for each  $x \in A + V, y \in B$ ,

$$\phi(x) < c_2 \leq \phi(y)$$

Specifically,  $c_2 = \sup_{x \in A+V} \phi(x)$ . Since  $\phi \in X^*$  is not constant,  $\phi$  is open and thus  $\phi(A + V)$  is open. Continuity of  $\phi$  implies that  $\phi(A)$  is compact. Therefore,  $\sup \phi(A) < \sup \phi(A + V)$ .

So there exists  $c_1 \in \phi(A + V)$  such that  $\sup \phi(A) < c_1$ . Hence there exists  $x_1 \in A + V$  such that  $\phi(x_1) = c_1$ . Then for each  $x \in A$  and  $y \in B$ ,

$$\begin{aligned}\phi(x) &\leq \sup \phi(A) \\ &< c_1 \\ &= \phi(x_1) \\ &< c_2 \\ &\leq \phi(y)\end{aligned}$$

If  $X$  is complex, then the previous part implies that there exists  $f : X \rightarrow \mathbb{R}$  and  $c_1, c_2 \in \mathbb{R}$  such that  $f$  is continuous, real-linear and for each  $x \in A$  and  $y \in B$ ,

$$f(x) < c_1 < c_2 \leq f(y)$$

A previous exercise implies that there exists a unique  $\phi \in X^*$  such that  $\operatorname{Re} \phi = f$ .  $\square$

**Exercise 5.5.13.** Let  $X$  be a locally convex space and  $M \subset X$  a closed subspace. If  $M \neq X$ , then there exists  $\phi \in X^*$  such that  $\phi \neq 0$  and  $\phi|_M = 0$ .

*Proof.* Assume that  $X$  is real. Suppose that  $M \neq X$ . Then there exists  $x_0 \in X$  such that  $x_0 \notin M$ . Since  $\{x_0\}$  is compact and convex,  $M$  is closed and convex and  $\{x_0\} \cap M = \emptyset$ , the second Hahn-Banach separation theorem implies that there exists  $\phi \in X^*$  such that for each  $x \in M$ ,

$$\phi(x_0) < \phi(x)$$

Since  $0 \in M$ ,

$$\begin{aligned}\phi(x_0) &< \phi(0) \\ &= 0\end{aligned}$$

so that  $\phi \neq 0$ . For the sake of contradiction, suppose that  $\phi|_M \neq 0$ . Then there exists  $x_1 \in M$  such that  $\phi(x_1) \neq 0$ . Then for each  $t \in \mathbb{R}$ ,

$$\begin{aligned}\phi(x_0) &< \phi(tx_1) \\ &= t\phi(x_1)\end{aligned}$$

Set  $t = \frac{\phi(x_0)}{\phi(x_1)}$ . Then

$$\begin{aligned}\phi(x_0) &< t\phi(x_1) \\ &= \phi(x_0)\end{aligned}$$

which is a contradiction. So  $\phi|_M = 0$ .  $\square$

**Exercise 5.5.14.** Let  $X$  be a locally convex space. Then  $X^*$  separates the points of  $X$ .

*Proof.* Let  $x, y \in X$ . The second Hahn-Banach separation theorem implies that there exists  $\phi \in X^*$  such that  $\phi(x) \neq \phi(y)$ .  $\square$

### 5.6. Quotient Spaces.

**Exercise 5.6.1.** Let  $X$  be a topological vector space and  $M \subset X$  a subspace. Then  $\pi : X \rightarrow X/M$  is open.

*Proof.* Define the action  $\phi : M \times X \rightarrow X$  by  $m \cdot x = x + m$ . Then  $o_x = x + M$ . Since for each  $m \in M$ , the map  $x \mapsto x + m$  is continuous, an exercise in the section on the quotient topology implies that  $\pi : X \rightarrow X/M$  is open.  $\square$

**Exercise 5.6.2.** Let  $(X, \mathcal{T})$  be a topological vector space and  $M \subset X$  a subspace. Then  $(X/M, \mathcal{T}_{X/M})$  is a topological vector space.

*Proof.* Denote addition on  $X$  and  $X/M$  by  $A : X^2 \rightarrow X$  and  $\bar{A} : (X/M)^2 \rightarrow X/M$  respectively. Similarly, denote scalar multiplication on  $X$  and  $X/M$  by  $\Lambda : \mathbb{C} \times X \rightarrow X$  and  $\bar{\Lambda} : \mathbb{C} \times (X/M) \rightarrow X/M$  respectively.

- Let  $\bar{x}, \bar{y} \in X/M$ . Let  $U \in \mathcal{N}_{\bar{x}+\bar{y}}$ . Since  $\pi : X \rightarrow X/M$  is continuous, we have that  $\pi^{-1}(U) \in \mathcal{N}_{x+y}$ . Since addition  $A : X^2 \rightarrow X$  is continuous,

$$\begin{aligned} A^{-1}(\pi^{-1}(U)) &= (\pi \circ A)^{-1}(U) \\ &\in \mathcal{N}_{(x,y)} \end{aligned}$$

Since  $\mathcal{B} = \{A \times B : A, B \subset X \text{ and } A, B \text{ are open}\}$  is a basis for the product topology on  $X^2$ , there exist  $V_x \times V_y \in \mathcal{B}$  such that  $(x, y) \in V_x \times V_y \subset (\pi \circ A)^{-1}(U)$ . Thus  $V_x \in \mathcal{N}_x, V_y \in \mathcal{N}_y$  and  $V_x \times V_y \in \mathcal{N}_{(x,y)}$ . Recall that  $\pi \times \pi : X^2 \rightarrow (X/M)^2$  is defined by  $\pi \times \pi(x, y) = (\pi(x), \pi(y))$ . For  $x, y \in X$ , we have that

$$\begin{aligned} \bar{A} \circ (\pi \times \pi)(x, y) &= \bar{A}(\bar{x}, \bar{y}) \\ &= \bar{x} + \bar{y} \\ &= \pi(x) + \pi(y) \\ &= \pi(x + y) \\ &= \pi \circ A(x, y) \end{aligned}$$

So  $\bar{A} \circ (\pi \times \pi) = \pi \circ A$ . Since  $\pi$  is open, an exercise in the section on the product topology implies that  $\pi \times \pi$  is open and therefore  $\pi \times \pi(V_x \times V_y) \in \mathcal{N}_{(\bar{x}, \bar{y})}$ . Hence

$$\begin{aligned} \bar{A} \circ (\pi \times \pi)(V_x \times V_y) &\subset \bar{A} \circ (\pi \times \pi)((\pi \circ A)^{-1}(U)) \\ &= \bar{A} \circ (\pi \times \pi)((\bar{A} \circ (\pi \times \pi))^{-1}(U)) \\ &\subset U \end{aligned}$$

So for each  $U \in \mathcal{N}_{\bar{x}+\bar{y}}$ , there exists  $\pi \times \pi(V_x \times V_y) \in \mathcal{N}_{(\bar{x}, \bar{y})}$  such that  $\bar{A}(\pi \times \pi(V_x \times V_y)) \subset U$ . Hence  $\bar{A}$  is continuous at  $(\bar{x}, \bar{y})$ . Since  $\bar{x}, \bar{y} \in X/M$  are arbitrary,  $\bar{A}$  is continuous.

- Let  $\lambda \in \mathbb{C}$  and  $\bar{x} \in X/M$ . Let  $U \in \mathcal{N}_{\lambda\bar{x}}$ . Since  $\pi$  is continuous,  $\pi^{-1}(U) \in \mathcal{N}_{\lambda x}$ . Since scalar multiplication  $\Lambda : \mathbb{C} \times X \rightarrow X$  is continuous,

$$\begin{aligned} \Lambda^{-1}(\pi^{-1}(U)) &= (\pi \circ \Lambda)^{-1}(U) \\ &\in \mathcal{N}_{(\lambda, x)} \end{aligned}$$

Since  $\mathcal{B} = \{A \times B : A \subset \mathbb{C}, B \subset X \text{ and } A, B \text{ are open}\}$  is a basis for the product topology on  $\mathbb{C} \times X$ , there exist  $V_\lambda \times V_x \in \mathcal{B}$  such that  $(\lambda, x) \in V_\lambda \times V_x \subset (\pi \circ \Lambda)^{-1}(U)$ . Thus  $V_\lambda \in \mathcal{N}_\lambda, V_x \in \mathcal{N}_x$  and  $V_\lambda \times V_x \in \mathcal{N}_{(\lambda, x)}$ . As in the previous part,  $\pi \circ \Lambda =$

$\bar{\Lambda} \circ (\text{id}_{\mathbb{C}} \times \pi)$  and  $\text{id}_{\mathbb{C}}$  is open. Hence  $\text{id}_{\mathbb{C}} \times \pi$  is open and  $\text{id}_{\mathbb{C}} \times \pi(V_{\lambda} \times V_x) \in \mathcal{N}_{(\lambda, \bar{x})}$ . As in the previous part we have that

$$\begin{aligned} \bar{\Lambda} \circ (\text{id}_{\mathbb{C}} \times \pi)(V_{\lambda} \times V_x) &\subset \bar{\Lambda} \circ (\text{id}_{\mathbb{C}} \times \pi)((\pi \circ \Lambda)^{-1}(U)) \\ &= \bar{\Lambda} \circ (\text{id}_{\mathbb{C}} \times \pi)((\bar{\Lambda} \circ (\text{id}_{\mathbb{C}} \times \pi))^{-1}(U)) \\ &\subset U \end{aligned}$$

So for each  $U \in \mathcal{N}_{\lambda \bar{x}}$ , there exists  $\text{id}_{\mathbb{C}} \times \pi(V_{\lambda} \times V_x) \in \mathcal{N}_{(\lambda, \bar{x})}$  such that  $\bar{\Lambda}(\text{id}_{\mathbb{C}} \times \pi(V_{\lambda} \times V_x)) \subset U$ . Hence  $\bar{\Lambda}$  is continuous at  $(\lambda, \bar{x})$ . Since  $\lambda \in \mathbb{C}$  and  $\bar{x} \in X/M$  are arbitrary,  $\bar{\Lambda}$  is continuous. □

**Exercise 5.6.3.** Let  $X$  be a topological vector space and  $M \subset X$  a subspace. If  $M$  is closed, then  $X/M$  is Hausdorff.

*Proof.* Suppose that  $M$  is closed. Define the action  $\phi : M \times X \rightarrow X$  by  $m \cdot x = m + x$ . Denote by  $\sim$ , the equivalence relation induced by  $\phi$  (i.e.  $x \sim y$  iff  $x - y \in M$ ). A previous exercise implies that  $\pi : X \rightarrow X/M$  is open. Let  $(x_{\alpha}, y_{\alpha})_{\alpha \in A} \subset \sim$  be a net and  $(x, y) \in X \times X$ . Suppose that  $(x_{\alpha}, y_{\alpha}) \rightarrow (x, y)$ . Then  $x_{\alpha} \rightarrow x$  and  $y_{\alpha} \rightarrow y$ . Therefore  $x_{\alpha} - y_{\alpha} \rightarrow x - y$ . Since for each  $\alpha \in A$ ,  $x_{\alpha} - y_{\alpha} \in M$  and  $M$  is closed, we have that  $x - y \in M$ . Hence  $(x, y) \in \sim$  and  $\sim$  is closed. Since  $\pi$  is open, a previous exercise in the section on separation and countability implies that  $X/M$  is Hausdorff. □

**Definition 5.6.4.**

**Exercise 5.6.5.**

**Exercise 5.6.6.** Let  $X, Y$  be topological vector spaces. and  $\phi : X \rightarrow Y$  linear. Then  $\ker \phi$  is closed iff  $\phi$  is continuous.

*Proof.* Suppose that  $\phi$  is continuous. Since  $\{0\} \subset Y$  is closed,  $\ker \phi = \phi^{-1}(\{0\})$  is closed. Conversely, suppose that  $\ker \phi$  is closed. □

### 5.7. Duality.

**Definition 5.7.1.** Let  $X$  be a topological vector space and  $x \in X$ . Define  $\hat{x} : X^* \rightarrow \mathbb{C}$  by  $\hat{x}(f) = f(x)$ . We define  $\hat{X} = \{\hat{x} : x \in X\}$ .

**Definition 5.7.2.** Let  $X$  be a topological vector space. We define the **weak topology on  $X$** , denoted  $\mathcal{T}_w$ , by  $\mathcal{T}_w = \tau_X(X^*)$  (i.e. the initial topology on  $X$  generated by  $X^*$ ).

**Definition 5.7.3.** Let  $X$  be a topological vector space,  $(x_\alpha)_{\alpha \in A} \subset X$  and  $x \in X$ . Then  $(x_\alpha)_{\alpha \in A}$  is said to **converge weakly to  $x$** , denoted  $x_\alpha \xrightarrow{w} x$  if  $(x_\alpha)_{\alpha \in A}$  converges to  $x$  in the weak topology.

**Exercise 5.7.4.** Let  $X$  be a topological vector,  $(x_\alpha)_{\alpha \in A} \subset X$  a net and  $x \in X$ . Then  $x_\alpha \xrightarrow{w} x$  iff for each  $\lambda \in X^*$ ,  $\lambda(x_\alpha) \rightarrow \lambda(x)$ .

*Proof.* Immediate by Exercise 4.3.17. □

**Definition 5.7.5.** Let  $X$  be a topological vector space. We define the **weak-\* topology on  $X^*$** , denoted  $\mathcal{T}_{w*}$ , by  $\mathcal{T}_{w*} = \tau_X(\hat{X})$  (i.e. the initial topology on  $X^*$  generated by  $\hat{X}$ ).

**Definition 5.7.6.** Let  $X$  be a topological vector space,  $(\lambda_\alpha)_{\alpha \in A} \subset X^*$  and  $\lambda \in X^*$ . Then  $(\lambda_\alpha)_{\alpha \in A}$  is said to **converge in weak-\* to  $\lambda$** , denoted  $\lambda_\alpha \xrightarrow{w*} \lambda$  if  $(\lambda_\alpha)_{\alpha \in A}$  converges to  $\lambda$  in the weak-\* topology.

**Exercise 5.7.7.** Let  $X$  be a topological vector,  $(\lambda_\alpha)_{\alpha \in A} \subset X^*$  a net and  $\lambda \in X^*$ . Then  $\lambda_\alpha \xrightarrow{w*} \lambda$  iff for each  $x \in X$ ,  $\lambda_\alpha(x) \rightarrow \lambda(x)$ .

*Proof.* Immediate by Exercise 4.3.17. □

**Exercise 5.7.8.** Let  $X$  be a topological vector space.

- (1) If  $X^*$  separates the points of  $X$ , then  $(X, \mathcal{T}_w)$  is a locally convex space
- (2)  $(X^*, \mathcal{T}_{w*})$  is a locally convex space

*Proof.*

- (1) Suppose that  $X^*$  separates the points of  $X$ . For  $\lambda \in X^*$ , define  $p_\lambda : X \rightarrow [0, \infty)$  by  $p_\lambda = |\lambda|$ . Set  $\mathcal{P}_w = \{p_\lambda : \lambda \in X^*\}$ . Then  $\mathcal{P}_w$  separates the points of  $X$ . Let  $(x_\alpha)_{\alpha \in A} \subset X$  be a net and  $x \in X$ . Suppose that  $x_\alpha \xrightarrow{w} x$ . Let  $\lambda \in X^*$ . Then

$$\begin{aligned} p_\lambda(x_\alpha - x) &= |\lambda(x_\alpha - x)| \\ &= |\lambda(x_\alpha) - \lambda(x)| \\ &\rightarrow 0 \end{aligned}$$

So  $x_\alpha \rightarrow x$  in  $\tau_X(\pi_p : p \in \mathcal{P}_w)$ .

Conversely, suppose that  $x_\alpha \rightarrow x$  in  $\tau_X(\pi_p : p \in \mathcal{P}_w)$ . Then for each  $x \in X$ ,

$$\begin{aligned} |\lambda(x_\alpha) - \lambda(x)| &= p_\lambda(x_\alpha - x) \\ &\rightarrow 0 \end{aligned}$$

So that  $\lambda(x_\alpha) \rightarrow \lambda(x)$  and  $x_\alpha \xrightarrow{w} x$ . Hence  $\mathcal{T}_w = \tau_X(\pi_p : p \in \mathcal{P}_w)$  and  $(X, \mathcal{T}_w)$  is a locally convex space.



- (2) For  $x \in X$ , define  $p_x : X^* \rightarrow [0, \infty)$  by  $p_x = |\hat{x}|$ . Set  $\mathcal{P}_{w^*} = \{p_x : x \in X\}$ . Let  $\phi \in X^*$ . Suppose that  $\phi \neq 0$ . Then there exists  $x \in X$  such that

$$\begin{aligned}\hat{x}(\phi) &= \phi(x) \\ &\neq 0\end{aligned}$$

So  $\mathcal{P}_{w^*}$  separates the points of  $X^*$ . Let  $(\lambda_\alpha)_{\alpha \in A} \subset X^*$  be a net and  $\lambda \in X^*$ . Suppose that  $\lambda_\alpha \xrightarrow{w^*} \lambda$ . Let  $x \in X$ . Then

$$\begin{aligned}p_x(\lambda_\alpha - \lambda) &= |\hat{x}(\lambda_\alpha - \lambda)| \\ &= |\hat{x}(\lambda_\alpha) - \hat{x}(\lambda)| \\ &\rightarrow 0\end{aligned}$$

So  $\lambda_\alpha \rightarrow \lambda$  in  $\tau_{X^*}(\pi_p : p \in \mathcal{P}_{w^*})$ .

Conversely, suppose that  $\lambda_\alpha \rightarrow \lambda$  in  $\tau_{X^*}(\pi_p : p \in \mathcal{P}_{w^*})$ . Then for each  $x \in X$ ,

$$\begin{aligned}|\hat{x}(\lambda_\alpha) - \hat{x}(\lambda)| &= p_x(\lambda_\alpha - \lambda) \\ &\rightarrow 0\end{aligned}$$

So that  $\hat{x}(\lambda_\alpha) \rightarrow \hat{x}(\lambda)$  and  $\lambda_\alpha \xrightarrow{w^*} \lambda$ . Hence  $\mathcal{T}_{w^*} = \tau_{X^*}(\pi_p : p \in \mathcal{P}_{w^*})$  and  $(X^*, \mathcal{T}_{w^*})$  is a locally convex space. □

**Note 5.7.9.** Let  $X$  be a topological vector space. When we equip  $X^*$  with the weak-\* topology, we write  $X^{**}$  in place of  $(X^*)^*$ .

**Exercise 5.7.10.** Let  $X$  be a topological vector space. Then  $X^{**} = \hat{X}$ .

**Hint:** Hahn-Banach theorem

*Proof.* Let  $f \in X^{**}$ . Define  $p_f = |f|$ . Then  $p_f$  is a continuous seminorm on  $X^*$ . Therefore  $B_{p_f}(0, 1)$  is open. A previous exercise implies that there exists  $p \in \mathcal{P}_{w^*}$  and  $r > 0$  such that

$$\begin{aligned}B_{r^{-1}p}(0, 1) &= B_p(0, r) \\ &\subset B_{p_f}(0, 1)\end{aligned}$$

A previous exercise implies that  $p_f \leq r^{-1}p$ . By definition of  $\mathcal{P}_{w^*}$ , there exists  $x \in X$  such that  $p = |\hat{x}|$ . Thus

$$\begin{aligned}p_f &= |f| \\ &\leq r^{-1}p \\ &= |r^{-1}\hat{x}|\end{aligned}$$

Therefore  $\ker \hat{x} \subset \ker f$ . □

## 6. BANACH SPACES

## 6.1. Introduction.

**Note 6.1.1.** In the following, we will consider vector spaces over  $\mathbb{C}$ . There are analogous results for real vector spaces as well, just replace every  $\mathbb{C}$  with  $\mathbb{R}$ .

**Definition 6.1.2.** Let  $X$  be a normed vector space. Then  $X$  is said to be a **Banach space** if  $X$  is complete.

**Definition 6.1.3.** Let  $X$  be a normed vector space and  $(x_i)_{i=1}^n \subset X$ . The series  $\sum_{i=1}^{\infty} x_i$  is said to **converge** if the sequence  $s_n := \sum_{i=1}^n x_i$  converges. The series  $\sum_{i=1}^{\infty} x_i$  is said to **converge absolutely** if  $\sum_{i \in \mathbb{N}} \|x_i\| < \infty$ .

**Exercise 6.1.4.** Let  $X$  be a normed vector space. Then  $X$  is complete iff for each  $(x_i)_{i \in \mathbb{N}} \subset X$ ,  $\sum_{i=1}^{\infty} x_i$  converges absolutely implies that  $\sum_{i=1}^{\infty} x_i$  converges.

**Hint:** Given a Cauchy sequence  $(x_n)_{n \in \mathbb{N}}$ , obtain a subsequence  $(x_{n_j})_{j \in \mathbb{N}} \subset (x_n)_{n \in \mathbb{N}}$  such that for each  $j \in \mathbb{N}$ ,  $\|x_{n_{j+1}} - x_{n_j}\| < 2^{-j}$ . Define a new sequence  $(y_j)_{j \in \mathbb{N}} \subset X$  by

$$y_j = \begin{cases} x_{n_1} & j = 1 \\ x_{n_j} - x_{n_{j-1}} & j \geq 2 \end{cases}$$

*Proof.* Suppose that  $X$  is complete. Let  $(x_i)_{i \in \mathbb{N}} \subset X$ . Suppose that  $\sum_{i=1}^{\infty} x_i$  converges absolutely. Let  $\epsilon > 0$ . Choose  $N \in \mathbb{N}$  such that for each  $m, n \in \mathbb{N}$ , if  $m, n \geq N$  and  $m < n$ , then  $\sum_{i=m+1}^n \|x_i\| < \epsilon$ . Let  $m, n \in \mathbb{N}$ . Suppose that  $m < n$ . Then

$$\begin{aligned} \|s_n - s_m\| &= \left\| \sum_{i=1}^n x_i - \sum_{i=1}^m x_i \right\| \\ &= \left\| \sum_{i=m+1}^n x_i \right\| \\ &\leq \sum_{i=m+1}^n \|x_i\| \\ &< \epsilon \end{aligned}$$

Thus  $(s_n)_{n \in \mathbb{N}}$  is Cauchy. Since  $X$  is complete,  $\sum_{i=1}^{\infty} x_i$  converges. Conversely, Suppose that for each  $(x_i)_{i \in \mathbb{N}} \subset X$ ,  $\sum_{i=1}^{\infty} x_i$  converges absolutely implies that  $\sum_{i=1}^{\infty} x_i$  converges. Let  $(x_i)_{i \in \mathbb{N}} \subset X$  be Cauchy. Proceed inductively to create a strictly increasing sequence  $(n_i)_{i \in \mathbb{N}} \subset \mathbb{N}$  such that for each  $m, n \in \mathbb{N}$ , if  $m, n \geq n_i$ , then  $\|x_m - x_n\| < 2^{-i}$ . Define  $(y_i)_{i \in \mathbb{N}} \subset X$  by

$$y_i = \begin{cases} x_{n_1} & i = 1 \\ x_{n_i} - x_{n_{i-1}} & i \geq 2 \end{cases}$$

Then  $\sum_{i=1}^k y_i = x_{n_k}$  and

$$\begin{aligned} \sum_{i \in \mathbb{N}} \|y_i\| &= \|x_{n_1}\| + \sum_{i \in \mathbb{N}} \|x_{n_i} - x_{n_{i-1}}\| \\ &\leq \|x_{n_1}\| + 2 \sum_{i \in \mathbb{N}} 2^{-i} \\ &= \|x_{n_1}\| + 2 \end{aligned}$$

Hence  $(x_{n_k})_{k \in \mathbb{N}} = (\sum_{i=1}^k y_i)_{i \in \mathbb{N}}$  converges. Since  $(x_i)_{i \in \mathbb{N}}$  is cauchy and has a convergent subsequence, it converges. So  $X$  is complete.  $\square$

**Exercise 6.1.5.** Let  $X$  be a normed vector space. Then addition  $X \times X \rightarrow X$  and scalar multiplication  $\mathbb{C} \times X \rightarrow X$  are continuous and  $\|\cdot\| : X \rightarrow [0, \infty)$  is continuous.

*Proof.* Let  $\epsilon > 0$ . Choose  $\delta = \frac{\epsilon}{2}$ . Let  $(x_1, y_1), (x_2, y_2) \in X \times X$ . Suppose that

$$\max\{\|x_1 - x_2\|, \|y_1 - y_2\|\} < \delta$$

Then

$$\begin{aligned} \|(x_1 + y_1) - (x_2 + y_2)\| &= \|(x_1 - x_2) + (y_1 - y_2)\| \\ &\leq \|x_1 - x_2\| + \|y_1 - y_2\| \\ &< 2\delta \\ &= \epsilon \end{aligned}$$

Hence addition is uniformly continuous.

Let  $(\lambda_1, x_1) \in \mathbb{C} \times X$  and  $\epsilon > 0$ . Choose  $\delta = \min\{\frac{\epsilon}{2(|\lambda_1| + \|x_1\| + 1)}, \frac{\sqrt{\epsilon}}{\sqrt{2}}\}$ . Let  $(\lambda_2, x_2) \in \mathbb{C} \times X$ . Suppose that

$$\max\{|\lambda_1 - \lambda_2|, \|x_1 - x_2\|\} < \delta$$

Then

$$\begin{aligned} \|\lambda_1 x_1 - \lambda_2 x_2\| &= \|\lambda_1 x_1 - \lambda_1 x_2 + \lambda_1 x_2 - \lambda_2 x_2\| \\ &= \|\lambda_1(x_1 - x_2) + (\lambda_1 - \lambda_2)x_2\| \\ &\leq |\lambda_1|\|x_1 - x_2\| + |\lambda_1 - \lambda_2|\|x_2\| \\ &\leq |\lambda_1|\|x_1 - x_2\| + |\lambda_1 - \lambda_2|(\|x_1 - x_2\| + \|x_1\|) \\ &< |\lambda_1|\delta + \delta(\delta + \|x_1\|) \\ &= (|\lambda_1| + \|x_1\|)\delta + \delta^2 \\ &< \frac{\epsilon}{2} + \frac{\epsilon}{2} \\ &= \epsilon \end{aligned}$$

Since  $(\lambda_1, x_1) \in \mathbb{C} \times X$  is arbitrary, scalar multiplication is continuous.

Let  $\epsilon > 0$ . Choose  $\delta = \epsilon$ . Let  $x, y \in X$ . Suppose that  $\|x - y\| < \delta$ . Then

$$\begin{aligned} |\|x\| - \|y\|| &\leq \|x - y\| \\ &< \delta \\ &= \epsilon \end{aligned}$$

So  $\|\cdot\| : X \rightarrow [0, \infty)$  is uniformly continuous.  $\square$

## 6.2. Bounded Operators.

**Definition 6.2.1.** Let  $X, Y$  be a normed vector spaces and  $T : X \rightarrow Y$  linear. Then  $T$  is said to be **bounded** if  $T(\overline{B(0, 1)})$  is bounded. We define

$$L(X, Y) = \{T : X \rightarrow Y : T \text{ is linear and bounded}\}$$

When  $X = Y$ , we write  $L(X)$ .

**Exercise 6.2.2.** Let  $X, Y$  be a normed vector spaces and  $T : X \rightarrow Y$  linear. Then  $T$  is bounded iff there exists  $C \geq 0$  such that for each  $x \in X$ ,

$$\|Tx\| \leq C\|x\|$$

*Proof.* Suppose that  $T$  is bounded. If  $T = 0$ , choose  $C = 0$ . Suppose that  $T \neq 0$ . Set  $A = \{\|Tx\| : \|x\| = 1\}$ . Since  $T \neq 0$ , there exists  $x_0 \in X$  such that  $\|x_0\| = 1$  so that  $A \neq \emptyset$ . Boundedness of  $T$  implies that  $A$  is bounded. Set  $C = \sup A$ . Let  $x \in X$ . If  $x = 0$ , then  $Tx = 0$  and  $\|Tx\| \leq C\|x\|$ . Suppose that  $x \neq 0$ . Then  $Tx = \|x\|T(\|x\|^{-1}x)$ . Since  $\|\|x\|^{-1}x\| = 1$ , we have that

$$\begin{aligned} \|Tx\| &= \|T(\|x\|^{-1}x)\|\|x\| \\ &\leq C\|x\| \end{aligned}$$

Conversely, suppose that there exists  $C \geq 0$  such that for each  $x \in X$ ,  $\|Tx\| \leq C\|x\|$ . Let  $x \in \overline{B(0, 1)}$ . Then

$$\begin{aligned} \|Tx\| &\leq C\|x\| \\ &\leq C \end{aligned}$$

So that  $T(\overline{B(0, 1)})$  is bounded. □

**Exercise 6.2.3.** Set  $X = C^1([0, 1])$  and  $Y = C([0, 1])$ . Equip both  $X$  and  $Y$  with the sup norm. Define  $T : X \rightarrow Y$  by  $Tf = f'$ . Then  $T$  is not bounded.

*Proof.* For the sake of contradiction, suppose that  $T$  is bounded. Then there exists  $C \geq 0$  such that for each  $f \in X$ ,  $\|Tf\| \leq C\|f\|$ . Choose  $n \in \mathbb{N}$  such that  $n > C$ . Define  $f \in X$  by  $f(x) = x^n$ . Then

$$\begin{aligned} n &= \|Tf\| \\ &\leq C\|f\| \\ &= C \end{aligned}$$

which is a contradiction. Hence  $T$  is not bounded. □

**Exercise 6.2.4.** Let  $X, Y$  be a normed vector spaces and  $T : X \rightarrow Y$  a linear map. Then  $T$  is bounded iff there exists  $r, s > 0$  such that  $T(B(0, r)) \subset B(0, s)$

*Proof.* Suppose that  $T$  is bounded. Then there exists  $C \geq 0$  such that for each  $x \in X$ ,  $\|Tx\| \leq C\|x\|$ . Thus  $T(B(0, 1)) \subset B(0, C + 1)$ . Conversely. Suppose that there exists  $r, s > 0$  such that  $T(B(0, r)) \subset B(0, s)$ . Define  $C = \frac{2s}{r}$ . Let  $x \in X$ . Put  $\alpha = \frac{r}{2\|x\|}$ . Then

$\alpha x \in B(0, r)$ . So  $T(\alpha x) = \alpha T(x) \in B(0, s)$ . Hence

$$\begin{aligned}\|T(\alpha x)\| &= \|\alpha T(x)\| \\ &= |\alpha| \|T(x)\| \\ &= \frac{r}{2\|x\|} \|T(x)\| \\ &< s.\end{aligned}$$

Thus

$$\|Tx\| < \frac{2s}{r} \|x\| = C\|x\|$$

So  $T$  is bounded. □

**Exercise 6.2.5.** Let  $X, Y$  be normed vector spaces and  $T : X \rightarrow Y$ . Suppose that  $T$  is linear. Then there exists  $x_0 \in X$  such that  $T$  is continuous at  $x_0$  iff  $T$  is continuous at 0.

*Proof.* Suppose that there exists  $x_0 \in X$  such that  $T$  is continuous at  $x_0$ . Since  $T$  is linear,  $T(0) = 0$ . Let  $(x_n)_{n \in \mathbb{N}} \subset X$ . Suppose that  $x_n \rightarrow 0$ . Then  $x_n + x_0 \rightarrow x_0$ . Hence

$$\begin{aligned}T(x_n) + T(x_0) &= T(x_n + x_0) \\ &\rightarrow T(x_0)\end{aligned}$$

This implies that

$$\begin{aligned}T(x_n) &\rightarrow 0 \\ &= T(0)\end{aligned}$$

Therefore  $T$  is continuous at 0.

Conversely, if  $T$  is continuous at 0, then trivially, there exists  $x_0 \in X$  such that  $T$  is continuous at  $x_0$ . □

**Exercise 6.2.6.** Let  $X, Y$  be normed vector spaces and  $T : X \rightarrow Y$  a linear map. Then the following are equivalent:

- (1)  $T$  is continuous
- (2)  $T$  is continuous at  $x = 0$
- (3)  $T$  is bounded

*Proof.*

- (1)  $\implies$  (2):

Trivial

- (2)  $\implies$  (3):

Suppose that  $T$  is continuous at  $x = 0$ . Then there exists  $\delta > 0$  such that for each  $x \in X$ , if  $\|x\| < \delta$ , then  $\|Tx\| < 1$ . Choose  $C = \frac{2}{\delta}$ . If  $x = 0$ , then  $\|Tx\| \leq C\|x\|$ .

Suppose that  $\|x\| \neq 0$ . Define  $y = \frac{\delta}{2\|x\|}x$ . Then  $\|y\| < \delta$ . So

$$\begin{aligned}1 &> \|Ty\| \\ &= \frac{\delta}{2\|x\|} \|Tx\|\end{aligned}$$

Thus

$$\begin{aligned}\|Tx\| &< \frac{2}{\delta}\|x\| \\ &= C\|x\|\end{aligned}$$

Hence  $T$  is bounded.

• (3)  $\implies$  (1)

Suppose that  $T$  is bounded. Then there exists  $C \geq 0$  such that for each  $x \in X$ ,  $\|Tx\| \leq C\|x\|$ . Let  $\epsilon > 0$ . Choose  $\delta = \frac{\epsilon}{C+1}$ . Let  $x, y \in X$ . Suppose that  $\|x - y\| < \delta$ . Then

$$\begin{aligned}\|Tx - Ty\| &= \|T(x - y)\| \\ &\leq C\|x - y\| \\ &< (C + 1)\delta \\ &= \epsilon\end{aligned}$$

So  $T$  is continuous. □

**Definition 6.2.7.** Let  $X, Y$  be normed vector spaces. Define  $\|\cdot\| : L(X, Y) \rightarrow [0, \infty)$  by

$$\|T\| = \inf\{C \geq 0 : \text{for each } x \in X, \|Tx\| \leq C\|x\|\}$$

We call  $\|\cdot\|$  the **operator norm on  $L(X, Y)$**

**Exercise 6.2.8.** Let  $X, Y$  be normed vector spaces. If  $X \neq \{0\}$ , then the operator norm on  $L(X, Y)$  is given by:

$$(1) \|T\| = \sup_{\|x\|=1} \|Tx\|$$

$$(2) \|T\| = \sup_{x \neq 0} \|x\|^{-1} \|Tx\|$$

$$(3) \|T\| = \inf\{C \geq 0 : \text{for each } x \in X, \|Tx\| \leq C\|x\|\}$$

*Proof.* Since  $X \neq \{0\}$ , the supremums in (1) and (2) are well defined. Let  $T \in L(X, Y)$ . By linearity of  $T$ , the sets over which the supremums are taken in (1) and (2) are the same. So (1) and (2) are equal.

Now, set  $M = \sup_{\|x\|=1} \|Tx\|$  and  $m = \inf\{C \geq 0 : \text{for each } x \in X, \|Tx\| \leq C\|x\|\}$ . Let  $x \in X$ .

If  $\|x\| = 0$ , then  $\|Tx\| \leq M\|x\|$ . Suppose that  $\|x\| \neq 0$ . Then

$$\begin{aligned}\|Tx\| &= \left( \|T(x/\|x\|)\| \right) \|x\| \\ &\leq M\|x\|\end{aligned}$$

Hence  $M \in \{C \geq 0 : \text{for each } x \in X, \|Tx\| \leq C\|x\|\}$  and  $m \leq M$ . Let  $C \in \{C \geq 0 : \text{for each } x \in X, \|Tx\| \leq C\|x\|\}$ . Suppose that  $\|x\| = 1$ . Then  $\|Tx\| \leq C\|x\| = C$ . So  $M \leq C$ . Therefore  $M \leq m$ . So  $M = m$  and the supremum in (1) is the same as the infimum in (3). □

**Note 6.2.9.** From here on, unless stated otherwise, we assume  $X \neq 0$ .

**Exercise 6.2.10.** Let  $X, Y$  be normed vector spaces and  $T \in L(X, Y)$ . Then for each  $x \in X$ ,  $\|Tx\| \leq \|T\|\|x\|$

*Proof.* This is just part of the previous exercise. Let  $x \in X$ . If  $x = 0$ , then  $\|Tx\| \leq \|T\|\|x\|$ . Suppose that  $x \neq 0$ . Then  $\|Tx\| = T(x/\|x\|)\|x\| \leq \|T\|\|x\|$   $\square$

**Exercise 6.2.11.** Let  $X, Y$  be normed vector spaces. Then the operator norm is a norm on  $L(X, Y)$ .

*Proof.* Let  $S, T \in L(X, Y)$  and  $\alpha \in \mathbb{C}$ . For each  $x \in X$ , we have that

$$\begin{aligned} \|(S + T)x\| &= \|Sx + Tx\| \\ &\leq \|Sx\| + \|Tx\| \\ &\leq \|S\|\|x\| + \|T\|\|x\| \\ &= (\|S\| + \|T\|)\|x\| \end{aligned}$$

So  $\|S + T\| \leq \|S\| + \|T\|$ .

Using the definition of  $\|T\|$ , we see that

$$\begin{aligned} \|\alpha T\| &= \sup_{\|x\|=1} \|(\alpha T)x\| \\ &= \sup_{\|x\|=1} |\alpha| \|Tx\| \\ &= |\alpha| \sup_{\|x\|=1} \|Tx\| \\ &= |\alpha| \|T\| \end{aligned}$$

So  $\|\alpha S\| = |\alpha| \|S\|$ .

Suppose that  $\|T\| = 0$ . Let  $x \in X$ . Then  $\|Tx\| \leq \|T\|\|x\| = 0$ . So  $Tx = 0$ . Since  $x \in X$  is arbitrary, we have that  $T = 0$ .  $\square$

**Exercise 6.2.12.** Let  $X, Y, Z$  be normed vector spaces,  $T \in L(X, Y)$  and  $S \in L(Y, Z)$ . Define  $ST : X \rightarrow Z$  by  $STx = S(Tx)$ . Then  $ST \in L(X, Z)$  and  $\|ST\| \leq \|S\|\|T\|$ .

*Proof.* Clearly  $ST$  is linear. Let  $x \in X$ . Then

$$\begin{aligned} \|STx\| &= \|S(Tx)\| \\ &\leq \|S\|\|Tx\| \\ &\leq \|S\|\|T\|\|x\| \end{aligned}$$

So  $\|ST\| \leq \|S\|\|T\|$ .  $\square$

**Definition 6.2.13.** Let  $X, Y$  be normed vector spaces and  $T \in L(X, Y)$ . Then  $T$  is said to be **invertible** or an **isomorphism** if  $T$  is a bijection and  $T^{-1} \in L(Y, X)$ .

**Definition 6.2.14.** Let  $X$  be a normed vector space. Define  $GL(X) := \{T \in L(X, X) : T \text{ is invertible}\}$ .

**Exercise 6.2.15.** Let  $X, Y$  be normed vector spaces. If  $Y$  is complete, then so is  $L(X, Y)$ .

*Proof.* Suppose that  $Y$  is complete. Let  $(T_n)_{n \in \mathbb{N}} \subset L(X, Y)$ . Suppose that  $(T_n)_{n \in \mathbb{N}}$  is Cauchy. Since for each  $m, n \in \mathbb{N}$ ,  $|\|T_m\| - \|T_n\|| \leq \|T_m - T_n\|$ , we have that  $(\|T_n\|)_{n \in \mathbb{N}} \subset [0, \infty)$  is Cauchy. Hence  $\lim_{n \rightarrow \infty} \|T_n\|$  exists.

Let  $x \in X$  and  $m, n \in \mathbb{N}$ . Then

$$\begin{aligned} \|T_m x - T_n x\| &= \|(T_m - T_n)x\| \\ &\leq \|T_m - T_n\| \|x\| \end{aligned}$$

So  $(T_n x)_{n \in \mathbb{N}} \subset Y$  is Cauchy and hence converges. Define  $T : X \rightarrow Y$  by  $Tx = \lim_{n \rightarrow \infty} T_n x$ .

Since addition and scalar multiplication are continuous,  $T$  is linear. Let  $x \in X$  and  $\epsilon > 0$ . Choose  $N \in \mathbb{N}$  such that for each  $n \in \mathbb{N}$ , if  $n \geq N$ , then  $\|Tx - T_n x\| < \epsilon$ . Then for each  $n \in \mathbb{N}$ , if  $n \geq N$  we have that

$$\begin{aligned} \|Tx\| &\leq \|Tx - T_n x\| + \|T_n x\| \\ &< \epsilon + \|T_n x\| \\ &\leq \epsilon + \|T_n\| \|x\| \end{aligned}$$

Thus  $\|Tx\| \leq \epsilon + (\lim_{n \rightarrow \infty} \|T_n\|) \|x\|$ . Since  $\epsilon > 0$  is arbitrary,  $\|Tx\| \leq (\lim_{n \rightarrow \infty} \|T_n\|) \|x\|$ . Thus  $T \in L(X, Y)$  and  $\|T\| \leq \lim_{n \rightarrow \infty} \|T_n\|$ .

Note that since addition, scalar multiplication and  $\|\cdot\|$  are continuous, we have that for each  $n \in \mathbb{N}$  and  $x \in X$ ,  $\|(T_n - T_m)x\|$  converges to  $\|(T_n - T)x\|$  because

$$\begin{aligned} \lim_{m \rightarrow \infty} \|(T_n - T_m)x\| &= \lim_{m \rightarrow \infty} \|T_n x - T_m x\| \\ &= \|T_n x - \lim_{m \rightarrow \infty} T_m x\| \\ &= \|T_n x - Tx\| \\ &= \|(T_n - T)x\| \end{aligned}$$

Let  $\epsilon > 0$ . Choose  $N \in \mathbb{N}$  such that for each  $m, n \in \mathbb{N}$  if  $n, m \geq N$ , then  $\|T_n - T_m\| < \epsilon$ . Then for each  $n \in \mathbb{N}$  if  $n \geq N$ , then for each  $x \in X$ ,

$$\|(T_n - T_m)x\| \leq \|T_n - T_m\| \|x\| < \epsilon \|x\|$$

Combining this with the previous fact, we see that for each  $n \in \mathbb{N}$ , if  $n \geq N$ , then for each  $x \in X$ ,

$$\|(T_n - T)x\| \leq \epsilon \|x\|$$

In particular, for each  $n \in \mathbb{N}$ , if  $n \geq N$ , then

$$\|T_n - T\| = \sup_{\|x\|=1} \|(T_n - T)x\| \leq \epsilon$$

This implies that  $T_n$  converges to  $T$  in  $L(X, Y)$ . Since

$$|\|T_n\| - \|T\|| \leq \|T_n - T\|$$



it is clear that  $\lim_{n \rightarrow \infty} \|T_n\| = \|T\|$

□

### 6.3. Direct Sums.

**Definition 6.3.1.** Let  $X, Y$  be normed vector spaces and  $p \in [1, \infty]$ . Let  $\|\cdot\|'_p : \mathbb{R}^2 \rightarrow [0, \infty)$  denote the usual  $l^p$  norm. We define  $\|\cdot\|_p : X \oplus Y \rightarrow [0, \infty)$  by

$$\|(x, y)\|_p = \|(\|x\|, \|y\|)\|'_p$$

**Exercise 6.3.2.** Let  $X, Y$  be normed vector spaces. Then

- (1) for each  $p \in [1, \infty]$ ,  $\|\cdot\|_p : X \oplus Y \rightarrow [0, \infty)$  is a norm on  $X \oplus Y$
- (2)  $\{\|\cdot\|_p : p \in [1, \infty]\}$  are equivalent.

*Proof.*

- (1) Let  $p \in [1, \infty]$ ,  $(x_1, y_1), (x_2, y_2) \in X \oplus Y$  and  $\lambda \in \mathbb{C}$ .

- Clearly if  $(x_1, y_1) = (0, 0)$ , then  $\|S\|_p = 0$ . Conversely, suppose that  $\|(x_1, y_1)\|_p = 0$ . Then  $\|x_1\| = 0$  and  $\|y_1\| = 0$ . So  $x_1 = 0$  and  $y_1 = 0$ . Therefore  $S = 0$ .

•

$$\begin{aligned} \|\lambda(x_1, y_1)\|_p &= \|(\|\lambda x_1\|, \|\lambda y_1\|)\|'_p \\ &= \|(|\lambda|\|x_1\|, |\lambda|\|y_1\|)\|'_p \\ &= \| |\lambda| (\|x_1\|, \|y_1\|) \|'_p \\ &= |\lambda| \|(\|x_1\|, \|y_1\|)\|'_p \\ &= |\lambda| \|(x_1, y_1)\|_p \end{aligned}$$

•

$$\begin{aligned} \|(x_1, y_1) + (x_2, y_2)\|_p &= \|(\|x_1 + x_2\|, \|y_1 + y_2\|)\|'_p \\ &\leq \|(\|x_1\| + \|x_2\|, \|y_1\| + \|y_2\|)\|'_p \\ &= \|(\|x_1\|, \|y_1\|) + (\|x_2\|, \|y_2\|)\|'_p \\ &\leq \|(\|x_1\|, \|y_1\|)\|'_p + \|(\|x_2\|, \|y_2\|)\|'_p \\ &= \|(x_1, y_1)\|_p + \|(x_2, y_2)\|_p \end{aligned}$$

- (2) All norms on  $\mathbb{R}^2$  are equivalent.

□

**Exercise 6.3.3.** Let  $X, Y$  be Banach spaces. Then  $X \oplus Y$  equipped with  $\|\cdot\|_p : X \oplus Y \rightarrow [0, \infty)$  is a Banach space.

*Proof.*

□

**Exercise 6.3.4.** Let  $X, Y$  and  $Z$  be Banach spaces and  $p \in [0, \infty]$ . Equip  $Y \oplus Z$  with  $\|\cdot\|_p$ . Let  $T \in L(X, Y \oplus Z)$  with  $T = (T_Y, T_Z)$ . Then  $T_Y \in L(X, Y)$  and  $T_Z \in L(X, Z)$ .

*Proof.* Let  $x \in X$ . Then  $\|T_Y(x)\|, \|T_Z(x)\| \leq$

**FINISH!!!**

□

**Definition 6.3.5.** Let  $X, Y$  and  $Z$  be Banach spaces and  $p \in [0, \infty]$ . Let  $\|\cdot\|'_p : \mathbb{R}^2 \rightarrow [0, \infty)$  denote the usual  $l^p$  norm. Equip  $Y \oplus Z$  with  $\|\cdot\|_p$ . Let  $T \in L(X, Y \oplus Z)$  with  $T = (T_Y, T_Z)$ . Define  $\|T\|_p : L(X, Y \oplus Z) \rightarrow [0, \infty)$  by

$$\|T\|_p = \|(\|T_Y\|, \|T_Z\|)\|'_p$$

**Exercise 6.3.6.** Let  $X, Y$  and  $Z$  be Banach spaces and  $p \in [0, \infty]$ . Then  $\|\cdot\|_p : L(X, Y \oplus Z) \rightarrow [0, \infty)$  is a norm on  $L(X, Y \oplus Z)$ .

*Proof.* Let  $\lambda \in \mathbb{C}$  and  $S, T \in L(X, Y \oplus Z)$  with  $S = (S_Y, S_Z)$  and  $T = (T_Y, T_Z)$ .

- Clearly if  $S = 0$ , then  $\|S\|_p = 0$ . Conversely, suppose that  $\|S\|_p = 0$ . Then  $\|S_Y\| = 0$  and  $\|S_Z\| = 0$ . So  $S_Y = 0$  and  $S_Z = 0$ . Therefore  $S = 0$ .
- 

$$\begin{aligned} \|\lambda S\|_p &= \|(\|\lambda S_Y\|, \|\lambda S_Z\|)\|'_p \\ &= \|(|\lambda| \|S_Y\|, |\lambda| \|S_Z\|)\|'_p \\ &= \| |\lambda| (\|S_Y\|, \|S_Z\|) \|'_p \\ &= |\lambda| \|(\|S_Y\|, \|S_Z\|)\|'_p \\ &= |\lambda| \|S\|_p \end{aligned}$$

•

$$\begin{aligned} \|S + T\|_p &= \|(\|S_Y + T_Y\|, \|S_Z + T_Z\|)\|'_p \\ &\leq \|(\|S_Y\| + \|T_Y\|, \|S_Z\| + \|T_Z\|)\|'_p \\ &= \|(\|S_Y\|, \|S_Z\|) + (\|T_Y\|, \|T_Z\|)\|'_p \\ &\leq \|(\|S_Y\|, \|S_Z\|)\|'_p + \|(\|T_Y\|, \|T_Z\|)\|'_p \\ &= \|S\|_p + \|T\|_p \end{aligned}$$

So  $\|\cdot\|_p : L(X, Y \oplus Z) \rightarrow [0, \infty)$  is a norm on  $L(X, Y \oplus Z)$ . □

**Exercise 6.3.7.** Let  $X, Y$  and  $Z$  be Banach spaces and  $p \in [0, \infty]$ . Equip  $Y \oplus Z$  with  $\|\cdot\|_p$ . Let  $T \in L(X, Y \oplus Z)$  with  $T = (T_Y, T_Z)$ . Then  $\|T\| \leq 2^{1/p} \|T\|_p$ .

*Proof.* Let  $x \in X$ . If  $p < \infty$ , then

$$\begin{aligned} \|T(x)\|_p &= \|(T_Y(x), T_Z(x))\|_p \\ &= \|(\|T_Y(x)\|, \|T_Z(x)\|)\|'_p \\ &= \left( \|T_Y(x)\|^p + \|T_Z(x)\|^p \right)^{1/p} \\ &\leq \left( \|T_Y\|^p \|x\|^p + \|T_Z\|^p \|x\|^p \right)^{1/p} \\ &\leq \left[ (\|T_Y\|^p + \|T_Z\|^p) \|x\|^p + (\|T_Y\|^p + \|T_Z\|^p) \|x\|^p \right]^{1/p} \\ &= \left[ 2(\|T_Y\|^p + \|T_Z\|^p) \|x\|^p \right]^{1/p} \\ &= 2^{1/p} \|T\|_p \|x\| \end{aligned}$$

Hence  $\|T\| \leq 2^{1/p} \|T\|_p$ . If  $p = \infty$ , then

$$\begin{aligned}
 \|T(x)\|_\infty &= \max(\|T_Y(x)\|, \|T_Z(x)\|) \\
 &\leq \max(\|T_Y\| \|x\|, \|T_Z\| \|x\|) \\
 &\leq \max \left[ \max(\|T_Y\|, \|T_Z\|) \|x\|, \max(\|T_Y\|, \|T_Z\|) \|x\| \right] \\
 &= \max(\|T_Y\|, \|T_Z\|) \|x\| \\
 &= \|T\|_\infty \|x\|
 \end{aligned}$$

Hence

$$\begin{aligned}
 \|T\| &\leq \|T\|_\infty \\
 &= 2^{1/\infty} \|T\|_\infty
 \end{aligned}$$

□

**Exercise 6.3.8.** Let  $X$  and  $X_1, \dots, X_n$  be Banach spaces and  $p \in [0, \infty]$ . Equip  $\bigoplus_{j=1}^n X_j$  with

$\|\cdot\|_p$ . Let  $T \in L(X, \bigoplus_{j=1}^n X_j)$ . Then  $\|T\| \leq n^{1/p} \|T\|_p$ .

*Proof.* Similar to the previous exercise.

□

#### 6.4. Quotient Spaces.

**Definition 6.4.1.** Let  $X$  be a normed vector space and  $M \subset X$  a closed subspace. Define  $\|\cdot\| : X/M \rightarrow [0, \infty)$  by

$$\|x + M\| := \inf_{y \in M} \|x + y\|$$

We call  $\|\cdot\|$  the **subspace norm on  $X/M$**

**Exercise 6.4.2.** Let  $X$  be a normed vector space and  $M \subsetneq X$  a proper, closed subspace of  $X$ . Then

- (1) The previously defined subspace norm on  $X/M$  is well defined and is a norm.
- (2) For each  $\epsilon > 0$ , there exists  $x \in X$  such that  $\|x\| = 1$  and  $\|x + M\| \geq 1 - \epsilon$ .
- (3) The projection map  $\pi : X \rightarrow X/M$  defined by  $\pi(x) = x + M$  is continuous and  $\|\pi\| = 1$ .
- (4) If  $X$  is complete, then  $X/M$  is complete.

*Proof.*

- (1) Let  $x, y \in X$  and  $\alpha \in \mathbb{C}$ . Suppose that  $x + M = y + M$ . Then there exists  $m \in M$  such that  $x = y + m$ . Since  $M$  is a subspace, the map  $T : M \rightarrow M$  given by  $Tx = x + m$  is a bijection. So

$$\inf_{z \in M} \|y + m + z\| = \inf_{z \in M} \|y + z\|$$

which implies that

$$\begin{aligned} \|x + M\| &= \inf_{z \in M} \|x + z\| \\ &= \inf_{z \in M} \|y + m + z\| \\ &= \inf_{z \in M} \|y + z\| \\ &= \|y + M\| \end{aligned}$$

So  $\|\cdot\| : X/M \rightarrow [0, \infty)$  is well defined.

We observe that for each  $z, w \in M$ ,

$$\|x + y + z\| \leq \|x + w\| + \|y + w + z\|$$

Taking infimums over  $M$  with respect to  $z$  in this inequality implies that for each  $w \in M$ ,

$$\begin{aligned} \inf_{z \in M} \|x + y + z\| &\leq \inf_{z \in M} (\|x + w\| + \|y + w + z\|) \\ &= \|x + w\| + \inf_{z \in M} \|y + w + z\| \end{aligned}$$

Again we use the fact that for each  $w \in M$ ,

$$\inf_{z \in M} \|y + w + z\| = \inf_{z \in M} \|y + z\|$$

This implies that for each  $w \in M$ ,

$$\inf_{z \in M} \|x + y + z\| \leq \|x + w\| + \inf_{z \in M} \|y + z\|$$

Therefore, taking infimums over  $M$  with respect to  $w$  in this inequality yields

$$\begin{aligned}
\|x + y + M\| &= \inf_{z \in M} \|x + y + z\| \\
&\leq \inf_{w \in M} \left( \|x + w\| + \inf_{z \in M} \|y + z\| \right) \\
&= \inf_{w \in M} \|x + w\| + \inf_{z \in M} \|y + z\| \\
&= \|x + M\| + \|y + M\|
\end{aligned}$$

If  $\alpha = 0$ , then  $\alpha x = 0$ . Choosing  $z = 0 \in M$  gives  $\|\alpha x + M\| = 0 = |\alpha| \|x + M\|$ . Suppose that  $\alpha \neq 0$ . Then the map  $T : M \rightarrow M$  given by  $Tx = \alpha^{-1}x$  is a bijection and thus  $\inf_{z \in M} \|x + \alpha^{-1}z\| = \inf_{z \in M} \|x + z\|$ . Hence we have that

$$\begin{aligned}
\|\alpha x + M\| &= \inf_{z \in M} \|\alpha x + z\| \\
&= \inf_{z \in M} |\alpha| \|x + \alpha^{-1}z\| \\
&= |\alpha| \inf_{z \in M} \|x + \alpha^{-1}z\| \\
&= |\alpha| \inf_{z \in M} \|x + z\| \\
&= |\alpha| \|x + M\|
\end{aligned}$$

Suppose that  $\|x\| = 0$ . Choose a sequence  $(z_n)_{n \in \mathbb{N}} \subset M$  such that

$$\begin{aligned}
\lim_{n \rightarrow \infty} \|x - z_n\| &= \inf_{z \in M} \|x + z\| \\
&= 0
\end{aligned}$$

Then  $\lim_{n \rightarrow \infty} z_n = x$ . Since  $M$  is closed,  $x \in M$ . Hence  $x + M = 0 + M$ .

- (2) Since  $M$  is a proper subspace, there exists  $v \in X$  such that  $v \notin M$ . Then  $\|v + M\| \neq 0$ . Let  $\epsilon > 0$ . Then  $(1 - \epsilon)^{-1} \|v + M\| > \|v + M\|$ . So there exists  $z \in M$  such that

$$0 < \|v + M\| \leq \|v + z\| < (1 - \epsilon)^{-1} \|v + M\|$$

Choose  $x = \|v + z\|^{-1}(v + z)$ . Then  $\|x\| = 1$  and

$$\begin{aligned}
\|x + M\| &= \|v + z\|^{-1} \|v + z + M\| \\
&= \|v + z\|^{-1} \|v + M\| \\
&> 1 - \epsilon
\end{aligned}$$

- (3) Let  $x \in X$ . Taking  $z = 0$ , we see that  $\|\pi(x)\| = \|x + M\| \leq \|x + z\| = \|x\|$ . So  $\pi$  is bounded and in particular,

$$\sup_{\|x\|=1} \|\pi(x)\| \leq 1$$

From (2) we see that

$$\sup_{\|x\|=1} \|\pi(x)\| \geq 1$$

Hence  $\|\pi\| = 1$ .

- (4) Suppose that  $X$  is complete. Let  $(x_i + M)_{i \in \mathbb{N}} \subset X/M$ . Suppose that  $\sum_{i \in \mathbb{N}} \|x_i + M\| < \infty$ . Let  $\epsilon > 0$ . Then for each  $i \in \mathbb{N}$ , there exists  $z_i \in M$  such that  $\|x_i + z_i\| < \|x_i + M\| + \epsilon 2^{-i}$ . Define the sequence  $(a_i)_{i \in \mathbb{N}} \subset X$  by  $a_i = x_i + z_i$ . Then we have

$$\begin{aligned} \sum_{i \in \mathbb{N}} \|a_i\| &= \sum_{i \in \mathbb{N}} \|x_i + z_i\| \\ &\leq \sum_{i \in \mathbb{N}} \left( \|x_i + M\| + \epsilon 2^{-i} \right) \\ &= \sum_{i \in \mathbb{N}} \|x_i + M\| + \epsilon \end{aligned}$$

Since  $\epsilon > 0$  is arbitrary, it follows that

$$\sum_{i \in \mathbb{N}} \|a_i\| \leq \sum_{i \in \mathbb{N}} \|x_i + M\| < \infty$$

Since  $X$  is complete,  $\sum_{i=1}^{\infty} a_i$  converges in  $X$ . Define  $(s_n)_{n \in \mathbb{N}} \subset X$  and  $s \in X$  by  $s_n = \sum_{i=1}^n a_i$  and  $s = \sum_{i=1}^{\infty} a_i$ . Since  $\lim_{n \rightarrow \infty} s_n = s$ , and  $\pi : X \rightarrow X/M$  is continuous, it follows that  $\lim_{n \rightarrow \infty} \pi(s_n) = \pi(s)$ . Since

$$\begin{aligned} \pi(s_n) &= \sum_{i=1}^n a_i + M \\ &= \sum_{i=1}^n x_i + M \end{aligned}$$

We have that  $\sum_{i=1}^{\infty} x_i + M$  converges which implies that  $X/M$  is complete.

□

**Exercise 6.4.3.** Let  $X, Y$  be normed vector spaces and  $T \in L(X, Y)$ . Then

- (1)  $\ker T$  is closed
- (2) there exists a unique map  $S : X/\ker T \rightarrow T(X)$  such that  $T = S \circ \pi$ . Furthermore  $S$  is a bounded linear bijection and  $\|S\| = \|T\|$ .

*Proof.*

- (1) Since  $T$  is continuous and  $\ker T = T^{-1}(\{0\})$ , we have that  $\ker T$  is closed.
- (2) Suppose that there exists  $S_1, S_2 \in L(X/\ker T, T(X))$  such that  $T = S_1 \circ \pi$  and  $T = S_2 \circ \pi$ . Let  $x \in X$ . Then

$$S_1(x + \ker T) = S_1(\pi(x)) = T(x) = S_2(\pi(x)) = S_2(x + \ker T)$$

So  $S_1 = S_2$ . Therefore such a map is unique.

Define  $S : X/\ker T \rightarrow T(X)$  by  $S(x + \ker T) = T(x)$ . Then  $S$  is clearly a linear bijection that satisfies  $T = S \circ \pi$ . Let  $x \in X$  and  $z \in \ker T$ . Then

$$\begin{aligned}\|S(x + \ker T)\| &= \|T(x)\| \\ &= \|T(x + z)\| \\ &\leq \|T\|\|x + z\|\end{aligned}$$

Thus

$$\|S(x + \ker T)\| \leq \|T\| \inf_{z \in \ker T} \|x + z\| = \|T\|\|x + \ker T\|$$

So  $S$  is bounded and  $\|S\| \leq \|T\|$ . This implies that

$$\|T\| = \|S \circ \pi\| \leq \|S\|\|\pi\| = \|S\|$$

Thus  $\|S\| = \|T\|$ .

□

**Exercise 6.4.4.** Let  $X, Y$  be normed vector spaces. Define  $\phi : L(X, Y) \times X \rightarrow Y$  by  $\phi(T, x) = Tx$ . Then  $\phi$  is continuous.

*Proof.* Let  $(T_1, x_1) \in L(X, Y) \times X$  and  $\epsilon > 0$ . Choose  $\delta = \min\{\frac{\epsilon}{2(\|x_1\| + \|T_1\| + 1)}, \frac{\sqrt{\epsilon}}{\sqrt{2}}\}$ . Let  $(t_2, x_2) \in L(X, Y) \times X$ . Suppose that

$$\|(T_1, x_1) - (T_2, x_2)\| = \max\{\|T_1 - T_2\|, \|x_1 - x_2\|\} < \delta$$

Then

$$\begin{aligned}\|\phi(T_1, x_1) - \phi(T_2, x_2)\| &= \|T_1x_1 - T_2x_2\| \\ &= \|T_1x_1 - T_2x_1 + T_2x_1 - T_2x_2\| \\ &\leq \|(T_1 - T_2)x_1\| + \|T_2(x_1 - x_2)\| \\ &\leq \|T_1 - T_2\|\|x_1\| + \|T_2\|\|x_1 - x_2\| \\ &\leq \|T_1 - T_2\|\|x_1\| + (\|T_1 - T_2\| + \|T_1\|)\|x_1 - x_2\| \\ &< \delta\|x_1\| + (\delta + \|T_1\|)\delta \\ &= \delta(\|T_1\| + \|x_1\|) + \delta^2 \\ &< \frac{\epsilon}{2} + \frac{\epsilon}{2} \\ &= \epsilon\end{aligned}$$

So  $\phi$  is continuous.

□

**Exercise 6.4.5.** Let  $X$  be a normed vector space and  $M \subset X$  a subspace. Then  $\overline{M}$  is a subspace.

*Proof.* Let  $x, y \in \overline{M}$  and  $\alpha \in \mathbb{C}$ . Then there exist sequences  $(x_n)_{n \in \mathbb{N}} \subset M$  and  $(y_n)_{n \in \mathbb{N}} \subset M$  such that  $x_n \rightarrow x$  and  $y_n \rightarrow y$ . Since  $M$  is a subspace,  $(x_n + y_n)_{n \in \mathbb{N}} \subset M$  and  $(\alpha x_n)_{n \in \mathbb{N}} \subset M$ . Since addition and scalar multiplication are continuous, we have that  $x_n + y_n \rightarrow x + y$  and  $\alpha x_n \rightarrow \alpha x$ . Thus  $x + y \in \overline{M}$  and  $\alpha x \in \overline{M}$  and hence  $\overline{M}$  is a subspace. □



### 6.5. Applications of the Hahn-Banach Theorem.

**Definition 6.5.1.** Let  $X$  be a normed vector space over  $\mathbb{C}$ , and  $T : X \rightarrow \mathbb{C}$ . Then  $T$  is said to be a **bounded linear functional on  $X$**  if  $T \in L(X, \mathbb{C})$ . We define the **dual space of  $X$** , denoted  $X^*$ , by  $X^* = L(X, \mathbb{C})$ .

**Note 6.5.2.** We define  $X^*$  similarly when  $X$  is a normed vector space over  $\mathbb{R}$ .

**Definition 6.5.3.** Let  $X$  be a normed vector space and  $p : X \rightarrow \mathbb{R}$  a sublinear functional. Then  $p$  is said to be **bounded** if there exists  $M > 0$  such that for each  $x \in X$ ,  $p(x) \leq M\|x\|$ .

**Exercise 6.5.4.** Let  $X$  be a normed vector space and  $p : X \rightarrow \mathbb{R}$  a sublinear functional. Then  $p$  is bounded iff  $p$  is Lipschitz.

*Proof.* Suppose that  $p$  is bounded. Then there exists  $M > 0$  such that for each  $x \in X$ ,  $p(x) \leq M\|x\|$ . Let  $x, y \in X$ . Then the previous exercise implies that

$$\begin{aligned} -M\|x - y\| &= -M\|y - x\| \\ &\leq -p(y - x) \\ &\leq p(x) - p(y) \\ &\leq p(x - y) \\ &\leq M\|x - y\| \end{aligned}$$

So that

$$|p(x) - p(y)| \leq M\|x - y\|$$

and  $p$  is Lipschitz. Conversely, suppose that  $p$  is Lipschitz. Then there exists  $M > 0$  such that for each  $x, y \in X$ ,  $|p(x) - p(y)| \leq M\|x - y\|$ . Let  $x \in X$ . Then

$$\begin{aligned} p(x) &\leq |p(x)| \\ &= |p(x) - p(0)| \\ &\leq M\|x - 0\| \\ &\leq M\|x\| \end{aligned}$$

So  $p$  is bounded. □

**Exercise 6.5.5.** Let  $X$  be a normed vector space,  $p : X \rightarrow \mathbb{R}$  a bounded sublinear functional and  $\phi : X \rightarrow \mathbb{R}$  a linear functional. If  $\phi \leq p$ , then  $\phi \in X^*$ .

*Proof.* Since  $p$  is Lipschitz, there exists  $M > 0$  such that for each  $x \in X$ ,

$$\begin{aligned} p(x) &\leq |p(x)| \\ &\leq M\|x\| \end{aligned}$$

Let  $x \in X$ . Then

$$\begin{aligned} \phi(x) &\leq p(x) \\ &\leq |p(x)| \\ &\leq M\|x\| \end{aligned}$$

and therefore

$$\begin{aligned}
 -M\|x\| &= -M\|-x\| \\
 &\leq -p(-x) \\
 &\leq -\phi(-x) \\
 &= \phi(x)
 \end{aligned}$$

So that  $|\phi(x)| \leq M\|x\|$  and  $\phi \in X^*$ . □

**Exercise 6.5.6.** Let  $X$  be a normed vector space and  $p : X \rightarrow \mathbb{R}$  a bounded sublinear functional. Then there exists  $\phi \in X^*$  such that for each  $x \in X$ ,  $\phi(x) \leq p(x)$ .

*Proof.* A previous exercise implies there exists  $\phi : X \rightarrow \mathbb{R}$  such that  $\phi$  is linear and  $\phi \leq p$ . The previous exercise implies that  $\phi \in X^*$ . □

**Exercise 6.5.7. Equivalency of linearity (Bounded Case)**

Let  $X$  be a normed vector space and  $p : X \rightarrow \mathbb{R}$  a bounded sublinear functional. Then the following are equivalent:

- (1) there exists a unique  $\phi \in X^*$  such that  $\phi \leq p$
- (2) for each  $x \in X$ ,  $-p(-x) = p(x)$
- (3)  $p$  is linear

*Proof.* Basically the same as last time. □

**Exercise 6.5.8.** Let  $X$  be a normed vector space,  $M \subset X$  a subspace and  $f \in M^*$ . Then there exists  $F \in X^*$  such that  $\|F\| = \|f\|$  and  $F|_M = f$ .

*Proof.* If  $f = 0$ , Choose  $F = 0$ . Suppose  $f \neq 0$ . Then  $\|f\| \neq 0$  and there exists  $x_0 \in M$  such that  $x_0 \neq 0$ . Thus  $\|f\| \neq 0$ . Define  $p : X \rightarrow [0, \infty)$  by  $p(x) = \|f\|\|x\|$ . Then  $p$  is a sublinear functional on  $X$  and for each  $x \in M$ ,  $|f(x)| \leq p(x)$ . So there exists a linear functional  $F : X \rightarrow \mathbb{C}$  such that for each  $x \in X$ ,  $|F(x)| \leq p(x) = \|f\|\|x\|$  and  $F|_M = f$ . Thus  $F \in X^*$  with  $\|F\| \leq \|f\|$ . Also

$$\|F\| = \sup_{\substack{x \in X \\ \|x\|=1}} |F(x)| \geq \sup_{\substack{x \in M \\ \|x\|=1}} |F(x)| = \sup_{\substack{x \in M \\ \|x\|=1}} |f(x)| = \|f\|$$

So  $\|F\| = \|f\|$ . □

**Exercise 6.5.9.** Let  $X$  be a normed vector space,  $M \subsetneq X$  a proper closed subspace and  $x \in X \setminus M$ . Then there exists  $F \in X^*$  such that  $F|_M = 0$ ,  $\|F\| = 1$  and  $F(x) = \|x + M\| \neq 0$ . **Hint:** Consider  $f : M + \mathbb{C}x \rightarrow \mathbb{C}$  defined by  $f(m + \lambda x) = \lambda\|x + M\|$ .

*Proof.* Define  $f : M + \mathbb{C}x \rightarrow \mathbb{C}$  as above. Clearly  $f$  is linear and  $f|_M = 0$ . Let  $m \in M$  and  $\lambda \in \mathbb{C}$ . If  $\lambda = 0$ , then  $|f(m + \lambda x)| = 0 \leq \|m + \lambda x\|$ . Suppose that  $\lambda \neq 0$ . Then

$$\begin{aligned}
 |f(m + \lambda x)| &= |\lambda|\|x + M\| \\
 &= \|\lambda x + M\| \\
 &= \inf_{z \in M} \|z + \lambda x\| \\
 &\leq \|m + \lambda x\|
 \end{aligned}$$

So  $f \in (M + \mathbb{C}x)^*$  and  $\|f\| \leq 1$ . Let  $\epsilon > 0$ . A previous exercise tells us that there exist  $m \in M, \lambda \in \mathbb{C}$  such that  $\|m + \lambda x\| = 1$  and  $\|m + \lambda x + M\| > 1 - \epsilon$ . Then

$$\begin{aligned} |f(m + \lambda x)| &= |\lambda| \|x + M\| \\ &= \|\lambda x + M\| \\ &= \|m + \lambda x + M\| \\ &> 1 - \epsilon \end{aligned}$$

So

$$\|f\| = \sup_{\substack{z \in M + \mathbb{C}x \\ \|z\|=1}} |f(z)| \geq 1$$

Hence  $\|f\| = 1$ . The same exercise also tells us that  $f(x) = \|x + M\| \neq 0$ . Using the previous exercise, there exists  $F \in X^*$  such that  $\|F\| = \|f\| = 1$  and  $F|_{M + \mathbb{C}x} = f$ .  $\square$

**Exercise 6.5.10.** Let  $X$  be a normed vector space and  $x \in X$ . If  $x \neq 0$ , then there exists  $F \in X^*$  such that  $\|F\| = 1$  and  $F(x) = \|x\|$ .

*Proof.* Define  $f : \mathbb{C}x \rightarrow \mathbb{C}$  by  $f(\lambda x) = \lambda \|x\|$ . Then  $f$  is linear and  $f(x) = \|x\|$ . Clearly

$$\sup_{\substack{z \in \mathbb{C}x \\ \|z\|=1}} |f(z)| = 1$$

So  $f \in (\mathbb{C}x)^*$  and  $\|f\| = 1$ . By a previous exercise, there exists  $F \in X^*$  such that  $\|F\| = \|f\| = 1$  and  $F|_{\mathbb{C}x} = f$ .  $\square$

**Exercise 6.5.11.** Let  $X$  be a normed vector space and  $x \in X$ . Then  $x = 0$  iff for each  $\phi \in X^*$ ,  $\phi(x) = 0$ .

*Proof.* Clear by previous exercise.  $\square$

**Exercise 6.5.12.** Let  $X$  be a normed vector space. Then  $X^*$  separates the points of  $X$ .

*Proof.* Let  $x, y \in X$ . Suppose that  $x \neq y$ . Then  $x - y \neq 0$ . The previous exercise implies that there exists  $F \in X^*$  such that  $\|F\| = 1$  and

$$F(x) - F(y) = F(x - y) = \|x - y\| \neq 0$$

Thus  $F(x) \neq F(y)$  and  $X^*$  separates the points of  $X$ .  $\square$

**Exercise 6.5.13.** Let  $X$  be a normed vector space and  $f : X \rightarrow \mathbb{C}$  a linear functional on  $X$ . Then  $f$  is bounded iff  $\ker f$  is closed.

*Proof.* Suppose that  $f$  is continuous. Since  $\{0\}$  is closed, we have that  $\ker f = f^{-1}(\{0\})$  is closed. Conversely, suppose that  $\ker f$  is closed. If  $\ker f = X$ , then  $f = 0$  and  $f$  is continuous. Suppose that  $\ker f \neq X$ . Then  $\ker f$  is a proper, closed subspace of  $X$ . A previous exercise tells us that there exists  $x \in X$  such that  $\|x\| = 1$  and  $\|x + \ker f\| > \frac{1}{2}$ . Let  $y \in X$ . Suppose that  $\|y\| < \frac{1}{2}$ . Then for each  $z \in \ker f$ ,

$$\begin{aligned} \|z - (x + y)\| &= \|(z - x) - y\| \\ &\geq \|z - x\| - \|y\| \\ &> \frac{1}{2} - \frac{1}{2} \\ &= 0 \end{aligned}$$

So  $x + y \notin \ker f$ . Therefore  $f(B(x, \frac{1}{2})) \cap \{0\} = \emptyset$ . If  $f(B(x, \frac{1}{2}))$  is unbounded, then  $f(B(x, \frac{1}{2})) = \mathbb{C}$  by linearity. This is a contradiction since  $0 \notin f(B(x, \frac{1}{2}))$ . So There exists  $s > 0$  such that  $f(B(x, \frac{1}{2})) \subset B(0, s)$  and thus  $f$  is bounded.  $\square$

**Exercise 6.5.14.** Let  $X$  be a normed vector space.

- (1) Let  $M \subsetneq X$  be a proper closed subspace of  $X$  and  $x \in X \setminus M$ . Then  $M + \mathbb{C}x$  is closed.
- (2) Let  $M \subset X$  be a finite dimensional subspace of  $X$ . Then  $M$  is closed.

*Proof.* (1) Let  $y \in X$  and  $(y_n)_{n \in \mathbb{N}} \subset M + \mathbb{C}x$ . Suppose that  $y_n \rightarrow y$ . If  $y \in M$ , then  $y \in M + \mathbb{C}x$ . Suppose that  $y \notin M$ . For each  $n \in \mathbb{N}$ , there exists  $m_n \in M$  and  $\lambda_n \in \mathbb{C}$  such that  $y_n = m_n + \lambda_n x$ . A previous exercise tells us that there exists  $F \in X^*$  such that  $\|F\| = 1$ ,  $F|_M = 0$  and  $F(x) = \|x + M\| \neq 0$ . Since  $F$  is continuous,  $F(y_n) \rightarrow F(y)$ . Since for each  $n \in \mathbb{N}$ ,

$$F(y_n) = F(m_n + \lambda_n x) = F(m_n) + \lambda_n(Fx) = \lambda_n F(x)$$

we have that  $\lambda_n F(x) \rightarrow F(y)$ . Since  $F(x) \neq 0$ , this implies that  $\lambda_n \rightarrow F(x)^{-1}F(y)$ . It follows that  $\lambda_n x \rightarrow F(x)^{-1}F(y)x$ . Since for each  $n \in \mathbb{N}$ ,  $m_n = y_n - \lambda_n x$ , we know that  $m_n \rightarrow y - F(x)^{-1}F(y)x$ . Since  $(m_n)_{n \in \mathbb{N}} \subset M$  and  $M$  is closed, we have that  $y - F(x)^{-1}F(y)x \in M$  and therefore  $y \in M + \mathbb{C}x$ . Hence  $M + \mathbb{C}x$  is closed.

- (2) If  $M = X$ , then  $M$  is closed. Suppose that  $M \neq X$ . Let  $(x_i)_{i=1}^n$  be a basis for  $M$ . Define  $N_0 = \{0\}$  and for each  $i = 1, 2, \dots, n$ , define  $N_i = N_{i-1} + \mathbb{C}x_i$ . Since  $N_0$  is a proper closed subspace of  $X$  and  $x_1 \in X \setminus N_0$ , (1) implies that  $N_1$  is closed. Proceed inductively to obtain that  $M = N_n$  is closed.  $\square$

**Exercise 6.5.15.** Let  $X$  be an infinite-dimensional normed vector space.

- (1) There exists a sequence  $(x_n)_{n \in \mathbb{N}} \subset X$  such that for each  $m, n \in \mathbb{N}$ ,  $\|x_n\| = 1$  and if  $m \neq n$ , then  $\|x_m - x_n\| > \frac{1}{2}$ .
- (2)  $X$  is not locally compact.

*Proof.*

- (1) Define  $N_0 = \{0\}$ . Then  $N_0$  is a closed proper subspace of  $X$ . Choose  $x_1 \in X$  such that  $\|x_1\| = 1$ . Using the results of previous exercises, we proceed inductively. For each  $n \geq 2$  we define  $N_{n-1} = \text{span}(x_1, x_2, \dots, x_{n-1})$ . Then  $N_{n-1}$  is a closed proper subspace of  $X$ . Thus we may choose  $x_n \in X$  such that  $\|x_n\| = 1$  and  $\|x_n + N_{n-1}\| > \frac{1}{2}$ . Let  $m, n \in \mathbb{N}$ . Suppose that  $m < n$ . Then  $x_m \in N_{n-1}$ . Thus  $\|x_n - x_m\| \geq \|x_n + N_{n-1}\| > \frac{1}{2}$ .

- (2) Suppose that  $X$  is locally compact. Then  $\overline{B(0, 1)}$  is compact and therefore sequentially compact. Using  $(x_n)_{n \in \mathbb{N}} \subset \overline{B(0, 1)}$  defined in (1), we see that there exists a subsequence  $(x_{n_k})_{k \in \mathbb{N}}$ ,  $x \in \overline{B(0, 1)}$  such that  $x_{n_k} \rightarrow x$ . Then  $(x_{n_k})_{k \in \mathbb{N}}$  is Cauchy. So there exists  $N \in \mathbb{N}$  such that for each  $j, k \in \mathbb{N}$ , if  $j, k \geq N$ , then  $\|x_{n_j} - x_{n_k}\| < \frac{1}{2}$ . Then  $\|x_{n_N} - x_{n_{N+1}}\| < \frac{1}{2}$ . This is a contradiction since by construction,  $\|x_{n_N} - x_{n_{N+1}}\| > \frac{1}{2}$ . Thus  $X$  is not locally compact.



### 6.6. The Baire Category and Closed Graph Theorems.

**Theorem 6.6.1.** Let  $X, Y$  be Banach spaces and  $T \in L(X, Y)$ . If  $T$  is surjective, then  $T$  is open.

**Corollary 6.6.2.** Let  $X, Y$  be Banach spaces and  $T \in L(X, Y)$ . If  $T$  is a bijection, then  $T^{-1} \in L(X, Y)$ .

**Definition 6.6.3.** Let  $X, Y$  be sets and  $f : X \rightarrow Y$ . We define the **graph of  $f$** ,  $\Gamma(f)$ , by  $\Gamma(f) = \{(x, y) \in X \times Y : f(x) = y\}$ .

**Theorem 6.6.4.** Let  $X, Y$  be Banach spaces and  $T : X \rightarrow Y$  a linear map. If  $\Gamma(T)$  is closed, then  $T \in L(X, Y)$ .

**Note 6.6.5.** We recall that  $\Gamma(T)$  is closed iff for each  $(x_n)_{n \in \mathbb{N}} \subset X$ ,  $x \in X$  and  $y \in Y$ ,  $x_n \rightarrow x$  and  $T(x_n) \rightarrow y$  implies that  $T(x) = y$ .

**Theorem 6.6.6.** Let  $X, Y$  be Banach spaces and  $S \subset L(X, Y)$ . If for each  $x \in X$ ,

$$\sup_{T \in S} \|Tx\| < \infty$$

then

$$\sup_{T \in S} \|T\| < \infty$$

**Exercise 6.6.7.** Let  $\mu$  be counting measure on  $(\mathbb{N}, \mathcal{P}(\mathbb{N}))$ . Define  $h : \mathbb{N} \rightarrow \mathbb{N}$  and  $\nu$  on  $(\mathbb{N}, \mathcal{P}(\mathbb{N}))$  by  $h(n) = n$  and  $d\nu = h d\mu$ . Define  $X = L^1(\nu)$  and  $Y = L^1(\mu)$ . Equip both  $X$  and  $Y$  with the  $L^1$  norm with respect to  $\mu$ .

- (1) We have that  $X$  is a proper subspace of  $Y$  and therefore  $X$  is not complete.
- (2) Define  $T : X \rightarrow Y$  by  $Tf(n) = nf(n)$ . Then  $T$  is linear,  $\Gamma(T)$  is closed, and  $T$  is unbounded.
- (3) Define  $S : Y \rightarrow X$  by  $Sg(n) = \frac{1}{n}g(n)$ . Then  $S \in L(Y, X)$ ,  $S$  is surjective and  $S$  is not open.

*Proof.*

- (1) Note that for each  $f : \mathbb{N} \rightarrow \mathbb{C}$ ,

$$\begin{aligned} \|f\|_{\mu,1} &= \sum_{n=1}^{\infty} |f(n)| \\ &\leq \sum_{n=1}^{\infty} n|f(n)| \\ &= \|f\|_{\nu,1} \end{aligned}$$

Hence  $X$  is a subspace of  $Y$ . Define  $f : \mathbb{N} \rightarrow \mathbb{C}$  by  $f(n) = \frac{1}{n^2}$ . Then

$$\|f\|_{\mu,1} = \sum_{n=1}^{\infty} \frac{1}{n^2} < \infty$$

So  $f \in Y$ . However

$$\|f\|_{\nu,1} = \sum_{n=1}^{\infty} \frac{1}{n} = \infty$$

So  $f \notin X$ . Thus  $X$  is a proper subspace of  $Y$ . Let  $g \in Y$  and  $\epsilon > 0$ . Since the simple functions are dense in  $L^1(\mu)$ , there exists  $\phi \in L^1(\mu)$  such that  $\phi$  is simple and  $\|g - \phi\|_{\mu,1} < \epsilon$ . Then there exist  $(c_i)_{i=1}^k \subset \mathbb{C}$  and  $(E_i)_{i=1}^k \subset \mathcal{P}(\mathbb{N})$  such that for each  $i, j \in \{1, 2, \dots, k\}$ ,  $E_i$  is finite,  $i \neq j$  implies that  $E_i \cap E_j = \emptyset$  and

$$\phi = \sum_{i=1}^k c_i \chi_{E_i}$$

Define  $c = \max\{|c_i| : i = 1, 2, \dots, k\}$  and  $m = \max \left[ \bigcup_{i=1}^k E_i \right]$ . Then

$$\begin{aligned} \|\phi\|_{\nu,1} &= \sum_{n=1}^m n |\phi(n)| \\ &\leq \sum_{n=1}^m mc \\ &= cm^2 \\ &< \infty \end{aligned}$$

Hence  $\phi \in X$  and  $X$  is dense in  $Y$ . Since  $X$  is a dense, proper subspace, it is not closed. Since  $Y$  is complete and  $X \subset Y$  is not closed, we have that  $X$  is not complete.

- (2) Clearly  $T$  is linear. Let  $(f_j)_{j \in \mathbb{N}} \subset X$ ,  $f \in X$  and  $g \in Y$ . Suppose that  $f_j \xrightarrow{L^1(\mu)} f$  and  $Tf_j \xrightarrow{L^1(\mu)} g$ .

Note that for each  $j \in \mathbb{N}$  and  $n \in \mathbb{N}$ ,

$$|f_j(n) - f(n)| \leq \sum_{n=1}^{\infty} |f_j(n) - f(n)| = \|f_j - f\|_{\mu,1}$$

and

$$|nf_j(n) - g(n)| \leq \sum_{n=1}^{\infty} |nf_j(n) - g(n)| = \|Tf_j - g\|_{\mu,1}$$

Thus for each  $n \in \mathbb{N}$ ,  $f_j(n) \xrightarrow{j} f(n)$  and  $nf_j(n) \xrightarrow{j} g(n)$ . This implies that for each  $n \in \mathbb{N}$ ,  $nf(n) = g(n)$ . Thus  $Tf = g$  which implies that  $\Gamma(T)$  is closed. Suppose, for the sake of contradiction, that  $T$  is bounded. Then there exists  $C \geq 0$  such that for each  $f \in X$ ,  $\|Tf\|_{\mu,1} \leq C\|f\|_{\mu,1}$ . Choose  $n \in \mathbb{N}$  such that  $n > C$ . Define  $f : \mathbb{N} \rightarrow \mathbb{C}$  by  $f = \chi_{\{n\}}$ . As established above,  $S^+ \subset L^1(\mu)$ . Then  $\|f\|_{\mu,1} = 1$  and

$$\begin{aligned} \|Tf\|_{\mu,1} &= n \\ &> C \\ &= C\|f\|_{\mu,1} \end{aligned}$$

which is a contradiction. So  $T$  is unbounded.

(3) Clearly  $S$  is linear. Let  $g \in Y$ . Then

$$\begin{aligned}\|Sg\|_{\mu,1} &= \sum_{n=1}^{\infty} \frac{1}{n} |g(n)| \\ &\leq \sum_{n=1}^{\infty} |g(n)| \\ &= \|g\|_{\mu,1}\end{aligned}$$

So  $S$  is bounded and  $\|S\| \leq 1$ . Thus  $S \in L(Y, X)$ . Let  $f \in X$ . Define  $g : \mathbb{N} \rightarrow \mathbb{C}$  by  $g(n) = nf(n)$ . By definition,  $g \in Y$  and we have that

$$\begin{aligned}Sg(n) &= \frac{1}{n} g(n) \\ &= f(n)\end{aligned}$$

Hence  $Sg = f$  and thus  $S$  is surjective. Let  $g \in Y$ . Suppose that  $Sg = 0$ . Then

$$\sum_{n=1}^{\infty} \frac{1}{n} |g(n)| = \|Sg\| = 0$$

Thus for each  $n \in \mathbb{N}$ ,  $g(n) = 0$ . Hence  $\ker S = \{0\}$  and  $S$  is injective. Note that for each  $A \subset Y$ ,  $S(A) = T^{-1}(A)$ . If  $S$  is open, then  $T$  is continuous which as shown above is a contradiction. So  $g$  is not open. □

**Exercise 6.6.8.** Let  $X = C^1([0, 1])$  and  $Y = C([0, 1])$ . Equip both  $X$  and  $Y$  with the uniform norm.

(1) Then  $X$  is not complete

(2) Define  $T : X \rightarrow Y$  by  $Tf = f'$ . Then  $\Gamma(T)$  is closed and  $T$  is not bounded.

*Proof.* (1) Recall that for each  $a, b \geq 0$  and  $p \in \mathbb{N}$ ,

$$(a^{\frac{1}{p}} + b^{\frac{1}{p}})^p = \sum_{n=0}^p \binom{p}{n} a^{\frac{n}{p}} b^{\frac{p-n}{p}} \geq a + b$$

Thus  $(a + b)^{\frac{1}{p}} \leq a^{\frac{1}{p}} + b^{\frac{1}{p}}$ .

For each  $n \in \mathbb{N}$ , define  $f_n : [0, 1] \rightarrow \mathbb{C}$  by  $f_n(x) = \sqrt{(x - \frac{1}{2})^2 + \frac{1}{n^2}}$ . Then  $(f_n)_{n \in \mathbb{N}} \subset X$ . Define  $f : [0, 1] \rightarrow \mathbb{C}$  by  $f(x) = |x - \frac{1}{2}|$ . Then  $f \in Y \cap X^c$ . Note that for each  $n \in \mathbb{N}$ ,  $f \leq f_n$ . Our observation above implies that for each  $x \in X$ ,

$$\begin{aligned}f_n(x) &= \left[ (x - \frac{1}{2})^2 + \frac{1}{n^2} \right]^{\frac{1}{2}} \\ &\leq |x - \frac{1}{2}| + \frac{1}{n}\end{aligned}$$

Thus  $0 \leq f_n - f \leq \frac{1}{n}$ . This implies that  $f_n \xrightarrow{u} f$ . Since  $f \notin X$ ,  $X$  is not complete.



- (2) Let  $(f_n)_{n \in \mathbb{N}} \subset X$ ,  $f \in X$  and  $g \in Y$ . Suppose that  $f_n \xrightarrow{u} f$  and  $Tf_n \xrightarrow{u} g$ . Let  $x \in [0, 1]$ . Then  $f_n(x) \rightarrow f(x)$  and  $f_n(0) \rightarrow f(0)$  and  $f'_n \xrightarrow{u} g$ . Applying the DCT to this sequence of integrable functions that converges uniformly to an integrable function on a finite measure space (a previous exercise) we have that

$$\begin{aligned} f_n(x) - f_n(0) &= \int_{[0,x]} f'_n dm \\ &\rightarrow \int_{[0,x]} g dm \end{aligned}$$

Since  $f_n(x) - f_n(0) \rightarrow f(x) - f(0)$ , we know that

$$f(x) - f(0) = \int_{[0,x]} g dm$$

Thus  $Tf = g$  and  $\Gamma(T)$  is closed.

By Exercise 6.2.3,  $T$  is not bounded. □

**Exercise 6.6.9.** Let  $X, Y$  be Banach spaces and  $T \in L(X, Y)$ . Then  $X/\ker T \cong T(X)$  iff  $T(X)$  is closed.

*Proof.* Since  $X$  is a Banach space and  $T$  is continuous, we have that  $\ker T$  is closed and  $X/\ker T$  is a Banach space. Suppose that  $X/\ker T \cong T(X)$ . Then  $T(X)$  is complete. Since  $Y$  is complete, this implies that  $T(X)$  is closed.

Conversely Suppose that  $T(X)$  is closed. Then  $T(X)$  is complete. Define  $S : X/\ker T \rightarrow T(X)$  by  $S(x + \ker T) = T(x)$ . A previous exercise tells us that the map  $S : X/\ker T \rightarrow T(X)$  defined by  $S(x + \ker T) = T(x)$  is a bounded linear bijection. Since  $T(X)$  is complete and  $S$  is surjective,  $S^{-1}$  is bounded and thus  $S$  is an isomorphism. □

**Exercise 6.6.10.** Let  $X$  be a separable Banach space. Define  $B_X = \{x \in X : \|x\| < 1\}$ . Let  $(x_n)_{n \in \mathbb{N}} \subset B_X$  a dense subset of the unit ball and  $\mu$  the counting measure on  $(\mathbb{N}, \mathcal{P}(\mathbb{N}))$ . Define  $T : L^1(\mu) \rightarrow X$  by

$$Tf = \sum_{n=1}^{\infty} f(n)x_n$$

Then

- (1)  $T$  is well defined and  $T \in L(L^1(\mu), X)$
- (2)  $T$  is surjective
- (3) There exists a closed subspace  $K \subset L^1(\mu)$  such that  $L^1(\mu)/K \cong X$

*Proof.* (1) Let  $f \in L^1(\mu)$ . Since  $X$  is complete and

$$\begin{aligned} \sum_{n=1}^{\infty} \|f(n)x_n\| &= \sum_{n=1}^{\infty} |f(n)| \|x_n\| \\ &\leq \sum_{n=1}^{\infty} |f(n)| \\ &< \infty \end{aligned}$$

we have that  $\sum_{n=1}^{\infty} f(n)x_n$  converges and thus  $Tf \in X$ . Hence  $T$  is well defined.

Clearly  $T$  is linear. Let  $f \in L^1(\mu)$ . Then

$$\begin{aligned} \|Tf\| &= \left\| \sum_{n=1}^{\infty} f(n)x_n \right\| \\ &\leq \sum_{n=1}^{\infty} \|f(n)x_n\| \\ &\leq \sum_{n=1}^{\infty} |f(n)| \\ &= \|f\|_1 \end{aligned}$$

So  $T$  is bounded with  $\|T\| \leq 1$ .

- (2) Let  $x \in X$ . Suppose that  $\|x\| < 1$ . Then  $x \in B_X$ . So there exists  $n_1 \in \mathbb{N}$  such that  $\|x - x_{n_1}\| < \frac{1}{2}$ . Then  $2(x - x_{n_1}) \in B_X$ . Since for each  $j \in \mathbb{N}$ ,  $B_X \setminus (x_n)_{n=1}^j$  is dense in  $B_X$ , there exists  $n_2 \in \mathbb{N}$  such that  $x_{n_2} \notin (x_n)_{n=1}^{n_1}$  and  $\|2(x - x_{n_1}) - x_{n_2}\| < \frac{1}{2}$  which implies that  $\|x - (x_{n_1} - \frac{1}{2}x_{n_2})\| < \frac{1}{4}$ .

Proceed inductively to obtain a subsequence  $(x_{n_k})_{k \in \mathbb{N}}$  such that for each  $k \geq 2$ ,  $x_{n_k} \notin (x_n)_{n=1}^{n_{k-1}}$  and  $\|x - \sum_{j=1}^k 2^{1-j}x_{n_j}\| < \frac{1}{2^k}$ . Then  $x = \sum_{k=1}^{\infty} 2^{1-k}x_{n_k}$ .

Define  $f : \mathbb{N} \rightarrow \mathbb{C}$  by  $f = \sum_{k=1}^{\infty} 2^{1-k}\chi_{\{n_k\}}$ . Then  $\|f\|_1 = \sum_{k=1}^{\infty} 2^{1-k} < \infty$ , so  $f \in L^1(\mu)$  and  $Tf = \sum_{k=1}^{\infty} 2^{1-k}x_{n_k} = x$ . Now, suppose that  $\|x\| \geq 1$ , then  $\frac{1}{2\|x\|}x \in B_X$ .

The above argument shows that there exists  $f \in L^1(\mu)$  such that  $Tf = \frac{1}{2\|x\|}x$ . Then

$$2\|x\|f \in L^1(\mu) \text{ and } T(2\|x\|f) = 2\|x\|Tf = x.$$

So for each  $x \in X$ , there exists  $f \in L^1(\mu)$  such that  $Tf = x$  and thus  $T$  is surjective.

- (3) Since  $X$  is a Banach space and  $T$  is surjective, the previous exercise implies that  $L^1(\mu)/\ker T \cong X$ .

□

## 6.7. Duality.

**Note 6.7.1.** Let  $X$  be a normed vector space. Then  $X^*$  is a normed vector space. In general the weak-\* topology on  $X^*$  is not necessarily the same as the norm topology on  $X^*$ . In the context of normed vector spaces, we will write  $X^{**}$  to denote  $(X^*)^*$  when  $X^*$  is equipped with the norm topology and  $\hat{X}$  to denote  $(X^*)^*$  when  $X^*$  is equipped with the weak-\* topology.

**Exercise 6.7.2.** Let  $X$  be a normed vector space and  $x \in X$ . Define  $\hat{x} : X^* \rightarrow \mathbb{C}$  by  $\hat{x}(f) = f(x)$ . Then  $\hat{x} \in X^{**}$  and  $\|\hat{x}\| = \|x\|$ .

**Hint:** Hahn-Banach theorem

*Proof.* Let  $f, g \in X^*$  and  $\lambda \in \mathbb{C}$ . Then

$$\hat{x}(f + \lambda g) = (f + \lambda g)(x) = f(x) + \lambda g(x) = \hat{x}(f) + \lambda \hat{x}(g)$$

So  $\hat{x}$  is linear. For each  $f \in X^*$ ,

$$|\hat{x}(f)| = |f(x)| \leq \|x\| \|f\|$$

Hence  $\hat{x} \in X^{**}$  with  $\|\hat{x}\| \leq \|x\|$ . If  $x = 0$ , then  $\hat{x} = 0$  and  $\|\hat{x}\| = \|x\|$ . Suppose that  $x \neq 0$ . Then a previous exercise implies that there exists  $F \in X^*$  such that  $\|F\| = 1$  and  $F(x) = \|x\|$ . Then we have that

$$\sup_{\substack{f \in X^* \\ \|f\|=1}} |\hat{x}(f)| = \sup_{\substack{f \in X^* \\ \|f\|=1}} |f(x)| \geq |F(x)| = \|x\|$$

Hence  $\|\hat{x}\| = \|x\|$ . □

**Exercise 6.7.3.** Let  $X$  be a normed vector space. If  $X$  is separable, then there exist  $(\phi_n)_{n \in \mathbb{N}} \subset X^*$  such that for each  $n \in \mathbb{N}$ ,  $\|\phi_n\| = 1$  and for each  $x \in X$ ,

$$\|x\| = \sup_{n \in \mathbb{N}} |\phi_n(x)|$$

**Hint:** Let  $(x_n)_{n \in \mathbb{N}} \subset X$  be a dense subset. A previous exercise on the Hahn-Banach theorem implies that for each  $n$ , there exists  $\phi_n \in X^*$  such that  $\|\phi_n\| = 1$  and  $\phi_n(x_n) = \|x_n\|$ . Then for each  $x \in X$ ,

$$\|x\| = \sup_{n \in \mathbb{N}} |\phi_n(x)|$$

*Proof.* Suppose that  $X$  is separable. Then there exists  $(x_n)_{n \in \mathbb{N}} \subset X$  such that  $(x_n)_{n \in \mathbb{N}}$  is dense in  $X$ . A previous exercise on the Hahn-Banach theorem implies that for each  $n$ , there exists  $\phi_n \in X^*$  such that  $\|\phi_n\| = 1$  and  $\phi_n(x_n) = \|x_n\|$ . Let  $x \in X$ . Then

$$\begin{aligned} \|x\| &= \|\hat{x}\| \\ &= \sup_{\substack{\phi \in X^* \\ \|\phi\|=1}} \|\hat{x}(\phi)\| \\ &= \sup_{\substack{\phi \in X^* \\ \|\phi\|=1}} \|\phi(x)\| \\ &\geq \sup_{n \in \mathbb{N}} \|\phi_n(x)\| \end{aligned}$$

Let  $\epsilon > 0$ . Choose  $N \in \mathbb{N}$  such that  $\|x - x_N\| < \epsilon/2$ . Then

$$\begin{aligned}
 \|x\| &\leq \|x - x_N\| + \|x_N\| \\
 &= \|x - x_N\| + |\phi_N(x_N)| \\
 &\leq \|x - x_N\| + |\phi_N(x_N - x)| + |\phi_N(x)| \\
 &\leq \|x - x_N\| + \|\phi_N\| \|x_N - x\| + |\phi_N(x)| \\
 &\leq 2\|x - x_N\| + |\phi_N(x)| \\
 &< 2\frac{\epsilon}{2} + |\phi_N(x)| \\
 &\leq \epsilon + \sup_{n \in \mathbb{N}} |\phi_n(x)|
 \end{aligned}$$

Since  $\epsilon > 0$  is arbitrary,  $\|x\| \leq \sup_{n \in \mathbb{N}} |\phi_n(x)|$ . So  $\|x\| = \sup_{n \in \mathbb{N}} |\phi_n(x)|$ . □

**Exercise 6.7.4.** Let  $X$  be a normed vector space. Define  $\phi : X \rightarrow X^{**}$  by  $\phi(x) = \hat{x}$ . Then  $\phi$  is a linear isometry.

*Proof.* Let  $x, y \in X$  and  $\lambda \in \mathbb{C}$ . Then for each  $f \in X^*$ , we have that

$$\begin{aligned}
 \phi(x + \lambda y)(f) &= \widehat{x + \lambda y}(f) \\
 &= f(x + \lambda y) \\
 &= f(x) + \lambda f(y) \\
 &= \hat{x}(f) + \lambda \hat{y}(f) \\
 &= \phi(x)(f) + \lambda \phi(y)(f)
 \end{aligned}$$

So  $\phi(x + \lambda y) = \phi(x) + \lambda \phi(y)$  and  $\phi$  is linear. The previous exercise tells us that

$$\begin{aligned}
 \|\phi(x) - \phi(y)\| &= \|\phi(x - y)\| \\
 &= \|\widehat{x - y}\| = \|x - y\|
 \end{aligned}$$

So  $\phi$  is an isometry. □

**Definition 6.7.5.** Let  $X$  be a normed vector space and define  $\phi : X \rightarrow X^{**}$  as above. We define  $\hat{X} = \phi(X) \subset X^{**}$ . Since  $\hat{X}$  and  $X$  are isomorphic, we may identify  $X$  as a subset of  $X^{**}$ .

**Definition 6.7.6.** Let  $X$  be a normed vector space and define  $\phi : X \rightarrow X^{**}$  as above. Then  $X$  is said to be **reflexive** if  $\phi$  is surjective. In this case  $\phi$  is then an isomorphism

**Definition 6.7.7.** Let  $X, Y$  be normed vector spaces and  $T \in L(X, Y)$ . Define the **adjoint of  $T$** , denoted  $T^* : Y^* \rightarrow X^*$ , by  $T^*(f) = f \circ T$ .

**Exercise 6.7.8.** Let  $X, Y$  be normed vector spaces and  $T \in L(X, Y)$ .

- (1) Then  $T^* \in L(Y^*, X^*)$ .
- (2) Applying the result from (1) twice, we have that  $T^{**} \in L(X^{**}, Y^{**})$ . We have that for each  $x \in X$ ,  $T^{**}(\hat{x}) = \widehat{T(x)}$ .
- (3)  $T^*$  is injective iff  $T(X)$  is dense in  $Y$ .
- (4) If  $T^*(Y^*)$  is dense in  $X^*$ , then  $T$  is injective. The converse is true if  $X$  is reflexive.

*Proof.*

(1) Let  $f \in Y^*$ . Then  $\|T^*(f)\| = \|f \circ T\| \leq \|T\|\|f\|$ . So  $T^* \in L(Y^*, X^*)$  with  $\|T^*\| \leq \|T\|$ .

(2) Let  $x \in X$ . Let  $f \in Y^*$ . Then

$$\begin{aligned} T^{**}(\hat{x})(f) &= \hat{x} \circ T^*(f) \\ &= \hat{x}(T^*(f)) \\ &= \hat{x}(f \circ T) \\ &= f \circ T(x) \\ &= f(T(x)) \\ &= \widehat{T(x)}(f) \end{aligned}$$

Hence  $T^{**}(\hat{x}) = \widehat{T(x)}$ .

(3) Suppose that  $T(X)$  is not dense in  $Y$ . Then  $\overline{T(X)} \neq Y$ . So  $T(X)$  is a proper closed subspace of  $Y$  and there exists  $y \in Y$  such that  $y \notin \overline{T(X)}$ . By a previous exercise, there exists  $f \in Y^*$  such that  $f(y) = \|y + \overline{T(X)}\| \neq 0$ ,  $\|f\| = 1$  and  $f|_{\overline{T(X)}} = 0$ . Let  $x \in X$ . Then  $T^*(f)(x) = f \circ T(x) = 0$ . Hence  $T^*(f) = 0 = T^*(0)$ . Since  $f \neq 0$ ,  $T^*$  is not injective.

Now suppose that  $T(X)$  is dense in  $Y$ . Let  $f, g \in Y^*$ . Define  $h \in Y^*$  by  $h = f - g$ . Suppose that  $T^*(f) = T^*(g)$ . Then  $T^*(h) = 0$ . So for each  $x \in X$ ,  $h(T(x)) = 0$ . Let  $y \in Y$  and  $\epsilon > 0$ . By continuity, there exists  $\delta > 0$  such that for each  $y' \in Y$ , if  $\|y - y'\| < \delta$ , then  $\|h(y) - h(y')\| < \epsilon$ . Since  $T(X)$  is dense in  $Y$ , there exists  $x \in X$  such that  $\|y - T(x)\| < \delta$ . Thus

$$\begin{aligned} \|h(y)\| &\leq \|h(y) - h(T(x))\| + \|h(T(x))\| \\ &= \|h(y) - h(T(x))\| \\ &< \epsilon \end{aligned}$$

Since  $\epsilon > 0$  is arbitrary,  $\|h(y)\| = 0$ . This implies that  $h(y) = 0$  and therefore  $f(y) = g(y)$ . Since  $y \in Y$  is arbitrary,  $f = g$  and  $T^*$  is injective.

(4) For the sake of contradiction, suppose that  $T^*(Y^*)$  is dense in  $X^*$  and  $T$  is not injective. Then there exist  $x_1, x_2 \in X$  such that  $x_1 \neq x_2$  and  $T(x_1) = T(x_2)$ . Define  $x = x_1 - x_2$ . Then  $x \neq 0$  and  $T(x) = 0$ . A previous exercise implies that there exists  $F \in X^*$  such that  $F(x) = \|x\| \neq 0$  and  $\|F\| = 1$ . Let  $\epsilon > 0$ . Choose  $g \in Y^*$  such that  $\|F - T^*(g)\| < \epsilon$ . Then

$$\begin{aligned} \|x\| &= |F(x)| \\ &\leq |F(x) - T^*(g)(x)| + |T^*(g)(x)| \\ &< \epsilon\|x\| + |g(T(x))| \\ &= \epsilon\|x\| \end{aligned}$$

Since  $\epsilon > 0$  is arbitrary, we have that  $\|x\| = 0$  which is a contradiction. Hence if  $T^*(Y^*)$  is dense in  $X^*$ , then  $T$  is injective.

Now, suppose that  $X$  is reflexive and  $T$  is injective. Let  $\phi_1, \phi_2 \in X^{**}$ . Suppose that  $T^{**}(\phi_1) = T^{**}(\phi_2)$ . Then  $T^{**}(\phi_1 - \phi_2) = 0$ . Since  $X$  is reflexive, there exist  $x_1, x_2 \in X$  such that  $\phi_1 = \hat{x}_1$  and  $\phi_2 = \hat{x}_2$ . Define  $x = x_1 - x_2$ . Then  $T^{**}(\hat{x}) = 0$ . So for each  $f \in Y^*$ ,

$$\begin{aligned} T^{**}(\hat{x})(f) &= \hat{x} \circ T^*(f) \\ &= \hat{x}(T^*(f)) \\ &= \hat{x}(f \circ T) \\ &= f \circ T(x) \\ &= f(T(x)) \\ &= 0 \end{aligned}$$

Suppose that  $T(x) \neq 0$ . Then a previous exercise implies that there exists  $g \in Y^*$  such that  $g(T(x)) = \|T(x)\| \neq 0$  and  $\|g\| = 1$ . This is a contradiction since  $g(T(x)) = 0$ . So  $T(x) = 0$ . Since  $T$  is injective, this implies that  $x = 0$ . Hence  $\hat{x} = 0$  and thus  $\phi_1 = \phi_2$ . Thus  $T^{**}$  is injective. By (3), we have that  $T^*(Y^*)$  is dense in  $X^*$ . □

**Exercise 6.7.9.** Let  $X$  be a normed vector space. Then  $X$  is reflexive iff  $X^*$  is reflexive.

*Proof.* Suppose that  $X$  is reflexive. Let  $\alpha \in X^{***}$ . Define  $f : X \rightarrow \mathbb{C}$  by  $f(x) = \alpha(\hat{x})$ . Clearly  $f$  is linear and a previous exercise tells us that for each  $x \in X$ ,

$$\begin{aligned} |f(x)| &\leq \|\alpha\| \|\hat{x}\| \\ &= \|\alpha\| \|x\| \end{aligned}$$

So  $f \in X^*$ . Let  $\phi \in X^{**}$ . Since  $X$  is reflexive, there exists  $x \in X$  such that  $\phi = \hat{x}$ . Then

$$\begin{aligned} \alpha(\phi) &= \alpha(\hat{x}) \\ &= f(x) \\ &= \hat{x}(f) \\ &= \hat{f}(\hat{x}) \\ &= \hat{f}(\phi) \end{aligned}$$

Hence  $\alpha = \hat{f}$ . Thus the map  $X^* \rightarrow X^{***}$  given by  $f \mapsto \hat{f}$  is surjective and so  $X^*$  is reflexive.

Conversely, suppose that  $X^*$  is reflexive. Since  $\phi : X \rightarrow X^{**}$  given by  $\phi(x) = \hat{x}$  is an isometry,  $\hat{X} \subset X^{**}$  is closed. For the sake of contradiction, suppose that  $\hat{X} \neq X^{**}$ . Then there exists  $\alpha \in X^{**}$  such that  $\alpha \notin \hat{X}$ . Thus there exists  $F \in X^{***}$  such that  $\|F\| = 1$ ,  $F(\alpha) = \|\alpha + \hat{X}\| \neq 0$  and  $F|_{\hat{X}} = 0$ . Since  $X^*$  is reflexive, there exists  $f \in X^*$  such that  $F = \hat{f}$ . A previous exercise tells us that  $\|f\| = \|\hat{f}\| = \|F\| = 1$ . Since for each  $x \in X$ ,  $f(x) = \hat{x}(f) = \hat{f}(\hat{x}) = F(\hat{x}) = 0$ , we have that  $f = 0$ . Thus  $\|f\| = 0$ , a contradiction. So  $\hat{X} = X^{**}$  and  $X$  is reflexive. □

**Definition 6.7.10.** Let  $X$  be a normed vector space,  $M \subset X$  and  $N \subset X^*$ . We define the **annihilator** of  $M$  and the annihilator of  $N$ , denoted by  $M^\perp \subset X^*$  and  ${}^\perp N \subset X$  respectively, by

$$\begin{aligned} M^\perp &= \{\phi \in X^* : \text{for each } x \in M, \phi(x) = 0\} \\ {}^\perp N &= \{x \in X : \text{for each } \phi \in N, \phi(x) = 0\} \end{aligned}$$

**Exercise 6.7.11.** Let  $X$  be a normed vector space,  $M \subset X$  and  $N \subset X^*$ . Then

(1)

$$M^\perp = \bigcap_{x \in M} \ker \hat{x}$$

(2)

$${}^\perp N = \bigcap_{\phi \in N} \ker \phi$$

*Proof.*

(1)

$$\begin{aligned} M^\perp &= \{\phi \in X^* : \text{for each } x \in M, \phi(x) = 0\} \\ &= \bigcap_{x \in M} \{\phi \in X^* : \phi(x) = 0\} \\ &= \bigcap_{x \in M} \{\phi \in X^* : \hat{x}(\phi) = 0\} \\ &= \bigcap_{x \in M} \ker \hat{x} \end{aligned}$$

(2)

$$\begin{aligned} {}^\perp N &= \{x \in X : \text{for each } \phi \in N, \phi(x) = 0\} \\ &= \bigcap_{\phi \in N} \{x \in X : \phi(x) = 0\} \\ &= \bigcap_{\phi \in N} \ker \phi \end{aligned}$$

□

**Exercise 6.7.12.** Let  $X$  be a normed vector space,  $M \subset X$  and  $N \subset X^*$ . Then

(1)  $M^\perp$  is weak-\* closed

(2)  ${}^\perp N$  is closed

*Proof.*

(1) Let  $(\phi_n)_{n \in \mathbb{N}} \subset M^\perp$  and  $\phi \in X^*$ . Suppose that  $\phi_n \xrightarrow{w^*} \phi$ . Then for each  $x \in X$ ,  $\phi_n(x) \rightarrow \phi(x)$ . Let  $x \in M$ . By definition, for each  $n \in \mathbb{N}$ ,  $\phi_n(x) = 0$ . Thus  $\phi_n(x) \rightarrow 0$  which implies that  $\phi(x) = 0$  and  $\phi \in \ker \hat{x}$ . Since  $x \in M$  is arbitrary,

$$\begin{aligned} \phi &\in \bigcap_{x \in M} \ker \hat{x} \\ &= M^\perp \end{aligned}$$

- (2) Let  $(x_n)_{n \in \mathbb{N}} \subset {}^\perp N$  and  $x \in X$ . Suppose that  $x_n \rightarrow x$ . Let  $\phi \in N$ . Continuity implies that  $\phi(x_n) \rightarrow \phi(x)$ . By definition, for each  $n \in \mathbb{N}$ ,  $\phi(x_n) = 0$ . Thus  $\phi(x_n) \rightarrow 0$  which implies that  $\phi(x) = 0$ . So  $x \in \ker \phi$ . Since  $\phi \in N$  is arbitrary,

$$\begin{aligned} x &\in \bigcap_{\phi \in N} \ker \phi \\ &= {}^\perp N \end{aligned}$$

□

**Exercise 6.7.13.** Let  $X$  be a normed vector space,  $M \subset X$  and  $N \subset X^*$ . Then

- (1)  ${}^\perp(M^\perp) = \bar{M}$ , i.e. the norm closure of  $M$   
 (2)  $({}^\perp N)^\perp = \text{cl}_{w^*}(N)$ , i.e. the weak-\* closure of  $N$ .

*Proof.*

- (1) Let  $x \in M$ , then by definition, for each  $\phi \in M^\perp$ ,  $\phi(x) = 0$ . Again by definition,  $x \in {}^\perp(M^\perp)$ . So  $M \subset {}^\perp(M^\perp)$ . Since  ${}^\perp(M^\perp)$  is closed,  $\bar{M} \subset {}^\perp(M^\perp)$ . For the sake of contradiction, suppose that  ${}^\perp(M^\perp) \not\subset \bar{M}$ . Then there exists  $x \in {}^\perp(M^\perp)$  such that  $x \notin \bar{M}$ . Exercise 6.5.9 implies that there exists  $\phi \in X^*$  such that  $\phi|_{\bar{M}} = 0$ ,  $\|\phi\| = 1$  and  $\phi(x) = \|x + \bar{M}\| > 0$ . By definition,  $\phi \in M^\perp$ . Since  $\phi(x) \neq 0$ , we have that  $x \notin {}^\perp(M^\perp)$ . This is a contradiction and so  ${}^\perp(M^\perp) \subset \bar{M}$ .
- (2)

□



**6.8. Compact Operators.****Definition 6.8.1.**

### 6.9. Multilinear Maps.

**Definition 6.9.1.** Let  $X_1, \dots, X_n, Y$  be normed vector spaces and  $T : \prod_{i=1}^n X_i \rightarrow Y$  multilinear. Then  $T$  is said to be **bounded** if there exists  $C \geq 0$  such that for each  $x_1, \dots, x_n \in X$ ,

$$\|T(x_1, \dots, x_n)\| \leq C\|x_1\| \cdots \|x_n\|$$

We define

$$L^n(X_1, \dots, X_n; Y) = \left\{ T : \prod_{i=1}^n X_i \rightarrow Y : T \text{ is multilinear and bounded} \right\}$$

If  $X_1 = \dots = X_n = X$ , we write  $L^n(X, Y)$  in place of  $L^n(X, \dots, X; Y)$ . If  $X_1 = \dots = X_n = Y = X$ , we write  $L^n(X)$ .

**Note 6.9.2.** For the remainder of this section we will primarily consider  $L^2(X_1, X_2; Y)$  to avoid notational clutter, but all results immediately generalize to  $L^n(X_1, \dots, X_n; Y)$

**Exercise 6.9.3.** Let  $X_1, X_2$  and  $Y$  be normed vector spaces and  $T : X_1 \times X_2 \rightarrow Y$  bilinear. Then the following are equivalent:

- (1)  $T$  is continuous
- (2)  $T$  is continuous at  $(0, 0)$
- (3)  $T$  is bounded

*Proof.*

- (1)  $\implies$  (2):

Trivial

- (2)  $\implies$  (3):

Suppose that  $T$  is continuous at  $(0, 0)$ . For the sake of contradiction, suppose that  $T$  is not bounded. Then for each  $C \geq 0$ , there exist  $(x_1, x_2) \in X_1 \times X_2$  such that  $\|T(x_1, x_2)\| > C\|x_1\|\|x_2\|$ . Hence there exist  $(a_n)_{n \in \mathbb{N}} \subset X_1$  and  $(b_n)_{n \in \mathbb{N}} \subset X_2$  such that for each  $n \in \mathbb{N}$ ,  $\|T(a_n, b_n)\| > n^2\|a_n\|\|b_n\|$ . Hence for each  $n \in \mathbb{N}$ ,  $\|a_n\|, \|b_n\| > 0$ . Define

$$(a'_n)_{n \in \mathbb{N}} \subset X_1$$

and  $(b'_n)_{n \in \mathbb{N}} \subset X_2$  by  $a'_n = \frac{a_n}{n\|a_n\|}$  and  $b'_n = \frac{b_n}{n\|b_n\|}$ . Then  $(a'_n, b'_n) \rightarrow (0, 0)$ . Continuity implies that  $T(a'_n, b'_n) \rightarrow 0$ . By construction, for each  $n \in \mathbb{N}$ ,

$$\begin{aligned} \|T(a'_n, b'_n)\| &= \frac{1}{n^2\|a_n\|\|b_n\|} \|T(a_n, b_n)\| \\ &> \frac{n^2\|a_n\|\|b_n\|}{n^2\|a_n\|\|b_n\|} \\ &= 1 \end{aligned}$$

which is a contradiction. So  $T$  is bounded.

- (3)  $\implies$  (1):

Suppose that  $T$  is bounded. Then there exists  $C > 0$  such that for each  $(x_1, x_2) \in X_1 \times X_2$ ,  $\|T(x_1, x_2)\| \leq C\|x_1\|\|x_2\|$ . Let  $(a, b) \in X_1 \times X_2$  and  $(a_n, b_n)_{n \in \mathbb{N}} \subset X_1 \times X_2$ .

Suppose that  $(a_n, b_n) \rightarrow (a, b)$ . Then  $a_n \rightarrow a$ ,  $b_n \rightarrow b$  and  $(a_n)_{n \in \mathbb{N}}$ ,  $(b_n)_{n \in \mathbb{N}}$  are bounded. So there exists  $B \geq 0$  such that for each  $n \in \mathbb{N}$   $\|b_n\| \leq B$ . Hence

$$\begin{aligned} \|T(a_n, b_n) - T(a, b)\| &= \|T(a_n, b_n) - T(a, b_n) + T(a, b_n) - T(a, b)\| \\ &\leq \|T(a_n, b_n) - T(a, b_n)\| + \|T(a, b_n) - T(a, b)\| \\ &= \|T(a_n - a, b_n)\| + \|T(a, b_n - b)\| \\ &\leq C(\|a_n - a\|\|b_n\| + \|a\|\|b_n - b\|) \\ &\leq C(\|a_n - a\|B + \|a\|\|b_n - b\|) \\ &\rightarrow 0 \end{aligned}$$

Thus  $T$  is continuous. □

**Definition 6.9.4.** Let  $X_1, X_2$  and  $Y$  be normed vector spaces and  $T \in L^2(X_1, X_2; Y)$ . We define the **operator norm** on  $L^2(X_1, X_2; Y)$ , denoted  $\|\cdot\| : L^2(X_1, X_2; Y) \rightarrow [0, \infty)$ , by

$$\|T\| = \inf\{C \geq 0 : \text{for each } (x_1, x_2) \in X_1 \times X_2, \|T(x_1, x_2)\| \leq C\|x_1\|\|x_2\|\}$$

**Exercise 6.9.5.** Let  $X_1, X_2$  and  $Y$  be normed vector spaces. If  $X_1 \neq \{0\}$  and  $X_2 \neq \{0\}$ , then the operator norm on  $L(X, Y)$  is given by:

$$\begin{aligned} (1) \quad \|T\| &= \sup_{\|x_1\|=1, \|x_2\|=1} \|T(x_1, x_2)\| \\ (2) \quad \|T\| &= \sup_{x_1 \neq 0, x_2 \neq 0} \|x_1\|^{-1} \|x_2\|^{-1} \|T(x_1, x_2)\| \\ (3) \quad \|T\| &= \inf\{C \geq 0 : \text{for each } (x_1, x_2) \in X_1 \times X_2, \|T(x_1, x_2)\| \leq C\|x_1\|\|x_2\|\} \end{aligned}$$

*Proof.* Since  $X_1 \neq \{0\}$  and  $X_2 \neq \{0\}$ , the supremums in (1) and (2) are well defined. Let  $T \in L^2(X_1, X_2; Y)$ . Bilinearity of  $T$  implies that the sets over which the supremums are taken in (1) and (2) are the same. So (1) and (2) are equal.

Now, set

$$M = \sup_{\|x_1\|=1, \|x_2\|=1} \|T(x_1, x_2)\|$$

and

$$m = \inf\{C \geq 0 : \text{for each } (x_1, x_2) \in X_1 \times X_2, \|T(x_1, x_2)\| \leq C\|x_1\|\|x_2\|\}$$

Let  $(x_1, x_2) \in X_1 \times X_2$ . If  $\|x_1\| = 0$  or  $\|x_2\| = 0$ , then  $T(x_1, x_2) = 0$  and  $\|T(x_1, x_2)\| \leq M\|x_1\|\|x_2\|$ . Suppose that  $\|x_1\| \neq 0$  and  $\|x_2\| \neq 0$ . Then

$$\begin{aligned} \|T(x_1, x_2)\| &= \left( \|T(\|x_1\|^{-1}x_1, \|x_2\|^{-1}x_2)\| \right) \|x_1\|\|x_2\| \\ &\leq M\|x_1\|\|x_2\| \end{aligned}$$

Hence  $M \in \{C \geq 0 : \text{for each } x \in X, \|Tx\| \leq C\|x\|\}$  and  $m \leq M$ . Let  $C \in \{C \geq 0 : \text{for each } (x_1, x_2) \in X_1 \times X_2, \|T(x_1, x_2)\| \leq C\|x_1\|\|x_2\|\}$ . Suppose that  $\|x_1\| = 1$  and  $\|x_2\| = 1$ . Then  $\|T(x_1, x_2)\| \leq C\|x_1\|\|x_2\| = C$ . So  $M \leq C$ . Therefore  $M \leq m$ . So  $M = m$  and the supremum in (1) is the same as the infimum in (3). □

**Exercise 6.9.6.** Let  $X_1, X_2$  and  $Y$  be normed vector spaces. Then  $\|\cdot\| : L^2(X_1, X_2; Y) \rightarrow [0, \infty)$  is a norm.

*Proof.* □

**Exercise 6.9.7.** Let  $X_1, X_2, Y$  be normed vector spaces and  $T_1 \in L(X_1, L(X_2, Y))$ . Define  $T : X_1 \times X_2 \rightarrow Y$  by  $T(x_1, x_2) = T_1(x_1)(x_2)$ . Then  $T \in L^2(X_1, X_2; Y)$ .

*Proof.* It is straightforward to show that  $T$  is multilinear. For  $x_1 \in X_1$  and  $x_2 \in X_2$ ,

$$\begin{aligned} \|T(x_1, x_2)\| &= \|T_1(x_1)(x_2)\| \\ &\leq \|T_1(x_1)\| \|x_2\| \\ &\leq \|T_1\| \|x_1\| \|x_2\| \end{aligned}$$

So  $T \in L^2(X_1, X_2; Y)$ . □

**Exercise 6.9.8.** Let  $X_1, X_2, Y$  be normed vector spaces and  $T \in L^2(X_1, X_2; Y)$ . Define the map  $T_1 : X_1 \rightarrow Y^{X_2}$  by  $T_1(x_1)(\cdot) = T(x_1, \cdot)$ . Then  $T_1 \in L(X_1, L(X_2, Y))$ .

*Proof.* Let  $x_1 \in X_1$ . By definition of  $T$ ,  $T_1(x_1)$  is linear. Since  $T$  is bounded, there exists  $C \geq 0$  such that for each  $a_1 \in X_1$ ,  $a_2 \in X_2$ ,  $T(a_1, a_2) \leq C\|a_1\|\|a_2\|$ . Then for each  $x_2 \in X_2$ ,

$$\begin{aligned} \|T_1(x_1)(x_2)\| &= \|T(x_1, x_2)\| \\ &\leq (C\|x_1\|)\|x_2\| \end{aligned}$$

So  $T_1(x_1) \in L(X_2, Y)$  with  $\|T_1(x_1)\| \leq C\|x_1\|$ . Since  $x_1 \in X_1$  was arbitrary,  $T_1 : X_1 \rightarrow L(X_2, Y)$ . By definition of  $T$ ,  $T_1$  is linear. The preceding argument tells us that for each  $x_1 \in X_1$ ,

$$\|T_1(x_1)\| \leq C\|x_1\|$$

So  $T_1 \in L(X_1, L(X_2, Y))$  with  $\|T_1\| \leq C$ . □

**Exercise 6.9.9.** Let  $X_1, X_2$  be normed vector spaces. Define a map  $\phi : L^2(X_1, X_2; Y) \rightarrow L(X_1, L(X_2, Y))$  by  $\phi(T)(x_1)(x_2) = T(x_1, x_2)$ . Then  $T$  is an isometric isomorphism.

*Proof.* □

**Definition 6.9.10.** Let  $X_1, X_2$  be normed vector spaces,  $\phi_1 \in X_1^*$  and  $\phi_2 \in X_2^*$ . Define  $\phi_1 \otimes \phi_2 : X_1 \times X_2$  by  $\phi_1 \otimes \phi_2(x_1, x_2) = \phi_1(x_1)\phi_2(x_2)$ .

**Exercise 6.9.11.** Let  $X_1, X_2$  be normed vector spaces,  $\phi_1 \in X_1^*$  and  $\phi_2 \in X_2^*$ . Then  $\phi_1 \otimes \phi_2 \in L^2(X_1, X_2; \mathbb{C})$ .

*Proof.* Clear. □

**Exercise 6.9.12.** Let  $X_1, X_2$  be normed vector spaces and  $(x_1, x_2) \in X_1 \times X_2$ . If for each  $\phi_1 \in X_1^*$  and  $\phi_2 \in X_2^*$ ,  $\phi_1 \otimes \phi_2(x_1, x_2) = 0$ , then  $x_1 = 0$  or  $x_2 = 0$ .

*Proof.* Suppose that  $x_1 \neq 0$  and  $x_2 \neq 0$ . The previous section implies that there exist  $\phi_1 \in X_1^*$  and  $\phi_2 \in X_2^*$  such that  $\phi_1(x_1) = \|x_1\| \neq 0$  and  $\phi_2(x_2) = \|x_2\| \neq 0$ . Then

$$\begin{aligned} \phi_1 \otimes \phi_2(x_1, x_2) &= \phi_1(x_1)\phi_2(x_2) \\ &\neq 0 \end{aligned}$$

□

### 6.10. Banach Algebras.

**Definition 6.10.1.** Let  $X$  be a Banach space and an associative algebra. Then  $X$  is said to be a **Banach algebra** if for each  $S, T \in X$ ,  $\|ST\| \leq \|S\|\|T\|$ .

**Definition 6.10.2.** Let  $X$  be a Banach algebra and  $I \in X$ . Then  $I$  is said to be an **identity** if for each  $T \in X$ ,  $IT = TI = T$ .

**Definition 6.10.3.** Let  $X$  be a Banach algebra. and  $I \in X$ . Then  $I$  is said to be an **identity** if  $I \neq 0$  and for each  $T \in X$ ,  $IT = TI = T$ .

**Definition 6.10.4.** Let  $X$  be a Banach algebra. Then  $X$  is said to be **unital** if there exists  $I \in X$  such that  $I$  is an identity.

**Exercise 6.10.5.** Let  $X$  be a unital Banach algebra. Then there exists a unique  $I \in X$  such that  $I$  is an identity.

*Proof.* Clear. □

**Note 6.10.6.** We denote the unique identity element by  $I$ .

**Definition 6.10.7.** Let  $X$  be a unital Banach algebra and  $T, S \in X$ . Then  $S$  is said to be an **inverse** of  $T$  if  $TS = ST = I$ .

**Definition 6.10.8.** Let  $X$  be a unital Banach algebra and  $T \in X$ . Then  $T$  is said to be **invertible** if there exists  $S \in X$  such that  $S$  is an inverse of  $T$ .

**Exercise 6.10.9.** Let  $X$  be a unital Banach algebra and  $T \in X$ . If  $T$  is invertible, then there exists a unique  $S \in X$  such that  $S$  is an inverse of  $T$ .

*Proof.* Clear. □

**Note 6.10.10.** We denote the unique inverse of  $T$  by  $T^{-1}$ .

**Exercise 6.10.11. Fundamental Example:**

Let  $X$  be a Banach space. Then  $GL(X)$  is a unital Banach algebra.

*Proof.* Clear. □

**Definition 6.10.12.** Let  $X$  be a unital Banach algebra. We define  $GL(X) = \{T \in X : T \text{ is invertible}\}$ .

**Exercise 6.10.13.** Let  $X$  be a unital Banach algebra. Then  $GL(X)$  is a group.

*Proof.* Clear. □

**Exercise 6.10.14.** Let  $X$  be a unital Banach algebra. Then  $1 \leq \|I\|$ .

*Proof.* Since  $I \neq 0$ ,  $\|I\| \neq 0$ . By definition,

$$\|I\| = \|II\| \leq \|I\|\|I\|$$

Hence  $1 \leq \|I\|$ . □

**Exercise 6.10.15.** Let  $X$  be a Banach algebra. Then multiplication is continuous.

*Proof.* Let  $(S_1, T_1) \in X \times X$  and  $\epsilon > 0$ . Choose  $\delta = \min\{\frac{\epsilon}{2(\|S_1\| + \|T_1\| + 1)}, \frac{\sqrt{\epsilon}}{\sqrt{2}}\}$ . Let  $(S_2, T_2) \in X \times X$ . Suppose that

$$\|(S_1, T_1) - (S_2, T_2)\| = \max\{\|S_1 - S_2\|, \|T_1 - T_2\|\} < \delta$$

Then

$$\begin{aligned} \|S_1T_1 - S_2T_2\| &= \|S_1T_1 - S_2T_1 + S_2T_1 - S_2T_2\| \\ &\leq \|S_1 - S_2\|\|T_1\| + \|S_2\|\|T_1 - T_2\| \\ &\leq \|S_1 - S_2\|\|T_1\| + (\|S_1 - S_2\| + \|S_1\|)\|T_1 - T_2\| \\ &\leq \delta\|T_1\| + (\delta + \|S_1\|)\delta \\ &= \delta(\|S_1\| + \|T_1\|) + \delta^2 \\ &< \frac{\epsilon}{2} + \frac{\epsilon}{2} \\ &= \epsilon \end{aligned}$$

□

**Exercise 6.10.16.** Let  $X$  be a unital Banach algebra. Then

- (1) For each  $T \in X$ , if  $\|I - T\| < 1$ , then  $T \in GL(X)$  and

$$T^{-1} = \sum_{n=0}^{\infty} (I - T)^n$$

- (2) For each  $S, T \in X$ , if  $S \in GL(X)$  and  $\|S - T\| < \|S^{-1}\|^{-1}$ , then  $T \in GL(X)$ .  
 (3)  $GL(X)$  is open.

*Proof.*

- (1) Let  $T \in X$ . Suppose that  $\|I - T\| < 1$ . Then

$$\sum_{n=0}^{\infty} \|(I - T)^n\| \leq \sum_{n=0}^{\infty} \|I - T\|^n < \infty$$

Since  $X$  is a complete,  $\sum_{n=0}^{\infty} (I - T)^n$  converges in  $X$ .

Define  $(S_k)_{k=0}^{\infty} \subset X$  and  $S \in X$  by  $S_k = \sum_{n=0}^k (I - T)^n$  and

$S = \sum_{n=0}^{\infty} (I - T)^n$ . Then for each  $k \in \mathbb{N}$ ,

$$\begin{aligned} S_kT &= S_k - S_k(I - T) \\ &= (I - T)^0 - (I - T)^{k+1} \\ &= I - (I - T)^{k+1} \end{aligned}$$

and  $\|S_kT - I\| \leq \|I - T\|^{k+1}$ . Since multiplication on Banach algebras is continuous, we have that

$$ST = (\lim_{k \rightarrow \infty} S_k)T = \lim_{k \rightarrow \infty} S_kT = I$$

Similarly  $TS = I$ . Thus  $T \in GL(X)$  and  $T^{-1} = S \in X$ .

(2) Let  $S, T \in X$ . Suppose that  $S \in GL(X)$  and  $\|S - T\| < \|S^{-1}\|^{-1}$ . Then

$$\begin{aligned}\|I - S^{-1}T\| &= \|S^{-1}(S - T)\| \\ &\leq \|S^{-1}\|\|S - T\| \\ &< 1\end{aligned}$$

So  $S^{-1}T \in GL(X)$ . Thus  $T = S(S^{-1}T) \in GL(X)$ .

(3) Let  $T \in GL(X)$ . Choose  $\delta = \|T^{-1}\|^{-1}$ . By (2),  $B(T, \delta) \subset GL(X)$ .

□

## 7. HILBERT SPACES

## 7.1. Introduction.

**Definition 7.1.1.** Let  $H$  be a vector space and  $\langle \cdot, \cdot \rangle : H \rightarrow \mathbb{C}$ . Then  $\langle \cdot, \cdot \rangle$  is said to be an **inner product** on  $H$  if for each  $x, y, z \in H$  and  $c \in \mathbb{C}$

- (1)  $\langle x, y + cz \rangle = \langle x, y \rangle + c\langle x, z \rangle$
- (2)  $\langle x, y \rangle = \langle y, x \rangle^*$
- (3)  $\langle x, x \rangle \geq 0$
- (4) if  $\langle x, x \rangle = 0$ , then  $x = 0$ .

**Note 7.1.2.** In mathematics, inner products are conventionally defined to be linear in the first argument. However, in my opinion, the convention in physics of defining inner products to be linear in the second argument makes more sense.

**Exercise 7.1.3.** Let  $H$  be an inner product space,  $(x_j)_{j=1}^n, (y_j)_{j=1}^n \subset H$  and  $(\alpha_j)_{j=1}^n, (\beta_j)_{j=1}^n \subset \mathbb{C}$ . Then

$$\left\langle \sum_{i=1}^n \alpha_i x_i, \sum_{j=1}^n \beta_j y_j \right\rangle = \sum_{i=1}^n \sum_{j=1}^n \alpha_i^* \beta_j \langle x_i, y_j \rangle$$

*Proof.* Clear. □

**Definition 7.1.4.** Let  $H$  be an inner product space. Define the **induced norm**, denoted  $\| \cdot \| : H \rightarrow \mathbb{C}$ , by

$$\|x\| = \langle x, x \rangle^{\frac{1}{2}}$$

**Exercise 7.1.5. Cauchy-Schwarz Inequality**

Let  $H$  be an inner product space. Then for each  $x, y \in H$ ,  $|\langle x, y \rangle| \leq \|x\| \|y\|$  and  $|\langle x, y \rangle| = \|x\| \|y\|$  iff  $x \in \text{span}(y)$ .

**Hint:** For  $x, y \in H$ , put  $z = \text{sgn} \langle x, y \rangle^* y$  and Consider  $f : \mathbb{R} \rightarrow [0, \infty)$  defined by  $f(t) = \|x - tz\|^2$

*Proof.* Let  $x, y \in H$ . If  $y = 0$ , then the claim holds trivially. Suppose that  $y \neq 0$ . Put  $z = \text{sgn} \langle x, y \rangle^* y$ . So  $\langle x, z \rangle = |\langle x, y \rangle|$  and  $\|z\| = \|y\|$ . Define  $f : \mathbb{R} \rightarrow [0, \infty)$  by

$$f(t) = \|x - tz\|^2$$

. Then for each  $t \in \mathbb{R}$ ,

$$\begin{aligned} 0 &\leq f(t) \\ &= \|x - tz\|^2 \\ &= \|x\|^2 + |t|^2 \|z\|^2 - 2 \text{Re}(t \langle x, z \rangle) \\ &= \|x\|^2 + t^2 \|y\|^2 - 2t |\langle x, y \rangle| \end{aligned}$$



Thus  $f$  is a quadratic with a minimum at  $t_0 = \frac{|\langle x, y \rangle|}{\|y\|^2}$ . Hence

$$\begin{aligned} 0 &\leq f(t_0) \\ &= \|x\|^2 + \frac{|\langle x, y \rangle|}{\|y\|^2} - 2 \frac{|\langle x, y \rangle|}{\|y\|^2} \\ &= \|x\|^2 - \frac{|\langle x, y \rangle|}{\|y\|^2} \end{aligned}$$

Which implies that

$$|\langle x, y \rangle|^2 \leq \|x\|^2 \|y\|^2$$

and hence the claim holds. Clearly if  $x \in \text{span}(y)$ , then equality holds. Conversely, if equality holds, then  $x - z = 0$  which implies that  $x \in \text{span}(y)$ .  $\square$

**Exercise 7.1.6.** Let  $H$  be an inner product space. Then the induced norm,  $\|\cdot\| : H \rightarrow \mathbb{C}$ , is a norm.

*Proof.* Let  $x, y \in H$  and  $c \in \mathbb{C}$ . Then

- (1) By definition, if  $\|x\| = 0$ , then  $\langle x, x \rangle = 0$ , which implies that  $x = 0$ .
- (2) Note that

$$\begin{aligned} \|cx\|^2 &= \langle cx, cx \rangle \\ &= c * c \langle x, x \rangle \\ &= |c|^2 \|x\|^2 \end{aligned}$$

So  $\|cx\| = |c| \|x\|$

- (3) The Cauchy-Schwarz inequality implies that

$$\begin{aligned} \|x + y\|^2 &= \|x\|^2 + \|y\|^2 + 2 \text{Re}(\langle x, y \rangle) \\ &\leq \|x\|^2 + \|y\|^2 + 2|\langle x, y \rangle| \\ &\leq \|x\|^2 + \|y\|^2 + 2\|x\| \|y\| \\ &= (\|x\| + \|y\|)^2 \end{aligned}$$

Hence  $\|x + y\| \leq \|x\| + \|y\|$ .

$\square$

**Definition 7.1.7.** Let  $H$  be an inner product space,  $x, y \in H$  and  $S \subset H$ . Then

- (1)  $x$  and  $y$  are said to be **orthogonal** if  $\langle x, y \rangle = 0$ .
- (2)  $S$  is said to be **orthogonal** if for each  $x, y \in S$ ,  $x, y$  are orthogonal.

**Exercise 7.1.8. (Pythagorean theorem):**

Let  $H$  be an inner product space and  $(x_j)_{j=1}^n \subset H$  an orthogonal set. Then

$$\left\| \sum_{j=1}^n x_j \right\|^2 = \sum_{j=1}^n \|x_j\|^2$$

*Proof.* We have that

$$\begin{aligned}
 \left\| \sum_{j=1}^n x_j \right\|^2 &= \left\langle \sum_{i=1}^n x_i, \sum_{j=1}^n x_j \right\rangle \\
 &= \sum_{i=1}^n \sum_{j=1}^n \langle x_i, x_j \rangle \\
 &= \sum_{j=1}^n \langle x_j, x_j \rangle \\
 &= \sum_{j=1}^n \|x_j\|^2
 \end{aligned}$$

□

**Exercise 7.1.9.** Let  $H$  be an inner product space and  $S \subset H$ . Suppose that  $0 \notin S$ . If  $S$  is orthogonal, then  $S$  is linearly independent.

*Proof.* Let  $x_1, \dots, x_n \in S$  and  $c_1, \dots, c_n \in \mathbb{C}$ . Suppose that  $\sum_{j=1}^n c_j x_j = 0$ . Since  $(c_j x_j)_{j=1}^n$  is orthogonal, the Pythagorean theorem implies that

$$\begin{aligned}
 0 &= \left\| \sum_{i=1}^n c_i x_i \right\|^2 \\
 &= \sum_{j=1}^n |c_j|^2 \|x_j\|^2
 \end{aligned}$$

So for each  $j \in \{1, \dots, n\}$ ,  $c_j = 0$  and  $S$  is linearly independent. □

**Definition 7.1.10.** Let  $H$  be an inner product space and  $S \subset H$ . Then  $S$  is said to be **orthonormal** if  $S$  is orthogonal and for each  $x \in S$ ,  $\|x\| = 1$ .

**Exercise 7.1.11. Bessel's Inequality:**

Let  $H$  be an inner product space and  $S \subset H$ . If  $S$  is orthonormal, then for each  $x \in H$ ,

$$\sum_{u \in S} |\langle u, x \rangle|^2 \leq \|x\|^2$$

and in particular,  $\{u \in S : \langle u, x \rangle \neq 0\}$  is countable.

*Proof.* Suppose that  $S$  is orthonormal. Let  $x \in H$  and  $F \subset S$  finite. Then the Pythagorean theorem implies that

$$\begin{aligned}
 0 &\leq \left\| x - \sum_{u \in F} \langle u, x \rangle u \right\|^2 \\
 &= \|x\|^2 + \left\| \sum_{u \in F} \langle u, x \rangle u \right\|^2 - 2 \operatorname{Re} \left\langle x, \sum_{u \in F} \langle u, x \rangle u \right\rangle \\
 &= \|x\|^2 + \sum_{u \in F} |\langle u, x \rangle|^2 \|u\|^2 - 2 \sum_{u \in F} |\langle u, x \rangle|^2 \\
 &= \|x\|^2 - \sum_{u \in F} |\langle u, x \rangle|^2
 \end{aligned}$$

So

$$\sum_{u \in F} |\langle u, x \rangle|^2 \leq \|x\|^2$$

By definition of the sum,

$$\sum_{u \in S} |\langle u, x \rangle|^2 \leq \|x\|^2$$

Basic integration theory then tells us that  $\{u \in S : \langle u, x \rangle \neq 0\}$  is countable.  $\square$

**Definition 7.1.12.** Let  $H$  be an inner product space. Then  $H$  is said to be a **Hilbert space** if  $H$  is complete with respect to the induced norm on  $H$ .

**Exercise 7.1.13.** Let  $H$  be a Hilbert space and  $S \subset H$ . Suppose that  $S$  is orthonormal. Then the following are equivalent:

- (1) For each  $x \in H$ , if for each  $u \in S$ ,  $\langle u, x \rangle = 0$ , then  $x = 0$ .
- (2) For each  $x \in H$ , there exist  $(u_j)_{j \in \mathbb{N}} \subset S$  such that  $x = \sum_{j \in \mathbb{N}} \langle u_j, x \rangle u_j$  and for each  $u \notin (u_j)_{j \in \mathbb{N}}$ ,  $\langle u, x \rangle = 0$ .
- (3) For each  $x \in H$ ,  $\|x\|^2 = \sum_{u \in S} |\langle u, x \rangle|^2$ .

*Proof.*

- (1)  $\implies$  (2):

Suppose that for each  $x \in H$ , if for each  $u \in S$ ,  $\langle u, x \rangle = 0$ , then  $x = 0$ . Let  $x \in H$ . Put  $S_* = \{u \in S : \langle u, x \rangle \neq 0\}$ . The previous exercise implies that  $S_*$  is countable. Write  $S_* = (u_j)_{j=1}^\infty$ . The previous exercise tells us that  $\sum_{j \in \mathbb{N}} |\langle u_j, x \rangle|^2 \leq \|x\|^2$  and hence converges. Thus for  $\epsilon > 0$ , there exist  $N \in \mathbb{N}$  such that for each  $m, n \in \mathbb{N}$ ,  $m, n \geq N$  implies that if  $m < n$ , then

$$\sum_{j=m+1}^n |\langle u_j, x \rangle|^2 < \epsilon$$

Define  $(y_n)_{n \in \mathbb{N}} \subset H$  by

$$y_n = \sum_{j=1}^n \langle u_j, x \rangle u_j$$

Then for each  $m, n \in \mathbb{N}$ ,  $m, n \geq N$  implies that if  $m < n$ , then

$$\begin{aligned} \|y_n - y_m\|^2 &= \left\| \sum_{j=1}^n \langle u_j, x \rangle u_j - \sum_{j=1}^m \langle u_j, x \rangle u_j \right\|^2 \\ &= \left\| \sum_{j=m+1}^n \langle u_j, x \rangle u_j \right\|^2 \\ &= \sum_{j=m+1}^n |\langle u_j, x \rangle|^2 \\ &< \epsilon \end{aligned}$$

So  $(y_n)_{n \in \mathbb{N}}$  is Cauchy. Since  $H$  is complete, there exists  $y \in H$  such that  $y_n \rightarrow y$ . By definition,

$$y = \sum_{j \in \mathbb{N}} \langle u_j, x \rangle u_j$$

Continuity of  $\langle \cdot, \cdot \rangle : H \times H \rightarrow \mathbb{C}$  implies that

(1) for each  $u \in S \setminus S_*$ ,

$$\begin{aligned} \langle u, x - y \rangle &= \langle u, x \rangle - \langle u, y \rangle \\ &= \langle u, x \rangle - \lim_{n \rightarrow \infty} \langle u, y_n \rangle \\ &= \langle u, x \rangle - \lim_{n \rightarrow \infty} \sum_{j=1}^n \langle u_j, x \rangle \langle u, u_j \rangle \\ &= 0 - 0 \\ &= 0 \end{aligned}$$

(2) for each  $k \in \mathbb{N}$ ,

$$\begin{aligned} \langle u_k, x - y \rangle &= \langle u_k, x \rangle - \langle u_k, y \rangle \\ &= \langle u_k, x \rangle - \lim_{n \rightarrow \infty} \langle u_k, y_n \rangle \\ &= \langle u_k, x \rangle - \lim_{n \rightarrow \infty} \sum_{j=1}^n \langle u_j, x \rangle \langle u_k, u_j \rangle \\ &= \langle u_k, x \rangle - \langle u_k, x \rangle \\ &= 0 \end{aligned}$$

So for each  $u \in S$ ,  $\langle u, x - y \rangle = 0$ . By assumption,  $x - y = 0$  and hence

$$x = \sum_{j \in \mathbb{N}} \langle u_j, x \rangle u_j$$

• (2)  $\implies$  (3):

Suppose that for each  $x \in H$ , there exist  $(u_j)_{j \in \mathbb{N}} \subset S$  such that  $x = \sum_{j \in \mathbb{N}} \langle u_j, x \rangle u_j$  and

for each  $u \notin (u_j)_{j \in \mathbb{N}}$ ,  $\langle u, x \rangle = 0$ . Then continuity of  $\|\cdot\| : H \rightarrow [0, \infty)$  implies that

$$\begin{aligned} \|x\|^2 &= \left\| \lim_{n \rightarrow \infty} \sum_{j=1}^n \langle u_j, x \rangle u_j \right\|^2 \\ &= \lim_{n \rightarrow \infty} \left\| \sum_{j=1}^n \langle u_j, x \rangle u_j \right\|^2 \\ &= \lim_{n \rightarrow \infty} \sum_{j=1}^n |\langle u_j, x \rangle|^2 \\ &= \sum_{j \in \mathbb{N}} |\langle u_j, x \rangle|^2 \\ &= \sum_{u \in S} |\langle u, x \rangle|^2 \end{aligned}$$

• (3)  $\implies$  (4):

Suppose that for each  $x \in H$ ,  $\|x\|^2 = \sum_{u \in S} |\langle u, x \rangle|^2$ . Let  $x \in H$ . Suppose that for each  $u \in S$ ,  $\langle u, x \rangle = 0$ . Then

$$\begin{aligned} \|x\|^2 &= \sum_{u \in S} |\langle u, x \rangle|^2 \\ &= 0 \end{aligned}$$

So  $x = 0$

□

**Definition 7.1.14.** Let  $H$  be a Hilbert space and  $S \subset H$ . Then  $S$  is said to be an **orthonormal basis of  $H$**  if

- (1)  $S$  is orthonormal
- (2) for each  $x \in H$ , if for each  $u \in S$ ,  $\langle u, x \rangle = 0$ , then  $x = 0$

## 7.2. Operators and Functionals.

### Definition 7.2.1. (Adjoint of an Operator):

Let  $H$  be a Hilbert space and  $A, B \in L(H)$ . Then  $B$  is said to be the **adjoint** of  $A$  if for each  $x_1, x_2 \in H$ ,

$$\langle x_1, Ax_2 \rangle = \langle Bx_1, x_2 \rangle$$

In this case, we write

$$B = A^*$$

**Note 7.2.2.** In physics, the adjoint of  $A$  is typically denoted by  $A^\dagger$ .

**Exercise 7.2.3.** Let  $H$  be a Hilbert space,  $A, B \in L(H)$  and  $\lambda \in \mathbb{C}$ , then

- (1)  $(A^*)^* = A$
- (2)  $(A + B)^* = A^* + B^*$
- (3)  $(AB)^* = B^*A^*$
- (4)  $(\lambda A)^* = \lambda^*A^*$
- (5)  $A$  and  $B$  commute iff  $A^*$  and  $B^*$  commute.

*Proof.* Let  $x_1, x_2 \in H$ . Then

(1)

$$\begin{aligned} \langle Ax_1, x_2 \rangle &= \langle x_2, Ax_1 \rangle^* \\ &= \langle A^*x_2, x_1 \rangle^* \quad (\text{by definition}) \\ &= \langle x_1, A^*x_2 \rangle \end{aligned}$$

(2)

$$\begin{aligned} \langle x_1, (A + B)x_2 \rangle &= \langle x_1, Ax_2 \rangle + \langle x_1, Bx_2 \rangle \\ &= \langle A^*x_1, x_2 \rangle + \langle B^*x_1, x_2 \rangle \\ &= \langle (A^* + B^*)x_1, x_2 \rangle \end{aligned}$$

(3)

$$\begin{aligned} \langle x_1, ABx_2 \rangle &= \langle A^*x_1, Bx_2 \rangle \\ &= \langle B^*A^*x_1, x_2 \rangle \end{aligned}$$

(4)

$$\begin{aligned} \langle x_1, \lambda Ax_2 \rangle &= \lambda \langle x_1, Ax_2 \rangle \\ &= \lambda \langle A^*x_1, x_2 \rangle \\ &= \langle \lambda^*A^*x_1, x_2 \rangle \end{aligned}$$

(5) If  $A$  and  $B$  commute, then

$$\begin{aligned} A^*B^* &= (BA)^* \\ &= (AB)^* \\ &= B^*A^* \end{aligned}$$

Conversely, if  $A^*$  and  $B^*$  commute then

$$\begin{aligned} AB &= (B^*A^*)^* \\ &= (A^*B^*)^* \\ &= BA \end{aligned}$$

□

**Definition 7.2.4.** Let  $H$  be a Hilbert space and  $Q \in L(H)$ . Then  $Q$  is said to be **self-adjoint** if

$$Q = Q^*$$

**Exercise 7.2.5.** Let  $H$  be a Hilbert space and  $Q \in L(H)$ . If  $Q$  is a self-adjoint then

- (1) the eigenvalues of  $Q$  are real.
- (2) the eigenvectors of  $Q$  corresponding to distinct eigenvalues are orthogonal.

*Proof.* Suppose that  $Q$  is self-adjoint.

- (1) Let  $\lambda$  be an eigenvalue of  $Q$  with corresponding eigenvector  $x$ . Then

$$\begin{aligned} \lambda \langle x, x \rangle &= \langle x, Qx \rangle \\ &= \langle Qx, x \rangle \\ &= \lambda^* \langle x, x \rangle \end{aligned}$$

Thus  $\lambda = \lambda^*$  and is real

- (2) Let  $\lambda_1$  and  $\lambda_2$  be eigenvalues of  $Q$  with corresponding eigenvectors  $x_1$  and  $x_2$ . Suppose that  $\lambda_1 \neq \lambda_2$ . Then

$$\begin{aligned} \lambda_2 \langle x_1, x_2 \rangle &= \langle x_1, Qx_2 \rangle \\ &= \langle Qx_1, x_2 \rangle \\ &= \lambda_1 \langle x_1, x_2 \rangle \end{aligned}$$

So  $(\lambda_2 - \lambda_1) \langle x_1, x_2 \rangle = 0$ . Which implies that  $\langle x_1, x_2 \rangle = 0$

□

**Exercise 7.2.6.** Let  $H$  be a Hilbert space,  $A, B \in L(H)$  and  $\lambda \in \mathbb{R}$ . Suppose that  $A, B$  are self-adjoint. If  $A$  and  $B$  commute and then  $\lambda AB$  is self-adjoint.

*Proof.*

$$\begin{aligned} (\lambda AB)^* &= \lambda^* (AB)^* \\ &= \lambda B^* A^* \\ &= \lambda BA \\ &= \lambda AB \end{aligned}$$

□

**Definition 7.2.7. (Adjoint of a Vector):**

Let  $H$  be a Hilbert space and  $x \in H$ . We define the **adjoint** of  $x$ , denoted  $x^* \in H^*$ , by  $x^*y = \langle x, y \rangle$ .

**Note 7.2.8.** In mathematics, where linearity of the inner product is in the first argument,  $x^*$  is typically referred to by  $u_x \in H^*$  where  $u_x(y) = \langle y, x \rangle$ . In physics, where the inner product with linearity in the second argument,  $x^*\phi$  is usually written in the so-called “bra-ket” notation as  $\langle x|\phi \rangle$  which works smoothly since it aligns with the linearity of  $u_x(\phi_1 + \lambda\phi_2)$  and the conjugate-linearity of  $u_{x_1 + \lambda x_2}(\phi)$ . In this way, it generalizes the notation for  $\langle x, y \rangle = x^T y$  for  $\mathbb{R}^n$  to  $\langle x, y \rangle = x^* y$  for  $\mathbb{C}^n$ .

**Exercise 7.2.9.** Let  $H$  be a Hilbert space,  $x, y \in H$  and  $\lambda \in \mathbb{C}$ . Then

- (1)  $(x + y)^* = x^* + y^*$
- (2)  $(\lambda x)^* = \lambda^* x^*$

*Proof.* Clear. □

**Definition 7.2.10.** Let  $H$  be a Hilbert space,  $x, y \in H$  and  $A \in L(H)$ . We define

- (1)  $x^* A \in H^*$  by  $(x^* A)y = x^*(Ay)$
- (2)  $xy^* \in L(H)$  by  $(xy^*)z = (y^*z)x$

**Exercise 7.2.11.** Let  $H$  be a Hilbert space,  $A \in L(H)$  and  $x \in H$ . Then

$$(Ax)^* = x^* A^*$$

*Proof.* Let  $y \in H$ . Then

$$\begin{aligned} (Ax)^* y &= \langle Ax, y \rangle \\ &= \langle x, A^* y \rangle \\ &= x^* A^* y \end{aligned}$$

□

**Definition 7.2.12. (Commutator):**

Let  $H$  be a Hilbert space and  $A, B \in L(H)$ . The **commutator** of  $A$  and  $B$ , denoted  $[A, B]$ , is defined by

$$[A, B] = AB - BA$$

**Exercise 7.2.13.** Let  $H$  be a Hilbert space and  $A, B, C \in L(H)$ . Then

- (1)  $[AB, C] = A[B, C] + [A, C]B$
- (2)  $[A, BC] = B[A, C] + [A, B]C$

*Proof.*

(1)

$$\begin{aligned} [AB, C] &= ABC - CAB \\ &= ABC - ACB + ACB - CAB \\ &= A(BC - CB) + (AC - CA)B \\ &= A[B, C] + [A, C]B \end{aligned}$$

(2) Similar to (1).

□



### 7.3. Tensor Products.

**Note 7.3.1.** This section assumes familiarity with the algebraic tensor product of two vector spaces. See section ??? of [1] for details.

**Definition 7.3.2.** Let  $X, Y$  and  $Z$  be Banach spaces and  $\phi \in L^2(X, Y; Z)$ . Then  $(Z, \phi)$  is said to be a **tensor product** of  $X$  with  $Y$  if

- (1)  $\text{span } \phi(X \times Y)$  is dense in  $Z$
- (2) for each Banach space  $W$  and  $\psi \in L^2(X, Y; W)$ , there exists a unique  $\psi' \in L(Z, W)$  such that  $\psi' \circ \phi = \psi$ , i.e. such that the following diagram commutes:

$$\begin{array}{ccc} X \times Y & \xrightarrow{\phi} & Z \\ & \searrow \psi & \downarrow \psi' \\ & & W \end{array}$$

If  $(Z, \phi)$  is a tensor product of  $X$  with  $Y$ . We often write  $Z = X \otimes Y$  and for each  $x \in X$ ,  $y \in Y$ , we often write  $\phi(x, y) = x \otimes y$ .

**Exercise 7.3.3.** Let  $X$  and  $Y$  be Banach spaces,  $U \subset X$  and  $V \subset Y$ . Set  $W = \{u \otimes v : u \in U \text{ and } v \in V\} \subset X \otimes Y$ . If  $U$  and  $V$  are linearly independent, then  $W$  is linearly independent.

**Hint:** For  $\phi \in X^*$ ,  $\psi \in Y^*$ , define  $T \in L^2(X, Y; \mathbb{C})$  by  $T(x, y) = \phi(x)\psi(y)$ .

*Proof.* Let  $w = \sum_{u \in U} \sum_{v \in V} \lambda_{u,v} u \otimes v$ . Suppose that  $w = 0$ . Let  $\phi \in X^*$  and  $\psi \in Y^*$ . Define  $T \in L^2(X, Y; \mathbb{C})$  by  $T(x, y) = \phi(x)\psi(y)$ . By definition of the tensor product, there exists a unique  $T' \in L(X \otimes Y, \mathbb{C})$  such that for each  $x \in X$  and  $y \in Y$ ,  $T'(x \otimes y) = T(x, y)$ . Then

$$\begin{aligned} 0 &= T'(w) \\ &= T'\left(\sum_{u \in U} \sum_{v \in V} \lambda_{u,v} u \otimes v\right) \\ &= \sum_{u \in U} \sum_{v \in V} \lambda_{u,v} T'(u \otimes v) \\ &= \sum_{u \in U} \sum_{v \in V} \lambda_{u,v} T(u, v) \\ &= \sum_{u \in U} \sum_{v \in V} \lambda_{u,v} \phi(u) \psi(v) \\ &= \phi\left(\sum_{u \in U} \sum_{v \in V} \lambda_{u,v} \psi(v) u\right) \end{aligned}$$

Since  $\phi \in X^*$  is arbitrary, a previous exercise in the section on linear functionals implies that

$$\begin{aligned} 0 &= \sum_{u \in U} \sum_{v \in V} \lambda_{u,v} \psi(v) u \\ &= \sum_{u \in U} \left( \sum_{v \in V} \lambda_{u,v} \psi(v) \right) u \end{aligned}$$

Linear independence of  $U$  implies that for each  $u \in U$ ,

$$\begin{aligned} 0 &= \sum_{v \in V} \lambda_{u,v} \psi(v) \\ &= \psi \left( \sum_{v \in V} \lambda_{u,v} v \right) \end{aligned}$$

Since  $\psi \in Y^*$  is arbitrary, for each  $u \in U$ ,

$$\sum_{v \in V} \lambda_{u,v} v = 0$$

Linear independence of  $V$  implies that for each  $u \in U, v \in V, \lambda_{u,v} = 0$ . Hence  $W$  is linearly independent.  $\square$

**Exercise 7.3.4. Uniqueness:**

Let  $X, Y$  and  $Z$  be Banach spaces and  $\phi \in L^2(X, Y; Z)$ . Suppose that  $(Z, \phi)$  is a tensor product of  $X$  with  $Y$ . Then  $(Z, \phi)$  is unique up to isomorphism.

*Proof.* Let  $W$  be a Banach space and  $\psi \in L^2(X, Y; W)$ . Suppose that  $(W, \psi)$  is a tensor product of  $X$  with  $Y$ . Since  $(Z, \phi)$  is a tensor product of  $X$  with  $Y$ , there exists a unique  $\psi' \in L(Z, W)$  such that  $\psi' \circ \phi = \psi$ . Since  $(W, \psi)$  is a tensor product of  $X$  with  $Y$ , there exists a unique  $\phi' \in L(W, Z)$  such that  $\phi' \circ \psi = \phi$ . Thus the following diagram commutes:

$$\begin{array}{ccc} & & W \\ & \nearrow \psi & \downarrow \phi' \\ X \times Y & \xrightarrow{\phi} & Z \\ & \searrow \psi & \downarrow \psi' \\ & & W \end{array}$$

On the other hand, since  $(W, \psi)$  is a tensor product of  $X$  with  $Y$ , there exists a unique  $\Psi \in L(W)$  such that  $\Psi \circ \psi = \psi$ . Thus the following diagram commutes:

$$\begin{array}{ccc} X \times Y & \xrightarrow{\psi} & W \\ & \searrow \psi & \downarrow \Psi \\ & & W \end{array}$$

Since  $I_W \in L(W)$  and  $I_W \circ \psi = \psi$ , uniqueness of  $\Psi$  implies that  $\Psi = I_W$ . From the first diagram, we see that  $\psi' \circ \phi'$  satisfies  $(\psi' \circ \phi') \circ \psi = \psi$ . Since  $\psi' \circ \phi' \in L(W)$ , uniqueness of  $\Psi$  implies that  $\Psi = \psi' \circ \phi'$ . Thus  $\psi' \circ \phi' = I_W$ .

Similarly, we could have initially considered the following diagram:

$$\begin{array}{ccc} & & Z \\ & \nearrow \phi & \downarrow \psi' \\ X \times Y & \xrightarrow{\psi} & W \\ & \searrow \phi & \downarrow \phi' \\ & & Z \end{array}$$

Playing a similar game, we could use the fact that there exists a unique  $\Phi \in L(Z)$  such that  $\Phi \circ \phi = \phi$  to obtain the following diagram:

$$\begin{array}{ccc} X \times Y & \xrightarrow{\phi} & Z \\ & \searrow \phi & \downarrow \Phi \\ & & Z \end{array}$$

As before, uniqueness enables us to conclude that  $\phi' \circ \psi' = I_Z$ . Thus  $\psi'$  and  $\phi'$  are isomorphisms and  $Z \cong W$ .  $\square$

**Note 7.3.5.** The following definitions and exercises will cover the explicit construction of a tensor product of Banach spaces.

**Definition 7.3.6.** Let  $X$  and  $Y$  be Banach spaces. Define  $X \otimes^{\text{alg}} Y = \text{span}\{x \otimes y : x \in X \text{ and } y \in Y\}$  to be the algebraic tensor product of  $X$  with  $Y$  (see section ??? of [1] for details).

**Exercise 7.3.7.** Let  $X$  and  $Y$  be Banach spaces and  $x \otimes y \in X \otimes^{\text{alg}} Y$ . If for each  $\phi \in X^*$  and  $\psi \in Y^*$ ,  $\phi \otimes \psi(x, y) = 0$ , then  $x \otimes y = 0$ .

*Proof.* The previous section tells us that for each  $\phi \in X^*$  and  $\psi \in Y^*$ ,  $\phi \otimes \psi(x, y) = 0$ , then  $x = 0$  or  $y = 0$ . This implies that  $x \otimes y = 0$ .  $\square$

**Definition 7.3.8. The Projective Norm:**

Define  $\|\cdot\|_\pi : X \otimes^{\text{alg}} Y \rightarrow [0, \infty)$  by

$$\|u\|_\pi = \inf \left\{ \sum_{j=1}^n \|x_j\| \|y_j\| : (x_j)_{j=1}^n \subset X, (y_j)_{j=1}^n \subset Y \text{ and } u = \sum_{j=1}^n x_j \otimes y_j \right\}$$

**Exercise 7.3.9.** Let  $X$  and  $Y$  be Banach spaces. Then  $\|\cdot\|_\pi : X \otimes^{\text{alg}} Y \rightarrow [0, \infty)$  is a norm on  $X \otimes^{\text{alg}} Y$ .

*Proof.*

- Let  $\lambda \in \mathbb{C}$ ,  $u \in X \otimes^{\text{alg}} Y$ . If  $\lambda = 0$ , then  $\lambda u = 0u = 0 \otimes 0$  and clearly  $\|\lambda u\|_\pi = 0 = |\lambda| \|u\|_\pi$ . Suppose that  $\lambda \neq 0$ . Let  $\epsilon > 0$ . Then there exist  $(x_j)_{j=1}^n \subset X$  and  $(y_j)_{j=1}^n \subset Y$  such that  $u = \sum_{j=1}^n x_j \otimes y_j$  and  $\sum_{j=1}^n \|x_j\| \|y_j\| < \|u\|_\pi + \epsilon/|\lambda|$ . Then  $\lambda u = \sum_{j=1}^n (\lambda x_j) \otimes y_j$ . Therefore

$$\begin{aligned} \|\lambda u\|_\pi &\leq \sum_{j=1}^n \|\lambda x_j\| \|y_j\| \\ &\leq |\lambda| \sum_{j=1}^n \|x_j\| \|y_j\| \\ &< |\lambda| \left( \|u\|_\pi + \frac{\epsilon}{|\lambda|} \right) \\ &= |\lambda| \|u\|_\pi + \epsilon \end{aligned}$$

Since  $\epsilon > 0$  is arbitrary,  $\|\lambda u\|_\pi \leq |\lambda|\|u\|_\pi$ . For the sake of contradiction, suppose that  $\|\lambda u\|_\pi < |\lambda|\|u\|_\pi$ . Then there exists  $(x_j)_{j=1}^n \subset X$  and  $(y_j)_{j=1}^n \subset Y$  such that  $\lambda u = \sum_{j=1}^n x_j \otimes y_j$  and  $\sum_{j=1}^n \|x_j\|\|y_j\| < |\lambda|\|u\|_\pi$ . Hence  $u = \sum_{j=1}^n (\lambda^{-1}x_j) \otimes y_j$ . This implies that

$$\begin{aligned} \|u\|_\pi &\leq \sum_{j=1}^n \|\lambda^{-1}x_j\|\|y_j\| \\ &= |\lambda|^{-1} \sum_{j=1}^n \|x_j\|\|y_j\| \\ &< |\lambda|^{-1}|\lambda|\|u\|_\pi \\ &= \|u\|_\pi \end{aligned}$$

which is a contradiction. Therefore  $\|\lambda u\|_\pi \geq |\lambda|\|u\|_\pi$  which implies that  $\|\lambda u\|_\pi = |\lambda|\|u\|_\pi$

- Let  $u, v \in X \otimes^{\text{alg}} Y$  and  $\epsilon > 0$ . Then there exist  $(x_j)_{j=1}^n, (a_k)_{k=1}^m \subset X$  and  $(y_j)_{j=1}^n, (b_k)_{k=1}^m \subset Y$  such that  $u = \sum_{j=1}^n x_j \otimes y_j$ ,  $v = \sum_{k=1}^m a_k \otimes b_k$ ,  $\sum_{j=1}^n \|x_j\|\|y_j\| < \|u\|_\pi + \epsilon/2$  and  $\sum_{k=1}^m \|a_k\|\|b_k\| < \|v\|_\pi + \epsilon/2$ . Then  $u + v = \sum_{j=1}^n x_j \otimes y_j + \sum_{k=1}^m a_k \otimes b_k$  which implies that

$$\begin{aligned} \|u + v\|_\pi &\leq \sum_{j=1}^n \|x_j\|\|y_j\| + \sum_{k=1}^m \|a_k\|\|b_k\| \\ &< \|u\|_\pi + \epsilon/2 + \|v\|_\pi + \epsilon/2 \\ &= \|u\|_\pi + \|v\|_\pi + \epsilon \end{aligned}$$

Since  $\epsilon > 0$  is arbitrary,  $\|u + v\|_\pi \leq \|u\|_\pi + \|v\|_\pi$ .

- Let  $u \in X \otimes^{\text{alg}} Y$ . Suppose that  $\|u\| = 0$ . Let  $\phi \in X^*$  and  $\psi \in Y^*$  and  $\epsilon > 0$ . Then there exist  $(x_j)_{j=1}^n \subset X$  and  $(y_j)_{j=1}^n \subset Y$  such that  $u = \sum_{j=1}^n x_j \otimes y_j$  and

$$\sum_{j=1}^n \|x_j\|\|y_j\| < \frac{\epsilon}{\|\phi\|\|\psi\| + 1}$$

Then

$$\begin{aligned}
 \sum_{j=1}^n |\phi \otimes \psi(x_j, y_j)| &= \sum_{j=1}^n |\phi(x_j)\psi(y_j)| \\
 &\leq \sum_{j=1}^n \|\phi\| \|x_j\| \|\psi\| \|y_j\| \\
 &= \|\phi\| \|\psi\| \sum_{j=1}^n \|x_j\| \|y_j\| \\
 &< \|\phi\| \|\psi\| \frac{\epsilon}{\|\phi\| \|\psi\| + 1} \\
 &< \epsilon
 \end{aligned}$$

Then for each  $j \in \{1, \dots, n\}$ ,  $|\phi \otimes \psi(x_j, y_j)| < \epsilon$ . **FINISH!!!** Try using sequences and continuity and a common refinement of representation and averaging

□

**Exercise 7.3.10. Existence:**

*Proof.*

□

## 8. DIFFERENTIATION

## 8.1. The Gateaux Derivative.

**Note 8.1.1.** In this section, we assume all Banach spaces to be over  $\mathbb{R}$ .

**Definition 8.1.2.** Let  $X, Y$  be Banach spaces,  $A \subset X$  open,  $f : A \rightarrow Y$ ,  $x_0 \in A$  and  $x \in X$ . Then  $f$  is said to be

- (1) **right-hand-differentiable at  $x_0$  in the direction  $x$**  if the limit

$$\lim_{t \rightarrow 0^+} \frac{f(x_0 + tx) - f(x_0)}{t}$$

exists. If  $f$  is right-hand-differentiable at  $x_0$  in the direction  $x$ , we define the **right-hand derivative** of  $f$  at  $x_0$  in the direction  $x$ , denoted by  $d^+f(x_0; x)$ , to be the above limit.

- (2) **left-hand-differentiable at  $x_0$  in the direction  $x$**  if the limit

$$\lim_{t \rightarrow 0^-} \frac{f(x_0 + tx) - f(x_0)}{t}$$

exists. If  $f$  is left-hand-differentiable at  $x_0$  in the direction  $x$ , we define the **left-hand derivative** of  $f$  at  $x_0$  in the direction  $x$ , denoted by  $d^-f(x_0; x)$ , to be the above limit.

- (3) **differentiable at  $x_0$  in the direction  $x$**  if the limit

$$\lim_{t \rightarrow 0} \frac{f(x_0 + tx) - f(x_0)}{t}$$

exists. If  $f$  is differentiable at  $x_0$  in the direction  $x$ , we define the **derivative** of  $f$  at  $x_0$  in the direction  $x$ , denoted by  $df(x_0; x)$ , to be the above limit.

**Exercise 8.1.3.** Let  $X, Y$  be Banach spaces,  $A \subset X$  open,  $f : A \rightarrow \mathbb{R}$  and  $x_0 \in A$ . Then  $df(x_0; 0) = 0$ .

*Proof.* Clear. □

**Definition 8.1.4. The Gateaux Derivative:**

Let  $X, Y$  be Banach spaces,  $A \subset X$  open,  $f : A \rightarrow Y$  and  $x_0 \in A$ . Then  $f$  is said to be

- (1) **right-hand Gateaux differentiable at  $x_0$**  if for each  $x \in X$ ,  $d^+f(x_0; x)$  exists. We define the **right-hand Gateaux derivative** of  $f$  at  $x_0$ , denoted  $d^+f(x_0) : X \rightarrow Y$ , to be

$$d^+f(x_0)(x) = d^+f(x_0; x)$$

- (2) **left-hand Gateaux differentiable at  $x_0$**  if for each  $x \in X$ ,  $d^-f(x_0; x)$  exists. We define the **left-hand Gateaux derivative** of  $f$  at  $x_0$ , denoted  $d^-f(x_0) : X \rightarrow Y$ , to be

$$d^-f(x_0)(x) = d^-f(x_0; x)$$

- (3) **Gateaux differentiable at  $x_0$**  if for each  $x \in X$ ,  $df(x_0; x)$  exists. We define the **Gateaux derivative** of  $f$  at  $x_0$ , denoted  $df(x_0) : X \rightarrow Y$ , to be

$$df(x_0)(x) = df(x_0; x)$$

**Definition 8.1.5.** Let  $Y$  be a Banach space,  $A \subset \mathbb{R}$  open and  $f : A \rightarrow Y$ . Then  $f$  is said to be **Gateaux differentiable** if for each  $x \in A$ ,  $f$  is Gateaux differentiable at  $x$ . If  $f$  is Gateaux differentiable, we define  $df : A \rightarrow Y^X$  by  $x_0 \mapsto df(x_0)$ .

**Exercise 8.1.6.** Let  $X, Y$  be Banach spaces,  $A \subset X$  open,  $f, g : A \rightarrow Y$ ,  $\lambda \in \mathbb{R}$  and  $x_0 \in A$ . If  $f, g$  are Gateaux differentiable at  $x_0$ , then  $f + \lambda g$  is Gateaux differentiable at  $x_0$  and  $d[f + \lambda g](x_0) = df(x_0) + \lambda dg(x_0)$ .

*Proof.* Similar to the case of the derivative from Calc I. □

**Exercise 8.1.7.** Let  $X, Y$  be Banach spaces,  $A \subset X$  open,  $f : A \rightarrow Y$  and  $x_0 \in A$ . Suppose that  $f$  is Gateaux differentiable at  $x_0$ . Then for each  $\lambda \in \mathbb{R}$  and  $x \in X$ ,

$$df(x_0)(\lambda x) = \lambda df(x_0)(x)$$

*Proof.* Let  $\lambda \in \mathbb{R}$  and  $x \in X$ . Then

$$\begin{aligned} df(x_0)(\lambda x) &= \lim_{t \rightarrow 0} \frac{f(x_0 + t\lambda x) - f(x_0)}{t} \\ &= \lim_{t \rightarrow 0} \lambda \frac{f(x_0 + t\lambda x) - f(x_0)}{\lambda t} \\ &= \lambda \lim_{t \rightarrow 0} \frac{f(x_0 + t\lambda x) - f(x_0)}{\lambda t} \\ &= \lambda \lim_{t \rightarrow 0} \frac{f(x_0 + tx) - f(x_0)}{t} \\ &= \lambda df(x_0)(x) \end{aligned}$$

□

**Exercise 8.1.8.** Let  $X, Y$  be Banach spaces,  $A \subset X$  open,  $f : A \rightarrow Y$ . If  $f$  is constant, then  $f$  is Gateaux differentiable and for each  $x_0 \in A, x \in X$ ,

$$df(x_0)(x) = 0$$

*Proof.* Suppose that  $f$  is constant. Then there exists  $c \in Y$  such that for each  $x \in A$ ,  $f(x) = c$ . Let  $x_0 \in A, x \in X$ . Then

$$\begin{aligned} df(x_0)(x) &= \lim_{t \rightarrow 0} \frac{f(x_0 + tx) - f(x_0)}{t} \\ &= \lim_{t \rightarrow 0} \frac{c - c}{t} \\ &= 0 \end{aligned}$$

□

**Exercise 8.1.9.** Let  $X, Y$  be Banach spaces,  $A \subset X$  open,  $f : A \rightarrow Y$ . If  $f$  is linear, then  $f$  is Gateaux differentiable and for each  $x_0 \in A, x \in X$ ,

$$df(x_0)(x) = f(x)$$

*Proof.* Suppose that  $f$  is linear. Let  $x_0 \in A, x \in X$ . Then

$$\begin{aligned} df(x_0)(x) &= \lim_{t \rightarrow 0} \frac{f(x_0 + tx) - f(x_0)}{t} \\ &= \lim_{t \rightarrow 0} \frac{f(x_0) + tf(x) - f(x_0)}{t} \\ &= f(x) \end{aligned}$$

□

**Exercise 8.1.10.** There exist Banach spaces  $X, Y$ , and  $f : X \rightarrow Y$  such that  $f$  is Gateaux differentiable and  $f$  is nowhere continuous.

**Hint:** use Exercise 8.1.9

*Proof.* Set  $X = C^1([0, 1])$  and  $Y = C([0, 1])$ . Equip both  $X$  and  $Y$  with the sup norm. Define  $T : X \rightarrow Y$  by  $Tf = f'$ . Then Exercise 6.2.3 implies that  $T$  is not bounded. Since  $T$  is linear, Exercise 8.1.9 implies that  $T$  is Gateaux differentiable. Since  $T$  is not bounded, Exercise 6.2.6 implies that  $T$  is not continuous at 0. Then Exercise 6.2.5 tells us that  $T$  is nowhere continuous.  $\square$

**Exercise 8.1.11.** Set  $A = \{(x, y) \in \mathbb{R}^2 : y = -x^2 \text{ and } x \neq 0\}$ . Define  $f : \mathbb{R}^2 \setminus A \rightarrow \mathbb{R}$  by

$$f(x, y) = \begin{cases} 0 & (x, y) = (0, 0) \\ \frac{x^4 y}{x^6 + y^3} & \text{otherwise} \end{cases}$$

Then  $f$  is Gateaux differentiable at  $(0, 0)$  and  $f$  is not continuous at  $(0, 0)$ .

**Hint:** Consider the set  $B = \{(x, x^2) : x \in \mathbb{R}\} \subset \mathbb{R}^2 \setminus A$ .

*Proof.*  $\square$

**Exercise 8.1.12.** Let  $Y$  be a Banach space,  $A \subset \mathbb{R}$  open,  $f : A \rightarrow Y$  and  $x_0 \in A$ . Suppose that  $f$  is Gateaux differentiable at  $x_0$ . Then  $df(x_0) \in L(\mathbb{R}, Y)$ .

*Proof.* Let  $x, y, \lambda \in \mathbb{R}$ .

(1) The previous exercise implies

$$\begin{aligned} df(x_0)(x + \lambda y) &= df(x_0)((x + \lambda y)1) \\ &= (x + \lambda y)df(x_0)(1) \\ &= xdf(x_0)(1) + \lambda ydf(x_0)(1) \\ &= df(x_0)(x) + \lambda df(x_0)(y) \end{aligned}$$

So  $df(x_0) : \mathbb{R} \rightarrow Y$  is linear.

(2) Since

$$\begin{aligned} \|df(x_0)(x)\| &= \|xdf(x_0)(1)\| \\ &= |x|\|df(x_0)(1)\| \end{aligned}$$

We have that  $df(x_0) : \mathbb{R} \rightarrow Y$  is bounded with  $\|df(x_0)\| \leq \|df(x_0)(1)\|$ .  $\square$

**Exercise 8.1.13.** Let  $X$  be a Banach space,  $A \subset X$  open,  $f : A \rightarrow \mathbb{R}$  and  $x_0 \in A$ . If  $f$  is Gateaux differentiable at  $x_0$  and  $f$  has a local extremum at  $x_0$ , then  $df(x_0) = 0$ .

*Proof.* Suppose that  $f$  is Gateaux differentiable at  $x_0$  and  $f$  has a local minimum point at  $x_0$ . Then there exists  $\delta > 0$  such that  $B(x_0, \delta) \subset A$  and for each  $y \in B(x_0, \delta)$ ,  $f(x_0) \leq f(y)$ . For the sake of contradiction, suppose that  $df(x_0) \neq 0$ . Then there exists  $x \in X$  such that  $x \neq 0$  and  $df(x_0)(x) \neq 0$ .

First, suppose that  $df(x_0)(x) < 0$ . Choose  $\epsilon = -df(x_0)(x) > 0$ . Then there exists  $t_0 > 0$  such that for each  $t \in B^*(0, t_0)$ ,  $x_0 + tx \in B(x_0, \delta)$  and

$$\left| \frac{f(x_0 + tx) - f(x_0)}{t} - df(x_0)(x) \right| < \epsilon$$



This implies that for each  $t \in B^*(0, t_0)$ ,

$$\begin{aligned} \frac{f(x_0 + tx) - f(x_0)}{t} &< \epsilon + df(x_0)(x) \\ &= 0 \end{aligned}$$

and hence  $f(x_0 + tx) < f(x_0)$ , which is a contradiction.

Now, suppose that  $df(x_0)(x) > 0$ . Then

$$\begin{aligned} df(x_0)(-x) &= -df(x_0)(x) \\ &< 0 \end{aligned}$$

Similarly to above, this implies that there exists  $t_0 > 0$  such that for each  $t \in B^*(0, t_0)$ ,  $x_0 - tx \in B(x_0, \delta)$  and  $f(x_0 - tx) < f(x_0)$  which is a contradiction. So  $df(x_0)(x) = 0$  and  $df(x_0) = 0$ .

If  $f$  has a local maximum at  $x_0$ , then  $-f$  has a local minimum point at  $x_0$ . Then

$$\begin{aligned} df(x_0) &= -d[-f](x_0) \\ &= -0 \\ &= 0 \end{aligned}$$

□

**Exercise 8.1.14.** Let  $X, Y, Z$  be Banach spaces,  $A \subset X$  open,  $B \subset Y$  open,  $f : A \rightarrow Y$ ,  $g : B \rightarrow Z$  and  $x_0 \in A$ . Suppose that  $f$  is affine. If  $g$  is Gateaux differentiable at  $f(x_0)$ , then  $g \circ f$  is Gateaux differentiable at  $x_0$  and

$$d(g \circ f)(x_0)(x) = dg(f(x_0))(df(x_0)(x))$$

*Proof.* Suppose that  $g$  is Gateaux differentiable at  $f(x_0)$ . Since  $f$  is affine, there exists  $h : A \rightarrow Y$  and  $c \in Y$  such that  $f = h + c$ . Then

$$\begin{aligned} df(x_0) &= dh(x_0) \\ &= h \end{aligned}$$

Let  $x \in X$ . Choose  $\delta > 0$  such that for each  $t \in B(0, \delta) \subset \mathbb{R}$ ,  $f(x_0) + th(x) \in B$ . Then for each  $t \in B^*(0, \delta)$ ,

$$\begin{aligned} g \circ f(x_0 + tx) &= g\left(f(x_0) + t \frac{f(x_0 + tx) - f(x_0)}{t}\right) \\ &= g(f(x_0) + th(x)) \end{aligned}$$

This implies that

$$\begin{aligned} d(g \circ f)(x_0) &= \lim_{t \rightarrow 0} \frac{g \circ f(x_0 + tx) - g(f(x_0))}{t} \\ &= \lim_{t \rightarrow 0} \frac{g(f(x_0) + th(x)) - g(f(x_0))}{t} \\ &= dg(f(x_0))(h(x)) \\ &= dg(f(x_0))(df(x_0)(x)) \end{aligned}$$

□

## 8.2. The Frechet Derivative.

**Exercise 8.2.1.** Let  $X, Y$  be a normed vector spaces and  $\phi : X \rightarrow Y$  linear. If  $\phi(h) = o(\|h\|)$  as  $h \rightarrow 0$ , then  $\phi = 0$ .

*Proof.* Let  $h_0 \in X$ . If  $h_0 = 0$ , then  $\phi(h_0) = 0$ . Suppose that  $h_0 \neq 0$ . Define  $(h_n)_{n \in \mathbb{N}} \subset X$  by

$$h_n = \frac{h_0}{n}$$

Then  $h_n \rightarrow 0$ . By continuity of  $\phi$  and our initial assumption we have that

$$\begin{aligned} \|h_0\|^{-1}\phi(h_0) &= \phi\left(\frac{h_0}{\|h_0\|}\right) \\ &= \phi\left(\frac{h_n}{\|h_n\|}\right) \\ &= \frac{\phi(h_n)}{\|h_n\|} \\ &\rightarrow 0 \end{aligned}$$

which implies that  $\|h_0\|^{-1}\phi(h_0) = 0$ . So  $\phi(h_0) = 0$  and hence  $\phi = 0$ .  $\square$

**Exercise 8.2.2.** Let  $X, Y$  be a normed vector spaces,  $A \subset X$  open,  $f : A \rightarrow Y$  and  $x_0 \in A$ . Suppose that there exists  $\phi : X \rightarrow Y$  such that  $\phi$  is linear and

$$f(x_0 + h) = f(x_0) + \phi(h) + o(\|h\|) \quad \text{as } h \rightarrow 0$$

then  $\phi$  is unique.

*Proof.* Suppose that there exists  $\psi : X \rightarrow Y$  such that  $\psi$  is linear and such that

$$f(x_0 + h) = f(x_0) + \psi(h) + o(\|h\|) \quad \text{as } h \rightarrow 0$$

Then  $\phi(h) - \psi(h) = o(h)$ . Since  $\phi - \psi$  is linear, the previous exercise implies that  $\phi = \psi$ .  $\square$

**Note 8.2.3.** Recall that for Banach spaces  $X$  and  $Y$ , there isomorphic isometry

$$L(X, L(X, \dots, L(X, Y)) \dots) \rightarrow L^n(X, Y)$$

given by  $\phi \mapsto \psi_\phi$  where

$$\psi_\phi(x_1, x_2, \dots, x_n) = \phi(x_1)(x_2), \dots, (x_n)$$

### Definition 8.2.4. Frechet Derivative:

Let  $X, Y$  be a Banach spaces,  $A \subset X$  open,  $f : A \rightarrow Y$  and  $x_0 \in A$ .

- (1) • Then  $f$  is said to be **Frechet differentiable at**  $x_0$  if there exists  $Df(x_0) \in L(X, Y)$  such that,

$$f(x_0 + h) = f(x_0) + Df(x_0)(h) + o(\|h\|) \quad \text{as } h \rightarrow 0$$

- If  $f$  is Frechet differentiable at  $x_0$ , we define the **Frechet derivative of  $f$  at  $x_0$**  to be  $Df(x_0)$ .
- We say that  $f$  is **Frechet differentiable** if for each  $x \in A$ ,  $f$  is Frechet differentiable at  $x$ .
- If  $f$  is Frechet differentiable, we define the **Frechet derivative of  $f$** , denoted  $Df : A \rightarrow L(X, Y)$ , by  $x \mapsto D^{(1)}f(x)$ .

- (2) Continuing inductively, we set  $D^0f = f$  and for  $n \geq 2$ ,

- $f$  is said to be  **$n$ -th order Frechet differentiable at  $x_0$**  if  $f$  is  $(n-1)$ -th order Frechet differentiable and  $D^{n-1}f$  is Frechet differentiable at  $x_0$ .
- If  $f$  is  $n$ -th order Frechet differentiable at  $x_0$ , we define  $D^n f(x_0) \in L^n(X, Y)$  by

$$D^n f(x_0) = D[D^{n-1}f](x_0)$$

- We say that  $f$  is  **$n$ -th order Frechet differentiable** if  $f$  is  $(n-1)$ -th order Frechet differentiable and for each  $x \in A$ ,  $D^{n-1}f$  is Frechet differentiable at  $x$ .
- If  $f$  is  $n$ -th order Frechet differentiable, we define the  **$n$ -th order Frechet derivative of  $f$** , denoted  $D^n f : A \rightarrow L^n(X, Y)$  by  $x \mapsto D^n f(x)$

(3) If  $f$  is  $n$ -th order differentiable, then  $f$  is said to be **continuously  $n$ -th order differentiable** if  $D^n f$  is continuous. We define

$$C^n(A, Y) = \{f : A \rightarrow Y : f \text{ is continuously } n\text{-th order differentiable}\}$$

**Exercise 8.2.5.** Let  $X, Y$  be a banach spaces,  $A \subset X$  open,  $f, g : A \rightarrow Y$ ,  $\lambda \in \mathbb{R}$  and  $x_0 \in A$ . If  $f$  and  $g$  are Frechet differentiable at  $x_0$ , then  $f + \lambda g$  is Frechet differentiable at  $x_0$  and  $D(f + \lambda g)(x_0) = Df(x_0) + \lambda Dg(x_0)$ .

*Proof.* Suppose that  $f$  and  $g$  are Frechet differentiable at  $x_0$ . Then

$$f(x_0 + h) = f(x_0) + Df(x_0)(h) + o(\|h\|) \quad \text{as } h \rightarrow 0$$

and

$$g(x_0 + h) = g(x_0) + Dg(x_0)(h) + o(\|h\|) \quad \text{as } h \rightarrow 0$$

This implies that

$$\begin{aligned} (f + \lambda g)(x_0 + h) &= f(x_0 + h) + \lambda g(x_0 + h) \\ &= f(x_0) + Df(x_0)(h) + o(\|h\|) + \lambda g(x_0) + \lambda Dg(x_0)(h) + o(\|h\|) \\ &= (f + \lambda g)(x_0) + [Df(x_0) + \lambda Dg(x_0)](h) + o(\|h\|) \quad \text{as } h \rightarrow 0 \end{aligned}$$

Since  $Df(x_0) + \lambda Dg(x_0) \in L(X, Y)$ ,  $f + \lambda g$  is Frechet differentiable at  $x_0$  and  $D(f + \lambda g)(x_0) = Df(x_0) + \lambda Dg(x_0)$ .  $\square$

**Exercise 8.2.6.** Let  $X, Y$  be a banach spaces,  $A \subset X$  open,  $f : A \rightarrow Y$  and  $x_0 \in A$ . If  $f$  is Frechet differentiable at  $x_0$ , then  $f$  is continuous at  $x_0$ .

*Proof.* Suppose that  $f$  is Frechet differentiable at  $x_0$ . Then  $f(x) - f(x_0) = Df(x_0)(x - x_0) + o(\|x - x_0\|)$  as  $x \rightarrow x_0$ . Hence  $\|f(x) - f(x_0)\| \leq \|Df(x_0)\| \|x - x_0\| + o(\|x - x_0\|)$  as  $x \rightarrow x_0$ . This implies that  $f(x) \rightarrow f(x_0)$  as  $x \rightarrow x_0$  and therefore  $f$  is continuous at  $x_0$ .  $\square$

**Exercise 8.2.7.** Let  $X, Y$  be a banach spaces,  $A \subset X$  open,  $f : A \rightarrow Y$  and  $x_0 \in A$ . If  $f$  is Frechet differentiable at  $x_0$ , then  $f$  is Gateaux differentiable at  $x_0$  and  $df(x_0) = Df(x_0)$ .

*Proof.* Suppose that  $f$  is Frechet differentiable at  $x_0$ . Then  $f(x_0 + h) = f(x_0) + Df(x_0)(h) + o(\|h\|)$  as  $h \rightarrow 0$ . Let  $x \in X$ . Then  $f(x_0 + tx) - f(x_0) = tDf(x_0)(x) + o(t)$  as  $t \rightarrow 0$ . This implies that  $f$  is differentiable at  $x_0$  in the direction  $x$  and

$$\begin{aligned} df(x_0)(x) &= \lim_{t \rightarrow 0} \frac{f(x_0 + tx) - f(x_0)}{t} \\ &= Df(x_0)(x) \end{aligned}$$

Since  $x \in X$  is arbitrary,  $f$  is Gateaux differentiable at  $x_0$  and  $df(x_0) = Df(x_0)$ .  $\square$

**Exercise 8.2.8.** Let  $X$  be a Banach space,  $A \subset X$  open,  $f : A \rightarrow \mathbb{R}$  and  $x_0 \in A$ . If  $f$  is Frechet differentiable at  $x_0$  and  $f$  has a local extremum at  $x_0$ , then  $Df(x_0) = 0$ .

*Proof.* Suppose that  $f$  is Frechet differentiable at  $x_0$  and  $f$  has a local extremum at  $x_0$ . Two previous exercises imply that  $f$  is Gateaux differentiable at  $x_0$  and

$$\begin{aligned} Df(x_0) &= df(x_0) \\ &= 0 \end{aligned}$$

□

**Definition 8.2.9.** Let  $X, Y$  be a Banach spaces,  $A \subset X$  open,  $f : A \rightarrow Y$  and  $x_0 \in A$ . Suppose that  $f$  is Frechet differentiable at  $x_0$ . Define  $R_f(x_0) : A - x_0 \rightarrow Y$  by

$$R_f(x_0)(h) = f(x_0 + h) - f(x_0) - Df(x_0)(h)$$

**Exercise 8.2.10.** Let  $X, Y$  be a Banach spaces,  $A \subset X$  open,  $f : A \rightarrow Y$  and  $x_0 \in A$ . If  $f$  is Frechet differentiable at  $x_0$ , then

$$f(x_0 + h) - f(x_0) = O(\|h\|) \text{ as } h \rightarrow 0$$

*Proof.* Suppose that  $f$  is Frechet differentiable at  $x_0$ . Then  $R_f(h) = o(\|h\|)$  as  $h \rightarrow 0$ . Hence there exists  $\delta > 0$  such that  $B(0, \delta) \subset A - x_0$  and for each  $h \in B(0, \delta)$ ,  $\|R_f(h)\| \leq \|h\|$ . Hence for each  $h \in B(0, \delta)$

$$\begin{aligned} \|f(x_0 + h) - f(x_0)\| &= \|Df(x_0)(h) + R_f(x_0)(h)\| \\ &\leq \|Df(x_0)(h)\| + \|R_f(x_0)(h)\| \\ &\leq \|Df(x_0)\| \|h\| + \|h\| \\ &= (\|Df(x_0)\| + 1) \|h\| \end{aligned}$$

□

**Exercise 8.2.11. Chain Rule:**

Let  $X, Y, Z$  be a Banach spaces,  $A \subset X$  open,  $B \subset Y$  open,  $f : A \rightarrow Y$ ,  $g : B \rightarrow Z$  and  $x_0 \in A$ . Suppose that  $f(x_0) \in B$ . If  $f$  is Frechet differentiable at  $x_0$  and  $g$  is Frechet differentiable at  $f(x_0)$ , then  $g \circ f$  is Frechet differentiable at  $x_0$  and

$$D(g \circ f)(x_0) = Dg(f(x_0)) \circ Df(x_0)$$

*Proof.* Suppose that  $f$  is Frechet differentiable at  $x_0$  and  $g$  is Frechet differentiable at  $f(x_0)$ .

- The previous exercise implies that there exists  $\delta^* > 0$  and  $K > 0$  such that for each  $h \in B(0, \delta^*)$ ,  $\|f(x_0 + h) - f(x_0)\| \leq K\|h\|$ . Let  $\epsilon > 0$ . Since  $R_g(f(x_0))(h') = o(\|h'\|)$  as  $h' \rightarrow 0$ , there exists  $\delta' > 0$  such that for each  $h' \in B(0, \delta')$ ,  $\|R_g(f(x_0))(h')\| \leq \frac{\epsilon}{K} \|h'\|$ .

Choose  $\delta = \min(\delta'/K, \delta^*)$ . Let  $h \in B(0, \delta)$ . Then

$$\begin{aligned} \|f(x_0 + h) - f(x_0)\| &\leq K\|h\| \\ &< \delta' \end{aligned}$$

This implies that

$$\begin{aligned}\|R_g(f(x_0))(f(x_0+h) - f(x_0))\| &\leq \frac{\epsilon}{K} \|f(x_0+h) - f(x_0)\| \\ &\leq \frac{\epsilon}{K} K \|h\| \\ &\leq \epsilon \|h\|\end{aligned}$$

So  $R_g(f(x_0))(f(x_0+h) - f(x_0)) = o(\|h\|)$  as  $h \rightarrow 0$ .

- Since  $\|Dg(f(x_0))(R_f(x_0)(h))\| \leq \|Dg(f(x_0))\| \|R_f(x_0)(h)\|$  and  $R_f(x_0)(h) = o(h)$  as  $h \rightarrow 0$ , we have that  $Dg(f(x_0))(R_f(x_0)(h)) = o(h)$  as  $h \rightarrow 0$ .
- Combining the previous two observations, we have that  $Dg(f(x_0))(R_f(x_0)(h)) + R_g(f(x_0))(f(x_0+h) - f(x_0)) = o(\|h\|)$  as  $h \rightarrow 0$ .
- All together, we obtain

$$\begin{aligned}g \circ f(x_0+h) &= g(f(x_0)) + f(x_0+h) - f(x_0) \\ &= g(f(x_0)) + Dg(f(x_0))(f(x_0+h) - f(x_0)) + R_g(f(x_0))(f(x_0+h) - f(x_0)) \\ &= g(f(x_0)) + Dg(f(x_0))(Df(x_0)(h) + R_f(x_0)(h)) \\ &\quad + R_g(f(x_0))(f(x_0+h) - f(x_0)) \\ &= g(f(x_0)) + Dg(f(x_0))(Df(x_0)(h)) + Dg(f(x_0))(R_f(x_0)(h)) \\ &\quad + R_g(f(x_0))(f(x_0+h) - f(x_0)) \\ &= g \circ f(x_0) + Dg(f(x_0)) \circ Df(x_0)(h) + o(\|h\|) \text{ as } h \rightarrow 0\end{aligned}$$

So  $g \circ f$  is Frechet differentiable at  $x_0$  and  $D(g \circ f)(x_0) = Dg(f(x_0)) \circ Df(x_0)$ . □

**Exercise 8.2.12.** Let  $Y$  be a Banach space,  $A \subset \mathbb{R}$  open and  $f : A \rightarrow Y$ . Then  $f$  is Gateaux differentiable iff  $f$  is Frechet differentiable.

*Proof.* Suppose that  $f$  is Gateaux differentiable. Let  $x_0 \in A$ . A previous exercise implies that  $df(x_0) \in L(\mathbb{R}, Y)$ . By definition,

$$\lim_{h \rightarrow 0} \left\| \frac{f(x_0+h) - f(x_0)}{h} - df(x_0)(1) \right\| = 0$$

This is equivalent to saying that

$$f(x_0+h) = f(x_0) + df(x_0)(h) + o(|h|) \quad \text{as } h \rightarrow 0$$

So  $f$  is Frechet differentiable at  $x_0$  and  $Df(x_0) = df(x_0)$ . □

### 8.3. The Calc I Derivative.

#### Definition 8.3.1. Calc I Derivative:

Let  $Y$  be a Banach space,  $A \subset \mathbb{R}$  open,  $f : A \rightarrow Y$  and  $x_0 \in A$ .

- (1) • If  $f$  is Frechet differentiable at  $x_0$ , we define the **calc I derivative of  $f$  at  $x_0$** , denoted

$$f'(x_0) \text{ or } \frac{df}{dt}(x_0)$$

by

$$\begin{aligned} f'(x_0) &= \lim_{t \rightarrow 0} \frac{f(x_0 + t) - f(x_0)}{t} \\ &= df(x_0)(1) \\ &= Df(x_0)(1) \end{aligned}$$

- If  $f$  is Frechet differentiable, we define  $f' : A \rightarrow Y$  by  $x \mapsto f'(x)$ .
- (2) Continuing inductively, we set  $f^{(0)} = f$  and for  $n \geq 1$ ,
- if  $f^{(n-1)}$  is Frechet differentiable at  $x_0$ , we define the **( $n$ )-th order calc I derivative of  $f$  at  $x_0$** , denoted  $f^{(n)}(x_0)$ , by

$$f^{(n)}(x_0) = [f^{(n-1)}]'(x_0)$$

- if  $f^{(n-1)}$  is Frechet differentiable, we define  $f^{(n)} : A \rightarrow Y$  by

$$f^{(n)} = [f^{(n-1)}]'$$

**Exercise 8.3.2.** Let  $Y$  be a Banach space,  $A \subset \mathbb{R}$  open and  $f : A \rightarrow Y$ . If  $f$  is  $n$ -th order Frechet differentiable, then for each  $x_0 \in A$  and  $k \in \{1, \dots, n\}$ ,

$$f^{(k)}(x_0) = D^k f(x_0)(1^{\oplus k})$$

*Proof.* Let  $x_0 \in A$ . We proceed by induction. The base case is true by definition. Let  $k \in \{1, \dots, n\}$ . Suppose the claim is true for  $k-1$ . Then

$$f^{(k-1)}(x_0) = D^{k-1} f(x_0)(1^{\oplus(k-1)})$$

Since  $f$  is  $n$ -th order Frechet differentiable,

$$D^{k-1} f(x_0 + h) = D^{k-1} f(x_0) + D^k f(x_0)(h) + o(\|h\|) \quad \text{as } h \rightarrow 0$$

This implies that

$$\begin{aligned} f^{(k-1)}(x_0 + h) &= D^{k-1} f(x_0 + h)(1^{\oplus(k-1)}) \\ &= D^{k-1} f(x_0)(1^{\oplus(k-1)}) + D^k f(x_0)(h)(1^{\oplus(k-1)}) + o(\|h\|) \quad \text{as } h \rightarrow 0 \end{aligned}$$

Therefore for each  $h \in \mathbb{R}$ ,

$$Df^{(k-1)}(x_0)(h) = D^k f(x_0)(h)(1^{\oplus(k-1)})$$

and by definition,

$$\begin{aligned} f^{(k)}(x_0) &= [f^{(k-1)}]'(x_0) \\ &= Df^{(k-1)}(x_0)(1) \\ &= D^k f(x_0)(1^{\oplus k}) \end{aligned}$$

□

**Exercise 8.3.3.** Let  $X, Y$  be Banach spaces,  $A \subset X$  open,  $f \in C^n(A, Y)$ ,  $x_0 \in A$ , and  $h \in X$ . Suppose that  $\{x_0 + th : t \in [0, 1]\} \subset A$ . Define and  $g : (0, 1) \rightarrow Y$  by

$$g(t) = f(x_0 + th)$$

Then for each  $k \in \{1 \dots, n\}$  and  $t \in (0, 1)$ ,

$$g^{(k)}(t) = D^k f(x_0 + th)(h^{\oplus k})$$

*Proof.* We proceed by induction. It is straightforward to show that the claim is true for  $k = 1$ .

Let  $k \in \{1 \dots, n\}$ . Suppose that  $g^{(k-1)}(t) = D^{k-1} f(x_0 + th)(h^{\oplus(k-1)})$ . Since  $f \in C^k(A, Y)$ ,

$$D^{k-1} f(x_0 + s_0 h + th) = D^{k-1} f(x_0 + s_0 h) + D^k f(x_0 + s_0 h)(th) + o(\|t\|) \quad \text{as } t \rightarrow 0$$

The previous exercise implies that

$$\begin{aligned} g^{(k-1)}(s_0 + t) &= D^{k-1} g(s_0 + t)(1^{\oplus(k-1)}) \\ &= D^{k-1} f(x_0 + s_0 h + th)(h^{\oplus(k-1)}) \\ &= D^{k-1} f(x_0 + s_0 h)(h^{\oplus(k-1)}) + D^k f(x_0 + s_0 h)(th)(h^{\oplus(k-1)}) + o(\|t\|) \quad \text{as } t \rightarrow 0 \end{aligned}$$

Hence

$$Dg^{(k-1)}(s_0)(t) = D^k f(x_0 + s_0 h)(th)(h^{\oplus(k-1)})$$

and

$$\begin{aligned} g^{(k)}(t) &= Dg^{(k-1)}(t)(1) \\ &= D^k f(x_0 + th)(h^{\oplus k}) \end{aligned}$$

□

#### 8.4. Mean Value Theorem.

**Exercise 8.4.1.** Let  $X$  be a Banach space,  $A \subset X$  open and convex, and  $f : A \rightarrow \mathbb{R}$ . If  $f$  is continuous and Gateaux differentiable, then for each  $x, y \in A$ , there exists  $t^* \in (0, 1)$  such that  $f(x) - f(y) = df(t^*x + (1 - t^*)y)(x - y)$ .

*Proof.* Suppose that  $f$  is continuous and Gateaux differentiable. Let  $x, y \in A$ . The claim is clearly true when  $f(x) = f(y)$ . Suppose that  $f(x) \neq f(y)$ . Define  $h : [0, 1] \rightarrow X$  by  $h(t) = tx + (1 - t)y$ . Set  $g = f \circ h : [0, 1] \rightarrow \mathbb{R}$ . Then  $g$  is continuous on  $[0, 1]$  and Exercise 8.1.14 implies that  $g$  is Gateaux differentiable on  $(0, 1)$ . Then Exercise 8.2.12 Exercise 8.1.14 and the mean value theorem implies that there exists  $t^* \in (0, 1)$  such that

$$\begin{aligned} f(x) - f(y) &= g(1) - g(0) \\ &= g'(t^*) \\ &= dg(t^*)(1) \\ &= df(h(t^*))(dh(t^*)(1)) \\ &= df(h(t^*))(h'(t^*)) \\ &= df(t^*x + (1 - t^*)y)(x - y) \end{aligned}$$

□

**Exercise 8.4.2.** Let  $X$  be a Banach space,  $A \subset X$  open and convex, and  $f : A \rightarrow \mathbb{R}$ . If  $f$  is Frechet differentiable, then for each  $x, y \in A$ , there exists  $t^* \in (0, 1)$  such that  $f(x) - f(y) = Df(t^*x + (1 - t^*)y)(x - y)$ .

*Proof.* Suppose that  $f$  is Frechet differentiable. Then  $f$  is continuous and Gateaux differentiable. Now apply the previous exercise. □

#### Exercise 8.4.3. Mean Value Theorem:

Let  $X, Y$  be Banach spaces,  $A \subset X$  open and convex and  $f : A \rightarrow Y$ . If  $f$  is Frechet differentiable, then for each  $x, y \in A$ , there exists  $t^* \in (0, 1)$  such that

$$\|f(x) - f(y)\| \leq \|Df(t^*x + (1 - t^*)y)\| \|x - y\|$$

**Hint:** For  $x, y \in A$  with  $f(x) \neq f(y)$ , using a Hahn-Banach argument, find  $\lambda \in Y^*$  such that  $\|\lambda\| = 1$  and  $\lambda(f(x) - f(y)) = \|f(x) - f(y)\|$ .

*Proof.* Suppose that  $f$  is Frechet differentiable. Let  $x, y \in A$ . The claim is clearly true when  $f(x) = f(y)$ . Suppose that  $f(x) \neq f(y)$ . An exercise in the section on linear functionals implies that there exists  $\lambda \in Y^*$  such that  $\lambda(f(x) - f(y)) = \|f(x) - f(y)\|$  and  $\|\lambda\| = 1$ . Define  $g : [0, 1] \rightarrow \mathbb{R}$  by

$$g(t) = \lambda(f(tx + (1 - t)y))$$

Then  $g$  is continuous and (Frechet) differentiable on  $(0, 1)$  with

$$Dg(t)(h) = \lambda \circ Df(tx + (1 - t)y)((x - y)h)$$

which implies that

$$\begin{aligned} g'(t) &= Dg(t)(1) \\ &= \lambda \circ Df(tx + (1 - t)y)((x - y)) \end{aligned}$$



The mean value theorem implies that there exists  $t^* \in (0, 1)$  such that

$$\begin{aligned}\|f(x) - f(y)\| &= \lambda(f(x) - f(y)) \\ &= g(1) - g(0) \\ &= g'(t^*) \\ &= \lambda \circ Df(t^*x + (1 - t^*)y)((x - y))\end{aligned}$$

Taking absolute values, we see that

$$\begin{aligned}\|f(x) - f(y)\| &= |\lambda \circ Df(t^*x + (1 - t^*)y)((x - y))| \\ &\leq \|\lambda\| \|Df(t^*x + (1 - t^*)y)\| \|x - y\| \\ &\leq \|Df(t^*x + (1 - t^*)y)\| \|x - y\|\end{aligned}$$

□

**Exercise 8.4.4.** Let  $X, Y$  be Banach spaces,  $A \subset X$  open and convex and  $f : A \rightarrow Y$ . Suppose that  $f$  is Frechet differentiable. If for each  $x \in A$ ,  $Df(x) = 0$ , then  $f$  is constant.

*Proof.* Suppose that for each  $x \in A$ ,  $Df(x) = 0$ . Let  $x, y \in A$ . Then the mean value theorem implies that there exists  $t \in (0, 1)$  such that

$$\begin{aligned}\|f(x) - f(y)\| &\leq \|Df(tx + (1 - t)y)\| \|x - y\| \\ &= 0\end{aligned}$$

So  $f(x) = f(y)$ .

□

**Exercise 8.4.5.** Let  $X, Y$  be Banach spaces,  $A \subset X$  open and convex and  $f, g : A \rightarrow Y$ . Suppose that  $f$  and  $g$  are Frechet differentiable. If  $Df = Dg$ , then there exists  $c \in Y$  such that  $f = g + c$ .

*Proof.* Suppose that  $Df = Dg$ . Then  $D(f - g) = 0$  and the previous exercise implies that  $f - g$  is constant. □

**Exercise 8.4.6.** Let  $X, Y$  be Banach spaces,  $A \subset \mathbb{R}$  open and  $f : A \rightarrow Y$ . Suppose that  $f$  is Frechet differentiable. Then  $f' \in C(A, Y)$  iff  $f \in C^1(A, Y)$ .

*Proof.* Suppose that  $f' \in C(A, Y)$ . Let  $x, y \in A$  and  $h \in \mathbb{R}$ . Then

$$\begin{aligned}\|(Df(x) - Df(y))(h)\| &= \|Df(x)(h) - Df(y)(h)\| \\ &= \|hf'(x) - hf'(y)\| \\ &= \|h(f'(x) - f'(y))\| \\ &= \|f'(x) - f'(y)\| |h|\end{aligned}$$

So  $\|Df(x) - Df(y)\| \leq \|f'(x) - f'(y)\|$ . Hence continuity of  $f'$  implies continuity of  $Df$  and  $f \in C^1(A, Y)$ . Conversely, suppose that  $f \in C^1(A, Y)$ . Let  $x, y \in A$ . Then

$$\begin{aligned}\|f'(x) - f'(y)\| &= \|Df(x)(1) - Df(y)(1)\| \\ &= \|(Df(x) - Df(y))(1)\| \\ &\leq \|Df(x) - Df(y)\|\end{aligned}$$

Hence continuity of  $Df$  implies continuity of  $f'$  and  $f' \in C(A, Y)$ .

□

### 8.5. Taylor's Theorem.

**Note 8.5.1.** This section makes use of the Bochner integral. For reference, see .

**Exercise 8.5.2.** Let  $Y$  be a separable Banach space,  $f : [a, b] \rightarrow Y$  continuous so that  $f$  is Bochner-integrable. Define  $F : (a, b) \rightarrow Y$  by

$$F(x) = \int_{(a,x]} f dm$$

Then  $F \in C^1((a, b), Y)$  and for each  $x_0 \in (a, b)$  and  $F'(x_0) = f(x_0)$ .

*Proof.* Let  $x_0 \in (a, b)$  and  $h \in (0, b - x_0)$ . Then continuity implies that

$$\begin{aligned} \frac{1}{\|h\|} \left\| \int_{(x_0, x_0+h]} f - f(x_0) dm \right\| &\leq \frac{1}{\|h\|} \max_{x \in (x_0, x_0+h]} \|f(x) - f(x_0)\| \|h\| \\ &= \max_{x \in [x_0, x_0+h]} \|f(x) - f(x_0)\| \\ &\rightarrow 0 \text{ as } h \rightarrow 0 \end{aligned}$$

So

$$\int_{(x_0, x_0+h]} f - f(x_0) dm = o(\|h\|) \quad \text{as } h \rightarrow 0$$

Therefore

$$\begin{aligned} F(x_0 + h) &= \int_{(a, x_0+h]} f dm \\ &= \int_{(a, x_0]} f dm + \int_{(x_0, x_0+h]} f dm \\ &= \int_{(a, x_0]} f dm + hf(x_0) + \int_{(x_0, x_0+h]} f - f(x_0) dm \\ &= F(x_0) + hf(x_0) + o(\|h\|) \quad \text{as } h \rightarrow 0 \end{aligned}$$

The case is similar for  $h \in (x_0 - b, 0)$ . Since the map  $h \mapsto f(x_0)h$  is bounded,  $F$  is Frechet differentiable at  $x_0$  and  $DF(x_0)(h) = f(x_0)h$ . This implies that  $F'(x_0) = f(x_0)$  and the previous exercise implies tells us that continuity of  $f$  implies continuity of  $DF$ . So  $F \in C^1(A, Y)$ .  $\square$

**Exercise 8.5.3. Fundamental Theorem of Calculus:** Let  $Y$  be a separable Banach space and  $f \in C^1((a, b), Y)$ . Then for each  $x, x_0 \in (a, b)$ ,  $x_0 < x$  implies that

- (1)  $f'$  is Bochner integrable on  $(x_0, x]$
- (2)

$$f(x) - f(x_0) = \int_{(x_0, x]} f' dm$$

*Proof.* (1) Since  $f \in C^1((a, b), Y)$ , a previous exercise tells us that  $f' \in C_Y(a, b)$ . Let  $x, x_0 \in (a, b)$ . Suppose that  $x_0 < x$ . Choose  $c, d \in (a, b)$  such that  $a < c < x_0 < x < d < b$ . Then  $f'$  is continuous on  $[c, d]$  and hence Bochner-integrable on  $(c, d]$  and  $(x_0, x]$ .

(2) Define  $g : (c, d) \rightarrow Y$  by

$$g(\xi) = \int_{(c, \xi]} f' dm$$

Then the previous exercise implies that  $g \in C_Y^1(c, d)$  and for each  $t \in (c, d)$ ,  $g'(t) = f'(t)$ . Let  $t \in (c, d)$  and  $h \in \mathbb{R}$ . Then

$$\begin{aligned} Dg(t)(h) &= hg'(t) \\ &= hf'(t) \\ &= Df(t)(h) \end{aligned}$$

So  $Dg = Df$  on  $(c, d)$ . A previous exercise implies that there exists  $c \in Y$  such that  $f = g + c$  on  $(c, d)$ . Then

$$\begin{aligned} f(x) - f(x_0) &= g(x) + c - (g(x_0) + c) \\ &= g(x) - g(x_0) \\ &= \int_{(c, x]} f' dm - \int_{(c, x_0]} f' dm \\ &= \int_{(x_0, x]} f' dm \end{aligned}$$

□

**Exercise 8.5.4.** Let  $Y$  be a Banach space,  $A \subset \mathbb{R}$  open and  $g : A \rightarrow Y$ . If  $g$  is  $n$ -th order Frechet differentiable, then

$$\frac{d}{dt} \sum_{k=0}^{n-1} \frac{(1-t)^k}{k!} g^{(k)}(t) = \frac{(1-t)^{n-1}}{(n-1)!} g^{(n)}(t)$$

*Proof.* Taking the derivative yields a telescoping series. □

**Exercise 8.5.5. Taylor's Theorem I:**

Let  $X$  be a Banach space,  $Y$  a separable Banach space,  $A \subset X$  open and convex,  $f \in C^{n+1}(A, Y)$ ,  $x_0 \in A$ , and  $h \in A - x_0$ . Then

$$f(x_0 + h) = \sum_{k=0}^n \frac{1}{k!} D^k f(x_0)(h^{\oplus k}) + R(x_0)(h)$$

where  $R(x_0) : A - x_0 \rightarrow Y$  is defined by

$$R(x_0)(h) = \frac{1}{n!} \int_{(0,1)} (1-t)^n D^{n+1} f(x_0 + th)(h^{\oplus(n+1)}) dm(t)$$

and  $R(x_0)(h) = o(\|h\|^n)$  as  $h \rightarrow 0$ .

**Hint:** Define  $g : (0, 1) \rightarrow Y$  by

$$g(t) = f(x_0 + th)$$

Then use the previous exercise and the fundamental theorem of calculus.

*Proof.* Let  $h \in X$ . Suppose that  $x_0 + h \in A$ . Define  $g : (0, 1) \rightarrow Y$  by

$$g(t) = f(x_0 + th)$$

For each  $k \in \{1, \dots, n+1\}$ , a previous exercise implies that  $g^{(k)}(t) = D^k f(x_0 + th)(h^{\oplus k})$ , so  $g^{(k)}(0) = D^k f(x_0)(h^{\oplus k})$ . The previous exercise and the fundamental theorem of calculus tell us that

$$\begin{aligned}
 f(x_0 + h) - \sum_{k=0}^n \frac{1}{k!} D^k f(x_0)(h^{\oplus k}) &= g(1) - \sum_{k=0}^n \frac{1}{k!} g^{(k)}(0) \\
 &= \int_{(0,1)} \left[ \frac{d}{dt} \sum_{k=0}^n \frac{(1-t)^k}{k!} g^{(k)}(t) \right] dm(t) \\
 &= \int_{(0,1)} \frac{(1-t)^n}{n!} g^{(n+1)}(t) dm(t) \\
 &= \frac{1}{n!} \int_{(0,1)} (1-t)^n D^{n+1} f(x_0 + th)(h^{\oplus(n+1)}) dm(t) \\
 &= R(x_0)(h)
 \end{aligned}$$

Note that

$$\frac{1}{n+1} = \frac{1}{n!} \int_{(0,1)} (1-t)^n dm(t)$$

Since  $D^{n+1}f$  is continuous at  $x_0$ , there exists  $\delta_1 > 0$  such that for each  $h \in B(0, \delta_1)$ ,  $x_0 + h \in A$  and

$$\|D^{n+1}f(x_0 + h) - D^{n+1}f(x_0)\| < 1$$

Let  $\epsilon > 0$ . Choose  $\delta_2 > 0$  such that

$$\frac{1}{n+1} \left( \|D^{n+1}f(x_0)\| + 1 \right) \delta_2 < \epsilon$$

Set  $\delta = \min(\delta_1, \delta_2)$ . Let  $h \in B(0, \delta)$ . Then

$$\begin{aligned}
 \|R(x_0)(h)\| &= \left\| \int_{(0,1)} \frac{1}{n!} \int_{(0,1)} (1-t)^n D^{n+1}f(x_0 + th)(h^{\oplus(n+1)}) dm(t) \right\| \\
 &\leq \frac{1}{n!} \int_{(0,1)} \|(1-t)^n D^{n+1}f(x_0 + th)(h^{\oplus(n+1)})\| dm(t) \\
 &\leq \max_{t \in [0,1]} \|D^{n+1}f(x_0 + th)\| \|h\|^{n+1} \frac{1}{n!} \int_{(0,1)} (1-t)^n dm(t) \\
 &\leq \frac{1}{n+1} \left( \|D^{n+1}f(x_0)\| + \max_{t \in [0,1]} \|D^{n+1}f(x_0 + th) - D^{n+1}f(x_0)\| \right) \|h\|^{n+1} \\
 &< \frac{1}{n+1} \left( \|D^{n+1}f(x_0)\| + 1 \right) \|h\|^{n+1} \\
 &< \epsilon \|h\|^n
 \end{aligned}$$

So  $R(x_0)(h) = o(\|h\|^n)$  as  $h \rightarrow 0$ . □

### Exercise 8.5.6. Taylor's Theorem II:

Let  $X$  be a Banach space,  $Y$  a separable Banach space,  $A \subset X$  open and convex,  $f \in$

$C^n(A, Y)$ ,  $x_0 \in A$ , and  $h \in A - x_0$ . Then there exists  $R(x_0) : A - x_0 \rightarrow Y$  such that

$$f(x_0 + h) = \sum_{k=0}^n \frac{1}{k!} D^k f(x_0)(h^{\oplus k}) + R(x_0)(h)$$

and  $R(x_0)(h) = o(\|h\|^n)$  as  $h \rightarrow 0$ .

**Hint:** use Taylor's theorem and expand the derivative inside the integral.

*Proof.* This is clear by definition for  $n = 1$ . Suppose that  $n \geq 2$ . Taylor's theorem implies that

$$f(x_0 + h) = \sum_{k=0}^{n-2} \frac{1}{k!} D^k f(x_0)(h^{\oplus k}) + S(x_0)(h)$$

where  $S(x_0) : A - x_0 \rightarrow Y$  is defined by

$$S(x_0)(h) = \frac{1}{(n-2)!} \int_{(0,1)} (1-t)^{n-2} D^{n-1} f(x_0 + th)(h^{\oplus(n-1)}) dm(t)$$

and  $S(x_0; h) = o(\|h\|^n)$  as  $h \rightarrow 0$ . Define  $T^{n-1}(x_0) : A - x_0 \rightarrow L^{n-1}(X; Y)$  by

$$T^{n-1}(x_0)(h) = D^{n-1} f(x_0 + h) - D^{n-1} f(x_0) - D^n f(x_0)(h)$$

so that

$$D^{n-1} f(x_0 + h) = D^{n-1} f(x_0) + D^n f(x_0)(h) + T^{n-1}(x_0)(h)$$

and  $T^{n-1}(x_0)(h) = o(\|h\|)$  as  $h \rightarrow 0$ .

Define  $R(x_0) : A - x_0 \rightarrow Y$  by

$$R(x_0)(h) = \frac{1}{(n-2)!} \int_{(0,1)} (1-t)^{n-2} T^{n-1}(x_0)(th)(h^{\oplus(n-1)}) dm(t)$$

Note that

•

$$\int_0^1 (1-t)^{n-2} dt = \frac{1}{n-1}$$

•

$$\int_0^1 (1-t)^{n-2} t dt = \frac{1}{n(n-1)}$$

Let  $\epsilon > 0$ . Choose  $\delta > 0$  such that for each  $h \in B(0, \delta)$ ,  $h \in A - x_0$  and

$$\|T^{n-1}(x_0)(h)\| \leq \epsilon n! \|h\|$$

Let  $h \in B(0, \delta)$ . Then

$$\begin{aligned} \|R(x_0)(h)\| &= \left\| \frac{1}{(n-2)!} \int_{(0,1)} (1-t)^{n-2} T^{n-1}(x_0)(th)(h^{\oplus(n-1)}) dm(t) \right\| \\ &\leq \frac{1}{(n-2)!} \int_{(0,1)} (1-t)^{n-2} \|T^{n-1}(x_0)(th)(h^{\oplus(n-1)})\| dm(t) \\ &\leq \frac{1}{(n-2)!} \int_{(0,1)} (1-t)^{n-2} \|T^{n-1}(x_0)(th)\| \|h\|^{n-1} dm(t) \\ &\leq \frac{\epsilon}{(n-2)!} n! \|h\|^n \int_{(0,1)} (1-t)^{n-2} t dm(t) \\ &= \epsilon \|h\|^n \end{aligned}$$

So that  $R(x_0)(h) = o(\|h\|^n)$  as  $h \rightarrow 0$ .

Then

$$\begin{aligned}
 S(x_0)(h) &= \frac{1}{(n-2)!} \int_{(0,1)} (1-t)^{n-2} D^{n-1} f(x_0 + th)(h^{\oplus(n-1)}) dm(t) \\
 &= \frac{1}{(n-2)!} \int_{(0,1)} (1-t)^{n-2} D^{n-1} f(x_0)(h^{\oplus(n-1)}) dm(t) \\
 &\quad + \frac{1}{(n-2)!} \int_{(0,1)} (1-t)^{n-2} t D^n f(x_0)(h)(h^{\oplus(n-1)}) dm(t) \\
 &\quad + \frac{1}{(n-2)!} \int_{(0,1)} (1-t)^{n-2} T^{n-1}(x_0)(th)(h^{\oplus(n-1)}) dm(t) \\
 &= \frac{1}{(n-1)!} D^{n-1} f(x_0)(h^{\oplus(n-1)}) + \frac{1}{n!} D^n f(x_0)(h^{\oplus n}) + R_f(x_0)(h)
 \end{aligned}$$

Hence

$$\begin{aligned}
 f(x_0 + h) &= \sum_{k=0}^{n-2} \frac{1}{k!} D^k f(x_0)(h^{\oplus k}) + S(x_0)(h) \\
 &= \sum_{k=0}^n \frac{1}{k!} D^k f(x_0)(h^{\oplus k}) + R(x_0)(h)
 \end{aligned}$$

□

**Exercise 8.5.7. Taylor's Theorem III:**

Let  $X$  be a Banach space,  $A \subset X$  open and convex,  $f \in C^n(A)$ ,  $x_0 \in A$ , and  $h \in A - x_0$ . Then there exists  $t^* \in (0, 1)$  such that

$$f(x_0 + h) = \sum_{k=0}^{n-1} \frac{1}{k!} D^k f(x_0)(h^{\oplus k}) + \frac{1}{(n-1)!} (1-t^*)^{n-1} D^n f(x_0 + t^*h)(h^{\oplus n})$$

**Hint:** use Taylor's theorem and the mean value theorem.

*Proof.* Taylors Theorem implies that

$$f(x_0 + h) = \sum_{k=0}^{n-1} \frac{1}{k!} D^k f(x_0)(h^{\oplus k}) + R(x_0)(h)$$

where

$$R(x_0)(h) = \frac{1}{(n-1)!} \int_{(0,1)} (1-t)^{n-1} D^n f(x_0 + th)(h^{\oplus n}) dm(t)$$

Define  $F \in C^1([0, 1])$  by

$$F(t) = \int_{(0,t]} \frac{1}{(n-1)!} (1-s)^{n-1} D^n f(x_0 + sh)(h^{\oplus n}) dm(s)$$

Then the fundamental theorem of calculus implies that

$$F'(t) = \frac{1}{(n-1)!} (1-t)^{n-1} D^n f(x_0 + th)(h^{\oplus n})$$

The mean value theorem implies that there exists  $t^* \in (0, 1)$  such that

$$\begin{aligned} R(x_0)(h) &= F(1) - F(0) \\ &= F'(t^*) \\ &= \frac{1}{(n-1)!} (1-t^*)^{n-1} D^n f(x_0 + t^*h)(h^{\oplus n}) \end{aligned}$$

□

**Exercise 8.5.8.** Let  $X$  be a Banach space,  $A \subset X$  open and convex and  $f \in C^2(A)$ ,  $x_0 \in A$ . If  $f$  has a local minimum at  $x_0$ , then  $D^2f(x_0)$  is positive semidefinite.

*Proof.* Suppose that  $f$  has a local minimum at  $x_0$ , then  $Df(x_0) = 0$ . Let  $x \in X$ . Then

$$\begin{aligned} 0 &\leq f(x+h) - f(x_0) \\ &= \frac{1}{2} D^2f(x_0)(h, h) + o(\|h\|^2) \quad \text{as } h \rightarrow 0 \end{aligned}$$

Let  $h \in X$ . Then

$$0 \leq \frac{1}{2} t^2 D^2f(x_0)(h, h) + o(t^2) \quad \text{as } t \rightarrow 0$$

This implies that  $D^2f(x_0)(h, h) \geq 0$ . So  $D^2f(x_0)$  is positive semidefinite. □

### 8.6. The Gradient.

**Definition 8.6.1.** Let  $H$  be a Hilbert space,  $f : H \rightarrow \mathbb{R}$  and  $x_0 \in H$ . Suppose that  $f$  is Frechet differentiable at  $x_0$ . Then  $Df(x_0) \in H^*$ . We define the **gradient of  $f$  at  $x_0$** , denoted  $\nabla f(x_0) \in H$ , via the Riesz representation theorem to be the unique element of  $H$  satisfying

$$\langle \nabla f(x_0), y \rangle = Df(x_0)(y) \quad \text{for each } y \in H$$



## 9. CONVEXITY

## 9.1. Introduction.

**Note 9.1.1.** In this section, we assume all vector spaces are real.

**Definition 9.1.2.** Let  $X$  be a vector space and  $A \subset X$ . Then  $A$  is said to be **convex** if for each  $x, y \in A$ , and  $t \in [0, 1]$ ,  $tx + (1 - t)y \in A$ .

**Definition 9.1.3.** Let  $X$  be a vector space and  $f : A \rightarrow \mathbb{R}$ . Then  $f$  is said to be **convex** if for each  $x, y \in A$ ,  $t \in [0, 1]$ ,

$$f(tx + (1 - t)y) \leq tf(x) + (1 - t)f(y)$$

**Definition 9.1.4.** Let  $X$  be a vector space and  $f : A \rightarrow \mathbb{R}$ . Then  $f$  is said to be **strictly convex** if for each  $x, y \in A$ ,  $t \in (0, 1)$ ,  $x \neq y$  implies that

$$f(tx + (1 - t)y) < tf(x) + (1 - t)f(y)$$

**Exercise 9.1.5.** Let  $X$  be a vector space,  $f \in X^*$  and  $g : X \rightarrow \mathbb{R}$  constant. Then  $f$  and  $g$  are convex.

*Proof.* Let  $x, y \in X$  and  $t \in [0, 1]$ . Put  $c = g(0)$ . Then

$$f(tx + (1 - t)y) = tf(x) + (1 - t)f(y)$$

and

$$\begin{aligned} g(tx + (1 - t)y) &= c \\ &= tc + (1 - t)c \\ &= tg(x) + (1 - t)g(y) \end{aligned}$$

So  $f$  and  $g$  are convex. □

**Exercise 9.1.6.** Let  $f : [0, \infty) \rightarrow [0, \infty)$  be convex. If  $f(0) \leq 0$ , then for each  $x \in [0, \infty)$ ,  $t \in [0, 1]$ ,  $f(tx) \leq tf(x)$ .

*Proof.* Suppose that  $f(0) \leq 0$ . Let  $x \in [0, \infty)$  and  $t \in [0, 1]$ . Then

$$\begin{aligned} f(tx) &= f(tx + (1 - t)0) \\ &\leq tf(x) + (1 - t)f(0) \\ &\leq tf(x) \end{aligned}$$

□

**Exercise 9.1.7. Superadditivity:**

Let  $f : [0, \infty) \rightarrow [0, \infty)$  be convex. If  $f(0) = 0$ , then for each  $x, y \in [0, \infty)$ ,

$$f(x) + f(y) \leq f(x + y)$$

**Hint:**  $f(x) = f\left(\frac{x}{x+y}(x+y)\right)$

*Proof.* Suppose that  $f(0) \leq 0$ . Let  $x, y \in [0, \infty)$ . If  $x + y = 0$ , then  $x = y = 0$  and  $f(x) + f(y) = 0 = f(x + y)$ . Suppose that  $x + y \neq 0$ . Then the previous exercise implies that

$$\begin{aligned} f(x) + f(y) &= f\left(\frac{x}{x+y}(x+y)\right) + f\left(\frac{y}{x+y}(x+y)\right) \\ &\leq \frac{x}{x+y}f(x+y) + \frac{y}{x+y}f(x+y) \\ &= f(x+y) \end{aligned}$$

□

**Exercise 9.1.8.** Let  $X$  be a vector space,  $A \subset X$  convex,  $f, g : A \rightarrow \mathbb{R}$  and  $\lambda \geq 0$ . If  $f, g$  are convex, then

- (1)  $f + g$  is convex
- (2)  $\lambda f$  is convex

*Proof.* Suppose that  $f$  and  $g$  are convex. Let  $x, y \in A$  and  $t \in [0, 1]$ . Then

$$\begin{aligned} (f + \lambda g)(tx + (1-t)y) &= f(tx + (1-t)y) + \lambda g(tx + (1-t)y) \\ &\leq tf(x) + (1-t)f(y) + t\lambda g(x) + (1-t)\lambda g(y) \\ &= t(f(x) + \lambda g(x)) + (1-t)(f(y) + \lambda g(y)) \\ &= t(f + \lambda g)(x) + (1-t)(f + \lambda g)(y) \end{aligned}$$

□

**Definition 9.1.9.** Let  $X$  be a vector space and  $f : X \rightarrow \mathbb{R}$ . Then  $f$  is said to be **affine** if there exists  $\phi \in X^*$ ,  $a \in \mathbb{R}$  constant such that  $f = \phi + a$ .

**Exercise 9.1.10.** Let  $X$  be a vector space and  $f : X \rightarrow \mathbb{R}$ . If  $f$  is affine, then  $f$  is convex.

*Proof.* Suppose that  $f$  is affine. Then there exists  $\phi \in X^*$ ,  $a \in \mathbb{R}$  constant such that  $f = \phi + a$ . Then  $\phi$  is convex and  $g : X \rightarrow \mathbb{R}$  defined by  $g(x) = a$  is convex. So  $f = \phi + g$  is convex. □

**Exercise 9.1.11.** Let  $X$  be a vector space,  $A \subset X$  convex,  $f : \mathbb{R} \rightarrow \mathbb{R}$  and  $g : A \rightarrow \mathbb{R}$ . If  $f$  is convex and increasing and  $g$  is convex, then  $f \circ g$  is convex.

*Proof.* Let  $t \in [0, 1]$  and  $x, y \in A$ . Then convexity of  $g$  implies that

$$g(tx + (1-t)y) \leq tg(x) + (1-t)g(y)$$

and we have

$$\begin{aligned} f \circ g(tx + (1-t)y) &= f(g(tx + (1-t)y)) \\ &\leq f(tg(x) + (1-t)g(y)) && (f \text{ increasing}) \\ &\leq tf(g(x)) + (1-t)f(g(y)) && (f \text{ convex}) \\ &= tf \circ g(x) + (1-t)f \circ g(y) \end{aligned}$$

So  $f \circ g$  is convex. □

**Exercise 9.1.12.** Let  $X$  be a vector space,  $A \subset X$  convex,  $f : A \rightarrow \mathbb{R}$  convex and  $x_0 \in A$ . Then  $f$  has a local minimum point at  $x_0$  iff  $f$  has a global minimum point at  $x_0$ .

*Proof.* If  $f$  has a global minimum point at  $x_0$ , then  $f$  has a local minimum point at  $x_0$ . Conversely, suppose that  $f$  has a local minimum point at  $x_0$ . Then there exists  $\delta > 0$  such that for each  $x \in B(x_0, \delta) \cap A$ ,  $f(x_0) \leq f(x)$ . For the sake of contradiction, suppose that  $f$  does not have a global minimum point at  $x_0$ . Then there exists  $x' \in A$  such that  $f(x') < f(x_0)$ . Put  $t_0 = \min(\frac{\delta}{\|x' - x_0\| + 1}, 1) > 0$ . Let  $t \in (0, t_0)$ , then

$$\begin{aligned} \|(tx' + (1-t)x_0) - x_0\| &= t\|x' - x_0\| \\ &< \frac{\|x' - x_0\|\delta}{\|x' - x_0\| + 1} \\ &< \delta \end{aligned}$$

so that  $tx' + (1-t)x_0 \in B(x_0, \delta) \cap A$  and hence  $f(x_0) \leq f(tx' + (1-t)x_0)$ . Therefore

$$\begin{aligned} f(x_0) &\leq f(tx' + (1-t)x_0) \\ &\leq tf(x') + (1-t)f(x_0) \quad (\text{convexity of } f) \\ &< tf(x_0) + (1-t)f(x_0) \\ &= f(x_0) \end{aligned}$$

which is a contradiction. Hence  $f$  has a global minimum point at  $x_0$ .  $\square$

**Exercise 9.1.13.** Let  $X$  be a vector space,  $A \subset X$  convex,  $f : A \rightarrow \mathbb{R}$  strictly convex and  $x_0 \in X$ . If  $f$  has a local minimum point at  $x_0$ , then  $f$  has a unique global minimum point at  $x_0$ .

*Proof.* Suppose that  $f$  has a local minimum point at  $x_0$ . The previous exercise implies that  $f$  has a global minimum point at  $x_0$ . For the sake of contradiction suppose that there exists  $x_1 \in X$  such that  $f$  has a global minimum point at  $x_1$  and  $x_0 \neq x_1$ . This implies  $f(x_0) = f(x_1)$ . Set  $t = 1/2$ . Strict convexity implies that

$$\begin{aligned} f(tx_0 + (1-t)x_1) &< tf(x_0) + (1-t)f(x_1) \\ &= f(x_0) \end{aligned}$$

which is a contradiction since  $f$  has a global minimum point at  $x_0$ .  $\square$

**Definition 9.1.14.** Let  $X, Y$  be vector spaces,  $A \subset X \oplus Y$ . For  $y \in Y$ , define

$$A^y = \{x \in X : (x, y) \in A\}$$

and  $f^y : A^y \rightarrow \mathbb{R}$  by

$$f^y(x) = f(x, y)$$

**Exercise 9.1.15.** Let  $X, Y$  be vector spaces,  $A \subset X \oplus Y$  convex and  $f : A \rightarrow \mathbb{R}$  convex. Then for each  $y \in \pi_2(A)$ ,

- (1)  $A^y$  is convex
- (2)  $f^y$  is convex

where  $\pi_2 : X \times Y \rightarrow Y$ , the canonical projection of  $X \times Y$  onto  $Y$  given by  $\pi_2(x, y) = y$ .

*Proof.* Let  $y \in \pi_2(A)$ ,  $x_1, x_2 \in A^y$  and  $t \in [0, 1]$ . Then by definition,  $(x_1, y), (x_2, y) \in A$ .

- (1) Convexity of  $A$  implies that  $(tx_1 + (1-t)x_2, y) \in A$ . Hence  $tx_1 + (1-t)x_2 \in A^y$  and  $A^y$  is convex.

(2) Convexity of  $f$  implies that

$$\begin{aligned} f^y(tx_1 + (1-t)x_2) &= f(tx_1 + (1-t)x_2, y) \\ &= f(t(x_1, y) + (1-t)(x_2, y)) \\ &\leq tf(x_1, y) + (1-t)f(x_2, y) \\ &= tf^y(x_1) + (1-t)f^y(x_2) \end{aligned}$$

and so  $f^y$  is convex. □

**Exercise 9.1.16.** Let  $X, Y$  be vector spaces and  $A \subset X, B \subset Y$ . If  $A$  and  $B$  are convex, then  $A \times B \subset X \oplus Y$  is convex.

*Proof.* Suppose that  $A$  and  $B$  are convex. Let  $(x_1, y_1), (x_2, y_2) \in A \times B$  and  $t \in [0, 1]$ . Convexity of  $A$  and  $B$  implies that  $tx_1 + (1-t)x_2 \in A$  and  $ty_1 + (1-t)y_2 \in B$ . Therefore

$$\begin{aligned} t(x_1, y_1) + (1-t)(x_2, y_2) &= (tx_1 + (1-t)x_2, ty_1 + (1-t)y_2) \\ &\in A \times B \end{aligned}$$
□

**Exercise 9.1.17.** Let  $X, Y$  be vector spaces and  $A \subset X, B \subset Y$  convex (implying that  $A \times B$  is convex) and  $f : A \times B \rightarrow \mathbb{R}$  convex. Suppose that for each  $y \in B$ ,  $\{f(x, y) : x \in A\}$  is bounded below. Then  $\inf_{y \in B} f^y$  is convex

*Proof.* Put  $g = \inf_{y \in B} f^y$ . Let  $x_1, x_2 \in A, y_1, y_2 \in B$  and  $t \in [0, 1]$ . Put  $y' = ty_1 + (1-t)y_2$ . Then convexity of  $f$  implies that

$$\begin{aligned} g(tx_1 + (1-t)x_2) &\leq f^{y'}(tx_1 + (1-t)x_2) \\ &= f(tx_1 + (1-t)x_2, ty_1 + (1-t)y_2) \\ &= f(t(x_1, y_1) + (1-t)(x_2, y_2)) \\ &\leq tf(x_1, y_1) + (1-t)f(x_2, y_2) \\ &= tf^{y_1}(x_1) + (1-t)f^{y_2}(x_2) \end{aligned}$$

Since  $y_1 \in B$  is arbitrary, we have that

$$g(tx_1 + (1-t)x_2) \leq tg(x_1) + (1-t)f^{y_2}(x_2)$$

Similarly, since  $y_2 \in B$  is arbitrary, we have that

$$g(tx_1 + (1-t)x_2) \leq tg(x_1) + (1-t)g(x_2)$$

and  $f$  is convex. □

**Exercise 9.1.18.** Let  $X$  be a vector space,  $A \subset X$  convex and  $(f_\lambda)_{\lambda \in \Lambda} \subset \mathbb{R}^A$ . Suppose that for each  $\lambda \in \Lambda$ ,  $f_\lambda$  is convex. Define

- (1)  $A^* = \{x \in A : \sup_{\lambda \in \Lambda} f_\lambda(x) < \infty\}$
- (2)  $f^* : A^* \rightarrow \mathbb{R}$  by  $f^*(x) = \sup_{\lambda \in \Lambda} f_\lambda(x)$

Then

- (1)  $A^*$  is convex
- (2)  $f^*$  is convex

*Proof.* (1) Let  $x, y \in A$  and  $t \in [0, 1]$ . By definition,  $\sup_{\lambda \in \Lambda} f_\lambda(x), \sup_{\lambda \in \Lambda} f_\lambda(y) < \infty$ . Therefore

$$\begin{aligned} \sup_{\lambda \in \Lambda} f_\lambda(tx + (1-t)y) &\leq \sup_{\lambda \in \Lambda} [tf_\lambda(x) + (1-t)f_\lambda(y)] \\ &\leq t \sup_{\lambda \in \Lambda} f_\lambda(x) + (1-t) \sup_{\lambda \in \Lambda} f_\lambda(y) \\ &< \infty \end{aligned}$$

So  $tx + (1-t)y \in A$ .

- (2) By definition, the previous part implies that for each  $x, y \in A^*$ ,  $f^*(tx + (1-t)y) \leq tf^*(x) + (1-t)f^*(y)$ . So  $f^* : A^* \rightarrow \mathbb{R}$  is convex. □

**Exercise 9.1.19.** Let  $X$  be a normed vector space,  $A \subset X$  open and convex,  $f : A \rightarrow \mathbb{R}$  convex and  $x_0 \in A$ . If  $f$  is continuous at  $x_0$ , then  $f$  is locally Lipschitz at  $x_0$ .

**Hint:** Given  $x_1, x_2$  near  $x_0$  Choose a  $z$  near  $x_0$  s.t.  $x_1$  is a convex combination of  $x_2$  and  $z$ . Then repeat but with  $x_2$  as a convex combination of  $x_1$  and  $z$

*Proof.* By continuity,  $f$  is locally bounded at  $x_0$ . So there exist  $M, \delta > 0$  such that  $B(x_0, \delta) \subset A$  and for each  $x \in B(x_0, \delta)$ ,  $|f(x)| \leq M$ . Put  $\delta' = \frac{\delta}{2}$  and choose  $U = B(x_0, \delta')$ . Then  $U \subset A$  and  $U \in \mathcal{N}_{x_0}$ .

Let  $x_1, x_2 \in U$ . Suppose that  $x_1 \neq x_2$ . Define  $\alpha = \|x_1 - x_2\| > 0$ ,  $p = \frac{\alpha}{\alpha + \delta'}$ ,  $q = 1 - p$  and  $z = p^{-1}(x_1 - qx_2)$ . Then  $x_1 = pz + qx_2$  and

$$\begin{aligned} \|z - x_1\| &= \|(p^{-1} - 1)x_1 - p^{-1}qx_2\| \\ &= \frac{1-p}{p}\alpha \\ &= \frac{\delta'}{\alpha}\alpha \\ &= \delta' \end{aligned}$$

Therefore

$$\begin{aligned} \|z - x_0\| &\leq \|z - x_1\| + \|x_1 - x_0\| \\ &< \delta' + \delta' \\ &= \delta \end{aligned}$$

So  $z \in B(x_0, \delta)$ , which implies that

$$\begin{aligned} |f(z) - f(x_2)| &\leq |f(z) - f(x_2)| \\ &\leq |f(z)| + |f(x_2)| \\ &\leq 2M \end{aligned}$$

Since  $x_1 = pz + qx_2$ , convexity of  $f$  implies that  $f(x_1) \leq pf(z) + qf(x_2)$ . Hence

$$\begin{aligned}
 f(x_1) - f(x_2) &\leq pf(z) - pf(x_2) \\
 &= p(f(z) - f(x_2)) \\
 &\leq p2M \\
 &= \frac{\alpha}{\alpha + \delta'} 2M \\
 &\leq \alpha 2M \\
 &= 2M\|x_1 - x_2\|
 \end{aligned}$$

Similarly, choosing  $z = p^{-1}(x_2 - qx_1)$ , yields  $f(x_2) - f(x_1) \leq 2M\|x_1 - x_2\|$  which implies that

$$|f(x_1) - f(x_2)| \leq 2M\|x_1 - x_2\|$$

and  $f$  is Lipschitz on  $U$ . □

## 9.2. The Subdifferential.

**Exercise 9.2.1.** Let  $X$  be a Banach space,  $A \subset X$  open and convex,  $f : A \rightarrow \mathbb{R}$  convex,  $x_0 \in A$  and  $x \in X$ . Define  $T = \{t \in \mathbb{R} : x_0 + tx \in A\}$ . Then there exist  $a, b \in (0, \infty]$  such that  $T = (-a, b)$ .

*Proof.* Continuity of scalar multiplication and addition implies that  $T$  is an open neighborhood of 0. Let  $t > 0$  and  $s \in [0, t]$ . Then  $\frac{s}{t} \in [0, 1]$  and by convexity of  $A$ ,  $x_0 + tx \in A$  implies that

$$\begin{aligned} x_0 + sx &= \frac{s}{t}(x_0 + tx) + \left(1 - \frac{s}{t}\right)x_0 \\ &\in A \end{aligned}$$

Thus  $[0, t] \subset T$ . Similarly,  $x_0 - tx \in A$  implies that  $[-t, 0] \subset T$ .

Define  $a, b \in (0, \infty]$  by  $a = \sup\{t > 0 : x_0 - tx \in A\}$  and  $b = \sup\{t > 0 : x_0 + tx \in A\}$ . Then  $(-a, b) = T$ .  $\square$

**Definition 9.2.2.** Let  $X$  be a Banach space,  $A \subset X$  open and convex,  $f : A \rightarrow \mathbb{R}$  convex,  $x_0 \in A$  and  $x \in X$ . Define  $T$  as in the previous exercise and choose  $t_0 > 0$  such that  $(-t_0, t_0) \subset T$ . For  $t \in (0, t_0)$ , define the difference quotient  $q : (-t_0, t_0) \setminus \{0\} \rightarrow \mathbb{R}$  by

$$q(t) = \frac{f(x_0 + tx) - f(x_0)}{t}$$

**Exercise 9.2.3.** Let  $X$  be a Banach space,  $A \subset X$  open and convex,  $f : A \rightarrow \mathbb{R}$  convex,  $x_0 \in A$  and  $x \in X$ . Define  $t_0$  as above. Then

- (1)  $q(t)$  is increasing on  $(0, t_0)$
- (2)  $q(-t)$  decreasing on  $(0, t_0)$

**Hint:** As an example, look at the graph of  $f(x) = x^2$ . For the algebra, start at the desired end inequality and work backwards

*Proof.*

- (1) Let  $s, t \in (0, t_0)$  and suppose that  $s \leq t$ . Then  $x_0 + sx, x_0 + tx \in A$ . Note that since  $0 < s \leq t$ ,  $\frac{s}{t} \in (0, 1]$  and  $1 - \frac{s}{t} = \frac{t-s}{t} \in (0, 1]$ . Also, since  $A$  is convex, we have that

$$\left(\frac{t-s}{t}\right)x_0 + \left(\frac{s}{t}\right)(x_0 + tx) \in A$$

Convexity of  $f$  implies that

$$\begin{aligned} f(x_0 + sx) &= f\left(\left(\frac{t-s}{t}\right)x_0 + \left(\frac{s}{t}\right)(x_0 + tx)\right) \\ &\leq \left(\frac{t-s}{t}\right)f(x_0) + \left(\frac{s}{t}\right)f(x_0 + tx) \end{aligned}$$

This implies that

$$tf(x_0 + sx) \leq (t-s)f(x_0) + sf(x_0 + tx)$$

and after rearranging, we get

$$tf(x_0 + sx) - tf(x_0) \leq sf(x_0 + tx) - sf(x_0)$$

and so finally, dividing both sides by  $st$ , we obtain

$$\begin{aligned} q(s) &= \frac{f(x_0 + sx) - f(x_0)}{s} \\ &\leq \frac{f(x_0 + tx) - f(x_0)}{t} \\ &= q(t) \end{aligned}$$

as desired.

(2) Similar to (1). □

**Exercise 9.2.4.** Let  $X$  be a Banach space,  $A \subset X$  open and convex,  $f : A \rightarrow \mathbb{R}$  convex,  $x_0 \in A$  and  $x \in X$ . Define  $t_0$  as before. Then for each  $t \in (0, t_0)$ ,

$$q(-t) \leq q(t)$$

**Hint:** for sufficiently small  $t$ , convexity of  $f$  implies that  $f(x_0) \leq \frac{1}{2}f(x_0 - 2tx) + \frac{1}{2}f(x_0 + 2tx)$

*Proof.* Choose  $t_0$  as in the previous exercise. Since convexity of  $f$  implies that for each  $t \in (0, t_0/2)$ ,

$$f(x_0) \leq \frac{1}{2}f(x_0 - 2tx) + \frac{1}{2}f(x_0 + 2tx)$$

we have that for each  $t \in (0, t_0/2)$ ,

$$\begin{aligned} q(-2t) &= \frac{f(x_0 - 2tx) - f(x_0)}{-2t} \\ &\leq \frac{f(x_0 + 2tx) - f(x_0)}{2t} \\ &= q(2t) \end{aligned}$$

So for each  $t \in (0, t_0)$ ,  $q(-t) \leq q(t)$ . □

**Exercise 9.2.5.** Let  $X$  be a Banach space,  $A \subset X$  open and convex,  $f : A \rightarrow \mathbb{R}$  convex and  $x_0 \in A$ . Then

- (1)  $f$  is left-hand and right-hand Gateaux differentiable at  $x_0$  with  $d^-f(x_0) \leq d^+f(x_0)$
- (2) for each  $x \in X$ ,  $d^-f(x_0)(x) = -d^+f(x_0)(-x)$

*Proof.*

- (1) Let  $x \in X$ . Choose  $t_0 > 0$  as in the previous two exercises. Let  $t, u \in (0, t_0)$ . Choose  $s \in (0, \min(u, t))$ . The previous two exercises imply that

$$\begin{aligned} q(-u) &\leq q(-s) \\ &\leq q(s) \\ &\leq q(t) \end{aligned}$$

and therefore  $q(t)$  is an upper bound for  $\{q(-u) : u \in (0, t_0)\}$  and  $d^-f(x_0)(x) = \sup_{u \in (0, t_0)} q(-u)$  exists with  $d^-f(x_0)(x) \leq q(t)$ .



Since  $t \in (0, t_0)$  is arbitrary,  $d^-f(x_0)(x)$  is a lower bound for  $\{q(t) : t \in (0, t_0)\}$ . Therefore

$$d^+f(x_0)(x) = \inf_{t \in (0, t_0)} q(t)$$

exists with  $d^+f(x_0)(x) \geq d^-f(x_0)(x)$ .

(2) By definition, we have

$$\begin{aligned} d^-f(x_0)(x) &= \lim_{t \rightarrow 0^+} \frac{f(x_0 - tx) - f(x_0)}{-t} \\ &= - \lim_{t \rightarrow 0^+} \frac{f(x_0 - tx) - f(x_0)}{t} \\ &= -d^+f(x_0)(-x) \end{aligned}$$

□

**Exercise 9.2.6.** Let  $X$  be a Banach space,  $A \subset X$  open and convex,  $f : A \rightarrow \mathbb{R}$  convex and  $x_0 \in A$ . Then  $d^+f(x_0) : X \rightarrow \mathbb{R}$  is a sublinear functional.

*Proof.* Let  $x, y \in X$  and  $k \geq 0$ . If  $k = 0$ , then clearly

$$d^+f(x_0)(kx) = kd^+(x_0)(x)$$

If  $k > 0$ . Then

$$\begin{aligned} d^+f(x_0)(kx) &= \lim_{t \rightarrow 0^+} \frac{f(x_0 + tkx) - f(x_0)}{t} \\ &= k \lim_{t \rightarrow 0^+} \frac{f(x_0 + tkx) - f(x_0)}{tk} \\ &= kd^+f(x_0)(x) \end{aligned}$$

Define  $t_0 > 0$  as before and let  $t \in (0, \frac{t_0}{2})$ . Note that

$$x_0 + tx + ty = \frac{1}{2}(x_0 + 2tx) + \frac{1}{2}(x_0 + 2ty)$$

Convexity of  $f$  implies that

$$f(x_0 + tx + ty) \leq \frac{1}{2}f(x_0 + 2tx) + \frac{1}{2}f(x_0 + 2ty)$$

which implies that

$$\frac{f(x_0 + tx + ty) - f(x_0)}{t} \leq \frac{f(x_0 + 2tx) - f(x_0)}{2t} + \frac{f(x_0 + 2ty) - f(x_0)}{2t}$$

Therefore

$$\begin{aligned}
d^+f(x_0)(x+y) &= \lim_{t \rightarrow 0^+} \frac{f(x_0 + t(x+y)) - f(x_0)}{t} \\
&= \lim_{t \rightarrow 0^+} \frac{f(x_0 + tx + ty) - f(x_0)}{t} \\
&\leq \lim_{t \rightarrow 0^+} \left[ \frac{f(x_0 + 2tx) - f(x_0)}{2t} + \frac{f(x_0 + 2ty) - f(x_0)}{2t} \right] \\
&= \lim_{t \rightarrow 0^+} \frac{f(x_0 + 2tx) - f(x_0)}{2t} + \lim_{t \rightarrow 0^+} \frac{f(x_0 + 2ty) - f(x_0)}{2t} \\
&= d^+f(x_0)(x) + d^+f(x_0)(y)
\end{aligned}$$

□

**Exercise 9.2.7.** Let  $X$  be a Banach space,  $A \subset X$  open and convex,  $f : A \rightarrow \mathbb{R}$  convex and  $x_0 \in A$ . Then for each  $x \in A$ ,

$$d^+f(x_0)(x - x_0) \leq f(x) - f(x_0)$$

*Proof.* Let  $x \in A$ . Define  $T = \{t \in \mathbb{R} : x_0 + t(x - x_0) \in A\}$  similarly to earlier. Clearly  $1 \in T$  and

$$\begin{aligned}
d^+f(x_0)(x - x_0) &= \inf_{t \in (0,1]} \frac{f(x_0 + t(x - x_0)) - f(x_0)}{t} \\
&\leq f(x) - f(x_0)
\end{aligned}$$

□

**Exercise 9.2.8.** Let  $X$  be a Banach space,  $A \subset X$  open and convex,  $f : A \rightarrow \mathbb{R}$  convex and  $x_0 \in A$ . If  $f$  is continuous at  $x_0$ , then  $d^+f(x_0)$  is Lipschitz (equivalently bounded).

*Proof.* Suppose that  $f$  is continuous at  $x_0$ . A previous exercise about convex functions tells us that  $f$  is locally Lipschitz at  $x_0$ , so there exists  $\delta, M > 0$  such that for each  $x_1, x_2 \in B(x_0, \delta)$ ,  $|f(x_1) - f(x_2)| \leq M\|x_1 - x_2\|$ . Let  $x \in X$  and define  $t_0 = \frac{\delta}{\|x\|+1}$  so that for each  $t \in (0, t_0)$ ,

$$\begin{aligned}
\|(x_0 + tx) - x_0\| &= t\|x\| \\
&\leq t_0\|x\| \\
&= \frac{\delta\|x\|}{\|x\| + 1} \\
&< \delta
\end{aligned}$$

and  $x_0 + tx \in B(x_0, \delta)$ . Then for each  $t \in (0, t_0)$ ,

$$\begin{aligned}
d^+f(x_0)(x) &\leq \frac{f(x_0 + tx) - f(x_0)}{t} \\
&\leq \frac{|f(x_0 + tx) - f(x_0)|}{t} \\
&\leq t^{-1}M\|(x_0 + tx) - x_0\| \\
&= M\|x\|
\end{aligned}$$

Thus  $d^+f(x_0)$  is a bounded sublinear functional and a previous exercise in the section on sublinear functionals implies this is equivalent to  $d^+f(x_0)$  being Lipschitz. □

**Exercise 9.2.9.** Let  $X$  be a Banach space,  $A \subset X$  open and convex,  $f : A \rightarrow \mathbb{R}$  convex and  $x_0 \in A$ . If  $f$  is continuous at  $x_0$ , then there exists  $\phi \in X^*$  such that  $\phi \leq d^+f(x_0)$ .

*Proof.* Suppose that  $f$  is continuous at  $x_0$ . The previous exercise implies that  $d^+f(x_0)$  is Lipschitz (equivalently bounded). A previous exercise in the section discussing sublinear functionals tells us that boundedness of  $d^+f(x_0)$  implies that there exists  $\phi \in X^*$  such that  $\phi \leq d^+f(x_0)$ .  $\square$

**Definition 9.2.10. Subdifferential:**

Let  $X$  be a Banach space,  $A \subset X$  open and convex,  $f : A \rightarrow \mathbb{R}$  convex and  $x_0 \in A$ . We define the **subdifferential of  $f$  at  $x_0$** , denoted  $\partial f(x_0)$ , to be

$$\partial f(x_0) = \{\phi \in X^* : \text{for each } x \in A, f(x_0) + \phi(x - x_0) \leq f(x)\}$$

**Exercise 9.2.11.** Let  $X$  be a Banach space,  $A \subset X$  open and convex,  $f : A \rightarrow \mathbb{R}$  convex and  $x_0 \in A$ . If  $f$  is continuous at  $x_0$ , then  $\partial f(x_0) \neq \emptyset$ .

*Proof.* Suppose that  $f$  is continuous at  $x_0$ . The previous exercise tells us that there exists  $\phi \in X^*$  such that  $\phi \leq d^+f(x_0)$ . Let  $x \in A$ . A previous exercise implies that

$$\begin{aligned} \phi(x - x_0) &\leq d^+f(x_0)(x - x_0) \\ &\leq f(x) - f(x_0) \end{aligned}$$

Then  $f(x_0) + \phi(x - x_0) \leq f(x)$ .  $\square$

**Exercise 9.2.12.** Let  $X$  be a Banach space,  $A \subset X$  open and convex,  $f : A \rightarrow \mathbb{R}$  convex,  $\phi \in X^*$  and  $x_0 \in A$ . Then

(1) for each  $x \in A$ ,

$$\phi(x - x_0) \leq f(x) - f(x_0)$$

iff

$$\phi \leq d^+f(x_0)$$

(2)  $\partial f(x_0) = \{\phi \in X^* : \phi \leq d^+f(x_0)\}$

*Proof.*

(1) Suppose that for each  $x \in A$ ,  $\phi(x - x_0) \leq f(x) - f(x_0)$ . Let  $x \in X$ . Define  $t_0$  as before. Then for each  $t \in (0, t_0)$ ,

$$\begin{aligned} t\phi(x) &= \phi((x_0 + tx) - x_0) \\ &\leq f(x_0 + tx) - f(x_0) \end{aligned}$$

This implies that  $\phi(x) \leq d^+f(x_0)(x)$ .

Conversely, suppose that  $\phi \leq d^+f(x_0)$ . Let  $x \in A$ . A previous exercise implies that,

$$\begin{aligned} \phi(x - x_0) &\leq d^+f(x_0)(x - x_0) \\ &\leq f(x) - f(x_0) \end{aligned}$$

(2) Clear.  $\square$

**Exercise 9.2.13.** Let  $X$  be a Banach space,  $A \subset X$  open and convex,  $f : A \rightarrow \mathbb{R}$  convex and  $x_0 \in A$ . If  $f$  is continuous at  $x_0$ , then the following are equivalent:

(1)  $f$  is Gateaux differentiable at  $x_0$

- (2)  $d^+f(x_0)$  is linear
- (3)  $\#\partial f(x_0) = 1$

*Proof.* Suppose that  $f$  is continuous at  $x_0$ . Then  $d^+f(x_0)$  is Lipschitz and bounded.

- (1)  $\implies$  (2):

Suppose that  $f$  is Gateaux differentiable at  $x_0$ . Let  $x \in X$ . Then a previous exercise implies that

$$\begin{aligned} -df^+(x_0)(-x) &= df^-f(x_0)(x) \\ &= df^+f(x_0)(x) \end{aligned}$$

An exercise in the section on sublinear functionals implies that  $df^+f(x_0)$  is linear.

- (2)  $\implies$  (3):

Suppose that  $df^+f(x_0)$  is linear. Let  $\phi \in \partial f(x_0)$ . The previous exercise implies that  $\phi \leq df^+f(x_0)$ . Equivalence of linearity in the section on sublinear functionals implies that  $d^+f(x_0) = \phi$ .

- (3)  $\implies$  (1):

Suppose that  $\#\partial f(x_0) = 1$ . Since  $\partial f(x_0) = \{\phi \in X^* : \phi \leq d^+f(x_0)\}$ , equivalence of linearity in the section on sublinear functionals implies that  $d^+f(x_0)$  is linear. This implies that  $d^+f(x_0) = d^-f(x_0)$  and which implies that  $f$  is Gateaux differentiable at  $x_0$ . □

**Exercise 9.2.14.** Let  $X$  be a Banach space,  $A \subset X$  open and convex,  $f, g : A \rightarrow \mathbb{R}$  convex,  $\lambda \geq 0$  and  $x_0 \in A$ . Then

$$\partial f(x_0) + \lambda \partial g(x_0) \subset \partial[f + \lambda g](x_0)$$

*Proof.* Let  $\zeta \in \partial f(x_0) + \lambda \partial g(x_0)$ . Then there exist  $\phi \in \partial f(x_0)$  and  $\psi \in \partial g(x_0)$  such that  $\zeta = \phi + \lambda\psi$ . A previous exercise implies that  $\phi \leq d^+f(x_0)$  and  $\lambda\psi \leq \lambda d^+g(x_0) = d^+[\lambda g](x_0)$ . Hence

$$\begin{aligned} \zeta &= \phi + \lambda\psi \\ &\leq d^+f(x_0) + d^+[\lambda g](x_0) \\ &= d^+[f + \lambda g](x_0) \end{aligned}$$

So  $\zeta \in \partial[f + \lambda g](x_0)$  □

**Exercise 9.2.15.** Let  $X$  be a Banach space,  $A \subset X$  open and convex,  $f : A \rightarrow \mathbb{R}$  convex and  $x_0 \in A$ . If  $f$  is continuous at  $x_0$ , then  $f$  has a global minimum point at  $x_0$  iff  $0 \in \partial f(x_0)$ .

*Proof.* Suppose that  $f$  has a global minimum point at  $x_0$ . Let  $x \in X$ . Then

$$\begin{aligned} d^+f(x_0)(x) &= \lim_{t \rightarrow 0^+} \frac{f(x_0 + tx) - f(x_0)}{t} \\ &\geq 0 \end{aligned}$$

So  $0 \leq d^+f(x_0)$  and  $0 \in \partial f(x_0)$ .

Conversely, suppose that  $0 \in \partial f(x_0)$ . Let  $x \in A$ . Then

$$\begin{aligned} 0 &= 0(x - x_0) \\ &\leq f(x) - f(x_0) \end{aligned}$$

So that  $f(x_0) \leq f(x)$  which implies that  $f$  has a global minimum point at  $x_0$ .  $\square$

**Exercise 9.2.16.** Let  $X$  be a Banach space,  $A \subset X$  open and convex,  $f : A \rightarrow \mathbb{R}$  convex and  $x_0 \in A$ . If  $f$  is Frechet differentiable at  $x_0$ , then  $\partial f(x_0) = \{Df(x_0)\}$ .

*Proof.* Clear.  $\square$

**Exercise 9.2.17.** Let  $X$  be a Banach space,  $A \subset X$  open and convex,  $f : A \rightarrow \mathbb{R}$  convex and  $x_0 \in A$ . Suppose that  $f$  is Frechet differentiable at  $x_0$ . If  $Df(x_0) = 0$ , then  $f$  has a global minimum point at  $x_0$ .

*Proof.* Suppose that  $Df(x_0) = 0$ . Since  $\partial f(x_0) = \{Df(x_0)\}$ , a previous exercise implies that  $f$  has a global minimum point at  $x_0$ .  $\square$

**Exercise 9.2.18.** Let  $X$  be a Banach space,  $A \subset X$  open and convex,  $f : A \rightarrow \mathbb{R}$  convex and  $x_0 \in A$ . Suppose that  $f$  is Frechet differentiable at  $x_0$ . Then for each  $x \in A$ ,  $f(x) \geq f(x_0) + Df(x_0)(x - x_0)$

*Proof.* Since  $Df(x_0) \in \partial f(x_0)$ , for each  $x \in A$ ,  $Df(x_0)(x - x_0) \leq f(x) - f(x_0)$ .  $\square$

**Exercise 9.2.19.** Let  $X$  be a Banach space,  $A \subset X$  open and convex,  $f : A \rightarrow \mathbb{R}$ . Suppose that  $f$  is Frechet differentiable. Then  $f$  is convex iff for each  $x_0, x \in A$ ,  $f(x) \geq f(x_0) + Df(x_0)(x - x_0)$ .

*Proof.* Suppose that  $f$  is convex. Then the previous exercise implies that for each  $x_0, x \in A$ ,  $f(x) \geq f(x_0) + Df(x_0)(x - x_0)$ . Conversely, suppose that for each  $x_0, x \in A$ ,  $f(x) \geq f(x_0) + Df(x_0)(x - x_0)$ . Let  $x_0, x, y \in A$ . Then  $f(x) \geq f(x_0) + Df(x_0)(x - x_0)$  and  $f(y) \geq f(x_0) + Df(x_0)(y - x_0)$ .

**FINISH!!!**  $\square$

**Exercise 9.2.20.** Let  $X$  be a Banach space,  $A \subset X$  open and convex, and  $f \in C^2(A)$ . Then  $f$  is convex iff for each  $x_0 \in A$ ,  $D^2f(x_0)$  is positive semidefinite.

**Hint:** Define  $g : A \rightarrow \mathbb{R}$  by  $g(x) = f(x) - Df(x_0)(x - x_0)$  and show  $g$  is convex and use Taylor's Theorem

*Proof.* Suppose that  $f$  is convex. Let  $x_0 \in X$ . Define  $g : A \rightarrow \mathbb{R}$  by  $g(x) = f(x) - Df(x_0)(x - x_0)$ . Since  $g$  is the sum of a convex function and an affine function,  $g$  is convex. Since  $f \in C^2(A)$ , we have that  $g \in C^2(A)$  and it is straightforward to show that for each  $x \in A$ ,  $Dg(x) = Df(x) - Df(x_0)$  and  $D^2g(x) = D^2f(x)$ . In particular,  $Dg(x_0) = 0$ . Hence  $g$  has a global minimum point at  $x_0$ . This implies that  $D^2f(x_0)$  is positive semidefinite. Conversely, suppose that for each  $x_0 \in A$ ,  $D^2f(x_0)$  is positive semidefinite. Let

**FINISH!!!**  $\square$

### 9.3. Conjugacy.

**Definition 9.3.1.** Let  $X$  be a Banach space,  $A \subset X$  and  $f : A \rightarrow \mathbb{R}$ . Define

- (1)  $A^* \subset X^*$  and  $f^* : A^* \rightarrow \mathbb{R}$
- (2)  $A^{**} \subset X$  and  $f^{**} : A^{**} \rightarrow \mathbb{R}$

by

(1)

$$A^* = \left\{ \phi \in X^* : \sup_{x \in A} [\phi(x) - f(x)] < \infty \right\}$$

and

$$f^*(\phi) = \sup_{x \in A} [\phi(x) - f(x)]$$

(2)

$$A^{**} = \left\{ x \in X : \sup_{\phi \in A^*} [\hat{x}(\phi) - f^*(\phi)] < \infty \right\}$$

and

$$f^{**}(x) = \sup_{\phi \in A^*} [\hat{x}(\phi) - f^*(\phi)]$$

**Note 9.3.2.** If  $X$  is a Hilbert space, we may define  $A^* \subset X$  and  $f^* : A^* \rightarrow \mathbb{R}$  via the Riesz representation theorem by

$$A^* = \left\{ y \in X : \sup_{x \in A} [\langle y, x \rangle - f(x)] < \infty \right\}$$

and  $f^* : A^* \rightarrow \mathbb{R}$  and

$$f^*(y) = \sup_{x \in A} [\langle y, x \rangle - f(x)]$$

**Exercise 9.3.3.** Let  $X$  be a Banach space,  $A \subset X$  and  $f : A \rightarrow \mathbb{R}$ . Then

- (1)  $A^*$  is convex and  $f^* : A^* \rightarrow \mathbb{R}$  is convex and weak\* lower semicontinuous.
- (2)  $A^{**}$  is convex and  $f^{**} : A^{**} \rightarrow \mathbb{R}$  is convex and weakly lower semicontinuous.

*Proof.*

- (1) For  $x \in A$ , define  $g_x : X^* \rightarrow \mathbb{R}$  by  $g_x(\phi) = \hat{x}(\phi) - f(x)$ . Then for each  $x \in A$ ,  $g_x$  is convex and weak\* lower semicontinuous since it is affine and weak\* continuous. Exercise 9.1.18 implies that  $A^* = \{\phi \in X^* : \sup_{x \in A} g_x(\phi) < \infty\}$  is convex and  $f^* = \sup_{x \in A} g_x$  is convex.
- (2) For  $\phi \in A^*$ , define  $h_\phi : X \rightarrow \mathbb{R}$  by  $h_\phi(x) = \phi(x) - f^*(\phi)$ . Then for each  $\phi \in A^*$ ,  $h_\phi$  is convex and weakly lower semicontinuous since it is affine and weakly continuous. Exercise 9.1.18 implies that  $A^{**} = \{x \in X : \sup_{\phi \in A^*} h_\phi(x) < \infty\}$  is convex and  $f^{**} = \sup_{\phi \in A^*} h_\phi$  is convex.

□

**Exercise 9.3.4.** Let  $X$  be a Banach space,  $A \subset X$  and  $f : A \rightarrow \mathbb{R}$ . Then for each  $x \in A$  and  $\phi \in A^*$ ,  $f^*(\phi) \geq \phi(x) - f(x)$ .

*Proof.* Clear by definition.  $\square$

**Exercise 9.3.5.** Let  $X$  be a Banach space,  $A \subset X$  and  $f : A \rightarrow \mathbb{R}$ . Then  $A \subset A^{**}$ .

*Proof.* Let  $x \in A$ . Then the previous exercise implies that

$$\begin{aligned} \sup_{\phi \in A^*} [\phi(x) - f^*(\phi)] &\leq f(x) \\ &< \infty \end{aligned}$$

So  $x \in A^{**}$ .  $\square$

**Exercise 9.3.6.** Let  $X$  be a Banach space,  $A \subset X$  convex,  $f : A \rightarrow \mathbb{R}$  convex and lower semicontinuous and  $x_0 \in A$ .

- (1) if  $x_0 \in A$ , then for each  $\epsilon > 0$ , there exists  $\phi \in A^*$  such that for each  $x \in A$ ,  $f(x) > f(x_0) + \phi(x - x_0) - \epsilon$
- (2) if  $x_0 \notin A$ , then for each  $M \in \mathbb{R}$ , there exists  $\phi \in A^*$  such that for each  $x \in A$ ,  $f(x) > M + \phi(x - x_0)$

**Hint:** Apply second Hahn-Banach separation theorem to  $\{(x_0, f(x_0) - \epsilon)\}$  and  $\text{epi } f$ .

*Proof.*

- (1) Suppose that  $x_0 \in A$ . Let  $\epsilon > 0$ . Since  $f$  is convex and lower semicontinuous,  $\text{epi } f \subset X \times \mathbb{R}$  is convex and closed,  $\{(x_0, f(x_0) - \epsilon)\} \subset X \times \mathbb{R}$  is convex and compact and  $\{(x_0, f(x_0) - \epsilon)\} \cap \text{epi } f = \emptyset$ . Thus, there exists  $\lambda \in \mathbb{R}$ ,  $\psi \in X^*$  and  $k \in \mathbb{R}$  such that for each  $x \in A$  and  $r \geq f(x)$ ,

$$\psi(x) + \lambda r < k < \psi(x_0) + \lambda(f(x_0) - \epsilon)$$

Taking  $(x, r) = (x_0, f(x_0))$  implies that  $0 < -\lambda\epsilon$  and hence that  $\lambda < 0$ . Set  $\phi = |\lambda|^{-1}\psi$ . For  $x \in A$ , set  $r = f(x)$ . Then

$$\begin{aligned} \psi(x) - |\lambda|f(x) &< \psi(x_0) - |\lambda|(f(x_0) - \epsilon) \\ \iff |\lambda|^{-1}\psi(x) - f(x) &< |\lambda|^{-1}\psi(x_0) - (f(x_0) - \epsilon) \\ \iff \phi(x) - f(x) &< \phi(x_0) - (f(x_0) - \epsilon) \\ \iff f(x) &> f(x_0) + \phi(x - x_0) - \epsilon \end{aligned}$$

Since for each  $x \in A$ ,  $\phi(x) - f(x) < \phi(x_0) - f(x_0) + \epsilon$ , we have that

$$\begin{aligned} \sup_{a \in A} [\phi(x) - f(x)] &\leq \phi(x_0) - f(x_0) + \epsilon \\ &< \infty \end{aligned}$$

So  $\phi \in A^*$ .

- (2) Suppose that  $x_0 \notin A$ . Let  $M \in \mathbb{R}$ . Repeat the previous argument for  $(x_0, M)$  and  $\text{epi } f$ .  $\square$

**Exercise 9.3.7.** Let  $X$  be a Banach space,  $A \subset X$  convex and  $f : A \rightarrow \mathbb{R}$  convex and lower semicontinuous. Then

- (1)  $A = A^{**}$
- (2)  $f = f^{**}$

*Proof.*

- (1) A previous exercise implies that  $A \subset A^{**}$ . Let  $x_0 \in X$ . Suppose that  $x_0 \notin A$ . Let  $M \in \mathbb{R}$ . The previous exercise implies that there exists  $\phi_0 \in A^*$  such that for each  $x \in A$ ,  $f(x) > M + \phi_0(x - x_0)$ . Then

$$\begin{aligned}\phi_0(x_0) - f^*(\phi_0) &= \phi_0(x_0) - \sup_{x \in A} [\phi_0(x) - f(x)] \\ &= \phi_0(x_0) + \inf_{x \in A} [f(x) - \phi_0(x)] \\ &\geq \phi_0(x_0) + (M - \phi_0(x_0)) \\ &= M\end{aligned}$$

Therefore

$$\begin{aligned}\sup_{\phi \in A^*} [\phi(x_0) - f^*(\phi)] &\geq \phi_0(x_0) - f^*(\phi_0) \\ &\geq M\end{aligned}$$

Since  $M \in \mathbb{R}$  is arbitrary,

$$\sup_{\phi \in A^*} [\phi(x_0) - f^*(\phi)] = \infty$$

and  $x_0 \notin A^{**}$ . So  $A^c \subset (A^{**})^c$ , which implies that  $A^{**} \subset A$ . Thus  $A^{**} = A$ .

- (2) Part (1) and a previous exercise imply that  $f^{**} \leq f$ . Suppose that  $f \not\leq f^{**}$ . Then there exists  $x_0 \in A$  such that  $f(x_0) > f^{**}(x_0)$ . Choose  $\epsilon > 0$  such that  $f(x_0) > f^{**}(x_0) + 2\epsilon$ . A previous exercise implies that there exists  $\phi \in A^*$  such that for each  $x \in A$ ,  $f(x) > f(x_0) + \phi(x - x_0) - \epsilon$ . Choose  $a \in A$  such that  $f^*(\phi) - \epsilon < \phi(a) - f(a)$ . Then

$$\begin{aligned}f(x_0) &> f^{**}(x_0) + 2\epsilon \\ &\geq \phi(x_0) - f^*(\phi) + 2\epsilon \\ &> \phi(x_0 - a) + f(a) + \epsilon \\ &> \phi(x_0 - a) + f(x_0) + \phi(a - x_0) - \epsilon + \epsilon \\ &= f(x_0)\end{aligned}$$

which is a contradiction. So  $f \leq f^{**}$  and hence  $f = f^{**}$ .

□

**Definition 9.3.8.** Let

**Definition 9.3.9.**  $\partial f$

**Exercise 9.3.10.**



## 10. TOPOLOGICAL GROUPS

## 10.1. Topological Groups.

**Note 10.1.1.** This section establishes some basic results about topological groups and gives examples of common topological groups in analysis, specifically automorphism groups of metric spaces.

**Definition 10.1.2.** Let  $G$  be a group and  $\mathcal{T}$  a topology on  $G$ . Then  $(G, \mathcal{T})$  is said to be a **topological group** if the maps

- (1)  $G \times G \rightarrow G$  given by  $(x, y) \mapsto xy$
- (2)  $G \rightarrow G$  given by  $x \mapsto x^{-1}$

are continuous.

**Note 10.1.3.** For the remainder of this chapter, measurability is in reference to  $(G, \mathcal{B}(\mathcal{T}))$ . That is, the measurable sets are the Borel sets.

**Definition 10.1.4.** Let  $G$  be a topological group. We define

$$\text{Homeo}(G) = \{\phi : G \rightarrow G : \phi \text{ is a homeomorphism}\}$$

**Note 10.1.5.** Let  $G$  be a topological group. Then  $\text{Homeo}(G)$  is a group.

**Definition 10.1.6.** Let  $G$  be a group. Define  $\iota : G \rightarrow G$  by  $\iota(x) = x^{-1}$ .

**Exercise 10.1.7.** Let  $G$  be a topological group. Then  $\iota \in \text{Homeo}(G)$ .

*Proof.* By assumption  $\iota$  is continuous. We know from basic group theory that  $\iota$  is a bijection with  $\iota^{-1} = \iota$ .  $\square$

**Definition 10.1.8.** Let  $G$  be a group and  $S \subset G$ , then  $S$  is said to be **symmetric** if  $\iota(S) = S$ , ( i.e.  $S^{-1} = S$ ).

**Definition 10.1.9.** Let  $G$  be a topological group and  $\phi : G \rightarrow G$ . Then  $\phi$  is said to be an **automorphism** of  $G$  if  $\phi$  is a homomorphism and a homeomorphism. We define

$$\text{Aut}(G) = \{\phi : G \rightarrow G : \phi \text{ is an automorphism}\}$$

**Exercise 10.1.10.** Let  $G$  be a topological group. Then  $\iota \in \text{Aut}(G)$  iff  $G$  is abelian.

*Proof.* Basic group theory tells us that  $\iota$  is a homomorphism iff  $G$  is abelian.  $\square$

**Definition 10.1.11.** Let  $G$  be a group and  $g \in G$ . Define  $l_g : G \rightarrow G$  and  $r_g : G \rightarrow G$  by  $l_g(x) = gx$  and  $r_g(x) = xg^{-1}$ .

**Exercise 10.1.12.** Let  $G$  be a topological group and  $g \in G$ . Then  $l_g, r_g \in \text{Homeo}(G)$ .

*Proof.* By assumption  $l_g$  and  $r_g$  are continuous. We know from basic group theory that  $l_g$  and  $r_g$  are bijections with  $l_g^{-1} = l_{g^{-1}}$  and  $r_g^{-1} = r_{g^{-1}}$  so  $l_g$  and  $r_g$  are homeomorphisms.  $\square$

**Exercise 10.1.13.** Let  $G$  be a topological group. Define  $\phi, \psi : G \rightarrow \text{Homeo}(G)$  by  $\phi(g) = l_g$  and  $\psi(g) = r_g$ . Then  $\phi, \psi$  are homomorphisms.

*Proof.* Let  $g_1, g_2 \in G$ . Then

$$l_{g_1} \circ l_{g_2}(x) = l_{g_1}(g_2x) = g_1g_2x = l_{g_1g_2}(x)$$

and

$$r_{g_1} \circ r_{g_2}(x) = r_{g_1}(xg_2^{-1}) = xg_2^{-1}g_1^{-1} = x(g_1g_2)^{-1} = r_{g_1g_2}(x)$$

$\square$

**Exercise 10.1.14.** Let  $G$  be a topological group. Then for each  $U \subset G$  and  $g \in G$ , if  $U$  is open, then  $gU$ ,  $Ug$  and  $U^{-1}$  are open.

*Proof.* Let  $U \subset G$  and  $g \in G$ . Suppose that  $U$  is open. Since  $l_g, r_g$  and  $\iota$  are homeomorphisms,  $l_g(U) = gU$ ,  $r_g(U) = Ug$  and  $\iota(U) = U^{-1}$  are open.  $\square$

**Definition 10.1.15.** Let  $G$  be a topological group,  $y \in G$  and  $f \in L^0$ . Define  $L_y, R_y : L^0(G) \rightarrow L^0(G)$  by  $L_y f = f \circ l_y^{-1}$  and  $R_y f = f \circ r_y^{-1}$ , that is,  $L_y f(x) = f(y^{-1}x)$  and  $R_y f(x) = f(xy)$ .

**Exercise 10.1.16.** Let  $G$  be a topological group and  $y \in G$ . Then  $L_y, R_y \in \text{Sym}(L^0(G))$ .

*Proof.* It is straight forward to show that  $L_y^{-1} = L_{y^{-1}}$  and  $R_y^{-1} = R_{y^{-1}}$ .  $\square$

**Exercise 10.1.17.** Let  $G$  be a topological group. Define  $\phi, \psi : G \rightarrow \text{Sym}(L^0(G))$  by  $\phi(y) = L_y$  and  $\psi(y) = R_y$ . Then  $\phi$  and  $\psi$  are homomorphisms.

*Proof.* Let  $y, z \in G$  and  $f \in L^0(G)$ . Then

$$\begin{aligned} L_y \circ L_z(f) &= L_y(L_z(f)) \\ &= L_y(f \circ l_z^{-1}) \\ &= (f \circ l_z^{-1}) \circ l_y^{-1} \\ &= f \circ (l_z^{-1} \circ l_y^{-1}) \\ &= f \circ (l_y \circ l_z)^{-1} \\ &= f \circ l_{yz}^{-1} \\ &= L_{yz}(f) \end{aligned}$$

The case is similar for  $R_y$  and  $R_z$ .  $\square$

**Exercise 10.1.18.** Let  $G$  be a topological group,  $U \in \mathcal{B}(G)$  and  $y \in G$ . Then  $L_y \chi_U = \chi_{yU}$  and  $R_y \chi_U = \chi_{Uy^{-1}}$ .

*Proof.* Let  $x \in G$ . Then

$$\begin{aligned} L_y \chi_U(x) = 1 &\iff y^{-1}x \in U \\ &\iff x \in yU \\ &\iff \chi_{yU}(x) = 1 \end{aligned}$$

The case is similar for  $R_y$   $\square$

**Exercise 10.1.19.** Let  $G$  be a topological group,  $y \in G$  and  $f \in L^0(G)$ . Then  $\text{supp}(L_y f) = y \text{supp}(f)$  and  $\text{supp}(R_y f) = \text{supp}(f)y^{-1}$

*Proof.* Put  $A = \{x \in G : L_y f(x) \neq 0\}$  and  $B = \{x \in G : f(x) \neq 0\}$ . Then

$$\begin{aligned} x \in A &\iff L_y f(x) \neq 0 \\ &\iff f(y^{-1}x) \neq 0 \\ &\iff y^{-1}x \in B \\ &\iff x \in yB \end{aligned}$$

Thus  $A = yB$  which implies that  $\text{cl } A = y \text{cl } B$ . Therefore  $\text{supp}(L_y f) = y \text{supp}(f)$ .  $\square$

**Exercise 10.1.20.** Let  $G$  be a topological group and  $y \in G$ . Then  $L_y, R_y$  are linear and if we restrict to the bounded measurable functions, then  $L_y, R_y \in L(B(G))$  and  $\|L_y\|, \|R_y\| = 1$ .

*Proof.* Let  $f, g \in L^0(G)$  and  $\lambda \in \mathbb{C}$ . Then

$$\begin{aligned} L_y(\lambda f + g)(x) &= (\lambda f + g)(y^{-1}x) \\ &= \lambda f(y^{-1}x) + g(y^{-1}x) \\ &= \lambda L_y f(x) + L_y g(x) \end{aligned}$$

So  $L_y$  is linear. Next, we restrict to  $B(G) \cap L^0$ . We note that

$$\{|f(y^{-1}x)| : x \in y \operatorname{supp}(f)\} = \{|f(x)| : x \in \operatorname{supp}(f)\}$$

This implies that

$$\begin{aligned} \|L_y f\|_u &= \sup_{x \in \operatorname{supp}(L_y f)} |L_y f(x)| \\ &= \sup_{x \in y \operatorname{supp}(f)} |f(y^{-1}x)| \\ &= \sup_{x \in \operatorname{supp}(f)} |f(x)| \\ &= \|f\|_u \end{aligned}$$

So  $L_y$  is bounded. Hence  $L_y \in L(L^0)$ . The case is similar for  $R_y$ . □

**Definition 10.1.21.** Let  $G$  be a topological group. We say that  $G$  is a **locally compact group** if  $G$  is locally compact and Hausdorff.

## 10.2. Automorphism Groups of Metric Spaces.

**Definition 10.2.1.** Let  $(X, \tau)$  be a topological space. Define

$$\text{Aut}(X) = \{\sigma : X \rightarrow X : \sigma \text{ is a homeomorphism}\}$$

**Exercise 10.2.2.** Let  $(X, d)$  be a compact metric space. Then  $(\text{Aut}(X), d_u)$  is a topological group.

*Proof.* Let  $(\sigma_n)_{n \in \mathbb{N}}, (\tau_n)_{n \in \mathbb{N}} \subset \text{Aut}(X)$  and  $\sigma, \tau \in \text{Aut}(X)$ . Suppose that  $\sigma_n \xrightarrow{u} \sigma$  and  $\tau_n \xrightarrow{u} \tau$ .

- (1) Let  $\epsilon > 0$ . Since  $X$  is compact and  $\sigma$  is continuous,  $\sigma$  is uniformly continuous. Then there exists  $\delta > 0$  such that for each  $x, y \in X$ ,  $d(x, y) < \delta$  implies that  $d(\sigma(x), \sigma(y)) \leq \epsilon/2$ . Choose  $N_\sigma \in \mathbb{N}$  such that for each  $n \in \mathbb{N}$ ,  $n \geq N_\sigma$  implies that  $d_u(\sigma_n, \sigma) < \epsilon/2$ . Choose  $N_\tau \in \mathbb{N}$  such that for each  $n \in \mathbb{N}$ ,  $n \geq N_\tau$  implies that  $d_u(\tau_n, \tau) < \delta$ . Put  $N = \max(N_\sigma, N_\tau)$ . Let  $n \in \mathbb{N}$  and  $x \in X$ . Suppose that  $n \geq N$ . Then

$$\begin{aligned} d(\sigma_n \circ \tau_n(x), \sigma \circ \tau(x)) &\leq d(\sigma_n(\tau_n(x)), \sigma(\tau_n(x))) + d(\sigma(\tau_n(x)), \sigma(\tau(x))) \\ &< \epsilon/2 + \epsilon/2 \\ &= \epsilon \end{aligned}$$

So  $d_u(\sigma_n \circ \tau_n, \sigma \circ \tau) \leq \epsilon$  and  $\circ : \text{Aut}(X)^2 \rightarrow \text{Aut}(X)$  is continuous.

- (2) Suppose that  $\sigma = \text{id}_X$ . Let  $\epsilon > 0$ . Then there exists  $N \in \mathbb{N}$  such that for each  $n \in \mathbb{N}$ ,  $n \geq N$  implies that  $d_u(\sigma_n, \text{id}_X) < \epsilon$ . Let  $n \in \mathbb{N}$ . Suppose that  $n \geq N$ . Then

$$\begin{aligned} \sup_{x \in X} d(\sigma_n^{-1}(x), x) &= \sup_{x \in \sigma_n(X)} d(\sigma_n^{-1}(x), x) \\ &= \sup_{x \in X} d(\sigma_n^{-1}(\sigma_n(x)), \sigma_n(x)) \\ &= \sup_{x \in X} d(x, \sigma_n(x)) \\ &< \epsilon \end{aligned}$$

So  $\sigma_n^{-1} \xrightarrow{u} \text{id}_X$ . Now suppose that  $\sigma \neq \text{id}_X$ . Since  $\sigma_n \xrightarrow{u} \sigma$ , part (1) implies that  $\sigma^{-1} \circ \sigma_n \xrightarrow{u} \text{id}_X$ . Applying the result from above, we get that  $\sigma_n^{-1} \circ \sigma \xrightarrow{u} \text{id}_X$ . Applying part (1) again implies that  $\sigma_n^{-1} \xrightarrow{u} \sigma^{-1}$ . So the map  $\sigma \mapsto \sigma^{-1}$  is continuous.

Hence  $\text{Aut}(X)$  is a topological group.  $\square$

**Definition 10.2.3.** Let  $(X, d)$  be a metric space. Define

$$\text{Aut}(X, d) = \{\sigma : X \rightarrow X : \sigma \text{ is an isometric isomorphism}\}$$

**Exercise 10.2.4.** Let  $(X, d)$  be a compact metric space. Then  $(\text{Aut}(X, d), d_u)$  is a compact subgroup of  $(\text{Aut}(X), d_u)$ .

*Proof.* Clearly,  $(\text{Aut}(X, d), d_u)$  is a topological subgroup. To show compactness, use the Arzela Ascoli theorem.  $\square$

**Definition 10.2.5.** Let  $(X, \tau)$  be a topological space and  $\mu : \mathcal{B}(X) \rightarrow \mathbb{R}$  a Borel measure. Define

$$\text{Aut}(X, \mu) = \{\sigma \in \text{Aut}(X) : \sigma_*\mu = \mu\}$$

**Exercise 10.2.6.** Let  $(X, d)$  be a compact metric space and  $\mu : \mathcal{B}(X) \rightarrow \mathbb{R}$  an outer-regular Borel measure. Then  $\text{Aut}(X, \mu)$  is a closed subgroup of  $\text{Aut}(X)$ .

*Proof.* It is clear that  $\text{Aut}(X, \mu)$  is a subgroup of  $\text{Aut}(X)$ . Let  $(\sigma_n)_{n \in \mathbb{N}} \subset \text{Aut}(X, \mathcal{B}(X), \mu)$  and  $\sigma \in \text{Aut}(X)$ . Suppose that  $\sigma_n \xrightarrow{u} \sigma$ . Let  $E \subset X$  be closed,  $U \subset X$  open and suppose that  $E \subset U$ . An exercise in the section on metric spaces tells us that there exists  $N \in \mathbb{N}$  such that for each  $n \in \mathbb{N}$ ,  $n \geq N$  implies that  $\sigma(E) \subset \sigma_n(U)$ . Then

$$\begin{aligned} \mu(\sigma(E)) &\leq \mu(\sigma_N(U)) \\ &= \mu(U) \end{aligned}$$

Therefore, since  $\mu$  is outer regular,  $\mu(\sigma(E)) \leq \mu(E)$ . Since  $\sigma_n^{-1} \xrightarrow{u} \sigma^{-1}$ , we may apply the above argument to obtain that

$$\begin{aligned} \mu(E) &= \mu(\sigma^{-1}(\sigma(E))) \\ &\leq \mu(\sigma(E)) \end{aligned}$$

Hence  $\mu(E) = \mu(\sigma(E))$ . Applying the whole argument above thus far to  $\sigma^{-1}$ , we see that  $\mu(E) = \mu(\sigma^{-1}(E))$ . Since  $E \subset X$  is an arbitrary closed set and  $\mathcal{B}(X) = \sigma(E \subset X : E \text{ is closed})$ , we have that  $\mu = \sigma_*\mu$ . Thus  $\sigma \in \text{Aut}(X, \mu)$  which implies that  $\text{Aut}(X, \mu)$  is closed.  $\square$

**Definition 10.2.7.** Let  $(X, d)$  be a compact metric space and  $\mu : \mathcal{B}(X) \rightarrow \mathbb{R}$  an outer-regular Borel measure. Define  $\text{Aut}(X, d, \mu) = \text{Aut}(X, d) \cap \text{Aut}(X, \mu)$ .

**Exercise 10.2.8.** Let  $(X, d)$  be a compact metric space and  $\mu : \mathcal{B}(X) \rightarrow \mathbb{R}$  an outer-regular Borel measure. Then  $\text{Aut}(X, d, \mu)$  is compact.

*Proof.* Since  $\text{Aut}(X, d)$  is compact and  $\text{Aut}(X, \mu)$  is closed,  $\text{Aut}(X, d, \mu)$  is compact.  $\square$

## 11. GROUP ACTIONS ON METRIC SPACES

## 11.1. Introduction.

**Note 11.1.1.** For a set  $X$ , a group  $G$  and a (left) group action  $\phi : G \times X \rightarrow X$ , we will write  $\phi(g, x)$  as  $g \cdot x$ . We denote the projection map by  $\pi : X \rightarrow X/G$ .

**Definition 11.1.2.** Let  $X$  be a set,  $G$  a group,  $\phi : G \times X \rightarrow X$  a group action and  $g \in G$ . Define  $l_g : X \rightarrow X$  by

$$l_g(x) = g \cdot x$$

**Definition 11.1.3.** Let  $X$  be a topological space,  $G$  a group and  $\phi : G \times X \rightarrow X$  a group action. Then  $\phi$  is said to be  $X$ -continuous if for each  $g \in G$ ,  $l_g$  is continuous.

**Exercise 11.1.4.** Let  $X$  be a topological space,  $G$  a group and  $\phi : G \times X \rightarrow X$  an  $X$ -continuous group action. Then for each  $g \in G$ ,  $l_g \in \text{Homeo}(X)$ .

*Proof.* Let  $g \in G$ , then  $l_g$  and  $l_g^{-1} = l_{g^{-1}}$  are continuous, so  $l_g \in \text{Homeo}(X)$ . □

**Definition 11.1.5.** Let  $(X, d)$  be a metric space,  $G$  a group, and  $\phi : G \times X \rightarrow X$  a group action. Then  $\phi$  is said to be an **isometric group action** if for each  $g \in G$ ,  $l_g : X \rightarrow X$  is an isometry.

**Exercise 11.1.6.** Let  $(X, d)$  be a metric space,  $G$  a group, and  $\phi : G \times X \rightarrow X$  an isometric group action. Then  $\phi$  is  $X$ -continuous.

*Proof.* Clear since isometries are continuous. □

**Definition 11.1.7.** Let  $X$  be a set,  $G$  a group and  $\phi : G \times X \rightarrow X$  an  $X$ -continuous group action. Let  $g \in G$ . Define  $L_g : \mathbb{C}^X \rightarrow \mathbb{C}^X$  by

$$\begin{aligned} L_g(f)(x) &= f \circ l_g^{-1} \\ &= f \circ l_{g^{-1}} \end{aligned}$$

**Definition 11.1.8.** Let  $X$  be a set,  $G$  a group,  $\phi : G \times X \rightarrow X$  a group action and  $f : X \rightarrow \mathbb{C}$ . Then  $f$  is said to be  **$G$ -invariant** if for each  $g \in G$ ,  $L_g f = f$ .

**Exercise 11.1.9.** Let  $X$  be a set,  $G$  a group,  $\phi : G \times X \rightarrow X$  a group action and  $f : X \rightarrow \mathbb{C}$ . Then  $f$  is  $G$ -invariant iff for each  $g \in G$   $x \in X$ ,  $f(g \cdot x) = f(x)$ .

*Proof.* Clear. □

**Definition 11.1.10.** Let  $X$  be a set,  $G$  a group,  $\phi : G \times X \rightarrow X$  a group action and  $f : X \rightarrow \mathbb{C}$ . Suppose that  $f$  is  $G$ -invariant. Define  $\bar{f} : X/G \rightarrow \mathbb{C}$  by  $\bar{f}(\bar{x}) = f(x)$ .

**Exercise 11.1.11.** Let  $X$  be a set,  $G$  a group,  $\phi : G \times X \rightarrow X$  a group action and  $f : X \rightarrow \mathbb{C}$ . Suppose that  $f$  is  $G$ -invariant. Then  $f = \bar{f} \circ \pi$ .

*Proof.* Clear. □

## 11.2. Induced Metrics on Orbit Spaces.

**Note 11.2.1.** This section establishes the criteria for the existence of a metric on the orbit space of a metric space under a group action.

**Definition 11.2.2.** Let  $(X, d)$  be a metric space,  $G$  a group, and  $\phi : G \times X \rightarrow X$  a group action. We define  $\bar{d} : X/G \times X/G \rightarrow [0, \infty)$  by

$$\bar{d}(\bar{x}, \bar{y}) = \inf_{\substack{a \in \bar{x} \\ b \in \bar{y}}} d(a, b)$$

**Exercise 11.2.3.** Let  $(X, d)$  be a metric space,  $G$  a group, and  $\phi : G \times X \rightarrow X$  an isometric group action. Then for each  $x, y \in X$ ,

$$\bar{d}(\bar{x}, \bar{y}) = \inf_{g \in G} d(g \cdot x, y)$$

*Proof.* Let  $x, y \in X$ ,  $a \in \bar{x}$  and  $b \in \bar{y}$ . Then there exists there exists  $g_a, g_b \in G$  such that  $a = g_a \cdot x$  and  $b = g_b \cdot y$ . Set  $g = g_b^{-1} g_a$ . Since the map  $z \mapsto g_b^{-1} \cdot z$  is an isometry,

$$\begin{aligned} d(a, b) &= d(g_a \cdot x, g_b \cdot y) \\ &= d(g_b^{-1} g_a \cdot x, y) \\ &= d(g \cdot x, y) \end{aligned}$$

Let  $\epsilon > 0$ . Then there exist  $a^* \in \bar{x}$  and  $b^* \in \bar{y}$  such that  $d(a^*, b^*) < \bar{d}(\bar{x}, \bar{y}) + \epsilon$ . The above argument implies that there exists  $g^* \in G$  such that

$$\begin{aligned} \inf_{g \in G} d(g \cdot x, y) &\leq d(g^* \cdot x, y) \\ &= d(a^*, b^*) \\ &< \bar{d}(\bar{x}, \bar{y}) + \epsilon \end{aligned}$$

Since  $\epsilon > 0$  is arbitrary,

$$\inf_{g \in G} d(g \cdot x, y) \leq \bar{d}(\bar{x}, \bar{y})$$

Conversely, since  $\{(g \cdot x, y) : g \in G\} \subset \{(a, b) : a \in \bar{x}, b \in \bar{y}\}$ , we have that

$$\inf_{g \in G} d(g \cdot x, y) \geq \bar{d}(\bar{x}, \bar{y})$$

□

**Exercise 11.2.4.** Let  $(X, d)$  be a metric space,  $G$  a group, and  $\phi : G \times X \rightarrow X$  an isometric group action. Then for each  $x, y, z \in X$ ,

$$\bar{d}(\bar{x}, \bar{y}) \leq \bar{d}(\bar{x}, \bar{z}) + \bar{d}(\bar{z}, \bar{y})$$

*Proof.* Let  $x, y, z \in X$ . An exercise in section (2.1) implies that  $d(\bar{x}, \bar{y}) \leq d(\bar{x}, \bar{z}) + d(\bar{z}, \bar{y})$ . The previous exercise implies that

$$\begin{aligned} d(\bar{x}, \bar{z}) &= \inf_{a \in \bar{x}} d(a, z) \\ &= \inf_{g \in G} d(g \cdot x, z) \\ &= \bar{d}(\bar{x}, \bar{z}) \end{aligned}$$

Similarly,  $d(z, \bar{y}) = \bar{d}(\bar{z}, \bar{y})$ . Then

$$\begin{aligned} d(\bar{x}, \bar{y}) &\leq d(\bar{x}, z) + d(z, \bar{y}) \\ &= \bar{d}(\bar{x}, \bar{z}) + \bar{d}(\bar{z}, \bar{y}) \end{aligned}$$

□

**Exercise 11.2.5.** Let  $(X, d)$  be a metric space,  $G$  a group, and  $\phi : G \times X \rightarrow X$  an isometric group action. If for each  $x \in X$ ,  $\bar{x}$  is closed, then for each  $x, y \in X$ ,  $\bar{d}(\bar{x}, \bar{y}) = 0$  implies that  $\bar{x} = \bar{y}$ .

*Proof.* Suppose that for each  $x \in X$ ,  $\bar{x}$  is closed. Let  $x, y \in X$ . Suppose that  $\bar{d}(\bar{x}, \bar{y}) = 0$ . Then  $\inf_{g \in G} d(g \cdot x, y) = 0$ . Hence there exists  $(g_n)_{n \in \mathbb{N}} \subset G$  such that  $g_n \cdot x \rightarrow y$ . Since  $(g_n \cdot x)_{n \in \mathbb{N}} \subset \bar{x}$  and  $\bar{x}$  is closed,  $y \in \bar{x}$ . Thus  $\bar{x} = \bar{y}$ . □

**Exercise 11.2.6.** Let  $(X, d)$  be a metric space,  $G$  a group, and  $\phi : G \times X \rightarrow X$  an isometric group action. If for each  $x \in X$ ,  $\bar{x}$  is closed, then  $\bar{d}$  is a metric on  $X/G$ .

*Proof.* Clear by preceding exercises. □

**Exercise 11.2.7.** Let  $(X, d)$  be a metric space,  $(G, \tau)$  a topological group, and  $\phi : G \times X \rightarrow X$  an isometric group action. Suppose that  $G$  is compact and for each  $x \in X$ , the map  $g \mapsto g \cdot x$  is continuous. Then  $\bar{d}$  is a metric on  $X/G$ .

*Proof.* Let  $x \in X$ . Since  $G$  is compact and the map  $g \mapsto g \cdot x$  is continuous,  $\bar{x} = G \cdot x$  is compact and therefore closed. The previous exercise implies that  $\bar{d}$  is a metric. □

**Exercise 11.2.8.** Let  $(X, d)$  be a metric space,  $G$  a group, and  $\phi : G \times X \rightarrow X$  an isometric group action. Suppose that  $\bar{d}$  is a metric on  $X/G$ . Then the projection map  $\pi : X \rightarrow X/G$  is Lipschitz and therefore continuous.

*Proof.* Let  $x, y \in X$ . Then

$$\begin{aligned} \bar{d}(\pi(x), \pi(y)) &= \bar{d}(\bar{x}, \bar{y}) \\ &= \inf_{g \in G} d(g \cdot x, y) \\ &\leq d(x, y) \end{aligned}$$

□

**Exercise 11.2.9.** Let  $(X, d)$  be a metric space,  $G$  a group, and  $\phi : G \times X \rightarrow X$  an isometric group action. Suppose that  $\bar{d}$  is a metric on  $X/G$ . Let  $(x_n)_{n \in \mathbb{N}} \subset X$  and  $x \in X$ . Then  $\bar{x}_n \xrightarrow{\bar{d}} \bar{x}$  iff there exists a sequence  $(g_n)_{n \in \mathbb{N}}$  such that  $g_n \cdot x_n \xrightarrow{d} x$ .

*Proof.* Suppose that  $\bar{x}_n \xrightarrow{\bar{d}} \bar{x}$ . For  $n \in \mathbb{N}$ , choose  $g_n \in G$  such that  $d(g_n \cdot x_n, x) < \bar{d}(\bar{x}_n, \bar{x}) + 2^{-n}$ . Then  $d(g_n \cdot x_n, x) \rightarrow 0$  and  $g_n \cdot x_n \xrightarrow{d} x$ .

Conversely, suppose that there exists a sequence  $(g_n)_{n \in \mathbb{N}}$  such that  $g_n \cdot x_n \xrightarrow{d} x$ . Since  $\pi : X \rightarrow X/G$  is continuous, we have that

$$\begin{aligned} g_n \cdot x_n \xrightarrow{d} x &\implies \pi(g_n \cdot x_n) \xrightarrow{\bar{d}} \pi(x) \\ &\implies \bar{x}_n \xrightarrow{\bar{d}} \bar{x} \end{aligned}$$

□



**Exercise 11.2.10.** Let  $X$  be a set,  $d_1, d_2 : X^2 \rightarrow [0, \infty)$  metrics,  $G$  a group and  $\phi : G \times X \rightarrow X$  an isometric group action. Suppose that  $d_1$  and  $d_2$  are topologically equivalent.

- (1) Then  $\bar{d}_1$  is a metric on  $X/G$  iff  $\bar{d}_2$  is a metric on  $X/G$
- (2) If  $\bar{d}_1$  and  $\bar{d}_2$  are metrics, then  $\bar{d}_1$  and  $\bar{d}_2$  are topologically equivalent.

*Proof.*

- (1) •  $\implies$  Suppose that  $\bar{d}_1$  is a metric. Let  $x, y \in X$ . Suppose that  $\bar{d}_2(\bar{x}, \bar{y}) = 0$ . Then there exist  $(g_n)_{n \in \mathbb{N}} \subset G$  such that  $d_2(g_n \cdot x, y) \rightarrow 0$ . Since  $d_1$  and  $d_2$  are topologically equivalent,  $d_1(g_n \cdot x, y) \rightarrow 0$ . Thus  $\bar{d}_1(\bar{x}, \bar{y}) = 0$ . Since  $\bar{d}_1$  is a metric,  $\bar{x} = \bar{y}$ . Hence  $\bar{d}_2$  is a metric.
- $\impliedby$  Similar.
- (2) Suppose that  $\bar{d}_1$  and  $\bar{d}_2$  are metrics. Let  $(\bar{x}_n)_{n \in \mathbb{N}} \subset X/G$  and  $\bar{x} \in X/G$ .
  - Suppose that  $\bar{x}_n \xrightarrow{\bar{d}_1} \bar{x}$ . Then there exists a sequence  $(g_n)_{n \in \mathbb{N}}$  such that  $g_n \cdot x_n \xrightarrow{d_1} x$ . Since  $d_1$  and  $d_2$  are topologically equivalent,  $g_n \cdot x_n \xrightarrow{d_2} x$ . This implies that  $\bar{x}_n \xrightarrow{\bar{d}_2} \bar{x}$ .
  - Suppose that  $\bar{x}_n \xrightarrow{\bar{d}_2} \bar{x}$ . Then similarly to above,  $\bar{x}_n \xrightarrow{\bar{d}_1} \bar{x}$ .

□

**Exercise 11.2.11.** Let  $X$  be a set,  $d_1, d_2 : X^2 \rightarrow [0, \infty)$  metrics on  $X$ ,  $G$  a group and  $\phi : G \times X \rightarrow X$  an isometric group action. Suppose that  $d_1$  and  $d_2$  are equivalent. If  $\bar{d}_1$  and  $\bar{d}_2$  are metrics, then  $\bar{d}_1$  and  $\bar{d}_2$  are equivalent.

*Proof.* Suppose that  $\bar{d}_1$  and  $\bar{d}_2$  are metrics. Since  $d_1, d_2$  are equivalent, there exist  $C_1, C_2 > 0$  such that for each  $x, y \in X$ ,  $C_1 d_1(x, y) \leq d_2(x, y) \leq C_2 d_1(x, y)$ . Let  $x, y \in X$ . Then

$$\begin{aligned} C_1 \bar{d}_1(\bar{x}, \bar{y}) &= C_1 \inf_{g \in G} d_1(g \cdot x, y) \\ &= \inf_{g \in G} C_1 d_1(g \cdot x, y) \\ &\leq \inf_{g \in G} d_2(g \cdot x, y) \\ &= \bar{d}_2(\bar{x}, \bar{y}) \end{aligned}$$

and

$$\begin{aligned} \bar{d}_2(\bar{x}, \bar{y}) &= \inf_{g \in G} d_2(g \cdot x, y) \\ &\leq \inf_{g \in G} C_2 d_1(g \cdot x, y) \\ &= C_2 \inf_{g \in G} d_1(g \cdot x, y) \\ &= C_2 \bar{d}_1(\bar{x}, \bar{y}) \end{aligned}$$

So that  $C_1 \bar{d}_1 \leq \bar{d}_2 \leq C_2 \bar{d}_1$

□

**Exercise 11.2.12.** Let  $(X, d)$  be a metric space,  $G$  a group and  $\phi : G \times X \rightarrow X$  an isometric group action. Suppose that  $\bar{d}$  is a metric. Then  $\pi : X \rightarrow X/G$  is a quotient map.

*Proof.*

- Clearly  $\pi$  is surjective.

- Let  $C \subset X/G$ . Suppose that  $C$  is closed. Since  $\pi$  is continuous, if  $\pi^{-1}(C)$  is closed. Conversely, suppose that  $\pi^{-1}(C)$  is closed. Let  $(\bar{x}_\alpha)_\alpha \subset C$  be a net and  $\bar{x} \in X/G$ . Suppose that  $\bar{x}_\alpha \rightarrow \bar{x}$ . Then there exists  $(g_\alpha)_{\alpha \in A} \subset G$  such that  $g_\alpha \cdot x_\alpha \rightarrow x$ . Since  $(g_\alpha \cdot x_\alpha)_{\alpha \in A} \subset \pi^{-1}(C)$ ,  $x \in \pi^{-1}(C)$ . Hence  $\bar{x} \in C$  and  $C$  is closed. Then Exercise 4.6.4 implies that  $\pi$  is a quotient map. □

**Exercise 11.2.13.** Let  $(X, d)$  be a metric space,  $G$  a group and  $\phi : G \times X \rightarrow X$  an isometric group action. Suppose that  $\bar{d}$  is a metric. Then  $\pi : X \rightarrow X/G$  is open.

*Proof.* Let  $U \subset X$ . Suppose that  $U$  is open. Then

$$\pi^{-1}(\pi(U)) = \bigcup_{g \in G} g \cdot U$$

Since for each  $g \in G$ ,  $l_g \in \text{Homeo}(X)$ , we have that for each  $g \in G$ ,  $g \cdot U$  is open. Therefore  $\bigcup_{g \in G} g \cdot U$  is open. Hence  $\pi^{-1}(\pi(U))$  is open. Then Exercise 4.6.6 implies that  $\pi$  is open. □

**Exercise 11.2.14.** Let  $(X, d)$  be a metric space,  $G$  a group and  $\phi : G \times X \rightarrow X$  an isometric group action. Suppose that  $\bar{d}$  is a metric. Then  $\bar{d}$  metrizes the quotient topology  $\pi_*\tau(d)$  on  $X/G$ .

*Proof.* Immediate by the previous exercise and Exercise 4.6.14. □

**Exercise 11.2.15.** Let  $(X, d)$  be a metric space,  $G$  a group, and  $\phi : G \times X \rightarrow X$  an isometric group action. Let  $f : X \rightarrow \mathbb{C}$ . Suppose that  $f$  is  $G$ -invariant and  $\bar{d}$  is a metric. If  $f \in C(X)$ , then  $\bar{f} \in C(X/G)$ .

**Hint:** Exercise 4.6.14

*Proof.* Suppose that  $f \in C(X)$ . Exercise 4.6.14 implies that  $\bar{f} : X \rightarrow \mathbb{C}$  is the unique map such that  $\bar{f} \circ \pi = f$  and  $\bar{f}$  is continuous. □

**Exercise 11.2.16.** Let  $(X, d)$  be a metric space,  $G$  a group, and  $\phi : G \times X \rightarrow X$  an isometric group action. Let  $f : X \rightarrow \mathbb{C}$ . Suppose that  $f$  is  $G$ -invariant and  $\bar{d}$  is a metric. If  $f \in C(X)$ , then  $\bar{f} \in C(X/G)$ .

**Hint:** Exercise 4.6.14

*Proof.* Suppose that  $f \in C(X)$ . Exercise 4.6.14 implies that  $\bar{f} : X \rightarrow \mathbb{C}$  is the unique map such that  $\bar{f} \circ \pi = f$  and  $\bar{f}$  is continuous. □

### 11.3. Fundamental Examples.

**Note 11.3.1.** This section uses results from the previous two sections to establish metrics on some fundamental orbit spaces of metric spaces under a group action.

**Exercise 11.3.2. Procrustes Distance:**

Consider the metric space  $(\mathbb{C}^{n \times d}, \|\cdot\|_F)$ , topological group  $(U_d, \|\cdot\|_F)$  and the (right) action  $\phi : X \times U_d \rightarrow X$  by  $X \cdot U = XU$ . Then

- (1)  $\phi$  is a continuous isometric group action
- (2)  $U_d$  is compact
- (3)  $\bar{d}$  is a metric on  $\mathbb{C}^{n \times d}/U_d$

*Proof.* Clear. □

**Exercise 11.3.3.** Let  $X$  be a compact metric space and  $\mu : \mathcal{B}(X) \rightarrow [0, \infty]$  a Borel measure. Define the (right) group action  $\phi : L^1(\mu) \times \text{Aut}(X, \mu) \rightarrow L^1(\mu)$  by

$$f \cdot \sigma = f \circ \sigma$$

Then  $\phi$  is an isometric group action.

*Proof.* Let  $\sigma \in \text{Aut}(X, \mu)$  and  $f \in L^1(\mu)$ . Then

$$\begin{aligned} \|f \cdot \sigma\|_1 &= \int_X |f \circ \sigma| d\mu \\ &= \int_X |f| \circ \sigma d\mu \\ &= \int_{\sigma(X)} |f| d\sigma_*\mu \\ &= \int_{\sigma(X)} |f| d\mu \\ &= \int_X |f| d\mu \\ &= \|f\|_1 \end{aligned}$$

□

**Exercise 11.3.4.** Let  $X$  be a compact metric space and  $\mu : \mathcal{B}(X) \rightarrow [0, \infty]$  a Radon measure. Define the (right) group action  $\phi : L^1(\mu) \times \text{Aut}(X, \mu) \rightarrow L^1(\mu)$  by

$$f \cdot \sigma = f \circ \sigma$$

Then for each  $f \in L^1(\mu)$ , the map  $\sigma \mapsto f \cdot \sigma$  is continuous.

*Proof.* Let  $f \in L^1(\mu)$ ,  $(\sigma_n)_{n \in \mathbb{N}} \subset \text{Aut}(X, \mu)$  and  $\sigma \in \text{Aut}(X, \mu)$ . Suppose that  $\sigma_n \xrightarrow{u} \sigma$ . Since  $\mu$  is Radon,  $C_c(X)$  is dense in  $L^1(\mu)$  and therefore, there exists  $\phi \in C_c(X)$  such that  $\|\phi - f\| < \epsilon/3$ . Since  $X$  is compact and  $\mu$  is Radon,  $\mu(X) < \infty$ . Since  $\phi$  is uniformly continuous, Exercise 3.1.22 implies that  $\phi \circ \sigma_n \xrightarrow{u} \phi \circ \sigma$ . So there exists  $N \in \mathbb{N}$  such that for each  $n \in \mathbb{N}$ ,  $n \geq N$  implies that  $\|\phi \circ \sigma_n - \phi \circ \sigma\|_u < \frac{\epsilon}{3(\mu(X)+1)}$ . Let  $n \in \mathbb{N}$ . Suppose that

$n \geq \mathbb{N}$ . Then

$$\begin{aligned}
 \|f \circ \sigma_n - f \circ \sigma\|_1 &\leq \|f \circ \sigma_n - \phi \circ \sigma_n\|_1 + \|\phi \circ \sigma_n - \phi \circ \sigma\|_1 + \|\phi \circ \sigma - f \circ \sigma\|_1 \\
 &= \|(f - \phi) \circ \sigma_n\|_1 + \|\phi \circ \sigma_n - \phi \circ \sigma\|_1 + \|(\phi - f) \circ \sigma\|_1 \\
 &= \|f - \phi\|_1 + \|\phi \circ \sigma_n - \phi \circ \sigma\|_1 + \|\phi - f\|_1 \\
 &= \|f - \phi\|_1 + \|\phi \circ \sigma_n - \phi \circ \sigma\|_{u\mu(X)} + \|\phi - f\|_1 \\
 &< \frac{\epsilon}{3} + \frac{\epsilon}{3} + \frac{\epsilon}{3} \\
 &= \epsilon
 \end{aligned}$$

So that  $f \circ \sigma_n \xrightarrow{u} f \circ \sigma$  which implies that the map  $\sigma \mapsto f \cdot \sigma$  is continuous.  $\square$

**Exercise 11.3.5. Cut Distance:**

Let  $X$  be a compact metric space and  $\mu : \mathcal{B}(X) \rightarrow [0, \infty]$  a Radon measure. Define the (right) group action  $\phi : L^1(\mu) \times \text{Aut}(X, \mu) \rightarrow L^1(\mu)$  by

$$f \cdot \sigma = f \circ \sigma$$

Then

- (1)  $\phi$  is an isometric group action
- (2)  $\text{Aut}(X, d, \mu)$  is compact
- (3) for each  $f \in L^1(\mu)$ , the map  $\sigma \mapsto f \cdot \sigma$  is continuous.
- (4)  $\bar{d}$  is a metric on  $L^1(\mu) / \text{Aut}(X, d, \mu)$

*Proof.* Clear by the preceding exercises.  $\square$

**Note 11.3.6.** The preceding distance is not quite the Cut distance, as the Cut norm only considers a subset of measurable sets for a function of two variables, but with some work, maybe I can show it is a distance.

## 12. APPENDIX

## 12.1. Summation.

**Definition 12.1.1.** Let  $f : X \rightarrow [0, \infty)$ , Then we define

$$\sum_{x \in X} f(x) := \sup_{\substack{F \subset X \\ F \text{ finite}}} \sum_{x \in F} f(x)$$

This definition coincides with the usual notion of summation when  $X$  is countable. For  $f : X \rightarrow \mathbb{C}$ , we can write  $f = g + ih$  where  $g, h : X \rightarrow \mathbb{R}$ . If

$$\sum_{x \in X} |f(x)| < \infty,$$

then the same is true for  $g^+, g^-, h^+, h^-$ . In this case, we may define

$$\sum_{x \in X} f(x)$$

in the obvious way.

The following note justifies the notation  $\sum_{x \in X} f(x)$  where  $f : X \rightarrow \mathbb{C}$ .

**Note 12.1.2.** Let  $f : X \rightarrow \mathbb{C}$  and  $\alpha : X \rightarrow X$  a bijection. If  $\sum_{x \in X} |f(x)| < \infty$ , then  $\sum_{x \in X} f(\alpha(x)) = \sum_{x \in X} f(x)$ .

## 12.2. Asymptotic Notation.

**Definition 12.2.1.** Let  $X$  be a topological space,  $Y, Z$  be normed vector spaces,  $f : X \rightarrow Y$ ,  $g : X \rightarrow Z$  and  $x_0 \in X \cup \{\infty\}$ . Then we write

$$f = o(g) \quad \text{as } x \rightarrow x_0$$

if for each  $\epsilon > 0$ , there exists  $U \in \mathcal{N}_{x_0}$  such that for each  $x \in U$ ,

$$\|f(x)\| \leq \epsilon \|g(x)\|$$

**Exercise 12.2.2.** Let  $X$  be a topological space,  $Y, Z$  be normed vector spaces,  $f : X \rightarrow Y$ ,  $g : X \rightarrow Z$  and  $x_0 \in X \cup \{\infty\}$ . If there exists  $U \in \mathcal{N}_{x_0}$  such that for each  $x \in U \setminus \{x_0\}$ ,  $g(x) > 0$ , then

$$f = o(g) \text{ as } x \rightarrow x_0 \quad \text{iff} \quad \lim_{x \rightarrow x_0} \frac{\|f(x)\|}{\|g(x)\|} = 0$$

**Exercise 12.2.3.** Let  $X$  and  $Y$  be normed vector spaces,  $A \subset X$  open and  $f : A \rightarrow Y$ . Suppose that  $0 \in A$ . If  $f(h) = o(\|h\|)$  as  $h \rightarrow 0$ , then for each  $h \in X$ ,  $f(th) = o(|t|)$  as  $t \rightarrow 0$ .

*Proof.* Suppose that  $f(h) = o(\|h\|)$  as  $h \rightarrow 0$ . Let  $h \in X$  and  $\epsilon > 0$ . Choose  $\delta' > 0$  such that for each  $h' \in B(0, \delta')$ ,  $h' \in A$  and

$$\|f(h')\| \leq \frac{\epsilon}{\|h\| + 1} \|h'\|$$

Choose  $\delta > 0$  such that for each  $t \in B(0, \delta)$ ,  $th \in B(0, \delta')$ . Let  $t \in B(0, \delta)$ . Then

$$\begin{aligned} \|f(th)\| &\leq \frac{\epsilon}{\|h\| + 1} |t| \|h\| \\ &< \epsilon |t| \end{aligned}$$

So  $f(th) = o(|t|)$  as  $t \rightarrow 0$ . □

**Definition 12.2.4.** Let  $X$  be a topological space,  $Y, Z$  be normed vector spaces,  $f : X \rightarrow Y$ ,  $g : X \rightarrow Z$  and  $x_0 \in X \cup \{\infty\}$ . Then we write

$$f = O(g) \quad \text{as } x \rightarrow x_0$$

if there exists  $U \in \mathcal{N}_{x_0}$  and  $M \geq 0$  such that for each  $x \in U$ ,

$$\|f(x)\| \leq M \|g(x)\|$$

## REFERENCES

- [1] [Introduction to Algebra](#)
- [2] [Introduction to Analysis](#)
- [3] [Introduction to Fourier Analysis](#)
- [4] [Introduction to Measure and Integration](#)