

Presentation

Carson James

May 24, 2022

Definition

We define

$\Lambda_+^{n \times r} = \{\Sigma \in \mathbb{R}^{n \times r} : \Sigma \text{ is diagonal and positive semi-definite}\}$ and
 $O_n = \{U \in \mathbb{R}^{n \times n} : U \text{ is orthogonal}\}.$

Model 1:

1. Fix $M \in \mathbb{R}^{n \times n_M}$ columns of M are orthogonal and set $P_M = M(M^T M)^{-1}M^T$?
2. Choose $\Sigma_Z \in \Lambda_+^{n_M \times r}$, $\Sigma_X \in \Lambda_+^{n \times p}$, $U_Z \in O_r$ and $U_X \in O_p$.
3. Set $V_Z^T = \Sigma_Z U_Z$ and $V_X^T = \Sigma_X U_X$.
4. Set $J_Z = M V_Z^T$ and $J_X = M V_X^T$.
5. Choose $I_Z \in \mathbb{R}^{n \times r}$, $I_X \in \mathbb{R}^{n \times p}$ such that $\mathcal{C}(I_Z) \cup \mathcal{C}(I_X) \subset \mathcal{C}(I - P_M)$.
6. Choose $E_X \in \mathbb{R}_{n \times p}$ with $(E_X)_{i,j} \sim N(0, \sigma^2)$
7. Set $Z = J_Z + I_Z$ and $X = J_X + I_X + E_X$

Then $\mathcal{C}(M) \perp \mathcal{C}(I_Z), \mathcal{C}(I_X)$.

Model 2: We consider a modification of the planted partition model which is a submodel of the stochastic block model with n nodes and r blocks (for now $r = 2$).

- Choose $U \in \mathbb{R}^{n \times 2}$ such that for each $i \in \{1, \dots, n\}$,

$$U_{i,j} = \begin{cases} 1 & \text{node } i \text{ is in block } j \\ 0 & \text{else} \end{cases}$$

as in the stochastic block model. We choose

$$U = \begin{pmatrix} 1 & 0 \\ \vdots & \vdots \\ 1 & 0 \\ 0 & 1 \\ \vdots & \vdots \\ 0 & 1 \end{pmatrix}$$

- ▶ Choose $W \in \mathbb{R}^{n \times 1}$, with $W = (1, -1, \dots, 1, -1)^T$ and $\mathcal{C}(W) \perp \mathcal{C}(U)$.
- ▶ Choose $0 < b < a < 1$ and set

$$Q = \begin{pmatrix} a & b \\ b & a \end{pmatrix}$$

to be the block probability matrix from the planted partition model.

- ▶ Set $B = Q^{1/2}$.
- ▶ Choose $\alpha \in (0, 1)$ and set $Z = (1 - \alpha)(0, W) + \alpha(UB, 0)$
- ▶ Set $X = (0, W)$

Then

- ▶ $U^T W = 0$
- ▶ $BB^T \in [0, 1]^{2 \times 2}$ and $(UB)(UB)^T \in [0, 1]^{n \times n}$
- ▶ if α is close enough to 1, then $Z \in [0, 1]^{n \times n}$
- ▶ here $J_Z = J_X = (0, W)$ and $I_Z = (UB, 0)$, $I_X = (0, 0)$.

Model 3: We consider another modification of the planted partition model which is a submodel of the stochastic block model with n nodes and r blocks (for now $r = 2$).

- Choose $U \in \mathbb{R}^{n \times 2}$ such that for each $i \in \{1, \dots, n\}$,

$$U_{i,j} = \begin{cases} 1 & \text{node } i \text{ is in block } j \\ 0 & \text{else} \end{cases}$$

as in the stochastic block model. We choose

$$U = \begin{pmatrix} 1 & 0 \\ \vdots & \vdots \\ 1 & 0 \\ 0 & 1 \\ \vdots & \vdots \\ 0 & 1 \end{pmatrix}$$

- ▶ Choose $W \in \mathbb{R}^{n \times 1}$, with $W = (1, -1, \dots, 1, -1)^T$ and $\mathcal{C}(W) \perp \mathcal{C}(U)$.
- ▶ Choose $0 < b < a < 1$ and set

$$Q = \begin{pmatrix} a & b \\ b & a \end{pmatrix}$$

to be the block probability matrix from the planted partition model.

- ▶ Set $B = Q^{1/2}$.
- ▶ Choose $K^T \in O(p)$ and $I_X \in \mathbb{R}^{n \times p}$ such that $\mathcal{C}(I_X) \perp \mathcal{C}(W)$.
- ▶ Choose $E_X \in \mathbb{R}^{n \times p}$ with $(E_X)_{i,j} \sim N(0, \sigma^2)$
- ▶ Define $J_Z = W \begin{pmatrix} \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \end{pmatrix}$, $I_Z = UB$, $J_X = (W \ 0) K^T$
- ▶ Choose $\alpha \in (0, 1)$ and set $Z = (1 - \alpha)J_Z + \alpha I_Z$
- ▶ Set $X = J_X + I_X + E_X$

Then

- ▶ $U^T W = 0$
- ▶ $BB^T \in [0, 1]^{2 \times 2}$ and $(UB)(UB)^T \in [0, 1]^{n \times n}$
- ▶ if α is close enough to 1, then $Z \in [0, 1]^{n \times n}$

Analysis of Initial Solution:

Consider the following model for the data:

$$A_{ij} \sim \text{Ber}(ZZ^T)_{ij}), \quad i > j, i, j \in [n],$$

$$X_{uv} = (W)_{uv} + \epsilon_{uv}, \quad u \in [n], v \in [p].$$

Assume $\epsilon_{uv} \stackrel{iid}{\sim} N(0, \sigma^2)$. Write

$$Z = [M, R_Z]\Gamma,$$

$$W = [M, R_W]U^T,$$

where $M \in \mathbb{R}^{n \times r_M}$, $R_Z \in \mathbb{R}^{n \times r_Z}$ and $R_W \in \mathbb{R}^{n \times r_W}$ are matrices with orthogonal columns, and $\Gamma \in \mathbb{R}^{(r_M+r_Z) \times (r_M+r_Z)}$ and $U \in \mathbb{R}^{p \times (r_M+r_W)}$ some other matrices.

Define $\hat{V}^{(1)}$ as the matrix of $(r_M + r_Z)$ leading eigenvectors of A , and $\hat{V}^{(2)}$ as the matrix of $(r_M + r_W)$ left leading singular vectors of X . Then define \hat{M} as the matrix of r_M left leading singular vectors of $[\hat{V}^{(1)}, \hat{V}^{(2)}]$. Set

$$\epsilon = \sqrt{\frac{1}{2} \left(\frac{\delta(ZZ^T)}{\lambda_{\min}^2(\Gamma\Gamma^T)} + \frac{n}{\lambda_{\min}^2(U^T U)} \right)}$$

Conjecture

With the assumptions as above, and some regularity conditions (TBD) (maybe if $\epsilon = o(1)$ as $n \rightarrow \infty$), there exists some orthogonal matrix O such that

$$\mathbb{E} \|\hat{M} - MO\|_F = O\left(\frac{r_M^{1/2}}{\sqrt{n}}\right).$$

Idea

Define \tilde{M} to be the matrix consisting of the r_M left leading singular vectors of $\tilde{\Pi}$. Then

$$\begin{aligned}\min_{W \in O_{r_M}} \|\hat{M} - MW\|_F &\leq \|\hat{M}\hat{M}^T - MM^T\|_F \\ &\leq \|\hat{M}\hat{M}^T - \tilde{M}\tilde{M}^T\|_F + \|\tilde{M}\tilde{M}^T - MM^T\|_F\end{aligned}$$

Then we control both of these errors.

To control the first error, we need the following lemma:

Lemma

Let $X \in \mathbb{R}^{n \times p}$ with $X_{i,j} \sim N(0, \sigma^2)$ and $a > 1$. Set

$C_{n,p} = \frac{a}{a-1} \frac{3}{2} \left[(\sqrt{n} + \sqrt{p}) + \frac{5}{\log(3/2)} \sqrt{\log(n \wedge p)} \right] \sigma$. Then for each $t \geq C_{n,p}$,

$$P(\|X\| \geq t) \leq \exp \left(- \frac{t^2}{2(a\sigma)^2} \right)$$

and for each $q \geq 1$,

$$\mathbb{E}(\|X\|^q) = O(\sigma^q (\sqrt{n} + \sqrt{p})^q)$$