

1

2

Introduction to Differential Geometry

Contents

Notation	vii
Preface	1
1 Review of Fundamentals	3
1.1 Set Theory	3
1.2 Calculus	4
1.2.1 Differentiation	4
1.2.2 Integration	5
1.3 Topology	6
2 Multilinear Algebra	7
2.1 (r, s) -Tensors	7
2.2 Covariant k -Tensors	10
3 Smooth Manifolds	19
3.1 Topological Manifolds	19
3.2 Smooth Manifolds	26
3.3 Smooth Maps	29
3.4 Partitions of Unity	33
3.5 The Tangent Space	34
3.6 The Cotangent Space	39
3.7 Maps of Full Rank	41
3.8 Submanifolds	42
4 Vector Bundles and Tensor Fields	45
4.1 The Vector Bundle	45
4.2 The cotangent Bundle	48
4.3 The (r, s) -Tensor Bundle	48
4.4 Vector Fields	49
4.5 1-Forms	50
4.6 (r, s) -Tensor Fields	51
4.7 Differential Forms	53
5 Extra	57
5.1 Integration of Differential Forms	59

Notation

$\mathcal{M}_+(X, \mathcal{A})$	finite measures on (X, \mathcal{A})
v	velocity

Preface

cc-by-nc-sa

Chapter 1

Review of Fundamentals

1.1 Set Theory

Definition 1.1.0.1. Let $\{A_i\}_{i \in I}$ be a collection of sets. The **disjoint union of** $\{A_i\}_{i \in I}$, denoted $\coprod_{i \in I} A_i$, is defined by

$$\coprod_{i \in I} A_i = \bigcup_{i \in I} \{i\} \times A_i$$

We define the **natural projection map**, denoted $\pi : \coprod_{i \in I} A_i \rightarrow I$, by $\pi(i, a) = i$.

Definition 1.1.0.2. Let E and M be sets, $\pi : E \rightarrow B$ a surjection and $\sigma : B \rightarrow E$. Then σ is said to be a section of (E, M, π) if $\pi \circ \sigma = \text{id}_M$.

Note 1.1.0.3. Let $\{A_i\}_{i \in I}$ be a collection of sets and $\sigma : I \rightarrow \coprod_{i \in I} A_i$. We will typically be interested in sections σ of $\left(\coprod_{i \in I} A_i, I, \pi \right)$.

Exercise 1.1.0.4. Let $\{A_i\}_{i \in I}$ be a collection of sets and $\sigma : I \rightarrow \coprod_{i \in I} A_i$. Then σ is a section of $\coprod_{i \in I} A_i$ iff for each $i \in I$, $\sigma(i) \in A_i$

Proof. Clear. □

1.2 Calculus

1.2.1 Differentiation

Definition 1.2.1.1. Let $n \geq 1$. For $i = 1, \dots, n$, define $x^i : \mathbb{R}^n \rightarrow \mathbb{R}$ by $x^i(a^1, \dots, a^n) = a^i$. The functions $(x^i)_{i=1}^n$ are called the **standard coordinate functions on \mathbb{R}^n** .

Definition 1.2.1.2. Let $U \subset \mathbb{R}^n$ be open, $f : U \rightarrow \mathbb{R}$ and $a \in U$. Then f is said to be **differentiable with respect to x^i at a** if

$$\lim_{h \rightarrow 0} \frac{f(a + he^i) - f(a)}{h}$$

exists. If f is differentiable with respect to x^i at a , we define the **partial derivative of f with respect to x^i at a** , denoted

$$\frac{\partial f}{\partial x^i}(a) \text{ or } \frac{\partial}{\partial x^i} f$$

to be the limit above.

Definition 1.2.1.3. Let $U \subset \mathbb{R}^n$ be open and $f : U \rightarrow \mathbb{R}$. Then f is said to be **differentiable with respect to x^i** if for each $a \in U$, f is differentiable with respect to x^i at a .

Exercise 1.2.1.4. Let $U \subset \mathbb{R}^n$ be open, $f : U \rightarrow \mathbb{R}$ and $a \in U$. Suppose that $\frac{\partial^2 f}{\partial x^i \partial x^j}$ and $\frac{\partial^2 f}{\partial x^j \partial x^i}$ exist and are continuous at a . Then

$$\frac{\partial^2 f}{\partial x^i \partial x^j}(a) = \frac{\partial^2 f}{\partial x^j \partial x^i}(a)$$

Proof. □

Definition 1.2.1.5. Let $U \subset \mathbb{R}^n$ be open and $f : U \rightarrow \mathbb{R}$. Then f is said to be **smooth** if for each $i_1, \dots, i_k \in \{1, \dots, n\}$, $\frac{\partial^k f}{\partial x^{i_1} \dots \partial x^{i_k}}$ exists and is continuous on U .

Definition 1.2.1.6. Let $U \subset \mathbb{R}^n$, $f : U \rightarrow \mathbb{R}$. Then f is said to be **smooth** if there exists $U' \subset \mathbb{R}^n$ and $f' : U' \rightarrow \mathbb{R}$ such that $U \subset U'$, U' is open, $f'|_U = f$ and f' is smooth. The set of smooth functions on U is denoted $C^\infty(U)$.

Theorem 1.2.1.7. Taylor's Theorem:

Let $U \subset \mathbb{R}^n$ be open and convex, $p \in U$, $f \in C^\infty(U)$ and $T \in \mathbb{N}$. Then there exist $(g_\alpha)_{|\alpha|=T+1} \subset C^\infty(U)$ such that for each $x \in U$,

$$f(x) = \sum_{k=0}^T \left[\sum_{|\alpha|=k} (x-p)^\alpha \partial^\alpha f(p) \right] + \sum_{|\alpha|=T+1} (x-p)^\alpha g_\alpha(x)$$

and for each $|\alpha| = T+1$,

$$g_\alpha(p) = \frac{1}{(T+1)!} \partial^\alpha f(p)$$

Proof. See analysis notes □

Definition 1.2.1.8. Let $U \subset \mathbb{R}^n$ and $F : U \rightarrow \mathbb{R}^m$. Let x^1, \dots, x^n be the standard coordinate functions on \mathbb{R}^n and y_1, \dots, y_m be the standard coordinate functions on \mathbb{R}^m . For $i \in \{1, \dots, m\}$, we define the **i th component of F** , denoted $F^i : U \rightarrow \mathbb{R}$, by

$$F^i = y^i \circ F$$

Thus $F = (F_1, \dots, F_m)$

Definition 1.2.1.9. Let $U \subset \mathbb{R}^n$ be open and $F : U \rightarrow \mathbb{R}^m$. Then F is said to be **smooth** if for each $i \in \{1, \dots, m\}$, the i th component of F , $F^i : U \rightarrow \mathbb{R}$, is smooth.

Definition 1.2.1.10. Let $U \subset \mathbb{R}^n$ and $F : U \rightarrow \mathbb{R}^m$. Then F is said to be **smooth** if for each $x \in U$, there exists $U_x \in \mathcal{N}_x$ and $\tilde{F} : U_x \rightarrow \mathbb{R}^m$ such that U_x is open, \tilde{F} is smooth and $\tilde{F}|_{U \cap U_x} = F|_{U \cap U_x}$.

Definition 1.2.1.11. Let $U \subset \mathbb{R}^n$ and $V \subset \mathbb{R}^m$ and $F : U \rightarrow V$. Then F is said to be a **diffeomorphism** if F is a bijection and F, F^{-1} are smooth.

Exercise 1.2.1.12. Let $U \subset \mathbb{R}^n$ and $V \subset \mathbb{R}^m$ and $F : U \rightarrow V$. If F is a diffeomorphism, then F is a homeomorphism.

Proof. Suppose that F is a diffeomorphism. By definition, F is a bijection and F and F^{-1} are smooth. Thus, F and F^{-1} are continuous and F is a homeomorphism. \square

Definition 1.2.1.13. Let $U \subset \mathbb{R}^n$ be open, $p \in U$ and $F : U \rightarrow \mathbb{R}^m$. We define the **Jacobian of F at p** , denoted $\frac{\partial F}{\partial x}(p) \in \mathbb{R}^{m \times n}$, by

$$\left(\frac{\partial F}{\partial x}(p) \right)_{i,j} = \frac{\partial F^i}{\partial x^j}(p)$$

Exercise 1.2.1.14. Inverse Function Theorem:

Let $U, V \subset \mathbb{R}^n$ be open and $F : U \rightarrow V$.

Exercise 1.2.1.15. Let $U, V \subset \mathbb{R}^n$ and $F : U \rightarrow V$. Then F is a diffeomorphism iff for each $p \in U$, there exists a relatively open neighborhood $N \subset U$ of p such that $F|_N : N \rightarrow F(N)$ is a diffeomorphism

Proof. content... \square

1.2.2 Integration

1.3 Topology

Definition 1.3.0.1. Let $(X, \mathcal{T}_X), (Y, \mathcal{T}_Y)$ be topological spaces and $f : X \rightarrow Y$. Then f is said to be **continuous** if for each $U \in \mathcal{T}$, $f^{-1}(U) \in \mathcal{T}_X$.

Definition 1.3.0.2. Let $(X, \mathcal{T}_X), (Y, \mathcal{T}_Y)$ be topological spaces and $f : X \rightarrow Y$. Then f is said to be a **homeomorphism** if f is a bijection and f, f^{-1} are continuous.

Definition 1.3.0.3. Let X, Y be topological spaces. Then X and Y are said to be **homeomorphic** if there exists $f : X \rightarrow Y$ such that f is a homeomorphism. If X and Y are homeomorphic, we write $X \cong Y$.

Theorem 1.3.0.4. Let $m, n \in \mathbb{N}$. If $m \neq n$, then $\mathbb{R}^m \not\cong \mathbb{R}^n$

Chapter 2

Multilinear Algebra

2.1 (r, s) -Tensors

Definition 2.1.0.1. Let V_1, \dots, V_k, W be vector spaces and $\alpha : \prod_{i=1}^n V_i \rightarrow W$. Then α is said to be **multilinear** if for each $i \in \{1, \dots, k\}$, $v \in V$, $c \in \mathbb{R}$ and $v_1, \dots, v_k \in V$,

$$\alpha(v_1, \dots, v_i + cv, \dots, v_k) = \alpha(v_1, \dots, v_i, \dots, v_k) + c\alpha(v_1, \dots, v, \dots, v_k)$$

We define

$$L(V_1, \dots, V_k; W) = \left\{ \alpha : \prod_{i=1}^n V_i \rightarrow W : \alpha \text{ is multilinear} \right\}$$

Note 2.1.0.2. For the remainder of this section we let V denote an n -dimensional vector space with basis $\{e^1, \dots, e^n\}$ with dual space V^* and dual basis $\{\epsilon_1, \dots, \epsilon_n\}$ defined by $\epsilon^i(e^j) = \delta_{i,j}$. We identify V with V^{**} by the isomorphism $V \rightarrow V^{**}$ defined by $v \mapsto \hat{v}$ where $\hat{v}(\alpha) = \alpha(v)$ for each $\alpha \in V^*$.

Definition 2.1.0.3. Let $\alpha : (V^*)^r \times V^s \rightarrow \mathbb{R}$. Then α is said to be an (r, s) -tensor on V if $\alpha \in L(\underbrace{V^*, \dots, V^*}_r, \underbrace{V, \dots, V}_s; \mathbb{R})$. The set of all (r, s) -tensors on V is denoted $T_s^r(V)$.

When $r = s = 0$, we set $T_s^r = \mathbb{R}$.

Exercise 2.1.0.4. We have that $T_s^r(V)$ is a vector space.

Proof. Clear. □

Exercise 2.1.0.5. Under the identification of V with V^{**} as noted above, we have that $V = T_0^1(V)$.

Proof. By definition,

$$\begin{aligned} V &= V^{**} \\ &= L(V^*; \mathbb{R}) \\ &= T_0^1(V) \end{aligned}$$

□

Definition 2.1.0.6. Let $\alpha \in T_{s_1}^{r_1}(V)$ and $\beta \in T_{s_2}^{r_2}(V)$. We define the **tensor product of α with β** , denoted $\alpha \otimes \beta \in T_{s_1+s_2}^{r_1+r_2}(V)$, by

$$\alpha \otimes \beta(v^*, w^*, v, w) = \alpha(v^*, v)\beta(w^*, w)$$

for each $v^* \in (V^*)^{r_1}$, $w^* \in (V^*)^{r_2}$, $v \in V^{s_1}$ and $w \in V^{s_2}$.

When $r_1 = s_1 = r_2 = s_2 = 0$ (so that $\alpha, \beta \in \mathbb{R}$), we set $\alpha \otimes \beta = \alpha\beta$.

Definition 2.1.0.7. We define the **tensor product**, denoted $\otimes : T_{s_1}^{r_1}(V) \times T_{s_2}^{r_2}(V) \rightarrow T_{s_1+s_2}^{r_1+r_2}(V)$ by

$$(\alpha, \beta) \mapsto \alpha \otimes \beta$$

Exercise 2.1.0.8. The tensor product $\otimes : T_{s_1}^{r_1}(V) \times T_{s_2}^{r_2}(V) \rightarrow T_{s_1+s_2}^{r_1+r_2}(V)$ is well defined.

Proof. Tedious but straightforward. □

Exercise 2.1.0.9. The tensor product $\otimes : T_{s_1}^{r_1}(V) \times T_{s_2}^{r_2}(V) \rightarrow T_{s_1+s_2}^{r_1+r_2}(V)$ is associative.

Proof. Let $\alpha \in T_{s_1}^{r_1}(V)$, $\beta \in T_{s_2}^{r_2}(V)$ and $\gamma \in T_{s_3}^{r_3}(V)$. Then for each $u^* \in (V^*)^{r_1}$, $v^* \in (V^*)^{r_2}$, $w^* \in (V^*)^{r_3}$, $u \in V^{s_1}$, $v \in V^{s_2}$, $w \in V^{s_3}$,

$$\begin{aligned} (\alpha \otimes \beta) \otimes \gamma(u^*, v^*, w^*, u, v, w) &= (\alpha \otimes \beta)(u^*, v^*, u, v) \gamma(w^*, w) \\ &= [\alpha(u^*, u) \beta(v^*, v)] \gamma(w^*, w) \\ &= \alpha(u^*, u) [\beta(v^*, v) \gamma(w^*, w)] \\ &= \alpha(u^*, u) (\beta \otimes \gamma)(v^*, w^*, v, w) \\ &= \alpha \otimes (\beta \otimes \gamma)(u^*, v^*, w^*, u, v, w) \end{aligned}$$

So that

$$(\alpha \otimes \beta) \otimes \gamma = \alpha \otimes (\beta \otimes \gamma)$$

□

Exercise 2.1.0.10. The tensor product $\otimes : T_{s_1}^{r_1}(V) \times T_{s_2}^{r_2}(V) \rightarrow T_{s_1+s_2}^{r_1+r_2}(V)$ is bilinear.

Proof.

1. Linearity in the first argument:

Let $\alpha, \beta \in T_{s_1}^{r_1}(V)$, $\gamma \in T_{s_2}^{r_2}(V)$, $\lambda \in \mathbb{R}$, $v^* \in (V^*)^{r_1}$, $w^* \in (V^*)^{r_2}$, $v \in V^{s_1}$ and $w \in V^{s_2}$. To see that the tensor product is linear in the first argument, we note that

$$\begin{aligned} [(\alpha + \lambda\beta) \otimes \gamma](v^*, w^*, v, w) &= (\alpha + \lambda\beta)(v^*, v) \gamma(w^*, w) \\ &= [\alpha(v^*, v) + \lambda\beta(v^*, v)] \gamma(w^*, w) \\ &= \alpha(v^*, v) \gamma(w^*, w) + \lambda\beta(v^*, v) \gamma(w^*, w) \\ &= \alpha \otimes \gamma(v^*, w^*, v, w) + \lambda(\beta \otimes \gamma)(v^*, w^*, v, w) \\ &= [\alpha \otimes \gamma + \lambda(\beta \otimes \gamma)](v^*, w^*, v, w) \end{aligned}$$

So that

$$(\alpha + \lambda\beta) \otimes \gamma = \alpha \otimes \gamma + \lambda(\beta \otimes \gamma)$$

2. Linearity in the second argument:

Similar to (1).

□

Definition 2.1.0.11.

1. Define $\mathcal{I}_{\otimes k} = \{(i_1, i_2, \dots, i_k) \in \mathbb{N}^k : i_1, \dots, i_k \leq n\}$. Each element $I \in \mathcal{I}_k$ is called an **unordered multi-index of length k** . Recall that $\#\mathcal{I}_{\otimes k} = n^k$.
2. Define $\mathcal{I}_{\wedge k} = \{(i_1, i_2, \dots, i_k) \in \mathbb{N}^k : i_1 < i_2 < \dots < i_k \leq n\}$. Each element $I \in \mathcal{I}_k$ is called an **ordered multi-index of length k** . Recall that $\#\mathcal{I}_{\wedge k} = \binom{n}{k}$.

Note 2.1.0.12. For the remainder of this section we will write \mathcal{I}_k in place of $\mathcal{I}_{\otimes k}$.

Definition 2.1.0.13. Let $I = \{(i_1, i_2, \dots, i_k) \in \mathcal{I}_k$.

1. Define $\epsilon^I \in (V^*)^k$ and $e_I \in V^k$ by

$$\epsilon^I = (\epsilon^{i_1}, \dots, \epsilon^{i_k})$$

and

$$e^I = (e^{i_1}, \dots, e^{i_k})$$

2. Define $e^{\otimes I} \in T_0^k(V)$ and $\epsilon^{\otimes I} \in T_k^0(V)$ by

$$e^{\otimes I} = e^{i_1} \otimes \cdots \otimes e^{i_k}$$

and

$$\epsilon^{\otimes I} = \epsilon^{i_1} \otimes \cdots \otimes \epsilon^{i_k}$$

Exercise 2.1.0.14. Let $\alpha, \beta \in T_s^r(V)$. If for each $I \in \mathcal{I}_r, J \in \mathcal{I}_s$, $\alpha(\epsilon^I, e^J) = \beta(\epsilon^I, e^J)$, then $\alpha = \beta$.

Proof. Suppose that for each $I \in \mathcal{I}_r, J \in \mathcal{I}_s$, $\alpha(\epsilon^I, e^J) = \beta(\epsilon^I, e^J)$. Let $v_1^*, \dots, v_r^* \in V^*$ and $v_1, \dots, v_s \in V$. For each $i \in \{1, \dots, r\}$ and $j \in \{1, \dots, s\}$, write

$$v_i^* = \sum_{k_i=1}^n a_{i,k_i} \epsilon^{k_i}$$

and

$$v_j = \sum_{l_j=1}^n b_{j,l_j} e^{l_j}$$

Then

$$\begin{aligned} \alpha(v_1^*, \dots, v_r^*, v_1, \dots, v_s) &= \sum_{k_1, \dots, k_r=1}^n \sum_{l_1, \dots, l_s=1}^n \prod_{i=1}^r \prod_{j=1}^s a_{i,k_i} b_{j,l_j} \alpha(\epsilon^{k_1}, \dots, \epsilon^{k_r}, e^{l_1}, \dots, e^{l_s}) \\ &= \sum_{k_1, \dots, k_r=1}^n \sum_{l_1, \dots, l_s=1}^n \prod_{i=1}^r \prod_{j=1}^s a_{i,k_i} b_{j,l_j} \beta(\epsilon^{k_1}, \dots, \epsilon^{k_r}, e^{l_1}, \dots, e^{l_s}) \\ &= \beta(v_1^*, \dots, v_r^*, v_1, \dots, v_s) \end{aligned}$$

So that $\alpha = \beta$. □

Exercise 2.1.0.15. Let $I, K \in \mathcal{I}_r$ and $J, L \in \mathcal{I}_s$. Then $e^{\otimes I} \otimes \epsilon^{\otimes J}(\epsilon^K, e^L) = \delta_{I,K} \delta_{J,L}$.

Proof. Write $I = (i_1, \dots, i_r), K = (k_1, \dots, k_r)$ and $J = (j_1, \dots, j_s), L = (l_1, \dots, l_s)$. Then

$$\begin{aligned} e^{\otimes I} \otimes \epsilon^{\otimes J}(\epsilon^K, e^L) &= e^{\otimes I}(\epsilon^K) \epsilon^{\otimes J}(e^L) \\ &= e^{i_1} \otimes \cdots \otimes e^{i_r}(\epsilon^{k_1}, \dots, \epsilon^{k_r}) \epsilon^{j_1} \otimes \cdots \otimes \epsilon^{j_s}(e^{l_1}, \dots, e^{l_s}) \\ &= \left[\prod_{m=1}^r e^{i_m}(\epsilon^{k_m}) \right] \left[\prod_{n=1}^s \epsilon^{j_n}(e^{l_n}) \right] \\ &= \left[\prod_{m=1}^r \delta_{i_m, k_m} \right] \left[\prod_{n=1}^s \delta_{j_n, l_n} \right] \\ &= \delta_{I,K} \delta_{J,L} \end{aligned}$$

□

Exercise 2.1.0.16. The set $\{e^{\otimes I} \otimes \epsilon^{\otimes J} : I \in \mathcal{I}_r, J \in \mathcal{I}_s\}$ is a basis for $T_s^r(V)$ and $\dim T_s^r(V) = n^{r+s}$.

Proof. Let $(a_J^I)_{I \in \mathcal{I}_r, J \in \mathcal{I}_s} \subset \mathbb{R}$. Let $\alpha = \sum_{(I,J) \in \mathcal{I}_r \times \mathcal{I}_s} a_J^I e^{\otimes I} \otimes \epsilon^{\otimes J}$. Suppose that $\alpha = 0$. Then for each $(I, J) \in \mathcal{I}_r \times \mathcal{I}_s$, $\alpha(\epsilon^I, e^J) = a_J^I = 0$. Thus $\{e^{\otimes I} \otimes \epsilon^{\otimes J} : I \in \mathcal{I}_r, J \in \mathcal{I}_s\}$ is linearly independent. Let $\beta \in T_s^r(V)$. For $(I, J) \in \mathcal{I}_r \times \mathcal{I}_s$, put $b_J^I = \beta(\epsilon^I, e^J)$. Define $\mu = \sum_{(I,J) \in \mathcal{I}_r \times \mathcal{I}_s} b_J^I e^{\otimes I} \otimes \epsilon^{\otimes J} \in T_s^r(V)$. Then for each $(I, J) \in \mathcal{I}_r \times \mathcal{I}_s$, $\mu(\epsilon^I, e^J) = b_J^I = \beta(\epsilon^I, e^J)$. Hence $\mu = \beta$ and therefore $\beta \in \text{span}\{e^{\otimes I} \otimes \epsilon^{\otimes J}\}$. □

2.2 Covariant k -Tensors

Definition 2.2.0.1. Let $\alpha : V^k \rightarrow \mathbb{R}$. Then α is said to be a **covariant k -tensor on V** if $\alpha \in T_k^0(V)$. We denote the set of covariant k -tensors by $T^k(V)$.

Definition 2.2.0.2. For $\sigma \in S_k$ and $\alpha \in T^k(V)$, define the $\sigma\alpha : V^k \rightarrow \mathbb{R}$ by

$$\sigma\alpha(v_1, \dots, v_k) = \alpha(v_{\sigma(1)}, \dots, v_{\sigma(k)})$$

The map $\alpha \mapsto \sigma\alpha$ is called the **permutation action** of S_k on $T^k(V)$

Exercise 2.2.0.3. The permutation action of S_k on $T^k(V)$ is a group action.

Proof.

1. Clearly for each $\sigma \in S_k$ and $\alpha \in T^k(V)$, $\sigma\alpha \in T^k(V)$.
2. Clearly for each $\alpha \in T^k(V)$, $e\alpha = \alpha$.
3. Let $\tau, \sigma \in S_k$ and $\alpha \in T^k(V)$. Then for each $v_1, \dots, v_k \in V$,

$$\begin{aligned} (\tau\sigma)\alpha(v_1, \dots, v_k) &= \alpha(v_{\tau\sigma(1)}, \dots, v_{\tau\sigma(k)}) \\ &= \tau\alpha(v_{\sigma(1)}, \dots, v_{\sigma(k)}) \\ &= \tau(\sigma\alpha)(v_1, \dots, v_k) \end{aligned}$$

□

Exercise 2.2.0.4. Let $\sigma \in S_k$. Then $L_\sigma : T^k(V) \rightarrow T^k(V)$ given by $L_\sigma(\alpha) = \sigma\alpha$ is a linear transformation.

Proof. Let $\alpha, \beta \in T^k(V)$, $c \in \mathbb{R}$ and $v_1, \dots, v_k \in V$. Then

$$\begin{aligned} \sigma(c\alpha + \beta)(v_1, \dots, v_k) &= (c\alpha + \beta)(v_{\sigma(1)}, \dots, v_{\sigma(k)}) \\ &= c\alpha(v_{\sigma(1)}, \dots, v_{\sigma(k)}) + \beta(v_{\sigma(1)}, \dots, v_{\sigma(k)}) \\ &= c\sigma\alpha(v_1, \dots, v_k) + \sigma\beta(v_1, \dots, v_k) \end{aligned}$$

So $\sigma(c\alpha + \beta) = c\sigma\alpha + \sigma\beta$.

□

Definition 2.2.0.5. Let $\alpha \in T^k(V)$. Then α is said to be

- **symmetric** if for each $\sigma \in S_k$, $\sigma\alpha = \alpha$
- **antisymmetric** if for each $\sigma \in S_k$, $\sigma\alpha = \text{sgn}(\sigma)\alpha$
- **alternating** if for each $v_1, \dots, v_k \in V$, if there exists $i, j \in \{1, \dots, k\}$ such that $v_i = v_j$, then $\alpha(v_1, \dots, v_k) = 0$.

We denote the set of symmetric k -tensors on V by $\Sigma^k(V)$. We denote the set of alternating k -tensors on V by $\Lambda^k(V)$.

Exercise 2.2.0.6. Let $\alpha \in T^k(V)$. Then α is antisymmetric iff α is alternating.

Proof. Suppose that α is antisymmetric. Let $v_1, \dots, v_k \in V$. Suppose that there exists $i, j \in \{1, \dots, k\}$ such that $v_i = v_j$. Define $\sigma \in S_k$ by $\sigma = (i, j)$. Then

$$\begin{aligned} \alpha(v_1, \dots, v_i, \dots, v_j, \dots, v_k) &= \alpha(v_1, \dots, v_j, \dots, v_i, \dots, v_k) \\ &= \sigma(\alpha)(v_1, \dots, v_i, \dots, v_j, \dots, v_k) \\ &= \text{sgn}(\sigma)\alpha(v_1, \dots, v_i, \dots, v_j, \dots, v_k) \\ &= -\alpha(v_1, \dots, v_i, \dots, v_j, \dots, v_k) \end{aligned}$$

Therefore $2\alpha(v_1, \dots, v_i, \dots, v_j, \dots, v_k) = 0$ which implies that $\alpha(v_1, \dots, v_i, \dots, v_j, \dots, v_k) = 0$. Hence α is alternating.

Conversely, suppose that α is alternating. Let $i, j \in \{1, \dots, k\}$ and $v_1, \dots, v_k \in V$. Then

$$\begin{aligned} 0 &= \alpha(v_1, \dots, v_i + v_j, \dots, v_i + v_j, \dots, v_k) \\ &= \alpha(v_1, \dots, v_i, \dots, v_j, \dots, v_k) + \alpha(v_1, \dots, v_j, \dots, v_i, \dots, v_k) \end{aligned}$$

Since $i, j \in \{1, \dots, k\}$ and $v_1, \dots, v_k \in V$ are arbitrary, we have that for each $\tau \in S_k$, τ is a transposition implies that

$$\begin{aligned} \tau\alpha &= -\alpha \\ &= \text{sgn}(\tau)\alpha \end{aligned}$$

Let $n \in \mathbb{N}$. Suppose that for each $\tau_1, \dots, \tau_{n-1} \in S_k$ if for each $j \in \{1, \dots, n-1\}$, τ_j is a transposition, then $(\tau_1 \cdots \tau_{n-1})\alpha = \text{sgn}(\tau_1 \cdots \tau_{n-1})\alpha$. Let $\tau_1, \dots, \tau_n \in S_k$. Suppose that for each $j \in \{1, \dots, n\}$, τ_j is a transposition. Then

$$\begin{aligned} (\tau_1 \cdots \tau_n)\alpha &= (\tau_1 \cdots \tau_{n-1})(\tau_n\alpha) \\ &= (\tau_1 \cdots \tau_{n-1})(\text{sgn}(\tau_n)\alpha) \\ &= (\text{sgn}(\tau_n)(\tau_1 \cdots \tau_{n-1})\alpha) \\ &= (\text{sgn}(\tau_n) \text{sgn}(\tau_1 \cdots \tau_{n-1})\alpha) \\ &= \text{sgn}(\tau_1 \cdots \tau_n)\alpha \end{aligned}$$

By induction, for each $n \in \mathbb{N}$ and $\tau_1, \dots, \tau_n \in S_k$, if for each $j \in \{1, \dots, n\}$, τ_j is a transposition, then $(\tau_1 \cdots \tau_n)\alpha = \text{sgn}(\tau_1 \cdots \tau_n)\alpha$. Now let $\sigma \in S_k$. Then there exist $n \in \mathbb{N}$ and $\tau_1, \dots, \tau_n \in S_k$ such that $\sigma = \tau_1 \cdots \tau_n$ and for each $j \in \{1, \dots, n\}$, τ_j is a transposition. Hence

$$\begin{aligned} \sigma\alpha &= (\tau_1 \cdots \tau_n)\alpha \\ &= \text{sgn}(\tau_1 \cdots \tau_n)\alpha \\ &= \text{sgn}(\sigma)\alpha \end{aligned}$$

Therefore α is antisymmetric. □

Definition 2.2.0.7. Define the **symmetric operator** $S : T^k(V) \rightarrow \Sigma^k(V)$ by

$$\text{Sym}(\alpha) = \frac{1}{k!} \sum_{\sigma \in S_k} \sigma\alpha$$

Define the **alternating operator** $A : T^k(V) \rightarrow \Lambda^k(V)$ by

$$\text{Alt}(\alpha) = \frac{1}{k!} \sum_{\sigma \in S_k} \text{sgn}(\sigma)\sigma\alpha$$

Exercise 2.2.0.8.

1. For $\alpha \in T^k(V)$, $\text{Sym}(\alpha)$ is symmetric.
2. For $\alpha \in T^k(V)$, $\text{Alt}(\alpha)$ is alternating.

Proof.

1. Let $\alpha \in T^k(V)$ and $\sigma \in S_k$. Then

$$\begin{aligned} \sigma \text{Sym}(\alpha) &= \sigma \left[\frac{1}{k!} \sum_{\tau \in S_k} \tau\alpha \right] \\ &= \frac{1}{k!} \sum_{\tau \in S_k} \sigma\tau\alpha \\ &= \frac{1}{k!} \sum_{\tau \in S_k} \tau\alpha \\ &= \text{Sym}(\alpha) \end{aligned}$$

2. Let $\alpha \in T^k(V)$ and $\sigma \in S_k$. Then

$$\begin{aligned}
 \sigma \operatorname{Alt}(\alpha) &= \sigma \left[\frac{1}{k!} \sum_{\tau \in S_k} \operatorname{sgn}(\tau) \tau \alpha \right] \\
 &= \frac{1}{k!} \sum_{\tau \in S_k} \operatorname{sgn}(\tau) \sigma \tau \alpha \\
 &= \frac{1}{k!} \sum_{\tau \in S_k} \operatorname{sgn}(\sigma) \operatorname{sgn}(\sigma \tau) \sigma \tau \alpha \\
 &= \operatorname{sgn}(\sigma) \frac{1}{k!} \sum_{\tau \in S_k} \operatorname{sgn}(\sigma \tau) \sigma \tau \alpha \\
 &= \operatorname{sgn}(\sigma) \frac{1}{k!} \sum_{\tau \in S_k} \operatorname{sgn}(\tau) \tau \alpha \\
 &= \operatorname{sgn}(\sigma) \operatorname{Alt}(\alpha)
 \end{aligned}$$

□

Exercise 2.2.0.9.

1. For $\alpha \in \Sigma^k(V)$, $\operatorname{Sym}(\alpha) = \alpha$.
2. For $\alpha \in \Lambda^k(V)$, $\operatorname{Alt}(\alpha) = \alpha$.

Proof.

1. Let $\alpha \in \Sigma^k(V)$. Then

$$\begin{aligned}
 \operatorname{Sym}(\alpha) &= \frac{1}{k!} \sum_{\sigma \in S_k} \sigma \alpha \\
 &= \frac{1}{k!} \sum_{\sigma \in S_k} \alpha \\
 &= \alpha
 \end{aligned}$$

2. Let $\alpha \in \Lambda^k(V)$. Then

$$\begin{aligned}
 \operatorname{Alt}(\alpha) &= \frac{1}{k!} \sum_{\sigma \in S_k} \operatorname{sgn}(\sigma) \sigma \alpha \\
 &= \frac{1}{k!} \sum_{\sigma \in S_k} \operatorname{sgn}(\sigma)^2 \alpha \\
 &= \alpha
 \end{aligned}$$

□

Exercise 2.2.0.10. The symmetric operator $S : T^k(V) \rightarrow \Sigma^k(V)$ and the alternating operator $A : T^k(V) \rightarrow \Lambda^k(V)$ are linear.

Proof. Clear. □

Definition 2.2.0.11. Let $\alpha \in \Lambda^k(V)$ and $\beta \in \Lambda^l(V)$. The **exterior product** of α and β is defined to be the map $\alpha \wedge \beta \in \Lambda^{k+l}(V)$ given by

$$\alpha \wedge \beta = \frac{(k+l)!}{k!l!} \operatorname{Alt}(\alpha \otimes \beta)$$

Thus $\wedge : \Lambda^k(V) \times \Lambda^l(V) \rightarrow \Lambda^{k+l}(V)$.

Exercise 2.2.0.12. The exterior product $\wedge : \Lambda^k(V) \times \Lambda^l(V) \rightarrow \Lambda^{k+l}(V)$ is bilinear.

Proof. Clear. □

Exercise 2.2.0.13. Let $\alpha \in T^k(V)$ and $\beta \in T_l(V)$. Then

$$1. \text{ Alt}(\text{Alt}(\alpha) \otimes \beta) = \text{Alt}(\alpha \otimes \beta)$$

$$2. \text{ Alt}(\alpha \otimes \text{Alt}(\beta)) = \text{Alt}(\alpha \otimes \beta)$$

Proof. First note that if we fix $\mu \in S_{k+l}$, then for each $\tau \in S_k$, choosing $\sigma = \mu\tau^{-1}$ yields $\sigma\tau = \mu$. For each $\mu \in S_{k+l}$, the map $\phi_\mu : S_k \rightarrow S_{k+l}$ given by $\phi_\mu(\tau) = \mu\tau^{-1}$ is injective. Thus for each $\mu \in S_{k+l}$, we have that $\#\{(\sigma, \tau) \in S_{k+l} \times S_k : \mu = \sigma\tau\} = k!$

1. Then

$$\begin{aligned} \text{Alt}(\text{Alt}(\alpha) \otimes \beta) &= \frac{1}{(k+l)!} \sum_{\sigma \in S_{k+l}} \text{sgn}(\sigma) \sigma \left[\text{Alt}(\alpha) \otimes \beta \right] \\ &= \frac{1}{(k+l)!} \sum_{\sigma \in S_{k+l}} \text{sgn}(\sigma) \sigma \left[\left(\frac{1}{k!} \sum_{\tau \in S_k} \text{sgn}(\tau) \tau \alpha \right) \otimes \beta \right] \\ &= \frac{1}{(k+l)!} \sum_{\sigma \in S_{k+l}} \text{sgn}(\sigma) \sigma \left[\frac{1}{k!} \sum_{\tau \in S_k} \text{sgn}(\tau) (\tau \alpha) \otimes \beta \right] \\ &= \frac{1}{(k+l)!} \sum_{\sigma \in S_{k+l}} \text{sgn}(\sigma) \sigma \left[\frac{1}{k!} \sum_{\tau \in S_k} \text{sgn}(\tau) \tau (\alpha \otimes \beta) \right] \\ &= \frac{1}{k!(k+l)!} \sum_{\sigma \in S_{k+l}} \sum_{\tau \in S_k} \text{sgn}(\sigma\tau) \sigma\tau (\alpha \otimes \beta) \\ &= \frac{k!}{k!(k+l)!} \sum_{\mu \in S_{k+l}} \text{sgn}(\mu) \mu (\alpha \otimes \beta) \\ &= \frac{1}{(k+l)!} \sum_{\mu \in S_{k+l}} \text{sgn}(\mu) \mu (\alpha \otimes \beta) \\ &= \text{Alt}(\alpha \otimes \beta) \end{aligned}$$

2. Similar to (1). □

Exercise 2.2.0.14. The exterior product $\wedge : \Lambda^k(V) \times \Lambda^l(V) \rightarrow \Lambda^{k+l}(V)$ is associative.

Proof. Let $\alpha \in \Lambda^k(V)$, $\beta \in \Lambda^l(V)$ and $\gamma \in \Lambda^m(V)$. Then

$$\begin{aligned}
 (\alpha \wedge \beta) \wedge \gamma &= \left[\frac{(k+l)!}{k!l!} \text{Alt}(\alpha \otimes \beta) \right] \wedge \gamma \\
 &= \frac{(k+l+m)!}{(k+l)!m!} \text{Alt} \left(\left[\frac{(k+l)!}{k!l!} \text{Alt}(\alpha \otimes \beta) \right] \otimes \gamma \right) \\
 &= \frac{(k+l+m)!}{(k+l)!m!} \frac{(k+l)!}{k!l!} \text{Alt}(\text{Alt}(\alpha \otimes \beta) \otimes \gamma) \\
 &= \frac{(k+l+m)!}{m!} \frac{1}{k!l!} \text{Alt}((\alpha \otimes \beta) \otimes \gamma) \\
 &= \frac{(k+l+m)!}{k!(l+m)!} \frac{(l+m)!}{l!m!} \text{Alt}(\alpha \otimes (\beta \otimes \gamma)) \\
 &= \frac{(k+l+m)!}{k!(l+m)!} \frac{(l+m)!}{l!m!} \text{Alt}(\alpha \otimes \text{Alt}(\beta \otimes \gamma)) \\
 &= \frac{(k+l+m)!}{k!(l+m)!} \text{Alt}(\alpha \otimes \frac{(l+m)!}{l!m!} \text{Alt}(\beta \otimes \gamma)) \\
 &= \frac{(k+l+m)!}{k!(l+m)!} \text{Alt}(\alpha \otimes (\beta \wedge \gamma)) \\
 &= \alpha \wedge (\beta \wedge \gamma)
 \end{aligned}$$

□

Exercise 2.2.0.15. Let $\alpha_i \in \Lambda^{k_i}(V)$ for $i = 1, \dots, m$. Then

$$\bigwedge_{i=1}^m \alpha_i = \frac{(\sum_{i=1}^m k_i)!}{\prod_{i=1}^m k_i!} \text{Alt} \left(\bigotimes_{i=1}^m \alpha_i \right)$$

Proof. To see that the statement is true in the case $m = 3$, the proof of the previous exercise tells us that indeed

$$\alpha_1 \wedge \alpha_2 \wedge \alpha_3 = \frac{(k_1 + k_2 + k_3)!}{k_1!k_2!k_3!} \text{Alt}(\alpha_1 \otimes \alpha_2 \otimes \alpha_3)$$

Now, suppose that the statement is true for each $3 \leq m \leq m_0$. Then the proof of the previous exercise tells us the

$$\begin{aligned}
 \bigwedge_{i=1}^{m_0+1} \alpha_i &= \left(\bigwedge_{i=1}^{m_0-1} \alpha_i \right) \wedge \alpha_{m_0} \wedge \alpha_{m_0+1} \\
 &= \frac{(\sum_{i=1}^{m_0-1} k_i + k_{m_0} + k_{m_0+1})!}{(\sum_{i=1}^{m_0-1} k_i)!k_{m_0}!k_{m_0+1}!} \text{Alt} \left(\left[\bigwedge_{i=1}^{m_0-1} \alpha_i \right] \otimes \alpha_{m_0} \otimes \alpha_{m_0+1} \right) \\
 &= \frac{(\sum_{i=1}^{m_0-1} k_i + k_{m_0} + k_{m_0+1})!}{(\sum_{i=1}^{m_0-1} k_i)!k_{m_0}!k_{m_0+1}!} \text{Alt} \left(\left[\frac{(\sum_{i=1}^{m_0-1} k_i)!}{\prod_{i=1}^{m_0-1} k_i!} \text{Alt} \left(\bigotimes_{i=1}^{m_0-1} \alpha_i \right) \right] \otimes \alpha_{m_0} \otimes \alpha_{m_0+1} \right) \\
 &= \frac{(\sum_{i=1}^{m_0+1} k_i)!}{\prod_{i=1}^{m_0+1} k_i!} \text{Alt} \left(\text{Alt} \left[\bigotimes_{i=1}^{m_0-1} \alpha_i \right] \otimes \alpha_{m_0} \otimes \alpha_{m_0+1} \right) \\
 &= \frac{(\sum_{i=1}^{m_0+1} k_i)!}{\prod_{i=1}^{m_0+1} k_i!} \text{Alt} \left(\left[\bigotimes_{i=1}^{m_0-1} \alpha_i \right] \otimes \alpha_{m_0} \otimes \alpha_{m_0+1} \right) \\
 &= \frac{(\sum_{i=1}^{m_0+1} k_i)!}{\prod_{i=1}^{m_0+1} k_i!} \text{Alt} \left(\bigotimes_{i=1}^{m_0+1} \alpha_i \right)
 \end{aligned}$$

□

Exercise 2.2.0.16. Define $\tau \in S_{k+l}$ by

$$\tau = \begin{pmatrix} 1 & 2 & \cdots & l & l+1 & l+2 & \cdots & l+k \\ 1+k & 2+k & \cdots & l+k & 1 & 2 & \cdots & k \end{pmatrix}$$

Then the inversion number of τ is kl . (Hint: inversion number)

Proof.

$$\begin{aligned} N(\tau) &= \sum_{i=1}^l k \\ &= kl \end{aligned}$$

Since $\text{sgn}(\tau) = (-1)^{N(\tau)}$ we know that $\text{sgn}(\tau) = (-1)^{kl}$. □

Exercise 2.2.0.17. Let $\alpha \in \Lambda^k(V)$, $\beta \in \Lambda^l(V)$. Then

$$\alpha \wedge \beta = (-1)^{kl} \beta \wedge \alpha$$

Proof. Define $\tau \in S_{k+l}$ as in the previous exercise. Note that For $\sigma \in S_{k+l}$ and $v_1, \dots, v_{k+l} \in V$, we have that

$$\begin{aligned} \sigma\tau(\beta \otimes \alpha)(v_1, \dots, v_l, v_{l+1}, \dots, v_{l+k}) &= \beta \otimes \alpha(v_{\sigma\tau(1)}, \dots, v_{\sigma\tau(l)}, v_{\sigma\tau(l+1)}, \dots, v_{\sigma\tau(l+k)}) \\ &= \beta(v_{\sigma\tau(1)}, \dots, v_{\sigma\tau(l)})\alpha(v_{\sigma\tau(l+1)}, \dots, v_{\sigma\tau(l+k)}) \\ &= \beta(v_{\sigma(1+k)}, \dots, v_{\sigma(l+k)})\alpha(v_{\sigma(1)}, \dots, v_{\sigma(k)}) \\ &= \alpha(v_{\sigma(1)}, \dots, v_{\sigma(k)})\beta(v_{\sigma(1+k)}, \dots, v_{\sigma(l+k)}) \\ &= \alpha \otimes \beta(v_{\sigma(1)}, \dots, v_{\sigma(k)}, v_{\sigma(1+k)}, \dots, v_{\sigma(l+k)}) \\ &= \sigma(\alpha \otimes \beta)(v_1, \dots, v_k, v_{1+k}, \dots, v_{l+k}) \end{aligned}$$

Thus $\sigma\tau(\beta \otimes \alpha) = \sigma(\alpha \otimes \beta)$. Then

$$\begin{aligned} \beta \wedge \alpha &= \frac{(k+l)!}{k!l!} \text{Alt}(\beta \otimes \alpha) \\ &= \frac{(k+l)!}{k!l!} \frac{1}{(k+l)!} \sum_{\sigma \in S_{k+l}} \text{sgn}(\sigma) \sigma(\beta \otimes \alpha) \\ &= \frac{(k+l)!}{k!l!} \frac{1}{(k+l)!} \sum_{\sigma \in S_{k+l}} \text{sgn}(\sigma\tau) \sigma\tau(\beta \otimes \alpha) \\ &= \text{sgn}(\tau) \frac{(k+l)!}{k!l!} \frac{1}{(k+l)!} \sum_{\sigma \in S_{k+l}} \text{sgn}(\sigma) \sigma(\alpha \otimes \beta) \\ &= \text{sgn}(\tau) \frac{(k+l)!}{k!l!} \text{Alt}(\alpha \otimes \beta) \\ &= \text{sgn}(\tau) \alpha \wedge \beta \\ &= (-1)^{kl} \alpha \wedge \beta \end{aligned}$$

□

Exercise 2.2.0.18. Let $\alpha \in \Lambda^k(V)$. If k is odd, then $\alpha \wedge \alpha = 0$.

Proof. Suppose that k is odd. The previous exercise tells us that

$$\begin{aligned} \alpha \wedge \alpha &= (-1)^{k^2} \alpha \wedge \alpha \\ &= -\alpha \wedge \alpha \end{aligned}$$

Thus $\alpha \wedge \alpha = 0$. □

Exercise 2.2.0.19. Fundamental Example:

Let $\alpha_1, \dots, \alpha_m \in \Lambda^1(V)$ and $v_1, \dots, v_m \in V$. Then

$$\left(\bigwedge_{i=1}^m \alpha_i \right) (v_1, \dots, v_m) = \det(\alpha_i(v_j))$$

Proof. The previous exercises tell us that

$$\begin{aligned} \left(\bigwedge_{i=1}^m \alpha_i \right) (v_1, \dots, v_m) &= m! \operatorname{Alt} \left(\bigotimes_{i=1}^m \alpha_i \right) (v_1, \dots, v_m) \\ &= m! \left[\frac{1}{m!} \sum_{\sigma \in S_m} \operatorname{sgn}(\sigma) \sigma \left(\bigotimes_{i=1}^m \alpha_i \right) \right] (v_1, \dots, v_m) \\ &= \sum_{\sigma \in S_m} \operatorname{sgn}(\sigma) \left(\bigotimes_{i=1}^m \alpha_i \right) (v_{\sigma(1)}, \dots, v_{\sigma(m)}) \\ &= \sum_{\sigma \in S_m} \operatorname{sgn}(\sigma) \prod_{i=1}^m \alpha_i(v_{\sigma(i)}) \\ &= \det(\alpha_i(v_j)) \end{aligned}$$

□

Note 2.2.0.20. Recall that $\mathcal{I}_{\wedge k} = \{(i_1, i_2, \dots, i_k) \in \mathbb{N}^k : i_1 < i_2 < \dots < i_k \leq n\}$ and that $\#\mathcal{I}_{\wedge k} = \binom{n}{k}$. For the remainder of this section, we will write \mathcal{I}_k in place of $\mathcal{I}_{\wedge k}$.

Definition 2.2.0.21. Let $I = \{(i_1, i_2, \dots, i_k) \in \mathcal{I}_k\}$. Define $\epsilon^{\wedge I} \in \Lambda^k(V)$ by

$$\epsilon^{\wedge I} = \epsilon^{i_1} \wedge \dots \wedge \epsilon^{i_k}$$

Exercise 2.2.0.22. Let $I = (i_1, \dots, i_k)$ and $J = (j_1, \dots, j_k) \in \mathcal{I}_k$. Then $\epsilon^{\wedge I}(e^J) = \delta_{I,J}$.

Proof. Put $A = \begin{pmatrix} \epsilon^{i_1}(e^{j_1}) & \dots & \epsilon^{i_1}(e^{j_k}) \\ \vdots & & \vdots \\ \epsilon^{i_k}(e^{j_1}) & \dots & \epsilon^{i_k}(e^{j_k}) \end{pmatrix}$. A previous exercise tells us that $\epsilon^{\wedge I}(e^J) = \det A$. If $I = J$, then

$A = I_{k \times k}$ and therefore $\epsilon^{\wedge I}(e^J) = 1$. Suppose that $I \neq J$. Put $l_0 = \min\{l : 1 \leq l \leq k, i_l \neq j_l\}$. If $i_{l_0} < j_{l_0}$, then all entries on the l_0 th row of A are 0. If $i_{l_0} > j_{l_0}$, then all entries on the l_0 th column of A are 0. □

Exercise 2.2.0.23. Let $\alpha, \beta \in \Lambda^k(V)$. If for each $I \in \mathcal{I}_k$, $\alpha(e^I) = \beta(e^I)$, then $\alpha = \beta$.

Proof. Suppose that for each $I \in \mathcal{I}_k$, $\alpha(e^I) = \beta(e^I)$. Let $v_1, \dots, v_k \in V$. For $i = 1, \dots, k$, write $v_i =$

$\sum_{j_i=1}^n a_{i,j_i} e^{j_i}$. Then

$$\begin{aligned}
 \alpha(v_1, \dots, v_k) &= \sum_{j_1, \dots, j_k=1}^n \left(\prod_{i=1}^k a_{i,j_i} \right) \alpha(e^{j_1}, \dots, e^{j_k}) \\
 &= \sum_{j_1 \neq \dots \neq j_k}^n \left(\prod_{i=1}^k a_{i,j_i} \right) \alpha(e^{j_1}, \dots, e^{j_k}) \\
 &= \sum_{J \in \mathcal{I}_k} \left[\sum_{\sigma \in S_J} \operatorname{sgn}(\sigma) \left(\prod_{i=1}^k a_{i, \sigma(j_i)} \right) \right] \alpha(e^J) \\
 &= \sum_{J \in \mathcal{I}_k} \left[\sum_{\sigma \in S_J} \operatorname{sgn}(\sigma) \left(\prod_{i=1}^k a_{i, \sigma(j_i)} \right) \right] \beta(e^J) \\
 &= \sum_{j_1, \dots, j_k=1}^n \left(\prod_{i=1}^k a_{i,j_i} \right) \beta(e^{j_1}, \dots, e^{j_k}) \\
 &= \beta(v_1, \dots, v_k)
 \end{aligned}$$

□

Exercise 2.2.0.24. The set $\{\epsilon^{\wedge I} : I \in \mathcal{I}_k\}$ is a basis for $\Lambda^k(V)$ and $\dim \Lambda^k(V) = \binom{n}{k}$.

Proof. Let $(a_I)_{I \in \mathcal{I}_k} \subset \mathbb{R}$. Let $\alpha = \sum_{I \in \mathcal{I}_k} a_I \epsilon^{\wedge I}$. Suppose that $\alpha = 0$. Then for each $J \in \mathcal{I}_k$, $\alpha(e^J) = a_J = 0$.

Thus $\{\epsilon^{\wedge I} : I \in \mathcal{I}_k\}$ is linearly independent. Let $\beta \in \Lambda^k(V)$. For $I \in \mathcal{I}_k$, put $b_I = \beta(e^I)$. Define $\mu = \sum_{I \in \mathcal{I}_k} b_I \epsilon^{\wedge I} \in \Lambda^k(V)$. Then for each $J \in \mathcal{I}_k$, $\mu(e^J) = b_J = \beta(e^J)$. Hence $\mu = \beta$ and therefore $\beta \in \operatorname{span}\{\epsilon^{\wedge I} : I \in \mathcal{I}_k\}$. □

Chapter 3

Smooth Manifolds

3.1 Topological Manifolds

Exercise 3.1.0.1. We have that \mathbb{R} is homeomorphic to $(0, \infty)$

Proof. Define $f : \mathbb{R} \rightarrow (0, \infty)$ by $f(x) = e^x$. Then f is a homeomorphism. □

Definition 3.1.0.2. Let $n \in \mathbb{N}$. We define the **upper half space** of \mathbb{R}^n , denoted \mathbb{H}^n , by

$$\mathbb{H}^n = \{(x_1, x_2, \dots, x_n) \in \mathbb{R}^n : x_n \geq 0\}$$

and we define

$$\partial\mathbb{H}^n = \{(x_1, x_2, \dots, x_n) \in \mathbb{R}^n : x_n = 0\}$$

$$\text{Int } \mathbb{H}^n = \{(x_1, x_2, \dots, x_n) \in \mathbb{R}^n : x_n > 0\}$$

We endow \mathbb{H}^n , $\partial\mathbb{H}^n$ and $\text{Int } \mathbb{H}^n$ with the subspace topology inherited from \mathbb{R}^n .

Definition 3.1.0.3. We define $\mathbb{R}^0 = \{0\}$ and $\mathbb{H}^0 = \emptyset$ endowed with the discrete topology.

Exercise 3.1.0.4. Let $n \in \mathbb{N}$.

1. $\partial\mathbb{H}^n$ is homeomorphic to \mathbb{R}^{n-1}
2. $\text{Int } \mathbb{H}^n$ is homeomorphic to \mathbb{R}^n

Proof.

1. Define $f : \partial\mathbb{H}^n \rightarrow \mathbb{R}^{n-1}$ by

$$f(x_1, \dots, x_{n-1}, 0) = (x_1, \dots, x_{n-1})$$

Then f is a homeomorphism.

2. Define $f : \mathbb{R}^n \rightarrow \text{Int } \mathbb{H}^n$ by $f(x_1, \dots, x_{n-1}, x_n) = (x_1, \dots, x_{n-1}, e^{x_n})$. Then f is a homeomorphism. □

Definition 3.1.0.5. Let M be a topological space and $n \in \mathbb{N}_0$. Let $U \subset M$ and $V \subset \mathbb{R}^n$ and $\phi : U \rightarrow V$. Then (U, ϕ) is said to be a **n -coordinate chart on M** if

- U is open in M
- V is open in \mathbb{R}^n or V is open in \mathbb{H}^n
- ϕ is a homeomorphism

We denote the set of all n -coordinate charts on M by $X^n(M)$.

Definition 3.1.0.6. Let M be a topological space and $n \in \mathbb{N}$. Then M is said to be **locally Euclidean of dimension n** if for each $p \in M$, there exists $(U, \phi) \in X^n(M)$ such that $p \in U$.

Definition 3.1.0.7. Let M be a topological space and $n \in \mathbb{N}$. Then M is said to be an **n -dimensional topological manifold** if

1. M is Hausdorff
2. M is second-countable
3. M is locally Euclidean of dimension n

Exercise 3.1.0.8. Let M be an n -dimensional topological manifold, $(U, \phi) \in X(M)$ and $U' \subset U$. If U' is open in M , then $(U', \phi|_{U'}) \in X^n(U)$.

Proof. Suppose that U' is open in M . Set $\phi' = \phi|_{U'}$.

1. Since $U' \subset U$ and U' is open in M , we have that U' is open in U
2. By assumption $\phi(U)$ is open in \mathbb{R}^n or $\phi(U)$ is open in \mathbb{H}^n . Since ϕ is a homeomorphism and U' is open in U , we have that $\phi(U')$ is open in $\phi(U)$. Therefore $\phi'(U')$ is open in \mathbb{R}^n or $\phi'(U')$ is open in \mathbb{H}^n .
3. Since $\phi : U \rightarrow V$ is a homeomorphism, $\phi' : U' \rightarrow \phi'(U')$ is a homeomorphism.

So $(U', \phi') \in X^n(U)$. □

Exercise 3.1.0.9. Topological Open Submanifolds:

Let M be an n -dimensional topological manifold and $U \subset M$ open. Then U is an n -dimensional topological manifold.

Proof.

1. Since M is Hausdorff, U is Hausdorff.
2. M is second-countable, U is second countable.
3. Let $p \in U$. Since then there exists $(V, \psi) \in X^n(M)$ such that $p \in V$. Set $V' = U \cap V$ and $\psi' = \psi|_{U \cap V}$. The previous exercise implies that $(V', \psi') \in X^n(U)$. Therefore U is locally Euclidean of dimension n .

Hence U is an n -dimensional topological manifold. □

Theorem 3.1.0.10. Topological Invariance of Dimension:

Let M be an n -dimensional topological manifold and N a p -dimensional topological manifold. If M and N are homeomorphic, then $n = p$.

Note 3.1.0.11. In light of the previous theorem, unless specified otherwise, we write $X(M)$ in place of $X^n(M)$ and refer to n -coordinate charts as coordinate charts.

Definition 3.1.0.12. Let M be an n -dimensional topological manifold and $(U, \phi) \in X(M)$. Then (U, ϕ) is said to be an

- **interior chart** if $\phi(U)$ is open in \mathbb{R}^n
- **boundary chart** if $\phi(U)$ is open in \mathbb{H}^n and $\phi(U) \cap \partial\mathbb{H}^n \neq \emptyset$

We denote the set of all interior charts on M and the set of all boundary charts on M by $X_{\text{Int}}(M)$ and $X_{\partial}(M)$ respectively.

Exercise 3.1.0.13. Let M be an n -dimensional topological manifold. Then

1. $X(M) = X_{\text{Int}}(M) \cup X_{\partial}(M)$
2. $X_{\text{Int}}(M) \cap X_{\partial}(M) = \emptyset$

Proof.

1. By definition, $X_{\text{Int}}(M) \cup X_{\partial}(M) \subset X(M)$. Let $(U, \phi) \in X(M)$. Since (U, ϕ) is a coordinate chart on M , $\phi(U)$ is open in \mathbb{R}^n or $\phi(U)$ is open in \mathbb{H}^n . If $\phi(U)$ is open in \mathbb{R}^n , then

$$\begin{aligned} (U, \phi) &\in X_{\text{Int}}(M) \\ &\subset X_{\text{Int}}(M) \cup X_{\partial}(M) \end{aligned}$$

Suppose that $\phi(U)$ is open in \mathbb{H}^n . If $\phi(U) \cap \partial\mathbb{H}^n = \emptyset$, then $\phi(U)$ is open in \mathbb{R}^n and

$$\begin{aligned} (U, \phi) &\in X_{\text{Int}}(M) \\ &\subset X_{\text{Int}}(M) \cup X_{\partial}(M) \end{aligned}$$

Suppose that $\phi(U) \cap \partial\mathbb{H}^n \neq \emptyset$. Then

$$\begin{aligned} (U, \phi) &\in X_{\partial}(M) \\ &\subset X_{\text{Int}}(M) \cup X_{\partial}(M) \end{aligned}$$

So $X(M) \subset X_{\text{Int}}(M) \cup X_{\partial}(M)$.

2. For the sake of contradiction, suppose that $X_{\text{Int}}(M) \cup X_{\partial}(M) \neq \emptyset$. Then there exists $(U, \phi) \in X(M)$ such that $(U, \phi) \in X_{\text{Int}}(M)$ and $(U, \phi) \in X_{\partial}(M)$. Therefore $\phi(U)$ is open in \mathbb{R}^n , $\phi(U)$ is open in \mathbb{H}^n and $\phi(U) \cap \partial\mathbb{H}^n \neq \emptyset$. Since $\phi(U)$ is open in \mathbb{R}^n and $\phi(U) \subset \mathbb{H}^n$, $\phi(U) \subset \text{Int } \mathbb{H}^n$ and therefore $\phi(U) \cap \partial\mathbb{H}^n = \emptyset$ which is a contradiction.

□

Definition 3.1.0.14. Let M be an n -dimensional topological manifold. We define the

- **interior** of M , denoted $\text{Int } M$, by

$$\text{Int } M = \{p \in M : \text{there exists } (U, \phi) \in X_{\text{Int}}(M) \text{ such that } p \in U\}$$

- **boundary** of M , denoted ∂M , by

$$\partial M = \{p \in M : \text{there exists } (V, \psi) \in X_{\partial}(M) \text{ such that } p \in V \text{ and } \psi(p) \in \partial\mathbb{H}^n\}$$

Exercise 3.1.0.15. Let M be an n -dimensional topological manifold, $(U, \phi) \in X_{\partial}(M)$ and $p \in U$. If $\phi(p) \notin \partial\mathbb{H}^n$, then $p \in \text{Int } M$.

Proof. Suppose that $\phi(p) \notin \partial\mathbb{H}^n$. Then $\phi(p) \in \text{Int } \mathbb{H}^n$. Hence there exists $B' \subset \phi(U)$ such that B' is open in \mathbb{R}^n and $\phi(p) \in B'$. Set $U' = \phi^{-1}(B')$ and $\phi' = \phi|_{U'}$. Then U' is open in M and $\phi' : U' \rightarrow B'$ is a homeomorphism. Hence $(U', \phi') \in X_{\text{Int}}(M)$. Since $\phi(p) \in B'$, we have that $p \in U'$. By definition, $p \in \text{Int } M$. □

Exercise 3.1.0.16. Let M be an n -dimensional topological manifold. Then

1. $M = \text{Int } M \cup \partial M$
2. $\text{Int } M \cap \partial M = \emptyset$

Hint: simply connected

Proof.

1. By definition, $\text{Int } M \cup \partial M \subset M$. Let $p \in M$. Since M is a manifold, there exists $(U, \phi) \in X(M)$ such that $p \in U$. A previous exercise implies that $(U, \phi) \in X_{\text{Int}}(M) \cup X_{\partial}(M)$. If $(U, \phi) \in X_{\text{Int}}(M)$, then by definition,

$$\begin{aligned} p &\in \text{Int } M \\ &\subset \text{Int } M \cup \partial M \end{aligned}$$

Suppose that $(U, \phi) \in X_{\partial}(M)$. If $\phi(p) \in \partial \mathbb{H}^n$, then by definition,

$$\begin{aligned} p &\in \partial M \\ &\subset \text{Int } M \cup \partial M \end{aligned}$$

Suppose that $\phi(p) \notin \partial \mathbb{H}^n$. The previous exercise implies that $p \in \text{Int } M$. Therefore,

$$\begin{aligned} p &\in \text{Int } M \\ &\subset \text{Int } M \cup \partial M \end{aligned}$$

Hence $M \subset \text{Int } M \cup \partial M$.

2. For the sake of contradiction, suppose that $\text{Int } M \cap \partial M \neq \emptyset$. Then there exists $p \in M$ such that $p \in \text{Int } M \cap \partial M$. By definition, there exists $(U, \phi) \in X_{\text{Int}}(M)$, $(V, \psi) \in X_{\partial}(M)$ such that $p \in U \cap V$ and $\psi(p) \in \partial \mathbb{H}^n$. Note that $\psi(U \cap V)$ is open in \mathbb{H}^n , $\phi(U \cap V)$ is open in \mathbb{R}^n and $\phi|_{U \cap V} \circ (\psi|_{U \cap V})^{-1} : \psi^{-1}(U \cap V) \rightarrow \phi(U \cap V)$ is a homeomorphism. Since $\psi(U \cap V)$ is open in \mathbb{H}^n , there exists an $B_{\psi} \subset \psi(U \cap V)$ such that B_{ψ} is open in \mathbb{H}^n , B_{ψ} is simply connected and $\psi(p) \in B_{\psi}$. Set $B_{\phi} = \phi \circ \psi^{-1}(B_{\psi})$. Since $\phi(U \cap V)$ is open in \mathbb{R}^n , B_{ϕ} is open in \mathbb{R}^n . Since B_{ψ} is simply connected and $\phi|_{U \cap V} \circ (\psi|_{U \cap V})^{-1} : \psi^{-1}(U \cap V) \rightarrow \phi(U \cap V)$ is a homeomorphism, B_{ϕ} is simply connected. Set $B'_{\phi} = B_{\phi} \setminus \{\phi(p)\}$ and $B'_{\psi} = B_{\psi} \setminus \{\psi(p)\}$. Then $\phi \circ \psi^{-1} : B'_{\psi} \rightarrow B'_{\phi}$ is a homeomorphism. Since $\psi(p) \in \partial \mathbb{H}^n$, B'_{ψ} is simply connected. Since B_{ϕ} is open in \mathbb{R}^n , B'_{ϕ} is not simply connected. This is a contradiction since B'_{ϕ} is homeomorphic to B'_{ψ} . So $\partial M \cap \text{Int } M = \emptyset$.

□

Exercise 3.1.0.17. Let M be an n -dimensional topological manifold, $(U, \phi) \in X(M)$ and $p \in U$. If $p \in \partial M$, then $(U, \phi) \in X_{\partial}(M)$.

Hint: simply connected

Proof. Suppose that $p \in \partial M$. Then there exists a $(V, \psi) \in X_{\partial}(M)$ such that $p \in V$ and $\psi(p) \in \partial \mathbb{H}^n$. Note that $\psi(U \cap V)$ is open in \mathbb{H}^n , $\phi(U \cap V)$ is open in \mathbb{R}^n and $\phi|_{U \cap V} \circ (\psi|_{U \cap V})^{-1} : \psi^{-1}(U \cap V) \rightarrow \phi(U \cap V)$ is a homeomorphism.

Since $\psi(U \cap V)$ is open in \mathbb{H}^n , there exists $B_{\psi} \subset \psi(U \cap V)$ such that B_{ψ} is open in \mathbb{H}^n , B_{ψ} is simply connected and $\psi(p) \in B_{\psi}$. Set $B_{\phi} = \phi \circ \psi^{-1}(B_{\psi})$.

For the sake of contradiction, suppose that $(U, \phi) \in X_{\text{Int}}(M)$. Then $\phi(U)$ is open in \mathbb{R}^n . Hence $\phi(U \cap V)$ is open in \mathbb{R}^n and B_{ϕ} is open in \mathbb{R}^n . Since $\phi|_{U \cap V} \circ (\psi|_{U \cap V})^{-1} : \psi^{-1}(U \cap V) \rightarrow \phi(U \cap V)$ is a homeomorphism, B_{ϕ} is simply connected. Set $B'_{\phi} = B_{\phi} \setminus \{\phi(p)\}$ and $B'_{\psi} = B_{\psi} \setminus \{\psi(p)\}$. Since $\psi(p) \in \partial \mathbb{H}^n$, B'_{ψ} is simply connected. Since B_{ϕ} is open in \mathbb{R}^n , B'_{ϕ} is not simply connected. This is a contradiction since B'_{ϕ} is homeomorphic to B'_{ψ} . So $(U, \phi) \notin X_{\text{Int}}(M)$. Since $(X_{\text{Int}}(M))^c = X_{\partial}(M)$, we have that $(U, \phi) \in X_{\partial}(M)$.

□

Exercise 3.1.0.18. Let M be an n -dimensional topological manifold, $(U, \phi) \in X_{\partial}(M)$ and $p \in U$. Then

1. $p \in \partial M$ iff $\phi(p) \in \partial \mathbb{H}^n$
2. $p \in \text{Int } M$ iff $\phi(p) \in \text{Int } \mathbb{H}^n$

Proof.

1. Suppose that $p \in \partial M$. For the sake of contradiction, suppose that $\phi(p) \notin \partial \mathbb{H}^n$. Then $\phi(p) \in \text{Int } \mathbb{H}^n$. Hence there exists $B' \subset \phi(U)$ such that B' is open in \mathbb{R}^n and $\phi(p) \in B'$. Set $U' = \phi^{-1}(B')$ and $\phi' = \phi|_{U'}$. Then $p \in U'$ and $(U', \phi') \in X_{\text{Int}}(M)$. Since $p \in U'$, the previous exercise implies that $(U', \phi') \in X_{\partial}(M)$. This is a contradiction since $X_{\text{Int}}(M) \cap X_{\partial}(M) = \emptyset$. So $\phi(p) \in \partial \mathbb{H}^n$. Conversely, suppose that $\phi(p) \in \partial \mathbb{H}^n$. By definition, $p \in \partial M$.
2. A previous exercise implies that $\text{Int } M = (\partial M)^c$. Part (1) implies that

$$\begin{aligned} p &\in (\partial M)^c \\ &= \text{Int } M \end{aligned}$$

if and only if

$$\begin{aligned} \phi(p) &\in (\partial \mathbb{H}^n)^c \\ &= \text{Int } \mathbb{H}^n \end{aligned}$$

□

Exercise 3.1.0.19. Let M be an n -dimensional topological manifold and $p \in M$. Then $p \in \partial M$ iff for each $(U, \phi) \in X(M)$, $p \in U$ implies that $(U, \phi) \in X_{\partial}(M)$ and $\phi(p) \in \partial \mathbb{H}^n$.

Proof. Suppose that $p \in \partial M$. Let $(U, \phi) \in X(M)$. Suppose that $p \in U$. The previous two exercises imply that $(U, \phi) \in X_{\partial}(M)$ and $\phi(p) \in \partial \mathbb{H}^n$.

Conversely, suppose that for each $(U, \phi) \in X(M)$, $p \in U$ implies that $(U, \phi) \in X_{\partial}(M)$ and $\phi(p) \in \partial \mathbb{H}^n$. Since M is a manifold, there exists $(U, \phi) \in X(M)$ such that $p \in U$. By assumption, $(U, \phi) \in X_{\partial}(M)$ and $\phi(p) \in \partial \mathbb{H}^n$. Since $(U, \phi) \in X_{\partial}(M)$, a previous exercise implies that $\phi(p) \in \partial \mathbb{H}^n$ iff $p \in \partial M$. Since $\phi(p) \in \partial \mathbb{H}^n$, we have that $p \in \partial M$. □

Exercise 3.1.0.20. Let M be an n -dimensional topological manifold. Let $(U, \phi) \in X_{\partial}(M)$. Then

1. $\phi(U \cap \partial M) = \phi(U) \cap \partial \mathbb{H}^n$
2. $\phi(U \cap \text{Int } M) = \phi(U) \cap \text{Int } \mathbb{H}^n$

Proof.

1. Since $(U, \phi) \in X_{\partial}(M)$, a previous exercise implies that for each $p \in U$, $p \in \partial M$ iff $\phi(p) \in \partial \mathbb{H}^n$. Let $q \in \phi(U \cap \partial M)$. Then there exists $p \in U \cap \partial M$ such that $\phi(p) = q$. Since $p \in \partial M$, $\phi(p) \in \partial \mathbb{H}^n$. Hence

$$\begin{aligned} q &= \phi(p) \\ &\in \phi(U) \cap \partial \mathbb{H}^n \end{aligned}$$

Since $q \in \phi(U \cap \partial M)$ is arbitrary, $\phi(U \cap \partial M) \subset \phi(U) \cap \partial \mathbb{H}^n$.

Let $q \in \phi(U) \cap \partial \mathbb{H}^n$. Then there exists $p \in U$ such that $q = \phi(p)$. Since $\phi(p) \in \partial \mathbb{H}^n$, we have that $p \in \partial M$. Hence $p \in U \cap \partial M$ and

$$\begin{aligned} q &= \phi(p) \\ &\in \phi(U \cap \partial M) \end{aligned}$$

Since $q \in \phi(U) \cap \partial \mathbb{H}^n$ is arbitrary, $\phi(U) \cap \partial \mathbb{H}^n \subset \phi(U \cap \partial M)$. Thus $\phi(U \cap \partial M) = \phi(U) \cap \partial \mathbb{H}^n$.

2. Since $(U, \phi) \in X_{\partial}(M)$, a previous exercise implies that for each $p \in U$, $p \in \text{Int } M$ iff $\phi(p) \in \text{Int } \mathbb{H}^n$. Let $q \in \phi(U \cap \text{Int } M)$. Then there exists $p \in U \cap \text{Int } M$ such that $\phi(p) = q$. Since $p \in \text{Int } M$, $\phi(p) \in \text{Int } \mathbb{H}^n$. Hence

$$\begin{aligned} q &= \phi(p) \\ &\in \phi(U) \cap \text{Int } \mathbb{H}^n \end{aligned}$$

Since $q \in \phi(U \cap \text{Int } M)$ is arbitrary, $\phi(U \cap \text{Int } M) \subset \phi(U) \cap \text{Int } \mathbb{H}^n$.

Let $q \in \phi(U) \cap \text{Int } \mathbb{H}^n$. Then there exists $p \in U$ such that $q = \phi(p)$. Since $\phi(p) \in \text{Int } \mathbb{H}^n$, we have that $p \in \text{Int } M$. Hence $p \in U \cap \text{Int } M$ and

$$\begin{aligned} q &= \phi(p) \\ &\in \phi(U \cap \text{Int } M) \end{aligned}$$

Since $q \in \phi(U) \cap \text{Int } \mathbb{H}^n$ is arbitrary, $\phi(U) \cap \text{Int } \mathbb{H}^n \subset \phi(U \cap \text{Int } M)$. Thus $\phi(U \cap \text{Int } M) = \phi(U) \cap \text{Int } \mathbb{H}^n$. □

Exercise 3.1.0.21. Let M be an n -dimensional topological manifold. Then

1. $\text{Int } M$ is open
2. ∂M is closed

Proof.

1. Let $p \in \text{Int } M$. Then there exists $(U, \phi) \in X_{\text{Int}}(M)$ such that $p \in U$. By definition of coordinate charts, U is open. By definition of $\text{Int } M$, for each $q \in U$, $q \in \text{Int } M$. Hence $U \subset \text{Int } M$. Since $p \in \text{Int } M$ is arbitrary, we have that for each $p \in \text{Int } M$, there exists $U \subset \text{Int } M$ such that U is open. Hence $\text{Int } M$ is open.
2. Since $\partial M = (\text{Int } M)^c$, and $\text{Int } M$ is open, we have that ∂M is closed. □

Exercise 3.1.0.22. Let M be an n -dimensional topological manifold, $(U, \phi) \in X_{\partial}(M)$ and $\lambda : \partial \mathbb{H}^n \rightarrow \mathbb{R}^{n-1}$ a homeomorphism. Define $\bar{U} \subset \partial M$ and $\bar{\phi} : \bar{U} \rightarrow \lambda(\phi(\bar{U}))$ by $\bar{U} = U \cap \partial M$ and $\bar{\phi} = \lambda|_{\phi(\bar{U})} \circ \phi|_{\bar{U}}$. Then $(\bar{U}, \bar{\phi}) \in X^{n-1}(\partial M)$.

Proof.

1. Since U is open in M , $\bar{U} = U \cap \partial M$ is open in ∂M .
2. Since $(U, \phi) \in X_{\partial}(M)$, $\phi(U)$ is open in \mathbb{H}^n . A previous exercise implies that $\phi(\bar{U}) = \phi(U) \cap \partial \mathbb{H}^n$ which is open in $\partial \mathbb{H}^n$. Since $\lambda : \partial \mathbb{H}^n \rightarrow \mathbb{R}^{n-1}$ is a homeomorphism, we have that $\lambda(\phi(\bar{U}))$ is open in \mathbb{R}^{n-1} .
3. Since $\phi|_{\bar{U}} : \bar{U} \rightarrow \phi(U) \cap \partial \mathbb{H}^n$ and $\lambda|_{\phi(\bar{U})} : \phi(\bar{U}) \rightarrow \lambda(\phi(\bar{U}))$ are homeomorphisms, we have that $\bar{\phi} = \lambda|_{\phi(\bar{U})} \circ \phi|_{\bar{U}}$ is a homeomorphism.

Hence $(\bar{U}, \bar{\phi}) \in X^{n-1}(\partial M)$. □

Exercise 3.1.0.23. Let M be an n -dimensional topological manifold. Then

1. ∂M is an $(n-1)$ -dimensional topological manifold
2. $\partial(\partial M) = \emptyset$

Proof.

1. (a) Since M is Hausdorff, ∂M is Hausdorff.
 (b) Since M is second-countable, ∂M is second countable.
 (c) Let $p \in \partial M$. Then there exists $(U, \phi) \in X_{\partial}(M)$ such that $\phi(p) \in \partial \mathbb{H}^n$. Choose $\lambda : \partial \mathbb{H}^n \rightarrow \mathbb{R}^{n-1}$ such that λ is a homeomorphism and define $(\bar{U}, \bar{\phi}) \in X^{n-1}(\partial M)$ as in the previous exercise. Then $p \in \bar{U}$. Hence ∂M is locally Euclidean of dimension $n-1$.

Hence ∂M is an $(n-1)$ -dimensional topological manifold.

2. Let $p \in \partial M$. Then there exists $(U, \phi) \in X_{\partial}(M)$ such that $\phi(p) \in \partial \mathbb{H}^n$. Choose $\lambda : \partial \mathbb{H}^n \rightarrow \mathbb{R}^{n-1}$ such that λ is a homeomorphism and define $(\bar{U}, \bar{\phi}) \in X(\partial M)$ as in the previous exercise. Then $p \in \bar{U}$. Since $\bar{\phi}(\bar{U})$ is open in \mathbb{R}^{n-1} , we have that $(\bar{U}, \bar{\phi}) \in X_{\text{Int}}(\partial M)$. Therefore $p \in \text{Int}(\partial M)$. Since $p \in \partial M$ is arbitrary, $\text{Int } \partial M = \partial M$. Hence

$$\begin{aligned}\partial(\partial M) &= (\text{Int}(\partial M))^c \\ &= (\partial M)^c \\ &= \emptyset\end{aligned}$$

□

Exercise 3.1.0.24. Let M be an n -dimensional topological manifold and $U \subset M$ open. Then $\partial U = \partial M \cap U$.

Proof. Let $p \in \partial U$. Then there exists $(V, \psi) \in X_{\partial}(U)$ such that $p \in V$ and $\psi(p) \in \partial \mathbb{H}^n$.

FINISH!!!

□


Exercise 3.1.0.25. Graph of Continuous Function:

Let $f \in C(\mathbb{R})$. Set $M = \{(x, y) \in \mathbb{R}^2 : f(x) = y\}$ (i.e. the graph of f). Then M is a 1-dimensional manifold.

Proof. Set $U = \mathbb{R}$ and define $\phi : U \rightarrow M$ by $\phi(x) = (x, f(x))$. Then $\phi^{-1} = \pi_1$. Since f is continuous, ϕ is continuous. Since π_1 is continuous, ϕ is a homeomorphism. □

Exercise 3.1.0.26. Nodal Cubic:

Let $M = \{(x, y) \in \mathbb{R}^2 : y^2 = x^2 - x^3\}$. We equip M with the subspace topology.



Then M is not a 1-dimensional topological manifold.

Hint: connected components

Proof. Suppose that M is a 1-dimensional manifold. Set $p = (0,0)$. Then there exists $(U, \phi) \in X(M)$ such that $p \in U$. Since $\phi(U)$ is open (in \mathbb{R} or \mathbb{H}), there exists a $B \subset \phi(U)$ such that B is open (in \mathbb{R} or \mathbb{H}), B is connected and $\phi(p) \in B$. Set $V = \phi^{-1}(B)$, $V' = V \setminus \{p\}$ and $B' = B \setminus \{\phi(p)\}$. Then $\phi : V \rightarrow B$ and $\phi' : V' \rightarrow B'$ are homeomorphisms. Since B is open (in \mathbb{R} or \mathbb{H}) and connected, B' has at most two connected components. Then V' This is a contradiction since V' has four connected components and B' and V' are homeomorphic. □

3.2 Smooth Manifolds

Definition 3.2.0.1. Let M be an n -dimensional topological manifold and $(U, \phi), (V, \psi) \in X(M)$. Then (U, ϕ) and (V, ψ) are said to be **smoothly compatible** if

$$\psi|_{U \cap V} \circ (\phi|_{U \cap V})^{-1} : \phi(U \cap V) \rightarrow \psi(U \cap V) \text{ is a diffeomorphism}$$

Definition 3.2.0.2. Let M be an n -dimensional topological manifold.

1. Let $\mathcal{A} \subset X(M)$. Then \mathcal{A} is said to be an **atlas on M** if $\bigcup_{(U, \phi) \in \mathcal{A}} U = M$.
2. Let \mathcal{A} be an atlas on M . Then \mathcal{A} is said to be **smooth** if for each $(U, \phi), (V, \psi) \in \mathcal{A}$, (U, ϕ) and (V, ψ) are smoothly compatible.
3. Let \mathcal{A} be a smooth atlas on M . Then \mathcal{A} is said to be **maximal** if for each smooth atlas \mathcal{B} on M , $\mathcal{A} \subset \mathcal{B}$ implies that $\mathcal{A} = \mathcal{B}$. A maximal smooth atlas on M is called a **smooth structure on M** .
4. Let \mathcal{A} be an atlas on M . Then (M, \mathcal{A}) is said to be an **n -dimensional smooth manifold** if \mathcal{A} is a smooth structure on M .

Exercise 3.2.0.3. Let M be an n -dimensional topological manifold and \mathcal{B} a smooth atlas on M . Then there exists a unique smooth structure \mathcal{A} on M such that $\mathcal{B} \subset \mathcal{A}$.

Proof. Define

$$\mathcal{A} = \{(U, \phi) \in X(M) : \text{for each } (V, \psi) \in \mathcal{B}, (U, \phi) \text{ and } (V, \psi) \text{ are smoothly compatible}\}$$

Clearly $\mathcal{B} \subset \mathcal{A}$. Let (U, ϕ) and $(V, \psi) \in \mathcal{A}$. Define $F : \phi(U \cap V) \rightarrow \psi(U \cap V)$ by

$$F = \psi|_{U \cap V} \circ (\phi|_{U \cap V})^{-1}$$

Let $q \in \phi(U \cap V)$. Set $p = \phi^{-1}(q)$. Since $p \in U \cap V \subset M$, there exists $(W, \chi) \in \mathcal{B}$ such that $p \in W$. By definition of \mathcal{A} , $\psi|_{W \cap V} \circ (\chi|_{W \cap V})^{-1} : \chi(W \cap V) \rightarrow \psi(W \cap V)$ and $\chi|_{U \cap W} \circ (\phi|_{U \cap W})^{-1} : \phi(U \cap W) \rightarrow \chi(U \cap W)$ are diffeomorphisms. Set $N = U \cap W \cap V$. Then $q \in \phi(N) \subset \phi(U \cap V)$ and

$$\begin{aligned} F|_{\phi(N)} &= \psi|_N \circ (\phi|_N)^{-1} \\ &= [\psi|_N \circ (\chi|_N)^{-1}] \circ [\chi|_N \circ (\phi|_N)^{-1}] \end{aligned}$$

is a diffeomorphism. Thus, for each $q \in \phi(U \cap V)$, there exists $N' \subset \phi(U \cap V)$ such that $F|_{N'}$ is a diffeomorphism. Hence F is a diffeomorphism and $(U, \phi), (V, \psi)$ are smoothly compatible. Therefore \mathcal{A} is a smooth atlas.

To see that \mathcal{A} is maximal, let \mathcal{B}' be a smooth atlas on M . Suppose that $\mathcal{A} \subset \mathcal{B}'$ and let $(U, \phi) \in \mathcal{B}'$. By definition, for each chart $(V, \psi) \in \mathcal{B}'$, (U, ϕ) and (V, ψ) are smoothly compatible. Since $\mathcal{B} \subset \mathcal{A} \subset \mathcal{B}'$, we have that $(U, \phi) \in \mathcal{A}$. So $\mathcal{A} = \mathcal{B}'$ and \mathcal{A} is a maximal smooth atlas on M . \square

Exercise 3.2.0.4. Let (M, \mathcal{A}) be an n -dimensional smooth manifold, $(U, \phi) \in \mathcal{A}$ and $U' \subset U$. If U' is open, then $(U', \phi|_{U'}) \in \mathcal{A}$.

Proof. Set $\phi' = \phi|_{U'}$. A previous exercise implies that $(U', \phi') \in X(U)$. Define $\mathcal{B} = \mathcal{A} \cup \{(U', \phi')\}$. Let $(V, \psi) \in \mathcal{B}$. If $(V, \psi) = (U', \phi')$, then

$$\phi' \circ \psi^{-1} = \text{id}_{U'}$$

which is a diffeomorphism. Thus $(U', \phi'), (V, \psi)$ are smoothly compatible. Suppose that $(V, \psi) \in \mathcal{A}$. Since \mathcal{A} is smooth, $\psi|_{U \cap V} \circ (\phi|_{U \cap V})^{-1} : \phi(U \cap V) \rightarrow \psi(U \cap V)$ is a diffeomorphism. Therefore $\psi|_{U' \cap V} \circ (\phi'|_{U' \cap V})^{-1} : \phi'(U' \cap V) \rightarrow \psi(U' \cap V)$ is a diffeomorphism and $(U', \phi'), (V, \psi)$ are smoothly compatible. Since $(V, \psi) \in \mathcal{A}$ is arbitrary, \mathcal{B} is smooth. Since \mathcal{A} is maximal and $\mathcal{A} \subset \mathcal{B}$, we have that $\mathcal{A} = \mathcal{B}$ and $(U', \phi') \in \mathcal{A}$. \square

Exercise 3.2.0.5. Let (M, \mathcal{A}) be a n -dimensional smooth manifold and $U \subset M$ open. Set $\mathcal{B} = \{(U', \phi') \in \mathcal{A} : U' \subset U\}$. Then \mathcal{B} is a smooth atlas on U .

Proof. Since $\mathcal{B} \subset \mathcal{A}$, we have that $\mathcal{B} \subset X(U)$. Let $p \in U$. Then there exists $(V, \psi) \in \mathcal{A}$ such that $p \in V$. The previous exercise implies that $(U \cap V, \psi|_{U \cap V}) \in \mathcal{A}$. By definition, $(U \cap V, \psi|_{U \cap V}) \in \mathcal{B}$. Since $p \in U$ is arbitrary, we have that for each $p \in U$, there exists $(U', \phi) \in \mathcal{B}$ such that $p \in U'$. Hence \mathcal{B} is an atlas on M . Since $\mathcal{B} \subset \mathcal{A}$, \mathcal{B} is smooth. \square

Definition 3.2.0.6. Smooth Open Submanifold:

Let (M, \mathcal{A}) be an n -dimensional smooth manifold and $U \subset M$ open. A previous exercise implies that U is an n -dimensional topological manifold. We define $\mathcal{A}|_U \subset X(U)$ to be the smooth structure on U such that $\{(U', \phi') \in \mathcal{A} : U' \subset U\} \subset \mathcal{A}|_U$. Then $(U, \mathcal{A}|_U)$ is said to be a **smooth open submanifold of (M, \mathcal{A})** .

Exercise 3.2.0.7. Let (M, \mathcal{A}) be a smooth n -dimensional manifold and $U \subset M$ open. Then $\partial U = \partial M \cap U$.

Proof. \square

Definition 3.2.0.8. Let (M, \mathcal{A}) be a n -dimensional smooth manifold. Define $\pi : \partial \mathbb{H}^n \rightarrow \mathbb{R}^{n-1}$ by

$$\pi(x_1, \dots, x_{n-1}, 0) = (x_1, \dots, x_{n-1})$$

For $(U, \phi) \in \mathcal{A}$, set $\bar{U} = U \cap \partial M$ and $\bar{\phi} = \pi \circ \phi|_{\bar{U}}$.

Define $\mathcal{A}|_{\partial M} = \{(\bar{U}, \bar{\phi}) : (U, \phi) \in \mathcal{A} \text{ and } U \cap \partial M \neq \emptyset\}$.

Exercise 3.2.0.9. Let (M, \mathcal{A}) be a n -dimensional smooth manifold. Then

1. $\mathcal{A}|_{\partial M}$ is a smooth structure on ∂M
2. $(\partial M, \mathcal{A}|_{\partial M})$ is a $(n-1)$ -dimensional smooth manifold
3. $\partial(\partial M) = \emptyset$

Proof.

1. A previous exercise implies that π is a homeomorphism and it is clear that π is a diffeomorphism. Let $p \in \partial M$. Then there exists a boundary chart (U, ϕ) on M such that $p \in U$. Thus $p \in \bar{U}$ and the previous exercise implies that

$$\begin{aligned} \phi(\bar{U}) &= \phi(U \cap \partial M) \\ &= \phi(U) \cap \partial \mathbb{H}^n \end{aligned}$$

Since U is open in M and $\phi(U)$ is open in \mathbb{H}^n , we have that \bar{U} is open in ∂M and $\phi(\bar{U})$ is open in $\partial \mathbb{H}^n$ which implies that $\pi(\phi(\bar{U}))$ is open in \mathbb{R}^{n-1} . Since $\phi : U \rightarrow \phi(U)$ is a homeomorphism, $\phi|_{\bar{U}} : \bar{U} \rightarrow \phi(\bar{U})$ is a homeomorphism. Hence $\bar{\phi} : \bar{U} \rightarrow \pi(\phi(\bar{U}))$ is a homeomorphism and $(\bar{U}, \bar{\phi})$ is an interior chart on ∂M . Therefore $\mathcal{A}|_{\partial M}$ is an atlas on ∂M . Let $(\bar{U}, \bar{\phi}), (\bar{V}, \bar{\psi}) \in \mathcal{A}|_{\partial M}$. Then

$$\bar{\phi} \circ \bar{\psi}^{-1} = \pi \circ \phi \circ \psi^{-1} \circ \pi^{-1}$$

which is a diffeomorphism. So $(\bar{U}, \bar{\phi})$ and $(\bar{V}, \bar{\psi})$ are smoothly compatible. Hence \mathcal{A} is smooth.

2. Subspaces of Hausdorff, second countable spaces are Hausdorff and second countable, so ∂M with the subspace topology is Hausdorff and second countable. Since $\mathcal{A}|_{\partial M}$ is an atlas on ∂M , $(\partial M, \mathcal{A}|_{\partial M})$ is a $(n-1)$ -dimensional manifold.
3. Since for each $(\bar{U}, \bar{\phi}) \in \mathcal{A}|_{\partial M}$, $\bar{\phi}(\bar{U})$ is open in \mathbb{R}^n , $(\bar{U}, \bar{\phi})$ is an interior chart. Hence $\partial(\partial M) = \emptyset$. \square

Note 3.2.0.10. For the rest of this section, we assume that (M, \mathcal{A}) is a n -dimensional smooth manifold and we denote the standard coordinate functions on \mathbb{R}^n by u^1, \dots, u^n . For a coordinate chart $(U, \phi) \in \mathcal{A}$ and $i \in \{1, \dots, n\}$, we will typically denote the i th coordinate of ϕ by x^i , that is, $x^i = u^i(\phi)$.

Exercise 3.2.0.11. Let (M, \mathcal{A}) be a n -dimensional smooth manifold and $S \subset M$ open. For $(U, \phi) \in \mathcal{A}$, define $U' \subset S$ and $\phi' : \tilde{U} \rightarrow \phi(U')$ by $U' = U \cap S$ and $\phi' = \phi|_{U \cap S}$. Set $\mathcal{A}' = \{(U', \phi') : (U, \phi) \in \mathcal{A}\}$. Then (S, \mathcal{A}') is a n -dimensional smooth manifold.

Proof. Since M is Hausdorff and second countable, so is S . Clearly $S = \bigcup_{(U', \phi') \in \mathcal{A}'} U'$ and for each $(U', \phi') \in \mathcal{A}'$, ϕ' is a homeomorphism and $\phi'(U') \subset \mathbb{H}^n$. Let $(U', \phi'), (V', \psi') \in \mathcal{A}'$. Then there exist $(U, \phi), (V, \psi) \in \mathcal{A}$ such that $U' = U \cap S$, $V' = V \cap S$. Then $\psi' \circ \phi'^{-1} : \phi(U \cap V \cap S) \rightarrow \psi(U \cap V \cap S)$ is given by

$$\psi' \circ \phi'^{-1} : \phi(U' \cap V') \rightarrow \psi(U' \cap V') = \psi \circ \phi^{-1} : \phi(U \cap V \cap S) \rightarrow \psi(U \cap V \cap S)$$

which is smooth. So \mathcal{A}' is a smooth atlas. Let \mathcal{B}' be a smooth atlas on S . Suppose that $\mathcal{A}' \subset \mathcal{B}'$. □

Exercise 3.2.0.12. Let (M, \mathcal{A}) a smooth manifold and $U \subset M$ open. For $(V, \psi) \in \mathcal{A}$, define $\bar{V} \subset M$ and $\bar{\psi} : \bar{V} \rightarrow \psi(\bar{V})$ by $\bar{V} = V \cap U$ and $\bar{\psi} = \psi|_{V \cap U}$. Define

$$\mathcal{A} \cap U = \{(\bar{V}, \bar{\psi}) : (V, \psi) \in \mathcal{A}\}$$

Then

1. \mathcal{A}_U is an atlas on U
2. \mathcal{A}_U is a smooth structure on U
3. (U, \mathcal{A}_U) is an n -dimensional smooth manifold.

Definition 3.2.0.13. Let (M, \mathcal{A}) an n -dimensional smooth manifold and $U \subset M$ open. Define (U, \mathcal{A}_U) as in the previous exercise.

3.3 Smooth Maps

Definition 3.3.0.1. Let $f : M \rightarrow \mathbb{R}$. Then f is said to be smooth if for each coordinate chart $(U, \phi) \in \mathcal{A}$, $f \circ \phi^{-1}$ is smooth. The set of all smooth functions on M is denoted $C^\infty(M)$.

Exercise 3.3.0.2. We have that $C^\infty(M)$ is a vector space.

Proof. Clear. □

Definition 3.3.0.3. Let (M, \mathcal{A}) be a smooth manifold, $(U, \phi) \in \mathcal{A}$. We define

$$C^\infty(U) = \{f : U \rightarrow \mathbb{R} : f \circ \phi^{-1} \in C^\infty(\phi(U))\}$$

Note 3.3.0.4. Later we will give a subset $S \subset M$ the structure of a smooth manifold such that Definition 3.3.0.1 and Definition 3.3.0.3 coincide.

Exercise 3.3.0.5. Let (M, \mathcal{A}) be a smooth manifold, $(U, \phi) \in \mathcal{A}$ with $\phi = (x^1, \dots, x^n)$, $p \in U$ and $f \in C^\infty(M)$. Then $f|_U \in C^\infty(U)$.

Proof. **FINISH!!!** □

Definition 3.3.0.6. Let (M, \mathcal{A}) be a smooth manifold, $(U, \phi) \in \mathcal{A}$ with $\phi = (x^1, \dots, x^n)$, $f \in C^\infty(U)$ and $i \in \{1, \dots, n\}$. We define the **partial derivative of f with respect to x^i** , denoted

$$\partial f / \partial x^i : U \rightarrow \mathbb{R} \quad \text{or} \quad \partial_i f : U \rightarrow \mathbb{R}$$

by

$$\frac{\partial f}{\partial x^i}(p) = \frac{\partial}{\partial u^i}[f \circ \phi^{-1}](\phi(p))$$

or equivalently,

$$\frac{\partial f}{\partial x^i} = \left(\frac{\partial}{\partial u^i}[f \circ \phi^{-1}] \right) \circ \phi$$

Exercise 3.3.0.7. Let (M, \mathcal{A}) be a smooth manifold, $(U, \phi) \in \mathcal{A}$ with $\phi = (x^1, \dots, x^n)$, $f \in C^\infty(U)$ and $i \in \{1, \dots, n\}$. Then $\partial / \partial x^i : C^\infty(U) \rightarrow C^\infty(U)$ is linear.

Proof. **FINISH!!!** □

Exercise 3.3.0.8. Let (M, \mathcal{A}) be a smooth manifold, $(U, \phi) \in \mathcal{A}$ with $\phi = (x^1, \dots, x^n)$, $f \in C^\infty(U)$ and $i, j \in \{1, \dots, n\}$. Then

$$\frac{\partial}{\partial x^i} \frac{\partial}{\partial x^j} f = \left(\frac{\partial}{\partial u^i} \frac{\partial}{\partial u^j}[f \circ \phi^{-1}] \right) \circ \phi$$

Proof.

$$\begin{aligned} \frac{\partial}{\partial x^i} \frac{\partial}{\partial x^j} f &= \frac{\partial}{\partial x^i} \left(\frac{\partial}{\partial x^j} f \right) \\ &= \frac{\partial}{\partial x^i} \left(\left[\frac{\partial}{\partial u^j}[f \circ \phi^{-1}] \right] \circ \phi \right) \\ &= \left(\frac{\partial}{\partial u^i} \left[\left(\left[\frac{\partial}{\partial u^j}[f \circ \phi^{-1}] \right] \circ \phi \right) \circ \phi^{-1} \right] \right) \circ \phi \\ &= \left(\frac{\partial}{\partial u^i} \left[\frac{\partial}{\partial u^j}[f \circ \phi^{-1}] \right] \right) \circ \phi \\ &= \left(\frac{\partial}{\partial u^i} \frac{\partial}{\partial u^j}[f \circ \phi^{-1}] \right) \circ \phi \end{aligned}$$

□

Exercise 3.3.0.9. Let (M, \mathcal{A}) be a smooth manifold, $(U, \phi) \in \mathcal{A}$ with $\phi = (x^1, \dots, x^n)$ and $i, j \in \{1, \dots, n\}$. Then

$$\frac{\partial}{\partial x^i} \frac{\partial}{\partial x^j} = \frac{\partial}{\partial x^j} \frac{\partial}{\partial x^i}$$

Proof. Let $f \in C^\infty(U)$. Since $f \circ \phi^{-1}$ is smooth,

$$\frac{\partial}{\partial u^i} \frac{\partial}{\partial u^j} [f \circ \phi^{-1}] = \frac{\partial}{\partial u^j} \frac{\partial}{\partial u^i} [f \circ \phi^{-1}]$$

The previous exercise implies that

$$\begin{aligned} \frac{\partial}{\partial x^i} \frac{\partial}{\partial x^j} f &= \left(\frac{\partial}{\partial u^i} \frac{\partial}{\partial u^j} [f \circ \phi^{-1}] \right) \circ \phi \\ &= \left(\frac{\partial}{\partial u^j} \frac{\partial}{\partial u^i} [f \circ \phi^{-1}] \right) \circ \phi \\ &= \frac{\partial}{\partial x^j} \frac{\partial}{\partial x^i} f \end{aligned}$$

□

Exercise 3.3.0.10. Let (M, \mathcal{A}) be a smooth manifold, $(U, \phi) \in \mathcal{A}$ with $\phi = (x^1, \dots, x^n)$ and $f \in C^\infty(U)$. Then for each $\alpha \in \mathbb{N}_0^n$,

$$\partial^\alpha f = (\partial^\alpha [f \circ \phi^{-1}]) \circ \phi$$

Proof. The claim is clearly true when $|\alpha| = 0$ or by definition if $|\alpha| = 1$. Let $n \in \mathbb{N}$ and suppose the claim is true for each $|\alpha| \in \{1, \dots, n-1\}$. Then there exists $i \in \{1, \dots, n\}$ such that $\alpha_i \geq 1$. Hence

$$\begin{aligned} \partial^\alpha f &= \partial^{e^i} (\partial^{\alpha-e^i} f) \\ &= \partial^{e^i} (\partial^{\alpha-e^i} [f \circ \phi^{-1}] \circ \phi) \\ &= (\partial^{e^i} [(\partial^{\alpha-e^i} [f \circ \phi^{-1}] \circ \phi) \circ \phi^{-1}]) \circ \phi \\ &= (\partial^{e^i} [\partial^{\alpha-e^i} [f \circ \phi^{-1}]]) \circ \phi \\ &= (\partial^\alpha [f \circ \phi^{-1}]) \circ \phi \end{aligned}$$

□

Exercise 3.3.0.11. Taylor's Theorem:

Let (M, \mathcal{A}) be a smooth manifold, $(U, \phi) \in \mathcal{A}$ with $\phi = (x^1, \dots, x^n)$ and $\phi(U)$ convex, $p \in U$, $f \in C^\infty(U)$ and $T \in \mathbb{N}$. Then there exist $(g_\alpha)_{|\alpha|=T+1} \subset C^\infty(U)$ such that

$$f = \sum_{k=0}^T \left[\sum_{|\alpha|=k} (x-p)^\alpha \partial^\alpha f(x_0) \right] + \sum_{|\alpha|=T+1} (x^i - x^i(p))^\alpha g_\alpha$$

and for each $|\alpha| = T+1$,

$$g_\alpha(p) = \frac{1}{(T+1)!} \partial^\alpha f(p)$$

Proof. Since $\phi(U)$ is open and convex and $f \circ \phi^{-1} \in C^\infty(\phi(U))$, Taylors thorem in section 2.1 implies that there exist $(\tilde{g}_\alpha)_{|\alpha|=T+1} \subset C^\infty(\phi(U))$ such that for each $q \in U$,

$$f \circ \phi^{-1}(\phi(q)) = \sum_{k=0}^T \left[\sum_{|\alpha|=k} (x^i(q) - x^i(p))^\alpha \partial^\alpha [f \circ \phi^{-1}](\phi(p)) \right] + \sum_{|\alpha|=T+1} (x^i(q) - x^i(p))^\alpha \tilde{g}_\alpha(\phi(q))$$

and for each $|\alpha| = T + 1$,

$$\begin{aligned}\tilde{g}_\alpha(\phi(p)) &= \frac{1}{(T+1)!} \partial^\alpha [f \circ \phi^{-1}](\phi(p)) \\ &= \frac{1}{(T+1)!} \partial^\alpha f(p)\end{aligned}$$

For $|\alpha| = T + 1$, set $g_\alpha = \tilde{g} \circ \phi$. Then

$$\begin{aligned}f(q) &= f \circ \phi^{-1}(\phi(q)) \\ &= \sum_{k=0}^T \left[\sum_{|\alpha|=k} (x^i(q) - x^i(p))^\alpha \partial^\alpha [f \circ \phi^{-1}](\phi(p)) \right] + \sum_{|\alpha|=T+1} (x^i(q) - x^i(p))^\alpha \tilde{g}_\alpha(\phi(q)) \\ &= \sum_{k=0}^T \left[\sum_{|\alpha|=k} (x^i(q) - x^i(p))^\alpha \partial^\alpha f(p) \right] + \sum_{|\alpha|=T+1} (x^i(q) - x^i(p))^\alpha g_\alpha(q)\end{aligned}$$

□

Definition 3.3.0.12. Let (N, \mathcal{B}) be a smooth manifold and $F : M \rightarrow N$. Then F is said to be

- **smooth** if for each $(U, \phi) \in \mathcal{A}$ and $(V, \psi) \in \mathcal{B}$,

$$\psi \circ F \circ \phi^{-1} : \phi(U \cap F^{-1}(V)) \rightarrow \psi(F(U) \cap V)$$

is smooth

- a **diffeomorphism** if F is a bijection and F, F^{-1} are smooth.

Exercise 3.3.0.13. Let (M, \mathcal{A}) and (N, \mathcal{B}) be smooth manifold and $F : M \rightarrow N$. If F is smooth, then F is continuous.

Proof. Suppose that F is smooth. Let $p \in M$. Choose $(U, \phi) \in \mathcal{A}$ and $(V, \psi) \in \mathcal{B}$ such that $p \in U$ and $F(p) \in V$. Put $\tilde{U} = U \cap F^{-1}(V)$ and $\tilde{V} = F(U) \cap V$.

Define $\tilde{\phi} : \tilde{U} \rightarrow \phi(\tilde{U})$ and $\tilde{\psi} : \tilde{V} \rightarrow \psi(\tilde{V})$ by

$$\tilde{\phi} = \phi|_{\tilde{U}}, \quad \tilde{\psi} = \psi|_{\tilde{V}}$$

Then $\tilde{\phi}$ and $\tilde{\psi}$ are homeomorphisms, $p \in \tilde{U}$ and $F(\tilde{U}) \subset \tilde{V}$. Define $\tilde{F} : \phi(\tilde{U}) \rightarrow \psi(\tilde{V})$ by

$$\tilde{F} = \tilde{\psi} \circ F \circ \tilde{\phi}^{-1}$$

By definition, \tilde{F} is smooth and therefore continuous. Since ϕ and ψ are homeomorphisms and $F|_{\tilde{U}} = \tilde{\psi}^{-1} \circ \tilde{F} \circ \tilde{\phi}$, we have that $F|_{\tilde{U}}$ is continuous. In particular, F is continuous at p and since $p \in M$ is arbitrary, F is continuous. □

Exercise 3.3.0.14. Let (M, \mathcal{A}) and (N, \mathcal{B}) be smooth manifold and $F : M \rightarrow N$. If F is a diffeomorphism, then F is a homeomorphism.

Proof. Suppose that F is a diffeomorphism. By definition, F and F^{-1} are smooth. The previous exercise implies that F and F^{-1} are continuous. Hence F is a homeomorphism. □

Exercise 3.3.0.15. Let (N, \mathcal{B}) be a smooth manifold and $F : M \rightarrow N$ a diffeomorphism. Then for each $(U, \phi) \in \mathcal{A}$, $(F(U), \phi \circ F^{-1}) \in \mathcal{B}$.

Proof. Let $(V, \psi) \in \mathcal{B}$.

1. Since ϕ and F^{-1} are homeomorphisms, $\phi \circ F^{-1} : F(U) \cap V \rightarrow \phi(U \cap F^{-1}(V))$ is a homeomorphism

2. Since F is a diffeomorphism,

$$\phi \circ F^{-1} \circ \psi^{-1} : \psi(F(U) \cap V) \rightarrow \phi(U \cap F^{-1}(V))$$

and

$$\psi \circ F \circ \phi^{-1} : \phi(F^{-1}(V) \cap U) \rightarrow \psi(V \cap F(U))$$

are smooth.

Therefore $(F(U), \phi \circ F^{-1})$ and (V, ψ) are smoothly compatible. Since \mathcal{B} is maximal, $(F(U), \phi \circ F^{-1}) \in \mathcal{B}$. □

Definition 3.3.0.16. Let (N, \mathcal{B}) be a smooth n -dimensional manifold, $F : M \rightarrow N$ smooth and $(V, \psi) \in \mathcal{B}$ with $\psi = (y^1, \dots, y^n)$. For $i \in \{1, \dots, n\}$, We define the **i -th component of F with respect to (V, ψ)** , denoted $F^i : V \rightarrow \mathbb{R}$, by

$$F^i = y^i \circ F$$

3.4 Partitions of Unity

Definition 3.4.0.1. Let $p \in M$, $U \in \mathcal{N}_a$ open and $\rho \in C_c^\infty(M)$. Then ρ is said to be a **bump function at p supported in U** if

1. $\rho \geq 0$
2. there exists $V \in \mathcal{N}_p$ such that V is open and $\rho|_V = 1$
3. $\text{supp } \rho \subset U$

Exercise 3.4.0.2. Define $f : \mathbb{R} \rightarrow \mathbb{R}$ by

$$f(t) = \begin{cases} e^{-\frac{1}{1-t^2}} & t \in (-1, 1) \\ 0 & t \notin (-1, 1) \end{cases}$$

Then $f \in C_c^\infty(\mathbb{R})$.

Proof.

□

3.5 The Tangent Space

Exercise 3.5.0.1. Let $p \in M$. We have that $C_p^\infty(M)$ is a vector space.

Proof. Clear. □

Definition 3.5.0.2. Let $(U, \phi) \in \mathcal{A}$ with $\phi = (x^1, \dots, x^n)$ and $p \in U$. For $i \in \{1, \dots, n\}$, define the partial derivative with respect to x^i at p , denoted

$$\left. \frac{\partial}{\partial x^i} \right|_p : C_p^\infty(M) \rightarrow \mathbb{R}, \text{ or } \partial_i|_p : C_p^\infty(M) \rightarrow \mathbb{R}$$

by

$$\left. \frac{\partial}{\partial x^i} \right|_p f = \frac{\partial f}{\partial x^i}(p)$$

Exercise 3.5.0.3. Let $(U, \phi) \in \mathcal{A}$ with $\phi = (x^1, \dots, x^n)$ and $p \in U$. Then for each $i, j \in \{1, \dots, n\}$, we have that

$$\left. \frac{\partial}{\partial x^i} \right|_p x^j(p) = \delta_{i,j}$$

Proof. Let $i, j \in \{1, \dots, n\}$. Then

$$\begin{aligned} \left. \frac{\partial}{\partial x^i} \right|_p x^j &= \left. \frac{\partial}{\partial u^i} \right|_{\phi(p)} x^j \circ \phi^{-1} \\ &= \left. \frac{\partial}{\partial u^i} \right|_{\phi(p)} u^j \circ \phi \circ \phi^{-1} \\ &= \left. \frac{\partial}{\partial u^i} \right|_{\phi(p)} u^j \\ &= \delta_{i,j} \end{aligned}$$

□

Exercise 3.5.0.4. Change of Coordinates:

Let $(U, \phi), (V, \psi) \in \mathcal{A}$ with $\phi = (x^1, \dots, x^n)$ and $\psi = (y^1, \dots, y^n)$, $p \in U \cap V$ and $f \in C_p^\infty(M)$. Then for each $i \in \{1, \dots, n\}$,

$$\left. \frac{\partial}{\partial y^i} \right|_p = \sum_{j=1}^n \left. \frac{\partial}{\partial x^j} \right|_p y^j(p) \left. \frac{\partial}{\partial x^i} \right|_p$$

Proof. Put $h = \phi \circ \psi^{-1}$ and write $h = (h_1, \dots, h_n)$. Then $\phi = h \circ \psi$ and $\psi^{-1} = \phi^{-1} \circ h$. By definition and the chain rule, we have that

$$\begin{aligned} \left. \frac{\partial}{\partial y^i} \right|_p f &= \left. \frac{\partial}{\partial u^i} \right|_{\psi(p)} f \circ \psi^{-1} \\ &= \left. \frac{\partial}{\partial u^i} \right|_{\psi(p)} f \circ \phi^{-1} \circ h \\ &= \sum_{j=1}^n \left(\left. \frac{\partial}{\partial u^j} \right|_{h \circ \psi(p)} f \circ \phi^{-1} \right) \left(\left. \frac{\partial}{\partial u^i} \right|_{\psi(p)} h_j \right) \\ &= \sum_{j=1}^n \left(\left. \frac{\partial}{\partial u^j} \right|_{\phi(p)} f \circ \phi^{-1} \right) \left(\left. \frac{\partial}{\partial u^i} \right|_{\psi(p)} x^j \circ \psi^{-1} \right) \\ &= \sum_{j=1}^n \left(\left. \frac{\partial}{\partial x^j} \right|_p f \right) \left(\left. \frac{\partial}{\partial y^i} \right|_p x^j \right) \end{aligned}$$

□

Definition 3.5.0.5. Let $p \in M$ and $v : C_p^\infty(M) \rightarrow \mathbb{R}$. Then v is said to be **Leibnizian** if for each $f, g \in C_p^\infty(M)$,

$$v(fg) = v(f)g(p) + f(p)v(g)$$

and v is said to be a **derivation at p** if for each $f, g \in C_p^\infty(M)$ and $a \in \mathbb{R}$,

1. v is linear
2. v is Leibnizian

We define the **tangent space of M at p** , denoted T_pM , by

$$T_pM = \{v : C_p^\infty(M) \rightarrow \mathbb{R} : v \text{ is a derivation at } p\}$$

Exercise 3.5.0.6. Let $f \in C_p^\infty(M)$ and $v \in T_pM$. If f is constant, then $vf = 0$.

Proof. Suppose that $f = 1$. Then $f^2 = f$ and $v(f^2) = 2v(f)$. So $v(f) = 2v(f)$ which implies that $v(f) = 0$. If $f \neq 1$, then there exists $c \in \mathbb{R}$ such that $f = c$. Since v is linear, $v(f) = cv(1) = 0$. \square

Exercise 3.5.0.7. Let $(U, \phi) \in \mathcal{A}$ with $\phi = (x^1, \dots, x^n)$ and $p \in U$. Then

$$\left\{ \frac{\partial}{\partial x^1} \Big|_p, \dots, \frac{\partial}{\partial x^n} \Big|_p \right\}$$

is a basis for T_pM and $\dim T_pM = n$.

Proof. Clearly $\frac{\partial}{\partial x^1} \Big|_p, \dots, \frac{\partial}{\partial x^n} \Big|_p \in T_pM$. Let $a_1, \dots, a_n \in \mathbb{R}$. Suppose that

$$v = \sum_{i=1}^n a_i \frac{\partial}{\partial x^i} \Big|_p = 0$$

Then

$$\begin{aligned} 0 &= vx^j \\ &= \sum_{i=1}^n a_i \frac{\partial}{\partial x^i} \Big|_p x^j \\ &= a_j \end{aligned}$$

Hence $\left\{ \frac{\partial}{\partial x^1} \Big|_p, \dots, \frac{\partial}{\partial x^n} \Big|_p \right\}$ is independent.

Now, let $v \in T_pM$ and $f \in C_p^\infty(M)$. By Taylor's theorem, there exist $g_1, \dots, g_n \in C_p^\infty(M)$ such that

$$f = f(p) + \sum_{i=1}^n (x^i - x^i(p))g_i$$

and for each $i \in \{1, \dots, n\}$,

$$g_i(p) = \frac{\partial}{\partial x^i} \Big|_p f$$

Then

$$\begin{aligned} v(f) &= \sum_{i=1}^n v(x^i - x^i(p))g_i(p) + \sum_{i=1}^n (x^i(p) - x^i(p))v(g_i) \\ &= \sum_{i=1}^n v(x^i)g_i(p) \\ &= \sum_{i=1}^n v(x^i) \frac{\partial}{\partial x^i} \Big|_p f \\ &= \left[\sum_{i=1}^n v(x^i) \frac{\partial}{\partial x^i} \Big|_p \right] f \end{aligned}$$

So

$$v = \sum_{i=1}^n v(x^i) \frac{\partial}{\partial x^i} \Big|_p$$

and

$$v \in \text{span} \left\{ \frac{\partial}{\partial x^1} \Big|_p, \dots, \frac{\partial}{\partial x^n} \Big|_p \right\}$$

□

Definition 3.5.0.8. Let (N, \mathcal{B}) be a smooth manifold, $F : M \rightarrow N$ smooth and $p \in M$. We define the **differential of F at p** , denoted $dF_p : T_p M \rightarrow T_{F(p)} N$, by

$$\left[dF_p(v) \right] (f) = v(f \circ F)$$

for $v \in T_p M$ and $f \in C_{F(p)}^\infty(N)$.

Exercise 3.5.0.9. Let (N, \mathcal{B}) be a smooth manifold, $F : M \rightarrow N$ smooth and $p \in M$. Then for each $v \in T_p M$, $dF_p(v)$ is a derivation.

Proof. Let $v \in T_p M$, $f, g \in C_{F(p)}^\infty(N)$ and $c \in \mathbb{R}$. Then

1.

$$\begin{aligned} dF_p(v)(f + cg) &= v((f + cg) \circ F) \\ &= v(f \circ F + cg \circ F) \\ &= v(f \circ F) + cv(g \circ F) \\ &= dF_p(v)(f) + cdF_p(v)(g) \end{aligned}$$

So $dF_p(v)$ is linear.

2.

$$\begin{aligned} dF_p(v)(fg) &= v(fg \circ F) \\ &= v((f \circ F) * (g \circ F)) \\ &= v(f \circ F) * (g \circ F)(p) + (f \circ F)(p) * v(g \circ F) \\ &= dF_p(v)(f) * g(F(p)) + f(F(p)) * dF_p(v)(g) \end{aligned}$$

So $dF_p(v)$ is Leibnizian and hence $dF_p(v) \in T_{F(p)} N$

□

Exercise 3.5.0.10. Let (N, \mathcal{B}) be a smooth manifold, $F : M \rightarrow N$ smooth and $p \in M$. If F is a diffeomorphism, then dF_p is an isomorphism.

Proof. Suppose that F is a diffeomorphism. Since F is a homeomorphism, $\dim N = n$. Choose $(U, \phi) \in \mathcal{A}$ such that $p \in U$. A previous exercise tells us that $(F(U), \phi \circ F^{-1}) \in \mathcal{B}$. Write $\phi = (x^1, \dots, x^n)$ and $\phi \circ F^{-1} = (y^1, \dots, y^n)$. Let $f \in C_{F(p)}^\infty(N)$. Then

$$\begin{aligned} \frac{\partial}{\partial y^i} \Big|_{F(p)} f &= \frac{\partial}{\partial u^i} \Big|_{\phi \circ F^{-1}(F(p))} f \circ (\phi \circ F^{-1})^{-1} \\ &= \frac{\partial}{\partial u^i} \Big|_{\phi(p)} f \circ F \circ \phi^{-1} \\ &= \frac{\partial}{\partial x^i} \Big|_p f \circ F \end{aligned}$$

Therefore

$$\begin{aligned} \left[dF_p \left(\frac{\partial}{\partial x^i} \Big|_p \right) \right] (f) &= \frac{\partial}{\partial x^i} \Big|_p f \circ F \\ &= \frac{\partial}{\partial y^i} \Big|_{F(p)} f \end{aligned}$$

Hence

$$dF_p \left(\frac{\partial}{\partial x^i} \Big|_p \right) = \frac{\partial}{\partial y^i} \Big|_{F(p)}$$

Since $\left\{ \frac{\partial}{\partial x^1} \Big|_p, \dots, \frac{\partial}{\partial x^n} \Big|_p \right\}$ is a basis for $T_p M$ and $\left\{ \frac{\partial}{\partial y^1} \Big|_{F(p)}, \dots, \frac{\partial}{\partial y^n} \Big|_{F(p)} \right\}$ is a basis for $T_{F(p)} N$, dF_p is an isomorphism. \square

Exercise 3.5.0.11. Let (M, \mathcal{A}) be a smooth m -dimensional manifold, (N, \mathcal{B}) a n -dimensional smooth manifold, $F : M \rightarrow N$ smooth, $(U, \phi) \in \mathcal{A}$ with $\phi = (x^1, \dots, x^m)$, $(V, \psi) \in \mathcal{B}$ with $\psi = (y^1, \dots, y^n)$ and $p \in U$. Define the ordered bases $B_\phi = \left\{ \frac{\partial}{\partial x^1} \Big|_p, \dots, \frac{\partial}{\partial x^m} \Big|_p \right\}$ and $B_\psi = \left\{ \frac{\partial}{\partial y^1} \Big|_{F(p)}, \dots, \frac{\partial}{\partial y^n} \Big|_{F(p)} \right\}$. Then the matrix representation of dF_p with respect to the bases B_ϕ and B_ψ is

$$dF_p^{i,j} = \frac{\partial F^i}{\partial x^j}(p)$$

Proof. Let $(dF_p)_{B_\phi, B_\psi} = (a_{i,j})_{i,j} \in \mathbb{R}^{n \times m}$. Then for each $j \in \{1, \dots, m\}$,

$$dF_p \left(\frac{\partial}{\partial x^j} \Big|_p \right) = \sum_{i=1}^n a_{i,j} \frac{\partial}{\partial y^i} \Big|_{F(p)}$$

This implies that

$$\begin{aligned} dF_p \left(\frac{\partial}{\partial x^j} \Big|_p \right) (y^k) &= \sum_{i=1}^n a_{i,j} \frac{\partial}{\partial y^i} \Big|_{F(p)} (y^k) \\ &= \sum_{i=1}^n a_{i,j} \delta_{i,k} \\ &= a_{k,j} \end{aligned}$$

By definition,

$$\begin{aligned} dF_p \left(\frac{\partial}{\partial x^j} \Big|_p \right) (y^k) &= \frac{\partial}{\partial x^j} \Big|_p y^k \circ F \\ &= \frac{\partial}{\partial x^j} \Big|_p F^k \\ &= \frac{\partial F^k}{\partial x^j}(p) \end{aligned}$$

\square

Note 3.5.0.12. Since $\text{rank } dF_p$ is independent of basis, it is independent of coordinate charts $(U, \phi) \in \mathcal{A}$ and $(V, \psi) \in \mathcal{B}$.

Definition 3.5.0.13. Let (N, \mathcal{B}) be a smooth manifold, $F : M \rightarrow N$ a diffeomorphism. Define the **push forward of F** , denoted

$$F_* : M \rightarrow \coprod_{p \in M} \text{Iso}(T_p M, T_{F(p)} N)$$

by

$$p \mapsto dF_p$$

3.6 The Cotangent Space

Definition 3.6.0.1. Let $p \in M$. We define the **cotangent space of M at p** , denoted T_p^*M , by

$$T_p^*M = (T_pM)^*$$

Definition 3.6.0.2. Let $f \in C^\infty(M)$. We define the **differential of f at p** , denoted $df_p : T_pM \rightarrow \mathbb{R}$, by

$$df_p(v) = vf$$

Exercise 3.6.0.3. Let $f \in C^\infty(M)$ and $p \in M$. Then $df_p \in T_p^*M$.

Proof. Let $v_1, v_2 \in T_pM$ and $\lambda \in \mathbb{R}$. Then

$$\begin{aligned} df_p(v_1 + \lambda v_2) &= (v_1 + \lambda v_2)f \\ &= v_1f + \lambda v_2f \\ &= df_p(v_1) + \lambda df_p(v_2) \end{aligned}$$

So that df_p is linear and hence $df_p \in T_p^*M$. □

Exercise 3.6.0.4. Let $(U, \phi) \in \mathcal{A}$ with $\phi = (x^1, \dots, x^n)$ and $p \in U$. Then for each $i, j \in \{1, \dots, n\}$,

$$dx_p^i \left(\frac{\partial}{\partial x^j} \Big|_p \right) = \delta_{i,j}$$

In particular, $\{dx_p^1, \dots, dx_p^n\}$ is the dual basis to $\left\{ \frac{\partial}{\partial x^1} \Big|_p, \dots, \frac{\partial}{\partial x^n} \Big|_p \right\}$ and $T_p^*M = \text{span}\{dx_p^1, \dots, dx_p^n\}$.

Proof. Let $i, j \in \{1, \dots, n\}$. Then by definition,

$$\begin{aligned} \left[dx_p^i \left(\frac{\partial}{\partial x^i} \Big|_p \right) \right]_p &= \frac{\partial}{\partial x^i} \Big|_p x^i \\ &= \delta_{i,i} \end{aligned}$$

□

Exercise 3.6.0.5. Let $f \in C^\infty(M)$, (U, ϕ) a chart on M with $\phi = (x^1, \dots, x^n)$ and $p \in U$. Then

$$df_p = \sum_{i=1}^n \frac{\partial f}{\partial x^i}(p) dx_p^i$$

Proof. Since $\{dx_p^1, \dots, dx_p^n\}$ is a basis for T_p^*M , for each there exist $a_1(p), \dots, a_n(p) \in \mathbb{R}$ such that $df_p = \sum_{i=1}^n a_i(p) dx_p^i$. Therefore, we have that

$$\begin{aligned} df_p \left(\frac{\partial}{\partial x^i} \Big|_p \right) &= \sum_{i=1}^n a_i(p) dx_p^i \left(\frac{\partial}{\partial x^i} \Big|_p \right) \\ &= a_j(p) \end{aligned}$$

By definition, we have that

$$\begin{aligned} df_p \left(\frac{\partial}{\partial x^i} \Big|_p \right) &= \frac{\partial f}{\partial x^i} \Big|_p \\ &= \frac{\partial f}{\partial x^j}(p) \end{aligned}$$

So $a_j(p) = \frac{\partial f}{\partial x^j}(p)$ and

$$df_p = \sum_{i=1}^n \frac{\partial f}{\partial x^i}(p) dx_p^i$$

□

3.7 Maps of Full Rank

Definition 3.7.0.1. Let (M, \mathcal{A}) and (N, \mathcal{B}) be smooth manifolds, $F : M \rightarrow N$ a smooth map and $p \in M$. We define the **rank of F at p** , denoted $\text{rank}_p F$, by $\text{rank}_p F = \text{rank } dF_p$. We say that F has **constant rank** if for each $p, q \in M$, $\text{rank}_p F = \text{rank}_q F$. If F has constant rank, we define the **rank of F** , denoted $\text{rank } F$, by $\text{rank } F = \text{rank}_p F$.

Definition 3.7.0.2. Let (M, \mathcal{A}) and (N, \mathcal{B}) be smooth manifolds, $F : M \rightarrow N$ a smooth map. Then F is said to be

- an **immersion** if for each $p \in M$, $dF_p : T_p M \rightarrow T_{F(p)} N$ is injective
- a **submersion** if for each $p \in M$, $dF_p : T_p M \rightarrow T_{F(p)} N$ is surjective

Exercise 3.7.0.3. Let (M, \mathcal{A}) and (N, \mathcal{B}) be smooth manifolds, $F : M \rightarrow N$ a smooth map.

Definition 3.7.0.4. Let (M, \mathcal{A}) and (N, \mathcal{B}) be smooth manifolds and $F : M \rightarrow N$ smooth. Then F is said to be an **embedding** if

1. F is an immersion
2. $F : M \rightarrow F(M)$.

Note 3.7.0.5. Here the topology on $F(M)$ is the subspace topology.

3.8 Submanifolds

Exercise 3.8.0.1. Let (M, \mathcal{A}) be a smooth manifold and $S \subset M$ open. For $(U, \phi) \in \mathcal{A}$, define $\tilde{U} \subset S$ and $\tilde{\phi} : \tilde{U} \rightarrow \phi(\tilde{U})$ by $\tilde{U} = U \cap S$ and $\tilde{\phi} = \phi|_{U \cap S}$. Set $\mathcal{B} = \{(\tilde{U}, \tilde{\phi}) : (U, \phi) \in \mathcal{A}\}$. Then \mathcal{B} is a smooth structure on S .

Proof.

□

Definition 3.8.0.2. Let (M, \mathcal{A}) and (N, \mathcal{B}) be smooth manifolds. Suppose that $M \subset N$. Then (M, \mathcal{A}) is said to be

1. an **immersed submanifold** of (N, \mathcal{B}) if $\text{id} : M \rightarrow N$ is a smooth immersion
2. an **embedded submanifold** of (N, \mathcal{B}) if $\text{id} : M \rightarrow N$ is a smooth embedding

Note 3.8.0.3. Essentially, embedded submanifolds are immersed submanifolds with the subspace topology.

Note 3.8.0.4. For the remainder of this section, we assume that $k \leq n$.

Definition 3.8.0.5. Let $U \subset \mathbb{R}^n$ and $S \subset U$. Then S is said to be a **k -slice** of U if $S = \{u \in U : u^{k+1}, \dots, u^n = 0\}$.

Exercise 3.8.0.6. Let $U \subset \mathbb{R}^n$ and $S \subset U$. Suppose that S is a k -slice of U . Define $\pi : \mathbb{R}^n \rightarrow \mathbb{R}^k$ by

$$\pi(u^1, \dots, u^k, \dots, u^n) = (u^1, \dots, u^k)$$

Then $\pi|_S \rightarrow \pi(S)$ is a diffeomorphism.

Proof. Clear. □

Definition 3.8.0.7. Let (M, \mathcal{A}) be a smooth manifold, $(U, \phi) \in \mathcal{A}$ and $S \subset U$. Then S is said to be a **k -slice** of U if $\phi(S)$ is a k -slice of $\phi(U)$.

Definition 3.8.0.8. Let (M, \mathcal{A}) be a smooth manifold, $S \subset M$ and $(U, \phi) \in \mathcal{A}$. Then (U, ϕ) is said to be a **k -slice chart for S** if $U \cap S$ is a k -slice of U .

Exercise 3.8.0.9. Let (M, \mathcal{A}) be a smooth manifold, $S \subset M$ and $(U, \phi) \in \mathcal{A}$ with $\phi = (x^1, \dots, x^n)$. If (U, ϕ) is a k -slice chart for S , then $\phi|_S = (x^1|_S, \dots, x^k|_S, 0, \dots, 0)$.

Proof. Clear. □

Definition 3.8.0.10. Let (M, \mathcal{A}) be a smooth manifold and $S \subset M$. Then S is said to satisfy the **local k -slice condition** if for each $p \in S$, there exists $(U, \phi) \in \mathcal{A}$ such that $p \in U$ and (U, ϕ) is a k -slice chart of S .

Exercise 3.8.0.11. Let (M, \mathcal{A}) be a n -dimensional smooth manifold and $S \subset M$ a subspace. If S satisfies the local k -slice condition, then there exists a smooth structure $\tilde{\mathcal{A}}$ on S such that $(S, \tilde{\mathcal{A}})$ is an embedded submanifold of M .

Proof. Suppose that S satisfies the local k -slice condition. Define $\pi : \mathbb{R}^n \rightarrow \mathbb{R}^k$ as above. Let $(U, \phi) \in \mathcal{A}$. Suppose that (U, ϕ) is a k -slice chart for S . Define $\tilde{U} = U \cap S$ and $\tilde{\phi} : \tilde{U} \rightarrow \pi \circ \phi(\tilde{U})$ by

$$\tilde{\phi} = \pi \circ \phi|_{\tilde{U}}$$

By definition, $\phi(\tilde{U})$ is a k -slice of $\phi(U)$. A previous exercise implies that $\pi|_{\phi(\tilde{U})} \rightarrow \pi \circ \phi(\tilde{U})$ is a diffeomorphism and hence a homeomorphism. Thus $\tilde{\phi}$ is a homeomorphism.

Define

$$\tilde{\mathcal{B}} = \{(\tilde{U}, \tilde{\phi}) : (U, \phi) \text{ is a } k\text{-slice for } S\}$$

Let $p \in S$. By assumption, there exists $(U, \phi) \in \mathcal{A}$ such that $p \in U$ and (U, ϕ) is a k -slice chart of S . Then $(\tilde{U}, \tilde{\phi}) \in \tilde{\mathcal{B}}$ and $\tilde{\mathcal{A}}$ is an atlas on S . By construction of $\tilde{\mathcal{B}}$, S is locally half Euclidean of dimension k . Since M is second countable Hausdorff, so is S in the subspace topology. Thus $(S, \tilde{\mathcal{B}})$ is a k -dimensional manifold. Let $(\tilde{U}, \tilde{\phi}), (\tilde{V}, \tilde{\psi}) \in \tilde{\mathcal{B}}$. Then

$$\tilde{\phi} \circ \tilde{\psi}^{-1}|_{\tilde{U} \cap \tilde{V}} = \pi|_{\phi(\tilde{U} \cap \tilde{V})} \circ \phi|_{\tilde{U} \cap \tilde{V}} \circ \psi|_{\tilde{U} \cap \tilde{V}}^{-1} \circ \pi|_{\psi(\tilde{U} \cap \tilde{V})}^{-1}$$

which is a diffeomorphism. So $(\tilde{U}, \tilde{\phi})$ and $(\tilde{V}, \tilde{\psi})$ smoothly compatible. Hence $\tilde{\mathcal{B}}$ is smooth. An exercise in section 4.1 implies that there exists a unique smooth structure $\tilde{\mathcal{A}}$ on S such that $\tilde{\mathcal{B}} \subset \tilde{\mathcal{A}}$. So $(S, \tilde{\mathcal{A}})$ is a smooth k -dimensional manifold.

Clearly $\text{id} : S \rightarrow S$ is a homeomorphism. Let $(V, \psi) \in \mathcal{A}$ and $(\tilde{U}, \tilde{\phi}) \in \tilde{\mathcal{A}}$.

Finish!! □

Definition 3.8.0.12.

Exercise 3.8.0.13.

Chapter 4

Vector Bundles and Tensor Fields

4.1 The Vector Bundle

Definition 4.1.0.1. Let E , M and F be smooth manifolds and $\pi : E \rightarrow M$ a smooth surjection, $U \subset M$ open and $\Phi : \pi^{-1}(U) \rightarrow U \times F$. Then (U, Φ) is said to be a **smooth local trivialization of E over U** if

1. Φ is a diffeomorphism
2. $\pi_U \circ \Phi = \pi|_{\pi^{-1}(U)}$ (where $\pi_U : U \times F \rightarrow U$ denotes projection onto U)

Exercise 4.1.0.2. Let E , M and F be sets and $\pi : E \rightarrow M$ and $\Phi : \pi^{-1}(U) \rightarrow U \times F$ a bijection. If $\pi_U \circ \Phi = \pi|_{\pi^{-1}(U)}$, then for each $A \subset U$,

$$\Phi(\pi^{-1}(A)) = A \times F$$

Hint: consider $\Phi^{-1}(A \times F)$

Proof. Let $A \subset U$. Since $\pi_U^{-1}(A) = A \times F$, we have that

$$\begin{aligned}\Phi^{-1}(A \times F) &= \Phi^{-1}(\pi_U^{-1}(A)) \\ &= (\pi_U \circ \Phi)^{-1}(A) \\ &= (\pi|_{\pi^{-1}(U)})^{-1}(A) \\ &= \pi^{-1}(A) \cap \pi^{-1}(U) \\ &= \pi^{-1}(A \cap U) \\ &= \pi^{-1}(A)\end{aligned}$$

Since Φ is a bijection, we have that

$$\begin{aligned}\Phi(\pi^{-1}(A)) &= \Phi \circ \Phi^{-1}(A \times F) \\ &= A \times F\end{aligned}$$

□

Definition 4.1.0.3. Let E and M be smooth manifolds and $\pi : E \rightarrow M$ a smooth surjection. Then (E, M, π) is said to be a **smooth vector bundle of rank n** if

1. for each $p \in M$, $\pi^{-1}(\{p\})$ is a n -dimensional real vector space.
2. for each $p \in M$, there exist open $U \in \mathcal{N}_p$ and $\Phi : \pi^{-1}(U) \rightarrow U \times \mathbb{R}^n$ such that (U, Φ) is a smooth local trivialization of E over U .

3. for each $p \in M$,

$$\Phi|_{\pi^{-1}(\{p\})} : \pi^{-1}(\{p\}) \rightarrow \{p\} \times \mathbb{R}^n$$

is an isomorphism.

Exercise 4.1.0.4. Let M be a n -dimensional smooth manifold. Set $E = M \times \mathbb{R}^n$ and define $\pi : E \rightarrow M$ by $\pi(p, x) = p$. Then (E, M, π) is a smooth vector bundle of rank n .

Proof.

1. For each $p \in M$, $\pi^{-1}(\{p\}) = \{p\} \times \mathbb{R}^n$ which may be given the obvious vector space structure.
2. Let $p \in M$. Set $U = M$. Then $\pi^{-1}(U) = E$. Define $\Phi : \pi^{-1}(U) \rightarrow U \times \mathbb{R}^n$ by $\Phi = \text{id}_E$. Then (U, Φ) is a smooth local trivialization of E over U .
3. Let $p \in M$. Then $\Phi|_{\pi^{-1}(\{p\})} : \pi^{-1}(\{p\}) \rightarrow \{p\} \times \mathbb{R}^n$ is clearly an isomorphism.

□

Theorem 4.1.0.5. Let E and M be smooth manifolds and $\pi : E \rightarrow M$ a smooth surjection.

Definition 4.1.0.6. We define the **tangent bundle of M** , denoted TM , by

$$TM = \coprod_{p \in M} T_p M$$

We denote the natural projection map by $\pi : TM \rightarrow M$.

Definition 4.1.0.7. Let $(U, \phi) \in \mathcal{A}$ with $\phi = (x^1, \dots, x^n)$. Define $\tilde{U} \subset TM$ and $\tilde{\phi} : \tilde{U} \rightarrow \phi(U) \times \mathbb{R}^n$ by

- $\tilde{U} = \pi^{-1}(U)$
-

$$\begin{aligned} \tilde{\phi} \left(\sum_{i=1}^n v^i \frac{\partial}{\partial x^i} \Big|_p \right) &= (\phi(p), v) \\ &= (x^1(p), \dots, x^n(p), v^1, \dots, v^n) \end{aligned}$$

Exercise 4.1.0.8. Let $(U, \phi) \in \mathcal{A}$ with $\phi = (x^1, \dots, x^n)$. Then $\tilde{\phi} : \tilde{U} \rightarrow \phi(U) \times \mathbb{R}^n$ is a bijection.

4.2 The cotangent Bundle

Definition 4.2.0.1. We define the **cotangent bundle of M** , denoted T^*M , by

$$T^*M = \coprod_{p \in M} T_p^*M$$

4.3 The (r, s) -Tensor Bundle

Definition 4.3.0.1. 1. the **cotangent bundle of M** , denoted T^*M , by

$$T^*M = \coprod_{p \in M} T_p^*M$$

2. the **(r, s) -tensor bundle of M** , denoted $T_s^r M$, by

$$T_s^r M = \coprod_{p \in M} T_s^r(T_p M)$$

3. the **k -alternating tensor bundle of M** , denoted $\Lambda^k(M)$, by

$$\Lambda^k M = \coprod_{p \in M} \Lambda^k(T_p M)$$

4.4 Vector Fields

Definition 4.4.0.1. Let $X : M \rightarrow TM$. Then X is said to be a **vector field on M** if for each $p \in M$, $X_p \in T_p M$.

For $f \in C^\infty(M)$, we define $Xf : M \rightarrow \mathbb{R}$ by

$$(Xf)_p = X_p(f)$$

and X is said to be **smooth** if for each $f \in C^\infty(M)$, Xf is smooth.

We denote the set of smooth vector fields on M by $\Gamma^1(M)$.

Definition 4.4.0.2. Let $f \in C^\infty(M)$ and $X, Y \in \Gamma^1(M)$. We define

- $fX \in \Gamma^1(M)$ by

$$(fX)_p = f(p)X_p$$

- $X + Y \in \Gamma^1(M)$ by

$$(X + Y)_p = X_p + Y_p$$

Exercise 4.4.0.3. The set $\Gamma^1(M)$ is a $C^\infty(M)$ -module.

Proof. Clear. □

Exercise 4.4.0.4. Let $X \in \Gamma^1(M)$ and $(U, \phi) \in \mathcal{A}$ with $\phi = (x^1, \dots, x^n)$. Then

$$X|_U = \sum_{i=1}^n (Xx^i) \frac{\partial}{\partial x^i}$$

Proof. Let $p \in M$. Then $X_p \in T_p M$ and $\left\{ \frac{\partial}{\partial x^1} \Big|_p, \dots, \frac{\partial}{\partial x^n} \Big|_p \right\}$ is a basis of $T_p M$. So there exist $f_1(p), \dots, f_n(p) \in \mathbb{R}$ such that $X_p = \sum_{i=1}^n f_i(p) \frac{\partial}{\partial x^i} \Big|_p$. Let $j \in \{1, \dots, n\}$. Then,

$$\begin{aligned} X_p(x^j) &= \sum_{i=1}^n f_i(p) \frac{\partial}{\partial x^i} x^j(p) \\ &= f_j(p) \end{aligned}$$

Hence $Xx^j = f_j$ and $X|_U = \sum_{i=1}^n (Xx^i) \frac{\partial}{\partial x^i}$. □

Exercise 4.4.0.5. Let $(U, \phi) \in \mathcal{A}$ with $\phi = (x^1, \dots, x^n)$. Then for each $i \in \{1, \dots, n\}$,

$$\frac{\partial}{\partial x^i} \in \Gamma(U)$$

Proof. Let $i \in \{1, \dots, n\}$ and $f \in C^\infty(M)$. Define $g : M \rightarrow \mathbb{R}$ by $g = \frac{\partial}{\partial x^i} f$. Let $(V, \psi) \in \mathcal{A}$. Then for each $x \in \psi(U \cap V)$,

$$\begin{aligned} g \circ \psi^{-1}(x) &= \frac{\partial}{\partial x^i} \Big|_{\psi^{-1}(x)} f \\ &= \frac{\partial}{\partial u^i} \Big|_{\phi \circ \psi^{-1}(x)} f \circ \phi^{-1} \\ &= \frac{\partial}{\partial u^i} [f \circ \phi^{-1}](\phi \circ \psi^{-1}(x)) \end{aligned}$$

Since $f \circ \phi^{-1}$ and $\phi \circ \psi^{-1}$ are smooth, $g \circ \psi^{-1}$ is smooth and hence g is smooth. Since $f \in C^\infty(M)$ was arbitrary, by definition, $\frac{\partial}{\partial x^i}$ is smooth. □

4.5 1-Forms

Definition 4.5.0.1. Let $\omega : M \rightarrow T^*M$. Then ω is said to be a **1-form on M** if for each $p \in M$, $\omega_p \in T_p^*M$. For each $X \in \Gamma^1(M)$, we define $\omega(X) : M \rightarrow \mathbb{R}$ by

$$\omega(X)_p = \omega_p(X_p)$$

and ω is said to be **smooth** if for each $X \in \Gamma^1(M)$, $\omega(X)$ is smooth. The set of smooth 1-forms on M is denoted $\Gamma_1(M)$.

Definition 4.5.0.2. Let $f \in C^\infty(M)$ and $\alpha, \beta \in \Gamma_1(M)$. We define

- $f\alpha \in \Gamma_1(M)$ by

$$(f\alpha)_p = f(p)\alpha_p$$

- $\alpha + \beta \in \Gamma_1(M)$ by

$$(\alpha + \beta)_p = \alpha_p + \beta_p$$

Exercise 4.5.0.3. The set $\Gamma_1(M)$ is a $C^\infty(M)$ -module.

Proof. Clear. □

Exercise 4.5.0.4.

4.6 (r, s) -Tensor Fields

Definition 4.6.0.1. Let $\alpha : M \rightarrow T_s^r M$. Then α is said to be a (r, s) -**tensor field on M** if for each $p \in M$, $\alpha_p \in T_p^r(T_p M)$.

For each $\omega \in \Gamma_1(M)^r$ and $X \in \Gamma^1(M)^s$, we define $\alpha(\omega, X) : M \rightarrow \mathbb{R}$ by

$$\alpha(\omega, X)_p = \alpha_p(\omega_p, X_p)$$

and α is said to be **smooth** if for each $\omega \in \Gamma_1(M)^r$ and $X \in \Gamma^1(M)^s$, $\alpha(\omega, X)$ is smooth. The set of smooth (r, s) -tensor fields on M is denoted $\Gamma_s^r(M)$.

Definition 4.6.0.2. Let $f \in C^\infty(M)$ and $\alpha, \beta \in \Gamma_s^r(M)$. We define

- $f\alpha : M \rightarrow T_s^r M$ by

$$(f\alpha)_p = f(p)\alpha_p$$

- $\alpha + \beta : M \rightarrow T_s^r M$ by

$$(\alpha + \beta)_p = \alpha_p + \beta_p$$

Exercise 4.6.0.3. Let $f \in C^\infty(M)$ and $\alpha, \beta \in \Gamma_s^r(M)$. Then

1. $f\alpha \in \Gamma_s^r(M)$ by

$$(f\alpha)_p = f(p)\alpha_p$$

2. $\alpha + \beta \in \Gamma_s^r(M)$ by

$$(\alpha + \beta)_p = \alpha_p + \beta_p$$

Proof. Clear. □

Exercise 4.6.0.4. The set $\Gamma_s^r(M)$ is a $C^\infty(M)$ -module.

Proof. Clear. □

Definition 4.6.0.5. Let $\alpha_1 \in \Gamma_{s_1}^{r_1}(M)$ and $\alpha_2 \in \Gamma_{s_2}^{r_2}(M)$. We define the **tensor product of α with β** , denoted $\alpha \otimes \beta : M \rightarrow T_{s_1+s_2}^{r_1+r_2} M$, by

$$(\alpha \otimes \beta)_p = \alpha_p \otimes \beta_p$$

Exercise 4.6.0.6. Let $\alpha_1 \in \Gamma_{s_1}^{r_1}(M)$ and $\alpha_2 \in \Gamma_{s_2}^{r_2}(M)$. Then $\alpha_1 \otimes \alpha_2 \in \Gamma_{s_1+s_2}^{r_1+r_2}(M)$

Proof. Let $\omega_1 \in \Gamma_1(M)^{r_1}$, $\omega_2 \in \Gamma_1(M)^{r_2}$, $X_1 \in \Gamma^1(M)^{s_1}$ and $X_2 \in \Gamma^1(M)^{s_2}$. By definition,

$$\alpha_1 \otimes \alpha_2(\omega_1, \omega_2, X_1, X_2) = \alpha_1(\omega_1, X_1)\alpha_2(\omega_2, X_2)$$

This implies that $\alpha_1 \otimes \alpha_2$ is smooth since α_1 and α_2 are smooth by assumption. □

Definition 4.6.0.7. We define the **tensor product**, denoted $\otimes : \Gamma_{s_1}^{r_1}(M) \times \Gamma_{s_2}^{r_2}(M) \rightarrow \Gamma_{s_1+s_2}^{r_1+r_2}(M)$ by

$$(\alpha_1, \alpha_2) \mapsto \alpha_1 \otimes \alpha_2$$

Exercise 4.6.0.8. The tensor product $\otimes : \Gamma_{s_1}^{r_1}(M) \times \Gamma_{s_2}^{r_2}(M) \rightarrow \Gamma_{s_1+s_2}^{r_1+r_2}(M)$ is associative.

Proof. Clear. □

Exercise 4.6.0.9. The tensor product $\otimes : \Gamma_{s_1}^{r_1}(M) \times \Gamma_{s_2}^{r_2}(M) \rightarrow \Gamma_{s_1+s_2}^{r_1+r_2}(M)$ is $C^\infty(M)$ -bilinear.

Proof. Clear. □

Definition 4.6.0.10. Let (N, \mathcal{B}) be a smooth manifold, $F : M \rightarrow N$ a smooth map and $\alpha \in \Gamma_k^0(N)$. We define the **pullback of α by F** , denoted $F^*\alpha \in \Gamma_k^0(M)$, by

$$(F^*\alpha)_p(v_1, \dots, v_k) = \alpha_{F(p)}(dF_p(v_1), \dots, dF_p(v_k))$$

for $p \in M$ and $v_1, \dots, v_k \in T_p M$

Exercise 4.6.0.11. Let (M, \mathcal{A}) , (N, \mathcal{B}) and (L, \mathcal{C}) be smooth manifolds, $F : M \rightarrow N$ and $G : N \rightarrow L$ smooth maps, $\alpha \in \Gamma_k^0(N)$, $\beta \in \Gamma_l^0(N)$, $\gamma \in \Gamma_k^0(L)$ and $f \in C^\infty(N)$. Then

1. $F^*(f\alpha) = (f \circ F)F^*\alpha$
2. $F^*(\alpha \otimes \beta) = F^*\alpha \otimes F^*\beta$
3. $F^*(\alpha + \beta) = F^*\alpha + F^*\beta$
4. $(G \circ F)^*\gamma = F^*(G^*\gamma)$
5. $id_N^*\alpha = \alpha$

Proof.

1.

$$\begin{aligned} [F^*(f\alpha)]_p(v_1, \dots, v_k) &= (f\alpha)_{F(p)}(dF_p(v_1), \dots, dF_p(v_k)) \\ &= f(F(p))\alpha_{F(p)}(dF_p(v_1), \dots, dF_p(v_k)) \\ &= (f \circ F)(p)(F^*\alpha)_p(v_1, \dots, v_k) \end{aligned}$$

So that $F^*(f\alpha) = (f \circ F)F^*\alpha$

2.

$$F^*$$

□

Definition 4.6.0.12.

Exercise 4.6.0.13.

Proof.

□

Exercise 4.6.0.14. Let $\alpha \in \Gamma_s^r(M)$ and $(U, \phi) \in \mathcal{A}$ with $\phi = (x^1, \dots, x^n)$. Then there exist $(f_J^I)_{I \in \mathcal{I}_r, J \in \mathcal{I}_s} \subset C^\infty(M)$ such that

$$\alpha|_U = \sum_{(I, J) \in \mathcal{I}_r \times \mathcal{I}_s} f_J^I \partial_{x^{\otimes I}} \otimes dx^{\otimes J}$$

Proof. Let $p \in M$. Then $\omega_p \in T_s^r(T_p M)$ and $\left\{ \partial_{x^{\otimes I}}|_p \otimes dx_p^{\otimes J} \right\}$ is a basis of $T_s^r(T_p M)$. So there exist $(f_J^I(p))_{I \in \mathcal{I}_r, J \in \mathcal{I}_s} \subset \mathbb{R}$ such that

$$\omega_p = \sum_{(I, J) \in \mathcal{I}_r \times \mathcal{I}_s} f_J^I(p) \partial_{x^{\otimes I}}|_p \otimes dx_p^{\otimes J}$$

Let $(K, L) \in \mathcal{I}_r \times \mathcal{I}_s$. Then

$$\begin{aligned} \alpha_p(dx_p^K, \partial_{x^L}|_p) &= \sum_{(I, J) \in \mathcal{I}_r \times \mathcal{I}_s} f_J^I(p) \partial_{x^{\otimes I}}|_p \otimes dx_p^{\otimes J}(dx_p^K, \partial_{x^L}|_p) \\ &= \sum_{(I, J) \in \mathcal{I}_r \times \mathcal{I}_s} f_J^I(p) \partial_{x^{\otimes I}}|_p(dx_p^K) dx_p^{\otimes J}(\partial_{x^L}|_p) \\ &= f_L^K(p) \end{aligned}$$

By assumption, the map $p \mapsto \alpha(dx_p^K, \partial_{x^L}|_p)$ is smooth, so that $f_L^K \in C^\infty(U)$.

□

Definition 4.6.0.15.

4.7 Differential Forms

Definition 4.7.0.1. We define

$$\Lambda^k(TM) = \coprod_{p \in M} \Lambda^k(T_p M)$$

Definition 4.7.0.2. Let $\omega : M \rightarrow \Lambda^k(TM)$. Then ω is said to be a **k -form on M** if for each $p \in M$, $\omega_p \in \Lambda^k(T_p M)$. For each $X \in \Gamma^1(M)^k$, we define $\omega(X) : M \rightarrow \mathbb{R}$ by

$$\omega(X)_p = \omega_p(X_p)$$

and ω is said to be **smooth** if for each $X \in \Gamma^1(M)^k$, $\omega(X)$ is smooth. The set of smooth k -forms on M is denoted $\Omega_k(M)$.

Note 4.7.0.3. Observe that

1. $\Omega_k(M) \subset \Gamma_k^0(M)$
2. $\Omega_0(M) = C^\infty(M)$

Exercise 4.7.0.4. The set $\Omega_k(M)$ is a $C^\infty(M)$ -submodule of $\Gamma_k^0(M)$.

Proof. Clear. □

Definition 4.7.0.5. Define the **exterior product**

$$\wedge : \Omega_k(M) \times \Omega_l(M) \rightarrow \Omega_{k+l}(M)$$

by

$$(\alpha \wedge \beta)_p = (\alpha)_p \wedge (\beta)_p$$

Note 4.7.0.6. For $f \in \Omega_0(M)$ and $\alpha \in \Omega_k(M)$, we have that $f \wedge \alpha = f\alpha$.

Exercise 4.7.0.7. The exterior product $\wedge : \Omega_k(M) \times \Omega_l(M) \rightarrow \Omega_{k+l}(M)$ is well defined.

Proof. Let $\alpha \in \Omega_k(M)$, $\beta \in \Omega_l(M)$, $(x^i)_{i=1}^k \subset \Gamma^1(M)$, $(y^j)_{j=1}^l \subset \Gamma^1(M)$ and $p \in M$. Then

$$\begin{aligned} \alpha \wedge \beta(X_1, \dots, X_{k+l})_p &= (\alpha \wedge \beta)_p(X_1(p), \dots, X_{k+l}(p)) \\ &= \frac{(k+l)!}{k!l!} \text{Alt}(\alpha_p \otimes \beta_p)(X_1(p), \dots, X_{k+l}(p)) \\ &= \frac{1}{k!l!} \sum_{\sigma \in S_{k+l}} \text{sgn}(\sigma) \sigma(\alpha_p \otimes \beta_p)(X_1(p), \dots, X_{k+l}(p)) \\ &= \frac{1}{k!l!} \sum_{\sigma \in S_{k+l}} \text{sgn}(\sigma) (\alpha_p \otimes \beta_p)(X_{\sigma(1)}(p), \dots, X_{\sigma(k+l)}(p)) \\ &= \frac{1}{k!l!} \sum_{\sigma \in S_{k+l}} \text{sgn}(\sigma) \alpha_p(X_{\sigma(1)}(p), \dots, X_{\sigma(k)}(p)) \beta(X_{\sigma(k+1)}(p), \dots, X_{\sigma(k+l)}(p)) \\ &= \frac{1}{k!l!} \sum_{\sigma \in S_{k+l}} \text{sgn}(\sigma) \alpha_p(X_{\sigma(1)}(p), \dots, X_{\sigma(k)}(p)) \beta(X_{\sigma(k+1)}(p), \dots, X_{\sigma(k+l)}(p)) \end{aligned}$$

□

Exercise 4.7.0.8. The exterior product $\wedge : \Omega_k(M) \times \Omega_l(M) \rightarrow \Omega_{k+l}(M)$ is $C^\infty(M)$ -bilinear.

Proof.

1. $C^\infty(M)$ -linearity in the first argument:

Let $\alpha \in \Omega_k(M)$, $\beta, \gamma \in \Omega_l(M)$, $f \in C^\infty(M)$ and $p \in M$. Bilinearity of $\wedge : \Lambda^k(T_p M) \times \Lambda^l(T_p M) \rightarrow \Lambda^{k+l}(T_p M)$ implies that

$$\begin{aligned} [(\beta + f\gamma) \wedge \alpha]_p &= (\beta + f\gamma)_p \wedge \alpha_p \\ &= (\beta_p + f(p)\gamma_p) \wedge \alpha_p \\ &= \beta_p \wedge \alpha_p + f(p)(\gamma_p \wedge \alpha_p) \\ &= [\beta \wedge \alpha + f(\gamma \wedge \alpha)]_p \end{aligned}$$

So that

$$(\beta + f\gamma) \wedge \alpha = \beta \wedge \alpha + f(\gamma \wedge \alpha)$$

and $\wedge : \Omega_k(M) \times \Omega_l(M) \rightarrow \Omega_{k+l}(M)$ is $C^\infty(M)$ -linear in the first argument.

2. $C^\infty(M)$ -linearity in the second argument:

Similar to (1).

□

Note 4.7.0.9. All of the results from multilinear algebra apply here.

Definition 4.7.0.10. We define the **exterior derivative** $d : \Omega_k(M) \rightarrow \Omega_{k+1}(M)$ inductively by

1. $d(d\alpha) = 0$ for $\alpha \in \Omega_p(M)$
2. $df(X) = Xf$ for $f \in \Omega_0(M)$
3. $d(\alpha \wedge \beta) = d\alpha \wedge \beta + (-1)^p \alpha \wedge d\beta$ for $\alpha \in \Omega_p(M)$ and $\beta \in \Omega_q(M)$
4. extending linearly

Exercise 4.7.0.11. Let (U, ϕ) be a chart on M with $\phi = (x^1, \dots, x^n)$. Then on U , for each $i, j \in \{1, \dots, n\}$,

$$dx^i \left(\frac{\partial}{\partial x^j} \right) = \delta_{i,j}$$

In particular, for each $p \in U$, $\{dx_p^1, \dots, dx_p^n\}$ is the dual basis to $\left\{ \frac{\partial}{\partial x^1} \Big|_p, \dots, \frac{\partial}{\partial x^n} \Big|_p \right\}$ and $T_p^* M = \text{span}\{dx_p^1, \dots, dx_p^n\}$.

Proof. Let $p \in U$ and $i, j \in \{1, \dots, n\}$. Then by definition,

$$\begin{aligned} \left[dx^i \left(\frac{\partial}{\partial x^j} \right) \right]_p &= \left(\frac{\partial}{\partial x^j} x^i \right)_p \\ &= \frac{\partial}{\partial x^j} \Big|_p x^i \\ &= \delta_{i,j} \end{aligned}$$

□

Exercise 4.7.0.12. Let $f \in C^\infty(M)$ and (U, ϕ) be a chart on M with $\phi = (x^1, \dots, x^n)$. Then

$$df|_U = \sum_{i=1}^n \frac{\partial f}{\partial x^i} dx^i$$

Proof. Let $p \in U$. Since $\{dx^1, \dots, dx^n\}$ is a basis for $\Lambda(T_p M)$, for each there exist $a_1(p), \dots, a_n(p) \in \mathbb{R}$ such that $df_p = \sum_{i=1}^n a^i(p) dx_p^i$. Therefore, we have that

$$\begin{aligned} df_p \left(\frac{\partial}{\partial x^i} \Big|_p \right) &= \sum_{i=1}^n a^i(p) dx_p^i \left(\frac{\partial}{\partial x^i} \Big|_p \right) \\ &= a_j(p) \end{aligned}$$

By definition, we have that

$$\begin{aligned} df_p \left(\frac{\partial}{\partial x^i} \Big|_p \right) &= \frac{\partial}{\partial x^i} \Big|_p f \\ &= \frac{\partial f}{\partial x^j}(p) \end{aligned}$$

So $a_j(p) = \frac{\partial f}{\partial x^j}(p)$ and

$$df_p = \sum_{i=1}^n \frac{\partial f}{\partial x^j}(p) dx_p^i$$

Therefore

$$df|_U = \sum_{i=1}^n \frac{\partial f}{\partial x^i} dx^i$$

□

Exercise 4.7.0.13. Let $f \in \Omega_0(M)$. If f is constant, then $df = 0$.

Proof. Suppose that f is constant. Let $p \in M$. Choose $(U, \phi) \in \mathcal{A}$ such that $p \in U$. Write $\phi = (x_1, \dots, x_n)$. Then for each $i \in \{1, \dots, n\}$,

$$\frac{\partial}{\partial x^i} \Big|_p f = 0$$

This implies that

$$\begin{aligned} df_p &= \sum_{i=1}^n \frac{\partial f}{\partial x^j}(p) dx_p^i \\ &= 0 \end{aligned}$$

□

Exercise 4.7.0.14.

Definition 4.7.0.15. Let $(U, \phi) \in \mathcal{A}$ with $\phi = (x^1, \dots, x^n)$ and $I = (i_1, \dots, i_k) \in \mathcal{I}_k$. We define

$$dx^i = dx_{i_1} \wedge \dots \wedge dx_{i_k} \in \Omega_k(M)$$

and we define

$$\frac{\partial}{\partial x^i} = \left(\frac{\partial}{\partial x^{i_1}}, \dots, \frac{\partial}{\partial x^{i_k}} \right)$$

Note 4.7.0.16. We have that

1.

$$dx^i \left(\frac{\partial}{\partial x^j} \right) = \delta_{I,J}$$

2. Since $\frac{\partial}{\partial x^i} \in \Gamma(U)^k$, by definition, for each $\omega \in \Omega_k(U)$,

$$\omega\left(\frac{\partial}{\partial x^i}\right) \in C^\infty(U)$$

Exercise 4.7.0.17. Let $\omega \in \Omega_k(M)$ and (U, ϕ) be a chart on M with $\phi = (x^1, \dots, x^n)$. Then

$$\omega = \sum_{I \in \mathcal{I}_k} \omega\left(\frac{\partial}{\partial x^i}\right) dx^i$$

Proof. Let $p \in U$. Since $\{dx_p^i : I \in \mathcal{I}_k\}$ is a basis for $\Lambda^k(T_p M)$, there exists $(f_I(p))_{I \in \mathcal{I}} \subset \mathbb{R}$ such that $\omega_p = \sum_{I \in \mathcal{I}_k} f_I(p) dx_p^i$. So for each $J \in \mathcal{I}_k$,

$$\begin{aligned} \omega\left(\frac{\partial}{\partial x^j}\right) &= \sum_{I \in \mathcal{I}_k} f_I dx^i \left(\frac{\partial}{\partial x^j}\right) \\ &= f_J \end{aligned}$$

□

Exercise 4.7.0.18. Let $\omega \in \Omega_k(M)$ and (U, ϕ) be a chart on M with $\phi = (x^1, \dots, x^n)$. If $\omega = \sum_{I \in \mathcal{I}_k} f_I dx^i$, then

$$d\omega = \sum_{I \in \mathcal{I}_k} \sum_{i=1}^n \frac{\partial f_I}{\partial x^i} dx^i \wedge dx^i$$

Proof. First we note that

$$\begin{aligned} d(f_I dx^i) &= df_I \wedge dx^i + (-1)^0 f d(dx^i) \\ &= df_I \wedge dx^i \\ &= \left(\sum_{i=1}^n \frac{\partial f_I}{\partial x^i} dx^i \right) \wedge dx^i \\ &= \sum_{i=1}^n \frac{\partial f_I}{\partial x^i} dx^i \wedge dx^i \end{aligned}$$

Then we extend linearly. □

Definition 4.7.0.19. Let (N, \mathcal{B}) be a smooth manifold and $F : M \rightarrow N$ be a diffeomorphism. Define the **pullback of F** , denoted $F^* : \Omega_k(N) \rightarrow \Omega_k(M)$ by

$$(F^*\omega)_p(v_1, \dots, v_k) = \omega_{F(p)}(dF_p(v_1), \dots, dF_p(v_k))$$

for $\omega \in \Omega_k(N)$, $p \in M$ and $v_1, \dots, v_k \in T_p M$

Chapter 5

Extra

Definition 5.0.0.1. When working in \mathbb{R}^n , we introduce the formal objects dx^1, dx_2, \dots, dx^n . Let $I = (i_1, i_2, \dots, i_k) \in \mathcal{I}_{k,n}$ and $\phi : \mathbb{R}^k \rightarrow \mathbb{R}^n$. Write $\phi = (\phi_1, \phi_2, \dots, \phi_n)$. We formally define $dx^i = dx_{i_1} \wedge dx_{i_2} \wedge \dots \wedge dx_{i_k}$ and $\phi_I = (\phi_{i_1}, \phi_{i_2}, \dots, \phi_{i_k})$.

Definition 5.0.0.2. Let $k \in \{0, 1, \dots, n\}$. We define a $C^\infty(\mathbb{R}^n)$ -module of dimension $\binom{n}{k}$, denoted $\Gamma^k(\mathbb{R}^n)$ to be

$$\Phi_k(\mathbb{R}^n) = \begin{cases} C^\infty(\mathbb{R}^n) & k = 0 \\ \text{span}\{dx^i : I \in \mathcal{I}_{k,n}\} & k \geq 1 \end{cases}$$

For each $\omega \in \Phi_k(\mathbb{R}^n)$ and $\chi \in \Gamma^l(\mathbb{R}^n)$, we may form their **exterior product**, denoted by $\omega \wedge \chi \in \Gamma^{k+l}(\mathbb{R}^n)$. Thus the exterior product is a map $\wedge : \Phi_k(\mathbb{R}^n) \times \Gamma^l(\mathbb{R}^n) \rightarrow \Gamma^{k+l}(\mathbb{R}^n)$. The exterior product is characterized by the following properties:

1. the exterior product is bilinear
2. for each $\omega \in \Phi_k(\mathbb{R}^n)$ and $\chi \in \Gamma^l(\mathbb{R}^n)$, $\omega \wedge \chi = -\chi \wedge \omega$
3. for each $\omega \in \Phi_k(\mathbb{R}^n)$, $\omega \wedge \omega = 0$
4. for each $f \in C^\infty(\mathbb{R}^n)$ and $\omega \in \Phi_k(\mathbb{R}^n)$, $f \wedge \omega = f\omega$

We call $\Phi_k(\mathbb{R}^n)$ the differential k -forms on \mathbb{R}^n . Let ω be a k -form on \mathbb{R}^n . If $k \geq 1$, then for each $I \in \mathcal{I}_{k,n}$, there exists $f_I \in C^\infty(\mathbb{R}^n)$ such that $\omega = \sum_{I \in \mathcal{I}_{k,n}} f_I dx^i$

Note 5.0.0.3. The terms dx^1, dx_2, \dots, dx^n are a sort of place holder for the coordinates of a point $x = (x_1, x_2, \dots, x_n) \in \mathbb{R}^n$. When we work with functions $\phi : \mathbb{R}^k \rightarrow \mathbb{R}^n$, we will have different coordinates and to avoid confusion, we will write $\{du^1, du_2, \dots, du_k\}$ when referencing the coordinates on \mathbb{R}^k and $\{dx^1, dx_2, \dots, dx^n\}$ when referencing the coordinates on \mathbb{R}^n .

Exercise 5.0.0.4. Let $B_{n \times n} = (b_{i,j}) \in [C^\infty(M)]^{n \times n}$ be an $n \times n$ matrix. Then

$$\bigwedge_{i=1}^n \left(\sum_{j=1}^n b_{i,j} dx^j \right) = (\det B) dx^1 \wedge dx_2 \wedge \dots \wedge dx^n$$

Proof. Bilinearity of the exterior product implies that

$$\begin{aligned}
 \bigwedge_{i=1}^n \left(\sum_{j=1}^n b_{i,j} dx^j \right) &= \left(\sum_{j=1}^n b_{1,j} dx^j \right) \wedge \left(\sum_{j=1}^n b_{2,j} dx^j \right) \wedge \cdots \wedge \left(\sum_{j=1}^n b_{n,j} dx^j \right) \\
 &= \sum_{j_1, \dots, j_n=1}^n \left(\prod_{i=1}^n b_{i,j_i} \right) dx_{j_1} \wedge dx_{j_2} \wedge \cdots \wedge dx_{j_n} \\
 &= \sum_{j_1 \neq \dots \neq j_n} \left(\prod_{i=1}^n b_{i,j_i} \right) dx_{j_1} \wedge dx_{j_2} \wedge \cdots \wedge dx_{j_n} \\
 &= \left[\sum_{\sigma \in S_n} \text{sgn}(\sigma) \left(\prod_{i=1}^n b_{i,\sigma(i)} \right) \right] dx_1 \wedge dx_2 \wedge \cdots \wedge dx_n \\
 &= (\det B) dx_1 \wedge dx_2 \wedge \cdots \wedge dx_n
 \end{aligned}$$

□

Definition 5.0.0.5. Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be a 0-form on \mathbb{R}^n . We define a 1-form, denoted df , on \mathbb{R}^n by

$$df = \sum_{i=1}^n \frac{\partial f}{\partial x^i} dx^i$$

Let $\omega = \sum_{I \in \mathcal{I}_{k,n}} f_I dx^I$ be a k -form on \mathbb{R}^n . We can define a differential $k+1$ -form, denoted $d\omega$, on \mathbb{R}^n by

$$d\omega = \sum_{I \in \mathcal{I}_{k,n}} df_I \wedge dx^I$$

Exercise 5.0.0.6. On \mathbb{R}^3 , put

1. $\omega_0 = f_0$,
2. $\omega_1 = f_1 dx^1 + f_2 dx^2 + f_3 dx^3$,
3. $\omega_2 = f_1 dx^2 \wedge dx^3 - f_2 dx^1 \wedge dx^3 + f_3 dx^1 \wedge dx^2$

Show that

1. $d\omega_0 = \frac{\partial f_0}{\partial x^1} dx^1 + \frac{\partial f_0}{\partial x^2} dx^2 + \frac{\partial f_0}{\partial x^3} dx^3$
2. $d\omega_1 = \left(\frac{\partial f_3}{\partial x^2} - \frac{\partial f_2}{\partial x^3} \right) dx^2 \wedge dx^3 + \left(\frac{\partial f_3}{\partial x^1} - \frac{\partial f_1}{\partial x^3} \right) dx^1 \wedge dx^3 + \left(\frac{\partial f_2}{\partial x^1} - \frac{\partial f_1}{\partial x^2} \right) dx^1 \wedge dx^2$
3. $d\omega_2 = \left(\frac{\partial f_1}{\partial x^1} + \frac{\partial f_2}{\partial x^2} + \frac{\partial f_3}{\partial x^3} \right) dx^1 \wedge dx^2 \wedge dx^3$

Proof. Straightforward. □

Exercise 5.0.0.7. Let $I \in \mathcal{I}_{k,n}$. Then there is a unique $I_* \in \mathcal{I}_{n-k,n}$ such that $dx^I \wedge dx_{I_*} = dx^1 \wedge dx^2 \wedge \cdots \wedge dx^n$.

Definition 5.0.0.8. We define a linear map $*$: $\Phi_k(\mathbb{R}^n) \rightarrow \Gamma^{n-k}(\mathbb{R}^n)$ called the **Hodge *-operator** by

$$* \sum_{I \in \mathcal{I}_{k,n}} f_I dx^I = \sum_{I \in \mathcal{I}_{k,n}} f_I dx_{I_*}$$

Definition 5.0.0.9. Let $\phi : \mathbb{R}^k \rightarrow \mathbb{R}^n$ be smooth. Write $\phi = (\phi_1, \phi_2, \dots, \phi_n)$. We define $\phi^* : \Phi_k(\mathbb{R}^n) \rightarrow \Phi_k(\mathbb{R}^k)$ via the following properties:

1. for each 0-form f on \mathbb{R}^n , $\phi^* f = f \circ \phi$

2. for $i = 1, \dots, n$, $\phi^* dx^i = d\phi_i$
3. for an s -form ω , and a t -form χ on \mathbb{R}^n , $\phi^*(\omega \wedge \chi) = (\phi^*\omega) \wedge (\phi^*\chi)$
4. for l -forms ω, χ on \mathbb{R}^n , $\phi^*(\omega + \chi) = \phi^*\omega + \phi^*\chi$

Exercise 5.0.0.10. Let $M \subset \mathbb{R}^n$ be a k -dimensional smooth submanifold of \mathbb{R}^n , $\phi : U \rightarrow V$ a smooth parametrization of M , $\omega = \sum_{I \in \mathcal{I}_{k,n}} f_I dx^I$ a k -form on \mathbb{R}^n . Then

$$\phi^*\omega = \left(\sum_{I \in \mathcal{I}_{k,n}} (f_I \circ \phi)(\det v\phi_I) \right) du^1 \wedge du_2 \wedge \dots \wedge du_k$$

Proof. By definition,

$$\begin{aligned} \phi^*\omega &= \phi^* \sum_{I \in \mathcal{I}_{k,n}} f_I dx^I \\ &= \sum_{I \in \mathcal{I}_{k,n}} (\phi^* f_I) \phi^* dx^I \\ &= \sum_{I \in \mathcal{I}_{k,n}} (f_I \circ \phi) d\phi_I \end{aligned}$$

A previous exercise tells us that for each $I \in \mathcal{I}_{k,n}$,

$$\begin{aligned} d\phi_I &= d\phi_{i_1} \wedge d\phi_{i_2} \wedge \dots \wedge d\phi_{i_n} \\ &= \left(\sum_{j=1}^n \frac{\partial \phi_{i_1}}{\partial u^j} du^j \right) \wedge \left(\sum_{j=1}^n \frac{\partial \phi_{i_2}}{\partial u^j} du^j \right) \wedge \dots \wedge \left(\sum_{j=1}^n \frac{\partial \phi_{i_k}}{\partial u^j} du^j \right) \\ &= (\det v\phi_I) du^1 \wedge du_2 \wedge \dots \wedge du_k \end{aligned}$$

Therefore

$$\begin{aligned} \phi^*\omega &= \sum_{I \in \mathcal{I}_{k,n}} (f_I \circ \phi) d\phi_I \\ &= \sum_{I \in \mathcal{I}_{k,n}} (f_I \circ \phi)(\det v\phi_I) du^1 \wedge du_2 \wedge \dots \wedge du_k \\ &= \left(\sum_{I \in \mathcal{I}_{k,n}} (f_I \circ \phi)(\det v\phi_I) \right) du^1 \wedge du_2 \wedge \dots \wedge du_k \end{aligned}$$

□

5.1 Integration of Differential Forms

Definition 5.1.0.1. Let $U \subset \mathbb{R}^k$ be open and $\omega = f dx^1 \wedge dx_2 \wedge \dots \wedge dx_k$ a k -form on \mathbb{R}^k . Define

$$\int_U \omega = \int_U f dx$$

Definition 5.1.0.2. Let $M \subset \mathbb{R}^n$ be a k -dimensional oriented smooth submanifold of \mathbb{R}^n , ω a k -form on \mathbb{R}^n and $\phi : U \rightarrow V$ a local smooth, orientation-preserving parametrization of M . Define

$$\int_V \omega = \int_U \phi^*\omega$$

Exercise 5.1.0.3.

Theorem 5.1.0.4. Stokes Theorem:

Let $M \subset \mathbb{R}^n$ be a k -dimensional oriented smooth submanifold of \mathbb{R}^n and ω a $k-1$ -form on \mathbb{R}^n . Then

$$\int_{\partial M} \omega = \int_M d\omega$$