

# INTRODUCTION TO BAYESIAN STATISTICS

CARSON JAMES

## CONTENTS

1. Introduction	1
2. Sampling	1
2.1. Inverse CDF Sampling	1
2.2. Importance Sampling	1
2.3. Rejection Sampling	1

## 1. INTRODUCTION

**Definition 1.0.1.** We define

$$\mathcal{D}(\mathbb{R}^n) = \{f \in L^1(\mathbb{R}^d) : f \geq 0 \text{ and } \|f\|_1 = 1\}$$

## 2. SAMPLING

**2.1. Inverse CDF Sampling.**

**2.2. Importance Sampling.**

**2.3. Rejection Sampling.**

**Exercise 2.3.1.** Let  $f, g \in \mathcal{D}(\mathbb{R}^d)$  and  $A \in \mathcal{B}(\mathbb{R}^d)$ . Suppose that  $m^d(A) > 0$ . If  $X \sim f$ , then  $X|X \in A \sim \|fI_A\|_1^{-1} fI_A$ .

*Proof.* Let  $C \in \mathcal{B}(\mathbb{R}^d)$ . Then

$$\begin{aligned} P(X \in C|X \in A) &= P(X \in C \cap A)P(X \in A)^{-1} \\ &= \|fI_A\|_1^{-1} \int_C fI_A dm^d \end{aligned}$$

So  $f_{X|X \in A} = \|fI_A\|_1^{-1} fI_A$ . □

**Exercise 2.3.2.** Let  $A, B \in \mathcal{B}(\mathbb{R}^d)$ . Suppose that  $A \subset B$  and  $0 < m^d(A)$  and  $m^d(B) < \infty$ . If  $X \sim \text{Uni}(B)$ , then  $X|X \in A \sim \text{Uni}(A)$ .

*Proof.* Clear using the previous exercise with  $f = I_B$ . □

**Exercise 2.3.3. (Fundamental Theorem of Simulation):**

Let  $f \in \mathcal{D}(\mathbb{R}^d)$  and  $c > 0$ . Define

$$G_c = \{(x, v) \in \mathbb{R}^{d+1} : 0 < v < cf(x)\}$$

(1) If  $X \sim f$  and  $U \sim \text{Uni}(0, 1)$  are independent, then  $(X, cUf(X)) \sim \text{Uni}(G_c)$ .

(2) If  $(X, V) \sim \text{Uni}(G_c)$ , then  $X \sim f$ .

*Proof.* First we note that  $m^{d+1}(G_c) = c$ .

(1) Suppose that  $X \sim f$  and  $U \sim \text{Uni}(0, 1)$  are independent and put  $Y = cUf(X)$ . Then  $Y|X = x \sim cUf(x) \sim \text{Uni}(0, cf(x))$  and we have that for each  $x \in \text{supp } X$  and  $y \in (0, cf(x))$ ,

$$\begin{aligned} f_{X,Y}(x, y) &= f_{Y|X}(y|x)f(x) \\ &= \frac{1}{cf(x)}f(x) \\ &= \frac{1}{c} \end{aligned}$$

So  $(X, Y) \sim \text{Uni}(G_c)$

(2) Suppose that  $(X, V) \sim \text{Uni}(G_c)$ . Then  $f_{X,V}(x, v) = \frac{1}{c}I_{G_c}(x, v)$ . So

$$\begin{aligned} f_X(x) &= \int_{\mathbb{R}} \frac{1}{c}I_{G_c}(x, v)dm(v) \\ &= \int_0^{cf(x)} \frac{1}{c}dv \\ &= f(x) \end{aligned}$$

So  $X \sim f$ .

□

**Exercise 2.3.4.** Let  $f, g \in \mathcal{D}(\mathbb{R}^d)$ ,  $c_f, c_g > 0$  and  $M > 0$ . Put  $\tilde{f} = c_f f$  and  $\tilde{g} = c_g g$ . Suppose that  $\tilde{f} \leq M\tilde{g}$ . If  $Y \sim g$  and  $U \sim \text{Uni}(0, 1)$  are independent, then  $Y|U \leq \frac{\tilde{f}(Y)}{M\tilde{g}(Y)} \sim f$  and  $P(U \leq \frac{\tilde{f}(Y)}{M\tilde{g}(Y)}) = \frac{c_f}{c_g M}$

*Proof.* Put

$$G_g = \{(y, v) \in \mathbb{R}^{d+1} : 0 < v < M\tilde{g}(y)\}$$

and

$$G_f = \{(y, v) \in \mathbb{R}^{d+1} : 0 < v < \tilde{f}(y)\}$$

Then  $G_f \subset G_g$ ,  $m^d(G_g) = c_g M$  and  $m^d(G_f) = c_f$ . By the first part of the fundamental theorem of simulation, we know that

$$(Y, MUc_g g(Y)) \sim \text{Uni}(G_g)$$

Since  $\{(Y, MUc_g g(Y)) \in G_f\} = \{U \leq \frac{c_f f(Y)}{M c_g g(Y)}\}$ , a previous exercise tells us that

$$(Y, MUc_g g(Y))|U \leq \frac{c_f f(Y)}{M c_g g(Y)} \sim \text{Uni}(G_f)$$

Then the second part of the fundamental theorem of simulation tells us that

$$Y|U \leq \frac{c_f f(Y)}{M c_g g(Y)} \sim f$$

Finally we have that

$$\begin{aligned} P\left(U \leq \frac{c_f f(Y)}{M c_g g(Y)}\right) &= P[(Y, M U c_g g(Y)) \in G_f] \\ &= \frac{c_f}{c_g M} \end{aligned}$$

□

**Definition 2.3.5. (Rejection Sampling Algorithm):**

Let  $f, g \in \mathcal{D}(\mathbb{R}^d)$ ,  $c_f, c_g > 0$  and  $M > 0$ . Put  $\tilde{f} = c_f f$  and  $\tilde{g} = c_g g$ . Suppose that  $\tilde{f} \leq M \tilde{g}$ . We define the **rejection sampling algorithm** as follows:

- (1) sample  $Y \sim g$  and  $U \sim \text{Uni}(0, 1)$  independently
- (2) if  $U \leq \frac{\tilde{f}(Y)}{M \tilde{g}(Y)}$ , accept  $Y$ , else return to (1).

If we sample  $(X_n)_{n \in \mathbb{N}}$  independently using the rejection sampler, then the previous exercises imply that  $(X_n)_{n \in \mathbb{N}} \stackrel{iid}{\sim} f$  and the acceptance rate is  $\frac{c_f}{c_g M}$ .

*Note 2.3.1.* Phrasing the rejection sampler in terms of  $\tilde{f}$  and  $\tilde{g}$  instead of  $f$  and  $g$  is useful because we may not always be able to solve for the normalizing constants.