

Introduction to Analysis

Carson James

Contents

Notation	xi
Preface	1
1 Set Theory	3
1.1 To Do	3
1.2 Sets	3
1.2.1 Union and Intersection	3
1.2.2 Covers	5
1.3 Functions	6
1.3.1 Introduction	6
1.3.2 Bijections	8
1.3.3 Nets	10
1.3.4 Sequences	10
1.4 Subsets	11
1.5 Products of Sets	12
1.6 Coproducts of Sets	17
1.7 Quotients of Sets	19
1.8 Equalizers of Maps	20
1.9 Projective Limits of Sets	21
2 Ordered Sets	23
2.1 Preorders	23
2.2 Directed Sets	25
2.3 Posets	26
3 Real and Complex Numbers	27
3.1 Real Numbers	27
3.2 Extended Real Numbers	28
3.3 Complex Numbers	28
I Topology	29
4 Topology	31
4.1 Introduction	31
4.2 Continuous Maps	43
4.3 Nets	51
4.3.1 Common Directed Sets	51
4.3.2 Nets in Topological Spaces	52
4.4 Filters	59
4.5 Subspace Topology	60
4.5.1 Introduction	60

4.5.2	Discrete Spaces	67
4.5.3	Topological Embeddings	67
4.6	Product Topology	68
4.6.1	Bases of the product topology	68
4.6.2	Characteristics of the Product Topology	70
4.6.3	Maps and the Product Topology	71
4.7	Coproduct Topology	74
4.8	Quotient Topology	79
4.8.1	Introduction	79
4.8.2	Category of Topological Spaces with Equivalence Relations	85
4.9	Equalizers of Continuous Maps	87
4.10	Projective Limits of Topological Spaces	88
4.11	Open Covers	92
4.11.1	Introduction	92
4.11.2	Subspace Topology	93
4.12	Separation Axioms	94
4.12.1	Introduction	94
4.12.2	Separation and Subspaces	97
4.12.3	Separation and Product Spaces	97
4.12.4	Separation and Quotient Spaces	98
4.13	Countability Axioms	102
4.13.1	First-Countability	102
4.13.2	Second-Countability	105
	Second-Countability and Subspaces	107
	Second-Countability and Product Spaces	108
	Second-Countability and Coproduct Spaces	108
	Second-Countability and Quotient Spaces	108
	Second-Countability and Projective Limits	108
4.14	Compactness	110
4.14.1	Compactness and Subspaces	110
4.14.2	The Finite Intersection Property	114
4.14.3	Compactness and Product Spaces	118
4.14.4	Compactness and Coproduct Spaces	121
4.14.5	Compactness and Quotient Spaces	121
4.14.6	Compactness and Projective Systems	122
4.14.7	Compactness and Continuity	122
4.14.8	Sequential Compactness	124
4.15	Locally Compact Hausdorff Spaces	126
4.15.1	Introduction	126
4.15.2	Arzela-Ascoli Theorem	127
4.15.3	Stone-Weierstrass Theorem	127
4.16	Tychonoff Spaces	128
4.17	Compactification	129
4.18	Zero-Dimensional Spaces	130
4.18.1	Zero-Dimensional Subspaces	130
4.18.2	Zero-Dimensional Product Spaces	130
4.18.3	Zero-Dimensional Coproduct Spaces	131
4.19	Semi-continuity	132
5	Locale Theory	135
5.1	Sober Spaces	135
5.2	Frames and Locales	137
5.2.1	Introdction	137
5.2.2	Sublocales	140

6	Metric Spaces	141
6.1	Introduction	141
6.1.1	Metrics	141
6.1.2	Metric Topology	146
6.1.3	Uniform and Lipschitz Continuity	151
6.1.4	Stuff to Reorder or Move	155
6.1.5	Isometries	159
6.2	Countability and Separation	162
6.2.1	Countability	162
6.2.2	Separation	165
6.3	Top -Equivalent Metrics	166
6.3.1	Subadditive Functions	166
6.3.2	Top Metric Preserving Functions	166
6.4	Subspaces	170
6.4.1	Introduction	170
6.4.2	Discrete Subsets	172
6.5	Product Spaces	174
6.6	Coproduct Spaces	176
6.7	Completeness	178
6.7.1	Introduction	178
6.7.2	Completeness and Subspaces	185
6.7.3	Completeness and Product Spaces	186
6.7.4	Completeness and Coproduct Spaces	188
6.8	Compactness	189
6.9	The Baire Category Theorem	196
6.10	Metric Spaces of Functions	198
6.10.1	Bounded Maps	198
6.10.2	Bounded Continuous Maps	199
6.10.3	Subspaces	200
6.10.4	Completeness	201
6.10.5	Continuous Maps on LCH spaces	203
6.10.6	Continuous Maps Compactifications	203
6.11	Metrizible Spaces	204
6.11.1	Introduction	204
6.11.2	Metrizability of Subspaces	205
6.11.3	Metrizability of Product Spaces	206
6.11.4	Metrizability of Coproduct Spaces	206
6.11.5	Metrizability of Projective Limits	206
6.12	Polish Spaces	207
6.12.1	Introduction	207
6.13	Ultrametric Spaces	209
6.13.1	Introduction	209
6.13.2	Partitions	212
6.14	Zero-dimensional Metric Spaces	221
6.14.1	Second-Countability	221
6.14.2	Polish Spaces	223
6.14.3	Ultrametric Spaces	224
6.14.4	Cantor Space	224
7	Analysis on $(C(X), \ \cdot\ _\infty)$	229
7.1	LEFTOVERS	230

II	Topological Vector Spaces	233
8	Topological Vector Spaces	235
8.1	Introduction	235
8.1.1	Topology	235
8.1.2	Continuous Linear Functionals	237
8.1.3	Products	239
8.2	Sublinear Functionals	241
8.3	Seminorms	244
8.4	Minkowski Functionals	251
8.5	Locally Convex Spaces	258
8.6	Continuous Linear Maps	263
8.7	Subspaces	264
8.8	Products	266
8.9	Direct Sums	267
8.10	Quotient Spaces	268
8.11	Compactness	270
8.12	Duality	272
8.12.1	Weak and Weak-* Topology	272
8.12.2	Annihilators	276
8.12.3	Adjoint	277
9	Banach Spaces	279
9.1	Introduction	279
9.2	Bounded Operators	282
9.3	Direct Sums	288
9.4	Quotient Spaces	291
9.5	Applications of the Hahn-Banach Theorem	295
9.6	Applications of the Baire Category Theorem	299
9.7	Duality	304
9.8	Compact Operators	312
9.9	Multilinear Maps	313
9.10	Tensor Products of Banach Spaces	318
9.11	Injective Tensor Product	318
9.12	Projective Tensor Product	319
9.13	Unbounded Operators	323
9.13.1	Closed Operators	324
10	Hilbert Spaces	325
10.1	TODO	325
10.2	Introduction	325
10.2.1	Inner product Spaces	325
10.2.2	Riesz Representation Theorem	343
10.3	Operators on Hilbert Spaces	348
10.3.1	Adjoint Operators	348
10.3.2	Unitary Maps	349
10.3.3	Spectral Theory (move to section on compact operators on hilbert space or banach algebra)	351
10.4	Subspaces of Hilbert Spaces	354
10.4.1	Introduction	354
10.4.2	Orthogonal Projections	354
10.5	Direct Sums of Hilbert spaces	358
10.6	Tensor Products	362
10.7	Rigged Hilbert Spaces	370
10.8	MISC, unitary transformations	371

III	Differentiation	373
11	Differentiation	375
11.1	TODO	376
11.2	The Gateaux Derivative	377
11.3	The Frechet Derivative	381
11.4	The Calc I Derivative	385
11.5	Mean Value Theorem	387
11.6	Taylor's Theorem	389
11.7	Implicit and Inverse Function Theorems	394
11.8	The Gradient	396
12	Banach Lattices	397
12.1	Introduction	397
12.1.1	Ordered Vector Spaces	397
12.1.2	Riesz Spaces	400
13	Banach Algebras	401
13.1	Introduction	401
13.2	Spectral Theory	408
13.2.1	*-Algebras	417
13.3	Exponential Map	418
14	Differential Equations	419
14.1	Introduction	419
14.2	Semigroup Theory	423
15	Convex Analysis	425
15.1	Subadditive Functions	425
15.2	Convex Sets	428
15.2.1	Introduction	428
15.2.2	Cones	433
15.3	Convex Functions	434
15.4	The Subdifferential	439
15.5	Conjugacy	445
15.5.1	Convex Conjugate	445
15.5.2	Convex Envelope	448
16	Topological Groups	449
16.1	Introduction	449
16.2	Group Actions	452
16.2.1	Introduction	452
16.2.2	Homogeneous Spaces	452
16.2.3	Common Examples	453
16.3	Quotient Groups	454
16.4	Automorphism Groups of Metric Spaces	455
17	Group Actions	457
17.1	Introduction	457
17.2	Group Actions on Metric Spaces	459
17.3	Fundamental Examples	463
17.4	Optimization over Compact Groups	465

Summation	467
.1 Summation of Positive Functions	467
.2 Summation of Complex-Valued Functions	468
Asymptotic Notation	469
Categories	473
Rings	475
.3 Introduction	475
.4 Quotient Rings	475

Notation

$\mathcal{M}_+(X, \mathcal{A})$	finite measures on $(X, \mathcal{A})$
v	velocity

Preface

cc-by-nc-sa

Chapter 1

Set Theory

1.1 To Do

- Need to merge these notes with set theory appendices and intro to math notes, to do this, need to make exercise references work out, which is the issue.

figure out how to structure discussion on orderings, the issue is that projective systems of sets rely on directed sets, directed sets need products of sets, order theory and the specialization topology need topology and separation axioms, maybe each section should list prereqs and put sections with more prereqs towards end of chapter, then we could have a chapter on orderings that cover directed sets that comes after the set theory chapter and in the section on projective systems, include an intro with the prereq material from the section on directed sets, or maybe just make orderings a separate set of notes about order theory and reference these notes. Honestly chapters should be indexed by a poset instead of a totally ordered set.

1.2 Sets

1.2.1 Union and Intersection

Definition 1.2.1.1. Let X, A be sets and $(E_\alpha)_{\alpha \in A} \subset \mathcal{P}(X)$. We define the

- **union of** $(E_\alpha)_{\alpha \in A}$, denoted $\bigcup_{\alpha \in A} E_\alpha \in \mathcal{P}(X)$, by

$$\bigcup_{\alpha \in A} E_\alpha := \{x \in X : \text{there exists } \alpha \in A \text{ such that } x \in E_\alpha\}$$

- **intersection of** $(E_\alpha)_{\alpha \in A}$, denoted $\bigcap_{\alpha \in A} E_\alpha \in \mathcal{P}(X)$, by

$$\bigcap_{\alpha \in A} E_\alpha := \{x \in X : \text{for each } \alpha \in A, x \in E_\alpha\}$$

Exercise 1.2.1.2. Let X, A be sets and $(E_\alpha)_{\alpha \in A} \subset \mathcal{P}(X)$. If $A = \emptyset$, then

1. $\bigcup_{\alpha \in A} E_\alpha = \emptyset$
2. $\bigcap_{\alpha \in A} E_\alpha = X$

Proof. Suppose that $A = \emptyset$.

1. Let $x \in X$. For the sake of contradiction, suppose that $x \in \bigcup_{\alpha \in A} E_\alpha$. Then there exists $\alpha \in A$ such that $x \in E_\alpha$. This is a contradiction since $A = \emptyset$. Hence $x \in \left(\bigcup_{\alpha \in A} E_\alpha \right)^c$. Since $x \in X$ is arbitrary, we have that for each $x \in X$, $x \in \left(\bigcup_{\alpha \in A} E_\alpha \right)^c$. Therefore $X \subset \left(\bigcup_{\alpha \in A} E_\alpha \right)^c$. Thus

$$\begin{aligned} \bigcup_{\alpha \in A} E_\alpha &\subset X^c \\ &= \emptyset. \end{aligned}$$

Since trivially, $\emptyset \subset \bigcup_{\alpha \in A} E_\alpha$, we have that $\bigcup_{\alpha \in A} E_\alpha = \emptyset$.

2. Let $x \in X$. Since $A = \emptyset$, it is vacuously true that for each $\alpha \in A$, $x \in E_\alpha$. Thus $x \in \bigcap_{\alpha \in A} E_\alpha$. Since $x \in X$ is arbitrary, we have that for each $x \in X$, $x \in \bigcap_{\alpha \in A} E_\alpha$. Hence $X \subset \bigcap_{\alpha \in A} E_\alpha$. Since trivially $\bigcap_{\alpha \in A} E_\alpha \subset X$, we have that $\bigcap_{\alpha \in A} E_\alpha = X$.

□

Exercise 1.2.1.3. Let X, A be sets and $(E_\alpha)_{\alpha \in A} \subset \mathcal{P}(X)$. Then

1. $\left(\bigcup_{\alpha \in A} E_\alpha \right)^c = \bigcap_{\alpha \in A} E_\alpha^c$
2. $\left(\bigcap_{\alpha \in A} E_\alpha \right)^c = \bigcup_{\alpha \in A} E_\alpha^c$

Proof.

1. We note that for each $x \in X$,

$$\begin{aligned} x \in \left(\bigcup_{\alpha \in A} E_\alpha \right)^c &\iff x \notin \bigcup_{\alpha \in A} E_\alpha \\ &\iff \text{it is not the case that } x \in \bigcup_{\alpha \in A} E_\alpha \\ &\iff \text{it is not the case that there exists } \alpha \in A \text{ such that } x \in E_\alpha \\ &\iff \text{for each } \alpha \in A, x \notin E_\alpha \\ &\iff \text{for each } \alpha \in A, x \in E_\alpha^c \\ &\iff x \in \bigcap_{\alpha \in A} E_\alpha^c. \end{aligned}$$

Thus $\left(\bigcup_{\alpha \in A} E_\alpha \right)^c = \bigcap_{\alpha \in A} E_\alpha^c$.

2. **The previous part** implies that

$$\begin{aligned} \left(\bigcup_{\alpha \in A} E_\alpha^c \right)^c &= \bigcap_{\alpha \in A} (E_\alpha^c)^c \\ &= \bigcap_{\alpha \in A} E_\alpha. \end{aligned}$$

Therefore

$$\begin{aligned} \left(\bigcap_{\alpha \in A} E_\alpha \right)^c &= \left[\left(\bigcup_{\alpha \in A} E_\alpha^c \right)^c \right]^c \\ &= \bigcup_{\alpha \in A} E_\alpha^c. \end{aligned}$$

□

Exercise 1.2.1.4. Let X, A be sets, $E \in \mathcal{P}(X)$ and $(F_\alpha)_{\alpha \in A} \subset \mathcal{P}(X)$. Then

1. $E \cap \left(\bigcup_{\alpha \in A} F_\alpha \right) = \bigcup_{\alpha \in A} (E \cap F_\alpha)$
2. $E \cup \left(\bigcap_{\alpha \in A} F_\alpha \right) = \bigcap_{\alpha \in A} (E \cup F_\alpha)$

Proof.

1. We note that for each $x \in X$,

$$\begin{aligned}
 x \in E \cap \left(\bigcup_{\alpha \in A} F_\alpha \right) &\iff x \in E \text{ and } x \in \bigcup_{\alpha \in A} F_\alpha \\
 &\iff x \in E \text{ and there exists } \alpha \in A \text{ such that } x \in F_\alpha \\
 &\iff \text{there exists } \alpha \in A \text{ such that } x \in E \text{ and } x \in F_\alpha \\
 &\iff \text{there exists } \alpha \in A \text{ such that } x \in E \cap F_\alpha \\
 &\iff x \in \bigcup_{\alpha \in A} (E \cap F_\alpha).
 \end{aligned}$$

$$\text{Therefore } E \cap \left(\bigcup_{\alpha \in A} F_\alpha \right) = \bigcup_{\alpha \in A} (E \cap F_\alpha).$$

2. We note that for each $x \in X$,

$$\begin{aligned}
 x \in E \cup \left(\bigcap_{\alpha \in A} F_\alpha \right) &\iff x \in E \text{ or } x \in \bigcap_{\alpha \in A} F_\alpha \\
 &\iff x \in E \text{ or for each } \alpha \in A, x \in F_\alpha \\
 &\iff \text{for each } \alpha \in A, x \in E \text{ or } x \in F_\alpha \\
 &\iff \text{for each } \alpha \in A, x \in E \cup F_\alpha \\
 &\iff x \in \bigcap_{\alpha \in A} (E \cup F_\alpha).
 \end{aligned}$$

□

1.2.2 Covers

Definition 1.2.2.1. Let X be a set and $E \subset X$.

Let $\mathcal{U} \subset \mathcal{P}(X)$. Then $\mathcal{U}$ is said to be a **cover of E in X** if $E \subset \bigcup_{U \in \mathcal{U}} U$.

We define $\text{Cover}_X(E) \subset \mathcal{T}$ by

$$\text{Cover}(E, \mathcal{T}) := \{\mathcal{U} \subset \mathcal{P}(X) : \mathcal{U} \text{ is a cover of } E \text{ in } X\}.$$

Note 1.2.2.2. When the context is clear,

- we may say $\mathcal{U}$ is a “cover of E ” instead of a “cover of E in X ”
- we may write $\text{Cover}(E)$ in place of $\text{Cover}_X(E)$.

Exercise 1.2.2.3. Let X be a set, $A \subset X$ and $\mathcal{U}, \mathcal{V} \in \text{Cover}_X(A)$. Suppose that $\mathcal{V}$ is finite. If for each $\mathcal{U}_0 \subset \mathcal{U}$, $\mathcal{U}_0$ is finite implies that $\mathcal{U}_0 \notin \text{Cover}_X(A)$, then there exists $V \in \mathcal{V}$ such that for each $\mathcal{U}_0 \subset \mathcal{U}$, $\mathcal{U}_0$ is finite implies that $\mathcal{U}_0 \notin \text{Cover}_X(V)$.

Proof. Suppose that for each $\mathcal{U}_0 \subset \mathcal{U}$, $\mathcal{U}_0$ is finite implies that $\mathcal{U}_0 \notin \text{Cover}_X(A)$. Suppose that for each $V \in \mathcal{V}$, there exists $\mathcal{U}_V \subset \mathcal{U}$ such that $\mathcal{U}_V$ is finite and $\mathcal{U}_V \in \text{Cover}_X(V)$. Define $\mathcal{U}_0 \subset \mathcal{U}$ by $\mathcal{U}_0 := \bigcup_{V \in \mathcal{V}} \mathcal{U}_V$. Since $\mathcal{V}$ is finite and for each $V \in \mathcal{V}$, $\mathcal{U}_V$ is finite, we have that $\mathcal{U}_0$ is finite. By construction,

$$\begin{aligned} A &\subset \bigcup_{V \in \mathcal{V}} V \quad (\text{since } \mathcal{V} \in \text{Cover}_X(A)) \\ &\subset \bigcup_{V \in \mathcal{V}} \left[\bigcup_{U \in \mathcal{U}_V} U \right] \\ &= \bigcup_{U \in \mathcal{U}_0} U. \end{aligned}$$

Hence $\mathcal{U}_0 \in \text{Cover}_X(A)$.

- Therefore, $\left(\text{for each } V \in \mathcal{V}, \text{ there exists } \mathcal{U}_V \subset \mathcal{U} \text{ such that } \mathcal{U}_V \text{ is finite and } \mathcal{U}_V \in \text{Cover}_X(V) \right)$ implies that $\left(\text{there exists } \mathcal{U}_0 \subset \mathcal{U} \text{ such that } \mathcal{U}_0 \text{ is finite and } \mathcal{U}_0 \in \text{Cover}_X(A) \right)$.
- By contrapositive, $\left(\text{for each } \mathcal{U}_0 \subset \mathcal{U}, \mathcal{U}_0 \text{ is finite implies that } \mathcal{U}_0 \notin \text{Cover}_X(A) \right)$ implies that $\left(\text{there exists } V \in \mathcal{V} \text{ such that for each } \mathcal{U}_0 \subset \mathcal{U}, \mathcal{U}_0 \text{ is finite implies that } \mathcal{U}_0 \notin \text{Cover}_X(V) \right)$.

□

1.3 Functions

1.3.1 Introduction

Exercise 1.3.1.1. Let X, Y be sets, $f : X \rightarrow Y$ and $A \subset X$. Then $A \subset f^{-1}(f(A))$.

Proof. Let $x \in A$. Then $f(x) \in f(A)$. Set $B := f(A)$. Since $f(x) \in B$, we have that

$$\begin{aligned} x &\in f^{-1}(B) \\ &= f^{-1}(f(A)) \end{aligned}$$

Since $x \in A$ is arbitrary, we have that $A \subset f^{-1}(f(A))$.

□

Exercise 1.3.1.2. Let X, Y be sets, $f : X \rightarrow Y$ and $B \subset Y$. Then $f(f^{-1}(B)) = B \cap f(X)$.

Proof. Let $y \in f(f^{-1}(B))$. Then there exists $x \in f^{-1}(B)$ such that $f(x) = y$. Thus

$$\begin{aligned} y &= f(x) \\ &\in B \end{aligned}$$

Since

$$\begin{aligned} f^{-1}(B) &= \{x \in X : f(x) \in B\} \\ &\subset X \end{aligned}$$

$$\begin{aligned} y &= f(x) \\ &\in f(X) \end{aligned}$$

Hence $y \in B \cap f(X)$. Since $y \in f(f^{-1}(B))$ is arbitrary, $f(f^{-1}(B)) \subset B \cap f(X)$.

Conversely, let $y \in B \cap f(X)$. Since $y \in f(X)$, there exists $x \in X$ such that $f(x) = y$. Since $y \in B$, $x \in f^{-1}(B)$. Hence

$$\begin{aligned} y &= f(x) \\ &\in f(f^{-1}(B)) \end{aligned}$$

Since $y \in B \cap f(X)$ is arbitrary, $B \cap f(X) \subset f(f^{-1}(B))$. Thus $f(f^{-1}(B)) = B \cap f(X)$. □

Exercise 1.3.1.3. Let X, Y be sets, $f : X \rightarrow Y$ and $\mathcal{A} \subset \mathcal{P}(X)$. Suppose that $\mathcal{A} \neq \emptyset$. Then

$$f\left(\bigcup_{A \in \mathcal{A}} A\right) = \bigcup_{A \in \mathcal{A}} f(A)$$

Proof.

- Let $y \in f\left(\bigcup_{A \in \mathcal{A}} A\right)$. Then there exists $x \in \bigcup_{A \in \mathcal{A}} A$ such that $y = f(x)$. Then there exists $A_0 \in \mathcal{A}$ such that $x \in A_0$. Therefore

$$\begin{aligned} y &= f(x) \\ &\in f(A_0) \\ &\subset \bigcup_{A \in \mathcal{A}} f(A). \end{aligned}$$

Since $y \in f\left(\bigcup_{A \in \mathcal{A}} A\right)$ is arbitrary, we have that for each $y \in f\left(\bigcup_{A \in \mathcal{A}} A\right)$, $y \in \bigcup_{A \in \mathcal{A}} f(A)$. Hence $f\left(\bigcup_{A \in \mathcal{A}} A\right) \subset \bigcup_{A \in \mathcal{A}} f(A)$.

- Let $y \in \bigcup_{A \in \mathcal{A}} f(A)$. Then there exists $A_0 \in \mathcal{A}$ such that $y \in f(A_0)$. Therefore, there exists $x \in A_0$ such that $y = f(x)$. Since

$$\begin{aligned} x &\in A_0 \\ &\subset \bigcup_{A \in \mathcal{A}} A, \end{aligned}$$

we have that

$$\begin{aligned} y &= f(x) \\ &\in f\left(\bigcup_{A \in \mathcal{A}} A\right). \end{aligned}$$

Since $y \in \bigcup_{A \in \mathcal{A}} f(A)$ is arbitrary, we have that for each $y \in \bigcup_{A \in \mathcal{A}} f(A)$, $y \in f\left(\bigcup_{A \in \mathcal{A}} A\right)$. Hence $f\left(\bigcup_{A \in \mathcal{A}} A\right) \supset \bigcup_{A \in \mathcal{A}} f(A)$. □

Exercise 1.3.1.4. Let X, Y be sets, $f : X \rightarrow Y$ and $\mathcal{A} \subset \mathcal{P}(X)$. Then

$$f\left(\bigcap_{A \in \mathcal{A}} A\right) \subset \bigcap_{A \in \mathcal{A}} f(A)$$

Proof. Let $y \in f\left(\bigcap_{A \in \mathcal{A}} A\right)$. Then there exists $x \in \bigcap_{A \in \mathcal{A}} A$ such that $y = f(x)$. Thus, for each $A \in \mathcal{A}$, $x \in A$. Hence for each $A \in \mathcal{A}$,

$$\begin{aligned} y &= f(x) \\ &\in f(A). \end{aligned}$$

Therefore $y \in \bigcap_{A \in \mathcal{A}} f(A)$. Since $y \in f\left(\bigcap_{A \in \mathcal{A}} A\right)$ is arbitrary, we have that for each $y \in f\left(\bigcap_{A \in \mathcal{A}} A\right)$, $y \in \bigcap_{A \in \mathcal{A}} f(A)$. Thus $f\left(\bigcap_{A \in \mathcal{A}} A\right) \subset \bigcap_{A \in \mathcal{A}} f(A)$. □

Exercise 1.3.1.5. Let X, Y be sets, $f : X \rightarrow Y$, $A \subset X$ and $B \subset Y$. Then

$$f(A \cap f^{-1}(B)) = f(A) \cap B$$

Proof. Let $y \in f(A \cap f^{-1}(B))$. Then there exists $a \in A \cap f^{-1}(B)$ such that $f(a) = y$. Then $a \in A$ and $f(a) \in B$. Hence

$$\begin{aligned} y &= f(a) \\ &\in f(A) \cap B. \end{aligned}$$

Since $y \in f(A \cap f^{-1}(B))$ is arbitrary, we have that $f(A \cap f^{-1}(B)) \subset f(A) \cap B$.

Conversely, let $y \in f(A) \cap B$. Then $y \in B$ and there exists $a \in A$ such that $f(a) = y$. Since $f(a) \in B$, we have that $a \in f^{-1}(B)$. Therefore

$$\begin{aligned} y &= f(a) \\ &\in f(A \cap f^{-1}(B)). \end{aligned}$$

Since $y \in f(A \cap f^{-1}(B))$ is arbitrary, we have that $f(A \cap f^{-1}(B)) \supset f(A) \cap B$.

Hence $f(A \cap f^{-1}(B)) = f(A) \cap B$. □

Exercise 1.3.1.6. Let X be a set, $K \subset X$, $(X_\alpha)_{\alpha \in A}$ a collection of sets and for each $\alpha \in A$, $f_\alpha : X \rightarrow X_\alpha$. For $\alpha \in A$, set $K_\alpha := f_\alpha(K)$. Then $K = \bigcap_{\alpha \in A} f_\alpha^{-1}(K_\alpha)$.

Proof.

- Let $x \in K$ and $\alpha \in A$. By definition,

$$\begin{aligned} f_\alpha(x) &\in f_\alpha(K) \\ &= K_\alpha \end{aligned}$$

Thus $x \in f_\alpha^{-1}(K_\alpha)$. Since $\alpha \in A$ is arbitrary, we have that for each $\alpha \in A$, $x \in f_\alpha^{-1}(K_\alpha)$. Thus $x \in \bigcap_{\alpha \in A} f_\alpha^{-1}(K_\alpha)$. Since $x \in K$ is arbitrary, $K \subset \bigcap_{\alpha \in A} f_\alpha^{-1}(K_\alpha)$.

- Let $\alpha \in A$. Exercise 1.3.1.1 implies that

$$\begin{aligned} K &\subset f_\alpha^{-1}(f_\alpha(K)) \\ &= f_\alpha^{-1}(K_\alpha). \end{aligned}$$

Since $\alpha \in A$ is arbitrary, for each $\alpha \in A$, $K \subset f_\alpha^{-1}(K_\alpha)$. Thus $K \subset \bigcap_{\alpha \in A} f_\alpha^{-1}(K_\alpha)$.

Therefore $K = \bigcap_{\alpha \in A} f_\alpha^{-1}(K_\alpha)$. □

1.3.2 Bijections

Definition 1.3.2.1. Let X, Y be sets and $f : X \rightarrow Y$. Then f is said to be a **surjection** if for each $y \in Y$, there exists $x \in X$ such that $f(x) = y$.

Exercise 1.3.2.2. Let X, Y be sets and $f : X \rightarrow Y$. Suppose that f is a surjection. Then for each $B \subset Y$, $f(f^{-1}(B)) = B$.

Proof. Let $B \subset Y$. Since f is surjective, $f(X) = Y$. A previous exercise implies that

$$\begin{aligned} f(f^{-1}(B)) &= B \cap f(X) \\ &= B \cap Y \\ &= B \end{aligned}$$

□

Exercise 1.3.2.3. Let X, Y, Z be sets, $f : X \rightarrow Y$ and $g, h : Y \rightarrow Z$. Suppose that f is a surjection. If $g \circ f = h \circ f$, then $g = h$.

Proof. Suppose that $g \circ f = h \circ f$. Let $y \in Y$. Since f is a surjection, there exists $x \in X$ such that $y = f(x)$. Then

$$\begin{aligned} g(y) &= g \circ f(x) \\ &= h \circ f(x) \\ &= h(y). \end{aligned}$$

Since $y \in Y$ is arbitrary, we have that for each $y \in Y$, $g(y) = h(y)$. Hence $g = h$. $\square$

Definition 1.3.2.4. Let X, Y be sets and $f : X \rightarrow Y$. Then f is said to be a **injection** if for each $a, b \in X$, $f(a) = f(b)$ implies that $a = b$.

Exercise 1.3.2.5. Let X, Y be sets and $f : X \rightarrow Y$. Suppose that f is an injection. Then for each $A \subset X$, $f^{-1}(f(A)) = A$.

Proof.

- Let $x \in f^{-1}(f(A))$. Then $f(x) \in f(A)$. Thus there exists $a \in A$ such that $f(x) = f(a)$. Since f is injective, $x = a$. Thus $x \in A$. Since $x \in f^{-1}(f(A))$ is arbitrary, we have that for each $x \in f^{-1}(f(A))$, $x \in A$. Hence $f^{-1}(f(A)) \subset A$.
- Exercise 1.3.1.1 implies that $A \subset f^{-1}(f(A))$.

Since $f^{-1}(f(A)) \subset A$ and $A \subset f^{-1}(f(A))$, we have that $f^{-1}(f(A)) = A$. $\square$

Exercise 1.3.2.6. Let X, Y be sets, $f : X \rightarrow Y$ and $\mathcal{A} \subset \mathcal{P}(X)$. Suppose that $\mathcal{A} \neq \emptyset$ and f is injective. Then

$$f\left(\bigcap_{A \in \mathcal{A}} A\right) = \bigcap_{A \in \mathcal{A}} f(A)$$

Proof. • Exercise 1.3.1.4 implies that $f\left(\bigcap_{A \in \mathcal{A}} A\right) \subset \bigcap_{A \in \mathcal{A}} f(A)$.

- Let $y \in \bigcap_{A \in \mathcal{A}} f(A)$. Then for each $A \in \mathcal{A}$, $y \in f(A)$. Thus, for each $A \in \mathcal{A}$, there exists $x_A \in A$ such that $y = f(x_A)$. Since $\mathcal{A} \neq \emptyset$, there exists $A_0 \in \mathcal{A}$. Let $A \in \mathcal{A}$. Then

$$\begin{aligned} f(x_A) &= y \\ &= f(x_{A_0}). \end{aligned}$$

Since f is injective, $x_A = x_{A_0}$. Since $A \in \mathcal{A}$ is arbitrary, we have that for each $A \in \mathcal{A}$, $x_A = x_{A_0}$. Thus for each $A \in \mathcal{A}$, $x_{A_0} \in A$. Hence $x_{A_0} \in \bigcap_{A \in \mathcal{A}} A$ and therefore

$$\begin{aligned} y &= f(x_{A_0}) \\ &\in f\left(\bigcap_{A \in \mathcal{A}} A\right). \end{aligned}$$

Since $y \in \bigcap_{A \in \mathcal{A}} f(A)$ is arbitrary, we have that for each $y \in \bigcap_{A \in \mathcal{A}} f(A)$, $y \in f\left(\bigcap_{A \in \mathcal{A}} A\right)$. Thus $\bigcap_{A \in \mathcal{A}} f(A) \subset f\left(\bigcap_{A \in \mathcal{A}} A\right)$.

Since $f\left(\bigcap_{A \in \mathcal{A}} A\right) \subset \bigcap_{A \in \mathcal{A}} f(A)$ and $\bigcap_{A \in \mathcal{A}} f(A) \subset f\left(\bigcap_{A \in \mathcal{A}} A\right)$, we have that $f\left(\bigcap_{A \in \mathcal{A}} A\right) = \bigcap_{A \in \mathcal{A}} f(A)$. $\square$

1.3.3 Nets

Definition 1.3.3.1. Let X be a set, A a directed set and $x : A \rightarrow Y$. Then x is said to be a **net** in X . We typically write $(x_\alpha)_{\alpha \in A}$.

Definition 1.3.3.2. Let X be a set, A a directed set and $\alpha_0 \in A$. We define the α_0 -**tail operator**, denoted $L_{\alpha_0} : X^A \rightarrow X^{[\alpha_0, \infty)}$, by $L(x) := x|_{[\alpha_0, \infty)}$, i.e. $L_{\alpha_0}((x_\alpha)_{\alpha \in A}) := (x_\alpha)_{\alpha \in [\alpha_0, \infty)}$.

Definition 1.3.3.3. Let X be a set, $(x_\alpha)_{\alpha \in A}$, $(y_\beta)_{\beta \in B} \subset X$ nets and $\phi : B \rightarrow A$. Then $((y_\beta)_{\beta \in B}, \phi)$ is said to be a **subnet of** $(x_\alpha)_{\alpha \in A}$ if

1. for each $\beta \in B$, $y_\beta = x_{\phi(\beta)}$
2. for each $\alpha_0 \in A$, there exists $\beta_0 \in B$ such that for each $\beta \in B$, $\beta \geq \beta_0$ implies that $\phi(\beta) \geq \alpha_0$

Note 1.3.3.4. We usually suppress ϕ and write α_β in place of $\phi(\beta)$.

1.3.4 Sequences

1.4 Subsets

Definition 1.4.0.1. Let X be a set and $A \subset X$. We define the **inclusion map from A to X** , denoted $\iota_A^X \in X^A$, by

$$\iota_A^X(a) := a, \quad a \in A.$$

Note 1.4.0.2. When the context is clear, we may write ι_A in place of ι_A^X .

Definition 1.4.0.3. Let X, Y be sets, $A \subset X$, $B \subset Y$, $f \in Y^X$ and $g \in B^X$. We define

- the **resitricition of f to A** , denoted $f|_A \in Y^A$, by $f|_A := f \circ \iota_A^X$.
- **FINISH!!!**

Definition 1.4.0.4. Let A, B, Y be sets, $f : A \rightarrow Y$ and $g : B \rightarrow Y$.

- Then
 - g is said to be **an extension of f** if $A \subset B$ and $g|_A = f$,
 - f is said to be **a restriction of g** if $A \subset B$ and $g|_A = f$.
- We define
 - $\text{Ext}(f) := \{g' \in Y^{B'} : B' \text{ is a set and } g' \text{ is an extension of } f\}$,
 - $\text{Rstr}(g) := \{f' \in Y^{A'} : A' \text{ is a set and } f' \text{ is a restriction of } g\}$.

Exercise 1.4.0.5. Let A, B, Y be sets, $f \in Y^A$ and $g \in Y^B$. Then $g \in \text{Ext}(f)$ iff $\Gamma(f) \subset \Gamma(g)$. **FINISH!!! (define $\Gamma(f)$)**

Proof.

- ($\implies$):
Suppose that $g \in \text{Ext}(f)$. Then $A \subset B$ and $g|_A = f$. Let $(x, y) \in \Gamma(f)$. Then

$$\begin{aligned} x &\in A \\ &\subset B. \end{aligned}$$

and

$$\begin{aligned} g(x) &= g(\iota_A^B(x)) \\ &= g \circ \iota_A^B(x) \\ &= f(x) \\ &= y \end{aligned}$$

Thus $(x, y) \in \Gamma(g)$. Since $(x, y) \in \Gamma(f)$ is arbitrary, we have that for each $(x, y) \in \Gamma(f)$, $(x, y) \in \Gamma(g)$. Hence $\Gamma(f) \subset \Gamma(g)$.

- ($\impliedby$):
Suppose that $\Gamma(f) \subset \Gamma(g)$. Let $x \in A$. Then

$$\begin{aligned} (x, T(x)) &\in \Gamma(f) \\ &\subset \Gamma(g). \end{aligned}$$

Hence $x \in B$ and $g(x) = T(x)$. Since $x \in A$ is arbitrary, we have that for each $x \in A$, $x \in B$ and

$$\begin{aligned} g \circ \iota_A^B(x) &= g(\iota_A^B(x)) \\ &= g(x) \\ &= T(x) \end{aligned}$$

Hence $A \subset B$ and

$$\begin{aligned} g|_A &= g \circ \iota_A^B \\ &= T. \end{aligned}$$

Thus $g \in \text{Ext}(f)$.

□

1.5 Products of Sets

Definition 1.5.0.1. Let A be a set and $(X_\alpha)_{\alpha \in A}$ a collection of sets. We define

- the **Cartesian product** of $(X_\alpha)_{\alpha \in A}$, denoted $\prod_{\alpha \in A} X_\alpha$, by

$$\prod_{\alpha \in A} X_\alpha := \{f : A \rightarrow \bigcup_{\alpha \in A} X_\alpha : \text{for each } \alpha \in A, f(\alpha) \in X_\alpha\}$$

- the α -th **projection map** of $\prod_{\alpha \in A} X_\alpha$ **onto** X_α , denoted $\pi_\alpha : \prod_{\alpha \in A} X_\alpha \rightarrow X_\alpha$, by

$$\pi_\alpha(f) := f(\alpha)$$

Exercise 1.5.0.2. Let A be a set and $(X_\alpha)_{\alpha \in A}, (Y_\alpha)_{\alpha \in A}$ collections of sets. If for each $\alpha \in A$, $X_\alpha \subset Y_\alpha$, then $\prod_{\alpha \in A} X_\alpha \subset \prod_{\alpha \in A} Y_\alpha$. **make this iff?**

Proof. **FINISH and cite in following exercises** □

Exercise 1.5.0.3. Let A be a set and $(X_\alpha)_{\alpha \in A}$ a collection of sets.

1. If $A = \emptyset$, then $\prod_{\alpha \in A} X_\alpha = \{\emptyset\}$.

- 2.

Proof.

1. Suppose that $A = \emptyset$. Then **ref ex here** implies that $\bigcup_{\alpha \in A} X_\alpha = \emptyset$. Since **ref ex here** implies that $\emptyset^\emptyset = \{\emptyset\}$, we have that

$$\begin{aligned} \prod_{\alpha \in A} X_\alpha &\subset \left(\bigcup_{\alpha \in A} X_\alpha \right)^A \\ &= \emptyset^\emptyset \\ &= \{\emptyset\}. \end{aligned}$$

Define $f_0 \in \left(\bigcup_{\alpha \in A} X_\alpha \right)^A$ by $f_0 := \emptyset$. Since it is vacuously true that for each $\alpha \in \emptyset$, $f_0(\alpha) \in X_\alpha$, we have that $f_0 \in \prod_{\alpha \in A} X_\alpha$. Hence

$$\begin{aligned} \{\emptyset\} &= \{f_0\} \\ &\in \left\{ f \in \left(\bigcup_{\alpha \in A} X_\alpha \right)^A : \text{for each } \alpha \in A, f(\alpha) \in X_\alpha \right\} \\ &= \prod_{\alpha \in A} X_\alpha. \end{aligned}$$

Since $\prod_{\alpha \in A} X_\alpha \subset \{\emptyset\}$ and $\{\emptyset\} \subset \prod_{\alpha \in A} X_\alpha$, we have that $\prod_{\alpha \in A} X_\alpha = \{\emptyset\}$.

- 2.

□

Exercise 1.5.0.4. Let A be a set and $(X_\alpha)_{\alpha \in A}$ a collection of sets. If $A \neq \emptyset$, then

1. $\prod_{\alpha \in A} X_\alpha = \emptyset$ iff there exists $\alpha \in A$ such that $X_\alpha = \emptyset$,

2. $\# \prod_{\alpha \in A} X_\alpha = 1$ iff for each $\alpha \in A$, $\#X_\alpha = 1$.

Proof. Suppose that $A \neq \emptyset$. Set $X := \prod_{\alpha \in A} X_\alpha$.

1. • ($\implies$):
Suppose that for each $\alpha \in A$, $X_\alpha \neq \emptyset$. Since $A \neq \emptyset$, the axiom of choice implies that $X \neq \emptyset$.
- ($\impliedby$):
Suppose that there exists $\alpha_0 \in A$ such that $X_{\alpha_0} = \emptyset$. For the sake of contradiction, suppose that $X \neq \emptyset$. Then there exists $f \in X$. Therefore

$$\begin{aligned} f(\alpha_0) &\in X_{\alpha_0} \\ &= \emptyset \end{aligned}$$

which is a contradiction. Hence $X = \emptyset$.

2. • ($\implies$):
For the sake of contradiction, suppose that $\#X = 1$ and there exists $\alpha_0 \in A$ such that $\#X_{\alpha_0} \neq 1$. Since $\#X = 1$, we have that $X \neq \emptyset$. Since $A \neq \emptyset$, **the prev part** implies that for each $\alpha \in A$, $X_\alpha \neq \emptyset$. In particular, $\#X_{\alpha_0} \neq 0$. Since $\#X_{\alpha_0} \notin \{0, 1\}$, we have that $\#X_{\alpha_0} \geq 2$. Therefore, there exists $y, z \in X_{\alpha_0}$ such that $y \neq z$. Define $(Y_\alpha)_{\alpha \in A}$ and $(Z_\alpha)_{\alpha \in A}$ by

$$Y_\alpha := \begin{cases} \{y\}, & \alpha = \alpha_0 \\ X_\alpha, & \alpha \neq \alpha_0 \end{cases}$$

and

$$Z_\alpha := \begin{cases} \{z\}, & \alpha = \alpha_0 \\ X_\alpha, & \alpha \neq \alpha_0 \end{cases}$$

Then for each $\alpha \in A$, $Y_\alpha, Z_\alpha \subset X_\alpha$ and $Y_\alpha, Z_\alpha \neq \emptyset$. **make prev ex** implies that $Y, Z \subset X$. Since $A \neq \emptyset$, the axiom of choice implies that $Y, Z \neq \emptyset$. Thus there exists $f_Y \in Y$ and $f_Z \in Z$. Since $Y, Z \subset X$, $\{f_Y, f_Z\} \subset X$. Since

$$\begin{aligned} f_Y(\alpha_0) &\in Y_{\alpha_0} \\ &= \{y\}, \end{aligned}$$

we have that $f_Y(\alpha_0) = y$. Similarly, $f_Z(\alpha_0) = z$. Therefore

$$\begin{aligned} f_Y(\alpha_0) &= y \\ &\neq z \\ &= f_Z(\alpha_0). \end{aligned}$$

Thus $f_Y \neq f_Z$. Hence $\#X \geq 2$. This is a contradiction since $\#X = 1$. Hence $\#X = 1$ implies that for each $\alpha \in A$, $\#X_\alpha = 1$.

- ($\impliedby$):
Suppose that for each $\alpha \in A$, $\#X_\alpha = 1$. **The prev part** implies that $X \neq \emptyset$. Thus there exists $f \in X$.
- $f \in X$, we have that $\{f\} \subset X$.
 - Let $f' \in X$ and $\alpha \in A$. Since $\#X_\alpha = 1$, there exists $x \in X_\alpha$ such that $X_\alpha = \{x\}$. Then

$$\begin{aligned} \{f(\alpha), f'(\alpha)\} &\subset X_\alpha \\ &= \{x\}. \end{aligned}$$

Thus

$$\begin{aligned} f'(\alpha) &= x \\ &= f(\alpha). \end{aligned}$$

Since $\alpha \in A$ is arbitrary, we have that for each $\alpha \in A$, $f'(\alpha) = f(\alpha)$. Hence $f' = f$. Since $f' \in X$ is arbitrary, we have that for each $f' \in X$, $f' \in \{f\}$. Thus $X \subset \{f\}$.

Since $X \subset \{f\}$ and $\{f\} \subset X$, we have that $X = \{f\}$. Therefore $\#X = 1$.

□

Exercise 1.5.0.5. Let Λ, B be sets and $(A_\lambda)_{\lambda \in \Lambda}$ a collection of sets. Then

$$\left(\bigcup_{\lambda \in \Lambda} A_\lambda \right) \times B = \bigcup_{\lambda \in \Lambda} (A_\lambda \times B)$$

Proof. Let $(x, y) \in \left(\bigcup_{\lambda \in \Lambda} A_\lambda \right) \times B$. Then $x \in \bigcup_{\lambda \in \Lambda} A_\lambda$ and $y \in B$. Therefore, there exists $\lambda \in \Lambda$ such that $x \in A_\lambda$. Hence

$$\begin{aligned} (x, y) &\in A_\lambda \times B \\ &\subset \bigcup_{\lambda \in \Lambda} (A_\lambda \times B) \end{aligned}$$

Thus $\left(\bigcup_{\lambda \in \Lambda} A_\lambda \right) \times B \subset \bigcup_{\lambda \in \Lambda} (A_\lambda \times B)$.

Conversely, let $(x, y) \in \bigcup_{\lambda \in \Lambda} (A_\lambda \times B)$. Then there exists $\lambda \in \Lambda$ such that $(x, y) \in A_\lambda \times B$. Then

$$\begin{aligned} x &\in A_\lambda \\ &\subset \bigcup_{\lambda \in \Lambda} A_\lambda \end{aligned}$$

and $y \in B$. Hence $(x, y) \in \left(\bigcup_{\lambda \in \Lambda} A_\lambda \right) \times B$. So $\bigcup_{\lambda \in \Lambda} (A_\lambda \times B) \subset \left(\bigcup_{\lambda \in \Lambda} A_\lambda \right) \times B$. □

Definition 1.5.0.6. Let $(X_\alpha)_{\alpha \in A}$ and $(Y_\alpha)_{\alpha \in A}$ be collections of sets and $(f_\alpha)_{\alpha \in A} \in \prod_{\alpha \in A} Y_\alpha^{X_\alpha}$, i.e. for each $\alpha \in A$, $f_\alpha : X_\alpha \rightarrow Y_\alpha$.

We define the **product of** $(f_\alpha)_{\alpha \in A}$, denoted $\prod_{\alpha \in A} f_\alpha : \prod_{\alpha \in A} X_\alpha \rightarrow \prod_{\alpha \in A} Y_\alpha$ by

$$\left(\left[\prod_{\alpha \in A} f_\alpha \right] (x) \right)_\beta = f_\beta(x_\beta)$$

Exercise 1.5.0.7. Let $(X_\alpha)_{\alpha \in A}$ and $(Y_\alpha)_{\alpha \in A}$ be collections of sets and $(f_\alpha)_{\alpha \in A} \in \prod_{\alpha \in A} Y_\alpha^{X_\alpha}$, i.e. for each $\alpha \in A$, $f_\alpha : X_\alpha \rightarrow Y_\alpha$.

Denote the α -th projection maps on $\prod_{\alpha \in A} X_\alpha$ and $\prod_{\alpha \in A} Y_\alpha$ by π_α^X and π_α^Y respectively. Then for each $\alpha \in A$, $\pi_\alpha^Y \circ \left[\prod_{\alpha \in A} f_\alpha \right] = f_\alpha \circ \pi_\alpha^X$, i.e. the following diagram commutes:

$$\begin{array}{ccc} \prod_{\alpha \in A} X_\alpha & \xrightarrow{\prod_{\alpha \in A} f_\alpha} & \prod_{\alpha \in A} Y_\alpha \\ \pi_\alpha^X \downarrow & & \downarrow \pi_\alpha^Y \\ X_\alpha & \xrightarrow{f_\alpha} & Y_\alpha \end{array}$$

Proof. Set $X := \prod_{\alpha \in A} X_\alpha$, $Y := \prod_{\alpha \in A} Y_\alpha$ and define $f : X \rightarrow Y$ by $f := \prod_{\alpha \in A} f_\alpha$. Let $\alpha \in A$ and $x \in X$. Then

$$\begin{aligned} \pi_\alpha^Y \circ f(x) &= (f(x))_\alpha \\ &= f_\alpha(x_\alpha) \\ &= f_\alpha \circ \pi_\alpha^X(x) \end{aligned}$$

Since $\alpha \in A$ and $x \in X$ are arbitrary, for each $\alpha \in A$, $\pi_\alpha^Y \circ f = f_\alpha \circ \pi_\alpha^X$. □

Exercise 1.5.0.8. Let $(X_\alpha)_{\alpha \in A}$ and $(Y_\alpha)_{\alpha \in A}$ be collections of sets and $(f_\alpha)_{\alpha \in A} \in \prod_{\alpha \in A} Y_\alpha^{X_\alpha}$, i.e. for each $\alpha \in A$, $f_\alpha : X_\alpha \rightarrow Y_\alpha$.

Suppose that for each $\alpha \in A$, f_α is a bijection. Then $\prod_{\alpha \in A} f_\alpha$ is a bijection and $\left[\prod_{\alpha \in A} f_\alpha \right]^{-1} = \prod_{\alpha \in A} f_\alpha^{-1}$.

Proof. Set $X := \prod_{\alpha \in A} X_\alpha$, $Y := \prod_{\alpha \in A} Y_\alpha$ and define $f : X \rightarrow Y$ and $g : Y \rightarrow X$ by $f := \prod_{\alpha \in A} f_\alpha$ and $g := \prod_{\alpha \in A} f_\alpha^{-1}$. Denote the α -th projection maps on $\prod_{\alpha \in A} X_\alpha$ and $\prod_{\alpha \in A} Y_\alpha$ by π_α^X and π_α^Y respectively. Let $\alpha \in A$. Then

$$\begin{aligned} \pi_\alpha^X \circ g \circ f &= g_\alpha \circ \pi_\alpha^Y \circ f \\ &= g_\alpha \circ f_\alpha \circ \pi_\alpha^X \\ &= f_\alpha^{-1} \circ f_\alpha \circ \pi_\alpha^X \\ &= \pi_\alpha^X \end{aligned}$$

and

$$\begin{aligned} \pi_\alpha^Y \circ f \circ g &= f_\alpha \circ \pi_\alpha^X \circ g \\ &= f_\alpha \circ g_\alpha \circ \pi_\alpha^Y \\ &= f_\alpha \circ f_\alpha^{-1} \circ \pi_\alpha^Y \\ &= \pi_\alpha^Y \end{aligned}$$

Since $\alpha \in A$ is arbitrary, we have that for each $\alpha \in A$, $\pi_\alpha^X \circ g \circ f = \pi_\alpha^X$ and $\pi_\alpha^Y \circ f \circ g = \pi_\alpha^Y$. Hence $g \circ f = \text{id}_X$ and $f \circ g = \text{id}_Y$. Thus f is a bijection and $f^{-1} = g$. $\square$

Definition 1.5.0.9. Let X be a set. We define

- the **diagonal of** $X \times X$, denoted $\Delta_X \subset X \times X$, by $\Delta_X := \{(x, x) : x \in X\}$,
- the **diagonal of** $X^\mathbb{N}$, denoted $\Delta_{X^\mathbb{N}} \subset X^\mathbb{N}$, by $\Delta_{X^\mathbb{N}} := \{x \in X^\mathbb{N} : \text{for each } m, n \in \mathbb{N}, \pi_m(x) = \pi_n(x)\}$.

Exercise 1.5.0.10. Let X be a set, $(B_n)_{n \in \mathbb{N}} \subset \mathcal{P}(X)$, $(Z_n)_{n \in \mathbb{N}}$ a collection of sets and $(f_n) \in \prod_{n \in \mathbb{N}} B_n^{Z_n}$. Suppose that for each $n \in \mathbb{N}$, f_n is a bijection. Set $Z_0 := \prod_{n \in \mathbb{N}} Z_n$. Define $f_0 : Z_0 \rightarrow X^\mathbb{N}$ and $Z \subset Z_0$ by $f_0 := \prod_{n \in \mathbb{N}} f_n$ and $Z := f_0^{-1}(\Delta_{X^\mathbb{N}})$. Then

1. for each $m, n \in \mathbb{N}$, $f_m \circ \pi_m|_Z = f_n \circ \pi_n|_Z$,
2. for each $n \in \mathbb{N}$, $f_n \circ \pi_n(Z) = \bigcap_{n \in \mathbb{N}} B_n$.

Proof.

1. Let $m, n \in \mathbb{N}$ and $z \in Z$. By construction, $f_0(z) \in \Delta_{X^\mathbb{N}}$. Therefore

$$\begin{aligned} f_n(\pi_n(z)) &= \pi_n(f_0(z)) \\ &= \pi_m(f_0(z)) \\ &= f_m(\pi_m(z)). \end{aligned}$$

Since $z \in Z$ is arbitrary, we have that $f_m \circ \pi_m|_Z = f_n \circ \pi_n|_Z$. Since $m, n \in \mathbb{N}$ are arbitrary, we have that for each $m, n \in \mathbb{N}$, $f_m \circ \pi_m|_Z = f_n \circ \pi_n|_Z$.

2. Set $B := \bigcap_{n \in \mathbb{N}} B_n$. Let $n \in \mathbb{N}$.

- Let $z \in Z$. The previous part implies that

$$\begin{aligned} f_n(\pi_n(z)) &= f_m(\pi_m(z)) \\ &\in B_m. \end{aligned}$$

Since $m \in \mathbb{N}$ is arbitrary, we have that for each $m \in \mathbb{N}$, $f_n(\pi_n(z)) \in B_m$. Hence

$$\begin{aligned} f_n(\pi_n(z)) &\in \bigcap_{m \in \mathbb{N}} B_m \\ &= B. \end{aligned}$$

Since $z \in Z$ is arbitrary, we have that $f_n \circ \pi_n(Z) \subset B$.

- Let $x \in B$. Then for each $m \in \mathbb{N}$, $x \in B_m$. Define $z \in Z_0$ by $z_m := f_m^{-1}(x)$. Then for each $m \in \mathbb{N}$,

$$\begin{aligned} \pi_m \circ f_0(z) &= f_m \circ \pi_m(z) \\ &= f_m(z_m) \\ &= f_m(f_m^{-1}(x)) \\ &= x \end{aligned}$$

In particular, for each $m \in \mathbb{N}$,

$$\begin{aligned} \pi_m \circ f_0(z) &= x \\ &= \pi_n \circ f_0(z). \end{aligned}$$

Thus $f_0(z) \in \Delta_{X^{\mathbb{N}}}$ and $z \in Z$. Therefore

$$\begin{aligned} x &= \pi_n \circ f_0(z) \\ &= f_n \circ \pi_n(z) \\ &\in f_n \circ \pi_n(Z). \end{aligned}$$

Since $x \in B$ is arbitrary, we have that $B \subset f_n \circ \pi_n(Z)$. □

Definition 1.5.0.11. Let A, X be sets, $(Y_\alpha)_{\alpha \in A}$ a collection of sets and $(f_\alpha)_{\alpha \in A} \in \prod_{\alpha \in A} Y_\alpha^X$, i.e. for each $\alpha \in A$, $f_\alpha : X \rightarrow Y_\alpha$. Set $Y = \prod_{\alpha \in A} Y_\alpha$. We define the **tuple of** $(f_\alpha)_{\alpha \in A}$, denoted $(f_\alpha)_{\alpha \in A} : X \rightarrow Y$, by $(f_\alpha)_{\alpha \in A}(x) := (f_\alpha(x))_{\alpha \in A}$.

Exercise 1.5.0.12. Let X be a set, $(Y_\alpha)_{\alpha \in A}$ a collection of sets, $(f_\alpha)_{\alpha \in A} \in \prod_{\alpha \in A} Y_\alpha^X$ and $E \subset X$. Then $(f_\alpha)_{\alpha \in A}|_E = (f_\alpha|_E)_{\alpha \in A}$.

Proof. Clear. **add details.** □

Definition 1.5.0.13. **Define slice maps here in terms of inclusion maps** Let X, Y be sets and $U \subset X \times Y$. For each $(x_0, y_0) \in U$, we define $U_{x_0} = \{y \in Y : (x_0, y) \in U\}$ and $U^{y_0} = \{x \in X : (x, y_0) \in U\}$.

Definition 1.5.0.14. Let X, Y and Z be sets, $U \subset X \times Y$ and $f : U \rightarrow Z$. For each $(x_0, y_0) \in U$, we define $f_{x_0} : U_{x_0} \rightarrow Z$ and $f^{y_0} : U^{y_0} \rightarrow Z$ by $f_{x_0} = f(x_0, \cdot)$ and $f^{y_0} = f(\cdot, y_0)$.

Exercise 1.5.0.15. Let X, Y and Z be sets, $U \subset X \times Y$, $f : U \rightarrow Z$ and $(x_0, y_0) \in U$. Then for each $V \subset Z$, $(f_{x_0})^{-1}(V) = (f^{-1}(V))_{x_0}$ and $(f^{y_0})^{-1}(V) = (f^{-1}(V))^{y_0}$.

Proof. Let $V \subset Z$. Then for each $x \in U^{y_0}$,

$$\begin{aligned} x \in (f^{y_0})^{-1}(V) &\iff f^{y_0}(x) \in V \\ &\iff f(x, y_0) \in V \\ &\iff (x, y_0) \in f^{-1}(V) \\ &\iff x \in (f^{-1}(V))^{y_0} \end{aligned}$$

So $(f^{y_0})^{-1}(V) = (f^{-1}(V))^{y_0}$. Similarly, $(f_{x_0})^{-1}(V) = (f^{-1}(V))_{x_0}$. □

Definition 1.5.0.16. Let X, Y, Z be sets. We define the **currying operator**, denoted $\text{cur} : Z^{X \times Y} \rightarrow (Z^Y)^X$, by $\text{cur}(f)(x)(y) = f(x, y)$.

Definition 1.5.0.17. Separating Points:

Let X be a set, $(Y_\alpha)_{\alpha \in A}$ a collection of sets and $\mathcal{F} \in \prod_{\alpha \in A} Y_\alpha^X$. Write $\mathcal{F} = (f_\alpha)_{\alpha \in A}$. Then $\mathcal{F}$ is said to **separate the points of** X if for each $x_1, x_2 \in X$, $x_1 \neq x_2$ implies that there exists $\alpha \in A$ such that $f_\alpha(x_1) \neq f_\alpha(x_2)$.

1.6 Coproducts of Sets

Definition 1.6.0.1. Let $(X_\alpha)_{\alpha \in A}$ be a collection of sets. We define the **disjoint union** of $(X_\alpha)_{\alpha \in A}$, denoted $\coprod_{\alpha \in A} X_\alpha$, by

$$\coprod_{\alpha \in A} X_\alpha := \{(\alpha, x) : \alpha \in A \text{ and } x \in X_\alpha\}$$

Definition 1.6.0.2. Let $(X_\alpha)_{\alpha \in A}$ be a collection of sets. For $\alpha \in A$, we define the **α -th embedding map** of X_α into $\coprod_{\alpha \in A} X_\alpha$, denoted $\iota_\alpha : X_\alpha \rightarrow \coprod_{\alpha \in A} X_\alpha$, by

$$\iota_\alpha(x) = (\alpha, x)$$

Exercise 1.6.0.3. Let A be a set and $(X_\alpha)_{\alpha \in A}$ be a collection of sets and $(E_\alpha)_{\alpha \in A} \in \prod_{\alpha \in A} \mathcal{P}(X_\alpha)$. Then

$$\bigcup_{\alpha \in A} \iota_\alpha(E_\alpha) = \coprod_{\alpha \in A} E_\alpha.$$

Proof. We have that for each $x \in \coprod_{\alpha \in A} X_\alpha$,

$$\begin{aligned} x \in \bigcup_{\alpha \in A} \iota_\alpha(E_\alpha) &\iff \text{there exists } \alpha \in A \text{ such that } x \in \iota_\alpha(E_\alpha) \\ &\iff \text{there exists } \alpha \in A \text{ such that } x \in \{\alpha\} \times E_\alpha \\ &\iff \text{there exists } \alpha \in A, y \in E_\alpha \text{ such that } x = (\alpha, y) \\ &\iff x \in \coprod_{\alpha \in A} E_\alpha. \end{aligned}$$

Thus $\bigcup_{\alpha \in A} \iota_\alpha(E_\alpha) = \coprod_{\alpha \in A} E_\alpha$. □

Exercise 1.6.0.4. Let A, B be sets, $(X_\alpha)_{\alpha \in A}$ be a collection of sets and for each $\alpha \in A$, $(E_{\alpha, \beta})_{\beta \in B} \in \prod_{\beta \in B} \mathcal{P}(X_\alpha)$. Then

$$\bigcup_{\beta \in B} \left(\coprod_{\alpha \in A} E_{\alpha, \beta} \right) = \coprod_{\alpha \in A} \left(\bigcup_{\beta \in B} E_{\alpha, \beta} \right).$$

Proof. We have that

$$\begin{aligned} x \in \bigcup_{\beta \in B} \left(\coprod_{\alpha \in A} E_{\alpha, \beta} \right) &\iff \text{there exists } \beta \in B \text{ such that } x \in \coprod_{\alpha \in A} E_{\alpha, \beta} \\ &\iff \text{there exists } \beta \in B, \alpha \in A \text{ and } y \in E_{\alpha, \beta} \text{ such that } x = (\alpha, y) \\ &\iff \text{there exists } \beta \in B, \alpha \in A \text{ such that } x \in \{\alpha\} \times E_{\alpha, \beta} \\ &\iff \text{there exists } \beta \in B, \alpha \in A \text{ such that } x \in \iota_\alpha(E_{\alpha, \beta}) \\ &\iff \text{there exists } \alpha \in A \text{ such that } x \in \bigcup_{\beta \in B} \iota_\alpha(E_{\alpha, \beta}) \\ &\iff \text{there exists } \alpha \in A \text{ such that } x \in \iota_\alpha \left(\bigcup_{\beta \in B} E_{\alpha, \beta} \right) \\ &\iff x \in \bigcup_{\alpha \in A} \iota_\alpha \left(\bigcup_{\beta \in B} E_{\alpha, \beta} \right) \\ &\iff x \in \coprod_{\alpha \in A} \left(\bigcup_{\beta \in B} E_{\alpha, \beta} \right) \quad (\text{by Exercise 1.6.0.3}). \end{aligned}$$

Thus

$$\bigcup_{\beta \in B} \left(\prod_{\alpha \in A} E_{\alpha, \beta} \right) = \prod_{\alpha \in A} \left(\bigcup_{\beta \in B} E_{\alpha, \beta} \right).$$

□

Definition 1.6.0.5. Let $(X_\alpha, \mathcal{T}_\alpha)_{\alpha \in A}$ and $(Y_\alpha, \mathcal{S}_\alpha)_{\alpha \in A}$ be collections of topological spaces and $(f_\alpha)_{\alpha \in A} \in \prod_{\alpha \in A} Y_\alpha^{X_\alpha}$, i.e. for each $\alpha \in A$, $f_\alpha : X_\alpha \rightarrow Y_\alpha$. Set $X = \prod_{\alpha \in A} X_\alpha$ and $Y = \prod_{\alpha \in A} Y_\alpha$. We define the **coproduct** of $(f_\alpha)_{\alpha \in A}$, denoted $\prod_{\alpha \in A} f_\alpha : X \rightarrow Y$, by $\prod_{\alpha \in A} f_\alpha(\beta, x) = (\beta, f_\beta(x))$.

Exercise 1.6.0.6. Let $(X_\alpha)_{\alpha \in A}$ and $(Y_\alpha)_{\alpha \in A}$ be collections of sets and $(f_\alpha)_{\alpha \in A} \in \prod_{\alpha \in A} Y_\alpha^{X_\alpha}$, i.e. for each $\alpha \in A$, $f_\alpha : X_\alpha \rightarrow Y_\alpha$. Set $f := \prod_{\alpha \in A} f_\alpha$. Then for each $\alpha \in A$, $f \circ \iota_\alpha^X = \iota_\alpha^Y \circ f_\alpha$, i.e. the following diagram commutes:

$$\begin{array}{ccc} \prod_{\alpha \in A} X_\alpha & \xrightarrow{f} & \prod_{\alpha \in A} Y_\alpha \\ \iota_\alpha^X \uparrow & & \uparrow \iota_\alpha^Y \\ X_\alpha & \xrightarrow{f_\alpha} & Y_\alpha \end{array}$$

Proof. Let $\alpha \in A$ and $x \in X_\alpha$. Then

$$\begin{aligned} f \circ \iota_\alpha^X(x) &= f(\alpha, x) \\ &= (\alpha, f_\alpha(x)) \\ &= \iota_\alpha^Y \circ f_\alpha(x). \end{aligned}$$

Since $x \in X$ are arbitrary, $f \circ \iota_\alpha^X = \iota_\alpha^Y \circ f_\alpha$. Since $\alpha \in A$ is arbitrary, we have that for each $\alpha \in A$, $f \circ \iota_\alpha^X = \iota_\alpha^Y \circ f_\alpha$. □

Exercise 1.6.0.7. Let $(X_\alpha)_{\alpha \in A}$ and $(Y_\alpha)_{\alpha \in A}$ be collections of sets and $(f_\alpha)_{\alpha \in A} \in \prod_{\alpha \in A} Y_\alpha^{X_\alpha}$, i.e. for each $\alpha \in A$, $f_\alpha : X_\alpha \rightarrow Y_\alpha$.

Suppose that for each $\alpha \in A$, f_α is a bijection. Then $\prod_{\alpha \in A} f_\alpha$ is a bijection and $\left[\prod_{\alpha \in A} f_\alpha \right]^{-1} = \prod_{\alpha \in A} f_\alpha^{-1}$.

Proof. Set $X := \prod_{\alpha \in A} X_\alpha$, $Y := \prod_{\alpha \in A} Y_\alpha$ and define $f : X \rightarrow Y$ and $g : Y \rightarrow X$ by $f := \prod_{\alpha \in A} f_\alpha$ and $g := \prod_{\alpha \in A} f_\alpha^{-1}$. Denote the α -th embedding maps into X and Y by ι_α^X and ι_α^Y respectively. Let $\alpha \in A$. Then

$$\begin{aligned} g \circ f \circ \iota_\alpha^X &= g \circ \iota_\alpha^Y \circ f_\alpha \\ &= \iota_\alpha^X \circ g_\alpha \circ f_\alpha \\ &= \iota_\alpha^X \circ f_\alpha^{-1} \circ f_\alpha \\ &= \iota_\alpha^X \end{aligned}$$

and

$$\begin{aligned} f \circ g \circ \iota_\alpha^Y &= f \circ \iota_\alpha^X \circ g_\alpha \\ &= \iota_\alpha^Y \circ f_\alpha \circ g_\alpha \\ &= \iota_\alpha^Y \circ f_\alpha \circ f_\alpha^{-1} \\ &= \iota_\alpha^Y \end{aligned}$$

Since $\alpha \in A$ is arbitrary, we have that for each $\alpha \in A$, $g \circ f \circ \iota_\alpha^X = \iota_\alpha^X$ and $f \circ g \circ \iota_\alpha^Y = \iota_\alpha^Y$. Hence $g \circ f = \text{id}_X$ and $f \circ g = \text{id}_Y$. Thus f is a bijection and $f^{-1} = g$. □

1.7 Quotients of Sets

Definition 1.7.0.1. Let X be a set and $\sim$ an equivalence relation on X . We define the **quotient set** of X by $\sim$, denoted $X/\sim$, by

$$X/\sim = \{\bar{x} : x \in X\}$$

Exercise 1.7.0.2.

1.8 Equalizers of Maps

Definition 1.8.0.1. Let X, Y be sets and $f, g : X \rightarrow Y$. We define the **equalizer of f and g** , denoted $\text{Eq}(f, g)$, by

$$\text{Eq}(f, g) := \{x \in X : f(x) = g(x)\}$$

Exercise 1.8.0.2.

1.9 Projective Limits of Sets

Note 1.9.0.1. Let $(X_j)_{j \in J}$ be a collection of sets. We denote the j -th projection map from $\prod_{j \in J} X_j$ onto X_j by $\text{proj}_j : \prod_{j \in J} X_j \rightarrow X_j$.

Definition 1.9.0.2. Let $(J, \leq)$ be a directed poset, $(X_j)_{j \in J}$ a collection of sets and for each $(j, k) \in \leq$, $\pi_{j,k} : X_k \rightarrow X_j$. Suppose that for each $j, k, l \in J$,

1. $\pi_{j,j} = \text{id}_{X_j}$,
2. $j \leq k$ and $k \leq l$ implies that $\pi_{j,k} \circ \pi_{k,l} = \pi_{j,l}$.

Then $((X_j)_{j \in J}, (\pi_{j,k})_{(j,k) \in \leq})$ is said to be a **projective system of sets**.

Definition 1.9.0.3. Let $(J, \leq)$ be a directed poset and $((X_j)_{j \in J}, (\pi_{j,k})_{(j,k) \in \leq})$ a projective system of sets. We define

- the **inverse limit** of $((X_j)_{j \in J}, (\pi_{j,k})_{(j,k) \in \leq})$, denoted $\varprojlim_{j \in J} X_j$, by

$$\varprojlim_{j \in J} X_j := \left\{ x \in \prod_{j \in J} X_j : \text{for each } (j, k) \in \leq, \pi_{j,k} \circ \text{proj}_k(x) = \text{proj}_j(x) \right\}$$

- the **j -th projection map of $\varprojlim_{j \in J} X_j$ onto X_j** , denoted $\pi_j : \varprojlim_{j \in J} X_j \rightarrow X_j$, by

$$\pi_j = \text{proj}_j \mid \varprojlim_{j \in J} X_j$$

Exercise 1.9.0.4. Let $(J, \leq)$ be a directed poset and $((X_j)_{j \in J}, (\pi_{j,k})_{(j,k) \in \leq})$ a projective system of sets. Then for each $j, k \in J$, $j \leq k$ implies that $\pi_{j,k} \circ \pi_k = \pi_j$.

Proof. Let $x \in \varprojlim_{j \in J} X_j$. Let $j, k \in J$. Suppose that $j \leq k$. By definition,

$$\begin{aligned} \pi_{j,k} \circ \pi_k(x) &= \pi_{j,k} \circ \text{proj}_k(x) \\ &= \text{proj}_j(x) \\ &= \pi_j(x). \end{aligned}$$

Since $x \in \varprojlim_{j' \in J} X_{j'}$ is arbitrary, we have that $\pi_{j,k} \circ \pi_k = \pi_j$. □

Exercise 1.9.0.5. Let $(J, \leq)$ be a directed poset and $((X_j)_{j \in J}, (\pi_{j,k})_{(j,k) \in \leq})$ a projective system of sets. Then

$$\varprojlim_{j \in J} X_j = \bigcap_{(j,k) \in \leq} \text{Eq}(\pi_{j,k} \circ \text{proj}_k, \text{proj}_j)$$

Proof. Set $X := \varprojlim_{j \in J} X_j$. By definition,

$$\begin{aligned} \varprojlim_{j \in J} X_j &= \bigcap_{(j,k) \in \leq} \{x \in X : \pi_{j,k} \circ \text{proj}_k(x) = \text{proj}_j(x)\} \\ &= \bigcap_{(j,k) \in \leq} \text{Eq}(\pi_{j,k} \circ \text{proj}_k, \text{proj}_j). \end{aligned}$$

□

Chapter 2

Ordered Sets

This chapter is deprecated and has been moved to intro to algebra notes, need to replace all references in this doc which link to this chapter over to there

2.1 Preorders

Definition 2.1.0.1. Preordered Set:

Let X be a set and $\leq \subset X \times X$ a binary relation on X . Then

- $\leq$ is said to be a **preorder on X** if
 1. for each $a \in X$, $a \leq a$
 2. for each $a, b, c \in X$, $a \leq b$ and $b \leq c$ implies that $a \leq c$
- $(X, \leq)$ is said to be a **preordered set** or **proset** if $\leq$ is a preorder on X .

Definition 2.1.0.2. Let $(A, \leq_A)$ and $(B, \leq_B)$ be directed sets. We define the

- **product preorder of $\leq_A$ and $\leq_B$ on $A \times B$** , denoted $\leq_A \otimes \leq_B$ by $(a_1, b_1) \leq_A \otimes \leq_B (a_2, b_2)$ iff $a_1 \leq_A a_2$ and $b_1 \leq_B b_2$.
- **product proset of $(A, \leq_A)$ and $(B, \leq_B)$** , denoted $(A, \leq_A) \otimes (B, \leq_B)$ by $(A, \leq_A) \otimes (B, \leq_B) := (A \times B, \leq_A \otimes \leq_B)$

Exercise 2.1.0.3. probably need to change notation since $\otimes$ might be reserved for something else. Let $(A, \leq_A)$ and $(B, \leq_B)$ be prosets. Then

1. $\leq_A \otimes \leq_B$ is a preorder on $A \times B$,
2. $(A, \leq_A) \otimes (B, \leq_B)$ is a proset.

Proof.

1. (a) Let $(a, b) \in A \times B$. Then $a \leq_A a$ and $b \leq_B b$. Therefore $(a, b) \leq_A \otimes \leq_B (a, b)$.
 (b) Let $(a_1, b_1), (a_2, b_2), (a_3, b_3) \in A \times B$. Suppose that $(a_1, b_1) \leq_A \otimes \leq_B (a_2, b_2)$ and $(a_2, b_2) \leq_A \otimes \leq_B (a_3, b_3)$. Then $a_1 \leq_A a_2$, $a_2 \leq_A a_3$, $b_1 \leq_B b_2$ and $b_2 \leq_B b_3$. Therefore $a_1 \leq_A a_3$ and $b_1 \leq_B b_3$. Hence $(a_1, b_1) \leq_A \otimes \leq_B (a_3, b_3)$.
 Hence $\leq_A \otimes \leq_B$ is a preorder on $A \times B$.
2. Since $\leq_A \otimes \leq_B$ is a preorder on $A \times B$, $(X, \leq)$ is a proset.

□

Definition 2.1.0.4. Let $(X, \leq)$ be a proset, $A \subset X$ and $x \in X$. Then x is said to be a

- **upper bound of A** if for each $a \in A$, $a \leq x$
- **lower bound of A** if for each $a \in A$, $a \geq x$

- **supremum of A or least upper bound of A** , denoted $x \in \sup A$, if

1. x is an upper bound of A
2. for each $y \in x$, y is an upper bound of A implies that $x \leq y$

If $\sup A = \{x\}$, we write $x = \sup A$.

- **infimum of A or greatest lower bound of A** , denoted $x \in \inf A$, if

1. x is a lower bound of A
2. for each $y \in x$, y is a lower bound of A implies that $x \geq y$.

If $\inf A = \{x\}$, we write $x = \inf A$.

Definition 2.1.0.5. Let $(X, \leq)$ be a poset, $A \subset X$. Then A is said to be

- **bounded above** if there exists $x \in X$ such that x is an upper bound of A .
- **bounded below** if there exists $x \in X$ such that x is a lower bound of A .

2.2 Directed Sets

Definition 2.2.0.1. Directed Set:

Let A be a set and $\leq \subset A \times A$ a binary relation on A . Then

- $\leq$ is said to be a **direction on A** if
 1. $\leq$ is a preorder on A
 2. for each $\alpha, \beta \in A$, there exists $\gamma \in A$ such that $\alpha, \beta \leq \gamma$
- $(A, \leq)$ is said to be a **directed set** if
 1. $A \neq \emptyset$
 2. $\leq$ is a direction on A

Exercise 2.2.0.2. Let $(A, \leq_A)$ and $(B, \leq_B)$ be directed sets. Then

1. $\leq_A \otimes \leq_B$ is a direction on $A \times B$,
2. $(A, \leq_A) \otimes (B, \leq_B)$ is a directed set.

Proof.

1. (a) Exercise 2.1.0.3 implies that $\leq_A \otimes \leq_B$ is a preorder of $A \times B$.
 (b) Let $(a_1, b_1), (a_2, b_2) \in A \times B$. Then there exist $a \in A$ and $b \in B$ such that $a_1, a_2 \leq_A a$ and $b_1, b_2 \leq_B b$. Hence $(a_1, b_1), (a_2, b_2) \leq_A \otimes \leq_B (a, b)$.

Hence $\leq_A \otimes \leq_B$ is a direction on $A \times B$.

2. (a) Since $A \neq \emptyset$ and $B \neq \emptyset$, we have that $A \times B \neq \emptyset$.
 (b) From above, $\leq_A \otimes \leq_B$ is a direction on $A \times B$.

Hence $(A \times B, \leq_A \otimes \leq_B)$ is a directed set.

□

2.3 Posets

Definition 2.3.0.1. Poset:

Let X be a set and $\leq \subset X \times X$ a binary relation on X . Then

- $\leq$ is said to be a **partial ordering on X** if for each $a, b, c \in X$,
 1. $\leq$ is a preorder on X
 2. $a \leq b$ and $b \leq a$ implies that $a = b$,
- $(X, \leq)$ is said to be a **partially ordered set** or **poset** if $\leq$ is a partial ordering on X .

Exercise 2.3.0.2. Let $(X, \leq)$ be a poset, $A \subset X$ and $x, y \in X$. If x and y are suprema of A , then $x = y$.

Proof. Suppose that x and y are suprema of A . Then x and y are upper bounds of A . Since x is a supremum of A and y is an upper bound of A , $x \leq y$. Since y is a supremum of A and x is an upper bound of A , $y \leq x$. Thus $x = y$. $\square$

Definition 2.3.0.3. Let $(X, \leq)$ be a poset. Then $(X, \leq)$ is said to satisfy the

- **least upper bound (LUB) property** if for each $A \subset X$, if $A \neq \emptyset$ and A is bounded above, then there exists $x \in X$ such that $x = \sup A$
- **greatest lower bound (GLB) property** if for each $A \subset X$, if $A \neq \emptyset$ and A is bounded below, then there exists $x \in X$ such that $x = \inf A$.

Exercise 2.3.0.4. LUB iff GLB

Proof. $\square$

Definition 2.3.0.5. Suplattice:

Let $(L, \leq)$ be a poset. Then $(L, \leq)$ is said to be a **suplattice** if for each $A \subset L$, there exists $x \in L$ such that $x = \sup A$.

Chapter 3

Real and Complex Numbers

3.1 Real Numbers

Note 3.1.0.1. As a starting point, we will take as fact the existence of the **natural numbers**

$$\mathbb{N} = \{1, 2, \dots\}$$

the **integers**

$$\mathbb{Z} = \{\dots, -2, -1, 0, 1, 2, \dots\}$$

and the **rational numbers**

$$\mathbb{Q} = \left\{ \frac{a}{b} : a \in \mathbb{Z}, b \in \mathbb{N} \right\}$$

Definition 3.1.0.2. Let X be a set and $\leq$ a relation on X . Then $\leq$ is said to be a **total order** if for each $a, b, c \in X$,

1. $a \leq a$
2. $a \leq b$ and $b \leq c$ implies that $a \leq c$
3. $a \leq b$ and $b \leq a$ implies that $a = b$
4. $a \leq b$ or $b \leq a$

Exercise 3.1.0.3. We define the relation $\leq$ on $\mathbb{Q}$ defined by

$$\frac{a}{b} \leq \frac{c}{d} \text{ iff } ad \leq bc$$

Then $\leq$ is a total order of $\mathbb{Q}$.

Proof. Let $\frac{a}{b}, \frac{c}{d}, \frac{e}{f} \in \mathbb{Q}$. Then

1. $\frac{a}{b} \leq \frac{a}{b}$ since $ab \leq ab$.
2. if $\frac{a}{b} \leq \frac{c}{d}$ and $\frac{c}{d} \leq \frac{e}{f}$, then $ad \leq bc$ and $cf \leq de$. Multiplying the first inequality by f and the second inequality by b , we obtain $adf \leq bcf \leq bde$. Dividing both sides by d yields $af \leq be$. Hence $\frac{a}{b} \leq \frac{e}{f}$.
3. if $\frac{a}{b} \leq \frac{c}{d}$ and $\frac{c}{d} \leq \frac{a}{b}$, then $ad \leq bc$ and $bc \leq ad$. This implies that $ad = bc$. Hence $\frac{a}{b} = \frac{c}{d}$.
- 4.

□

Definition 3.1.0.4. Let A be a set

Exercise 3.1.0.5. Let A, B be sets and $f : A \times B \rightarrow \mathbb{R}$. Then

$$\sup_{(a,b) \in A \times B} f(a,b) = \sup_{a \in A} \left[\sup_{b \in B} f(a,b) \right] = \sup_{b \in B} \left[\sup_{a \in A} f(a,b) \right]$$

Proof. For $(a,b) \in A \times B$, set $s_a = \sup_{b \in B} f(a,b)$ and $t_b = \sup_{a \in A} f(a,b)$. Let $(a,b) \in A \times B$. Then $\{a\} \times B \subset A \times B$. Therefore

$$\sup_{a \in A} s_a, \sup_{b \in B} t_b \leq \sup_{(x,y) \in A \times B} f(x,y)$$

Thus $\sup$

□

3.2 Extended Real Numbers

- define $\sup$ and $\inf$.

Definition 3.2.0.1.

- We define the **extended real numbers**, denoted $\overline{\mathbb{R}}$, by

$$\overline{\mathbb{R}} = \mathbb{R} \cup \{\pm\infty\}$$

- For $a, b \in \overline{\mathbb{R}}$, we define

$$\begin{aligned} a \pm \infty &= \pm\infty + a = \pm\infty, & a &\neq \mp\infty \\ a \cdot (\pm\infty) &= \pm\infty \cdot a = \pm\infty, & a &\in (0, +\infty] \\ a \cdot (\pm\infty) &= \pm\infty \cdot a = \mp\infty, & a &\in [-\infty, 0) \\ \frac{a}{\pm\infty} &= 0, & a &\in \mathbb{R} \\ \frac{\pm\infty}{a} &= \pm\infty, & a &\in (0, +\infty) \\ \frac{\pm\infty}{a} &= \mp\infty, & a &\in (-\infty, 0) \\ 0 \cdot (\pm\infty) &= \pm\infty \cdot 0 = 0 \end{aligned}$$

- We define $\leq_{\overline{\mathbb{R}}} \subset \overline{\mathbb{R}} \times \overline{\mathbb{R}}$ by

$$\leq_{\overline{\mathbb{R}}} = \leq_{\mathbb{R}} \cup \{(a,b) \in \overline{\mathbb{R}} \times \overline{\mathbb{R}} : a = -\infty \text{ or } b = \infty\}$$

Definition 3.2.0.2. Let $A \subset \overline{\mathbb{R}}$. We define the **supremum of A** , denoted $\sup A :=$.

3.3 Complex Numbers

We define $\text{Re}, \text{Im} : \mathbb{C} \rightarrow \mathbb{R}$ by $\text{Re}(a + ib) = a$ and $\text{Im}(a + ib) = b$.

Part I

Topology

Chapter 4

Topology

at this point, there is enough topology to make its own set of notes, covering topology, metric spaces, Frames and locales and pointless topology

One of the first fundamental ideas in algebra to which students are exposed is, roughly speaking, to think of an algebraic object, such as a group, ring, etc., in terms of its operations and relations. That is, we identify the structure of an object by what you can do with it. For instance in the context of groups, we often conceptualize a given group $(G, \cdot)$ in terms of its product $\cdot$.

Another fundamental idea to which students are exposed is, again roughly speaking, to learn about an algebraic object by viewing it from another perspective in the sense of viewing the object through different “lenses” which capture different aspects of its structure. For instance in the context of groups, we may study a given group $(G, \cdot)$ by studying its image under various homomorphisms (i.e. we process $(G, \cdot)$ in a way that preserves some aspect of its structure and study the processed output). By processing $(G, \cdot)$ and preserving only the information and structure in which we are interested, we can discard unrelated information, isolate the important information and simplify our study of $(G, \cdot)$.

In algebra, where the structure is given in terms of operations on the object, the appropriate notion of “structure preserving” map typically takes the form of a commutation relation between the map that processes the object and the operation. For example in the context of groups, a structure preserving map between two given groups $(G, \cdot)$ and $(H, *)$ is a map $f : G \rightarrow H$ which satisfies the relation that for each $g_1, g_2 \in G$, $f(g_1 \cdot g_2) = f(g_1) * f(g_2)$ so that the product structure of $(G, \cdot)$ is preserved through the processing map f in its image $f(G)$.

In this chapter, we examine topological spaces and, in analogy to algebraic objects, we discuss two ways to characterize the structure of a topological space, specifically open sets and nets, as well as two ways to characterize the notion of a map which preserves that structure.

4.1 Introduction

We will first begin by discussing a notion of a structure that allows us to describe objects which align with our usual, physical idea of what a “space” should be. Specifically, we define the concept of a topology in terms of open sets of a space. The notion of a topology in terms of open sets is perhaps unintuitive at first, but it affords one an almost geometric framework in which space can be thought.

Definition 4.1.0.1. Let X be a set and $\mathcal{T} \subset \mathcal{P}(X)$. Then $\mathcal{T}$ is said to be a **topology on X** if

1. $X, \emptyset \in \mathcal{T}$
2. for each $(U_\alpha)_{\alpha \in A} \subset \mathcal{T}$,

$$\bigcup_{\alpha \in A} U_\alpha \in \mathcal{T}$$

3. for each $(U_j)_{j=1}^n \subset \mathcal{T}$,

$$\bigcap_{j=1}^n U_j \in \mathcal{T}$$

Exercise 4.1.0.2. Let X be a set and $(\mathcal{T}_i)_{i \in I}$ a collection of topologies on X . Then $\bigcap_{i \in I} \mathcal{T}_i$ is a topology on X . change from i to α

Proof.

1. Since for each $i \in I$, $X, \emptyset \in \mathcal{T}_i$, we have that $X, \emptyset \in \bigcap_{i \in I} \mathcal{T}_i$.
2. Let $(U_\alpha)_{\alpha \in A} \subset \bigcap_{i \in I} \mathcal{T}_i$. Then for each $i \in I$, $(U_\alpha)_{\alpha \in A} \subset \mathcal{T}_i$. So for each $i \in I$, $\bigcup_{\alpha \in A} U_\alpha \in \mathcal{T}_i$. Thus $\bigcup_{\alpha \in A} U_\alpha \in \bigcap_{i \in I} \mathcal{T}_i$.
3. Let $(U_j)_{j=1}^n \subset \bigcap_{i \in I} \mathcal{T}_i$. Then for each $i \in I$, $(U_j)_{j=1}^n \subset \mathcal{T}_i$. So for each $i \in I$, $\bigcap_{j=1}^n U_j \in \mathcal{T}_i$. Thus $\bigcap_{j=1}^n U_j \in \bigcap_{i \in I} \mathcal{T}_i$.

So $\bigcap_{i \in I} \mathcal{T}_i$ is a topology on X . □

- introduce some definition for set of all topologies on X , maybe note that this is a poset ordered by inclusion and generated topology is value of left adjoint in galois connection from the power set to this poset (where the right adjoint is the forgetful $\mathcal{T} \in \text{set of topologies on } X \mapsto \mathcal{T} \in \mathcal{P}(X)$). We know a left adjoint is guaranteed since the right adjoint g preserves infima (i.e. intersections) so the left adjoint f of $\mathcal{E}$ is $\inf g^{-1}(\uparrow \mathcal{E})$, which one can easily show is the definition below)
- rework to say $\mathcal{T} \in \text{Top}(X)$ instead of $\mathcal{S}$?

Definition 4.1.0.3. Let X be a set and $\mathcal{E} \subset \mathcal{P}(X)$. Set

$$\mathcal{S} = \{\mathcal{T} \subset \mathcal{P}(X) : \mathcal{T} \text{ is a topology on } X \text{ and } \mathcal{E} \subset \mathcal{T}\}$$

We define the **topology generated by** $\mathcal{E}$ on X , denoted $\tau(\mathcal{E})$, by

$$\tau(\mathcal{E}) = \bigcap_{\mathcal{T} \in \mathcal{S}} \mathcal{T}$$

Exercise 4.1.0.4. Here introduce some properties of topology generated by sub collection of sets, like maybe $\mathcal{E}_1 \subset \mathcal{E}_2$ implies $\tau(\mathcal{E}_1) \subset \tau(\mathcal{E}_2)$?

Proof. **FINISH!!!** □

Definition 4.1.0.5. Let X be a set and $\mathcal{T} \subset \mathcal{P}(X)$ a topology on X , $x \in X$ and $\mathcal{B}_x \subset \mathcal{T}$. Then $\mathcal{B}_x$ is said to be a **local basis for** $\mathcal{T}$ at x if

1. for each $U \in \mathcal{B}_x$, $x \in U$
2. for each $V \in \mathcal{T}$, if $x \in V$, then there exists $U \in \mathcal{B}_x$ such that $U \subset V$

Exercise 4.1.0.6. Let X be a set, $\mathcal{T} \subset \mathcal{P}(X)$ a topology on X , $x \in X$ and $\mathcal{B}_x$ a local basis for $\mathcal{T}$ at x . Then $\mathcal{B}_x \neq \emptyset$.

Proof. Since $X \in \mathcal{T}$, $x \in X$ and $\mathcal{B}_x$ is a local basis for $\mathcal{T}$ at x , there exists $U \in \mathcal{B}_x$ such that $U \subset X$. Hence $\mathcal{B}_x \neq \emptyset$. □

Definition 4.1.0.7. Let X be a set and $\mathcal{T} \subset \mathcal{P}(X)$ a topology on X and $\mathcal{B} \subset \mathcal{T}$. Then $\mathcal{B}$ is said to be a **basis for** $\mathcal{T}$ if for each $V \in \mathcal{T}$ and $x \in V$, there exists $U \in \mathcal{B}$ such that $x \in U \subset V$.

Exercise 4.1.0.8. Let X be a set and $\mathcal{T} \subset \mathcal{P}(X)$ a topology on X and $\mathcal{B} \subset \mathcal{T}$. Then $\mathcal{B}$ is a basis for $\mathcal{T}$ iff for each $x \in X$, there exists $\mathcal{B}_x \subset \mathcal{B}$ such that $\mathcal{B}_x$ is a local basis for $\mathcal{T}$ at x .

Proof. Suppose that $\mathcal{B}$ is a basis for $\mathcal{T}$. Let $x \in X$. Define $\mathcal{B}_x = \{U \in \mathcal{B} : x \in U\}$.

1. By definition, for each $U \in \mathcal{B}_x$, $x \in U$
2. Let $V \in \mathcal{T}$. Suppose that $x \in V$. Since $\mathcal{B}$ is a basis, there exists $U \in \mathcal{B}$ such that $x \in U \subset V$. By definition, $U \in \mathcal{B}_x$.

Hence $\mathcal{B}_x$ is a local basis for $\mathcal{T}$ at x .

Conversely, suppose that for each $x \in X$, there exists $\mathcal{B}_x \subset \mathcal{B}$ such that $\mathcal{B}_x$ is a local basis for $\mathcal{T}$ at x . Let $V \in \mathcal{T}$ and $x \in V$. By assumption, there exists $\mathcal{B}_x \subset \mathcal{B}$ such that $\mathcal{B}_x$ is a local basis for $\mathcal{T}$ at x . Since $\mathcal{B}_x$ is a local basis for $\mathcal{T}$ at x , there exists $U \in \mathcal{B}_x \subset \mathcal{B}$ such that $x \in U \subset V$. Hence $\mathcal{B}$ is a basis for $\mathcal{T}$. □

Exercise 4.1.0.9. Let X be a set and $\mathcal{T} \subset \mathcal{P}(X)$ a topology on X and $\mathcal{B} \subset \mathcal{T}$. Then $\mathcal{B}$ is a basis for $\mathcal{T}$ iff for each $V \in \mathcal{T}$, there exists a collection $\mathcal{C} \subset \mathcal{B}$ such that

$$V = \bigcup_{U \in \mathcal{C}} U$$

Proof. Suppose that $\mathcal{B}$ is a basis for $\mathcal{T}$. Let $V \in \mathcal{T}$. Since $\mathcal{B}$ is a basis for $\mathcal{T}$, for each $x \in V$, there exists $U_x \in \mathcal{B}$ such that $x \in U_x \subset V$. Then $(U_x)_{x \in V} \subset \mathcal{B}$ satisfies $V = \bigcup_{x \in V} U_x$.

Conversely, suppose that for each $V \in \mathcal{T}$, there exists a collection $\mathcal{C} \subset \mathcal{B}$ such that $V = \bigcup_{U \in \mathcal{C}} U$. Let $V \in \mathcal{T}$ and $x \in V$. By assumption, there exists a collection $\mathcal{C} \subset \mathcal{B}$ such that $V = \bigcup_{U \in \mathcal{C}} U$. Since $x \in V$, there exists $U \in \mathcal{C}$ such that $x \in U$. Hence there exists $U \in \mathcal{B}$ such that $x \in U \subset V$. Then $\mathcal{B}$ is a basis for $\mathcal{T}$. $\square$

Exercise 4.1.0.10. Let X be a set and $\mathcal{T}_1, \mathcal{T}_2 \subset \mathcal{P}(X)$ topologies on X and $\mathcal{B} \subset \mathcal{T}_1$. Suppose that $\mathcal{T}_1 \subset \mathcal{T}_2$. If $\mathcal{B}$ is a basis for $\mathcal{T}_2$, then $\mathcal{B}$ is a basis for $\mathcal{T}_1$.

Proof. Suppose that $\mathcal{B}$ is a basis for $\mathcal{T}_2$. Let $V \in \mathcal{T}_1$. Then $V \in \mathcal{T}_2$. Since $\mathcal{B}$ is a basis for $\mathcal{T}_2$, the previous exercise implies that there exists a collection $(U_\alpha)_{\alpha \in A} \subset \mathcal{B}$ such that $V = \bigcup_{\alpha \in A} U_\alpha$. Thus the previous exercise implies that $\mathcal{B}$ is a basis for $\mathcal{T}_1$. $\square$

Exercise 4.1.0.11. Let X be a set and $\mathcal{B} \subset \mathcal{P}(X)$. Define

$$\mathcal{T}_{\mathcal{B}} = \{U \subset X : \text{for each } x \in U, \text{ there exists } V \in \mathcal{B} \text{ such that } x \in V \text{ and } V \subset U\}.$$

Then

1. $\mathcal{T}_{\mathcal{B}}$ is a topology on X iff
 - (a) for each $x \in X$, there exists $V \in \mathcal{B}$ such that $x \in V$
 - (b) for each $x \in X$ and $U, V \in \mathcal{B}$, if $x \in U \cap V$, then there exists $W \in \mathcal{B}$ such that $x \in W \subset U \cap V$
2. if $\mathcal{T}_{\mathcal{B}}$ is a topology on X , then $\mathcal{B}$ is a basis for $\mathcal{T}_{\mathcal{B}}$
3. if $\mathcal{T}_{\mathcal{B}}$ is a topology on X , then $\mathcal{T}_{\mathcal{B}} = \tau(\mathcal{B})$

Proof.

1. • $(\implies)$:
Suppose that $\mathcal{T}_{\mathcal{B}}$ is a topology on X .
 - (a) Let $x \in X$. Since $\mathcal{T}_{\mathcal{B}}$ is a topology on X , $X \in \mathcal{T}_{\mathcal{B}}$. Since $x \in X$, the definition of $\mathcal{T}_{\mathcal{B}}$ implies that there exists $V \in \mathcal{B}$ such that $x \in V \subset X$.
 - (b) Let $x \in X$ and $U, V \in \mathcal{B}$. Suppose that $x \in U \cap V$. Since $\mathcal{B} \subset \mathcal{T}$, we have that $U, V \in \mathcal{T}_{\mathcal{B}}$. Since $\mathcal{T}_{\mathcal{B}}$ is a topology on X , $U \cap V \in \mathcal{T}_{\mathcal{B}}$. By definition of $\mathcal{T}_{\mathcal{B}}$, there exists $W \in \mathcal{B}$ such that $x \in W \subset U \cap V$.
- $(\impliedby)$:
Conversely, suppose that (a) and (b) are satisfied.
 - Vacuously, $\emptyset \in \mathcal{T}_{\mathcal{B}}$. Condition (a) implies that $X \in \mathcal{T}_{\mathcal{B}}$.
 - Let $(U_\alpha)_{\alpha \in A} \subset \mathcal{T}_{\mathcal{B}}$ and $x \in \bigcup_{\alpha \in A} U_\alpha$. Then there exists $\alpha \in A$ such that $x \in U_\alpha$. Since $U_\alpha \in \mathcal{T}_{\mathcal{B}}$, the definition of $\mathcal{T}_{\mathcal{B}}$ implies that there exists $V \in \mathcal{B}$ such that

$$\begin{aligned} x &\in V \\ &\subset U_\alpha \\ &\subset \bigcup_{\alpha \in A} U_\alpha \end{aligned}$$

Since $x \in \bigcup_{\alpha \in A} U_\alpha$ is arbitrary, $\bigcup_{\alpha \in A} U_\alpha \in \mathcal{T}_{\mathcal{B}}$.

- Let $U_1, U_2 \in \mathcal{T}_B$ and $x \in U_1 \cap U_2$. The definition of $\mathcal{T}_B$ implies that for $j \in \{1, 2\}$, there exists $V_j \in \mathcal{B}$ such that $x \in V_j \subset U_j$. This implies that $x \in V_1 \cap V_2$ and by condition (b), there exists $W \in \mathcal{B}$ such that

$$\begin{aligned} x &\in W \\ &\subset V_1 \cap V_2 \\ &\subset U_1 \cap U_2 \end{aligned}$$

Since $x \in U_1 \cap U_2$ is arbitrary, $U_1 \cap U_2 \in \mathcal{T}_B$.

Thus $\mathcal{T}_B$ is a topology on X .

2. Suppose that $\mathcal{T}_B$ is a topology on X . Let $U \in \mathcal{T}_B$ and $x \in U$. By definition of $\mathcal{T}_B$, there exists $V \in \mathcal{B}$ such that $x \in V \subset U$. Since $U \in \mathcal{T}_B$ and $x \in U$ are arbitrary, $\mathcal{B}$ is a basis for $\mathcal{T}_B$.
3. Suppose that $\mathcal{T}_B$ is a topology on X . Since $\mathcal{B} \subset \mathcal{T}_B$, we have that $\tau(\mathcal{B}) \subset \mathcal{T}_B$. Let $U \in \tau(\mathcal{B})$. Conversely, let $U \in \mathcal{T}_B$. Since $\mathcal{T}_B$ is a topology on X , part (1) implies that $\mathcal{B}$ is a basis for $\mathcal{T}_B$. Then there exists $\mathcal{C} \subset \mathcal{B}$ such that

$$\begin{aligned} U &= \bigcup_{V \in \mathcal{C}} V \\ &\in \tau(\mathcal{B}) \end{aligned}$$

So $\mathcal{T}_B \subset \tau(\mathcal{B})$. Hence $\mathcal{T}_B = \tau(\mathcal{B})$. □

Exercise 4.1.0.12. Let X be a set and $\mathcal{B} \subset \mathcal{P}(X)$. Then $\mathcal{B}$ is a basis for $\tau(\mathcal{B})$ iff

1. for each $x \in X$, there exists $V \in \mathcal{B}$ such that $x \in V$
2. for each $x \in X$ and $U, V \in \mathcal{B}$, if $x \in U \cap V$, then there exists $W \in \mathcal{B}$ such that $x \in W \subset U \cap V$

Proof.

• ($\implies$) :

Suppose that $\mathcal{B}$ is a basis for $\tau(\mathcal{B})$.

1. Let $x \in X$. Since $X \in \tau(\mathcal{B})$ and $\mathcal{B}$ is a basis for $\tau(\mathcal{B})$, there exists $V \in \mathcal{B}$ such that $x \in V$.
2. Let $x \in X$ and $U, V \in \mathcal{B}$. Suppose that $x \in U \cap V$. Since $\tau(\mathcal{B})$ is a topology, $U \cap V \in \tau(\mathcal{B})$. Since $\mathcal{B}$ is a basis for $\tau(\mathcal{B})$, there exists $W \in \mathcal{B}$ such that $x \in W \subset U \cap V$.

• ($\impliedby$) :

Conversely, suppose that (1) and (2) are satisfied. The previous exercise implies that $\mathcal{B}$ is a basis for $\tau(\mathcal{B})$. □

Exercise 4.1.0.13. Let X be a set and $\mathcal{E} \subset \mathcal{P}(X)$. Define $\mathcal{B}, \mathcal{B}' \subset \mathcal{P}(X)$ by

$$\mathcal{B}' = \left\{ \bigcap_{j=1}^n V_j : n \in \mathbb{N} \text{ and } (V_j)_{j=1}^n \subset \mathcal{E} \right\}, \quad \mathcal{B} := \{X, \emptyset\} \cup \mathcal{B}'.$$

1. Then

- (a) $\mathcal{B}$ is a basis for $\tau(\mathcal{E})$
- (b)

$$\tau(\mathcal{E}) = \left\{ \bigcup_{\alpha \in A} V_\alpha : (V_\alpha)_{\alpha \in A} \subset \mathcal{B} \right\}$$

That is, each element of $\tau(\mathcal{E})$ is either $X, \emptyset$ or a union of finite intersections of elements of $\mathcal{E}$.

2. If there exist $(V_\alpha)_{\alpha \in A} \subset \mathcal{E}$ and $(U_j)_{j \in [n]} \subset \mathcal{E}$ such that $X = \bigcup_{\alpha \in A} V_\alpha$ and $\emptyset = \bigcap_{j \in [n]} U_j$. Then

- (a) $\mathcal{B}'$ is a basis for $\tau(\mathcal{E})$
- (b)

$$\tau(\mathcal{E}) = \left\{ \bigcup_{\alpha \in A} V_\alpha : (V_\alpha)_{\alpha \in A} \subset \mathcal{B} \right\}$$

That is, each element of $\tau(\mathcal{E})$ is a union of finite intersections of elements of $\mathcal{E}$.

Proof.

- 1. (a) Referring to Exercise 4.1.0.11, since $X \in \mathcal{B}$, condition (1) is satisfied and since for each $U, V \in \mathcal{B}$, $U \cap V \in \mathcal{B}$, condition (2) is satisfied. Hence there exists a topology $\mathcal{T}$ on X such that $\mathcal{B}$ is a basis for $\mathcal{T}$. Since $\mathcal{B} \subset \mathcal{T}$ and $\tau(\mathcal{E}) = \tau(\mathcal{B})$, we have that $\tau(\mathcal{E}) \subset \mathcal{T}$. Since $\mathcal{B}$ is a basis for $\mathcal{T}$ and $\mathcal{B} \subset \tau(\mathcal{E})$, Exercise 4.1.0.10 implies that $\mathcal{B}$ is a basis for $\tau(\mathcal{E})$.
- (b) Exercise 4.1.0.9 implies that

$$\tau(\mathcal{E}) = \left\{ \bigcup_{\alpha \in A} V_\alpha : (V_\alpha)_{\alpha \in A} \subset \mathcal{B} \right\}$$

2. Clear

□

Definition 4.1.0.14. Let X be a set and $\mathcal{T}$ a topology on X . Then $(X, \mathcal{T})$ is said to be a **topological space**. Let $U \subset X$. Then U is said to be **$\mathcal{T}$ -open** if $U \in \mathcal{T}$ and U is said to be **$\mathcal{T}$ -closed** if U^c is $\mathcal{T}$ -open.

Note 4.1.0.15. If we assume X is a topological space, we denote the implicit topology on X by $\mathcal{T}_X$.

Note 4.1.0.16. When the context is clear, we may say “open” or “closed” instead of “ $\mathcal{T}$ -open” or “ $\mathcal{T}$ -closed” respectively.

Definition 4.1.0.17. Let $(X, \mathcal{T})$ be a topological space and $S, N \subset X$. Then N is said to be a **$\mathcal{T}$ -neighborhood** of S if there exists $U \in \mathcal{T}$ such that $S \subset U$ and $U \subset N$. For $S \in X$, we define $\mathcal{N}(S, \mathcal{T}) := \{N \subset X : N \text{ is a } \mathcal{T}\text{-neighborhood of } S\}$

Note 4.1.0.18. When the context is clear, we write $\mathcal{N}(S)$ in place of $\mathcal{N}(S, \mathcal{T})$ and say “neighborhood” instead of “ $\mathcal{T}$ -neighborhood”.

Definition 4.1.0.19. Let $(X, \mathcal{T})$ be a topological space and $A \subset X$. We define

- $\text{IntSets}(A, \mathcal{T}) := \{U \subset X : U \subset A \text{ and } U \text{ is } \mathcal{T}\text{-open}\},$
- $\text{clSets}(A, \mathcal{T}) := \{C \subset X : A \subset C \text{ and } C \text{ is } \mathcal{T}\text{-closed}\},$

Note 4.1.0.20. When the context is clear, we write $\text{IntSets } A$ and $\text{clSets } A$ in place of $\text{IntSets}(A, \mathcal{T})$ and $\text{clSets}(A, \mathcal{T})$ respectively.

•

Definition 4.1.0.21. Let $(X, \mathcal{T})$ be a topological space and $A \subset X$. We define

- the **$\mathcal{T}$ -interior of A** , denoted $\text{Int}(A, \mathcal{T})$, by

$$\text{Int}(A, \mathcal{T}) = \bigcup_{U \in \text{IntSets}(A, \mathcal{T})} U,$$

- the **$\mathcal{T}$ -closure of A** , denoted $\text{cl}(A, \mathcal{T})$, by

$$\text{cl}(A, \mathcal{T}) = \bigcap_{C \in \text{clSets}(A, \mathcal{T})} C$$

Note 4.1.0.22. • When the context is clear, we write $\text{Int } A$ and $\text{cl } A$ in place of $\text{Int}(A, \mathcal{T})$ and $\text{cl}(A, \mathcal{T})$ respectively.

- For intuition, $\text{Int } A$ is the largest open subset of A and $\text{cl } A$ is the smallest closed superset of A .

Exercise 4.1.0.23. Let X be a topological space and $A \subset X$. Then

1. A is open iff $A = \text{Int } A$
2. A is closed iff $A = \text{cl } A$

Proof. Clear. □

Exercise 4.1.0.24. Let X be a topological space and $S, N \subset X$. Then $N \in \mathcal{N}(S)$ iff $S \subset \text{Int } N$.

Proof.

• $(\implies)$:

Suppose that $N \in \mathcal{N}(S)$. Then there exists $U \in \mathcal{T}_X$ such that $S \subset U$ and $U \subset N$. Hence $U \in \text{IntSets } N$ and therefore

$$\begin{aligned} S &\subset U \\ &\subset \bigcup_{V \in \text{IntSets } N} V \\ &= \text{Int } N. \end{aligned}$$

• $(\impliedby)$:

Suppose that $S \subset \text{Int } N$. Since $\text{Int } N \in \mathcal{T}_X$, $S \subset \text{Int } N$ and $\text{Int } N \subset N$, we have that $N \in \mathcal{N}(S)$. □

Exercise 4.1.0.25. Let $(X, \mathcal{T})$ be a topological space, $x \in X$ and $U, V \in \mathcal{N}(x, \mathcal{T})$. Then $U \cap V \in \mathcal{N}(x, \mathcal{T})$.

Proof. Since $U, V \in \mathcal{N}(x, \mathcal{T})$, Exercise 4.1.0.24 implies that $x \in \text{Int } U$ and $x \in \text{Int } V$. Then

$$\begin{aligned} x &\in \text{Int } U \cap \text{Int } V \\ &\subset U \cap V. \end{aligned}$$

Exercise 4.1.0.23 implies that $\text{Int } U, \text{Int } V \in \mathcal{T}$. Therefore $\text{Int } U \cap \text{Int } V \in \mathcal{T}$. Thus there exists $W \in \mathcal{T}$ such that $x \in W$ and $W \subset U \cap V$. By definition, $U \cap V \in \mathcal{N}(x, \mathcal{T})$. □

Exercise 4.1.0.26. Let X be a topological space and $A \subset X$. Then $(\text{Int } A)^c = \text{cl } A^c$.

Proof. We note that

1. for each $U \subset X$,

$$\begin{aligned} U \in \text{IntSets } A &\iff U \subset A \text{ and } U \text{ is open} \\ &\iff A^c \subset U^c \text{ and } U^c \text{ is closed} \\ &\iff U^c \in \text{clSets } A^c \end{aligned}$$

- 2.

$$\begin{aligned} C \in \text{clSets } A^c &\iff A^c \subset C \text{ and } C \text{ is closed} \\ &\iff C^c \subset A \text{ and } C^c \text{ is open} \\ &\iff C^c \in \text{IntSets } A \end{aligned}$$

Let $C \in \{U^c : U \in \text{IntSets } A\}$. Then there exists $U \in \text{IntSets } A$ such that $C = U^c$. By (1),

$$\begin{aligned} C &= U^c \\ &\in \text{clSets } A^c \end{aligned}$$

Since $C \in \{U^c : U \in \text{IntSets } A\}$ is arbitrary, $\{U^c : U \in \text{IntSets } A\} \subset \text{clSets } A^c$.

Conversely, let $C \in \text{clSets } A^c$. Then (2) implies that $C^c \in \text{IntSets } A$ and therefore

$$\begin{aligned} C &= (C^c)^c \\ &\in \{U^c : U \in \text{IntSets } A\} \end{aligned}$$

Since $C \in \text{clSets } A^c$ is arbitrary, $\text{clSets } A^c \subset \{U^c : U \in \text{IntSets } A\}$. Hence $\text{clSets } A^c = \{U^c : U \in \text{IntSets } A\}$ and therefore

$$\begin{aligned} (\text{Int } A)^c &= \left(\bigcup_{U \in \text{IntSets } A} U \right)^c \\ &= \bigcap_{U \in \text{IntSets } A} U^c \\ &= \bigcap_{C \in \text{clSets } A^c} C \\ &= \text{cl } A^c \end{aligned}$$

□

Exercise 4.1.0.27. Let X be a topological space and $A \subset X$. Then $(\text{cl } A)^c = \text{Int } A^c$.

Proof. Define $B = A^c$. The previous exercise implies that $(\text{Int } B)^c = \text{cl } B^c$. Therefore

$$\begin{aligned} \text{Int } A^c &= \text{Int } B \\ &= (\text{cl } B^c)^c \\ &= (\text{cl } A)^c \end{aligned}$$

□

Exercise 4.1.0.28. Let X be a topological space and $A \subset X$. Then A is open iff for each $x \in A$, there exists $U \in \mathcal{N}(x)$ such that U is open and $U \subset A$.

Proof. Suppose that A is open. Let $x \in A$. Then $A \in \mathcal{N}(x)$, A is open and $A \subset A$. Conversely, suppose that for each $x \in A$, there exists $U_x \in \mathcal{N}(x)$ such that U_x is open and $U_x \subset A$. Since

$$A = \bigcup_{x \in A} U_x,$$

we have that A is open. □

Exercise 4.1.0.29. Let X be a topological space, $A \subset X$ and $x \in X$. Then $x \in \text{cl } A$ iff for each $U \in \mathcal{N}(x)$, U is open implies that $A \cap U \neq \emptyset$.

Proof.

• ($\implies$)

Suppose that $x \in \text{cl } A$. Let $U \in \mathcal{N}(x)$. Suppose that U is open. For the sake of contradiction, suppose that $A \cap U = \emptyset$. Then $A \subset U^c$. Since U^c is closed, we have that

$$\begin{aligned} x &\in \text{cl } A \\ &\subset U^c \end{aligned}$$

This is a contradiction since $x \in U$. Hence $A \cap U \neq \emptyset$.

• ($\impliedby$)

Suppose that for each $U \in \mathcal{N}(x)$, U is open implies that $A \cap U \neq \emptyset$. For the sake of contradiction, suppose that $x \notin \text{cl } A$. By definition of closure, there exists $C \subset X$ such that C is closed, $A \subset C$ and $x \notin C$. Therefore C^c is open and $x \in C^c$. Thus $C^c \in \mathcal{N}(x)$. By assumption, $A \cap C^c \neq \emptyset$. This is a contradiction since $A \subset C$. So $x \in \text{cl } A$. □

Definition 4.1.0.30. Let $(X, \mathcal{T})$ be a topological space and $A \subset X$.

- Let $x \in X$. Then x is said to be a **$\mathcal{T}$ -limit point of A** if for each $U \in \mathcal{N}(x, \mathcal{T})$,

$$A \cap (U \setminus \{x\}) \neq \emptyset$$

- We define $\lim(A, \mathcal{T}) = \{x \in A : x \text{ is a limit point of } A\}$.

Exercise 4.1.0.31. Let $(X, \mathcal{T})$ be a topological space, $A \subset X$ and $x \in A$. Then $x \in \lim(A, \mathcal{T})$ iff for each $U \in \mathcal{N}^{\text{opn}}(x, \mathcal{T})$, $A \cap (U \setminus \{x\}) \neq \emptyset$.

Proof.

- ($\Rightarrow$):

Suppose that $x \in \lim(A, \mathcal{T})$. Then for each $U \in \mathcal{N}(x, \mathcal{T})$, $A \cap (U \setminus \{x\}) \neq \emptyset$. Let $U \in \mathcal{N}^{\text{opn}}(x, \mathcal{T})$. Since $\mathcal{N}^{\text{opn}}(x, \mathcal{T}) \subset \mathcal{N}(x, \mathcal{T})$, $U \in \mathcal{N}^{\text{opn}}(x, \mathcal{T})$. Thus $A \cap (U \setminus \{x\}) \neq \emptyset$. Since $U \in \mathcal{N}^{\text{opn}}(x, \mathcal{T})$ is arbitrary, we have that for each $U \in \mathcal{N}^{\text{opn}}(x, \mathcal{T})$, $A \cap (U \setminus \{x\}) \neq \emptyset$.

- ($\Leftarrow$):

Suppose that for each $U \in \mathcal{N}^{\text{opn}}(x, \mathcal{T})$, $A \cap (U \setminus \{x\}) \neq \emptyset$. Let $U \in \mathcal{N}(x, \mathcal{T})$. Then $\text{Int } U \in \mathcal{N}^{\text{opn}}(x, \mathcal{T})$. Hence $A \cap (\text{Int } U \setminus \{x\}) \neq \emptyset$. Thus there exists $a \in X$ such that $a \in A \cap (\text{Int } U \setminus \{x\})$. Then

$$\begin{aligned} a &\in A \cap (\text{Int } U \setminus \{x\}) &&= A \cap (\text{Int } U \cap \{x\}^c) \\ &= A \cap \text{Int } U \cap \{x\}^c \\ &\subset A \cap U \cap \{x\}^c \\ &= A \cap (U \cap \{x\}^c) &&= A \cap (U \setminus \{x\}). \end{aligned}$$

Thus $A \cap (U \setminus \{x\}) \neq \emptyset$. Since $U \in \mathcal{N}(x, \mathcal{T})$ is arbitrary, we have that for each $U \in \mathcal{N}(x, \mathcal{T})$, $A \cap (U \setminus \{x\}) \neq \emptyset$. □

Note 4.1.0.32. When context is clear, we write A' in place of $\lim(A, \mathcal{T})$.

Exercise 4.1.0.33. Let X be a topological space and $A \subset X$. Then $\text{cl } A = A \cup A'$.

Proof. Let $x \in A'$. For the sake of contradiction, suppose that $x \notin \text{cl } A$. By definition of closure, there exists $C \subset X$ such that C is closed, $A \subset C$ and $x \notin C$. Hence $x \in C^c \subset A^c$. Since C^c is open, $x \in \text{Int } A^c$. Since $x \in A'$ and $\text{Int } A^c \in \mathcal{N}(x)$, $[\text{Int } A^c \setminus \{x\}] \cap A \neq \emptyset$. This is a contradiction since $\text{Int } A^c \setminus \{x\} \subset A^c$. So $x \in \text{cl } A$ and $A' \subset \text{cl } A$. Since $A \subset \text{cl } A$, we have that $A \cup A' \subset \text{cl } A$. Conversely, let $x \in \text{cl } A$. For the sake of contradiction, suppose that $x \notin A \cup A'$. Then $x \in A^c \cap (A')^c$. Since $x \in (A')^c$, there exists $U \in \mathcal{N}(x)$ such that $(U \setminus \{x\}) \cap A = \emptyset$. Hence $U \setminus \{x\} \subset A^c$. Since $x \in A^c$,

$$\begin{aligned} \text{Int } U &\subset U \\ &= (U \setminus \{x\}) \cup \{x\} \\ &\subset A^c \end{aligned}$$

Which implies that $A \subset (\text{Int } U)^c$. Since $(\text{Int } U)^c$ is closed,

$$\begin{aligned} x &\in \text{cl } A \\ &\subset (\text{Int } U)^c \end{aligned}$$

which is a contradiction since $x \in \text{Int } U$. So $x \in A \cup A'$. Since $x \in \text{cl } A$ is arbitrary, $\text{cl } A \subset A \cup A'$. Therefore $\text{cl } A = A \cup A'$. □

Definition 4.1.0.34. Let $(X, \mathcal{T})$ be a topological space, $A \subset X$ and $x \in A$. Then x is said to a $\mathcal{T}$ -isolated point of A if $x \in \lim(A, \mathcal{T})^c$.

Note 4.1.0.35. When the context is clear, we may say “isolated point of A ” instead of “ $\mathcal{T}$ -isolated point of A ”.

Exercise 4.1.0.36. Let $(X, \mathcal{T})$ be a topological space, $A \subset X$ and $x \in A$. Then x is a $\mathcal{T}$ -isolated point of A iff there exists $U \in \mathcal{N}^{\text{opn}}(x, \mathcal{T})$ such that $U \cap A = \{x\}$.

Proof.

- ($\Rightarrow$) :

Suppose that x is a $\mathcal{T}$ -isolated point of A . Exercise ?? implies that there exists $U \in \mathcal{N}^{\text{opn}}(x, \mathcal{T})$ such that

$$\begin{aligned} A \cap U \cap \{x\}^c &= A \cap (U \setminus \{x\}) \\ &= \emptyset. \end{aligned}$$

Therefore

$$\begin{aligned} A \cap U &\subset (\{x\}^c)^c \\ &= \{x\}. \end{aligned}$$

Since $x \in A$ and $x \in U$, we have that

$$\begin{aligned} \{x\} &\subset A \cap U \\ &\subset \{x\}. \end{aligned}$$

Hence $A \cap U = \{x\}$.

- ($\Leftarrow$) :

Conversely, suppose that there exists $U \in \mathcal{N}^{\text{opn}}(x, \mathcal{T})$ such that $U \cap A = \{x\}$. Then

$$\begin{aligned} A \cap (U \setminus \{x\}) &= A \cap (U \cap \{x\}^c) \\ &= (A \cap U) \cap \{x\}^c &= (\{x\}) \cap \{x\}^c \\ &= \emptyset. \end{aligned}$$

Thus $x \in \text{lim}(A, \mathcal{T})^c$ and x is a $\mathcal{T}$ -isolated point of A .

-

□

Definition 4.1.0.37. Let $(X, \mathcal{T})$ be a topological space and $A \subset X$. Then A is said to be **discrete** if for each $x \in A$, x is an isolated point of A .

Definition 4.1.0.38. Let $(X, \mathcal{T})$ be a topological space and $A \subset X$. Then A is said to be **$\mathcal{T}$ -dense** in X if $\text{cl } A = X$.

Note 4.1.0.39. When the context is clear, we may say “dense in X ” instead of “ $\mathcal{T}$ -dense in X ”.

Exercise 4.1.0.40. Let X be a topological space and $A \subset X$. Then $A = \emptyset$ iff $\text{cl } A = \emptyset$.

Proof. Suppose $A = \emptyset$. Since A is closed,

$$\begin{aligned} \text{cl } A &= A \\ &= \emptyset \end{aligned}$$

Conversely, suppose that $\text{cl } A = \emptyset$. Since $A \subset \text{cl } A$, $A = \emptyset$.

□

Exercise 4.1.0.41. Let X be a topological space and $A \subset X$. Then A is dense in X iff for each $U \subset X$, U is open and $U \neq \emptyset$ implies that $A \cap U \neq \emptyset$.

Proof. Suppose that A is dense in X . Let $U \subset X$. Suppose that U is open. For the sake of contradiction, suppose that $A \cap U = \emptyset$. Then $U \subset A^c$. Thus $A \subset U^c$. Since U^c is closed, we have that

$$\begin{aligned} X &= \text{cl } A \\ &\subset U^c \end{aligned}$$

Therefore, $X = U^c$ and hence $U = \emptyset$. This is a contradiction. So for each $U \subset X$, U is open and $U \neq \emptyset$ implies that $A \cap U \neq \emptyset$. Conversely, suppose that for each $U \subset X$, if U is open and $U \neq \emptyset$, then $A \cap U \neq \emptyset$. Set $U = (\text{cl } A)^c$. Then U is open. For the sake of contradiction, suppose that $U \neq \emptyset$. By assumption there exists $x \in X$ such that

$$\begin{aligned} x &\in A \cap U \\ &= A \cap (\text{cl } A)^c \\ &\subset A \cap A^c \\ &= \emptyset \end{aligned}$$

which is a contradiction. Hence $U = \emptyset$. Then

$$\begin{aligned} X &= U^c \\ &= \text{cl } A \end{aligned}$$

so that A is dense in X . □

Definition 4.1.0.42. Let X be a topological space and $A \subset X$. Then A is said to be **nowhere dense** in X if $\text{Int cl } A = \emptyset$.

Exercise 4.1.0.43. Let X be a topological space and $A \subset X$. If A is nowhere dense in X , then $\text{cl } A$ is nowhere dense.

Proof. Suppose that A is nowhere dense in X . Then

$$\begin{aligned} \text{Int cl cl } A &= \text{Int cl } A \\ &= \emptyset \end{aligned}$$

Hence $\text{cl } A$ is nowhere dense. □

Exercise 4.1.0.44. Let X be a topological space and $A \subset X$. If A is nowhere dense in X , then A^c is dense.

Proof. Suppose that A is nowhere dense in X . Let $U \subset X$. Suppose that U is open and nonempty. For the sake of contradiction, suppose that $A^c \cap U = \emptyset$. Then

$$\begin{aligned} U &\subset (A^c)^c \\ &= A \\ &\subset \text{cl } A \end{aligned}$$

Since U is open, we have that

$$\begin{aligned} U &\subset \text{Int cl } A \\ &= \emptyset \end{aligned}$$

Therefore, $U = \emptyset$. This is a contradiction since U is nonempty. Hence $A^c \cap U \neq \emptyset$. Since U is arbitrary open nonempty subset of X , we have that for each $U \subset X$, if U is open and nonempty, then $A^c \cap U \neq \emptyset$. Thus A^c is dense. □

Definition 4.1.0.45. Let $(X, \mathcal{T})$ be a topological space, $A \subset X$ and $x \in X$. Then x is said to be a **$\mathcal{T}$ -condensation point** of A if for each $U \in \mathcal{N}(x, \mathcal{T})$, $U \cap A$ is uncountable.

Note 4.1.0.46. When the context is clear, we may say “condensation point” instead of “ $\mathcal{T}$ -condensation point”.

Exercise 4.1.0.47. Let $(X, \mathcal{T})$ be a topological space, $A \subset X$ and $x \in X$. If x is a $\mathcal{T}$ -condensation point of A , then x is a $\mathcal{T}$ -limit point of A

Proof. Suppose that x is a $\mathcal{T}$ -condensation point of A . Let $U \in \mathcal{N}(x, \mathcal{T})$. Since x is a $\mathcal{T}$ -condensation point of A , $U \cap A$ is uncountable. Hence

$$\begin{aligned} A \cap (U \setminus \{x\}) &= A \cap (U \cap \{x\}^c) \\ &= (A \cap U) \cap \{x\}^c \end{aligned}$$

is uncountable. In particular, $A \cap (U \setminus \{x\}) \neq \emptyset$. Since $U \in \mathcal{N}(x, \mathcal{T})$ is arbitrary, we have that for each $U \in \mathcal{N}(x, \mathcal{T})$, $A \cap (U \setminus \{x\}) \neq \emptyset$. Hence $x \in \lim_{\mathcal{T}} A$. $\square$

Exercise 4.1.0.48. Let $(X, \mathcal{T})$ be a topological space. Define $C \subset X$ by

$$C := \{x \in X : x \text{ is a } \mathcal{T}\text{-condensation point of } X\}.$$

Then

1. for each $x \in X$, $x \in C^c$ iff there exists $U \in \mathcal{T}$ such that $x \in U$, U is countable and $U \subset C^c$.
2. C is closed.

Proof.

1. Let $x \in X$.
 - $(\implies)$:
Suppose that $x \in C^c$. By definition, there exists $V \in \mathcal{N}(x)$ such that $x \in V$ and V is countable. Define $U \in \mathcal{T}$ by $U := \text{Int } V$. Since $V \in \mathcal{N}(x)$, we have that $x \in U$. Since $U \subset V$, we have that U is countable. Let $y \in U$. Since $y \in U$ and $U \cap X$ is countable, $y \in C^c$. Since $y \in U$ is arbitrary, we have that $U \subset C^c$.
 - $(\impliedby)$:
Suppose that there exists $U \in \mathcal{T}$ such that $x \in U$, U is countable and $U \subset C^c$. Then $U \in \mathcal{N}(x)$ and $U \cap X$ is countable. Thus $x \in C^c$.
2. The previous part implies that for each $x \in C^c$, there exists $U \in \mathcal{T}$ such that $x \in U$, U is countable and $U \subset C^c$. The axiom of choice implies that there exists $(U_x)_{x \in C^c} \subset \mathcal{T}$ such that for each $x \in C^c$, $x \in U_x$, U_x is countable and $U_x \subset C^c$. Thus

$$\begin{aligned} C^c &= \bigcup_{x \in C^c} U_x \\ &\in \mathcal{T}. \end{aligned}$$

Since C^c is open, we have that C is closed. $\square$

Exercise 4.1.0.49. Let $(X, \mathcal{T})$ be a topological space $x \in X$. The following are equivalent:

1. x is a $\mathcal{T}$ -condensation point of X ,
2. for each $U \in \mathcal{N}(x, \mathcal{T})$, U is uncountable,
3. for each $U \in \mathcal{N}(x, \mathcal{T})$, x is a $\mathcal{T}$ -condensation point of U ,

Proof.

1. $(1) \implies (2)$:
Suppose that x is a $\mathcal{T}$ -condensation point of X . Let $U \in \mathcal{N}(x, \mathcal{T})$. Since x is a $\mathcal{T}$ -condensation point of X , $U \cap X$ is uncountable. Since $U = U \cap X$, U is uncountable. Since $U \in \mathcal{N}(x, \mathcal{T})$ is arbitrary, we have that for each $U \in \mathcal{N}(x, \mathcal{T})$, U is uncountable.
2. $(2) \implies (3)$:
Suppose that for each $U \in \mathcal{N}(x, \mathcal{T})$, U is uncountable. Let $U, V \in \mathcal{N}(x, \mathcal{T})$. Exercise 4.1.0.25 implies that $U \cap V \in \mathcal{N}(x, \mathcal{T})$. Then by assumption, $U \cap V$ is uncountable. Since $V \in \mathcal{N}(x, \mathcal{T})$ is arbitrary, we have that for each $V \in \mathcal{N}(x, \mathcal{T})$, $U \cap V$ is uncountable. Hence x is a $\mathcal{T}$ -condensation point of U . Since $U \in \mathcal{N}(x, \mathcal{T})$ is arbitrary, we have that for each $U \in \mathcal{N}(x, \mathcal{T})$, x is a $\mathcal{T}$ -condensation point of U .

3. (3) $\implies$ (1):

Suppose that for each $U \in \mathcal{N}(x, \mathcal{T})$, x is a $\mathcal{T}$ -condensation point of U . Let $U \in \mathcal{N}(x, \mathcal{T})$. Since $X \in \mathcal{N}(x, \mathcal{T})$, and x is a $\mathcal{T}$ -condensation point of U , we have that $U \cap X$ is uncountable. Since $U \in \mathcal{N}(x, \mathcal{T})$ is arbitrary, we have that for each $U \in \mathcal{N}(x, \mathcal{T})$, $U \cap X$ is uncountable. Hence x is a $\mathcal{T}$ -condensation point of X . □

Definition 4.1.0.50. maybe make a notation for the set of G_δ/F_σ -sets Let $(X, \mathcal{T})$ be a topological space and $A \subset X$. Then

- A is said to be a G_δ -set of $(X, \mathcal{T})$ if there exist $(U_n)_{n \in \mathbb{N}} \subset \mathcal{T}$ such that $A = \bigcap_{n \in \mathbb{N}} U_n$
- A is said to be an F_σ -set of $(X, \mathcal{T})$ if A^c is a G_δ -set.

Note 4.1.0.51. When the context is clear, we may say A is a “ G_δ -set (resp. F_σ -set) of X ” instead of “ G_δ -set (resp. F_σ -set) of $(X, \mathcal{T})$ ”.

Exercise 4.1.0.52. Let $(X, \mathcal{T})$ be a topological space and $A \subset X$. Then A is an F_σ -set iff there exists $(C_n)_{n \in \mathbb{N}} \subset \mathcal{P}(X)$ such that for each $n \in \mathbb{N}$, C_n is $\mathcal{T}$ -closed and $A = \bigcup_{n \in \mathbb{N}} C_n$.

Proof. Clear □

Definition 4.1.0.53. Let X be a topological space, $E \subset X$.

- Let $x \in X$. Then x is said to be a **boundary point of E** if for each $U \in \mathcal{N}(x)$, $U \cap E \neq \emptyset$ and $U \cap E^c \neq \emptyset$.
- We define the **boundary of E** , denoted ∂E , by

$$\partial E = \{x \in X : x \text{ is a boundary point of } E\}$$

4.2 Continuous Maps

Now that we have introduced the definition of a topological space, we would like to characterize the transformations of one

Definition 4.2.0.1. Let $(X, \mathcal{T}_X)$ and $(Y, \mathcal{T}_Y)$ be topological spaces.

- Let $f : X \rightarrow Y$. Then f is said to be $(\mathcal{T}_X, \mathcal{T}_Y)$ -**continuous** if for each $B \in \mathcal{T}_Y$, $f^{-1}(B) \in \mathcal{T}_X$.
- We define $C((X, \mathcal{T}_X), (Y, \mathcal{T}_Y)) := \{f : X \rightarrow Y : f \text{ is } (\mathcal{T}_X, \mathcal{T}_Y)\text{-continuous}\}$.

Note 4.2.0.2. When the context is clear,

- we may write $C(X, Y)$ in place of $C((X, \mathcal{T}_X), (Y, \mathcal{T}_Y))$.
- we may write $C(X)$ in place of $C((X, \mathcal{T}_X), (\mathbb{C}, \mathcal{T}_{\mathbb{C}}))$.

Exercise 4.2.0.3. Let $(X, \mathcal{T}_X)$ and $(Y, \mathcal{T}_Y)$ be topological spaces and $f : X \rightarrow Y$. Then f is $(\mathcal{T}_X, \mathcal{T}_Y)$ -continuous iff for each $C \subset Y$, C is $\mathcal{T}_Y$ -closed implies that $f^{-1}(C)$ is $\mathcal{T}_X$ -closed.

Proof.

- $(\implies)$:
Suppose that f is $(\mathcal{T}_X, \mathcal{T}_Y)$ -continuous. Then f is $(\mathcal{T}_X, \mathcal{T}_Y)$ -continuous. Let $C \subset Y$. Suppose that C is $\mathcal{T}_Y$ -closed. Then $C^c \in \mathcal{T}_Y$. Since f is $(\mathcal{T}_X, \mathcal{T}_Y)$ -continuous, we have that

$$\begin{aligned} f^{-1}(C)^c &= f^{-1}(C^c) \\ &\in \mathcal{T}_X. \end{aligned}$$

Thus $f^{-1}(C)$ is $\mathcal{T}_X$ -closed. Since $C \subset Y$ with C closed in $(Y, \mathcal{T}_Y)$ is arbitrary, we have that for each $C \subset Y$, C is $\mathcal{T}_Y$ -closed implies that $f^{-1}(C)$ is $\mathcal{T}_X$ -closed.

- $(\impliedby)$:
Suppose that for each $C \subset Y$, C is $\mathcal{T}_Y$ -closed implies that $f^{-1}(C)$ is $\mathcal{T}_X$ -closed. Let $B \in \mathcal{T}_Y$. Then B^c is $\mathcal{T}_Y$ -closed. Hence $f^{-1}(B^c)$ is $\mathcal{T}_X$ -closed. Since $f^{-1}(B)^c = f^{-1}(B^c)$, we have that $f^{-1}(B)^c$ is $\mathcal{T}_X$ -closed. Hence $f^{-1}(B) \in \mathcal{T}_X$. Since $B \in \mathcal{T}_Y$ is arbitrary, we have that for each $B \in \mathcal{T}_Y$, $f^{-1}(B) \in \mathcal{T}_X$. Thus f is $(\mathcal{T}_X, \mathcal{T}_Y)$ -continuous.

□

Recall or give explanation on how topologies capture notion of arbitrary 'closeness'. Figure out how to explain this, copy to previous section. Ideas: if $U \in \mathcal{T} \cap \mathcal{N}(x)$ and $a \in U$, then a is U -close to x . $a = x$ (if hausdorff) when for each $U \in \mathcal{T} \cap \mathcal{N}(x)$, a is U -close, try to motivate topology axioms in terms of U -closeness, so if y is U_α -close to x for each $\alpha \in A$, then y is $\bigcup_{\alpha \in A} U_\alpha$ -close to x ,

draw a picture of this with sets in $\mathbb{R}^2$ with usual notion of distance, where U -closeness could correspond to supremum distance of U to x ? If y is U_1, U_2 -close to x , then y is $U_1 \cap U_2$ -close to x , here for finite neighborhoods, if y is inside each U_1, U_2 , then y is inside $U_1 \cap U_2$, drawing the sup distance in $\mathbb{R}^2$ would agree intuitively.

a map that preserves the notion of closeness should send points getting closer to points getting closer. Once candidate condition for continuity that satisfies this is $A \subset B$ implies that $f(A) \subset f(B)$. Show this is not sufficient for our intuition with some concrete counter examples and that the issue is that even though closer points would go to closer points, the inputs in the domain could be getting arbitrarily close, but the outputs don't get arbitrarily close to $f(x)$, i.e. come up with $(U_\alpha)_{\alpha \in A} \subset \mathcal{T}_X \cap \mathcal{N}(x)$ and $V \in \mathcal{N}(f(x)) \cap \mathcal{T}_Y$ such that $\alpha \leq \alpha'$ implies that $U_{\alpha'} \subset U_\alpha$, $\bigcap_{\alpha} U_\alpha = \{x\}$ and for each $\alpha \in A$, $f(U_\alpha) \not\subset V$. Translate this idea using the word "tolerances"

show that the negation of this issue is that if a sceptic gives you a tolerance $V \in \mathcal{N}(f(x))$ (i.e. for each V , doesn't matter which one they give), you can find (i.e. there exists) $U \in \mathcal{N}(x)$ such that $f(U) \subset V$.

Move this to nets section: Alternatively to say a map is continuous at a point is the same as demonstrating to someone that the map sends inputs getting arbitrarily close to x to outputs getting arbitrarily close to $f(x)$. That is if we have a net $x_\alpha \rightarrow x$, then $f(x_\alpha) \rightarrow f(x)$. in the sense that if that sceptic gives me a closeness tolerance for the outputs closeness in terms of tolerances

Definition 4.2.0.4. Let $(X, \mathcal{T}_X)$ and $(Y, \mathcal{T}_Y)$ be topological spaces, $f : X \rightarrow Y$ and $x \in X$. Then f is said to be $(\mathcal{T}_X, \mathcal{T}_Y)$ -**continuous at x** if for each $V \in \mathcal{N}(f(x), \mathcal{T}_Y)$, there exists $U \in \mathcal{N}(x, \mathcal{T}_X)$ such that $f(U) \subset V$.

Exercise 4.2.0.5. Let $(X, \mathcal{T}_X)$ and $(Y, \mathcal{T}_Y)$ be topological spaces, $f : X \rightarrow Y$ and $x \in X$. Then f is $(\mathcal{T}_X, \mathcal{T}_Y)$ -continuous at x iff for each $V \in \mathcal{N}(f(x), \mathcal{T}_Y)$, $f^{-1}(V) \in \mathcal{N}(x, \mathcal{T}_X)$.

Hint: for $U \in \mathcal{N}(x, \mathcal{T}_X)$ and $V \in \mathcal{N}(f(x), \mathcal{T}_Y)$, consider $f^{-1}(f(U))$ and $f(f^{-1}(V))$

Proof.

• ($\implies$):

Suppose that f is $(\mathcal{T}_X, \mathcal{T}_Y)$ -continuous at x . Let $V \in \mathcal{N}(f(x), \mathcal{T}_Y)$. Then there exists $U \in \mathcal{N}(x, \mathcal{T}_X)$ such that $f(U) \subset V$. Thus

$$\begin{aligned} x &\in \text{Int } U \\ &\subset U \\ &\subset f^{-1}(f(U)) \quad (\text{by ref set theory ex}) \\ &\subset f^{-1}(V) \quad (\text{by ref set theory ex}) \end{aligned}$$

So $f^{-1}(V) \in \mathcal{N}(x, \mathcal{T}_X)$.

• ($\impliedby$):

Conversely, suppose that for each $V \in \mathcal{N}(f(x), \mathcal{T}_Y)$, $f^{-1}(V) \in \mathcal{N}(x, \mathcal{T}_X)$. Let $V \in \mathcal{N}(f(x), \mathcal{T}_Y)$. Define $U \subset X$ by $U := f^{-1}(V)$. By assumption,

$$\begin{aligned} U &= f^{-1}(V) \\ &\in \mathcal{N}(x, \mathcal{T}_X). \end{aligned}$$

Then

$$f(U) = f(f^{-1}(V)) \subset V \quad (\text{by ref set theory ex})$$

Since $V \in \mathcal{N}(f(x), \mathcal{T}_Y)$ is arbitrary, we have that for each $V \in \mathcal{N}(f(x), \mathcal{T}_Y)$, there exists $U \in \mathcal{N}(x, \mathcal{T}_X)$ such that $f(U) \subset V$. Thus f is $(\mathcal{T}_X, \mathcal{T}_Y)$ -continuous at x . □

Exercise 4.2.0.6. Let $(X, \mathcal{T}_X)$ and $(Y, \mathcal{T}_Y)$ be topological spaces and $f : X \rightarrow Y$. Then f is $(\mathcal{T}_X, \mathcal{T}_Y)$ -continuous iff for each $x \in X$, f is $(\mathcal{T}_X, \mathcal{T}_Y)$ -continuous at x .

Proof.

• ($\implies$):

Suppose that f is $(\mathcal{T}_X, \mathcal{T}_Y)$ -continuous. Let $x \in X$ and $V \in \mathcal{N}(f(x), \mathcal{T}_Y)$. Then

$$\begin{aligned} f(x) &\in \text{Int}(V, \mathcal{T}_Y) \\ &\in \mathcal{T}_Y. \end{aligned}$$

Define $U \subset X$ by $U := f^{-1}(\text{Int}(V, \mathcal{T}_Y))$. Since f is $(\mathcal{T}_X, \mathcal{T}_Y)$, we have that $U \in \mathcal{T}_X$. Since $V \in \mathcal{N}(f(x), \mathcal{T}_Y)$, we have that $f(x) \in \text{Int}(V, \mathcal{T}_Y)$. Therefore

$$\begin{aligned} x &\in f^{-1}(\text{Int}(V, \mathcal{T}_Y)) \\ &= U. \end{aligned}$$

Hence $U \in \mathcal{N}(x, \mathcal{T}_X)$. Then

$$\begin{aligned} f(U) &= f(f^{-1}(\text{Int}(V, \mathcal{T}_Y))) \\ &\subset \text{Int}(V, \mathcal{T}_Y) \quad (\text{by ref set theory ex}) \\ &\subset V. \end{aligned}$$

Since $V \in \mathcal{N}(f(x), \mathcal{T}_Y)$ is arbitrary, we have that for each $V \in \mathcal{N}(f(x), \mathcal{T}_Y)$, there exists $U \in \mathcal{N}(x, \mathcal{T}_X)$ such that $f(U) \subset V$. Therefore f is $(\mathcal{T}_X, \mathcal{T}_Y)$ -continuous at x .

• ($\Leftarrow$):

Conversely, suppose that for each $x \in X$, f is $(\mathcal{T}_X, \mathcal{T}_Y)$ -continuous at x . Let $V \in \mathcal{T}_Y$ and $x \in f^{-1}(V)$. Then $V \in \mathcal{N}(f(x), \mathcal{T}_Y)$. Since f is $(\mathcal{T}_X, \mathcal{T}_Y)$ -continuous at x , we have that $f^{-1}(V) \in \mathcal{N}(x, \mathcal{T}_X)$. Then $x \in \text{Int}(f^{-1}(V), \mathcal{T}_X)$. Since $x \in f^{-1}(V)$ is arbitrary, we have that for each $x \in f^{-1}(V)$, $x \in \text{Int}(f^{-1}(V), \mathcal{T}_X)$. Hence

$$\begin{aligned} f^{-1}(V) &\subset \text{Int}(f^{-1}(V), \mathcal{T}_X) \\ &\subset f^{-1}(V). \end{aligned}$$

Thus

$$\begin{aligned} f^{-1}(V) &= \text{Int}(f^{-1}(V), \mathcal{T}_X) \\ &\in \mathcal{T}_X. \end{aligned}$$

Since $V \in \mathcal{T}_Y$ is arbitrary, we have that for each $V \in \mathcal{T}_Y$, $f^{-1}(V) \in \mathcal{T}_X$. So f is $(\mathcal{T}_X, \mathcal{T}_Y)$ -continuous. □

Definition 4.2.0.7. Let $(X, \mathcal{T}_X)$ and $(Y, \mathcal{T}_Y)$ be topological spaces and $f : X \rightarrow Y$. We define the

1. **push-forward of $\mathcal{T}_X$** , denoted $f_*\mathcal{T}_X$, by

$$f_*\mathcal{T}_X := \{V \subset Y : f^{-1}(V) \in \mathcal{T}_X\},$$

2. **pull-back of $\mathcal{T}_Y$** , denoted $f^*\mathcal{T}_Y$, by

$$f^*\mathcal{T}_Y := \{f^{-1}(V) : V \in \mathcal{T}_Y\}.$$

Exercise 4.2.0.8. Let $(X, \mathcal{T}_X)$ and $(Y, \mathcal{T}_Y)$ be topological spaces and $f : X \rightarrow Y$. Then

1. $f_*\mathcal{T}_X$ is a topology on Y
2. $f^*\mathcal{T}_Y$ is a topology on X

Proof.

1. • Since $f^{-1}(Y) = X \in \mathcal{T}_X$ and $f^{-1}(\emptyset) = \emptyset \in \mathcal{T}_X$, $Y, \emptyset \in f_*\mathcal{T}_X$.
• Let $(U_\alpha)_{\alpha \in A} \subset f_*\mathcal{T}_X$. Then for each $\alpha \in A$, $f^{-1}(U_\alpha) \in \mathcal{T}_X$. This implies that

$$\begin{aligned} f^{-1}\left(\bigcup_{\alpha \in A} U_\alpha\right) &= \bigcup_{\alpha \in A} f^{-1}(U_\alpha) \\ &\in \mathcal{T}_X \end{aligned}$$

Hence $\bigcup_{\alpha \in A} U_\alpha \in f_*\mathcal{T}_X$.

- Let $(U_j)_{j=1}^n \subset f_*\mathcal{T}_X$. Then for each $j \in 1, \dots, n$, $f^{-1}(U_j) \in \mathcal{T}_X$. This implies that

$$\begin{aligned} f^{-1}\left(\bigcap_{j=1}^n U_j\right) &= \bigcap_{j=1}^n f^{-1}(U_j) \\ &\in \mathcal{T}_X \end{aligned}$$

Hence $\bigcap_{j=1}^n U_j \in f_*\mathcal{T}_X$.

So $f_*\mathcal{T}_X$ is a topology on Y .

2. Similar to (1). □

Exercise 4.2.0.9. Let $(X, \mathcal{T}_X)$ and $(Y, \mathcal{T}_Y)$ be topological spaces, $f : X \rightarrow Y$ and $\mathcal{E} \subset \mathcal{P}(Y)$. Suppose that $\mathcal{T}_Y = \tau(\mathcal{E})$. Then f is continuous iff for each $B \in \mathcal{E}$, $f^{-1}(B) \in \mathcal{T}_X$.

Proof. Suppose that f is continuous. Since $\mathcal{E} \subset \mathcal{T}_Y$, clearly for each $B \in \mathcal{E}$, $f^{-1}(B) \in \mathcal{T}_X$.

Conversely, suppose that for each $B \in \mathcal{E}$, $f^{-1}(B) \in \mathcal{T}_X$. Then $\mathcal{E} \subset f_*\mathcal{T}_X$. Since $f_*\mathcal{T}_X$ is a topology on Y , we have that $\mathcal{T}_Y = \tau(\mathcal{E}) \subset f_*\mathcal{T}_X$. So f is continuous. $\square$

Definition 4.2.0.10. Let X be a set, $(Y_\alpha, \mathcal{T}_\alpha)_{\alpha \in A}$ a collection of topological spaces and $\mathcal{F} \in \prod_{\alpha \in A} Y_\alpha^X$ (i.e. $\mathcal{F} = (f_\alpha)_{\alpha \in A}$ where for each $\alpha \in A$, $f_\alpha : X \rightarrow Y_\alpha$). We define the **initial topology on X generated by $\mathcal{F}$** , denoted $\tau_X(\mathcal{F})$, by

$$\tau_X(\mathcal{F}) = \tau_X(\{f_\alpha^{-1}(B) : B \in \mathcal{T}_\alpha \text{ and } \alpha \in A\})$$

Note 4.2.0.11. The initial topology topology generated by $\mathcal{F}$ is also called the **weak topology generated by $\mathcal{F}$** and if $\mathcal{F} = \{f\}$, then $\tau_X(\mathcal{F}) = f^*\mathcal{T}_Y$.

Exercise 4.2.0.12. Let $(X, \mathcal{T})$ be a topological, $(Y_\alpha, \mathcal{T}_\alpha)_{\alpha \in A}$ a collection of topological spaces and $\mathcal{F} \in \prod_{\alpha \in A} Y_\alpha^X$ (i.e. $\mathcal{F} = (f_\alpha)_{\alpha \in A}$ where for each $\alpha \in A$, $f_\alpha : X \rightarrow Y_\alpha$). If for each $\alpha \in A$, f_α is $(\mathcal{T}, \mathcal{T}_\alpha)$ -continuous, then $\tau_X(\mathcal{F}) \subset \mathcal{T}$.

Proof. Let $\mathcal{T} \subset \mathcal{P}(X)$. Suppose that $\mathcal{T}$ is a topology on X and for each $\alpha \in A$, f_α is $(\mathcal{T}, \mathcal{T}_\alpha)$ -continuous. Set $\mathcal{V} := \{f_\alpha^{-1}(V) : V \in \mathcal{T}_\alpha \text{ and } \alpha \in A\}$. By definition, $\tau_X(\mathcal{F}) = \tau_X(\mathcal{V})$. Since for each $\alpha \in A$, f_α is $(\mathcal{T}, \mathcal{T}_\alpha)$ -continuous, we have that for each $\alpha \in A$ and $V \in \mathcal{T}_\alpha$, $f_\alpha^{-1}(V) \in \mathcal{T}$. Hence $\mathcal{V} \subset \mathcal{T}$. Therefore

$$\begin{aligned} \tau_X(\mathcal{F}) &= \tau_X(\mathcal{V}) \\ &\subset \mathcal{T}. \end{aligned}$$

$\square$

Note 4.2.0.13. Essentially, $\tau_X(\mathcal{F})$ is the smallest topology on X such that for each $\alpha \in A$, $f_\alpha : X \rightarrow Y_\alpha$ is continuous.

Exercise 4.2.0.14. Let $(Y_\alpha, \mathcal{T}_\alpha)_{\alpha \in A}$ be a collection of topological spaces, X a set, $(Z, \mathcal{C})$ a topological space, $\mathcal{F} = (f_\alpha)_{\alpha \in A} \in \prod_{\alpha \in A} Y_\alpha^X$ and $g : Z \rightarrow X$. Then g is $(\mathcal{C}, \tau_X(\mathcal{F}))$ -continuous iff for each $\alpha \in A$, $f_\alpha \circ g$ is $(\mathcal{C}, \mathcal{T}_\alpha)$ -continuous:

$$\begin{array}{ccc} Y_\alpha & \xleftarrow{f_\alpha} & X \\ & \nwarrow f_\alpha \circ g & \uparrow g \\ & & Z \end{array}$$

Proof. If g is $(\mathcal{C}, \tau_X(\mathcal{F}))$ -continuous, then clearly for each $\alpha \in A$, $f_\alpha \circ g$ is $(\mathcal{C}, \mathcal{T}_\alpha)$ -continuous.

Conversely, suppose that for each $\alpha \in A$, $f_\alpha \circ g$ is $(\mathcal{C}, \mathcal{T}_\alpha)$ -continuous. Let $\alpha \in A$ and $V \in \mathcal{T}_\alpha$. Continuity implies that,

$$\begin{aligned} g^{-1}(f_\alpha^{-1}(V)) &= (f_\alpha \circ g)^{-1}(V) \\ &\in \mathcal{C} \end{aligned}$$

Since $\alpha \in A$ and $V \in \mathcal{T}_\alpha$ are arbitrary, we have that for each $\alpha \in A$ and $V \in \mathcal{T}_\alpha$, $g^{-1}(f_\alpha^{-1}(V)) \in \mathcal{C}$. Since $\tau_X(\mathcal{F}) = \tau(\{f_\alpha^{-1}(V) : \alpha \in A \text{ and } V \in \mathcal{T}_\alpha\})$, Exercise 4.2.0.9 implies that g is $(\mathcal{C}, \tau_X(\mathcal{F}))$ -continuous. $\square$

Exercise 4.2.0.15. Let $(X, \mathcal{T})$ be a topological space. Set $\mathcal{F} = \text{Hom}_{\mathbf{Top}}((X, \mathcal{T}), (X, \mathcal{T}))$. Then $\tau_X(\mathcal{F}) = \mathcal{T}$.

Proof. Set $\mathcal{E} = \{f^{-1}(V) : V \in \mathcal{T}_Y \text{ and } f \in \mathcal{F}\}$. Since for each $f \in \mathcal{F}$, f is $(\mathcal{T}, \mathcal{T})$ -continuous, $\mathcal{E} \subset \mathcal{T}$.

Conversely, since $\text{id}_X \in \mathcal{F}$, we have that for each $U \in \mathcal{T}$,

$$\begin{aligned} U &= \text{id}_X^{-1}(U) \\ &\in \mathcal{E} \end{aligned}$$

So that $\mathcal{T} \subset \mathcal{E}$. Hence $\mathcal{E} = \mathcal{T}$ and

$$\begin{aligned} \tau_X(\mathcal{F}) &= \tau_X(\mathcal{E}) \\ &= \tau_X(\mathcal{T}) \\ &= \mathcal{T} \end{aligned}$$

$\square$

Definition 4.2.0.16. Let $(X_\alpha, \mathcal{T}_\alpha)_{\alpha \in A}$ be a collection of topological spaces, Y a set and $\mathcal{F} \in \prod_{\alpha \in A} Y^{X_\alpha}$ (i.e. $\mathcal{F} = (f_\alpha)_{\alpha \in A}$ where for each $\alpha \in A$, $f_\alpha : X_\alpha \rightarrow Y$). We define the **final topology on Y generated by $\mathcal{F}$** , denoted $\tau_Y(\mathcal{F})$, by

$$\tau_Y(\mathcal{F}) := \tau_Y(\{V \subset Y : \text{for each } \alpha \in A, f_\alpha^{-1}(V) \in \mathcal{T}_\alpha\})$$

Note 4.2.0.17. If $\mathcal{F} = \{f\}$, then $\tau_Y(\mathcal{F}) = f_*\mathcal{T}$.

Exercise 4.2.0.18. Let $(X_\alpha, \mathcal{T}_\alpha)_{\alpha \in A}$ be a collection of topological spaces, Y a set and $\mathcal{F} \in \prod_{\alpha \in A} Y^{X_\alpha}$ (i.e. $\mathcal{F} = (f_\alpha)_{\alpha \in A}$ where for each $\alpha \in A$, $f_\alpha : X_\alpha \rightarrow Y$). Then for each $\mathcal{T} \subset \mathcal{P}(Y)$ if $\mathcal{T}$ is a topology on Y and for each $\alpha \in A$, f_α is $(\mathcal{T}_\alpha, \mathcal{T})$ -continuous, then $\mathcal{T} \subset \tau_Y(\mathcal{F})$.

Proof. Let $\mathcal{T} \subset \mathcal{P}(Y)$. Suppose that $\mathcal{T}$ is a topology on Y and for each $\alpha \in A$, f_α is $(\mathcal{T}_\alpha, \mathcal{T})$ -continuous. Set $\mathcal{V} := \{V \subset Y : \text{for each } \alpha \in A, f_\alpha^{-1}(V) \in \mathcal{T}_\alpha\}$. By definition, $\tau_Y(\mathcal{F}) = \tau_Y(\mathcal{V})$. Let $V \in \mathcal{T}$. By assumption, for each $\alpha \in A$, f_α is $(\mathcal{T}_\alpha, \mathcal{T})$ -measurable. Thus for each $\alpha \in A$, $f_\alpha^{-1}(V) \in \mathcal{T}_\alpha$. Therefore $V \in \mathcal{V}$. Since $V \in \mathcal{T}$ is arbitrary, we have that

$$\begin{aligned} \mathcal{T} &\subset \mathcal{V} \\ &\subset \tau_Y(\mathcal{V}) \\ &= \tau_Y(\mathcal{F}). \end{aligned}$$

□

Note 4.2.0.19. Essentially, $\tau_X(\mathcal{F})$ is the largest topology on X such that for each $\alpha \in A$, $f_\alpha : X_\alpha \rightarrow Y$ is continuous.

Exercise 4.2.0.20. Let $(X_\alpha, \mathcal{T}_\alpha)_{\alpha \in A}$ be a collection of topological spaces, Y a set, $(Z, \mathcal{C})$ a topological space, $\mathcal{F} = (f_\alpha)_{\alpha \in A} \in \prod_{\alpha \in A} Y^{X_\alpha}$ and $g : Y \rightarrow Z$. Then g is $(\tau_Y(\mathcal{F}), \mathcal{C})$ -continuous iff for each $\alpha \in A$, $g \circ f_\alpha$ is $(\mathcal{T}_\alpha, \mathcal{C})$ -continuous:

$$\begin{array}{ccc} X_\alpha & \xrightarrow{f_\alpha} & Y \\ & \searrow g \circ f_\alpha & \downarrow g \\ & & Z \end{array}$$

Proof. If g is $(\tau_Y(\mathcal{F}), \mathcal{C})$ -continuous, then clearly for each $\alpha \in A$, $g \circ f_\alpha$ is $(\mathcal{T}_\alpha, \mathcal{C})$ -continuous.

Conversely, suppose that for each $\alpha \in A$, $g \circ f_\alpha$ is $(\mathcal{T}_\alpha, \mathcal{C})$ -continuous. Let $\alpha \in A$ and $V \in \mathcal{C}$. Continuity implies that

$$\begin{aligned} f_\alpha^{-1}(g^{-1}(V)) &= (g \circ f_\alpha)^{-1}(V) \\ &\in \mathcal{T}_\alpha \end{aligned}$$

Since $\alpha \in A$ is arbitrary, we have that by definition, $g^{-1}(V) \in \tau_Y(\mathcal{F})$. Since $V \in \mathcal{C}$ is arbitrary, g is $(\tau_Y(\mathcal{F}), \mathcal{C})$ -continuous. □

Definition 4.2.0.21. Let $(X, \mathcal{T}_X)$ and $(Y, \mathcal{T}_Y)$ be topological spaces and $f : X \rightarrow Y$. Then

1. f is said to be **open** if for each $A \in \mathcal{T}_X$, $f(A) \in \mathcal{T}_Y$.
2. f is said to be **closed** if for each $A \subset X$, if $A^c \in \mathcal{T}_X$, then $f(A)^c \in \mathcal{T}_Y$.

Exercise 4.2.0.22. Let $(X, \mathcal{T})$, $(Y, \mathcal{T}_Y)$ be topological spaces, $\mathcal{B} \subset \mathcal{T}$ a basis for $\mathcal{T}$ and $f : X \rightarrow Y$. Then f is open iff for each $U \in \mathcal{B}$, $f(U) \in \mathcal{T}_Y$.

Hint: $f\left(\bigcup_{\alpha \in A} A_\alpha\right) = \bigcup_{\alpha \in A} f(A_\alpha)$.

Proof. Clearly if f is open, then for each $U \in \mathcal{B}$, $f(U) \in \mathcal{T}_Y$.

Conversely, suppose that for each $U \in \mathcal{B}$, $f(U) \in \mathcal{T}_Y$. Let $U \in \mathcal{T}$. Then there exists $(U_\alpha)_{\alpha \in A} \subset \mathcal{B}$ such that $U = \bigcup_{\alpha \in A} U_\alpha$. Then

$$\begin{aligned} f(U) &= \bigcup_{\alpha \in A} f(U_\alpha) \\ &\in \mathcal{T}_Y \end{aligned}$$

Since $U \in \mathcal{T}$ is arbitrary, f is open. □

Exercise 4.2.0.23. Let $(X, \mathcal{T}_X)$, $(Y, \mathcal{T}_Y)$ be topological spaces, $f : X \rightarrow Y$ and $\mathcal{B}_X$ a basis for $\mathcal{T}_X$. Suppose that f is surjective, $(\mathcal{T}_X, \mathcal{T}_Y)$ -continuous and $(\mathcal{T}_X, \mathcal{T}_Y)$ -open. Then $\{f(A) : A \in \mathcal{B}_X\}$ is a basis for $\mathcal{T}_Y$.

Proof. Set $\mathcal{B}_Y := \{f(A) : A \in \mathcal{B}_X\}$. Since f is open, $\mathcal{B}_Y \subset \mathcal{T}_Y$. Let $V \in \mathcal{T}_Y$. Set $U = f^{-1}(V)$. Since f is continuous, $U \in \mathcal{T}_X$. Since $\mathcal{B}_X$ is a basis for $\mathcal{T}_X$, there exist $\mathcal{B}'_X \subset \mathcal{B}_X$ such that $U = \bigcup_{A \in \mathcal{B}'_X} A$. Define $\mathcal{B}'_Y \subset \mathcal{B}_Y$ by $\mathcal{B}'_Y = \{f(A) : A \in \mathcal{B}'_X\}$. Since f is surjective, we have that $f(f^{-1}(V)) = f(V)$ and therefore

$$\begin{aligned} V &= f(f^{-1}(V)) \\ &= f(U) \\ &= f\left(\bigcup_{A \in \mathcal{B}'_X} A\right) \\ &= \bigcup_{A \in \mathcal{B}'_X} f(A) \\ &= \bigcup_{B \in \mathcal{B}'_Y} B \end{aligned}$$

Since $V \in \mathcal{T}_Y$ is arbitrary, we have that for each $V \in \mathcal{T}_Y$, there exists $\mathcal{B}'_Y \subset \mathcal{B}_Y$ such that $V = \bigcup_{B \in \mathcal{B}'_Y} B$. Thus $\mathcal{B}_Y$ is a basis for $\mathcal{T}_Y$. $\square$

Exercise 4.2.0.24. Doob-Dynkin Lemma:

Let $(X_1, \mathcal{T}_1)$, $(X_2, \mathcal{T}_2)$ and $(X_3, \mathcal{T}_3)$ be topological spaces and $f : X_1 \rightarrow X_2$ and $g : X_1 \rightarrow X_3$. Suppose that f is surjective and $\mathcal{T}_1$ - $\mathcal{T}_2$ continuous and g is $\mathcal{T}_1$ - $\mathcal{T}_3$ continuous and $(X_3, \mathcal{T}_3)$ is a T_1 space. Then g is $f^*\mathcal{T}_2$ - $\mathcal{T}_3$ continuous iff there exists a unique $\phi : X_2 \rightarrow X_3$ such that ϕ is $\mathcal{T}_2$ - $\mathcal{T}_3$ continuous and $g = \phi \circ f$.

Hint: For each $t \in X_3$, set $A_t = g^{-1}(\{t\}) \in \mathcal{F}_{(f^*\mathcal{T}_2)}$ and choose $B_t \in \mathcal{T}_2$ such that $A_t = f^{-1}(B_t)$. Set $\phi(y) = t$ for $y \in B_t \cap f(X_1)$ and $t \in g(X_1)$.

Proof. Suppose that there exists a unique $\phi : X_2 \rightarrow X_3$ such that ϕ is $\mathcal{T}_2$ - $\mathcal{T}_3$ measurable and $g = \phi \circ f$. Since f is $f^*\mathcal{T}_2$ - $\mathcal{T}_2$ continuous, we have that $g = \phi \circ f$ is $f^*\mathcal{T}_2$ - $\mathcal{T}_3$ continuous.

Conversely, suppose that g is $f^*\mathcal{T}_2$ - $\mathcal{T}_3$ continuous.

• **(Existence)**

For each $t \in X_3$, set $A_t = g^{-1}(\{t\})$. Since $(X_3, \mathcal{T}_3)$ is a T_1 space, for each $t \in X_3$, $A_t \in \mathcal{F}_{f^*\mathcal{T}_2}$ and thus, there exists $B_t \in \mathcal{F}_{\mathcal{T}_2}$ such that $A_t = f^{-1}(B_t)$.

Note that

- for each $t \in g(X_1)$, there exists $x \in A_t$ such that $g(x) = t$. Hence $f(x) \in B_t$.
- for $t_1, t_2 \in g(X_1)$, $t_1 \neq t_2$ implies that

$$\begin{aligned} f^{-1}(B_{t_1} \cap B_{t_2}) &= A_{t_1} \cap A_{t_2} \\ &= g^{-1}(\{t_1\} \cap \{t_2\}) \\ &= \emptyset \end{aligned}$$

and since f is surjective,

$$\begin{aligned} B_{t_1} \cap B_{t_2} &= f(f^{-1}(B_{t_1} \cap B_{t_2})) \\ &= f(\emptyset) \\ &= \emptyset \end{aligned}$$

– we have that

$$\begin{aligned}
 f^{-1}\left(\bigcup_{t \in g(X_1)} B_t\right) &= \bigcup_{t \in g(X_1)} A_t \\
 &= \bigcup_{t \in g(X_1)} g^{-1}(\{t\}) \\
 &= g^{-1}(g(X_1)) \\
 &= X_1
 \end{aligned}$$

Since f is surjective, we have that

$$\begin{aligned}
 X_2 &= f(X_1) \\
 &= f\left(f^{-1}\left(\bigcup_{t \in g(X_1)} B_t\right)\right) \\
 &= \bigcup_{t \in g(X_1)} B_t
 \end{aligned}$$

Therefore,

- for each $t \in g(X_1)$, $B_t \neq \emptyset$
- $(A_t)_{t \in g(X_1)}$ is a partition of X_1
- $(B_t)_{t \in g(X_1)}$ is a partition of X_2

Define $\phi : X_2 \rightarrow X_3$ by $\phi(y) = t$ for $t \in g(X_1)$ and $y \in B_t$. Then the previous observations imply that ϕ is well defined and $\phi(X_2) = g(X_1)$. Since for each $t \in g(X_1)$ and $x \in A_t$, $f(x) \in B_t$ and $g(x) = t$, we have that $\phi \circ f(x) = t = g(x)$. So $\phi \circ f = g$.

To show that ϕ is continuous, let $C \in \mathcal{T}_3$. Choose $B \in \mathcal{T}_2$ such that $g^{-1}(C) = f^{-1}(B)$. Let $y \in \phi^{-1}(C) \subset X_2$. Set $t = \phi(y) \in C$ and choose $x \in X_1$ such that $y = f(x)$. Since

$$\begin{aligned}
 g(x) &= \phi \circ f(x) \\
 &= \phi(y) \\
 &= t \\
 &\in C
 \end{aligned}$$

$x \in g^{-1}(C) = f^{-1}(B)$. Therefore, $y = f(x) \in B$. So $\phi^{-1}(C) \subset B$.

Let $y \in B$. Choose $x \in X_1$ such that $f(x) = y$. Then $x \in f^{-1}(B) = g^{-1}(C)$. So

$$\begin{aligned}
 \phi(y) &= \phi \circ f(x) \\
 &= g(x) \\
 &\in C
 \end{aligned}$$

and $y \in \phi^{-1}(C)$. So $B \subset \phi^{-1}(C)$. Hence $\phi^{-1}(C) = B \in \mathcal{T}_2$ and ϕ is $\mathcal{T}_2$ - $\mathcal{T}_3$ continuous.

• **(Uniqueness)**

Let $\psi : X_2 \rightarrow X_3$. Suppose that ψ is $\mathcal{T}_2$ - $\mathcal{T}_3$ continuous and $g = \psi \circ f$. Let $y \in X_2$. Then there exists $x \in X_1$ such that $y = f(x)$. Then

$$\begin{aligned}
 \psi(y) &= \psi \circ f(x) \\
 &= g(x) \\
 &= \phi \circ f(x) \\
 &= \phi(y)
 \end{aligned}$$

So $\psi = \phi$.

□

Exercise 4.2.0.25. Let $(X_1, \mathcal{T}_1)$, $(X_2, \mathcal{T}_2)$ and $(X_3, \mathcal{T}_3)$ be topological spaces and $f : X_1 \rightarrow X_2$ and $g : X_1 \rightarrow X_3$. Suppose that f is $\mathcal{T}_1$ - $\mathcal{T}_2$ continuous and g is $\mathcal{T}_1$ - $\mathcal{T}_3$ continuous and $(X_3, \mathcal{T}_3)$ is a T_1 space. Then g is $f^*\mathcal{T}_2$ - $\mathcal{T}_3$ continuous iff there exists a unique $\phi : f(X_1) \rightarrow X_3$ such that ϕ is $\mathcal{T}_2 \cap f(X_1) - \mathcal{T}_3$ continuous and $g = \phi \circ f$.

Proof. A previous exercise implies that $f : X_1 \rightarrow f(X_1)$ is $\mathcal{T}_1 - \mathcal{T}_2 \cap f(X_1)$ continuous. Now apply the previous exercise. □

Definition 4.2.0.26. Let X be a topological space, $x_0 \in X$ and $f : X \rightarrow \mathbb{R}$. We define the **limit inferior of f as $x \rightarrow x_0$** (resp. **limit inferior of f as $x \rightarrow x_0$**), denoted $\liminf_{x \rightarrow x_0} f(x)$ (resp. $\liminf_{x \rightarrow x_0} f(x)$), by

$$\liminf_{x \rightarrow x_0} f(x) = \sup_{V \in \mathcal{N}(x_0)} \inf_{x \in V \setminus \{x_0\}} f(x)$$

resp.

$$\limsup_{x \rightarrow x_0} f(x) = \inf_{V \in \mathcal{N}(x_0)} \sup_{x \in V \setminus \{x_0\}} f(x)$$

Exercise 4.2.0.27. Let X be a topological space, $x_0 \in X$ and $f : X \rightarrow \mathbb{R}$. Then f is continuous at x_0 iff $\liminf_{x \rightarrow x_0} f(x) = \limsup_{x \rightarrow x_0} f(x) = f(x_0)$

Proof. Suppose that
FINISH!!!

□

4.3 Nets

In this section, we detail a second notion of structure which aligns with our physical notion of a space. Specifically, we discuss the nets and their convergence.

give axiomatization of topology via nets, see 'General Topology' by Kelly

4.3.1 Common Directed Sets

Note 4.3.1.1. We recall the definition of a directed set from Definition 2.2.0.1.

Definition 4.3.1.2. Let X be a set. Define the **reverse inclusion ordering** on $\mathcal{N}(x)$, denoted $\leq$, by $U \leq V$ iff $V \subset U$.

Exercise 4.3.1.3. Let X be a topological space and $x \in X$. Then $\mathcal{N}(x)$ ordered by reverse inclusion is a directed set.

Proof.

1. Clearly, for each $U \in \mathcal{N}(x)$, $U \leq U$.
2. Let $U, V, W \in \mathcal{N}(x)$. Suppose that $U \leq V$ and $V \leq W$. Then $W \subset V \subset U$ which implies that $W \subset U$ and hence $U \leq W$.
3. Let $U, V \in \mathcal{N}(x)$. Set $W = U \cap V$. Then $W \in \mathcal{N}(x)$ and $U, V \leq W$.

So $\mathcal{N}(x)$ is a directed set. □

Definition 4.3.1.4. Let $(A, \leq)$ be a directed set and $\alpha_0 \in A$. We define $[\alpha_0, \infty) \subset A$, by $[\alpha_0, \infty) := \{\alpha \in A : \alpha \geq \alpha_0\}$ and $\leq_{[\alpha_0, \infty)} := \leq \cap ([\alpha_0, \infty) \times [\alpha_0, \infty))$.

Exercise 4.3.1.5. Let $(A, \leq)$ be a directed set and $\alpha_0 \in A$. Then $([\alpha_0, \infty), \leq_{[\alpha_0, \infty)})$ is a directed set.

Proof. Set $B := [\alpha_0, \infty)$.

1. Let $\alpha \in B$. Since $B \subset A$, $\alpha \in A$. Since $(A, \leq)$ is a directed set, $\alpha \leq \alpha$. Since $\alpha \in B$, we have that $\alpha \leq_B \alpha$. Since $\alpha \in B$ is arbitrary, we have that for each $\alpha \in B$, $\alpha \leq_B \alpha$.
2. Let $\alpha, \beta, \gamma \in B$. Suppose that $\alpha \leq_B \beta$ and $\beta \leq \gamma$. Since $\leq_B \subset \leq$, we have that $\alpha \leq \beta$ and $\beta \leq \gamma$. Since $(A, \leq)$ is a directed set, $\alpha \leq \gamma$. Since $\alpha, \gamma \in B$, we have that $\alpha \leq_B \gamma$. Since $\alpha, \beta, \gamma \in B$ are arbitrary, we have that for each $\alpha, \beta, \gamma \in B$, $\alpha \leq_B \beta$ and $\beta \leq_B \gamma$ implies that $\alpha \leq_B \gamma$.
3. Let $\alpha, \beta \in B$. Since $B \subset A$, there exists $\gamma \in A$ such that $\alpha, \beta \leq \gamma$. Since $\alpha \in B$, $\alpha \geq \alpha_0$. Since $\gamma \geq \alpha$, we have that $\gamma \geq \alpha_0$. Hence $\gamma \in B$. Since $\alpha, \beta, \gamma \in B$, we have that $\alpha, \beta \leq_B \gamma$. Since $\alpha, \beta \in B$ are arbitrary, we have that for each $\alpha, \beta \in B$, there exists $\gamma \in B$ such that $\alpha, \beta \leq_B \gamma$.
4. Since $\alpha_0 \in B$, $B \neq \emptyset$.

So $(B, \leq_B)$ is a directed set. □

Definition 4.3.1.6. move to metric space chapter Let X be a metric space and $x_0 \in X$. Define the **reverse distance from x_0 ordering** on $X \setminus \{x_0\}$, denoted $\leq_{x_0}$, by $x \leq_{x_0} y$ iff $d(x, x_0) \geq d(y, x_0)$.

Exercise 4.3.1.7. move to metrix space chapter Let X be a metric space and $x_0 \in X$. Then $(X \setminus \{x_0\}, \leq_{x_0})$ is a directed set.

Proof.

1. Let $x \in X \setminus \{x_0\}$. Since $d(x, x_0) \geq d(x, x_0)$, $x \leq_{x_0} x$.
2. Let $x, y, z \in X \setminus \{x_0\}$. Suppose that $x \leq_{x_0} y$ and $y \leq_{x_0} z$. Then $d(x, x_0) \geq d(y, x_0)$ and $d(y, x_0) \geq d(z, x_0)$. Hence $d(x, x_0) \geq d(z, x_0)$ so that $x \leq z$.
3. Let $x, y \in X \setminus \{x_0\}$. Set

$$\begin{aligned} z &= \arg \min_{a \in \{x, y\}} d(a, x_0) \\ &\in X \setminus \{x_0\} \end{aligned}$$

Then $x, y \leq_{x_0} z$. □

4.3.2 Nets in Topological Spaces

maybe move the basic definition of net to the set theory section and also introduce sequences there

Definition 4.3.2.1. Let X be a topological space, $(x_\alpha)_{\alpha \in A} \subset X$ a net and $U \subset X$. Then $(x_\alpha)_{\alpha \in A}$ is said to be

- **eventually in U** if there exists $\beta \in A$ such that for each $\alpha \in A$ $\alpha \geq \beta$ implies that $x_\alpha \in U$
- **frequently in U** if for each $\alpha \in A$, there exists $\beta \in A$ such that $\beta \geq \alpha$ and $x_\beta \in U$

Exercise 4.3.2.2. Let X be a topological space, $(x_\alpha)_{\alpha \in A} \subset X$ a net and $U \subset X$. Then $(x_\alpha)_{\alpha \in A}$ is eventually in U iff there exists $\alpha_0 \in A$ such that $(x_{\alpha \in [\alpha_0, \infty)})_{\alpha \in A} \subset U$.

Proof.

- $(\implies)$:
Suppose that $(x_\alpha)_{\alpha \in A}$ is eventually in U . Then there exists $\alpha_0 \in A$ such that for each $\alpha \in A$, $\alpha \geq \alpha_0$ implies that $x_\alpha \in U$. Then $(x_\alpha)_{\alpha \in [\alpha_0, \infty)} \subset U$.
- $(\impliedby)$:
Suppose that there exists $\alpha_0 \in A$ such that $(x_\alpha)_{\alpha \in [\alpha_0, \infty)} \subset U$. Then for each $\alpha \in A$, $\alpha \geq \alpha_0$ implies that $x_\alpha \in U$. Hence $(x_\alpha)_{\alpha \in A}$ is eventually in U .

□

Exercise 4.3.2.3. (make exercise about the tail net and being frequently in U)

Proof.

□

Definition 4.3.2.4. Let X be a topological space, $(x_\alpha)_{\alpha \in A} \subset X$ a net and $x \in X$. Then $(x_\alpha)_{\alpha \in A}$ is said to **converge to x** , denoted $x_\alpha \rightarrow x$, if for each $U \in \mathcal{N}(x)$, $(x_\alpha)_{\alpha \in A}$ is eventually in U .

Definition 4.3.2.5. Let X be a topological space and $(x_\alpha)_{\alpha \in A} \subset X$ a net. Then $(x_\alpha)_{\alpha \in A}$ is said to **converge** if there exists $x \in X$ such that $x_\alpha \rightarrow x$.

Exercise 4.3.2.6. move or delete? Let X be a metric space and $x_0 \in X$. Set $A = X \setminus \{x_0\}$. Order A by reverse distance from x_0 . Define $(x_\alpha)_{\alpha \in A} \subset X$ by $x_\alpha = \alpha$. Then $x_\alpha \rightarrow x_0$.

Proof. Let $U \in \mathcal{N}(x_0)$. Since $x_0 \in \text{Int } U$, there exists $\delta > 0$ such that $B(x_0, \delta) \subset \text{Int } U$. Choose $\beta \in B^*(x_0, \delta)$. Let $\alpha \in A$. Suppose that $\alpha \geq \beta$. Then $d(\alpha, x_0) \leq d(\beta, x_0) < \delta$. Hence

$$\begin{aligned} x_\alpha &= \alpha \\ &\in B^*(x_0, \delta) \\ &\subset U \end{aligned}$$

Since $U \in \mathcal{N}(x_0)$ is arbitrary, $x_\alpha \rightarrow x_0$

□

Exercise 4.3.2.7. Let X be a topological space, $S \subset X$ and $x \in X$. Then $x \in S'$ iff there exists a net $(x_\alpha)_{\alpha \in A} \subset S \setminus \{x\}$ such that $x_\alpha \rightarrow x$.

Proof. Suppose that $x \in S'$. Set $A = \mathcal{N}(x)$, ordered by reverse inclusion. Since $x \in S'$, for each $\alpha \in A$, there exists $x_\alpha \in (\alpha \setminus \{x\}) \cap S$. Then $(x_\alpha)_{\alpha \in A} \subset S \setminus \{x\}$. Let $V \in \mathcal{N}(x)$. Choose $\beta = V$. Let $\alpha \in \mathcal{N}(x)$. Suppose that $\alpha \geq \beta$. Then

$$\begin{aligned} x_\alpha &\in (\alpha \setminus \{x\}) \cap S \\ &\subset \alpha \\ &\subset \beta \\ &= V \end{aligned}$$

So $(x_\alpha)_{\alpha \in \mathcal{N}(x)}$ is eventually in V . Since $V \in \mathcal{N}(x)$ is arbitrary, $x_\alpha \rightarrow x$.

Conversely, suppose that there exists a net $(x_\alpha)_{\alpha \in A} \subset S \setminus \{x\}$ such that $x_\alpha \rightarrow x$. Let $U \in \mathcal{N}(x)$. Since $(x_\alpha)_{\alpha \in A}$ is eventually in U , there exists $\beta \in A$ such that $x_\beta \in U$. Then $x_\beta \in (U \setminus \{x\}) \cap S$ and $(U \setminus \{x\}) \cap S \neq \emptyset$. Since $U \in \mathcal{N}(x)$ is arbitrary, $x \in S'$. □

Exercise 4.3.2.8. Let X be a topological space, $S \subset X$ and $x \in X$. Then $x \in \text{cl } S$ iff there exists a net $(x_\alpha)_{\alpha \in A} \subset S$ such that $x_\alpha \rightarrow x$.

Proof. Suppose that $x \in \text{cl } S$. Since $\text{cl } S = S \cup S'$, $x \in S$ or $x \in S'$. If $x \in S$, define $(x_n)_{n \in \mathbb{N}} \subset S$ by $x_n = x$. Then $x_n \rightarrow x$. If $x \in S'$, the previous exercise implies that there exists a net $(x_\alpha)_{\alpha \in A} \subset S \setminus \{x\} \subset S$ such that $x_\alpha \rightarrow x$. $\square$

Exercise 4.3.2.9. Let $(X, \mathcal{T})$ be a topological space, $U \in \mathcal{T}$ and $x \in X$. Then $x \in U$ iff $U \not\subset (\text{cl}\{x\})^c$.

Proof.

$$\begin{aligned} U \not\subset (\text{cl}\{x\})^c &\iff \text{there exists } y \in U \text{ such that } y \notin (\text{cl}\{x\})^c \\ &\iff \text{there exists } y \in U \text{ such that } y \in \text{cl}\{x\} \\ &\iff U \cap \text{cl}\{x\} \neq \emptyset. \end{aligned}$$

• ($\implies$):

Suppose that $x \in U$. Then $x \in U \cap \text{cl}\{x\}$. Hence $U \cap \text{cl}\{x\} \neq \emptyset$. Thus $U \not\subset (\text{cl}\{x\})^c$.

• ($\impliedby$):

Suppose that $U \not\subset (\text{cl}\{x\})^c$. Then $U \cap \text{cl}\{x\} \neq \emptyset$. Thus there exists $y \in X$ such that $y \in U \cap \text{cl}\{x\}$. Since $y \in \text{cl}\{x\}$ **a prev ex in nets section** implies that there exists a net $(x_\alpha)_{\alpha \in A} \subset \{x\}$ such that $x_\alpha \rightarrow y$. Since $(x_\alpha)_{\alpha \in A} \subset \{x\}$, we have that for each $\alpha \in A$, $x_\alpha = x$. Since $U \in \mathcal{T}$, $y \in U$ and $x_\alpha \rightarrow y$, we have that $(x_\alpha)_{\alpha \in A}$ is eventually in U . Thus there exists $\alpha_0 \in A$ such that for each $\alpha \in A$, $\alpha \geq \alpha_0$ implies that $x_\alpha \in U$. In particular,

$$\begin{aligned} x &= x_{\alpha_0} \\ &\in U. \end{aligned}$$

$\square$

Exercise 4.3.2.10. Let X be a topological space and $E \subset X$. Then

1. $\partial E = \text{cl } E \cap \text{cl } E^c$
2. $\partial E = \text{cl } E \setminus \text{Int } E$

Proof.

1. Let $x \in \partial E$. Then for each $U \in \mathcal{N}(x)$, $U \cap E \neq \emptyset$ and $U \cap E^c \neq \emptyset$. The axiom of choice implies that there exist nets $(a_U)_{U \in \mathcal{N}(x)} \subset E$ and $(b_U)_{U \in \mathcal{N}(x)} \subset E^c$ such that for each $U \in \mathcal{N}(x)$, $a_U, b_U \in U$. Then $a_U, b_U \rightarrow x$. Hence $x \in (\text{cl } E) \cap (\text{cl } E^c)$. Since $x \in \partial E$ is arbitrary, we have that $\partial E \subset (\text{cl } E) \cap (\text{cl } E^c)$.

Conversely, let $x \in (\text{cl } E) \cap (\text{cl } E^c)$. Then there exists $(a_\alpha)_{\alpha \in A} \subset E$ and $(b_\beta)_{\beta \in B} \subset E^c$ such that $a_\alpha \rightarrow x$ and $b_\beta \rightarrow x$. Let $U \in \mathcal{N}(x)$. Then there exists $\alpha_0 \in A$ and $\beta_0 \in B$ such that for each $\alpha \in A$ and $\beta \in B$, $\alpha \geq \alpha_0$ implies that $a_\alpha \in U$ and $\beta \geq \beta_0$ implies that $b_\beta \in U$. In particular, $a_{\alpha_0} \in U \cap E$ and $b_{\beta_0} \in U \cap E^c$. Hence $U \cap E \neq \emptyset$ and $U \cap E^c \neq \emptyset$. Since $U \in \mathcal{N}(x)$ is arbitrary, we have that for each $U \in \mathcal{N}(x)$, $U \cap E \neq \emptyset$ and $U \cap E^c \neq \emptyset$. Thus $x \in \partial E$. Since $x \in (\text{cl } E) \cap (\text{cl } E^c)$ is arbitrary, $(\text{cl } E) \cap (\text{cl } E^c) \subset \partial E$.

Therefore $\partial E = (\text{cl } E) \cap (\text{cl } E^c)$.

2. Exercise 4.1.0.27 and part (1) implies that

$$\begin{aligned} \partial E &= (\text{cl } E) \cap (\text{cl } E^c) \\ &= (\text{cl } E) \cap (\text{Int } E)^c \\ &= (\text{cl } E) \setminus \text{Int } E \end{aligned}$$

$\square$

Exercise 4.3.2.11. Topology in Terms of Nets:

Let X be a topological space and $U \subset X$. Then U is open iff for each net $(x_\alpha)_{\alpha \in A} \subset X$ and $x \in U$, $x_\alpha \rightarrow x$ implies that $(x_\alpha)_{\alpha \in A}$ is eventually in U .

Proof. Suppose that U is open. Let $(x_\alpha)_{\alpha \in A} \subset X$ be a net and $x \in U$. Suppose that $x_\alpha \rightarrow x$. Since $U \in \mathcal{N}(x)$, $(x_\alpha)_{\alpha \in A}$ is eventually in U .

Conversely, suppose that for each net $(x_\alpha)_{\alpha \in A} \subset X$ and $x \in U$, $x_\alpha \rightarrow x$ implies that $(x_\alpha)_{\alpha \in A}$ is eventually in U . For the sake of contradiction, suppose that U^c is not closed. Then there exists $x \in \text{cl } U^c$ such that $x \notin U^c$. Thus $x \in U$. Since $x \in \text{cl } U^c$, a previous exercise implies that there exists a net $(x_\alpha)_{\alpha \in A} \subset U^c$ such that $x_\alpha \rightarrow x$. By assumption, $(x_\alpha)_{\alpha \in A}$ is eventually in U . This is a contradiction since $(x_\alpha)_{\alpha \in A} \subset U^c$. Hence U^c is closed and hence U is open. $\square$

Exercise 4.3.2.12. Let X be a topological space, $U \in \mathcal{T}$ and $E \subset X$. If $U \cap \text{cl } E \neq \emptyset$, then $U \cap E \neq \emptyset$.

Proof. Suppose that $U \cap \text{cl } E \neq \emptyset$. Then there exists $x \in X$ such that $x \in U \cap \text{cl } E$. Since $x \in \text{cl } E$, there exists a net $(x_\alpha)_{\alpha \in A} \subset E$ such that $x_\alpha \rightarrow x$. Since $U \in \mathcal{N}(x)$, $(x_\alpha)_{\alpha \in A}$ is eventually in U . Thus there exists $\alpha_0 \in A$ such that for each $\alpha \geq \alpha_0$, $x_\alpha \in U$. In particular $x_{\alpha_0} \in U \cap E$. Hence $U \cap E \neq \emptyset$. $\square$

Exercise 4.3.2.13. Let $(X, \mathcal{T}_X)$ and $(Y, \mathcal{T}_Y)$ be topological spaces, $f : X \rightarrow Y$ and $x \in X$. Then f is continuous at x iff for each net $(x_\alpha)_{\alpha \in A} \subset X$, $x_\alpha \rightarrow x$ implies that $f(x_\alpha) \rightarrow f(x)$.

Proof.

• ($\Rightarrow$):

Suppose that f is continuous at x . Let $(x_\alpha)_{\alpha \in A} \subset X$ be a net. Suppose that $x_\alpha \rightarrow x$. Let $V \in \mathcal{N}(f(x))$. Continuity implies that $f^{-1}(V) \in \mathcal{N}(x)$. Since $x_\alpha \rightarrow x$, $(x_\alpha)_{\alpha \in A}$ is eventually in $f^{-1}(V)$. So there exists $\beta \in A$ such that for each $\alpha \in A$, $\alpha \geq \beta$ implies that $x_\alpha \in f^{-1}(V)$. Let $\alpha \in A$. Suppose that $\alpha \geq \beta$. Then $f(x_\alpha) \in V$. So $(f(x_\alpha))_{\alpha \in A}$ is eventually in V . Since $V \in \mathcal{N}(f(x))$ is arbitrary, $f(x_\alpha) \rightarrow f(x)$.

• ($\Leftarrow$):

Conversely, suppose that f is not continuous at x . Then there exists $V \in \mathcal{N}(f(x))$ such that $f^{-1}(V) \notin \mathcal{N}(x)$. Then $x \notin \text{Int}(f^{-1}(V))$. So $x \in (\text{Int}(f^{-1}(V)))^c = \text{cl } f^{-1}(V^c)$. This implies that there exists a net $(x_\alpha)_{\alpha \in A} \subset f^{-1}(V^c)$ such that $x_\alpha \rightarrow x$. Since for each $\alpha \in A$, $f(x_\alpha) \in V^c$, $f(x_\alpha)$ is not eventually in V . So $f(x_\alpha) \not\rightarrow f(x)$. $\square$

HERERERERER

Exercise 4.3.2.14. Let $(X, \mathcal{T}_X)$ and $(Y, \mathcal{T}_Y)$ be topological spaces and $f : X \rightarrow Y$. Then f is continuous iff for each net $(x_\alpha)_{\alpha \in A} \subset X$ and $x \in X$, $x_\alpha \rightarrow x$ implies that $f(x_\alpha) \rightarrow f(x)$.

Proof. Clear by prev ex (fill out). $\square$

Exercise 4.3.2.15. Let $(Y_\alpha, \mathcal{T}_\alpha)_{\alpha \in A}$ be a collection of topological spaces, X a set and $\mathcal{F} \in \prod_{\alpha \in A} Y_\alpha^X$ with $\mathcal{F} = (f_\alpha)_{\alpha \in A}$. Equip X with $\tau_X(\mathcal{F})$. Let $(x_\gamma)_{\gamma \in \Gamma} \subset X$ be a net and $x \in X$. Then $x_\gamma \rightarrow x$ iff for each $\alpha \in A$, $f_\alpha(x_\gamma) \rightarrow f_\alpha(x)$. (maybe reword without $\mathcal{F}$ and similar instances elsewhere)

Proof. Suppose that $x_\gamma \rightarrow x$. Let $\alpha \in A$. Since f_α is continuous, the previous exercise implies that $f_\alpha(x_\gamma) \rightarrow f_\alpha(x)$. Conversely, Suppose that for each $\alpha \in A$, $f_\alpha(x_\gamma) \rightarrow f_\alpha(x)$. Let $U \in \mathcal{N}(x)$. Since $\text{Int } U \in \tau_X(\mathcal{F})$, Exercise 4.1.0.13 implies there exist $V_1 \in \mathcal{T}_{\alpha_1}, \dots, V_n \in \mathcal{T}_{\alpha_n}$ such that $\bigcap_{j=1}^n f_{\alpha_j}^{-1}(V_j) \subset \text{Int } U$ and $x \in \bigcap_{j=1}^n f_{\alpha_j}^{-1}(V_j)$. Let $j \in \{1, \dots, n\}$. Since $f_{\alpha_j}^{-1}(V_j) \in \mathcal{N}(x)$, $V_j \in \mathcal{N}(f(x))$. By assumption, $f_{\alpha_j}(x_\gamma)$ is eventually in V_j . Thus there exist $\gamma'_j \in \Gamma$ such that for each $\gamma \geq \gamma'_j$, $f_{\alpha_j}(x_\gamma) \in V_j$, or equivalently, $x_\gamma \in f_{\alpha_j}^{-1}(V_j)$. Since Γ is directed, there exists $\gamma' \in \Gamma$ such that for each $j \in \{1, \dots, n\}$, $\gamma' \geq \gamma'_j$. Let $\gamma \in \Gamma$. Suppose that $\gamma \geq \gamma'$. Then

$$\begin{aligned} x_\gamma &\in \bigcap_{j=1}^n f_{\alpha_j}^{-1}(V_j) \\ &\subset \text{Int } U \\ &\subset U \end{aligned}$$

So $(x_\gamma)_{\gamma \in \Gamma}$ is eventually in U . Since $U \in \mathcal{N}(x)$ is arbitrary, $x_\gamma \rightarrow x$. $\square$

Exercise 4.3.2.16. reorganize to an iff

Let X be a set and $\mathcal{T}_1, \mathcal{T}_2$ topologies on X . Then the following are equivalent:

1. $\mathcal{T}_1 = \mathcal{T}_2$
2. for each net $(x_\alpha)_{\alpha \in A} \subset X$ and $x \in X$, $x_\alpha \rightarrow x$ in $\mathcal{T}_1$ iff $x_\alpha \rightarrow x$ in $\mathcal{T}_2$.

Proof.

- (1) $\implies$ (2):
Clear.

- (2) $\implies$ (1):

Let $U \in \mathcal{T}_1$ and $x \in U^c$. Since U^c is closed in $\mathcal{T}_1$, there exists a net $(x_\alpha)_{\alpha \in A} \subset U^c$ such that $x_\alpha \rightarrow x$ in $\mathcal{T}_1$. By assumption, $x_\alpha \rightarrow x$ in $\mathcal{T}_2$. So U^c is closed in $\mathcal{T}_2$ and $U \in \mathcal{T}_2$. Hence $\mathcal{T}_1 \subset \mathcal{T}_2$.

Similarly, $\mathcal{T}_2 \subset \mathcal{T}_1$.

□

Exercise 4.3.2.17. Let X, Y be topological spaces and $\phi : X \rightarrow Y$ a homeomorphism. Then for each $E \subset X$,

1. $\text{cl } \phi(E) = \phi(\text{cl } E)$
2. $\text{Int } \phi(E) = \phi(\text{Int } E)$

Proof.

1. Let $E \subset X$. Since $E \subset \text{cl } E$, we have that $\phi(E) \subset \phi(\text{cl } E)$. Since $\text{cl } E$ is closed, $\phi(\text{cl } E)$ is closed and thus $\text{cl } \phi(E) \subset \phi(\text{cl } E)$. Conversely, let $x \in \phi(\text{cl } E)$. Then $\phi^{-1}(x) \in \text{cl } E$. Then there exists a net $(y_\alpha)_{\alpha \in A} \subset E$ such that $y_\alpha \rightarrow \phi^{-1}(x)$. Then $(\phi(y_\alpha))_{\alpha \in A} \subset \phi(E)$ and $\phi(y_\alpha) \rightarrow x$. Thus $x \in \text{cl } \phi(E)$ and $\phi(\text{cl } E) \subset \text{cl } \phi(E)$.
2. Similar

□

Definition 4.3.2.18. Let X be a topological space, $(x_\alpha)_{\alpha \in A} \subset X$ a net and $x \in X$. Then x is said to be a **cluster point** or **accumulation point** of $(x_\alpha)_{\alpha \in A}$ if for each $U \in \mathcal{N}(x)$, $(x_\alpha)_{\alpha \in A}$ is frequently in U .

Note 4.3.2.19. We recall the definition of a subnet from Definition 1.3.3.3

Exercise 4.3.2.20. Let X be a topological space, $(x_\alpha)_{\alpha \in A} \subset X$ a net and $x \in X$. Then the following are equivalent:

1. x is a cluster point of $(x_\alpha)_{\alpha \in A}$
2. there exists a subnet $(x_{\alpha_\beta})_{\beta \in B}$ of $(x_\alpha)_{\alpha \in A}$ such that $x_{\alpha_\beta} \rightarrow x$
3. $x \in \bigcap_{\alpha \in A} \text{cl}\{x_\beta : \beta \geq \alpha\}$

Hint: Order $\mathcal{N}(x)$ by reverse inclusion and consider the product directed set $B = A \times \mathcal{N}(x)$. If x is a cluster point of $(x_\alpha)_{\alpha \in A}$, then for each $\beta = (\gamma, U) \in B$, there exists $\alpha_\beta \in A$ such that $\alpha_\beta \geq \gamma$ and $\alpha_\beta \in U$.

Proof.

- (1) $\implies$ (2):

Suppose that x is a cluster point of $(x_\alpha)_{\alpha \in A}$. Set $B = A \times \mathcal{N}(x)$. Since x is a cluster point of $(x_\alpha)_{\alpha \in A}$, for each $(\gamma, U) \in B$, there exists $\alpha_{(\gamma, U)} \in A$ such that $\alpha_{(\gamma, U)} \geq \gamma$ and $x_{\alpha_{(\gamma, U)}} \in U$. Let $\alpha_0 \in A$. Choose $\beta_0 = (\alpha_0, X) \in B$. Let $\beta = (\gamma, U) \in B$. Suppose that $\beta \geq \beta_0$. Then $\gamma \geq \alpha_0$ and

$$\begin{aligned} \alpha_\beta &= \alpha_{(\gamma, U)} \\ &\geq \gamma \\ &\geq \alpha_0 \end{aligned}$$

So that $(x_{\alpha_\beta})_{\beta \in B}$ is a subnet of $(x_\alpha)_{\alpha \in A}$. Let $U_0 \in \mathcal{N}(x)$. Choose $\alpha_0 \in A$ and set $\beta_0 = (\alpha_0, U_0)$. Let $\beta = (\gamma, U) \in B$. Suppose that $\beta \geq \beta_0$. Then

$$\begin{aligned} x_{\alpha_\beta} &= x_{\alpha_{(\gamma, U)}} \\ &\in U \\ &\subset U_0 \end{aligned}$$

Since $U_0 \in \mathcal{N}(x)$ is arbitrary, $x_{\alpha_\beta} \rightarrow x$.

• (2) $\implies$ (3):

Suppose that there exists a subnet $(x_{\alpha_\beta})_{\beta \in B}$ of $(x_\alpha)_{\alpha \in A}$ such that $x_{\alpha_\beta} \rightarrow x$. Let $\alpha \in A$. Then there exists $\beta_0 \in B$ such that for each $\beta \in B$, $\beta \geq \beta_0$ implies that $\alpha_\beta \geq \alpha$. Therefore, for each $\beta \in B$, $\beta \geq \beta_0$ implies that $x_{\alpha_\beta} \in E_\alpha$. So $(x_{\alpha_\beta})_{\beta \in B}$ is eventually in E_α . Since $x_{\alpha_\beta} \rightarrow x$ and $(x_{\alpha_\beta})_{\beta \in B}$ is eventually in E_α , Exercise 4.3.2.8 implies that $x \in \text{cl } E_\alpha$. Since $\alpha \in A$ is arbitrary, we have that $x \in \bigcap_{\alpha \in A} \text{cl } E_\alpha$.

• (3) $\implies$ (1):

Suppose that $x \in \bigcap_{\alpha \in A} \text{cl } E_\alpha$. Let $U \in \mathcal{N}(x)$. Since

$$\begin{aligned} x &\in [\text{Int } U] \cap \bigcap_{\alpha \in A} \text{cl } E_\alpha \\ &= \bigcap_{\alpha \in A} ([\text{Int } U] \cap \text{cl } E_\alpha) \end{aligned}$$

we have that for each $\alpha \in A$, $[\text{Int } U] \cap \text{cl } E_\alpha \neq \emptyset$. Exercise 4.3.2.12 implies that for each $\alpha \in A$,

$$\begin{aligned} \emptyset &\neq [\text{Int } U] \cap E_\alpha \\ &\subset U \cap E_\alpha \end{aligned}$$

Let $\alpha \in A$. Since $U \cap E_\alpha \neq \emptyset$, there exists $x_0 \in X$ such that $x_0 \in U \cap E_\alpha$. Since $x_0 \in E_\alpha$, there exists $\alpha_0 \in A$ such that $\alpha_0 \geq \alpha$ and

$$\begin{aligned} x_{\alpha_0} &= x_0 \\ &\in U \end{aligned}$$

Thus $(x_\alpha)_{\alpha \in A}$ is frequently in U . Since $U \in \mathcal{N}(x)$ is arbitrary, we have that for each $U \in \mathcal{N}(x)$, $(x_\alpha)_{\alpha \in A}$ is frequently in U . Thus x is a cluster point of $(x_\alpha)_{\alpha \in A}$. □

Exercise 4.3.2.21. Let X be a topological space, $(x_\alpha)_{\alpha \in A} \subset X$ a net and $x \in X$. If $x_\alpha \rightarrow x$, then for each subnet $(x_{\alpha_\beta})_{\beta \in B}$ of $(x_\alpha)_{\alpha \in A}$, $x_{\alpha_\beta} \rightarrow x$.

Proof. Suppose that $x_\alpha \rightarrow x$. Let $(x_{\alpha_\beta})_{\beta \in B}$ be a subnet of $(x_\alpha)_{\alpha \in A}$ and $U \in \mathcal{N}(x)$. Since $x_\alpha \rightarrow x$, there exists $\alpha_0 \in A$ such that for each $\alpha \geq \alpha_0$, $x_\alpha \in U$. Since $(x_{\alpha_\beta})_{\beta \in B}$ is a subnet of $(x_\alpha)_{\alpha \in A}$, there exists $\beta_0 \in B$ such that for each $\beta \in B$, $\beta \geq \beta_0$ implies that $\alpha_{\beta_e} \geq \alpha_0$. Then for each $\beta \in B$, $\beta \geq \beta_0$ implies that $x_{\alpha_\beta} \in U$. Since $U \in \mathcal{N}(x)$ is arbitrary, $x_{\alpha_\beta} \rightarrow x$. □

Exercise 4.3.2.22. Let X be a topological space, $(x_\alpha)_{\alpha \in A} \subset X$ a net and $x \in X$. Then $x_\alpha \rightarrow x$ iff for each subnet $(x_{\alpha_\beta})_{\beta \in B}$ of $(x_\alpha)_{\alpha \in A}$, there exists a subnet $(x_{\alpha_{\beta_\gamma}})_{\gamma \in \Gamma}$ of $(x_{\alpha_\beta})_{\beta \in B}$ such that $x_{\alpha_{\beta_\gamma}} \rightarrow x$.

Proof. **FINISH!!!** □

Definition 4.3.2.23. Let $(x_\alpha)_{\alpha \in A} \subset \mathbb{R}$ a net.

- We define the **limit inferior** of $(x_\alpha)_{\alpha \in A}$, denoted $\liminf_{\alpha \in A} x_\alpha \in [0, \infty]$, by

$$\liminf_{\alpha \in A} x_\alpha = \sup_{\beta \in A} \left[\inf_{\alpha \geq \beta} x_\alpha \right]$$

- We define the **limit superior** of $(x_\alpha)_{\alpha \in A}$, denoted $\limsup_{\alpha \in A} x_\alpha \in [0, \infty]$, by

$$\limsup_{\alpha \in A} x_\alpha = \inf_{\beta \in A} \left[\sup_{\alpha \geq \beta} x_\alpha \right]$$

Exercise 4.3.2.24. Let $(x_\alpha)_{\alpha \in A} \subset \mathbb{R}$ a net. Then

$$\liminf_{\alpha \in A} x_\alpha \leq \limsup_{\alpha \in A} x_\alpha$$

Proof. Set $s := \liminf_{\alpha \in A} x_\alpha$ and $S := \limsup_{\alpha \in A} x_\alpha$. Let $\epsilon > 0$. Then there exists $\beta_1, \beta_2 \in A$ such that for each $\alpha \in A$, $\alpha \geq \beta_1$ implies that $\sup_{\alpha \geq \beta_1} x_\alpha < S + \epsilon/2$ and $\inf_{\alpha \geq \beta_2} x_\alpha > s - \epsilon/2$. Set $\beta_0 := \max(\beta_1, \beta_2)$. Then

$$\begin{aligned} s - \frac{\epsilon}{2} &< \inf_{\alpha \geq \beta_2} x_\alpha \\ &\leq \inf_{\alpha \geq \beta_0} x_\alpha \\ &\leq \sup_{\alpha \geq \beta_0} x_\alpha \\ &\leq \sup_{\alpha \geq \beta_1} x_\alpha \\ &< S + \frac{\epsilon}{2} \end{aligned}$$

Therefore $-\epsilon < S - s$. Since $\epsilon > 0$ is arbitrary, we have that $0 \leq S - s$. Hence

$$\begin{aligned} \liminf_{\alpha \in A} x_\alpha &= s \\ &\leq S \\ &= \limsup_{\alpha \in A} x_\alpha \end{aligned}$$

□

Exercise 4.3.2.25. Let $(x_\alpha)_{\alpha \in A} \subset \mathbb{R}$ a net and $x \in \mathbb{R}$. Then $x_\alpha \rightarrow x$ iff

$$\liminf_{\alpha \in A} x_\alpha = \limsup_{\alpha \in A} x_\alpha = x$$

Proof. Suppose that $x_\alpha \rightarrow x$. Let $\epsilon > 0$. Then there exist $\beta \in A$ such that for each $\alpha \in A$, $\alpha \geq \beta$ implies that $x_\alpha \in B(x, \epsilon)$. So $\inf_{\alpha \geq \beta} x_\alpha \geq x - \epsilon$ and $\sup_{\alpha \geq \beta} x_\alpha \leq x + \epsilon$. Therefore

$$\begin{aligned} \liminf_{\alpha \in A} x_\alpha &= \sup_{\beta \in A} \left[\inf_{\alpha \geq \beta} x_\alpha \right] \\ &\geq x - \epsilon \end{aligned}$$

and

$$\begin{aligned} \limsup_{\alpha \in A} x_\alpha &= \inf_{\beta \in A} \left[\sup_{\alpha \geq \beta} x_\alpha \right] \\ &\leq x + \epsilon \end{aligned}$$

Since $\epsilon > 0$ is arbitrary,

$$\limsup_{\alpha \in A} x_\alpha \leq x \leq \liminf_{\alpha \in A} x_\alpha$$

Since $\liminf_{\alpha \in A} x_\alpha \leq \limsup_{\alpha \in A} x_\alpha$, we have that $\liminf_{\alpha \in A} x_\alpha = \limsup_{\alpha \in A} x_\alpha = x$.

□

Exercise 4.3.2.26. Let $(x_\alpha)_{\alpha \in A} \subset \mathbb{R}$ a net and $x \in \mathbb{R}$. Then

1. $\liminf_{\alpha \in A} -x_\alpha = -\limsup_{\alpha \in A} x_\alpha$
2. $(x_\alpha)_{\alpha \in A} \subset \mathbb{R} \setminus \{0\}$ implies that $\liminf_{\alpha \in A} x_\alpha^{-1} = \left(\limsup_{\alpha \in A} x_\alpha \right)^{-1}$.
3. generalize to any order reversing bijection of a totally ordered set (including $\overline{\mathbb{R}}$)

Proof.

1. We have that

$$\begin{aligned}
 \liminf_{\alpha \in A} -x_\alpha &= \sup_{\beta \in A} \left[\inf_{\alpha \geq \beta} -x_\alpha \right] \\
 &= \sup_{\beta \in A} \left[-\sup_{\alpha \geq \beta} x_\alpha \right] \\
 &= -\inf_{\alpha \in A} \left[\sup_{\alpha \geq \beta} x_\alpha \right] \\
 &= -\limsup_{\alpha \in A} x_\alpha
 \end{aligned}$$

2. Suppose that $(x_\alpha)_{\alpha \in A} \subset \mathbb{R} \setminus \{0\}$. Then

$$\begin{aligned}
 \liminf_{\alpha \in A} x_\alpha^{-1} &= \sup_{\beta \in A} \left[\inf_{\alpha \geq \beta} x_\alpha^{-1} \right] \\
 &= \sup_{\beta \in A} \left[\sup_{\alpha \geq \beta} x_\alpha \right]^{-1} \\
 &= \left(\inf_{\alpha \in A} \left[\sup_{\alpha \geq \beta} x_\alpha \right] \right)^{-1} \\
 &= \left(\limsup_{\alpha \in A} x_\alpha \right)^{-1}
 \end{aligned}$$

□

4.4 Filters

Note 4.4.0.1. This section assumes familiarity with basic lattice theory [cite lattice chapter in algebra notes](#).

Definition 4.4.0.2. Let d **FINISH!!!**

Definition 4.4.0.3. need to define $\liminf_{x \rightarrow x_0} f(x)$, $\liminf_{\substack{x \rightarrow x_0 \\ x \neq x_0}} f(x)$ and $\lim_{x \rightarrow x_0}$ for functions, might have to check out filters and move to later section [wikipage on limits of functions in terms of filters](#) Note for $\lim_{x \rightarrow x_0} f(x)$, if this limit is assumed so that x_0 does not matter, we can define on equivalence classes of functions defined on punctured neighborhoods of x_0 , see similar idea in appendices of little o notation Definition .2.0.2 **FINISH!!!!**
FINISH!!!

Definition 4.4.0.4. Let X be a topological space, $\mathcal{F}$ a proper filter on X and $x \in X$. Then $\mathcal{F}$ is said to **converge to** x if for each $U \in \mathcal{N}_x$, $U \in \mathcal{F}$.

Definition 4.4.0.5. Let X, Y be a topological spaces, $f : X \rightarrow Y$ and $\mathcal{F}$ a proper filter on X . We define the **pushforward filter on Y induced by $\mathcal{F}$** , denoted $f_*\mathcal{F}$, by $f_*\mathcal{F} := \{V \in Y : f^{-1}(V) \in \mathcal{F}\}$. **this should really just be about filters in $(\mathcal{P}(X), \subset)$**

Exercise 4.4.0.6. Let X, Y be a sets, $f : X \rightarrow Y$ and $\mathcal{F}$ a proper filter on X . Then

Definition 4.4.0.7. Let X, Y be a topological spaces, $f : X \rightarrow Y$, $\mathcal{F}$ a proper filter on X and $y \in Y$. Then f is said to **converge to y along $\mathcal{F}$** if $f_*(\mathcal{F})$ converges to y .

Exercise 4.4.0.8. Let X be a topological space, $\mathcal{F}$ a proper filter on X and $a \in Y$.

Exercise 4.4.0.9. Let X be a topological space, $f : X \rightarrow \mathbb{R}$, $(x_\alpha)_{\alpha \in A} \subset X$ a net and $x \in X$. Suppose that $x_\alpha \rightarrow x$ and for each $\alpha \in A$, $x_\alpha \neq x$. Then

1. $\liminf_{t \rightarrow x} f(t) \leq \liminf f(x_\alpha)$
2. $\limsup_{t \rightarrow x} f(t) \geq \limsup f(x_\alpha)$

Proof.

1. Let $V \in \mathcal{N}(x)$. Then there exists $\beta_0 \in A$ such that for each $\alpha \in A$, $\alpha \geq \beta_0$ implies that $x_\alpha \in V \setminus \{x\}$. Thus

$$\begin{aligned} \inf_{t \in V \setminus \{x\}} f(t) &\leq \inf_{\alpha \geq \beta_0} f(x_\alpha) \\ &\leq \sup_{\beta \in A} \left[\inf_{\alpha \geq \beta} f(x_\alpha) \right] \\ &= \liminf_{\alpha \in A} f(x_\alpha) \end{aligned}$$

and since $V \in \mathcal{N}(x)$ is arbitrary, we have that

$$\begin{aligned} \liminf_{t \rightarrow x} f(t) &= \sup_{V \in \mathcal{N}(x)} \left[\inf_{t \in V \setminus \{x\}} f(t) \right] \\ &\leq \liminf_{\alpha \in A} f(x_\alpha) \end{aligned}$$

2. Similar to (1).

□

4.5 Subspace Topology

4.5.1 Introduction

Definition 4.5.1.1. Let X be a set and $A \subset X$. We define the **inclusion map from A to X** , denoted $\iota : A \rightarrow X$, by $\iota(x) = x$.

Definition 4.5.1.2. Let $(X, \mathcal{T})$ be a topological space and $A \subset X$. We define the **subspace topology on A** , denoted $\mathcal{T} \cap A$, by

$$\mathcal{T} \cap A = \iota^* \mathcal{T}.$$

Note 4.5.1.3. Exercise 4.2.0.8 implies that $\mathcal{T} \cap A$ is a topology on A .

Exercise 4.5.1.4. Let $(X, \mathcal{T})$ be a topological space and $A \subset X$. Then

1. $\mathcal{T} \cap A = \{U \cap A : U \in \mathcal{T}\}$,
2. for each $E \subset A$, $E \in \mathcal{T} \cap A$ iff there exists $U \in \mathcal{T}$ such that $E = U \cap A$.

Proof.

1. Since for each $U \subset X$, $\iota^{-1}(U) = U \cap A$, we have that

$$\begin{aligned} \mathcal{T} \cap A &= \iota^* \mathcal{T} \\ &= \{\iota^{-1}(U) : U \in \mathcal{T}\} \\ &= \{U \cap A : U \in \mathcal{T}\} \end{aligned}$$

2. Clear

□

Exercise 4.5.1.5. Let X be a set, $\mathcal{E} \subset \mathcal{P}(X)$ and $A \subset X$. Then

$$\tau_X(\mathcal{E}) \cap A = \tau_A(\mathcal{E} \cap A).$$

Hint: $\tau_X(\mathcal{E}) \subset (\iota_A)_* \tau_A(\mathcal{E} \cap A)$

Proof.

- Clearly $\mathcal{E} \cap A \subset \tau_X(\mathcal{E}) \cap A$. Since $\tau_X(\mathcal{E}) \cap A$ is a topology on A , we have that

$$\begin{aligned} \tau_A(\mathcal{E} \cap A) &\subset \tau_A[\tau_X(\mathcal{E}) \cap A] \\ &= \tau_X(\mathcal{E}) \cap A. \end{aligned}$$

- Let $V_0 \in \mathcal{E}$. Then

$$\begin{aligned} \iota_A^{-1}(V_0) &= V_0 \cap A \\ &\in \mathcal{E} \cap A \\ &\subset \tau_A(\mathcal{E} \cap A). \end{aligned}$$

Hence $V_0 \in (\iota_A)_* \tau_A(\mathcal{E} \cap A)$. Since $V_0 \in \mathcal{E}$ is arbitrary, we have that $\mathcal{E} \subset (\iota_A)_* \tau_A(\mathcal{E} \cap A)$. Thus

$$\begin{aligned} \tau_X(\mathcal{E}) &\subset \tau_X[(\iota_A)_* \tau_A(\mathcal{E} \cap A)] \\ &= (\iota_A)_* \tau_A(\mathcal{E} \cap A). \end{aligned}$$

Let $V \in \tau_X(\mathcal{E}) \cap A$. Then there exists $V_0 \in \tau_X(\mathcal{E})$ such that $V = V_0 \cap A$. Therefore

$$\begin{aligned} V_0 &\in \tau_X(\mathcal{E}) \\ &\subset (\iota_A)_* \tau_A(\mathcal{E} \cap A). \end{aligned}$$

By definition of $(\iota_A)_*\tau_A(\mathcal{E} \cap A)$, we have that

$$\begin{aligned} V &= V_0 \cap A \\ &= \iota_A^{-1}(V_0) \\ &\in \tau_A(\mathcal{E} \cap A). \end{aligned}$$

Since $V \in \tau_X(\mathcal{E}) \cap A$ is arbitrary, we have that $\tau_X(\mathcal{E}) \cap A \subset \tau_A(\mathcal{E} \cap A)$.

Since $\tau_A(\mathcal{E} \cap A) \subset \tau_X(\mathcal{E}) \cap A$ and $\tau_X(\mathcal{E}) \cap A \subset \tau_A(\mathcal{E} \cap A)$, we have that $\tau_X(\mathcal{E}) \cap A = \tau_A(\mathcal{E} \cap A)$. □

Exercise 4.5.1.6. Let $(X, \mathcal{T})$ be a topological space, $A \subset X$ and $B \subset A$. Then $\mathcal{T} \cap B = (\mathcal{T} \cap A) \cap B$.

Proof.

- Let $U \in (\mathcal{T} \cap A) \cap B$. Then there exists $U_A \in \mathcal{T} \cap A$ such that $U = U_A \cap B$. Similarly, there exists $U_X \in \mathcal{T}$ such that $U_A = U_X \cap A$. Therefore

$$\begin{aligned} U &= U_A \cap B \\ &= (U_X \cap A) \cap B \\ &= U_X \cap (A \cap B) \\ &= U_X \cap B \\ &\in \mathcal{T} \cap B \end{aligned}$$

Since $U \in (\mathcal{T} \cap A) \cap B$ is arbitrary, we have that $(\mathcal{T} \cap A) \cap B \subset \mathcal{T} \cap B$.

- Conversely, let $U \in \mathcal{T} \cap B$. Then there exists $U_X \in \mathcal{T}$ such that $U = U_X \cap B$. Then

$$\begin{aligned} U &= U_X \cap B \\ &= U_X \cap (A \cap B) \\ &= (U_X \cap A) \cap B \\ &\in (\mathcal{T} \cap A) \cap B \end{aligned}$$

Since $U \in U_X \cap B$ is arbitrary, we have that $U_X \cap B \subset (\mathcal{T} \cap A) \cap B$.

Therefore $\mathcal{T} \cap B = (\mathcal{T} \cap A) \cap B$. □

Note 4.5.1.7. The previous exercise just indicates that the subspace topology on B does not depend on choice of the containing space as long as each containing space is a subspace of the original space **maybe rephrase this in terms of projective system or something**.

Exercise 4.5.1.8. Let $(X, \mathcal{T})$ be a topological space, $A \subset X$ and $S \subset A$. Then $\mathcal{N}(x, \mathcal{T} \cap A) = \mathcal{N}(x, \mathcal{T}) \cap A$.

Proof.

- Let $U \in \mathcal{N}(S, \mathcal{T} \cap A)$. Exercise 4.1.0.24 implies that $S \subset \text{Int}_{\mathcal{T} \cap A} U$. Since $\text{Int}_{\mathcal{T} \cap A} U \in \mathcal{T} \cap A$, there exists $V \in \mathcal{T}$ such that $\text{Int}_{\mathcal{T} \cap A} U = V \cap A$. Define $E \subset A$ by $E := U \setminus \text{Int}_{\mathcal{T} \cap A} U$. Then

$$\begin{aligned} U &= (\text{Int}_{\mathcal{T} \cap A} U) \cup E \\ &= (V \cap A) \cup (E \cap A) \\ &= [(V \cap A) \cup E] \cap [(V \cap A) \cup A] \\ &= [(V \cup E) \cap (A \cup E)] \cap [(V \cup A) \cap (A \cup A)] \\ &= [(V \cup E) \cap (A \cup E)] \cap [(V \cup A) \cap A] \\ &= [(V \cup E) \cap (A \cup E)] \cap A \\ &= (V \cup E) \cap [(A \cup E) \cap A] \\ &= (V \cup E) \cap A. \end{aligned}$$

Since $V \in \mathcal{T}$, $S \subset V$ and $V \subset V \cup E$, we have that $V \cup E \in \mathcal{N}(S, \mathcal{T})$. Thus

$$\begin{aligned} U &= (V \cup E) \cap A \\ &\in \mathcal{N}(S, \mathcal{T}) \cap A. \end{aligned}$$

Since $U \in \mathcal{N}(S, \mathcal{T} \cap A)$ is arbitrary, we have that for each $U \in \mathcal{N}(S, \mathcal{T} \cap A)$, $U \in \mathcal{N}(S, \mathcal{T}) \cap A$. Hence $\mathcal{N}(S, \mathcal{T} \cap A) \subset \mathcal{N}(S, \mathcal{T}) \cap A$.

- Let $U \in \mathcal{N}(S, \mathcal{T}) \cap A$. Then there exists $V \in \mathcal{N}(S, \mathcal{T})$ such that $U = V \cap A$. We note that $S \subset (\text{Int}_{\mathcal{T}} V) \cap A$ and $(\text{Int}_{\mathcal{T}} V) \cap A \in \mathcal{T} \cap A$. Define $E \subset X$ by $E := V \setminus \text{Int}_{\mathcal{T}} V$. Then $[(\text{Int}_{\mathcal{T}} V) \cap A] \cup (E \cap A) \in \mathcal{N}(S, \mathcal{T} \cap A)$ and therefore

$$\begin{aligned} U &= V \cap A \\ &= [(\text{Int}_{\mathcal{T}} V) \cup E] \cap A \\ &= [(\text{Int}_{\mathcal{T}} V) \cap A] \cup (E \cap A) \\ &\in \mathcal{N}(S, \mathcal{T} \cap A). \end{aligned}$$

Since $U \in \mathcal{N}(S, \mathcal{T}) \cap A$ is arbitrary, we have that for each $U \in \mathcal{N}(S, \mathcal{T}) \cap A$, $U \in \mathcal{N}(S, \mathcal{T} \cap A)$. Hence $\mathcal{N}(S, \mathcal{T} \cap A) \subset \mathcal{N}(S, \mathcal{T}) \cap A$. $\square$

Exercise 4.5.1.9. Let $(X, \mathcal{T})$ be a topological space, $E \subset X$, $A \subset E$ and $x \in E$. Then x is a $\mathcal{T} \cap E$ -condensation point of A iff x is a $\mathcal{T}$ -condensation point of A .

Proof.

- $(\implies)$:

Suppose that x is a $\mathcal{T} \cap E$ -condensation point of A . Let $U \in \mathcal{N}(x, \mathcal{T})$. Exercise 4.5.1.8 implies that $U \cap E \in \mathcal{N}(x, \mathcal{T} \cap E)$. Since x is a $\mathcal{T} \cap E$ -condensation point of A , we have that $(U \cap E) \cap A$ is uncountable. Since $A \subset E$, we have that

$$\begin{aligned} U \cap A &= U \cap (A \cap E) \\ &= (U \cap E) \cap A \end{aligned}$$

and therefore $U \cap A$ is uncountable. Since $U \in \mathcal{N}(x, \mathcal{T})$ is arbitrary, we have that for each $U \in \mathcal{N}(x, \mathcal{T})$, $U \cap A$ is uncountable. Hence x is a $\mathcal{T}$ -condensation point of A .

- $(\impliedby)$:

Suppose that x is a $\mathcal{T}$ -condensation point of A . Let $U \in \mathcal{N}(x, \mathcal{T} \cap E)$. Exercise 4.5.1.8 implies that $U \in \mathcal{N}(x, \mathcal{T}) \cap E$. Thus there exists $V \in \mathcal{N}(x, \mathcal{T})$ such that $U = V \cap E$. Since x is a $\mathcal{T}$ -condensation point of A , we have that $V \cap A$ is uncountable. Since

$$\begin{aligned} U \cap A &= (V \cap E) \cap A \\ &= V \cap (A \cap E) \\ &= V \cap A, \end{aligned}$$

we have that $U \cap A$ is uncountable. Since $U \in \mathcal{N}(x, \mathcal{T} \cap E)$ is arbitrary, we have that for each $U \in \mathcal{N}(x, \mathcal{T} \cap E)$, $U \cap A$ is uncountable. Hence x is a $\mathcal{T} \cap E$ -condensation point of A . $\square$

Exercise 4.5.1.10. Let $(X, \mathcal{T})$ be a topological space, $A \subset X$, $(x_\gamma)_{\gamma \in \Gamma} \subset A$ a net and $x \in A$. Then $x_\gamma \rightarrow x$ in $(A, \mathcal{T} \cap A)$ iff $x_\gamma \rightarrow x$ in $(X, \mathcal{T})$.

Proof.

- Suppose that $x_\gamma \rightarrow x$ in $(A, \mathcal{T} \cap A)$. Since $\iota : A \rightarrow X$ is $(\mathcal{T} \cap A, \mathcal{T})$ -continuous,

$$\begin{aligned} x_\gamma &= \iota(x_\gamma) \\ &\rightarrow \iota(x) \\ &= x \end{aligned}$$

So that $x_\gamma \rightarrow x$ in $(X, \mathcal{T})$.

- Conversely, suppose that $x_\gamma \rightarrow x$ in $(X, \mathcal{T})$. Let $V \in \mathcal{N}(x)$ in $(A, \mathcal{T} \cap A)$. Then $x \in \text{Int } V$ in $(A, \mathcal{T} \cap A)$. Hence there exists $U \in \mathcal{T}$ such that $\text{Int } V = U \cap A$. Thus $U \in \mathcal{N}(x)$ in $(X, \mathcal{T})$. Since $x_\gamma \rightarrow x$ in $(X, \mathcal{T})$ and $x \in U$, we have that $(x_\gamma)_{\gamma \in \Gamma}$ is eventually in U . Then $(x_\gamma)_{\gamma \in \Gamma}$ is eventually in $U \cap A = \text{Int } V \subset V$. So $x_\gamma \rightarrow x$ in $(A, \mathcal{T} \cap A)$.

□

Exercise 4.5.1.11. Let $(X, \mathcal{T}_X), (Y, \mathcal{T}_Y)$ be topological spaces, $B \subset Y$ and $f : X \rightarrow Y$. Suppose that $f(X) \subset B$. Then $f|_X^B$ is $(\mathcal{T}_X, \mathcal{T}_Y \cap B)$ -continuous iff f is $(\mathcal{T}_X, \mathcal{T}_Y)$ -continuous.

Proof.

- $(\implies)$:
Suppose that $f|_X^B$ is $(\mathcal{T}_X, \mathcal{T}_Y \cap B)$ -continuous. Let $(x_\gamma)_{\gamma \in \Gamma} \subset X$ a net and $x \in X$. Suppose that $x_\gamma \rightarrow x$. Since $f|_X^B$ is $(\mathcal{T}_X, \mathcal{T}_Y \cap B)$ -continuous, we have that

$$\begin{aligned} f(x_\gamma) &= f|_X^B(x_\gamma) \\ &\xrightarrow{(\mathcal{T}_Y \cap B)} f|_X^B(x) \\ &= f(x) \end{aligned}$$

Exercise 4.5.1.10 implies that $f(x_\gamma) \rightarrow f(x)$ in $(Y, \mathcal{T}_Y)$. Since $(x_\gamma)_{\gamma \in \Gamma} \subset X$ is an arbitrary net and $x \in X$ is arbitrary, we have that f is $(\mathcal{T}_X, \mathcal{T}_Y)$ -continuous.

- $(\impliedby)$:
Similar to (1).

□

Exercise 4.5.1.12. Let $(X, \mathcal{T}_X), (Y, \mathcal{T}_Y)$ be topological spaces and $f : X \rightarrow Y$. Then

1. f is $(\mathcal{T}_X, \mathcal{T}_Y)$ -continuous iff for each $x \in X$, there exists $U \in \mathcal{T}_X$ such that $x \in U$ and $f|_U$ is $(\mathcal{T}_X \cap U, \mathcal{T}_Y)$ -continuous.
2. f is $(\mathcal{T}_X, \mathcal{T}_Y)$ -continuous iff for each $E \subset X$, $f|_E$ is $(\mathcal{T}_X \cap E, \mathcal{T}_Y)$ -continuous.

Proof.

1. • $(\implies)$:
Suppose that f is continuous. Let $x \in X$. Define $U \in \mathcal{T}_X$ by $U := X$. Then $x \in U$ and $f|_U$ is $(\mathcal{T}_X \cap U, \mathcal{T}_Y)$ -continuous.
- $(\impliedby)$:
Suppose that for each $x \in X$, there exists $U \in \mathcal{T}_X$ such that $x \in U$ and $f|_U$ is $(\mathcal{T}_X \cap U, \mathcal{T}_Y)$ -continuous. Let $x \in X$ and $V \in \mathcal{T}_Y$. Suppose that $f(x) \in V$. By assumption, there exists $U_0 \in \mathcal{T}_X$ such that $x \in U_0$ and $f|_{U_0}$ is $(\mathcal{T}_X \cap U_0, \mathcal{T}_Y)$ -continuous. Define $U \subset X$ by $U = f|_{U_0}^{-1}(V)$. Since $V \in \mathcal{T}_Y$ and $f|_{U_0}$ is $(\mathcal{T}_X \cap U_0, \mathcal{T}_Y)$ -continuous, $U \in \mathcal{T}_X \cap U_0$. Since $U_0 \in \mathcal{T}_X$, $\mathcal{T}_X \cap U_0 \subset \mathcal{T}_X$ and therefore $U \in \mathcal{T}_X$. We note that $x \in U$. Since $f|_{U_0}^{-1}(V) = U_0 \cap f^{-1}(V)$, we have that

$$\begin{aligned} f(U) &= f(f|_{U_0}^{-1}(V)) \\ &= f(U_0 \cap f^{-1}(V)) \\ &\subset V \end{aligned}$$

Since $V \in \mathcal{T}_Y$ with $f(x) \in V$ is arbitrary, we have that for each $V \in \mathcal{T}_Y$, $f(x) \in V$ implies that there exists $U \in \mathcal{T}_X$ such that $x \in U$ and $f(U) \subset V$. Thus f is continuous at x . Since $x \in X$ is arbitrary, f is continuous.

2. • $(\implies)$:
Suppose that f is $(\mathcal{T}_X, \mathcal{T}_Y)$ -continuous. Let $E \subset X$. Then for each $V \in \mathcal{T}_Y$, $f^{-1}(V) \in \mathcal{T}_X$ and

$$\begin{aligned} f|_E^{-1}(V) &= (f \circ \iota_E)^{-1}(V) &= \iota_E^{-1}(f^{-1}(V)) \\ &= E \cap f^{-1}(V) \\ &\in \mathcal{T}_X \cap E \quad (\text{by Exercise 4.5.1.4}). \end{aligned}$$

Thus $f|_E$ is $(\mathcal{T}_X \cap E, \mathcal{T}_Y)$ -continuous. Since E is arbitrary, we have that for each $E \subset X$, $f|_E$ is $(\mathcal{T}_X \cap E, \mathcal{T}_Y)$ -continuous.

- ($\Leftarrow$):

Suppose that for each $E \subset X$, $f|_E$ is $(\mathcal{T}_X \cap E, \mathcal{T}_Y)$ -continuous. Define $E \subset X$ by $E := X$. Since $\mathcal{T} \cap E = \mathcal{T}$ and $f|_E = f$, we have that f is $(\mathcal{T}_X, \mathcal{T}_Y)$ -continuous.

□

Exercise 4.5.1.13. Let $(X, \mathcal{T})$ be a topological space, $A \subset X$ and $F \subset A$. Then F is $\mathcal{T} \cap A$ -closed iff there exists $C \subset X$ such that C is $\mathcal{T}$ -closed and $F = C \cap A$.

Proof. Suppose that F is $\mathcal{T} \cap A$ -closed. Then $A \setminus F$ is $\mathcal{T} \cap A$ -open. Hence there exists $U \in \mathcal{T}$ such that $A \setminus F = U \cap A$. Set $C = U^c$. Then C is closed in X and

$$\begin{aligned} F &= A \setminus (A \setminus F) \\ &= A \setminus (U \cap A) \\ &= A \cap [(U \cap A)^c] \\ &= A \cap (U^c \cup A^c) \\ &= (A \cap U^c) \cup (A \cap A^c) \\ &= A \cap U^c \\ &= A \cap C \end{aligned}$$

Conversely, suppose that there exists $C \subset X$ such that C is closed in X and $F = A \cap C$. Since $C^c \in \mathcal{T}$, we have that

$$\begin{aligned} A \setminus F &= A \cap F^c \\ &= A \cap [(A \cap C)^c] \\ &= A \cap (A^c \cup C^c) \\ &= (A \cap A^c) \cup (A \cap C^c) \\ &= A \cap C^c \\ &\in \mathcal{T} \cap A \end{aligned}$$

Thus $A \setminus F$ is $\mathcal{T} \cap A$ -open which implies that F is $\mathcal{T} \cap A$ -closed. □

Exercise 4.5.1.14. Let $(X, \mathcal{T}_X)$ and $(Y, \mathcal{T}_Y)$ be topological spaces and $f : X \rightarrow Y$. Then f is $(\mathcal{T}_X, \mathcal{T}_Y)$ -continuous iff f is $(\mathcal{T}_X, \mathcal{T}_Y \cap f(X))$ -continuous.

duplicate of Exercise 4.5.1.11, maybe keep statement in Exercise 4.5.1.11, but keep or add this proof in Exercise 4.5.1.14

Proof.

- ($\Rightarrow$):

Suppose that f is $(\mathcal{T}_X, \mathcal{T}_Y)$ -continuous. Let $B \in \mathcal{T}_Y \cap f(X)$. Then there exists $V \in \mathcal{T}_Y$ such that $B = V \cap f(X)$. Then

$$\begin{aligned} f^{-1}(B) &= f^{-1}(V \cap f(X)) \\ &= f^{-1}(V) \cap f^{-1}(f(X)) \\ &= f^{-1}(V) \cap X \\ &= f^{-1}(V) \\ &\in \mathcal{T}_X \end{aligned}$$

Since $B \in \mathcal{T}_Y \cap f(X)$ is arbitrary, f is $(\mathcal{T}_X, \mathcal{T}_Y \cap f(X))$ -continuous.

- ($\Leftarrow$):

Conversely, suppose that f is $(\mathcal{T}_X, \mathcal{T}_Y \cap f(X))$ -continuous. Let $V \in \mathcal{T}_Y$. Then $V \cap f(X) \in \mathcal{T}_Y \cap f(X)$ and

$$\begin{aligned} f^{-1}(V) &= f^{-1}(V \cap f(X)) \\ &\in \mathcal{T}_X \end{aligned}$$

Since $V \in \mathcal{T}_Y$ is arbitrary, f is $(\mathcal{T}_X, \mathcal{T}_Y)$ -continuous.

□

Exercise 4.5.1.15. Basis for Subspace Topology:

Let $(X, \mathcal{T})$ be a topological space, $\mathcal{B} \subset \mathcal{T}$ a basis for $\mathcal{T}$ on X and $A \subset X$. Then $\mathcal{B} \cap A$ is a basis for $\mathcal{T} \cap A$.

Proof. Let $E \in \mathcal{T} \cap A$. Then there exists $E' \in \mathcal{T}$ such that $E = E' \cap A$. Since $\mathcal{B}$ is a basis for $\mathcal{T}$ on X , there exists $\mathcal{U}' \subset \mathcal{B}$ such that $E' = \bigcup_{U' \in \mathcal{U}'} U'$. Define $\mathcal{U} \subset \mathcal{T} \cap A$ by $\mathcal{U} := \mathcal{U}' \cap A$. Then

$$\begin{aligned} \mathcal{U} &= \mathcal{U}' \cap A \\ &\subset \mathcal{B} \cap A \end{aligned}$$

and

$$\begin{aligned} E &= E' \cap A \\ &= \left(\bigcup_{U' \in \mathcal{U}'} U' \right) \cap A \\ &= \bigcup_{U' \in \mathcal{U}'} (U' \cap A) \\ &= \bigcup_{U \in \mathcal{U}' \cap A} U \\ &= \bigcup_{U \in \mathcal{U}} U \end{aligned}$$

Since $E \in \mathcal{T} \cap A$ is arbitrary, we have that for each $E \in \mathcal{T} \cap A$, there exists $\mathcal{U} \subset \mathcal{B} \cap A$ such that $E = \bigcup_{U \in \mathcal{U}} U$. Hence $\mathcal{B} \cap A$ is a basis for $\mathcal{T} \cap A$ on A . □

Exercise 4.5.1.16. Let $(X, \mathcal{T}_X)$, $(Y, \mathcal{T}_Y)$ be topological spaces and $f : X \rightarrow Y$. If f is $(\mathcal{T}_X, \mathcal{T}_Y)$ -open, then for each $U \in \mathcal{T}_X$, $f|_U$ is $(\mathcal{T}_X \cap U, \mathcal{T}_Y)$ -open.

Proof. Suppose that f is $(\mathcal{T}_X, \mathcal{T}_Y)$ -open. Let $U \in \mathcal{T}_X$ and $V \in \mathcal{T}_X \cap U$. Since $U \in \mathcal{T}_X$,

$$\begin{aligned} V &\in \mathcal{T} \cap U \\ &\subset \mathcal{T}. \end{aligned}$$

Since f is $(\mathcal{T}_X, \mathcal{T}_Y)$ -open and $V \subset U$, we have that

$$\begin{aligned} f|_U(V) &= f(V) \\ &\in \mathcal{T}_Y. \end{aligned}$$

Since $V \in \mathcal{T}_X \cap U$ is arbitrary, we have that for each $V \in \mathcal{T}_X \cap U$, $f|_U(V) \in \mathcal{T}_Y$. Thus $f|_U$ is $(\mathcal{T}_X \cap U, \mathcal{T}_Y)$ -open. Since $U \in \mathcal{T}_X$ is arbitrary, we have that for each $U \in \mathcal{T}_X$, $f|_U$ is $(\mathcal{T}_X \cap U, \mathcal{T}_Y)$ -open. □

Exercise 4.5.1.17. Let $(X, \mathcal{T}_X)$, $(Y, \mathcal{T}_Y)$ be topological spaces and $f : X \rightarrow Y$. Then f is $(\mathcal{T}_X, \mathcal{T}_Y)$ -open iff for each $x \in X$, there exists $U \in \mathcal{T}_X$ such that $x \in U$ and $f|_U$ is $(\mathcal{T}_X \cap U, \mathcal{T}_Y)$ -open.

Proof.

- ($\Rightarrow$):
Suppose that f is $(\mathcal{T}_X, \mathcal{T}_Y)$ -open. Let $x \in X$. Set $U := X$. Then $X \in \mathcal{T}_X$, $x \in U$ and $f|_U$ is $(\mathcal{T}_X \cap U, \mathcal{T}_Y)$ -open.
- ($\Leftarrow$):
Suppose that for each $x \in X$, there exists $U \in \mathcal{T}_X$ such that $x \in U$ and $f|_U$ is $(\mathcal{T}_X \cap U, \mathcal{T}_Y)$ -open. Let $V \in \mathcal{T}_X$. By assumption,

for each $x \in V$, there exists $U_x \in \mathcal{T}_X$ such that $f|_{U_x}$ is $(\mathcal{T}_X \cap U_x, \mathcal{T}_Y)$ -open. Then for each $x \in V$, $V \cap U_x \in \mathcal{T}_X \cap U_x$ and $V = \bigcup_{x \in V} (V \cap U_x)$. Since for each $x \in V$, $f|_{U_x}$ is $(\mathcal{T}_X \cap U_x, \mathcal{T}_Y)$ -open, we have that

$$\begin{aligned} f(V) &= f\left(\bigcup_{x \in V} (V \cap U_x)\right) \\ &= \bigcup_{x \in V} f(V \cap U_x) \\ &= \bigcup_{x \in V} f|_{U_x}(V \cap U_x) \\ &\in \mathcal{T}_Y \end{aligned}$$

Since $V \in \mathcal{T}_X$ is arbitrary, we have that for each $V \in \mathcal{T}_X$, $f(V) \in \mathcal{T}_Y$. Hence f is open. □

Exercise 4.5.1.18. Let $(X, \mathcal{T}_X), (Y, \mathcal{T}_Y)$ be topological spaces, $f : X \rightarrow Y$ and $E \subset X$. Suppose that f is $(\mathcal{T}_X, \mathcal{T}_Y)$ -closed. If E is closed in $(X, \mathcal{T}_Y)$, then $f|_E$ is $(\mathcal{T}_X \cap E, \mathcal{T}_Y)$ -closed.

Proof. Suppose that E is closed in $(X, \mathcal{T}_Y)$. Let $F \subset E$. Suppose that F is closed in $(E, \mathcal{T}_X \cap E)$. Exercise 4.5.1.13 implies that there exists $C \subset X$ such that C is closed in $(X, \mathcal{T}_X)$ and $F = C \cap E$. Since C, E are closed in $(X, \mathcal{T}_X)$, F is closed in $(X, \mathcal{T}_X)$. Since f is closed, $f(F)$ is closed in $(Y, \mathcal{T}_Y)$. Since $f|_E(F) = f(F)$, we have that $f|_E(F)$ is closed in $(Y, \mathcal{T}_Y)$. Since $F \subset E$ with F closed in $(E, \mathcal{T}_X \cap E)$ is arbitrary, we have that for each $F \subset E$, F is closed in $(E, \mathcal{T}_X \cap E)$ implies that $f|_E(F)$ is closed in $(Y, \mathcal{T}_Y)$. Hence $f|_E$ is $(\mathcal{T}_X \cap E, \mathcal{T}_Y)$ -closed. □

Exercise 4.5.1.19. Let $(X, \mathcal{T}_X), (A, \mathcal{T}_A)$ be topological spaces. Suppose that $A \subset X$. If $\iota_A|_A$ is a $(\mathcal{T}_A, \mathcal{T}_X \cap A)$ -homeomorphism, then $\mathcal{T}_A = \mathcal{T}_X \cap A$.

Proof. Suppose that $\iota_A|_A$ is a $(\mathcal{T}_A, \mathcal{T}_X \cap A)$ -homeomorphism.

- Let $U \in \mathcal{T}_A$. Since $\iota_A(U) = U$ and $\iota_A|_A$ is $(\mathcal{T}_A, \mathcal{T}_X \cap A)$ -open, we have that

$$\begin{aligned} U &= \iota_A(U) \\ &= \iota_A|_A(U) \\ &\in \mathcal{T}_X \cap A. \end{aligned}$$

Since $U \in \mathcal{T}_A$ is arbitrary, we have that $\mathcal{T}_A \subset \mathcal{T}_X \cap A$.

- Let $U \in \mathcal{T}_X \cap A$. Since $\iota_A|_A$ is $(\mathcal{T}_A, \mathcal{T}_X \cap A)$ -continuous and $U \subset A$, we have that we have that

$$\begin{aligned} U &= \iota_A^{-1}(U) \\ &= (\iota_A|_A)^{-1}(U) \\ &\in \mathcal{T}_A. \end{aligned}$$

Since $U \in \mathcal{T}_X \cap A$ is arbitrary, we have that $\mathcal{T}_X \cap A \subset \mathcal{T}_A$.

Hence $\mathcal{T}_A = \mathcal{T}_X \cap A$. □

Exercise 4.5.1.20. universal property

Proof. **FINISH!!!** □

4.5.2 Discrete Spaces

Exercise 4.5.2.1. Let $(X, \mathcal{T})$ be a topological space, $A \subset X$ and $x \in A$. Then x is an isolated point of A iff $\{x\} \in \mathcal{T} \cap A$

Proof.

• $(\implies)$:

Suppose that x is an isolated point of A . Then there exists $U \in \mathcal{T}$ such that

$$\begin{aligned}\{x\} &= U \cap A \\ &\in \mathcal{T} \cap A\end{aligned}$$

• $(\impliedby)$:

Conversely, suppose that $\{x\} \in \mathcal{T} \cap A$. Then there exists $U \in \mathcal{T}$ such that $\{x\} = U \cap A$. Since $x \in U$, $U \in \mathcal{N}^{\text{op}}(x, \mathcal{T})$. Hence x is an isolated point of A .

•

□

Exercise 4.5.2.2. Let $(X, \mathcal{T})$ be a topological space and $A \subset X$. Then A is discrete iff $\mathcal{T} \cap A = \mathcal{P}(A)$.

Proof. Suppose that A is discrete. Then for each $x \in A$, $\{x\} \in \mathcal{T} \cap A$. Let $U \in \mathcal{P}(A)$. Then

$$\begin{aligned}U &= \bigcup_{x \in U} \{x\} \\ &\in \mathcal{T} \cap A\end{aligned}$$

Since $U \in \mathcal{P}(A)$ is arbitrary, we have that $\mathcal{P}(A) \subset \mathcal{T} \cap A$. Since $\mathcal{T} \cap A \subset \mathcal{P}(A)$, we have that $\mathcal{T} \cap A = \mathcal{P}(A)$. Conversely, suppose that $\mathcal{T} \cap A = \mathcal{P}(A)$. Let $x \in A$. Then

$$\begin{aligned}\{x\} &\in \mathcal{P}(A) \\ &= \mathcal{T} \cap A\end{aligned}$$

Hence x is an isolated point of A . Since $x \in A$ is arbitrary, A is discrete.

□

4.5.3 Topological Embeddings

Definition 4.5.3.1. Let $(X, \mathcal{T}_X), (Y, \mathcal{T}_Y)$ be a topological space and $f : X \rightarrow Y$. Then f is said to be a **Top**-embedding if $f : X \rightarrow f(X)$ is a $(\mathcal{T}_X, \mathcal{T}_Y \cap f(X))$ -homeomorphism.

Exercise 4.5.3.2. Let $(X, \mathcal{T}_X), (Y, \mathcal{T}_Y)$ be a topological space and $f : X \rightarrow Y$. Suppose that f is $(\mathcal{T}_X, \mathcal{T}_Y)$ -continuous and injective. Then

1. f is open implies that f is a **Top**-embedding
2. f is closed implies that f is a **Top**-embedding

Proof.

1. Suppose that f is open. Since f is $(\mathcal{T}_X, \mathcal{T}_Y)$ -continuous, f is $(\mathcal{T}_X, \mathcal{T}_Y \cap f(X))$ -continuous. Since f is injective, $f : X \rightarrow f(X)$ is bijective. Since f is $(\mathcal{T}_X, \mathcal{T}_Y)$ -open, f is $(\mathcal{T}_X, \mathcal{T}_Y \cap f(X))$ -open. Thus $f : X \rightarrow f(X)$ is a $(\mathcal{T}_X, \mathcal{T}_Y \cap f(X))$ -homeomorphism.
2. Similar to (1) by Exercise 4.2.0.3.

□

4.6 Product Topology

4.6.1 Bases of the product topology

Definition 4.6.1.1. Let $(X_\alpha, \mathcal{T}_\alpha)_{\alpha \in A}$ be a collection of topological spaces. We define the **product topology** on $\prod_{\alpha \in A} X_\alpha$, denoted $\bigotimes_{\alpha \in A} \mathcal{T}_\alpha$, by

$$\bigotimes_{\alpha \in A} \mathcal{T}_\alpha = \tau(\pi_\alpha : \alpha \in A)$$

i.e. $\bigotimes_{\alpha \in A} \mathcal{T}_\alpha$ is the initial (weak) topology on $\prod_{\alpha \in A} X_\alpha$ generated by the projection maps $(\pi_\alpha)_{\alpha \in A}$.

Exercise 4.6.1.2. Let $(X_\alpha, \mathcal{T}_\alpha)_{\alpha \in A}$ be a collection of topological spaces. Define

$$\mathcal{B} = \left\{ \prod_{\alpha \in A} B_\alpha : \text{for each } \alpha \in A, B_\alpha \in \mathcal{T}_\alpha \text{ and } \#\{\alpha \in A : B_\alpha \neq X_\alpha\} < \infty \right\}$$

Then $\mathcal{B}$ is a basis for $\bigotimes_{\alpha \in A} \mathcal{T}_\alpha$.

Proof. Set $X = \prod_{\alpha \in A} X_\alpha$ and $\mathcal{T}_X = \bigotimes_{\alpha \in A} \mathcal{T}_\alpha$. Set

$$\mathcal{E} = \{\pi_\alpha^{-1}(B_\alpha) : \alpha \in A, B_\alpha \in \mathcal{T}_\alpha\}$$

By definition, $\mathcal{T}_X = \tau_X(\mathcal{E})$. Let $\alpha \in A$ and $B_\alpha \in \mathcal{T}_\alpha$. For $\beta \in A$, set

$$C_\beta = \begin{cases} B_\beta & \beta = \alpha \\ X_\beta & \beta \neq \alpha \end{cases}$$

Then

$$\pi_\alpha^{-1}(B_\alpha) = \prod_{\beta \in A} C_\beta$$

Hence $\mathcal{B} = \left\{ \bigcap_{j=1}^n V_j : (V_j)_{j=1}^n \subset \mathcal{E} \right\} \subset \mathcal{T}_X$. Exercise 4.1.0.13 implies that $\mathcal{B}$ is a basis for $\mathcal{T}_X$. □

Exercise 4.6.1.3. Let $(X_\alpha, \mathcal{T}_\alpha)_{\alpha \in A}$ be a collection of topological spaces. Then for each $\alpha \in A$, $\pi_\alpha : \prod_{\alpha \in A} X_\alpha \rightarrow X_\alpha$ is open.

Proof. Let $\alpha \in A$. Define

$$\mathcal{B} = \left\{ \prod_{\alpha \in A} B_\alpha : \text{for each } \alpha \in A, B_\alpha \in \mathcal{T}_\alpha \text{ and } \#\{\alpha \in A : B_\alpha \neq X_\alpha\} < \infty \right\}$$

Then Exercise 4.6.1.2 implies that $\mathcal{B}$ is a basis for $\bigotimes_{\alpha \in A} \mathcal{T}_\alpha$. Let $U \in \bigotimes_{\alpha \in A} \mathcal{T}_\alpha$. Then for each $\alpha' \in A$, there exist $B_{\alpha'} \in \mathcal{T}_{\alpha'}$ such that $U = \prod_{\alpha' \in A} B_{\alpha'}$ and $\#\{\alpha' \in A : B_{\alpha'} \neq X_{\alpha'}\} < \infty$. Then

$$\begin{aligned} \pi_\alpha(U) &= B_\alpha \\ &\in \mathcal{T}_\alpha \end{aligned}$$

Since $U \in \bigotimes_{\alpha \in A} \mathcal{T}_\alpha$ is arbitrary, we have that for each $U \in \bigotimes_{\alpha \in A} \mathcal{T}_\alpha$, $\pi_\alpha(U) \in \mathcal{T}_\alpha$. Exercise 4.2.0.22 implies that π_α is open. Since $\alpha \in A$ is arbitrary, we have that for each $\alpha \in A$, π_α is open. □

Exercise 4.6.1.4. Let $(X_\alpha, \mathcal{T}_\alpha)_{\alpha \in A}$ be a collection of topological spaces, $x \in \prod_{\alpha \in A} X_\alpha$ and for each $\alpha \in A$, $\mathcal{B}_{x_\alpha} \subset \mathcal{T}_\alpha$. Suppose that for each $\alpha \in A$, $\mathcal{B}_{x_\alpha}$ is a local basis for $\mathcal{T}_\alpha$ at x_α . Define

$$\mathcal{B}_x = \left\{ \prod_{\alpha \in A} U_\alpha : [\text{for each } \alpha \in A, U_\alpha \in \mathcal{T}_\alpha \text{ and } U_\alpha \neq X_\alpha \text{ implies that } U_\alpha \in \mathcal{B}_{x_\alpha}] \text{ and } \#\{\alpha \in A : U_\alpha \neq X_\alpha\} < \infty \right\}$$

Then $\mathcal{B}_x$ is a local basis for $\bigotimes_{\alpha \in A} \mathcal{T}_\alpha$ at x .

Proof. Set $X = \prod_{\alpha \in A} X_\alpha$ and $\mathcal{T} = \bigotimes_{\alpha \in A} \mathcal{T}_\alpha$.

1. By construction, for each $V \in \mathcal{B}_x$, $x \in V$.
2. Let $U \in \mathcal{T}$. Suppose that $x \in U$. Set

$$\mathcal{B} = \left\{ \prod_{\alpha \in A} B_\alpha : \text{for each } \alpha \in A, B_\alpha \in \mathcal{T}_\alpha \text{ and } \#\{\alpha \in A : B_\alpha \neq X_\alpha\} < \infty \right\}$$

The previous exercise implies that $\mathcal{B}$ is a basis for $\bigotimes_{\alpha \in A} \mathcal{T}_\alpha$. Thus for each $\alpha \in A$, there exists $B_\alpha \in \mathcal{T}_\alpha$ such that $\#\{\alpha \in A : B_\alpha \neq X_\alpha\} < \infty$ and $x \in \prod_{\alpha \in A} B_\alpha \subset U$. Set $J = \{\alpha \in A : B_\alpha \neq X_\alpha\}$. Since $x \in \prod_{\alpha \in A} B_\alpha$, for each $\alpha \in A$, $x_\alpha \in B_\alpha$. Since for each $\alpha \in A$, $\mathcal{B}_{x_\alpha}$ is a local basis for $\mathcal{T}_\alpha$ at x_α , the axiom of choice implies that there exists $(U_\alpha)_{\alpha \in A} \in \prod_{\alpha \in A} \mathcal{T}_\alpha$ such that for each if $\alpha \in J$, $U_\alpha \in \mathcal{B}_{x_\alpha}$ and $x_\alpha \in U_\alpha \subset B_\alpha$ and for each $\alpha \in J^c$, $U_\alpha = X_\alpha$. By definition, $\prod_{\alpha \in A} U_\alpha \in \mathcal{B}_x$. By construction,

$$\begin{aligned} x &\in \prod_{\alpha \in A} U_\alpha \\ &\subset \prod_{\alpha \in A} B_\alpha \\ &\subset U \end{aligned}$$

Since $U \in \mathcal{T}$ such that $x \in U$ is arbitrary, we have that $\mathcal{B}_x$ is a local basis for $\mathcal{T}$ at x . □

Exercise 4.6.1.5. Let $(X_j, \mathcal{T}_j)_{j=1}^n$ be a collection of topological spaces. Set

$$\mathcal{B} = \left\{ \prod_{j=1}^n A_j : \text{for each } j \in \{1, \dots, n\}, A_j \in \mathcal{T}_j \right\}$$

Then $\mathcal{B}$ is a basis for the product topology on $\prod_{j=1}^n X_j$.

Proof. Clear by previous exercise. □

Exercise 4.6.1.6. Basis for Product Topology:

Let $(X_\alpha, \mathcal{T}_\alpha)_{\alpha \in A}$ be a collection of topological spaces and for each $\alpha \in A$, $\mathcal{B}_\alpha$ a basis for $\mathcal{T}_\alpha$. Set $X = \prod_{\alpha \in A} X_\alpha$ and denote the product topology on X by $\mathcal{T}_X$. Set

$$\mathcal{B} = \left\{ \prod_{\alpha \in A} U_\alpha : \text{there exists } J \subset A \text{ such that } \#J < \infty, \right. \\ \left. \text{for each } \alpha \in J, U_\alpha \in \mathcal{B}_\alpha \text{ and for each } \alpha \in J^c, U_\alpha = X_\alpha \right\}$$

Then $\mathcal{B}$ is a basis for $\mathcal{T}_X$.

Proof. Set

$$\mathcal{B}' = \left\{ \prod_{\alpha \in A} V_\alpha : \text{for each } \alpha \in A, V_\alpha \in \mathcal{T}_\alpha \text{ and } \#\{\alpha \in A : V_\alpha \neq X_\alpha\} < \infty \right\}$$

The previous exercise implies that $\mathcal{B}'$ is a basis for $\mathcal{T}_X$. Then $\mathcal{B} \subset \mathcal{B}' \subset \mathcal{T}_X$. Let $V \in \mathcal{T}$ and $x \in V$. Write $x = (x_\alpha)_{\alpha \in A}$. Since $\mathcal{B}'$ is a basis for $\mathcal{T}_X$, for each $\alpha \in A$, there exists $V_\alpha \in \mathcal{T}_\alpha$ such that for finitely many $\alpha \in A$, $V_\alpha \neq X_\alpha$ and $x \in \prod_{\alpha \in A} V_\alpha \subset V$. Define $J \subset A$ by $J = \{\alpha \in A : V_\alpha \neq X_\alpha\}$. Then $\#J < \infty$. Let $\alpha \in J$. Then $x_\alpha \in V_\alpha$. Since $\mathcal{B}_\alpha$ is a basis for $\mathcal{T}_\alpha$, there exists $U'_\alpha \in \mathcal{B}_\alpha$ such that $x_\alpha \in U'_\alpha \subset V_\alpha$. For $\alpha \in A$, define $U_\alpha \in \mathcal{T}_\alpha$ by

$$U_\alpha = \begin{cases} U'_\alpha & \alpha \in J \\ X_\alpha & \alpha \in J^c \end{cases}$$

Set $U = \prod_{\alpha \in A} U_\alpha$. Then $U \in \mathcal{B}$ and

$$\begin{aligned} x &\in U \\ &= \prod_{\alpha \in A} U_\alpha \\ &\subset \prod_{\alpha \in A} V_\alpha \\ &\subset V \end{aligned}$$

Hence $\mathcal{B}$ is a basis for $\mathcal{T}_X$. □

4.6.2 Characteristics of the Product Topology

Exercise 4.6.2.1. Let $(X_\alpha, \mathcal{T}_\alpha)_{\alpha \in A}$ be a collection of topological spaces and for each $\alpha \in A$, $E_\alpha \subset X_\alpha$. Then

$$\text{cl} \left(\prod_{\alpha \in A} E_\alpha \right) = \prod_{\alpha \in A} \text{cl } E_\alpha$$

Hint: Exercise 4.1.0.29

Proof. Since for each $\alpha \in A$, $\pi_\alpha : \prod_{\alpha \in A} X_\alpha \rightarrow X_\alpha$ is continuous and $\text{cl } E_\alpha$ is closed, we have that for each $\alpha \in A$, $\pi_\alpha^{-1}(\text{cl } E_\alpha)$ is closed and thus

$$\prod_{\alpha \in A} \text{cl } E_\alpha = \bigcap_{\alpha \in A} \pi_\alpha^{-1}(\text{cl } E_\alpha)$$

is closed. Since for each $\alpha \in A$, $E_\alpha \subset \text{cl } E_\alpha$, we have that

$$\prod_{\alpha \in A} E_\alpha \subset \prod_{\alpha \in A} \text{cl } E_\alpha$$

which implies that

$$\text{cl} \left(\prod_{\alpha \in A} E_\alpha \right) \subset \prod_{\alpha \in A} \text{cl } E_\alpha$$

Conversely, let $x \in \prod_{\alpha \in A} \text{cl } E_\alpha$ and $U \in \mathcal{N}(x)$. Suppose that U is open. Set

$$\mathcal{B} = \left\{ \prod_{\alpha \in A} B_\alpha : \text{for each } \alpha \in A, B_\alpha \in \mathcal{T}_\alpha \text{ and } \#\{\alpha \in A : B_\alpha \neq X_\alpha\} < \infty \right\}$$

A previous exercise implies that $\mathcal{B}$ is a basis for the product topology on $\prod_{\alpha \in A} X_\alpha$. So for each $\alpha \in A$, there exists $U_\alpha \in \mathcal{T}_\alpha$ such that $\#\{\alpha \in A : U_\alpha \neq X_\alpha\} < \infty$ and $x \in \prod_{\alpha \in A} U_\alpha \subset U$. Then for each $\alpha \in A$, $x_\alpha \in \text{cl } E_\alpha \cap U_\alpha$. Let $\alpha \in A$. Since $x_\alpha \in \text{cl } E_\alpha$ and

$U_\alpha \in \mathcal{N}(x_\alpha)$ is an open neighborhood of x_α , Exercise 4.1.0.29 implies that $E_\alpha \cap U_\alpha \neq \emptyset$. Since $\alpha \in A$ is arbitrary, for each $\alpha \in A$, $E_\alpha \cap U_\alpha \neq \emptyset$. The axiom of choice implies that there exists

$$\begin{aligned} y &\in \prod_{\alpha \in A} (E_\alpha \cap U_\alpha) \\ &= \left(\prod_{\alpha \in A} E_\alpha \right) \cap \prod_{\alpha \in A} U_\alpha \\ &\subset \left(\prod_{\alpha \in A} E_\alpha \right) \cap U \end{aligned}$$

Hence $\left(\prod_{\alpha \in A} E_\alpha \right) \cap U \neq \emptyset$. Since $U \in \mathcal{N}(x)$ is an arbitrary open neighborhood of x , Exercise 4.1.0.29 implies that $x \in \text{cl } \prod_{\alpha \in A} E_\alpha$.

Since $x \in \prod_{\alpha \in A} \text{cl } E_\alpha$ is arbitrary, $\prod_{\alpha \in A} \text{cl } E_\alpha \subset \text{cl } \prod_{\alpha \in A} E_\alpha$.

Hence $\prod_{\alpha \in A} \text{cl } E_\alpha = \text{cl } \prod_{\alpha \in A} E_\alpha$ □

Exercise 4.6.2.2. Let $(X_\alpha, \mathcal{T}_\alpha)_{\alpha \in A}$ be a collection of topological spaces, $(a_\gamma)_{\gamma \in \Gamma} \subset \prod_{\alpha \in A} X_\alpha$ a net and $a \in \prod_{\alpha \in A} X_\alpha$. Then $a_\gamma \rightarrow a$ in $(\prod_{\alpha \in A} X_\alpha, \bigotimes_{\alpha \in A} \mathcal{T}_\alpha)$ iff for each $\alpha \in A$, $\pi_\alpha(a_\gamma) \xrightarrow{\gamma} \pi_\alpha(a)$ in $(X_\alpha, \mathcal{T}_\alpha)$.

Proof. Clear by Exercise 4.3.2.15. □

Exercise 4.6.2.3. Define $(e_n)_{n \in \mathbb{N}} \subset \mathbb{R}^{\mathbb{N}}$ by $\pi_k(e_n) := \delta_{n,k}$. Then $e_n \rightarrow 0$ in $(\mathbb{R}^{\mathbb{N}}, \mathcal{T}_{\mathbb{R}}^{\otimes \mathbb{N}})$.

Proof. Let $k \in \mathbb{N}$. Then

$$\begin{aligned} \pi_k(e_n) &= \delta_{n,k} \\ &\xrightarrow{n} 0. \end{aligned}$$

Since $k \in \mathbb{N}$ is arbitrary, we have that for each $k \in \mathbb{N}$, $\pi_k(e_n) \rightarrow 0$. Exercise 4.6.2.2 implies that $e_n \rightarrow 0$ in $(\mathbb{R}^{\mathbb{N}}, \mathcal{T}_{\mathbb{R}}^{\otimes \mathbb{N}})$. □

4.6.3 Maps and the Product Topology

Exercise 4.6.3.1. Let X be a topological space, $(Y_\alpha, \mathcal{T}_\alpha)_{\alpha \in A}$ a collection of topological spaces and $f : X \rightarrow \prod_{\alpha \in A} Y_\alpha$. Then f is continuous iff for each $\alpha \in A$, $\pi_\alpha \circ f$ is continuous.

Proof. Immediate by a Exercise 4.2.0.14. □

Exercise 4.6.3.2. Let $(X_\alpha, \mathcal{T}_\alpha)_{\alpha \in A}$ and $(Y_\alpha, \mathcal{S}_\alpha)_{\alpha \in A}$ be collections of topological spaces and $(f_\alpha)_{\alpha \in A} \in \prod_{\alpha \in A} Y_\alpha^{X_\alpha}$, i.e. for each $\alpha \in A$, $f_\alpha : X_\alpha \rightarrow Y_\alpha$. If for each $\alpha \in A$, f_α is continuous, then $\prod_{\alpha \in A} f_\alpha$ is continuous.

Proof. Set $X := \prod_{\alpha \in A} X_\alpha$, $Y := \prod_{\alpha \in A} Y_\alpha$ and define $f : X \rightarrow Y$ by $f := \prod_{\alpha \in A} f_\alpha$. Denote the α -th projection maps on X and Y by π_α^X and π_α^Y respectively. Set $\mathcal{T} := \bigotimes_{\alpha \in A} \mathcal{T}_\alpha$ and $\mathcal{S} := \bigotimes_{\alpha \in A} \mathcal{S}_\alpha$. Suppose that for each $\alpha \in A$, f_α is continuous. Exercise 1.5.0.7 implies that for each $\alpha \in A$, $\pi_\alpha^Y \circ f = f_\alpha \circ \pi_\alpha^X$. Let $\alpha \in A$. Then $f_\alpha \circ \pi_\alpha^X$ is continuous. Hence $\pi_\alpha^Y \circ f$ is continuous. Since $\alpha \in A$ is arbitrary, we have that for each $\alpha \in A$, $\pi_\alpha^Y \circ f$ is continuous. Exercise 4.6.3.1 implies that f is continuous. □

Exercise 4.6.3.3. Let $(X_\alpha, \mathcal{T}_\alpha)_{\alpha \in A}$ and $(Y_\alpha, \mathcal{S}_\alpha)_{\alpha \in A}$ be collections of topological spaces and $(f_\alpha)_{\alpha \in A} \in \prod_{\alpha \in A} Y_\alpha^{X_\alpha}$, i.e. for each $\alpha \in A$, $f_\alpha : X_\alpha \rightarrow Y_\alpha$. If $\#\{\alpha \in A : f_\alpha \text{ is not surjective}\} < \infty$ and for each $\alpha \in A$, f_α is open, then $\prod_{\alpha \in A} f_\alpha$ is open.

Proof. Set $X := \prod_{\alpha \in A} X_\alpha$, $Y := \prod_{\alpha \in A} Y_\alpha$ and define $f : X \rightarrow Y$ by $f := \prod_{\alpha \in A} f_\alpha$. Denote the α -th projection maps on X and Y by π_α^X and π_α^Y respectively. Set $\mathcal{T} := \bigotimes_{\alpha \in A} \mathcal{T}_\alpha$ and $\mathcal{S} := \bigotimes_{\alpha \in A} \mathcal{S}_\alpha$. Suppose that $\#\{\alpha \in A : f_\alpha \text{ is not surjective}\} < \infty$ and for each $\alpha \in A$, f_α is open. Set

$$\mathcal{B}_X := \left\{ \prod_{\alpha \in A} U_\alpha : \text{for each } \alpha \in A, U_\alpha \in \mathcal{T}_\alpha \text{ and } \#\{\alpha \in A : U_\alpha \neq X_\alpha\} < \infty \right\}$$

$$\mathcal{B}_Y := \left\{ \prod_{\alpha \in A} V_\alpha : \text{for each } \alpha \in A, V_\alpha \in \mathcal{S}_\alpha \text{ and } \#\{\alpha \in A : V_\alpha \neq Y_\alpha\} < \infty \right\}$$

A previous exercise implies that $\mathcal{B}_X$ is a basis for $\mathcal{T}$ and $\mathcal{B}_Y$ is a basis for $\mathcal{S}$. Let $U \in \mathcal{B}_X$. Then for each $\alpha \in A$ there exist $U_\alpha \in \mathcal{T}_\alpha$ such that $U = \prod_{\alpha \in A} U_\alpha$. Define

- $B_1 := \{\alpha \in A : f_\alpha \text{ is not surjective}\}$
- $B_2 := \{\alpha \in A : U_\alpha \neq X_\alpha\}$
- $B_3 := \{\alpha \in A : f_\alpha(U_\alpha) \neq Y_\alpha\}$

Let $\alpha \in A$. Suppose that $\alpha \in B_1^c \cap B_2^c$. Then f_α is surjective and $U_\alpha = X_\alpha$. Thus $f_\alpha(U_\alpha) = Y_\alpha$ and $\alpha \in B_3^c$. Therefore if $\alpha \in B_3$, then $\alpha \in B_1 \cup B_2$. Since $\alpha \in A$ is arbitrary, $B_3 \subset B_1 \cup B_2$. By assumption, $\#B_1 < \infty$ and $\#B_2 < \infty$. Thus

$$\begin{aligned} \#B_3 &\leq \#(B_1 \cup B_2) \\ &\leq \#B_1 + \#B_2 \\ &< \infty \end{aligned}$$

Since for each $\alpha \in A$, f_α is open, we have that for each $\alpha \in A$, $f_\alpha(U_\alpha) \in \mathcal{S}_\alpha$. Thus

$$\begin{aligned} f\left(\prod_{\alpha \in A} U_\alpha\right) &= \prod_{\alpha \in A} f_\alpha(U_\alpha) \\ &\in \mathcal{B}_Y \\ &\subset \mathcal{S} \end{aligned}$$

Since $U \in \mathcal{B}_X$ is arbitrary, we have that for each $U \in \mathcal{B}_X$, $f(U) \in \mathcal{S}$. Since $\mathcal{B}_X$ is a basis for $\mathcal{T}$, an exercise about open maps in the section on continuous maps implies that f is open. $\square$

Exercise 4.6.3.4. Let $(X_\alpha, \mathcal{T}_\alpha)_{\alpha \in A}$ and $(Y_\alpha, \mathcal{S}_\alpha)_{\alpha \in A}$ be collections of topological spaces and for each $\alpha \in A$, $U_\alpha \subset X_\alpha$, $V_\alpha \subset Y_\alpha$ and $f_\alpha : U_\alpha \rightarrow V_\alpha$. If for each $\alpha \in A$, f_α is $(\mathcal{T}_\alpha \cap U_\alpha)$ -continuous, then $\prod_{\alpha \in A} f_\alpha$ is $\left(\left[\bigotimes_{\alpha \in A} \mathcal{T}_\alpha\right] \cap \left[\prod_{\alpha \in A} U_\alpha\right], \left[\bigotimes_{\alpha \in A} \mathcal{S}_\alpha\right] \cap \left[\prod_{\alpha \in A} V_\alpha\right]\right)$ -continuous.

Proof. Denote the α -th projection maps on X and Y by π_α^X and π_α^Y respectively. Let $(x_\gamma)_{\gamma \in \Gamma} \subset \prod_{\alpha \in A} U_\alpha$ and $x \in \prod_{\alpha \in A} U_\alpha$. Suppose

that $x_\gamma \rightarrow x$ in $\left(\prod_{\alpha \in A} U_\alpha, \left[\bigotimes_{\alpha \in A} \mathcal{T}_\alpha\right] \cap \left[\prod_{\alpha \in A} U_\alpha\right]\right)$. An exercise in the section on the subspace topology implies that $x_\gamma \rightarrow x$ in $\left(\prod_{\alpha \in A} X_\alpha, \bigotimes_{\alpha \in A} \mathcal{T}_\alpha\right)$. Let $\alpha \in A$. Since π_α^X is $\left(\bigotimes_{\beta \in A} \mathcal{T}_\beta, \mathcal{T}_\alpha\right)$ -continuous, we have that $\pi_\alpha^X(x_\gamma) \rightarrow \pi_\alpha^X(x)$ in $(X_\alpha, \mathcal{T}_\alpha)$. Another application of the same exercise implies that $\pi_\alpha^X(x_\gamma) \rightarrow \pi_\alpha^X(x)$ in $(U_\alpha, \mathcal{T}_\alpha \cap U_\alpha)$. Since f_α is $\left(\mathcal{T}_\alpha \cap U_\alpha, \mathcal{S}_\alpha \cap V_\alpha\right)$ -continuous,

$$\begin{aligned} \pi_\alpha^Y \circ \left[\prod_{\alpha \in A} f_\alpha\right](x_\gamma) &= f_\alpha \circ \pi_\alpha^X(x_\gamma) \\ &\rightarrow f_\alpha \circ \pi_\alpha^X(x) \\ &= \pi_\alpha^Y \circ \left[\prod_{\alpha \in A} f_\alpha\right](x) \text{ in } (V_\alpha, \mathcal{S}_\alpha \cap V_\alpha) \end{aligned}$$

Another application of the exercise implies that $\pi_\alpha^Y \circ \left[\prod_{\alpha \in A} f_\alpha\right](x_\gamma) \rightarrow \pi_\alpha^Y \circ \left[\prod_{\alpha \in A} f_\alpha\right](x)$ in $(Y_\alpha, \mathcal{S}_\alpha)$. Since $(x_\gamma)_{\gamma \in \Gamma} \subset \prod_{\alpha \in A} U_\alpha$ and $x \in \prod_{\alpha \in A} U_\alpha$ with $x_\gamma \rightarrow x$ in $\left(\prod_{\alpha \in A} U_\alpha, \left[\bigotimes_{\alpha \in A} \mathcal{T}_\alpha\right] \cap \left[\prod_{\alpha \in A} U_\alpha\right]\right)$ is arbitrary, we have that $\prod_{\alpha \in A} f_\alpha$ is $\left(\left[\bigotimes_{\alpha \in A} \mathcal{T}_\alpha\right] \cap \left[\prod_{\alpha \in A} U_\alpha\right], \bigotimes_{\alpha \in A} \mathcal{S}_\alpha\right)$ -continuous. Another application of the exercise implies that $\prod_{\alpha \in A} f_\alpha$ is $\left(\left[\bigotimes_{\alpha \in A} \mathcal{T}_\alpha\right] \cap \left[\prod_{\alpha \in A} U_\alpha\right], \left[\bigotimes_{\alpha \in A} \mathcal{S}_\alpha\right] \cap \left[\prod_{\alpha \in A} V_\alpha\right]\right)$ -continuous. $\square$

Exercise 4.6.3.5. Let $(X, \mathcal{T}_X)$ be a topological space, $(Y_\alpha, \mathcal{T}_{Y_\alpha})_{\alpha \in A}$ a collection of topological spaces, $(f_\alpha)_{\alpha \in A} \in \prod_{\alpha \in A} Y_\alpha^X$. If for each $\alpha \in A$, f_α is $(\mathcal{T}_X, \mathcal{T}_{Y_\alpha})$ -continuous, then $(f_\alpha)_{\alpha \in A}$ is $(\mathcal{T}_X, \bigotimes_{\alpha \in A} \mathcal{T}_{Y_\alpha})$ -continuous.

Proof. Suppose that for each $\alpha \in A$, f_α is $(\mathcal{T}_X, \mathcal{T}_{Y_\alpha})$ -continuous. Set $f := (f_\alpha)_{\alpha \in A}$. Since for each $\alpha \in A$, $\pi_\alpha \circ f = f_\alpha$, Exercise 4.6.3.1 implies that f is $(\mathcal{T}_X, \bigotimes_{\alpha \in A} \mathcal{T}_{Y_\alpha})$ -continuous. $\square$

Definition 4.6.3.6. define slice maps and then define the below sets in terms of them in the set theory section.

Exercise 4.6.3.7. Let X and Y be topological spaces and $U \subset X \times Y$ open. Then for each $(x_0, y_0) \in U$, U^{x_0} and U^{y_0} are open.

Proof. Let $(x_0, y_0) \in U$. Define $\phi : X \rightarrow X \times Y$ by $\phi(x) = (x, y_0)$. Since $\pi_X \circ \phi = \text{id}_X$ and $\pi_Y \circ \phi$ is constant, $\pi_X \circ \phi$ and $\pi_Y \circ \phi$ are continuous. Therefore, ϕ is continuous. Then U^{y_0} is open since U is open and $\phi^{-1}(U) = U^{y_0}$. Similarly, U_{x_0} is open. $\square$

Exercise 4.6.3.8. Let X, Y and Z be topological spaces, $U \subset X \times Y$ open and $f : U \rightarrow Z$. Equip U with the subspace topology. Suppose that f is continuous. Let $(x_0, y_0) \in U$. Equip U_{x_0} and U^{y_0} with the subspace topology. Then $f_{x_0} : U_{x_0} \rightarrow Z$ and $f^{y_0} : U^{y_0} \rightarrow Z$ are continuous.

Proof. Let $(x_0, y_0) \in U$. Let $V \subset Z$. Suppose that V is open. Continuity of f implies that $f^{-1}(V)$ is open in U . Since U is open in $X \times Y$, $f^{-1}(V)$ is open in $X \times Y$. A previous exercise in the section on product sets implies that $(f^{y_0})^{-1}(V) = (f^{-1}(V))^{y_0}$. The previous exercise implies that $(f^{-1}(V))^{y_0}$ is open in X . So $(f^{y_0})^{-1}(V)$ is open in X . Since $(f^{y_0})^{-1}(V) \subset U^{y_0}$, $(f^{y_0})^{-1}(V)$ is open in U^{y_0} . Thus $f^{y_0} : U^{y_0} \rightarrow Z$ is continuous. Similarly, $f_{x_0} : U_{x_0} \rightarrow Z$ is continuous. $\square$

4.7 Coproduct Topology

Definition 4.7.0.1. Let $(X_\alpha, \mathcal{T}_\alpha)_{\alpha \in A}$ be a collection of topological spaces. We define the **coproduct topology** on $\coprod_{\alpha \in A} X_\alpha$, denoted

$\bigoplus_{\alpha \in A} \mathcal{T}_\alpha$, by

$$\bigoplus_{\alpha \in A} \mathcal{T}_\alpha = \tau(\iota_\alpha : \alpha \in A)$$

i.e. $\bigoplus_{\alpha \in A} \mathcal{T}_\alpha$ is the final topology on $\coprod_{\alpha \in A} X_\alpha$ generated by the embedding maps $(\iota_\alpha)_{\alpha \in A}$.

Exercise 4.7.0.2. Let $(X_\alpha, \mathcal{T}_\alpha)_{\alpha \in A}$ be a collection of topological spaces. Then

$$\bigoplus_{\alpha \in A} \mathcal{T}_\alpha = \left\{ V \subset \coprod_{\alpha \in A} X_\alpha : \text{for each } \alpha \in A, \iota_\alpha^{-1}(V) \in \mathcal{T}_\alpha \right\}$$

Proof. Set $X := \coprod_{\alpha \in A} X_\alpha$ and $\mathcal{T} := \left\{ V \subset \coprod_{\alpha \in A} X_\alpha : \text{for each } \alpha \in A, \iota_\alpha^{-1}(V) \in \mathcal{T}_\alpha \right\}$.

- 1. Clearly $\emptyset, X \in \mathcal{T}$.
- 2. Let $(U_\gamma)_{\gamma \in \Gamma} \subset \mathcal{T}$. Then by definition, for each $\gamma \in \Gamma$ and $\alpha \in A$, $\iota_\alpha^{-1}(U_\gamma) \in \mathcal{T}_\alpha$. Hence

$$\begin{aligned} \iota_\alpha^{-1}\left(\bigcup_{\gamma \in \Gamma} U_\gamma\right) &= \bigcup_{\gamma \in \Gamma} \iota_\alpha^{-1}(U_\gamma) \\ &\in \mathcal{T}_\alpha \end{aligned}$$

- 3. Let $(U_j)_{j=1}^n \subset \mathcal{T}$. Then by definition, for each $\alpha \in A$ and $j \in [n]$, $\iota_\alpha^{-1}(U_j) \in \mathcal{T}_\alpha$. Hence

$$\begin{aligned} \iota_\alpha^{-1}\left(\bigcap_{j=1}^n U_j\right) &= \bigcap_{j=1}^n \iota_\alpha^{-1}(U_j) \\ &\in \mathcal{T}_\alpha \end{aligned}$$

So $\mathcal{T}$ is a topology on X .

- Since $\mathcal{T}$ is a topology on X , we have that $\mathcal{T} = \tau_X(\mathcal{T})$. By Definition 4.2.0.16,

$$\begin{aligned} \mathcal{T} &= \tau_X(\mathcal{T}) \\ &= \tau_X(\iota_\alpha : \alpha \in A) \\ &= \bigoplus_{\alpha \in A} \mathcal{T}_\alpha. \end{aligned}$$

□

Exercise 4.7.0.3. Let $(X_\alpha, \mathcal{T}_\alpha)_{\alpha \in A}$ be a collection of measurable spaces. Then

$$\bigoplus_{\alpha \in A} \mathcal{T}_\alpha = \left\{ \coprod_{\alpha \in A} B_\alpha : B_\alpha \in \mathcal{T}_\alpha \right\}$$

Proof. Set

- $\mathcal{F} = \{V \subset \coprod_{\alpha \in A} X_\alpha : \text{for each } \alpha \in A, \iota_\alpha^{-1}(V) \in \mathcal{T}_\alpha\}$
- $\mathcal{G} = \left\{ \coprod_{\alpha \in A} B_\alpha : \text{for each } \alpha \in A, B_\alpha \in \mathcal{T}_\alpha \right\}$

Let $V \in \mathcal{G}$. Then for each $\alpha \in A$, there exists $B_\alpha \in \mathcal{T}_\alpha$ such that $V = \prod_{\alpha \in A} B_\alpha$. Therefore, for each $\alpha \in A$,

$$\begin{aligned}\iota_\alpha^{-1}(V) &= \iota_\alpha^{-1}\left(\prod_{\alpha \in A} B_\alpha\right) \\ &= B_\alpha \\ &\in \mathcal{T}_\alpha\end{aligned}$$

Hence $V \in \mathcal{F}$. Since $V \in \mathcal{G}$ is arbitrary, $\mathcal{G} \subset \mathcal{F}$.

Conversely, let $V \in \mathcal{F}$. Then for each $\alpha \in A$, $\iota_\alpha^{-1}(V) \in \mathcal{T}_\alpha$. For each $\alpha \in A$, define $B_\alpha \in \mathcal{T}_\alpha$ by $B_\alpha = \iota_\alpha^{-1}(V)$. Then

$$\begin{aligned}V &= \prod_{\alpha \in A} B_\alpha \\ &\in \mathcal{G}\end{aligned}$$

Since $V \in \mathcal{F}$ is arbitrary, $\mathcal{F} \subset \mathcal{G}$. The previous exercise implies that

$$\begin{aligned}\mathcal{G} &= \mathcal{F} \\ &= \bigoplus_{\alpha \in A} \mathcal{T}_\alpha\end{aligned}$$

□

Exercise 4.7.0.4. Let $(X_\alpha, \mathcal{T}_\alpha)_{\alpha \in A}$ be a collection of topological spaces. Then for each $\alpha \in A$, $\iota_\alpha : X_\alpha \rightarrow \prod_{\alpha' \in A} X_{\alpha'}$ is $\left(\mathcal{T}_\alpha, \bigoplus_{\alpha' \in A} \mathcal{T}_{\alpha'}\right)$ -open.

Proof. Let $\alpha, \beta \in A$ and $U \in \mathcal{T}_\alpha$. Then

$$\begin{aligned}\iota_\beta^{-1}(\iota_\alpha(U)) &= \iota_\beta^{-1}(\{\alpha\} \times U) \\ &= \begin{cases} U, & \beta = \alpha \\ \emptyset, & \beta \neq \alpha \end{cases} \\ &\in \mathcal{T}_\beta\end{aligned}$$

Since $\beta \in A$ is arbitrary, we have that for each $\beta \in A$, $\iota_\beta^{-1}(\iota_\alpha(U)) \in \mathcal{T}_\beta$. Exercise 4.7.0.2 implies that $\iota_\alpha(U) \in \bigoplus_{\beta \in A} \mathcal{T}_\beta$. Since $U \in \mathcal{T}_\alpha$

is arbitrary, we have that $\left(\mathcal{T}_\alpha, \bigoplus_{\alpha' \in A} \mathcal{T}_{\alpha'}\right)$ -open. □

Exercise 4.7.0.5. Let $(X_\alpha, \mathcal{T}_\alpha)_{\alpha \in A}$ be a collection of topological spaces. Then for each $\alpha \in A$, $\iota_\alpha : X_\alpha \rightarrow \prod_{\alpha' \in A} X_{\alpha'}$ is $\left(\mathcal{T}_\alpha, \bigoplus_{\alpha' \in A} \mathcal{T}_{\alpha'}\right)$ -closed.

Proof. Let $\alpha_0 \in A$ and $C \subset X_{\alpha_0}$. Suppose that C is closed in $(X_{\alpha_0}, \mathcal{T}_{\alpha_0})$. Define $(B_\alpha)_{\alpha \in A} \in \prod_{\alpha \in A} \mathcal{T}_\alpha$ by

$$B_\alpha := \begin{cases} C^c, & \alpha = \alpha_0 \\ X_\alpha, & \alpha \neq \alpha_0. \end{cases}$$

Then

$$\begin{aligned}[\iota_{\alpha_0}(C)]^c &= [\{\alpha_0\} \times C]^c \\ &= \prod_{\alpha \in A} B_\alpha \\ &\in \bigoplus_{\alpha \in A} \mathcal{T}_\alpha.\end{aligned}$$

Hence $\iota_{\alpha_0}(C)$ is closed in $(\coprod_{\alpha \in A} X_\alpha, \bigoplus_{\alpha \in A} \mathcal{T}_\alpha)$. Since $C \subset X_{\alpha_0}$ with C closed in $(X_{\alpha_0}, \mathcal{T}_{\alpha_0})$ is arbitrary, we have that for each $C \subset X_{\alpha_0}$, C closed in $(X_{\alpha_0}, \mathcal{T}_{\alpha_0})$ implies that $\iota_{\alpha_0}(C)$ is closed in $(\coprod_{\alpha \in A} X_\alpha, \bigoplus_{\alpha \in A} \mathcal{T}_\alpha)$. Hence ι_{α_0} is $(\mathcal{T}_{\alpha_0}, \bigoplus_{\alpha' \in A} \mathcal{T}_{\alpha'})$ -closed. Since $\alpha_0 \in A$ is arbitrary, we have that for each $\alpha \in A$, ι_α is $(\mathcal{T}_\alpha, \bigoplus_{\alpha' \in A} \mathcal{T}_{\alpha'})$ -closed. $\square$

Exercise 4.7.0.6. Let $(X_\alpha, \mathcal{T}_\alpha)_{\alpha \in A}$ be a collection of topological spaces and for each $\alpha \in A$, $\mathcal{B}_\alpha \subset \mathcal{T}_\alpha$. Suppose that for each $\alpha \in A$, $\mathcal{B}_\alpha$ is a basis for $\mathcal{T}_\alpha$. Then $\{\iota_\alpha(U) : \alpha \in A \text{ and } U \in \mathcal{B}_\alpha\}$ is a basis for $\bigoplus_{\alpha \in A} \mathcal{T}_\alpha$.

Proof. Set $\mathcal{B} := \{\iota_\alpha(U) : \alpha \in A \text{ and } U \in \mathcal{B}_\alpha\}$ and $\mathcal{T} := \bigoplus_{\alpha \in A} \mathcal{T}_\alpha$. Exercise 4.7.0.4 implies that $\mathcal{B} \subset \mathcal{T}$. Let $W \in \mathcal{T}$ and $(\alpha_0, x_0) \in W$. Set $U := \iota_{\alpha_0}^{-1}(W)$. By assumption, $x_0 \in U$. Since ι_{α_0} is $(\mathcal{T}_{\alpha_0}, \mathcal{T})$ -continuous, we have that $U \in \mathcal{T}_{\alpha_0}$. Exercise 4.7.0.4 implies that

$$\begin{aligned} \{\alpha_0\} \times U &= \iota_{\alpha_0}(U) \\ &\in \mathcal{T} \end{aligned}$$

By construction,

$$\begin{aligned} (\alpha_0, x_0) &\in \{\alpha_0\} \times U \\ &= \iota_{\alpha_0}(U) \\ &= \iota_{\alpha_0}(\iota_{\alpha_0}^{-1}(W)) \\ &\subset W. \end{aligned}$$

Since $W \in \mathcal{T}$ and $(\alpha_0, x_0) \in W$ are arbitrary, we have that for each $W \in \mathcal{T}$ and $(\alpha_0, x_0) \in W$, there exists $B \in \mathcal{B}$ such that $(\alpha_0, x_0) \in B \subset W$. Therefore $\mathcal{B}$ is a basis for $\mathcal{T}$. $\square$

Note 4.7.0.7. Let Γ be a directed set and $\gamma_0 \in \Gamma$. From Definition 1.3.3.2 we recall the γ_0 -tail operator $L_{\gamma_0} : X^\Gamma \rightarrow X^{[\gamma_0, \infty)}$.

Exercise 4.7.0.8. Let $(X_\alpha, \mathcal{T}_\alpha)_{\alpha \in A}$ be a collection of topological spaces, $(\alpha_\gamma, x_\gamma)_{\gamma \in \Gamma} \subset \prod_{\alpha \in A} X_\alpha$ a net and $(\alpha_0, x_0) \in \prod_{\alpha \in A} X_\alpha$. Then

$(\alpha_\gamma, x_\gamma) \rightarrow (\alpha_0, x_0)$ in $(\prod_{\alpha \in A} X_\alpha, \bigoplus_{\alpha \in A} \mathcal{T}_\alpha)$ iff there exists $\gamma_0 \in \Gamma$ such that

1. for each $\gamma \in \Gamma$, $\gamma \geq \gamma_0$ implies that $\alpha_\gamma = \alpha_0$ and $x_\gamma \in X_{\alpha_0}$
2. $[L_{\gamma_0}(x)]_\gamma \rightarrow x_0$ in $(X_{\alpha_0}, \mathcal{T}_{\alpha_0})$.

Proof. Set $X := \prod_{\alpha \in A} X_\alpha$ and $\mathcal{T} := \bigoplus_{\alpha \in A} \mathcal{T}_\alpha$.

• $(\implies)$:

1. Suppose that $(\alpha_\gamma, x_\gamma) \rightarrow (\alpha_0, x_0)$ in Exercise 4.7.0.4 implies that $\{\alpha_0\} \times X_{\alpha_0} \in \mathcal{N}(\alpha_0, x_0)$. Since $(\alpha_\gamma, x_\gamma) \rightarrow (\alpha_0, x_0)$, $(\alpha_\gamma, x_\gamma)_{\gamma \in \Gamma}$ is eventually in $\{\alpha_0\} \times X_{\alpha_0}$. Exercise 4.3.2.2 implies that there exists $\gamma_0 \in \Gamma$ such that $(\alpha_\gamma, x_\gamma)_{\gamma \in [\gamma_0, \infty)} \subset \{\alpha_0\} \times X_{\alpha_0}$. Hence for each $\gamma \in \Gamma$, $\gamma \geq \gamma_0$ implies that $\alpha_\gamma = \alpha_0$ and $(x_\gamma)_{\gamma \in [\gamma_0, \infty)} \subset X_{\alpha_0}$.
2. Define $(y_\gamma)_{\gamma \in [\gamma_0, \infty)} \subset X_{\alpha_0}$ by $y := L_{\gamma_0}(x)$. Let $U \in \mathcal{T}_{\alpha_0}$. Suppose that $x_0 \in U$. Exercise 4.7.0.4 implies that $\{\alpha_0\} \times U \in \mathcal{N}(\alpha_0, x_0)$. Since $(\alpha_\gamma, x_\gamma) \rightarrow (\alpha_0, x_0)$ in $(X, \mathcal{T})$, $(\alpha_\gamma, x_\gamma)_{\gamma \in \Gamma}$ is eventually in $\{\alpha_0\} \times U$. Exercise 4.3.2.2 implies that there exists $\gamma_1 \in \Gamma$ such that $(\alpha_\gamma, x_\gamma)_{\gamma \in [\gamma_1, \infty)} \subset \{\alpha_0\} \times U$. Since A is a directed set, there exists $\gamma_2 \in \Gamma$ such that $\gamma_0, \gamma_1 \leq \gamma_2$. Hence for each $\gamma \in [\gamma_2, \infty)$, $y_\gamma \in U$. So $(y_\gamma)_{\gamma \in [\gamma_0, \infty)}$ is eventually in U . Since $U \in \mathcal{T}_{\alpha_0}$ with $x_0 \in U$ is arbitrary, we have that $y_\gamma \rightarrow x_0$.

• $(\impliedby)$:

Suppose that there exists $\gamma_0 \in \Gamma$ such that

1. for each $\gamma \in \Gamma$, $\gamma \geq \gamma_0$ implies that $\alpha_\gamma = \alpha_0$ and $x_\gamma \in X_{\alpha_0}$
2. $[L_{\gamma_0}(x)]_\gamma \rightarrow x_0$ in $(X_{\alpha_0}, \mathcal{T}_{\alpha_0})$.

Let $W_0 \in \mathcal{N}(\alpha_0, x_0)$. Set $W := \text{Int } W_0$. Then $W \in \mathcal{T}$ and $(\alpha_0, x_0) \in W$. Exercise 4.7.0.6 implies that there exists $U \in \mathcal{T}_{\alpha_0}$ such that $(\alpha_0, x_0) \in \{\alpha_0\} \times U \subset W$. By assumption, there exists $\gamma_1 \in \Gamma$ such that for each $\gamma \in \Gamma$, $\gamma \geq \gamma_1$ implies that $\alpha_\gamma = \alpha_0$ and $x_\gamma \in X_{\alpha_0}$. Since $[L_{\gamma_1}(x)]_\gamma \rightarrow x_0$ in $(X_{\alpha_0}, \mathcal{T}_{\alpha_0})$ and $U \in \mathcal{T}_{\alpha_0}$, there exists $\gamma_2 \in [\gamma_1, \infty)$ such that for each $\gamma \in [\gamma_1, \infty)$, $\gamma \geq \gamma_2$ implies that $x_\gamma \in U$. Since Γ is a directed set, there exists $\gamma_0 \in \Gamma$ such that $\gamma \geq \gamma_1, \gamma_2$. Let $\gamma \in \Gamma$. Suppose that $\gamma \geq \gamma_0$. Then

$$\begin{aligned} (\alpha_\gamma, x_\gamma) &= (\alpha_0, x_\gamma) \\ &\in \{\alpha_0\} \times U \\ &\subset W \\ &\subset W_0 \end{aligned}$$

Hence $(\alpha_\gamma, x_\gamma)$ is eventually in W_0 . Since $W_0 \in \mathcal{N}(\alpha_0, x_0)$ is arbitrary, we have that for each $W_0 \in \mathcal{N}(\alpha_0, x_0)$, $(\alpha_\gamma, x_\gamma)$ is eventually in W_0 . Thus $(\alpha_\gamma, x_\gamma) \rightarrow (\alpha_0, x_0)$ in $(X, \mathcal{T})$. □

Exercise 4.7.0.9. Let $(X_\alpha, \mathcal{T}_\alpha)_{\alpha \in A}$ be a collection of topological spaces, $(Y, \mathcal{S})$ a topological space and $f : \prod_{\alpha \in A} X_\alpha \rightarrow Y$. Then f is

$\left(\bigoplus_{\alpha \in A} \mathcal{T}_\alpha, \mathcal{S} \right)$ -continuous iff for each $\alpha \in A$, $f \circ \iota_\alpha$ is $(\mathcal{T}_\alpha, \mathcal{S})$ -continuous.

Proof. Clear by Exercise 4.2.0.20. add more details □

Exercise 4.7.0.10. Let $(X_\alpha, \mathcal{T}_\alpha)_{\alpha \in A}$ and $(Y_\alpha, \mathcal{S}_\alpha)_{\alpha \in A}$ be collections of continuous spaces and $(f_\alpha)_{\alpha \in A} \in \prod_{\alpha \in A} Y_\alpha^{X_\alpha}$, i.e. for each $\alpha \in A$, $f_\alpha : X_\alpha \rightarrow Y_\alpha$. If for each $\alpha \in A$, f_α is $(\mathcal{T}_\alpha, \mathcal{S}_\alpha)$ -continuous, then $\prod_{\alpha \in A} f_\alpha$ is $(\bigoplus_{\alpha \in A} \mathcal{T}_\alpha, \bigoplus_{\alpha \in A} \mathcal{S}_\alpha)$ -continuous.

Proof. Set $X := \prod_{\alpha \in A} X_\alpha$, $Y := \prod_{\alpha \in A} Y_\alpha$, $\mathcal{T} := \bigoplus_{\alpha \in A} \mathcal{T}_\alpha$ and $\mathcal{S} := \bigoplus_{\alpha \in A} \mathcal{S}_\alpha$. Suppose that for each $\alpha \in A$, f_α is $(\mathcal{T}_\alpha, \mathcal{S}_\alpha)$ -continuous. Set $f = \prod_{\alpha \in A} f_\alpha$. Denote the α -th embedding maps on X and Y by ι_α^X and ι_α^Y respectively. Let $\alpha \in A$. Exercise 1.6.0.6 implies that $f \circ \iota_\alpha^X = \iota_\alpha^Y \circ f_\alpha$. Since $\iota_\alpha^Y \circ f_\alpha$ is $(\mathcal{T}_\alpha, \mathcal{S})$ -continuous, we have that $f \circ \iota_\alpha^X$ is $(\mathcal{T}_\alpha, \mathcal{S})$ -continuous. Since $\alpha \in A$ is arbitrary, we have that for each $\alpha \in A$, $f \circ \iota_\alpha^X$ is $(\mathcal{T}_\alpha, \mathcal{S})$ -continuous. Exercise 4.7.0.9 implies that f is $(\mathcal{T}, \mathcal{S})$ -continuous. □

Exercise 4.7.0.11. Let $(X, \mathcal{T})$ be a topological space and $(E_\alpha)_{\alpha \in A} \subset \mathcal{T}$. Define $\phi : \prod_{\alpha \in A} E_\alpha \rightarrow \bigcup_{\alpha \in A} E_\alpha$ by $\phi(\alpha, x) := x$. If $(E_\alpha)_{\alpha \in A}$ is disjoint, then ϕ is a $\left(\bigoplus_{\alpha \in A} [\mathcal{T} \cap E_\alpha], \mathcal{T} \cap \left[\bigcup_{\alpha \in A} E_\alpha \right] \right)$ -homeomorphism.

Proof. Suppose that $(E_\alpha)_{\alpha \in A}$ is disjoint.

• **(bijectivity) :**

– **(injectivity) :**

Let $(\alpha, x), (\beta, y) \in \prod_{\alpha \in A} E_\alpha$. Suppose that $\phi(\alpha, x) = \phi(\beta, y)$. Then $x = y$. Thus $x \in E_\alpha \cap E_\beta$ and therefore $E_\alpha \cap E_\beta \neq \emptyset$. Since $(E_{\alpha'})_{\alpha' \in A}$ is disjoint, we have that $\alpha = \beta$. Hence $(\alpha, x) = (\beta, y)$. Since $(\alpha, x), (\beta, y) \in \prod_{\alpha \in A} E_\alpha$ are arbitrary, we have that for each $(\alpha, x), (\beta, y) \in \prod_{\alpha \in A} E_\alpha$, $\phi(\alpha, x) = \phi(\beta, y)$ implies that $(\alpha, x) = (\beta, y)$. Thus ϕ is injective.

– **(surjectivity) :**

Let $x \in \bigcup_{\alpha \in A} E_\alpha$. Then there exists $\alpha \in A$ such that $x \in E_\alpha$. Then $(\alpha, x) \in \prod_{\alpha \in A} E_\alpha$ and $\phi(\alpha, x) = x$. Since $x \in \bigcup_{\alpha \in A} E_\alpha$ is arbitrary, we have that for each $x \in \bigcup_{\alpha \in A} E_\alpha$, there exists $a \in \prod_{\alpha \in A} E_\alpha$ such that $\phi(a) = x$. Hence ϕ is surjective.

So ϕ is a bijection.

• **(continuity) :**

For each $\alpha \in A$, define $\iota_{E_\alpha} : E_\alpha \rightarrow \bigcup_{\alpha \in A} E_\alpha$ by $\iota_{E_\alpha}(x) = x$.

- Let $\alpha \in A$. Since $\phi \circ \iota_\alpha = \iota_{E_\alpha}$ and ι_{E_α} is $\left(\mathcal{T} \cap E_\alpha, \mathcal{T} \cap \left[\bigcup_{\alpha \in A} E_\alpha\right]\right)$ -continuous (maybe give more details), we have that $\phi \circ \iota_\alpha$ is $\left(\mathcal{T} \cap E_\alpha, \mathcal{T} \cap \left[\bigcup_{\alpha \in A} E_\alpha\right]\right)$ -continuous. Since $\alpha \in A$ is arbitrary, we have that for each $\alpha \in A$, $\phi \circ \iota_\alpha$ is $\left(\mathcal{T} \cap E_\alpha, \mathcal{T} \cap \left[\bigcup_{\alpha \in A} E_\alpha\right]\right)$ -continuous. Exercise 4.7.0.9 implies that ϕ is $\left(\bigoplus_{\alpha \in A} (\mathcal{T} \cap E_\alpha), \mathcal{T} \cap \left[\bigcup_{\alpha \in A} E_\alpha\right]\right)$ -continuous.
- Let $B \in \bigoplus_{\alpha \in A} (\mathcal{T} \cap E_\alpha)$. Exercise 4.7.0.3 implies that for each $\alpha \in A$, there exist $B_\alpha \in \mathcal{T} \cap E_\alpha$ such that $B = \coprod_{\alpha \in A} B_\alpha$. Then for each $\alpha \in A$, there exists $C_\alpha \in \mathcal{T}$ such that $B_\alpha = C_\alpha \cap E_\alpha$. Since $(E_\alpha)_{\alpha \in A} \subset \mathcal{T}$, we have that for each $\alpha \in A$, $C_\alpha \cap E_\alpha \in \mathcal{T}$. Hence for each $\alpha \in A$,

$$\begin{aligned} B_\alpha &= C_\alpha \cap E_\alpha \\ &= (C_\alpha \cap E_\alpha) \cap \left[\bigcup_{\alpha' \in \mathbb{N}} E_{\alpha'}\right] \\ &\in \mathcal{T} \cap \left[\bigcup_{\alpha' \in \mathbb{N}} E_{\alpha'}\right]. \end{aligned}$$

Therefore

$$\begin{aligned} \phi(B) &= \phi\left(\coprod_{\alpha \in A} B_\alpha\right) \\ &= \phi\left(\bigcup_{\alpha \in A} \iota_\alpha(B_\alpha)\right) \\ &= \bigcup_{\alpha \in A} \phi \circ \iota_\alpha(B_\alpha) \\ &= \bigcup_{\alpha \in A} \iota_{E_\alpha}(B_\alpha) \\ &= \bigcup_{\alpha \in A} B_\alpha \\ &\in \mathcal{T} \cap \left[\bigcup_{\alpha \in A} E_\alpha\right]. \end{aligned}$$

Since $B \in \bigoplus_{\alpha \in A} (\mathcal{T} \cap E_\alpha)$ is arbitrary, we have that for each $B \in \bigoplus_{\alpha \in A} (\mathcal{T} \cap E_\alpha)$, $\phi(B) \in \mathcal{T}$.

Hence ϕ is a homeomorphism. □

Exercise 4.7.0.12. Let $(X, \mathcal{T})$ be a topological space and $(E_\alpha)_{\alpha \in A} \subset \mathcal{P}(X)$. Define $\phi : \coprod_{\alpha \in A} E_\alpha \rightarrow \bigcup_{\alpha \in A} E_\alpha$ by $\phi(\alpha, x) := x$. Then ϕ is $\left(\bigoplus_{\alpha \in A} (\mathcal{T} \cap E_\alpha), \mathcal{T} \cap \left(\bigcup_{\alpha \in A} E_\alpha\right)\right)$ -continuous.

Proof. Set $E := \bigcup_{\alpha \in A} E_\alpha$. Let $\alpha \in A$ and $x \in E_\alpha$. Then

$$\begin{aligned} \phi \circ \iota_\alpha(x) &= \phi(\alpha, x) \\ &= x \\ &= \text{id}_{E_\alpha}(x). \end{aligned}$$

Since $x \in E_\alpha$ is arbitrary, we have that for each $x \in E_\alpha$, $\phi \circ \iota_\alpha(x) = \text{id}_{E_\alpha}(x)$. Hence $\phi \circ \iota_\alpha = \text{id}_{E_\alpha}$. We note that $\phi \circ \iota_\alpha$ is $(\mathcal{T} \cap E_\alpha, \mathcal{T} \cap E_\alpha)$ -continuous. Since Exercise 4.5.1.6 implies that $(\mathcal{T} \cap E) \cap E_\alpha = \mathcal{T} \cap E_\alpha$, Exercise 4.5.1.11 implies that id_{E_α} is $(\mathcal{T} \cap E_\alpha, \mathcal{T} \cap E)$ -continuous. Since $\alpha \in A$ is arbitrary, we have that for each $\alpha \in A$, $\phi \circ \iota_\alpha$ is $(\mathcal{T} \cap E_\alpha, \mathcal{T} \cap E)$ -continuous. Exercise 4.7.0.9 implies that ϕ is $\left(\bigoplus_{\alpha \in A} (\mathcal{T} \cap E_\alpha), \mathcal{T} \cap E\right)$ -continuous. □

4.8 Quotient Topology

4.8.1 Introduction

Definition 4.8.1.1. Let $(X, \mathcal{T}_X)$, $(Y, \mathcal{T}_Y)$ be topological spaces and $f : X \rightarrow Y$. Then f is said to be an $(\mathcal{T}_X, \mathcal{T}_Y)$ -**quotient map** if

1. f is surjective
2. $\mathcal{T}_Y = f_*\mathcal{T}_X$

Note 4.8.1.2. We typically avoid specifying the topologies when they are clear from the context.

Exercise 4.8.1.3. Let $(X, \mathcal{T}_X)$, $(Y, \mathcal{T}_Y)$ be topological spaces and $f : X \rightarrow Y$. Suppose that f is surjective. Then

1. f is a quotient map iff

$$\text{for each } V \subset Y, V \in \mathcal{T}_Y \text{ iff } f^{-1}(V) \in \mathcal{T}_X$$

2. f is a quotient map iff

$$\text{for each } C \subset Y, C \text{ is closed in } Y \text{ iff } f^{-1}(C) \text{ is closed in } X$$

Proof.

1. • ($\Rightarrow$)

Suppose that f is a quotient map.

Let $V \subset Y$. Suppose that V is open, then

$$\begin{aligned} V &\in \mathcal{T}_Y \\ &= f_*\mathcal{T}_X \\ &= \{V' \subset Y : f^{-1}(V') \in \mathcal{T}_X\} \end{aligned}$$

Thus $f^{-1}(V) \in \mathcal{T}_X$. Hence $f^{-1}(V)$ is open.

Conversely, suppose that $f^{-1}(V)$ is open. Then $f^{-1}(V) \in \mathcal{T}_X$. Since

$$\begin{aligned} \mathcal{T}_Y &= f_*\mathcal{T}_X \\ &= \{V' \subset Y : f^{-1}(V') \in \mathcal{T}_X\}, \end{aligned}$$

we have that $V \in \mathcal{T}_Y$. Hence V is open.

Thus V is open iff $f^{-1}(V)$ is open. Since $V \subset Y$ is arbitrary, we have that for each $V \subset Y$, V is open iff $f^{-1}(V)$ is open.

- ($\Leftarrow$)

Suppose that for each $V \subset Y$, V is open iff $f^{-1}(V)$ is open. Then

$$\begin{aligned} \mathcal{T}_Y &= \{V \subset Y : f^{-1}(V) \in \mathcal{T}_X\} \\ &= f_*\mathcal{T}_X \end{aligned}$$

So f is a quotient map.

2. • ($\Rightarrow$)

Suppose that f is a quotient map.

Let $C \subset Y$. Suppose that C is closed, then $C^c \in \mathcal{T}_Y$. Continuity implies that

$$\begin{aligned} f^{-1}(C)^c &= f^{-1}(C^c) \\ &\in \mathcal{T}_X \end{aligned}$$

Thus $f^{-1}(C)$ is closed.

Conversely, suppose that $f^{-1}(C)$ is closed. Then

$$\begin{aligned} f^{-1}(C^c) &= f^{-1}(C)^c \\ &\in \mathcal{T}_X \end{aligned}$$

Since f is a quotient map, $C^c \in \mathcal{T}_Y$. Hence C is closed.

Thus C is closed iff $f^{-1}(C)$ is closed. Since $C \subset Y$ is arbitrary, we have that for each $C \subset Y$, C is closed iff $f^{-1}(C)$ is closed.

• ($\Leftarrow$)

Suppose that for each $C \subset Y$, C is closed iff $f^{-1}(C)$ is closed.

Let $V \subset Y$. Suppose that V is open. Then V^c is closed. By assumption $f^{-1}(V^c)$ is closed and therefore

$$\begin{aligned} f^{-1}(V) &= f^{-1}(V^c)^c \\ &\in \mathcal{T}_X. \end{aligned}$$

Thus $f^{-1}(V)$ is open.

Conversely, suppose that $f^{-1}(V)$ is open. Then $f^{-1}(V)^c$ is closed. Since $f^{-1}(V^c) = f^{-1}(V)^c$, we have that $f^{-1}(V^c)$ is closed. By assumption, V^c is closed. Thus V is open.

Thus V is open iff $f^{-1}(V)$ is open. Since $V \subset Y$ is arbitrary, we have that for each $V \subset Y$, V is open iff $f^{-1}(V)$ is open.

Part (1) implies that f is a quotient map.

□

Exercise 4.8.1.4. Let $(X, \mathcal{T}_X)$, $(Y, \mathcal{T}_Y)$ be topological spaces and $f : X \rightarrow Y$. If f is a quotient map, then f is continuous.

Proof. Suppose that f is a quotient map. Let $V \subset Y$. Suppose that V is open. [The previous exercise](#) implies that $f^{-1}(V)$ is open. Since $V \subset Y$ with V open is arbitrary, we have that for each $V \subset Y$, V is open implies that $f^{-1}(V)$ is open. Hence f is continuous. □

Exercise 4.8.1.5. Let $(X, \mathcal{T}_X)$, $(Y, \mathcal{T}_Y)$, $(Z, \mathcal{C})$ be topological spaces, $f : X \rightarrow Y$ and $g : Y \rightarrow Z$. If f is a quotient map, then g is continuous iff $g \circ f$ is continuous.

Proof. Suppose that f is a quotient map. Then $\mathcal{T}_Y = f_*\mathcal{T}_X$. [An exercise in the section on continuous maps](#) implies that g is continuous iff $g \circ f$ is continuous. □

Exercise 4.8.1.6. Let $(X, \mathcal{T}_X)$, $(Y_1, \mathcal{T}_{Y_1})$, $(Y_2, \mathcal{T}_{Y_2})$ be topological spaces, $f_1 : X \rightarrow Y_1$, $f_2 : X \rightarrow Y_2$ and $\phi : Y_1 \rightarrow Y_2$. Suppose that f_1 and f_2 are quotient maps and ϕ is a bijection. If $\phi \circ f_1 = f_2$, then ϕ is a homeomorphism.

Proof. Since f_1 and f_2 are quotient maps, they are continuous. Suppose that $\phi \circ f_1 = f_2$. Since f_2 is continuous, the previous exercise implies that ϕ is continuous. Since ϕ is a bijection, $f_1 = \phi^{-1} \circ f_2$. Similarly, since f_1 is continuous, the previous exercise implies that ϕ^{-1} is continuous. Hence ϕ is a homeomorphism. □

Exercise 4.8.1.7. [Restate the last exercise categorically: Let \$U : \mathbf{Top} \rightarrow \mathbf{Set}\$ be the forgetful functor. If \$\phi \in \text{Iso}_{U\(X\)/U\(\mathbf{Top}\)}\(U\(f_1\), U\(f_2\)\)\$ then there exists \$\phi' \in \text{Iso}_{X/\mathbf{Top}}\(f_1, f_2\)\$ such that \$U\(\phi'\) = \phi\$, adjoint functor?...](#)

Proof. **FINISH!!!**

□

Exercise 4.8.1.8. Let $(X, \mathcal{T}_X)$, $(Y, \mathcal{T}_Y)$ be topological spaces and $f : X \rightarrow Y$. Suppose that f is $(\mathcal{T}_X, \mathcal{T}_Y)$ -continuous and surjective. If f is $(\mathcal{T}_X, \mathcal{T}_Y)$ -open or f is $(\mathcal{T}_X, \mathcal{T}_Y)$ -closed, then f is a $(\mathcal{T}_X, \mathcal{T}_Y)$ -quotient map.

Proof.

- Suppose that f is $(\mathcal{T}_X, \mathcal{T}_Y)$ -open. Let $V \subset Y$.
Suppose that $V \in \mathcal{T}_Y$. Continuity implies that $f^{-1}(V) \in \mathcal{T}_X$. Conversely, suppose that $f^{-1}(V) \in \mathcal{T}_X$. Since f is open $f(f^{-1}(V)) \in \mathcal{T}_Y$. Surjectivity implies that $V = f(f^{-1}(V))$. So $V \in \mathcal{T}_Y$. Exercise 4.8.1.3 then implies that f is a quotient map.
- Suppose that f is closed. Then similarly to above, f is a quotient map.

□

Exercise 4.8.1.9. Let $(X, \mathcal{T}_X)$, $(Y, \mathcal{T}_Y)$ be topological spaces and $f : X \rightarrow Y$. Suppose that f is a quotient map. Then

1. f is open iff

for each $U \subset X$, U is open implies that $f^{-1}(f(U))$ is open

2. f is closed iff

for each $C \subset X$, C is closed implies that $f^{-1}(f(C))$ is closed

Proof.

1. • ($\implies$)
 Suppose that f is open.
 Let $U \subset X$. Suppose that U is open. Since f is open, $f(U)$ is open. Continuity implies that $f^{-1}(f(U))$ is open.
- ($\impliedby$)
 Suppose that for each $U \subset X$, U is open implies that $f^{-1}(f(U))$ is open.
 Let $U \subset X$. Suppose that U is open. By assumption, $f^{-1}(f(U))$ is open. Since f is a quotient map, $f(U)$ is open. Since $U \subset X$ with U open is arbitrary, we have that for each $U \subset X$, U is open implies $f(U)$ is open. Thus f is open.
2. • ($\implies$)
 Suppose that f is closed.
 Let $C \subset X$. Suppose that C is closed. Since f is closed, $f(C)$ is closed. Continuity implies that $f^{-1}(f(C))$ is closed.
- ($\impliedby$)
 Suppose that for each $C \subset X$, C is closed implies that $f^{-1}(f(C))$ is closed.
 Let $C \subset X$. Suppose that C is closed. By assumption, $f^{-1}(f(C))$ is closed. Since f is a quotient map, $f(C)$ is closed. Since $C \subset X$ with C closed is arbitrary, we have that for each $C \subset X$, C is closed implies $f(C)$ is closed. Thus f is closed.

□

Exercise 4.8.1.10. Let $(X, \mathcal{T})$ be a topological space, Y a set and $f : X \rightarrow Y$. Suppose that f is surjective. Then $f : X \rightarrow Y$ is a $(\mathcal{T}, f_*\mathcal{T})$ quotient map.

Proof. Clear by definition. □

Exercise 4.8.1.11. Let $(X, \mathcal{T})$ be a topological space, $\sim$ an equivalence relation on X and $\pi : X \rightarrow X/\sim$ the projection map given by $x \mapsto \bar{x}$. Then π is a $(\mathcal{T}, \pi_*\mathcal{T})$ -quotient map.

Proof. Since π is surjective, the previous exercise implies that π is a $(\mathcal{T}, \pi_*\mathcal{T})$ -quotient map. □

Definition 4.8.1.12. Let $(X, \mathcal{T})$ be a topological space, $\sim$ an equivalence relation on X and $\pi : X \rightarrow X/\sim$ the projection map given by $x \mapsto \bar{x}$. We define the **quotient topology on $X/\sim$** on $X/\sim$, denoted $\mathcal{T}_{X/\sim}$, by

$$\mathcal{T}_{X/\sim} = \pi_*\mathcal{T}$$

Definition 4.8.1.13. Let $(X, \mathcal{T}_X)$, $(Y, \mathcal{T}_Y)$ be topological spaces, $\sim_X$ an equivalence relation on X , $\sim_Y$ and equivalence relation on Y and $f : X \rightarrow Y$. Then f is said to be $(\sim_X, \sim_Y)$ -**invariant** if for each $x, y \in X$, $x \sim_X y$ implies that $f(x) \sim_Y f(y)$.

Exercise 4.8.1.14. Let $(X, \mathcal{T}_X)$, $(Y, \mathcal{T}_Y)$ be topological spaces, $\sim_X, \sim_Y$ equivalence relations on X and Y respectively, $\pi_X : X \rightarrow X/\sim_X$, $\pi_Y : Y \rightarrow Y/\sim_Y$ the respective projection maps and $f : X \rightarrow Y$ continuous. If f is $(\sim_X, \sim_Y)$ -invariant, then there exists a unique $\bar{f} : X/\sim_X \rightarrow Y/\sim_Y$ such that $\bar{f}$ is continuous and $\bar{f} \circ \pi_X = \pi_Y \circ f$, i.e. the following diagram commutes:

$$\begin{array}{ccc} X & \xrightarrow{f} & Y \\ \pi_X \downarrow & & \downarrow \pi_Y \\ X/\sim_X & \xrightarrow{\bar{f}} & Y/\sim_Y \end{array}$$

and $\bar{f}$ is $(\mathcal{T}_{X/\sim_X}, \mathcal{T}_{Y/\sim_Y})$ -continuous.

Proof. Suppose that f is $(\sim_X, \sim_Y)$ -invariant.

- **Existence:**

Define $\bar{f} : X/\sim_Y \rightarrow Y/\sim_Y$ by $\bar{f}(\bar{x}) = \overline{f(x)}$. Let $a, b \in X$. Then

$$\begin{aligned} \bar{a} = \bar{b} &\implies a \sim_X b \\ &\implies f(a) \sim_Y f(b) \\ &\implies \overline{f(a)} = \overline{f(b)} \\ &\implies \bar{f}(\bar{a}) = \bar{f}(\bar{b}) \end{aligned}$$

So $\bar{f}$ is well defined. By construction $\bar{f} \circ \pi_X = \pi_Y \circ f$.

- **Uniqueness:**

Let $g : X/\sim_X \rightarrow Y/\sim_Y$. Suppose that $g \circ \pi_X = \pi_Y \circ f$, i.e. the following diagram commutes:

$$\begin{array}{ccc} X & \xrightarrow{f} & Y \\ \pi_X \downarrow & & \downarrow \pi_Y \\ X/\sim_X & \xrightarrow{g} & Y/\sim_Y \end{array}$$

Then

$$\begin{aligned} g \circ \pi_X &= \pi_Y \circ f \\ &= \bar{f} \circ \pi_X \end{aligned}$$

i.e. the following diagram commutes:

$$\begin{array}{ccc} X & \xrightarrow{\pi_X} & Y \\ \pi_X \downarrow & & \downarrow \bar{f} \\ X/\sim_X & \xrightarrow{g} & Y/\sim_Y \end{array}$$

Since π_X is surjective, Exercise 1.3.2.3 implies that $\bar{f} = g$.

- **Continuity:**

Let $V \in \mathcal{T}_{Y/\sim_Y}$. Continuity of f and π_Y implies that

$$\begin{aligned} \pi_X^{-1}(\bar{f}^{-1}(V)) &= (\bar{f} \circ \pi_X)^{-1}(V) \\ &= (\pi_Y \circ f)^{-1}(V) \\ &= f^{-1}(\pi_Y^{-1}(V)) \\ &\in \mathcal{T}_X \end{aligned}$$

By definition of the quotient topology, $\bar{f}^{-1}(V) \in \mathcal{T}_{X/\sim_X}$. So $\bar{f}$ is $(\mathcal{T}_{X/\sim_X}, \mathcal{T}_{Y/\sim_Y})$ -continuous. □

Definition 4.8.1.15. Let $(X, \mathcal{T}_X)$, $(Y, \mathcal{T}_Y)$ be topological spaces, $\sim_X$ an equivalence relation on X and $f : X \rightarrow Y$. Then f is said to be **$\sim$ -invariant** if f is $(\sim, =_Y)$ -invariant.

Exercise 4.8.1.16. Let $(X, \mathcal{T}_X)$, $(Y, \mathcal{T}_Y)$ be topological spaces, $\sim \subset X \times X$ an equivalence relation on X , $\pi : X \rightarrow X/\sim$ the projection map and $f : X \rightarrow Y$ continuous. If f is $\sim$ -invariant, then there exists a unique $\bar{f} : X/\sim \rightarrow Y$ such that $\bar{f}$ is continuous and $\bar{f} \circ \pi = f$, i.e. the following diagram commutes:

$$\begin{array}{ccc} X & & \\ \pi \downarrow & \searrow f & \\ X/\sim & \xrightarrow{\bar{f}} & Y \end{array}$$

and $\bar{f}$ is $(\mathcal{T}_{X/\sim}, \mathcal{T}_Y)$ -continuous.

Proof. Set $\sim_X := \sim$ and $\sim_Y := =_Y$ and use Exercise 4.8.1.14. [add details](#) □

Exercise 4.8.1.17. Let $(X, \mathcal{T})$, $(Y, \mathcal{T}_Y)$ be topological spaces, $\sim$ an equivalence relation on X , $\pi : X \rightarrow X/\sim$ the projection map and $f : X \rightarrow Y$. Suppose f is a quotient map. If for each $a, b \in X$, $a \sim b$ iff $f(a) = f(b)$, then there exists a unique $\bar{f} : X/\sim \rightarrow Y$ such that $\bar{f}$ is a $(\mathcal{T}_{X/\sim}, \mathcal{T}_Y)$ -homeomorphism and $\bar{f} \circ \pi = f$.

Proof. Suppose that for each $a, b \in X$, $a \sim b$ iff $f(a) = f(b)$. Then f is $\sim$ -invariant. Since f is a quotient map, f is continuous. Exercise 4.8.1.16 implies that there exists a unique $\bar{f} : X/\sim \rightarrow Y$ such that $\bar{f}$ is continuous and $\bar{f} \circ \pi = f$. Let $y \in Y$. Since f is a quotient map, f is surjective. Therefore there exists $x \in X$ such that $f(x) = y$. Thus

$$\begin{aligned}\bar{f}(\bar{x}) &= f(x) \\ &= y\end{aligned}$$

Since $y \in Y$ is arbitrary, $\bar{f}$ is surjective. Let $\bar{a}, \bar{b} \in X/\sim$. Suppose that $\bar{f}(\bar{a}) = \bar{f}(\bar{b})$. Then

$$\begin{aligned}f(a) &= \bar{f}(\bar{a}) \\ &= \bar{f}(\bar{b}) \\ &= f(b)\end{aligned}$$

By assumption, $a \sim b$. Hence $\bar{a} = \bar{b}$. Since $\bar{a}, \bar{b} \in X/\sim$ are arbitrary, $\bar{f}$ is injective. Therefore, $\bar{f}$ is a bijection. Exercise 4.8.1.6 implies that $\bar{f}$ is a homeomorphism. □

Definition 4.8.1.18. Let $(X, \mathcal{T})$, $(Y, \mathcal{T}_Y)$ be topological spaces and $f : X \rightarrow Y$. We define the relation $\sim_f \subset X \times X$ by $x \sim_f y$ iff $f(x) = f(y)$.

Exercise 4.8.1.19. Let $(X, \mathcal{T})$, $(Y, \mathcal{T}_Y)$ be topological spaces and $f : X \rightarrow Y$. Then

1. $\sim_f$ is an equivalence relation on X ,
2. if f is a quotient map, then there exists a unique $\bar{f} : X/\sim_f \rightarrow Y$ such that $\bar{f}$ is a $(\mathcal{T}_{X/\sim_f}, \mathcal{T}_Y)$ -homeomorphism and $\bar{f} \circ \pi = f$.

Proof.

1. Clear.
2. Suppose that f is a quotient map. Exercise 4.8.1.17 implies that then there exists a unique $\bar{f} : X/\sim_f \rightarrow Y$ such that $\bar{f}$ is a $(\mathcal{T}_{X/\sim_f}, \mathcal{T}_Y)$ -homeomorphism and $\bar{f} \circ \pi = f$. □

Exercise 4.8.1.20. Let $(X, \mathcal{T})$, $(Y, \mathcal{T}_Y)$ and $(Z, \mathcal{T}_Z)$ be topological spaces, $f : X \rightarrow Y$ and $g : X \rightarrow Z$. Suppose that f is a quotient map. If g is $\sim_f$ -invariant, then there exists a unique $\bar{g} : Y \rightarrow Z$ such that $\bar{g}$ is continuous and $\bar{g} \circ f = g$.

Proof. Suppose that g is $\sim_f$ -invariant.

• **Existence:**

Exercise 4.8.1.16 implies that there exists a unique $\bar{g}_0 : X/\sim_f \rightarrow Z$ such that $\bar{g}_0$ is continuous and $\bar{g}_0 \circ \pi_{\sim_f} = g$. Exercise 4.8.1.19 implies that there exists a unique $\bar{f} : X/\sim_f \rightarrow Y$ such that $\bar{f}$ is a $(\mathcal{T}_{X/\sim_f}, \mathcal{T}_Y)$ -homeomorphism and $\bar{f} \circ \pi_{\sim_f} = f$. Define $\bar{g} : Y \rightarrow Z$ by $\bar{g} := \bar{g}_0 \circ \bar{f}^{-1}$. Since $\bar{g}_0$ and $\bar{f}^{-1}$ are continuous, we have that $\bar{g}$ is continuous. By construction,

$$\begin{aligned}\bar{g} \circ f &= (\bar{g}_0 \circ \bar{f}^{-1}) \circ (\bar{f} \circ \pi_{\sim_f}) \\ &= \bar{g}_0 \circ (\bar{f}^{-1} \circ \bar{f}) \circ \pi_{\sim_f} \\ &= \bar{g}_0 \circ \pi_{\sim_f} \\ &= g.\end{aligned}$$

Thus there exists $\bar{g} : Y \rightarrow Z$ such that $\bar{g}$ is continuous and $\bar{g} \circ f = g$.

• **Uniqueness:**

Let $\bar{g}' : Y \rightarrow Z$. Suppose that $\bar{g}'$ is continuous and $\bar{g}' \circ f = g$. Then


$$\begin{aligned}\bar{g}' \circ f &= g \\ &= \bar{g} \circ f.\end{aligned}$$

Since f is a bijection, we have that

$$\begin{aligned}\bar{g}' &= (\bar{g}' \circ f) \circ f^{-1} \\ &= (\bar{g} \circ f) \circ f^{-1} \\ &= \bar{g}.\end{aligned}$$

Since $\bar{g}' : Y \rightarrow Z$ such that $\bar{g}'$ continuous and $\bar{g}' \circ f = g$ is arbitrary, we have that for each $\bar{g}' : Y \rightarrow Z$, $\bar{g}$ is continuous and $\bar{g}' \circ f = g$ implies that $\bar{g}' = \bar{g}$. □

Exercise 4.8.1.21. Let X, X_1, X_2 be topological spaces and $f_1 : X \rightarrow X_1, f_2 : X \rightarrow X_2$ and $f_{1,2} : X_2 \rightarrow X_1$. Suppose that f_1 and f_2 are quotient maps. If $f_{1,2} \circ f_1 = f_2$, then there exists unique $\bar{f}_1 : X/\sim_{f_1} \rightarrow X_1, \bar{f}_2 : X/\sim_{f_2} \rightarrow X_2, \bar{f}_{1,2} : X/\sim_{f_2} \rightarrow X/\sim_{f_1}$ such that $\bar{f}_1, \bar{f}_2$ are homeomorphisms, $\bar{f}_{1,2}$ is continuous, $\bar{f}_1 \circ \pi_1 = f_1, \bar{f}_2 \circ \pi_2 = f_2, \bar{f}_{1,2} \circ \bar{f}_1 = f_{1,2}$ and $\bar{f}_{1,2} \circ \pi_1 = \pi_2$, i.e. the following diagram commutes:



Proof.

• **Existence and Uniqueness of $\bar{f}_k$:**

Since f_1, f_2 are quotient maps, Exercise 4.8.1.19 implies that there exist unique $\bar{f}_1 : X/\sim_{f_1} \rightarrow X_1, \bar{f}_2 : X/\sim_{f_2} \rightarrow X_2$ such that $\bar{f}_1, \bar{f}_2$ are homeomorphisms, $\bar{f}_1 \circ \pi_1 = f_1$ and $\bar{f}_2 \circ \pi_2 = f_2$.

• **Existence of $\bar{f}_{1,2}$:**

Define $\bar{f}_{1,2} : X/\sim_{f_1} \rightarrow X/\sim_{f_2}$ by $\bar{f}_{1,2} := \bar{f}_2^{-1} \circ f_{1,2} \circ \bar{f}_1$. Since $\bar{f}_2^{-1}, f_{1,2}, \bar{f}_1$ are continuous, $\bar{f}_{1,2}$ is continuous. By construction,

$$\begin{aligned}\bar{f}_2 \circ \bar{f}_{1,2} &= \bar{f}_2 \circ (\bar{f}_2^{-1} \circ f_{1,2} \circ \bar{f}_1) \\ &= (\bar{f}_2 \circ \bar{f}_2^{-1}) \circ (f_{1,2} \circ \bar{f}_1) \\ &= \text{id}_{X_2} \circ (f_{1,2} \circ \bar{f}_1) \\ &= f_{1,2} \circ \bar{f}_1.\end{aligned}$$

and therefore

$$\begin{aligned}\bar{f}_{1,2} \circ \pi_1 &= (\bar{f}_2^{-1} \circ f_{1,2} \circ \bar{f}_1) \circ \pi_1 \\ &= (\bar{f}_2^{-1} \circ f_{1,2}) \circ (\bar{f}_1 \circ \pi_1) \\ &= (\bar{f}_2^{-1} \circ f_{1,2}) \circ f_1 \\ &= \bar{f}_2^{-1} \circ (f_{1,2} \circ f_1) \\ &= \bar{f}_2^{-1} \circ f_2 \\ &= \pi_2\end{aligned}$$

- **Uniqueness of $\bar{f}_{1,2}$:**

Let $g_{1,2} : X/\sim_{f_1} \rightarrow X/\sim_{f_2}$. Suppose that $\bar{f}_2 \circ g_{1,2} = f_{1,2} \circ \bar{f}_1$. Then

$$\begin{aligned} g_{1,2} &= \bar{f}_2^{-1} \circ f_{1,2} \circ \bar{f}_1 \\ &= \bar{f}_{1,2}. \end{aligned}$$

Thus $\bar{f}_{1,2}$ is unique. □

4.8.2 Category of Topological Spaces with Equivalence Relations

Definition 4.8.2.1. We define the category **TopEq**

- $\text{Obj}(\mathbf{TopEq}) = \{(X, \sim) : X \text{ is a topological space and } \sim \text{ is an equivalence relation on } X\}$
- $\text{Hom}_{\mathbf{TopEq}}((X, \sim_X), (Y, \sim_Y)) = \{f : X \rightarrow Y : f \text{ is continuous and } f \text{ is } (\sim_X, \sim_Y)\text{-invariant}\}.$

Definition 4.8.2.2. We define $F : \mathbf{TopEq} \rightarrow \mathbf{Top}$ by

- $F(X, \sim) = X/\sim$
- $F(f) = \bar{f}$

Exercise 4.8.2.3. We have that $F : \mathbf{TopEq} \rightarrow \mathbf{Top}$ is a functor

Proof.

1. Let $(X, \sim_X), (Y, \sim_Y) \in \mathbf{TopEq}$ and $f \in \text{Hom}_{\mathbf{TopEq}}((X, \sim_X), (Y, \sim_Y))$. The previous exercise implies that $F(f) \in \text{Hom}_{\mathbf{Top}}(X/\sim_X, Y/\sim_Y)$.
2. Let $(X, \sim_X), (Y, \sim_Y), (Z, \sim_Z) \in \mathbf{TopEq}$, $f \in \text{Hom}_{\mathbf{TopEq}}((X, \sim_X), (Y, \sim_Y))$ and $g \in \text{Hom}_{\mathbf{TopEq}}((Y, \sim_Y), (Z, \sim_Z))$. Then

$$\begin{aligned} \pi_Z \circ (g \circ f) &= (\pi_Z \circ g) \circ f \\ &= (\bar{g} \circ \pi_Y) \circ f \\ &= \bar{g} \circ (\pi_Y \circ f) \\ &= \bar{g} \circ (\bar{f} \circ \pi_X) \\ &= (\bar{g} \circ \bar{f}) \circ \pi_X \end{aligned}$$

Uniqueness implies that $F(g \circ f) = F(g) \circ F(f)$. □

Exercise 4.8.2.4. Let $(X, \sim_X), (Y, \sim_Y), (Z, =) \in \text{Obj}(\mathbf{TopEq})$, $\iota \in \text{Hom}_{\mathbf{TopEq}}((X, \sim_X), (Y, \sim_Y))$, $f \in \text{Hom}_{\mathbf{TopEq}}((X, \sim_X), (Z, =))$ and $g \in \text{Hom}_{\mathbf{TopEq}}((Y, \sim_Y), (Z, =))$. Then $f = g \circ \iota$ iff $\bar{f} = \bar{g} \circ \bar{\iota}$, in which case the following diagram commutes:

$$\begin{array}{ccc} & & Z \\ & \nearrow f & \nwarrow g \\ X & \xrightarrow{\iota} & Y \\ \pi_X \downarrow & & \downarrow \pi_Y \\ X/\sim_X & \xrightarrow{\bar{\iota}} & Y/\sim_Y \end{array}$$

Proof. Suppose that $f = g \circ \iota$. Functoriality implies that

$$\begin{aligned}\bar{f} &= F(f) \\ &= F(g \circ \iota) \\ &= F(g) \circ F(\iota) \\ &= \bar{g} \circ \bar{\iota}\end{aligned}$$

Conversely, suppose that $\bar{f} = \bar{g} \circ \bar{\iota}$. Then

$$\begin{aligned}f &= \bar{f} \circ \pi_X \\ &= \bar{g} \circ \bar{\iota} \circ \pi_X \\ &= \bar{g} \circ \pi_Y \circ \iota \\ &= g \circ \iota\end{aligned}$$

□

Exercise 4.8.2.5. Let G be a group, X a topological space and $\phi : G \times X \rightarrow X$ a group action. Suppose that for each $g \in G$, the map $\phi_g \in \text{Sym}(X)$ defined by $\phi_g(x) = g \cdot x$ is continuous. Then $\pi : X \rightarrow X/G$ is open. **move this before category theory stuff**

Proof. Suppose that for each $g \in G$, ϕ_g is continuous. Let $g \in G$. Since $(\phi_g)^{-1} = \phi_{g^{-1}}$, ϕ_g is a homeomorphism. Hence for each $g \in G$ and $U \subset X$, U is open iff $g \cdot U$ is open. Let $U \subset X$. Suppose that U is open. Then $\pi^{-1}(\pi(U)) = \bigcup_{g \in G} g \cdot U$ is open. Since π is a quotient map, Exercise 4.8.1.9 implies that π is open. □

4.9 Equalizers of Continuous Maps

Note 4.9.0.1. Let X, Y be sets and $f, g : X \rightarrow Y$. We denote the equalizer of f and g by $\text{Eq}(f, g)$ as in Definition 1.8.0.1.

Exercise 4.9.0.2. Let $(X, \mathcal{T}_X), (Y, \mathcal{T}_Y)$ be topological spaces and $f, g : X \rightarrow Y$ $(\mathcal{T}_X, \mathcal{T}_Y)$ -continuous. If $(Y, \mathcal{T}_Y)$ is Hausdorff, then $\text{Eq}(f, g)$ is closed in $(X, \mathcal{T}_X)$.

Proof. Suppose that $(Y, \mathcal{T}_Y)$ is Hausdorff. Exercise 4.12.1.7 implies that Δ_Y is closed in $(Y \times Y, \mathcal{T}_Y \otimes \mathcal{T}_Y)$. Since f, g are $(\mathcal{T}_X, \mathcal{T}_Y)$ -continuous, Exercise 4.6.3.5 implies that $(f, g) : X \rightarrow Y \times Y$ is $(\mathcal{T}_X, \mathcal{T}_Y \otimes \mathcal{T}_Y)$ -continuous. Since $\text{Eq}(f, g) = (f, g)^{-1}(\Delta_Y)$, we have that $\text{Eq}(f, g)$ is closed in $(X, \mathcal{T}_X)$. $\square$

4.10 Projective Limits of Topological Spaces

Note 4.10.0.1. Let $((X_j)_{j \in J}, (\pi_{j,k})_{(j,k) \in \leq})$ a projective system of sets.

- We denote the j -th projection map from $\prod_{j \in J} X_j$ onto X_j by $\text{proj}_j : \prod_{j \in J} X_j \rightarrow X_j$.
- We denote the j -th projection map from $\varprojlim_{j \in J} X_j$ into X_j by $\pi_j : \varprojlim_{j \in J} X_j \rightarrow X_j$ as in Definition 1.9.0.3

define general inverse limits and show universal property and specific concrete examples

Definition 4.10.0.2. Let $(J, \leq)$ be a directed poset, $(X_j, \mathcal{T}_j)_{j \in J}$ a collection of topological spaces and for each $(j, k) \in \leq$, $\pi_{j,k} : X_k \rightarrow X_j$ a $(\mathcal{T}_k, \mathcal{T}_j)$ -continuous map. Suppose that for each $j, k, l \in J$,

1. $\pi_{j,j} = \text{id}_{X_j}$,
2. $j \leq k$ and $k \leq l$ implies that $\pi_{j,k} \circ \pi_{k,l} = \pi_{j,l}$.

Then $((X_j, \mathcal{T}_j)_{j \in J}, (\pi_{j,k})_{(j,k) \in \leq})$ is said to be a **Top-projective system**.

Definition 4.10.0.3. Let $(J, \leq)$ be a directed poset, $((X_j, \mathcal{T}_j)_{j \in J}, (\pi_{j,k})_{(j,k) \in \leq})$ a projective system of topological spaces. We define the **projective limit topology** on $\varprojlim_{j \in J} X_j$, denoted $\varprojlim_{j \in J} \mathcal{T}_j$, by

$$\varprojlim_{j \in J} \mathcal{T}_j := \tau(\pi_j : j \in J).$$

Exercise 4.10.0.4. Let $(J, \leq)$ be a directed poset and $((X_j, \mathcal{T}_j)_{j \in J}, (\pi_{j,k})_{(j,k) \in \leq})$ a projective system of topological spaces. Then

$$\varprojlim_{j \in J} \mathcal{T}_j = \left[\bigotimes_{j \in J} \mathcal{T}_j \right] \cap \varprojlim_{j \in J} X_j.$$

Proof.

- Define $\mathcal{E} \subset \varprojlim_{j \in J} \mathcal{T}_j$ by

$$\mathcal{E} := \{\pi_j^{-1}(V) : j \in J \text{ and } V \in \mathcal{T}_j\}.$$

By definition, $\varprojlim_{j \in J} \mathcal{T}_j = \tau(\mathcal{E})$. Let $U \in \mathcal{E}$. Then there exists $j \in J$ and $V \in \mathcal{T}_j$ such that $U = \pi_j^{-1}(V)$. Since $\pi_j = \text{proj}_j|_{\varprojlim_{j \in J} X_j}$

and $\text{proj}_j^{-1}(V) \in \left[\bigotimes_{j \in J} \mathcal{T}_j \right]$, we have that

$$\begin{aligned} U &= \pi_j^{-1}(V) \\ &= (\text{proj}_j|_{\varprojlim_{j \in J} X_j})^{-1}(V) \\ &= \text{proj}_j^{-1}(V) \cap \varprojlim_{j \in J} X_j \\ &\in \left[\bigotimes_{j \in J} \mathcal{T}_j \right] \cap \varprojlim_{j \in J} X_j. \end{aligned}$$

Since $U \in \mathcal{E}$ is arbitrary, we have that $\mathcal{E} \subset \left[\bigotimes_{j \in J} \mathcal{T}_j \right] \cap \varprojlim_{j \in J} X_j$. Since by definition, $\varprojlim_{j \in J} \mathcal{T}_j = \tau(\mathcal{E})$, we have that

$$\begin{aligned} \varprojlim_{j \in J} \mathcal{T}_j &= \tau(\mathcal{E}) \\ &\subset \tau\left(\left[\bigotimes_{j \in J} \mathcal{T}_j \right] \cap \varprojlim_{j \in J} X_j\right) \\ &= \left[\bigotimes_{j \in J} \mathcal{T}_j \right] \cap \varprojlim_{j \in J} X_j. \end{aligned}$$

- Define $\mathcal{F} \subset \bigotimes_{j \in J} \mathcal{T}_j$ by $\mathcal{F} := \{\text{proj}_j^{-1}(V) : j \in J \text{ and } V \in \mathcal{T}_j\}$. Let $F \in \mathcal{F} \cap \varprojlim_{j \in J} X_j$. Then there exists $F_0 \in \mathcal{F}$ such that $F = F_0 \cap \varprojlim_{j \in J} X_j$. Thus there exists $j \in J$ and $V \in \mathcal{T}_j$ such that $F_0 = \text{proj}_j^{-1}(V)$. Therefore

$$\begin{aligned}
F &= F_0 \cap \varprojlim_{j \in J} X_j \\
&= \text{proj}_j^{-1}(V) \cap \varprojlim_{j \in J} X_j \\
&= (\text{proj}_j|_{\varprojlim_{j \in J} X_j})^{-1}(V) \\
&= \pi_j^{-1}(V) \\
&\in \mathcal{E}.
\end{aligned}$$

Since $F \in \mathcal{F} \cap \varprojlim_{j \in J} X_j$ is arbitrary, we have that $\mathcal{F} \cap \varprojlim_{j \in J} X_j \subset \mathcal{E}$. Exercise 4.5.1.5 implies that

$$\begin{aligned}
\left[\bigotimes_{j \in J} \mathcal{T}_j \right] \cap \varprojlim_{j \in J} X_j &= \tau(\mathcal{F}) \cap \varprojlim_{j \in J} X_j \\
&= \tau(\mathcal{F} \cap \varprojlim_{j \in J} X_j) \\
&\subset \tau(\mathcal{E}) \\
&= \varprojlim_{j \in J} \mathcal{T}_j.
\end{aligned}$$

Since $\varprojlim_{j \in J} \mathcal{T}_j \subset \left[\bigotimes_{j \in J} \mathcal{T}_j \right] \cap \varprojlim_{j \in J} X_j$ and $\left[\bigotimes_{j \in J} \mathcal{T}_j \right] \cap \varprojlim_{j \in J} X_j \subset \varprojlim_{j \in J} \mathcal{T}_j$, we have that

$$\varprojlim_{j \in J} \mathcal{T}_j = \left[\bigotimes_{j \in J} \mathcal{T}_j \right] \cap \varprojlim_{j \in J} X_j.$$

□

Exercise 4.10.0.5. Let $(J, \leq)$ be a directed poset, $((X_j, \mathcal{T}_j)_{j \in J}, (\pi_{j,k})_{(j,k) \in \leq})$ a projective system of topological spaces. If for each $j \in J$, $(X_j, \mathcal{T}_j)$ is Hausdorff, then $\varprojlim_{j \in J} X_j$ is closed in $(\prod_{j \in J} X_j, \bigotimes_{j \in J} \mathcal{T}_j)$.

Proof. Suppose that for each $j \in J$, $(X_j, \mathcal{T}_j)$ is Hausdorff. Set $X := \varprojlim_{j \in J} X_j$ and $\mathcal{T} := \varprojlim_{j \in J} \mathcal{T}_j$. Exercise 1.9.0.5 implies that $X = \bigcap_{(j,k) \in \leq} \text{Eq}(\pi_{j,k} \circ \text{proj}_k, \text{proj}_j)$. Let $(j, k) \in \leq$. Since $\pi_{j,k}$ is $(\mathcal{T}_k, \mathcal{T}_j)$ -continuous and π_k is $(\mathcal{T}, \mathcal{T}_k)$ -continuous, we have that $\pi_{j,k} \circ \pi_k$ is $(\mathcal{T}, \mathcal{T}_j)$ -continuous. Since $\pi_{j,k} \circ \pi_k$ and π_j are $(\mathcal{T}, \mathcal{T}_j)$ -continuous and $(X_j, \mathcal{T}_j)$ is Hausdorff, Exercise 4.9.0.2 implies that $\text{Eq}(\pi_{j,k} \circ \text{proj}_k, \text{proj}_j)$ is $\mathcal{T}$ -closed. Since $(j, k) \in \leq$ is arbitrary, we have that for each $(j, k) \in \leq$, $\text{Eq}(\pi_{j,k} \circ \text{proj}_k, \text{proj}_j)$ is $\mathcal{T}$ -closed. Since $X = \bigcap_{(j,k) \in \leq} \text{Eq}(\pi_{j,k} \circ \text{proj}_k, \text{proj}_j)$, we have that X is closed in $(\prod_{j \in J} X_j, \bigotimes_{j \in J} \mathcal{T}_j)$. □

Definition 4.10.0.6. Let $(J, \leq)$ be a directed poset $((X_j)_{j \in J}, (\pi_{j,k})_{(j,k) \in \leq})$ be a projective system of topological spaces. Set $(X, (\pi_j)_{j \in J}) := \varprojlim_{j \in J} ((X_j)_{j \in J}, (\pi_{j,k})_{(j,k) \in \leq})$.

- The projective limit $(X, (\pi_j)_{j \in J})$ is said to be **perfect** (change this to a more descriptiv term) if for each $j \in J$, π_j is a quotient map.
- For each $j \in J$, we define
 - the **equivalence relation on X induced by π_j** , denoted $\sim_j \subset X \times X$, by $x \sim_j y$ iff $\pi_j(x) = \pi_j(y)$,

- the **quotient of X induced by π_j** , denoted X_j^Q , by $X_j^Q := X/\sim_j$,
- the **quotient map induced by π_j** , denoted $\pi_j^Q : X \rightarrow X_j^Q$, by $\pi_j^Q(x) := \{y \in X : y \sim_j x\}$.
- the **factor of π_j by π_j^Q** , denoted $\bar{\pi} : X_j^Q \rightarrow X_j$, to be the unique map from X_j^Q into X_j such that $\bar{\pi} \circ \pi_j^Q = \pi_j$ as defined in Exercise 4.8.1.16

Exercise 4.10.0.7. Let $(J, \leq)$ be a directed poset $((X_j)_{j \in J}, (\pi_{j,k})_{(j,k) \in \leq})$ be a projective system of topological spaces. Set $(X, (\pi_j)_{j \in J}) := \varprojlim_{j \in J} ((X_j)_{j \in J}, (\pi_{j,k})_{(j,k) \in \leq})$. Suppose that $(X, (\pi_j)_{j \in J})$ is perfect. Then

1. for each $(j, k) \in \leq$, $\bar{\pi}_j$ is a homeomorphism and there exist unique $\pi_{j,k}^Q : X_k^Q \rightarrow X_j^Q$ such that $\pi_{j,k}^Q$ is continuous, $\bar{\pi}_j \circ \pi_{j,k}^Q = \pi_{j,k}$, $\pi_{j,k} \circ \bar{\pi}_k = \bar{\pi}_j \circ \pi_{j,k}^Q$ and $\pi_{j,k}^Q \circ \pi_k^Q = \pi_j^Q$, i.e. the following diagram commutes:

$$\begin{array}{ccc}
 X_k & \xrightarrow{\pi_{j,k}} & X_j \\
 \bar{\pi}_k \uparrow & & \uparrow \bar{\pi}_j \\
 X_k^Q & \xrightarrow{\pi_{j,k}^Q} & X_j^Q \\
 \pi_k \swarrow & & \searrow \pi_j \\
 & X &
 \end{array}$$

2. $((X_j^Q)_{j \in J}, (\pi_{j,k}^Q)_{(j,k) \in \leq})$ is a projective system of topological spaces.
3. $\varprojlim_{j \in J} X_j^Q$ is homeomorphic to X . or define inverse limit generally and show that X is an inverse limit for the new system $(X_j^Q, (\pi_{j,k}^Q)_{(j,k) \in \leq})$.

Proof.

1. Let $j \in J$. Since $(X, (\pi_{j'})_{j' \in J})$ is perfect, π_j is a quotient map. Since $\pi_{j,k} \circ \pi_k = \pi_j$, Exercise 4.8.1.21 implies that there exist unique $\bar{\pi}_j : X_j^Q \rightarrow X_j$, $\pi_{j,k}^Q : X_k^Q \rightarrow X_j^Q$ such that $\bar{\pi}_j$ is a homeomorphism, $\pi_{j,k}^Q$ is continuous, $\bar{\pi}_j \circ \pi_{j,k}^Q = \pi_{j,k}$, $\pi_{j,k} \circ \bar{\pi}_k = \bar{\pi}_j \circ \pi_{j,k}^Q$ and $\pi_{j,k}^Q \circ \pi_k^Q = \pi_j^Q$.
2. Let $j, k, l \in J$. Suppose that $j \leq k$ and $k \leq l$. Since $((X_{j'})_{j' \in J}, (\pi_{j',k'})_{(j',k') \in \leq})$ is a projective system of topological spaces, $\pi_{j,j} = \text{id}_{X_j}$ and $\pi_{j,k} \circ \pi_{k,l} = \pi_{j,l}$.

(a) Since

$$\begin{aligned}
 \pi_{j,j} \circ \bar{\pi}_j &= \text{id}_{X_j} \circ \bar{\pi}_j \\
 &= \bar{\pi}_j \\
 &= \bar{\pi}_j \circ \text{id}_{X_j^Q},
 \end{aligned}$$

uniqueness of $\pi_{j,j}^Q$ implies that $\pi_{j,j}^Q = \text{id}_{X_j^Q}$.

(b) By construction

$$\begin{aligned}
 \pi_{j,k}^Q \circ \pi_{k,l}^Q &= (\bar{\pi}_j^{-1} \circ \pi_{j,k} \circ \bar{\pi}_k) \circ (\bar{\pi}_k^{-1} \circ \pi_{k,l} \circ \bar{\pi}_l) \\
 &= (\bar{\pi}_j^{-1} \circ \pi_{j,k}) \circ (\bar{\pi}_k \circ \bar{\pi}_k^{-1}) \circ (\pi_{k,l} \circ \bar{\pi}_l) \\
 &= (\bar{\pi}_j^{-1} \circ \pi_{j,k}) \circ \text{id}_{X_k} \circ (\pi_{k,l} \circ \bar{\pi}_l) \\
 &= (\bar{\pi}_j^{-1} \circ \pi_{j,k}) \circ (\pi_{k,l} \circ \bar{\pi}_l) \\
 &= \bar{\pi}_j^{-1} \circ (\pi_{j,k} \circ \pi_{k,l}) \circ \bar{\pi}_l \\
 &= \bar{\pi}_j^{-1} \circ \pi_{j,l} \circ \bar{\pi}_l \\
 &= \pi_{j,l}^Q
 \end{aligned}$$

Therefore $((X/\sim_j)_{j \in J}, (\pi_{j,k}^Q)_{(j,k) \in \leq})$ is a projective system of topological spaces.

(c) **FINISH!!!**

□

Note 4.10.0.8. Here we started with a given projective system (diagram of objects with morphisms). These objects and morphisms induce new objects and morphisms which are tied via isomorphisms to the originals. These new objects and morphisms have the same structure as the original (projective system). what is this phenomenon in general?

Exercise 4.10.0.9. **move to compactness section** Let $(J, \leq)$ be a directed poset $((X_j)_{j \in J}, (\pi_{j,k})_{(j,k) \in \leq})$ be a projective system of topological spaces. Set $(X, (\pi_j)_{j \in J}) := \varprojlim_{j \in J} ((X_j)_{j \in J}, (\pi_{j,k})_{(j,k) \in \leq})$. For each $j \in J$, define $C_j \subset C(X)$ by

$$C_j := \{f \in C(X) : \text{for each } x, y \in X, \pi_j(x) = \pi_j(y) \text{ implies that } f(x) = f(y)\}$$

and define $C \subset C(X)$ by $C := \bigcup_{j \in J} C_j$. Suppose that for each $j \in J$, X_j is a compact Hausdorff space and π_j is surjective. Then

1. $C_j = \{f \in C(X) : f \text{ is } \sim_j\text{-invariant}\}$
2. C is a subalgebra of $C(X)$,
3. C is dense in $C(X)$
4. for each $f \in C_j$, there exists a unique $\bar{f}_j \in C(X_j^Q)$ such that $\bar{f}_j \circ \pi_j^Q = f$.

Proof.

1. Clear.
2. Let $f_1, f_2 \in C$ and $\lambda \in \mathbb{C}$. Then there exist $j_1, j_2 \in J$ such that $f_1 \in C_{j_1}$ and $f_2 \in C_{j_2}$. Since J is directed, there exists $j_0 \in J$ such that $j_0 \geq j_1, j_2$. Let $x, y \in X$. Suppose that $\pi_{j_0}(x) = \pi_{j_0}(y)$. Then

$$\begin{aligned} \pi_{j_1}(x) &= \pi_{j_1, j_0} \circ \pi_{j_0}(x) \\ &= \pi_{j_1, j_0} \circ \pi_{j_0}(y) \\ &= \pi_{j_1}(y) \end{aligned}$$

and similarly, $\pi_{j_2}(x) = \pi_{j_2}(y)$. Therefore $f_1(x) = f_1(y)$ and $f_2(x) = f_2(y)$. Hence

$$\begin{aligned} (f_1 + \lambda f_2)(x) &= f_1(x) + \lambda f_2(x) \\ &= f_1(y) + \lambda f_2(y) \\ &= (f_1 + \lambda f_2)(y) \end{aligned}$$

and

$$\begin{aligned} (f_1 \cdot f_2)(x) &= f_1(x)f_2(x) \\ &= f_1(y)f_2(y) \\ &= (f_1 \cdot f_2)(y). \end{aligned}$$

Since $x, y \in X$ with $\pi_{j_0}(x) = \pi_{j_0}(y)$ are arbitrary, we have that for each $x, y \in X$, $\pi_{j_0}(x) = \pi_{j_0}(y)$ implies that $(f_1 + \lambda f_2)(x) = (f_1 + \lambda f_2)(y)$ and $(f_1 \cdot f_2)(x) = (f_1 \cdot f_2)(y)$. Thus

$$\begin{aligned} f_1 + \lambda f_2 &\in C_{j_0} \\ &\subset C \end{aligned}$$

and

$$\begin{aligned} f_1 \cdot f_2 &\in C_{j_0} \\ &\subset C \end{aligned}$$

Since $f_1, f_2 \in C$ and $\lambda \in \mathbb{C}$ are arbitrary, for each $f_1, f_2 \in C$ and $\lambda \in \mathbb{C}$, $f_1 + \lambda f_2 \in C$ and $f_1 \cdot f_2 \in C$. Hence C is a subalgebra of $C(X)$.

3. Since for each $j \in J$, X_j is a compact Hausdorff space, Exercise ?? implies that X is a compact Hausdorff space and Exercise ?? implies that $(X, (\pi_j)_{j \in J})$ is perfect. Let $j \in J$. Since $(X, (\pi_{j'})_{j' \in J})$ is perfect, Exercise 4.10.0.7 implies that $\bar{\pi} : X/\sim_j \rightarrow X_j$ is a homeomorphism. Since X_j is a compact Hausdorff space and X_j is homeomorphic to $X/\sim_j$, we have that $X/\sim_j$ is a compact Hausdorff space. **Exercise 4.12.4.3 implies that $\sim_j$ is closed in $X \times X$.** Let $x, y \in X$. Suppose that $x \neq y$. Exercise ?? (**Make exercise in set theory section about $x = y$ iff for each j , $\pi_j(x) = \pi_j(y)$**) implies that there exists $j \in J$ such that $\pi_j(x) \neq \pi_j(y)$. Since $\bar{\pi} : X_j^Q \rightarrow X_j$ is a homeomorphism,

$$\begin{aligned}\pi_j^Q(x) &= \bar{\pi}_j^{-1} \circ \pi_j(x) \\ &\neq \bar{\pi}_j^{-1} \circ \pi_j(y) \\ &= \pi_j^Q(y).\end{aligned}$$

Since $X/\sim_j$ is a compact Hausdorff space, Exercise ?? (**make exercise in section on uryshon thoerem to get continuous map separting points on compact hausdorff space**) implies that there exists $\phi \in C(X/\sim_j)$ such that $\phi(\pi_j^Q(x)) \neq \phi(\pi_j^Q(y))$. Define $f \in C(X)$ by $f := \phi \circ \pi_j^Q$. By construction $f \in C_j \subset C$ and $f(x) \neq f(y)$. Since $x, y \in X$ with $x \neq y$ are arbitrary, we have that for each $x, y \in X$, $x \neq y$ implies that there exists $f \in C$ such that $f(x) \neq f(y)$. Thus C separates the points of X . **finish off stone-weierstrass theorem assumptions to get denseness**

4. Let $f \in C_j$. Since f is continuous and $\sim_j$ -invariant, Exercise 4.8.1.16 implies that there exists a unique $\bar{f}_j \in C(X_j^Q)$ such that $\bar{f}_j \circ \pi_j^Q = f$. □

4.11 Open Covers

4.11.1 Introduction

Definition 4.11.1.1. **replace 'open cover' with $\text{Cover}^{\text{opn}}$** Let $(X, \mathcal{T})$ be a topological space and $E \subset X$.

- Let $\mathcal{U} \subset \mathcal{P}(X)$. Then $\mathcal{U}$ is said to be a **$\mathcal{T}$ -open cover of E** , or equivalently, an **open cover of E in $(X, \mathcal{T})$** if
 1. $\mathcal{U} \subset \mathcal{T}$
 2. $\mathcal{U} \in \text{Cover}(E)$.
- We define $\text{Cover}^{\text{opn}}(E, \mathcal{T}) \subset \mathcal{P}(\mathcal{T})$ by

$$\text{Cover}^{\text{opn}}(E, \mathcal{T}) := \{\mathcal{U} \subset \mathcal{P}(X) : \mathcal{U} \text{ is an open cover of } E \text{ in } (X, \mathcal{T})\}$$

Note 4.11.1.2. When the context is clear,

- we may say “open cover of E ” instead of “ $\mathcal{T}$ -open cover of E ”
- we may write $\text{Cover}^{\text{opn}}(E)$ in place of $\text{Cover}^{\text{opn}}(E, \mathcal{T})$.

Exercise 4.11.1.3. **might need to define preimage and image maps in analysis notes set theory section f^*, f_* , then recall definition in algebra notes section on galois connections from set theory notes once compiled.** Let $(X, \mathcal{T}_X)$ and $(Y, \mathcal{T}_Y)$ be topological spaces, $f : X \rightarrow Y$ $(\mathcal{T}_X, \mathcal{T}_Y)$ -continuous and $E \subset Y$. Then for each $\mathcal{V} \in \text{Cover}^{\text{opn}}(E, \mathcal{T}_Y)$, $f^*(\mathcal{V}) \in \text{Cover}^{\text{opn}}(f^*(E), \mathcal{T}_X)$.

Proof. Let $\mathcal{V} \in \text{Cover}^{\text{opn}}(E, \mathcal{T}_Y)$. Define $\mathcal{U} \subset \mathcal{P}(X)$ by $\mathcal{U} := f^*(\mathcal{V})$.

- Since f is $(\mathcal{T}_X, \mathcal{T}_Y)$ -continuous and $\mathcal{V} \subset \mathcal{T}_Y$, we have that $\mathcal{U} \subset \mathcal{T}_X$.
- Since $E \subset \bigcup_{V \in \mathcal{V}} V$, we have that

$$\begin{aligned}f^{-1}(E) &\subset f^{-1}\left(\bigcup_{V \in \mathcal{V}} V\right) \\ &= \bigcup_{V \in \mathcal{V}} f^{-1}(V) \\ &= \bigcup_{U \in \mathcal{U}} U.\end{aligned}$$

Hence $\mathcal{U} \in \text{Cover}^{\text{opn}}(f^*E)$. □

4.11.2 Subspace Topology

Exercise 4.11.2.1. Let $(X, \mathcal{T})$ be a topological space, $\mathcal{U} \subset \mathcal{T}$, $A \subset X$ and $E \subset A$. If $\mathcal{U}$ is an open cover of E in $(X, \mathcal{T})$, then $\mathcal{U} \cap A$ is an open cover of E in $(A, \mathcal{T} \cap A)$.

Proof. Suppose that $\mathcal{U}$ is an open cover of E in $(X, \mathcal{T})$.

1. Since $\mathcal{U} \subset \mathcal{T}$, $\mathcal{U} \cap A \subset \mathcal{T} \cap A$.
2. Since $\mathcal{U}$ is a cover of E in X , $E \subset \bigcup_{U \in \mathcal{U}} U$. Since $E \subset A$, we have that

$$\begin{aligned} E &= E \cap A \\ &\subset \left[\bigcup_{U \in \mathcal{U}} U \right] \cap A \\ &= \bigcup_{U \in \mathcal{U}} [U \cap A] \\ &= \bigcup_{U \in \mathcal{U} \cap A} U. \end{aligned}$$

Hence $\mathcal{U} \cap A$ is a cover of E in A .

Since $\mathcal{U} \cap A \subset \mathcal{T} \cap A$ and $\mathcal{U} \cap A$ is a cover of E in A , we have that $\mathcal{U} \cap A$ is an open cover of E in $(A, \mathcal{T} \cap A)$. □

4.12 Separation Axioms

4.12.1 Introduction

Reformat using $(X, \mathcal{T})$ and writing $U \in \mathcal{T}$ instead of saying U is open.

Definition 4.12.1.1. Let X be a topological space. Then X is said to be

1. **T_0 of Kolmogorov** if for each $x, y \in X$, if $x \neq y$, then there exists $U \subset X$ such that
 - (a) U is open
 - (b) $(x, y) \in U \times U^c$ or $(x, y) \in U^c \times U$
2. **T_1** if for each $x, y \in X$, if $x \neq y$, then there exists $U \in \mathcal{N}(x)$ such that U is open and $y \notin U$.
3. **T_2 or Hausdorff** if for each $x, y \in X$, if $x \neq y$, then there exist $U \in \mathcal{N}(x)$ and $V \in \mathcal{N}(y)$ such that U and V are open and $U \cap V = \emptyset$.
4. **T_3 or regular** if X is T_1 and for each $A \subset X$ and $x \in A^c$, if A is closed, then there exists $U \in \mathcal{N}(A)$ and $V \in \mathcal{N}(x)$ such that U and V are open and $U \cap V = \emptyset$.
5. **T_4 or normal** if X is T_1 and for each $A, B \subset X$, if A and B are closed and $A \cap B = \emptyset$, then there exists $U \in \mathcal{N}(A)$ and $V \in \mathcal{N}(B)$ such that U and V are open and $U \cap V = \emptyset$.

Note 4.12.1.2. Some authors do not require the T_1 assumption for regularity or normality.

Exercise 4.12.1.3. Let X be a topological space. Then the following are equivalent:

1. X is T_1
2. for each $x \in X$, $\{x\}$ is closed
3. for each $A \subset X$, $A = \bigcap_{U \in \mathcal{N}(A)} U$

Proof.

- (1) $\implies$ (2):
Suppose that X is T_1 . Let $x \in X$. Since X is T_1 , for each $a \in \{x\}^c$, there exists $U_a \in \mathcal{N}(a)$ such that U_a is open and $U_a \subset \{x\}^c$.
Therefore

$$\{x\}^c = \bigcup_{a \in \{x\}^c} U_a$$

which is open. Hence

$$\{x\} = \bigcap_{a \in \{x\}^c} U_a^c$$

which is closed.

- (2) $\implies$ (3):
Suppose that for each $x \in X$, $\{x\}$ is closed. Clearly, $A \subset \bigcap_{U \in \mathcal{N}(A)} U$. Since for each $x \in A^c$, $\{x\}^c \in \mathcal{N}(A)$, we have that

$$\begin{aligned} \bigcap_{U \in \mathcal{N}(A)} U &\subset \bigcap_{x \in A^c} \{x\}^c \\ &= \left(\bigcup_{x \in A^c} \{x\} \right)^c \\ &= (A^c)^c \\ &= A \end{aligned}$$

- (3) $\implies$ (1):

Suppose that for each $A \subset X$, $A = \bigcap_{U \in \mathcal{N}(A)} U$. Let $x, y \in X$. Suppose that $x \neq y$. Since $\{x\} = \bigcap_{V \in \mathcal{N}(x)} V$, $y \notin \bigcap_{V \in \mathcal{N}(x)} V$. Thus there exists $V \in \mathcal{N}(x)$ such that $y \notin V$. Set $U = \text{Int } V$. Then $U \in \mathcal{N}(x)$, U is open and $y \notin U$. Since $x, y \in X$ are arbitrary, X is T_1 . □

Exercise 4.12.1.4. Let $(X, \mathcal{T})$ be a topological space and $A \subset X$. Suppose that $(X, \mathcal{T})$ is T_1 . If A is countable, then A is an F_σ -set.

Proof. Suppose that A is countable. Exercise 4.12.1.3 implies that for each $x \in A$, $\{x\}$ is $\mathcal{T}$ -closed. Since $A = \bigcup_{x \in A} \{x\}$ and A is countable, we have that A is an F_σ -set. □

Exercise 4.12.1.5. Let X be a topological space. Then

1. X is T_1 implies that X is T_0
2. X is T_2 implies that X is T_1
3. X is T_3 implies that X is T_2
4. X is T_4 implies that X is T_3

Proof. Clear by definition and the previous exercise. □

Note 4.12.1.6. Let X be a set, we recall Definition 1.5.0.9 of Δ_X and $\Delta_{X^\mathbb{N}}$.

Exercise 4.12.1.7. Let X be a topological space. Then the following are equivalent:

1. X is Hausdorff
2. for each net $(x_\alpha)_{\alpha \in A} \subset X$ and $x, y \in X$, if $x_\alpha \rightarrow x$ and $x_\alpha \rightarrow y$, then $x = y$.
3. Δ_X is closed in $X \times X$.

Proof.

- (1) $\implies$ (2):

Suppose that X is Hausdorff. Let $(x_\alpha)_{\alpha \in A} \subset X$ be a net and $x, y \in X$. Suppose that $x_\alpha \rightarrow x$ and $x_\alpha \rightarrow y$. For the sake of contradiction, suppose that $x \neq y$. Then there exist $U \in \mathcal{N}(x)$ and $V \in \mathcal{N}(y)$ such that U and V are open and $U \cap V = \emptyset$. Since $x_\alpha \rightarrow x$, $(x_\alpha)_{\alpha \in A}$ is eventually in U and there exists $\beta_x \in A$ such that for each $\alpha \in A$, $\alpha \geq \beta_x$ implies that $x_\alpha \in U$. Since $x_\alpha \rightarrow y$, $(x_\alpha)_{\alpha \in A}$ is eventually in V and there exists $\beta_y \in A$ such that for each $\alpha \in A$, $\alpha \geq \beta_y$ implies that $x_\alpha \in V$. Since A is directed, there exists $\beta \in A$ such that $\beta \geq \beta_x, \beta_y$. Hence

$$\begin{aligned} x_\beta &\in U \cap V \\ &= \emptyset \end{aligned}$$

which is a contradiction. So $x = y$.

- (2) $\implies$ (3):

Let $(x_\alpha, y_\alpha)_{\alpha \in A} \subset \Delta_X$ be a net and $(x, y) \in X \times X$. Then for each $\alpha \in A$, $x_\alpha = y_\alpha$. Suppose that $(x_\alpha, y_\alpha) \rightarrow (x, y)$. So $x_\alpha \rightarrow x$ and $x_\alpha \rightarrow y$. Hence $x = y$ and $(x, y) \in \Delta_X$. Thus Δ_X is closed.

- (3) $\implies$ (1):

Suppose that Δ_X is closed. Let $x, y \in X$. Suppose that $x \neq y$. Then $(x, y) \in \Delta_X^c$. Recall that $\mathcal{B} = \{A \times B : A, B \subset X \text{ and } A, B \text{ are open}\}$ is a basis for the product topology on $X \times X$. Since Δ_X^c is open and $(x, y) \in \Delta_X^c$, there exist $A \times B \in \mathcal{B}$ such that $(x, y) \in A \times B \subset \Delta_X^c$. Suppose that $A \cap B \neq \emptyset$. Then there exists $z \in A \cap B$. Hence $(z, z) \in A \times B$. This is a contradiction since $A \times B \subset \Delta_X^c$. Thus $x \in A$, $y \in B$ and $A \cap B = \emptyset$ and A, B are open. Since $x, y \in X$ are arbitrary, X is Hausdorff. □

Exercise 4.12.1.8. Let $(X, \mathcal{T})$ be a topological space. If X is Hausdorff, then $\Delta_{X^{\mathbb{N}}}$ is closed in $(X^{\mathbb{N}}, \mathcal{T}^{\otimes \mathbb{N}})$.

Proof. Suppose that X is Hausdorff. Let $(x_\alpha)_{\alpha \in A} \subset \Delta_{X^{\mathbb{N}}}$ and $x \in X^{\mathbb{N}}$. Suppose that $x_\alpha \rightarrow x$. Then for each $n \in \mathbb{N}$, $\pi_n(x_\alpha) \xrightarrow{\alpha} \pi_n(x)$. Since $(x_\alpha)_{\alpha \in A} \subset \Delta_{X^{\mathbb{N}}}$, we have that for each $\alpha \in A$ and $m, n \in \mathbb{N}$, $\pi_m(x_\alpha) = \pi_n(x_\alpha)$. Let $n \in \mathbb{N}$. Then

$$\begin{aligned}\pi_n(x_\alpha) &= \pi_1(x_\alpha) \\ &\xrightarrow{\alpha} \pi_1(x).\end{aligned}$$

Since $\pi_n(x_\alpha) \xrightarrow{\alpha} \pi_n(x)$, $\pi_n(x_\alpha) \xrightarrow{\alpha} \pi_1(x)$ and X is Hausdorff, Exercise 4.12.1.7 implies that $\pi_1(x) = \pi_n(x)$. Since $n \in \mathbb{N}$ is arbitrary, we have that for each $m, n \in \mathbb{N}$,

$$\begin{aligned}\pi_m(x) &= \pi_1(x) \\ &= \pi_n(x).\end{aligned}$$

Hence $x \in \Delta_{X^{\mathbb{N}}}$. Since $(x_\alpha)_{\alpha \in A} \subset \Delta_{X^{\mathbb{N}}}$ and $x \in X^{\mathbb{N}}$ with $x_\alpha \rightarrow x$ are arbitrary, we have that for each $(x_\alpha)_{\alpha \in A} \subset \Delta_{X^{\mathbb{N}}}$ and $x \in X^{\mathbb{N}}$, if $x_\alpha \rightarrow x$, then $x \in \Delta_{X^{\mathbb{N}}}$. Hence $\Delta_{X^{\mathbb{N}}}$ is closed in $(X^{\mathbb{N}}, \mathcal{T}^{\otimes \mathbb{N}})$. $\square$

Exercise 4.12.1.9. Let X be a set, $(Y_\alpha, \mathcal{T}_\alpha)_{\alpha \in A}$ a collection of topological spaces and $\mathcal{F} \in \prod_{\alpha \in A} Y_\alpha^X$. Suppose that for each $\alpha \in A$, $(Y_\alpha, \mathcal{T}_\alpha)$ is Hausdorff. If $\mathcal{F}$ separates the points of X , then $(X, \tau_X(\mathcal{F}))$ is Hausdorff.

Proof. Write $\mathcal{F} = (f_\alpha)_{\alpha \in A}$. Suppose that $\mathcal{F}$ separates the points of X . Let $x_1, x_2 \in X$. Suppose that $x_1 \neq x_2$. Then there exists $\alpha \in A$ such that $f_\alpha(x_1) \neq f_\alpha(x_2)$. Since $(Y_\alpha, \mathcal{T}_\alpha)$ is Hausdorff, there exists $U_0, V_0 \in \mathcal{T}_\alpha$ such that $f_\alpha(x_1) \in U_0$, $f_\alpha(x_2) \in V_0$ and $U_0 \cap V_0 = \emptyset$. Define $U, V \in \tau_X(\mathcal{F})$ by $U := f_\alpha^{-1}(U_0)$ and $V := f_\alpha^{-1}(V_0)$. Then $x_1 \in U$, $x_2 \in V$ and

$$\begin{aligned}U \cap V &= f_\alpha^{-1}(U_0) \cap f_\alpha^{-1}(V_0) \\ &= f_\alpha^{-1}(U_0 \cap V_0) \\ &= f_\alpha^{-1}(\emptyset) \\ &= \emptyset.\end{aligned}$$

Since $x_1, x_2 \in X$ with $x_1 \neq x_2$ are arbitrary, we have that for each $x_1, x_2 \in X$, $x_1 \neq x_2$ implies that there exists $U, V \in \tau_X(\mathcal{F})$ such that $x_1 \in U$, $x_2 \in V$ and $U \cap V = \emptyset$. Hence $(X, \tau_X(\mathcal{F}))$ is Hausdorff. $\square$

Exercise 4.12.1.10. Let X be a topological space. Suppose that X is T_1 . Then X is regular iff for each $x \in X$ and $U \in \mathcal{N}(x)$, U is open implies that there exists $V \in \mathcal{N}(x)$ such that $\text{cl } V \subset U$.

Proof.

• ($\implies$):

Suppose that X is regular. Let $x \in X$ and $U \in \mathcal{N}(x)$. Suppose that U is open. Then U^c is closed. Since $x \notin U^c$, there exists $V_x \in \mathcal{N}(x)$ and $V_{U^c} \in \mathcal{N}(U^c)$ such that V_x and V_{U^c} are open and $V_x \cap V_{U^c} = \emptyset$. Therefore, $V_{U^c}^c$ is closed and $V_x \subset V_{U^c}^c \subset U$. Hence

$$\begin{aligned}x &\in V_x \\ &\subset \text{cl } V_x \\ &\subset \text{cl } V_{U^c}^c \\ &= V_{U^c}^c \\ &\subset U\end{aligned}$$

• ($\impliedby$):

Suppose that for each $x \in X$ and $U \in \mathcal{N}(x)$, U is open implies that there exists $V \in \mathcal{N}(x)$ such that $\text{cl } V \subset U$. Let $x \in X$ and $A \subset X$. Suppose that A is closed and $x \notin A$. Then A^c is open and $x \in A^c$. By assumption, there exists $V \in \mathcal{N}(x)$ such that $\text{cl } V \subset A^c$. Set $U_x = \text{Int } V$ and $U_A = \text{Int } V^c$. Then

$$\begin{aligned}A &\subset (\text{cl } V)^c \\ &= \text{Int } V^c \\ &= U_A\end{aligned}$$

so that U_x and U_A are open, $U_x \in \mathcal{N}(x)$, $U_A \in \mathcal{N}(A)$ and $U_x \cap U_A = \emptyset$. Hence X is regular.

Exercise 4.12.1.11. lemma for Uryshohns lemma

Proof. FINISH!!!

Exercise 4.12.1.12. Urysohn's Lemma for Normal Spaces:

Let X be a topological space. Suppose that X is normal. Let $A, B \subset X$. Suppose that A and B are closed and $A \cap B = \emptyset$. Then there exists $f \in C(X, [0, 1])$ such that $f|_A = 0$ and $f|_B = 1$.

Proof. FINISH!!!

Exercise 4.12.1.13. Tietze Extension Theorem for Normal Spaces:

Let X be a topological space. Suppose that X is normal. Let $A, B \subset X$. Suppose that A and B are closed and $A \cap B = \emptyset$. Then there exists $f \in C(X, [0, 1])$ such that $f|_A = 0$ and $f|_B = 1$.

Proof. FINISH!!!

4.12.2 Separation and Subspaces

Exercise 4.12.2.1. Let $(X, \mathcal{T})$ be a topological space and $A \subset X$. If $(X, \mathcal{T})$ is T_1 , then $(A, \mathcal{T} \cap A)$ is T_1 .

Proof. Suppose that $(X, \mathcal{T})$ is T_1 . Let $x \in A$. Since $(X, \mathcal{T})$ is T_1 , $\{x\}$ is closed in X . Thus $\{x\} = \{x\} \cap A$ is closed in $(A, \mathcal{T} \cap A)$. Since $x \in A$ is arbitrary, $(A, \mathcal{T} \cap A)$ is T_1 . $\square$

Exercise 4.12.2.2. Let $(X, \mathcal{T})$ be a topological space and $A \subset X$. If $(X, \mathcal{T})$ is Hausdorff, then $(A, \mathcal{T} \cap A)$ is Hausdorff.

Proof. Suppose that $(X, \mathcal{T})$ is Hausdorff. Let $x, y \in A$. Since $(X, \mathcal{T})$ is Hausdorff, there exist $U' \in \mathcal{N}(x)$, $V' \in \mathcal{N}(y)$ such that $U, V \in \mathcal{T}$ and $U' \cap V' = \emptyset$. Set $U = U' \cap A$ and $V = V' \cap A$. Then $U, V \in \mathcal{T} \cap A$, $x \in U$, $y \in V$ and

$$\begin{aligned} U \cap V &= (U' \cap A) \cap (V' \cap A) \\ &= (U' \cap V') \cap A \\ &= \emptyset \end{aligned}$$

Since $x, y \in A$ are arbitrary, $(A, \mathcal{T} \cap A)$ is Hausdorff. $\square$

Exercise 4.12.2.3. Let $(X, \mathcal{T})$ be a topological space and $A \subset X$. If $(X, \mathcal{T})$ is regular, then $(A, \mathcal{T} \cap A)$ is regular.

Proof. Suppose that $(X, \mathcal{T})$ is regular. Let $x \in A$ and $U \in \mathcal{N}(x)(\mathcal{T} \cap A)$. Suppose that $U \in \mathcal{T} \cap A$. Then there exists $U' \in \mathcal{T}$ such that $U = U' \cap A$. Since $(X, \mathcal{T})$ is regular, there exist $V' \in \mathcal{N}(x)(\mathcal{T})$ such that, $\text{cl}_{\mathcal{T}} V' \subset U'$. Set $V = V' \cap A$. Then $V \in \mathcal{N}(x)(\mathcal{T} \cap A)$

$$\begin{aligned} \text{cl}_{\mathcal{T} \cap A} V &= \text{cl}_{\mathcal{T}} V' \cap A \\ &\subset U' \cap A \end{aligned}$$

Since $x \in A$ are arbitrary, $(A, \mathcal{T} \cap A)$ is regular.

FINISH!!! $\square$

4.12.3 Separation and Product Spaces

Exercise 4.12.3.1. Let $(X_\alpha, \mathcal{T}_\alpha)_{\alpha \in A}$ be a collection of topological spaces. Set $X = \prod_{\alpha \in A} X_\alpha$ and denote the product topology on X by $\mathcal{T}_X$. If for each $\alpha \in A$, $(X_\alpha, \mathcal{T}_\alpha)$ is T_1 , then $(X, \mathcal{T}_X)$ is T_1 .

Proof. Suppose that for each $\alpha \in A$, $(X_\alpha, \mathcal{T}_\alpha)$ is T_1 . Let $(x_\alpha)_{\alpha \in A}, (y_\alpha)_{\alpha \in A} \in X$. Suppose that $(x_\alpha)_{\alpha \in A} \neq (y_\alpha)_{\alpha \in A}$. Then there exists $\alpha_0 \in A$ such that $x_{\alpha_0} \neq y_{\alpha_0}$. Then there exists $U_{\alpha_0} \in \mathcal{T}_{\alpha_0}$ such that $x_{\alpha_0} \in U_{\alpha_0}$ and $y_{\alpha_0} \notin U_{\alpha_0}$. Set $U = \pi_{\alpha_0}^{-1}(U_{\alpha_0})$. Then $U \in \mathcal{T}_X$, $(x_\alpha)_{\alpha \in A} \in U$ and $(y_\alpha)_{\alpha \in A} \notin U$. Since $(x_\alpha)_{\alpha \in A}, (y_\alpha)_{\alpha \in A} \in X$ are arbitrary, $(X, \mathcal{T}_X)$ is T_1 . $\square$

Exercise 4.12.3.2. Let $(X_\alpha, \mathcal{T}_\alpha)_{\alpha \in A}$ be a collection of topological spaces. Set $X = \prod_{\alpha \in A} X_\alpha$ and $\mathcal{T} := \bigotimes_{\alpha \in A} \mathcal{T}_\alpha$. If for each $\alpha \in A$, $(X_\alpha, \mathcal{T}_\alpha)$ is Hausdorff, then $(X, \mathcal{T})$ is Hausdorff.

Proof. Suppose that for each $\alpha \in A$, $(X_\alpha, \mathcal{T}_\alpha)$ is Hausdorff. Let $(x_\alpha)_{\alpha \in A}, (y_\alpha)_{\alpha \in A} \in X$. Suppose that $(x_\alpha)_{\alpha \in A} \neq (y_\alpha)_{\alpha \in A}$. Then there exists $\alpha_0 \in A$ such that $x_{\alpha_0} \neq y_{\alpha_0}$. Then there exists $U_{\alpha_0}, V_{\alpha_0} \in \mathcal{T}_{\alpha_0}$ such that $x_{\alpha_0} \in U_{\alpha_0}$, $y_{\alpha_0} \in V_{\alpha_0}$ and $U_{\alpha_0} \cap V_{\alpha_0} = \emptyset$. Set $U = \pi_{\alpha_0}^{-1}(U_{\alpha_0})$ and $V = \pi_{\alpha_0}^{-1}(V_{\alpha_0})$. Then $U, V \in \mathcal{T}$, $(x_\alpha)_{\alpha \in A} \in U$, $(y_\alpha)_{\alpha \in A} \in V$ and

$$\begin{aligned} U \cap V &= \pi_{\alpha_0}^{-1}(U_{\alpha_0}) \cap \pi_{\alpha_0}^{-1}(V_{\alpha_0}) \\ &= \pi_{\alpha_0}^{-1}(U_{\alpha_0} \cap V_{\alpha_0}) \\ &= \pi_{\alpha_0}^{-1}(\emptyset) \\ &= \emptyset \end{aligned}$$

Since $(x_\alpha)_{\alpha \in A}, (y_\alpha)_{\alpha \in A} \in X$ are arbitrary, $(X, \mathcal{T})$ is Hausdorff. □

Exercise 4.12.3.3. Let $(X_\alpha, \mathcal{T}_\alpha)_{\alpha \in A}$ be a collection of topological spaces. Set $X = \prod_{\alpha \in A} X_\alpha$ and denote the product topology on X by $\mathcal{T}_X$. If for each $\alpha \in A$, $(X_\alpha, \mathcal{T}_\alpha)$ is regular, then $(X, \mathcal{T}_X)$ is regular.

Proof. Let $x \in X$ and $U \in \mathcal{N}(x)$. Suppose that U is open. Set

$$\mathcal{B} = \left\{ \prod_{\alpha \in A} B_\alpha : \text{for each } \alpha \in A, B_\alpha \in \mathcal{T}_\alpha \text{ and } \#\{\alpha \in A : B_\alpha \neq X_\alpha\} < \infty \right\}$$

Then $\mathcal{B}$ is a basis for $\mathcal{T}_X$. So for each $\alpha \in A$, there exist $U_\alpha \in \mathcal{T}_\alpha$ such that $\#\{\alpha \in A : B_\alpha \neq X_\alpha\} < \infty$ and $x \in \prod_{\alpha \in A} U_\alpha \subset U$. Set $J = \{\alpha \in A : B_\alpha \neq X_\alpha\}$. Let $\alpha \in A$. Suppose that $\alpha \in J$. Then $x_\alpha \in U_\alpha$. Since $U_\alpha \in \mathcal{N}(x)$ is an open neighborhood of x_α and X_α is regular, the previous exercise implies that there exists $V_\alpha \in \mathcal{N}(x_\alpha)$ such that $\text{cl } V_\alpha \subset U_\alpha$. If $\alpha \in J^c$, set $V_\alpha = X_\alpha$. Define $V = \prod_{\alpha \in A} V_\alpha$. Then $V \in \mathcal{N}(x)$ and an exercise in the section on the product topology implies that

$$\begin{aligned} \text{cl } V &= \text{cl } \prod_{\alpha \in A} V_\alpha \\ &= \prod_{\alpha \in A} \text{cl } V_\alpha \\ &\subset \prod_{\alpha \in A} U_\alpha \\ &\subset U \end{aligned}$$

□

4.12.4 Separation and Quotient Spaces

Definition 4.12.4.1. Let $(X, \mathcal{T})$ be a topological space. Define $\sim_0 \subset X \times X$ by

$$\sim_0 = \{(x, y) \in X \times X : \text{for each } U \in \mathcal{T}, x \in U \text{ iff } y \in U\}.$$

Exercise 4.12.4.2. Let $(X, \mathcal{T})$ be a topological space. Then

1. $\sim_0$ is an equivalence relation on X ,
2. for each $x, y \in X$, $x \not\sim_0 y$ iff there exists $U \in \mathcal{T}$ such that $(x, y) \in U \times U^c$ or $(x, y) \in U^c \times U$,
3. for each $x, y \in X$, $x \sim_0 y$ iff $\text{cl}\{x\} = \text{cl}\{y\}$,
4. X is T_0 iff for each $x, y \in X$, $x \neq y$ iff $x \not\sim_0 y$,
5. (a) for each $U \in \mathcal{T}$, $\pi^{-1}(\pi(U)) = U$
 (b) $\pi : X \rightarrow X/\sim_0$ is open,
6. $X/\sim_0$ is T_0 ,

7. X is T_0 iff $X/\sim_0 \cong X$.

Proof.

1. Let $x, y, z \in X$.

- Clearly $x \sim_0 x$.
- Clearly $x \sim_0 y$ implies that $y \sim_0 x$.
- Suppose that $x \sim_0 y$ and $y \sim_0 z$. Let $U \in \mathcal{T}$.
 - Suppose that $x \in U$. Since $x \sim_0 y$, we have that $y \in U$. Since $y \sim_0 z$, we have that $z \in U$. Thus $x \in U$ implies that $z \in U$.
 - Similarly, $z \in U$ implies that $x \in U$.

Thus $x \in U$ iff $z \in U$. Since $U \in \mathcal{T}$ is arbitrary, we have that for each $U \in \mathcal{T}$, $x \in U$ iff $z \in U$. Therefore $x \sim_0 z$.

Since $x, y, z \in X$ are arbitrary, we have that $\sim_0$ is an equivalence relation on X .

2. Let $x, y \in X$. By definition,

$$\begin{aligned}
 x \not\sim_0 y &\iff \neg[\forall U \in \mathcal{T}, (x \in U \iff y \in U)] \\
 &\iff \exists U \in \mathcal{T} \text{ s.t. } \neg[(x \in U \implies y \in U) \text{ and } (y \in U \implies x \in U)] \\
 &\iff \exists U \in \mathcal{T} \text{ s.t. } \neg(x \in U \implies y \in U) \text{ or } \neg(y \in U \implies x \in U) \\
 &\iff \exists U \in \mathcal{T} \text{ s.t. } (x \in U \wedge y \notin U) \text{ or } (y \in U \wedge x \notin U) \\
 &\iff \exists U \in \mathcal{T} \text{ s.t. } (x, y) \in U \times U^c \text{ or } (x, y) \in U^c \times U
 \end{aligned}$$

3. Let $x, y \in X$.

- $(\implies)$:
 Suppose that $x \sim_0 y$. Define $\text{clSets } x := \{C \subset X : C \text{ is closed and } x \in C\}$ and $\text{clSets } y := \{C \subset X : C \text{ is closed and } y \in C\}$. Let $C \in \text{clSets } x$. For the sake of contradiction, suppose that $y \notin C$. Then $y \in C^c$. Since $x \sim_0 y$, $C^c \in \mathcal{T}$ and $y \in C^c$, we have that $x \in C^c$. This is a contradiction since $x \in C$. Thus $y \in C$ and therefore $C \in \text{clSets } y$. Since $C \in \text{clSets } x$ is arbitrary, we have that $\text{clSets } x \subset \text{clSets } y$. Similarly, $\text{clSets } y \subset \text{clSets } x$. Thus $\text{clSets } y = \text{clSets } x$. Therefore

$$\begin{aligned}
 \text{cl}\{x\} &= \bigcap_{C \in \text{clSets } x} C \\
 &= \bigcap_{C \in \text{clSets } y} C \\
 &= \text{cl}\{y\}
 \end{aligned}$$

- $(\impliedby)$:
 Suppose that $x \not\sim_0 y$. Then part (2) implies that there exists $U \in \mathcal{T}$ such that $(x, y) \in U \times U^c$ or $(x, y) \in U^c \times U$.
 - Suppose that $(x, y) \in U \times U^c$. Then $U^c \in \text{clSets } y$ and therefore

$$\begin{aligned}
 \text{cl}\{y\} &= \bigcap_{\substack{C \in \text{clSets } y \\ C \subset U^c}} C
 \end{aligned}$$

Since $x \in U$, $\{x\} \cap \text{cl}\{y\} = \emptyset$. Since $x \in \text{cl}\{x\}$, we have that $\text{cl}\{x\} \not\subset \text{cl}\{y\}$. Hence $\text{cl}\{x\} \neq \text{cl}\{y\}$.

– Similarly, if $(x, y) \in U^c \times U$, then $\text{cl}\{x\} \neq \text{cl}\{y\}$.

4. • $(\implies)$:

Suppose that X is T_0 . Let $x, y \in X$. Since X is T_0 , we have that

$$\begin{aligned}
 x \neq y &\iff \exists U \in \mathcal{T} \text{ s.t. } (x, y) \in U \times U^c \text{ or } (x, y) \in U^c \times U \\
 &\iff x \not\sim_0 y
 \end{aligned}$$

- ($\Leftarrow$) :

Suppose that for each $x, y \in X$, $x \neq y$ iff $x \not\sim_0 y$. Let $x, y \in X$. Suppose that $x \neq y$. By assumption, $x \not\sim_0 y$. Thus there exists $U \in \mathcal{T}$ such that $(x, y) \in U \times U^c$ or $(x, y) \in U^c \times U$. Since $x, y \in X$ are arbitrary, we have that X is T_0 .

5. (a) Let $U \in \mathcal{T}$ and $x \in \pi^{-1}(\pi(U))$. Then $\pi(x) \in \pi(U)$. Thus there exists $y \in U$ such that $\pi(x) = \pi(y)$. By definition of π , $x \sim_0 y$. By definition of $\sim_0$, $x \in U$. Since $x \in \pi^{-1}(\pi(U))$ is arbitrary, we have that $\pi^{-1}(\pi(U)) \subset U$. Exercise 1.3.1.1 implies that $U \subset \pi^{-1}(\pi(U))$. Thus $\pi^{-1}(\pi(U)) = U$. Since $U \in \mathcal{T}$ is arbitrary, we have that for each $U \in \mathcal{T}$, $\pi^{-1}(\pi(U)) = U$.
- (b) In particular, for each $U \in \mathcal{T}$,

$$\begin{aligned}\pi^{-1}(\pi(U)) &= U \\ &\in \mathcal{T}\end{aligned}$$

and since π is a quotient map, Exercise 4.8.1.9 implies that π is open.

6. Let $a, b \in X/\sim_0$. For the sake of contradiction, suppose that $a \sim_0 b$ and $a \neq b$. Since π is surjective, there exist $x, y \in X$ such that $\pi(x) = a$ and $\pi(y) = b$. Since $\pi(x) \neq \pi(y)$, $x \not\sim_0 y$. Therefore there exists $U \in \mathcal{T}$ such that $(x, y) \in U \times U^c$ or $(x, y) \in U^c \times U$.

- Suppose that $(x, y) \in U \times U^c$. Since $x \in U$ and π is open, we have that

$$\begin{aligned}a &= \pi(x) \\ &\in \pi(U)\end{aligned}$$

Since $a \sim_0 b$, $a \in \pi(U)$ and $\pi(U) \in \mathcal{T}_{X/\sim_0}$, we have that $b \in \pi(U)$. Therefore

$$\begin{aligned}y &\in \pi^{-1}(\pi(U)) \\ &= U\end{aligned}$$

This is a contradiction since $y \in U^c$.

- Similarly, if $(x, y) \in U^c \times U$, then $x \in U$ which is a contradiction.

Hence $a \sim_0 b$ implies that $a = b$.

Conversely, since $\sim_0$ is an equivalence relation, $a = b$ implies that $a \sim_0 b$.

Thus $a = b$ iff $a \sim_0 b$. Since $a, b \in X/\sim_0$ are arbitrary, we have that for each $a, b \in X/\sim_0$, $a = b$ iff $a \sim_0 b$. Part (4) implies that $X/\sim_0$ is T_0 .

7. FINISH!!!

□

Exercise 4.12.4.3. Let X be a topological space and $\sim$ an equivalence relation on X . If $\pi : X \rightarrow X/\sim$ is open, then $X/\sim$ is Hausdorff iff $\sim$ is closed in $X \times X$.

Proof. Suppose that $\pi : X \rightarrow X/\sim$ is open.

- ($\Rightarrow$) :

Suppose that $X/\sim$ is Hausdorff. Let $(x_\alpha, y_\alpha)_{\alpha \in A} \subset \sim$ be a net and $(x, y) \in X \times X$. Suppose that $x_\alpha, y_\alpha \rightarrow (x, y)$. Then $x_\alpha \rightarrow x$ and $y_\alpha \rightarrow y$. Since $\pi : X \rightarrow X/\sim$ is continuous, $\pi(x_\alpha) \rightarrow \pi(x)$ and $\pi(y_\alpha) \rightarrow \pi(y)$. Since for each $\alpha \in A$, $x_\alpha \sim y_\alpha$, we have that

$$\begin{aligned}\pi(x_\alpha) &= \pi(y_\alpha) \\ &\rightarrow \pi(y)\end{aligned}$$

Since $X/\sim$ is Hausdorff, $\pi(x) = \pi(y)$. Hence $x \sim y$ and $(x, y) \in \sim$. Thus $\sim$ is closed in $X \times X$.

• ($\Leftarrow$):

Conversely, suppose that $\sim$ is closed in $X \times X$ is closed. Let $\bar{x}, \bar{y} \in X/\sim$. Suppose that $\bar{x} \neq \bar{y}$. Then $(x, y) \in \sim^c$. Recall that $\mathcal{B} = \{A \times B : A, B \subset X \text{ and } A, B \text{ are open}\}$ is a basis for $X \times X$. Since $\sim^c$ is open and $(x, y) \in \sim^c$, there exist $A, B \subset X$ such that A, B are open and $(x, y) \in A \times B \subset \sim^c$. Thus $x \in A$ and $y \in B$. Since π is open, $\pi(A) = \bar{A}$ and $\pi(B) = \bar{B}$ are open. Suppose for the sake of contradiction that $\pi(A) \cap \pi(B) \neq \emptyset$. Then there exists $z \in X$ such that $\bar{z} \in \pi(A) \cap \pi(B)$. Therefore there exist $z_A \in A$ and $z_B \in B$ such that $z_A \sim z$ and $z \sim z_B$. Then $(z_A, z_B) \in A \times B$ and $(z_A, z_B) \in \sim$. This is a contradiction since $A \times B \subset \sim^c$. So $\pi(A) \cap \pi(B) = \emptyset$. Thus $\bar{x} \in \pi(A)$, $\bar{y} \in \pi(B)$, $\pi(A)$, $\pi(B)$ are open and $\pi(A) \cap \pi(B) = \emptyset$. Since $\bar{x}, \bar{y} \in X/\sim$ are arbitrary, $X/\sim$ is Hausdorff.

□

4.13 Countability Axioms

4.13.1 First-Countability

Definition 4.13.1.1. Let $(X, \mathcal{T})$ be a topological space. Then $(X, \mathcal{T})$ is said to be **first-countable** if for each $x \in X$, there exists $\mathcal{B}_x \subset \mathcal{T}$ such that

1. $\mathcal{B}_x$ is a local basis for $\mathcal{T}$ at x
2. $\mathcal{B}_x$ is countable

Exercise 4.13.1.2. Let $(X, \mathcal{T})$ be a topological space. Suppose that $(X, \mathcal{T})$ is first-countable. Then for each $x \in X$, there exists $(U_{x,n})_{n \in \mathbb{N}} \subset \mathcal{T}$ such that

1. $(U_{x,n})_{n \in \mathbb{N}}$ is a local basis for $\mathcal{T}$ at x
2. for each $n \in \mathbb{N}$, $U_{x,n+1} \subset U_{x,n}$

Proof.

1. Let $x \in X$. Since $(X, \mathcal{T})$ is first-countable, there exists $(E_{x,j})_{j \in \mathbb{N}} \subset \mathcal{T}$ such that $(E_{x,j})_{j \in \mathbb{N}}$ is a local basis for $\mathcal{T}$ at x . Define $(U_{x,n})_{n \in \mathbb{N}} \subset \mathcal{T}$ by

$$U_{x,n} = \bigcap_{j=1}^n E_{x,j}$$

- Since $(E_{x,j})_{j \in \mathbb{N}}$ is a local basis for $\mathcal{T}$ at x , for each $j \in \mathbb{N}$, $x \in E_{x,j}$. Therefore for each $n \in \mathbb{N}$,

$$\begin{aligned} x &\in \bigcap_{j=1}^n E_{x,j} \\ &= U_{x,n} \end{aligned}$$

- Let $V \in \mathcal{T}$. Suppose that $x \in V$. Since $(E_{x,j})_{j \in \mathbb{N}}$ is a local basis for $\mathcal{T}$ at x , there exists $n \in \mathbb{N}$ such that $E_{x,n} \subset V$. Then

$$\begin{aligned} U_{x,n} &= \bigcap_{j=1}^n E_{x,j} \\ &\subset E_{x,n} \\ &\subset V \end{aligned}$$

Thus $(U_{x,n})_{n \in \mathbb{N}}$ is a local basis for $\mathcal{T}$ at x .

2. By construction, for each $n \in \mathbb{N}$,

$$\begin{aligned} U_{x,n+1} &= \bigcap_{j=1}^{n+1} E_{x,j} \\ &\subset \bigcap_{j=1}^n E_{x,j} \\ &= U_{x,n} \end{aligned}$$

□

Exercise 4.13.1.3. Let $(X, \mathcal{T})$ be a topological space and $A \subset X$. Suppose that $(X, \mathcal{T})$ is first-countable. Then for each $x \in X$, $x \in \text{cl } A$ iff there exists $(x_n)_{n \in \mathbb{N}} \subset A$ such that $x_n \rightarrow x$

Proof. Let $x \in X$.

- ($\Rightarrow$):

Suppose that $x \in \text{cl } A$. Since $(X, \mathcal{T})$ is first-countable, Exercise 4.13.1.2 implies that there exists $(U_n)_{n \in \mathbb{N}} \subset \mathcal{T}$ such that

1. $(U_n)_{n \in \mathbb{N}}$ is a local basis for $\mathcal{T}$ at x
2. for each $n \in \mathbb{N}$, $U_{n+1} \subset U_n$

Exercise 4.1.0.29 implies that for each $n \in \mathbb{N}$, $U_n \cap A \neq \emptyset$. Define $F : \mathbb{N} \rightarrow \mathcal{P}(A)$ by $F(n) := U_n \cap A$. Since for each $n \in \mathbb{N}$, $F(n) \neq \emptyset$, the axiom of choice implies that there exists $(x_n)_{n \in \mathbb{N}} \in A^{\mathbb{N}}$ such that $(x_n)_{n \in \mathbb{N}}$ is a selection of F . Let $U \in \mathcal{N}(x)$. Since $(U_n)_{n \in \mathbb{N}}$ is a local basis for $\mathcal{T}$ at x , $x \in \text{Int } U$ and $\text{Int } U \in \mathcal{T}$, there exists $n_0 \in \mathbb{N}$ such that $U_{n_0} \subset \text{Int } U$. Then for each $n \in \mathbb{N}$, $n \geq n_0$ implies that

$$\begin{aligned} x_n &\in U_n \cap A \\ &\subset U_n \\ &\subset U_{n_0} \\ &\subset \text{Int } U \\ &\subset U. \end{aligned}$$

Thus $(x_n)_{n \in \mathbb{N}}$ is eventually in U . Since $U \in \mathcal{N}(x)$ is arbitrary, we have that for each $U \in \mathcal{N}(x)$, $(x_n)_{n \in \mathbb{N}}$ is eventually in U . Hence $x_n \rightarrow x$.

- ($\Leftarrow$):

If there exists $(x_n)_{n \in \mathbb{N}} \subset A$ such that $x_n \rightarrow x$, then Exercise 4.3.2.8 implies that $x \in \text{cl } A$.

□

Exercise 4.13.1.4. Let X be a topological space. Suppose that X is first-countable. Then for each $C \subset X$, C is closed iff for each $(a_n)_{n \in \mathbb{N}} \subset C$ and $a \in X$, $a_n \rightarrow a$ implies that $a \in C$.

Proof. Let $C \subset X$.

- ($\Rightarrow$):

Suppose that C is closed. Exercise 4.3.2.8 implies that for each $(a_n)_{n \in \mathbb{N}} \subset C$ and $a \in X$, $a_n \rightarrow a$ implies that $a \in C$.

- ($\Leftarrow$):

Suppose that for each $(a_n)_{n \in \mathbb{N}} \subset C$ and $a \in X$, $a_n \rightarrow a$ implies that $a \in C$. Then for each $x \in X$,

$$\begin{aligned} x \in \text{cl } C &\iff \text{there exists } (x_n)_{n \in \mathbb{N}} \subset C \text{ such that } x_n \rightarrow x \quad (\text{by Exercise 4.13.1.3}) \\ &\implies x \in C \quad (\text{by assumption}). \end{aligned}$$

Thus

$$\begin{aligned} \text{cl } C &\subset C \\ &\subset \text{cl } C. \end{aligned}$$

Hence $\text{cl } C = C$ and C is closed.

□

Exercise 4.13.1.5. Let $(X, \mathcal{T}_X), (Y, \mathcal{T}_Y)$ be topological spaces and $f : X \rightarrow Y$. Suppose that $(X, \mathcal{T}_X)$ is first-countable. Then f is continuous iff for each sequence $(x_n)_{n \in \mathbb{N}} \subset X$ and $x \in X$, $x_n \rightarrow x$ implies that $f(x_n) \rightarrow f(x)$.

Proof.

- ($\Rightarrow$):

Suppose that f is continuous. Let $(x_n)_{n \in \mathbb{N}} \subset X$ be a sequence and $x \in X$. Suppose that $x_n \rightarrow x$. Since $(x_n)_{n \in \mathbb{N}}$ is a net, a previous exercise implies that $f(x_n) \rightarrow f(x)$.

• ($\Leftarrow$):

Conversely, suppose that for each sequence $(x_n)_{n \in \mathbb{N}} \subset X$ and $x \in X$, $x_n \rightarrow x$ implies that $f(x_n) \rightarrow f(x)$. Since $(X, \mathcal{T})$ is first-countable, the previous exercise implies that there exists $(U_{x,n})_{n \in \mathbb{N}} \subset \mathcal{T}$ such that

1. $(U_{x,n})_{n \in \mathbb{N}}$ is a local basis for $\mathcal{T}$ at x
2. for each $n \in \mathbb{N}$, $U_{x,n+1} \subset U_{x,n}$

For the sake of contradiction, suppose that f is not continuous. Then there exists $x \in X$ such that f is not continuous at x . Thus there exists $V \in \mathcal{N}(f(x))$ such that for each $U \in \mathcal{N}(x)$, $f(U) \not\subset V$. In particular, for each $n \in \mathbb{N}$, $f(U_{x,n}) \cap V^c \neq \emptyset$ and therefore $U_{x,n} \cap f^{-1}(V^c) \neq \emptyset$. The axiom of choice implies that there exists a sequence $(x_n)_{n \in \mathbb{N}} \subset X$ such that for each $n \in \mathbb{N}$, $x_n \in U_{x,n}$ and $f(x_n) \in V^c$.

Let $U \in \mathcal{N}(x)$. Since $(U_{x,n})_{n \in \mathbb{N}}$ is a local basis for $\mathcal{T}$ at x , there exists $N \in \mathbb{N}$ such that $U_{x,N} \subset \text{Int } U$. Then for each $n \in \mathbb{N}$, $n \geq N$ implies that

$$\begin{aligned} x_n &\in U_{x,n} \\ &\subset U_{x,N} \\ &\subset \text{Int } U \\ &\subset U \end{aligned}$$

Hence $(x_n)_{n \in \mathbb{N}}$ is eventually in U . Since $U \in \mathcal{N}(x)$ is arbitrary, $x_n \rightarrow x$. By assumption, $f(x_n) \rightarrow f(x)$. This is a contradiction since for each $n \in \mathbb{N}$, $f(x_n) \in V^c$ and therefore it is not the case that $(f(x_n))_{n \in \mathbb{N}}$ is eventually in V . Hence f is continuous. $\square$

Exercise 4.13.1.6. Let $(X_n, \mathcal{T}_n)_{n \in \mathbb{N}} \subset \text{Obj}(\mathbf{Top})$. If for each $n \in \mathbb{N}$, $(X_n, \mathcal{T}_n)$ is first-countable, then $\left(\prod_{n \in \mathbb{N}} X_n, \bigotimes_{n \in \mathbb{N}} \mathcal{T}_n \right)$ is first-countable.

Proof. Set $X = \prod_{n \in \mathbb{N}} X_n$ and $\mathcal{T} = \bigotimes_{n \in \mathbb{N}} \mathcal{T}_n$. Let $x \in X$. Since for each $n \in \mathbb{N}$, X_n is first-countable, we have that for each $n \in \mathbb{N}$, there exists $\mathcal{B}_{x_n} \subset \mathcal{T}_n$ such that

1. $\mathcal{B}_{x_n}$ is a local basis for $\mathcal{T}_n$ at x_n
2. $\mathcal{B}_{x_n}$ is countable

Set

$$\mathcal{B}_x = \left\{ \prod_{n \in \mathbb{N}} U_n : \text{[for each } n \in \mathbb{N}, U_n \in \mathcal{T}_n \text{ and } U_n \neq X_n \text{ implies that } U_n \in \mathcal{B}_{x_n}] \text{ and } \#\{n \in \mathbb{N} : U_n \neq X_n\} < \infty \right\}$$

Then $\mathcal{B}_x$ is countable and an exercise in the section on the product topology implies that $\mathcal{B}_x$ is a local basis for $\mathcal{T}$ at x . Since $x \in X$ is arbitrary, we have that for each $x \in X$, there exists $\mathcal{B}_x \subset \mathcal{T}$ such that

1. $\mathcal{B}_x$ is a local basis for $\mathcal{T}$ at x
2. $\mathcal{B}_x$ is countable

Hence $(X, \mathcal{T})$ is first-countable. $\square$

Exercise 4.13.1.7. Let $(X, \mathcal{T})$ be a topological space, $(x_n)_{n \in \mathbb{N}} \subset X$ and $x \in X$. Suppose that $(X, \mathcal{T})$ is first countable and x is a cluster point of $(x_n)_{n \in \mathbb{N}}$. Then there exists $(x_{n_k})_{k \in \mathbb{N}} \subset (x_n)_{n \in \mathbb{N}}$ such that $x_{n_k} \rightarrow x$.

Proof. redo or give alternative proof while citing Exercise 4.3.2.20

- Since $(X, \mathcal{T})$ is first-countable, there exists $(U_n)_{n \in \mathbb{N}} \subset \mathcal{T}$ such that $(U_n)_{n \in \mathbb{N}}$ is a local basis for $\mathcal{T}$ at x . Define $(V_n)_{n \in \mathbb{N}} \subset \mathcal{T}$ by $V_n := \bigcap_{j=1}^n U_j$. Then for each $n \in \mathbb{N}$, $V_n \in \mathcal{N}(x)$.
- We define $(A_j)_{j,k \in \mathbb{N}} \subset \mathcal{P}(\mathbb{N})$ by $A_{k,j} := \{n \in \mathbb{N} : n \geq j \text{ and } x_n \in V_k\}$. Since x is a cluster point of $(x_n)_{n \in \mathbb{N}}$, for each $j, k \in \mathbb{N}$, $A_{k,j} \neq \emptyset$.

- We define $(n_k)_{k \in \mathbb{N}} \subset \mathbb{N}$ by

$$n_k := \begin{cases} \min A_{1,1}, & k = 1 \\ \min A_{k, n_{k-1}+1}, & k \geq 2 \end{cases}$$

By construction for each $k \in \mathbb{N}$, $n_k > n_{k-1}$. Thus $(x_{n_k})_{k \in \mathbb{N}} \subset (x_n)_{n \in \mathbb{N}}$.

- Let $U \in \mathcal{N}(x)$. Since $(U_n)_{n \in \mathbb{N}}$ is a local basis for $\mathcal{T}$ at x , there exists $k_0 \in \mathbb{N}$ such that

$$\begin{aligned} x &\in V_{k_0} \\ &\subset U_{k_0} \\ &\subset \text{Int } U \\ &\subset U \end{aligned}$$

Let $k \in \mathbb{N}$. Suppose that $k \geq k_0$. Since

$$\begin{aligned} n_k &> n_{k-1} \\ &\geq n_{k_0}, \end{aligned}$$

we have that $V_{n_k} \subset V_{n_{k_0}}$ and

$$\begin{aligned} x_{n_k} &\in V_{n_k} \\ &\subset V_{k_0} \\ &\subset U. \end{aligned}$$

Hence there exists $k_0 \in \mathbb{N}$ such that for each $k \in \mathbb{N}$, $k \geq k_0$ implies that $x_{n_k} \in U$. Thus (x_{n_k}) is eventually in U . Since $U \in \mathcal{N}(x)$ is arbitrary, we have that for each $U \in \mathcal{N}(x)$, (x_{n_k}) is eventually in U . Therefore $x_{n_k} \rightarrow x$ in $(X, \mathcal{T})$. □

4.13.2 Second-Countability

Definition 4.13.2.1. Let $(X, \mathcal{T})$ be a topological space. Then $(X, \mathcal{T})$ is said to be **second-countable** if there exists $\mathcal{B} \subset \mathcal{T}$ such that

1. $\mathcal{B}$ is a basis for $\mathcal{T}$
2. $\mathcal{B}$ is countable

Exercise 4.13.2.2. Let $(X, \mathcal{T}_X)$, $(Y, \mathcal{T}_Y)$ be topological spaces and $f : X \rightarrow Y$. Suppose that f is surjective, continuous and open. If X is second-countable, then Y is second-countable.

Exercise 4.13.2.3. **second-countable implies first-countable**

Proof. **FINISH!!!** □

Proof. Suppose that X is second-countable. Then there exists $\mathcal{B}_X \subset \mathcal{T}_X$ such that $\mathcal{B}_X$ is a basis for $\mathcal{T}_X$ and $\mathcal{B}_X$ is countable. Set $\mathcal{B}_Y = \{f(A) : A \in \mathcal{B}_X\}$. Since $\mathcal{B}_X$ is countable, $\mathcal{B}_Y$ is countable. Since f is surjective, continuous and open, a previous exercise implies that $\mathcal{B}_Y$ is a basis for $\mathcal{T}_Y$. Hence $(Y, \mathcal{T}_Y)$ is second-countable. □

Exercise 4.13.2.4. Let $(X, \mathcal{T}_X)$, $(Y, \mathcal{T}_Y)$ be topological spaces and $f : X \rightarrow Y$. Suppose that f is a homeomorphism. Then $(X, \mathcal{T}_X)$ is second-countable iff $(Y, \mathcal{T}_Y)$ is second-countable.

Proof.

- $(\implies)$:

Suppose that $(X, \mathcal{T}_X)$ is second-countable. Since f is surjective, continuous and open, the previous exercise implies that $(Y, \mathcal{T}_Y)$ is second-countable.

• ($\Leftarrow$) :

Conversely, suppose that $(Y, \mathcal{T}_Y)$ is second-countable. Since $f^{-1} : Y \rightarrow X$ is surjective, continuous and open, the previous exercise implies that $(X, \mathcal{T}_X)$ is second-countable. □

Definition 4.13.2.5. Let $(X, \mathcal{T})$ be a topological space. Then $(X, \mathcal{T})$ is said to be **separable** if there exists $S \subset X$ such that S is dense in X and S is countable.

Exercise 4.13.2.6. Let $(X, \mathcal{T})$ be a topological space. If $(X, \mathcal{T})$ is second-countable, then $(X, \mathcal{T})$ is separable.

Proof. Suppose that $(X, \mathcal{T})$ is second-countable. Then there exists $\mathcal{B} \subset \mathcal{T}$ such that $\mathcal{B}$ is a basis for $\mathcal{T}$ and $\mathcal{B}$ is countable. The axiom of choice implies that there exists $(x_U)_{U \in \mathcal{B}} \subset X$ such that each $U \in \mathcal{B}$, $x_U \in U$. Let $V \in \mathcal{T}$. Suppose that $V \neq \emptyset$. Then there exists $x \in V$. Since $\mathcal{B}$ is a basis for $\mathcal{T}$, there exists $U \in \mathcal{B}$ such that $x \in U \subset V$. Hence $x_U \in (x_U)_{U \in \mathcal{B}} \cap V$ which implies that $(x_U)_{U \in \mathcal{B}} \cap V \neq \emptyset$. Since $V \in \mathcal{T}$ such that $V \neq \emptyset$ is arbitrary, we have that for each $V \in \mathcal{T}$, $V \neq \emptyset$ implies that $(x_U)_{U \in \mathcal{B}} \cap V \neq \emptyset$. Exercise 4.1.0.41 implies that $(x_U)_{U \in \mathcal{B}}$ is dense in X . Since $\mathcal{B}$ is countable, $(X, \mathcal{T})$ is separable. □

Exercise 4.13.2.7. Let $(X, \mathcal{T}), (Y, \mathcal{T}_Y)$ be a topological spaces and $f : X \rightarrow Y$ a $(\mathcal{T}_X, \mathcal{T}_Y)$ -homeomorphism. Then $(X, \mathcal{T})$ is second-countable iff $(Y, \mathcal{T}_Y)$ is second-countable.

Proof.

• ($\Rightarrow$) :

Suppose that $(X, \mathcal{T})$ is second-countable. Then there exists $\mathcal{B}_X \subset \mathcal{T}_X$ such that $\mathcal{B}_X$ is a basis for $\mathcal{T}_X$ and $\mathcal{B}_X$ is countable. Define $\mathcal{B}_Y := \{f(U) : U \in \mathcal{B}_X\}$. Since f is a $(\mathcal{T}_X, \mathcal{T}_Y)$ -homeomorphism, we have that f is open and $\mathcal{B}_Y \subset \mathcal{T}_Y$. Since $\mathcal{B}_X$ is countable, $\mathcal{B}_Y$ is countable. Let $V \in \mathcal{T}_Y$ and $y \in V$. Set $U := f^{-1}(V)$ and $x := f^{-1}(y)$. Since f is continuous, $U \in \mathcal{T}_X$. By construction, $x \in U$. Since $\mathcal{B}_X$ is a basis for $\mathcal{T}_X$, there exists $B_X \in \mathcal{B}_X$ such that $x \in B_X$ and $B_X \subset U$. Set $B_Y := f(B_X)$. By definition, $B_Y \in \mathcal{B}_Y$ and

$$\begin{aligned} y &= f(x) \\ &\in f(B_X) \\ &\subset f(U) \\ &= V \end{aligned}$$

and

$$\begin{aligned} y &\in f(B_X) \\ &= B_Y. \end{aligned}$$

Since $V \in \mathcal{T}_Y$ and $y \in V$ are arbitrary, we have that for each $V \in \mathcal{T}_Y$ and $y \in V$, there exists $B_Y \in \mathcal{B}_Y$ such that $y \in B_Y \subset V$. Hence $\mathcal{B}_Y$ is a basis for $\mathcal{T}_Y$.

• ($\Leftarrow$) :

Similar to ($\Rightarrow$).

□

Definition 4.13.2.8. Let X be a topological space. Then X is said to be **Lindelöf** if for each open cover $\mathcal{U}$ of X , there exists a subcover $\mathcal{U}' \subset \mathcal{U}$ of X such that $\mathcal{U}'$ is countable.

NEED TO DEFINE SUBCOVER

FINISH!!!

Exercise 4.13.2.9. Let $(X, \mathcal{T})$ be a topological space. If $(X, \mathcal{T})$ is second-countable, then $(X, \mathcal{T})$ is Lindelöf.

Proof. Suppose that X is second-countable. Then there exists $\mathcal{B} \subset \mathcal{T}$ such that $\mathcal{B}$ is a basis for $\mathcal{T}$ and $\mathcal{B}$ is countable. Let $\mathcal{U}$ be an open cover of X . For $B \in \mathcal{B}$, define $\mathcal{U}_B \subset \mathcal{U}$ by $\mathcal{U}_B = \{U \in \mathcal{U} : B \subset U\}$. Set $\Gamma = \{B \in \mathcal{B} : \mathcal{U}_B \neq \emptyset\}$. The axiom of choice implies that there exists $(V_B)_{B \in \Gamma} \subset \mathcal{U}$ such that for each $B \in \Gamma$, $V_B \in \mathcal{U}_B$. Set $\mathcal{U}' = (V_B)_{B \in \Gamma}$. Let $x \in X$. Since $\mathcal{U}$ is an open cover of X , there exists $U \in \mathcal{U}$ such that $x \in U$. Since $\mathcal{B}$ is a basis for $\mathcal{T}$, there exists $B \in \mathcal{B}$ such that $x \in B \subset U$. Thus $B \in \Gamma$ and $x \in V_B$. So $\mathcal{U}'$ is a subcover of $\mathcal{U}$. Since $\mathcal{B}$ is countable, $\mathcal{U}'$ is countable. Since $\mathcal{U}$ is an arbitrary open cover of X , we have that for each open cover $\mathcal{U}$ of X , there exists a countable subcover $\mathcal{U}' \subset \mathcal{U}$. Hence $(X, \mathcal{T})$ is Lindelöf. □

Exercise 4.13.2.10. Let $(X, \mathcal{T})$ be a topological space. Define $C \subset X$ by

$$C := \{x \in X : x \text{ is a } \mathcal{T}\text{-condensation point of } X\}.$$

If $(X, \mathcal{T})$ is second-countable, then C^c is countable.

Hint: Exercise 4.1.0.48

Proof. Suppose that $(X, \mathcal{T})$ is second-countable. Then there exists $\mathcal{B} \subset \mathcal{T}$ such that $\mathcal{B}$ is a basis for $\mathcal{T}$ and $\mathcal{B}$ is countable. Let $x \in C^c$. Exercise 4.1.0.48 implies that there exists $U \in \mathcal{T}$ such that $x \in U$, U is countable and $U \subset C^c$. Since $\mathcal{B}$ is a basis for $\mathcal{T}$, there exists $B \in \mathcal{B}$ such that $x \in B$ and $B \subset U$. Then B and $B \subset C^c$ is countable. Since $x \in C^c$ is arbitrary, we have that for each $x \in C^c$, there exists $B \in \mathcal{B}$ such that $x \in B$, B is countable and $B \subset C^c$. The axiom of choice implies that there exists $(B_x)_{x \in C^c} \subset \mathcal{B}$ such that for each $x \in C^c$, $x \in B_x$, B_x is countable and $B_x \subset C^c$. Define $\mathcal{B}_0 \subset \mathcal{B}$ by $\mathcal{B}_0 := \{B \in \mathcal{B} : B \text{ is countable}\}$. Then $\bigcup_{B \in \mathcal{B}_0} B$ is countable.

Since

$$\begin{aligned} C^c &= \bigcup_{x \in C^c} B_x \\ &\subset \bigcup_{B \in \mathcal{B}_0} B, \end{aligned}$$

we have that C^c is countable. □

Second-Countability and Subspaces

Exercise 4.13.2.11. Let $(X, \mathcal{T})$ be a topological space and $A \subset X$. If $(X, \mathcal{T})$ is second-countable, then $(A, \mathcal{T} \cap A)$ is second-countable.

Proof. Suppose that $(X, \mathcal{T})$ is second-countable. Then there exists $\mathcal{B} \subset \mathcal{T}_d$ such that $\mathcal{B}$ is a basis for $\mathcal{T}_d$ and $\mathcal{B}$ is countable. Exercise 4.5.1.15 implies that $\mathcal{B} \cap A$ is a basis for $\mathcal{T} \cap A$. Since $\mathcal{T} \cap A$ is countable, $(A, \mathcal{T} \cap A)$ is second-countable. □

Exercise 4.13.2.12. Let $(X, \mathcal{T})$ be a topological space. Then $(X, \mathcal{T})$ is second-countable iff there exist $(U_n)_{n \in \mathbb{N}} \subset \mathcal{T}$ such that

1. $X = \bigcup_{n \in \mathbb{N}} U_n$,
2. for each $n \in \mathbb{N}$, $(U_n, \mathcal{T} \cap U_n)$ is second-countable.

Proof.

• $(\implies)$:

Suppose that $(X, \mathcal{T})$ is second-countable. Define $(U_n)_{n \in \mathbb{N}} \subset \mathcal{T}$ by

$$U_n := \begin{cases} X, & n = 1 \\ \emptyset, & n > 1. \end{cases}$$

Then

1. by construction, $X = \bigcup_{n \in \mathbb{N}} U_n$,
2. Exercise 4.13.2.11 implies that for each $n \in \mathbb{N}$, $(U_n, \mathcal{T} \cap U_n)$ is second-countable.

• $(\impliedby)$:

Suppose that there exist $(U_n)_{n \in \mathbb{N}} \subset \mathcal{T}$ such that

1. $X = \bigcup_{n \in \mathbb{N}} U_n$,
2. for each $n \in \mathbb{N}$, $(U_n, \mathcal{T} \cap U_n)$ is second-countable.

Since for each $n \in \mathbb{N}$, $(U_n, \mathcal{T} \cap U_n)$ is second-countable, we have that for each $n \in \mathbb{N}$, there exists $\mathcal{B}_n \subset \mathcal{T} \cap U_n$ such that $\mathcal{B}_n$ is a basis for $\mathcal{T} \cap U_n$ and $\mathcal{B}_n$ is countable. Since $(U_n)_{n \in \mathbb{N}} \subset \mathcal{T}$, we have that for each $n \in \mathbb{N}$,

$$\begin{aligned}\mathcal{B}_n &\subset \mathcal{T} \cap U_n \\ &\subset \mathcal{T}\end{aligned}$$

Define $\mathcal{B} \subset \mathcal{T}$ by $\mathcal{B} = \bigcup_{n \in \mathbb{N}} \mathcal{B}_n$. Then $\mathcal{B}$ is countable. Let $V \in \mathcal{T}$ and $n \in \mathbb{N}$. Then $V \cap U_n \in \mathcal{T} \cap U_n$. Since $V \cap U_n \in \mathcal{T} \cap U_n$, there exist $(B_{n,j})_{j \in \mathbb{N}} \subset \mathcal{B}_n$ such that $V \cap U_n = \bigcup_{j \in \mathbb{N}} B_{n,j}$. Then $(B_{n,j})_{n,j \in \mathbb{N}} \subset \mathcal{B}$ and

$$\begin{aligned}V &= V \cap X \\ &= V \cap \left(\bigcup_{n \in \mathbb{N}} U_n \right) \\ &= \bigcup_{n \in \mathbb{N}} V \cap U_n \\ &= \bigcup_{n \in \mathbb{N}} \bigcup_{j \in \mathbb{N}} B_{n,j}\end{aligned}$$

Since $V \in \mathcal{T}$ is arbitrary, we have that for each $V \in \mathcal{T}$, there exists $\mathcal{B}' \subset \mathcal{B}$ such that $V = \bigcup_{B \in \mathcal{B}'} B$. Hence $\mathcal{B}$ is a basis for $\mathcal{T}$. Since $\mathcal{B}$ is countable, $(X, \mathcal{T})$ is second-countable. □

Second-Countability and Product Spaces

Exercise 4.13.2.13. Let $(X_\alpha, \mathcal{T}_\alpha)_{\alpha \in A}$ be a collection of topological spaces. Set $X = \prod_{\alpha \in A} X_\alpha$ and $\mathcal{T}_X := \bigotimes_{\alpha \in A} \mathcal{T}_\alpha$. Suppose that A is countable. If for each $\alpha \in A$, $(X_\alpha, \mathcal{T}_\alpha)$ is second-countable, then $(X, \mathcal{T}_X)$ is second-countable.

Proof. Suppose that for each $\alpha \in A$, $(X_\alpha, \mathcal{T}_\alpha)$ is second-countable. Then for each $\alpha \in A$, there exists $\mathcal{B}_\alpha \subset \mathcal{T}_\alpha$ such that $\mathcal{B}_\alpha$ is a basis for $\mathcal{T}_\alpha$ and $\mathcal{B}_\alpha$ is countable. Set

$$\mathcal{B} = \left\{ \prod_{\alpha \in A} U_\alpha : \text{there exists } J \subset A \text{ such that } \#J < \infty, \right. \\ \left. \text{for each } \alpha \in J, U_\alpha \in \mathcal{B}_\alpha \text{ and for each } \alpha \in J^c, U_\alpha = X_\alpha \right\}$$

An exercise in the section on the product topology implies that $\mathcal{B}$ is a basis for $\mathcal{T}_X$. Since A is countable and for each $\alpha \in A$, $\mathcal{B}_\alpha$ is countable, we have that $\mathcal{B}$ is countable. Hence $\mathcal{T}_X$ is second-countable. □

Second-Countability and Coproduct Spaces

Exercise 4.13.2.14. Let $(X_\alpha, \mathcal{T}_\alpha)_{\alpha \in A}$ be a collection of topological spaces. Set $X = \coprod_{\alpha \in A} X_\alpha$ and $\mathcal{T} := \bigotimes_{\alpha \in A} \mathcal{T}_\alpha$. Suppose that A is countable. If for each $\alpha \in A$, $(X_\alpha, \mathcal{T}_\alpha)$ is second-countable, then $(X, \mathcal{T})$ is second-countable.

Proof. Suppose that for each $\alpha \in A$, $(X_\alpha, \mathcal{T}_\alpha)$ is second-countable. Then for each $\alpha \in A$, there exists $\mathcal{B}_\alpha \subset \mathcal{T}_\alpha$ such that $\mathcal{B}_\alpha$ is a basis for $\mathcal{T}_\alpha$ and $\mathcal{B}_\alpha$ is countable. Define $\mathcal{B} := \bigcup_{\alpha \in A} \mathcal{B}_\alpha$. Since A is countable and for each $\alpha \in A$, $\mathcal{B}_\alpha$ is countable, we have that $\mathcal{B}$ is countable. Exercise 4.7.0.6 implies that $\mathcal{B}$ is a basis for $\mathcal{T}$. Hence $(X, \mathcal{T})$ is second-countable. □

Second-Countability and Quotient Spaces

Second-Countability and Projective Limits

make $\varprojlim_{j \in J} X$ refer to set construction, always suppress π_j as in products, but π_j always refer to $\text{proj}|_X$, and define a projective limit of a projective system and show that $\varprojlim X_j$ is a projective limit in various categories when equipped with the corresponding structure

Exercise 4.13.2.15. Let $(J, \leq)$ be a directed set and $((X_j, \mathcal{T}_j)_{j \in J}, (\pi_{i,j})_{(i,j) \in \leq})$ be a **Top**-projective system. Set $(X, (\pi_j)_{j \in J}) := \varprojlim_{j \in J} ((X_j, \mathcal{T}_j)_{j \in J}, (\pi_{i,j})_{(i,j) \in \leq})$. Suppose that J is countable and for each $j \in J$, $(X_j, \mathcal{T}_j)$ is second-countable. Then $(X, \mathcal{T})$ is second-countable.

Proof. Since J is countable, Exercise 4.13.2.13 implies that $(X, \mathcal{T})$ is second-countable. Since $X \subset \prod_{j \in J} X_j$ and $\mathcal{T} = \left(\bigotimes_{j \in J} \mathcal{T}_j \right) \cap X$, Exercise 4.13.2.11 implies that $(X, \mathcal{T})$ is second-countable. $\square$

4.14 Compactness

4.14.1 Compactness and Subspaces

rework using $\mathcal{K}(X, \mathcal{T})$ as the set of compact subsets and $\text{Cover}^{\text{opn}}(X, \mathcal{T})$ instead of saying 'open cover'

Definition 4.14.1.1. Let $(X, \mathcal{T})$ be a topological space. Then $(X, \mathcal{T})$ is said to be **compact** if for each $\mathcal{U} \subset \mathcal{P}(X)$, $\mathcal{U}$ is an open cover of X in $(X, \mathcal{T})$ implies that there exists $\mathcal{U}_0 \subset \mathcal{U}$ such that $\mathcal{U}_0$ is an open cover of X in $(X, \mathcal{T})$ and $\mathcal{U}_0$ is finite.

Definition 4.14.1.2. Let $(X, \mathcal{T})$ be topological space.

- Let $E \subset X$. Then E is said to be **$\mathcal{T}$ -compact** or **compact in $(X, \mathcal{T})$** if $(E, \mathcal{T} \cap E)$ is compact
- We define $\mathcal{K}(X, \mathcal{T}) \subset \mathcal{P}(X)$ by

$$\mathcal{K}(X, \mathcal{T}) := \{K \subset X : K \text{ is compact in } (X, \mathcal{T})\}.$$

Note 4.14.1.3. When the context is clear, we may write $\mathcal{K}(X)$ in place of $\mathcal{K}(X, \mathcal{T})$.

Exercise 4.14.1.4. Let $(X, \mathcal{T})$ be a topological space, $E \subset X$ and $A \subset E$. Then $A \in \mathcal{K}(E, \mathcal{T} \cap E)$ iff $A \in \mathcal{K}(X, \mathcal{T})$.

Proof. We note that since $A \subset E$, $(\mathcal{T} \cap E) \cap A = \mathcal{T} \cap A$. Suppose that $A \in \mathcal{K}(E, \mathcal{T} \cap E)$. By definition, $(A, (\mathcal{T} \cap E) \cap A)$ is compact. Since $(A, (\mathcal{T} \cap E) \cap A) = (A, \mathcal{T} \cap A)$, by definition, $A \in \mathcal{K}(X, \mathcal{T})$.

Coversely, suppose that $A \in \mathcal{K}(X, \mathcal{T})$. By definition, $(A, \mathcal{T} \cap A)$ is compact. Similarly, since $(A, \mathcal{T} \cap A) = (A, (\mathcal{T} \cap E) \cap A)$, $A \in \mathcal{K}(E, \mathcal{T} \cap E)$. $\square$

Exercise 4.14.1.5. Let $(X, \mathcal{T})$ be a topological space and $K \subset X$. Then K is compact in $(X, \mathcal{T})$ iff for each $\mathcal{U} \subset \mathcal{P}(X)$, $\mathcal{U}$ is an open cover of K in $(X, \mathcal{T})$ implies that there exists $\mathcal{U}_0 \subset \mathcal{U}$ such that $\mathcal{U}_0$ is an open cover of K in $(X, \mathcal{T})$ and $\mathcal{U}_0$ is finite.

Proof.

- $(\implies)$:
Suppose that K is compact in $(X, \mathcal{T})$. Let $\mathcal{U} \subset \mathcal{P}(X)$. Suppose that $\mathcal{U}$ is an open cover of K in $(X, \mathcal{T})$. Then $\mathcal{U} \subset \mathcal{T}$ and $K \subset \bigcup_{U \in \mathcal{U}} U$. Therefore $\mathcal{U} \cap K \subset \mathcal{T} \cap K$ and

$$\begin{aligned} K &\subset \left[\bigcup_{U \in \mathcal{U}} U \right] \cap K \\ &= \bigcup_{U \in \mathcal{U}} U \cap K \end{aligned}$$

Hence $\mathcal{U} \cap K$ is an open cover of K in $(K, \mathcal{T} \cap K)$. Since K is compact in $(X, \mathcal{T})$, Exercise 4.14.1.4 implies that $(K, \mathcal{T} \cap K)$ is compact. By Definition 4.14.1.1, there exists $\mathcal{V}_0 \subset \mathcal{U} \cap K$ such that $\mathcal{V}_0$ is an open cover of K in $(K, \mathcal{T} \cap K)$ and $\mathcal{V}_0$ is finite. Since $\mathcal{V}_0 \subset \mathcal{U} \cap K$, for each $V \in \mathcal{V}_0$, there exists $U_V \in \mathcal{U}$ such that $V = U_V \cap K$. Set $\mathcal{U}_0 = \{U_V : V \in \mathcal{V}_0\}$. Then $\mathcal{U}_0 \subset \mathcal{U}$.

1. Since $\mathcal{U} \subset \mathcal{T}$

$$\begin{aligned} \mathcal{U}_0 &\subset \mathcal{U} \\ &\subset \mathcal{T} \end{aligned}$$

2. Since $\mathcal{V}_0$ is an open cover of K in $(K, \mathcal{T} \cap K)$, we have that

$$\begin{aligned} K &\subset \bigcup_{V \in \mathcal{V}_0} V \\ &= \bigcup_{U \in \mathcal{U}_0} U \cap K \\ &\subset \bigcup_{U \in \mathcal{U}_0} U \end{aligned}$$

By Definition ??, $\mathcal{U}_0$ is an open cover of K in $(X, \mathcal{T})$. By construction, $\mathcal{U}_0$ is finite.

• ($\Leftarrow$):

Suppose that for each $\mathcal{U} \subset \mathcal{P}(X)$, if $\mathcal{U}$ is an open cover of K in $(X, \mathcal{T})$, then there exists $\mathcal{U}_0 \subset \mathcal{U}$ such that $\mathcal{U}_0$ is an open cover of K in $(X, \mathcal{T})$ and $\mathcal{U}_0$ is finite. Let $\mathcal{V} \subset \mathcal{P}(K)$. Suppose that $\mathcal{V}$ is an open cover of K in $(K, \mathcal{T} \cap K)$. Then $\mathcal{V} \subset \mathcal{T} \cap K$ and $K \subset \bigcup_{V \in \mathcal{V}} V$. By definition of $\mathcal{T} \cap K$, for each $V \in \mathcal{V}$, there exists $U \in \mathcal{T}$ such that $V = U \cap K$. The axiom of choice implies that there exists $(U_V)_{V \in \mathcal{V}} \subset \mathcal{T}$ such that for each $V \in \mathcal{V}$, $V = U_V \cap K$. Therefore

$$\begin{aligned} K &\subset \bigcup_{V \in \mathcal{V}} V \\ &= \bigcup_{V \in \mathcal{V}} U_V \cap K \\ &\subset \bigcup_{V \in \mathcal{V}} U_V \end{aligned}$$

Hence $(U_V)_{V \in \mathcal{V}}$ is an open cover of K in $(X, \mathcal{T})$. By assumption, there exists $\mathcal{V}_0 \subset \mathcal{V}$ such that $(U_V)_{V \in \mathcal{V}_0}$ is an open cover of K in $(X, \mathcal{T})$ and $\mathcal{V}_0$ is finite.

1. Since $\mathcal{V} \subset \mathcal{T} \cap K$, we have that

$$\begin{aligned} \mathcal{V}_0 &\subset \mathcal{V} \\ &\subset \mathcal{T} \cap K \end{aligned}$$

2. Since $(U_V)_{V \in \mathcal{V}_0}$ is an open cover of K in $(X, \mathcal{T})$, we have that

$$\begin{aligned} K &= K \cap K \\ &\subset \left[\bigcup_{V \in \mathcal{V}_0} U_V \right] \cap K \\ &= \bigcup_{V \in \mathcal{V}_0} U_V \cap K \\ &= \bigcup_{V \in \mathcal{V}_0} V \end{aligned}$$

Therefore $\mathcal{V}_0$ is an open cover of K in $(K, \mathcal{T} \cap K)$. Since $\mathcal{V} \subset \mathcal{P}(K)$ such that $\mathcal{V}$ is an open cover of K in $(K, \mathcal{T} \cap K)$ is arbitrary, we have that for each $\mathcal{V} \subset \mathcal{P}(K)$, if $\mathcal{V}$ is an open cover of K in $(K, \mathcal{T} \cap K)$, then there exists $\mathcal{V}_0 \subset \mathcal{V}$ such that $\mathcal{V}_0$ is an open cover of K in $(K, \mathcal{T} \cap K)$ and $\mathcal{V}_0$ is finite. Hence $(K, \mathcal{T} \cap K)$ is compact. By definition, K is compact in $(X, \mathcal{T})$. □

Exercise 4.14.1.6. Let $(X, \mathcal{T})$ be a topological space and $K, L \subset X$. If K and L are compact, then $K \cup L$ is compact.

Proof. Suppose that K and L are compact. Let $\mathcal{U} \subset \mathcal{P}(X)$. Suppose that $\mathcal{U}$ is an open cover of $K \cup L$ in $(X, \mathcal{T})$. Since $K, L \subset K \cup L$, $\mathcal{U}$ is an open cover of K and L . Since K, L are compact, there exist $\mathcal{U}_K, \mathcal{U}_L \subset \mathcal{U}$ such that $\mathcal{U}_K$ is an open cover of K , $\mathcal{U}_L$ is an open cover of L and $\mathcal{U}_K, \mathcal{U}_L$ are finite. Define $\mathcal{U}_0 \subset \mathcal{U}$ by $\mathcal{U}_0 := \mathcal{U}_K \cup \mathcal{U}_L$. Then $\mathcal{U}_0$ is an open cover of $K \cup L$ and $\mathcal{U}_0$ is finite. Since $\mathcal{U} \subset \mathcal{P}(X)$ with $\mathcal{U}$ an open cover of $K \cup L$ arbitrary, we have that for each $\mathcal{U} \subset \mathcal{P}(X)$, if $\mathcal{U}$ is an open cover of $K \cup L$, there exists $\mathcal{U}_0 \subset \mathcal{U}$ such that $\mathcal{U}_0$ is an open cover of $K \cup L$ and $\mathcal{U}_0$ is finite. Thus $K \cup L$ is compact. □

Exercise 4.14.1.7. Let $(X, \mathcal{T})$ be a topological space, $U \in \mathcal{T}$ and $x \in U$. Then $U \setminus \{x\}$ is not compact.

Exercise 4.14.1.8. Let $(X, \mathcal{T})$ be a topological space and $K \subset X$. Suppose that $(X, \mathcal{T})$ is Hausdorff. If K is compact in X , then K is closed in X .

Proof. Suppose that K is compact. Let $y \in K^c$. Since $(X, \mathcal{T})$ is Hausdorff, for each $x \in K$, there exists $U_x, V_x \in \mathcal{T}$, such that $x \in U_x$, $y \in V_x$ and $U_x \cap V_x = \emptyset$. Thus $(U_x)_{x \in K}$ is an open cover of K in $(X, \mathcal{T})$. Since K is compact, there exist $x_1, \dots, x_n \in K$ such that $(U_{x_j})_{j=1}^n$ is an open cover of K in $(X, \mathcal{T})$. Set $V = \bigcap_{j=1}^n V_{x_j}$. Then $V \in \mathcal{T}$ and $y \in V$. Since for each $j \in \{1, \dots, n\}$, $V \subset V_{x_j}$, we have that

$$\begin{aligned} V \cap K &\subset V \cap \left[\bigcup_{j=1}^n U_{x_j} \right] \\ &= \bigcup_{j=1}^n (V \cap U_{x_j}) \\ &\subset \bigcup_{j=1}^n (V_{x_j} \cap U_{x_j}) \\ &= \bigcup_{j=1}^n \emptyset \\ &= \emptyset \end{aligned}$$

Thus $V \subset K^c$. Since $y \in K^c$ is arbitrary, we have that for each $y \in K^c$, there exists $V \in \mathcal{T}$ such that $y \in V$ and $V \subset K^c$. Hence K^c is open. Thus K is closed. $\square$

Exercise 4.14.1.9. Let $(X, \mathcal{T})$ be a topological space and $E \subset X$. If $(X, \mathcal{T})$ is compact and E is $\mathcal{T}$ -closed, then E is compact in $(X, \mathcal{T})$.

Proof. Suppose that $(X, \mathcal{T})$ is compact and E is $\mathcal{T}$ -closed. Let $\mathcal{U} \subset \mathcal{P}(X)$. Suppose that $\mathcal{U}$ is an open cover of E in $(X, \mathcal{T})$. Since E is $\mathcal{T}$ -closed, $E^c \in \mathcal{T}$. Set $\mathcal{U}' = \mathcal{U} \cup \{E^c\}$. Then $\mathcal{U}' \subset \mathcal{T}$ and $\mathcal{U}'$ is an open cover of X in $(X, \mathcal{T})$. Since $(X, \mathcal{T})$ is compact, there exists $\mathcal{U}'_0 \subset \mathcal{U}'$ such that $\mathcal{U}'_0$ is an open cover of X in $(X, \mathcal{T})$ and $\mathcal{U}'_0$ is finite. Set $\mathcal{U}_0 = \mathcal{U}'_0 \setminus \{E^c\}$. Then $\mathcal{U}_0$ is an open cover for E in $(X, \mathcal{T})$. Since $\mathcal{U}$ such that $\mathcal{U}$ is an open cover of E in $(X, \mathcal{T})$ is arbitrary, Exercise 4.14.1.5 implies that E is compact in $(X, \mathcal{T})$.

[GIVE MORE DETAILS](#)

[FINISH!!!](#) $\square$

Definition 4.14.1.10. Let X be a topological space and $E \subset X$. Then E is said to be **precompact** if $\text{cl } E$ is compact.

Exercise 4.14.1.11. Let $(X, \mathcal{T})$ be a topological space, $K \subset X$ and $x \in K^c$. Suppose that $(X, \mathcal{T})$ is Hausdorff. If K is compact in $(X, \mathcal{T})$, then there exist $U, V \in \mathcal{T}$ such that $U \cap V = \emptyset$, $x \in U$ and $K \subset V$.

Proof. Suppose that K is compact in $(X, \mathcal{T})$. Since $(X, \mathcal{T})$ is Hausdorff, we have that for each $y \in K$, there exist $U_y, V_y \in \mathcal{T}$ such that $U_y \cap V_y = \emptyset$, $x \in U_y$ and $y \in V_y$. Define $\mathcal{U}, \mathcal{V} \subset \mathcal{P}(X)$ by $\mathcal{U} := \{U_y : y \in K\}$ and $\mathcal{V} := \{V_y : y \in K\}$. Then $\mathcal{V}$ is an open cover of K in $(X, \mathcal{T})$. Since K is compact in $(X, \mathcal{T})$, there exists $\mathcal{V}_0 \subset \mathcal{V}$ such that $\mathcal{V}_0$ is finite and $\mathcal{V}_0$ is an open cover of K in $(X, \mathcal{T})$. By definition of $\mathcal{V}$, there exist $y_1, \dots, y_n \in K$ such that $\mathcal{V}_0 = \{V_{y_j}\}_{j=1}^n$. Define $\mathcal{U}_0 \subset \mathcal{U}$ by $\mathcal{U}_0 = \{U_{y_j}\}_{j=1}^n$. Define $U, V \subset \mathcal{T}$ by $U := \bigcap_{j=1}^n U_{y_j}$ and $V := \bigcup_{j=1}^n V_{y_j}$. Since $\mathcal{V}_0$ is an open cover of K in $(X, \mathcal{T})$, we have that $K \subset V$. Since for each $y \in K$, $x \in U_y$, we have that $x \in U$. Since for each $y \in K$, $U_y \cap V_y = \emptyset$, we have that

$$\begin{aligned} U \cap V &= \left(\bigcap_{j=1}^n U_{y_j} \right) \cap \left(\bigcup_{k=1}^n V_{y_k} \right) \\ &= \bigcup_{k=1}^n \left[\left(\bigcap_{j=1}^n U_{y_j} \right) \cap V_{y_k} \right] \\ &\subset \bigcup_{k=1}^n U_{y_k} \cap V_{y_k} \\ &= \bigcup_{k=1}^n \emptyset \\ &= \emptyset \end{aligned}$$

□

Exercise 4.14.1.12. Let $(X, \mathcal{T})$ be a topological space and $U \in \mathcal{T}$. Suppose that $(X, \mathcal{T})$ is Hausdorff. If $\text{cl } U$ is compact, then for each $x \in U$, there exists $K \in \mathcal{N}(x)$ such that $K \subset U$ and K is compact.

Proof. Suppose that $\text{cl } U$ is compact. Since $U \in \mathcal{T}$, $U \cap \partial U = \emptyset$. Thus

$$\begin{aligned} x &\in U \\ &\subset (\partial U)^c \end{aligned}$$

$x \notin \partial U$. Since $\text{cl } U$ is compact, ∂U is closed and $\partial U \subset \text{cl } U$, we have that ∂U is compact. [The previous exercise](#) implies that there exist $V, W \in \mathcal{T}$ such that $V \cap W = \emptyset$, $x \in V$ and $\partial U \subset W$. Since $V \subset W^c$ and W^c is closed, we have that $\text{cl } V \subset W^c$. Since $\partial U \subset W$, we have that $(\text{cl } V) \cap \partial U = \emptyset$. Hence $\text{cl } V \subset U$. Set $K = \text{cl } V$. Since K is closed, $K \subset \text{cl } U$ and $\text{cl } U$ is compact, we have that K is compact. By construction,

$$\begin{aligned} x &\in V \\ &= \text{Int } K \end{aligned}$$

so $K \in \mathcal{N}(x)$. □

Exercise 4.14.1.13. Let $(X, \mathcal{T})$ be a topological space. If $(X, \mathcal{T})$ is compact and Hausdorff, then $(X, \mathcal{T})$ is normal.

Proof. Suppose that $(X, \mathcal{T})$ is compact and Hausdorff. Since $(X, \mathcal{T})$ is Hausdorff, $(X, \mathcal{T})$ is $\mathbf{T}_1$. Let $E, F \subset X$. Suppose that E, F are closed and $E \cap F = \emptyset$. Since $(X, \mathcal{T})$ is compact and E, F are closed, E, F are compact. [A previous exercise](#) implies that for each $x \in E$, there exists $U_x, V_x \in \mathcal{T}$ such that $U_x \cap V_x = \emptyset$, $x \in U_x$ and $F \subset V_x$. The axiom of choice implies that there exist $(U_x)_{x \in E}, (V_x)_{x \in E} \subset \mathcal{T}$ such that for each $x \in X$, $U_x \cap V_x = \emptyset$, $x \in U_x$ and $F \subset V_x$. Then $(U_x)_{x \in E}$ is an open cover of E . Since E is compact, there exist $x_1, \dots, x_n \in E$ such that $(U_{x_j})_{j=1}^n$ is an open cover of E . Define $U, V \in \mathcal{T}$ by $U := \bigcup_{j=1}^n U_{x_j}$ and $V := \bigcap_{k=1}^n V_{x_k}$.

Then

$$\begin{aligned} U \cap V &= \left(\bigcup_{j=1}^n U_{x_j} \right) \cap \left(\bigcap_{k=1}^n V_{x_k} \right) \\ &= \bigcup_{j=1}^n \left[U_{x_j} \cap \left(\bigcap_{k=1}^n V_{x_k} \right) \right] \\ &\subset \bigcup_{j=1}^n (U_{x_j} \cap V_{x_j}) \\ &= \bigcup_{j=1}^n \emptyset \\ &= \emptyset \end{aligned}$$

By construction,

$$\begin{aligned} E &\subset \bigcup_{j=1}^n U_{x_j} \\ &= U \end{aligned}$$

and

$$\begin{aligned} F &\subset \bigcap_{k=1}^n V_{x_k} \\ &= V \end{aligned}$$

Since $E, F \subset X$ with E, F closed and $E \cap F = \emptyset$ are arbitrary, we have that for each $E, F \subset X$, E, F are closed and $E \cap F = \emptyset$ implies that there exist $U, V \in \mathcal{T}$ such that $U \cap V = \emptyset$, $E \subset U$, $F \subset V$. So $(X, \mathcal{T})$ is normal. □

4.14.2 The Finite Intersection Property

Definition 4.14.2.1. Let $(X, \mathcal{T})$ be a set and $\mathcal{A} \subset \mathcal{P}(X)$. Then $\mathcal{A}$ is said to have the **finite intersection property** if for each $\mathcal{B} \subset \mathcal{A}$, $\mathcal{B}$ is finite implies that $\bigcap_{B \in \mathcal{B}} B \neq \emptyset$. We define

$$\text{FIP}(X, \mathcal{T}) = \{\mathcal{A} \subset \mathcal{P}(X) : \mathcal{A} \text{ has the finite intersection property}\}$$

and order $\text{FIP}(X, \mathcal{T})$ by inclusion.

Note 4.14.2.2. When the context is clear, we write $\text{FIP}(X)$ in place of $\text{FIP}(X, \mathcal{T})$.

Exercise 4.14.2.3. Let X be a set. Then $\text{FIP}(X)$ ordered by inclusion is a poset.

Proof. Clear. □

Exercise 4.14.2.4. Let X be a set and $\mathcal{A}_0 \in \text{FIP}(X)$. Then there exists $\mathcal{A} \in \text{FIP}(X)$ such that $\mathcal{A}$ is maximal in $[\mathcal{A}_0, \infty)$.

Proof. Let $\mathcal{C} \subset [\mathcal{A}_0, \infty)$. Suppose that $\mathcal{C}$ is a chain.

- Suppose that $\mathcal{C} = \emptyset$. Set $\mathcal{S} := \mathcal{A}_0$. Then $\mathcal{S} \in [\mathcal{A}_0, \infty)$ and it is vacuously true that for each $\mathcal{E} \in \mathcal{C}$, $\mathcal{E} \subset \mathcal{S}$. Since $\mathcal{C} \subset [\mathcal{A}_0, \infty)$ with $\mathcal{C}$ a chain is arbitrary, we have that for each $\mathcal{C} \subset [\mathcal{A}_0, \infty)$, if $\mathcal{C}$ is a chain, then there exists $\mathcal{S} \in [\mathcal{A}_0, \infty)$ such that $\mathcal{S}$ is an upper bound for $\mathcal{C}$.
- Suppose that $\mathcal{C} \neq \emptyset$. Define $\mathcal{S} \in \mathcal{P}(X)$ by $\mathcal{S} := \bigcup_{\mathcal{E} \in \mathcal{C}} \mathcal{E}$. Since $\mathcal{C} \neq \emptyset$, there exists $\mathcal{E}_0 \in [\mathcal{A}_0, \infty)$ such that $\mathcal{E}_0 \in \mathcal{C}$. Since $\mathcal{A}_0 \subset \mathcal{E}_0$, $\mathcal{A}_0 \subset \mathcal{S}$. Let $\mathcal{B} \subset \mathcal{S}$. Suppose that $\mathcal{B}$ is finite. Since $\mathcal{B} \subset \mathcal{S}$ and $\mathcal{S} = \bigcup_{\mathcal{E} \in \mathcal{C}} \mathcal{E}$, we have that for each $B \in \mathcal{B}$, there exists $\mathcal{E}_B \in \mathcal{C}$ such that $B \in \mathcal{E}_B$. Since $\mathcal{B}$ is finite and $\mathcal{C}$ is totally ordered, there exists $B_0 \in \mathcal{B}$ such that $\mathcal{E}_{B_0} = \max_{B \in \mathcal{B}} \mathcal{E}_B$. Therefore for each $B \in \mathcal{B}$,

$$\begin{aligned} B &\in \mathcal{E}_B \\ &\subset \mathcal{E}_{B_0} \end{aligned}$$

Hence $\mathcal{B} \subset \mathcal{E}_{B_0}$ which implies that

$$\bigcap_{B \in \mathcal{E}_{B_0}} B \subset \bigcap_{B \in \mathcal{B}} B.$$

Since $\mathcal{E}_{B_0} \in \mathcal{C}$, and $\mathcal{C} \subset \text{FIP}(X)$, we have that $\mathcal{E}_{B_0} \in \text{FIP}(X)$. Since $\mathcal{E}_{B_0} \in \text{FIP}(X)$, $\mathcal{B} \subset \mathcal{E}_{B_0}$ and $\mathcal{B}$ is finite, we have that $\bigcap_{B \in \mathcal{E}_{B_0}} B \neq \emptyset$. Thus there exists $x \in X$ such that $x \in \bigcap_{B \in \mathcal{E}_{B_0}} B$. Since $\bigcap_{B \in \mathcal{E}_{B_0}} B \subset \bigcap_{B \in \mathcal{B}} B$, we have that $x \in \bigcap_{B \in \mathcal{B}} B$. Hence $\bigcap_{B \in \mathcal{B}} B \neq \emptyset$. Since $\mathcal{B} \subset \mathcal{S}$ with $\mathcal{B}$ finite is arbitrary, we have that for each $\mathcal{B} \subset \mathcal{S}$, $\mathcal{B}$ is finite implies that $\bigcap_{B \in \mathcal{B}} B \neq \emptyset$. Thus $\mathcal{S} \in \text{FIP}(X)$. Since $\mathcal{A}_0 \subset \mathcal{S}$ and $\mathcal{S} \in \text{FIP}(X)$, we have that $\mathcal{S} \in [\mathcal{A}_0, \infty)$. By construction, for each $\mathcal{E} \in \mathcal{C}$, $\mathcal{E} \subset \mathcal{S}$ so that $\mathcal{S}$ is an upper bound for $\mathcal{C}$. Since $\mathcal{C} \subset [\mathcal{A}_0, \infty)$ with $\mathcal{C}$ a chain is arbitrary, we have that for each $\mathcal{C} \subset [\mathcal{A}_0, \infty)$, if $\mathcal{C}$ is a chain, then there exists $\mathcal{S} \in [\mathcal{A}_0, \infty)$ such that $\mathcal{S}$ is an upper bound for $\mathcal{C}$. Zorn's lemma implies that there exists $\mathcal{A} \in [\mathcal{A}_0, \infty]$ such that $\mathcal{A}$ is maximal. □

Exercise 4.14.2.5. Let X be a set and $\mathcal{A} \in \text{FIP}(X)$. Suppose that $\mathcal{A}$ is maximal. Then

1. for each $\mathcal{B} \subset \mathcal{A}$, if $\mathcal{B}$ is finite, then $\bigcap_{B \in \mathcal{B}} B \in \mathcal{A}$,
 2. for each $B \subset X$, if for each $A \in \mathcal{A}$, $B \cap A \neq \emptyset$, then $B \in \mathcal{A}$.
- Hint:** use part (1)

Proof.

1. Let $\mathcal{B} \subset \mathcal{A}$. Suppose that $\mathcal{B}$ is finite. Set $B_0 := \bigcap_{B \in \mathcal{B}} B$ and $\mathcal{A}_0 = \mathcal{A} \cup \{B_0\}$. Let $\mathcal{C} \subset \mathcal{A}_0$. Suppose that $\mathcal{C}$ is finite.

- Suppose that $B_0 \notin \mathcal{C}$. Since $\mathcal{A} \in \text{FIP}(X)$, $\mathcal{C} \subset \mathcal{A}$, $\mathcal{C}$ is finite,

$$\bigcap_{C \in \mathcal{C}} C = \bigcap_{C \in \mathcal{C} \cap \mathcal{A}} C \neq \emptyset$$

- Now suppose that $B_0 \in \mathcal{C}$. Since $\mathcal{C}$ is finite and $\mathcal{B}$ is finite, we have that $(\mathcal{C} \cap \mathcal{A}) \cup \mathcal{B}$ is finite. Since $\mathcal{A} \in \text{FIP}(X)$ and $(\mathcal{C} \cap \mathcal{A}) \cup \mathcal{B} \subset \mathcal{A}$,

$$\begin{aligned} \bigcap_{C \in \mathcal{C}} C &= \left(\bigcap_{C \in \mathcal{C} \cap \mathcal{A}} C \right) \cap B_0 \\ &= \left(\bigcap_{C \in \mathcal{C} \cap \mathcal{A}} C \right) \cap \left(\bigcap_{B \in \mathcal{B}} B \right) \\ &= \bigcap_{C \in (\mathcal{C} \cap \mathcal{A}) \cup \mathcal{B}} C \\ &\neq \emptyset \end{aligned}$$

Therefore $\bigcap_{C \in \mathcal{C}} C \neq \emptyset$. Since $\mathcal{C} \subset \mathcal{A}_0$ with $\mathcal{C}$ finite is arbitrary, we have that for each $\mathcal{C} \subset \mathcal{A}_0$, $\mathcal{C}$ is finite implies that $\bigcap_{C \in \mathcal{C}} C \neq \emptyset$. Hence $\mathcal{A}_0 \in \text{FIP}(X)$. Since $\mathcal{A} \in \text{FIP}(X)$ is maximal and $\mathcal{A} \subset \mathcal{A}_0$, we have that $\mathcal{A} = \mathcal{A}_0$ and therefore $B_0 \in \mathcal{A}$.

2. Let $B \subset X$. Suppose that for each $A \in \mathcal{A}$, $B \cap A \neq \emptyset$. Define $\mathcal{A}_0 = \mathcal{A} \cup \{B\}$. Let $\mathcal{C} \subset \mathcal{A}_0$. Suppose that $\mathcal{C}$ is finite.

- If $B \notin \mathcal{C}$, then

$$\bigcap_{C \in \mathcal{C}} C = \bigcap_{C \in \mathcal{C} \cap \mathcal{A}} C \neq \emptyset$$

- Suppose that $B \in \mathcal{C}$. Since $\mathcal{C} \cap \mathcal{A}$ is finite, part (1) implies that $\bigcap_{C \in \mathcal{C} \cap \mathcal{A}} C \in \mathcal{A}$. Then by assumption, $\left(\bigcap_{C \in \mathcal{C} \cap \mathcal{A}} C \right) \cap B \neq \emptyset$. Therefore

$$\begin{aligned} \bigcap_{C \in \mathcal{C}} C &= \bigcap_{C \in (\mathcal{C} \cap \mathcal{A}) \cup \{B\}} C \\ &= \left(\bigcap_{C \in \mathcal{C} \cap \mathcal{A}} C \right) \cap B \\ &\neq \emptyset \end{aligned}$$

Therefore $\bigcap_{C \in \mathcal{C}} C \neq \emptyset$. Since $\mathcal{C} \subset \mathcal{A}_0$ with $\mathcal{C}$ finite is arbitrary, we have that for each $\mathcal{C} \subset \mathcal{A}_0$, $\mathcal{C}$ is finite implies that $\bigcap_{C \in \mathcal{C}} C \neq \emptyset$. Hence $\mathcal{A}_0 \in \text{FIP}(X)$. Since $\mathcal{A} \in \text{FIP}(X)$ is maximal and $\mathcal{A} \subset \mathcal{A}_0$, we have that $\mathcal{A} = \mathcal{A}_0$ and therefore $B \in \mathcal{A}$. □

Note 4.14.2.6. Recall the definition of $\text{clSets}(A, \mathcal{T})$ in Definition 4.1.0.19.

Exercise 4.14.2.7. Let $(X, \mathcal{T})$ be a topological space. Then $(X, \mathcal{T})$ is compact iff for each $\mathcal{C} \subset \text{clSets } \emptyset(X, \mathcal{T})$, $\mathcal{C} \in \text{FIP}(X, \mathcal{T})$ implies that $\bigcap_{C \in \mathcal{C}} C \neq \emptyset$.

Hint: consider $\{C^c : C \in \mathcal{C}\}$ and whether it is an open cover

Proof.

- ($\Rightarrow$):

Suppose that $(X, \mathcal{T})$ is compact. Let $\mathcal{C} \subset \text{clSets } \emptyset$. Suppose that $\mathcal{C} \in \text{FIP}(X, \mathcal{T})$. For the sake of contradiction, suppose that $\bigcap_{C \in \mathcal{C}} C = \emptyset$. Define $\mathcal{U} \subset \mathcal{T}$ by $\mathcal{U} := \{C^c : C \in \mathcal{C}\}$. Then

$$\begin{aligned} \bigcup_{U \in \mathcal{U}} U &= \bigcup_{C \in \mathcal{C}} C^c \\ &= \left(\bigcap_{C \in \mathcal{C}} C \right)^c \\ &= \emptyset^c \\ &= X \end{aligned}$$

Thus $\mathcal{U}$ is an open cover of X . Since $(X, \mathcal{T})$ is compact, there exists $\mathcal{U}' \subset \mathcal{U}$ such that $\mathcal{U}'$ is finite and $\mathcal{U}'$ is an open cover of X . Define $\mathcal{C}' \subset \mathcal{C}$ by $\mathcal{C}' := \{U^c : U \in \mathcal{U}'\}$. Then

$$\begin{aligned} \bigcap_{C \in \mathcal{C}'} C &= \bigcap_{U \in \mathcal{U}'} U^c \\ &= \left(\bigcup_{U \in \mathcal{U}'} U \right)^c \\ &= X^c \\ &= \emptyset \end{aligned}$$

However, since $\mathcal{C} \in \text{FIP}(X, \mathcal{T})$, $\mathcal{C}' \subset \mathcal{C}$ and $\mathcal{C}'$ is finite, we have that $\bigcap_{C \in \mathcal{C}'} C \neq \emptyset$. This is a contradiction. Hence $\bigcap_{C \in \mathcal{C}} C \neq \emptyset$.

Since $\mathcal{C} \subset \text{clSets } \emptyset$ such that $\mathcal{C} \in \text{FIP}(X, \mathcal{T})$ is arbitrary, we have that for each $\mathcal{C} \subset \text{clSets } \emptyset$, $\mathcal{C} \in \text{FIP}(X, \mathcal{T})$ implies that $\bigcap_{C \in \mathcal{C}} C \neq \emptyset$.

- ($\Leftarrow$):

Suppose that for each $\mathcal{C} \subset \text{clSets } \emptyset$, $\mathcal{C} \in \text{FIP}(X, \mathcal{T})$ implies that $\bigcap_{C \in \mathcal{C}} C \neq \emptyset$. Let $\mathcal{U} \subset \mathcal{P}(X)$. Suppose that $\mathcal{U}$ is an open cover of X in $(X, \mathcal{T})$. Then $\mathcal{U} \subset \mathcal{T}$ and $X = \bigcup_{U \in \mathcal{U}} U$. Define $\mathcal{C} \subset \text{clSets } \emptyset$ by $\mathcal{C} = \{U^c : U \in \mathcal{U}\}$. Then

$$\begin{aligned} \bigcap_{C \in \mathcal{C}} C &= \bigcap_{U \in \mathcal{U}} U^c \\ &= \left(\bigcup_{U \in \mathcal{U}} U \right)^c \\ &= X^c \\ &= \emptyset \end{aligned}$$

By assumption, $\mathcal{C} \notin \text{FIP}(X, \mathcal{T})$. Thus there exists $\mathcal{C}' \subset \mathcal{C}$ such that $\mathcal{C}'$ is finite and $\bigcap_{C \in \mathcal{C}'} C = \emptyset$. Define $\mathcal{U}' \subset \mathcal{U}$ by $\mathcal{U}' := \{C^c : C \in \mathcal{C}'\}$. Then $\mathcal{U}'$ is finite and

$$\begin{aligned} X &= \emptyset^c \\ &= \left(\bigcap_{C \in \mathcal{C}'} C \right)^c \\ &= \bigcup_{C \in \mathcal{C}'} C^c \\ &= \bigcup_{U \in \mathcal{U}'} U \end{aligned}$$

Hence $\mathcal{U}'$ is an open cover of X in $(X, \mathcal{T})$. Since $\mathcal{U} \subset \mathcal{P}(X)$ such that $\mathcal{U}$ is an open cover of X in $(X, \mathcal{T})$, we have that for each $\mathcal{U} \subset \mathcal{P}(X)$, $\mathcal{U}$ is an open cover of X in $(X, \mathcal{T})$ implies that there exists $\mathcal{U}' \subset \mathcal{U}$ such that $\mathcal{U}'$ is finite and $\mathcal{U}'$ is an open cover of X in $(X, \mathcal{T})$. Thus $(X, \mathcal{T})$ is compact. □

Exercise 4.14.2.8. Let $(X, \mathcal{T})$ be a topological space. Then the following are equivalent:

1. $(X, \mathcal{T})$ is compact
2. for each net $(x_\alpha)_{\alpha \in A} \subset X$, there exists $x \in X$ such that x is a cluster point of $(x_\alpha)_{\alpha \in A}$.
3. for each net $(x_\alpha)_{\alpha \in A} \subset X$, there exists a subnet $(x_{\alpha_\beta})_{\beta \in B}$ of $(x_\alpha)_{\alpha \in A}$ and $x \in X$ such that $x_{\alpha_\beta} \rightarrow x$.

Hint:

- (1) $\implies$ (2):
For $\alpha \in A$, set $E_\alpha := \{x_{\alpha'} : \alpha' \geq \alpha\}$. Then $\{\text{cl } E_\alpha : \alpha \in A\} \in \text{FIP}(X, \mathcal{T})$.
- (3) $\implies$ (1):
If $(X, \mathcal{T})$ is not compact, choose open cover $\mathcal{U}$ of X such that for each $\mathcal{U}_0 \subset \mathcal{U}$, $\mathcal{U}_0$ is finite implies that $\mathcal{U}_0$ is not an open cover of X . Consider $\mathcal{F}_\mathcal{U} = \{\mathcal{U}' \subset \mathcal{U} : \mathcal{U}' \text{ is finite}\}$ ordered by inclusion. Then there exists a net $(x_{\mathcal{U}'})_{\mathcal{U}' \in \mathcal{F}_\mathcal{U}} \subset X$ such that for each $x_{\mathcal{U}'} \notin \bigcup_{U \in \mathcal{U}'} U$.

Proof.

- (1) $\implies$ (2):
Suppose that $(X, \mathcal{T})$ is compact. Let $(x_\alpha)_{\alpha \in A} \subset X$ be a net. For $\alpha_0 \in A$, define $E_{\alpha_0} = \{x_{\alpha'} : \alpha' \geq \alpha_0\}$. Then for each $\alpha_1, \alpha_2 \in A$, $\alpha_1 \leq \alpha_2$ implies that $E_{\alpha_2} \subset E_{\alpha_1}$. Since A is directed, for each $\alpha \in A$, $E_\alpha \neq \emptyset$ and for each $A_0 \subset A$, A_0 is finite implies that there exists $\alpha_0 \in A$ such that for each $\alpha \in A_0$, $\alpha_0 \geq \alpha$.
Define $\mathcal{E} \subset \mathcal{P}(X)$ by $\mathcal{E} := \{\text{cl } E_\alpha : \alpha \in A\}$. Let $A_0 \subset A$. Suppose that A_0 is finite. Then there exists $\alpha_0 \in A$ such that for each $\alpha \in A_0$, $\alpha_0 \geq \alpha$. Then for each $\alpha \in A$, $E_{\alpha_0} \subset E_\alpha$. Thus

$$\begin{aligned} \emptyset &\neq E_{\alpha_0} \\ &\subset \bigcap_{\alpha \in A_0} E_\alpha \\ &\subset \bigcap_{\alpha \in A_0} \text{cl } E_\alpha \end{aligned}$$

Since $A_0 \subset A$ with A_0 finite is arbitrary, we have that for each $A_0 \subset A$, A_0 is finite implies that $\bigcap_{\alpha \in A_0} \text{cl } E_\alpha \neq \emptyset$. Thus $\mathcal{E} \in \text{FIP}(X, \mathcal{T})$. Since $(X, \mathcal{T})$ is compact, [the previous exercise](#) implies that $\bigcap_{\alpha \in A} \text{cl } E_\alpha \neq \emptyset$. Thus there exists $x \in X$ such that $x \in \bigcap_{\alpha \in A} \text{cl } E_\alpha$. Exercise 4.3.2.20 implies that x is a cluster point of $(x_\alpha)_{\alpha \in A}$.

- (2) $\implies$ (3):
Immediate by Exercise 4.3.2.20.
- (3) $\implies$ (1):
Suppose that $(X, \mathcal{T})$ is not compact. Then there exists $\mathcal{U} \subset \mathcal{P}(X)$ such that $\mathcal{U}$ is an open cover of X in $(X, \mathcal{T})$ and for each $\mathcal{U}_0 \subset \mathcal{U}$, $\mathcal{U}_0$ is finite implies that $\mathcal{U}_0$ is not an open cover of X in $(X, \mathcal{T})$. Define $\mathcal{F}_\mathcal{U} \subset \mathcal{P}(X)$ by $\mathcal{F}_\mathcal{U} := \{\mathcal{U}_0 \subset \mathcal{U} : \mathcal{U}_0 \text{ is finite}\}$. We define $\leq \subset \mathcal{F}_\mathcal{U} \times \mathcal{F}_\mathcal{U}$ by inclusion so that $\mathcal{U}_1 \leq \mathcal{U}_2$ iff $\mathcal{U}_1 \subset \mathcal{U}_2$. Then $(\mathcal{F}_\mathcal{U}, \leq)$ is a directed set. By construction for each $\mathcal{U}' \in \mathcal{F}_\mathcal{U}$, $\left(\bigcup_{U \in \mathcal{U}'} U\right)^c \neq \emptyset$. The axiom of choice implies that there exists a net $(x_{\mathcal{U}'})_{\mathcal{U}' \in \mathcal{F}_\mathcal{U}} \subset X$ such that for each $\mathcal{U}' \in \mathcal{F}_\mathcal{U}$, $x_{\mathcal{U}'} \in \left(\bigcup_{U \in \mathcal{U}'} U\right)^c$.

For the sake of contradiction suppose that there exists a subnet $(x_{\mathcal{U}'_\beta})_{\beta \in B}$ of $(x_{\mathcal{U}'})_{\mathcal{U}' \in \mathcal{F}_U}$ and $x \in X$ such that $x_{\mathcal{U}'_\beta} \rightarrow x$. Since $\mathcal{U}$ is an open cover of X in $(X, \mathcal{T})$, there exists $U_0 \in \mathcal{U}$ such that $x \in U_0$. Since $x_{\mathcal{U}'_\beta} \rightarrow x$ and $U_0 \in \mathcal{N}(x)$, there exists $\beta_0 \in B$ such that for each $\beta \in B$, $\beta \geq \beta_0$ implies that $x_{\mathcal{U}'_\beta} \in U_0$. Define $\mathcal{U}_0 \in \mathcal{F}_U$ by $\mathcal{U}_0 := \{U_0\}$. Since $(x_{\mathcal{U}'_\beta})_{\beta \in B}$ is a subnet of $(x_{\mathcal{U}'})_{\mathcal{U}' \in \mathcal{F}_U}$, there exists $\beta_1 \in B$ such that for each $\beta \in B$, $\beta \geq \beta_1$ implies that $\mathcal{U}'_\beta \geq \mathcal{U}_0$. Since B is a directed set, there exists $\beta_2 \in B$ such that $\beta_2 \geq \beta_0, \beta_1$.

Since $\beta_2 \geq \beta_0$, we have that $x_{\mathcal{U}'_{\beta_2}} \in U_0$. Since $\beta_2 \geq \beta_1$, we have that $\mathcal{U}'_{\beta_2} \geq \mathcal{U}_0$. Hence $\mathcal{U}_0 \subset \mathcal{U}'_{\beta_2}$ and therefore

$$\begin{aligned} U_0 &= \bigcup_{U \in \mathcal{U}_0} U \\ &\subset \bigcup_{U \in \mathcal{U}'_{\beta_2}} U \end{aligned}$$

By construction,

$$\begin{aligned} x_{\mathcal{U}'_{\beta_2}} &\in \left(\bigcup_{U \in \mathcal{U}'_{\beta_2}} U \right)^c \\ &\subset \left(\bigcup_{U \in \mathcal{U}_0} U \right)^c \\ &= U_0^c \end{aligned}$$

This is a contradiction. Thus for each subnet $(x_{\mathcal{U}'_\beta})_{\beta \in B}$ of $(x_{\mathcal{U}'})_{\mathcal{U}' \in \mathcal{F}_U}$ and $x \in X$, we have that $x_{\mathcal{U}'_\beta} \not\rightarrow x$. Therefore there exists a net $(x_\alpha)_{\alpha \in A} \subset X$ such that for each subnet $(x_{\alpha_\beta})_{\beta \in B}$ of $(x_\alpha)_{\alpha \in A}$ and $x \in X$, $x_{\alpha_\beta} \not\rightarrow x$. By contrapositive, we have that (3) $\implies$ (1).

□

4.14.3 Compactness and Product Spaces

Exercise 4.14.3.1. Tychonoff's Theorem:

Let $(X_\alpha, \mathcal{T}_\alpha)_{\alpha \in A}$ be a collection of topological spaces. Suppose that for each $\alpha \in A$, $(X_\alpha, \mathcal{T}_\alpha)$ is compact. Then $\left(\prod_{\alpha \in A} X_\alpha, \bigotimes_{\alpha \in A} \mathcal{T}_\alpha \right)$ is compact.

Hint:

Proof. Set $X := \prod_{\alpha \in A} X_\alpha$ and $\mathcal{T} := \bigotimes_{\alpha \in A} \mathcal{T}_\alpha$. Let $\mathcal{C} \subset \text{clSets } \emptyset$. Suppose that $\mathcal{C} \in \text{FIP}(X, \mathcal{T})$. A previous exercise implies that there exists $\mathcal{D} \in \text{FIP}(X, \mathcal{T})$ such that $\mathcal{D}$ is maximal in $[\mathcal{C}, \infty)$. Let $\alpha \in A$. Set $\mathcal{D}_\alpha := \{\pi_\alpha(D) : D \in \mathcal{D}\}$.

Let $\mathcal{D}_{\alpha,0} \subset \mathcal{D}_\alpha$. Suppose that $\mathcal{D}_{\alpha,0}$ is finite. Then there exist $\mathcal{D}_0 \subset \mathcal{D}$ such that $\mathcal{D}_0$ is finite and $\mathcal{D}_{\alpha,0} = \{\pi_\alpha(D) : D \in \mathcal{D}_0\}$. Since $\mathcal{D} \in \text{FIP}(X, \mathcal{T})$, $\bigcap_{D \in \mathcal{D}_0} D \neq \emptyset$. Therefore

$$\begin{aligned} \emptyset &\neq \pi_\alpha \left(\bigcap_{D \in \mathcal{D}_0} D \right) \\ &\subset \bigcap_{D \in \mathcal{D}_0} \pi_\alpha(D) \\ &= \bigcap_{D \in \mathcal{D}_{\alpha,0}} D \end{aligned}$$

Since $\mathcal{D}_{\alpha,0} \subset \mathcal{D}_\alpha$ with $\mathcal{D}_{\alpha,0}$ finite is arbitrary, we have that for each $\mathcal{D}_{\alpha,0} \subset \mathcal{D}_\alpha$, $\mathcal{D}_{\alpha,0}$ is finite implies that $\bigcap_{D \in \mathcal{D}_{\alpha,0}} D \neq \emptyset$. Hence

$\mathcal{D}_\alpha \in \text{FIP}(X, \mathcal{T})$.

For the sake of contradiction, suppose that $\bigcap_{D \in \mathcal{D}} \text{cl } \pi_\alpha(D) = \emptyset$. Then $X = \bigcup_{D \in \mathcal{D}} [\text{cl } \pi_\alpha(D)]^c$. Since $\{[\text{cl } \pi_\alpha(D)]^c : D \in \mathcal{D}\} \subset \mathcal{T}_\alpha$ and $(X_\alpha, \mathcal{T}_\alpha)$ is compact, there exists $\mathcal{D}_0 \subset \mathcal{D}$ such that $\mathcal{D}_0$ is finite and $X = \bigcup_{D \in \mathcal{D}_0} [\text{cl } \pi_\alpha(D)]^c$. Therefore $\bigcap_{D \in \mathcal{D}_0} \text{cl } \pi_\alpha(D) = \emptyset$.

Since $\mathcal{D}_\alpha \in \text{FIP}(X, \mathcal{T})$ and $\{\text{cl } \pi_\alpha(D) : D \in \mathcal{D}_0\} \subset \mathcal{D}_\alpha$ is finite, we have that $\bigcap_{D \in \mathcal{D}_0} \text{cl } \pi_\alpha(D) \neq \emptyset$. This is a contradiction. Hence $\bigcap_{D \in \mathcal{D}} \text{cl } \pi_\alpha(D) \neq \emptyset$. Since $\alpha \in A$ is arbitrary, we have that for each $\alpha \in A$, $\bigcap_{D \in \mathcal{D}} \text{cl } \pi_\alpha(D) \neq \emptyset$.

The axiom of choice implies that there exists $x \in X$ such that for each $\alpha \in A$, $x_\alpha \in \bigcap_{D \in \mathcal{D}} \text{cl } \pi_\alpha(D)$. Set

$$\mathcal{E}_x := \{\pi_\alpha^{-1}(E_\alpha) : \alpha \in A, E_\alpha \in \mathcal{T}_\alpha, x_\alpha \in E_\alpha\}$$

Let $\alpha \in A$ and $E_\alpha \in \mathcal{T}_\alpha$. Suppose that $x \in \pi_\alpha^{-1}(E_\alpha)$. Then $x_\alpha \in E_\alpha$. Let $D \in \mathcal{D}$. Since $x_\alpha \in E_\alpha \cap \text{cl } \pi_\alpha(D)$, $E_\alpha \cap \text{cl } \pi_\alpha(D) \neq \emptyset$. Exercise 4.3.2.12 implies that $E_\alpha \cap \pi_\alpha(D) \neq \emptyset$. Therefore $\pi_\alpha^{-1}(E_\alpha) \cap D \neq \emptyset$. Since $D \in \mathcal{D}$ is arbitrary, we have that for each $D \in \mathcal{D}$, $\pi_\alpha^{-1}(E_\alpha) \cap D \neq \emptyset$. Since $\mathcal{D} \in \text{FIP}(X, \mathcal{T})$, a previous exercise implies that $\pi_\alpha^{-1}(E_\alpha) \in \mathcal{D}$. Since $\alpha \in A$ and $E_\alpha \in \mathcal{T}_\alpha$ are arbitrary, we have that $\mathcal{E}_x \subset \mathcal{D}$. Set

$$\mathcal{B}_x := \left\{ \bigcap_{j=1}^n V_j : (V_j)_{j=1}^n \subset \mathcal{E}_x \right\}$$

Then an exercise in the section on the product topology implies that $\mathcal{B}_x \subset \mathcal{T}_X$ and $\mathcal{B}_x$ is a local basis for $\mathcal{T}$ at x . Since $\mathcal{D} \in \text{FIP}(X, \mathcal{T})$, a previous exercise implies that $\mathcal{B}_x \subset \mathcal{D}$.

Let $D \in \mathcal{D}$ and $E \in \mathcal{B}_x$. Since $\mathcal{D} \in \text{FIP}(X, \mathcal{T})$ and $D, E \in \mathcal{D}$, we have that $D \cap E \neq \emptyset$. Since $E \in \mathcal{B}_x$ is arbitrary we have that for each $E \in \mathcal{B}_x$, $D \cap E \neq \emptyset$. An exercise in the introduction to topology section implies that $x \in \text{cl } D$. Since $D \in \mathcal{D}$ is arbitrary, we have that for each $D \in \mathcal{D}$, $x \in \text{cl } D$. Thus $x \in \bigcap_{D \in \mathcal{D}} \text{cl } D$ and therefore $\bigcap_{D \in \mathcal{D}} \text{cl } D \neq \emptyset$. Since $\mathcal{C} \subset \text{clSets } \emptyset$, for each $C \in \mathcal{C}$, $C = \text{cl } C$.

By construction, $\mathcal{C} \subset \mathcal{D}$, which implies that

$$\begin{aligned} \emptyset &\neq \bigcap_{D \in \mathcal{D}} \text{cl } D \\ &\subset \bigcap_{C \in \mathcal{C}} \text{cl } C \\ &= \bigcap_{C \in \mathcal{C}} C \end{aligned}$$

Since $\mathcal{C} \subset \text{clSets } \emptyset$ with $\mathcal{C} \in \text{FIP}(X, \mathcal{T})$ is arbitrary, we have that for each $\mathcal{C} \subset \text{clSets } \emptyset$, $\mathcal{C} \in \text{FIP}(X, \mathcal{T})$ implies that $\bigcap_{C \in \mathcal{C}} C \neq \emptyset$. The previous exercise implies that $(X, \mathcal{T})$ is compact. $\square$

Exercise 4.14.3.2. Let $(X_\alpha, \mathcal{T}_\alpha)_{\alpha \in A}$ be a collection of topological spaces. Set $X := \prod_{\alpha \in A} X_\alpha$ and $\mathcal{T}_X := \bigotimes_{\alpha \in A} \mathcal{T}_\alpha$. Let $\alpha_0 \in A$. Set $A' := A \setminus \{\alpha_0\}$, $Y := \prod_{\alpha \in A'} X_\alpha$ and $\mathcal{T}_Y := \bigotimes_{\alpha \in A'} \mathcal{T}_\alpha$. Let $E \subset X$ and $a \in \pi_{\alpha_0}(E)^c$. Suppose that for each $\alpha \in A$, $(X_\alpha, \mathcal{T}_\alpha)$ is compact and E is closed in $(X, \mathcal{T}_X)$.

1. For each $y \in Y$, there exists $x^y \in X$ such that $\pi_{\alpha_0}(x^y) = a$, for each $\alpha \in A'$, $\pi_\alpha(x^y) = y_\alpha$ and $x^y \in E^c$.
2. For each $y \in Y$, there exists $U_{\alpha_0}^y \in \mathcal{T}_{\alpha_0}$, $V^y \in \mathcal{T}_Y$ and $W^y \in \mathcal{T}_X$ such that $U_{\alpha_0}^y = \pi_{\alpha_0}(W^y)$, $V^y = \prod_{\alpha \in A'} \pi_\alpha(W^y)$, $a \in U_{\alpha_0}^y$, $y \in V^y$ and $x^y \in W^y \subset E^c$.

Hint: E is closed

3. There exist $U_{\alpha_0} \in \mathcal{T}_{\alpha_0}$ such that

- (a) $a \in U_{\alpha_0}$,

Hint: consider the intersection of a finite open cover of Y .

- (b) $U_{\alpha_0} \subset \pi_{\alpha_0}(E)^c$

Hint: proof by contradiction

Proof.

1. Let $y \in Y$. Define $x^y \in X$ by $x_\alpha^y := \begin{cases} a, & \alpha = \alpha_0 \\ y_\alpha, & \alpha \neq \alpha_0. \end{cases}$ For the sake of contradiction, suppose that $x^y \in E$. Then

$$\begin{aligned} a &= x_{\alpha_0}^y \\ &= \pi_{\alpha_0}(x^y) \\ &\in \pi_{\alpha_0}(E). \end{aligned}$$

This is a contradiction since $a \in (\pi_{\alpha_0}(E))^c$. Hence $x^y \in E^c$.

2. Since E is closed in $(X, \mathcal{T}_X)$, $E^c \in \mathcal{T}_X$. Exercise 4.6.1.2 then implies that there exist $(U_\alpha^y)_{\alpha \in A} \in \prod_{\alpha \in A} \mathcal{T}_\alpha$ such that $\prod_{\alpha \in A} U_\alpha^y \in \mathcal{T}_X$, $x^y \in \prod_{\alpha \in A} U_\alpha^y \subset E^c$ and $\#\{\alpha \in A : U_\alpha^y \neq X_\alpha\} < \infty$. Set $W^y := \prod_{\alpha \in A} U_\alpha^y$ and $V^y := \prod_{\alpha \in A'} U_\alpha^y$. Since $\#\{\alpha \in A : U_\alpha^y \neq X_\alpha\} < \infty$, we have that $\#\{\alpha \in A' : U_\alpha^y \neq X_\alpha\} < \infty$ and therefore $V^y \in \mathcal{T}_Y$. Then

$$\begin{aligned} a &= x_{\alpha_0}^y \\ &\in U_{\alpha_0}^y \end{aligned}$$

and $y \in V^y$. By construction, $U_{\alpha_0}^y = \pi_{\alpha_0}(W^y)$ and $V^y = \prod_{\alpha \in A'} \pi_\alpha(W^y)$. Since $y \in Y$ is arbitrary, we have that for each $y \in Y$, there exists $U_{\alpha_0}^y \in \mathcal{T}_{\alpha_0}$, $V^y \in \mathcal{T}_Y$ and $W^y \in \mathcal{T}_X$ such that $U_{\alpha_0}^y = \pi_{\alpha_0}(W^y)$, $V^y = \prod_{\alpha \in A'} \pi_\alpha(W^y)$, $a \in U_{\alpha_0}^y$, $y \in V^y$ and $x^y \in W^y \subset E^c$.

3. (a) Exercise 4.14.3.1 implies that $(Y, \mathcal{T}_Y)$ is compact. Since $Y \subset \bigcup_{y \in Y} V^y$ and $(Y, \mathcal{T}_Y)$ is compact, there exists $y_1, \dots, y_n \in Y$

such that $Y \subset \bigcup_{j=1}^n V^{y_j}$. Set $U_{\alpha_0} := \bigcap_{j=1}^n U_{\alpha_0}^{y_j}$. Then $U_{\alpha_0} \in \mathcal{T}_{\alpha_0}$ and $a \in U_{\alpha_0}$.

- (b) For the sake of contradiction, suppose that $U_{\alpha_0} \not\subset \pi_{\alpha_0}(E)^c$. Then there exists $b \in U_{\alpha_0}$ such that $b \in \pi_{\alpha_0}(E)$. Since $b \in \pi_{\alpha_0}(E)$, there exists $z \in E$ such that $\pi_{\alpha_0}(z) = b$. Since $Y \subset \bigcup_{j=1}^n V^{y_j}$, there exists $j_0 \in [n]$ such that $(z_\alpha)_{\alpha \in A'} \in V^{y_{j_0}}$. Since

$$\begin{aligned} z_{\alpha_0} &= b \\ &\in U_{\alpha_0} \\ &= \bigcap_{j=1}^n U_{\alpha_0}^{y_j} \\ &\subset U_{\alpha_0}^{y_{j_0}}, \end{aligned}$$

$$\begin{aligned} (z_\alpha)_{\alpha \in A'} &\in V^{y_{j_0}} \\ &= \prod_{\alpha \in A'} U_\alpha^{y_{j_0}} \end{aligned}$$

and $W^{y_{j_0}} = \prod_{\alpha \in A} U_\alpha^{y_{j_0}}$, we have that $z \in W^{y_{j_0}}$. Since $z \in E$,

$$\begin{aligned} z &\in W^{y_{j_0}} \cap E \\ &= \emptyset, \end{aligned}$$

which is a contradiction. Hence $U_{\alpha_0} \subset \pi_{\alpha_0}(E)^c$.

□

Exercise 4.14.3.3. Let $(X_\alpha, \mathcal{T}_\alpha)_{\alpha \in A}$ be a collection of topological spaces. Suppose that for each $\alpha \in A$, $(X_\alpha, \mathcal{T}_\alpha)$ is compact. Then for each $\alpha \in A$, $\pi_\alpha : \prod_{\alpha \in A} X_\alpha \rightarrow X_\alpha$ is closed.

Hint: Exercise 4.14.3.2

Proof. Set $X := \prod_{\alpha \in A} X_\alpha$ and $\mathcal{T}_X := \bigotimes_{\alpha \in A} \mathcal{T}_\alpha$. Let $\alpha_0 \in A$ and $E \subset X$. Suppose that E is closed in $(X, \mathcal{T}_X)$. Let $a \in \pi_{\alpha_0}(E)^c$.

Exercise 4.14.3.2 implies that there exists $U_{\alpha_0} \in \mathcal{T}_{\alpha_0}$ such that $a \in U_{\alpha_0}$ and $U_{\alpha_0} \subset \pi_{\alpha_0}(E)^c$. Since $a \in \pi_{\alpha_0}(E)^c$ is arbitrary, we have that for each $a \in \pi_{\alpha_0}(E)^c$, there exists $U_{\alpha_0} \in \mathcal{T}_{\alpha_0}$ such that $a \in U_{\alpha_0}$ and $U_{\alpha_0} \subset \pi_{\alpha_0}(E)^c$. Therefore $\pi_{\alpha_0}(E)^c \in \mathcal{T}_{\alpha_0}$ and $\pi_{\alpha_0}(E)$ is closed in $(X_{\alpha_0}, \mathcal{T}_{\alpha_0})$. Since $E \subset X$ with E closed in $(X, \mathcal{T}_X)$ is arbitrary, we have that for each $E \subset X$, E closed in $(X, \mathcal{T}_X)$ implies that $\pi_{\alpha_0}(E)$ is closed in $(X_{\alpha_0}, \mathcal{T}_{\alpha_0})$. Hence π_{α_0} is closed. Since $\alpha_0 \in A$ is arbitrary, we have that for each $\alpha \in A$, $\pi_\alpha : \prod_{\alpha \in A} X_\alpha \rightarrow X_\alpha$

is closed. □

4.14.4 Compactness and Coproduct Spaces

4.14.5 Compactness and Quotient Spaces

Exercise 4.14.5.1. Let $(X, \mathcal{T})$ be a topological space and $\sim \subset X \times X$ an equivalence relation. If $(X, \mathcal{T})$ is compact, then $(X/\sim, \mathcal{T}/\sim)$ is compact.

Proof. Suppose that $(X, \mathcal{T})$ is compact. Let $\mathcal{V} \subset \mathcal{P}(X/\sim)$. Suppose that $\mathcal{V}$ is an open cover of $X/\sim$ in $(X, \mathcal{T}/\sim)$. Then $\mathcal{V} \subset \mathcal{T}/\sim$ and $X/\sim \subset \bigcup_{V \in \mathcal{V}} V$. Define $\mathcal{U} \subset \mathcal{P}(X)$ by $\mathcal{U} := \{\pi^{-1}(V) : V \in \mathcal{V}\}$. Since π is $(\mathcal{T}, \mathcal{T}/\sim)$ -continuous and $\mathcal{V} \subset \mathcal{T}/\sim$, we have that

$$\begin{aligned} \mathcal{U} &= \{\pi^{-1}(V) : V \in \mathcal{V}\} \\ &\in \mathcal{T}. \end{aligned}$$

Since $X/\sim \subset \bigcup_{V \in \mathcal{V}} V$, we have that

$$\begin{aligned} X &= \pi^{-1}(X/\sim) \\ &\subset \pi^{-1}\left(\bigcup_{V \in \mathcal{V}} V\right) \\ &= \bigcup_{V \in \mathcal{V}} \pi^{-1}(V) \\ &= \bigcup_{U \in \mathcal{U}} U \end{aligned}$$

Thus $\mathcal{U}$ is an open cover of X in $(X, \mathcal{T})$. Since $(X, \mathcal{T})$ is compact, there exists $\mathcal{U}_0 \subset \mathcal{U}$ such that $\mathcal{U}_0$ is an open cover of X in $(X, \mathcal{T})$ and $\mathcal{U}_0$ is finite. For each $U \in \mathcal{U}_0$, define $A_U \subset \mathcal{V}$ by $A_U := \{V \in \mathcal{V} : \pi^{-1}(V) = U\}$. By construction, for each $U \in \mathcal{U}_0$, $A_U \neq \emptyset$. Since $\mathcal{U}_0$ is finite, $\prod_{U \in \mathcal{U}_0} A_U \neq \emptyset$. Thus there exists $(V_U)_{U \in \mathcal{U}_0} \in \prod_{U \in \mathcal{U}_0} A_U$. Define $\mathcal{V}_0 \subset \mathcal{P}(X/\sim)$ by $\mathcal{V}_0 := \{V_U : U \in \mathcal{U}_0\}$. Then $\mathcal{V}_0$ is finite. By construction,

$$\begin{aligned} \mathcal{V}_0 &\subset \mathcal{V} \\ &\subset \mathcal{T}/\sim \end{aligned}$$

and for each $U \in \mathcal{U}_0$, $\pi^{-1}(V_U) = U$. Since π is surjective, we have that for each $U \in \mathcal{U}_0$,

$$\begin{aligned} V_U &= V_U \cap X/\sim \\ &= \pi(\pi^{-1}(V_U)) \\ &= \pi(U). \end{aligned}$$

Therefore

$$\begin{aligned}
 X/\sim &= \pi(X) \\
 &\subset \pi\left(\bigcup_{U \in \mathcal{U}_0} U\right) \\
 &= \bigcup_{U \in \mathcal{U}_0} \pi(U) \\
 &= \bigcup_{U \in \mathcal{U}_0} V_U \\
 &= \bigcup_{V \in \mathcal{V}_0} V.
 \end{aligned}$$

Hence $\mathcal{V}_0$ is an open cover of $X/\sim$. Since $\mathcal{V} \subset \mathcal{P}(X)$ such that $\mathcal{V}$ is an open cover of $X/\sim$ is arbitrary, we have that for each $\mathcal{V} \subset \mathcal{P}(X)$, if $\mathcal{V}$ is an open cover of $X/\sim$ in $(X/\sim, \mathcal{T}/\sim)$, then there exists $\mathcal{V}_0 \subset \mathcal{V}$ such that $\mathcal{V}_0$ is an open cover of $X/\sim$ in $(X/\sim, \mathcal{T}/\sim)$ and $\mathcal{V}_0$ is finite. Thus $(X/\sim, \mathcal{T}/\sim)$ is compact. $\square$

4.14.6 Compactness and Projective Systems

Exercise 4.14.6.1. Let $(J, \leq)$ be a directed poset, $((X_j, \mathcal{T}_j)_{j \in J}, (\pi_{j,k})_{(j,k) \in \leq})$ a projective system of topological spaces. Set $X := \varprojlim_{j \in J} X_j$ and $\mathcal{T} := \varprojlim_{j \in J} \mathcal{T}_j$. If for each $j \in J$, $(X_j, \mathcal{T}_j)$ is compact Hausdorff, then for each $j \in J$, $\pi_j : X \rightarrow X_j$ is $(\mathcal{T}, \mathcal{T}_j)$ -closed.

Proof. Suppose that for each $j \in J$, $(X_j, \mathcal{T}_j)$ is compact Hausdorff. Let $j_0 \in J$. Set $X_0 := \prod_{j \in J} X_j$ and $\mathcal{T}_0 := \bigotimes_{j \in J} \mathcal{T}_j$. Since for each $j \in J$, $(X_j, \mathcal{T}_j)$ is compact, Exercise 4.14.3.3 implies that for each $j \in J$, $\text{proj}_j : X_0 \rightarrow X_j$ is $(\mathcal{T}_0, \mathcal{T}_j)$ -closed. Since for each $j \in J$, $(X_j, \mathcal{T}_j)$ is Hausdorff, Exercise 4.10.0.5 implies that X is closed in $(X_0, \mathcal{T}_0)$. Since $\pi_{j_0} = \text{proj}_{j_0}|_X$, Exercise 4.5.1.18 then implies that $\pi_{j_0} : X \rightarrow X_{j_0}$ is $(\mathcal{T}, \mathcal{T}_{j_0})$ -closed. Since $j_0 \in J$ is arbitrary, we have that for each $j \in J$, $\pi_j : X \rightarrow X_j$ is $(\mathcal{T}, \mathcal{T}_j)$ -closed. $\square$

Exercise 4.14.6.2. Let $(J, \leq)$ be a directed poset, $((X_j, \mathcal{T}_j)_{j \in J}, (\pi_{j,k})_{(j,k) \in \leq})$ a projective system of topological spaces. Set $((X, \mathcal{T}), (\pi_j)_{j \in J}) := \varprojlim_{j \in J} ((X_j, \mathcal{T}_j)_{j \in J}, (\pi_{j,k})_{(j,k) \in \leq})$. If for each $j \in J$, $(X_j, \mathcal{T}_j)$ is a compact Hausdorff space, then $(X, \mathcal{T})$ is a compact Hausdorff space.

Proof. Suppose that for each $j \in J$, $(X_j, \mathcal{T}_j)$ is compact Hausdorff. Set $X_0 := \prod_{j \in J} X_j$ and $\mathcal{T}_0 := \bigotimes_{j \in J} \mathcal{T}_j$. Since for each $j \in J$, $(X_j, \mathcal{T}_j)$ is Hausdorff, Exercise 4.12.3.1 implies that $(X_0, \mathcal{T}_0)$ is Hausdorff. Since $\mathcal{T} = \mathcal{T}_0 \cap X$, Exercise 4.12.2.2 implies that $(X, \mathcal{T})$ is Hausdorff. Since for each $j \in J$, $(X_j, \mathcal{T}_j)$ is compact, Exercise 4.14.3.1 implies that $(X_0, \mathcal{T}_0)$ is compact. Exercise 4.10.0.5 implies that X is closed in $(X_0, \mathcal{T}_0)$. Since X is closed in $(X_0, \mathcal{T}_0)$, Exercise 4.14.1.9 implies that $(X, \mathcal{T})$ is compact. $\square$

Exercise 4.14.6.3. Let $(J, \leq)$ be a directed poset $((X_j)_{j \in J}, (\pi_{j,k})_{(j,k) \in \leq})$ be a projective system of topological spaces. Set $(X, (\pi_j)_{j \in J}) := \varprojlim_{j \in J} ((X_j)_{j \in J}, (\pi_{j,k})_{(j,k) \in \leq})$. If for each $j \in J$, X_j is a compact Hausdorff space and π_j is surjective, then $(X, (\pi_j)_{j \in J})$ is perfect.

Proof. Suppose that for each $j \in J$, X_j is a compact Hausdorff space and π_j is surjective. Let $j \in J$. Since for each $j' \in J$, $X_{j'}$ is compact Hausdorff, Exercise 4.14.6.1 implies that $\pi_{j'}$ is closed. Since π_j is continuous, closed and surjective, Exercise 4.8.1.8 implies that π_j is a quotient map. Since $j \in J$ is arbitrary, we have that for each $j \in J$, π_j is a quotient map. Thus X is perfect. $\square$

4.14.7 Compactness and Continuity

Exercise 4.14.7.1. Let $(X, \mathcal{T}_X)$ and $(Y, \mathcal{T}_Y)$ be topological spaces and $f : X \rightarrow Y$. Suppose that f is $(\mathcal{T}_X, \mathcal{T}_Y)$ -continuous. Then for each $K \subset X$, if K is compact in $(X, \mathcal{T}_X)$, then $f(K)$ is compact in $(Y, \mathcal{T}_Y)$.

Proof. Let $K \subset X$. Suppose that K is compact in $(X, \mathcal{T}_X)$. Let $\mathcal{V} \subset \mathcal{P}(Y)$. Suppose that $\mathcal{V}$ is an open cover of $f(K)$ in $(Y, \mathcal{T}_Y)$. By definition, $\mathcal{V} \subset \mathcal{T}_Y$ and $f(K) \subset \bigcup_{V \in \mathcal{V}} V$. Define $\mathcal{U} \subset \mathcal{P}(X)$ by $\mathcal{U} = \{f^{-1}(V) : V \in \mathcal{V}\}$. Since f is continuous and $\mathcal{V} \subset \mathcal{T}_Y$, $\mathcal{U} \subset \mathcal{T}_X$ and

by construction

$$\begin{aligned}
 K &\subset f^{-1}(f(K)) \\
 &\subset f^{-1}\left(\bigcup_{V \in \mathcal{V}} V\right) \\
 &= \bigcup_{V \in \mathcal{V}} f^{-1}(V) \\
 &= \bigcup_{U \in \mathcal{U}} U
 \end{aligned}$$

Hence $\mathcal{U}$ is an open cover of K in $(X, \mathcal{T})$. Since K is compact, there exists $\mathcal{U}_0 \subset \mathcal{U}$ such that $\mathcal{U}_0$ is an open cover of K in $(X, \mathcal{T}_X)$ and $\mathcal{U}_0$ is finite. For $U \in \mathcal{U}_0$, set $A_U = \{V \in \mathcal{V} : f^{-1}(V) = U\}$. By construction, for each $U \in \mathcal{U}_0$, $A_U \neq \emptyset$. Since $\mathcal{U}_0$ is finite, $\prod_{U \in \mathcal{U}_0} A_U \neq \emptyset$. Thus there exists $V \in \prod_{U \in \mathcal{U}_0} A_U$. Set $\mathcal{V}_0 = \{V_U : U \in \mathcal{U}_0\}$. By construction $\mathcal{V}_0 \subset \mathcal{V}$ and for each $U \in \mathcal{U}_0$, $f^{-1}(V_U) = U$. Therefore

$$\begin{aligned}
 f(K) &\subset f\left(\bigcup_{U \in \mathcal{U}_0} U\right) \\
 &= \bigcup_{U \in \mathcal{U}_0} f(U) \\
 &= \bigcup_{U \in \mathcal{U}_0} f(f^{-1}(V_U)) \\
 &= \bigcup_{U \in \mathcal{U}_0} V_U \cap f(X) \\
 &\subset \bigcup_{U \in \mathcal{U}_0} V_U \\
 &= \bigcup_{V \in \mathcal{V}_0} V
 \end{aligned}$$

Hence $\mathcal{V}_0$ is an open cover of $f(K)$. Since $\mathcal{V} \subset \mathcal{P}(X)$ with $\mathcal{V}$ an open cover of $f(K)$ is arbitrary, we have that $f(K)$ is compact. $\square$

Definition 4.14.7.2. Let $(X, \mathcal{T}_X), (Y, \mathcal{T}_Y)$ be topological spaces and $f : X \rightarrow Y$. Then f is said to be **proper** if for each $K \subset X$, K is compact in $(Y, \mathcal{T}_Y)$ implies that $f^{-1}(K)$ is compact in $(X, \mathcal{T})$.

Exercise 4.14.7.3. Closed Map Lemma:

Let $(X, \mathcal{T}_X), (Y, \mathcal{T}_Y)$ be topological spaces and $f \in \text{Hom}_{\mathbf{Top}}((X, \mathcal{T}_X), (Y, \mathcal{T}_Y))$. Suppose that $(X, \mathcal{T}_X)$ is compact, $(Y, \mathcal{T}_Y)$ is Hausdorff. Then

1. f is closed,
2. f is proper,
3. f is surjective implies that f is a quotient map,
4. f is injective implies that f is a **Top**-embedding,
5. f is bijective implies that $f \in \text{Iso}_{\mathbf{Top}}((X, \mathcal{T}_X), (Y, \mathcal{T}_Y))$.

Proof.

1. Let $C \subset X$. Suppose that C is $\mathcal{T}_X$ -closed. Since $(X, \mathcal{T}_X)$ is compact, Exercise 4.14.1.9 implies that $C \in \mathcal{K}(X, \mathcal{T}_X)$. Since f is $(\mathcal{T}_X, \mathcal{T}_Y)$ -continuous and $C \in \mathcal{K}(X, \mathcal{T}_X)$, Exercise 4.14.7.1 implies that $f(C) \in \mathcal{K}(Y, \mathcal{T}_Y)$. Since $(Y, \mathcal{T}_Y)$ is Hausdorff, Exercise 4.14.1.8 implies that $f(C)$ is $\mathcal{T}_Y$ -closed. Since $C \subset X$ with C closed in $(X, \mathcal{T}_X)$ is arbitrary, we have that for each $C \subset X$, C is closed in $(X, \mathcal{T}_X)$ implies that $f(C)$ is closed in $(Y, \mathcal{T}_Y)$.

2. Let $K \in \mathcal{K}(Y, \mathcal{T}_Y)$ and $\mathcal{U} \in \text{Cover}^{\text{opn}}(f^{-1}(K), \mathcal{T}_X)$. Since $(Y, \mathcal{T}_Y)$ is Hausdorff, Exercise 4.14.1.8 implies that K is $\mathcal{T}_Y$ -closed. Exercise 4.2.0.3 then implies that $f^{-1}(K)$ is $\mathcal{T}_X$ -closed. Since $(X, \mathcal{T}_X)$ is compact, Exercise 4.14.1.9 implies that $f^{-1}(K) \in \mathcal{K}(X, \mathcal{T})$ is compact in $(X, \mathcal{T})$. Since $f \in \text{Hom}_{\mathbf{Top}}((X, \mathcal{T}_X), (Y, \mathcal{T}_Y))$, Exercise 4.11.1.3 implies that $f^*(\mathcal{V}) \in \text{Cover}^{\text{opn}}(f^{-1}(K))$. Define $\mathcal{U} \in \mathcal{T}_X$ by $\mathcal{U} := f^*(\mathcal{V})$.
3. Suppose that f is surjective. Exercise 4.8.1.8 implies that f is a quotient map.
4. Suppose that f is injective. Exercise 4.5.3.2 implies that f is a **Top**-embedding.
5. Clear by previous parts.

□

Exercise 4.14.7.4. Let $(X, \mathcal{T})$ be a topological space, $(Y_\alpha, \mathcal{T}_\alpha)_{\alpha \in A}$ a collection of topological spaces, $\mathcal{F} \in \prod_{\alpha \in A} Y_\alpha^X$ and $E \in \mathcal{K}(X)$. Suppose that for each $\alpha \in A$, f is $(\mathcal{T}, \mathcal{T}_\alpha)$ -continuous. If $(X, \tau_X(\mathcal{F}))$ is Hausdorff, then

1. $\tau_X(\mathcal{F}) \cap E = \mathcal{T} \cap E$,
2. for each net $(x_\alpha)_{\alpha \in \Gamma} \subset E$ and $x \in E$, $x_\alpha \rightarrow x$ in $(X, \mathcal{T})$ iff $x_\alpha \rightarrow x$ in $(X, \tau_X(\mathcal{F}))$.

Proof. Suppose that $(X, \tau_X(\mathcal{F}))$ is Hausdorff.

1. Define $F : X \rightarrow X$ by $F(x) := x$. Then F is $(\mathcal{T}, \mathcal{T})$ -continuous. Since for each $\alpha \in A$, f is $(\mathcal{T}, \mathcal{T}_\alpha)$ -continuous, Exercise 4.2.0.12 implies that $\tau_X(\mathcal{F}) \subset \mathcal{T}$. Thus F is $(\mathcal{T}, \tau_X(\mathcal{F}))$ -continuous. Define $f : E \rightarrow E$ by $f := T|_E^E$. Exercise 4.5.1.12 and Exercise 4.5.1.11 imply that f is $(\mathcal{T} \cap E, \tau_X(\mathcal{F}) \cap E)$ -continuous. Since $(E, \mathcal{T} \cap E)$ is compact, $(E, \tau_X(\mathcal{F}))$ is Hausdorff, f is $(\mathcal{T} \cap E, \tau_X(\mathcal{F}) \cap E)$ -continuous and f is a bijection, Exercise 4.14.7.3 implies that f is a homeomorphism. Therefore $\tau_X(\mathcal{F}) \cap E = \mathcal{T} \cap E$.
2. Let $(x_\alpha)_{\alpha \in \Gamma} \subset E$ be a net and $x \in E$.

- ($\implies$):
Suppose that $x_\alpha \rightarrow x$ in $(X, \mathcal{T})$. Since F is $(\mathcal{T}, \tau_X(\mathcal{F}))$ -continuous, we have that

$$\begin{aligned} x_\alpha &= F(x_\alpha) \\ &\xrightarrow{\tau_X(\mathcal{F})} F(x) \\ &= x. \end{aligned}$$

- ($\impliedby$):
Suppose that $x_\alpha \rightarrow x$ in $(X, \tau_X(\mathcal{F}))$. Exercise 4.5.1.10 implies that $x_\alpha \rightarrow x$ in $(E, \tau_X(\mathcal{F}) \cap E)$. **The prev part** implies that $x_\alpha \rightarrow x$ in $(E, \mathcal{T} \cap E)$. Exercise 4.5.1.10 again implies that $x_\alpha \rightarrow x$ in $(X, \mathcal{T})$.

□

4.14.8 Sequential Compactness

Definition 4.14.8.1. Let $(X, \mathcal{T})$ be a topological space. Then $(X, \mathcal{T})$ is said to be **sequentially compact** if for each $(x_n)_{n \in \mathbb{N}} \subset X$, there exists $(x_{n_k})_{k \in \mathbb{N}} \subset (x_n)_{n \in \mathbb{N}}$ and $x \in X$ such that $x_{n_k} \rightarrow x$.

Exercise 4.14.8.2. Let $(X, \mathcal{T})$ be a topological space. Suppose that $(X, \mathcal{T})$ is first-countable and $(X, \mathcal{T})$ is not sequentially compact. Then

1. there exists $(x_n)_{n \in \mathbb{N}} \subset X$ such that for each $x \in X$, there exists $U_x \in \mathcal{T}$ and $n_x \in \mathbb{N}$ such that $x \in U_x$ and for each $n \in \mathbb{N}$, $n \geq n_x$ implies that $x_n \in U_x^c$,
Hint: Exercise 4.13.1.7
2. $(X, \mathcal{T})$ is not compact

Proof.

1. Since $(X, \mathcal{T})$ is not sequentially compact, there exists $(x_n)_{n \in \mathbb{N}} \subset X$ such that for each $(x_{n_k})_{k \in \mathbb{N}} \subset (x_n)_{n \in \mathbb{N}}$ and $x \in X$, $x_{n_k} \not\rightarrow x$. Exercise 4.13.1.7 implies that for each $x \in X$, x is not a cluster point of $(x_n)_{n \in \mathbb{N}}$. Thus for each $x \in X$, there exists $U_x \in \mathcal{T}$ and $n_x \in \mathbb{N}$ such that $x \in U_x$ and for each $n \in \mathbb{N}$, $n \geq n_x$ implies that $x_n \in U_x^c$.
2. For the sake of contradiction suppose that $(X, \mathcal{T})$ is compact. Define $\mathcal{U} \subset \mathcal{T}$ by $\mathcal{U} := \{U_x : x \in X\}$. Then $\mathcal{U}$ is an open cover of X in $(X, \mathcal{T})$. Since $(X, \mathcal{T})$ is compact, there exists $J \in \mathbb{N}$ and $(a_j)_{j \in [J]} \subset X$ such that $(U_{a_j})_{j \in [J]}$ is an open cover of X in $(X, \mathcal{T})$. Set $n_0 := \max(n_{a_j} : j \in [J])$. Let $n \in \mathbb{N}$. Suppose that $n \geq n_0$. Then for each $j \in [J]$,

$$\begin{aligned} n &\geq n_0 \\ &\geq n_{a_j}. \end{aligned}$$

Thus for each $j \in [J]$, $x_n \in U_{a_j}^c$. Hence

$$\begin{aligned} x_n &\in \bigcap_{j \in [J]} U_{a_j}^c \\ &= \left(\bigcup_{j \in [J]} U_{a_j} \right)^c \\ &= X^c \\ &= \emptyset. \end{aligned}$$

This is a contradiction. Therefore $(X, \mathcal{T})$ is not compact. □

Exercise 4.14.8.3. Let $(X, \mathcal{T})$ be a topological space and $(x_n)_{n \in \mathbb{N}} \subset X$. Suppose that $(X, \mathcal{T})$ is first-countable. If $(X, \mathcal{T})$ is compact, then $(X, \mathcal{T})$ is sequentially compact.

Proof. Clear by the previous exercise. (add detail?) □

4.15 Locally Compact Hausdorff Spaces

Maybe move this section towards the end? I think we need to define the sup-norm first and metric results first, so it should be a 'later' topic, like compactifications and zero-dimensional spaces, in fact at this point, each section that is not linear, should have all prereqs listed and where to find them

4.15.1 Introduction

Definition 4.15.1.1. Let X be a topological space. Then

- X is said to be **locally compact** if for each $x \in X$, there exists $K \in \mathcal{N}(x)$ such that K is compact
- X is said to be a **locally compact Hausdorff (LCH) space** if X is locally compact and X is Hausdorff.

Exercise 4.15.1.2. Let X be a LCH space and $U \subset X$. Suppose that U is open. Then for each $x \in U$, there exists $K \in \mathcal{N}(x)$ such that $K \subset U$ and K is compact.

Proof. Let $x \in U$. Since X is locally compact, there exists $K_0 \in \mathcal{N}(x)$ such that K_0 is compact. Set $U_0 = (\text{Int } K_0) \cap U$. Since $\text{cl } U_0 \subset K_0$, $\text{cl } U_0$ is closed and K_0 is compact, we have that $\text{cl } U_0$ is compact. Since U_0 is open and $x \in U_0$, [an exercise in the section on compactness](#) implies that there exists $K \in \mathcal{N}(x)$ such that $K \subset U$ and K is compact. $\square$

Exercise 4.15.1.3. Let X be a LCH space and $U \subset X$ and $K \subset U$. If U is open and K is compact, then there exists $V \subset X$ such that V is open, $K \subset V$, $\text{cl } V \subset U$ and V is precompact.

Proof. Suppose that U is open and K is compact. [The previous exercise](#) implies that for each $x \in K$, there exist $N \in \mathcal{N}(x)$ such that $N \subset U$ and N is compact. The axiom of choice implies that there exists $(N_x)_{x \in K} \subset \mathcal{P}(X)$ such that for each $x \in K$, $N_x \in \mathcal{N}(x)$, $N_x \subset U$ and N_x is compact. Then $(\text{Int } N_x)_{x \in K}$ is an open cover of K . Since K is compact, there exist $x_1, \dots, x_n \in K$ such that $(\text{Int } N_{x_j})_{j=1}^n$ is an open cover of K . Set $V = \bigcup_{j=1}^n \text{Int } N_{x_j}$. Then V is open and since $(\text{Int } N_{x_j})_{j=1}^n$ is an open cover of K , we have that

$$\begin{aligned} K &\subset \bigcup_{j=1}^n \text{Int } N_{x_j} \\ &= V \end{aligned}$$

By construction, $\text{cl } V = \bigcap_{j=1}^n N_{x_j}$ which is compact, so V is precompact. Finally

$$\begin{aligned} \text{cl } V &= \bigcap_{j=1}^n N_{x_j} \\ &\subset U \end{aligned}$$

$\square$

Exercise 4.15.1.4. Urysohn's Lemma for LCH Spaces:

change notation from $C_c(X)$ to $f : X \rightarrow [0, 1]$ such that blah blah Let X be a LCH space, $U \subset X$ and $K \subset U$. If U is open and K is compact, then there exists $f \in C_c(X, [0, 1])$ such that $f|_K = 1$ and $\text{supp } f \subset U$.

Proof. Suppose that U is open and K is compact. [The previous exercise](#) implies that there exists $V \subset X$ such that V is open, $K \subset V$, $\text{cl } V \subset U$ and V is precompact. **FINISH!!!** $\square$

Exercise 4.15.1.5. Let X, Y be topological spaces and $f : X \rightarrow Y$. Suppose that Y is a LCH space and f is continuous. If f is proper, then f is closed.

Hint: Let $(y_\alpha)_{\alpha \in A} \subset f(C)$ be a net and $y \in Y$. Suppose that $y_\alpha \rightarrow y$ and consider $K_y \in \mathcal{N}(y)$, then $f(C \cap f^{-1}(K_y)) = f(C) \cap K_y$.

Proof. Suppose that f is proper. Let $C \subset X$. Suppose that C is closed in X . Let $(y_\alpha)_{\alpha \in A} \subset f(C)$ be a net and $y \in Y$. Suppose that $y_\alpha \rightarrow y$.

Since Y is LCH, there exists $K_y \in \mathcal{N}(y)$ such that K_y is compact. Since Y is Hausdorff, Exercise 4.14.1.8 implies that K_y is closed. Since f is continuous, $f^{-1}(K_y)$ is closed. Hence $C \cap f^{-1}(K_y)$ is closed. Since f is proper, $f^{-1}(K_y)$ is compact. Exercise 4.14.1.9 implies that $C \cap f^{-1}(K_y)$ is compact. Since f is continuous, $f(C \cap f^{-1}(K_y))$ is compact. Exercise 1.3.1.5 implies that $f(C \cap f^{-1}(K_y)) = f(C) \cap K_y$. Since $f(C) \cap K_y$ is compact and Y is Hausdorff, Exercise 4.14.1.8 implies that $f(C) \cap K_y$ is closed in Y .

Since $K_y \in \mathcal{N}(y)$ and $y_\alpha \rightarrow y$, there exists $\alpha_0 \in A$ such that for each $\alpha \in A$, $\alpha \geq \alpha_0$ implies that $y_\alpha \in K_y$. Hence for each $\alpha \in A$, $\alpha \geq \alpha_0$ implies that $y_\alpha \in f(C) \cap K_y$. Since $f(C) \cap K_y$ is closed in Y , we have that

$$\begin{aligned} y &\in f(C) \cap K_y \\ &\subset f(C). \end{aligned}$$

Since $(y_\alpha)_{\alpha \in A} \subset f(C)$ a net and $y \in Y$ such that $y_\alpha \rightarrow y$ are arbitrary, we have that for each net $(y_\alpha)_{\alpha \in A} \subset f(C)$ and $y \in Y$, $y_\alpha \rightarrow y$ implies that $y \in f(C)$. Hence $f(C)$ is closed in Y . Since $C \subset X$ such that C is closed in X is arbitrary, we have that for each $C \subset X$, C is closed in X implies that $f(C)$ is closed in Y . Therefore f is closed. $\square$

4.15.2 Arzela-Ascoli Theorem

4.15.3 Stone-Weierstrass Theorem

Definition 4.15.3.1. Let X be a compact Hausdorff space and $\mathcal{A} \subset C(X, \mathbb{C})$ a subalgebra. Then $\mathcal{A}$ is said to **separate the points of X** if for each $x, y \in X$, $x \neq y$ implies that there exists $f \in \mathcal{A}$ such that $f(x) \neq f(y)$.

Exercise 4.15.3.2. Equip $\mathbb{R}^2$ with pointwise addition and multiplication.

1. For each $\mathcal{A} \in \{\{(0,0)\}, \text{span}\{(1,0)\}, \text{span}\{(0,1)\}, \text{span}\{(1,1), \mathbb{R}^2\}\}$, $\mathcal{A}$ is a subalgebra of $\mathbb{R}^2$.
2. For each $\mathcal{A} \subset \mathbb{R}^2$, $\mathcal{A}$ is a subalgebra implies that $\mathcal{A} \in \{\{(0,0)\}, \text{span}\{(1,0)\}, \text{span}\{(0,1)\}, \text{span}\{(1,1), \mathbb{R}^2\}\}$.

Hint: If $(s, t) \in \mathcal{A}$, then $(s^2, t^2) \in \mathcal{A}$.

Proof.

1. Clear.
2. Since $\dim \mathbb{R}^2 = 2$, we have that $\dim \mathcal{A} \in \{0, 1, 2\}$. If $\dim \mathcal{A} = 0$, then $\mathcal{A} = \{(0,0)\}$. If $\dim \mathcal{A} = 2$, then $\mathcal{A} = \mathbb{R}^2$. Suppose that $\dim \mathcal{A} = 1$. Then there exists $(s, t) \in \mathcal{A}$ such that $\mathcal{A} = \text{span}\{(s, t)\}$. Thus $(s, t) \neq (0,0)$. Since $(s, t) \in \mathcal{A}$, $(s^2, t^2) \in \mathcal{A}$. Since $(s, t) \neq (0,0)$, $(s^2, t^2) \neq (0,0)$. Since $\mathcal{A} = \text{span}\{(s, t)\}$, there exists $\lambda \in \mathbb{R}$ such that $(s^2, t^2) = \lambda(s, t)$. Thus $s(s - \lambda) = 0$ and $t(t - \lambda) = 0$. Since $(s^2, t^2) \neq (0,0)$, $\lambda \neq 0$. Hence $(s, t) \in \{(\lambda, 0), (0, \lambda), (\lambda, \lambda)\}$. Therefore

$$\begin{aligned} \mathcal{A} &= \text{span}\{(s, t)\} \\ &\in \{\text{span}\{(1,0)\}, \text{span}\{(0,1)\}, \text{span}\{(1,1)\}\}. \end{aligned}$$

$\square$

Exercise 4.15.3.3. For each $\epsilon > 0$, there exists $p \in \mathbb{R}[x]$ such that $p(0) = 0$ and for each $x \in [-1, 1]$, $\left| |x| - p(x) \right| < \epsilon$.

Hint: Define $f : (-1, 1) \rightarrow \mathbb{R}$ by $f(t) := (1 - t)^{1/2}$. Consider the Taylor series for f at $t = 0$.

Proof. **FINISH!!!**

$\square$

Exercise 4.15.3.4. Let X be a compact Hausdorff space and $\mathcal{A} \subset C(X, \mathbb{C})$ a closed subalgebra.

Exercise 4.15.3.5. Real Stone-Weierstrass Theorem Let X be a compact Hausdorff space and $\mathcal{A} \subset C(X, \mathbb{C})$ a closed subalgebra. Then $\mathcal{A} = C(X, \mathbb{R})$ or there exists $x_0 \in X$ such that $\mathcal{A} = \{f \in C(X, \mathbb{R}) : f(x_0) = 0\}$.

Proof. content...

$\square$

FINISH!!!

4.16 Tychonoff Spaces

4.17 Compactification

Definition 4.17.0.1. Let X and Y be topological spaces and $\phi : X \rightarrow Y$. Then (Y, ϕ) is said to be a **compactification of X** if

1. Y is compact
2. $\phi(X)$ is dense in Y
3. $\phi|_{\phi(X)} : X \rightarrow \phi(X)$ is a homeomorphism

Definition 4.17.0.2. Let $X, X' \in \text{Obj}(\mathbf{Top})$ and $\iota \in \text{Hom}_{\mathbf{Top}}(X, Y)$. Then (X', ι) is said to be a **Stone-Čech compactification of X** if

1. (X', ι) is a compactification of X
2. for each compactification (Y, ϕ) of X , there exists a unique $\phi' \in \text{Hom}_{\mathbf{Top}}(X', Y)$ such that $\phi' \circ \iota = \phi$, i.e. the following diagram commutes:

$$\begin{array}{ccc} X & \xrightarrow{\iota} & X' \\ & \searrow \phi & \downarrow \phi' \\ & & Y \end{array}$$

Exercise 4.17.0.3. show that a Stone-Čech compactification is just a colimit of the Diagram D given by $D(*) := X$ in some category.

4.18 Zero-Dimensional Spaces

Definition 4.18.0.1. Let $(X, \mathcal{T})$ be a topological space. Then $(X, \mathcal{T})$ is said to be **zero-dimensional** if there exists $\mathcal{B} \subset \mathcal{T}$ such that $\mathcal{B}$ is a basis for $\mathcal{T}$ and for each $B \in \mathcal{B}$, B is closed.

Exercise 4.18.0.2. Let $(X_1, \mathcal{T}_1)$ and $(X_2, \mathcal{T}_2)$ be topological spaces and $\phi : X_1 \rightarrow X_2$. If ϕ is a homeomorphism, then $(X_1, \mathcal{T}_1)$ is zero-dimensional iff $(X_2, \mathcal{T}_2)$ is zero-dimensional.

Proof. **FINISH!!!** □

4.18.1 Zero-Dimensional Subspaces

Exercise 4.18.1.1. Let $(X, \mathcal{T})$ be a topological space and $A \subset X$. If $(X, \mathcal{T})$ is a zero-dimensional space, then $(A, \mathcal{T} \cap A)$ is zero-dimensional.

Proof. Suppose that $(X, \mathcal{T})$ is zero-dimensional. Then there exists $\mathcal{B}_0 \subset \mathcal{T}$ such that $\mathcal{B}_0$ is a basis for $\mathcal{T}$ and for each $B_0 \in \mathcal{B}_0$, B_0 is open and closed. Exercise 4.5.1.15 implies that $\mathcal{B}_0 \cap A$ is a basis for $\mathcal{T} \cap A$. Let $B \in \mathcal{B} \cap A$. By definition, there exists $B_0 \in \mathcal{B}_0$ such that $B = B_0 \cap A$. Since B_0 is open and $\mathcal{T}$ -closed, Exercise 4.5.1.13 implies that B is closed in $(A, \mathcal{T} \cap A)$. Since $B \in \mathcal{B}$ is arbitrary, we have that for each $B \in \mathcal{B}$, B is closed. Thus there exists $\mathcal{B} \subset \mathcal{T} \cap A$ such that $\mathcal{B}$ is a basis for $\mathcal{T} \cap A$ and for each $B \in \mathcal{B}$, B is closed. Hence $(A, \mathcal{T} \cap A)$ is zero-dimensional. □

4.18.2 Zero-Dimensional Product Spaces

(change $n \in \mathbb{N}$ to $\alpha \in A$ or something, doesn't need to be countable.)

Exercise 4.18.2.1. Let $(X_n, \mathcal{T}_n)_{n \in \mathbb{N}}$ be a collection of topological spaces. If for each $n \in \mathbb{N}$, $(X_n, \mathcal{T}_n)$ is zero-dimensional, then $(\prod_{n \in \mathbb{N}} X_n, \bigotimes_{n \in \mathbb{N}} \mathcal{T}_n)$ is zero-dimensional.

Proof. Suppose that for each $n \in \mathbb{N}$, $(X_n, \mathcal{T}_n)$ is zero-dimensional. Then for each $n \in \mathbb{N}$, there exists $\mathcal{B}_n \subset \mathcal{T}_n$ such that $\mathcal{B}_n$ is a basis for $\mathcal{T}_n$ and for each $n \in \mathbb{N}$, $B \in \mathcal{B}_n$, B is open and closed in $(X_n, \mathcal{T}_n)$. Set

$$\mathcal{B} := \left\{ \prod_{n \in \mathbb{N}} B_n : \text{there exists } J \subset \mathbb{N} \text{ such that } \#J < \infty, \right. \\ \left. \text{for each } n \in J, B_n \in \mathcal{B}_n \text{ and for each } n \in J^c, B_n = X_n \right\}.$$

Exercise 4.6.1.6 implies that $\mathcal{B}$ is a basis for $\bigotimes_{n \in \mathbb{N}} \mathcal{T}_n$. Let $B \in \mathcal{B}$. By definition of $\mathcal{B}$, there exists $J \subset \mathbb{N}$ and $(B_n)_{n \in \mathbb{N}} \in \prod_{n \in \mathbb{N}} \mathcal{T}_n$ such that $\#J < \infty$, for each $n \in J$, $B_n \in \mathcal{B}_n$ and for each $n \in J^c$, $B_n = X_n$. Let $n \in \mathbb{N}$. If $n \in J$, then $B_n \in \mathcal{B}_n$ and therefore B_n is closed in $(X_n, \mathcal{T}_n)$. If $n \in J^c$, then $B_n = X_n$ and therefore B_n is closed in $(X_n, \mathcal{T}_n)$. Therefore for each $n \in \mathbb{N}$, B_n is closed in $(X_n, \mathcal{T}_n)$. Hence for each $n \in \mathbb{N}$, $\pi_n^{-1}(B_n)$ is closed in $(\prod_{n \in \mathbb{N}} X_n, \bigotimes_{n \in \mathbb{N}} \mathcal{T}_n)$. Since J is finite and

$$B = \prod_{n \in \mathbb{N}} B_n \\ = \bigcap_{n \in J} \pi_n^{-1}(B_n),$$

we have that B is closed in $(\prod_{n \in \mathbb{N}} X_n, \bigotimes_{n \in \mathbb{N}} \mathcal{T}_n)$. Since $B \in \mathcal{B}$ is arbitrary, we have that for each $B \in \mathcal{B}$, B is closed in $(\prod_{n \in \mathbb{N}} X_n, \bigotimes_{n \in \mathbb{N}} \mathcal{T}_n)$. Hence there exists $\mathcal{B} \subset \bigotimes_{n \in \mathbb{N}} \mathcal{T}_n$ such that $\mathcal{B}$ is a basis for $\bigotimes_{n \in \mathbb{N}} \mathcal{T}_n$ and for each $B \in \mathcal{B}$, B is closed in $(\prod_{n \in \mathbb{N}} X_n, \bigotimes_{n \in \mathbb{N}} \mathcal{T}_n)$. Thus $(\prod_{n \in \mathbb{N}} X_n, \bigotimes_{n \in \mathbb{N}} \mathcal{T}_n)$ is zero-dimensional. □

4.18.3 Zero-Dimensional Coproduct Spaces

(change $n \in \mathbb{N}$ to $\alpha \in A$ or something, doesn't need to be countable.)

Exercise 4.18.3.1. Let $(X_n, \mathcal{T}_n)_{n \in \mathbb{N}}$ be a collection of topological spaces. If for each $n \in \mathbb{N}$, $(X_n, \mathcal{T}_n)$ is zero-dimensional, then $(\coprod_{n \in \mathbb{N}} X_n, \bigoplus_{n \in \mathbb{N}} \mathcal{T}_n)$ is zero-dimensional.

Proof. Suppose that for each $n \in \mathbb{N}$, $(X_n, \mathcal{T}_n)$ is zero-dimensional. Then for each $n \in \mathbb{N}$, there exists $\mathcal{B}_n \subset \mathcal{T}_n$ such that $\mathcal{B}_n$ is a basis for $\mathcal{T}_n$ and for each $B \in \mathcal{B}_n$, B is closed. Define $\mathcal{B} \subset \bigoplus_{n \in \mathbb{N}} \mathcal{T}_n$ by $\mathcal{B} := \{\iota_n(U) : n \in \mathbb{N} \text{ and } U \in \mathcal{B}_n\}$. Exercise 4.7.0.6 implies that $\mathcal{B}$ is a basis for $\bigoplus_{n \in \mathbb{N}} \mathcal{T}_n$. Exercise 4.7.0.5 implies that for each $n \in \mathbb{N}$, ι_n is closed. Hence for each $B \in \mathcal{B}$, B is closed. Thus $(\coprod_{n \in \mathbb{N}} X_n, \bigoplus_{n \in \mathbb{N}} \mathcal{T}_n)$ is zero-dimensional. $\square$

Exercise 4.18.3.2. Let $(X, \mathcal{T})$ be a topological space and $(A_n)_{n \in \mathbb{N}} \subset \mathcal{P}(X)$. Suppose that $(A_n)_{n \in \mathbb{N}}$ is disjoint. If $(X, \mathcal{T})$ is zero dimensional, then $(\bigcup_{n \in \mathbb{N}} A_n, \mathcal{T} \cap [\bigcup_{n \in \mathbb{N}} A_n])$ is zero-dimensional.

Proof. Suppose that Since for each $n \in \mathbb{N}$, $A_n \subset X$ and $(X, \mathcal{T})$ is zero-dimensional, Exercise 4.18.1.1 implies that for each $n \in \mathbb{N}$, $(A_n, \mathcal{T} \cap A_n)$ is zero-dimensional. Exercise 4.18.3.1 then implies that $(\coprod_{n \in \mathbb{N}} A_n, \bigoplus_{n \in \mathbb{N}} (\mathcal{T} \cap A_n))$ is zero-dimensional. Define $\phi : \coprod_{\alpha \in A} E_\alpha \rightarrow \bigcup_{\alpha \in A} E_\alpha$ by $\phi(\alpha, x) := x$. Since $(A_n)_{n \in \mathbb{N}}$ is disjoint, Exercise 4.7.0.11 implies that ϕ is a $\left(\bigoplus_{n \in \mathbb{N}} [\mathcal{T} \cap A_n], \mathcal{T} \cap \left[\bigcup_{n \in \mathbb{N}} A_n \right] \right)$ -homeomorphism. Exercise 4.18.0.2 then implies that $(\bigcup_{n \in \mathbb{N}} A_n, \mathcal{T} \cap [\bigcup_{n \in \mathbb{N}} A_n])$ is zero-dimensional. $\square$

4.19 Semi-continuity

Definition 4.19.0.1. Let X be a topological space and $f : X \rightarrow [-\infty, \infty]$.

- Let $x_0 \in X$. Then f is said to be **lower semicontinuous at x_0** if

$$\liminf_{\substack{x \rightarrow x_0 \\ x \neq x_0}} f(x) \geq f(x_0)$$

- Then f is said to be **lower semicontinuous** if for each $x_0 \in X$, f is lower semicontinuous at x_0 .

Exercise 4.19.0.2. Let X be a topological space and $f : X \rightarrow (\infty, \infty]$. Then f is lower semicontinuous iff for each $\alpha \in \mathbb{R}$, $f^{-1}((\alpha, \infty])$ is open.

Proof.

- $(\implies)$:

Suppose that f is lower semicontinuous. Let $\alpha \in \mathbb{R}$ and $x_0 \in f^{-1}((\alpha, \infty])$.

- Suppose that $f(x_0) = \infty$. Since f is lower semicontinuous,

$$\begin{aligned} \sup_{V \in \mathcal{N}(x_0)} \left[\inf_{x \in V \setminus \{x_0\}} f(x) \right] &= \liminf_{\substack{x \rightarrow x_0 \\ x \neq x_0}} f(x) \\ &\geq f(x_0) \\ &= \infty \end{aligned}$$

Thus there exists $V_\alpha \in \mathcal{N}(x_0)$ such that

$$\begin{aligned} \inf_{x \in V_\alpha \setminus \{x_0\}} f(x) &\geq \alpha + 1 \\ &> \alpha \end{aligned}$$

Hence for each $x \in V_\alpha \setminus \{x_0\}$,

$$\begin{aligned} f(x) &\geq \inf_{t \in V_\alpha \setminus \{x_0\}} f(t) \\ &> \alpha \end{aligned}$$

Since $f(x_0) = \infty$, $f(x_0) > \alpha$. Hence

$$\begin{aligned} \text{Int } V_\alpha &\subset V_\alpha \\ &\subset f^{-1}((\alpha, \infty]) \end{aligned}$$

Thus there exists $V \in \mathcal{N}(x_0)$ such that V is open and $V \subset f^{-1}((\alpha, \infty])$.

- Suppose that $f(x_0) < \infty$. Set $\epsilon := f(x_0) - \alpha$. By definition,

$$\begin{aligned} \sup_{V \in \mathcal{N}(x_0)} \left[\inf_{x \in V \setminus \{x_0\}} f(x) \right] &\geq f(x_0) \\ &> f(x_0) - \epsilon \end{aligned}$$

Choose $V_\epsilon \in \mathcal{N}(x_0)$ such that

$$\begin{aligned} \inf_{x \in V_\epsilon \setminus \{x_0\}} f(x) &> f(x_0) - \epsilon \\ &= \alpha \end{aligned}$$

Then

$$\begin{aligned} \text{Int } V_\epsilon &\subset V_\epsilon \\ &\subset f^{-1}((\alpha, \infty]) \end{aligned}$$

Thus there exists $V \in \mathcal{N}(x_0)$ such that V is open and $V \subset f^{-1}((\alpha, \infty])$.

Since $x_0 \in f^{-1}((\alpha, \infty])$ is arbitrary, we have that for each $x_0 \in f^{-1}((\alpha, \infty])$, there exists $V \subset f^{-1}((\alpha, \infty])$ such that V is open and $x_0 \in V$. Hence $f^{-1}((\alpha, \infty])$ is open. Since $\alpha \in \mathbb{R}$ is arbitrary, we have that for each $\alpha \in \mathbb{R}$, $f^{-1}((\alpha, \infty])$ is open.

• ($\Leftarrow$) :

Suppose that for each $\alpha \in \mathbb{R}$, $f^{-1}((\alpha, \infty])$ is open. Let $x_0 \in X$.

– Suppose that $f(x_0) = \infty$. For $M > 0$, define $V_M = f^{-1}((M, \infty])$. Then for each $M > 0$, $V_M \in \mathcal{N}(x_0)$ and

$$\begin{aligned} \liminf_{\substack{x \rightarrow x_0 \\ x \neq x_0}} f(x) &= \sup_{V \in \mathcal{N}(x_0)} \left[\inf_{x \in V \setminus \{x_0\}} f(x) \right] \\ &\geq \sup_{M > 0} \left[\inf_{x \in V_M \setminus \{x_0\}} f(x) \right] \\ &\geq \sup_{n \in \mathbb{N}} M \\ &= \infty \\ &= f(x_0) \end{aligned}$$

– Suppose that $f(x_0) < \infty$. Set $\alpha := f(x_0)$. For $n \in \mathbb{N}$, define $V_n = f^{-1}((f(x_0) - 1/n, \infty])$. Then for each $n \in \mathbb{N}$, $V_n \in \mathcal{N}(x_0)$ and

$$\begin{aligned} \liminf_{\substack{x \rightarrow x_0 \\ x \neq x_0}} f(x) &= \sup_{V \in \mathcal{N}(x_0)} \left[\inf_{x \in V \setminus \{x_0\}} f(x) \right] \\ &\geq \sup_{n \in \mathbb{N}} \left[\inf_{x \in V_n \setminus \{x_0\}} f(x) \right] \\ &\geq \sup_{n \in \mathbb{N}} f(x_0) - 1/n \\ &= f(x_0) \end{aligned}$$

Hence So f is lower semicontinuous at x_0 . Since $x_0 \in X$ is arbitrary, we have that for each $x_0 \in X$, f is lower semicontinuous at x_0 . Therefore f is lower semicontinuous. □

Definition 4.19.0.3. Let X be a topological space and $f : X \rightarrow \mathbb{R}$. We define the **epigraph of f** , denoted $\text{epi } f$, by

$$\text{epi } f = \{(x, y) \in X \times \mathbb{R} : f(x) \leq y\}$$

Exercise 4.19.0.4. Let X be a topological space and $f : X \rightarrow \mathbb{R}$. Then f is lower semicontinuous iff $\text{epi } f$ is closed.

Proof. Suppose that f is lower semicontinuous. Let $(x_\alpha, y_\alpha)_{\alpha \in A} \subset \text{epi } f$ be a net and $(x, y) \in X \times \mathbb{R}$. Then for each $\alpha \in A$, $f(x_\alpha) \leq y_\alpha$. Suppose that $(x_\alpha, y_\alpha) \rightarrow (x, y)$. Then $x_\alpha \rightarrow x$ and $y_\alpha \rightarrow y$. Therefore

$$\begin{aligned} f(x) &\leq \liminf_{\substack{t \rightarrow x \\ t \neq x}} f(t) \\ &\leq \liminf_{\alpha \in A} f(x_\alpha) \\ &\leq \liminf_{\alpha \in A} y_\alpha \\ &= y \end{aligned}$$

So $(x, y) \in \text{epi } f$ and $\text{epi } f$ is closed.

Conversely, suppose that $\text{epi } f$ is closed. □

Exercise 4.19.0.5. Let X be a topological space and $(f_\lambda)_{\lambda \in \Lambda} \subset (-\infty, \infty]^X$. Suppose that for each $\lambda \in \Lambda$, f_λ is lower semicontinuous. Set $f = \sup_{\lambda \in \Lambda} f_\lambda$. Then f is lower semicontinuous.

Proof. Let $\alpha \in \mathbb{R}$ and $x \in X$. Then

$$\begin{aligned}
 x \in f^{-1}((\alpha, \infty]) &\iff \sup_{\lambda \in \Lambda} f_{\lambda}(x) > \alpha \\
 &\iff \text{there exists } \lambda \in \Lambda \text{ such that } f_{\lambda}(x) > \alpha \\
 &\iff \text{there exists } \lambda \in \Lambda \text{ such that } x \in f_{\lambda}^{-1}((\alpha, \infty]) \\
 &\iff x \in \bigcup_{\lambda \in \Lambda} f_{\lambda}^{-1}((\alpha, \infty])
 \end{aligned}$$

Since for each $\lambda \in \Lambda$, $f_{\lambda}^{-1}((\alpha, \infty])$ is open, $f^{-1}((\alpha, \infty]) = \bigcup_{\lambda \in \Lambda} f_{\lambda}^{-1}((\alpha, \infty])$ is open. So f is lower semicontinuous. $\square$

Chapter 5

Locale Theory

Note 5.0.0.1. This chapter assumes familiarity with lattice theory. See [algebra notes](#)

5.1 Sober Spaces

Exercise 5.1.0.1. Let $(X, \mathcal{T})$ be a topological space. Then

1. for each $\mathcal{U} \subset \mathcal{T}$,

$$(a) \sup(\mathcal{U}, \subset) = \bigcup_{U \in \mathcal{U}} U$$

$$(b) \inf(\mathcal{U}, \subset) = \text{Int} \bigcap_{U \in \mathcal{U}} U$$

2. $(\mathcal{T}, \subset)$ is a complete lattice

3. $(\mathcal{T}, \subset, \emptyset, X)$ is a bounded distributive lattice.

Proof. make and ref exercise as fundamental example in algebra lattice notes □

Exercise 5.1.0.2. Let $(X, \mathcal{T})$ be a topological space. Then for each $x \in X$, $(\text{cl}\{x\})^c \in \text{prime}_\wedge(\mathcal{T}, \subset)$.

Proof. Let $x \in X$. Let $U, V \in \mathcal{T}$. We note that

$$\begin{aligned} U \cap V \subset (\text{cl}\{x\})^c &\iff \text{cl}\{x\} \subset (U \cap V)^c \\ &\iff \text{cl}\{x\} \subset U^c \cup V^c \\ &\implies x \in U^c \cup V^c \\ &\iff x \in U^c \text{ or } x \in V^c \\ &\iff \{x\} \subset U^c \text{ or } \{x\} \subset V^c \\ &\iff \text{cl}\{x\} \subset U^c \text{ or } \text{cl}\{x\} \subset V^c \\ &\iff U \subset (\text{cl}\{x\})^c \text{ or } V \subset (\text{cl}\{x\})^c. \end{aligned}$$

Since $U, V \in \mathcal{T}$ are arbitrary, we have that for each $U, V \in \mathcal{T}$, $U \cap V \subset (\text{cl}\{x\})^c$ implies that $U \subset (\text{cl}\{x\})^c$ or $V \subset (\text{cl}\{x\})^c$. Thus $(\text{cl}\{x\})^c \in \text{prime}_\wedge(\mathcal{T}, \subset)$. □

Definition 5.1.0.3. move def of $\mathcal{N}^{\text{opn}}(X, \mathcal{T})$ to intro to topology section (actually name it the ‘open neighborhood filter’) because I think $\mathcal{U}(x)$ or $\mathcal{U}_x$ is used in later chapters, then change it there. Let $(X, \mathcal{T})$ be a topological space.

- We define $\text{Punct}(X, \mathcal{T}) \subset \mathcal{T}$ by $\text{Punct}(X, \mathcal{T}) := \{\text{cl}(\{x\}, \mathcal{T})^c : x \in X\}$.
- Let $S \subset X$. We define $\mathcal{N}^{\text{opn}}(S, \mathcal{T}) \subset \mathcal{T}$ by $\mathcal{N}^{\text{opn}}(S, \mathcal{T}) := \mathcal{N}(S, \mathcal{T}) \cap \mathcal{T}$

Note 5.1.0.4. When the context is clear, we write $\text{Punct}(X)$ and $\mathcal{N}^{\text{opn}}(S)$ in place of $\text{Punct}(X, \mathcal{T})$ and $\mathcal{N}^{\text{opn}}(S)$ respectively.

Definition 5.1.0.5. Let $(X, \mathcal{T})$ be a topological space. Then $(X, \mathcal{T})$ is said to be **sober** if $\text{prime}_\wedge(\mathcal{T}, \subset) = \text{Punct}(X)$.

Exercise 5.1.0.6. Let $(X, \mathcal{T})$ be a topological space. Then $\{\mathcal{N}^{\text{opn}}(x) : x \in X\} \subset \mathcal{F}_{cp}(\mathcal{T})$

Proof. Let $x \in X$.

- 1. Since $X \in \mathcal{N}^{\text{opn}}(x)$, $\mathcal{N}^{\text{opn}}(x) \neq \emptyset$.
- 2. Let $U, V \in \mathcal{N}^{\text{opn}}(x)$. Then $U, V \in \mathcal{T}$ and $x \in U, V$. Thus

$$\begin{aligned} U \wedge V &= U \cap V \\ &\in \mathcal{T} \end{aligned}$$

and

$$\begin{aligned} x &\in U \cap V \\ &= U \wedge V. \end{aligned}$$

Hence $U \wedge V \in \mathcal{N}^{\text{opn}}(x)$. Since $U, V \in \mathcal{N}^{\text{opn}}(x)$ are arbitrary, we have that for each $U, V \in \mathcal{N}^{\text{opn}}(x)$, $U \wedge V \in \mathcal{N}^{\text{opn}}(x)$.

- 3. Let $V \in \mathcal{T}$ and $U \in \mathcal{N}^{\text{opn}}(x)$. Then

$$\begin{aligned} x &\in U \\ &\subset V \cup U \\ &= V \vee U. \end{aligned}$$

Since $V, U \in \mathcal{T}$, we have that

$$\begin{aligned} V \vee U &= V \cup U \\ &\in \mathcal{T}. \end{aligned}$$

Hence $V \vee U \in \mathcal{N}^{\text{opn}}(x)$. Since $V \in \mathcal{T}$ and $U \in \mathcal{N}^{\text{opn}}(x)$ are arbitrary, we have that for each $V \in \mathcal{T}$ and $U \in \mathcal{N}^{\text{opn}}(x)$, $V \vee U \in \mathcal{N}^{\text{opn}}(x)$.

Thus $\mathcal{N}^{\text{opn}}(x) \in \mathcal{F}(\mathcal{T})$.

- Let $\mathcal{V} \subset \mathcal{T}$. Suppose that $\bigvee \mathcal{V} \in \mathcal{N}^{\text{opn}}(x)$. Since $\bigvee \mathcal{V} = \bigcup \mathcal{V}$ and $\mathcal{N}^{\text{opn}}(x) = \{U \in \mathcal{T} : x \in U\}$, we have that $x \in \bigcup \mathcal{V}$. Then there exists $V \in \mathcal{V}$ such that $x \in V$. Hence $V \in \mathcal{N}^{\text{opn}}(x)$. Since $\mathcal{V} \subset \mathcal{T}$ such that $\bigvee \mathcal{V} \in \mathcal{N}^{\text{opn}}(x)$ is arbitrary, we have that for each $\mathcal{V} \subset \mathcal{T}$, $\bigvee \mathcal{V} \in \mathcal{N}^{\text{opn}}(x)$ implies that there exists $V \in \mathcal{V}$ such that $V \in \mathcal{N}^{\text{opn}}(x)$. Therefore $\mathcal{N}^{\text{opn}}(x) \in \mathcal{F}_{cp}(\mathcal{T})$. Since $x \in X$ is arbitrary, we have that for each $x \in X$, $\mathcal{N}^{\text{opn}}(x) \in \mathcal{F}_{cp}(\mathcal{T})$. Thus $\{\mathcal{N}^{\text{opn}}(x) : x \in X\} \subset \mathcal{F}_{cp}(\mathcal{T})$

□

Exercise 5.1.0.7. Let $(X, \mathcal{T})$ be a topological space. Then $(X, \mathcal{T})$ is sober iff $\mathcal{F}_{cp}(\mathcal{T}) = \{\mathcal{N}^{\text{opn}}(x) : x \in X\}$.

Hint: Let $\mathcal{U} \in \mathcal{F}_{cp}(\mathcal{T})$. Then there exists $x \in X$ such that $\bigwedge \mathcal{U}^c = (\text{cl}\{x\})^c$ and for each $U \in \mathcal{T}$, $U \in \mathcal{U}$ iff $x \in U$.

Proof.

- ($\implies$):
Suppose that $(X, \mathcal{T})$ is sober. Let $\mathcal{U} \in \mathcal{F}_{cp}(\mathcal{T})$. **An ex in algebra notes on complete lattices** implies that $\bigvee \mathcal{U}^c \in \text{prime}_\wedge(\mathcal{T}, \subset)$ and $\bigvee \mathcal{U}^c \in \mathcal{U}^c$. Since $(X, \mathcal{T})$ is sober, there exists $x \in X$ such that $\bigvee \mathcal{U}^c = (\text{cl}\{x\})^c$. Since $\mathcal{U} \in \mathcal{F}(\mathcal{T})$, $\mathcal{U}$ is an upper set. Hence $\mathcal{U}^c$ is a lower set. Let $U \in \mathcal{T}$.

- If $U \in \mathcal{U}^c$, then $U \subset \bigwedge \mathcal{U}^c$.
- Suppose that $U \subset \bigwedge \mathcal{U}^c$. Since $\bigvee \mathcal{U}^c \in \mathcal{U}^c$ and $\mathcal{U}^c$ is a lower set, we have that $U \in \mathcal{U}^c$.

Hence $U \in \mathcal{U}^c$ iff $U \subset \bigwedge \mathcal{U}^c$. A previous ex in the intro section on topology implies that (make/ref ex) $U \subset (\text{cl}\{x\})^c$ iff $x \in U^c$. Then

$$\begin{aligned} U \notin \mathcal{U} &\iff U \in \mathcal{U}^c \\ &\iff U \subset \bigwedge \mathcal{U}^c \\ &\iff U \subset (\text{cl}\{x\})^c \\ &\iff x \in U^c \\ &\iff x \notin U. \end{aligned}$$

Hence

$$\begin{aligned} U \in \mathcal{U} &\iff x \in U \\ &\iff U \in \mathcal{N}^{\text{opn}}(x). \end{aligned}$$

Since $U \in \mathcal{T}$ is arbitrary, we have that for each $U \in \mathcal{T}$, $U \in \mathcal{U}$ iff $U \in \mathcal{N}^{\text{opn}}(x)$. Hence $\mathcal{U} = \mathcal{N}^{\text{opn}}(x)$. Thus

• ($\Leftarrow$):

Suppose that $(X, \mathcal{T})$ is not sober. Then there exists $U_0 \in \text{prime}_\wedge(\mathcal{T})$ such that for each $x \in X$, $U_0 \neq (\text{cl}\{x\})^c$. Since $U_0 \in \text{prime}_\wedge(\mathcal{T})$, an ex in algebra notes on completely prime lattices implies that $(\downarrow U_0)^c \in \mathcal{F}_{cp}(\mathcal{T})$. Since $U_0 \in \text{prime}_\wedge(\mathcal{T})$, an ex in algebra notes on completely prime lattices implies that $\bigvee (\downarrow U_0)^c \in \mathcal{F}_{cp}(\mathcal{T})$. For the sake of contradiction, suppose that there exists $x \in X$ such that $(\downarrow U_0)^c = \mathcal{N}^{\text{opn}}(x)$. an ex in the section on intro to nets implies that for each $U \in \mathcal{T}$,

$$\begin{aligned} U \not\subset U_0 &\iff U \in (\downarrow U_0)^c \\ &\iff U \in \mathcal{N}^{\text{opn}}(x) \\ &\iff x \in U \\ &\iff U \not\subset (\text{cl}\{x\})^c \quad (\text{Exercise 4.3.2.9}). \end{aligned}$$

Thus for each $U \in \mathcal{T}$, $U \not\subset U_0$ iff $U \not\subset (\text{cl}\{x\})^c$. Hence $U_0 = (\text{cl}\{x\})^c$, which is a contradiction. Thus for each $x \in X$, $(\downarrow U_0)^c \neq \mathcal{N}^{\text{opn}}(x)$. Since $(\downarrow U_0)^c \in \mathcal{F}_{cp}(\mathcal{T})$, we have that there exists $F \in \mathcal{F}_{cp}(\mathcal{T})$ such that $F \not\subset \{\mathcal{N}^{\text{opn}}(x) : x \in X\}$. Therefore $\mathcal{F}_{cp}(\mathcal{T}) \not\subset \{\mathcal{N}^{\text{opn}}(x) : x \in X\}$. Hence $\mathcal{F}_{cp}(\mathcal{T}) \neq \{\mathcal{N}^{\text{opn}}(x) : x \in X\}$. □

Exercise 5.1.0.8. Let L be a complete lattice.

5.2 Frames and Locales

5.2.1 Introduction

Definition 5.2.1.1. Let L be a complete lattice. Then L is said to be a

- **frame** if for each $A \subset L$ and $b \in L$,

$$\left(\bigvee_{a \in A} a \right) \wedge b = \bigvee_{a \in A} (a \wedge b).$$

maybe define and use the following notation:

$$\left(\bigvee A \right) \wedge b = \bigvee (A \wedge b).$$

- **coframe** if L^* is a frame.

Definition 5.2.1.2. Let $(L, \leq_L)$, $(M, \leq_M)$ be frames and $f : L \rightarrow M$. Then f is said to be a $(\leq_L, \leq_M)$ -frame homomorphism if

1. f is $(\leq_L, \leq_M)$ -arbitrary join preserving

2. f is $(\leq_L, \leq_M)$ -finite meet preserving

Definition 5.2.1.3. We define the **category of frames**, denoted **Frm**, by

- $\text{Obj}(\mathbf{Frm}) := \{(L, \leq) : (L, \leq) \text{ is a frame}\}$
- for $(L, \leq_L), (M, \leq_M) \in \text{Obj}(\mathbf{Frm})$,

$$\text{Hom}_{\mathbf{Frm}}((L, \leq_L), (M, \leq_M)) := \{f \in \text{Hom}_{\mathbf{Set}}(L, M) : f \text{ is a } (\leq_L, \leq_M)\text{-frame homomorphism}\}$$

- for $(L, \leq_L), (M, \leq_M), (K, \leq_K) \in \text{Obj}(\mathbf{Frm})$ and $f \in \text{Hom}_{\mathbf{Frm}}((L, \leq_L), (M, \leq_M))$, $g \in \text{Hom}_{\mathbf{Frm}}((M, \leq_M), (K, \leq_K))$,

$$g \circ_{\mathbf{Frm}} f := g \circ_{\mathbf{Set}} f$$

Exercise 5.2.1.4. **Frm** is a category

Note 5.2.1.5. We recall **ref ex** about maps f that preserve arb joins/meets having right/left adjoints f^*/f_* from algebra notes

Exercise 5.2.1.6. Define **Loc** by

- $\text{Obj}(\mathbf{Loc}) := \text{Obj}(\mathbf{Frm})$
- For each $L, M \in \text{Obj}(\mathbf{Loc})$,

$$\text{Hom}_{\mathbf{Loc}}(L, M) := \{f^* : f \in \text{Hom}_{\mathbf{Frm}}(M, L)\}$$

- For each $L, M, K \in \text{Obj}(\mathbf{Loc})$ and $f \in \text{Hom}_{\mathbf{Loc}}(L, M)$, $g \in \text{Hom}_{\mathbf{Loc}}(M, K)$,

$$g \circ_{\mathbf{Loc}} f := g \circ_{\mathbf{Set}} f$$

•

Exercise 5.2.1.7. We have that

1. $\mathbf{Loc} \in \text{Obj}(\mathbf{Cat})$
2. $\mathbf{Loc} \cong_{\mathbf{Cat}} \mathbf{Frm}^{\text{op}}$.

Proof.

1. **FINISH!!!**

2. • Define $F : \mathbf{Frm}^{\text{op}} \rightarrow \mathbf{Loc}$ by

- $F(L) := L$,
- $F(f) := f^*$.
- Let $L, M \in \text{Obj}(\mathbf{Frm}^{\text{op}})$ and $f \in \text{Hom}_{\mathbf{Frm}^{\text{op}}}(L, M)$. Then $f \in \text{Hom}_{\mathbf{Frm}}(M, L)$. Hence

$$\begin{aligned} F(f) &= f^* \\ &\in \text{Hom}_{\mathbf{Loc}}(L, M) \\ &= \text{Hom}_{\mathbf{Loc}}(F(L), F(M)). \end{aligned}$$

- Let $L, M, N \in \text{Obj}(\mathbf{Frm}^{\text{op}})$, $f \in \text{Hom}_{\mathbf{Frm}^{\text{op}}}(L, M)$ and $g \in \text{Hom}_{\mathbf{Frm}^{\text{op}}}(M, N)$. Then $f \in \text{Hom}_{\mathbf{Frm}}(M, L)$ and $g \in \text{Hom}_{\mathbf{Frm}}(N, M)$. Hence $f \circ_{\mathbf{Frm}} g \in \text{Hom}_{\mathbf{Frm}}()$. Therefore

$$\begin{aligned} F(g \circ_{\mathbf{Frm}^{\text{op}}} f) &= F(f \circ_{\mathbf{Frm}} g) \\ &= F(f \circ_{\mathbf{Set}} g) \\ &= (f \circ_{\mathbf{Set}} g)^* \\ &= g^* \circ_{\mathbf{Set}} f^* \quad (\text{by ref ex in algebra notes}) \\ &= F(g) \circ_{\mathbf{Set}} F(f) \\ &= F(g) \circ_{\mathbf{Loc}} F(f). \end{aligned}$$

- Let $L \in \text{Obj}(\mathbf{Frm}^{\text{op}})$. Then

$$\begin{aligned}
 F(\text{id}_L^{\mathbf{Frm}^{\text{op}}}) &= F(\text{id}_L) \\
 &= \text{id}_L^* \\
 &= \text{id}_L \quad (\text{by ex in algebra notes}) \\
 &= \text{id}_{F(L)} \\
 &= \text{id}_{F(L)}^{\mathbf{Loc}}.
 \end{aligned}$$

Thus F is a functor.

- Write $F = (F_0, F_1)$.
 - Since $F_0 = \text{id}_{\text{Obj}(\mathbf{Loc})}$, F_0 is a bijection.
 - make exercise in algebra notes on complete lattices that the left adjoints are bijective to right adjoints, define operation $f \mapsto f^*$ and $g \mapsto g_*$ for f a left adjoint and g a right adjoint, show that these are inverses and thus bijections

Since F_0 and F_1 are bijections, ref exercise in notes on category theory implies that $F \in \text{Iso}_{\text{Cat}}(\mathbf{Frm}^{\text{op}}, \mathbf{Loc})$ (rewrite because $\mathbf{Frm}$ and $\mathbf{Loc}$ are not small, but maybe its okay if we are considering U -small categories $\mathbf{Loc}$ and $\mathbf{Frm}$ in the first place).

□

Exercise 5.2.1.8. Let $L, M \in \text{Obj}(\mathbf{Frm})$. Then

1. $\text{RegEpi}_{\mathbf{Frm}}(L, M) = \text{Epi}_{\mathbf{Set}}(L, M) \cap \text{Hom}_{\mathbf{Frm}}(L, M)$
2. $\text{RegMono}_{\mathbf{Loc}}(M, L) = \text{Mono}_{\mathbf{Set}}(M, L) \cap \text{Hom}_{\mathbf{Loc}}(M, L)$

Proof.

1.
 - Let $m \in \text{RegEpi}_{\mathbf{Frm}}(L, M)$. Then $m \in \text{Hom}_{\mathbf{Frm}}(L, M)$. (need exercise showing that if $m \in \text{Hom}_{\mathbf{Frm}}(L, M)$, then $m(L) \in \text{Obj}(\mathbf{Frm})$ and $m|^{m(L)} \in \text{Hom}_{\mathbf{Frm}}(L, m(L))$, then using the below idea, since $m|^{m(L)}$ is surjective frame homo, it is regular epi, and we might be able to show that if $g \circ f$ is regular epi and f is regular epi, then g is regular epi, idk, but to show that $m(L) \in \text{Obj}(\mathbf{Frm})$, we need to show that if $(M, \leq_M)$ is a poset and $K \subset M$ and $B \subset K$, if $\sup(B, \leq_M)$ exists and belongs to K , then $\sup(B, \leq_M|_K)$ exists and equals $\sup(B, \leq_M)$)
 - Let $m \in \text{Epi}_{\mathbf{Set}}(L, M) \cap \text{Hom}_{\mathbf{Frm}}(L, M)$. Then m is surjective, m is arbitrary join preserving and m finite meet preserving. Define $f, g \in \text{End}_{\mathbf{Frm}}(L)$ by $f, g := \text{id}_L$.
 - (a) By construction,

$$\begin{aligned}
 m \circ_{\mathbf{Frm}} f &= m \circ_{\mathbf{Set}} \text{id}_L \\
 &= m \\
 &= m \circ_{\mathbf{Set}} \text{id}_L \\
 &= m \circ_{\mathbf{Frm}} g.
 \end{aligned}$$

- (b) Let $M' \in \text{Obj}(\mathbf{Frm})$ and $m' \in \text{Hom}_{\mathbf{Frm}}(L, M')$. Suppose that $m' \circ_{\mathbf{Frm}} f = m' \circ_{\mathbf{Frm}} g$. Then $m' \circ_{\mathbf{Set}} f = m' \circ_{\mathbf{Set}} g$.
 - Since

$$\begin{aligned}
 m &\in \text{Epi}_{\mathbf{Set}}(L, M) \\
 &= \text{RegEpi}_{\mathbf{Set}}(L, M),
 \end{aligned}$$

there exists a unique $\phi \in \text{Hom}_{\mathbf{Set}}(M, M')$ such that $\phi \circ_{\mathbf{Set}} m = m'$.

* Let $B \subset M$. Since m is surjective, there exists $A \subset L$ such that $B = m(A)$. Then

$$\begin{aligned}
 \phi\left(\bigvee B\right) &= \phi\left(\bigvee m(A)\right) \\
 &= \phi\left(m\left(\bigvee A\right)\right) \quad (\text{since } m \in \text{Hom}_{\mathbf{Frm}}) \\
 &= \phi \circ_{\mathbf{Set}} m\left(\bigvee A\right) \\
 &= m'\left(\bigvee A\right) \\
 &= \bigvee m'(A) \quad (\text{since } m' \in \text{Hom}_{\mathbf{Frm}}) \\
 &= \bigvee \phi \circ_{\mathbf{Set}} m(A) \\
 &= \bigvee \phi(m(A)) \\
 &= \bigvee \phi(B).
 \end{aligned}$$

Since $B \subset M$ is arbitrary, we have that for each $B \subset M$, $\phi(\bigvee B) = \bigvee \phi(B)$. Thus ϕ is arbitrary join preserving.

* Similarly ϕ is finite join preserving **add details?**.

Thus $\phi \in \text{Hom}_{\mathbf{Frm}}(M, M')$.

– Let $\phi' \in \text{Hom}_{\mathbf{Frm}}(M, M')$. Suppose that $\phi' \circ_{\mathbf{Frm}} m = m'$. Then $\phi' \in \text{Hom}_{\mathbf{Set}}(M, M')$ and $\phi' \circ_{\mathbf{Set}} m = m'$. Uniqueness of ϕ then implies that $\phi = \phi'$.

Thus there exists a unique $\phi \in \text{Hom}_{\mathbf{Frm}}(M, M')$ such that $\phi \circ_{\mathbf{Frm}} m = m'$. Since $M' \in \text{Obj}(\mathbf{Frm})$ and $m' \in \text{Hom}_{\mathbf{Frm}}(L, M')$ are arbitrary, we have that for each $M' \in \text{Obj}(\mathbf{Frm})$ and $m' \in \text{Hom}_{\mathbf{Frm}}(L, M')$, there exists a unique $\phi \in \text{Hom}_{\mathbf{Frm}}(M, M')$ such that $\phi \circ_{\mathbf{Frm}} m = m'$.

Therefore $(M, m) \in \text{Coequi}_{\mathbf{Frm}}(f, g)$. Hence $m \in \text{RegEpi}_{\mathbf{Frm}}(L, M)$. Since $m \in \text{Epi}_{\mathbf{Set}}(L, M) \cap \text{Hom}_{\mathbf{Frm}}(L, M)$ is arbitrary, we have that for each $m \in \text{Epi}_{\mathbf{Set}}(L, M) \cap \text{Hom}_{\mathbf{Frm}}(L, M)$, $m \in \text{RegEpi}_{\mathbf{Frm}}(L, M)$. Thus $\text{Epi}_{\mathbf{Set}}(L, M) \cap \text{Hom}_{\mathbf{Frm}}(L, M) \subset \text{RegEpi}_{\mathbf{Frm}}(L, M)$.

Since $\text{RegEpi}_{\mathbf{Frm}}(L, M) \subset \text{Epi}_{\mathbf{Set}}(L, M) \cap \text{Hom}_{\mathbf{Frm}}(L, M)$ and $\text{Epi}_{\mathbf{Set}}(L, M) \cap \text{Hom}_{\mathbf{Frm}}(L, M) \subset \text{RegEpi}_{\mathbf{Frm}}(L, M)$, we have that $\text{RegEpi}_{\mathbf{Frm}}(L, M) = \text{Epi}_{\mathbf{Set}}(L, M) \cap \text{Hom}_{\mathbf{Frm}}(L, M)$.

2. **Use duality**

□

5.2.2 Sublocales

go over regular/extremal monomorphisms, maybe show both are equivalent for **Frm**.

Definition 5.2.2.1.

Chapter 6

Metric Spaces

6.1 Introduction

In order so that we can talk about premetric structures in model theory, do the following:

- talk about pseudometric spaces,
- define uniform continuity for pseudometric spaces,
- show that the equivalence relation defined by $x \sim y$ iff $d(x, y) = 0$ induces a metric on quotient space
- show that if $f : (X, d_X) \rightarrow (Y, d_Y)$ is uniformly continuous, then f is $(\sim_X, \sim_Y)$ invariant and thus has a factor $\text{Factor}_{\sim_X, \sim_Y}(f) = \bar{f} : X / \sim_X \rightarrow Y / \sim_Y$.
- so any predicate/function/constant in a metric prestructure based on a pseudometric space induces one in the structure based on the induced metric space

6.1.1 Metrics

Definition 6.1.1.1. Let X be a set and $d : X \times X \rightarrow \mathbb{R}$. Then

- d is said to be
 - **semidefinite** if for each $x \in X$, $d(x, x) = 0$,
 - **definite** if for each $x, y \in X$, $d(x, y) = 0$ iff $x = y$,
 - **symmetric** if for each $x, y \in X$, $d(x, y) = d(y, x)$,
- d is said to satisfy the **triangle inequality** if for each $x, y, z \in X$, $d(x, y) \leq d(x, z) + d(z, y)$,
- d is said to be a
 - **pseudometric on X** if
 1. d is semidefinite,
 2. d is symmetric,
 3. d satisfies the triangle inequality,
 - **metric on X** if
 1. d is a pseudometric on X ,
 2. d is definite.

Note 6.1.1.2. If d is definite, then d is semidefinite.

Exercise 6.1.1.3. Let X be a set and $d : X \times X \rightarrow \mathbb{R}$ a pseudometric on X . Then for each $d \geq 0$.

Proof. Let $x, y \in X$. We note that

$$\begin{aligned} d(x, x) \leq d(x, y) + d(y, x) &\implies d(x, x) - d(x, y) \leq d(y, x) \\ &\implies -d(x, y) \leq d(y, x) \\ &\implies 0 \leq 2d(x, y) \\ &\implies d(x, y) \geq 0 \end{aligned}$$

Since $d(x, x) \leq d(x, y) + d(y, x)$, we have that $d(x, y) \geq 0$. Since $x, y \in X$ are arbitrary, we have that for each $x, y \in X$, $d(x, y) \geq 0$. Thus $d \geq 0$. $\square$

Definition 6.1.1.4. Let X be a set and $d : X \times X \rightarrow [0, \infty)$. Then (X, d) is said to be a

- **pseudometric space** if d is a pseudometric on X ,
- **metric space** if d is a metric on X .

Note 6.1.1.5. When the context is clear, we may write X in place of (X, d) .

Definition 6.1.1.6. Let (X, d) be a pseudometric space. We define the **equivalence relation on X induced by d** , denoted $\sim_d \subset X \times X$, by $\sim_d := \{(x, y) \in X \times X : d(x, y) = 0\}$.

Exercise 6.1.1.7. Let (X, d) be a pseudometric space. Then $\sim_d$ is an equivalence relation on X .

Proof.

1. Let $x \in X$. Since d is semidefinite, $d(x, x) = 0$. Thus $x \sim_d x$. Since $x \in X$ is arbitrary, we have that for each $x \in X$, $x \sim_d x$.
2. Since d is symmetric, for each $x, y \in X$,

$$\begin{aligned} x \sim_d y &\iff d(x, y) = 0 \\ &\iff d(y, x) = 0 \\ &\iff y \sim_d x. \end{aligned}$$

3. Let $x, y, z \in X$. Suppose that $x \sim_d y$ and $y \sim_d z$. Then $d(x, y) = 0$ and $d(y, z) = 0$. Since d satisfies the triangle inequality, we have that

$$\begin{aligned} 0 &\leq d(x, z) \\ &\leq d(x, y) + d(y, z) \\ &= 0 + 0 \\ &= 0. \end{aligned}$$

Thus $d(x, z) = 0$. Therefore $x \sim_d z$. Since $x, y, z \in X$ are arbitrary, we have that for each $x, y, z \in X$, $x \sim_d y$ and $y \sim_d z$ implies that $x \sim_d z$. $\square$

Definition 6.1.1.8. Let (X, d) be a pseudometric space. We define the **metric on $X/\sim_d$ induced by d** , denoted $d_{X/\sim_d} : X/\sim_d \times X/\sim_d \rightarrow [0, \infty)$, by $d_{X/\sim_d} := \text{factor}(d, \pi_{\sim_d} \times \pi_{\sim_d})$.

Note 6.1.1.9. By $d(\pi_{\sim_d}(x), \pi_{\sim_d}(y)) = d(x, y)$

Exercise 6.1.1.10. Let (X, d) be a pseudometric space. Then $d_{\sim_d} : X/\sim_d \times X/\sim_d \rightarrow [0, \infty)$ is a metric on $X/\sim_d$.

Proof.

1. Let $a, b \in X/\sim_d$. Then there exists $x, y \in X$ such that $a = \pi_{\sim_d}(x)$ and $b = \pi_{\sim_d}(y)$.

- ($\implies$):
If $d_{\sim_d}(a, b) = 0$, then

$$\begin{aligned} d(x, y) &= d_{\sim_d}(\pi_{\sim_d}(x), \pi_{\sim_d}(y)) \\ &= d_{\sim_d}(a, b) \\ &= 0. \end{aligned}$$

Hence $x \sim_d y$. Therefore

$$\begin{aligned} a &= \pi_{\sim_d}(x) \\ &= \pi_{\sim_d}(y) \\ &= b. \end{aligned}$$

- ($\impliedby$):
If $a = b$, then

$$\begin{aligned} d_{\sim_d}(a, b) &= d_{\sim_d}(a, a) \\ &= d_{\sim_d}(\pi_{\sim_d}(x), \pi_{\sim_d}(x)) \\ &= d(x, x) \\ &= 0. \end{aligned}$$

Since $a, b \in X/\sim_d$ are arbitrary, we have that for each $a, b \in X/\sim_d$, $d_{\sim_d}(a, b) = 0$ iff $a = b$. Therefore $d_{\sim_d}$ is definite.

2. Let $a, b \in X/\sim_d$. Then there exists $x, y \in X$ such that $a = \pi_{\sim_d}(x)$ and $b = \pi_{\sim_d}(y)$. Hence

$$\begin{aligned} d_{\sim_d}(a, b) &= d_{\sim_d}(\pi_{\sim_d}(x), \pi_{\sim_d}(y)) \\ &= d(x, y) \\ &= d(y, x) \\ &= d_{\sim_d}(\pi_{\sim_d}(y), \pi_{\sim_d}(x)) \\ &= d_{\sim_d}(b, a). \end{aligned}$$

Since $a, b \in X/\sim_d$ are arbitrary, we have that for each $a, b \in X/\sim_d$, $d_{\sim_d}(a, b) = d_{\sim_d}(b, a)$. Hence $d_{\sim_d}$ is symmetric.

3. Let $a, b, c \in X/\sim_d$. Then there exists $x, y, z \in X$ such that $a = \pi_{\sim_d}(x)$, $b = \pi_{\sim_d}(y)$ and $c = \pi_{\sim_d}(z)$. Hence

$$\begin{aligned} d_{\sim_d}(a, c) &= d_{\sim_d}(\pi_{\sim_d}(x), \pi_{\sim_d}(z)) \\ &= d(x, z) \\ &\leq d(x, y) + d(y, z) \\ &= d_{\sim_d}(\pi_{\sim_d}(x), \pi_{\sim_d}(y)) + d_{\sim_d}(\pi_{\sim_d}(y), \pi_{\sim_d}(z)) \\ &= d_{\sim_d}(a, b) + d_{\sim_d}(b, c). \end{aligned}$$

Therefore $d_{\sim_d}$ satisfies the triangle inequality.

Thus $d_{\sim_d}$ is a metric on $X/\sim_d$. □

Exercise 6.1.1.11. Show $d_{\mathbb{R}} : \mathbb{R} \rightarrow \mathbb{R}$ given by $d_{\mathbb{R}}(a, b) := |a - b|$ is a metric and make section on absolute value in intro to math notes

Proof. **FINISH!!!** □

Exercise 6.1.1.12. Reverse Triangle Inequality:

Let (X, d) be a pseudometric space. Then for each $x, y, z \in X$, $|d(x, y) - d(y, z)| \leq d(x, z)$.

Proof. Let $x, y, z \in X$. The triangle inequality implies that $d(x, y) \leq d(x, z) + d(z, y)$. Hence $d(x, y) - d(z, y) \leq d(x, z)$. Similarly, the triangle inequality implies that $d(y, z) \leq d(y, x) + d(x, z)$. Then $-d(y, z) \geq -d(y, x) - d(x, z)$. Therefore

$$-d(x, z) \leq d(x, y) - d(z, y) \leq d(x, z)$$

and $|d(x, y) - d(z, y)| \leq d(x, z)$. □

FINISH!!! change 'metric' to 'pseudo metric' in any exercises where possible to generalize

Definition 6.1.1.13. Let (X, d) be a metric space, $A \subset X$ and $x \in X$. We define the **distance between A and x** , denoted $d(x, A)$, by

$$d(x, A) = \inf\{d(x, a) : a \in A\}$$

Exercise 6.1.1.14. Let (X, d) be a metric space, $A \subset X$. Then for each $x, y \in X$, $|d(x, A) - d(y, A)| \leq d(x, y)$.

Proof. Let $x, y \in X$. Let $\epsilon > 0$. Then there exists $a \in A$ such that $d(y, a) < d(y, A) + \epsilon$. The triangle inequality implies that

$$\begin{aligned} d(x, A) &\leq d(x, a) \\ &\leq d(x, y) + d(y, a) \\ &< d(x, y) + d(y, A) + \epsilon. \end{aligned}$$

Since $\epsilon > 0$ is arbitrary, we have that $d(x, A) \leq d(x, y) + d(y, A)$. Hence $d(x, A) - d(y, A) \leq d(x, y)$.

Similarly, we have that $d(y, A) - d(x, A) \leq d(x, y)$. Therefore $|d(y, A) - d(x, A)| \leq d(x, y)$. □

Definition 6.1.1.15. Let (X, d) be a metric space and $A, B \subset X$. We define the **distance between A and B** , denoted $d(A, B)$, by

$$d(A, B) = \inf_{\substack{a \in A \\ b \in B}} d(a, b)$$

Exercise 6.1.1.16. Let (X, d) be a metric space. Then for each $A, B \subset X$ and $c \in X$,

$$d(A, B) \leq d(A, c) + d(c, B)$$

Proof. Let $A, B \subset X$, $c \in X$ and $\epsilon > 0$. Choose $a \in A$ and $b \in B$ such that $d(a, c) < d(A, c) + \epsilon/2$ and $d(c, b) < d(c, B) + \epsilon/2$. Then

$$\begin{aligned} d(A, B) &\leq d(a, b) \\ &\leq d(a, c) + d(c, b) \\ &< d(A, c) + \frac{\epsilon}{2} + d(c, B) + \frac{\epsilon}{2} \\ &= d(A, c) + d(c, B) + \epsilon \end{aligned}$$

Since $\epsilon > 0$ is arbitrary, $d(A, B) \leq d(A, c) + d(c, B)$. □

Definition 6.1.1.17. Let (X, d) be a metric space and $A \subset X$.

- Then A is said to be d -bounded if there exists $M \in [0, \infty)$ such that for each $x, y \in A$, $d(x, y) \leq M$.
- We define $\text{diam}(A, d) \in [0, \infty]$, by $\text{diam}(A, d) := \inf\{M \in [0, \infty) : \text{for each } x, y \in A, d(x, y) \leq M\}$.

Note 6.1.1.18. When the context is clear, we may say “bounded” instead of “ d -bounded” and we may write $\text{diam}(A)$ in place of $\text{diam}(A, d)$.

Exercise 6.1.1.19. Let (X, d) be a metric space and $A \subset X$. Then

1. $\text{diam}(A) < \infty$ implies that for each $x, y \in A$, $d(x, y) \leq \text{diam}(A)$,
2. A is bounded iff $\text{diam}(A) < \infty$.

Proof. Define $E \subset [0, \infty)$ by $E := \{M \in [0, \infty) : \text{for each } x, y \in A, d(x, y) \leq M\}$.

1. Suppose that $\text{diam}(A) < \infty$. Then $\text{diam}(A) \in [0, \infty)$. Let $x, y \in A$ and $\epsilon > 0$. Since

$$\begin{aligned}\text{diam}(A) + \epsilon &> \text{diam}(A) \\ &= \inf E,\end{aligned}$$

there exists $M \in E$ such that $M < \text{diam}(A) + \epsilon$. Since $M \in E$, we have that

$$\begin{aligned}d(x, y) &\leq M \\ &< \text{diam}(A) + \epsilon.\end{aligned}$$

Since $\epsilon > 0$ is arbitrary, we have that for each $\epsilon > 0$, $d(x, y) < \text{diam}(A) + \epsilon$. Hence $d(x, y) \leq \text{diam}(A)$. Since $x, y \in A$ are arbitrary, we have that for each $x, y \in A$, $d(x, y) \leq \text{diam}(A)$.

2. • ($\implies$):

Suppose that A is bounded. Since A is bounded, $E \neq \emptyset$. Since $0 \in \text{lb}(E)$, **reference least upper bound property** implies that there exists $M_0 \in [0, \infty)$ such that $M_0 = \inf E$. By definition,

$$\begin{aligned}\text{diam}(A) &= \inf E \\ &= M_0 \\ &< \infty.\end{aligned}$$

- ($\impliedby$):

Suppose that $\text{diam}(A) < \infty$. **The previous part (ref)** implies that $\text{diam}(A) \in E$. Thus $E \neq \emptyset$ and A is bounded. □

Exercise 6.1.1.20. Let (X, d) be a metric space and $A \subset X$. Suppose that $A \neq \emptyset$.

1. For each $a \in \mathbb{R}$, $a < \text{diam}(A)$ implies that there exists $x, y \in A$ such that $d(x, y) > a$,
2. If A is bounded, then $\text{diam}(A) = \sup\{d(x, y) : x, y \in A\}$.

Proof.

1. Let $a \in \mathbb{R}$. Suppose that $a < \text{diam}(A)$.

- Suppose that $a < 0$. Since $A \neq \emptyset$, there exists $x \in X$ such that $x \in A$. Then

$$\begin{aligned}d(x, x) &= 0 \\ &> a.\end{aligned}$$

- Suppose that $a \geq 0$. If $a \in E$, then

$$\begin{aligned}\text{diam}(A) &= \inf E \\ &\leq a \\ &< \text{diam}(A),\end{aligned}$$

which is a contradiction. Thus $a \notin E$. Hence there exists $x, y \in A$ such that $d(x, y) > a$.

2. Suppose that A is bounded. Define $s \in [0, \infty]$ and $E \subset [0, \infty)$ by $s := \sup\{d(x, y) : x, y \in A\}$ and $E := \{M \in [0, \infty) : \text{for each } x, y \in A, d(x, y) \leq M\}$.

- Let $M \in E$. Then $M \in \text{ub}\{d(x, y) : x, y \in A\}$. Therefore

$$\begin{aligned}s &= \sup\{d(x, y) : x, y \in A\} \\ &\leq M.\end{aligned}$$

Since $M \in E$ is arbitrary, we have that for each $M \in E$, $s \leq M$. Hence $s \in \text{lb } E$. Thus

$$\begin{aligned}s &\leq \inf E \\ &= \text{diam}(A).\end{aligned}$$

- Let $\epsilon > 0$. Since $\text{diam}(A) - \epsilon < \text{diam}(A)$, the prev part (ref) implies that there exists $x, y \in A$ such that $d(x, y) > \text{diam}(A) - \epsilon$. Then

$$\begin{aligned} \text{diam}(A) - \epsilon &< d(x, y) \\ &\leq \sup\{d(x, y) : x, y \in A\} \\ &= s. \end{aligned}$$

Since $\epsilon > 0$ is arbitrary, we have that $\text{diam}(A) \leq s$.

Since $s \leq \text{diam}(A)$ and $\text{diam}(A) \leq s$, we have that $\text{diam}(A) = s$.

□

6.1.2 Metric Topology

REWORK for pseudometrics, show that $\mathcal{T}_d$ is T_0 iff d is a metric on X

Definition 6.1.2.1. Let (X, d) be a pseudometric space.

- Let $x \in X$ and $r > 0$. We define the

– **open ball of radius r centered at x in (X, d)** , denoted $B_d(x, r)$, by

$$B_d(x, r) = \{y \in X : d(x, y) < r\}$$

– **closed ball of radius r centered at x in (X, d)** , denoted $\bar{B}_d(x, r)$, by

$$\bar{B}_d(x, r) = \{y \in X : d(x, y) \leq r\}$$

– **circle of radius r centered at x in (X, d)** , denoted $T_d(x, r)$, by

$$T_d(x, r) = \{y \in X : d(x, y) = r\}$$

- We define $\mathcal{B}_d := \{B_d(x, \delta) : x \in X \text{ and } \delta > 0\}$.

Note 6.1.2.2. When the context is clear we write $B(x, r)$, $\bar{B}(x, r)$ and $T(x, r)$ in place of $B_d(x, r)$, $\bar{B}_d(x, r)$ and $T_d(x, r)$ respectively.

Exercise 6.1.2.3. go back and explain that the reverse triangle inequality is essentially this fact and draw picture to illustrate Let (X, d) be a pseudometric space and $B \in \mathcal{B}_d$. Then for each $x \in B$, there exists $\delta > 0$ such that $B_d(x, \delta) \subset B$.

Proof. Let $x \in B$. Since $B \in \mathcal{B}_d$, there exists $x_0 \in X$ and $\epsilon > 0$ such that $B = B_d(x_0, \epsilon)$. Since $x \in B_d(x_0, \epsilon)$, we have that $d(x, x_0) < \epsilon$. Define $\delta > 0$ by $\delta := \epsilon - d(x_0, x)$. Let $y \in B_d(x, \delta)$. The reverse triangle inequality implies that

$$\begin{aligned} d(y, x_0) - d(x_0, x) &\leq |d(y, x_0) - d(x_0, x)| \\ &\leq d(y, x) \\ &< \delta. \end{aligned}$$

Therefore

$$\begin{aligned} d(y, x_0) &< \delta + d(x_0, x) \\ &= (\epsilon - d(x_0, x)) + d(x_0, x) \\ &= \epsilon. \end{aligned}$$

Thus $y \in B_d(x_0, \epsilon)$. Since $y \in B_d(x, \delta)$ is arbitrary, we have that for each $y \in B_d(x, \delta)$, $y \in B_d(x_0, \epsilon)$. Hence

$$\begin{aligned} B_d(x, \delta) &\subset B_d(x_0, \epsilon) \\ &= B. \end{aligned}$$

Since $x \in B$ is arbitrary, we have that for each $x \in B$, there exists $\delta > 0$ such that $B_d(x, \delta) \subset B$.

□

Exercise 6.1.2.4. Let (X, d) be a pseudometric space, $U \subset X$ and $x \in U$. Then there exists $B \in \mathcal{B}_d$ such that $x \in B$ and $B \subset U$ iff there exists $\delta > 0$ such that $B_d(x, \delta) \subset U$.

Proof.

- ($\implies$):
Suppose that there exists $B \in \mathcal{B}_d$ such that $x \in B$ and $B \subset U$. Exercise 6.1.2.3 implies that $\delta > 0$ such that $B_d(x, \delta) \subset B$. Then $B_d(x, \delta) \subset U$.
- ($\impliedby$):
Suppose that there exists $\delta > 0$ such that $B_d(x, \delta) \subset U$. Since $x \in B_d(x, \delta)$ and $B_d(x, \delta) \subset U$, we have that there exists $B \in \mathcal{B}_d$ such that $x \in B$ and $B \subset U$.

□

Definition 6.1.2.5. Let (X, d) be a pseudometric space. We define the **metric topology on X** , denoted $\mathcal{T}_d$, by

$$\mathcal{T}_d := \tau_X(\mathcal{B}_d).$$

Exercise 6.1.2.6. Let (X, d) be a pseudometric space. Then

1. $\mathcal{T}_d = \{U \subset X : \text{for each } x \in U, \text{ there exists } \delta > 0 \text{ such that } B(x, \delta) \subset U\}$,
2. $\mathcal{B}_d$ is a basis for $\mathcal{T}_d$.

Proof.

1. • Define $\mathcal{F} \subset \mathcal{P}(X)$ by $\mathcal{F} := \{U \subset X : \text{for each } x \in U, \text{ there exists } V \in \mathcal{B}_d \text{ such that } x \in V \text{ and } V \subset U\}$.
 - Clearly for each $x \in X$, there exists $V \in \mathcal{B}_d$ such that $x \in V$.
 - Let $x \in X$ and $U, V \in \mathcal{B}_d$. Suppose that $x \in U \cap V$. Since $x \in U \cap V$, we have that $x \in U$ and $x \in V$. Exercise 6.1.2.3 implies that there exists $\delta_U, \delta_V > 0$ such that $B_d(x, \delta_U) \subset U$ and $B_d(x, \delta_V) \subset V$. Define $\delta > 0$ and $W \in \mathcal{B}_d$ by $\delta := \min\{\delta_U, \delta_V\}$ and $W := B_d(x, \delta)$. Then $x \in W$ and $W \subset U \cap V$.

Exercise 4.1.0.11 then implies that $\mathcal{F}$ is a topology on X , $\mathcal{B}_d$ is a basis for $\mathcal{F}$ and $\mathcal{F} = \tau_X(\mathcal{B}_d)$.

- Define $\mathcal{G} \subset \mathcal{P}(X)$ by $\mathcal{G} := \{U \subset X : \text{for each } x \in U, \text{ there exists } \delta > 0 \text{ such that } B_d(x, \delta) \subset U\}$. We note that

$$\begin{aligned} U \in \mathcal{F} &\iff \text{for each } x \in U, \text{ there exists } B \in \mathcal{B}_d \text{ such that } x \in B \text{ and } B \subset U \\ &\iff \text{for each } x \in U, \text{ there exists } \delta > 0 \text{ such that } B_d(x, \delta) \subset U \quad (\text{by Exercise 6.1.2.4}) \\ &\iff U \in \mathcal{G}. \end{aligned}$$

Thus $\mathcal{F} = \mathcal{G}$. By definition,

$$\begin{aligned} \mathcal{T}_d &= \tau_X(\mathcal{B}_d) \\ &= \mathcal{F} \\ &= \mathcal{G}. \end{aligned}$$

2. Since $\mathcal{B}_d$ is a basis for $\mathcal{F}$ and $\mathcal{F} = \mathcal{T}_d$, we have that $\mathcal{B}_d$ is a basis for $\mathcal{T}_d$.

□

Exercise 6.1.2.7. Let (X, d) be a pseudometric space, $\mathcal{B}$ a basis for $\mathcal{T}_d$ and $\epsilon > 0$. Define $\mathcal{B}_\epsilon \subset \mathcal{T}_d$ by $\mathcal{B}_\epsilon := \{B \in \mathcal{B} : \text{diam}(B) \leq \epsilon\}$. Then $\mathcal{B}_\epsilon$ is a basis for $\mathcal{T}_d$.

Proof. Let $U \in \mathcal{T}_d$ and $x \in U$. Exercise 6.1.2.6 implies that there exists $\delta_0 > 0$ such that $B(x, \delta_0) \subset U$. Set $\delta := \min(\delta_0, \epsilon/2)$. Since $B(x, \delta) \in \mathcal{T}_d$ and $\mathcal{B}$ is a basis for $\mathcal{T}_d$, there exists $B \in \mathcal{B}$ such that $x \in B$ and $B \subset B(x, \delta)$. We note that

$$\begin{aligned} B &\subset B(x, \delta) \\ &\subset B(x, \delta_0) \\ &\subset U \end{aligned}$$

and

$$\begin{aligned} \text{diam}(B) &\leq \text{diam}(B(x, \delta)) \\ &\leq 2\delta \\ &\leq \epsilon. \end{aligned}$$

Since $\text{diam}(B) \leq \epsilon$, $B \in \mathcal{B}_\epsilon$. Since $U \in \mathcal{T}_d$ and $x \in U$ are arbitrary, we have that for each $U \in \mathcal{T}_d$ and $x \in U$, there exists $B \in \mathcal{B}_\epsilon$ such that $x \in B$ and $B \subset U$. Thus $\mathcal{B}_\epsilon$ is a basis for $\mathcal{T}_d$. $\square$

Exercise 6.1.2.8. Let (X, d) be a pseudometric space and $x \in X$. Set $\mathcal{B}_x = \{B(x, n^{-1}) : n \in \mathbb{N}\}$. Then $\mathcal{B}$ is a local basis for $\mathcal{T}_d$ at x .

Proof.

1. Clearly for each $U \in \mathcal{B}_x$, $x \in U$.
2. Let $V \in \mathcal{T}_d$. Suppose that $x \in V$. Exercise 6.1.2.6 implies that there exists $\delta > 0$ such that $B(x, \delta) \in V$. Choose $n \in \mathbb{N}$ such that $n^{-1} < \delta$. Set $U := B(x, n^{-1})$. Then $U \in \mathcal{B}_x$ and

$$\begin{aligned} U &= B(x, n^{-1}) \\ &\subset B(x, \delta) \\ &\subset V. \end{aligned}$$

Hence $\mathcal{B}_x$ is a local basis for $\mathcal{T}_d$ at x . $\square$

Exercise 6.1.2.9. Let (X, d) be a pseudometric space. Then for each $x \in X$ and $r \geq 0$,

1. $\bar{B}(x, r)$ is $\mathcal{T}_d$ -closed,
2. $T(x, r)$ is $\mathcal{T}_d$ -closed,

Proof. Let $x \in X$ and $r \geq 0$.

1. Let $y \in \bar{B}(x, r)^c$. Then $d(x, y) > r$. Define $\epsilon > 0$ by $\epsilon := d(x, y) - r$. Let $z \in B(y, \epsilon)$. Since $d(x, y) \leq d(x, z) + d(z, y)$, we have that

$$\begin{aligned} d(x, z) &\geq d(x, y) - d(z, y) \\ &> d(x, y) - \epsilon \\ &= r. \end{aligned}$$

Thus $z \in \bar{B}(x, r)^c$. Since $z \in B(y, \epsilon)$ is arbitrary, we have that for each $z \in B(y, \epsilon)$, $z \in \bar{B}(x, r)^c$. Hence $B(y, \epsilon) \subset \bar{B}(x, r)^c$. Since $y \in \bar{B}(x, r)^c$ is arbitrary, we have that for each $y \in \bar{B}(x, r)^c$, there exists $\epsilon > 0$ such that $B(y, \epsilon) \subset \bar{B}(x, r)^c$. Exercise 6.1.2.6 then implies that $\bar{B}(x, r)^c \in \mathcal{T}_d$. Thus $\bar{B}(x, r)$ is $\mathcal{T}_d$ -closed.

2. We note that

$$\begin{aligned} T(x, r) &= \bar{B}(x, r) \setminus B(x, r) \\ &= \bar{B}(x, r) \cap B(x, r)^c. \end{aligned}$$

Since $\bar{B}(x, r)$ is $\mathcal{T}_d$ -closed and $B(x, r)^c$ is $\mathcal{T}_d$ -closed, we have that $T(x, r)$ is $\mathcal{T}_d$ -closed. $\square$

Exercise 6.1.2.10. Let (X, d_X) , (Y, d_Y) be pseudometric spaces, $f : X \rightarrow Y$ and $x_0 \in X$. Then f is continuous at x_0 iff for each $\epsilon > 0$, there exists $\delta > 0$ such that for each $x \in X$, $d_X(x_0, x) < \delta$ implies that $d_Y(f(x_0), f(x)) < \epsilon$.

Proof.

- ($\implies$) :

Suppose that f is continuous at x_0 . Let $\epsilon > 0$. Set $V := B_{d_Y}(f(x_0), \epsilon)$. Since $V \in \mathcal{N}(f(x_0))$, continuity at x_0 and Exercise 4.2.0.5 imply that $f^{-1}(V) \in \mathcal{N}(x_0)$. Then $x_0 \in \text{Int } f^{-1}(V)$ and $\text{Int } f^{-1}(V) \in \mathcal{T}_d$. Exercise 6.1.2.6 then implies that there exists $\delta > 0$ such that $B_{d_X}(x_0, \delta) \subset \text{Int } f^{-1}(V)$. Set $U := B_{d_X}(x_0, \delta)$. Let $x \in X$. Suppose that $d_X(x_0, x) < \delta$. Then $x \in U$. Since $U \subset f^{-1}(V)$, we have that

$$\begin{aligned} f(x) &\in V \\ &= B_{d_Y}(f(x_0), \epsilon) \end{aligned}$$

and $d_Y(f(x_0), f(x)) < \epsilon$. Since $\epsilon > 0$ is arbitrary, we have that for each $\epsilon > 0$, there exists $\delta > 0$ such that for each $x \in X$, $d_X(x_0, x) < \delta$ implies that $d_Y(f(x_0), f(x)) < \epsilon$.

- ($\impliedby$) :

Suppose that for each $\epsilon > 0$, there exists $\delta > 0$ such that for each $x \in X$, $d_X(x_0, x) < \delta$ implies that $d_Y(f(x_0), f(x)) < \epsilon$.
FINISH!!!

□

Exercise 6.1.2.11. Let (X, d) be a pseudometric space. Then $(X, \mathcal{T}_d)$ is first-countable.

Proof. Let $x \in X$. Exercise 6.1.2.8 implies that for each $x \in X$, there exists $\mathcal{B}_x \subset \mathcal{T}_d$ such that

1. $\mathcal{B}_x$ is a local basis for $\mathcal{T}_d$ at x ,
2. $\mathcal{B}_x$ is countable.

Thus $(X, \mathcal{T}_d)$ is first-countable. □

Exercise 6.1.2.12. rework with sequences Let (X, d) be a pseudometric space, $(x_n)_{n \in \mathbb{N}} \subset X$ and $x \in X$. The following are equivalent:

1. $x_n \rightarrow x$ in $(X, \mathcal{T}_d)$
2. $d(x_n, x) \rightarrow 0$ in $(\mathbb{R}, \mathcal{T}_{\mathbb{R}})$.
3. for each $\epsilon > 0$, there exists $N \in \mathbb{N}$ such that for each $n \in \mathbb{N}$, $n \geq N$ implies that $d(x_n, x) < \epsilon$.

Proof.

- (1) $\implies$ (2):

Suppose that $x_n \rightarrow x$ in $(X, \mathcal{T}_d)$. Let $U \in \mathcal{N}(0, \mathcal{T}_{\mathbb{R}})$. Then there exists $\epsilon > 0$ such that $(-\epsilon, \epsilon) \subset U$. Since $B(x, \epsilon) \in \mathcal{N}(x)$ and $x_n \rightarrow x$ in $(X, \mathcal{T}_d)$, we have that $(x_n)_{n \in \mathbb{N}}$ is eventually in $B(x, \epsilon)$. Thus there exists $n_0 \in \mathbb{N}$ such that for each $n \in \mathbb{N}$, $n \geq n_0$ implies that $x_n \in B(x, \epsilon)$. Let $n \in \mathbb{N}$. Suppose that $n \geq n_0$. Then

$$\begin{aligned} d(x_n, x) &\in (-\epsilon, \epsilon) \\ &\subset U. \end{aligned}$$

Thus $(d(x_n, x))_{n \in \mathbb{N}}$ is eventually in U . Since $U \in \mathcal{N}(0, \mathcal{T}_{\mathbb{R}})$ is arbitrary, we have that for each $U \in \mathcal{N}(0, \mathcal{T}_{\mathbb{R}})$, $(d(x_n, x))_{n \in \mathbb{N}}$ is eventually in U . Hence $d(x_n, x) \rightarrow 0$.

- (2) $\implies$ (3):

Suppose that $d(x_n, x) \rightarrow 0$ in $(\mathbb{R}, \mathcal{T}_{\mathbb{R}})$. Let $\epsilon > 0$. Since $(-\epsilon, \epsilon) \in \mathcal{N}(0, \mathcal{T}_{\mathbb{R}})$ and $d(x_n, x) \rightarrow 0$ in $(\mathbb{R}, \mathcal{T}_{\mathbb{R}})$, we have that $(d(x_n, x))_{n \in \mathbb{N}}$ is eventually in $(-\epsilon, \epsilon)$. Thus there exists $N \in \mathbb{N}$ such that for each $n \in \mathbb{N}$, $n \geq N$ implies that $d(x_n, x) \in (-\epsilon, \epsilon)$. In particular, for each $n \in \mathbb{N}$, $n \geq N$ implies that $d(x_n, x) < \epsilon$.

- (3) $\implies$ (1):

Suppose that for each $\epsilon > 0$, there exists $N \in \mathbb{N}$ such that for each $n \in \mathbb{N}$, $n \geq N$ implies that $d(x_n, x) < \epsilon$. Let $U \in \mathcal{N}(x)$. Then $\text{Int } U \in \mathcal{N}^{\text{opn}}(x)$. Exercise 6.1.2.6 implies that there exists $x_0 \in X$ and $\delta > 0$ such that $x \in B(x_0, \delta)$ and $B(x_0, \delta) \subset \text{Int } U$.

Define $\epsilon > 0$ by $\epsilon := \delta/2$. By assumption, there exists $N \in \mathbb{N}$ such that for each $n \in \mathbb{N}$, $n \geq N$ implies that $d(x_n, x) < \epsilon$. Let $n \in \mathbb{N}$. Suppose that $n \geq N$. Then

$$\begin{aligned} d(x_n, x_0) &\leq d(x_n, x) + d(x, x_0) \\ &< \epsilon + \epsilon \\ &= 2\epsilon \\ &= \delta. \end{aligned}$$

Thus

$$\begin{aligned} x_n &\in B(x_0, \delta) \\ &\subset \text{Int } U \\ &\subset U. \end{aligned}$$

Since $n \in \mathbb{N}$ with $n \geq N$ is arbitrary, we have that for each $n \in \mathbb{N}$, $n \geq N$ implies that $x_n \in U$. Hence $(x_n)_{n \in \mathbb{N}}$ is eventually in U . Since $U \in \mathcal{N}(x)$ is arbitrary, we have that for each $U \in \mathcal{N}(x)$, $(x_n)_{n \in \mathbb{N}}$ is eventually in U . Thus $x_n \rightarrow x$ in $(X, \mathcal{T}_d)$. □

Exercise 6.1.2.13. Let (X, d) be a pseudometric space. Then d is $(\mathcal{T}_d \otimes \mathcal{T}_d, \mathcal{T}_{\mathbb{R}})$ -continuous. Think I need to make an exercise showing a metric or abs val induces $\mathcal{T}_R$

Proof. Let $(x_n, y_n)_{n \in \mathbb{N}} \subset X \times X$ and $(x, y) \in X \times X$. Suppose that $(x_n, y_n)_n \rightarrow (x, y)$ in $(X \times X, \mathcal{T}_d \otimes \mathcal{T}_d)$. ref product net theorem in product topology section implies that $x_n \rightarrow x$ and $y_n \rightarrow y$. Let $\epsilon > 0$. Then there exists $N_x \in \mathbb{N}$ such that for each $n \in \mathbb{N}$, $n \geq N_x$ implies that $d(x_n, x) < \epsilon/2$. Similarly, there exists $N_y \in \mathbb{N}$ such that for each $n \in \mathbb{N}$, $n \geq N_y$ implies that $d(y_n, y) < \epsilon/2$. Define $N \in \mathbb{N}$ by $N := \max\{N_x, N_y\}$. Then for each $n \in \mathbb{N}$, $n \geq N$ implies that

$$\begin{aligned} |d(x_n, y_n) - d(x, y)| &\leq |d(x_n, y_n) - d(x_n, y)| + |d(x_n, y) - d(x, y)| \\ &\leq d(y_n, y) + d(x_n, x) \quad (\text{by Exercise 6.1.1.12}) \\ &< \frac{\epsilon}{2} + \frac{\epsilon}{2} \\ &= \epsilon. \end{aligned}$$

Thus $d(x_n, y_n) \rightarrow d(x, y)$. Since $(x_n, y_n)_{n \in \mathbb{N}} \subset X \times X$ and $(x, y) \in X \times X$ with $(x_n, y_n) \rightarrow (x, y)$ in $(X \times X, \mathcal{T}_d \otimes \mathcal{T}_d)$ is arbitrary, we have that for each $(x_n, y_n)_{n \in \mathbb{N}} \subset X \times X$ and $(x, y) \in X \times X$, $(x_n, y_n) \rightarrow (x, y)$ in $(X \times X, \mathcal{T}_d \otimes \mathcal{T}_d)$ implies that $d(x_n, y_n) \rightarrow d(x, y)$. Exercise 6.1.2.11 and Exercise 4.13.1.5 imply that d is $(\mathcal{T}_d \otimes \mathcal{T}_d, \mathcal{T}_{\mathbb{R}})$ -continuous. □

Exercise 6.1.2.14. Let (X, d) be a pseudometric space and $A \subset X$. If $\text{diam } A < \infty$, then $\text{diam cl } A = \text{diam } A$.

Proof. Suppose that $\text{diam } A < \infty$.

- If $A = \emptyset$, then $\text{cl } A = \emptyset$ and

$$\begin{aligned} \text{diam cl } A &= 0 \\ &= \text{diam } A. \end{aligned}$$

- Suppose that $A \neq \emptyset$. rework after making section on $\mathbb{R}$ showing things like $x_n \leq a$ and $x_n \rightarrow x$ implies that $x \leq a$.

Since $A \subset \text{cl } A$, we have that $\{d(x, y) : x, y \in A\} \subset \{d(x, y) : x, y \in \text{cl } A\}$. Therefore

$$\begin{aligned} \text{diam } A &= \sup\{d(x, y) : x, y \in A\} \\ &\leq \sup\{d(x, y) : x, y \in \text{cl } A\}. \end{aligned}$$

Let $x, y \in \text{cl } A$. Exercise 6.1.2.11 and Exercise 4.13.1.3 imply that there exist $(x_n)_{n \in \mathbb{N}}, (y_n)_{n \in \mathbb{N}} \subset A$ such that $x_n \rightarrow x$ in $(X, \mathcal{T}_d)$ and $y_n \rightarrow y$ in $(X, \mathcal{T}_d)$. Exercise 6.1.2.13 implies that d is $(\mathcal{T}_d \otimes \mathcal{T}_d, \mathcal{T}_{\mathbb{R}})$ -continuous. Since for each $n \in \mathbb{N}$, $d(x_n, y_n) \leq \text{diam } A$, we have that $(d(x_n, y_n))_{n \in \mathbb{N}} \subset [0, \text{diam } A]$. Since $[0, \text{diam } A]$ is $\mathcal{T}_{\mathbb{R}}$ -closed and $d(x_n, y_n) \rightarrow d(x, y)$, Exercise 4.13.1.4 implies that

$d(x, y) \in [0, \text{diam } A]$. Hence $d(x, y) \leq \text{diam } A$. Since $x, y \in \text{cl } A$ are arbitrary, we have that for each $x, y \in \text{cl } A$, $d(x, y) \leq \text{diam } A$. Then $\text{diam } A \in \text{ub}\{d(x, y) : x, y \in A\}$ and

$$\begin{aligned} \text{diam } A &\leq \text{diam } \text{cl } A &&= \sup\{d(x, y) : x, y \in A\} \quad (\text{by Exercise 6.1.1.20}) \\ &\leq \text{diam } A. \end{aligned}$$

Hence $\text{diam } \text{cl } A = \text{diam } A$. □

Exercise 6.1.2.15. Let (X, d) be a pseudometric space, $C \subset X$ and $x \in X$. Suppose that C is closed. Then $d(x, C) = 0$ iff $x \in C$.

Proof.

- ($\implies$):
Suppose that $d(x, C) = 0$. Then for each $n \in \mathbb{N}$, there exists $c_n \in C$ such that $d(x, c_n) < 1/n$. Then $c_n \rightarrow x$. Since C is closed, $x \in C$.
- ($\impliedby$):
Conversely, suppose that $x \in C$. Then

$$\begin{aligned} d(x, C) &= \inf\{d(x, c) : c \in C\} \\ &\leq d(x, x) \\ &= 0 \end{aligned}$$

Hence $d(x, C) = 0$. □

6.1.3 Uniform and Lipschitz Continuity

Definition 6.1.3.1. Let (X, d_X) and (Y, d_Y) be pseudometric spaces and $f : X \rightarrow Y$. Then f is said to be (d_X, d_Y) -**uniformly continuous** if for each $\epsilon > 0$, there exists $\delta > 0$ such that for each $x \in X$, $f(B_{d_X}(x, \delta)) \subset B_{d_Y}(f(x), \epsilon)$.

Note 6.1.3.2. When the context is clear, we may say “uniformly continuous” instead of “ (d_X, d_Y) -uniformly continuous”.

Exercise 6.1.3.3. Let (X, d_X) and (Y, d_Y) be pseudometric spaces and $f : X \rightarrow Y$. Then f is (d_X, d_Y) -uniformly continuous iff for each $\epsilon > 0$, there exists $\delta > 0$ such that for each $x_1, x_2 \in X$, $d_X(x_1, x_2) < \delta$ implies that $d_Y(f(x_1), f(x_2)) < \epsilon$.

Proof.

- ($\implies$):
Suppose that f is (d_X, d_Y) -uniformly continuous. Let $\epsilon > 0$. By Definition 6.1.3.1, there exists $\delta > 0$ such that for each $x \in X$, $f(B_{d_X}(x, \delta)) \subset B_{d_Y}(f(x), \epsilon)$. Let $x_1, x_2 \in X$. Suppose that $d_X(x_1, x_2) < \delta$. Then $x_2 \in B_{d_X}(x_1, \delta)$. Therefore

$$\begin{aligned} f(x_2) &\in f(B_{d_X}(x_1, \delta)) \\ &\subset B_{d_Y}(f(x_1), \epsilon). \end{aligned}$$

Hence $d_Y(f(x_1), f(x_2)) < \epsilon$. Since $x_1, x_2 \in X$ with $d_X(x_1, x_2) < \delta$ are arbitrary, we have that for each $x_1, x_2 \in X$, $d_X(x_1, x_2) < \delta$ implies that $d_Y(f(x_1), f(x_2)) < \epsilon$. Since $\epsilon > 0$ is arbitrary, we have that for each $\epsilon > 0$, there exists $\delta > 0$ such that for each $x_1, x_2 \in X$, $d_X(x_1, x_2) < \delta$ implies that $d_Y(f(x_1), f(x_2)) < \epsilon$.

- ($\impliedby$):
Suppose that for each $\epsilon > 0$, there exists $\delta > 0$ such that for each $x_1, x_2 \in X$, $d_X(x_1, x_2) < \delta$ implies that $d_Y(f(x_1), f(x_2)) < \epsilon$. Let $\epsilon > 0$. By assumption, there exists $\delta > 0$ such that for each $x_1, x_2 \in X$, $d_X(x_1, x_2) < \delta$ implies that $d_Y(f(x_1), f(x_2)) < \epsilon$. Let $x \in X$ and $y' \in f(B_{d_X}(x, \delta))$. Then there exists $x' \in B_{d_X}(x, \delta)$ such that $y' = f(x')$. Thus $d_X(x, x') < \delta$. Therefore $d_Y(f(x), f(x')) < \epsilon$. Hence

$$\begin{aligned} y' &= f(x') \\ &\in B_{d_Y}(f(x), \epsilon). \end{aligned}$$

Since $y' \in f(B_{d_X}(x, \delta))$ is arbitrary, we have that for each $y' \in f(B_{d_X}(x, \delta))$, $y' \in B_{d_Y}(f(x), \epsilon)$. Thus $f(B_{d_X}(x, \delta)) \subset B_{d_Y}(f(x), \epsilon)$. Since $x \in X$ is arbitrary, we have that for each $x \in X$, $f(B_{d_X}(x, \delta)) \subset B_{d_Y}(f(x), \epsilon)$. Since $\epsilon > 0$ is arbitrary, we have that for each $\epsilon > 0$, there exists $\delta > 0$ such that for each $x \in X$, $f(B_{d_X}(x, \delta)) \subset B_{d_Y}(f(x), \epsilon)$. Hence f is (d_X, d_Y) -uniformly continuous. □

Definition 6.1.3.4. Let (X, d_X) and (Y, d_Y) be pseudometric spaces.

- Let $f : X \rightarrow Y$.

- We define the (d_X, d_Y) -**Lipschitz constants** of f , denoted $\text{LipConst}(f, (d_X, d_Y)) \subset \mathbb{R}$, by

$$\text{LipConst}(f, (d_X, d_Y)) := \{K \in \mathbb{R} : \text{for each } x_1, x_2 \in X, d_Y(f(x_1), f(x_2)) \leq K d_X(x_1, x_2)\}.$$

- We define the (d_X, d_Y) -**Lipschitz constant** of f , denoted $\text{Lip}(f, (d_X, d_Y)) \in \overline{\mathbb{R}}_+$, by

$$\text{Lip}(f, (d_X, d_Y)) := \inf \text{LipConst}(f, (d_X, d_Y))$$

- Then f is said to be

- * (d_X, d_Y) -**Lipschitz** if $\text{LipConst}(f, (d_X, d_Y)) \neq \emptyset$,
- * $(K, (d_X, d_Y))$ -**Lipschitz** if $K \in \text{LipConst}(f, (d_X, d_Y))$.

- Then f is said to be

Note 6.1.3.5. When the context is clear,

- we may say
 - “Lipschitz” instead of “ (d_X, d_Y) -Lipschitz”
 - “ K -Lipschitz” instead of “ $(K, (d_X, d_Y))$ -Lipschitz”
- we may write
 - $\text{LipConst}(f)$ in place of $\text{LipConst}(f, (d_X, d_Y))$,
 - $\text{Lip}(f)$ in place of $\text{Lip}(f, (d_X, d_Y))$
 - $C_{\text{Lip}}(X, Y)$ in place of $C_{\text{Lip}}((X, d_X), (Y, d_Y))$.

Exercise 6.1.3.6. Let (X, d_X) and (Y, d_Y) be pseudometric spaces and $f : X \rightarrow Y$. Then f is Lipschitz iff $\text{Lip}(f) < \infty$.

Proof. **FINISH!!!** □

Exercise 6.1.3.7. Let (X, d_X) and (Y, d_Y) be pseudometric spaces and $f : X \rightarrow Y$. If f is Lipschitz, then $\text{Lip}(f) \in \text{LipConst}(f)$.

Proof. Suppose that f is Lipschitz. Then $\text{LipConst}(f) \neq \emptyset$. Define $K_0 \in \mathbb{R}_+$ by $K_0 := \text{Lip}(f)$. Let $x_1, x_2 \in X$ and $\epsilon > 0$. Since $K_0 = \inf \text{LipConst}(f)$, there exists $K \in \text{LipConst}(f)$ such that $K < K_0 + \epsilon/(d_X(x_1, x_2) + 1)$. Then

$$\begin{aligned} d_Y(f(x_1), f(x_2)) &\leq K d_X(x_1, x_2) \\ &< \left(K_0 + \frac{\epsilon}{d_X(x_1, x_2) + 1} \right) d_X(x_1, x_2) \\ &= K_0 d_X(x_1, x_2) + \epsilon \frac{d_X(x_1, x_2)}{d_X(x_1, x_2) + 1} \\ &< K_0 d_X(x_1, x_2) + \epsilon. \end{aligned}$$

Since $\epsilon > 0$ is arbitrary, we have that for each $\epsilon > 0$, $d_Y(f(x_1), f(x_2)) < K_0 d_X(x_1, x_2) + \epsilon$. Hence $d_Y(f(x_1), f(x_2)) \leq K_0 d_X(x_1, x_2)$. Since $x_1, x_2 \in X$ are arbitrary, we have that for each $x_1, x_2 \in X$, $d_Y(f(x_1), f(x_2)) \leq K_0 d_X(x_1, x_2)$. Thus f is K_0 -Lipschitz and $K_0 \in \text{LipConst}(f)$. □

Exercise 6.1.3.8. Let (X, d_X) and (Y, d_Y) be metric spaces and $f \in C_{\text{Lip}}(X, Y)$. If $\#X \geq 2$, then

$$\text{Lip}(f) = \sup \left\{ \frac{d_Y(f(x_1), f(x_2))}{d_X(x_1, x_2)} : x_1, x_2 \in X \text{ and } x_1 \neq x_2 \right\}.$$

Proof. Suppose that $\#X \geq 2$. Define $S \subset [0, \infty)$ by

$$S := \left\{ \frac{d_Y(f(x_1), f(x_2))}{d_X(x_1, x_2)} : x_1, x_2 \in X \text{ and } x_1 \neq x_2 \right\}$$

Since $\#X \geq 2$, $S \neq \emptyset$.

- Let $s \in S$. Then there exist $x_1, x_2 \in X$ such that $x_1 \neq x_2$ and

$$s = \frac{d_Y(f(x_1), f(x_2))}{d_X(x_1, x_2)}.$$

The prev ex implies that $\text{Lip}(f) \in \text{LipConst}(f)$. Therefore

$$s \leq \text{Lip}(f).$$

Since $s \in S$ is arbitrary, we have that for each $s \in S$, $s \leq \text{Lip}(f)$. Hence $\text{Lip}(f) \in \text{ub}(S)$. Thus $\sup S \leq \text{Lip}(f)$.

- Let $x_1, x_2 \in X$.

- If $x_1 = x_2$, then

$$\begin{aligned} d_Y(f(x_1), f(x_2)) &= 0 \\ &\leq (\sup S) d_X(x_1, x_2). \end{aligned}$$

- Suppose that $x_1 \neq x_2$. Then

$$\begin{aligned} d_Y(f(x_1), f(x_2)) &= \frac{d_Y(f(x_1), f(x_2))}{d_X(x_1, x_2)} d_X(x_1, x_2) \\ &\leq (\sup S) d_X(x_1, x_2). \end{aligned}$$

Since $x_1, x_2 \in X$ are arbitrary, we have that for each $x_1, x_2 \in X$, $d_Y(f(x_1), f(x_2)) \leq (\sup S) d_X(x_1, x_2)$. Hence $\sup S \in \text{LipConst}(f)$ and

$$\begin{aligned} \text{Lip}(f) &= \min \text{LipConst}(f) \\ &\leq \sup S. \end{aligned}$$

Since $\sup S \leq \text{Lip}(f)$ and $\text{Lip}(f) \leq \sup S$, we have that $\sup S = \text{Lip}(f)$. □

Exercise 6.1.3.9. Let (X, d_X) and (Y, d_Y) be pseudometric spaces and $f : X \rightarrow Y$.

1. If f is Lipschitz, then f is uniformly continuous.
2. $C_{\text{Lip}}(X, Y) \subset C(X, Y)$.

Proof.

1. Suppose that f is Lipschitz. Let $\epsilon > 0$. Since f is Lipschitz, there exists $K > 0$ such that for each $x_1, x_2 \in X$, $d_Y(f(x_1), f(x_2)) \leq K d_X(x_1, x_2)$. Set $\delta := \epsilon/K$. Then $\delta > 0$. Let $x_1, x_2 \in X$. Suppose that $d_X(x_1, x_2) < \delta$. Then

$$\begin{aligned} d_Y(f(x_1), f(x_2)) &\leq K d_X(x_1, x_2) \\ &< K \delta \\ &= \frac{K \epsilon}{K} \\ &= \epsilon \end{aligned}$$

Since $\epsilon > 0$ is arbitrary, we have that for each $\epsilon > 0$, there exists $\delta > 0$ such that for each $x_1, x_2 \in X$, $d_X(x_1, x_2) < \delta$ implies that $d_Y(f(x_1), f(x_2)) < \epsilon$. Hence f is uniformly continuous.

2. Clear. □

Exercise 6.1.3.10. Let (X, d) be a pseudometric space, $A \subset X$. Then $d(\cdot, A)$ is 1-Lipschitz. □

Proof. Clear by Exercise 6.1.1.14. □

Exercise 6.1.3.11. Define $\phi : [0, \infty) \rightarrow [0, 1]$ by $\phi(t) := t/(1+t)$. Then ϕ is 1-Lipschitz.

Proof. Let $t_1, t_2 \in [0, \infty)$. Then

$$\begin{aligned} \phi(t_1) - \phi(t_2) &= \frac{t_1}{1+t_1} - \frac{t_2}{1+t_2} \\ &= \frac{t_1(1+t_2) - t_2(1+t_1)}{(1+t_1)(1+t_2)} \\ &= \frac{t_1 - t_2}{(1+t_1)(1+t_2)}. \end{aligned}$$

Therefore

$$\begin{aligned} |\phi(t_1) - \phi(t_2)| &= \left| \frac{t_1 - t_2}{(1+t_1)(1+t_2)} \right| \\ &= \frac{|t_1 - t_2|}{|1+t_1||1+t_2|} \\ &\leq |t_1 - t_2|. \end{aligned}$$

Since $t_1, t_2 \in [0, \infty)$ is arbitrary, we have that ϕ is Lipschitz. □

Definition 6.1.3.12. Let (X, d) be a pseudometric space and $f \in C_{\text{Lip}}(X, Y)$. Then f is said to be a (d_X, d_Y) -**contraction mapping** if $\text{Lip}(f) \in [0, 1)$.

Note 6.1.3.13. When the context is clear, we may say “contraction mapping” instead of “ (d_X, d_Y) -contraction mapping”.

Exercise 6.1.3.14. Let (X, d) be a complete metric space and $f \in C_{\text{Lip}}(X, X)$. Define $\rho \in \mathbb{R}_+$ by $\rho := \text{Lip}(f)$. Then for each $x_0 \in X$ and $n \in \mathbb{N}$,

$$d(f^n(x_0), x_0) \leq \frac{1 - \rho^n}{1 - \rho} d(x_0, f(x_0))$$

Hint: geometric series

Proof. Let $x_0 \in X$. Define $(x_n)_{n \in \mathbb{N}} \subset X$ by $x_n := f^n(x_0)$. Let $n \in \mathbb{N}$. Then

$$\begin{aligned} d(x_n, x_{n+1}) &= d(f^n(x_0), f^{n+1}(x_0)) \\ &\leq \rho^n d(x_0, x_1) \quad (\text{by make and ref ex}). \end{aligned}$$

Therefore

$$\begin{aligned} d(x_n, x_0) &\leq \sum_{j=0}^{n-1} d(x_j, x_{j+1}) \\ &\leq \sum_{j=0}^{n-1} \rho^j d(x_0, x_1) \\ &= d(x_0, x_1) \sum_{j=0}^{n-1} \rho^j \\ &= d(x_0, x_1) \frac{1 - \rho^n}{1 - \rho} \\ &= \frac{1 - \rho^n}{1 - \rho} d(x_0, x_1). \end{aligned}$$

Since $n \in \mathbb{N}$ is arbitrary, we have that for each $n \in \mathbb{N}$,

$$d(f^n(x_0), x_0) = d(x_n, x_0) \leq \frac{1 - \rho^n}{1 - \rho} d(x_0, x_1) = \frac{1 - \rho^n}{1 - \rho} d(x_0, f(x_0)).$$

Since $x_0 \in X$ is arbitrary, we have that for each $x_0 \in X$ and $n \in \mathbb{N}$,

$$d(f^n(x_0), x_0) \leq \frac{1 - \rho^n}{1 - \rho} d(x_0, f(x_0)).$$

□

6.1.4 Stuff to Reorder or Move

Definition 6.1.4.1. Let (X, d) be a metric space, $A \subset X$ and $\epsilon > 0$. We define the ϵ -**enlargement** of A , denoted A_ϵ , by

$$A_\epsilon = \{x \in X : d(x, A) < \epsilon\}$$

Exercise 6.1.4.2. Let (X, d) be a metric space, $A \subset X$ and $\epsilon > 0$. Then A_ϵ is open.

Proof. Let $x \in A_\epsilon$. By definition, $d(x, A) < \epsilon$. Set $\delta = (\epsilon - d(x, A))/2$. Then $\delta > 0$ and thus there exists $a \in A$ such that $d(x, a) < d(x, A) + \delta$. Let $y \in B(x, \delta)$. Therefore

$$\begin{aligned} d(y, A) &= \inf\{d(y, b) : b \in A\} \\ &\leq d(y, a) \\ &\leq d(y, x) + d(x, a) \\ &< \delta + d(x, A) + \delta \\ &= d(x, A) + 2\delta \\ &= d(x, A) + \epsilon - d(x, A) \\ &= \epsilon \end{aligned}$$

Hence $y \in A_\epsilon$. Since $y \in B(x, \delta)$ is arbitrary, $B(x, \delta) \subset A_\epsilon$. Since $x \in A_\epsilon$ is arbitrary, we have that for each $x \in A_\epsilon$, there exists $\delta > 0$ such that $B(x, \delta) \subset A_\epsilon$. Hence A_ϵ is open. □

Exercise 6.1.4.3. Let (X, d) be a metric space and $C, U \subset X$.

1. If C is closed, then $C = \bigcap_{n \in \mathbb{N}} C_{1/n}$.
2. If U is open, then $U = \bigcup_{n \in \mathbb{N}} [(U^c)_{1/n}]^c$.

Proof.

1. Suppose that C is closed. Since for each $n \in \mathbb{N}$, $C \subset C_{1/n}$, we have that $C \subset \bigcap_{n \in \mathbb{N}} C_{1/n}$.

For the sake of contradiction, suppose that $\bigcap_{n \in \mathbb{N}} C_{1/n} \not\subset C$. Then there exists $x \in \bigcap_{n \in \mathbb{N}} C_{1/n}$ such that $x \notin C$. Since C is closed, a previous exercise implies that $d(x, C) > 0$. Set $\epsilon = d(x, C)$. Since $\epsilon > 0$, there exists $N \in \mathbb{N}$ such that $1/N < \epsilon$. Since $x \in \bigcap_{n \in \mathbb{N}} C_{1/n}$, $x \in C_{1/N}$. Thus

$$\begin{aligned} d(x, C) &< 1/N \\ &< \epsilon \\ &= d(x, C) \end{aligned}$$

which is a contradiction. Hence $\bigcap_{n \in \mathbb{N}} C_{1/n} \subset C$. Thus $C = \bigcap_{n \in \mathbb{N}} C_{1/n}$.

2. Suppose that U is open. Then U^c is closed. The previous part implies that

$$\begin{aligned} U &= (U^c)^c \\ &= \left(\bigcap_{n \in \mathbb{N}} (U^c)_{1/n} \right)^c \\ &= \bigcup_{n \in \mathbb{N}} [(U^c)_{1/n}]^c \end{aligned}$$

□

Exercise 6.1.4.4. Let (X, d) be a metric space. Then

1. for each $U \subset X$, if U is open, then U is an F_σ set,
2. for each $C \subset X$, if C is closed, then C is a G_δ set.

Proof. Clear by [the previous exercise](#).

□

Exercise 6.1.4.5. Let (X, d) be a metric space, $(x_n)_{n \in \mathbb{N}} \subset X$, $x \in X$ and $r > 0$. If $x_n \rightarrow x$, then

1. for each $\epsilon > 0$ there exists $N \in \mathbb{N}$ such that for each $n \in \mathbb{N}$, $n \geq N$ implies that $B(x_n, r) \subset B(x, r + \epsilon)$.
2. for each $\epsilon > 0$ there exists $N \in \mathbb{N}$ such that for each $n \in \mathbb{N}$, $n \geq N$ implies that $\bar{B}(x_n, r) \subset \bar{B}(x, r + \epsilon)$.

Proof. Suppose that $x_n \rightarrow x$.

1. Let $\epsilon > 0$. Since $x_n \rightarrow x$, there exists $N \in \mathbb{N}$ such that for each $n \in \mathbb{N}$, $n \geq N$ implies that $d(x_n, x) < \epsilon$. Let $n \in \mathbb{N}$. Suppose that $n \geq N$. Let $y \in B(x_n, r)$. Then

$$\begin{aligned} d(y, x) &\leq d(y, x_n) + d(x_n, x) \\ &< r + \epsilon \end{aligned}$$

Thus $y \in B(x, r + \epsilon)$. Since $y \in B(x_n, r)$ is arbitrary, $B(x_n, r) \subset B(x, r + \epsilon)$.

2. Similar to (1).

□

Exercise 6.1.4.6. Let (X, d) be a pseudometric space and $E \subset X$. Then E is dense in $(X, \mathcal{T}_d)$ iff for each $x \in X$ and $\epsilon > 0$, there exists $x' \in E$ such that $d(x, x') < \epsilon$.

Proof.

• ($\implies$):

Suppose that E is dense in $(X, \mathcal{T})$. Let $x \in X$ and $\epsilon > 0$. Since $B(x, \epsilon) \in \mathcal{T}_d$ and $B(x, \epsilon) \neq \emptyset$, Exercise 4.1.0.41 implies that there exists $x' \in E$ such that $x' \in B(x, \epsilon)$. Then $d(x, x') < \epsilon$. Since $x \in X$ and $\epsilon > 0$ are arbitrary, we have that for each $x \in X$ and $\epsilon > 0$, there exists $x' \in E$ such that $d(x, x') < \epsilon$.

• ($\impliedby$):

Conversely, suppose that for each $x \in X$ and $\epsilon > 0$, there exists $x' \in E$ such that $d(x, x') < \epsilon$. Let $U \in \mathcal{T}_d$. Suppose that $U \neq \emptyset$. Then there exists $x \in X$ such that $x \in U$. Exercise 6.1.2.6 implies that there exists $\delta > 0$ such that $B_d(x, \delta) \subset U$. By assumption, there exists $x' \in E$ such that $d(x, x') < \delta$. Then

$$\begin{aligned} x' &\in E \cap B_d(x, \delta) \\ &\subset E \cap U. \end{aligned}$$

Hence $E \cap U \neq \emptyset$. Since $U \in \mathcal{T}_d$ with $U \neq \emptyset$ is arbitrary, we have that for each $U \in \mathcal{T}_d$, $U \neq \emptyset$ implies that $E \cap U \neq \emptyset$. Exercise 4.1.0.41 then implies that E is dense in $(X, \mathcal{T}_d)$.

□

Exercise 6.1.4.7. *needs to go after discussion of metric spaces being Hausdorff* Let (X, d) be a metric space and $U \in \mathcal{T}_d$. Define $C \subset X \times \mathbb{R}$ and $f : U \rightarrow C$ by $C := \{(x, d(x, U^c)^{-1}) : x \in U\}$ and $f(x) := (x, d(x, U^c)^{-1})$. Then

1. f is a homeomorphism,
2. C is closed in $(X \times \mathbb{R}, \mathcal{T}_d \otimes \mathcal{T}_{\mathbb{R}})$.

Proof.

1.
 - Exercise 6.1.3.10 and Exercise 6.1.2.15 imply that f is well-defined and continuous.
 - We note that $\text{proj}_1|_C \circ f = \text{id}_X$ and $f \circ \text{proj}_1|_C = \text{id}_C$. Thus f is a bijection with $f^{-1} = \text{proj}_1|_C$.
 - Since $\text{proj}_1 : X \times \mathbb{R} \rightarrow X$ is continuous, $f^{-1} = \text{proj}_1|_C$ is continuous. Hence f is a homeomorphism.
2. Let $(y_n)_{n \in \mathbb{N}} \subset C$ and $y \in X \times \mathbb{R}$. Suppose that $y_n \rightarrow y$. Since $C = f(U)$, there exist $(x_n)_{n \in \mathbb{N}} \subset U$ such that for each $n \in \mathbb{N}$, $y_n = f(x_n)$. Since $y \in X \times \mathbb{R}$, there exists $x \in X$ and $a \in \mathbb{R}$ such that $y = (x, a)$. Since $y_n \rightarrow y$, we have that

$$\begin{aligned} (x_n, d(x_n, U^c)^{-1}) &= y_n \\ &\rightarrow y \\ &= (x, a). \end{aligned}$$

Hence $x_n \rightarrow x$ and $d(x_n, U^c)^{-1} \rightarrow a$. For the sake of contradiction, suppose that $x \in U^c$. Since $x_n \rightarrow x$,

$$\begin{aligned} d(x_n, U^c) &\leq d(x_n, x) \\ &\rightarrow 0. \end{aligned}$$

Therefore

$$\begin{aligned} d(x_n, U^c)^{-1} &\rightarrow \infty \\ &\neq a. \end{aligned}$$

This is a contradiction. Hence $x \in U$. By continuity,

$$\begin{aligned} (x_n, d(x_n, U^c)^{-1}) &= f(x_n) \\ &\rightarrow f(x) \\ &= (x, d(x, U^c)^{-1}). \end{aligned}$$

In particular, $d(x_n, U^c)^{-1} \rightarrow d(x, U^c)^{-1}$. Since d is a metric, $(X, \mathcal{T}_d)$ is Hausdorff. Since $d(x_n, U^c)^{-1} \rightarrow a$ and $d(x_n, U^c)^{-1} \rightarrow d(x, U^c)^{-1}$, we have that $a = d(x, U^c)^{-1}$. Therefore

$$\begin{aligned} y &= (x, a) \\ &= (x, d(x, U^c)^{-1}) \\ &= f(x) \\ &\in f(U) \\ &= C. \end{aligned}$$

Since $(y_n)_{n \in \mathbb{N}} \subset C$ and $y \in X \times \mathbb{R}$ with $y_n \rightarrow y$ are arbitrary, we have that for each $(y_n)_{n \in \mathbb{N}} \subset C$ and $y \in X \times \mathbb{R}$, $y_n \rightarrow y$ implies that $y \in C$. Thus C is closed in $X \times \mathbb{R}$.

□

Exercise 6.1.4.8. *needs to go after discussion of metric spaces being Hausdorff* Let (X, d) be a metric space and $(U_n)_{n \in \mathbb{N}} \subset \mathcal{T}_d$. Define $E \subset X$, $C \subset X \times \mathbb{R}^{\mathbb{N}}$ and $f : E \rightarrow C$ by

$$E := \bigcap_{n \in \mathbb{N}} U_n, \quad C := \{(x, d(x, U_1^c)^{-1}, d(x, U_2^c)^{-1}, \dots) : x \in X\} \quad \text{and} \quad f(x) := (x, d(x, U_1^c)^{-1}, d(x, U_2^c)^{-1}, \dots).$$

Then

1. f is a homeomorphism,
2. C is closed in $X \times \mathbb{R}^{\mathbb{N}}$.

Proof.

1.
 - Exercise 6.1.3.10 and Exercise 6.1.2.15 imply that f is well-defined and continuous.
 - We note that $\text{proj}_1|_C \circ f = \text{id}_X$ and $f \circ \text{proj}_1|_C = \text{id}_C$. Thus f is a bijection with $f^{-1} = \text{proj}_1|_C$.
 - Since $\text{proj}_1 : X \times \mathbb{R} \rightarrow X$ is continuous, $f^{-1} = \text{proj}_1|_C$ is continuous. Hence f is a homeomorphism.
2. Let $(y_n)_{n \in \mathbb{N}} \subset C$ and $y \in X \times \mathbb{R}^{\mathbb{N}}$. Suppose that $y_n \rightarrow y$. Since $C = f(U)$, there exist $(x_n)_{n \in \mathbb{N}} \subset U$ such that for each $n \in \mathbb{N}$, $y_n = f(x_n)$. Since $y \in X \times \mathbb{R}^{\mathbb{N}}$, there exists $x \in X$ and $a \in \mathbb{R}^{\mathbb{N}}$ such that $y = (x, a)$. Since $y_n \rightarrow y$, we have that

$$\begin{aligned} (x_n, d(x_n, U_1^c)^{-1}, d(x_n, U_2^c)^{-1}, \dots) &= y_n \\ &\rightarrow y \\ &= (x, a_1, a_2, \dots). \end{aligned}$$

Hence $x_n \rightarrow x$ and for each $k \in \mathbb{N}$ $d(x_n, U_k^c)^{-1} \xrightarrow{n} a_k$. For the sake of contradiction, suppose that $x \in E^c$. Since

$$\begin{aligned} E^c &= \left(\bigcap_{k \in \mathbb{N}} U_k \right)^c \\ &= \bigcup_{k \in \mathbb{N}} U_k^c \end{aligned}$$

there exists $k_0 \in \mathbb{N}$ such that $x \in U_{k_0}^c$. Since $x_n \rightarrow x$,

$$\begin{aligned} d(x_n, U_{k_0}^c) &\leq d(x_n, x) \\ &\rightarrow 0. \end{aligned}$$

Therefore

$$\begin{aligned} d(x_n, U_{k_0}^c)^{-1} &\rightarrow \infty \\ &\neq a_{k_0}. \end{aligned}$$

This is a contradiction. Hence $x \in E$. By continuity,

$$\begin{aligned} (x_n, d(x_n, U_1^c)^{-1}, d(x_n, U_2^c)^{-1}, \dots) &= f(x_n) \\ &\rightarrow f(x) \\ &= (x, d(x, U_1^c)^{-1}, d(x, U_2^c)^{-1}, \dots). \end{aligned}$$

In particular, for each $k \in \mathbb{N}$, $d(x_n, U_k^c)^{-1} \xrightarrow{n} d(x, U_k^c)^{-1}$. Since d is a metric, $(X, \mathcal{T}_d)$ is Hausdorff. Since for each $k \in \mathbb{N}$, $k \in \mathbb{N}$ $d(x_n, U_k^c)^{-1} \xrightarrow{n} a_k$ and $d(x_n, U_k^c)^{-1} \xrightarrow{n} d(x, U_k^c)^{-1}$, we have that for each $k \in \mathbb{N}$, $a_k = d(x, U_k^c)^{-1}$. Therefore

$$\begin{aligned} y &= (x, a_1, a_2, \dots) \\ &= (x, d(x, U_1^c)^{-1}, d(x, U_2^c)^{-1}, \dots) \\ &= f(x) \\ &\in f(U) \\ &= C. \end{aligned}$$

Since $(y_n)_{n \in \mathbb{N}} \subset C$ and $y \in X \times \mathbb{R}^{\mathbb{N}}$ with $y_n \rightarrow y$ are arbitrary, we have that for each $(y_n)_{n \in \mathbb{N}} \subset C$ and $y \in X \times \mathbb{R}$, $y_n \rightarrow y$ implies that $y \in C$. Thus C is closed in $X \times \mathbb{R}$.

□

6.1.5 Isometries

CHANGE TO PSEUDOMETRIC SPACES????!!

FIX!!!

move pullback stuff to intro??

Definition 6.1.5.1. Let (X, d_X) and (Y, d_Y) be metric spaces and $f : X \rightarrow Y$. Then f is said to be a (d_X, d_Y) -**isometry** if for each $x_1, x_2 \in X$, $d_Y(f(x_1), f(x_2)) = d_X(x_1, x_2)$.

Exercise 6.1.5.2. Let (X, d_X) and (Y, d_Y) be metric spaces and $f : X \rightarrow Y$. If f is a (d_X, d_Y) -isometry, then f is injective.

Proof. Suppose that f is an (d_X, d_Y) -isometry. Let $x_1, x_2 \in X$. Suppose that $f(x_1) = f(x_2)$. Then

$$\begin{aligned} d_X(x_1, x_2) &= d_Y(f(x_1), f(x_2)) \\ &= 0. \end{aligned}$$

Hence $x_1 = x_2$. Since $x_1, x_2 \in X$ are arbitrary, we have that for each $x_1, x_2 \in X$, $f(x_1) = f(x_2)$ implies that $x_1 = x_2$. Thus f is injective. $\square$

Definition 6.1.5.3. Let X a set, (Y, d) be a metric space and $f : X \rightarrow Y$ an injection. We define the **pullback of d by f** , denoted $f^*d : X \times X \rightarrow [0, \infty)$, by

$$f^*d(x_1, x_2) := d(f(x_1), f(x_2))$$

Exercise 6.1.5.4. Let X a set, (Y, d) be a metric space and $f : X \rightarrow Y$ an injection. Then f^*d is a metric on X .

Proof. Let $x_1, x_2, x_3 \in X$. Then

1.

$$\begin{aligned} f^*d(x_1, x_2) &= d(f(x_1), f(x_2)) \\ &= d(f(x_2), f(x_1)) \\ &= f^*d(x_2, x_1) \end{aligned}$$

2.

$$\begin{aligned} f^*d(x_1, x_2) = 0 &\iff d(f(x_1), f(x_2)) = 0 \\ &\iff f(x_1) = f(x_2) \\ &\iff x_1 = x_2 \end{aligned}$$

3.

$$\begin{aligned} f^*d(x_1, x_3) &= d(f(x_1), f(x_3)) \\ &\leq d(f(x_1), f(x_2)) + d(f(x_2), f(x_3)) \\ &= f^*d(x_1, x_2) + f^*d(x_2, x_3) \end{aligned}$$

So f^*d is a metric on X . $\square$

Exercise 6.1.5.5. Let X a set, (Y, d) be a metric space and $f : X \rightarrow Y$ an injection. Then f is an (f^*d, d) -isometry.

Proof. Let $x_1, x_2 \in X$. Then

$$d(f(x_1), f(x_2)) = f^*d(x_1, x_2).$$

Since $x_1, x_2 \in X$ are arbitrary, we have that f is (f^*d, d) -isometry. $\square$

Exercise 6.1.5.6. Let $(X, d_X), (Y, d_Y)$ be metric spaces and $f : X \rightarrow Y$. Then f is a (d_X, d_Y) -isometry iff f is injective and $d_X = f^*d_Y$.

Proof.

• $(\implies)$:

Suppose that f is a (d_X, d_Y) -isometry. Exercise 6.1.5.2 implies that f is injective. Then for each $x_1, x_2 \in X$, we have that

$$\begin{aligned} d_X(x_1, x_2) &= d_Y(f(x_1), f(x_2)) \\ &= f^*d_Y(x_1, x_2) \end{aligned}$$

Since $x_1, x_2 \in X$ are arbitrary, we have that $d_X = f^*d_Y$.

• $(\impliedby)$:

Conversely, suppose that f is injective and $d_X = f^*d_Y$. Exercise 6.1.5.6 implies that f is a (f^*d_Y, d_Y) -isometry. Since $d_X = f^*d_Y$, we have that f is a (d_X, d_Y) -isometry. □

Exercise 6.1.5.7. Let $(X, d_X), (Y, d_Y)$ be metric spaces and $f : X \rightarrow Y$. Suppose that f is a (d_X, d_Y) -isometry and f is a bijection. Then f^{-1} is a (d_Y, d_X) -isometry.

Proof. Since f^{-1} is a bijection, f^{-1} is injective. Let $y_1, y_2 \in Y$. Exercise 6.1.5.6 implies that $f^*d_Y = d_X$. Set $x_1 := f^{-1}(y_1)$ and $x_2 := f^{-1}(y_2)$. Then

$$\begin{aligned} (f^{-1})^*d_X(y_1, y_2) &= d_X(f^{-1}(y_1), f^{-1}(y_2)) \\ &= d_X(x_1, x_2) \\ &= f^*d_Y(x_1, x_2) \\ &= d_Y(f(x_1), f(x_2)) \\ &= d_Y(y_1, y_2). \end{aligned}$$

Since $y_1, y_2 \in Y$ are arbitrary, we have that for each $y_1, y_2 \in Y$, $(f^{-1})^*d_X(y_1, y_2) = d_Y(y_1, y_2)$. Hence $d_Y = (f^{-1})^*d_X$. Exercise 6.1.5.6 then implies that f^{-1} is a (d_Y, d_X) -isometry. □

Exercise 6.1.5.8. can we just assume bijection and show homeo iff $\mathcal{T}_Y = \mathcal{T}_{(f^{-1})^*d_X}$??? Let $(X, \mathcal{T}_X)$ and $(Y, \mathcal{T}_Y)$ be topological spaces, $d_X : X \times X \rightarrow [0, \infty)$ a metric on X and $f : X \rightarrow Y$ a $(\mathcal{T}_X, \mathcal{T}_Y)$ -homeomorphism. Then $\mathcal{T}_Y = \mathcal{T}_{(f^{-1})^*d_X}$.

Proof. Set $d_Y := (f^{-1})^*d_X$. Let $(y_n)_{n \in \mathbb{N}} \subset Y$ and $y \in Y$. Define $(x_n)_{n \in \mathbb{N}} \subset X$ and $x \in X$ by $x_n := f^{-1}(y_n)$ and $x := f^{-1}(y)$.

• Suppose that $y_n \rightarrow y$ in $(Y, \mathcal{T}_Y)$. Since f^{-1} is a $(\mathcal{T}_Y, \mathcal{T}_X)$ -homeomorphism,

$$\begin{aligned} x_n &= f^{-1}(y_n) \\ &\rightarrow f^{-1}(y) \\ &= x \end{aligned}$$

in $(X, \mathcal{T}_X)$. Since $\mathcal{T}_X = \mathcal{T}_{d_X}$, we have that

$$\begin{aligned} d_Y(y_n, y) &= (f^{-1})^*d_X(y_n, y) \\ &= d_X(f^{-1}(y_n), f^{-1}(y)) \\ &= d_X(x_n, x) \\ &\rightarrow 0. \end{aligned}$$

Thus $y_n \rightarrow y$ in $(X, \mathcal{T}_{d_Y})$.

- Conversely, suppose that $y_n \rightarrow y$ in $(Y, \mathcal{T}_{d_Y})$. Then

$$\begin{aligned} d_X(x_n, x) &= d_X(f^{-1}(y_n), f^{-1}(y)) \\ &= (f^{-1})^* d_X(y_n, y) \\ &= d_Y(y_n, y) \\ &\rightarrow 0. \end{aligned}$$

Hence $x_n \rightarrow x$ in $(X, \mathcal{T}_{d_X})$. Since $\mathcal{T}_{d_X} = \mathcal{T}_X$ and f is a $(\mathcal{T}_X, \mathcal{T}_Y)$ -homeomorphism, we have that

$$\begin{aligned} y_n &= f(x_n) \\ &\rightarrow f(x) \\ &= y \end{aligned}$$

in $(Y, \mathcal{T}_Y)$.

Since $(y_n)_{n \in \mathbb{N}}$ and $y \in Y$ are arbitrary, we have that for each $(y_n)_{n \in \mathbb{N}}$ and $y \in Y$, $y_n \rightarrow y$ in $(Y, \mathcal{T}_Y)$ iff $y_n \rightarrow y$ in $(Y, \mathcal{T}_{d_Y})$. Exercise 4.3.2.16 implies that $\mathcal{T}_Y = \mathcal{T}_{d_Y}$. $\square$

6.2 Countability and Separation

move this section right after metric topology section

6.2.1 Countability

include first-countability exercise in this section

Exercise 6.2.1.1. Let (X, d) be a metric space. Then $(X, \mathcal{T}_d)$ is second-countable iff $(X, \mathcal{T}_d)$ is separable.

Proof.

- $(\implies)$:
Suppose that $(X, \mathcal{T}_d)$ is second-countable. Exercise 4.13.2.6 implies that $(X, \mathcal{T}_d)$ is separable.
- $(\impliedby)$:
Conversely, suppose that $(X, \mathcal{T}_d)$ is separable. Then there exists $(x_n)_{n \in \mathbb{N}} \subset X$ such that $(x_n)_{n \in \mathbb{N}}$ is dense in X . Define $\mathcal{B} := \{B(x_n, q) : (n, q) \in \mathbb{N} \times \mathbb{Q} \cap (0, 1)\}$. Then $\mathcal{B}$ is countable.
Let $U \in \mathcal{T}_d$ and $x \in U$. Since U is open, Exercise 6.1.2.6 implies that there exists $\delta_0 > 0$ such that $B(x, \delta_0) \subset U$. Set $\delta := \delta_0/2$. Since $B(x, \delta)$ is open, $B(x, \delta) \neq \emptyset$ and $(x_n)_{n \in \mathbb{N}}$ is dense in X , Exercise 4.1.0.41 implies that there exists $n \in \mathbb{N}$ such that $x_n \in B(x, \delta)$. Set $B := B(x_n, \delta)$. By construction $B \in \mathcal{B}$ and $x \in B$. Let $y \in B$. Then

$$\begin{aligned} d(x, y) &\leq d(x, x_n) + d(x_n, y) \\ &< \delta + \delta \\ &= 2\delta \\ &= \delta_0 \end{aligned}$$

Thus

$$\begin{aligned} y &\in B(x, \delta_0) \\ &\subset U \end{aligned}$$

Since $y \in B$ is arbitrary, we have that $B \subset U$. Therefore, there exists $B \in \mathcal{B}$ such that

$$\begin{aligned} x &\in B \\ &\subset U. \end{aligned}$$

Hence $\mathcal{B}$ is a basis for $\mathcal{T}_d$. Since $\mathcal{B}$ is countable, $(X, \mathcal{T}_d)$ is second-countable. □

Exercise 6.2.1.2. Let (X, d_X) and (Y, d_Y) be metric spaces and $f : X \rightarrow Y$ a $(\mathcal{T}_{d_X}, \mathcal{T}_{d_Y})$ -homeomorphism. Then $(X, \mathcal{T}_{d_X})$ is separable iff $(Y, \mathcal{T}_{d_Y})$ is separable.

Proof.

- $(\implies)$:
Suppose that $(X, \mathcal{T}_{d_X})$ is separable. Exercise 6.2.1.1 implies that $(X, \mathcal{T}_{d_X})$ is second-countable. Exercise 4.13.2.7 implies that $(Y, \mathcal{T}_{d_Y})$ is second-countable. Another application of Exercise 6.2.1.1 implies that $(Y, \mathcal{T}_{d_Y})$ is separable.
- $(\impliedby)$:
Similar to $(\implies)$. □

Exercise 6.2.1.3. Let (X, d) be a metric space. Suppose that $X \neq \emptyset$. If $(X, \mathcal{T}_d)$ is separable, then for each $n \in \mathbb{N}$, there exists $\mathcal{B} \subset \{\bar{B}(x, \delta) : x \in X, \delta > 0\}$ such that $\mathcal{B}$ is countable, $\mathcal{B}$ is disjoint, $X = \bigcup_{B \in \mathcal{B}} B$ and for each $B \in \mathcal{B}$, $\text{diam}(B) \leq 1/n$.

Proof. (find out how to rigorously handle these types of proofs where objects are defined inductively and the process may or may not terminate, i.e. proofs that are essentially a while-loop)

Suppose that $(X, \mathcal{T}_d)$ is separable. Let $n \in \mathbb{N}$. Since $(X, \mathcal{T}_d)$ is separable, there exists $S \subset X$ such that S is countable and S is dense in X . Since $X \neq \emptyset$, we have that $S \neq \emptyset$.

- Suppose that $\#S = 1$. Write $S = \{x\}$. Define $\mathcal{B} := \{\bar{B}(x, 1/2n)\}$. Then $\mathcal{B}$ is countable, $\mathcal{B}$ is disjoint, $X = \bigcup_{B \in \mathcal{B}} B$ and for each $B \in \mathcal{B}$, $\text{diam}(B) \leq 1/n$.
- Suppose that $\#S \in \mathbb{N} \setminus \{1\}$. Write $S = \{x_j\}_{j \in [J]}$ where $j_1 \neq j_2$ implies $x_{j_1} \neq x_{j_2}$. Define $r > 0$, $(r_j)_{j \in [J]} \in (0, \infty)^{[J]}$ and $\mathcal{B} \subset \mathcal{T}_d$ by $r := \min\{d(x, y) : x, y \in S \text{ and } x \neq y\}$, $r_j := 2^{-1} \min\{r, n^{-1}\}$ and $\mathcal{B} := \{\bar{B}(x_j, r_j)\}_{j \in [J]}$. Then $\mathcal{B}$ is countable, $\mathcal{B}$ is disjoint, $X = \bigcup_{B \in \mathcal{B}} B$ and for each $B \in \mathcal{B}$, $\text{diam}(B) \leq 1/n$.
- Suppose that $\#S = \infty$. Write $S = \{x_j\}_{j \in \mathbb{N}}$ where $j_1 \neq j_2$ implies $x_{j_1} \neq x_{j_2}$. We define $\mathcal{B} \subset \{\bar{B}(x, \delta) : x \in X, \delta > 0\}$ inductively as follows:
 - We define $j_1 := 1$, $r_1 := (2n)^{-1}$ and $B_1 := \bar{B}(x_{j_1}, r_1)$.
 - For $k \geq 2$, define $J_k \subset \mathbb{N}$ by $J_k := \left\{ j \in \mathbb{N} : x_j \notin \bigcup_{k' \in [k-1]} \bar{B}(x_{j_{k'}}, r_{k'}) \right\}$.
 - * If $J_k \neq \emptyset$, define $j_k := \min J_k$. Since $\left[\bigcup_{k' \in [k-1]} \bar{B}(x_{j_{k'}}, r_{k'}) \right]^c$ is open and $x_{j_k} \in \left[\bigcup_{k' \in [k-1]} \bar{B}(x_{j_{k'}}, r_{k'}) \right]^c$, there exists $r_k > 0$ such that $r_k \leq (2n)^{-1}$ and $B(x_{j_k}, r_k) \subset \left[\bigcup_{k' \in [k-1]} \bar{B}(x_{j_{k'}}, r_{k'}) \right]^c$. Define $B_k := \bar{B}(x_{j_k}, r_k)$.
 - * If $J_k = \emptyset$, set $\mathcal{B} := \{B_{k'}\}_{k' \in [k-1]}$. Then $\mathcal{B}$ is countable, $\mathcal{B}$ is disjoint, $X = \bigcup_{B \in \mathcal{B}} B$ and for each $B \in \mathcal{B}$, $\text{diam}(B) \leq 1/n$.

Define $\mathcal{B} := \{B_k\}_{k \in \mathbb{N}}$. Then $\mathcal{B}$ is countable, $\mathcal{B}$ is disjoint, $X = \bigcup_{B \in \mathcal{B}} B$ and for each $B \in \mathcal{B}$, $\text{diam}(B) \leq 1/n$.

Since $n \in \mathbb{N}$ is arbitrary, we have that for each $n \in \mathbb{N}$, there exists $\mathcal{B} \subset \{\bar{B}(x, \delta) : x \in X, \delta > 0\}$ such that $\mathcal{B}$ is countable, $\mathcal{B}$ is disjoint, $X = \bigcup_{B \in \mathcal{B}} B$ and for each $B \in \mathcal{B}$, $\text{diam}(B) \leq 1/n$. $\square$

Definition 6.2.1.4.

- for $n \in \mathbb{N}$ and $\gamma \in \mathbb{N}^n$, we define the **length** of γ , denoted $|\gamma|$, by $|\gamma| := n$,
- for $n \in \mathbb{N}$, we define the **projection of $\mathbb{N}^{[n+1]}$ onto $\mathbb{N}^n$** , denoted $\pi_{[n]}^{[n+1]} : \mathbb{N}^{n+1} \rightarrow \mathbb{N}^n$, by $\pi_{[n]}^{[n+1]}(\gamma_1, \dots, \gamma_{n+1}) := (\gamma_1, \dots, \gamma_n)$.

Exercise 6.2.1.5. Let (X, d) be a metric space. Suppose that $(X, \mathcal{T}_d)$ is separable and $X \neq \emptyset$. Define $\Gamma := \bigcup_{n \in \mathbb{N}} \mathbb{N}^n$. Then there exists $(C_\gamma)_{\gamma \in \Gamma} \subset \mathcal{P}(X)$ such that

1. for each $\gamma \in \Gamma$,
 - (a) C_γ is closed and $C_\gamma \neq \emptyset$,
 - (b) $\text{diam } C_\gamma \leq |\gamma|^{-1}$,
 - (c) for each $n \in \mathbb{N}$, $|\gamma| = n$ implies that $C_\gamma = \bigcup_{\gamma' \in (\pi_{[n]}^{[n+1]})^{-1}(\{\gamma\})} C_{\gamma'}$
2. $X = \bigcup_{|\gamma|=1} C_\gamma$

Proof. Since X is separable, there exists $(x_j^0)_{j \in \mathbb{N}} \subset X$ such that $(x_j^0)_{j \in \mathbb{N}}$ is dense in X . We define $(C_\gamma)_{\gamma \in \Gamma}$ inductively as follows:

- For $\gamma \in \mathbb{N}$, we define $C_\gamma := \bar{B}(x_\gamma^0, 2^{-1})$.

- Let $n \geq 2$ and $\gamma \in \mathbb{N}^n$. Define $\tilde{\gamma} \in \mathbb{N}^{n-1}$ by $\tilde{\gamma} := \pi_{[n-1]}^{[n]}(\gamma)$. Since $(x_j^0)_{j \in \mathbb{N}}$ is dense in X and $C_{\tilde{\gamma}} \neq \emptyset$, we have that $\#\{j \in \mathbb{N} : \bar{B}(x_j^0, (2n)^{-1}) \cap C_{\tilde{\gamma}} \neq \emptyset\} = \infty$. Define $(x_j^{\tilde{\gamma}})_{j \in \mathbb{N}} \subset X$ to be an enumeration of $\{x_j^0 : j \in \mathbb{N} \text{ and } \bar{B}(x_j^0, (2n)^{-1}) \cap C_{\tilde{\gamma}} \neq \emptyset\}$. Define $C_\gamma \subset C_{\tilde{\gamma}}$ by $C_\gamma := \bar{B}(x_{\gamma_n}^{\tilde{\gamma}}, (2n)^{-1}) \cap C_{\tilde{\gamma}}$.

1. Let $\gamma \in \Gamma$.

- (a) By construction, C_γ is closed and $C_\gamma \neq \emptyset$.
- (b) Set $n := |\gamma|$.
 - If $n = 1$, then

$$\begin{aligned} \text{diam } C_\gamma &= \text{diam } \bar{B}(x_{\gamma_1}^0, 2^{-1}) \\ &\leq 1 \\ &= |\gamma|^{-1}. \end{aligned}$$

- Suppose that $n \geq 2$. Set $\tilde{\gamma} := \pi_{[n-1]}^{[n]}(\gamma)$. Then

$$\begin{aligned} \text{diam } C_\gamma &= \text{diam } \bar{B}(x_{\gamma_n}^{\tilde{\gamma}}, (2n)^{-1}) \cap C_{\tilde{\gamma}} \\ &\leq \text{diam } \bar{B}(x_{\gamma_n}^{\tilde{\gamma}}, (2n)^{-1}) \\ &\leq n^{-1} \\ &= |\gamma|^{-1}. \end{aligned}$$

(c) Let $n \in \mathbb{N}$. Suppose that $|\gamma| = n$.

- By construction, for each $\gamma' \in (\pi_{[n]}^{[n+1]})^{-1}(\{\gamma\})$, $C_{\gamma'} \subset C_\gamma$. Hence

$$\bigcup_{\gamma' \in (\pi_{[n]}^{[n+1]})^{-1}(\{\gamma\})} C_{\gamma'} \subset C_\gamma.$$

Let $x \in C_\gamma$. Since $(x_j^0)_{j \in \mathbb{N}}$ is dense in X , there exists $k \in \mathbb{N}$ such that $d(x_k^0, x) \leq [2(n+1)]^{-1}$. Thus

$$\begin{aligned} \emptyset &\neq \{x\} \\ &\subset \bar{B}(x_k^0, [2(n+1)]^{-1}) \cap C_\gamma \end{aligned}$$

and by construction, there exists $j \in \mathbb{N}$ such that $x_j^\gamma = x_k^0$. Define $\gamma' \in (\pi_{[n]}^{[n+1]})^{-1}(\{\gamma\})$ by $\gamma' := (\gamma_1, \dots, \gamma_n, j)$. Then

$$\begin{aligned} x &\in \bar{B}(x_j^\gamma, [2(n+1)]^{-1}) \cap C_\gamma \\ &= \bar{B}(x_{\gamma'_{n+1}}^\gamma, [2(n+1)]^{-1}) \cap C_\gamma \\ &= C_{\gamma'} \\ &\subset \bigcup_{\gamma' \in (\pi_{[n]}^{[n+1]})^{-1}(\{\gamma\})} C_{\gamma'} \end{aligned}$$

Since $x \in C_\gamma$ is arbitrary, we have that for each $x \in C_\gamma$, $x \in \bigcup_{\gamma' \in (\pi_{[n]}^{[n+1]})^{-1}(\{\gamma\})} C_{\gamma'}$. Hence $C_\gamma \subset \bigcup_{\gamma' \in (\pi_{[n]}^{[n+1]})^{-1}(\{\gamma\})} C_{\gamma'}$.

Since $\bigcup_{\gamma' \in (\pi_{[n]}^{[n+1]})^{-1}(\{\gamma\})} C_{\gamma'} \subset C_\gamma$ and $C_\gamma \subset \bigcup_{\gamma' \in (\pi_{[n]}^{[n+1]})^{-1}(\{\gamma\})} C_{\gamma'}$, we have that $C_\gamma = \bigcup_{\gamma' \in (\pi_{[n]}^{[n+1]})^{-1}(\{\gamma\})} C_{\gamma'}$.

2. By construction, since $(x_j^0)_{j \in \mathbb{N}}$ is dense in X ,

$$\begin{aligned} X &= \bigcup_{j \in \mathbb{N}} \bar{B}(x_j^0, 2^{-1}) \\ &= \bigcup_{|\gamma|=1} \bar{B}(x_{\gamma_1}^0, 2^{-1}) \\ &= \bigcup_{|\gamma|=1} C_\gamma. \end{aligned}$$



6.2.2 Separation

discuss kolmogorov quotients of pseudometric spaces and a pseudometric space (X, d) being T_0 iff d is a metric
discuss metric spaces being Hausdorff

6.3 Top-Equivalent Metrics

6.3.1 Subadditive Functions

Exercise 6.3.1.1. move elsewhere Let $\phi : [0, \infty) \rightarrow [0, \infty)$. Suppose that

1. ϕ is continuous
2. $\phi^{-1}(\{0\}) = \{0\}$

Then for each $(s_n)_{n \in \mathbb{N}} \subset [0, \infty)$, $s_n \rightarrow 0$ iff $\phi(s_n) \rightarrow 0$.

6.3.2 Top Metric Preserving Functions

account for pseudometrics?

Definition 6.3.2.1. Let X be a set, $d_1, d_2 : X \times X \rightarrow [0, \infty)$ metrics on X . Then d_1 and d_2 are said to be **Top-equivalent**, denoted $d_1 \sim_{\text{Top}} d_2$, if $\mathcal{T}_{d_1} = \mathcal{T}_{d_2}$.

Exercise 6.3.2.2. Let X be a set, $d_1, d_2 : X \times X \rightarrow [0, \infty)$ metrics on X . Then d_1 and d_2 are **Top**-equivalent iff for each $(x_n)_{n \in \mathbb{N}} \subset X$ and $x \in X$, $x_n \xrightarrow{d_1} x$ iff $x_n \xrightarrow{d_2} x$.

Proof. FINISH!!! □

Definition 6.3.2.3. Let $\phi : [0, \infty) \rightarrow [0, \infty)$. Then ϕ is said to be **Top metric-preserving** if for each set X and metric d on X ,

1. $\phi \circ d$ is a metric on X
2. $\phi \circ d \sim_{\text{Top}} d$

Definition 6.3.2.4. Let (X, d) be a metric space and $\phi : [0, \infty) \rightarrow [0, \infty)$. Suppose that ϕ is said to be **Top** metric-preserving. We define the ϕ -iterate of d , denoted d^ϕ , by $d^\phi = \phi \circ d$.

Proof.

- ($\implies$):
Let $(s_n)_{n \in \mathbb{N}} \subset [0, \infty)$. Suppose that $s_n \rightarrow 0$. Since ϕ is continuous,

$$\begin{aligned} \phi(s_n) &\rightarrow \phi(0) \\ &= 0 \end{aligned}$$

- ($\impliedby$):

- make exercise about subnets or subsequences to establish $x_\alpha \not\rightarrow x$ iff there exists $U \in \mathcal{N}(x)$ and subnet x_{α_β} such that $(x_{\alpha_\beta})_\beta \subset U^c$

Conversely, suppose that $\phi(s_n) \rightarrow 0$. For the sake of contradiction, suppose that $s_n \not\rightarrow 0$. Then there exists $\epsilon > 0$ and a subsequence $(s_{n_k})_{k \in \mathbb{N}} \subset (s_n)_{n \in \mathbb{N}}$ such that $(s_{n_k})_{k \in \mathbb{N}} \subset [\epsilon, \infty)$. Define $s \in [\epsilon, \infty)$ by $s := \liminf_{k \rightarrow \infty} s_{n_k}$. Then there exists a subsequence $(s_{n_{k_j}})_{j \in \mathbb{N}}$ such that $s_{n_{k_j}} \rightarrow s$. Define $(t_j)_{j \in \mathbb{N}} \subset [\epsilon, \infty)$ by $t_j := s_{n_{k_j}}$. Since ϕ is continuous, $\phi(t_j) \rightarrow \phi(s)$. Since $(\phi(t_j))_{j \in \mathbb{N}}$ is a subsequence of $(\phi(s_n))_{n \in \mathbb{N}}$ and $\phi(s_n) \rightarrow 0$, we have that $\phi(t_j) \rightarrow 0$. Hence $\phi(s) = 0$. Then

$$\begin{aligned} s &\in \phi^{-1}(\{0\}) \\ &= \{0\} \end{aligned}$$

and $s = 0$. This is a contradiction since $s \in [\epsilon, \infty)$. Thus $s_n \rightarrow 0$. □

Exercise 6.3.2.5. Let $\phi : [0, \infty) \rightarrow [0, \infty)$. Suppose that ϕ is increasing. Then ϕ is **Top** metric-preserving iff

1. $\phi^{-1}(\{0\}) = \{0\}$
2. ϕ is subadditive
3. ϕ is continuous

Then ϕ is **Top** metric-preserving.

Proof.

• ($\implies$):

Suppose that ϕ is **Top** metric-preserving. Define $d : [0, \infty) \times [0, \infty) \rightarrow [0, \infty)$ by $d(x, y) := |x - y|$. Since ϕ is **Top** metric-preserving, d^ϕ is a metric on $[0, \infty)$.

1. Since d^ϕ is definite, we have that for each $x \in [0, \infty)$,

$$\begin{aligned} x \in \phi^{-1}(\{0\}) &\iff \phi(x) = 0 && \iff \phi(|x - 0|) = 0 \\ &\iff d^\phi(x, 0) && \\ &\iff x = 0. \end{aligned}$$

Hence $\phi^{-1}(\{0\}) = \{0\}$.

2. Since d^ϕ satisfies the triangle inequality, we have that for each $x, y \in [0, \infty)$,

$$\begin{aligned} \phi(x + y) &= \phi(|(x + y) - 0|) \\ &= d^\phi(x + y, 0) \\ &\leq d^\phi(x + y, x) + d^\phi(x, 0) \\ &= \phi(|(x + y) - x|) + \phi(|x - 0|) \\ &= \phi(y) + \phi(x). \end{aligned}$$

Thus ϕ is subadditive.

3. Let $(a_n)_{n \in \mathbb{N}} \subset [0, \infty)$ and $a \in [0, \infty)$. Suppose that $a_n \rightarrow a$. Since d^ϕ is a metric, [an ex in the metric topology section](#) implies that d^ϕ is $(\mathcal{T}_d \otimes \mathcal{T}_d, \mathcal{T}_{\mathbb{R}})$ continuous. Then

$$\begin{aligned} \phi(a_n) &= \phi(|a_n - 0|) \\ &= d^\phi(a_n, 0) \\ &\rightarrow d^\phi(a, 0) \\ &= \phi(|a - 0|) \\ &= \phi(a). \end{aligned}$$

Thus ϕ is continuous at a . Since $a \in [0, \infty)$ is arbitrary, we have that for each $a \in [0, \infty)$, ϕ is continuous at a . Hence ϕ is continuous.

• ($\impliedby$):

Let (X, d) be a metric space.

1. (a) Let $x, y \in X$. Suppose that $x = y$. Then $d(x, y) = 0$. Since $0 \in \phi^{-1}(\{0\})$, we have that

$$\begin{aligned} d^\phi(x, y) &= \phi(d(x, y)) \\ &= \phi(0) \\ &= 0 \end{aligned}$$

Conversely, suppose that $d^\phi(x, y) = 0$. Then $\phi(d(x, y)) = 0$ and therefore

$$\begin{aligned} d(x, y) &\in \phi^{-1}(\{0\}) \\ &= \{0\} \end{aligned}$$

Thus $d(x, y) = 0$. Since d is a metric on X , $x = y$. Hence $d^\phi(x, y) = 0$ iff $x = y$.

(b) Let $x, y, z \in X$. Since ϕ is increasing and subadditive, we have that

$$\begin{aligned} d_{\phi(x,z)} &= \phi(d(x,z)) \\ &\leq \phi(d(x,y) + d(y,z)) \\ &\leq \phi(d(x,y)) + \phi(d(y,z)) \\ &= d^\phi(x,y) + d^\phi(y,z) \end{aligned}$$

Therefore d^ϕ is a metric on X .

2. Let $(x_n)_{n \in \mathbb{N}} \subset X$ and $x \in X$. Suppose that $x_n \xrightarrow{d} x$. Then $d(x_n, x) \rightarrow 0$. Since ϕ is continuous,

$$\begin{aligned} d^\phi(x_n, x) &= \phi(d(x_n, x)) \\ &\rightarrow 0 \end{aligned}$$

So $x_n \xrightarrow{d^\phi} x$.

Conversely, suppose that $x_n \xrightarrow{d^\phi} x$. Then

$$\begin{aligned} \phi(d(x_n, x)) &= d^\phi(x_n, x) \\ &\rightarrow 0 \end{aligned}$$

The previous exercise implies that $d(x_n, x) \rightarrow 0$. Hence $x_n \xrightarrow{d} x$. Since $(x_n)_{n \in \mathbb{N}} \subset X$ and $x \in X$ are arbitrary, we have that $(x_n)_{n \in \mathbb{N}} \subset X$ and $x \in X$, $x_n \xrightarrow{d} x$ iff $x_n \xrightarrow{d^\phi} x$. Therefore $d^\phi \sim_{\mathbf{Top}} d$.

Since (X, d) is arbitrary, ϕ is **Top** metric-preserving. □

Exercise 6.3.2.6. Let (X, d) be a metric space and $\phi : [0, \infty) \rightarrow [0, \infty)$. Suppose that ϕ is strictly increasing and ϕ is **Top** metric-preserving. Then

1. for each $x \in X$ and $r > 0$,

(a) $B_{d^\phi}(x, \phi(r)) = B_d(x, r)$,

(b) $\bar{B}_{d^\phi}(x, \phi(r)) = \bar{B}_d(x, r)$,

2. (a) $\mathcal{B}_d \subset \mathcal{B}_{d^\phi}$,

(b) $\bar{\mathcal{B}}_d \subset \bar{\mathcal{B}}_{d^\phi}$.

Proof.

1. Let $x \in X$ and $r > 0$. Since ϕ is strictly increasing, for each $s, t \in [0, \infty)$, $s < t$ iff $\phi(s) < \phi(t)$ and $s \leq t$ iff $\phi(s) \leq \phi(t)$.

(a) For each $y \in X$,

$$\begin{aligned} y \in B_{d^\phi}(x, \phi(r)) &\iff d^\phi(x, y) < \phi(r) \\ &\iff \phi(d(x, y)) < \phi(r) \\ &\iff d(x, y) < r \\ &\iff y \in B_d(x, r). \end{aligned}$$

Thus $B_{d^\phi}(x, \phi(r)) = B_d(x, r)$.

(b) Similarly, for each $y \in X$,

$$\begin{aligned} y \in \bar{B}_{d^\phi}(x, \phi(r)) &\iff d^\phi(x, y) \leq \phi(r) \\ &\iff \phi(d(x, y)) \leq \phi(r) \\ &\iff d(x, y) \leq r \\ &\iff y \in \bar{B}_d(x, r). \end{aligned}$$

Thus $\bar{B}_{d^\phi}(x, \phi(r)) = \bar{B}_d(x, r)$.

2. (a) Let $B \in \bar{\mathcal{B}}_d$. Then there exists $x \in X$ and $r > 0$ such that $B = \bar{B}_d(x, r)$. Then

$$\begin{aligned} B &= \bar{B}_d(x, r) \\ &= \bar{B}_{d^\phi}(x, \phi(r)) \\ &\in \bar{\mathcal{B}}_{d^\phi}. \end{aligned}$$

Since $B \in \bar{\mathcal{B}}_d$ is arbitrary, we have that for each $B \in \bar{\mathcal{B}}_d$, $B \in \bar{\mathcal{B}}_{d^\phi}$.

- (b) **Similar to the prev part**

□

Exercise 6.3.2.7. Define $\phi : [0, \infty) \rightarrow [0, 1)$ by

$$\phi(t) = \frac{t}{1+t}$$

Then ϕ is **Top** metric-preserving.

Proof.

1. We note that $\phi \in C^\infty([0, \infty))$ and for each $t \in [0, \infty)$,

$$\phi'(t) = \frac{1}{(1+t)^2} \quad \text{and} \quad \phi''(t) = -\frac{2}{(1+t)^3}$$

In particular, ϕ is continuous.

2. Since $\phi' > 0$, ϕ is strictly increasing.
3. Since $\phi'' < 0$, ϕ is strictly concave. Since $\phi(0) = 0$, an exercise in the section on convex functions implies that ϕ is subadditive. [reference section on convex functions](#)
4. Clearly $\phi^{-1}(\{0\}) = \{0\}$.

So ϕ is **Top** metric-preserving.

□

Exercise 6.3.2.8. Let $a \in (0, \infty)$. Define $\phi_a : [0, \infty) \rightarrow [0, \infty)$ by

$$\phi_a(t) = t \wedge a$$

Then ϕ_a is **Top** metric-preserving.

Proof.

1. Clearly ϕ is continuous.
2. Clearly, ϕ is increasing.
3. Since the minimum of two concave functions is concave, ϕ is concave. Since $\phi(0) = 0$, an exercise in the section on convex functions implies that ϕ is subadditive. [reference section on convex functions](#)
4. Clearly $\phi^{-1}(\{0\}) = \{0\}$.

So ϕ_a is **Top** metric-preserving.

□

6.4 Subspaces

6.4.1 Introduction

Definition 6.4.1.1. Let (X, d) be a pseudometric space and $E \subset X$. We define the **subspace pseudometric on E induced by d** , denoted $d|_E : E \times E \rightarrow [0, \infty)$, by

$$d|_E := d_{E \times E}.$$

Exercise 6.4.1.2. Let (X, d) be a pseudometric space and $E \subset X$.

1. We have that

- (a) $d|_E$ is a pseudometric on E .
- (b) $\mathcal{T}_{d|_E} = \mathcal{T}_d \cap E$.

2. If d is a metric on X , then $d|_E$ is a metric on E .

Proof.

1. We have that

- (a) • For each $x \in E$,

$$\begin{aligned} d|_E(x, x) &= d(x, x) \\ &= 0 \end{aligned}$$

Thus d is semidefinite.

- For each $x, y \in E$,

$$\begin{aligned} d|_E(x, y) &= d(x, y) \\ &= d(y, x) \\ &= d|_E(y, x). \end{aligned}$$

Thus $d|_E$ is symmetric.

- For each $x, y, z \in E$,

$$\begin{aligned} d|_E(x, z) &= d(x, z) \\ &\leq d(x, y) + d(y, z) \\ &= d|_E(x, y) + d|_E(y, z). \end{aligned}$$

Thus $d|_E$ satisfies the triangle inequality.

Hence $d|_E$ is a pseudometric on E .

(b) Let $(x_n)_{n \in \mathbb{N}} \subset E$ and $x \in E$.

- Suppose that $x_n \rightarrow x$ in $(E, \mathcal{T}_{d|_E})$. Then $d|_E(x_n, x) \rightarrow 0$. Hence

$$\begin{aligned} d(x_n, x) &= d|_E(x_n, x) \\ &\rightarrow 0. \end{aligned}$$

Therefore $x_n \rightarrow x$ in $(X, \mathcal{T}_d)$. Exercise 4.5.1.10 implies that $x_n \rightarrow x$ in $(E, \mathcal{T}_d \cap E)$.

- Conversely, suppose that $x_n \rightarrow x$ in $(E, \mathcal{T}_d \cap E)$. Exercise 4.5.1.10 implies that $x_n \rightarrow x$ in $(X, \mathcal{T}_d)$. Then $d(x_n, x) \rightarrow 0$. Hence

$$\begin{aligned} d|_E(x_n, x) &= d(x_n, x) \\ &\rightarrow 0. \end{aligned}$$

Therefore $x_n \rightarrow x$ in $(E, \mathcal{T}_{d|_E})$.

Hence $x_n \rightarrow x$ in $(E, \mathcal{T}_{d|_E})$ iff $x_n \rightarrow x$ in $(E, \mathcal{T}_d \cap E)$. Exercise 4.3.2.16 implies that $\mathcal{T}_d \cap E = \mathcal{T}_{d|_E}$.

2. Suppose that d is a metric on X . Let $x, y \in E$. Suppose that $d|_E(x, y) = 0$. Then

$$\begin{aligned} 0 &= d|_E(x, y) \\ &= d(x, y). \end{aligned}$$

Since d is definite, $x = y$. Since $x, y \in E$ are arbitrary, we have that for each $x, y \in E$, $d|_E(x, y) = 0$ implies that $x = y$. Thus $d|_E$ is definite. Hence $d|_E$ is a metric on E .

□

Exercise 6.4.1.3. Let (X, d) be a metric space and $E \subset X$. Then for each $x \in E$ and $r > 0$,

1. $B_{d|_E}(x, r) = B_d(x, r) \cap E$,
2. $\bar{B}_{d|_E}(x, r) = \bar{B}_d(x, r) \cap E$.

Proof. Let $x \in E$ and $r > 0$.

1. • Let $y \in B_{d|_E}(x, r)$. Then $y \in E$ and

$$\begin{aligned} d(x, y) &= d|_E(x, y) \\ &< r. \end{aligned}$$

Thus $y \in B_d(x, r)$. Therefore $y \in B_d(x, r) \cap E$. Since $y \in B_{d|_E}(x, r)$ is arbitrary, we have that for each $y \in B_{d|_E}(x, r)$, $y \in B_d(x, r) \cap E$. Hence $B_{d|_E}(x, r) \subset B_d(x, r) \cap E$.

- Let $y \in B_d(x, r) \cap E$. Then

$$\begin{aligned} d|_E(x, y) &= d(x, y) \\ &< r. \end{aligned}$$

Hence $y \in B_{d|_E}(x, r)$. Since $y \in B_d(x, r) \cap E$ is arbitrary, we have that for each $y \in B_d(x, r) \cap E$, $y \in B_{d|_E}(x, r)$. Hence $B_d(x, r) \cap E \subset B_{d|_E}(x, r)$.

2. Similar to (1).

□

Exercise 6.4.1.4. Let (X, d) be a metric space and $E \subset X$. If $(X, \mathcal{T}_d)$ is separable, then $(E, \mathcal{T}_d \cap E)$ is separable.

Proof. Suppose that $(X, \mathcal{T}_d)$ is separable. Exercise 6.2.1.1 implies that $(X, \mathcal{T}_d)$ is second-countable. Exercise 4.13.2.11 implies that $(E, \mathcal{T}_d \cap E)$ is second-countable. Another application of Exercise 6.2.1.1 implies that $(E, \mathcal{T}_d \cap E)$ is separable. □

6.4.2 Discrete Subsets

Definition 6.4.2.1. Let (X, d) be a metric space and $S \subset X$. Then S is said to be **discrete** if for each $x \in X$, there exists $r > 0$ such that $B(x, r) \cap X = \{x\}$.

Exercise 6.4.2.2. Let (X, d) be a metric space and $S \subset X$. Then S is discrete iff $\mathcal{T}_d \cap S = \mathcal{T}_{d_{scrt}(X)}$.

Proof. **FINISH!!!** □

Exercise 6.4.2.3. **compare with discrete sets in topology section** Let (X, d) be a metric space, $A \subset X$ and $x \in A$. Then x is an isolated point of A iff there exists $r > 0$ such that $B(x, r) \cap A = \{x\}$.

Proof.

• $(\implies)$:

Suppose that x is an isolated point of A . Then there exists $U \subset X$ such that U is open in X and $U \cap A = \{x\}$. Since U is open, $x \in U$ and $\{B(x, r) : r > 0\}$ is a local basis for the topology on X at x , there exists $r > 0$ such that $B(x, r) \subset U$. Hence

$$\begin{aligned} B(x, r) \cap A &\subset U \cap A \\ &= \{x\} \end{aligned}$$

Since $x \in B(x, r) \cap A$, we have that $\{x\} \subset B(x, r) \cap A$. Hence $B(x, r) \cap A = \{x\}$.

• $(\impliedby)$:

Conversely, suppose that there exists $r > 0$ such that $B(x, r) \cap A = \{x\}$. Since $B(x, r)$ is open in X , x is an isolated point of A . □

Exercise 6.4.2.4. Let (X, d) be a metric space and $A \subset X$. Suppose that A is discrete. If X is separable, then A is countable.

Hint: If $E \subset X$ is countable and dense in X , then for each $x \in A$, there exists $y \in E$ and $q \in \mathbb{Q} \cap (0, \infty)$ such that $x \in B(y, q)$.

Proof. Suppose that X is separable. Let $x \in A$. Since X is separable, there exists $(x_n)_{n \in \mathbb{N}} \subset X$ such that $(x_n)_{n \in \mathbb{N}}$ is dense in X . Since A is discrete, x is an isolated point of A and **the previous exercise** implies that there exists $r > 0$ such that $B(x, r) \cap A = \{x\}$. Choose $q \in \mathbb{Q} \cap (0, r)$. Set $\epsilon = \min(r - q, q)$. Then $\epsilon > 0$. Since $(x_n)_{n \in \mathbb{N}}$ is dense in X , there exists $N \in \mathbb{N}$ such that $d(x_N, x) < \epsilon$. Let $y \in B(x_N, q)$. Then

$$\begin{aligned} d(y, x) &\leq d(y, x_N) + d(x_N, x) \\ &< q + \epsilon \\ &\leq q + (r - q) \\ &= r \end{aligned}$$

Thus $y \in B(x, r)$. Since $y \in B(x_N, q)$ is arbitrary, we have that $B(x_N, q) \subset B(x, r)$. In addition,

$$\begin{aligned} d(x_N, x) &< \epsilon \\ &\leq q \end{aligned}$$

which implies that $x \in B(x_N, q)$ and therefore $\{x\} \subset B(x_N, q) \cap A$. Conversely,

$$\begin{aligned} B(x_N, q) \cap A &\subset B(x, r) \cap A \\ &= \{x\} \end{aligned}$$

Hence $B(x_N, q) \cap A = \{x\}$. Since $x \in A$ is arbitrary, we have that for each $x \in A$,

$$\{B(x_n, q) : (n, q) \in \mathbb{N} \times (\mathbb{Q} \cap (0, \infty)) \text{ and } B(x_n, q) \cap A = \{x\}\} \neq \emptyset$$

For each $x \in A$, define

$$V(x) := \{B(x_n, q) : (n, q) \in \mathbb{N} \times (\mathbb{Q} \cap (0, \infty)) \text{ and } B(x_n, q) \cap A = \{x\}\}$$

Define

$$V := \{B(x_n, q) : (n, q) \in \mathbb{N} \times (\mathbb{Q} \cap (0, \infty))\}$$

Since $\bigcup_{x \in A} V(x) \subset V$ and V is countable, we have that $\bigcup_{x \in A} V(x)$ is countable. The axiom of choice implies that there exists $\phi : A \rightarrow \bigcup_{x \in A} V(x)$ such that for each $x \in A$, $\phi(x) \in V(x)$. Let $x, y \in A$. Suppose that $\phi(x) = \phi(y)$. By construction,

$$\begin{aligned} \{x\} &= \phi(x) \cap A \\ &= \phi(y) \cap A \\ &= \{y\} \end{aligned}$$

Hence $x = y$. Since $x, y \in A$ are arbitrary, ϕ is injective. Since $\phi : A \rightarrow \bigcup_{x \in A} V(x)$ is injective, and $\bigcup_{x \in A} V(x)$ is countable, we have that A is countable. □

6.5 Product Spaces

Definition 6.5.0.1. Let $(X_n, d_n)_{n \in \mathbb{N}}$ be a collection of pseudometric spaces. Define $\phi : [0, \infty) \rightarrow [0, 1)$ by $\phi(t) := t/(1+t)$. We define the **product pseudometric on** $\prod_{n \in \mathbb{N}} X_n$, denoted $\bigotimes d_n : X \times X \rightarrow [0, \infty)$ by

$$\bigotimes d_n((x_n)_{n \in \mathbb{N}}, (y_n)_{n \in \mathbb{N}}) = \sum_{n \in \mathbb{N}} 2^{-n} d_n^\phi(x_n, y_n).$$

Exercise 6.5.0.2. Let $(X_n, d_n)_{n \in \mathbb{N}}$ be a collection of pseudometric spaces. Set $X := \prod_{n \in \mathbb{N}} X_n$, $\mathcal{T} := \bigotimes_{n \in \mathbb{N}} \mathcal{T}_{d_n}$ and $d := \bigotimes_{n \in \mathbb{N}} d_n$.

1. We have that

- (a) d is a pseudometric on X ,
- (b) $\mathcal{T}_d = \mathcal{T}$.

2. If for each $n \in \mathbb{N}$, d_n is a metric on X_n , then d is a metric on X

Proof.

1. Set $X := \prod_{n \in \mathbb{N}} X_n$ and $d := \bigotimes_{n \in \mathbb{N}} d_n$.

- (a) • We have that for each $(x_n)_{n \in \mathbb{N}} \in X$,

$$\begin{aligned} d((x_n)_{n \in \mathbb{N}}, (x_n)_{n \in \mathbb{N}}) &= \sum_{n \in \mathbb{N}} 2^{-n} d_n^\phi(x_n, x_n) \\ &= \sum_{n \in \mathbb{N}} 2^{-n} 0 \\ &= 0. \end{aligned}$$

Thus d is semidefinite.

- We have that for each $(x_n)_{n \in \mathbb{N}}, (y_n)_{n \in \mathbb{N}} \in X$,

$$\begin{aligned} d((x_n)_{n \in \mathbb{N}}, (y_n)_{n \in \mathbb{N}}) &= \sum_{n \in \mathbb{N}} 2^{-n} d_n^\phi(x_n, y_n) \\ &= \sum_{n \in \mathbb{N}} 2^{-n} d_n^\phi(y_n, x_n) \\ &= d((y_n)_{n \in \mathbb{N}}, (x_n)_{n \in \mathbb{N}}). \end{aligned}$$

Thus d is symmetric.

- We have that for each $(x_n)_{n \in \mathbb{N}}, (y_n)_{n \in \mathbb{N}}, (z_n)_{n \in \mathbb{N}} \in X$,

$$\begin{aligned} d((x_n)_{n \in \mathbb{N}}, (y_n)_{n \in \mathbb{N}}) &= \sum_{n \in \mathbb{N}} 2^{-n} d_n^\phi(x_n, y_n) \\ &\leq \sum_{n \in \mathbb{N}} 2^{-n} [d_n^\phi(x_n, z_n) + d_n^\phi(z_n, y_n)] \\ &= \sum_{n \in \mathbb{N}} 2^{-n} d_n^\phi(x_n, z_n) + \sum_{n \in \mathbb{N}} 2^{-n} d_n^\phi(z_n, y_n) \\ &= d((x_n)_{n \in \mathbb{N}}, (z_n)_{n \in \mathbb{N}}) + d((z_n)_{n \in \mathbb{N}}, (y_n)_{n \in \mathbb{N}}). \end{aligned}$$

Thus d satisfies the triangle inequality.

Hence d is a pseudometric on X .

- (b) Let $(a_m)_{m \in \mathbb{N}} \subset X$ and $a \in X$. Then for each $m \in \mathbb{N}$, there exist $(x_{m,n})_{n \in \mathbb{N}} \in X$ such that $a_m = (x_{m,n})$ and there exists $(x_n)_{n \in \mathbb{N}} \in X$ such that $a = (x_n)_{n \in \mathbb{N}}$.

- Suppose that $a_m \rightarrow a$ in $(X, \mathcal{T})$. Let $\epsilon > 0$. Choose $N_0 \in \mathbb{N}$ such that $\sum_{n \geq N_0+1} 2^{-n} < \epsilon/2$. Since $a_m \rightarrow a$ in $(X, \mathcal{T})$, we have that for each $n \in \mathbb{N}$, $x_{m,n} \xrightarrow{m} x_n$ in $(X_n, \mathcal{T}_{d_n})$. Hence for each $n \in \mathbb{N}$, $d_n(x_{m,n}, x_n) \xrightarrow{m} 0$. Exercise 6.3.2.7 implies that for each $n \in \mathbb{N}$, $d_n^\phi(x_{m,n}, x_n) \xrightarrow{m} 0$. Let $n \in [N_0]$. Since $d_n^\phi(x_{m,n}, x_n) \xrightarrow{m} 0$, there exists $M_n \in \mathbb{N}$ such that for each $m \in \mathbb{N}$, $m \geq M_n$ implies that $d_n^\phi(x_{m,n}, x_n) < \epsilon/2$. Set $M := \max(M_1, \dots, M_{N_0})$. Let $m \in \mathbb{N}$. Suppose that $m \geq M$. Then

$$\begin{aligned}
 d(a_m, a) &= \sum_{n \in \mathbb{N}} 2^{-n} d_n^\phi(x_{m,n}, x_n) \\
 &= \sum_{n=1}^{N_0} 2^{-n} d_n^\phi(x_{m,n}, x_n) + \sum_{n \geq N_0+1} 2^{-n} d_n^\phi(x_{m,n}, x_n) \\
 &< \sum_{n=1}^{N_0} 2^{-n} \left(\frac{\epsilon}{2} \right) + \sum_{n \geq N_0+1} 2^{-n} \\
 &= \frac{\epsilon}{2} \sum_{n=1}^{N_0} 2^{-n} + \sum_{n \geq N_0+1} 2^{-n} \\
 &< \frac{\epsilon}{2} + \frac{\epsilon}{2} \\
 &= \epsilon
 \end{aligned}$$

Since $\epsilon > 0$ is arbitrary, we have that for each $\epsilon > 0$, there exists $M \in \mathbb{N}$ such that for each $m \in \mathbb{N}$, $m \geq M$ implies that $d(a_m, a) < \epsilon$. Hence $d(a_m, a) \rightarrow 0$ and therefore $a_m \rightarrow a$ in $(X, \mathcal{T}_d)$.

- Conversely, suppose that $a_m \rightarrow a$ in $(X, \mathcal{T}_d)$. Then $d(a_m, a) \rightarrow 0$. Let $n \in \mathbb{N}$. Then

$$\begin{aligned}
 2^{-n} d_n^\phi(x_{m,n}, x_n) &\leq d(a_m, a) \\
 &\rightarrow 0
 \end{aligned}$$

Hence $d_n^\phi(x_{m,n}, x_n) \rightarrow 0$. Thus $x_{m,n} \xrightarrow{m} x_n$ in $(X, \mathcal{T}_{d_n^\phi})$. Since $\mathcal{T}_{d_n^\phi} = \mathcal{T}_{d_n}$, we have that $x_{m,n} \xrightarrow{m} x_n$ in $(X, \mathcal{T}_{d_n})$. Since $n \in \mathbb{N}$ is arbitrary, we have that $a_m \rightarrow a$ in $(X, \mathcal{T})$.

Since $(a_m)_{m \in \mathbb{N}} \subset X$ and $a \in X$ are arbitrary, we have that for each $(a_m)_{m \in \mathbb{N}} \subset X$ and $a \in X$, $a_m \rightarrow a$ in $(X, \mathcal{T})$ iff $a_m \rightarrow a$ in $(X, \mathcal{T}_d)$. Exercise 4.3.2.16 implies that $\mathcal{T}_d = \mathcal{T}$.

2. Suppose that for each $n \in \mathbb{N}$, d_n is a metric on X_n . Let $(x_n)_{n \in \mathbb{N}}, (y_n)_{n \in \mathbb{N}} \in X$. Suppose that $d((x_n)_{n \in \mathbb{N}}, (y_n)_{n \in \mathbb{N}}) = 0$. Then for each $n \in \mathbb{N}$, $d_n^\phi(x_n, y_n) = 0$. Therefore for each $n \in \mathbb{N}$, $x_n = y_n$. Hence $(x_n)_{n \in \mathbb{N}} = (y_n)_{n \in \mathbb{N}}$. Since $(x_n)_{n \in \mathbb{N}}, (y_n)_{n \in \mathbb{N}} \in X$ are arbitrary, we have that for each $(x_n)_{n \in \mathbb{N}}, (y_n)_{n \in \mathbb{N}} \in X$, $d((x_n)_{n \in \mathbb{N}}, (y_n)_{n \in \mathbb{N}}) = 0$ implies that $(x_n)_{n \in \mathbb{N}} = (y_n)_{n \in \mathbb{N}}$. Thus d is definite and d is a metric on X . □

Note 6.5.0.3. Thus the product topology is basically the topology of waning importance. More importance is given to earlier entries than later entries in a point of the product space. should have another example of a metric compatible with the product, maybe the supremum/n one

Definition 6.5.0.4. Let (X, d_X) and (Y, d_Y) be metric spaces. We define define max, sum, and other equivalent metrics $d_{\max} : (X \times Y)^2 \rightarrow [0, \infty)$ by **FINISH!!!**

Exercise 6.5.0.5. all the above metrics are equivalent (top equivalent or more stronger metrically equivalent?)

Proof. **FINISH!!!!** □

Definition 6.5.0.6. Let (X, d) be a metric space. Then d is $(d_{X^2}, d_{\mathbb{R}})$ -uniformly continuous.

Proof. **FINISH!!!** □

Exercise 6.5.0.7. content...

6.6 Coproduct Spaces

Definition 6.6.0.1. Let $(X_\alpha, d_\alpha)_{\alpha \in A}$ be a collection of metric space. Set $X := \coprod_{n \in \mathbb{N}} X_n$. Define $\phi : [0, \infty) \rightarrow [0, 1)$ by $\phi(t) := t/(1+t)$.

We define the **coproduct metric on X** , denoted $d_X : X \times X \rightarrow [0, \infty)$, by

$$d_X((\alpha, x), (\beta, y)) := \begin{cases} d_\alpha^\phi(x, y), & \alpha = \beta \\ 1, & \alpha \neq \beta \end{cases}$$

Exercise 6.6.0.2. Let $(X_\alpha, d_\alpha)_{\alpha \in A}$ be a collection of pseudometric spaces. Set $X := \coprod_{\alpha \in A} X_\alpha$ and $\mathcal{T} := \bigotimes_{\alpha \in A} \mathcal{T}_{d_\alpha}$.

1. We have that

- (a) d_X is a pseudometric on X ,
- (b) $\mathcal{T}_{d_X} = \mathcal{T}$.

2. If for each $\alpha \in A$, d_α is a metric on X_α , then d_X is a metric on X .

Proof.

- 1. (a) i. Let $(\alpha, x) \in X$.
 - Then

$$\begin{aligned} d_X((\alpha, x), (\alpha, x)) &= d_\alpha^\phi(x, x) \\ &= 0. \end{aligned}$$

Since $(\alpha, x) \in X$ is arbitrary, we have that for each $(\alpha, x) \in X$, $d_X((\alpha, x), (\alpha, x)) = 0$. Thus d_X is semidefinite.

ii. Let $(\alpha_1, x_1), (\alpha_2, x_2) \in X$. Then

$$d_X((\alpha_1, x_1), (\alpha_2, x_2)) = \begin{cases} d_\alpha^\phi(x_1, x_2) & \alpha_1 = \alpha_2 \\ 1 & \alpha_1 \neq \alpha_2 \end{cases} = \begin{cases} d_\alpha^\phi(x_2, x_1) & \alpha_2 = \alpha_1 \\ 1 & \alpha_2 \neq \alpha_1 \end{cases} = d_X((\alpha_2, x_2), (\alpha_1, x_1)).$$

Since $(\alpha_1, x_1), (\alpha_2, x_2) \in X$ is arbitrary, we have that for each $(\alpha_1, x_1), (\alpha_2, x_2) \in X$, $d_X((\alpha_1, x_1), (\alpha_2, x_2)) = d_X((\alpha_2, x_2), (\alpha_1, x_1))$. Thus d_X is symmetric.

iii. Let $(\alpha_1, x_1), (\alpha_2, x_2), (\alpha_3, x_3) \in X$.

- Suppose that $\alpha_1 = \alpha_2$ and $\alpha_1 = \alpha_3$. Then $\alpha_2 = \alpha_3$ and

$$\begin{aligned} d_X((\alpha_1, x_1), (\alpha_2, x_2)) &= d_{\alpha_1}^\phi(x_1, x_2) \\ &\leq d_{\alpha_1}^\phi(x_1, x_3) + d_{\alpha_1}^\phi(x_3, x_2) \\ &= d_X((\alpha_1, x_1), (\alpha_3, x_3)) + d_X((\alpha_3, x_3), (\alpha_2, x_2)). \end{aligned}$$

- Suppose that $\alpha_1 \neq \alpha_2$ or $\alpha_2 \neq \alpha_3$.

– If $\alpha_1 \neq \alpha_2$, then $d_X((\alpha_1, x_1), (\alpha_2, x_2)) = 1$ and

$$\begin{aligned} d_X((\alpha_1, x_1), (\alpha_3, x_3)) &\leq 1 \\ &\leq d_X((\alpha_1, x_1), (\alpha_2, x_2)) + 1 \\ &= d_X((\alpha_1, x_1), (\alpha_2, x_2)) + d_X((\alpha_2, x_2), (\alpha_3, x_3)). \end{aligned}$$

– If $\alpha_2 \neq \alpha_3$, then $d_X((\alpha_2, x_2), (\alpha_3, x_3)) = 1$ and

$$\begin{aligned} d_X((\alpha_1, x_1), (\alpha_3, x_3)) &\leq 1 \\ &\leq 1 + d_X((\alpha_2, x_2), (\alpha_3, x_3)) \\ &= d_X((\alpha_1, x_1), (\alpha_2, x_2)) + d_X((\alpha_2, x_2), (\alpha_3, x_3)). \end{aligned}$$

Since $(\alpha_1, x_1), (\alpha_2, x_2), (\alpha_3, x_3) \in X$ are arbitrary, we have that for each $(\alpha_1, x_1), (\alpha_2, x_2), (\alpha_3, x_3) \in X$, $d_X((\alpha_1, x_1), (\alpha_3, x_3)) \leq d_X((\alpha_1, x_1), (\alpha_2, x_2)) + d_X((\alpha_2, x_2), (\alpha_3, x_3))$. Thus d_X satisfies the triangle inequality.

Thus d_X is a pseudometric on X .

(b) Let $(\alpha_j, x_j)_{j \in \mathbb{N}} \subset X$ and $(\alpha_0, x_0) \in X$.

- Suppose that $(\alpha_j, x_j) \rightarrow (\alpha_0, x_0)$ in $(X, \mathcal{T})$. Exercise 4.7.0.8 implies that there exists $j_0 \in \mathbb{N}$ such that
 - i. for each $j \in \mathbb{N}$, $j \geq j_0$ implies that $\alpha_j = \alpha_0$,
 - ii. $[L_{j_0}(x)]_j \rightarrow x_0 \in (X_{\alpha_0}, \mathcal{T}_{d_{\alpha_0}})$.

Hence $d_{\alpha_0}(x_{j+j_0}, x_0) \rightarrow 0$. Therefore

$$\begin{aligned} d_X((\alpha_{j+j_0}, x_{j+j_0}), (\alpha_0, x_0)) &= d^\phi(x_j, x_0) \\ &\rightarrow 0. \end{aligned}$$

Hence $(\alpha_j, x_j) \rightarrow (\alpha_0, x_0)$ in $(X, \mathcal{T}_{d_X})$.

- Conversely, suppose that $(\alpha_j, x_j) \rightarrow (\alpha_0, x_0)$ in $(X, \mathcal{T}_{d_X})$. Then $d_X((\alpha_j, x_j), (\alpha_0, x_0)) \rightarrow 0$. Therefore there exists $j_0 \in \mathbb{N}$ such that for each $j \in \mathbb{N}$, $j \geq j_0$ implies that $\alpha_j = \alpha_0$ and $d_{\alpha_0}^\phi(x_{j+j_0}, x_0) \rightarrow 0$. Therefore $d_{\alpha_0}(L_{j_0}(x), x_0) \rightarrow 0$. Exercise 4.7.0.8 implies that $(\alpha_j, x_j) \rightarrow (\alpha_0, x_0)$ in $(X, \mathcal{T})$.

Since $(\alpha_j, x_j) \rightarrow (\alpha_0, x_0)$ in $(X, \mathcal{T}_{d_X})$ iff $(\alpha_j, x_j) \rightarrow (\alpha_0, x_0)$ in $(X, \mathcal{T})$, Exercise 4.3.2.16 implies that $\mathcal{T} = \mathcal{T}_{d_X}$.

2. Suppose that for each $\alpha \in A$, d_α is a metric on X_α . Let $(\alpha_1, x_1), (\alpha_2, x_2) \in X$. Suppose that $d_X((\alpha_1, x_1), (\alpha_2, x_2)) = 0$. For the sake of contradiction, suppose that $\alpha_1 \neq \alpha_2$. Then

$$\begin{aligned} 0 &= d_X((\alpha_1, x_1), (\alpha_2, x_2)) \\ &= 1 \end{aligned}$$

which is a contradiction. Thus $\alpha_1 = \alpha_2$. Then

$$\begin{aligned} 0 &= d_X((\alpha_1, x_1), (\alpha_2, x_2)) \\ &= d_{\alpha_1}^\phi(x_1, x_2). \end{aligned}$$

Hence $x_1 = x_2$ and $(\alpha_1, x_1) = (\alpha_2, x_2)$. Thus d_X is definite. Since d_X is a pseudometric on X and d_X is definite, we have that d_X is a metric on X . □

Exercise 6.6.0.3. Let $(X_\alpha, d_\alpha)_{\alpha \in A}$ be a collection of metric space. Set $X := \coprod_{\alpha \in A} X_\alpha$ and $\mathcal{T} := \bigotimes_{\alpha \in A} \mathcal{T}_{d_\alpha}$. Then for each $\alpha \in A$, $\iota_\alpha : X_\alpha \rightarrow X$ is Lipschitz.

Proof. Define $\phi : [0, \infty) \rightarrow [0, 1]$ by $\phi(t) := t/(1+t)$. Exercise 6.1.3.11 implies that ϕ is Lipschitz. Hence there exists $K > 0$ such that for each $a, b \in [0, \infty)$, $|\phi(a) - \phi(b)| \leq K|a - b|$. Let $\alpha \in A$ and $x_1, x_2 \in X_\alpha$. Then

$$\begin{aligned} d_X(\iota_\alpha(x_1), \iota_\alpha(x_2)) &= d_X((\alpha, x_1), (\alpha, x_2)) \\ &= d_\alpha^\phi(x_1, x_2) \\ &= \phi(d_\alpha(x_1, x_2)) \\ &= \phi(|d_\alpha(x_1, x_2) - 0|) && \leq K|d_\alpha(x_1, x_2) - \phi(0)| \\ &= Kd_\alpha(x_1, x_2) \end{aligned}$$

Since $x_1, x_2 \in X_\alpha$ are arbitrary, we have that ι_α is Lipschitz. Since $\alpha \in A$ is arbitrary, we have that for each $\alpha \in A$, ι_α is Lipschitz. □

6.7 Completeness

6.7.1 Introduction

Definition 6.7.1.1. Let (X, d) be a metric space and $(x_\alpha)_{\alpha \in A} \subset X$ a net. Then $(x_\alpha)_{\alpha \in A}$ is said to be **Cauchy** in (X, d) if for each $\epsilon > 0$, there exists $\alpha_0 \in \mathbb{N}$ such that for each $\alpha_1, \alpha_2 \in A$, $\alpha_1, \alpha_2 \geq \alpha_0$ implies that $d(x_{\alpha_1}, x_{\alpha_2}) < \epsilon$.

Exercise 6.7.1.2. Let (X, d) be a metric space and $(x_\alpha)_{\alpha \in A} \subset X$ a net. If $(x_\alpha)_{\alpha \in A}$ is Cauchy, then for each subnet $(y_\beta)_{\beta \in B}$, $(y_\beta)_{\beta \in B}$ is Cauchy.

Proof. Suppose that $(x_\alpha)_{\alpha \in A}$ is Cauchy. Let $(y_\beta)_{\beta \in B}$ be a subnet. Write $y_\beta = x_{\alpha_\beta}$. Let $\epsilon > 0$. Since $(x_\alpha)_{\alpha \in A}$ is Cauchy, there exists $\alpha_0 \in A$ such that for each $\alpha_1, \alpha_2 \in A$, $\alpha_1, \alpha_2 \geq \alpha_0$ implies that $d(x_{\alpha_1}, x_{\alpha_2}) < \epsilon$. Since $(y_\beta)_{\beta \in B} \subset (x_\alpha)_{\alpha \in A}$ is a subnet, there exists $\beta_0 \in B$ such that for each $\beta \in B$, $\beta \geq \beta_0$ implies that $\alpha_\beta \geq \alpha_0$. Then for each $\beta_1, \beta_2 \in B$, $\beta_1, \beta_2 \geq \beta_0$ implies that

$$\begin{aligned} d(y_{\beta_1}, y_{\beta_2}) &= d(x_{\alpha_{\beta_1}}, x_{\alpha_{\beta_2}}) \\ &< \epsilon. \end{aligned}$$

Since $\epsilon > 0$ is arbitrary, we have that for each $\epsilon > 0$, there exists $\beta_0 \in B$ such that for each $\beta_1, \beta_2 \in B$, $\beta_1, \beta_2 \geq \beta_0$ implies that $d(y_{\beta_1}, y_{\beta_2}) < \epsilon$. Hence $(y_\beta)_{\beta \in B}$ is Cauchy. Since $(y_\beta)_{\beta \in B} \subset (x_\alpha)_{\alpha \in A}$ a subnet is arbitrary, we have that for each subnet $(y_\beta)_{\beta \in B} \subset (x_\alpha)_{\alpha \in A}$, $(y_\beta)_{\beta \in B}$ is Cauchy. $\square$

Exercise 6.7.1.3. Let (X, d) be a metric space and $(x_n)_{n \in \mathbb{N}} \subset X$ and $x \in X$. If $x_n \rightarrow x$, then $(x_n)_{n \in \mathbb{N}}$ is Cauchy.

Proof. Suppose that $x_n \rightarrow x$. Let $\epsilon > 0$. Then there exists $N \in \mathbb{N}$ such that for each $n \in \mathbb{N}$, $n \geq N$ implies that $d(x_n, x) < \epsilon/2$. Then for each $m, n \in \mathbb{N}$, $m, n \geq N$ implies that

$$\begin{aligned} d(x_m, x_n) &\leq d(x_m, x) + d(x, x_n) \\ &< \frac{\epsilon}{2} + \frac{\epsilon}{2} \\ &= \epsilon. \end{aligned}$$

Hence $(x_n)_{n \in \mathbb{N}}$ is Cauchy. $\square$

Exercise 6.7.1.4. Let (X, d) be a metric space and $(x_n)_{n \in \mathbb{N}} \subset X$. Suppose that $(x_n)_{n \in \mathbb{N}}$ is Cauchy. If there exists $(x_{n_k})_{k \in \mathbb{N}} \subset (x_n)_{n \in \mathbb{N}}$ and $x \in X$ such that $x_{n_k} \rightarrow x$, then $x_n \rightarrow x$. **NEED TO DEFINE SUBSEQUENCE (specifically how n_k is an increasing sequence from $\mathbb{N} \rightarrow \mathbb{N}$ and therefore $n_k \geq k$) in topology net section!!!**

Proof. Suppose that there exists $(x_{n_k})_{k \in \mathbb{N}} \subset (x_n)_{n \in \mathbb{N}}$ and $x \in X$ such that $x_{n_k} \rightarrow x$. Let $\epsilon > 0$. Since $x_{n_k} \rightarrow x$, there exists $K_0 \in \mathbb{N}$ such that for each $k \in \mathbb{N}$, $k \geq K_0$ implies that $d(x_{n_k}, x) < \epsilon/2$. Since $(x_n)_{n \in \mathbb{N}}$ is Cauchy, there exists $N_0 \in \mathbb{N}$ such that for each $m, n \in \mathbb{N}$, $m, n \geq N_0$ implies that $d(x_m, x_n) < \epsilon/2$. Set $K := \max(N_0, K_0)$. Then $K \geq K_0$ and since $(n_k)_{k \in \mathbb{N}}$ is strictly increasing, we have that

$$\begin{aligned} n_K &\geq K \\ &\geq N_0. \end{aligned}$$

Therefore for each $n \in \mathbb{N}$, $n \geq N_K$ implies that

$$\begin{aligned} d(x_n, x) &\leq d(x_n, x_{n_K}) + d(x_{n_K}, x) \\ &< \frac{\epsilon}{2} + \frac{\epsilon}{2} \\ &= \epsilon. \end{aligned}$$

Since $\epsilon > 0$ is arbitrary, we have that for each $\epsilon > 0$, there exists $N \in \mathbb{N}$ such that for each $n \in \mathbb{N}$, $n \geq N$ implies that $d(x_n, x) < \epsilon$. Hence $x_n \rightarrow x$. $\square$

Exercise 6.7.1.5. Let (X, d_X) and (Y, d_Y) be metric spaces and $f : X \rightarrow Y$. If f is uniformly continuous, then for each $(a_n)_{n \in \mathbb{N}} \subset X$, $(a_n)_{n \in \mathbb{N}}$ is Cauchy in (X, d_X) implies that $(f(a_n))_{n \in \mathbb{N}}$ is Cauchy in (Y, d_Y) .

Proof. Suppose that f is uniformly continuous. Let $(a_n)_{n \in \mathbb{N}} \subset X$. Suppose that $(a_n)_{n \in \mathbb{N}}$ is Cauchy in (X, d_X) . Let $\epsilon > 0$. Since f is uniformly continuous, there exists $\delta > 0$ such that for each $x_1, x_2 \in X$, $d_X(x_1, x_2) < \delta$ implies that $d_Y(f(x_1), f(x_2)) < \epsilon$. Since $(a_n)_{n \in \mathbb{N}}$ is Cauchy in (X, d_X) , there exists $N \in \mathbb{N}$ such that for each $m, n \in \mathbb{N}$, $m, n \geq N$ implies that $d_X(a_m, a_n) < \delta$. Let $m, n \in \mathbb{N}$. Suppose that $m, n \geq N$. Then $d_Y(f(a_m), f(a_n)) < \epsilon$. Since $\epsilon > 0$ is arbitrary, we have that for each $\epsilon > 0$, there exists $N \in \mathbb{N}$ such that for each $m, n \in \mathbb{N}$, $m, n \geq N$ implies that $d_Y(f(a_m), f(a_n)) < \epsilon$. Hence $(f(a_n))_{n \in \mathbb{N}}$ is Cauchy in (Y, d_Y) . $\square$

Exercise 6.7.1.6. Let (X, d) be a metric space and $\phi : [0, \infty) \rightarrow [0, \infty)$. Suppose that ϕ is **Top** metric-preserving and

1. ϕ is homeomorphism, (relax this to homeo near 0)
2. $\phi(0) = 0$.

Then for each $(x_n)_{n \in \mathbb{N}} \subset X$, $(x_n)_{n \in \mathbb{N}}$ is Cauchy in (X, d) iff $(x_n)_{n \in \mathbb{N}}$ is Cauchy in (X, d^ϕ) . (try showing that a homoemorphism with $\phi(0) = 0$ is strictly increasing)

Proof. Let $(x_n)_{n \in \mathbb{N}} \subset X$. Since ϕ is a homoemorphism and $\phi(0) = 0$, we have that ϕ is injective and strictly increasing.

• $(\implies)$:

Suppose that $(x_n)_{n \in \mathbb{N}}$ is Cauchy in (X, d) . Let $\epsilon > 0$. Since ϕ is injective and $\phi(0) = 0$, $\phi^{-1}(\epsilon) > 0$ and there exists $N \in \mathbb{N}$ such that for each $m, n \in \mathbb{N}$, $m, n \geq N$ implies that $d(x_m, x_n) < \phi^{-1}(\epsilon)$. Let $m, n \in \mathbb{N}$. Suppose that $m, n \geq N$. Since ϕ is increasing, we have that

$$\begin{aligned} d^\phi(x_m, x_n) &= \phi \circ d(x_m, x_n) \\ &\leq \phi(\phi^{-1}(\epsilon)) \\ &= \epsilon \end{aligned}$$

Thus $(x_n)_{n \in \mathbb{N}}$ is Cauchy in (X, d^ϕ) .

• $(\impliedby)$:

Conversely, suppose that $(x_n)_{n \in \mathbb{N}}$ is Cauchy in (X, d^ϕ) . Let $\epsilon > 0$. Since ϕ is injective and $\phi(0) = 0$, $\phi(\epsilon) > 0$ and there exists $N \in \mathbb{N}$ such that for each $m, n \in \mathbb{N}$, $m, n \geq N$ implies that $d^\phi(x_m, x_n) < \phi(\epsilon)$. Let $m, n \in \mathbb{N}$. Suppose that $m, n \geq N$. Since ϕ^{-1} is increasing, we have that

$$\begin{aligned} d(x_m, x_n) &= \phi^{-1} \circ \phi \circ d(x_m, x_n) \\ &= \phi^{-1} \circ d^\phi(x_m, x_n) \\ &\leq \phi^{-1}(\phi(\epsilon)) \\ &= \epsilon \end{aligned}$$

Thus $(x_n)_{n \in \mathbb{N}}$ is Cauchy in (X, d) . $\square$

Definition 6.7.1.7. Let (X, d) be a metric space. Then (X, d) is said to be **complete** if for each $(a_n)_{n \in \mathbb{N}} \subset X$, $(a_n)_{n \in \mathbb{N}}$ is Cauchy in (X, d) implies that there exists $a \in X$ such that $a_n \rightarrow a$ in $(X, \mathcal{T}_d)$.

Exercise 6.7.1.8. Let (X, d) be a metric space. If (X, d) is complete then for each net $(x_\alpha)_{\alpha \in A} \subset X$, $(x_\alpha)_{\alpha \in A}$ is Cauchy implies that $(x_\alpha)_{\alpha \in A}$ converges.

Proof. Suppose that (X, d) is Cauchy. Let $(x_\alpha)_{\alpha \in A} \subset X$ be a net. Suppose that $(x_\alpha)_{\alpha \in A}$ is Cauchy. Let $(y_\beta)_{\beta \in B} \subset (x_\alpha)_{\alpha \in A}$ be a subnet.

- If $\#B < \infty$, then make/ref ex in topology section on nets implies that $x_{\alpha_\beta} \rightarrow x_{\max B}$.
- Suppose that $\#B = \infty$. make/ref ex in subnet section of set theory part implies that there exists a subnet $(z_n)_{n \in \mathbb{N}} \subset (y_\beta)_{\beta \in B}$. Exercise ?? then implies that $(z_n)_{n \in \mathbb{N}} \subset (x_\alpha)_{\alpha \in A}$ is a subnet. ref prev ex implies that $(z_n)_{n \in \mathbb{N}}$ is Cauchy. Since (X, d) is complete, there exists $z \in X$ such that $z_n \rightarrow z$.

Since $(y_\beta)_{\beta \in B} \subset (x_\alpha)_{\alpha \in A}$ a subnet is arbitrary, we have that for each subnet $(y_\beta)_{\beta \in B} \subset (x_\alpha)_{\alpha \in A}$, there exists a subnet $(z_\gamma)_{\gamma \in \Gamma} \subset (x_\alpha)_{\alpha \in A}$ such that $(z_\gamma)_{\gamma \in \Gamma}$ converges. Exercise 4.3.2.22 then implies that $(x_\alpha)_{\alpha \in A}$ converges. Since $(x_\alpha)_{\alpha \in A} \subset X$ a net with $(x_\alpha)_{\alpha \in A}$ Cauchy is arbitrary, we have that for each net $(x_\alpha)_{\alpha \in A}$, $(x_\alpha)_{\alpha \in A}$ is Cauchy implies that $(x_\alpha)_{\alpha \in A}$ converges. $\square$

Exercise 6.7.1.9. Let $(X, d_X), (Y, d_Y)$ be metric spaces, $f : X \rightarrow Y$ a (d_X, d_Y) -isometry and $(x_n)_{n \in \mathbb{N}}$. Then $(x_n)_{n \in \mathbb{N}}$ is Cauchy in (X, d_X) iff $(f(x_n))_{n \in \mathbb{N}}$ is Cauchy in (Y, d_Y) .

Proof. Since f is a (d_X, d_Y) -isometry, Exercise 6.1.5.6 implies that f is injective and $d_X = f^*d_Y$.

• $(\implies)$:

Suppose that $(x_n)_{n \in \mathbb{N}}$ is Cauchy in (X, d_X) . Let $\epsilon > 0$. Since $(x_n)_{n \in \mathbb{N}}$ is Cauchy in (X, d_X) , there exists $N \in \mathbb{N}$ such that for each $m, n \in \mathbb{N}$, $m, n \geq N$ implies that $d_X(x_m, x_n) < \epsilon$. Let $m, n \in \mathbb{N}$. Suppose that $m, n \geq N$. Then

$$\begin{aligned} d_Y(f(x_m), f(x_n)) &= f^*d_Y(x_m, x_n) \\ &= d_X(x_m, x_n) \\ &< \epsilon. \end{aligned}$$

Since $\epsilon > 0$ is arbitrary, we have that for each $\epsilon > 0$, there exists $N \in \mathbb{N}$ such that for each $m, n \in \mathbb{N}$, $m, n \geq N$ implies that $d_Y(f(x_m), f(x_n)) < \epsilon$. Hence $(f(x_n))_{n \in \mathbb{N}}$ is Cauchy in (Y, d_Y) .

• $(\impliedby)$:

Conversely, suppose that $(f(x_n))_{n \in \mathbb{N}}$ is Cauchy in (Y, d_Y) . Let $\epsilon > 0$. Since $(f(x_n))_{n \in \mathbb{N}}$ is Cauchy in (Y, d_Y) , there exists $N \in \mathbb{N}$ such that for each $m, n \in \mathbb{N}$, $m, n \geq N$ implies that $d_Y(f(x_m), f(x_n)) < \epsilon$. Let $m, n \in \mathbb{N}$. Suppose that $m, n \geq N$. Then

$$\begin{aligned} d_X(x_m, x_n) &= f^*d_Y(x_m, x_n) \\ &= d_Y(f(x_m), f(x_n)) \\ &< \epsilon. \end{aligned}$$

Since $\epsilon > 0$ is arbitrary, we have that for each $\epsilon > 0$, there exists $N \in \mathbb{N}$ such that for each $m, n \in \mathbb{N}$, $m, n \geq N$ implies that $d_X(x_m, x_n) < \epsilon$. Hence $(x_n)_{n \in \mathbb{N}}$ is Cauchy in (X, d_X) . $\square$

Exercise 6.7.1.10. Let $(X, d_X), (Y, d_Y)$ be metric spaces and $f : X \rightarrow Y$ a surjective (d_X, d_Y) -isometry. Then (X, d_X) is complete iff (Y, d) is complete.

Proof. Exercise 6.1.5.6 implies that f is injective and $d_X = f^*d_Y$. Since f is surjective, f is a bijection. Exercise 6.1.5.7 implies that f^{-1} is a (d_Y, d_X) -isometry and Exercise 6.1.5.6 implies that $d_Y = (f^{-1})^*d_X$.

• $(\implies)$:

Suppose that (X, d_X) is complete. Let $(y_n)_{n \in \mathbb{N}} \subset Y$. Suppose that $(y_n)_{n \in \mathbb{N}}$ is Cauchy in (Y, d_Y) . Define $(x_n)_{n \in \mathbb{N}} \subset X$ by $x_n := f^{-1}(y_n)$. Since f^{-1} is a (d_Y, d_X) -isometry, Exercise 6.7.1.9 then implies that $(x_n)_{n \in \mathbb{N}}$ is Cauchy in (X, d_X) . Since (X, d_X) is complete, there exists $x \in X$ such that $x_n \rightarrow x$ in (X, d_X) . Set $y := f(x)$. Then

$$\begin{aligned} d_Y(y_n, y) &= (f^{-1})^*d_X(y_n, y) \\ &= d_X(f^{-1}(y_n), f^{-1}(y)) \\ &= d_X(x_n, x) \\ &\rightarrow 0 \end{aligned}$$

Exercise 6.1.2.12 implies that $y_n \rightarrow y$. Since $(y_n)_{n \in \mathbb{N}} \subset Y$ with $(y_n)_{n \in \mathbb{N}}$ Cauchy in (Y, d_Y) is arbitrary, we have that for each $(y_n)_{n \in \mathbb{N}} \subset Y$, $(y_n)_{n \in \mathbb{N}}$ is Cauchy in (Y, d_Y) implies that there exists $y \in Y$ such that $y_n \rightarrow y$. Hence (Y, d_Y) is complete.

• $(\impliedby)$:

Similar to $(\implies)$. $\square$

Exercise 6.7.1.11. Let $(X, d_X), (Y, d_Y)$ be metric spaces, $S \subset X$ and $f_0 : S \rightarrow Y$. Suppose that (Y, d_Y) is complete, S is dense in X and f_0 is uniformly continuous. Then there exists a unique $f : X \rightarrow Y$ such that f is uniformly continuous and $f|_S = f_0$.

Proof.

- For each $x \in X$, define $E_x \subset S^{\mathbb{N}}$ by $E_x := \{(x_n)_{n \in \mathbb{N}} \subset S : x_n \rightarrow x \text{ and } f_0(x_n) \text{ converges}\}$. Let $x \in X$. Since S is dense in X , there exists $(x_n)_{n \in \mathbb{N}} \subset S$ such that $x_n \rightarrow x$. Exercise 6.7.1.3 then implies that $(x_n)_{n \in \mathbb{N}}$ is Cauchy in (X, d_X) . Since f_0 is uniformly continuous, Exercise 6.7.1.5 implies that $(f_0(x_n))_{n \in \mathbb{N}}$ is Cauchy in (Y, d_Y) . Since (Y, d_Y) is complete, there exists $y_x \in Y$ such that $f_0(x_n) \rightarrow y_x$. Thus $E_x \neq \emptyset$. Since $x \in X$ is arbitrary, we have that for each $x \in X$, $E_x \neq \emptyset$. The axiom of choice implies that there exists $\phi \in \prod_{x \in X} E_x$. Define $f : X \rightarrow Y$ by $f(x) := \lim_{n \rightarrow \infty} f_0(\phi(x)_n)$.
- Let $\epsilon > 0$. Since f_0 is uniformly continuous, there exists $\delta_0 > 0$ such that for each $a, b \in S$, $d_X(a, b) < \delta_0$ implies $d_Y(f_0(a), f_0(b)) < \epsilon/3$. Define $\delta > 0$ by $\delta := \delta_0/3$. Let $a, b \in X$. Suppose that $d_X(a, b) < \delta$. Since $\phi(a)_n \rightarrow a$, $\phi(b)_n \rightarrow b$, there exist $M_a, M_b \in \mathbb{N}$ such that for each $n \in \mathbb{N}$, $n \geq M_a$ implies that $d_X(\phi(a)_n, a) < \delta$ and $n \geq M_b$ implies that $d_X(\phi(b)_n, b) < \delta$. Define $M \in \mathbb{N}$ by $M := \max\{M_a, M_b\}$. Let $n \in \mathbb{N}$. Suppose that $n \geq M$. Since $d_X(a, b) < \delta$, we have that

$$\begin{aligned} d_X(\phi(a)_n, \phi(b)_n) &\leq d_X(\phi(a)_n, a) + d_X(a, b) + d_X(b, \phi(b)_n) \\ &< \delta + \delta + \delta \\ &= \delta_0. \end{aligned}$$

Therefore $d_Y(f_0(\phi(a)_n), f_0(\phi(b)_n)) < \epsilon/3$. Since $n \in \mathbb{N}$ with $n \geq M$ is arbitrary, we have that for each $n \in \mathbb{N}$, $n \geq M$ implies that $d_Y(f_0(\phi(a)_n), f_0(\phi(b)_n)) < \epsilon/3$. Since $f_0(\phi(a)_n) \rightarrow f(a)$ and $f_0(\phi(b)_n) \rightarrow f(b)$, there exist $N_a, N_b \in \mathbb{N}$ such that for each $n \in \mathbb{N}$, $n \geq N_a$ implies that $d_Y(f_0(\phi(a)_n), f(a)) < \epsilon/3$ and $n \geq N_b$ implies that $d_Y(f_0(\phi(b)_n), f(b)) < \epsilon/3$. Define $N \in \mathbb{N}$ by $N := \max\{M, N_a, N_b\}$. Then

$$\begin{aligned} d_Y(f(a), f(b)) &\leq d_Y(f(a), f_0(\phi(a)_N)) + d_Y(f_0(\phi(a)_N), f_0(\phi(b)_N)) + d_Y(f_0(\phi(b)_N), f(b)) \\ &< \frac{\epsilon}{3} + \frac{\epsilon}{3} + \frac{\epsilon}{3} \\ &= \epsilon. \end{aligned}$$

Since $\epsilon > 0$ is arbitrary, we have that for each $\epsilon > 0$, there exists $\delta > 0$ such that for each $a, b \in X$, $d_X(a, b) < \delta$ implies that $d_Y(f(a), f(b)) < \epsilon$. Hence f is uniformly continuous.

- – Let $x \in S$, $(a_n)_{n \in \mathbb{N}}$ and $(b_n)_{n \in \mathbb{N}} \in E_x$. By definition, $(a_n)_{n \in \mathbb{N}}, (b_n)_{n \in \mathbb{N}} \subset S$, $a_n, b_n \rightarrow x$. As in the previous part, there exists $y_a, y_b \in Y$ such that $f_0(a_n) \rightarrow y_a$ and $f_0(b_n) \rightarrow y_b$. Let $\epsilon > 0$. Since f_0 is uniformly continuous, there exists $\delta > 0$ such that for each $a, b \in S$, $d_X(a, b) < \delta$ implies $d_Y(f_0(a), f_0(b)) < \epsilon/3$. Since $f_0(a_n) \rightarrow y_a$ and $f_0(b_n) \rightarrow y_b$, there exist $N_a, N_b \in \mathbb{N}$ such that for each $n \in \mathbb{N}$, $n \geq N_a$ implies that $d_Y(f_0(a_n), y_a) < \epsilon/3$ and $n \geq N_b$ implies that $d_Y(f_0(b_n), y_b) < \epsilon/3$. Since $a_n \rightarrow x$ and $b_n \rightarrow x$, we have that

$$\begin{aligned} d_X(a_n, b_n) &\leq d_X(a_n, x) + d_X(x, b_n) \\ &\rightarrow 0. \end{aligned}$$

Thus there exists $N_0 \in \mathbb{N}$ such that for each $n \in \mathbb{N}$, $n \geq N_0$ implies that $d_X(a_n, b_n) < \delta$. Set $N := \max\{N_0, N_a, N_b\}$. Then

$$\begin{aligned} d_Y(y_a, y_b) &\leq d_Y(y_a, f_0(a_N)) + d_Y(f_0(a_N), f_0(b_N)) + d_Y(f_0(b_N), y_b) \\ &< \frac{\epsilon}{3} + \frac{\epsilon}{3} + \frac{\epsilon}{3} \\ &= \epsilon. \end{aligned}$$

Since $\epsilon > 0$ is arbitrary, we have that for each $\epsilon > 0$, $d_Y(y_a, y_b) < \epsilon$. Hence $d_Y(y_a, y_b) = 0$ and

$$\begin{aligned} \lim_{n \rightarrow \infty} f_0(a_n) &= y_a \\ &= y_b \\ &= \lim_{n \rightarrow \infty} f_0(b_n). \end{aligned}$$

Since $x \in S$ and $(a_n)_{n \in \mathbb{N}}, (b_n)_{n \in \mathbb{N}} \in E_x$ are arbitrary, we have that for each $x \in S$ and $(a_n)_{n \in \mathbb{N}}, (b_n)_{n \in \mathbb{N}} \in E_x$, $\lim_{n \rightarrow \infty} f_0(a_n) = \lim_{n \rightarrow \infty} f_0(b_n)$.

- Let $x \in S$. Define $(x_n)_{n \in \mathbb{N}} \in E_x$ by $x_n := x$. Then from above, we have that

$$\begin{aligned} f(x) &= \lim_{n \rightarrow \infty} f_0(\phi(x)_n) \\ &= \lim_{n \rightarrow \infty} f_0(x_n) \\ &= f_0(x). \end{aligned}$$

Since $x \in S$ is arbitrary, we have that for each $x \in S$, $f(x) = f_0(x)$. Hence $f|_S = f_0$.

- Let $g : X \rightarrow Y$. Suppose that g is uniformly continuous and $g|_S = f_0$. Let $x \in X$. Since $E_x \neq \emptyset$, there exists $(x_n)_{n \in \mathbb{N}} \in E_x$. Then from above, we have that

$$\begin{aligned} f(x) &= \lim_{n \rightarrow \infty} f_0(\phi(x)_n) \\ &= \lim_{n \rightarrow \infty} f_0(x_n) \\ &= \lim_{n \rightarrow \infty} g(x_n) \\ &= g(x). \end{aligned}$$

Since $x \in X$ is arbitrary, we have that for each $x \in X$, $f(x) = g(x)$. Thus $g = f$. Hence f is unique. □

Exercise 6.7.1.12. Cantor's Nested Set Theorem:

Let (X, d) be a complete metric space and $(C_n)_{n \in \mathbb{N}} \subset \mathcal{P}(X)$. Suppose that

- for each $n \in \mathbb{N}$, C_n is closed, $C_n \neq \emptyset$ and $C_{n+1} \subset C_n$,
- $\text{diam } C_n \rightarrow 0$.

Then there exists $x \in X$ such that $\bigcap_{n \in \mathbb{N}} C_n = \{x\}$.

Hint: Use $(C_n)_{n \in \mathbb{N}}$ to obtain a Cauchy sequence.

Proof.

- Since for each $n \in \mathbb{N}$, $C_n \neq \emptyset$, the axiom of choice implies that there exists $(x_n)_{n \in \mathbb{N}}$ such that for each $n \in \mathbb{N}$, $x_n \in C_n$. Let $\epsilon > 0$. Since $\text{diam } C_n \rightarrow 0$, there exists $N \in \mathbb{N}$ such that for each $n \in \mathbb{N}$, $n \geq N$ implies that $\text{diam } C_n < \epsilon$. Let $m, n \in \mathbb{N}$. Suppose that $m, n \geq N$.

- Suppose that $m \geq n$. Then $x_n \in C_n$ and

$$\begin{aligned} x_m &\in C_m \\ &\subset C_n. \end{aligned}$$

Hence

$$\begin{aligned} d(x_m, x_n) &\leq \text{diam } C_n \\ &< \epsilon. \end{aligned}$$

- Similarly, if $n > m$, then $d(x_m, x_n) < \epsilon$.

Since $m, n \in \mathbb{N}$ such that $m, n \geq N$ are arbitrary, we have that for each $m, n \in \mathbb{N}$, $m, n \geq N$ implies that $d(x_m, x_n) < \epsilon$. Since $\epsilon > 0$ is arbitrary, we have that for each $\epsilon > 0$, there exists $N \in \mathbb{N}$ such that for each $m, n \in \mathbb{N}$, $m, n \geq N$ implies that $d(x_m, x_n) < \epsilon$. Thus $(x_n)_{n \in \mathbb{N}}$ is Cauchy. Since (X, d) is complete, there exists $x \in X$ such that $x_n \rightarrow x$.

- – For the sake of contradiction, suppose that $x \notin \bigcap_{n \in \mathbb{N}} C_n$. Then $x \in \bigcup_{n \in \mathbb{N}} C_n^c$. Hence there exists $N_1 \in \mathbb{N}$ such that $x \in C_{N_1}^c$. Since for each $n \in \mathbb{N}$, $C_{n+1} \subset C_n$, we have that for each $n \in \mathbb{N}$, $n \geq N_1$ implies that

$$\begin{aligned} x &\in C_{N_1}^c \\ &\subset C_n^c. \end{aligned}$$

Since C_{N_1} is closed, Exercise 6.1.2.15 implies that $d(x, C_{N_1}) > 0$. Set $\epsilon := d(x, C_{N_1})$. We note that for each $n \in \mathbb{N}$, $n \geq N_1$ implies that $\{d(x, y) : y \in C_n\} \subset \{d(x, y) : y \in C_{N_1}\}$ and therefore

$$\begin{aligned} d(x, C_n) &= \inf\{d(x, y) : y \in C_n\} \\ &\geq \inf\{d(x, y) : y \in C_{N_1}\} \\ &= d(x, C_{N_1}) \\ &= \epsilon. \end{aligned}$$

Since $x_n \rightarrow x$, there exists $N_2 \in \mathbb{N}$ such that for each $n \in \mathbb{N}$, $n \geq N_2$ implies that $d(x_n, x) < \epsilon$. Set $N := \max(N_1, N_2)$. Then

$$\begin{aligned} \epsilon &= d(x, C_{N_1}) \\ &\leq d(x, C_N) \\ &\leq d(x, x_N) \\ &< \epsilon, \end{aligned}$$

which is a contradiction. Hence $\{x\} \subset \bigcap_{n \in \mathbb{N}} C_n$.

- For the sake of contradiction, suppose that $\bigcap_{n \in \mathbb{N}} C_n \not\subset \{x\}$. Then there exists $x_0 \in \bigcap_{n \in \mathbb{N}} C_n$ such that $x_0 \neq x$. Set $\epsilon := d(x, x_0)$. Then $\epsilon > 0$. Since $\text{diam } C_n \rightarrow 0$, there exists $N \in \mathbb{N}$ such that for each $n \in \mathbb{N}$, $n \geq N$ implies that $\text{diam } C_n < \epsilon$. Since $x, x_0 \in \bigcap_{n \in \mathbb{N}} C_n$, we have that $x, x_0 \in C_N$ and therefore

$$\begin{aligned} \epsilon &= d(x, x_0) \\ &\leq \text{diam } C_N \\ &< \epsilon, \end{aligned}$$

which is a contradiction. Hence $\bigcap_{n \in \mathbb{N}} C_n \subset \{x\}$.

Since $\{x\} \subset \bigcap_{n \in \mathbb{N}} C_n$ and $\bigcap_{n \in \mathbb{N}} C_n \subset \{x\}$, we have that $\bigcap_{n \in \mathbb{N}} C_n = \{x\}$.

□

pulling from [9]

Exercise 6.7.1.13. Banach Fixed-Point Theorem: Let (X, d) be a complete metric space and $f \in C_{\text{Lip}}(X, X)$. Suppose that $X \neq \emptyset$. Define $\rho \in \mathbb{R}_+$ by $\rho := \text{Lip}(f)$. If f is a (d_X, d_Y) -contraction mapping, then

1. for each $x_0 \in X$,

- (a) for each $m, n \in \mathbb{N}$, $m \leq n$ implies that

$$d(x_m, x_n) < \rho^m \frac{d(x_0, x_1)}{1 - \rho},$$

- (b) $(f^n(x_0))_{n \in \mathbb{N}}$ is Cauchy

2. there exists a unique $x_* \in X$ such that

- (a) for each $x_0 \in X$, $f^n(x_0) \rightarrow x_*$

$$(b) \ f(x_*) = x_*$$

3. for each $x_0 \in X$ and $n \in \mathbb{N}$,

$$(a) \ d(x_n, x_*) \leq \rho^n(1 - \rho)^{-1}d(x_0, x_1),$$

$$(b) \ d(x_{n+1}, x_*) \leq \rho(1 - \rho)^{-1}d(x_n, x_{n+1}).$$

Proof.

1. Let $x_0 \in X$. Define $(x_n)_{n \in \mathbb{N}} \subset X$ by $x_n := f^n(x_0)$.

(a) For each $m, n \in \mathbb{N}$, $m \leq n$ implies that

$$\begin{aligned} d(x_m, x_n) &= d(f^m(x_0), f^m(x_{n-m})) \\ &\leq \rho^m d(x_0, x_{n-m}) \\ &\leq \rho^m \frac{1 - \rho^{n-m}}{1 - \rho} d(x_0, x_1) \quad (\text{by Exercise 6.1.3.14}) \\ &< \rho^m \frac{d(x_0, x_1)}{1 - \rho}. \end{aligned}$$

(b) Let $\epsilon > 0$. Since $\rho^n \rightarrow 0$, there exists $N \in \mathbb{N}$ such that for each $n \in \mathbb{N}$, $n \geq N$ implies that $\rho^n < \epsilon(1 - \rho)[d(x_0, x_1) + 1]^{-1}$. Let $m, n \in \mathbb{N}$. Suppose that $m, n \geq N$.

• If $m \leq n$, then

$$\begin{aligned} d(x_m, x_n) &< \rho^m \frac{d(x_0, x_1)}{1 - \rho} \quad (\text{by prev part}) \\ &< \left(\frac{(1 - \rho)\epsilon}{d(x_0, x_1) + 1} \right) \left(\frac{d(x_0, x_1)}{1 - \rho} \right) \\ &= \left(\frac{(1 - \rho)\epsilon}{1 - \rho} \right) \left(\frac{d(x_0, x_1)}{d(x_0, x_1) + 1} \right) \\ &< \epsilon. \end{aligned}$$

• If $m > n$, the similarly, $d(x_m, x_n) < \epsilon$.

Since $\epsilon > 0$ is arbitrary, we have that for each $\epsilon > 0$, there exists $N \in \mathbb{N}$ such that for each $m, n \in \mathbb{N}$, $m, n \geq N$ implies that $d(x_m, x_n) < \epsilon$. Thus $(x_n)_{n \in \mathbb{N}}$ is Cauchy. Since $x_0 \in X$ is arbitrary, we have that for each $x_0 \in X$, $(f^n(x_0))_{n \in \mathbb{N}}$ is Cauchy.

2. • **Existence:**

(a) Since $X \neq \emptyset$, there exists $x \in X$. The prev part implies that $(f^n(x))_{n \in \mathbb{N}}$ is Cauchy. Since (X, d) is complete, there exists $x_* \in X$ such that $f^n(x) \rightarrow x_*$. Let $x_0 \in X$ and $\epsilon > 0$. Since $\rho^n \rightarrow 0$, there exists $N_1 \in \mathbb{N}$ such that for each $n \in \mathbb{N}$, $n \geq N_1$ implies that $\rho^n < \epsilon[2(d(x_0, x) + 1)]^{-1}$. Since $f^n(x) \rightarrow x_*$, there exists $N_2 \in \mathbb{N}$ such that for each $n \in \mathbb{N}$, $n \geq N_2$ implies that $d(f^n(x), x_*) < \epsilon/2$. Define $N \in \mathbb{N}$ by $N := \max\{N_1, N_2\}$. Then for each $n \in \mathbb{N}$, $n \geq N$ implies that

$$\begin{aligned} d(f^n(x_0), x_*) &\leq d(f^n(x_0), f^n(x)) + d(f^n(x), x_*) \\ &\leq \rho^n d(x_0, x) + d(f^n(x), x_*) \\ &< \frac{\epsilon}{2} \left(\frac{d(x_0, x)}{d(x_0, x) + 1} \right) + \frac{\epsilon}{2} \\ &< \frac{\epsilon}{2} + \frac{\epsilon}{2} \\ &= \epsilon. \end{aligned}$$

Since $\epsilon > 0$ is arbitrary, we have that for each $\epsilon > 0$, there exists $N \in \mathbb{N}$ such that for each $n \in \mathbb{N}$, $n \geq N$ implies that $d(f^n(x_0), x_*) < \epsilon$. Hence $f^n(x_0) \rightarrow x_*$. Since $x_0 \in X$ is arbitrary, we have that for each $x_0 \in X$, $f^n(x_0) \rightarrow x_*$.

- (b) **The prev part** implies that for each $x_0 \in X$, $f^n(x_0) \rightarrow x_*$. In particular, $f^n(x_*) \rightarrow x_*$. Let $\epsilon > 0$. Since $f^n(x_*) \rightarrow x_*$, there exists $N \in \mathbb{N}$ such that for each $n \in \mathbb{N}$, $n \geq N$ implies that $d(f^n(x_*), x_*) < \epsilon/2$. Then

$$\begin{aligned} d(f(x_*), x_*) &\leq d(f(x_*), f^{N+1}(x_*)) + d(f^{N+1}(x_*), x_*) \\ &\leq \rho d(x_*, f^N(x_*)) + d(f^{N+1}(x_*), x_*) \\ &< \frac{\epsilon}{2} + \frac{\epsilon}{2} \\ &= \epsilon. \end{aligned}$$

Since $\epsilon > 0$ is arbitrary, we have that for each $\epsilon > 0$, $d(f(x_*), x_*) < \epsilon$. Hence

$$\begin{aligned} 0 &\leq d(f(x_*), x_*) \\ &\leq 0. \end{aligned}$$

Thus $d(f(x_*), x_*) = 0$ and $f(x_*) = x_*$.

• **Uniqueness:**

Let $x' \in X$. Suppose that

- for each $x_0 \in X$, $f^n(x_0) \rightarrow x'$
- $f(x') = x'$

In particular, $f^n(x') \rightarrow x'$. **The prev existence part** implies that $f^n(x') \rightarrow x_*$. Since $(X, \mathcal{T}_d)$ is Hausdorff, $x' = x_*$.

3. **FINISH!!!**

□

6.7.2 Completeness and Subspaces

Exercise 6.7.2.1. Let (X, d) be a metric space, $E \subset X$ and $(a_n)_{n \in \mathbb{N}} \subset E$. Then $(a_n)_{n \in \mathbb{N}}$ is Cauchy in $(E, d|_{E \times E})$ iff $(a_n)_{n \in \mathbb{N}}$ is Cauchy in (X, d) .

Proof. Set $d|_E := d|_{E \times E}$.

• **($\implies$):**

Suppose that $(a_n)_{n \in \mathbb{N}}$ is Cauchy in $(E, d|_{E \times E})$. Let $\epsilon > 0$. Since $(a_n)_{n \in \mathbb{N}}$ is Cauchy in $(E, d|_E)$, there exists $N \in \mathbb{N}$ such that for each $m, n \in \mathbb{N}$, $m, n \geq N$ implies that $d|_E(a_m, a_n) < \epsilon$. Thus for each $m, n \in \mathbb{N}$, $m, n \geq N$ implies that

$$\begin{aligned} d(a_m, a_n) &= d|_E(a_m, a_n) \\ &< \epsilon \end{aligned}$$

Since $\epsilon > 0$ is arbitrary, we have that for each $\epsilon > 0$, there exists $N \in \mathbb{N}$ such that for each $m, n \in \mathbb{N}$, $m, n \geq N$ implies that $d(a_m, a_n) < \epsilon$. Hence $(a_n)_{n \in \mathbb{N}}$ is Cauchy in (X, d) .

• **($\impliedby$):**

Suppose that $(a_n)_{n \in \mathbb{N}}$ is Cauchy in (X, d) . Let $\epsilon > 0$. Since $(a_n)_{n \in \mathbb{N}}$ is Cauchy in (X, d) , there exists $N \in \mathbb{N}$ such that for each $m, n \in \mathbb{N}$, $m, n \geq N$ implies that $d(a_m, a_n) < \epsilon$. Thus for each $m, n \in \mathbb{N}$, $m, n \geq N$ implies that

$$\begin{aligned} d|_E(a_m, a_n) &= d(a_m, a_n) \\ &< \epsilon \end{aligned}$$

Since $\epsilon > 0$ is arbitrary, we have that for each $\epsilon > 0$, there exists $N \in \mathbb{N}$ such that for each $m, n \in \mathbb{N}$, $m, n \geq N$ implies that $d|_E(a_m, a_n) < \epsilon$. Hence $(a_n)_{n \in \mathbb{N}}$ is Cauchy in $(E, d|_E)$.

□

Exercise 6.7.2.2. Let (X, d) be a metric space and $C \subset X$. Suppose that (X, d) is complete. Then C is $\mathcal{T}_d$ -closed iff $(C, d|_{C \times C})$ is complete.

Proof.

- ($\Rightarrow$):

Suppose that C is $\mathcal{T}_d$ -closed. Set $d_C := d|_{C \times C}$. Let $(a_n)_{n \in \mathbb{N}} \subset C$. Suppose that $(a_n)_{n \in \mathbb{N}}$ is Cauchy in (C, d_C) . Exercise 6.7.2.1 implies that $(a_n)_{n \in \mathbb{N}}$ is Cauchy in (X, d) . Since (X, d) is complete, there exists $a \in X$ such that $a_n \rightarrow a$ in $(X, \mathcal{T}_d)$. Since C is $\mathcal{T}_d$ -closed, $a \in C$. Exercise 4.5.1.10 implies that $a_n \rightarrow a$ in $(C, \mathcal{T}_d \cap C)$. Exercise 6.4.1.2 then implies that $a_n \rightarrow a$ in $(C, \mathcal{T}_{d_C})$. Since $(a_n)_{n \in \mathbb{N}} \subset C$ is arbitrary, we have that for each $(a_n)_{n \in \mathbb{N}} \subset C$, $(a_n)_{n \in \mathbb{N}}$ is Cauchy in (C, d_C) implies that there exists $a \in C$ such that $a_n \rightarrow a$ in $(C, \mathcal{T}_{d_C})$. Hence (C, d_C) is complete.

- ($\Leftarrow$):

Suppose that (C, d_C) is complete. Let $(a_n)_{n \in \mathbb{N}} \subset C$ and $a \in X$. Suppose that $a_n \rightarrow a$ in $(X, \mathcal{T}_d)$. ?? then implies that $(a_n)_{n \in \mathbb{N}}$ is Cauchy. Since (C, d_C) is complete, there exists $a_0 \in C$ such that $a_n \rightarrow a_0$ in $(C, \mathcal{T}_{d_C})$. Exercise 6.4.1.2 and Exercise 4.5.1.10 then implies that $a_n \rightarrow a_0$ in $(X, \mathcal{T}_d)$. Since $(X, \mathcal{T}_d)$ is Hausdorff, $a_n \rightarrow a$ in $(X, \mathcal{T}_d)$ and $a_n \rightarrow a_0$ in $(X, \mathcal{T}_d)$, Exercise 4.12.1.7 implies that

$$\begin{aligned} a &= a_0 \\ &\in C. \end{aligned}$$

Since $(a_n)_{n \in \mathbb{N}} \subset C$ and $a \in X$ with $a_n \rightarrow a$ in $(X, \mathcal{T}_d)$ are arbitrary, we have that for each $(a_n)_{n \in \mathbb{N}} \subset C$ and $a \in X$, $a_n \rightarrow a$ in $(X, \mathcal{T}_d)$ implies that $a \in C$. Since $(X, \mathcal{T}_d)$ is first-countable, Exercise 4.13.1.4 implies that C is $\mathcal{T}_d$ -closed. □

6.7.3 Completeness and Product Spaces

Exercise 6.7.3.1. Let $(X_n, d_n)_{n \in \mathbb{N}}$ be a collection of metric spaces. Set $X := \prod_{n \in \mathbb{N}} X_n$ and define $d_X : X \times X \rightarrow [0, \infty)$ as in Definition 6.5.0.1. Then for each $(a_m)_{m \in \mathbb{N}} \subset X$, $(a_m)_{m \in \mathbb{N}}$ is Cauchy in (X, d_X) iff for each $n \in \mathbb{N}$, $(\pi_n(a_m))_{m \in \mathbb{N}}$ is Cauchy in (X_n, d_n) .

Proof. Define $\phi : [0, \infty) \rightarrow [0, \infty)$ as in Definition 6.5.0.1. Let $(a_m)_{m \in \mathbb{N}} \subset X$.

- Suppose that $(a_m)_{m \in \mathbb{N}}$ is Cauchy in (X, d_X) . Let $n \in \mathbb{N}$ and $\epsilon > 0$. Since $(a_m)_{m \in \mathbb{N}}$ is Cauchy in (X, d_X) , there exists $M \in \mathbb{N}$ such that for each $m_1, m_2 \in \mathbb{N}$, $m_1, m_2 \geq M$ implies that $d_X(a_{m_1}, a_{m_2}) < 2^{-n}\epsilon$. Let $m_1, m_2 \in \mathbb{N}$. Suppose that $m_1, m_2 \geq M$. Then

$$\begin{aligned} d_n^\phi(\pi_n(a_{m_1}), \pi_n(a_{m_2})) &\leq 2^n d_X(a_{m_1}, a_{m_2}) \\ &< \frac{2^n \epsilon}{2^n} \\ &= \epsilon. \end{aligned}$$

Thus $(\pi_n(a_m))_{m \in \mathbb{N}}$ is Cauchy in (X_n, d_n^ϕ) . Exercise 6.7.1.6 and Exercise 6.3.2.7 imply that $(\pi_n(a_m))_{m \in \mathbb{N}}$ is Cauchy in (X_n, d_n) .

- Conversely, suppose that for each $n \in \mathbb{N}$, $(\pi_n(a_m))_{m \in \mathbb{N}}$ is Cauchy in (X_n, d_n) . Exercise 6.7.1.6 and Exercise 6.3.2.7 imply that for each $n \in \mathbb{N}$, $(\pi_n(a_m))_{m \in \mathbb{N}}$ is Cauchy in (X_n, d_n^ϕ) . Let $\epsilon > 0$. Choose $N_0 \in \mathbb{N}$ such that $\sum_{n \geq N_0+1} 2^{-n} < \epsilon/2$. Since for each $n \in \mathbb{N}$, $(\pi_n(a_m))_{m \in \mathbb{N}}$ is Cauchy in (X_n, d_n^ϕ) , we have that for each $n \in [N_0]$, there exists $M_n \in \mathbb{N}$ such that for each $m_1, m_2 \in \mathbb{N}$, $m_1, m_2 \geq M_n$ implies that $d_n^\phi(\pi_n(a_{m_1}), \pi_n(a_{m_2})) < \epsilon/2$. Set $M := \max(M_1, \dots, M_{N_0})$. Let $m_1, m_2 \in \mathbb{N}$. Suppose

that $m_1, m_2 \geq M$. Then

$$\begin{aligned}
 d_X(a_{m_1}, a_{m_2}) &= \sum_{n \in \mathbb{N}} 2^{-n} d_n^\phi(\pi_n(a_{m_1}), \pi_n(a_{m_2})) \\
 &= \sum_{n=1}^{N_0} 2^{-n} d_n^\phi(\pi_n(a_{m_1}), \pi_n(a_{m_2})) + \sum_{n \geq N_0+1} 2^{-n} d_n^\phi(\pi_n(a_{m_1}), \pi_n(a_{m_2})) \\
 &< \sum_{n=1}^{N_0} 2^{-n} \left(\frac{\epsilon}{2} \right) + \sum_{n \geq N_0+1} 2^{-n} \\
 &= \frac{\epsilon}{2} \sum_{n=1}^{N_0} 2^{-n} + \sum_{n \geq N_0+1} 2^{-n} \\
 &< \frac{\epsilon}{2} + \frac{\epsilon}{2} \\
 &= \epsilon.
 \end{aligned}$$

Since $\epsilon > 0$ is arbitrary, we have that for each $\epsilon > 0$, there exists $M \in \mathbb{N}$ such that for each $m_1, m_2 \in \mathbb{N}$, $m_1, m_2 \geq M$ implies that $d_X(a_{m_1}, a_{m_2}) < \epsilon$. Hence $(a_m)_{m \in \mathbb{N}}$ is Cauchy in (X, d_X) .

Thus (a_m) is Cauchy in (X, d_X) iff for each $n \in \mathbb{N}$, $(\pi_n(a_m))_{m \in \mathbb{N}}$ is Cauchy in (X_n, d_n) . Since $(a_m)_{m \in \mathbb{N}} \subset X$ is arbitrary, we have that for each $(a_m)_{m \in \mathbb{N}} \subset X$, $(a_m)_{m \in \mathbb{N}}$ is Cauchy in (X, d_X) iff for each $n \in \mathbb{N}$, $(\pi_n(a_m))_{m \in \mathbb{N}}$ is Cauchy in (X_n, d_n) . $\square$

Exercise 6.7.3.2. Let $(X_n, d_n)_{n \in \mathbb{N}}$ be a collection of metric spaces. Set $X := \prod_{n \in \mathbb{N}} X_n$ and define $d_X : X \times X \rightarrow [0, \infty)$ as in Definition 6.5.0.1. Then (X, d_X) is complete iff for each $n \in \mathbb{N}$, (X_n, d_n) is complete.

Proof. Define $\phi : [0, \infty) \rightarrow [0, 1)$ and $\mathcal{T}$ as in Definition 6.5.0.1.

- Suppose that (X, d_X) is complete. Let $n \in \mathbb{N}$ and $(x_m)_{m \in \mathbb{N}} \subset X_n$. Suppose that $(x_m)_{m \in \mathbb{N}}$ is Cauchy in (X_n, d_n) . Choose $a_0 \in X$. Define $(a_m)_{m \in \mathbb{N}} \subset X$ by

$$\pi_k(a_m) = \begin{cases} \pi_k(a_0), & k \neq n \\ x_m, & k = n \end{cases}$$

Let $\epsilon > 0$. Since $(x_m)_{m \in \mathbb{N}}$ is Cauchy in (X_n, d_n) , Exercise 6.7.1.6 implies that $(x_m)_{m \in \mathbb{N}}$ is Cauchy in (X_n, d_n^ϕ) . Thus there exists $M \in \mathbb{N}$ such that for each $m_1, m_2 \in \mathbb{N}$, $m_1, m_2 \geq M$ implies that $d_n^\phi(x_{m_1}, x_{m_2}) < 2^n \epsilon$. Let $m_1, m_2 \in \mathbb{N}$. Suppose that $m_1, m_2 \geq M$. Then

$$\begin{aligned}
 d_X(a_{m_1}, a_{m_2}) &= \sum_{k \in \mathbb{N}} 2^{-k} d_k^\phi(\pi_k(a_{m_1}), \pi_k(a_{m_2})) \\
 &= 2^{-n} d_n^\phi(\pi_n(a_{m_1}), \pi_n(a_{m_2})) + \sum_{k \neq n} 2^{-k} d_k^\phi(\pi_k(a_{m_1}), \pi_k(a_{m_2})) \\
 &= 2^{-n} d_n^\phi(x_{m_1}, x_{m_2}) + \sum_{k \neq n} 2^{-k} d_k^\phi(\pi_k(a_0), \pi_k(a_0)) \\
 &= 2^{-n} d_n^\phi(x_{m_1}, x_{m_2}) \\
 &< 2^{-n} (2^n \epsilon) \\
 &= \epsilon.
 \end{aligned}$$

Since $\epsilon > 0$ is arbitrary, we have that for each $\epsilon > 0$, there exists $M \in \mathbb{N}$ such that for each $m_1, m_2 \in \mathbb{N}$, $m_1, m_2 \geq M$ implies that $d_X(a_{m_1}, a_{m_2}) < \epsilon$. Therefore $(a_m)_{m \in \mathbb{N}}$ is Cauchy in (X, d_X) . Since (X, d_X) is complete, there exists $a \in X$ such that $a_m \rightarrow a$ in $(X, \mathcal{T}_{d_X})$. Exercise 6.5.0.2 implies that $a_m \rightarrow a$ in $(X, \mathcal{T})$. Define $x \in X_n$ by $x := \pi_n(a)$. Exercise 4.6.2.2 implies that

$$\begin{aligned}
 x_m &= \pi_n(a_m) \\
 &\rightarrow \pi_n(a) \\
 &= x
 \end{aligned}$$

in $(X_n, \mathcal{T}_{d_n})$. Since $(x_m)_{m \in \mathbb{N}} \subset X_n$ with $(x_m)_{m \in \mathbb{N}}$ Cauchy is arbitrary, we have that for each $(x_m)_{m \in \mathbb{N}} \subset X_n$, if $(x_m)_{m \in \mathbb{N}}$ is Cauchy, then there exists $x \in X_n$ such that $x_m \rightarrow x$ in $(X, \mathcal{T}_{d_n})$. Hence (X_n, d_n) is Complete. Since $n \in \mathbb{N}$ is arbitrary, we have that for each $n \in \mathbb{N}$, (X_n, d_n) is complete.

- Conversely, suppose that for each $n \in \mathbb{N}$, (X_n, d_n) is complete. Let $(a_m)_{m \in \mathbb{N}} \subset X$. Suppose that $(a_m)_{m \in \mathbb{N}}$ is Cauchy in (X, d_X) . Exercise 6.7.3.1 implies that for each $n \in \mathbb{N}$, $(\pi_n(a_m))_{m \in \mathbb{N}}$ is Cauchy in (X_n, d_n) . Since for each $n \in \mathbb{N}$, (X_n, d_n) is complete, we have that for each $n \in \mathbb{N}$, there exists $x_n \in X_n$ such that $\pi_n(a_m) \xrightarrow{m} x_n$ in $(X_n, \mathcal{T}_{d_n})$. Define $a \in X$ by $a := (x_n)_{n \in \mathbb{N}}$. Exercise 4.6.2.2 implies that $a_m \rightarrow a$ in $(X, \mathcal{T})$. Exercise 6.5.0.2 implies that $a_m \rightarrow a$ in $(X, \mathcal{T}_{d_X})$. Since $(a_m)_{m \in \mathbb{N}} \subset X$ is arbitrary, we have that for each $(a_m)_{m \in \mathbb{N}} \subset X$, if $(a_m)_{m \in \mathbb{N}}$ is Cauchy in (X, d_X) , then there exists $a \in X$ such that $a_m \rightarrow a$ in $(X, \mathcal{T}_{d_X})$. Hence (X, d_X) is complete. □

6.7.4 Completeness and Coproduct Spaces

Exercise 6.7.4.1. Let $(X_\alpha, d_\alpha)_{\alpha \in A}$ be a collection of metric spaces. Set $X := \coprod_{\alpha \in A} X_\alpha$ and define $\phi : [0, \infty) \rightarrow [0, 1)$ and $d_X : X \times X \rightarrow [0, \infty)$ as in Definition 6.6.0.1. Then for each $(\alpha_j, x_j)_{j \in \mathbb{N}} \subset X$, $(\alpha_j, x_j)_{j \in \mathbb{N}}$ is Cauchy in (X, d_X) iff there exists $\alpha_0 \in A$ and $j_0 \in \mathbb{N}$ such that

1. for each $j \in \mathbb{N}$, $j \geq j_0$ implies that $\alpha_j = \alpha_0$ and $x_j \in X_{\alpha_0}$,
2. $(x_{j'+j_0})_{j' \in \mathbb{N}}$ is Cauchy in $(X_{\alpha_0}, d_{\alpha_0})$.

Proof. Let $(\alpha_j, x_j)_{j \in \mathbb{N}} \subset X$.

- ($\implies$) :

Suppose that $(\alpha_j, x_j)_{j \in \mathbb{N}}$ is Cauchy in (X, d_X) . Set $\epsilon_0 := 1/2$. Since $(\alpha_j, x_j)_{j \in \mathbb{N}}$ is Cauchy in (X, d_X) , there exists $j_0 \in \mathbb{N}$ such that for each $j, k \in \mathbb{N}$, $j, k \geq j_0$ implies that $d_X((\alpha_j, x_j), (\alpha_k, x_k)) < \epsilon_0$. Set $\alpha_0 := \alpha_{j_0}$.

1. Since $\epsilon_0 < 1$, we have that for each $j \in \mathbb{N}$, $j \geq j_0$ implies that $\alpha_j = \alpha_0$ and therefore $x_j \in X_{\alpha_0}$.
2. Define $(y_j)_{j \in \mathbb{N}} \subset X_{\alpha_0}$ by $y_j := x_{j+j_0}$. Let $\epsilon > 0$. Since $(\alpha_j, x_j)_{j \in \mathbb{N}}$ is Cauchy in $(X, \mathcal{T}_X)$, there exists $N_0 \in \mathbb{N}$ such that for each $j, k \in \mathbb{N}$, $j, k \geq N_0$ implies that $d_X((\alpha_j, x_j), (\alpha_k, x_k)) < \epsilon$. Set $N := \max(j_0 + 1, N_0) - j_0$. Then for each $j, k \in \mathbb{N}$, $j, k \geq N$ implies that $j + j_0, k + j_0 \geq N_0$ and

$$\begin{aligned} d_{\alpha_0}^{\phi}(y_j, y_k) &= d_{\alpha_0}^{\phi}(x_{j+j_0}, x_{k+j_0}) \\ &= d_X((\alpha_j, x_{j+j_0}), (\alpha_k, x_{k+j_0})) \\ &< \epsilon. \end{aligned}$$

Thus $(y_j)_{j \in \mathbb{N}}$ is Cauchy in $(X_{\alpha_0}, d_{\alpha_0}^{\phi})$. Exercise 6.7.1.6 implies that $(y_j)_{j \in \mathbb{N}}$ is Cauchy in $(X_{\alpha_0}, d_{\alpha_0})$.

- ($\impliedby$) :

Conversely, suppose that there exists $\alpha_0 \in A$ and $j_0 \in \mathbb{N}$ such that

1. for each $j \in \mathbb{N}$, $j \geq j_0$ implies that $\alpha_j = \alpha_0$ and $x_j \in X_{\alpha_0}$,
2. $(x_{j'+j_0})_{j' \in \mathbb{N}}$ is Cauchy in $(X_{\alpha_0}, d_{\alpha_0})$.

Define $(y_j)_{j \in \mathbb{N}} \subset X_{\alpha_0}$ by $y_j := x_{j+j_0}$. Since $(y_j)_{j \in \mathbb{N}}$ is Cauchy in $(X_{\alpha_0}, d_{\alpha_0})$, we have that $(y_j)_{j \in \mathbb{N}}$ is Cauchy in $(X_{\alpha_0}, d_{\alpha_0}^{\phi})$. Let $\epsilon > 0$. Since $(y_j)_{j \in \mathbb{N}}$ is Cauchy in $(X_{\alpha_0}, d_{\alpha_0}^{\phi})$, there exists $N_0 \in \mathbb{N}$ such that for each $j, k \in \mathbb{N}$, $j, k \geq N_0$ implies that $d_{\alpha_0}^{\phi}(y_j, y_k) < \epsilon$. Set $N := \max(j_0, N_0) + j_0$. Then for each $j, k \in \mathbb{N}$, $j, k \geq N$ implies that $j - j_0, k - j_0 \geq N_0, j_0$. Thus

$$\begin{aligned} d_X((\alpha_j, x_j), (\alpha_k, x_k)) &= d^{\phi}(x_j, x_k) \\ &= d^{\phi}(y_{j-j_0}, y_{k-j_0}) \\ &< \epsilon. \end{aligned}$$

Hence $(\alpha_j, x_j)_{j \in \mathbb{N}}$ is Cauchy in (X, d_X) . □

Exercise 6.7.4.2. Let $(X_\alpha, d_\alpha)_{\alpha \in A}$ be a collection of metric spaces. Set $X := \coprod_{\alpha \in A} X_\alpha$ and define $\phi : [0, \infty) \rightarrow [0, 1)$ and $d_X : X \times X \rightarrow [0, \infty)$ as in Definition 6.6.0.1. Then (X, d_X) is complete iff for each $\alpha \in A$, (X_α, d_α) is complete.

Proof.

• $(\implies)$:

Suppose that (X, d_X) is complete. Let $\alpha \in A$ and $(x_n)_{n \in \mathbb{N}} \subset X_\alpha$. Suppose that $(x_n)_{n \in \mathbb{N}}$ is Cauchy in (X_α, d_α) . Exercise 6.6.0.3 implies that $\iota_\alpha : X_\alpha \rightarrow X$ is Lipschitz. Therefore ι_α is uniformly continuous and Exercise 6.7.1.5 implies that $(\iota_\alpha(x_n))_{n \in \mathbb{N}}$ is Cauchy in (X, d_X) . Since (X, d_X) is complete, there exists $(\alpha_0, x_0) \in X$ such that $(\iota_\alpha(x_n)) \rightarrow (\alpha_0, x_0)$ in (X, d_X) . Exercise 6.6.0.2 and Exercise 4.7.0.8 imply that $x_n \rightarrow x_0$. Since $(x_n)_{n \in \mathbb{N}} \subset X_\alpha$ with $(x_n)_{n \in \mathbb{N}}$ Cauchy in (X_α, d_α) is arbitrary, we have that for each $(x_n)_{n \in \mathbb{N}} \subset X_\alpha$, if $(x_n)_{n \in \mathbb{N}}$ is Cauchy, then there exists $x_0 \in X_\alpha$ such that $x_n \rightarrow x_0$. Hence (X_α, d_α) is complete. Since $\alpha \in A$ is arbitrary, we have that for each $\alpha \in A$, (X_α, d_α) is complete.

• $(\impliedby)$:

Conversely, suppose that for each $\alpha \in A$, (X_α, d_α) is complete. Let $(\alpha_n, x_n)_{n \in \mathbb{N}} \subset X$. Suppose that $(\alpha_n, x_n)_{n \in \mathbb{N}}$ is Cauchy in (X, d_X) . Exercise 6.7.4.1 implies that there exists $\alpha_0 \in A$ and $j_0 \in \mathbb{N}$ such that

1. for each $j \in \mathbb{N}$, $j \geq j_0$ implies that $\alpha_j = \alpha_0$ and $x_j \in X_{\alpha_0}$,
2. $(x_{j'+j_0})_{j' \in \mathbb{N}}$ is Cauchy in $(X_{\alpha_0}, d_{\alpha_0})$.

Define $(y_j)_{j \in \mathbb{N}} \subset X_{\alpha_0}$ by $y_j := x_{j+j_0}$. Since $(X_{\alpha_0}, d_{\alpha_0})$ is complete, and $(y_j)_{j \in \mathbb{N}}$ is Cauchy in $(X_{\alpha_0}, d_{\alpha_0})$, there exists $x_0 \in X_{\alpha_0}$ such that $y_n \rightarrow x_0$ in $(X, \mathcal{T}_{d_{\alpha_0}})$. Thus $y_n \rightarrow x_0$ in $(X, \mathcal{T}_{d_{\alpha_0}^\phi})$. Let $\epsilon > 0$. Since $y_n \rightarrow x_0$ in $(X, \mathcal{T}_{d_{\alpha_0}^\phi})$, there exists $N_0 \in \mathbb{N}$ such that for each $n \in \mathbb{N}$, $n \geq N_0$ implies that $d_{\alpha_0}^\phi(y_n, x_0) < \epsilon$. Set $N := \max(j_0, N_0) + j_0$. Let $n \in \mathbb{N}$. Suppose that $n \geq N$. Then $n - j_0 \geq j_0, N_0$ and

$$\begin{aligned} d_X((\alpha_n, x_n), (\alpha_0, x_0)) &= d_X((\alpha_0, x_n), (\alpha_0, x_0)) \\ &= d^\phi(x_n, x_0) \\ &= d^\phi(y_{n-j_0}, x_0) \\ &< \epsilon. \end{aligned}$$

Hence $(\alpha_n, x_n) \rightarrow (\alpha_0, x_0)$ in $(X, \mathcal{T}_{d_X})$. Since $(\alpha_n, x_n)_{n \in \mathbb{N}} \subset X$ with $(\alpha_n, x_n)_{n \in \mathbb{N}}$ Cauchy is arbitrary, we have that for each $(\alpha_n, x_n)_{n \in \mathbb{N}} \subset X$, $(\alpha_n, x_n)_{n \in \mathbb{N}}$ is Cauchy implies that there exists $(\alpha_0, x_0) \in X$ such that $(\alpha_n, x_n) \rightarrow (\alpha_0, x_0)$ in $(X, \mathcal{T}_{d_X})$. Thus $(X, \mathcal{T}_{d_X})$ is complete. □

6.8 Compactness

Definition 6.8.0.1. Let (X, d) be a metric space. Then (X, d) is said to be **totally bounded** if for each $\epsilon > 0$, there exists $\mathcal{U} \subset \mathcal{T}_d$ such that $\mathcal{U}$ is an open cover of X , $\mathcal{U}$ is finite and for each $U \in \mathcal{U}$, $\text{diam}(U) \leq \epsilon$.

Exercise 6.8.0.2. Let (X, d) be a metric space. Suppose that $X \neq \emptyset$. Then X is totally bounded iff for each $\epsilon > 0$, there exists $n \in \mathbb{N}$ and $(x_j)_{j=1}^n \subset X$ such that $X \subset \bigcup_{j=1}^n B(x_j, \epsilon)$.

Proof.

• $(\implies)$:

Suppose that X is totally bounded in (X, d) . Let $\epsilon > 0$. Define $\delta > 0$ by $\delta := \epsilon/2$. Then, there exists $\mathcal{U}' \subset \mathcal{T}_d$ such that $\mathcal{U}'$ is an open cover of X , $\mathcal{U}'$ is finite and for each $U \in \mathcal{U}'$, $\text{diam}(U) \leq \delta$. Define $\mathcal{U} \subset \mathcal{U}'$ by $\mathcal{U} := \{U \in \mathcal{U}' : U \neq \emptyset\}$. Since $X \neq \emptyset$, $\mathcal{U} \neq \emptyset$. Define $n \in \mathbb{N}$ by $n := \#\mathcal{U}$. Write $\mathcal{U} = (U_j)_{j=1}^n$. Since for each $j \in [n]$, $U_j \neq \emptyset$, there exists $(x_j)_{j=1}^n \subset X$ such that for

each $j \in [n]$, $x_j \in U_j$. Let $j \in [n]$ and $y \in U_j$. Then

$$\begin{aligned} d(x_j, y) &\leq \sup_{a, b \in U_j} d(a, b) \\ &= \text{diam}(U_j) \\ &\leq \delta \\ &= \frac{\epsilon}{2} \\ &< \epsilon. \end{aligned}$$

Hence $y \in B(x_j, \epsilon)$. Since $y \in U_j$ is arbitrary, we have that for each $y \in U_j$, $y \in B(x_j, \epsilon)$. Thus $U_j \subset B(x_j, \epsilon)$. Since $j \in [n]$ is arbitrary, we have that for each $j \in [n]$, $U_j \subset B(x_j, \epsilon)$. Then

$$\begin{aligned} X &\subset \bigcup_{j=1}^n U_j \\ &\subset \bigcup_{j=1}^n B(x_j, \epsilon). \end{aligned}$$

• ($\Leftarrow$) :

Suppose that for each $\epsilon > 0$, there exists $n \in \mathbb{N}$ and $(x_j)_{j=1}^n \subset X$ such that $X \subset \bigcup_{j=1}^n B(x_j, \epsilon)$. Let $\epsilon > 0$. Define $\delta > 0$ by

$\delta := \epsilon/2$. By assumption, there exists $n \in \mathbb{N}$ and $(x_j)_{j=1}^n \subset X$ such that $X \subset \bigcup_{j=1}^n B(x_j, \delta)$. We note that for each $j \in [n]$,

$$\begin{aligned} \text{diam } B(x_j, \delta) &\leq 2\delta \\ &= \epsilon. \end{aligned}$$

Define $\mathcal{U} \subset \mathcal{T}_d$ by $\mathcal{U} := \{B(x_j, \delta) : j \in [n]\}$. Then $\mathcal{U}$ is an open cover of X , $\mathcal{U}$ is finite and for each $U \in \mathcal{U}$, $\text{diam}(U) \leq \epsilon$. □

Definition 6.8.0.3. Let (X, d) be a metric space and $A \subset X$. Then A is said to be **totally bounded** in (X, d) if $(A, d|_{A^2})$ is totally bounded.

Exercise 6.8.0.4. Let (X, d) be a metric space and $A \subset X$. Suppose that $A \neq \emptyset$. Then A is totally bounded in (X, d) iff for each $\epsilon > 0$, there exists $n \in \mathbb{N}$ and $(x_j)_{j=1}^n \subset A$ such that $A \subset \bigcup_{j=1}^n B_d(x_j, \epsilon)$.

Proof.

• ($\Rightarrow$) :

Suppose that A is totally bounded in (X, d) . By definition, $(A, d|_A)$ is totally bounded. Let $\epsilon > 0$. Since $A \neq \emptyset$, Exercise 6.8.0.2 implies that there exists $n \in \mathbb{N}$ and $(x_j)_{j=1}^n \subset A$ such that $A \subset \bigcup_{j=1}^n B_{d|_A}(x_j, \epsilon)$. Exercise 6.4.1.3 then implies that

$$\begin{aligned} A &\subset \bigcup_{j=1}^n B_{d|_A}(x_j, \epsilon) \\ &= \bigcup_{j=1}^n B_d(x_j, \epsilon) \cap A \\ &\subset \bigcup_{j=1}^n B_d(x_j, \epsilon). \end{aligned}$$

Since $\epsilon > 0$ is arbitrary, we have that for each $\epsilon > 0$, there exists $n \in \mathbb{N}$ and $(x_j)_{j=1}^n \subset A$ such that $A \subset \bigcup_{j=1}^n B_d(x_j, \epsilon)$

• ($\Leftarrow$):

Suppose that for each $\epsilon > 0$, there exists $n \in \mathbb{N}$ and $(x_j)_{j=1}^n \subset A$ such that $A \subset \bigcup_{j=1}^n B_d(x_j, \epsilon)$. Let $\epsilon > 0$. Define $\delta > 0$ by $\delta := \epsilon/2$. By assumption, there exists $n \in \mathbb{N}$ and $(x_j)_{j=1}^n \subset A$ such that $A \subset \bigcup_{j=1}^n B_d(x_j, \delta)$. Define $\mathcal{U} \subset \mathcal{T}_{d|_A}$ by $\mathcal{U} := \{B_d(x_j, \delta) \cap A : j \in [n]\}$. Then

$$\begin{aligned} A &\subset \left[\bigcup_{j=1}^n B_d(x_j, \delta) \right] \cap A \\ &= \bigcup_{j=1}^n [B_d(x_j, \delta) \cap A] \\ &= \bigcup_{U \in \mathcal{U}} U. \end{aligned}$$

Let $U \in \mathcal{U}$. Then there exists $j \in [n]$ such that $U = B_d(x_j, \delta) \cap A$. Thus

$$\begin{aligned} \text{diam}_{d|_A} U &= \sup_{a, b \in U} d|_A(a, b) \\ &= \sup_{a, b \in B_d(x_j, \delta) \cap A} d(a, b) \\ &\leq \sup_{a, b \in B_d(x_j, \delta)} d(a, b) \\ &= \text{diam}_d B_d(x_j, \delta) \\ &\leq 2\delta \\ &= \epsilon. \end{aligned}$$

make and cite exercise about diameter and subspaces mentioned in Exercise 6.8.0.5 Hence $\mathcal{U}$ is an open cover of A , $\mathcal{U}$ is finite and for each $U \in \mathcal{U}$, $\text{diam}_{d|_A} U \leq \epsilon$. By definition, $(A, d|_A)$ is totally bounded and therefore A is totally bounded in (X, d) . $\square$

Exercise 6.8.0.5. Let (X, d) be a metric space and $A \subset X$. If (X, d) is totally bounded, then A is totally bounded in (X, d) .

Proof. Suppose that (X, d) is totally bounded. Let $\epsilon > 0$. Then there exists $\mathcal{U} \subset \mathcal{T}_d$ such that $\mathcal{U}$ is an open cover of X , $\mathcal{U}$ is finite and for each $U \in \mathcal{U}$, $\text{diam}(U) \leq \epsilon$.

- We note that Exercise 6.4.1.2 implies that $\mathcal{T}_d \cap A = \mathcal{T}_{d|_A}$ and Exercise 4.11.2.1 then implies that $\mathcal{U} \cap A$ is an open cover of A in $(A, \mathcal{T}_{d|_A})$.
- Since $\mathcal{U}$ is finite, $\mathcal{U} \cap A$ is finite.
- Since for each $U \in \mathcal{U}$, $\text{diam}_d(U) \leq \epsilon$, we have that for each $U \in \mathcal{U} \cap A$,

$$\begin{aligned} \text{diam}_{d|_A}(U) &= \sup_{a, b \in U} d|_A(a, b) \\ &= \sup_{a, b \in U} d(a, b) \\ &= \text{diam}_d(U). \end{aligned}$$

make exercise in metric subspace section about how in general for $U \in \mathcal{P}(X)$, $\text{diam}_{d|_A} U \cap A \leq \text{diam}_d U$ and for each $U \in \mathcal{P}(A)$, $\text{diam}_{d|_A}(U) = \text{diam}_d(U)$, then reference in this proof.

Therefore $(A, d|_A)$ is totally bounded and by definition, A is totally bounded in (X, d) . $\square$

Exercise 6.8.0.6. Let (X, d) be a metric space and $\mathcal{U} \subset \mathcal{T}_d$. Suppose that (X, d) is totally bounded, $X \neq \emptyset$ and $\mathcal{U} \in \text{Cover}^{\text{opn}}(X)$. If for each $\mathcal{U}_0 \subset \mathcal{U}$, $\mathcal{U}_0$ is finite implies that $\mathcal{U}_0 \notin \text{Cover}^{\text{opn}}(X)$, then there exists $(x_n)_{n \in \mathbb{N}} \subset X$, $(V_n)_{n \in \mathbb{N}} \subset \mathcal{T}_d$ and $(\mathcal{U}_n)_{n \in \mathbb{N}} \in \mathcal{P}(\mathcal{T}_d)^{\mathbb{N}}$ such that for each $n \in \mathbb{N}$,

- $x_n \in V_n$, $V_{n+1} \subset V_n$ and $\text{diam } V_n \leq 1/n$,
- $\mathcal{U}_n = \mathcal{U} \cap V_n$,
- for each $\mathcal{U}_0 \subset \mathcal{U}_n$, $\mathcal{U}_0$ is finite implies that $\mathcal{U}_0 \notin \text{Cover}^{\text{opn}}(V_n)$.

Proof. Suppose that for each $\mathcal{U}_0 \subset \mathcal{U}$, $\mathcal{U}_0$ is finite implies that $\mathcal{U}_0 \notin \text{Cover}^{\text{opn}}(X)$. We define $(x_n)_{n \in \mathbb{N}} \subset X$, $(V_n)_{n \in \mathbb{N}} \subset \mathcal{T}_d$ and $(\mathcal{U}_n)_{n \in \mathbb{N}} \in \mathcal{P}(\mathcal{T}_d)^{\mathbb{N}}$ inductively as follows:

• **Base Case:**

Set $r_1 := 2^{-1}$. Since (X, d) is totally bounded, Exercise 6.8.0.4 implies that there exists $k_1 \in \mathbb{N}$ and $(x_j^{(1)})_{j=1}^{k_1} \subset X$ such that $X \subset \bigcup_{j=1}^{k_1} B(x_j^{(1)}, r_1)$. Exercise 1.2.2.3 then implies that there exists $j_1 \in [k_1]$ such that for each $\mathcal{U}_0 \subset \mathcal{U}$, $\mathcal{U}_0$ is finite implies that $\mathcal{U}_0 \notin \text{Cover}^{\text{opn}}(B(x_{j_1}^{(1)}, r_1))$. Define $x_1 \in X$, $V_1 \in \mathcal{T}_d$ and $\mathcal{U}_1 \subset \mathcal{T}_d$ by $x_1 := x_{j_1}^{(1)}$, $V_1 := B(x_1, r_1)$ and $\mathcal{U}_1 := \mathcal{U} \cap V_1$. We note the following:

- By construction, $V_1 \in \mathcal{T}_d$ and

$$\begin{aligned} \text{diam } V_1 &\leq 2r_1 \\ &= 1. \end{aligned}$$

- Since (X, d) is totally bounded, Exercise 6.8.0.5 implies that V_1 is totally bounded in (X, d) .
- By construction, $x_1 \in V_1$ and therefore $V_1 \neq \emptyset$.
- By construction $\mathcal{U}_1 = \mathcal{U} \cap V_1$.
- Let $\mathcal{U}_0 \subset \mathcal{U}_1$. Suppose that $\mathcal{U}_0$ is finite. For the sake of contradiction, suppose that $\mathcal{U}_0 \in \text{Cover}^{\text{opn}}(V_1)$. Since $\mathcal{U}_1 = \mathcal{U} \cap V_1$, there exists $\mathcal{U}_0^* \subset \mathcal{U}$ such that $\mathcal{U}_0^*$ is finite and $\mathcal{U}_0 = \mathcal{U}_0^* \cap V_1$. Then

$$\begin{aligned} V_1 &\subset \bigcup_{U \in \mathcal{U}_0} U \\ &= \bigcup_{U \in \mathcal{U}_0^*} U \cap V_1 \\ &\subset \bigcup_{U \in \mathcal{U}_0^*} U. \end{aligned}$$

Hence $\mathcal{U}_0^* \in \text{Cover}^{\text{opn}}(V_1)$, which is a contradiction. Thus $\mathcal{U}_0 \notin \text{Cover}^{\text{opn}}(V_1)$. Since $\mathcal{U}_0 \subset \mathcal{U}_1$ such that $\mathcal{U}_0$ is finite is arbitrary, we have that for each $\mathcal{U}_0 \subset \mathcal{U}_1$, $\mathcal{U}_0$ is finite implies that $\mathcal{U}_0 \notin \text{Cover}^{\text{opn}}(V_1)$.

• **Induction Step:**

Let $n \geq 2$. Set $r_n := (2n)^{-1}$. Suppose that

- $V_{n-1} \in \mathcal{T}_d$,
- V_{n-1} is totally bounded in (X, d) ,
- $V_{n-1} \neq \emptyset$,
- $\mathcal{U}_{n-1} = \mathcal{U} \cap V_{n-1}$,
- for each $\mathcal{U}_0 \subset \mathcal{U}_{n-1}$, $\mathcal{U}_0$ is finite implies that $\mathcal{U}_0 \notin \text{Cover}^{\text{opn}}(V_{n-1})$.

Since V_{n-1} is totally bounded in (X, d) , Exercise 6.8.0.4 implies that there exists $k_n \in \mathbb{N}$ and $(x_j^{(n)})_{j=1}^{k_n} \subset V_{n-1}$ such that $V_{n-1} \subset \bigcup_{j=1}^{k_n} B(x_j^{(n)}, r_n)$. Since $\mathcal{U}_{n-1} = \mathcal{U} \cap V_{n-1}$, $V_{n-1} \in \mathcal{T}_d$ and $\mathcal{U} \in \text{Cover}^{\text{opn}}(X)$, we have that $\mathcal{U}_{n-1} \in \text{Cover}^{\text{opn}}(V_{n-1})$. Exercise 1.2.2.3 then implies that there exists $j_n \in [k_n]$ such that for each $\mathcal{U}_0 \subset \mathcal{U}_{n-1}$, $\mathcal{U}_0$ is finite implies that $\mathcal{U}_0 \notin \text{Cover}^{\text{opn}}(B(x_{j_n}^{(n)}, r_n))$. Define $x_n \in X$, $V_n \in \mathcal{T}_d$ and $\mathcal{U}_n \subset \mathcal{T}_d$ by $x_n := x_{j_n}^{(n)}$, $V_n := B(x_n, r_n) \cap V_{n-1}$ and $\mathcal{U}_n := \mathcal{U} \cap V_n$. We note the following:

- By construction $V_n \in \mathcal{T}_d$, $V_n \subset V_{n-1}$ and

$$\begin{aligned} \text{diam } V_n &\leq 2r_n \\ &= n^{-1}. \end{aligned}$$

- Since (X, d) is totally bounded, V_n is totally bounded in (X, d) .
- By construction, $x_n \in V_n$ and therefore $V_n \neq \emptyset$.
- By construction, $\mathcal{U}_n = \mathcal{U} \cap V_n$.
- Similarly to the base case, for each $\mathcal{U}_0 \subset \mathcal{U}_n$, $\mathcal{U}_0$ is finite implies that $\mathcal{U}_0 \notin \text{Cover}^{\text{opn}}(V_n)$. **maybe write out details**

□

Exercise 6.8.0.7. Let (X, d) be a metric space. The following are equivalent:

1. $(X, \mathcal{T}_d)$ is compact
2. $(X, \mathcal{T}_d)$ is sequentially compact
3. (X, d) is complete and totally bounded

rework with Exercise 6.8.0.4 in mind

Proof.

1. $(1) \implies (2)$:
Immediate by Exercise 4.14.8.3
2. $(2) \implies (3)$:
Suppose that $(X, \mathcal{T}_d)$ is sequentially compact.

• **(completeness):**

Let $(x_n)_{n \in \mathbb{N}} \subset X$. Suppose that $(x_n)_{n \in \mathbb{N}}$ is Cauchy. Since $(X, \mathcal{T}_d)$ is sequentially compact, there exists $(x_{n_k})_{k \in \mathbb{N}} \subset (x_n)_{n \in \mathbb{N}}$ and $x \in X$ such that $x_{n_k} \rightarrow x$. Exercise 6.7.1.4 implies that $x_n \rightarrow x$. Since $(x_n)_{n \in \mathbb{N}} \subset X$ with $(x_n)_{n \in \mathbb{N}}$ Cauchy is arbitrary, we have that for each $(x_n)_{n \in \mathbb{N}} \subset X$, $(x_n)_{n \in \mathbb{N}}$ is Cauchy implies that there exists $x \in X$ such that $x_n \rightarrow x$. Thus (X, d) is complete.

• **(total boundedness):**

If $X = \emptyset$, then X is totally bounded. Suppose that $X \neq \emptyset$. For the sake of contradiction, suppose that (X, d) is not totally bounded. Exercise 6.8.0.4 then implies that there exists $\epsilon > 0$ such that for each $n \in \mathbb{N}$ and $(x_j)_{j=1}^n \subset X$, $X \not\subset \bigcup_{j=1}^n B(x_j, \epsilon)$.

Define $(x_n)_{n \in \mathbb{N}} \subset X$ inductively as follows:

- Since $X \neq \emptyset$, there exists $a \in X$. Define $x_1 \in X$ by $x_1 := a$.
- Let $n \geq 2$. By assumption, $X \not\subset \bigcup_{j=1}^{n-1} B(x_j, \epsilon)$. Thus there exists $b \in X$ such that $x \in \bigcap_{j=1}^{n-1} B(x_j, \epsilon)^c$. Define $x_n \in X$ by $x_n := b$.

By construction, for each m, n , $m \neq n$ implies that $d(x_m, x_n) \geq \epsilon$. Since (X, d) is sequentially compact, there exists $(x_{n_k})_{k \in \mathbb{N}} \subset (x_n)_{n \in \mathbb{N}}$ and $x \in X$ such that $x_{n_k} \rightarrow x$. Exercise 6.7.1.3 implies that $(x_{n_k})_{k \in \mathbb{N}}$ is Cauchy. Thus there exists $K \in \mathbb{N}$ such that for each $k, l \in \mathbb{N}$, $k, l \geq K$ implies that $d(x_{n_k}, x_{n_l}) < \epsilon$. In particular,

$$\begin{aligned} \epsilon &\leq d(x_{n_K}, x_{n_{K+1}}) \\ &< \epsilon, \end{aligned}$$

which is a contradiction. Hence (X, d) is totally bounded.

3. $(3) \implies (1)$:

REWORK and make content about obtaining V_n, x_n a separate exercise to make proof shorter!!! Suppose that (X, d) is complete and totally bounded. If $X = \emptyset$, then $(X, \mathcal{T})$ is compact. Suppose that $X \neq \emptyset$. For the sake of contradiction, suppose that $(X, \mathcal{T}_d)$ is not compact. Then there exists $\mathcal{U} \in \text{Cover}^{\text{opn}}(X)$ such that for each $\mathcal{U}_0 \subset \mathcal{U}$, $\mathcal{U}_0$ is finite implies that $\mathcal{U}_0 \notin \text{Cover}^{\text{opn}}(X)$. Exercise 6.8.0.6 then implies that there exist $(x_n)_{n \in \mathbb{N}} \subset X$, $(V_n)_{n \in \mathbb{N}} \subset \mathcal{T}_d$ and $(\mathcal{U}_n)_{n \in \mathbb{N}} \in \mathcal{P}(\mathcal{T}_d)^{\mathbb{N}}$ such that for each $n \in \mathbb{N}$,

- $x_n \in V_n$, $V_{n+1} \subset V_n$ and $\text{diam } V_n \leq 1/n$,
- $\mathcal{U}_n = \mathcal{U} \cap V_n$,
- for each $\mathcal{U}_0 \subset \mathcal{U}_n$, $\mathcal{U}_0$ is finite implies that $\mathcal{U}_0 \notin \text{Cover}^{\text{opn}}(V_n)$.

Let $\epsilon > 0$. Choose $N \in \mathbb{N}$ such that $2/N < \epsilon$. Let $m, n \in \mathbb{N}$. Suppose that $m, n \geq N$. Then $x_m, x_n, x_N \in V_N$ and therefore

$$\begin{aligned} d(x_m, x_n) &\leq d(x_m, x_N) + d(x_N, x_n) \\ &\leq \text{diam}(V_N) + \text{diam}(V_N) \\ &\leq 1/N + 1/N \\ &= \frac{2}{N} \\ &< \epsilon. \end{aligned}$$

Since $\epsilon > 0$ is arbitrary, we have that for each $\epsilon > 0$, there exists $N \in \mathbb{N}$ such that for each $m, n \in \mathbb{N}$, $m, n \geq N$ implies that $d(x_m, x_n) < \epsilon$. Thus $(x_n)_{n \in \mathbb{N}}$ is Cauchy. Since (X, d) is complete, there exists $x \in X$ such that $x_n \rightarrow x$. Since for each $n \in \mathbb{N}$, $x_n \in V_n$ and $V_{n+1} \subset V_n$, we have that for each $n \in \mathbb{N}$, $x_n \in \text{cl } V_n$, $\text{cl } V_{n+1} \subset \text{cl } V_n$. Exercise 6.1.2.14 then implies that for each $n \in \mathbb{N}$,

$$\begin{aligned} \text{diam cl } V_n &= \text{diam } V_n \\ &\leq \frac{1}{n}. \end{aligned}$$

Exercise 6.7.1.12 implies that there exists $x_0 \in X$ such that $\bigcap_{n \in \mathbb{N}} \text{cl } V_n = \{x_0\}$. Then $x_n \rightarrow x_0$. (show details!!!, make ex about if $(x_\alpha)_{\alpha \in A}$ is eventually in U , then $x \in U$, then note that for each $N \in \mathbb{N}$, $(x_n)_{n \in \mathbb{N}}$ is eventually in V_N , so for each $N \in \mathbb{N}$, $x \in V_N$, thus $x \in \bigcap_{n \in \mathbb{N}} V_n$, dont even need cantor nestes set theorem.) Since $(X, \mathcal{T}_d)$ is Hausdorff, ref ex implies that $x = x_0$. Hence $x \in \bigcap_{n \in \mathbb{N}} V_n$. Since $\mathcal{U} \in \text{Cover}^{\text{opn}}(X)$, there exists $U \in \mathcal{U}$ such that $x \in U$. Since $U \in \mathcal{T}_d$, there exists $\epsilon > 0$ such that $B(x, \epsilon) \subset U$. Choose $N \in \mathbb{N}$ such that $2/N < \epsilon$. In particular $x \in V_N$ and for each $y \in V_N$,

$$\begin{aligned} d(y, x) &\leq d(y, x_N) + d(x_N, x) \\ &< \frac{1}{N} + \frac{1}{N} \\ &= \frac{2}{N} \\ &< \epsilon. \end{aligned}$$

Hence

$$\begin{aligned} V_N &\subset B(x, \epsilon) \\ &\subset U. \end{aligned}$$

and therefore $V_N \subset U \cap V_N$. Define $\mathcal{U}_0 \subset \mathcal{U}_N$ by $\mathcal{U}_0 := \{U \cap V_N\}$. Then $\mathcal{U}_0$ is finite and $\mathcal{U}_0 \in \text{Cover}^{\text{opn}}(V_N)$, which is a contradiction. Hence $(X, \mathcal{T}_d)$ is compact. □

Exercise 6.8.0.8. Let (X, d) be a metric space, $U \in \mathcal{T}_d$ and $x \in U$.

1. If $x \in U'$, then $U \setminus \{x\}$ is not compact.
2. If x is a condensation point of U , then $U \setminus \{x\}$ is not compact.

Proof.

1. Suppose that $x \in U'$. Set $U_0 := U \setminus \{x\}$. For the sake of contradiction, suppose that U_0 is compact. Exercise ?? implies that U_0 is sequentially compact. Since $x \in U'$, Exercise 4.3.2.7 implies that there exists $(x_n)_{n \in \mathbb{N}} \subset U_0$ such that $x_n \rightarrow x$. (Exercise 4.3.2.7 is for nets, need exercise to show existence of a subnet that is a sequence from a net) Since U_0 is sequentially compact, there exists $(x_{n_k})_{k \in \mathbb{N}} \subset (x_n)_n$ and $x_0 \in U_0$ such that $x_{n_k} \rightarrow x_0$. Since $x_n \rightarrow x$, we have that $x_{n_k} \rightarrow x$. Since X is Hausdorff,

$$\begin{aligned} x &= x_0 \\ &\in U_0, \end{aligned}$$

which is a contradiction. Thus U_0 is not compact.

2. Suppose that x is a condensation point of U . Exercise 4.1.0.47 implies that $x \in U'$. The previous part then implies that $U \setminus \{x\}$ is not compact.

□

6.9 The Baire Category Theorem

Exercise 6.9.0.1. Let X be a complete metric space and $(U_n)_{n \in \mathbb{N}} \subset \mathcal{P}(X)$. Suppose that for each $n \in \mathbb{N}$, U_n is open and dense in X . Then $\bigcap_{n \in \mathbb{N}} U_n$ is dense in X .

Hint: Let $W \subset X$. Suppose that W is open. Since U_1 is open and dense in X , Exercise 4.1.0.41 implies that $U_1 \cap W$ is open and nonempty. Hence there exists $x_1 \in U_1 \cap W$ and $r_1 \in (0, 2^{-1})$ such that $\bar{B}(x_1, r_1) \subset U_1 \cap W$. Inductively define $(x_n)_{n \in \mathbb{N}} \subset X$ and $(r_n)_{n \in \mathbb{N}} \subset (0, 1)$.

Proof. Set $U := \bigcap_{n \in \mathbb{N}} U_n$. Let $W \subset X$. Suppose that W is open and nonempty. Since U_1 is open and dense in X , Exercise 4.1.0.41 implies that $U_1 \cap W$ is open and nonempty. Hence there exists $x_1 \in U_1 \cap W$ and $r_1 \in (0, 2^{-1})$ such that $\bar{B}(x_1, r_1) \subset U_1 \cap W$. For $n \geq 2$, Exercise 4.1.0.41 implies that $U_n \cap B(x_{n-1}, r_{n-1})$ is open and nonempty. Hence there exists $x_n \in U_n \cap B(x_{n-1}, r_{n-1})$ and $r_n \in (0, 2^{-n})$ such that $\bar{B}(x_n, r_n) \subset U_n \cap B(x_{n-1}, r_{n-1})$. Note that for each $N, n \in \mathbb{N}$, if $n \geq N$, then by definition,

$$\begin{aligned} x_n &\in B(x_n, r_n) \\ &\subset U_n \cap B(x_{n-1}, r_{n-1}) \\ &\subset \left(\bigcap_{j=N+1}^n U_j \right) \cap B(x_N, r_N) \end{aligned}$$

Let $\epsilon > 0$. Choose $N \in \mathbb{N}$ such that $2^{1-N} < \epsilon$. Let $n, m \in \mathbb{N}$. Suppose that $n, m \geq N$. Then

$$\begin{aligned} d(x_n, x_m) &\leq d(x_n, x_N) + d(x_N, x_m) \\ &\leq 2^{-N} + 2^{-N} \\ &= 2^{1-N} \\ &< \epsilon \end{aligned}$$

Thus $(x_n)_{n \in \mathbb{N}}$ is Cauchy. Since X is complete, there exists $x \in X$ such that $x_n \rightarrow x$. Let $n \in \mathbb{N}$. Since $(x_n)_{n \geq N} \subset \bar{B}(x_1, r_1)$, we have that

$$\begin{aligned} x &\in \bar{B}(x_1, r_1) \\ &\subset W \end{aligned}$$

Similarly, for each $n \in \mathbb{N}$,

$$\begin{aligned} x_n &\in U_n \cap \bar{B}(x_n, r_n) \\ &\subset U_n \end{aligned}$$

which implies that $x \in U$. Hence $\bigcap_{n \in \mathbb{N}} U_n \cap W \neq \emptyset$. Since W is an arbitrary open nonempty subset of X , we have that for each $W \subset X$, if W is open and nonempty, then $U \cap W \neq \emptyset$. By definition, U is dense in X . $\square$

Exercise 6.9.0.2. Let X be a complete metric space and $(A_n)_{n \in \mathbb{N}} \subset \mathcal{P}(X)$. If for each $n \in \mathbb{N}$, A_n is nowhere dense, then $X \neq \bigcup_{n \in \mathbb{N}} A_n$.

Proof. Suppose that for each $n \in \mathbb{N}$, A_n is nowhere dense. Exercise 4.1.0.43 and Exercise 4.1.0.44 imply that for each $n \in \mathbb{N}$, $(\text{cl } A_n)^c$ is dense and open. For the sake of contradiction, suppose that $X = \bigcup_{n \in \mathbb{N}} A_n$. Then $X = \bigcup_{n \in \mathbb{N}} \text{cl } A_n$. Exercise 6.9.0.1 implies that $\emptyset = \bigcap_{n \in \mathbb{N}} (\text{cl } A_n)^c$ is dense in X . This is a contradiction. Hence $X \neq \bigcup_{n \in \mathbb{N}} A_n$. $\square$

Definition 6.9.0.3. Let X be a topological space. Set $\mathcal{D}_O(X) = \{U \subset X : U \text{ is open and dense in } X\}$. Then X is said to be a **Baire space** if for each $(U_n)_{n \in \mathbb{N}} \subset \mathcal{D}_O(X)$, $\bigcap_{n \in \mathbb{N}} U_n$ is dense in X .

Definition 6.9.0.4. Let X be a topological space. Set $\mathcal{D}_N(X) = \{U \subset X : U \text{ is nowhere dense in } X\}$. Let $E \subset X$. Then E is said to be **meager** in X if there exist $(A_n)_{n \in \mathbb{N}} \subset \mathcal{D}_N(X)$ such that $E = \bigcup_{n \in \mathbb{N}} A_n$.

Theorem 6.9.0.5. Baire Category Theorem:

Let X be a complete metric space. Then

1. X is a Baire space
2. X is not meager

Proof. Immediate by Exercise 6.9.0.1 and Exercise 6.9.0.2.

□

Definition 6.9.0.6. content...

6.10 Metric Spaces of Functions

6.10.1 Bounded Maps

Definition 6.10.1.1. Let X be a set, (Y, d) be a metric space.

- Let $f \in Y^X$. Then f is said to be **d -bounded** if $f(X)$ is d -bounded.
- We define $\text{Bnd}(X, (Y, d)) := \{f \in Y^X : f \text{ is } d\text{-bounded}\}$.

Note 6.10.1.2. When the context is clear,

- we may say “bounded” instead of d -bounded
- we may write $\text{Bnd}(X, Y)$ in place of $\text{Bnd}(X, (Y, d))$.

Exercise 6.10.1.3. Let X be a set and (Y, d) a metric space. Then for each $f, g \in \text{Bnd}(X, (Y, d))$, $d \circ (f, g) \in \text{Bnd}(X, ([0, \infty), d_{|\cdot|_{[0, \infty)}}))$. (need to make an exercise earlier showing that $|\cdot|_{\mathbb{R}}$ induces a metric., also need an exercise saying $A \subset \mathbb{R}$ is $d_{|\cdot|}$ -bounded iff A is bounded above and A is bounded below).

Proof. Define $F : X \rightarrow [0, \infty)$ by $F(x) := d \circ (f, g)$. Since $f, g \in \text{Bnd}(X, (Y, d))$, $f(X)$ is bounded and $g(X)$ is bounded. Thus there exist $M_f, M_g \in [0, \infty)$ such that for each $y_1^f, y_2^f \in f(X)$ and $y_1^g, y_2^g \in g(X)$, $d(y_1^f, y_2^f) \leq M_f$. Define $M \in [0, \infty)$ by $M := M_f + M_g$. Let $a, b \in F(X)$. Then there exist $x_1, x_2 \in X$ such that $a = F(x_1)$ and $b = F(x_2)$. Define $y_1^f, y_2^f \in f(X)$ and $y_1^g, y_2^g \in g(X)$ by $y_1^f := f(x_1), y_2^f := f(x_2), y_1^g := g(x_1), y_2^g := g(x_2)$. Then

$$\begin{aligned} d_{[0, \infty)}(a, b) &= |a - b| \\ &= |d(f(x_1), g(x_1)) - d(f(x_2), g(x_2))| \\ &\leq |d(f(x_1), g(x_1)) - d(g(x_1), f(x_2))| + |d(g(x_1), f(x_2)) - d(f(x_2), g(x_2))| \\ &\leq d(f(x_1), f(x_2)) + d(g(x_1), g(x_2)) \quad (\text{by reverse triangle inequality ref ex}) \\ &= d(y_1^f, y_2^f) + d(y_1^g, y_2^g) \\ &\leq M_f + M_g \\ &= M. \end{aligned}$$

Since $a, b \in F(X)$ are arbitrary, we have that for each $a, b \in F(X)$, $d_{[0, \infty)}(a, b) \leq M$. Thus there exists $M \in [0, \infty)$ such that for each $a, b \in F(X)$, $d_{[0, \infty)}(a, b) \leq M$. Hence $F(X)$ is bounded. Therefore $F \in \text{Bnd}(X, ([0, \infty), d_{|\cdot|_{[0, \infty)}}))$. $\square$

Definition 6.10.1.4. Let X be a set and (Y, d) a metric space. We define the **sup metric** or **uniform metric** on $\text{Bnd}(X, (Y, d))$, denoted $d_\infty : \text{Bnd}(X, (Y, d)) \times \text{Bnd}(X, (Y, d)) \rightarrow [0, \infty)$, by

$$d_\infty(f, g) := \text{diam}(d \circ (f, g)(X) \cup \{0\}, d_{|\cdot|_{[0, \infty)}}).$$

Exercise 6.10.1.5. Let X be a set and (Y, d) a metric space. If $X \neq \emptyset$, then for each $f, g \in \text{Bnd}(X, (Y, d))$,

$$d_\infty(f, g) := \sup_{x \in X} d(f(x), g(x)).$$

Proof. need to be careful of assumptions on nonempty sets and ref exercises on suprema/infima Define $F : X \rightarrow [0, \infty)$ by $F(x) := d \circ (f, g)$.

- Since F **FINISH!!!!**. Let $a, b \in d \circ (f, g)(X) \cup \{0\}$. Suppose that $b \geq a$. Then

$$\begin{aligned} d_{[0, \infty)}(a, b) &= |a - b| \\ &= b - a \\ &\leq b \\ &= d(f(x), g(x)) \\ &\leq \sup_{x \in X} d(f(x), g(x)). \end{aligned}$$

Since $a, b \in F(X)$ are arbitrary, we have that for each $a, b \in F(X)$, $d_{[0,\infty)}(a, b) \leq \sup_{x \in X} d(f(x), g(x))$. Hence

$$\begin{aligned} d_\infty(f, g) &= \text{diam}(F(X) \cup \{0\}, d_{|\cdot|_{[0,\infty)}}) \\ &\leq \sup_{x \in X} d(f(x), g(x)). \end{aligned}$$

- For each $x \in X$,

$$\begin{aligned} d(f(x), g(x)) &= F(x) \\ &= |F(x) - 0| \\ &\leq \text{diam}(F(X) \cup \{0\}, d_{|\cdot|_{[0,\infty)}}) \\ &= d_\infty(f, g). \end{aligned}$$

ref ex about suprema/infima in intro to math notes implies that $\sup_{x \in X} d(f(x), g(x)) \leq d_\infty(f, g)$.

Since $d_\infty(f, g) \leq \sup_{x \in X} d(f(x), g(x))$ and $\sup_{x \in X} d(f(x), g(x)) \leq d_\infty(f, g)$, we have that

$$d_\infty(f, g) = \sup_{x \in X} d(f(x), g(x)).$$

□

Exercise 6.10.1.6. Let X be a set and (Y, d) a metric space. Then d_∞ is a metric on $\text{Bnd}(X, (Y, d))$.

Proof.

1. Since d is symmetric, we have that for each $f, g \in \text{Bnd}(X, (Y, d))$,

$$\begin{aligned} d_\infty(f, g) &= \text{diam}(d \circ (f, g) \cup \{0\}) \\ &= \text{diam}(d \circ (g, f) \cup \{0\}) \\ &= d_\infty(g, f). \end{aligned}$$

2. For each

- 3.

FINISH!!!

□

6.10.2 Bounded Continuous Maps

Exercise 6.10.2.1. Let $(X, d_X), (Y, d_Y)$ be metric spaces, $(f_n)_{n \in \mathbb{N}} \subset \text{Bnd}(X, (Y, d))$ and $f \in \text{Bnd}(X, Y)$. If for each $n \in \mathbb{N}$, f_n is uniformly continuous and $f_n \xrightarrow{d_\infty} f$, then f is uniformly continuous.

Proof. Suppose that for each $n \in \mathbb{N}$, f_n is uniformly continuous and $f_n \xrightarrow{d_\infty} f$. Let $\epsilon > 0$. Since $f_n \xrightarrow{d_\infty} f$, there exists $N \in \mathbb{N}$ such that for each $n \in \mathbb{N}$, $n \geq N$ implies that $d_\infty(f_n, f) < \epsilon/3$. Since f_N is uniformly continuous, there exists $\delta > 0$ such that for each $x_1, x_2 \in X$, $d_X(x_1, x_2) < \delta$ implies that $d_Y(f_N(x_1), f_N(x_2)) < \epsilon/3$. Let $x_1, x_2 \in X$. Suppose that $d_X(x_1, x_2) < \delta$. Then

$$\begin{aligned} d_Y(f(x_1), f(x_2)) &\leq d_Y(f(x_1), f_N(x_1)) + d_Y(f_N(x_1), f_N(x_2)) + d_Y(f_N(x_2), f(x_2)) \\ &\leq d_\infty(f, f_N) + d_Y(f_N(x_1), f_N(x_2)) + d_\infty(f, f_N) \\ &< \frac{\epsilon}{3} + \frac{\epsilon}{3} + \frac{\epsilon}{3} \\ &= \epsilon. \end{aligned}$$

Since $\epsilon > 0$ is arbitrary, we have that for each $\epsilon > 0$, there exists $\delta > 0$ such that for each $x_1, x_2 \in X$, $d_X(x_1, x_2) < \delta$ implies that $d_Y(f(x_1), f(x_2)) < \epsilon$. Hence f is uniformly continuous. □

Definition 6.10.2.2. Let (X, MT) be a topological space and (Y, d) a metric space. We define

$$C_b((X, MT), (Y, d)) := C((X, \mathcal{T}), (Y, \mathcal{T}_d)) \cap \text{Bnd}(X, (Y, d)).$$

Note 6.10.2.3. When the context is clear,

- we write $C_b(X, Y)$ in place of $C_b((X, MT), (Y, d))$
- we write $C_b(X)$ in place of $C_b((X, \mathcal{T}), (\mathbb{C}, d_{\mathbb{C}}))$

Exercise 6.10.2.4. Let $(X, \mathcal{T})$ be a topological space and (Y, d) a metric space. Then

1. $C_b(X, Y)$ is closed in $\text{Bnd}(X, Y)$,
2. $\text{Bnd}(X, Y)$ is complete,
3. If (Y, d) is complete, then $C_b(X, Y)$ is complete.

Proof.

- 1.
- 2.
- 3.

FINISH!!!

□

6.10.3 Subspaces

Exercise 6.10.3.1. Let X be a topological space, (Y, d) a metric space and $E \subset X$. Suppose that $E \neq \emptyset$. If E is dense in X , then for each $f, g \in C_b(X)$, $d_{\infty}(f|_E, g|_E) = d_{\infty}(f, g)$.

Proof. Let $f, g \in C_b(X)$ and $\epsilon > 0$. Since $d_{\infty}(f, g) - \epsilon/3 < d_{\infty}(f, g)$, there exists $x \in X$ such that $d_{\infty}(f, g) - \epsilon/3d(f(x), g(x))$. Since $f \in C(X)$, there exists $U_f \in \mathcal{T}_X$ such that $x \in U_f$ and $f(U_f) \subset B(f(x), \epsilon/3)$. Similarly, $g \in C(X)$, there exists $U_g \in \mathcal{T}_X$ such that $x \in U_g$ and $g(U_g) \subset B(g(x), \epsilon/3)$. Define $U \in \mathcal{T}_X$ by $U := U_f \cap U_g$. Then $x \in U$, $f(U) \subset B(f(x), \epsilon/3)$ and $g(U) \subset B(g(x), \epsilon/3)$. Since E is dense in X , Exercise 4.1.0.41 implies that $E \cap U \neq \emptyset$. Thus there exists $x_0 \in X$ such that $x \in E \cap U$. Since $E \subset X$,

$$\begin{aligned} d_{\infty}(f|_E, g|_E) &= \sup_{x' \in E} d(f(x'), g(x')) \\ &\leq \sup_{x' \in X} d(f(x'), g(x')) \\ &= d_{\infty}(f, g). \end{aligned}$$

Therefore

$$\begin{aligned} d_{\infty}(f, g) &\leq d(f(x), g(x)) + \frac{\epsilon}{3} \\ &\leq d(f(x), f(x_0)) + d(f(x_0), g(x_0)) + d(g(x_0), g(x)) + \frac{\epsilon}{3} \\ &< \frac{\epsilon}{3} + d(f(x_0), g(x_0)) + \frac{\epsilon}{3} + \frac{\epsilon}{3} \\ &= d(f(x_0), g(x_0)) + \epsilon \\ &\leq \sup_{x' \in E} d(f(x'), g(x')) \\ &= d_{\infty}(f|_E, g|_E) \\ &\leq d_{\infty}(f, g). \end{aligned}$$

Hence $d_{\infty}(f|_E, g|_E) = d_{\infty}(f, g)$

□

6.10.4 Completeness

Exercise 6.10.4.1. Let X be a set and (Y, d) a metric space. If (Y, d) is complete, then $(\text{Bnd}(X, (Y, d)), d_\infty)$ is complete.

Proof. Suppose that (Y, d) is complete. Let $(f_n)_{n \in \mathbb{N}} \subset \text{Bnd}(X, (Y, d))$. Suppose that $(f_n)_{n \in \mathbb{N}}$ is Cauchy.

- Let $x \in X$. Since for each $m, n \in \mathbb{N}$,

$$\begin{aligned} d(f_m(x), f_n(x)) &\leq \sup_{x' \in X} d(f_m(x'), f_n(x')) \\ &= d_\infty(f_m, f_n). \end{aligned}$$

ref ex then implies that $(f_n(x))_{n \in \mathbb{N}}$ is Cauchy. Since (Y, d) is complete, there exists a unique $y \in Y$ such that $f_n(x) \xrightarrow{d} y$. Since $x \in X$ is arbitrary, we have that for each $x \in X$, $\#\{y \in Y : \text{there exists a unique } y \in Y \text{ such that } f_n(x) \xrightarrow{d} y\} = 1$. ref aoc in set theory section implies that there exists a unique $f : X \rightarrow Y$ such that for each $x \in X$, $f_n(x) \xrightarrow{d} f(x)$. ref or make ex about $f_n \rightarrow f$ pointwise and f_n bounded in $\text{Bnd}(X, Y)$ implies that f is bounded implies that $f \in \text{Bnd}(X, (Y, d))$.

- Let $\epsilon > 0$. Since $(f_n)_{n \in \mathbb{N}}$ is Cauchy, there exists $N \in \mathbb{N}$ such that for each $n \in \mathbb{N}$, $n \geq N$ implies that $d_\infty(f_m, f_n) < \epsilon/2$. Let $n \in \mathbb{N}$. Suppose that $n \geq N$. Let $x \in X$. ref ex about $d(x, \cdot)$ being continuous implies that

$$\begin{aligned} d(f_n(x), f(x)) &= d(f_n(x), \lim_{m \rightarrow \infty} f_m(x)) \\ &= \lim_{m \rightarrow \infty} d(f_n(x), f_m(x)) \\ &\leq \sup_{x' \in X} d(f_n(x'), f_m(x')) \\ &= d_\infty(f_n, f_m) \\ &< \frac{\epsilon}{2}. \end{aligned}$$

Since $x \in X$ is arbitrary, we have that for each $x \in X$, $d(f_n(x), f(x)) < \epsilon/2$. Therefore

$$\begin{aligned} d_\infty(f_n, f) &= \sup_{x' \in X} d(f_n(x'), f(x')) \\ &\leq \frac{\epsilon}{2} \\ &< \epsilon. \end{aligned}$$

Since $\epsilon > 0$ is arbitrary, we have that for each $\epsilon > 0$, there exists $N \in \mathbb{N}$ such that for each $n \in \mathbb{N}$, $n \geq N$ implies that $d_\infty(f_n, f) < \epsilon$. Hence $f_n \xrightarrow{d_\infty} f$.

Since $(f_n)_{n \in \mathbb{N}} \subset \text{Bnd}(X, (Y, d))$ with $(f_n)_{n \in \mathbb{N}}$ Cauchy is arbitrary, we have that for each $(f_n)_{n \in \mathbb{N}} \subset \text{Bnd}(X, (Y, d))$, $(f_n)_{n \in \mathbb{N}}$ is Cauchy implies that there exists $f \in \text{Bnd}(X, (Y, d))$ such that $f_n \xrightarrow{d_\infty} f$. Hence $(\text{Bnd}(X, (Y, d)), d_\infty)$ is complete. $\square$

Exercise 6.10.4.2. Moore-Osgood Theorem: Let (X, d) be a metric space, A, B directed sets and $(y_\alpha)_{\alpha \in A}, (z_\beta)_{\beta \in B}, (x_{\alpha, \beta})_{(\alpha, \beta) \in A \times B} \subset X$ nets. Suppose that

- $(x_{\alpha, \beta})_{\alpha \in A} \xrightarrow{\text{unif}} (y_\alpha)_{\alpha \in A}$,
- for each $\alpha \in A$, $x_{\alpha, \beta} \rightarrow z_\beta$.

If (X, d) is complete, then

1. $(z_\beta)_{\beta \in B}$ is Cauchy and there exists $x \in X$ such that $z_\beta \rightarrow x$.

Hint: continuity of $d : X \times X \rightarrow \mathbb{R}_+$

2. $y_\alpha \rightarrow x$
3. $x_{\alpha, \beta} \rightarrow x$.

Proof.

1. Since $(x_{\alpha,\beta})_{\alpha \in A} \xrightarrow{\text{unif}} (y_{\alpha})_{\alpha \in A}$, we have that $(x_{\alpha,\beta})_{\alpha \in A} \xrightarrow{\text{p.w.}} (y_{\alpha})_{\alpha \in A}$ and for each $\alpha \in A$, $x_{\alpha,\beta} \rightarrow y_{\alpha}$. Let $\epsilon > 0$. Since $(x_{\alpha,\beta})_{\alpha \in A} \xrightarrow{\text{unif}} (y_{\alpha})_{\alpha \in A}$, $d_{\infty}((x_{\alpha,\beta})_{\alpha \in A}, (y_{\alpha})_{\alpha \in A}) \rightarrow 0$. Thus there exists $\beta_0 \in B$ such that for each $\beta \in B$, $\beta \geq \beta_0$ implies that $d_{\infty}((x_{\alpha,\beta})_{\alpha \in A}, (y_{\alpha})_{\alpha \in A}) < \epsilon$. Let $\beta_1, \beta_2 \in B$. Suppose that $\beta_1, \beta_2 \geq \beta_0$. Then for each $\alpha \in A$,

$$\begin{aligned} d(x_{\alpha,\beta_1}, x_{\alpha,\beta_2}) &\leq d(x_{\alpha,\beta_1}, y_{\alpha}) + d(y_{\alpha}, x_{\alpha,\beta_2}) \\ &\leq d_{\infty}((x_{\alpha',\beta_1})_{\alpha' \in A}, (y_{\alpha'})_{\alpha' \in A}) + d_{\infty}((x_{\alpha',\beta_1})_{\alpha' \in A}, (y_{\alpha'})_{\alpha' \in A}) \\ &< \frac{\epsilon}{4} + \frac{\epsilon}{4} \\ &= \frac{\epsilon}{2}. \end{aligned}$$

Since for each $\beta \in B$, $x_{\alpha,\beta} \rightarrow z_{\beta}$, continuity of $d : X \times X \rightarrow \mathbb{R}_+$ (Exercise 17.3.0.5) **ref ex here** implies that

$$\begin{aligned} d(z_{\beta_1}, z_{\beta_2}) &= \lim_{\alpha \in A} d(x_{\alpha,\beta_1}, x_{\alpha,\beta_2}) \\ &\leq \frac{\epsilon}{2} \\ &< \epsilon. \end{aligned}$$

Since $\beta_1, \beta_2 \in B$ with $\beta_1, \beta_2 \geq \beta_0$ are arbitrary, we have that for each $\beta_1, \beta_2 \in B$, $\beta_1, \beta_2 \geq \beta_0$ implies that $d(z_{\beta_1}, z_{\beta_2}) < \epsilon$. Since $\epsilon > 0$ is arbitrary, we have that for each $\epsilon > 0$, there exists $\beta_0 \in B$ such that for each $\beta_1, \beta_2 \in B$, $\beta_1, \beta_2 \geq \beta_0$ implies that $d(z_{\beta_1}, z_{\beta_2}) < \epsilon$. Hence $(z_{\beta})_{\beta}$ is Cauchy. Since (X, d) is complete, there exists $x \in X$ such that $z_{\beta} \rightarrow x$.

2. Let $\epsilon > 0$. Since $z_{\beta} \rightarrow x$, there exists $\beta_1 \in B$ such that for each $\beta \in B$, $\beta \geq \beta_1$ implies that $d(z_{\beta}, x) < \epsilon/3$. Since $(x_{\alpha,\beta})_{\alpha \in A} \xrightarrow{\text{unif}} (y_{\alpha})_{\alpha \in A}$, there exists $\beta_2 \in B$ such that for each $\beta \in B$, $\beta \geq \beta_2$ implies that for each $\alpha \in A$, $d(x_{\alpha,\beta}, y_{\alpha}) < \epsilon/3$. Since B is directed, there exists $\beta_0 \in B$ such that $\beta_1, \beta_2 \leq \beta_0$. Since $x_{\alpha,\beta_0} \rightarrow z_{\beta_0}$, there exists $\alpha_0 \in A$ such that for each $\alpha \in A$, $\alpha \geq \alpha_0$ implies that $d(x_{\alpha,\beta_0}, z_{\beta_0}) < \epsilon/3$. Then for each $\alpha \in A$, $\alpha \geq \alpha_0$ implies that

$$\begin{aligned} d(y_{\alpha}, x) &\leq d(y_{\alpha}, x_{\alpha,\beta_0}) + d(x_{\alpha,\beta_0}, z_{\beta_0}) + d(z_{\beta_0}, x) \\ &< \frac{\epsilon}{3} + \frac{\epsilon}{3} + \frac{\epsilon}{3} \\ &= \epsilon. \end{aligned}$$

Since $\epsilon > 0$ is arbitrary, we have that for each $\epsilon > 0$, there exists $\alpha_0 \in A$ such that for each $\alpha \in A$, $\alpha \geq \alpha_0$ implies that $d(y_{\alpha}, x) < \epsilon$. Hence $y_{\alpha} \rightarrow x$.

3. Let $\epsilon > 0$. Since $y_{\alpha} \rightarrow x$, there exists $\alpha_0 \in A$ such that for each $\alpha \in A$, $\alpha \geq \alpha_0$ implies that $d(y_{\alpha}, x) < \epsilon/2$. Since $(x_{\alpha,\beta})_{\alpha \in A} \xrightarrow{\text{unif}} (y_{\alpha})_{\alpha \in A}$, there exists $\beta_0 \in B$, such that for each $\beta \in B$, $\beta \geq \beta_0$ implies that $d_{\infty}((x_{\alpha,\beta})_{\alpha \in A}, (y_{\alpha})_{\alpha \in A}) < \epsilon/2$. Let $(\alpha, \beta) \in A \times B$. Suppose that $(\alpha, \beta) \geq (\alpha_0, \beta_0)$. Then $\alpha \geq \alpha_0$ and $\beta \geq \beta_0$. Therefore

$$\begin{aligned} d(x_{\alpha,\beta}, x) &= d(x_{\alpha,\beta}, y_{\alpha}) + d(y_{\alpha}, x) \\ &\leq d_{\infty}((x_{\alpha,\beta})_{\alpha \in A}, (y_{\alpha})_{\alpha \in A}) + d(y_{\alpha}, x) \\ &< \frac{\epsilon}{2} + \frac{\epsilon}{2} \\ &= \epsilon. \end{aligned}$$

Since $(\alpha, \beta) \in A \times B$ with $(\alpha, \beta) \geq (\alpha_0, \beta_0)$ is arbitrary, we have that for each $(\alpha, \beta) \in A \times B$, $(\alpha, \beta) \geq (\alpha_0, \beta_0)$ implies that $d(x_{\alpha,\beta}, x) < \epsilon$. Since $\epsilon > 0$ is arbitrary, we have that for each $\epsilon > 0$, there exists $(\alpha_0, \beta_0) \in A \times B$ such that for each $(\alpha, \beta) \in A \times B$, $(\alpha, \beta) \geq (\alpha_0, \beta_0)$ implies that $d(x_{\alpha,\beta}, x) < \epsilon$. Hence $x_{\alpha,\beta} \rightarrow x$.

□

6.10.5 Continuous Maps on LCH spaces

Definition 6.10.5.1. Let X be a topological space.

- Let $f \in C(X)$.
 - We define the **support of f** , denoted $\text{supp}(f) \subset X$, by $\text{supp}(f) := \text{cl}\{x \in X : |f(x)| \neq 0\}$.
 - Then f is said to
 - * **have compact support** if $\text{supp}(f) \in \mathcal{K}(X)$,
 - * **vanish at infinity** if for each $\epsilon > 0$, $\{x \in X : |f(x)| \geq \epsilon\} \in \mathcal{K}(X)$.
- We define
 - $C_c(X) := \{f \in C(X) : f \text{ has compact support}\}$,
 - $C_0(X) := \{f \in C(X) : f \text{ vanishes at infinity}\}$.

Exercise 6.10.5.2. Let X be a LCH space. Then $\text{cl } C_c(X) = C_0(X)$.

Proof. **FINISH!!!**

□

6.10.6 Continuous Maps Compactifications

Exercise 6.10.6.1. Let X be a topological space, $(\eta(X), \iota)$ a Stone-Ćech compactification of X and $\mathcal{F} \subset C_b(X)$. If $\mathcal{F}$ is dense in $C_b(X)$, then $\eta(\mathcal{F})$ is dense in $C(\eta(X))$. **rephrase $\eta(f)$ to be defined for $f : X \rightarrow Y$ with $\text{Img } f \subset K \in \mathcal{K}(X)$ instead of $f : X \rightarrow K$.**

Proof. **Clean up, especially handling $\eta(\mathcal{F})$, since $\eta(f)$ is defined when $\text{cod}(f)$ is compact.** Suppose that $\mathcal{F}$ is dense in $C_b(X)$. Let $g \in C(\eta(X))$ and $\epsilon > 0$. Define $g' \in C_b(X)$ by $g' := g \circ \iota$. Since $g' \in C_b(X)$, there exists $R \geq 0$ such that $g'(X) \subset \bar{B}(0, R/2)$. Define $K \subset \mathbb{C}$ by $K := \bar{B}(0, R)$. Since $\mathcal{F}$ is dense in $C_b(X)$, there exists $f' \in \mathcal{F}$ such that $d_\infty(f', g') < \min\{\epsilon, R/2\}$. Then for each $x \in X$,

$$\begin{aligned} |f'(x)| &\leq |f'(x) - g'(x)| + |g'(x)| \\ &\leq d_\infty(f', g') + \frac{R}{2} \\ &< \frac{R}{2} + \frac{R}{2} \\ &= R. \end{aligned}$$

Hence $f'(X) \subset K$. Define $f_0 \in C(\eta(X), K)$ and $f \in \eta(\mathcal{F})$ by $f := \eta(f')$. Since $\iota(X)$ is dense in $\eta(X)$, **a prev ex** implies that

$$\begin{aligned} d_\infty(f, g) &= d_\infty(f|_{\iota(X)}, g|_{\iota(X)}) \\ &= d_\infty(f|_{\iota(X)} \circ \iota, g|_{\iota(X)} \circ \iota) \\ &= d_\infty(f \circ \iota, g \circ \iota) \\ &= d_\infty(f', g') \\ &< \epsilon. \end{aligned}$$

Since $g \in C(\eta(X))$ and $\epsilon > 0$ are arbitrary, we have that for each $g \in C(\eta(X))$ and $\epsilon > 0$, there exists $f \in \eta(\mathcal{F})$ such that $d_\infty(f, g) < \epsilon$. □

6.11 Metrizable Spaces

6.11.1 Introduction

Definition 6.11.1.1. Let $(X, \mathcal{T})$ be a topological space. Then $(X, \mathcal{T})$ is said to be **metrizable** if there exists a metric $d : X \times X \rightarrow [0, \infty)$ such that $\mathcal{T} = \mathcal{T}_d$.

Definition 6.11.1.2. We define the **Hilbert cube**, denoted $\mathbb{H}$, by $\mathbb{H} := [0, 1]^{\mathbb{N}}$.

Exercise 6.11.1.3. Let $(X, \mathcal{T})$ be a topological space. Suppose that $(X, \mathcal{T})$ is separable and metrizable. Then there exists $B \subset \mathbb{H}$ and $f : X \rightarrow B$ such that f is a homeomorphism.

Hint: Choose $(a_n)_{n \in \mathbb{N}} \subset X$ and $d : X \times X \rightarrow [0, 1]$ such that $(a_n)_{n \in \mathbb{N}}$ is dense in $(X, \mathcal{T})$ and $\mathcal{T}_d = \mathcal{T}$. Define $f : X \rightarrow \mathbb{H}$ by $f(x) := (d(x, a_j))_{j \in \mathbb{N}}$.

Proof. Since $(X, \mathcal{T})$ is metrizable, Exercise 17.3.0.5 (reference exercise here) implies that there exists a metric $d : X \times X \rightarrow [0, 1]$ such that $\mathcal{T} = \mathcal{T}_d$. Since $(X, \mathcal{T})$ is separable, there exists $(a_j)_{j \in \mathbb{N}}$ such that $(a_j)_{j \in \mathbb{N}}$ is dense in $(X, \mathcal{T})$. Define $B \subset \mathbb{H}$ and $f : X \rightarrow B$ by $B := \{(d(x, a_j))_{j \in \mathbb{N}} : x \in X\}$ and $f(x) := (d(x, a_j))_{j \in \mathbb{N}}$.

- Let $x, y \in X$. Suppose that $f(x) = f(y)$. Then $(d(x, a_j))_{j \in \mathbb{N}} = (d(y, a_j))_{j \in \mathbb{N}}$. Since $(a_j)_{j \in \mathbb{N}}$ is dense in $(X, \mathcal{T})$, there exists a subsequence $(a_{j_k})_{k \in \mathbb{N}} \subset (a_j)_{j \in \mathbb{N}}$ such that $a_{j_k} \rightarrow x$. Thus

$$\begin{aligned} d(y, a_{j_k}) &= d(x, a_{j_k}) \\ &\rightarrow 0. \end{aligned}$$

Thus $a_{j_k} \rightarrow y$. Since $(X, \mathcal{T})$ is Hausdorff, $a_{j_k} \rightarrow x$ and $a_{j_k} \rightarrow y$, we have that $x = y$. Thus $f(x) = f(y)$ implies that $x = y$. Since $x, y \in X$ are arbitrary, we have that for each $x, y \in X$, $f(x) = f(y)$ implies that $x = y$. Hence f is injective.

- By construction f is surjective.

Hence f is a bijection.

- Let $(x_n)_{n \in \mathbb{N}} \subset X$ and $x \in X$. Suppose that $x_n \rightarrow x$. Since $d : X \times X \rightarrow [0, 1]$ is continuous, we have that for each $j \in \mathbb{N}$, $d(x_n, a_j) \rightarrow d(x, a_j)$ in $[0, 1]$. Thus

$$\begin{aligned} f(x_n) &= (d(x_n, a_j))_{j \in \mathbb{N}} \\ &\rightarrow (d(x, a_j))_{j \in \mathbb{N}} \\ &= f(x) \end{aligned}$$

in $\mathbb{H}$. Thus $x_n \rightarrow x$ implies that $f(x_n) \rightarrow f(x)$. Since $(x_n)_{n \in \mathbb{N}} \subset X$ and $x \in X$ are arbitrary, we have that for each $(x_n)_{n \in \mathbb{N}} \subset X$ and $x \in X$, $x_n \rightarrow x$ implies that $f(x_n) \rightarrow f(x)$. Hence f is continuous.

- Let $(r_n)_{n \in \mathbb{N}} \subset B$ and $r \in B$. Suppose that $r_n \rightarrow r$. Define $(x_n)_{n \in \mathbb{N}} \subset X$ and $x \in X$ by $x_n := f^{-1}(r_n)$ and $x := f^{-1}(r)$. Since

$$\begin{aligned} (d(x_n, a_j))_{j \in \mathbb{N}} &= r_n \\ &\rightarrow r \\ &= (d(x, a_j))_{j \in \mathbb{N}}, \end{aligned}$$

we have that for each $j \in \mathbb{N}$,

$$\begin{aligned} d(x_n, a_j) &= \pi_j(r_n) \\ &\rightarrow \pi_j(r) \\ &= d(x, a_j). \end{aligned}$$

Since $(a_j)_{j \in \mathbb{N}}$ is dense in $(X, \mathcal{T})$, there exists a subsequence $(a_{j_k})_{k \in \mathbb{N}} \subset (a_j)_{j \in \mathbb{N}}$ such that $a_{j_k} \rightarrow x$. Let $\epsilon > 0$. Then $\epsilon/3 > 0$. Since $a_{j_k} \rightarrow x$, there exists $K \in \mathbb{N}$ such that for each $k \in \mathbb{N}$, $k \geq K$ implies that $d(a_{j_k}, x) < \epsilon/3$. Since $d(x_n, a_{j_k}) \rightarrow d(x, a_{j_k})$,

there exists $N \in \mathbb{N}$ such that for each $n \in \mathbb{N}$, $n \geq N$ implies that $|d(x_n, a_j) - d(x, a_j)| < \epsilon/3$. Let $n \in \mathbb{N}$. Suppose that $n \geq N$. Then

$$\begin{aligned}
 d(f^{-1}(r_n), f^{-1}(r)) &= d(x_n, x) \\
 &\leq d(x_n, a_{j_K}) + d(a_{j_K}, x) \\
 &< \left[d(x, a_{j_K}) + \frac{\epsilon}{3} \right] + d(a_{j_K}, x) \\
 &= 2d(x, a_{j_K}) + \frac{\epsilon}{3} \\
 &< \frac{2\epsilon}{3} + \frac{\epsilon}{3} \\
 &= \epsilon.
 \end{aligned}$$

Since $\epsilon > 0$ is arbitrary, we have that for each $\epsilon > 0$, there exists $N \in \mathbb{N}$ such that for each $n \in \mathbb{N}$, $n \geq N$ implies that $d(f^{-1}(r_n), f^{-1}(r)) < \epsilon$. Hence $f^{-1}(r_n) \rightarrow f^{-1}(r)$. Since $(r_n)_{n \in \mathbb{N}} \subset B$ and $r \in B$ are arbitrary, we have that for each $(r_n)_{n \in \mathbb{N}} \subset B$ and $r \in B$, $r_n \rightarrow r$ implies that $f^{-1}(r_n) \rightarrow f^{-1}(r)$. Hence f^{-1} is continuous.

Thus f is a homeomorphism. □

Definition 6.11.1.4. Let $(X, \mathcal{T})$ be a topological space. Then $(X, \mathcal{T})$ is said to be **completely metrizable** if there exists a metric $d : X \times X \rightarrow [0, \infty)$ such that $\mathcal{T} = \mathcal{T}_d$ and (X, d) is complete.

Exercise 6.11.1.5. Let $(X, \mathcal{T})$ be a topological space and $C \subset X$. Suppose that $(X, \mathcal{T})$ is completely metrizable. If C is closed, then $(C, \mathcal{T} \cap C)$ is completely metrizable. Then $(X, \mathcal{T})$ is said to be **completely metrizable** if there exists a metric $d : X \times X \rightarrow [0, \infty)$ such that $\mathcal{T} = \mathcal{T}_d$ and (X, d) is complete.

Proof. Suppose that C is closed. Since $(X, \mathcal{T})$ is completely metrizable, there exists a metric $d : X \times X \rightarrow [0, \infty)$ such that $\mathcal{T} = \mathcal{T}_d$ and (X, d) is complete. Exercise 6.7.2.2 implies that $(C, d|_{C \times C})$ is complete. Exercise 6.4.1.2 implies that $\mathcal{T} \cap C = \mathcal{T}_{d|_{C \times C}}$. Thus $(C, \mathcal{T} \cap C)$ is completely metrizable. □

Exercise 6.11.1.6. Let $(X, \mathcal{T}_X)$ and $(Y, \mathcal{T}_Y)$ be topological spaces and $f : X \rightarrow Y$ a $(\mathcal{T}_X, \mathcal{T}_Y)$ -homeomorphism. Then $(X, \mathcal{T}_X)$ is completely metrizable iff $(Y, \mathcal{T}_Y)$ is completely metrizable.

Proof.

• $(\implies)$:

Suppose that $(X, \mathcal{T}_X)$ is completely metrizable. Then there exists a metric $d_X : X \times X \rightarrow [0, \infty)$ such that $\mathcal{T}_X = \mathcal{T}_{d_X}$ and (X, d_X) is complete. Set $d_Y := (f^{-1})^* d_X$.

– Exercise 6.1.5.8 implies that $\mathcal{T}_Y = \mathcal{T}_{d_Y}$. So $(Y, \mathcal{T}_Y)$ is metrizable.

– Exercise 6.1.5.6 implies that f^{-1} is a (d_Y, d_X) -isometry. Since (X, d_X) is complete, Exercise 6.7.1.10 implies that (Y, d_Y) is complete.

Hence $(Y, \mathcal{T}_Y)$ is completely metrizable.

• $(\impliedby)$:

Similar to $(\implies)$.

□

6.11.2 Metrizable Subspaces

Exercise 6.11.2.1. Let $(X, \mathcal{T})$ be a topological space and $C \subset X$. Suppose that $(X, \mathcal{T})$ is completely metrizable. If C is $\mathcal{T}$ -closed, then $(C, \mathcal{T} \cap C)$ is completely metrizable.

Proof. Suppose that C is $\mathcal{T}$ -closed. Since $(X, \mathcal{T})$ is completely metrizable, there exists a metric $d : X \times X \rightarrow [0, \infty)$ such that $\mathcal{T} = \mathcal{T}_d$ and (X, d) is complete. Exercise 6.7.2.2 implies that $(C, d|_{C \times C})$ is complete. Exercise 6.4.1.2 implies that $\mathcal{T}_{d|_{C \times C}} = \mathcal{T}_d \cap C$. Hence $(C, \mathcal{T}_d \cap C)$ is completely metrizable. □

6.11.3 Metrizable of Product Spaces

Exercise 6.11.3.1.

Exercise 6.11.3.2. Let $(X_n, \mathcal{T}_n)_{n \in \mathbb{N}}$ be a collection of topological spaces. Suppose that for each $n \in \mathbb{N}$, $(X_n, \mathcal{T}_n)$ is completely metrizable. Then $(\prod_{n \in \mathbb{N}} X_n, \bigotimes_{n \in \mathbb{N}} \mathcal{T}_n)$ is completely metrizable.

Proof. Define $\phi : [0, \infty) \rightarrow [0, \infty)$, X and $\mathcal{T}$ as in Definition 6.5.0.1. Since for each $n \in \mathbb{N}$, $(X_n, \mathcal{T}_n)$ is completely metrizable, for each $n \in \mathbb{N}$, there exists a metric $d_n : X_n \times X_n \rightarrow [0, \infty)$ such that (X_n, d_n) is a complete metric space. Exercise 6.7.3.2 implies that (X, d_X) is complete. Since Exercise 6.5.0.2 implies that $\mathcal{T} = \mathcal{T}_{d_X}$, we have that $(X, \mathcal{T})$ is completely metrizable. $\square$

6.11.4 Metrizable of Coproduct Spaces

Exercise 6.11.4.1. Let $(X_\alpha, \mathcal{T}_\alpha)_{\alpha \in A}$ be a collection of topological spaces. Suppose that for each $\alpha \in A$, $(X_\alpha, \mathcal{T}_\alpha)$ is completely metrizable. Then $(\prod_{\alpha \in A} X_\alpha, \bigoplus_{\alpha \in A} \mathcal{T}_\alpha)$ is completely metrizable.

Proof. Define $\phi : [0, \infty) \rightarrow [0, \infty)$, X and $\mathcal{T}$ as in Definition 6.6.0.1. Since for each $\alpha \in A$, $(X_\alpha, \mathcal{T}_\alpha)$ is completely metrizable, for each $\alpha \in A$, there exists a metric $d_\alpha : X_\alpha \times X_\alpha \rightarrow [0, \infty)$ such that (X_α, d_α) is a complete metric space. Define $d_X : X \times X \rightarrow [0, \infty)$ as in Definition 6.6.0.1. Exercise 6.7.4.2 implies that (X, d_X) is complete. Since Exercise 6.6.0.2 implies that $\mathcal{T} = \mathcal{T}_{d_X}$, we have that $(X, \mathcal{T})$ is completely metrizable. $\square$

6.11.5 Metrizable of Projective Limits

Exercise 6.11.5.1. Let $(J, \leq)$ be a directed set and $((X_j, \mathcal{T}_j)_{j \in J}, (\pi_{i,j})_{(i,j) \in \leq})$ be a **Top**-projective system. Set $(X, (\pi_j)_{j \in J}) := \varprojlim_{j \in J} ((X_j, \mathcal{T}_j)_{j \in J}, (\pi_{i,j})_{(i,j) \in \leq})$. Suppose that J is countable and for each $j \in J$, $(X_j, \mathcal{T}_j)$ is completely metrizable. Then $(X, \mathcal{T})$ is completely metrizable.

Proof. Since J is countable, Exercise 6.11.3.2 implies that $(\prod_{j \in J} X_j, \bigotimes_{j \in J} \mathcal{T}_j)$ is completely metrizable. Since for each $j \in J$, $(X_j, \mathcal{T}_j)$ is metrizable, for each $j \in J$, $(X_j, \mathcal{T}_j)$ is Hasudorff. Exercise 4.10.0.5 implies that X is closed in $(\prod_{j \in J} X_j, \bigotimes_{j \in J} \mathcal{T}_j)$. Since $\mathcal{T} = \left(\bigotimes_{j \in J} \mathcal{T}_j \right) \cap X$, Exercise 6.11.2.1 implies that $(X, \mathcal{T})$ is completely metrizable. $\square$

6.12 Polish Spaces

6.12.1 Introduction

Definition 6.12.1.1. Let $(X, \mathcal{T})$ be a topological space. Then $(X, \mathcal{T})$ is said to be a **Polish** space if

1. $(X, \mathcal{T})$ is completely metrizable
2. $(X, \mathcal{T})$ is separable

Exercise 6.12.1.2. Let $(X, \mathcal{T})$ be a Polish space and $C \subset X$. If C is $\mathcal{T}$ -closed, then $(C, \mathcal{T} \cap C)$ is a Polish space.

Proof. Suppose that C is closed.

1. Since $(X, \mathcal{T})$ is completely metrizable and C is closed, Exercise 6.11.2.1 implies that $(C, \mathcal{T} \cap C)$ is completely metrizable.
2. Since $(X, \mathcal{T})$ is separable, Exercise 6.4.1.4 implies that $(C, \mathcal{T} \cap C)$ is separable.

Therefore $(C, \mathcal{T} \cap C)$ is a Polish space. □

Exercise 6.12.1.3. Let $(X_n, \mathcal{T}_n)_{n \in \mathbb{N}}$ be a collection of Polish spaces. Then $(\prod_{n \in \mathbb{N}} X_n, \bigotimes_{n \in \mathbb{N}} \mathcal{T}_n)$ is a Polish space.

Proof. Since for each $n \in \mathbb{N}$, $(X_n, \mathcal{T}_n)$ is a Polish space, we have that for each $n \in \mathbb{N}$, $(X_n, \mathcal{T}_n)$ is completely metrizable and $(X_n, \mathcal{T}_n)$ is separable. Set $X := \prod_{n \in \mathbb{N}} X_n$ and $\mathcal{T} := \bigotimes_{n \in \mathbb{N}} \mathcal{T}_n$.

1. Exercise 6.11.3.2 implies that $(X, \mathcal{T})$ is completely metrizable.
2. Since for each $n \in \mathbb{N}$, $(X_n, \mathcal{T}_n)$ is separable, Exercise 6.2.1.1 implies that for each $n \in \mathbb{N}$, $(X_n, \mathcal{T}_n)$ is second-countable. Exercise 4.13.2.13 then implies that $(X, \mathcal{T})$ is second-countable. Another application of Exercise 6.2.1.1 implies that $(X, \mathcal{T})$ is separable.

Thus $(X, \mathcal{T})$ is a Polish space. □

Exercise 6.12.1.4. Let $(X_n, \mathcal{T}_n)_{n \in \mathbb{N}}$ be a collection of Polish spaces. Then $(\prod_{n \in \mathbb{N}} X_n, \bigoplus_{n \in \mathbb{N}} \mathcal{T}_n)$ is Polish.

Proof. Set $X := \prod_{n \in \mathbb{N}} X_n$ and $\mathcal{T} := \bigoplus_{n \in \mathbb{N}} \mathcal{T}_n$. Since for each $n \in \mathbb{N}$, $(X_n, \mathcal{T}_n)$ is a Polish space, for each $n \in \mathbb{N}$, $(X_n, \mathcal{T}_n)$ is completely metrizable and second-countable. Exercise 6.11.4.1 implies that $(X, \mathcal{T})$ is completely metrizable and Exercise 4.13.2.14 implies that $(X, \mathcal{T})$ is second-countable. Hence $(X, \mathcal{T})$ is a Polish space. □

Exercise 6.12.1.5. Let $(J, \leq)$ be a directed set and $((X_j, \mathcal{T}_j)_{j \in J}, (\pi_{i,j})_{(i,j) \in \leq})$ be a **Top**-projective system. Set $(X, (\pi_j)_{j \in J}) := \varprojlim_{j \in J} ((X_j, \mathcal{T}_j)_{j \in J}, (\pi_{i,j})_{(i,j) \in \leq})$. Suppose that J is countable and for each $j \in J$, $(X_j, \mathcal{T}_j)$ is a Polish space. Then $(X, \mathcal{T})$ is a Polish space.

Proof. Since for each $j \in J$, $(X_j, \mathcal{T}_j)$ is a Polish space, for each $j \in J$, $(X_j, \mathcal{T}_j)$ is completely metrizable. Exercise 6.11.5.1 that $(X, \mathcal{T})$ is completely metrizable. Since for each $j \in J$, $(X_j, \mathcal{T}_j)$ is separable, Exercise 6.2.1.1 implies that for each $j \in J$, $(X_j, \mathcal{T}_j)$ is second-countable. Since J is countable, Exercise 4.13.2.15 implies that $(X, \mathcal{T})$ is second-countable. Another application of Exercise 6.2.1.1 implies that $(X, \mathcal{T})$ is separable. Hence $(X, \mathcal{T})$ is a Polish space. □

Exercise 6.12.1.6. Let $(X, \mathcal{T}_X)$ and $(Y, \mathcal{T}_Y)$ be topological spaces and $f : X \rightarrow Y$ a $(\mathcal{T}_X, \mathcal{T}_Y)$ -homeomorphism. Then $(X, \mathcal{T}_X)$ is a Polish space iff $(Y, \mathcal{T}_Y)$ is a Polish space.

Proof.

• $(\implies)$:

Suppose that $(X, \mathcal{T}_X)$ is a Polish space. Then $(X, \mathcal{T}_X)$ is completely metrizable and $(X, \mathcal{T}_X)$ is separable.

1. Exercise 6.11.1.6 implies that $(Y, \mathcal{T}_Y)$ is completely metrizable.
2. Since $(X, \mathcal{T}_X)$ and $(Y, \mathcal{T}_Y)$ are metrizable, there exist metrics $d_X : X \times X \rightarrow [0, \infty)$ and $d_Y : Y \times Y \rightarrow [0, \infty)$ on X and Y respectively such that $\mathcal{T}_X = \mathcal{T}_{d_X}$ and $\mathcal{T}_Y = \mathcal{T}_{d_Y}$. Since $(X, \mathcal{T}_X)$ is separable and f is a $(\mathcal{T}_X, \mathcal{T}_Y)$ -homeomorphism, we have that Exercise 6.2.1.2 implies that $(Y, \mathcal{T}_Y)$ is separable.

Thus $(Y, \mathcal{T}_Y)$ is a Polish space.

- $(\Leftarrow)$:

Similar to $(\Rightarrow)$.

□

Exercise 6.12.1.7. Let $(X, \mathcal{T})$ be a Polish space and $U \in \mathcal{T}$. Then $(U, \mathcal{T} \cap U)$ is a Polish space.

Hint: Exercise 6.1.4.7

Proof. Define $C \subset X \times \mathbb{R}$ and $f : U \rightarrow C$ as in Exercise 6.1.4.7. Exercise 6.1.4.7 implies that f is a $(\mathcal{T} \cap U, (\mathcal{T} \otimes \mathcal{T}_{\mathbb{R}}) \cap C)$ -homeomorphism and C is closed. Exercise 6.12.1.3 implies that $(X \times \mathbb{R}, \mathcal{T} \otimes \mathcal{T}_{\mathbb{R}})$ is a Polish space. Exercise 6.12.1.6 implies that $(C, (\mathcal{T} \otimes \mathcal{T}_{\mathbb{R}}) \cap C)$ is a Polish space. □

Exercise 6.12.1.8. Let $(X, \mathcal{T})$ be a Polish space and $E \subset X$. Then E is a G_δ -set iff $(E, \mathcal{T} \cap E)$ is a Polish space.

Hint: Exercise 6.1.4.8

Proof.

- $(\Rightarrow)$:

Suppose that E is a G_δ -set. Set $\mathcal{S} := \mathcal{T} \otimes (\mathcal{T}_{\mathbb{R}})^{\otimes \mathbb{N}}$. Since E is a G_δ -set, there exists $(U_n)_{n \in \mathbb{N}} \subset \mathcal{T}$ such that $E = \bigcap_{n \in \mathbb{N}} U_n$.

Exercise 6.1.4.8 implies that there exists $C \subset X \times \mathbb{R}^{\mathbb{N}}$ and $f : E \rightarrow C$ such that f is a $(\mathcal{T} \cap E, \mathcal{S} \cap C)$ -homeomorphism and C is closed in $X \times \mathbb{R}^{\mathbb{N}}$. Exercise 6.12.1.3 implies that $X \times \mathbb{R}^{\mathbb{N}}$ is a Polish space. Since C is closed in $X \times \mathbb{R}^{\mathbb{N}}$, Exercise 6.12.1.2 implies that $(C, \mathcal{S} \cap C)$ is a Polish space. Exercise 6.12.1.6 implies that $(E, \mathcal{T} \cap E)$ is a Polish space.

- $(\Leftarrow)$:

Conversely, suppose that $(E, \mathcal{T} \cap E)$ is a Polish space. **FINISH!!!**

□

Exercise 6.12.1.9. Let $(X, \mathcal{T})$ be a Polish space. Set $\Delta_{X^{\mathbb{N}}} := \{x \in X : \text{for each } m, n \in \mathbb{N}, \pi_m(x) = \pi_n(x)\}$. Then $(\Delta_{X^{\mathbb{N}}}, \mathcal{T}^{\otimes \mathbb{N}} \cap \Delta_{X^{\mathbb{N}}})$ is a Polish space. **Hint:** Exercise 4.12.1.8

Proof. Since $(X, \mathcal{T})$ is Polish, $(X, \mathcal{T})$ is Hausdorff. Exercise 6.12.1.3 implies that $(X^{\mathbb{N}}, \mathcal{T}^{\otimes \mathbb{N}})$ is a Polish space. and Exercise 4.12.1.8 implies that $\Delta_{X^{\mathbb{N}}}$ is closed in $(X^{\mathbb{N}}, \mathcal{T}^{\otimes \mathbb{N}})$. Exercise 6.12.1.2 then implies that $(\Delta_{X^{\mathbb{N}}}, \mathcal{T}^{\otimes \mathbb{N}} \cap \Delta_{X^{\mathbb{N}}})$ is a Polish space. □

Exercise 6.12.1.10. We have that $\mathbb{H}$ is a Polish space.

Proof. Since $[0, 1]$ is a Polish space, Exercise 6.12.1.3 implies that $\mathbb{H}$ is a Polish space. □

Exercise 6.12.1.11. Let $(X, \mathcal{T})$ be a Polish space. Then there exists $B \subset \mathbb{H}$ and $f : X \rightarrow B$ such that B is a G_δ set and f is a homeomorphism.

Hint: Exercise 6.11.1.3

Proof. Define $B \subset \mathbb{H}$ and $f : X \rightarrow B$ as in Exercise 6.11.1.3. Then f is a homeomorphism. Since X is a Polish space, Exercise 6.12.1.6 implies that B is a Polish space. Exercise 6.12.1.8 implies that B is a G_δ -set. □

6.13 Ultrametric Spaces

6.13.1 Introduction

Ultrametric spaces are given by sequences of partitions of X , $(\mathcal{P}_j)_{j \in \mathbb{N}}$, where for each j and $E \in \mathcal{P}_j$, there exists $\mathcal{E} \subset \mathcal{P}_{j+1}$ such that $E = \bigcup_{F \in \mathcal{E}} F$. Then set $d(x, y) = 2^{-n}$ if $n = \max\{j \in \mathbb{N} : \text{there exists } E \in \mathcal{P}_j \text{ such that } x, y \in E\}$.

Definition 6.13.1.1. Let X be a set and $d : X \times X \rightarrow [0, \infty)$. Then d is said to be an **ultrametric on X** if for each $x, y, z \in X$,

1. **(symmetry):** $d(x, y) = d(y, x)$
2. **(definiteness):** $d(x, y) = 0$ iff $x = y$
3. **(strong triangle inequality):** $d(x, z) \leq \max(d(x, y), d(y, z))$

Exercise 6.13.1.2. Let X be a set and d an ultrametric on X . Then d is a metric on X .

Proof. Let $x, y, z \in X$. Since $(d(x, y), d(y, z)) \geq 0$, we have that $d(x, y), d(y, z) \leq d(x, y) + d(y, z)$. Therefore

$$\begin{aligned} d(x, y) &\leq \max(d(x, y), d(y, z)) \\ &\leq d(x, y) + d(y, z) \end{aligned}$$

□

Definition 6.13.1.3. Let X be a set and d an ultrametric on X . Then (X, d) is said to be an **ultrametric space**.

Exercise 6.13.1.4. Isosceles Triangle Property:

Let (X, d) be an ultrametric space and $x, y, z \in X$.

1. If $d(x, y) < d(y, z)$, then

$$\begin{aligned} \max(d(x, y), d(x, z)) &= d(y, z) \\ &= d(x, z) \\ &= \max(d(x, y), d(y, z)). \end{aligned}$$

2. If $d(x, y) \neq d(y, z)$, then $d(x, z) = \max(d(x, y), d(y, z))$.

Proof.

1. Suppose that $d(x, y) < d(y, z)$. Then

$$\begin{aligned} d(x, z) &\leq \max(d(x, y), d(y, z)) \\ &= d(y, z) \end{aligned}$$

For the sake of contradiction, suppose that $d(x, z) \leq d(x, y)$. Since $d(x, y) < d(y, z)$, we have that

$$\begin{aligned} d(y, z) &\leq \max(d(x, y), d(x, z)) \\ &= d(x, y) \\ &< d(y, z) \end{aligned}$$


which is a contradiction. Hence $d(x, y) < d(x, z)$. Therefore, we have that

$$\begin{aligned} d(x, z) &\leq \max(d(x, y), d(y, z)) \\ &= d(y, z) \\ &\leq \max(d(y, x), d(x, z)) \\ &= d(x, z) \end{aligned}$$

Hence

$$\begin{aligned}\max(d(x, y), d(x, z)) &= d(y, z) \\ &= d(x, z) \\ &= \max(d(x, y), d(y, z))\end{aligned}$$

and we have the following isosceles triangle:



2. Suppose that $d(x, y) \neq d(y, z)$.

If $d(x, y) < d(y, z)$, then part (1) implies that $d(x, z) = \max(d(x, y), d(y, z))$.

Suppose that $d(y, z) < d(x, y)$. Then $d(z, y) < d(y, x)$. If we permute x and z in part (1), we see that

$$\begin{aligned}d(x, z) &= \max(d(z, y), d(y, x)) \\ &= \max(d(x, y), d(y, z)).\end{aligned}$$

□

Definition 6.13.1.5. Let (X, d) be an ultrametric space and $r > 0$. We define the

- **open r -ball relation** on X , denoted $\sim_r \subset X \times X$ by $x \sim_r y$ iff $d(x, y) < r$
- **closed r -ball relation** on X , denoted $\simeq_r \subset X \times X$ by $x \simeq_r y$ iff $d(x, y) \leq r$

Exercise 6.13.1.6. Let (X, d) be an ultrametric space and $r > 0$. Then

1. $\sim_r$ is an equivalence relation on X
2. $\simeq_r$ is an equivalence relation on X .

Proof.

1. (a) Let $x \in X$. Since

$$\begin{aligned}d(x, x) &= 0 \\ &< r\end{aligned}$$

we have that $x \sim_r x$.

- (b) Let $x, y \in X$. Suppose that $x \sim_r y$. Then $d(x, y) < r$. This implies that

$$\begin{aligned}d(y, x) &= d(x, y) \\ &< r\end{aligned}$$

So $y \sim_r x$.

- (c) Let $x, y, z \in X$. Suppose that $x \sim_r y$ and $y \sim_r z$. Then $d(x, y) < r$ and $d(y, z) < r$. The strong triangle inequality implies that

$$\begin{aligned}d(x, z) &\leq \max(d(x, y), d(y, z)) \\ &< r\end{aligned}$$

Hence $x \sim_r z$.

2. (a) Let $x \in X$. Since

$$\begin{aligned} d(x, x) &= 0 \\ &\leq r \end{aligned}$$

we have that $x \simeq_r x$.

- (b) Let $x, y \in X$. Suppose that $x \simeq_r y$. Then $d(x, y) \leq r$. This implies that

$$\begin{aligned} d(y, x) &= d(x, y) \\ &\leq r \end{aligned}$$

So $y \simeq_r x$.

- (c) Let $x, y, z \in X$. Suppose that $x \simeq_r y$ and $y \simeq_r z$. Then $d(x, y) \leq r$ and $d(y, z) \leq r$. The strong triangle inequality implies that

$$\begin{aligned} d(x, z) &\leq \max(d(x, y), d(y, z)) \\ &\leq r \end{aligned}$$

Hence $x \simeq_r z$.

□

Definition 6.13.1.7. Let (X, d) be an ultrametric space and $r > 0$. We denote

- the projection of X onto $X/\sim_r$ by $\pi_r^d : X \rightarrow X/\sim_r$
- the projection of X onto $X/\simeq_r$ by $\bar{\pi}_r^d : X \rightarrow X/\simeq_r$

Exercise 6.13.1.8. Let (X, d) be an ultrametric space, $x \in X$ and $r > 0$. Then

1. $\pi_r^d(x) = B(x, r)$
2. $\bar{\pi}_r^d(x) = \bar{B}(x, r)$.

Proof.

1. For each $y \in X$,

$$\begin{aligned} y \sim_r x &\iff d(x, y) < r \\ &\iff y \in B(x, r) \end{aligned}$$

so that $\pi_r^d(x) = B(x, r)$.

2. For each $y \in X$,

$$\begin{aligned} y \simeq_r x &\iff d(x, y) \leq r \\ &\iff y \in \bar{B}(x, r) \end{aligned}$$

so that $\bar{\pi}_r^d(x) = \bar{B}(x, r)$.

□

Exercise 6.13.1.9. Let (X, d) be an ultrametric space, $x, y \in X$ and $s \in [0, \infty)$. If $y \in \bar{B}(x, s)$, then for each $r \in [0, \infty)$, $r \leq s$ implies that $\bar{B}(y, r) \subset \bar{B}(x, s)$.

Proof. Suppose that $y \in \bar{B}(x, s)$. Let $r \in [0, \infty)$. Suppose that $r \leq s$. Let

$$\begin{aligned} z &\in \bar{B}(y, r) \\ &\subset \bar{B}(y, s) \end{aligned}$$

Then $z \simeq_s y$. Since $y \simeq_s x$, the previous exercise implies that $z \simeq_s x$. Hence $z \in \bar{B}(x, s)$. Since $z \in \bar{B}(y, r)$ is arbitrary, $\bar{B}(y, r) \subset \bar{B}(x, s)$. □

Exercise 6.13.1.10. Let (X, d) be an ultrametric space, $x \in X$ and $r > 0$. Then

1. $B(x, r)$ is closed and open,
2. $T(x, r)$ is closed and open,
3. $\bar{B}(x, r)$ is closed and open.

Proof.

1. By definition, $B(x, r)$ is open. Since $\sim_r$ is an equivalence relation, we have that

$$B(x, r)^c = \bigcup_{y \in B(x, r)^c} B(y, r)$$

which is open. Hence $B(x, r)$ is closed.

2. Exercise 6.1.2.9 implies that $T(x, r)$ is closed. Let $y \in T(x, r)$. By definition, $d(x, y) = r$. Let $z \in B(y, r)$. Then

$$\begin{aligned} d(y, z) &< r \\ &= d(x, y) \end{aligned}$$

Exercise 6.13.1.4 then implies that

$$\begin{aligned} d(x, z) &= \max(d(x, y), d(y, z)) \\ &= d(x, y) \\ &= r \end{aligned}$$

Hence $z \in T(x, r)$. Since $z \in B(y, r)$ is arbitrary, we have that for each $z \in B(y, r)$, $z \in T(x, r)$. Thus $B(y, r) \subset T(x, r)$. Since $y \in T(x, r)$ is arbitrary, we have that for each $y \in T(x, r)$, there exists $r > 0$ such that $B(y, r) \subset T(x, r)$. Thus $T(x, r)$ is open.

3. Exercise 6.1.2.9 implies that $\bar{B}(x, r)$ is closed. Since $B(x, r)$ and $T(x, r)$ are open and

$$\bar{B}(x, r) = B(x, r) \cup T(x, r),$$

we have that $\bar{B}(x, r)$ is open.

□

6.13.2 Partitions

Definition 6.13.2.1. Let X be a set. We define the **collection of partitions of X** , denoted $\text{Part}(X)$, by

$$\text{Part}(X) = \{\mathcal{P} \subset \mathcal{P}(X) : \mathcal{P} \text{ is a partition of } X\}$$

Let Γ be a nonempty preordered set and $\mathcal{P} : \Gamma \rightarrow \text{Part}(X)$.

- For each $r \in \Gamma$, we define the **r -th partition relation** on X , denoted $\sim_{\mathcal{P}_r}$, by $x \sim_{\mathcal{P}_r} y$ iff there exists $E \in \mathcal{P}_r$ such that $x \in E$ and $y \in E$.
- For $r \in \Gamma$, we denote the projection of X onto $X/\sim_{\mathcal{P}_r}$ by $\pi_r^{\mathcal{P}} : X \rightarrow X/\sim_{\mathcal{P}_r}$ so that

$$\mathcal{P}_r = \{\pi_r^{\mathcal{P}}(x) : x \in X\}$$

Then

- $\mathcal{P}$ is said to **separate points** if for each $x, y \in X$, $x \neq y$ implies that there exists $r \in \Gamma$ such that $\pi_r^{\mathcal{P}}(x) \neq \pi_r^{\mathcal{P}}(y)$
- $\mathcal{P}$ is said to **collect points** if for each $x, y \in X$, there exists $r \in \Gamma$ such that $\pi_r^{\mathcal{P}}(x) = \pi_r^{\mathcal{P}}(y)$.

- $\mathcal{P}$ is said to be **decreasing** if for each $r, s \in \Gamma$, $r \leq s$ implies that for each $x \in X$, there exists $\mathcal{F}_x \subset X$ such that

$$\pi_r^{\mathcal{P}}(x) = \bigcup_{y \in \mathcal{F}_x} \pi_s^{\mathcal{P}}(y)$$

Exercise 6.13.2.2. Let X be a set, Γ a nonempty preordered set and $\mathcal{P} : \Gamma \rightarrow \text{Part}(X)$. Suppose that $\mathcal{P}$ is decreasing. Let $x \in X$ and $r, s \in \Gamma$. If $r \leq s$, then $\pi_s^{\mathcal{P}}(x) \subset \pi_r^{\mathcal{P}}(x)$.

Proof. Suppose that $r \leq s$. Since $\mathcal{P}$ is decreasing, there exists $\mathcal{F}_x \subset X$ such that

$$\pi_r^{\mathcal{P}}(x) = \bigcup_{y \in \mathcal{F}_x} \pi_s^{\mathcal{P}}(y)$$

Since $x \in \pi_r^{\mathcal{P}}(x)$, there exists $y_0 \in \mathcal{F}_x$ such that $x \in \pi_s^{\mathcal{P}}(y_0)$. Then $\pi_s^{\mathcal{P}}(y_0) = \pi_s^{\mathcal{P}}(x)$ and

$$\begin{aligned} \pi_s^{\mathcal{P}}(x) &= \pi_s^{\mathcal{P}}(y_0) \\ &\subset \bigcup_{y \in \mathcal{F}_x} \pi_s^{\mathcal{P}}(y) \\ &= \pi_r^{\mathcal{P}}(x) \end{aligned}$$

□

Exercise 6.13.2.3. Let X be a set, Γ a nonempty preordered set and $\mathcal{P} : \Gamma \rightarrow \text{Part}(X)$. Suppose that $\mathcal{P}$ is decreasing. Then for each $x, y \in X$ and $s \in \Gamma$, if $\pi_s^{\mathcal{P}}(x) = \pi_s^{\mathcal{P}}(y)$, then for each $r \in \Gamma$, $r \leq s$ implies that $\pi_r^{\mathcal{P}}(x) = \pi_r^{\mathcal{P}}(y)$.

Proof. Let $x, y \in X$ and $s \in \Gamma$. Suppose that $\pi_s^{\mathcal{P}}(x) = \pi_s^{\mathcal{P}}(y)$. Let $r \in \Gamma$. Suppose that $r \leq s$. Since $\mathcal{P}$ is decreasing, there exists $\mathcal{F}_x \subset X$ such that $\pi_r^{\mathcal{P}}(x) = \bigcup_{z \in \mathcal{F}_x} \pi_s^{\mathcal{P}}(z)$. Since $\mathcal{P}_s$ is a partition of X , there exists $x' \in \mathcal{F}_x$ such that $\pi_s^{\mathcal{P}}(x') = \pi_s^{\mathcal{P}}(x)$. Since $\pi_s^{\mathcal{P}}(x) = \pi_s^{\mathcal{P}}(y)$, we have that

$$\begin{aligned} y &\in \bigcup_{z \in \mathcal{F}_x} \pi_s^{\mathcal{P}}(z) \\ &= \pi_r^{\mathcal{P}}(x) \end{aligned}$$

Since $\mathcal{P}_r$ is a partition of X , $\pi_r^{\mathcal{P}}(y) = \pi_r^{\mathcal{P}}(x)$. □

Exercise 6.13.2.4. Let X be a set, Γ a nonempty preordered set and $\mathcal{P} : \Gamma \rightarrow \text{Part}(X)$. Suppose that $\mathcal{P}$ is decreasing. Let $x, y \in X$ and $r, s \in \Gamma$. Suppose $r \leq s$. If $\pi_r^{\mathcal{P}}(x) \cap \pi_s^{\mathcal{P}}(y) \neq \emptyset$, then $\pi_s^{\mathcal{P}}(y) \subset \pi_r^{\mathcal{P}}(x)$.

Proof. Suppose that $\pi_r^{\mathcal{P}}(x) \cap \pi_s^{\mathcal{P}}(y) \neq \emptyset$. Then there exists $z \in X$ such that $z \in \pi_r^{\mathcal{P}}(x) \cap \pi_s^{\mathcal{P}}(y)$. Therefore $\pi_r^{\mathcal{P}}(z) = \pi_r^{\mathcal{P}}(x)$ and $\pi_s^{\mathcal{P}}(z) = \pi_s^{\mathcal{P}}(y)$. Since $r \leq s$, Exercise 6.13.2.3 implies that $\pi_r^{\mathcal{P}}(z) = \pi_r^{\mathcal{P}}(y)$. Since $\mathcal{P}$ is decreasing, we have that

$$\begin{aligned} \pi_s^{\mathcal{P}}(y) &\subset \pi_r^{\mathcal{P}}(y) \\ &= \pi_r^{\mathcal{P}}(z) \\ &= \pi_s^{\mathcal{P}}(z) \end{aligned}$$

□

Exercise 6.13.2.5. Let X be a set, Γ a nonempty preordered set and $\mathcal{P} : \Gamma \rightarrow \text{Part}(X)$. Suppose that $\mathcal{P}$ is decreasing. Let $x, y \in X$. Suppose that there exists $r \in \Gamma$ such that $\pi_r^{\mathcal{P}}(x) \neq \pi_r^{\mathcal{P}}(y)$. Then for each $s \geq r$, $\pi_s^{\mathcal{P}}(x) \neq \pi_s^{\mathcal{P}}(y)$.

Proof. Let $x, y \in X$. Let $s \geq r$. Since $\mathcal{P}$ is decreasing, there exist $\mathcal{F}_x, \mathcal{F}_y \subset X$ such that $\pi_r^{\mathcal{P}}(x) = \bigcup_{z \in \mathcal{F}_x} \pi_s^{\mathcal{P}}(z)$ and $\pi_r^{\mathcal{P}}(y) = \bigcup_{w \in \mathcal{F}_y} \pi_s^{\mathcal{P}}(w)$.

Since $\pi_r^{\mathcal{P}}(x) \cap \pi_r^{\mathcal{P}}(y) = \emptyset$, we have that for each $z \in \mathcal{F}_x$ and $w \in \mathcal{F}_y$, $\pi_s^{\mathcal{P}}(z) \cap \pi_s^{\mathcal{P}}(w) = \emptyset$. Since $\mathcal{P}_s$ is a partition of X , there exist $x' \in \mathcal{F}_x$ and $y' \in \mathcal{F}_y$ such that $\pi_s^{\mathcal{P}}(x') = \pi_s^{\mathcal{P}}(x)$ and $\pi_s^{\mathcal{P}}(y') = \pi_s^{\mathcal{P}}(y)$. Therefore $\pi_s^{\mathcal{P}}(x) \cap \pi_s^{\mathcal{P}}(y) = \emptyset$ and thus $\pi_s^{\mathcal{P}}(x) \neq \pi_s^{\mathcal{P}}(y)$. □

Definition 6.13.2.6. Let X be a set, Γ a nonempty poset with the LUB property, $\mathcal{P} : \Gamma \rightarrow \text{Part}(X)$ and $x, y \in X$.

- We define $A^{\mathcal{P}}(x, y) \subset \Gamma$ by

$$A^{\mathcal{P}}(x, y) := \{r \in \Gamma : \pi_r^{\mathcal{P}}(x) = \pi_r^{\mathcal{P}}(y)\}.$$

- If $A^{\mathcal{P}}(x, y) \neq \emptyset$ and $A^{\mathcal{P}}(x, y)$ is bounded above, we define $\alpha^{\mathcal{P}}(x, y) \in \Gamma$ by

$$\alpha^{\mathcal{P}}(x, y) := \sup A^{\mathcal{P}}(x, y).$$

Then $\mathcal{P}$ is said to be **left-continuous** if for each $x, y \in X$, if $A^{\mathcal{P}}(x, y) \neq \emptyset$ and $A^{\mathcal{P}}(x, y)$ is bounded above, then $\alpha^{\mathcal{P}}(x, y) \in A^{\mathcal{P}}(x, y)$

Definition 6.13.2.7. Let X be a set and $\mathcal{P} : \mathbb{N} \rightarrow \text{Part}(X)$. For $x, y \in X$, we define the **left-continuous extension of $\mathcal{P}$** denoted $\bar{\mathcal{P}} : [0, \infty) \rightarrow \text{Part}(X)$, by

$$\bar{\mathcal{P}}_r = \mathcal{P}_{\lceil r \rceil}$$

Is this even necessary? Is there an example $\mathcal{P} : \mathbb{N} \rightarrow \text{Part}(X)$ that is not already left continuous?, maybe

Exercise 6.13.2.8. Let X be a set and $\mathcal{P} : \mathbb{N} \rightarrow \text{Part}(X)$. Then $\bar{\mathcal{P}}$ is left-continuous.

Proof. Let $x, y \in X$. Suppose that $A^{\bar{\mathcal{P}}}(x, y) \neq \emptyset$ and $\alpha^{\bar{\mathcal{P}}}(x, y) < \infty$. Set $s = \alpha^{\bar{\mathcal{P}}}(x, y)$.

- For the sake of contradiction, suppose that $s \neq \lceil s \rceil$. Then $\lfloor s \rfloor < s < \lceil s \rceil$. Set $\epsilon = 2^{-1} \min(s - \lfloor s \rfloor, \lceil s \rceil - s)$. Then $\epsilon > 0$, $s - \epsilon \in (\lfloor s \rfloor, s)$ and $s + \epsilon \in (s, \lceil s \rceil)$. Since $s - \epsilon \in (\lfloor s \rfloor, s)$, there exists $r \in A^{\bar{\mathcal{P}}}(x, y)$ such that $r \in (s - \epsilon, s]$. Set $t = s + \epsilon$. Since $r, t \in (\lfloor s \rfloor, \lceil s \rceil)$, we have that

$$\lceil r \rceil, \lceil t \rceil = \lceil s \rceil$$

Therefore, the definition of $\bar{\mathcal{P}}$ implies that

$$\begin{aligned} \pi_t^{\bar{\mathcal{P}}}(x) &= \pi_{\lceil t \rceil}^{\mathcal{P}}(x) \\ &= \pi_{\lceil s \rceil}^{\mathcal{P}}(x) \end{aligned}$$

Similarly, $\pi_t^{\bar{\mathcal{P}}}(y) = \pi_{\lceil s \rceil}^{\mathcal{P}}(y)$, $\pi_r^{\bar{\mathcal{P}}}(x) = \pi_{\lceil s \rceil}^{\mathcal{P}}(x)$ and $\pi_r^{\bar{\mathcal{P}}}(y) = \pi_{\lceil s \rceil}^{\mathcal{P}}(y)$.

Since $r \in A^{\bar{\mathcal{P}}}(x, y)$, we have that $\pi_r^{\bar{\mathcal{P}}}(x) = \pi_r^{\bar{\mathcal{P}}}(y)$. By definition of $\bar{\mathcal{P}}$, we have that

$$\begin{aligned} \pi_t^{\bar{\mathcal{P}}}(x) &= \pi_{\lceil s \rceil}^{\mathcal{P}}(x) \\ &= \pi_r^{\bar{\mathcal{P}}}(x) \\ &= \pi_r^{\bar{\mathcal{P}}}(y) \\ &= \pi_{\lceil s \rceil}^{\mathcal{P}}(y) \\ &= \pi_t^{\bar{\mathcal{P}}}(y) \end{aligned}$$

Hence $t \in A^{\bar{\mathcal{P}}}(x, y)$. This is a contradiction since

$$\begin{aligned} \sup A^{\bar{\mathcal{P}}}(x, y) &= s \\ &< t \\ &\leq \sup A^{\bar{\mathcal{P}}}(x, y) \end{aligned}$$

Hence $s = \lceil s \rceil$ and $s \in \mathbb{N}$.

- Choose $r \in A^{\bar{\mathcal{P}}}(x, y)$ such that $r \in (s - 1, s]$. Then $\lceil r \rceil = s$, and

$$\begin{aligned} \pi_s^{\bar{\mathcal{P}}}(x) &= \pi_{\lceil s \rceil}^{\mathcal{P}}(x) \\ &= \pi_s^{\mathcal{P}}(x) \\ &= \pi_{\lceil r \rceil}^{\mathcal{P}}(x) \\ &= \pi_r^{\bar{\mathcal{P}}}(x) \end{aligned}$$

Similarly, $\pi_s^{\bar{\mathcal{P}}}(y) = \pi_r^{\bar{\mathcal{P}}}(y)$. Since $r \in A^{\bar{\mathcal{P}}}(x, y)$, $\pi_r^{\bar{\mathcal{P}}}(x) = \pi_r^{\bar{\mathcal{P}}}(y)$. Hence

$$\begin{aligned}\pi_s^{\bar{\mathcal{P}}}(x) &= \pi_r^{\bar{\mathcal{P}}}(x) \\ &= \pi_r^{\bar{\mathcal{P}}}(y) \\ &= \pi_s^{\bar{\mathcal{P}}}(y)\end{aligned}$$

Hence

$$\begin{aligned}\alpha^{\bar{\mathcal{P}}}(x, y) &= s \\ &\in A^{\bar{\mathcal{P}}}(x, y)\end{aligned}$$

Since $x, y \in X$ with $A^{\bar{\mathcal{P}}}(x, y) \neq \emptyset$ and $\alpha^{\bar{\mathcal{P}}}(x, y) < \infty$ are arbitrary, $\bar{\mathcal{P}}$ is left-continuous. $\square$

Definition 6.13.2.9. Let X be a set, Γ a nonempty poset with the LUB property and $\mathcal{P} : \Gamma \rightarrow \text{Part}(X)$. Then $\mathcal{P}$ is said to be **ultrametric-equivalent** if

1. $\mathcal{P}$ separates points
2. $\mathcal{P}$ collects points
3. $\mathcal{P}$ is decreasing
4. $\mathcal{P}$ is left-continuous

Exercise 6.13.2.10. Let X be a set and $\mathcal{P} : \mathbb{N} \rightarrow \text{Part}(X)$. Suppose that $\mathcal{P}$ separates points, collects points and is decreasing. Then $\bar{\mathcal{P}}$ is ultrametric-equivalent.

Proof.

1. Let $x, y \in X$. Suppose that $x \neq y$. Since $\mathcal{P}$ separates points, there exists $n \in \mathbb{N}$ such that

$$\begin{aligned}\pi_n^{\bar{\mathcal{P}}}(x) &= \pi_n^{\mathcal{P}}(x) \\ &\neq \pi_n^{\mathcal{P}}(y) \\ &= \pi_n^{\bar{\mathcal{P}}}(y)\end{aligned}$$

Since $x, y \in X$ with $x \neq y$ are arbitrary, $\bar{\mathcal{P}}$ separates points.

2. Let $x, y \in X$. Since $\mathcal{P}$ collects points, there exists $n \in \mathbb{N}$ such that

$$\begin{aligned}\pi_n^{\bar{\mathcal{P}}}(x) &= \pi_n^{\mathcal{P}}(x) \\ &= \pi_n^{\mathcal{P}}(y) \\ &= \pi_n^{\bar{\mathcal{P}}}(y)\end{aligned}$$

Since $x, y \in X$ are arbitrary, $\bar{\mathcal{P}}$ collects points.

3. Let $r, s \in (0, \infty)$. Suppose that $r \leq s$. Let $x \in X$. Since $r \leq s$, we have that $\lceil r \rceil \leq \lceil s \rceil$. Since $\mathcal{P}$ is decreasing, there exists $\mathcal{F}_x \subset X$ such that

$$\begin{aligned}\pi_r^{\bar{\mathcal{P}}}(x) &= \pi_{\lceil r \rceil}^{\mathcal{P}}(x) \\ &= \bigcup_{y \in \mathcal{F}_x} \pi_{\lceil s \rceil}^{\mathcal{P}}(y) \\ &= \bigcup_{y \in \mathcal{F}_x} \pi_s^{\bar{\mathcal{P}}}(y)\end{aligned}$$

Since $r, s \in (0, \infty)$ with $r \leq s$ and $x \in X$ are arbitrary, we have that $\bar{\mathcal{P}}$ is decreasing.

4. Exercise 6.13.2.8 implies that $\bar{\mathcal{P}}$ is left continuous.

Since $\bar{\mathcal{P}}$ separates points, collects points, is decreasing and is left continuous, $\bar{\mathcal{P}}$ is ultrametric-equivalent. $\square$

Exercise 6.13.2.11. Let X be a set, Γ a nonempty poset with the LUB property and $\mathcal{P} : \Gamma \rightarrow \text{Part}(X)$. Suppose that $\mathcal{P}$ is ultrametric-equivalent. Then for each $x, y \in X$, if $x \neq y$, then $\alpha^{\mathcal{P}}(x, y)$ exists and $A^{\mathcal{P}}(x, y) = (-\infty, \alpha^{\mathcal{P}}(x, y)]$

Proof. Let $x, y \in X$. Suppose that $x \neq y$. Since $\mathcal{P}$ collects points, there exists $s > 0$ such that $\pi_s^{\mathcal{P}}(x) = \pi_s^{\mathcal{P}}(y)$. Hence $A^{\mathcal{P}}(x, y) \neq \emptyset$. Since $\mathcal{P}$ separates points, there exists $r \in \Gamma$ such that $r \notin A^{\mathcal{P}}(x, y)$. If there exists $s \in A^{\mathcal{P}}(x, y)$ such that $r \leq s$, then Exercise 6.13.2.3 implies that $r \in A^{\mathcal{P}}(x, y)$ which is a contradiction. Thus for each $s \in A^{\mathcal{P}}(x, y)$, $s < r$. Hence r is an upper bound of $A^{\mathcal{P}}(x, y)$ and $A^{\mathcal{P}}(x, y)$ is bounded above. Since $A^{\mathcal{P}}(x, y)$ is nonempty and bounded above and Γ has the LUB property, $\alpha^{\mathcal{P}}(x, y)$ exists. By definition of the supremum, $A^{\mathcal{P}}(x, y) \subset (-\infty, \alpha^{\mathcal{P}}(x, y)]$. Since $\mathcal{P}$ is left-continuous, $\alpha^{\mathcal{P}}(x, y) \in A^{\mathcal{P}}(x, y)$. Exercise 6.13.2.3 then implies that $(-\infty, \alpha^{\mathcal{P}}(x, y)] \subset A^{\mathcal{P}}(x, y)$. Hence $A^{\mathcal{P}}(x, y) = (-\infty, \alpha^{\mathcal{P}}(x, y)]$. $\square$

Exercise 6.13.2.12. Fundamental Example:

Let X be a set, Γ a nonempty poset with the LUB property and $\mathcal{P} : \Gamma \rightarrow \text{Part}(X)$ and $(\epsilon_\gamma)_{\gamma \in \Gamma} \subset [0, \infty)$. Suppose that $(\epsilon_\gamma)_{\gamma \in \Gamma}$ is strictly decreasing and $\mathcal{P}$ is ultrametric-equivalent. Define $d^{\mathcal{P}} : X \times X \rightarrow (0, \infty)$ by

$$d^{\mathcal{P}}(x, y) = \begin{cases} \exp(-\epsilon_{\alpha^{\mathcal{P}}(x, y)}) & x \neq y \\ 0 & x = y \end{cases}$$

Then $d^{\mathcal{P}}$ is an ultrametric on X .

Proof. Let $x, y \in X$.

1. Suppose that $x \neq y$. Since $\sim_{\mathcal{P}_r}$ is symmetric,

$$\begin{aligned} \alpha^{\mathcal{P}}(x, y) &= \sup A^{\mathcal{P}}(x, y) \\ &= \sup A^{\mathcal{P}}(y, x) \\ &= \alpha^{\mathcal{P}}(y, x). \end{aligned}$$

Hence $d^{\mathcal{P}}(x, y) = d^{\mathcal{P}}(y, x)$. If $x = y$, then

$$\begin{aligned} d^{\mathcal{P}}(x, y) &= 0 \\ &= d^{\mathcal{P}}(y, x). \end{aligned}$$

2. By definition, $d^{\mathcal{P}}(x, y) = 0$ iff $x = y$.

3. If $x = z$, $x = y$ or $y = z$, then $d^{\mathcal{P}}(x, z) \leq \max(d^{\mathcal{P}}(x, y), d^{\mathcal{P}}(y, z))$. Suppose that $x \neq z$, $x \neq y$ and $y \neq z$. Then $d^{\mathcal{P}}(x, z) \neq 0$, $d^{\mathcal{P}}(x, y) \neq 0$ and $d^{\mathcal{P}}(y, z) \neq 0$.

Suppose that $d^{\mathcal{P}}(x, y) \leq d^{\mathcal{P}}(y, z)$. Then $\alpha^{\mathcal{P}}(y, z) \leq \alpha^{\mathcal{P}}(x, y)$. Exercise 6.13.2.11 implies that $\alpha^{\mathcal{P}}(y, z) \in A^{\mathcal{P}}(x, y) \cap A^{\mathcal{P}}(y, z)$. Hence

$$\begin{aligned} \pi_{\alpha^{\mathcal{P}}(y, z)}^{\mathcal{P}}(x) &= \pi_{\alpha^{\mathcal{P}}(y, z)}^{\mathcal{P}}(y) \\ &= \pi_{\alpha^{\mathcal{P}}(y, z)}^{\mathcal{P}}(z) \end{aligned}$$

Hence $\alpha^{\mathcal{P}}(y, z) \in A^{\mathcal{P}}(x, z)$. Therefore $\alpha^{\mathcal{P}}(y, z) \leq \alpha^{\mathcal{P}}(x, z)$ which implies that

$$\begin{aligned} d^{\mathcal{P}}(x, z) &\leq d^{\mathcal{P}}(y, z) \\ &= \max(d^{\mathcal{P}}(x, y), d^{\mathcal{P}}(y, z)) \end{aligned}$$

Similarly, if $d^{\mathcal{P}}(y, z) \leq d^{\mathcal{P}}(x, y)$, then

$$\begin{aligned} d^{\mathcal{P}}(x, z) &\leq d^{\mathcal{P}}(x, y) \\ &= \max(d^{\mathcal{P}}(x, y), d^{\mathcal{P}}(y, z)) \end{aligned}$$

Hence $d^{\mathcal{P}}$ satisfies the strong triangle inequality. Therefore $d^{\mathcal{P}}$ is an ultrametric on X . $\square$

Definition 6.13.2.13. Let (X, d) be an ultrametric space and Γ a directed poset. Write $d(X \times X) \setminus \{0\} = (\epsilon_\gamma)_{\gamma \in \Gamma}$. Suppose that $(\epsilon_\gamma)_{\gamma \in \Gamma}$ is strictly decreasing and $\epsilon_\gamma \rightarrow 0$. We define $\mathcal{P}^d : \Gamma \rightarrow \text{Part}(X)$ by

$$\mathcal{P}_\gamma^d = \{\bar{\pi}_{\epsilon_\gamma}^d(x) : x \in X\}.$$

Exercise 6.13.2.14. Let (X, d) be an ultrametric space and Γ a directed poset. Write $d(X \times X) \setminus \{0\} = (\epsilon_\gamma)_{\gamma \in \Gamma}$. Suppose that $(\epsilon_\gamma)_{\gamma \in \Gamma}$ is strictly decreasing and $\epsilon_\gamma \rightarrow 0$. Then $\mathcal{P}^d$

1. $\mathcal{P}^d$ separates points
2. $\mathcal{P}^d$ collects points,
3. $\mathcal{P}^d$ is decreasing.

Proof.

1. Let $x, y \in X$. Suppose that $x \neq y$. Then $d(x, y) > 0$. Thus there exists $s \in \Gamma$ such that $d(x, y) = \epsilon_s$. Since $(\epsilon_\gamma)_{\gamma \in \Gamma}$ is strictly decreasing and $\epsilon_\gamma \rightarrow 0$, there exists $r \in \Gamma$ such that $s < r$ and $\epsilon_r < \epsilon_s$. Then

$$\begin{aligned} d(x, y) &= \epsilon_s \\ &> \epsilon_r. \end{aligned}$$

Hence $x \not\sim_{\epsilon_r} y$ and therefore

$$\begin{aligned} \pi_r^{\mathcal{P}^d}(x) \cap \pi_r^{\mathcal{P}^d}(y) &= \bar{B}(x, \epsilon_r) \cap \bar{B}(y, \epsilon_r) \\ &= \emptyset. \end{aligned}$$

Since $x, y \in X$ with $x \neq y$ are arbitrary, we have that for each $x, y \in X$, there exists $r \in \Gamma$ such that $\pi_r^{\mathcal{P}^d}(x) \neq \pi_r^{\mathcal{P}^d}(y)$. Hence .

2. Let $x, y \in X$.

- Suppose that $x = y$. Since Γ is directed, $\Gamma \neq \emptyset$. Thus there exists $r \in \Gamma$ such that

$$\begin{aligned} d(x, y) &= 0 \\ &\leq \epsilon_r. \end{aligned}$$

Then

$$\begin{aligned} \pi_r^{\mathcal{P}^d}(x) &= \bar{B}(x, \epsilon_r) \\ &= \bar{B}(y, \epsilon_r) \\ &= \pi_r^{\mathcal{P}^d}(y). \end{aligned}$$

- Suppose that $x \neq y$. Then $d(x, y) > 0$. Thus there exists $r \in \Gamma$ such that $\epsilon_r = d(x, y)$. Then $x \simeq_{\epsilon_r} y$ and

$$\begin{aligned} \pi_r^{\mathcal{P}^d}(x) &= \bar{B}(x, \epsilon_r) \\ &= \bar{B}(y, \epsilon_r) \\ &= \pi_r^{\mathcal{P}^d}(y). \end{aligned}$$

Since $x, y \in X$ are arbitrary, we have that for each $x, y \in X$, there exists $r \in \Gamma$ such that $\pi_r^{\mathcal{P}^d}(x) = \pi_r^{\mathcal{P}^d}(y)$. Hence $\mathcal{P}^d$ collects points.

3. Let $r, s \in \Gamma$. Suppose that $r \leq s$. Since $(\epsilon_\gamma)_{\gamma \in \Gamma}$ is strictly decreasing, $\epsilon_s \leq \epsilon_r$. Let $x \in X$. Choose $\mathcal{F}_x = \pi_r^{\mathcal{P}^d}(x)$. Let $y \in \bar{B}(x, \epsilon_r)$. By definition of $\mathcal{P}^d$ and Exercise 6.13.1.8, we have that

$$\begin{aligned}\pi_r^{\mathcal{P}^d}(x) &= \bar{\pi}_{\epsilon_r}^d(x) \\ &= \bar{B}(x, \epsilon_r)\end{aligned}$$

and

$$\begin{aligned}\pi_s^{\mathcal{P}^d}(y) &= \bar{\pi}_{\epsilon_s}^d(y) \\ &= \bar{B}(y, \epsilon_s).\end{aligned}$$

Since $\epsilon_s \leq \epsilon_r$ and $y \in \bar{B}(x, \epsilon_r)$, Exercise 6.13.1.9 implies that

$$\begin{aligned}\bar{B}(y, \epsilon_s) &\subset \bar{B}(x, \epsilon_r) \\ &= \pi_r^{\mathcal{P}^d}(x)\end{aligned}$$

Since $y \in \bar{B}(x, \epsilon_r)$ is arbitrary, we have that for each $y \in \bar{B}(x, \epsilon_r)$, $\bar{B}(y, \epsilon_s) \subset \bar{B}(x, \epsilon_r)$. Therefore

$$\begin{aligned}\pi_r^{\mathcal{P}^d}(x) &= \bar{B}(x, \epsilon_r) \\ &= \bigcup_{y \in \bar{B}(x, \epsilon_r)} \bar{B}(y, \epsilon_s) \\ &= \bigcup_{y \in \mathcal{F}_x} \pi_s^{\mathcal{P}^d}(y)\end{aligned}$$

Since $x \in X$ is arbitrary, we have that for each $x \in X$, there exists $\mathcal{F}_x \subset X$ such that $\pi_r^{\mathcal{P}^d}(x) = \bigcup_{y \in \mathcal{F}_x} \pi_s^{\mathcal{P}^d}(y)$. Since $r, s \in \Gamma$ with $r \leq s$ are arbitrary, we have that for each $r, s \in \Gamma$, $r \leq s$ implies that for each $x \in X$, there exists $\mathcal{F}_x \subset X$ such that $\pi_r^{\mathcal{P}^d}(x) = \bigcup_{y \in \mathcal{F}_x} \pi_s^{\mathcal{P}^d}(y)$. Therefore $\mathcal{P}^d$ is decreasing. □

Exercise 6.13.2.15. Let (X, d) be an ultrametric space. If $(X, \mathcal{T}_d)$ is compact, then for each $(a_n)_{n \in \mathbb{N}} \subset d(X \times X)$, $(a_n)_{n \in \mathbb{N}}$ is injective implies that $a_n \rightarrow 0$.

Hint: Exercise 6.13.1.4

Proof. Suppose that $(X, \mathcal{T}_d)$ is compact. For the sake of contradiction, suppose that there exists $(a_n)_{n \in \mathbb{N}} \subset d(X \times X)$ such that $(a_n)_{n \in \mathbb{N}}$ is injective and $a_n \not\rightarrow 0$. Then there exists $\epsilon > 0$ and $(a_{n_j})_{j \in \mathbb{N}} \subset (a_n)_{n \in \mathbb{N}}$ such that for each $n \in \mathbb{N}$, $a_{n_j} \geq \epsilon$. Define $(b_j)_{j \in \mathbb{N}} \subset d(X \times X)$ by $b_j := a_{n_j}$. Since $(b_n)_{n \in \mathbb{N}} \subset d(X \times X)$, there exists $(x_n, y_n) \in X \times X$ such that for each $n \in \mathbb{N}$, $d(x_n, y_n) = b_n$. Since $(X, \mathcal{T}_d)$ is compact, $(X \times X, \mathcal{T}_d \otimes \mathcal{T}_d)$ is compact and there exists a subsequence $(x_{n_j}, y_{n_j})_{j \in \mathbb{N}} \subset (x_n, y_n)_{n \in \mathbb{N}}$ and $(x, y) \in X \times X$ such that $(x_{n_j}, y_{n_j}) \rightarrow (x, y)$. Since for each $j \in \mathbb{N}$, $b_{n_j} \geq \epsilon$, we have that

$$\begin{aligned}d(x, y) &= \lim_{j \rightarrow \infty} d(x_{n_j}, y_{n_j}) \\ &= \lim_{j \rightarrow \infty} b_{n_j} \\ &\geq \epsilon.\end{aligned}$$

Since $(x_{n_j}, y_{n_j}) \rightarrow (x, y)$, we have that $x_{n_j} \rightarrow x$ and $y_{n_j} \rightarrow y$. Then there exists $J_1, J_2 \in \mathbb{N}$ such that for each $j \in \mathbb{N}$, $j \geq J_1$ implies that $d(x_{n_j}, x) < \epsilon$ and $j \geq J_2$ implies that $d(y_{n_j}, y) < \epsilon$. Define $J \in \mathbb{N}$ by $J := \max\{J_1, J_2\}$. Let $j \in \mathbb{N}$. Suppose that $j \geq J$. Since

$$\begin{aligned}d(x_{n_j}, x), d(y_{n_j}, y) &< \epsilon \\ &\leq d(x, y),\end{aligned}$$

Exercise 6.13.1.4 implies that

$$\begin{aligned} d(x, y_{n_j}) &= \max\{d(x, y), d(y, y_{n_j})\} \\ &= d(x, y) \\ &\geq \epsilon \\ &> d(x_{n_j}, x). \end{aligned}$$

Another application of Exercise 6.13.1.4 implies that

$$\begin{aligned} d(x_{n_j}, y_{n_j}) &= \max\{d(x_{n_j}, x), d(x, y_{n_j})\} \\ &= d(x, y_{n_j}). \end{aligned}$$

Therefore

$$\begin{aligned} b_{n_j} &= d(x_{n_j}, y_{n_j}) \\ &= d(x, y_{n_j}) \\ &= d(x, y). \end{aligned}$$

Since $j \in \mathbb{N}$ with $j \geq J$ is arbitrary, we have that for each $j \in \mathbb{N}$, $j \geq J$ implies that $b_{n_j} = d(x, y)$. This is a contradiction since $(b_{n_j})_{j \in \mathbb{N}}$ is injective. Thus for each $(a_n)_{n \in \mathbb{N}} \subset d(X \times X)$, $(a_n)_{n \in \mathbb{N}}$ is injective implies that $a_n \rightarrow 0$. $\square$

Exercise 6.13.2.16. Let (X, d) be an ultrametric space. If $(X, \mathcal{T}_d)$ is compact, then $d(X \times X) \setminus \{0\}$ is discrete.

Proof. Set $A := d(X \times X) \setminus \{0\}$. For the sake of contradiction, suppose that A is not discrete. Then there exists $a \in A$ and $(a_n)_{n \in \mathbb{N}} \subset A \setminus \{a\}$ such that $(a_n)_{n \in \mathbb{N}}$ is injective and $a_n \rightarrow a$. Exercise 6.13.2.15 implies that $a_n \rightarrow 0$. Thus $a = 0$, which is a contradiction. Hence A is discrete. $\square$

Exercise 6.13.2.17. Let (X, d) be an ultrametric space. Suppose that $(X, \mathcal{T}_d)$ is compact. Write $d(X \times X) \setminus \{0\} = (\epsilon_n)_{n \in \mathbb{N}}$. Suppose that $(\epsilon_n)_{n \in \mathbb{N}}$ is strictly decreasing and $\epsilon_n \rightarrow 0$. Then $\mathcal{P}^d$ is ultrametric equivalent.

Proof.

- Exercise 6.13.2.14 implies that $\mathcal{P}^d$ separates points, collects points and is decreasing.
- Let $x, y \in X$. Suppose that $A^{\mathcal{P}^d}(x, y) \neq \emptyset$ and $A^{\mathcal{P}^d}(x, y)$ is bounded above. Since $A^{\mathcal{P}^d}(x, y) \subset \mathbb{N}$, we have that

$$\begin{aligned} \alpha^{\mathcal{P}^d}(x, y) &= \sup A^{\mathcal{P}^d}(x, y) \\ &= \max A^{\mathcal{P}^d}(x, y) \\ &\in A^{\mathcal{P}^d}(x, y). \end{aligned}$$

Since $x, y \in X$ such that $A^{\mathcal{P}^d}(x, y) \neq \emptyset$ and $A^{\mathcal{P}^d}(x, y)$ is bounded above are arbitrary, we have that for each $x, y \in X$, $A^{\mathcal{P}}(x, y) \neq \emptyset$ and $A^{\mathcal{P}}(x, y)$ is bounded above implies that $\alpha^{\mathcal{P}}(x, y) \in A^{\mathcal{P}}(x, y)$. Thus $\mathcal{P}^d$ is left-continuous.

Hence $\mathcal{P}^d : \mathbb{N} \rightarrow \text{Part}(X)$ is ultrametric-equivalent. $\square$

Exercise 6.13.2.18. Conjecture:

Let (X, d) be an ultrametric space. Then $d^{\mathcal{P}^d} \sim_{\text{Top}} d$.

FINISH!!!

Exercise 6.13.2.19. Conjecture:

Let (X, d) be an ultrametric space. If $(X, \mathcal{T}_d)$ is compact, then there exists $\mathcal{P} : \mathbb{N} \rightarrow \text{Part}(X)$ such that $\mathcal{P}$ separates points, collects points, is decreasing and $d^{\mathcal{P}} \sim_{\text{Top}} d$.

make exercise in earlier section that if $d(X \times X)$ is finite, then $\mathcal{T}_d = \mathcal{P}(X)$ i.e. d is basically the discrete metric.

Proof. Suppose that $(X, \mathcal{T}_d)$ is compact.

- Suppose that $d(X \times X)$ is finite. Set $n := \#d(X \times X)$ and write $d(X \times X) = (a_j)_{j \in [n]}$ where $a_1 \geq \dots \geq a_n$. Define $\mathcal{P} : \mathbb{N} \rightarrow \text{Part}(X)$ by $\mathcal{P}_j := \{\bar{B}(x, a_n) : x \in X\}$. **FINISH!!!**
- Suppose that $d(X \times X)$ is infinite. Since $(X, \mathcal{T}_d)$ is compact, there exists $a \in d(X \times X)$ such that $a = \max d(X \times X)$. Then there exists $(a_n)_{n \in \mathbb{N}} \subset d(X \times X)$ such that $a_1 = a$ and $(a_n)_{n \in \mathbb{N}}$ is strictly decreasing. Exercise 6.13.2.15 then implies that $a_n \rightarrow 0$. Define $\mathcal{P} : \mathbb{N} \rightarrow \text{Part}(X)$ by $\mathcal{P}_n := \{\bar{B}(x, a_n) : x \in X\}$.
 - Let $x, y \in X$. Suppose that $x \neq y$. Define $\epsilon > 0$ by $\epsilon := d(x, y)$. Since $a_n \rightarrow 0$, there exists $N \in \mathbb{N}$ such that for each $n \in \mathbb{N}$, $n \geq N$ implies that $a_n < \epsilon$. If $y \in \bar{B}(x, a_N)$, then

$$\begin{aligned} d(x, y) &\leq a_N \\ &< \epsilon \\ &= d(x, y), \end{aligned}$$

which is a contradiction. Hence $y \notin \bar{B}(x, a_N)$ and therefore

$$\begin{aligned} \pi_n^{\mathcal{P}}(x) &= \bar{B}(x, a_N) \\ &\neq \bar{B}(y, a_N) \\ &= \pi_1^{\mathcal{P}}(y). \end{aligned}$$

Since $x, y \in X$ with $x \neq y$ are arbitrary, we have that for each $x, y \in X$, $x \neq y$ implies that there exists $N \in \mathbb{N}$ such that $\pi_n^{\mathcal{P}}(x) \neq \pi_n^{\mathcal{P}}(y)$. Hence $\mathcal{P}$ separates points.

- Let $x, y \in X$. Since $a_1 = a$,

$$\begin{aligned} \pi_1^{\mathcal{P}}(x) &= \bar{B}(x, a) \\ &= X \\ &= \bar{B}(y, a) \\ &= \pi_1^{\mathcal{P}}(y). \end{aligned}$$

Since $x, y \in X$ are arbitrary, we have that for each $x, y \in X$, $\pi_1^{\mathcal{P}}(x) = \pi_1^{\mathcal{P}}(y)$. Thus $\mathcal{P}$ collects points.

- Let $m, n \in \mathbb{N}$. Suppose that $m \leq n$. Let $x \in X$. Define $\mathcal{F}_x := \bar{B}(x, a_m)$. Since $m \leq n$, $a_n \leq a_m$. Exercise 6.13.1.9 then implies that

$$\begin{aligned} \pi_m^{\mathcal{P}}(x) &= \bar{B}(x, a_m) \\ &= \bigcup_{y \in \mathcal{F}_x} \bar{B}(x, a_n) \\ &= \bigcup_{y \in \mathcal{F}_x} \pi_n^{\mathcal{P}}(y). \end{aligned}$$

Since $x \in X$ is arbitrary, we have that for each $x \in X$, there exists $\mathcal{F}_x \subset X$ such that $\pi_m^{\mathcal{P}}(x) = \bigcup_{y \in \mathcal{F}_x} \pi_n^{\mathcal{P}}(y)$. Since $m, n \in \mathbb{N}$ with $m \leq n$ are arbitrary, we have that for each $m, n \in \mathbb{N}$, $m \leq n$ implies that for each $x \in X$, there exists $\mathcal{F}_x \subset X$ such that $\pi_m^{\mathcal{P}}(x) = \bigcup_{y \in \mathcal{F}_x} \pi_n^{\mathcal{P}}(y)$. Thus $\mathcal{P}$ is decreasing.

Exercise 6.13.2.10 then implies that $\bar{\mathcal{P}}$ is ultrametric-equivalent and Exercise 6.13.2.12 implies that $d^{\mathcal{P}}$ is an ultrametric on X .
need to show $d^{\bar{\mathcal{P}}} \sim_{\text{Top}} d$.

□

want to characterize when a discrete valued metric is basically a tree.

6.14 Zero-dimensional Metric Spaces

6.14.1 Second-Countability

Note 6.14.1.1. Zero dimensional metric spaces are generalizations of countable sets. The following exercise captures some of the behavior of such spaces. [rewrite or write more here](#)

Exercise 6.14.1.2. Let (X, d) be a metric space, $U \in \mathcal{T}_d$ and $\epsilon > 0$. Suppose that $(X, \mathcal{T}_d)$ is separable and zero-dimensional and U is not compact. Then there exists $(V_n)_{n \in \mathbb{N}} \subset \mathcal{T}_d$ such that

- for each $n \in \mathbb{N}$, $V_n \neq \emptyset$, V_n is closed and $\text{diam } V_n \leq \epsilon$,
- $(V_n)_{n \in \mathbb{N}}$ is disjoint,
- $U = \bigcup_{n \in \mathbb{N}} V_n$

[space out with steps and give hints!!!](#)

Proof. Exercise 6.2.1.1 implies that $(X, \mathcal{T}_d)$ is second-countable. Then Exercise 4.13.2.11 implies that $(U, \mathcal{T}_d \cap U)$ is second-countable. Exercise 4.13.2.9 then implies that $(U, \mathcal{T}_d \cap U)$ is Lindelöf. Since $U \in \mathcal{T}_d$, we have that $\mathcal{T}_d \cap U \subset \mathcal{T}_d$. Since U is not compact, there exists $\mathcal{V} \subset \mathcal{T}_d \cap U$ such that $\mathcal{V}$ is an open cover of U and for each $\mathcal{V}' \subset \mathcal{V}$, $\mathcal{V}'$ is finite implies that $\mathcal{V}'$ is not an open cover of U . Since $(X, \mathcal{T}_d)$ is zero-dimensional, there exists $\mathcal{B} \subset \mathcal{T}_d$ such that $\mathcal{B}$ is a basis for $\mathcal{T}_d$ and for each $B \in \mathcal{B}$, B is closed. Define $\mathcal{B}_\epsilon \subset \mathcal{T}_d$ by $\mathcal{B}_\epsilon := \{B \in \mathcal{B} : \text{diam}(B) \leq \epsilon\}$. Exercise 6.1.2.7 implies that $\mathcal{B}_\epsilon$ is a basis for $\mathcal{T}_d$. Define $\mathcal{W} \subset \mathcal{T}_d \cap U$ by

$$\mathcal{W} := \{W \in \mathcal{T}_d \cap U : W \neq \emptyset, W \text{ is } \mathcal{T}_d\text{-closed, } \text{diam } W \leq \epsilon \text{ and there exists } V \in \mathcal{V} \text{ such that } W \subset V\}.$$

Let $x \in U$. Since $\mathcal{V} \subset \mathcal{T}_d \cap U$ is a $\mathcal{T}_d \cap U$ -open cover of U , there exists $V \in \mathcal{V}$ such that $x \in V$. Since $V \in \mathcal{T}_d$ and $\mathcal{B}_\epsilon$ is a basis for $\mathcal{T}_d$, there exists $B \in \mathcal{B}_\epsilon$ such that $x \in B$ and $B \subset V$. Thus $B \neq \emptyset$, $B \subset V$ and $\text{diam } B \leq \epsilon$. Hence $B \in \mathcal{W}$. Since $x \in U$ is arbitrary, we have that for each $x \in U$, there exists $B \in \mathcal{W}$ such that $x \in B$. Hence $\mathcal{W}$ is a $\mathcal{T}_d \cap U$ -open cover of U . Since $(U, \mathcal{T}_d \cap U)$ is Lindelöf, there exists $(W'_n)_{n \in \mathbb{N}} \subset \mathcal{W}$ such that $(W'_n)_{n \in \mathbb{N}}$ is an open cover of U . Define $(W_n)_{n \in \mathbb{N}} \subset \mathcal{T}_d \cap U$ by

$$W_n := \begin{cases} W'_1, & n = 1 \\ W'_n \setminus \bigcup_{j=1}^{n-1} W'_j, & n \geq 2. \end{cases}$$

Then for each $n \in \mathbb{N}$, W_n is closed,

$$\begin{aligned} \text{diam } W_n &\leq \text{diam } W'_n \\ &\leq \epsilon, \end{aligned}$$

and

$$\begin{aligned} U &= \bigcup_{n \in \mathbb{N}} W'_n \\ &= \bigcup_{n \in \mathbb{N}} W_n. \end{aligned}$$

Define $J \subset \mathbb{N}$ by $J := \{n \in \mathbb{N} : W_n \neq \emptyset\}$. For the sake of contradiction, suppose that $\#J < \infty$. Then there exists $N \in \mathbb{N}$ such that for each $n \in \mathbb{N}$, $n \geq N$ implies that $W_n = \emptyset$. Then

$$\begin{aligned} U &= \bigcup_{n \in \mathbb{N}} W_n \\ &= \bigcup_{n=1}^N W_n \\ &\subset \bigcup_{n=1}^N W'_n. \end{aligned}$$

Since for each $n \in \mathbb{N}$, $W'_n \subset U$, we have that $U = \bigcup_{n=1}^N W'_n$. By construction, for each $n \in \mathbb{N}$, there exists $V_n \in \mathcal{V}$ such that $W'_n \subset V_n$. Then

$$\begin{aligned} U &= \bigcup_{n=1}^N W'_n \\ &\subset \bigcup_{n=1}^N V_n. \end{aligned}$$

Therefore $(V_n)_{n=1}^N$ is an open cover of U . This is a contradiction. Hence $\#J = \infty$. We define $(j_n)_{n \in \mathbb{N}} \subset \mathbb{N}$ and $(V_n)_{n \in \mathbb{N}} \subset \mathcal{W}$ by

$$j_n := \begin{cases} 1, & n = 1 \\ \min\{j \in [j_{n-1}]^c : W_j \neq \emptyset\}, & n \geq 2 \end{cases}$$

and $V_n := W_{j_n}$. Then for each $n \in \mathbb{N}$, $V_n \neq \emptyset$, V_n is closed, $\text{diam } V_n \leq \epsilon$, $(V_n)_{n \in \mathbb{N}}$ is disjoint, and $U = \bigcup_{n \in \mathbb{N}} V_n$. $\square$

Note 6.14.1.3. We recall Definition 6.2.1.4 (finish statement).

Exercise 6.14.1.4. Let (X, d) be a metric space. Suppose that $(X, \mathcal{T}_d)$ is separable and zero-dimensional, $X \neq \emptyset$ and for each $x \in X$, x is a $\mathcal{T}_d$ -condensation point of X . Then there exists $(C_\gamma)_{\gamma \in \Gamma} \subset \mathcal{T}_d$ and $(x_\gamma)_{\gamma \in \Gamma} \in \prod_{\gamma \in \Gamma} C_\gamma$ such that

1.
 - there exists $x_0 \in X$ such that $X \setminus \{x_0\} = \bigcup_{|\gamma|=1} C_\gamma$,
 - $(C_\gamma)_{\gamma \in \mathbb{N}}$ is disjoint
2. for each $\gamma \in \Gamma$,
 - (a) C_γ is $\mathcal{T}_d$ -closed and $C_\gamma \neq \emptyset$,
 - (b) $\text{diam } C_\gamma \leq |\gamma|^{-1}$,
 - (c) for each $n \in \mathbb{N}$, $|\gamma| = n$ implies that
 - $C_\gamma \setminus \{x_\gamma\} = \bigcup_{\gamma' \in (\pi_{[n]}^{[n+1]})^{-1}(\{\gamma\})} C_{\gamma'}$
 - $(C_{\gamma'})_{\gamma' \in (\pi_{[n]}^{[n+1]})^{-1}(\{\gamma\})}$ is disjoint
 - (d) for each $x \in C_\gamma$, x is a $\mathcal{T}_d \cap C_\gamma$ -condensation point of C_γ .

possibly reformat to say the same more efficiently, maybe index by Γ including length 0 indices?

Proof. We define $(C_\gamma)_{\gamma \in \Gamma}$ and $(x_\gamma)_{\gamma \in \Gamma} \in \prod_{\gamma \in \Gamma} C_\gamma$ inductively as follows:

• **Base Case:**

Let $\gamma \in \mathbb{N}^1$. Since $X \neq \emptyset$, there exists $x_0 \in X$. Since $(X, \mathcal{T}_d)$ is Hausdorff, $\{x_0\}$ is closed. Define $X_0 \in \mathcal{T}_d$ by $X_0 := X \setminus \{x_0\}$. Since x_0 is a $\mathcal{T}_d$ -condensation point of X , Exercise 6.8.0.8 implies that X_0 is not compact. Exercise 6.14.1.2 then implies that there exists $(V_n)_{n \in \mathbb{N}} \subset \mathcal{T}_d$ such that

- for each $n \in \mathbb{N}$, $V_n \neq \emptyset$, V_n is $\mathcal{T}_d$ -closed and $\text{diam } V_n \leq 1$,
- $(V_n)_{n \in \mathbb{N}}$ is disjoint,
- $X_0 = \bigcup_{n \in \mathbb{N}} V_n$

We define $C_\gamma := V_{\gamma_1}$. Let $x \in C_\gamma$. Since $C_\gamma \in \mathcal{T}_d$, we have that $C_\gamma \in \mathcal{N}(x, \mathcal{T}_d)$. By assumption, x is a $\mathcal{T}_d$ -condensation point of X . Exercise 4.1.0.49 then implies that x is a $\mathcal{T}_d$ -condensation point of C_γ and Exercise 4.5.1.9 implies that x is a $\mathcal{T}_d \cap C_\gamma$ -condensation point of C_γ . Since $x \in C_\gamma$ is arbitrary, we have that for each $x \in C_\gamma$, x is a $\mathcal{T}_d \cap C_\gamma$ -condensation point of C_γ .

• **Induction Step:**

Let $n \geq 2$ and $\gamma \in \mathbb{N}^n$. Define $\tilde{\gamma} \in \mathbb{N}^{n-1}$ by $\tilde{\gamma} := \pi_{[n-1]}^{[n]}(\gamma)$. Suppose that

- $C_{\tilde{\gamma}} \in \mathcal{T}_d$, $C_{\tilde{\gamma}}$ is $\mathcal{T}_d$ -closed and $C_{\tilde{\gamma}} \neq \emptyset$,
- for each $x \in C_{\tilde{\gamma}}$, x is a $\mathcal{T}_d \cap C_{\tilde{\gamma}}$ -condensation point of $C_{\tilde{\gamma}}$

Define $d_{\tilde{\gamma}} : C_{\tilde{\gamma}}^2 \rightarrow [0, \infty)$ by $d_{\tilde{\gamma}} := d|_{C_{\tilde{\gamma}}}$. Since (X, d) is separable, Exercise 17.3.0.5 [cite ex here](#) implies that $(C_{\tilde{\gamma}}, d_{\tilde{\gamma}})$ is separable. Since $(X, \mathcal{T}_d)$ is zero-dimensional, Exercise 4.18.1.1 implies that $(C_{\tilde{\gamma}}, \mathcal{T}_d \cap C_{\tilde{\gamma}})$ is zero-dimensional. Since $C_{\tilde{\gamma}} \neq \emptyset$, there exists $x_{\tilde{\gamma}} \in C_{\tilde{\gamma}}$. Since $(C_{\tilde{\gamma}}, d_{\tilde{\gamma}})$ is Hausdorff, $\{x_{\tilde{\gamma}}\}$ is $\mathcal{T}_d \cap C_{\tilde{\gamma}}$ -closed. Define $C_{\tilde{\gamma},0} \in \mathcal{T}_d \cap C_{\tilde{\gamma}}$ by $C_{\tilde{\gamma},0} := C_{\tilde{\gamma}} \setminus \{x_{\tilde{\gamma}}\}$. By assumption, $x_{\tilde{\gamma}}$ is a $\mathcal{T}_d \cap C_{\tilde{\gamma}}$ -condensation point of $C_{\tilde{\gamma}}$. Exercise 6.8.0.8 then implies that $C_{\tilde{\gamma},0}$ is not compact. Exercise 6.14.1.2 implies that there exists $(V_n^{\tilde{\gamma}})_{n \in \mathbb{N}} \subset \mathcal{T}_d \cap C_{\tilde{\gamma}}$ such that

- for each $n \in \mathbb{N}$, $V_n^{\tilde{\gamma}} \neq \emptyset$, $V_n^{\tilde{\gamma}}$ is $\mathcal{T}_d \cap C_{\tilde{\gamma}}$ -closed and $\text{diam } V_n^{\tilde{\gamma}} \leq 1/n$,
- $(V_n^{\tilde{\gamma}})_{n \in \mathbb{N}}$ is disjoint,
- $C_{\tilde{\gamma},0} = \bigcup_{n \in \mathbb{N}} V_n^{\tilde{\gamma}}$.

We define $C_{\gamma} := V_n^{\tilde{\gamma}}$. Since $C_{\gamma} \in \mathcal{T}_d \cap C_{\tilde{\gamma}}$ and $C_{\tilde{\gamma}} \in \mathcal{T}_d$, we have that $C_{\gamma} \in \mathcal{T}_d$. Since C_{γ} is $\mathcal{T}_d \cap C_{\tilde{\gamma}}$ -closed and $C_{\tilde{\gamma}}$ is $\mathcal{T}_d$ -closed, Exercise 17.3.0.5 [cite ex](#) implies that C_{γ} is $\mathcal{T}_d$ -closed. Let $x \in C_{\gamma}$. Since $C_{\gamma} \in \mathcal{T}_d$, we have that $C_{\gamma} \in \mathcal{N}(x, \mathcal{T}_d)$. By assumption, x is a $\mathcal{T}_d$ -condensation point of X . Exercise 4.1.0.49 then implies that x is a $\mathcal{T}_d$ -condensation point of C_{γ} and Exercise 4.5.1.9 implies that x is a $\mathcal{T}_d \cap C_{\gamma}$ -condensation point of C_{γ} . Since $x \in C_{\gamma}$ is arbitrary, we have that for each $x \in C_{\gamma}$, x is a $\mathcal{T}_d \cap C_{\gamma}$ -condensation point of C_{γ} .

Thus there exist $(C_{\gamma})_{\gamma \in \Gamma} \subset \mathcal{T}_d$ and $(x_{\gamma})_{\gamma \in \Gamma} \in \prod_{\gamma \in \Gamma} C_{\gamma}$ such that

1. • there exists $x_0 \in X$ such that $X \setminus \{x_0\} = \bigcup_{|\gamma|=1} C_{\gamma}$,
 • $(C_{\gamma})_{\gamma \in \mathbb{N}}$ is disjoint
2. for each $\gamma \in \Gamma$,
 - (a) C_{γ} is $\mathcal{T}_d$ -closed and $C_{\gamma} \neq \emptyset$,
 - (b) $\text{diam } C_{\gamma} \leq |\gamma|^{-1}$,
 - (c) • $C_{\gamma} \setminus \{x_{\gamma}\} = \bigcup_{\gamma' \in (\pi_{[n]}^{[n+1]})^{-1}(\{\gamma\})} C_{\gamma'}$
 • $(C_{\gamma'})_{\gamma' \in (\pi_{[n]}^{[n+1]})^{-1}(\{\gamma\})}$ is disjoint
 - (d) for each $x \in C_{\gamma}$, x is a $\mathcal{T}_d \cap C_{\gamma}$ -condensation point of C_{γ} .

□

6.14.2 Polish Spaces

Exercise 6.14.2.1. Let X be a Polish space. Suppose that X is zero-dimensional. Define $X_0 \subset X$ by

$$X_0 := \{x \in X : x \text{ is a condensation point of } X\}.$$

Then X_0 is Polish, X_0 is zero dimensional and for each $x \in X_0$, x is a condensation point of X_0 .

Hint: Exercise 4.13.2.10 and Exercise 4.1.0.48

Proof. Since X is Polish, X is second-countable. Exercise 4.13.2.10 and Exercise 4.1.0.48 imply that X_0 is closed and X_0^c is countable. Since X_0 is closed and X is Polish, Exercise 6.12.1.2 implies that X_0 is Polish. Since $X_0 \subset X$ and X is zero-dimensional, Exercise 4.18.1.1 implies that X_0 is zero-dimensional. Let $x \in X_0$ and $U \in \mathcal{N}(x)$. By definition, x is a condensation point of X . Thus $U \cap X$ is uncountable. For the sake of contradiction, suppose that $U \cap X_0$ is countable. We note that

$$\begin{aligned} U \cap X &= [(U \cap X) \cap X_0] \cup [(U \cap X) \cap X_0^c] \\ &= (U \cap X_0) \cup (U \cap X \cap X_0^c). \end{aligned}$$

Since $U \cap X \cap X_0^c \subset X_0^c$ and X_0^c is countable, we have that $U \cap X \cap X_0^c$ is countable. Since $U \cap X_0$ is countable and $U \cap X = (U \cap X_0) \cup (U \cap X \cap X_0^c)$, we have that $U \cap X$ is countable, which is a contradiction. Hence $U \cap X_0$ is uncountable. Since $U \in \mathcal{N}(x)$ is arbitrary, we have that for each $U \in \mathcal{N}(x)$, $U \cap X_0$ is uncountable. Hence x is a condensation point of X_0 . Since $x \in X_0$ is arbitrary, we have that for each $x \in X_0$, x is a condensation point of X_0 . $\square$

6.14.3 Ultrametric Spaces

Exercise 6.14.3.1. Let (X, d) be an ultrametric space. Then $(X, \mathcal{T}_d)$ is zero-dimensional.

Proof. Exercise 6.1.2.6 implies that $\mathcal{B}_d$ is a basis for $\mathcal{T}_d$. An exercise in the section on ultrametric spaces implies that for each $B \in \mathcal{B}_d$, B is closed. Hence $(X, \mathcal{T}_d)$ is zero-dimensional. $\square$

6.14.4 Cantor Space

Definition 6.14.4.1. We define the **Cantor space**, denoted $\mathcal{C}$, by $\mathcal{C} := \{0, 1\}^{\mathbb{N}}$.

Exercise 6.14.4.2. We have that $\mathcal{C}$ is a zero-dimensional Polish space.

Proof. Clear by make finite set examples in sections on polish space section and discrete topology in zero-dimensional space section $\square$

Exercise 6.14.4.3. For each $x \in [0, 1]$, there exists $(x_n)_{n \in \mathbb{N}} \in \mathcal{C}$ such that $x = \sum_{j \in \mathbb{N}} x_j 2^{-j}$.

Hint: Set $x_1 = \begin{cases} 0, & x < 1/2 \\ 1, & x \geq 1/2 \end{cases}$ and proceed inductively.

Proof. Let $x \in [0, 1]$. Set

$$x_1 = \begin{cases} 0, & x < 1/2 \\ 1, & x \geq 1/2 \end{cases}$$

and for $j \geq 2$, set

$$x_j = \begin{cases} 0, & x - \sum_{k=1}^{j-1} x_k 2^{-k} < 2^{-j} \\ 1, & x - \sum_{k=1}^{j-1} x_k 2^{-k} \geq 2^{-j} \end{cases}$$

Note that for each $j \in \mathbb{N}$, $x - \sum_{k=1}^j x_k 2^{-k} \in [0, 2^{-j}]$. Hence $x = \sum_{j \in \mathbb{N}} x_j 2^{-j}$ $\square$

Exercise 6.14.4.4. Let $(x_n)_{n \in \mathbb{N}}, (y_n)_{n \in \mathbb{N}} \subset \mathcal{C}$. Suppose that $(x_n)_{n \in \mathbb{N}} \neq (y_n)_{n \in \mathbb{N}}$. Set $N = \min\{j \in \mathbb{N} : x_j \neq y_j\}$. Suppose that $x_N = 0$ and $y_N = 1$. Then

$$\sum_{j \in \mathbb{N}} x_j 2^{-j} = \sum_{j \in \mathbb{N}} y_j 2^{-j}$$

iff for each $j \in \mathbb{N}$, $j > N$ implies that $x_j = 1$ and $y_j = 0$.

Proof. Suppose that

$$\sum_{j \in \mathbb{N}} x_j 2^{-j} = \sum_{j \in \mathbb{N}} y_j 2^{-j}$$

By definition of N , for each $j \in \mathbb{N}$, $j < N$ implies that $x_j = y_j$. Hence

$$\sum_{j=N}^{\infty} x_j 2^{-j} = \sum_{j=N}^{\infty} y_j 2^{-j}$$

Since $x_N = 0$ and $y_N = 1$, we have that

$$\sum_{j=N+1}^{\infty} x_j 2^{-j} = 2^{-N} + \sum_{j=N+1}^{\infty} y_j 2^{-j}$$

Thus $2^{-N} = \sum_{j=N+1}^{\infty} (x_j - y_j)2^{-j}$. For the sake of contradiction, suppose that there exists $m > N$ and $x_m \neq 1$ or $y_m \neq 0$. Then

$$\begin{aligned} 2^{-N} &= \sum_{j=N+1}^{\infty} (x_j - y_j)2^{-j} \\ &< \sum_{j=N+1}^{\infty} (1 - 0)2^{-j} \\ &= 2^{-N} \end{aligned}$$

which is a contradiction. Hence for each $m \in \mathbb{N}$, $m > N$ implies that $x_m = 1$ and $y_m = 0$.

Conversely, suppose that for each $j \in \mathbb{N}$, $j > N$ implies that $x_j = 1$ and $y_j = 0$. Then

$$\begin{aligned} \sum_{j \in \mathbb{N}} x_j 2^{-j} &= \sum_{j=1}^{N-1} x_j + \sum_{j \geq N+1} 2^{-j} \\ &= \sum_{j=1}^{N-1} x_j + 2^{-N} \sum_{j \in \mathbb{N}} 2^{-j} \\ &= \sum_{j=1}^{N-1} y_j + 2^{-N} \\ &= \sum_{j=1}^N y_j 2^{-j} \\ &= \sum_{j \in \mathbb{N}} y_j 2^{-j} \end{aligned}$$

□

Definition 6.14.4.5.

- We define $Z \subset \mathcal{C}$ by

$$Z = \left\{ (x_n)_{n \in \mathbb{N}} \in \mathcal{C} : \#\{n \in \mathbb{N} : x_n = 0\} = \infty \right\} \cup \{(1, 1, 1, \dots)\}$$

- We define $\phi : Z \rightarrow [0, 1]$ by

$$\phi(x) = \sum_{n \in \mathbb{N}} x_n 2^{-n}$$

Exercise 6.14.4.6. We have that Z^c is an F_σ -set, Z is a G_δ -set and Z is a zero-dimensional Polish space.

Hint: Z^c is countable

Proof. Since $\mathcal{C}$ is zero-dimensional and $Z \subset \mathcal{C}$, Exercise 4.18.1.1 implies that Z is zero-dimensional. Since $Z^c \subset \left\{ (x_n)_{n \in \mathbb{N}} \in \mathcal{C} : \#\{n \in \mathbb{N} : x_n = 0\} < \infty \right\}$, we have that Z^c is countable. Exercise 4.12.1.4 then implies that Z^c is an F_σ -set. Thus Z is a G_δ -set. Exercise 6.12.1.8 then implies that Z is a Polish space. □

Exercise 6.14.4.7. We have that $\phi : Z \rightarrow [0, 1]$ is a continuous bijection.

Hint: Weierstrass M-test

Proof.

- Let $x \in [0, 1]$. Then Exercise 6.14.4.3 implies that there exists $(x_n)_{n \in \mathbb{N}} \in \mathcal{C}$ such that $x = \sum_{n \in \mathbb{N}} x_n 2^{-n}$. If for each $n \in \mathbb{N}$, $x_n = 1$, then $(x_n)_{n \in \mathbb{N}} \in Z$. Suppose that there exists $n \in \mathbb{N}$ such that $x_n = 0$. If $\#\{n \in \mathbb{N} : x_n = 0\} = \infty$, then $(x_n)_{n \in \mathbb{N}} \in Z$. Suppose that $\#\{n \in \mathbb{N} : x_n = 0\} < \infty$. Set $N = \max\{n \in \mathbb{N} : x_n = 0\}$. Define $(y_n)_{n \in \mathbb{N}} \in Z$ by

$$y_n = \begin{cases} x_n, & n \in \{1, \dots, N-1\} \\ 1, & n = N \\ 0, & n > N \end{cases}$$

Then Exercise 6.14.4.4 implies that $\phi((y_n)_{n \in \mathbb{N}}) = x$. Since $x \in [0, 1]$ is arbitrary, ϕ is surjective.

Let $(x_n)_{n \in \mathbb{N}}, (y_n)_{n \in \mathbb{N}} \in Z$. Suppose that $(x_n)_{n \in \mathbb{N}} \neq (y_n)_{n \in \mathbb{N}}$. If $\phi((x_n)_{n \in \mathbb{N}}) = \phi((y_n)_{n \in \mathbb{N}})$, then Exercise 6.14.4.4 implies that $(x_n)_{n \in \mathbb{N}} \notin Z$ or $(y_n)_{n \in \mathbb{N}} \notin Z$, which is a contradiction. Hence $\phi((x_n)_{n \in \mathbb{N}}) \neq \phi((y_n)_{n \in \mathbb{N}})$. Since $(x_n)_{n \in \mathbb{N}}, (y_n)_{n \in \mathbb{N}} \in Z$ are arbitrary, ϕ is injective. So ϕ is a bijection.

- For $n \in \mathbb{N}$, define $\phi_n : Z \rightarrow [0, 1]$ by $\phi_n := 2^{-n} \pi_n|_Z$. Then $\phi = \sum_{n \in \mathbb{N}} \phi_n$ and $\|\phi_n\|_\infty = 2^{-n}$. The Weierstrass M-test implies that ϕ is continuous.

□

Exercise 6.14.4.8. There exists $Z \subset \mathcal{C}$ and $\phi : Z \rightarrow [0, 1]$ such that Z is a zero-dimensional Polish space and ϕ is a continuous bijection.

Proof. Clear by Exercise 6.14.4.6 and Exercise 6.14.4.7. □

Definition 6.14.4.9. We define $a : \mathbb{N}^2 \rightarrow \mathbb{N}$ by $a(k, n) = 2^{k-1}(2n-1)$.

Exercise 6.14.4.10. We have that $a : \mathbb{N}^2 \rightarrow \mathbb{N}$ is a bijection.

Proof.

• **Injectivity**

Let $(k, n), (k', n') \in \mathbb{N}^2$. Suppose that $a(k, n) = a(k', n')$. Then $2^{k-1}(2n-1) = 2^{k'-1}(2n'-1)$. Set $l = 2n-1$ and $l' = 2n'-1$. Then $l = 2^{k'-k}l'$. Since $l, l' \in \mathbb{N}$ and $l' \equiv 1 \pmod{2}$, we have that $k' - k \geq 0$. Since $l \equiv 1 \pmod{2}$, we have that $k' - k = 0$. Therefore $l = l'$ which implies that $n = n'$. Therefore $(k, n) = (k', n')$. Since $(k, n), (k', n') \in \mathbb{N}^2$ are arbitrary, a is injective.

• **Surjectivity:**

Let $j \in \mathbb{N}$. Define $k_0, l_0 \in \mathbb{N}$ by $k_0 = \max\{k \in \mathbb{N} : \gcd(j, 2^{k-1}) = 2^{k-1}\}$ and $l_0 = j/2^{k_0-1}$. Since $l_0 \equiv 1 \pmod{2}$ there exists $n_0 \in \mathbb{N}$ such that $l_0 = 2n_0 - 1$. Thus, $a(k_0, n_0) = j$. Since $j \in \mathbb{N}$ is arbitrary, a is surjective.

□

Definition 6.14.4.11. We define $\eta_0, \eta_1 : \mathcal{C} \rightarrow \mathcal{C}$ and $H : \mathcal{C} \rightarrow \mathcal{C}^{\mathbb{N}}$ by

- $\eta_0(x) = (x_{2n})_{n \in \mathbb{N}}$
- $\eta_1(x) = (x_{2n-1})_{n \in \mathbb{N}}$
- for $k \in \mathbb{N}$, $[H(x)]_k = \eta_1 \circ (\eta_0)^{k-1}(x)$.

Exercise 6.14.4.12.

1. For each $x \in \mathcal{C}$, $[H(x)]_k = (x_{2^{k-1}(2n-1)})_{n \in \mathbb{N}}$
2. $H : \mathcal{C} \rightarrow \mathcal{C}^{\mathbb{N}}$ is a bijection
3. $H : \mathcal{C} \rightarrow \mathcal{C}^{\mathbb{N}}$ is a homeomorphism

Proof.

1. Let $x \in \mathcal{C}$ and $k \in \mathbb{N}$. Define $y \in \mathcal{C}$ by $y = (\eta_0)^{k-1}(x)$. Clearly $y = (x_{2^{k-1}n})_{n \in \mathbb{N}}$. Then

$$\begin{aligned} [H(x)]_k &= \eta_1 \circ (\eta_0)^{k-1}(x) \\ &= \eta_1(y) \\ &= (y_{2n-1})_{n \in \mathbb{N}} \\ &= (x_{2^{k-1}(2n-1)})_{n \in \mathbb{N}} \end{aligned}$$

2. • **Injectivity:**

Let $x, y \in \mathcal{C}$. Suppose that $H(x) = H(y)$. Let $j \in \mathbb{N}$. Define $(k, n) \in \mathbb{N}^2$ by $(k, n) = a^{-1}(j)$. Then $j = 2^{k-1}(2n-1)$ and

$$\begin{aligned} x_j &= x_{2^{k-1}(2n-1)} \\ &= ([H(x)]_k)_n \\ &= ([H(y)]_k)_n \\ &= y_{2^{k-1}(2n-1)} \\ &= y_j \end{aligned}$$

Since $j \in \mathbb{N}$ is arbitrary, $x = y$. Since $x, y \in \mathcal{C}$ are arbitrary, H is injective.

- **Surjectivity:**

Let $X \in \mathcal{C}^{\mathbb{N}}$. Define $(k_j, n_j)_{j \in \mathbb{N}} \in (\mathbb{N}^2)^{\mathbb{N}}$ and $x \in \mathcal{C}$ by $(k_j, n_j) = a^{-1}(j)$ and $x_j = (X_{k_j})_{n_j}$. Let $(k, n) \in \mathbb{N}$. Define $j \in \mathbb{N}$ by $j = a(k, n)$. Then

$$\begin{aligned} ([H(x)]_k)_n &= x_{2^{k-1}(2n-1)} \\ &= x_{a(k, n)} \\ &= x_j \\ &= (X_k)_n \end{aligned}$$

Hence $X = H(x)$. Since $X \in \mathcal{C}^{\mathbb{N}}$ is arbitrary, H is surjective.

Therefore H is a bijection.

3. Let $(x_\alpha)_{\alpha \in A} \subset \mathcal{C}$ be a net and $x \in \mathcal{C}$. Suppose that $x_\alpha \rightarrow x$. Then for each $n \in \mathbb{N}$,

$$\begin{aligned} (x_\alpha)_n &= \pi_n(x_\alpha) \\ &\rightarrow \pi_n(x) \\ &= x_n \end{aligned}$$

Let $k \in \mathbb{N}$. Then for each $n \in \mathbb{N}$,

$$\begin{aligned} \pi_n(\pi_k(H(x_\alpha))) &= (x_\alpha)_{2^{k-1}(2n-1)} \\ &\rightarrow x_{2^{k-1}(2n-1)} \\ &= \pi_n(\pi_k(H(x))) \end{aligned}$$

Thus

$$\pi_k(H(x_\alpha)) \rightarrow \pi_k(H(x))$$

Since $k \in \mathbb{N}$ is arbitrary, we have that

$$H(x_\alpha) \rightarrow H(x)$$

Hence H is continuous.

Conversely, let $(X_\alpha)_{\alpha \in A} \subset \mathcal{C}^{\mathbb{N}}$ be a net and $X \in \mathcal{C}^{\mathbb{N}}$. Suppose that $X_\alpha \rightarrow X$. Then for each $k, n \in \mathbb{N}$,

$$\begin{aligned} ([X_\alpha]_k)_n &= \pi_n(\pi_k(X_\alpha)) \\ &\rightarrow \pi_n(\pi_k(X)) \\ &= (X_k)_n \end{aligned}$$

Let $j \in \mathbb{N}$. Define $(k, n) \in \mathbb{N}^2$ by $(k, n) = a^{-1}(j)$. Then

$$\begin{aligned}\pi_j(H^{-1}(X_\alpha)) &= [H^{-1}(X_\alpha)]_j \\ &= ([X_\alpha]_k)_n \\ &\rightarrow (X_k)_n \\ &= [H^{-1}(X)]_j \\ &= \pi_j(H^{-1}(X))\end{aligned}$$

Since $j \in \mathbb{N}$ is arbitrary,

$$H^{-1}(X_\alpha) \rightarrow H^{-1}(X)$$

Hence H^{-1} is continuous. Thus H is a homeomorphism.

□

Chapter 7

Analysis on $(C(X), \|\cdot\|_\infty)$

7.1 LEFTOVERS

Definition 7.1.0.1. Let X be a metric space. Then X is said to be **separable** if there exists $D \subset X$ such that D is countable and for each $x \in X$ and $\epsilon > 0$, there exists $y \in D$ such that $d(x, y) < \epsilon$.

Exercise 7.1.0.2. Let X be a metric space. If X is separable, then for each $A \subset X$, if A is open, then

1. there there exist $(x_n)_{n \in \mathbb{N}} \subset X$ and $(r_n)_{n \in \mathbb{N}} \subset (0, \infty)$ such that

$$A = \bigcup_{n \in \mathbb{N}} B(x_n, r_n)$$

i.e. A is a countable union of open balls

2. there there exist $(x_n)_{n \in \mathbb{N}} \subset X$ and $(r_n)_{n \in \mathbb{N}} \subset (0, \infty)$ such that

$$A = \bigcup_{n \in \mathbb{N}} \bar{B}(x_n, r_n)$$

i.e. A is a countable union of closed balls.

Proof. Suppose that X is separable. Then there exists $(x_n)_{n \in \mathbb{N}} \subset X$ such that for each $x \in X$ and $\epsilon > 0$, there exists $N \in \mathbb{N}$ such that $d(x, x_N) < \epsilon$. Let $A \subset X$. Suppose that A is open.

1. Set

$$\mathcal{B} = \{B(x_n, r) : r \in \mathbb{Q} \text{ and } B(x_n, r) \subset A\}$$

Note that $\mathcal{B}$ is countable. Let $x \in A$. Since A is open, there exists $s \in \mathbb{R}$ such that $B(x, s) \subset A$. Then there exists $r \in \mathbb{Q} \cap (0, s)$. Choose $N \in \mathbb{N}$ such that $d(x, x_N) < r/2$. Let $y \in B(x_N, r/2)$, then

$$\begin{aligned} d(x, y) &\leq d(x, x_N) + d(x_N, y) \\ &< r/2 + r/2 \\ &= r \end{aligned}$$

Therefore

$$\begin{aligned} x &\in B(x_N, r/2) \\ &\subset B(x, r) \\ &\subset A \end{aligned}$$

Hence $B(x_N, r/2) \in \mathcal{B}$ and $x \in \bigcup_{B \in \mathcal{B}} B$. Since $x \in A$ is arbitrary, $A \subset \bigcup_{B \in \mathcal{B}} B$.

2. Similar, but take $r/4$ instead of $r/2$.

□

Definition 7.1.0.3. Let X be a set, $d_1, d_2 : X \times X \rightarrow [0, \infty)$ metrics on X . Then d_1 and d_2 are said to be **equivalent** if there exist $A, B > 0$ such that

$$Ad_1 \leq d_2 \leq Bd_1$$

Definition 7.1.0.4. Let (X, d_X) and (Y, d_Y) be metric spaces and $f : X \rightarrow Y$. Then f is said to be **Lipchitz** if there exists $K \geq 0$ such that for each $a, b \in X$,

$$d_Y(f(a), f(b)) \leq Kd_X(a, b)$$

Exercise 7.1.0.5. Let (X, d_X) and (Y, d_Y) be metric spaces and $f : X \rightarrow Y$. If f is Lipchitz, then f is uniformly continuous.

Proof. By definition, there exists $K \geq 0$ such that for each $a, b \in X$,

$$d_Y(f(a), f(b)) \leq K d_X(a, b)$$

Let $\epsilon > 0$. Choose $\delta = \epsilon/(K+1)$. Let $a, b \in X$. Suppose that $d_X(a, b) < \delta$. Then

$$\begin{aligned} d_Y(f(a), f(b)) &\leq K d_X(a, b) \\ &< K\delta \\ &= K \frac{\epsilon}{K+1} \\ &< \epsilon \end{aligned}$$

□

Definition 7.1.0.6. Let (X, d_X) and (Y, d_Y) be metric spaces and $f : X \rightarrow Y$.

- $x_0 \in X$. Then f is said to be **locally Lipchitz at** x_0 if there exists $U \in \mathcal{N}(x_0)$ such that f is Lipschitz on U .
- Then f is said to be **locally Lipchitz** if for each $x_0 \in X$, f is locally Lipschitz at x_0 .

Definition 7.1.0.7. Let X, Y be metric spaces and $T : X \rightarrow Y$. Then T is said to be an **isometry** if for each $x_1, x_2 \in X$, $d(Tx_1, Tx_2) = d(x_1, x_2)$.

Exercise 7.1.0.8. Let X, Y be metric spaces and $T : X \rightarrow Y$ an isometry. Then T is injective.

Proof. Let $x_1, x_2 \in X$. Suppose that $Tx_1 = Tx_2$. Then $0 = d(Tx_1, Tx_2) = d(x_1, x_2)$. So $x_1 = x_2$. Hence T is injective. □

Note 7.1.0.9. Let X, Y be metric spaces and $T : X \rightarrow Y$ an isometry. Then T is clearly continuous. If T is surjective, then T^{-1} is an isometry and therefore continuous. Hence T is a homeomorphism.

Definition 7.1.0.10. Let (X, d) be a metric space. Then (X, d) is said to be a **Polish space** if (X, d) is complete and separable.

Exercise 7.1.0.11. Let (X, d) be a compact metric space, $E \subset X$ closed, $U \subset X$ open. Suppose that $E \subset U$. Then there exists $\delta > 0$ such that for each $x \in E$, $B(x, \delta) \subset U$.

Proof. Since X is compact, E and U^c are compact. Then there exist $x_0 \in E$ and $y_0 \in U^c$ such that $d(E, U^c) = d(x_0, y_0)$. Since $E \cap U^c = \emptyset$, $x_0 \neq y_0$ and $d(E, U^c) > 0$. Put $\epsilon = d(E, U^c)$ and $\delta = \frac{\epsilon}{2}$. Let $x \in E$, $w \in B(x, \delta)$ and $y \in U^c$. Then

$$\begin{aligned} d(y, w) &\geq d(y, x) - d(x, w) \\ &> \epsilon - \delta \\ &= \epsilon - \frac{\epsilon}{2} \\ &= \frac{\epsilon}{2} \\ &> 0 \end{aligned}$$

So $y \neq w$. Since $y \in U^c$ and $w \in B(x, \delta) \subset U$, $B(x, \delta) \subset U$. □

Definition 7.1.0.12. Let S be a set, (X, d) a metric space and $B(S, X) = \{f : S \rightarrow X : f \text{ is bounded}\}$. We define the **supremum metric**, denoted $d_u : B(S, X) \times B(S, X) \rightarrow [0, \infty)$, by

$$d_u(f, g) = \sup_{x \in X} d(f(x), g(x))$$

Exercise 7.1.0.13. Let X be a set, (Y, d_Y) , (Z, d_Z) metric spaces, $(f_n)_{n \in \mathbb{N}} \subset B(X, Y)$, $f \in B(X, Y)$ and $g \in C(Y, Z)$. Suppose that g is uniformly continuous. If $f_n \xrightarrow{u} f$, then $g \circ f_n \xrightarrow{u} g \circ f$.

Proof. Suppose that $f_n \xrightarrow{u} f$. Let $\epsilon > 0$. Uniform continuity of g implies that there exists $\delta > 0$ such that for each $y_1, y_2 \in Y$, $d_Y(y_1, y_2) < \delta$ implies that $d_Z(g(y_1), g(y_2)) < \epsilon/2$. Uniform convergence implies that there exists $N \in \mathbb{N}$ such that for each $n \in \mathbb{N}$, $n \geq N$ implies that $d_u(f_n, f) < \delta/2$. Let $n \in \mathbb{N}$. Suppose that $n \geq N$. Let $x \in X$. Then $d_Y(f_n(x), f(x)) < \delta$. This implies that $d_Z(g(f_n(x)), g(f(x))) < \epsilon/2$. Hence $\sup_{x \in X} d_Z(g \circ f_n(x), g \circ f(x)) \leq \epsilon/2$. Thus $d_u(g \circ f_n, g \circ f) < \epsilon$. So $g \circ f_n \xrightarrow{u} g \circ f$. □

Definition 7.1.0.14. Let (X, d) be a metric space. Define

1. $\text{Aut}(X) = \{\sigma : X \rightarrow X : \sigma \text{ is a homeomorphism}\}$
2. $\text{Aut}(X, d) = \{\sigma : X \rightarrow X : \sigma \text{ is an isometric isomorphism}\}$

Exercise 7.1.0.15. Let (X, d) be a compact metric space, $E \subset X$ closed, $U \subset X$ open. Suppose that $E \subset U$. Let $(f_n)_{n \in \mathbb{N}} \in \text{Aut}(X)$, $f \in \text{Aut}(X)$. Suppose that $f_n \xrightarrow{u} f$. Then there exists $N \in \mathbb{N}$ such that for each $n \geq N$, $f(E) \subset f_n(U)$.

Proof. Since f is a homeomorphism, E is closed and U is open, $f(E)$ is compact and $f(U)$ is open and $f(E) \subset f(U)$. Then $d(f(E), f(U^c)) > 0$. Put $\epsilon = d(f(E), f(U^c))$. Choose $\delta = \epsilon/2$. Then there exists $N \in \mathbb{N}$ such that for each $n \in \mathbb{N}$, $n \geq N$ implies that $\sup_{z \in X} d(f(z), f_n(z)) < \delta$. Let $n \geq N$, $x \in E$ and $w \in B(f(x), \delta)$. For the sake of contradiction, suppose that $w \in f_n(U^c)$. Then there exist $p \in U^c$ such that $w = f_n(p)$. Put $z = f(p) \in f(U^c)$. Then

$$\begin{aligned} \epsilon &\leq d(f(x), z) \\ &\leq d(f(x), w) + d(w, z) \\ &= d(f(x), w) + d(f_n(p), f(p)) \\ &< \delta + \delta \\ &= \epsilon \end{aligned}$$

which is a contradiction. So $w \in f_n(U)$. Hence $B(f(x), \delta) \subset f_n(U)$

□

Part II

Topological Vector Spaces

Chapter 8

Topological Vector Spaces

8.1 Introduction

8.1.1 Topology

Definition 8.1.1.1. Let X be a vector space and $\mathcal{T}$ a topology on X . Then X is said to be a **topological vector space** if

1. addition $X \times X \rightarrow X$ is continuous
2. scalar multiplication $\mathbb{C} \times X \rightarrow X$ is continuous

Note 8.1.1.2. We usually suppress the topology $\mathcal{T}$.

Exercise 8.1.1.3. Let X be a vector space and $\mathcal{T}$ a topology on X . Then $(X, \mathcal{T})$ is a topological vector space iff for each $(x_\alpha)_{\alpha \in A}$, $(y_\alpha)_{\alpha \in A} \subset X$, $(\lambda_\alpha)_{\alpha \in A} \subset \mathbb{K}$ nets, $x, y \in X$ and $\lambda \in \mathbb{K}$, $x_\alpha + \lambda_\alpha y_\alpha \rightarrow x + \lambda y$.

Proof. Clear by Exercise 4.3.2.13. □

Exercise 8.1.1.4. Let X be a topological vector space, $y \in X$ and $\lambda \in \mathbb{C}^\times$. Define $f, g : X \rightarrow X$ by $f(x) = x + y$ and $g(x) = \lambda x$. Then f and g are homeomorphisms.

Proof. Since X is a topological vector space, f and g are continuous. Clearly f and g are bijections with $f^{-1}(x) = x - y$ and $g^{-1}(x) = \lambda^{-1}x$. Again, since X is a topological vector space, f^{-1} and g^{-1} are continuous. □

Exercise 8.1.1.5. Let X be a topological vector space. Then X is T_1 iff $\{0\}$ is closed.

Proof. Clear by Exercise 8.1.1.4 and Exercise 4.12.1.3. □

Exercise 8.1.1.6. We have that $(\mathbb{C}, \mathcal{T}_{\mathbb{C}})$ is a topological vector space.

Exercise 8.1.1.7. Let X be a topological vector space, $x, y \in X$ and $U \in \mathcal{N}(x)$. If U is open, then there exists $r > 0$ such that for each $t \in \mathbb{R}$, $|t| \leq r$ implies that $x + ty \in U$.

Proof. Suppose that U is open. For the sake of contradiction, suppose that for each $r > 0$, there exists $t \in \mathbb{R}$ such that $t \leq r$ and $x + ty \notin U$. Then for each $n \in \mathbb{N}$, there exists $t_n \in \mathbb{R}$ such that $|t_n| \leq 1/n$ and $x + t_n y \notin U^c$. Since $t_n \rightarrow 0$,

$$\begin{aligned} x + t_n y &\rightarrow x + 0y \\ &= x \end{aligned}$$

Since U^c is closed, $x \in U^c$. This is a contradiction. Hence there exists $r > 0$ such that for each $t \in \mathbb{R}$, $|t| \leq r$ implies that $x + ty \in U$. □

Exercise 8.1.1.8. Let X be a topological vector space and $A, B \subset X$. If A is open, then $A + B$ is open.

Proof. Suppose that A is open. Then for each $b \in B$, $A + b$ is open. Since

$$A + B = \bigcup_{b \in B} A + b$$

we have that $A + B$ is open. □

Exercise 8.1.1.9. Let X be a topological vector space and $A, B \subset X$. Suppose that A is compact, B is closed and $A \cap B = \emptyset$. Then there exists $U \in \mathcal{N}(0)$ such that U is open and $(A + U) \cap B = \emptyset$.

Proof. Set $\Gamma = \{U \in \mathcal{N}(0) : U \text{ is open}\}$ and order Γ by reverse inclusion, so that Γ is a directed set. For the sake of contradiction, suppose that for each $U \in \Gamma$, $(A + U) \cap B \neq \emptyset$. Then for each $\gamma \in \Gamma$, there exist $a_\gamma \in A$ and $u_\gamma \in \gamma$ such that $a_\gamma + u_\gamma \in B$. Let $V \in \mathcal{N}(0)$. Since $\text{Int } V \in \Gamma$

$$\begin{aligned} u_{\text{Int } V} &\in \text{Int } V \\ &\subset V \end{aligned}$$

Since $V \in \mathcal{N}(0)$ is arbitrary, $u_\gamma \rightarrow 0$. Since A is compact, there exists $a \in A$ and a subnet $(a_{\gamma_\zeta})_{\zeta \in Z}$ of $(a_\gamma)_{\gamma \in \Gamma}$ such that $a_{\gamma_\zeta} \rightarrow a$. Then $a_{\gamma_\zeta} + u_{\gamma_\zeta} \rightarrow a$. Since $(a_{\gamma_\zeta} + u_{\gamma_\zeta})_{\zeta \in Z} \subset B$ and B is closed, we have that $a \in B$. This is a contradiction since $A \cap B = \emptyset$. So there exists $U \in \mathcal{N}(0)$ such that U is open and $(A + U) \cap B = \emptyset$. □

Exercise 8.1.1.10. Let X be a topological vector space and $U \in \mathcal{N}(0)$. If U is open, then there exists $V \in \mathcal{N}(0)$ such that V is open and $V + V \subset U$.

Proof. Suppose that U is open. Set $\Gamma = \{V \in \mathcal{N}(0) : V \text{ is open}\}$ and order Γ by reverse inclusion, so that Γ is a directed set. For the sake of contradiction, suppose that for each $V \in \mathcal{N}(0)$, if V is open, then $V + V \not\subset U$. Then for each $\gamma \in \Gamma$, there exists $x_\gamma, y_\gamma \in \gamma$ such that $x_\gamma + y_\gamma \in U^c$. Let $W \in \mathcal{N}(0)$. Set $\beta = \text{Int } W$. Then $\beta \in \Gamma$. Then for each $\gamma \geq \beta$,

$$\begin{aligned} x_\gamma, y_\gamma &\in \gamma \\ &\subset \beta \\ &\subset W \end{aligned}$$

So that $(x_\gamma)_{\gamma \in \Gamma}$ and $(y_\gamma)_{\gamma \in \Gamma}$ are eventually in W . Since $W \in \mathcal{N}(0)$ is arbitrary, $x_\gamma \rightarrow 0$ and $y_\gamma \rightarrow 0$. Therefore $x_\gamma + y_\gamma \rightarrow 0$. Since for each $\gamma \in \Gamma$, $x_\gamma + y_\gamma \in U^c$ and U^c is closed, $0 \in U^c$. This is a contradiction, so there exists $V \in \mathcal{N}(0)$ such that V is open and $V + V \subset U$. □

Exercise 8.1.1.11. Let X be a topological vector space. Then for each $A \in \text{Conv}(X)$, $\text{cl } A \in \text{Conv}(X)$.

Proof. Let $x, y \in \text{cl } A$ and $t \in [0, 1]$. Then there exist nets $(x_\alpha)_{\alpha \in \Gamma}, (y_\alpha)_{\alpha \in \Gamma} \subset A$ such that $x_\alpha \rightarrow x$ and $y_\alpha \rightarrow y$. Since A is convex, $(tx_\alpha + (1-t)y_\alpha)_{\alpha \in \Gamma} \subset A$. Since X is a topological vector space, $tx_\alpha + (1-t)y_\alpha \rightarrow tx + (1-t)y$. Thus $tx + (1-t)y \in \text{cl } A$. Since $x, y \in \text{cl } A$ and $t \in [0, 1]$ are arbitrary, we have that for each $x, y \in \text{cl } A$ and $t \in [0, 1]$, $tx + (1-t)y \in \text{cl } A$. Hence $\text{cl } A \in \text{Conv}(X)$. □

Exercise 8.1.1.12. Let X be a topological vector space and $E \in \mathcal{K}(X) \cap \text{Conv}(X)$. If E is metrizable, then

1. $\text{extrm } E$ is a G_δ -set of E
2. $\text{extrm } E \in \mathcal{B}(X)$.

Proof. Write $\mathcal{T}_E := \mathcal{T}_X \cap E$. Suppose that E is metrizable. Then there exists $d : E \times E \rightarrow [0, \infty)$ such that d is a metric on E and $\mathcal{T}_d = \mathcal{T}_E$. We note that since $E \in \mathcal{K}(X)$, $E \times E \in \mathcal{K}(X \times X)$.

1. • By **defn**, for each $x \in E$, $x \notin \text{extrm } E$ iff there exists $a, b \in E$ such that $a \neq b$ and $2^{-1}(a + b) = x$. Define $F \subset E$ by $F := \{2^{-1}(a + b) : a, b \in E \text{ and } d(a, b) > 0\}$. Then $E \setminus (\text{extrm } E) = F$. For each $n \in \mathbb{N}$, define $F_n \subset F$ by $F_n := \{2^{-1}(a + b) : a, b \in E \text{ and } d(a, b) > n^{-1}\}$. Then $F = \bigcup_{n \in \mathbb{N}} F_n$.

- Let $n \in \mathbb{N}$. Since d is continuous, $d^{-1}([n^{-1}, \infty))$ is closed in $E \times E$, we have that $d^{-1}([n^{-1}, \infty)) \in \mathcal{K}(E \times E)$. Since Define $T : E \times E \rightarrow E$ by $T(a, b) := 2^{-1}(a + b)$. Since **addition, scalar mult** are continuous, T is continuous. Hence

$$\begin{aligned} F_n &= T(d^{-1}([n^{-1}, \infty))) \\ &\in \mathcal{K}(E). \end{aligned}$$

Since E is Hausdorff, F_n is closed in E . Thus $E \setminus F_n$ is open in E . Since $n \in \mathbb{N}$ is arbitrary, we have that for each $n \in \mathbb{N}$, $E \setminus F_n$ is open in E . Since

$$\begin{aligned} \text{extrm } E &= E \setminus [E \setminus (\text{extrm } E)] \\ &= E \setminus F \\ &= E \setminus \left(\bigcup_{n \in \mathbb{N}} F_n \right) \\ &= \bigcap_{n \in \mathbb{N}} (E \setminus F_n), \end{aligned}$$

we have that $\text{extrm } E$ is a G_δ -set of E .

2. Since E is closed in X , and for each $n \in \mathbb{N}$, F_n is closed in E , we have that for each $n \in \mathbb{N}$, F_n is closed in X . Hence F is an F_σ -set of X and $F \in \mathcal{B}(X)$. Since $E \in \mathcal{B}(X)$, we have that

$$\begin{aligned} \text{extrm } E &= E \setminus F \\ &\in \mathcal{B}(X). \end{aligned}$$

□

8.1.2 Continuous Linear Functionals

Definition 8.1.2.1. Let $(X, \mathcal{T}_X), (Y, \mathcal{T}_Y)$ be topological vector spaces. We define $L((X, \mathcal{T}_X); (Y, \mathcal{T}_Y)) := \{T : X \rightarrow Y : T \text{ is } \mathbb{K}\text{-linear and } T \text{ is continuous}\}$.

Note 8.1.2.2. When the context is clear, we may write $L(X, Y)$ in place of $L((X, \mathcal{T}_X); (Y, \mathcal{T}_Y))$.

Exercise 8.1.2.3. Let X, Y be topological vector spaces. Then $L(X; Y)$ is a vector space.

Proof. **FINISH!!!**

□

Definition 8.1.2.4. Let $(X, \mathcal{T})$ be a topological vector space over $\mathbb{K}$. We define the **topological dual space of** $(X, \mathcal{T})$, denoted $(X, \mathcal{T})^*$, by $(X, \mathcal{T})^* := L((X, \mathcal{T}); (\mathbb{K}, \mathcal{T}_{\mathbb{K}}))$.

Note 8.1.2.5. When the context is clear we write X^* in place of $(X, \mathcal{T})^*$.

Exercise 8.1.2.6. Let X, Y be topological vector spaces and $\phi : X \rightarrow Y$. Suppose that ϕ is linear. Then ϕ is continuous iff ϕ is continuous at 0.

Proof. If ϕ is continuous, then ϕ is continuous at 0.

Conversely, suppose that ϕ is continuous at 0. Let $(x_\alpha)_{\alpha \in A} \subset X$ be a net and $x \in X$. Suppose that $x_\alpha \rightarrow x$. Then $x_\alpha - x \rightarrow 0$. Hence

$$\begin{aligned} \phi(x_\alpha) - \phi(x) &= \phi(x_\alpha - x) \\ &\rightarrow \phi(0) \\ &= 0 \end{aligned}$$

Therefore $\phi(x_\alpha) \rightarrow \phi(x)$ and ϕ is continuous at x . Since $x \in X$ is arbitrary, ϕ is continuous.

□

Exercise 8.1.2.7. Let X be a topological vector space and $\phi : X \rightarrow \mathbb{C}$ linear. Then $\phi \in X^*$ iff $|\phi|$ is continuous.

Proof. Suppose that ϕ is continuous. Since $|\cdot| : \mathbb{C} \rightarrow [0, \infty)$ is continuous, $|\phi|$ is continuous.

Conversely, suppose that $|\phi|$ is continuous. Let $(x_\alpha)_{\alpha \in A} \subset X$ be a net and $x \in X$. Suppose that $x_\alpha \rightarrow x$. Then $x_\alpha - x \rightarrow 0$. Therefore

$$\begin{aligned} |\phi(x_\alpha) - \phi(x)| &= |\phi(x_\alpha - x)| \\ &\rightarrow |\phi(0)| \\ &= 0 \end{aligned}$$

So $\phi(x_\alpha) \rightarrow \phi(x)$ and ϕ is continuous. □

Exercise 8.1.2.8. Let X be a real topological vector space and $\phi \in X^*$. If ϕ is not constant, then ϕ is open.

Hint: There exists $x_* \in X$ such that $\phi(x_*) = 1$ and for each $U \subset X$ open and $x \in U$, there exists $r > 0$ such that for each $t \in \mathbb{R}$, $|t| \leq r$ implies that $x + tx_* \in U$.

Proof. Suppose that ϕ is not constant. Then there exists $x_0 \in X$ such that $\phi(x_0) \neq 0$. Set $x_* = \phi(x_0)^{-1}x_0$. Then

$$\begin{aligned} \phi(x_*) &= \phi(\phi(x_0)^{-1}x_0) \\ &= \phi(x_0)^{-1}\phi(x_0) \\ &= 1 \end{aligned}$$

Let $U \subset X$ be open and $y \in \phi(U)$. Then there exists $x \in U$ such that $\phi(x) = y$. Since U is open, a previous exercise implies that there exists $r > 0$ such that for each $t \in \mathbb{R}$, $|t| \leq r$ implies that $x + tx_* \in U$. Let $t \in (-r, r)$. Then $\phi(x + tx_*) \in \phi(U)$. Since

$$\begin{aligned} \phi(x + tx_*) &= \phi(x) + t\phi(x_*) \\ &= y + t \end{aligned}$$

we have that $(y - r, y + r) \subset \phi(U)$. Since $y \in U$ is arbitrary, $\phi(U)$ is open thus ϕ is open. □

Definition 8.1.2.9. Let X be a vector space and $\phi : X \rightarrow \mathbb{C}$. Then ϕ is said to be **real-linear** if for each $x, y \in X$ and $\lambda \in \mathbb{R}$, $\phi(x + \lambda y) = \phi(x) + \lambda\phi(y)$.

Exercise 8.1.2.10. Let X be a topological vector space and $\phi \in X^*$. Then $\text{Re } \phi$ is continuous and real-linear.

Proof. Clear. □

Exercise 8.1.2.11. Let X be a topological vector space and $f : X \rightarrow \mathbb{R}$. If f is continuous and real-linear, then there exists a unique $\phi \in X^*$ such that $\text{Re } \phi = f$.

Hint: For each $z \in \mathbb{C}$, $z = \pi_{\mathbb{R}}(z) - i\pi_{\mathbb{R}}(iz)$

Proof. Suppose that f is continuous and real-linear. Define $\phi : X \rightarrow \mathbb{C}$ by $\phi(x) = f(x) - if(ix)$. Then ϕ is continuous. Let $x, y \in X$ and $\lambda \in \mathbb{C}$. Write $\lambda = a + bi$. Then

$$\begin{aligned} \phi(x + \lambda y) &= f(x + \lambda y) - if(ix + \lambda y) \\ &= f(x + ay + iby) - if(ix + iay - by) \\ &= f(x) + af(y) + bf(iy) - if(ix) - iaf(iy) + ibf(y) \\ &= [f(x) - if(ix)] + a[f(y) - if(iy)] + ib[f(y) - if(iy)] \\ &= \phi(x) + a\phi(y) + ibf(y) \\ &= \phi(x) + \lambda\phi(y) \end{aligned}$$

So ϕ is linear and $\phi \in X^*$. Let $\psi \in X^*$. Suppose that $f = \text{Re } \psi$. Then for each $x \in X$,

$$\begin{aligned} \phi(x) &= f(x) - if(ix) \\ &= \text{Re } [\psi(x)] - i\text{Re } [\psi(ix)] \\ &= \text{Re } [\psi(x)] - \text{Re } [i\psi(x)] \\ &= \text{Re } [\psi(x)] + \text{Im } [\psi(x)] \\ &= \psi(x) \end{aligned}$$

So $\psi = \phi$ and ϕ is unique. □

Exercise 8.1.2.12. Let X be a vector space, $(Y_\alpha, \mathcal{T}_\alpha)_{\alpha \in A}$ a collection of topological vector spaces and $(\phi_\alpha)_{\alpha \in A} \in \prod_{\alpha \in A} \mathcal{L}(X; Y_\alpha)$. Set $\mathcal{T} := \sigma_X(\phi_\alpha : \alpha \in A)$. Then $(X, \mathcal{T})$ is a topological vector space.

Proof. Let $(x_{1,\gamma})_{\gamma \in \Gamma}, (x_{2,\gamma})_{\gamma \in \Gamma} \subset X$, $(\lambda_\gamma)_{\gamma \in \Gamma} \subset \mathbb{K}$ be nets, $x_1, x_2 \in X$ and $\lambda \in \mathbb{K}$. Suppose that $x_{1,\gamma} \rightarrow x_1$ and $x_{2,\gamma} \rightarrow x_2$ in $(X, \mathcal{T})$ and $\lambda_\gamma \rightarrow \lambda$. Let $\alpha \in A$. Since $\mathcal{T} = \sigma_X(\phi_\alpha : \alpha \in A)$, Exercise 4.3.2.15 implies that $\phi_\alpha(x_{1,\gamma}) \rightarrow \phi_\alpha(x_1)$ and $\phi_\alpha(x_{2,\gamma}) \rightarrow \phi_\alpha(x_2)$. Since $(Y_\alpha, \mathcal{T}_\alpha)$ is a topological vector space, we have that

$$\begin{aligned} \phi_\alpha(x_{1,\gamma} + \lambda_\gamma x_{2,\gamma}) &= \phi_\alpha(x_{1,\gamma}) + \lambda_\gamma \phi_\alpha(x_{2,\gamma}) \\ &\rightarrow \phi_\alpha(x_1) + \lambda \phi_\alpha(x_2) \\ &= \phi_\alpha(x_1 + \lambda x_2). \end{aligned}$$

Since $\alpha \in A$ is arbitrary, we have that for each $\alpha \in A$, $\phi_\alpha(x_{1,\gamma} + \lambda_\gamma x_{2,\gamma}) \rightarrow \phi_\alpha(x_1 + \lambda x_2)$. Exercise 4.3.2.15 implies that $x_{1,\gamma} + \lambda_\gamma x_{2,\gamma} \rightarrow x_1 + \lambda x_2$ in $(X, \mathcal{T})$. Since $(x_{1,\gamma})_{\gamma \in \Gamma}, (x_{2,\gamma})_{\gamma \in \Gamma} \subset X$, $(\lambda_\gamma)_{\gamma \in \Gamma} \subset \mathbb{K}$ nets, $x_1, x_2 \in X$ and $\lambda \in \mathbb{K}$ are arbitrary, we have that for each $(x_{1,\gamma})_{\gamma \in \Gamma}, (x_{2,\gamma})_{\gamma \in \Gamma} \subset X$, $(\lambda_\gamma)_{\gamma \in \Gamma} \subset \mathbb{K}$ nets, $x_1, x_2 \in X$ and $\lambda \in \mathbb{K}$, $x_{1,\gamma} \rightarrow x_1$, $x_{2,\gamma} \rightarrow x_2$ in $(X, \mathcal{T})$ and $\lambda_\gamma \rightarrow \lambda$ implies that $x_{1,\gamma} + \lambda_\gamma x_{2,\gamma} \rightarrow x_1 + \lambda x_2$ in $(X, \mathcal{T})$. Exercise 8.1.1.3 implies that $(X, \mathcal{T})$ is a topological vector space. $\square$

8.1.3 Products

Exercise 8.1.3.1. Let $(X_\alpha, \mathcal{T}_\alpha)_{\alpha \in A}$ a collection of topological vector spaces. Then $\left(\prod_{\alpha \in A} X_\alpha, \bigotimes_{\alpha \in A} \mathcal{T}_\alpha \right)$ is a topological vector space.

Proof. Clear by Exercise 8.1.2.12. $\square$

Exercise 8.1.3.2. We have that $(\mathbb{R}^{\mathbb{N}}, \mathcal{T}_{\mathbb{R}}^{\otimes \mathbb{N}})$ is a topological vector space.

Proof. Clear by Exercise 8.1.3.1. $\square$

Exercise 8.1.3.3. Define $(e_n)_{n \in \mathbb{N}} \subset \mathbb{R}^{\mathbb{N}}$ by $\pi_k(e_n) := \delta_{n,k}$.

1. We have that $\text{span}(e_n : n \in \mathbb{N})$ is dense in $(\mathbb{R}^{\mathbb{N}}, \mathcal{T}_{\mathbb{R}}^{\otimes \mathbb{N}})$.

2. Let $\phi \in (\mathbb{R}^{\mathbb{N}})^*$. Then $\#\{n \in \mathbb{N} : \phi(e_n) \neq 0\} < \infty$.

Hint: Exercise 4.6.2.3

3. We have that $(\mathbb{R}^{\mathbb{N}})^* = \text{span}\{\pi_j : j \in \mathbb{N}\}$.

Hint: Exercise 6.7.1.11

Proof.

1. Define $E \subset \mathbb{R}^{\mathbb{N}}$ by $E := \text{span}\{e_j : j \in \mathbb{N}\}$. Let $x \in \mathbb{R}^{\mathbb{N}}$. Define $(x_n)_{n \in \mathbb{N}} \subset E$ by $x_n := \sum_{j=1}^n \pi_j(x) e_j$. Let $k \in \mathbb{N}$. Then for each $n \in \mathbb{N}$, $n \geq k$ implies that

$$\begin{aligned} \pi_k(x_n) &= \sum_{j=1}^n \pi_j(x) \pi_k(e_j) \\ &= \sum_{j=1}^n \pi_j(x) \delta_{k,j} \\ &= \pi_k(x). \end{aligned}$$

Therefore $\pi_k(x_n) \rightarrow \pi_k(x)$. Since $k \in \mathbb{N}$ is arbitrary, we have that for each $k \in \mathbb{N}$, $\pi_k(x_n) \rightarrow \pi_k(x)$. Thus $x_n \rightarrow x$ in $(\mathbb{R}^{\mathbb{N}}, \mathcal{T}_{\mathbb{R}}^{\otimes \mathbb{N}})$. Hence $x \in \text{cl } E$. Since $x \in \mathbb{R}^{\mathbb{N}}$ is arbitrary, we have that $\mathbb{R}^{\mathbb{N}} \subset \text{cl } E$. Since clearly, $\text{cl } E \subset \mathbb{R}^{\mathbb{N}}$, we have that $\mathbb{R}^{\mathbb{N}} = \text{cl } E$. Thus E is dense in $\mathbb{R}^{\mathbb{N}}$.

2. Define $a \in \mathbb{R}^{\mathbb{N}}$ by $a_n := \phi(e_n)$. Set $J := \{n \in \mathbb{N} : a_n \neq 0\}$. For the sake of contradiction, suppose that $\#J = \infty$. Then there exists a subsequence $(a_{n_j})_{j \in \mathbb{N}} \subset (a_n)_{n \in \mathbb{N}}$ such that for each $j \in \mathbb{N}$, $a_{n_j} \neq 0$. Define $(x_j)_{j \in \mathbb{N}} \subset \mathbb{R}^{\mathbb{N}}$ by $x_j := a_{n_j}^{-1}e_{n_j}$. We note that for each $j \in \mathbb{N}$,

$$\begin{aligned}\phi(x_j) &= \phi(a_{n_j}^{-1}e_{n_j}) \\ &= a_{n_j}^{-1}\phi(e_{n_j}) \\ &= a_{n_j}^{-1}a_{n_j} \\ &= 1.\end{aligned}$$

Thus $\phi(x_j) \rightarrow 1$. Since for each $k \in \mathbb{N}$,

$$\begin{aligned}\pi_k(x_j) &= a_{n_j}^{-1}\pi_k(e_{n_j}) \\ &= a_{n_j}^{-1}\delta_{k,n_j} \\ &\xrightarrow{j} 0.\end{aligned}$$

Exercise 4.6.2.2 implies that $x_j \rightarrow 0$ in $(\mathbb{R}^{\mathbb{N}}, \mathcal{T}_{\mathbb{R}}^{\otimes \mathbb{N}})$. Since ϕ is continuous, $\phi(x_j) \rightarrow 0$. This is a contradiction. Hence $\#J < \infty$.

3. Define $F \subset (\mathbb{R}^{\mathbb{N}})^*$ by $F := \text{span}\{\pi_j : j \in \mathbb{N}\}$. Let $\phi \in (\mathbb{R}^{\mathbb{N}})^*$. The previous part implies that there exists $N \in \mathbb{N}$ such that for each $n \in \mathbb{N}$, $n > N$ implies that $\phi(e_n) = 0$. Define $\lambda \in (\mathbb{R}^{\mathbb{N}})^*$ by $\lambda := \phi - \sum_{n=1}^N \phi(e_n)\pi_n$. Then for each $l \in \mathbb{N}$, $l \leq N$ implies that

$$\begin{aligned}\lambda(e_l) &= \phi(e_l) - \sum_{n=1}^N \phi(e_n)\pi_n(e_l) \\ &= \phi(e_l) - \sum_{n=1}^N \phi(e_n)\delta_{n,l} \\ &= \phi(e_l) - \phi(e_l) \\ &= 0.\end{aligned}$$

and $l > N$ implies that

$$\begin{aligned}\lambda(e_l) &= \phi(e_l) - \sum_{n=1}^N \phi(e_n)\pi_n(e_l) \\ &= \phi(e_l) - \sum_{n=1}^N \phi(e_n)\delta_{n,l} \\ &= 0 - 0 \\ &= 0.\end{aligned}$$

Thus $\lambda|_E = 0$. Since E is dense in $\mathbb{R}^{\mathbb{N}}$ and λ is continuous. We recall that Exercise 6.5.0.2 implies that $\mathbb{R}^{\mathbb{N}}$ is a metric space. Exercise 6.7.1.11 then implies that $\lambda = 0$. Thus

$$\begin{aligned}\phi &= \sum_{n=1}^N \phi(e_n)\pi_n \\ &\in F.\end{aligned}$$

Since $\phi \in (\mathbb{R}^{\mathbb{N}})^*$ is arbitrary, we have that $(\mathbb{R}^{\mathbb{N}})^* \subset F$. Clearly $F \subset (\mathbb{R}^{\mathbb{N}})^*$. Hence $(\mathbb{R}^{\mathbb{N}})^* = F$.

□

8.2 Sublinear Functionals

Definition 8.2.0.1. Let X be a vector space and $p : X \rightarrow \mathbb{R}$. Then

- p is said to be
 - **subadditive** if for each $x, y \in X$, $p(x + y) \leq p(x) + p(y)$,
 - **positively homogeneous** if for each $x \in X$ and $\lambda \in [0, \infty)$, $p(\lambda x) = \lambda p(x)$
- p is said to be a **sublinear functional on X** if
 1. p is subadditive,
 2. p is positively homogeneous,

Exercise 8.2.0.2. Let X be a vector space and $p : X \rightarrow \mathbb{R}$ be a sublinear functional. Then $p(0) = 0$.

Proof. We have that

$$\begin{aligned} p(0) &= p(0 \cdot 0) \\ &= 0p(0) \\ &= 0 \end{aligned}$$

□

Proof. Clear

□

Exercise 8.2.0.3. Let X be a vector space and $p : X \rightarrow \mathbb{R}$ a sublinear functional. Then for each $x, y \in X$

1. $-p(-x) \leq p(x)$
2. $-p(y - x) \leq p(x) - p(y) \leq p(x - y)$

Proof. Let $x, y \in X$.

1. We have

$$\begin{aligned} 0 &= p(0) \\ &= p(x - x) \\ &\leq p(x) + p(-x) \end{aligned}$$

So $-p(-x) \leq p(x)$.

2. We have

$$\begin{aligned} p(x) &= p(x - y + y) \\ &\leq p(x - y) + p(y) \end{aligned}$$

So $p(x) - p(y) \leq p(x - y)$. Switching x and y gives us $p(y) - p(x) \leq p(y - x)$ and multiplying both sides by -1 yields $-p(y - x) \leq p(x) - p(y)$

Putting these two together, we see that

$$-p(y - x) \leq p(x) - p(y) \leq p(x - y)$$

□

Exercise 8.2.0.4. Hahn-Banach Theorem for Sublinear Functionals

Let X be a vector space, $p : X \rightarrow \mathbb{R}$ a sublinear functional, $M \subset X$ a subspace and $f : M \rightarrow \mathbb{R}$ a linear functional. If for each $x \in M$, $f(x) \leq p(x)$, then there exists a linear functional $F : X \rightarrow \mathbb{R}$ such that for each $x \in X$, $F(x) \leq p(x)$ and $F|_M = f$.

Proof. **FINISH!!!**

□

Exercise 8.2.0.5. Let X be a vector space and $p : X \rightarrow \mathbb{R}$ a sublinear functional. Then there exists a linear functional $F : X \rightarrow \mathbb{R}$ such that for each $x \in X$, $F(x) \leq p(x)$.

Proof. Take $M = \{0\}$ and $f \equiv 0$ and apply the Hahn-Banach theorem. $\square$

Exercise 8.2.0.6. Equivalency of linearity (General Case) Let X be a vector space and $p : X \rightarrow \mathbb{R}$ a sublinear functional. Then the following are equivalent:

1. there exists a unique $F \in X^*$ such that $F \leq p$
2. for each $x \in X$, $-p(-x) = p(x)$
3. p is linear

Hint: If there exists $x \in X$ such that $-p(-x) \neq p(x)$, define $f_1, f_2 : \text{span}(x) \rightarrow \mathbb{R}$ by $f_1(tx) = tp(x)$ and $f_2(tx) = -tp(-x)$

Proof.

- (1) $\implies$ (2):

Suppose that there exists a unique $F \in X^*$ such that $F \leq p$. For the sake of contradiction, suppose that there exists $x \in X$ such that $-p(-x) \neq p(x)$. Define $f_1, f_2 : \text{span}(x) \rightarrow \mathbb{R}$ by

$$f_1(tx) = tp(x)$$

and

$$f_2(tx) = -tp(-x)$$

Let $y \in \text{span}(x)$. Then there exists $t \in \mathbb{R}$ such that $y = tx$. Then for each $k \in \mathbb{R}$,

$$\begin{aligned} f_1(ky) &= f_1(ktx) \\ &= ktp(x) \\ &= kf_1(tx) \\ &= kf_1(y) \end{aligned}$$

Similarly, $f_2(ky) = kf_2(y)$ and so $f_1, f_2 \in \text{span}(x)^*$. If $t \geq 0$, then

$$\begin{aligned} f_1(y) &= f_1(tx) \\ &= tp(x) \\ &= p(tx) \\ &= p(y) \end{aligned}$$

If $t < 0$, then

$$\begin{aligned} f_1(y) &= f_1(tx) \\ &= tp(x) \\ &= -|t|p(x) \\ &= -p(|t|x) \\ &= -p(-tx) \\ &\leq p(tx) \\ &= p(y) \end{aligned}$$

So $f_1 \leq p$ on $\text{span}(x)$. Similarly, $f_2 \leq p$ on $\text{span}(x)$. The Hahn-Banach theorem implies that there exist $F_1, F_2 \in X^*$ such that $F_1, F_2 \leq p$ and $F_1 = f_1, F_2 = f_2$ on $\text{span}(x)$. By the assumption of uniqueness, $F_1 = F_2$. This is a contradiction since

$$\begin{aligned} F_1(x) &= p(x) \\ &\neq -p(-x) \\ &= F_2(x) \end{aligned}$$

So for each $x \in X$, $-p(-x) = p(x)$.

- (2) $\Rightarrow$ (3):

Suppose that for each $x \in X$, $-p(-x) = p(x)$. The previous exercise implies that there exists $F \in X^*$ such that $F \leq p$. Let $x \in X$. Then

$$\begin{aligned} -F(x) &= F(-x) \\ &\leq p(-x) \\ &= -p(x) \end{aligned}$$

So $p(x) \leq F(x)$ and $p \leq F$. Therefore $p = F$ and p is linear.

- (3) $\Rightarrow$ (1):

Suppose that p is linear. Let $F \in X^*$. Suppose that $F \leq p$. Let $x \in X$. Then as in the case for (2) $\Rightarrow$ (3), we have that

$$\begin{aligned} -F(x) &= F(-x) \\ &\leq p(-x) \\ &= -p(x) \end{aligned}$$

which implies that $p = F$. So p is the unique linear function $F \in X^*$ such that $F \leq p$.

□

8.3 Seminorms

Definition 8.3.0.1. Let X be a vector space.

- Let $p : X \rightarrow \mathbb{R}$. Then
 - p is said to be
 - * **absolutely homogeneous** if for each $x \in X$ and $\lambda \in \mathbb{K}$, $p(\lambda x) = |\lambda|p(x)$,
 - * **definite** if for each $x \in X$, $p(x) = 0$ implies that $x = 0$.
 - p is said to be a
 - * **seminorm on X** if
 1. p is subadditive,
 2. p is absolutely homogeneous,
 - * **norm on X** if
 1. p is a seminorm on X
 2. p is definite.
- We define $\text{SemiNorm}(X) := \{p \in \mathbb{R}^X : p \text{ is a seminorm on } X\}$.

Exercise 8.3.0.2. Let X be a vector space and $p \in \text{SemiNorm}(X)$, then p is a sublinear functional.

Proof. Clear. □

Definition 8.3.0.3. Let X be a vector space and $p \in \text{SemiNorm}(X)$. We define the **pseudometric induced by p** , denoted $d_p : X \times X \rightarrow [0, \infty)$, by $d_p(x, y) := p(x - y)$.

Exercise 8.3.0.4. Let X be a vector space and $p \in \text{SemiNorm}(X)$. Then

1. d_p is a pseudometric on X .
2. p is a norm on X implies that d_p is a metric on X .

Proof.

1. (a) For each $x \in X$,

$$\begin{aligned} d_p(x, x) &= p(x - x) \\ &= p(0) \\ &= 0 \quad (\text{by Exercise 8.3.0.2 and Exercise 8.2.0.2}). \end{aligned}$$

Thus d is semidefinite.

- (b) For each $x, y \in X$,

$$\begin{aligned} d_p(x, y) &= p(x - y) \\ &= p(-(y - x)) \\ &= |-1|p(y - x) \\ &= p(y - x) \\ &= d_p(y, x). \end{aligned}$$

Thus d is symmetric.

- (c) For each $x, y, z \in X$,

$$\begin{aligned} d_p(x, z) &= p(x - z) \\ &= p([x - y] + [y - z]) \\ &\leq p(x - y) + p(y - z) \\ &= d_p(x, y) + d_p(y, z). \end{aligned}$$

Thus d satisfies the triangle inequality,

Hence d_p is a pseudometric.

2. Suppose that p is a norm on X . Then for each $x, y \in X$,

$$\begin{aligned} d_p(x, y) = 0 &\implies p(x - y) = 0 \\ &\implies x - y = 0 \quad (\text{since } p \text{ is definite}) \\ &\implies x = y. \end{aligned}$$

Thus d_p is definite. Thus d_p is a metric on X . □

Exercise 8.3.0.5. Let X be a vector space and $p \in \text{SemiNorm}(X)$. Then $p \geq 0$.

Proof. For each $x \in X$,

$$\begin{aligned} p(x) &= p(x - 0) \\ &= d_p(x, 0) \\ &\geq 0 \quad (\text{by Exercise 6.1.1.3}). \end{aligned}$$

Thus $p \geq 0$. □

Exercise 8.3.0.6. Let X be a vector space and $p \in \text{SemiNorm}(X)$. Then

1. **Triangle Inequality:**

for each $x, y \in X$, $p(x - y) \leq p(x) + p(y)$

2. **Reverse Triangle Inequality:**

for each $x, y \in X$, $|p(x) - p(y)| \leq |p(x) - p(y)|$

Proof.

1. For each $x, y \in X$,

$$\begin{aligned} p(x - y) &= d_p(x, y) \\ &\leq d_p(x, 0) + d_p(0, y) \\ &= p(x - 0) + p(0 - y) \\ &= p(x) + p(y). \end{aligned}$$

2. For each $x, y \in X$,

$$\begin{aligned} |p(x) - p(y)| &= |d_p(x, 0) - d_p(0, y)| \\ &\leq d_p(x, y) \\ &= p(x - p(y)). \end{aligned}$$

Exercise 8.3.0.7. Let X be a vector space and $\phi \in X^*$. Then $|\phi|$ is a seminorm on X . □

Proof. Clear. □

Exercise 8.3.0.8. Let X, Y be a vector spaces, $T \in \mathcal{L}(X; Y)$ and p a seminorm on Y . Then $p \circ T$ is a seminorm on X . □

Proof. Clear. □

Exercise 8.3.0.9. Reverse Triangle Inequality:

Let X be a vector space and $p \in \text{SemiNorm}(X)$. Then for each $x, y \in X$, $|p(x) - p(y)| \leq p(x - y)$.

Proof. Let $x, y \in X$. Then

$$\begin{aligned} p(x) &= p(x - y + y) \\ &\leq p(x - y) + p(y) \end{aligned}$$

So $p(x) - p(y) \leq p(x - y)$. Similarly, $p(y) \leq p(y - x) + p(x)$ and so $p(x) - p(y) \leq p(x - y)$. Therefore $|p(x) - p(y)| \leq p(x - y)$. $\square$

Exercise 8.3.0.10. Let X be a vector space, $p \in \text{SemiNorm}(X)$ and $\phi \in X^*$. Then $\phi \leq p$ iff $|\phi| \leq p$.

Proof. • ($\implies$):

Suppose that $\phi \leq p$. Let $x \in X$. Then

$$\begin{aligned} -\phi(x) &= \phi(-x) \\ &\leq p(-x) \\ &= p(x) \end{aligned}$$

So $-p(x) \leq \phi(x)$. Hence $-p \leq \phi \leq p$. Thus $|\phi| \leq p$.

• ($\impliedby$):

Conversely, if $|\phi| \leq p$, then

$$\begin{aligned} \phi &\leq |\phi| \\ &\leq p. \end{aligned}$$

$\square$

Definition 8.3.0.11. Let X be a vector space and $p \in \text{SemiNorm}(X)$. We define the **kernel of p** , denoted $\ker p$, by $\ker p = p^{-1}(\{0\})$.

Exercise 8.3.0.12. Let X be a vector space and $p \in \text{SemiNorm}(X)$. Then $\ker p$ is a subspace of X .

Proof. Let $x, y \in \ker p$ and $\lambda \in \mathbb{K}$. Then $p(x) = 0$ and $p(y) = 0$. Thus

$$\begin{aligned} p(x + \lambda y) &\leq p(x) + p(\lambda y) \\ &= p(x) + |\lambda|p(y) \\ &= 0 \end{aligned}$$

Hence $x + \lambda y \in \ker p$. Since $x, y \in \ker p$ and $\lambda \in \mathbb{K}$ are arbitrary, we have that for each $x, y \in \ker p$ and $\lambda \in \mathbb{K}$, $x + \lambda y \in \ker p$. Thus $\ker p$ is a subspace of X . $\square$

Exercise 8.3.0.13. (ref def of $\sim_M$ for subspace M of vector space in algebra notes and ref defintion of $\sim_p$ from topology notes) Let X be a vector space and $p \in \text{SemiNorm}(X)$. Then

1. $\sim_{\ker p} \subset \sim_p$
2. if $\ker p \neq X$, then $\sim_{\ker p} \neq \sim_p$,
3. p is $\sim_{\ker p}$ -invariant.

Proof. 1. Let $(x, y) \in \sim_{\ker p}$. Then $x - y \in \ker p$. Thus there exists $n \in \ker p$ such that $x - y = n$. Therefore

$$\begin{aligned} p(x) &= p(y + n) \\ &\leq p(y) + p(n) \\ &= p(y) \\ &= p(x - n) \\ &\leq p(x) + p(-n) \\ &= p(x) + p(n) \\ &= p(x). \end{aligned}$$

Hence $p(x) = p(y)$ and $(x, y) \in \sim_p$. Since $(x, y) \in \sim_{\ker p}$ is arbitrary, we have that for each $(x, y) \in \sim_{\ker p}$, $(x, y) \in \sim_p$. Thus $\sim_{\ker p} \subset \sim_p$.

2. Suppose that $\ker p \neq X$. Then there exists $x \in X$ such that $x \notin \ker p$. Thus $p(x) \neq 0$. Since $\ker p$ is a subspace, $-x \notin \ker p$. Since

$$\begin{aligned} p(x - (-x))p(2x) \\ &= 2p(x) \\ &\neq 0, \end{aligned}$$

we have that $x - (-x) \notin \ker p$ and $x \not\sim_{\ker p} -x$. Since $p(-x) = p(x)$, we have that $x \sim_p -x$. Thus $(x, -x) \in \sim_p$ and $(x, -x) \notin \sim_{\ker p}$. Thus $\sim_{\ker p} \neq \sim_p$.

3. Let $x, y \in X$. Suppose that $x \sim_{\ker p} y$. **The previous part** implies that $x \sim_p y$. Thus $p(x) = p(y)$. Since $x, y \in X$ with $x \sim_{\ker p} y$ are arbitrary, we have that for each $x, y \in X$, $x \sim_{\ker p} y$ implies that $p(x) = p(y)$. Thus p is $\sim_{\ker p}$ -invariant. $\square$

Definition 8.3.0.14. Let X be a vector space and $p \in \text{SemiNorm}(X)$. We define the

- **canonical projection induced by p from X to $X/\ker p$** , denoted $\pi_p : X \rightarrow X/\ker p$, by $\pi_p(x) := x + \ker p$.
- **norm on $X/\ker p$ induced by p** , denoted $\bar{p} : X/\ker p \rightarrow [0, \infty)$, by $\bar{p} := \text{factor}(p, \pi_p)$.

Note 8.3.0.15. We recall that

- π_p is linear,
- $\bar{p}$ is the unique $f : X/\ker p \rightarrow [0, \infty)$ such that $f \circ \pi_p = p$ and in particular, for each $x \in X$, $\bar{p}(x + \ker p) = p(x)$.

Exercise 8.3.0.16. Let X be a vector space and $p \in \text{SemiNorm}(X)$. Then $\bar{p} : X/\ker p \rightarrow [0, \infty)$ is a norm

Proof.

1. (a) Let $a, b \in X/\ker p$. Then there exist $x, y \in X$ such that $a = x + \ker p$ and $b = y + \ker p$. Therefore

$$\begin{aligned} \bar{p}(a + b) &= \bar{p}((x + \ker p) + (y + \ker p)) \\ &= \bar{p}((x + y) + \ker p) \\ &= p(x + y) \\ &\leq p(x) + p(y) \\ &= \bar{p}(x + \ker p) + \bar{p}(y + \ker p) \\ &= \bar{p}(a) + \bar{p}(b). \end{aligned}$$

Since $a, b \in X/\ker p$ are arbitrary, we have that for each $a, b \in X/\ker p$, $\bar{p}(a + b) \leq \bar{p}(a) + \bar{p}(b)$. Thus $\bar{p}$ is subadditive.

- (b) Let $a \in X/\ker p$ and $\lambda \in \mathbb{K}$. Then there exists $x \in X$ such that $a = x + \ker p$. Hence

$$\begin{aligned} \bar{p}(\lambda a) &= \bar{p}(\lambda[x + \ker p]) \\ &= \bar{p}(\lambda x + \ker p) \\ &= p(\lambda x) \\ &= |\lambda|p(x) \\ &= |\lambda|\bar{p}(x + \ker p) \\ &= |\lambda|\bar{p}(a). \end{aligned}$$

Since $a \in X/\ker p$ and $\lambda \in \mathbb{K}$ are arbitrary, we have that for each $a \in X/\ker p$ and $\lambda \in \mathbb{K}$, $\bar{p}(\lambda a) = |\lambda|\bar{p}(a)$. Thus $\bar{p}$ is absolutely homogeneous.

Therefore $\bar{p}$ is a seminorm on $X/\ker p$.

2. Let $a \in X/\ker p$. Suppose that $\bar{p}(a) = 0$. Since $a \in X/\ker p$, there exists $x \in X$ such that $a = x + \ker p$. Then

$$\begin{aligned} p(x) &= \bar{p}(x + \ker p) \\ &= \bar{p}(a) \\ &= 0. \end{aligned}$$

Thus $x \in \ker p$. Hence

$$\begin{aligned} a &= x + \ker p \\ &= 0 + \ker p \\ &= 0_{X/\ker p}. \end{aligned}$$

Since $a \in X/\ker p$ with $\bar{p}(a) = 0$, we have that for each $a \in X/\ker p$, $\bar{p}(a) = 0$ implies that $a = 0_{X/\ker p}$. Hence $\bar{p}$ is definite.

Thus $\bar{p}$ is a norm on $X/\ker p$. □

Definition 8.3.0.17. Let X be a vector space, $p \in \text{SemiNorm}(X)$, $x \in X$ and $r > 0$. We define the

- **open semiball of p at x of radius r** , denoted $B_p(x, r)$, by

$$B_p(x, r) := B_{d_p}(x, r)$$

- **closed semiball of p at x of radius r** , denoted $\bar{B}_p(x, r)$, by

$$\bar{B}_p(x, r) := \bar{B}_{d_p}(x, r)$$

Note 8.3.0.18. By definition [ref definition in pseudometric space section](#)

- $B_p(x, r) = \{y \in X : p(x - y) < r\}$.
- $\bar{B}_p(x, r) = \{y \in X : p(x - y) \leq r\}$

Exercise 8.3.0.19. Let X be a vector space, $p \in \text{SemiNorm}(X)$, $x \in X$ and $r > 0$. Then $B_p(x, r) = x + rB_p(0, 1)$.

Proof. Let $y \in B_p(x, r)$. Then

$$\begin{aligned} p(r^{-1}(y - x)) &= r^{-1}p(y - x) \\ &< r^{-1}r \\ &= 1 \end{aligned}$$

So $r^{-1}(y - x) \in B_p(0, 1)$. By definition, there exists $u \in B_p(0, 1)$ such that $r^{-1}(y - x) = u$, which implies that

$$\begin{aligned} y &= x + ru \\ &\in x + rB_p(0, 1) \end{aligned}$$

Conversely, let $y \in x + rB_p(0, 1)$. By definition, there exists $u \in B_p(0, 1)$ such that $y = x + ru$. Then

$$\begin{aligned} p(y - x) &= p(ru) \\ &= rp(u) \\ &< r \end{aligned}$$

So $y \in B_p(x, r)$ □

Exercise 8.3.0.20. Let X be a vector space and $p, q \in \text{SemiNorm}(X)$. Then $p \leq q$ iff $B_q(0, 1) \subset B_p(0, 1)$.

Proof. Suppose that $p \leq q$. Let $x \in B_q(0, 1)$. Then

$$\begin{aligned} p(x) &\leq q(x) \\ &< 1 \end{aligned}$$

So $x \in B_p(0, 1)$.

Conversely, suppose that $B_q(0, 1) \subset B_p(0, 1)$. Let $x \in X$. If $p(x) = 0$, then $p(x) \leq q(x)$. Suppose that $p(x) > 0$. For the sake of contradiction, suppose that $p(x) > q(x)$. Then

$$\begin{aligned} q\left(\frac{x}{p(x)}\right) &= \frac{q(x)}{p(x)} \\ &< 1 \end{aligned}$$

Therefore, $x/p(x) \in B_q(0, 1) \subset B_p(0, 1)$. By definition,

$$\begin{aligned} \frac{p(x)}{p(x)} &= p\left(\frac{x}{p(x)}\right) \\ &< 1 \end{aligned}$$

which is a contradiction. So $p(x) \leq q(x)$. Since $x \in X$ is arbitrary, $p \leq q$. □

Exercise 8.3.0.21. Let X be a topological vector space and $p \in \text{SemiNorm}(X)$. Suppose that p is continuous. Then

1. $B_p(0, 1)$ is open
2. $\bar{B}_p(0, 1)$ is closed

Proof.

1. Since $B_p(0, 1) = p^{-1}([0, 1))$ and p is continuous, $B_p(0, 1)$ is open.
2. Since $\text{bar}B_p(0, 1) = p^{-1}([0, 1])$ and p is continuous, $\text{bar}B_p(0, 1)$ is closed. □

Exercise 8.3.0.22. Let X be a topological vector space and $p \in \text{SemiNorm}(X)$. Then the following are equivalent:

1. p is continuous
2. $B_p(0, 1)$ is open
3. $\bar{B}_p(0, 1) \in \mathcal{N}(0)$
4. p is continuous at 0.

Proof.

- (1) $\implies$ (2):
Clear by Exercise 8.3.0.21.
- (2) $\implies$ (3):
Clear since $B_p(0, 1) \subset \bar{B}_p(0, 1)$.
- (3) $\implies$ (4):
Let $(x_\alpha)_{\alpha \in A} \subset X$ be a net. Suppose that $x_\alpha \rightarrow 0$. Let $U \in \mathcal{N}(0)$. Suppose that $U \in \mathcal{N}(0)$. Then there exists $\epsilon > 0$ such that $\bar{B}(0, \epsilon) \subset U$. Since the map $f_\epsilon : X \rightarrow X$ defined by $f_\epsilon(x) = \epsilon x$ is a homeomorphism, $\bar{B}_p(0, \epsilon) = \epsilon \bar{B}_p(0, 1) \in \mathcal{N}(0)$. Hence there exists $\beta \in A$ such that for each $\alpha \geq \beta$, $x_\alpha \in \bar{B}_p(0, \epsilon)$. Let $\alpha \in A$. Suppose that $\alpha \geq \beta$. By definition, $p(x_\alpha) \leq \epsilon$. So $p(x_\alpha) \in \bar{B}(0, \epsilon) \subset U$. Hence $(p(x_\alpha))_{\alpha \in A}$ is eventually in U . Since $U \in \mathcal{N}(0)$ is arbitrary, $p(x_\alpha) \rightarrow 0$. So p is continuous at 0.

- (4) $\implies$ (1):

Let $(x_\alpha)_{\alpha \in A} \subset X$ be a net and $x \in X$. Suppose that $x_\alpha \rightarrow x$. Then $x_\alpha - x \rightarrow 0$. Therefore $p(x_\alpha - x) \rightarrow 0$. The reverse triangle inequality implies that $p(x_\alpha) \rightarrow p(x)$. So p is continuous.

□

Exercise 8.3.0.23. Let X be a topological vector space and $p \in \text{SemiNorm}(X)$. Then p is continuous iff there exists a continuous seminorm $q : X \rightarrow [0, \infty)$ such that $p \leq q$.

Proof. Suppose that p is continuous. Set $q = p$. Then q is a continuous and $p \leq q$.

Conversely, suppose that there exists a continuous seminorm $q : X \rightarrow [0, \infty)$ such that $p \leq q$. Then $\bar{B}_q(0, 1) \subset \bar{B}_p(0, 1)$. The previous exercise tells us that

$$\begin{aligned} q \text{ is continuous} &\iff \bar{B}_q(0, 1) \in \mathcal{N}(0) \\ &\implies \bar{B}_p(0, 1) \in \mathcal{N}(0) \\ &\iff p \text{ is continuous} \end{aligned}$$

□

Theorem 8.3.0.24. Hahn-Banach Theorem for Seminorms (move this somewhere else?)

Let X be a vector space, $p \in \text{SemiNorm}(X)$, $M \subset X$ a subspace and $f : M \rightarrow \mathbb{C}$ a linear functional. If for each $x \in M$, $|f(x)| \leq p(x)$, then there exists a linear functional $F : X \rightarrow \mathbb{C}$ such that for each $x \in X$, $|F(x)| \leq p(x)$ and $F|_M = f$.

8.4 Minkowski Functionals

Note 8.4.0.1. The rest of this chapter assumes familiarity with the basic ideas of convex sets (Section 15.2).

Definition 8.4.0.2. Let X be a vector space.

- Let $E \subset X$. Then E is said to be **balanced** if for each $x \in E$, $r \in \mathbb{K}$, $|r| \leq 1$ implies that $rx \in E$.
- We define $\text{Bal}(X) \subset \mathcal{P}(X)$ by $\text{Bal}(X) := \{E \subset X : E \text{ is balanced}\}$.

Exercise 8.4.0.3. Let X be a vector space.

1. For each $E \in \text{Bal}(X)$, $E \neq \emptyset$ implies that $0 \in E$.
2. For each $M \subset X$, M is a subspace of X implies that $M \in \text{Bal}(X)$.

Proof. Clear. □

Exercise 8.4.0.4. Let X be a vector space and $\mathcal{B} \subset \text{Bal}(X)$. Then

1. $\bigcap_{E \in \mathcal{B}} E \neq \emptyset$,
2. $\bigcap_{E \in \mathcal{B}} E \in \text{Bal}(X)$.

Proof. Suppose that $\mathcal{B} \neq \emptyset$. Define $E \subset X$ by $E := \bigcap_{E' \in \mathcal{B}} E'$.

1. **The prev ex** implies that for each $E' \in \mathcal{B}$, $0 \in E'$. Thus

$$\begin{aligned} 0 &\in \bigcap_{E' \in \mathcal{B}} E' \\ &= E. \end{aligned}$$

Hence $E \neq \emptyset$.

2. Let $x \in E$ and $r \in \mathbb{K}$. Suppose that $|r| \leq 1$. Let $E_0 \in \mathcal{B}$. Then

$$\begin{aligned} x &\in E \\ &= \bigcap_{E' \in \mathcal{B}} E' \\ &\subset E_0. \end{aligned}$$

Since $E_0 \in \text{Bal}(X)$, $x \in E_0$ and $|r| \leq 1$, we have that $rx \in E_0$. Since $E_0 \in \mathcal{B}$ is arbitrary, we have that for each $E_0 \in \mathcal{B}$, $rx \in E_0$. Thus

$$\begin{aligned} rx &\in \bigcap_{E' \in \mathcal{B}} E' \\ &= E. \end{aligned}$$

Since $x \in E$ and $r \in \mathbb{K}$ with $|r| \leq 1$ are arbitrary, we have that for each $x \in E$ and $r \in \mathbb{K}$, $|r| \leq 1$ implies that $rx \in E$. Hence $E \in \text{Bal}(X)$. □

Definition 8.4.0.5. Let X be a vector space and $A \subset X$. Define $\mathcal{B}_A \subset \text{Bal}(X)$ by $\mathcal{B}_A := \{E \in \text{Bal}(X) : A \subset E\}$. We define the **balanced hull of A** , denoted $\text{bal } A \subset X$, by

$$\text{bal } A := \bigcap_{E \in \mathcal{B}_A} E.$$

Note 8.4.0.6. We may think of $\text{bal } A$ as the smallest balanced subset of X that contains A .

Exercise 8.4.0.7. Let X be a vector space and $A \subset X$. Then

$$\text{bal } A = \bigcup_{\substack{r \in \mathbb{K} \\ |r| \leq 1}} rA.$$

Proof. Define $B_A \subset X$ by $B_A := \bigcup_{\substack{r \in \mathbb{K} \\ |r| \leq 1}} rA$.

- – We note that

$$\begin{aligned} A &= 1 \cdot A \\ &\subset \bigcup_{\substack{r \in \mathbb{K} \\ |r| \leq 1}} rA \\ &= B_A. \end{aligned}$$

- Let $x \in B_A$ and $r \in \mathbb{K}$. Suppose that $|r| \leq 1$. Since $x \in B_A$, there exists $r_0 \in \mathbb{K}$ such that $|r_0| \leq 1$ and $x \in r_0A$. Since

$$\begin{aligned} |rr_0| &= |r||r_0| \\ &\leq 1 \cdot 1 \\ &= 1, \end{aligned}$$

we have that

$$\begin{aligned} rx &\in r(r_0A) \\ &= (rr_0)A \\ &\in \bigcup_{\substack{r \in \mathbb{K} \\ |r| \leq 1}} r'A \\ &= B_A. \end{aligned}$$

Since $x \in B_A$ and $r \in \mathbb{K}$ with $|r| \leq 1$ are arbitrary, we have that for each $x \in B_A$ and $r \in \mathbb{K}$, $|r| \leq 1$ implies that $rx \in B_A$. Hence $B_A \in \text{Bal}(X)$.

- Since $A \subset B_A$ and $B_A \in \text{Bal}(X)$, we have that $\text{bal } A \subset B_A$.
- Let $x \in B_A$. Then there exists $r \in \mathbb{K}$ such that $|r| \leq 1$ and $x \in rA$. Thus there exists $a \in A$ such that $x = ra$. Since $A \subset \text{bal } A$, we have that $a \in \text{bal } A$. Since $\text{bal } A \in \text{Bal}(X)$ and $|r| \leq 1$, we have that

$$\begin{aligned} x &= ra \\ &\in \text{bal } A. \end{aligned}$$

Since $x \in B_A$ is arbitrary, we have that for each $x \in B_A$, $x \in \text{bal } A$. Thus $B_A \subset \text{bal}(A)$.

Since $\text{bal}(A) \subset B_A$ and $B_A \subset \text{bal}(A)$, we have that $\text{bal } A = B_A$. □

Exercise 8.4.0.8. Let X be a vector space, $A \in \text{Bal}(X)$. Then for each $x \in X$ and $\lambda \in \mathbb{K}$, $\lambda x \in A$ iff $|\lambda|x \in A$.

Proof. Let $x \in X$ and $\lambda \in \mathbb{K}$.

- ($\implies$):

If $\lambda = 0$, then the claim is clearly true. Suppose that $\lambda \neq 0$. Define $s \in \mathbb{K}$ by $s := \text{sgn}(\lambda)$. Suppose that $\lambda x \in A$. Since A is balanced and $|s| = |s^{-1}| = 1$,

$$\begin{aligned} |\lambda|x &= s^{-1}\lambda x \\ &\in A \end{aligned}$$

• ($\Leftarrow$):

Conversely, suppose that $|\lambda|x \in A$. Then

$$\begin{aligned}\lambda x &= s|\lambda|x \\ &\in A\end{aligned}$$

□

Exercise 8.4.0.9. Let X be a vector space and $A \subset X$. If $A \in \text{Bal}(X)$, then $\text{conv } A \in \text{Bal}(X)$.

Proof. Suppose that $A \in \text{Bal}(X)$. Let $x \in \text{conv } A$ and $r \in \mathbb{K}$. Suppose that $|r| \leq 1$. Then there exist $(a_j)_{j \in [n]} \subset A$ and $(\lambda_j)_{j \in [n]} \subset [0, 1]$ such that $x = \sum_{j \in [n]} \lambda_j a_j$ and $\sum_{j \in [n]} \lambda_j = 1$. Since $A \in \text{Bal}(X)$, for each $j \in \{1, \dots, n\}$,

$$\begin{aligned}ra_j &\in A \\ &\subset \text{conv } A\end{aligned}$$

Since $\text{conv } A$ is convex, we have that

$$\begin{aligned}rx &= r \sum_{j \in [n]} \lambda_j a_j \\ &= \sum_{j \in [n]} \lambda_j ra_j \\ &\in \text{conv } A\end{aligned}$$

Hence $\text{conv } A \in \text{Bal}(X)$. □

Definition 8.4.0.10. Let X be a vector space and $A \subset X$. Then A is said to be **absorbing** if for each $x \in X$, there exists $r > 0$ such that for each $c \in \mathbb{R}$, $|c| \geq r$ implies that $x \in cA$.

Exercise 8.4.0.11. Let X be a topological vector space and $A \in \mathcal{N}(0)$. Then A is absorbing.

Proof. Let $x \in A$. For the sake of contradiction, suppose that for each $r > 0$, there exists $c \in \mathbb{R}$ such that $|c| \geq r$ and $c^{-1}x \in A^c$. Then there exists a sequence $(c_n)_{n \in \mathbb{N}} \subset \mathbb{R}$ such that for each $n \in \mathbb{N}$, $c_n \geq n$ and $c_n^{-1}x \in A^c$. Since $c_n^{-1} \rightarrow 0$, $c_n^{-1}x \rightarrow 0$. Since $A \in \mathcal{N}(0)$, $(c_n^{-1}x)_{n \in \mathbb{N}}$ is eventually in A . This is a contradiction. So there exists $r > 0$ such that for each $c \in \mathbb{R}$, $|c| \geq r$ implies that $x \in cA$. Hence A is absorbing. □

Definition 8.4.0.12. Let X be a vector space and $A \subset X$. For $x \in X$, set

$$T_x^A = \{t > 0 : x \in tA\}$$

We define the **Minkowski functional**, denoted $p_A : X \rightarrow [0, \infty]$, by

$$p_A(x) = \inf T_x^A$$

Exercise 8.4.0.13. Let X be a vector space and $A \subset X$. Suppose that A is convex, absorbing and $0_X \in A$. Then

1. $p_A : X \rightarrow [0, \infty)$
2. $p(0_X) = 0$
3. p_A is a sublinear functional on X

Proof.

1. Since A is absorbing, there exists $r > 0$ such that for each $c \in \mathbb{R}$, $|c| \geq r$ implies that $x \in cA$. Therefore $p_A(x) \leq |c|$ and $p_A : X \rightarrow [0, \infty)$.

2. Since $0 \in A$,

$$\begin{aligned} p_A(0) &= \inf T_0^A \\ &= 0 \end{aligned}$$

3. • Let $\epsilon > 0$. Choose $t_x \in T_x^A$ and $t_y \in T_y^A$ such that $t_x < p_A(x) + \epsilon/2$ and $t_y < p_A(y) + \epsilon/2$. By definition, $t_x^{-1}x, t_y^{-1}y \in A$. Set $\theta = t_x(t_x + t_y)^{-1} \in (0, 1)$. Since A is convex,

$$\begin{aligned} (t_x + t_y)^{-1}(x + y) &= (t_x + t_y)^{-1}x + (t_x + t_y)^{-1}y \\ &= \theta t_x^{-1}x + (1 - \theta)t_y^{-1}y \\ &\in A \end{aligned}$$

Therefore, $t_x + t_y \in T_{x+y}^A$ and

$$\begin{aligned} p_A(x + y) &\leq t_x + t_y \\ &< p_A(x) + \frac{\epsilon}{2} + p_A(y) + \frac{\epsilon}{2} \\ &= p_A(x) + p_A(y) + \epsilon. \end{aligned}$$

Since $\epsilon > 0$ is arbitrary, $p_A(x + y) \leq p_A(x) + p_A(y)$.

• If $\lambda = 0$, then

$$\begin{aligned} p_A(\lambda x) &= p_A(0) \\ &= 0 \\ &= |\lambda|p_A(x) \end{aligned}$$

Suppose that $\lambda > 0$. Let $t > 0$. Then

$$\begin{aligned} p_A(\lambda x) &= \inf\{t > 0 : \lambda x \in tA\} \\ &= \inf\{t > 0 : x \in \lambda^{-1}tA\} \\ &= \inf\{\lambda s > 0 : x \in sA\} \\ &= \lambda \inf\{s > 0 : x \in sA\} \\ &= \lambda p_A(x) \end{aligned}$$

So p is a sublinear functional on X . □

Exercise 8.4.0.14. Let X be a vector space and $A \subset X$. Suppose that A is convex, absorbing and $0 \in A$. Then $p_A^{-1}[0, 1) \subset A$.

Proof. Let $x \in p_A^{-1}[0, 1)$. Then $p_A(x) < 1$. By definition, there exists $t \in (0, 1)$ such that $x \in tA$. Thus $t^{-1}x \in A$. Since $0 \in A$ and A is convex, we have that

$$\begin{aligned} x &= t(t^{-1}x) + (1 - t)0 \\ &\in A \end{aligned}$$

Since $x \in p_A^{-1}[0, 1)$ is arbitrary, $p_A^{-1}[0, 1) \subset A$. □

Exercise 8.4.0.15. Let X be a topological vector space and $A \subset X$. Suppose that A is open, convex, and $0 \in A$. Then $p_A^{-1}[0, 1) = A$.

Hint: for $x \in A$, consider the sequence $(1 + 1/n)x$

Proof. Since A is open and $0 \in A$, $A \in \mathcal{N}(0)$ which implies that A is absorbing. The previous exercise implies that $p_A^{-1}[0, 1) \subset A$. Conversely, let $x \in A$. Since A is open, $A \in \mathcal{N}(x)$. Since $1 + 1/n \rightarrow 1$, $(1 + 1/n)x \rightarrow x$. Therefore, there exists $N \in \mathbb{N}$ such that for each $n \in \mathbb{N}$, $n \geq N$ implies that $(1 + 1/n)x \in A$. In particular, $x \in (1 + 1/N)^{-1}A$. Hence $(1 + 1/N)^{-1} \in T_x^A$ and

$$\begin{aligned} p_A(x) &\leq (1 + 1/N)^{-1} \\ &< 1 \end{aligned}$$

So $x \in p_A^{-1}[0, 1)$ and $A \subset p_A^{-1}[0, 1)$. □

Exercise 8.4.0.16. Let X be a topological vector space, $A \subset X$ and $x_0 \in A^c$. Suppose that A is convex, $A \in \mathcal{N}(0)$ and A is open. Then there exists $F \in X^*$ such that $\operatorname{Re} F(x_0) = 1$ and $\operatorname{Re} F|_A < 1$.

Hint: Assume X is real.

1. **Existence:** Consider a special $f \in (\mathbb{R}x_0)^*$ and use p_A to apply the Hahn-Banach theorem.

2. **Continuity:** for $\epsilon > 0$, consider the neighborhood $U_\epsilon = \epsilon A \cap -\epsilon A$

Proof. Assume that X is real.

1. Define $f \in (\mathbb{R}x_0)^*$ by $f(tx_0) = t$. Then $f(x_0) = 1$. Since $A \in \mathcal{N}(0)$, $0 \in A$ and a previous exercise implies that A is absorbing. Since A is convex, absorbing and $0 \in A$, $p_A : X \rightarrow [0, \infty)$ is a sublinear functional on X . Since $x_0 \in A^c$, the previous exercise implies that $1 \leq p_A(x_0)$. Let $x \in \mathbb{R}x_0$. Then there exists $t \in \mathbb{R}$ such that $x = tx_0$.

• If $t \geq 0$, then

$$\begin{aligned} f(x) &= t \\ &\leq tp_A(x_0) \\ &= p_A(tx_0) \\ &= p_A(x) \end{aligned}$$

• If $t < 0$, then $-t > 0$ and an exercise from the section on sublinear functionals implies that

$$\begin{aligned} f(x) &= t \\ &= < 0 \\ &\leq p_A(x) \end{aligned}$$

So $f \leq p_A$ on $\mathbb{R}x_0$. The Hahn-Banach theorem implies that there exists $F : X \rightarrow \mathbb{R}$ such that F is linear, $F|_{\mathbb{R}x_0} = f$ and $F \leq p_A$. The previous exercise implies that $p_A|_A < 1$. Hence $F|_A < 1$.

2. Let $V \in \mathcal{N}(0_{\mathbb{R}})$. Choose $\epsilon > 0$ such that $B(0, \epsilon) \subset V$. Set $U_\epsilon = \epsilon A \cap -\epsilon A$. Then $U_\epsilon \in \mathcal{N}(0)$. Let $u \in U_\epsilon$. Then $\epsilon^{-1}u, -\epsilon^{-1}u \in A$. A previous exercise implies that $p_A^{-1}([0, 1)) = A$. Hence

$$\begin{aligned} \epsilon^{-1}F(u) &= F(\epsilon^{-1}u) \\ &\leq p_A(\epsilon^{-1}u) \\ &< 1 \end{aligned}$$

So $F(u) < \epsilon$. Similarly, $F(-u) < \epsilon$. So $-\epsilon < F(u) < \epsilon$ and

$$\begin{aligned} F(U_\epsilon) &\subset B(0, \epsilon) \\ &\subset V \end{aligned}$$

Since $V \in \mathcal{N}(0_{\mathbb{R}})$ is arbitrary, F is continuous at 0. Since F is linear and F is continuous at 0, F is continuous. Hence $F \in X^*$.

If X is complex, then the previous part implies that there exists $G : X \rightarrow \mathbb{R}$ such that G is continuous, real-linear, $G(x_0) = 1$ and $G|_A < 1$. A previous exercise implies that there exists a unique $F \in X^*$ such that $\operatorname{Re} F = G$. $\square$

Exercise 8.4.0.17. Hahn-Banach Separation Theorem 1:

Let X be a topological vector space and $A, B \subset X$. Suppose that A, B are nonempty, convex and disjoint. If A is open, then there exists $\phi \in X^*$ and $c \in \mathbb{R}$ such that for each $x \in A, y \in B$,

$$\operatorname{Re} \phi(x) < c \leq \operatorname{Re} \phi(y)$$

Hint: Assume X is real.

1. Choose $a_0 \in A$ and $b_0 \in B$ and set $x_0 = b_0 - a_0$ and $C = A - B + x_0$. Then there exists $\phi \in X^*$ such that $\phi(x_0) = 1$ and $\phi|_C < 1$.

2. For each $a \in A$, $b \in B$, $\phi(a) < \phi(b)$. Set $c = \sup_{a \in A} \phi(a)$. Since ϕ is not constant, ϕ is open.

Proof. Assume X is real.

1. Since A, B are nonempty, there exist $a_0 \in A$ and $b_0 \in B$. Set $x_0 = b_0 - a_0$. Previous exercises imply that $A - B$ is open and convex. Set $C = A - B + x_0$. Then C is open and convex. Since

$$\begin{aligned} 0 &= a_0 - b_0 + x_0 \\ &\in C \end{aligned}$$

$C \in \mathcal{N}(0)$. For the sake of contradiction, suppose that $x_0 \in C$. Then there exist $a \in A$, $b \in B$ such that $x_0 = a - b + x_0$. This implies that $a = b$. This is a contradiction since $A \cap B = \emptyset$. Hence $x_0 \notin C$. The previous exercise implies that there exists a $\phi \in X^*$ such that $\phi(x_0) = 1$ and $\phi|_C < 1$.

2. Let $x \in A$ and $y \in B$. Then

$$\begin{aligned} \phi(a) - \phi(b) + 1 &= \phi(a) - \phi(b) + \phi(x_0) \\ &= \phi(a - b + x_0) \\ &< 1 \end{aligned}$$

So $\phi(a) < \phi(b)$. Set $c = \sup_{a \in A} \phi(a)$. Since A is open and $\phi \in X^*$ is open. Thus for each $x \in A$, $y \in B$,

$$\phi(x) < c \leq \phi(y)$$

If X is complex, then the previous part implies that there exists $f : X \rightarrow \mathbb{R}$ and $c \in \mathbb{R}$ such that f is continuous, real-linear and for each $x \in A$ and $y \in B$,

$$f(x) < c \leq f(y)$$

A previous exercise implies that there exists a unique $\phi \in X^*$ such that $\operatorname{Re} \phi = f$. □

Definition 8.4.0.18. Let X be a vector space and $A \subset X$. Then A is said to be an **absorbing disk** if A is convex, absorbing and balanced.

Exercise 8.4.0.19. Let X be a topological vector space and $A \in \operatorname{Conv}(X) \cap \operatorname{Bal}(X)$. If A is open, then A is an absorbing disk.

Proof. Suppose that A is open. Since $A \in \operatorname{Bal}(X)$, $0 \in A$. Since A is open, $A \in \mathcal{N}(0)$. A previous exercise implies that A is absorbing. By definition, A is an absorbing disk. □

Exercise 8.4.0.20. Let X be a vector space, $p \in \operatorname{SemiNorm}(X)$ and $r > 0$. Then $B_p(0, r)$ is an absorbing disk.

Proof.

1. Let $a, b \in B_p(0, r)$ and $t \in [0, 1]$. Then $p(a - x) < r$ and $p(b) < r$. So

$$\begin{aligned} p([ta + (1 - t)b]) &\leq p(ta + p((1 - t)b) \\ &= tp(a) + (1 - t)p(b) \\ &< tr + (1 - t)r \\ &= r \end{aligned}$$

So $ta + (1 - t)b \in B_p(0, r)$ and $B_p(0, r)$ is convex.

2. Let $a \in X$. Set $s = (p(a) + 1)/r$. Then for each $t \geq s$, $tr \geq p(a) + 1$ so that

$$\begin{aligned} a &\in B_p(0, p(a) + 1) \\ &\subset B_p(0, tr) \\ &= tB_p(0, r) \end{aligned}$$

So $B_p(0, r)$ is absorbing.

3. Let $a \in B_p(0, r)$ and $u \in \mathbb{C}$. Suppose that $|u| \leq 1$. Then

$$\begin{aligned} p(ua) &= |u|p(a) \\ &< |u|r \\ &\leq r \end{aligned}$$

So $ua \in B_p(0, r)$ and $B_p(0, r) \in \text{Bal}(X)$.

Since $B_p(0, r)$ is convex, absorbing and balanced, it is an absorbing disk. □

Exercise 8.4.0.21. Let X be a vector space and $A \subset X$. Suppose that A is an absorbing disk. Then $p_A \in \text{SemiNorm}(X)$.

Proof. Since A is an absorbing disk, A is convex, absorbing and balanced. So $0 \in A$ and the previous exercise tells us that p is a sublinear functional on X . Let $x \in X$ and $\lambda \in \mathbb{C}$.

If $\lambda = 0$, then

$$\begin{aligned} p_A(\lambda x) &= p_A(0) \\ &= 0 \\ &= |\lambda|p_A(x) \end{aligned}$$

Suppose that $\lambda \neq 0$. Since $A \in \text{Bal}(X)$, for $t > 0$, $\lambda t^{-1}x \in A$ iff $|\lambda|t^{-1}x \in A$. So

$$\begin{aligned} p_A(\lambda x) &= \inf\{t > 0 : \lambda x \in tA\} \\ &= \inf\{t > 0 : x \in |\lambda|^{-1}tA\} \\ &= \inf\{|\lambda|s > 0 : x \in sA\} \\ &= |\lambda| \inf\{s > 0 : x \in sA\} \\ &= |\lambda|p_A(x) \end{aligned}$$

So p is a seminorm on X . □

Exercise 8.4.0.22. Let X be a topological vector space and $A \subset X$. Suppose that A is an absorbing disk and A is open. Then $B_{p_A}(0, 1) = A$.

Proof. Clear by previous exercise. □

Exercise 8.4.0.23. Let X be a topological vector space and $A \subset X$. Suppose that A is an absorbing disk. Then p_A is continuous iff A is open.

Proof. If A is open, then

$$\begin{aligned} A &= B_{p_A}(0, 1) \\ &\subset \bar{B}_{p_A}(0, 1) \end{aligned}$$

which implies that $\bar{B}_{p_A}(0, 1) \in \mathcal{N}(0)$. An exercise in the previous section implies that p_A is continuous. Conversely, if p_A is continuous, then an exercise in the previous section implies that $B_{p_A}(0, 1)$ is open. □

Exercise 8.4.0.24. Let $(X, \mathcal{T})$ be a topological vector space. Then for each $A \in \mathcal{T} \cap \text{Bal}(X) \cap \text{Conv}(X)$, A is an absorbing disk and p_A is continuous and $B_{p_A}(0, 1) = A$.

Proof. **FINISH!!!** clear by prev exercises □

8.5 Locally Convex Spaces

- **TO DO:**
- show that $\mathcal{P} \sim \{|\phi| : \phi \in X^*\}$, **FINISH!!!!**

Definition 8.5.0.1. Let X be a vector space and $\mathcal{P} \subset \text{SemiNorm}(X)$. We define the **seminorm topology on X induced by $\mathcal{P}$** , denoted $\mathcal{T}_{\mathcal{P}} \subset \mathcal{P}(X)$, by $\mathcal{T}_{\mathcal{P}} := \tau_X(B_p(x, r) : p \in \mathcal{P}, x \in X \text{ and } r > 0)$.

Exercise 8.5.0.2. Let X be a vector space and $\mathcal{P} \subset \text{SemiNorm}(X)$. Then $\mathcal{T}_{\mathcal{P}} = \tau_X(\pi_p : p \in \mathcal{P})$ **FINISH!!!!**.

Proof. Set $\mathcal{T} := \tau_X(\pi_p : p \in \mathcal{P})$.

- For each $p \in \mathcal{P}$, $x \in X$ and $r > 0$,

$$\begin{aligned} B_p(x, r) &= \pi_p^{-1}(B_{\bar{p}}(0, r)) \quad (\text{ref and make ex for this and later}) \\ &\in \tau_X(\pi_{p'} : p' \in \mathcal{P}) \\ &= \mathcal{T}. \end{aligned}$$

Therefore $\{B_p(x, r) : p \in \mathcal{P}, x \in X \text{ and } r > 0\} \subset \mathcal{T}$. Hence

$$\begin{aligned} \mathcal{T}_{\mathcal{P}} &= \tau_X(B_p(x, r) : p \in \mathcal{P}, x \in X \text{ and } r > 0) \\ &\subset \mathcal{T}. \end{aligned}$$

- Let $p \in \mathcal{P}$, $a \in X/\ker p$ and $r > 0$. Then there exists $x \in X$ such that $a = x + \ker p$. Therefore (MAKE EX showing that for $a \in X/\ker p$, $V \subset X/\ker p$, $\pi_p^{-1}(a+rV) = a + \bigcup_{b \in V} rb$ and $\pi^{-1}(\{a+b\}) = a+b$), (MAKE EX SHOWING $\pi_p^{-1}(B_{\bar{p}}(0, 1)) = B_p(0, 1)$) **FINISH!!!!**

$$\begin{aligned} \pi_p^{-1}(B_{\bar{p}}(a, r)) &= \pi_p^{-1}(a + rB_{\bar{p}}(0_{X/\ker p}, 1)) \\ &= \pi^{-1}(\{a\}) + r\pi^{-1}(B_{\bar{p}}(0_{X/\ker p}, 1)) \\ &= a + rB_p(0, 1) \\ &= \bigcup_{x \in a} x + rB_p(0, 1) \\ &= \bigcup_{x \in a} B_p(x, r) \\ &\in \tau_X(B_{p'}(x, r) : p' \in \mathcal{P}, x \in X \text{ and } r > 0) \\ &= \mathcal{T}_{\mathcal{P}}. \end{aligned}$$

Since $p \in \mathcal{P}$, $a \in X/\ker p$ and $r > 0$ are arbitrary, we have that for each $p \in \mathcal{P}$, $a \in X/\ker p$ and $r > 0$, $\pi_p^{-1}(B_{\bar{p}}(a, r)) \in \mathcal{T}_{\mathcal{P}}$. Thus $\{\pi_p^{-1}(B_{\bar{p}}(a, r)) : p' \in \mathcal{P}, x \in X \text{ and } r > 0\} \subset \mathcal{T}_{\mathcal{P}}$. Since for each $p \in \mathcal{P}$, $\mathcal{T}_p = \tau_{X/\ker p}(B_{\bar{p}}(a, r) : a \in X/\ker p \text{ and } r > 0)$, **ref ex on weak topology** implies that

$$\begin{aligned} \mathcal{T} &= \tau_X(\pi_p : p \in \mathcal{P}) \\ &= \tau_X(\pi_p^{-1}(B_{\bar{p}}(a, r)) : p \in \mathcal{P}, a \in X/\ker p, \text{ and } r > 0) \quad \text{ref ex here about weak topology} \\ &\subset \mathcal{T}_{\mathcal{P}}. \end{aligned}$$

Since $\mathcal{T}_{\mathcal{P}} \subset \mathcal{T}$ and $\mathcal{T} \subset \mathcal{T}_{\mathcal{P}}$, we have that $\mathcal{T}_{\mathcal{P}} = \mathcal{T}$. □

Exercise 8.5.0.3. Let X be a vector space and $\mathcal{P} \subset \text{SemiNorm}(X)$. If $\mathcal{P} \neq \{0\}$, then $\mathcal{T}_{\mathcal{P}} \neq \tau_X(\mathcal{P})$.

Proof. Suppose that $\mathcal{P} \neq \{0\}$. Then there exists $p \in \mathcal{P}$ such that $p \neq 0$. Thus $\ker p \neq X$. **ref ex in seminorm section** implies that $\ker p_0 \not\sim_{p_0}$. Thus there exists $x, y \in X$ such that $x \sim_{p_0} y$ such that $x \not\sim_{\ker p_0} y$. Therefore **FINISH!!!!, FIX!!!** □

Note 8.5.0.4. It appears that a decent number of sources define $\mathcal{T}_{\mathcal{P}}$ as the weak topology on X induced by $\mathcal{P}$. The previous exercise shows that this is not the case. **ref ex that $\ker p \neq \sim_p$, so that $x_\alpha \rightarrow a$**

Exercise 8.5.0.5. Let X be a vector space, $\mathcal{P} \subset \text{SemiNorm}(X)$, $(x_\alpha)_{\alpha \in A} \subset X$ a net and $x \in X$. Then $x_\alpha \rightarrow x$ in $(X, \mathcal{T}_{\mathcal{P}})$ iff for each $p \in \mathcal{P}$, $p(x_\alpha - x) \rightarrow 0$.

Proof. We note that **(turn this into ex and ref here)**

$$\begin{aligned} p(x_\alpha - x) \rightarrow 0 &\iff \bar{p} \circ \pi_p(x_\alpha - x) \rightarrow 0 \\ &\iff \bar{p}(\pi_p(x_\alpha - x)) \rightarrow 0 \\ &\iff \bar{p}(\pi_p(x_\alpha) - \pi_p(x)) \rightarrow 0 \\ &\iff \pi_p(x_\alpha) \xrightarrow{\mathcal{T}_{\bar{p}}} \pi_p(x). \end{aligned}$$

Thus

$$\begin{aligned} \text{for each } p \in \mathcal{P}, p(x_\alpha - x) \rightarrow 0 &\iff \text{for each } p \in \mathcal{P}, \pi_p(x_\alpha) \xrightarrow{\mathcal{T}_{\bar{p}}} \pi_p(x) \quad (\text{by prev ex}) \\ &\iff \text{for each } p \in \mathcal{P}, p(x_\alpha - x) \rightarrow 0. \end{aligned}$$

□

Exercise 8.5.0.6. Let X be a vector space and $\mathcal{P} \subset \text{SemiNorm}(X)$. Then $(X, \mathcal{T}_{\mathcal{P}})$ is a topological vector space.

Proof. Let $(x_\alpha)_{\alpha \in A}, (y_\alpha)_{\alpha \in A} \subset X$, $(\lambda_\alpha)_{\alpha \in A} \subset \mathbb{K}$ be nets, $x, y \in X$ and $\lambda \in \mathbb{K}$. Suppose that $x_\alpha \xrightarrow{\mathcal{T}_{\mathcal{P}}} x$, $y_\alpha \xrightarrow{\mathcal{T}_{\mathcal{P}}} y$ and $\lambda_\alpha \rightarrow \lambda$. Let $p \in \mathcal{P}$. **A prev ex** implies that $p([x_\alpha - x] + \lambda_\alpha[y_\alpha - y]) \rightarrow 0$ and $p(y_\alpha - y) \rightarrow 0$. Therefore

$$\begin{aligned} p([x_\alpha + \lambda_\alpha y_\alpha] - [x + \lambda y]) &= p([x_\alpha - x] + \lambda_\alpha[y_\alpha - y]) \\ &\leq p([x_\alpha - x] + |\lambda_\alpha|p(y_\alpha - y)) \\ &\rightarrow 0. \end{aligned}$$

Since $p \in \mathcal{P}$ is arbitrary, we have that for each $p \in \mathcal{P}$, $p([x_\alpha + \lambda_\alpha y_\alpha] - [x + \lambda y]) \rightarrow 0$. **The same prev ex** then implies that $x_\alpha + \lambda_\alpha y_\alpha \rightarrow x + \lambda y$. Since $(x_\alpha)_{\alpha \in A}, (y_\alpha)_{\alpha \in A} \subset X$, $(\lambda_\alpha)_{\alpha \in A} \subset \mathbb{K}$ nets, $x, y \in X$ and $\lambda \in \mathbb{K}$ with $x_\alpha \xrightarrow{\mathcal{T}_{\mathcal{P}}} x$, $y_\alpha \xrightarrow{\mathcal{T}_{\mathcal{P}}} y$ and $\lambda_\alpha \rightarrow \lambda$ are arbitrary, we have that for each $(x_\alpha)_{\alpha \in A}, (y_\alpha)_{\alpha \in A} \subset X$, $(\lambda_\alpha)_{\alpha \in A} \subset \mathbb{K}$ nets, $x, y \in X$ and $\lambda \in \mathbb{K}$, $x_\alpha \xrightarrow{\mathcal{T}_{\mathcal{P}}} x$, $y_\alpha \xrightarrow{\mathcal{T}_{\mathcal{P}}} y$ and $\lambda_\alpha \rightarrow \lambda$ implies that $x_\alpha + \lambda_\alpha y_\alpha \rightarrow x + \lambda y$. **ref/make prev ex in intro section** implies that $(X, \mathcal{T})$ is a topological vector space. □

Exercise 8.5.0.7. Let X be a vector space and $\mathcal{P} \subset \text{SemiNorm}(X)$.

1. Define $\mathcal{B}_0 \subset \mathcal{T}_{\mathcal{P}}$ by

$$\mathcal{B}_0 := \left\{ \bigcap_{p \in P} B_p(0, r) : P \subset \mathcal{P}, \#P < \infty \text{ and } r > 0 \right\}.$$

Then

- (a) $\mathcal{B} \subset \text{Bal}(X) \cap \text{Conv}(X)$
- (b) $\mathcal{B}_0$ is a local basis for $\mathcal{T}$ at 0.

2. Define $\mathcal{B} \subset \mathcal{T}_{\mathcal{P}}$ by

$$\mathcal{B} := \left\{ \bigcap_{p \in P} B_p(x, r) : P \subset \mathcal{P}, \#P < \infty, x \in X \text{ and } r > 0 \right\}.$$

Then

- (a) $\mathcal{B} \subset \text{Bal}(X) \cap \text{Conv}(X)$
- (b) $\mathcal{B}$ is a basis for $\mathcal{T}_{\mathcal{P}}$.

Proof.

1. ref topology section , (similar to prev part) ref exercises in prev section about convec,balanced sets
2. ref/make ex about local bases determining bases in topological vector spaces, then ref prev ex about seminorm topology yielding a tvs

□

Exercise 8.5.0.8. Let $(X, \mathcal{T})$ be topological vector space. Then there exists $\mathcal{P} \subset \text{SemiNorm}(X, \mathcal{T})$ such that $\mathcal{T} = \mathcal{T}_{\mathcal{P}}$ iff there exists $\mathcal{U}_0 \subset \mathcal{T} \cap \text{Bal}(X) \cap \text{Conv}(X)$ such that $\mathcal{U}_0$ is a local basis for $\mathcal{T}$ at 0.

Proof.

- ($\Rightarrow$):
Suppose that there exists $\mathcal{P} \subset \text{SemiNorm}(X, \mathcal{T})$ such that $\mathcal{T} = \mathcal{T}_{\mathcal{P}}$. Exercise 8.5.0.7 implies that there exists $\mathcal{U}_0 \subset \mathcal{T} \cap \text{Bal}(X) \cap \text{Conv}(X)$ such that $\mathcal{U}_0$ is a local basis for $\mathcal{T}$ at 0.
- ($\Leftarrow$):
Suppose that there exists $\mathcal{U}_0 \subset \mathcal{T} \cap \text{Bal}(X) \cap \text{Conv}(X)$ such that $\mathcal{U}_0$ is a local basis for $\mathcal{T}$ at 0. An ex in prev section about minkowski functionals implies that for each $U \in \mathcal{U}_0$, $p_U \in \text{SemiNorm}(X, \mathcal{T})$ and $B_{p_U}(0, 1) = U$. Define $\mathcal{P} \subset \text{SemiNorm}(X, \mathcal{T})$ by $\mathcal{P} := \{p_U : U \in \mathcal{U}_0\}$. Since $\mathcal{P} \subset \text{SemiNorm}(X, \mathcal{T})$, make/ref ex implies that $\mathcal{T}_{\mathcal{P}} \subset \mathcal{T}$. Since for each $U \in \mathcal{U}_0$,

$$\begin{aligned} U &= B_{p_U}(0, 1) \\ &\in \mathcal{T}_{\mathcal{P}}, \end{aligned}$$

we have that

$$\begin{aligned} \mathcal{U}_0 &\subset \mathcal{T}_{\mathcal{P}} \\ &\subset \mathcal{T}. \end{aligned}$$

Since $\mathcal{U}_0$ is a local basis for $\mathcal{T}$ at 0, make/ref ex in topology chapter about if $\mathcal{U}_0$ is local basis for $\mathcal{T}_2$ at x and $\mathcal{U}_0 \subset \mathcal{T}_1 \subset \mathcal{T}_2$, then $\mathcal{U}_0$ is a local basis for $\mathcal{T}_1$ at x implies that $\mathcal{U}_0$ is a local basis for $\mathcal{T}_{\mathcal{P}}$ at 0. ref ex about local bases for tvs implies that

$$\begin{aligned} \mathcal{T}_{\mathcal{P}} &= \tau_X(\{x + U : U \in \mathcal{U}_0 \text{ and } x \in X\}) \\ &= \mathcal{T}. \end{aligned}$$

□

Definition 8.5.0.9. Let X be a vector space and $\mathcal{P} \in \text{SemiNorm}(X)$. Then $\mathcal{P}$ is said to **separate points of X** if for each $x \in X$, if $x \neq 0$, then there exists $p \in \mathcal{P}$ such that $p(x) \neq 0$.

Exercise 8.5.0.10. Let X be a vector space and $\mathcal{P} \subset \text{SemiNorm}(X)$. Then $(X, \mathcal{T}_{\mathcal{P}})$ is Hausdorff iff $\mathcal{P}$ separates the points of X .

Proof.

- ($\Rightarrow$):
Suppose that $(X, \mathcal{T}_{\mathcal{P}})$ is Hausdorff. Let $x \in X$. Suppose that $x \neq 0$. Then there exists $U, V \in \mathcal{T}$ such that $0 \in U$, $x \in V$ and $U \cap V = \emptyset$. Exercise 8.5.0.7 implies that there exist $P_U, P_V \subset \mathcal{P}$, $r_U, r_V \subset$ such that $\#P_U, \#P_V < \infty$ and $\bigcap_{p \in P_U} B_p(0, r_U) \subset U$ and $\bigcap_{p \in P_V} B_p(x, r_V) \subset V$. For the sake of contradiction, suppose that for each $p \in \mathcal{P}$, $p(x) = 0$. Then for each $p \in P_U$, $x \in B_p(0, r_U)$. Thus

$$\begin{aligned} x &\in \bigcap_{p \in P_U} B_p(0, r_U) \\ &\subset U. \end{aligned}$$

Since $x \in V$,

$$\begin{aligned} x &\in U \cap V \\ &= \emptyset, \end{aligned}$$

which is a contradiction. Therefore there exists $p \in \mathcal{P}$ such that $p(x) \neq 0$. Hence $\mathcal{P}$ separates the points of X .

- ($\Leftarrow$):

Suppose that $\mathcal{P}$ separates the points of X . Let $x, y \in X$. Suppose that $x \neq y$. Then $x - y \neq 0$. Thus there exists $p \in \mathcal{P}$ such that $p(x - y) \neq 0$. **make and ref ex in pseudometric space section that if $d(x, y) > 0$, then x, y live in disjoint open balls. FINISH!!!**

□

Definition 8.5.0.11. Let X be vector space and $\mathcal{P}_1, \mathcal{P}_2 \subset \text{SemiNorm}(X)$. Then $\mathcal{P}_1$ is said to be **topologically equivalent to $\mathcal{P}_2$** , denoted $\mathcal{P}_1 \sim_{\text{Top}} \mathcal{P}_2$, if $\mathcal{T}_{\mathcal{P}_1} = \mathcal{T}_{\mathcal{P}_2}$.

Exercise 8.5.0.12. **rework, use previous exercise,** Let X be vector space, $\mathcal{P} \subset \text{SemiNorm}(X)$ and $\mathcal{U}_0 \subset \mathcal{T}_{\mathcal{P}} \cap \text{Bal}(X) \cap \text{Conv}(X)$. Suppose that $\mathcal{U}_0$ is a local basis for $\mathcal{T}_{\mathcal{P}}$ at 0. Define $\mathcal{P}' \subset \text{SemiNorm}(X, \mathcal{T}_{\mathcal{P}})$ by $\mathcal{P}' := \{p_U : U \in \mathcal{U}_0\}$. Then $\mathcal{P}' \sim_{\text{Top}} \mathcal{P}$.

Proof. Can we just copy the idea from the prev ex about getting $\mathcal{U}_0 \subset \mathcal{T}' \subset \mathcal{T}$ such that $\mathcal{U}_0$ is a local basis of $\mathcal{T}$ implying $\mathcal{U}_0$ is a local basis for $\mathcal{T}'$ and then $\mathcal{T}' = \mathcal{T}$? This should work for general tvs, and we could ref that here as a consequence in the lcs case □

Proof. An ex in prev section about minkowski functionals implies that for each $U \in \mathcal{U}_0$ and $p_U \in \text{SemiNorm}(X, \mathcal{T}_{\mathcal{P}})$ and $B_{p_U}(0, 1) = U$.

- Define $\mathcal{B}' \subset \mathcal{T}_{\mathcal{P}'}$ by $\mathcal{B}' := \{B_p(x, r) : p \in \mathcal{P}', x \in X \text{ and } r > 0\}$. By definition (ref def of $\mathcal{T}_{\mathcal{P}}$), $\mathcal{T}_{\mathcal{P}'} = \tau_X(\mathcal{B}')$. ref prev exercises in tvs intro section implies that

$$\begin{aligned} \mathcal{B}' &= \{B_p(x, r) : p \in \mathcal{P}', x \in X \text{ and } r > 0\} \\ &= \{x + rB_p(0, 1) : p \in \mathcal{P}', x \in X \text{ and } r > 0\} \\ &= \{x + rB_{p_U}(0, 1) : U \in \mathcal{U}_0, x \in X \text{ and } r > 0\} \\ &= \{x + rU : U \in \mathcal{U}_0, x \in X \text{ and } r > 0\} \\ &\subset \mathcal{T}_{\mathcal{P}} \quad (\text{ref ex about local bases in tvs}). \end{aligned}$$

In particular,

$$\begin{aligned} \mathcal{T}_{\mathcal{P}'} &= \tau_X(\mathcal{B}') \\ &\subset \mathcal{T}_{\mathcal{P}}. \end{aligned}$$

- Define $\mathcal{B} \subset \mathcal{T}_{\mathcal{P}}$ by $\mathcal{B} := \{x + U : U \in \mathcal{U}_0 : x \in X\}$. Since $\mathcal{U}_0$ is a local basis for $\mathcal{T}_{\mathcal{P}}$, (make) an exercise in tvs intro section implies that $\mathcal{B}$ is a basis for $\mathcal{T}_{\mathcal{P}}$ and $\mathcal{T}_{\mathcal{P}} = \tau(\mathcal{B})$. Since

$$\begin{aligned} \mathcal{B} &= \{x + U : U \in \mathcal{U}_0 : x \in X\} \\ &\subset \{x + rU : U \in \mathcal{U}_0, x \in X \text{ and } r > 0\} \\ &= \mathcal{B}', \end{aligned}$$

we have that

$$\begin{aligned} \mathcal{T}_{\mathcal{P}} &= \tau_X(\mathcal{B}) \\ &\subset \tau_X(\mathcal{B}') \\ &= \mathcal{T}_{\mathcal{P}'}. \end{aligned}$$

Since $\mathcal{T}_{\mathcal{P}'} \subset \mathcal{T}_{\mathcal{P}}$ and $\mathcal{T}_{\mathcal{P}} \subset \mathcal{T}_{\mathcal{P}'}$, we have that $\mathcal{T}_{\mathcal{P}'} = \mathcal{T}_{\mathcal{P}}$. Tus $\mathcal{P}' \sim_{\text{Top}} \mathcal{P}$. □

Exercise 8.5.0.13. Characterize $\mathcal{T}_{\mathcal{P}}$

Proof. □

Definition 8.5.0.14. Let $(X, \mathcal{T})$ be a topological vector space.

- Then $\mathcal{T}$ is said to be **locally convex** if
 1. $(X, \mathcal{T})$ is Hausdorff
 2. there exists $\mathcal{U}_0 \subset \mathcal{T} \cap \text{Bal}(X) \cap \text{Conv}(X)$ such that $\mathcal{U}_0$ is a local basis for $\mathcal{T}$ at 0.

- We define the **family of continuous seminorms on X associated to $\mathcal{T}$** , denoted $\mathcal{P}_{\mathcal{T}} \subset \text{SemiNorm}(X, \mathcal{T})$, by $\mathcal{P}_{\mathcal{T}} := \{p_A : A \in \mathcal{T} \cap \text{Bal}(X) \cap \text{Conv}(X)\}$. **rework this, make ex showing that if $U = B_p(0, 1)$, then $P_U = p$, then show that $\mathcal{T}_{\mathcal{T}}$ is the largest family of continuous seminorms, then call $\mathcal{P}_{\mathcal{T}}$ the 'largest' or 'maximal' family of continuous seminorms on X associated to (or induced by) $\mathcal{T}$.**

Note 8.5.0.15.

- When the context is clear we may say ‘locally convex space’ instead of ‘locally convex topological vector space’.
- It is not a universal convention to require the topology to be Hausdorff, so it is important to check the definition when consulting other sources.

Exercise 8.5.0.16. Let X be a vector space, $(Y_{\alpha}, \mathcal{T}_{\alpha})_{\alpha \in A}$ a collection of locally convex spaces and $(\phi_{\alpha})_{\alpha \in A} \in \prod_{\alpha \in A} \mathcal{L}(X, Y_{\alpha})$. Set $\mathcal{T} := \sigma_X(\phi_{\alpha} : \alpha \in A)$. Then $(X, \mathcal{T})$ is a locally convex space.

Proof. Since for each $\alpha \in A$, $(Y_{\alpha}, \mathcal{T}_{\alpha})$ is locally convex, **ref ex** implies that for each $\alpha \in A$, there exists a $\mathcal{P}_{\alpha} \subset \text{SemiNorm}(X, \mathcal{T}_{\alpha})$ on Y_{α} such that $\mathcal{T}_{\alpha} = \mathcal{T}_{\mathcal{P}_{\alpha}}$. Exercise 8.3.0.8 implies that for each $\alpha \in A$ and $q \in \mathcal{P}_{\alpha}$, $q \circ \phi_{\alpha} \in \text{SemiNorm}(X)$. Define $\mathcal{P} \subset \text{SemiNorm}(X)$ by $\mathcal{P} := \{q \circ \phi_{\alpha} : \alpha \in A \text{ and } q \in \mathcal{P}_{\alpha}\}$. Define $\mathcal{T}' \subset \mathcal{P}(X)$ by $\mathcal{T}' := \mathcal{T}_{\mathcal{P}}$.

- Let $p \in \mathcal{P}$. Then there exists $\alpha \in A$ and $q \in \mathcal{P}_{\alpha}$ such that $p = q \circ \phi_{\alpha}$. Since $\mathcal{T} = \sigma_X(\phi_{\alpha} : \alpha \in A)$, we have that ϕ_{α} is $(\mathcal{T}, \mathcal{T}_{\alpha})$ -continuous. Since $\mathcal{P}_{\alpha} \subset \text{SemiNorm}(Y_{\alpha}, \mathcal{T}_{\alpha})$, we have that

$$\begin{aligned} p &= q \circ \phi_{\alpha} \\ &\in \text{SemiNorm}(X, \mathcal{T}). \end{aligned}$$

Since $p \in \mathcal{P}$ is arbitrary, we have that for each $p \in \mathcal{P}$, $p \in \text{SemiNorm}(X, \mathcal{T})$. Thus $\mathcal{P} \subset \text{SemiNorm}(X, \mathcal{T})$. **ref/make ex** implies that $\mathcal{T}_{\mathcal{P}} \subset \mathcal{T}$.

- **FIX!!! REWORK!!!!** Let $(x_{\gamma})_{\gamma \in \Gamma} \subset X$ be a net and $x \in X$. Suppose that $x_{\gamma} \rightarrow x$ in $(X, \mathcal{T})$. **need exercise showing $(X, \mathcal{T})$ is a topological vector space.** Since $(X, \mathcal{T})$ is a topological vector space, $x_{\gamma} - x \rightarrow 0$ in $(X, \mathcal{T})$. Suppose that $x_{\gamma} \rightarrow x$ in $(X, \mathcal{T}')$. Let $\alpha \in A$ and $q \in \mathcal{P}_{\alpha}$. Since $q \circ \phi_{\alpha} \in \mathcal{P}$, Exercise 8.5.0.5 implies that

$$\begin{aligned} q(\phi_{\alpha}(x_{\gamma}) - \phi_{\alpha}(x)) &= q \circ \phi_{\alpha}(x_{\gamma} - x) \\ &\rightarrow 0. \end{aligned}$$

Since $q \in \mathcal{P}_{\alpha}$ is arbitrary, we have that for each $q \in \mathcal{P}_{\alpha}$, $q(\phi_{\alpha}(x_{\gamma}) - \phi_{\alpha}(x)) \rightarrow 0$. Exercise 8.5.0.5 then implies that $\phi_{\alpha}(x_{\gamma}) \rightarrow \phi_{\alpha}(x)$ in $(Y_{\alpha}, \mathcal{T}_{\alpha})$. Since $\alpha \in A$ is arbitrary, we have that for each $\alpha \in A$, $\phi_{\alpha}(x_{\gamma}) \rightarrow \phi_{\alpha}(x)$ in $(Y_{\alpha}, \mathcal{T}_{\alpha})$. Exercise 4.3.2.15 implies that $x_{\gamma} \rightarrow x$ in $(X, \mathcal{T})$.

Thus $x_{\gamma} \rightarrow x$ in $(X, \mathcal{T}')$ iff $x_{\gamma} \rightarrow x$ in $(X, \mathcal{T})$. Exercise 4.3.2.16 then implies that $\mathcal{T} = \mathcal{T}'$. □

8.6 Continuous Linear Maps

redo in terms of "boundedness", need to define bounded subsets, then continuous maps should send bounded sets to bounded sets

Definition 8.6.0.1. Let X, Y be topological vector spaces. We define

$$L(X; Y) = \{T : X \rightarrow Y : T \text{ is linear and continuous}\}$$

8.7 Subspaces

Exercise 8.7.0.1. Let $(X, \mathcal{T})$ be a locally convex space with associated family of seminorms $\mathcal{P}$ and $M \subset X$ a subspace. Define $\mathcal{P}_M = \{p|_M : p \in \mathcal{P}\}$. Then $(M, \mathcal{T} \cap M)$ is a locally convex space with associated family of seminorms $\mathcal{P}_M$.

Proof. Let $(x_\alpha)_{\alpha \in A} \subset M$ be a net and $x \in M$. Suppose that $x_\alpha \rightarrow x$ in $\mathcal{T} \cap M$. Then an exercise in the section on the subspace topology implies that $x_\alpha \rightarrow x$ in $\mathcal{T}$. Let $q \in \mathcal{P}_M$. Then there exists $p \in \mathcal{P}$ such that $q = p|_M$. Therefore

$$\begin{aligned} q(x_\alpha - x) &= p|_M(x_\alpha - x) \\ &= p(x_\alpha - x) \\ &\rightarrow 0 \end{aligned}$$

Hence $x_\alpha \rightarrow x$ in $\tau_X(\pi_q : q \in \mathcal{P}_M)$.

Conversely, suppose that $x_\alpha \rightarrow x$ in $\tau_X(\pi_q : q \in \mathcal{P}_M)$. Let $p \in \mathcal{P}$. Then

$$\begin{aligned} p(x_\alpha - x) &= p|_M(x_\alpha - x) \\ &\rightarrow 0 \end{aligned}$$

Hence $x_\alpha \rightarrow x$ in $\mathcal{T}$. So $x_\alpha \rightarrow x$ in $\mathcal{T} \cap M$. Therefore $\mathcal{T} \cap M = \tau_X(\pi_q : q \in \mathcal{P}_M)$. □

Exercise 8.7.0.2. Let X be a locally convex space, $M \subset X$ a subspace and $f \in M^*$. Then there exists $F \in X^*$ such that $F|_M = f$.

Proof. Set $p_f = |f|$. Since p_f is a continuous seminorm, $B_{p_f}(0, 1)$ is open in M . Therefore, there exists $U \subset X$ open such that $B_{p_f}(0, 1) = U \cap M$. A previous exercise implies that there exists $p \in \mathcal{P}$ and $r > 0$ such that $B_p(0, r) \subset U$. Set $A = B_p(0, r)$. Since A is open, $p_A : X \rightarrow [0, \infty)$ is continuous and $A = B_{p_A}(0, 1)$. Hence

$$\begin{aligned} B_{p_A|_M}(0, 1) &= A \cap M \subset U \cap M \\ &= B_{p_f}(0, 1) \end{aligned}$$

Therefore $p_f \leq p_A|_M$ and $|f| \leq p_A$ on M . The Hahn-Banach theorem implies that there exists $F : X \rightarrow \mathbb{C}$ such that F is linear, $F|_M = f$ and $|F| \leq p_A$. Since p_A is continuous, $|F|$ is continuous, which implies that F is continuous. So $F \in X^*$. □

Exercise 8.7.0.3. Hahn-Banach Separation Theorem 2:

Let X be a locally convex space and $A, B \subset X$. Suppose that A, B are nonempty, convex and disjoint. If A is compact and B is closed, then there exists $\phi \in X^*$ and $c_1, c_2 \in \mathbb{R}$ such that for each $x \in A, y \in B$,

$$\operatorname{Re} \phi(x) < c_1 < c_2 \leq \operatorname{Re} \phi(y)$$

Hint: Assume X is real. Since X is locally convex, there exists $V \subset U$ such that V is an open absorbing disk and $(A + V) \cap B = \emptyset$. Then apply the first Hahn-Banach separation theorem to $A + V$ and B .

Proof. Assume X is real. Suppose that A is compact and B is closed. A previous exercise implies that there exists $U \in \mathcal{N}(0)$ such that U is open and $(A + U) \cap B = \emptyset$. Since X is locally convex, there exists $V \subset U$ such that V is an open absorbing disk. Then $(A + V)$ is open and convex. By the first Hahn-Banach separation theorem, there exist $\phi \in X^*$ and $c_2 \in \mathbb{R}$ such that for each $x \in A + V, y \in B$,

$$\phi(x) < c_2 \leq \phi(y)$$

Specifically, $c_2 = \sup_{x \in A+V} \phi(x)$. Since $\phi \in X^*$ is not constant, ϕ is open and thus $\phi(A + V)$ is open. Continuity of ϕ implies that $\phi(A)$ is compact. Therefore, $\sup \phi(A) < \sup \phi(A + V)$. So there exists $c_1 \in \phi(A + V)$ such that $\sup \phi(A) < c_1$. Hence there exists $x_1 \in A + V$ such that $\phi(x_1) = c_1$. Then for each $x \in A$ and $y \in B$,

$$\begin{aligned} \phi(x) &\leq \sup \phi(A) \\ &< c_1 \\ &= \phi(x_1) \\ &< c_2 \\ &\leq \phi(y) \end{aligned}$$

If X is complex, then the previous part implies that there exists $f : X \rightarrow \mathbb{R}$ and $c_1, c_2 \in \mathbb{R}$ such that f is continuous, real-linear and for each $x \in A$ and $y \in B$,

$$f(x) < c_1 < c_2 \leq f(y)$$

A previous exercise implies that there exists a unique $\phi \in X^*$ such that $\operatorname{Re} \phi = f$. □

Exercise 8.7.0.4. Let X be a locally convex space and $M \subset X$ a closed subspace. If $M \neq X$, then there exists $\phi \in X^*$ such that $\phi \neq 0$ and $\phi|_M = 0$.

Proof. Assume that X is real. Suppose that $M \neq X$. Then there exists $x_0 \in X$ such that $x_0 \notin M$. Since $\{x_0\}$ is compact and convex, M is closed and convex and $\{x_0\} \cap M = \emptyset$, the second Hahn-Banach separation theorem implies that there exists $\phi \in X^*$ such that for each $x \in M$,

$$\phi(x_0) < \phi(x)$$

Since $0 \in M$,

$$\begin{aligned} \phi(x_0) &< \phi(0) \\ &= 0 \end{aligned}$$

so that $\phi \neq 0$. For the sake of contradiction, suppose that $\phi|_M \neq 0$. Then there exists $x_1 \in M$ such that $\phi(x_1) \neq 0$. Then for each $t \in \mathbb{R}$,

$$\begin{aligned} \phi(x_0) &< \phi(tx_1) \\ &= t\phi(x_1) \end{aligned}$$

Set $t = \frac{\phi(x_0)}{\phi(x_1)}$. Then

$$\begin{aligned} \phi(x_0) &< t\phi(x_1) \\ &= \phi(x_0) \end{aligned}$$

which is a contradiction. So $\phi|_M = 0$. □

Exercise 8.7.0.5. Let X be a locally convex space. Then X^* separates the points of X .

Proof. Let $x, y \in X$. The second Hahn-Banach separation theorem implies that there exists $\phi \in X^*$ such that $\phi(x) \neq \phi(y)$. □

8.8 Products

8.9 Direct Sums

8.10 Quotient Spaces

Note 8.10.0.1. We recall the following (ref linear algebra notes):

- Let X be a vector space. Then
 - the **subspace relation on X induced by M** , denoted $\sim_M \subset X^2$, by $\sim_M := \{(x_1, x_2) \in X^2 : x_1 - x_2 \in M\}$,
 - $\sim_M \in \text{Equiv}(X)$ and $X/M = \{x + M : x \in X\}$ and X/M is a vector space,
 - **quotient map of X onto X/M** , denoted $\pi_M : X \rightarrow X/M$, by $\pi_M(x) := x + M$.
 - $\pi_M \in \mathcal{L}(X, X/M)$,
- Let X, Y, Z be vector spaces. Then
 -
 - for each $T \in \mathcal{L}(X, Y)$, $M \subset \ker T$ iff T is $\sim_M$ -invariant,
 - for each $f : Y \rightarrow Z$ and $T \in \mathcal{L}(X, Y)$, S is $\sim_T$ -invariant iff S is $\sim_{\ker T}$ -invariant.
 - for each $f : Y \rightarrow Z$ and $T \in \mathcal{L}(X, Y)$, f is $\sim_T$ -invariant implies that there exists a unique $\bar{f} : \text{Img } T \rightarrow Z$ such that $\bar{f} \circ T = f$.
 -

Exercise 8.10.0.2. Let $(X, \mathcal{T})$ be a topological vector space and $M \subset X$ a subspace. Then $(X/M, \mathcal{T}_{X/M})$ is a topological vector space.

Proof. Denote addition on X and X/M by $A : X^2 \rightarrow X$ and $\bar{A} : (X/M)^2 \rightarrow X/M$ respectively. Similarly, denote scalar multiplication on X and X/M by $\Lambda : \mathbb{C} \times X \rightarrow X$ and $\bar{\Lambda} : \mathbb{C} \times (X/M) \rightarrow X/M$ respectively.

- Let $\bar{x}, \bar{y} \in X/M$. Let $U \in \mathcal{N}(\bar{x} + \bar{y})$. Since $\pi : X \rightarrow X/M$ is continuous, we have that $\pi^{-1}(U) \in \mathcal{N}(x + y)$. Since addition $A : X^2 \rightarrow X$ is continuous,

$$\begin{aligned} (\pi \circ A)^{-1}(U) &= A^{-1}(\pi^{-1}(U)) \\ &\in \mathcal{N}(x, y) \end{aligned}$$

Since $\mathcal{B} = \{A \times B : A, B \subset X \text{ and } A, B \text{ are open}\}$ is a basis for the product topology on X^2 , there exist $V_x \times V_y \in \mathcal{B}$ such that $(x, y) \in V_x \times V_y \subset (\pi \circ A)^{-1}(U)$. Thus $V_x \in \mathcal{N}(x)$, $V_y \in \mathcal{N}(y)$ and $V_x \times V_y \in \mathcal{N}(x, y)$. Recall that $\pi \times \pi : X^2 \rightarrow (X/M)^2$ is defined by $\pi \times \pi(x, y) = (\pi(x), \pi(y))$. For $x, y \in X$, we have that

$$\begin{aligned} \bar{A} \circ (\pi \times \pi)(x, y) &= \bar{A}(\bar{x}, \bar{y}) \\ &= \bar{x} + \bar{y} \\ &= \pi(x) + \pi(y) \\ &= \pi(x + y) \\ &= \pi \circ A(x, y) \end{aligned}$$

So $\bar{A} \circ (\pi \times \pi) = \pi \circ A$. Since π is open, an exercise in the section on the product topology implies that $\pi \times \pi$ is open and therefore $\pi \times \pi(V_x \times V_y) \in \mathcal{N}(\bar{x}, \bar{y})$. Hence

$$\begin{aligned} \bar{A} \circ (\pi \times \pi)(V_x \times V_y) &\subset \bar{A} \circ (\pi \times \pi)((\pi \circ A)^{-1}(U)) \\ &= \bar{A} \circ (\pi \times \pi)((\bar{A} \circ (\pi \times \pi))^{-1}(U)) \\ &\subset U \end{aligned}$$

So for each $U \in \mathcal{N}(\bar{x} + \bar{y})$, there exists $\pi \times \pi(V_x \times V_y) \in \mathcal{N}(\bar{x}, \bar{y})$ such that $\bar{A}(\pi \times \pi(V_x \times V_y)) \subset U$. Hence $\bar{A}$ is continuous at $(\bar{x}, \bar{y})$. Since $\bar{x}, \bar{y} \in X/M$ are arbitrary, $\bar{A}$ is continuous.

- Let $\lambda \in \mathbb{C}$ and $\bar{x} \in X/M$. Let $U \in \mathcal{N}(\lambda\bar{x})$. Since π is continuous, $\pi^{-1}(U) \in \mathcal{N}(\lambda x)$. Since scalar multiplication $\Lambda : \mathbb{C} \times X \rightarrow X$ is continuous,

$$\begin{aligned}\Lambda^{-1}(\pi^{-1}(U)) &= (\pi \circ \Lambda)^{-1}(U) \\ &\in \mathcal{N}(\lambda, x)\end{aligned}$$

Since $\mathcal{B} = \{A \times B : A \subset \mathbb{C}, B \subset X \text{ and } A, B \text{ are open}\}$ is a basis for the product topology on $\mathbb{C} \times X$, there exist $V_\lambda \times V_x \in \mathcal{B}$ such that $(\lambda, x) \in V_\lambda \times V_x \subset (\pi \circ \Lambda)^{-1}(U)$. Thus $V_\lambda \in \mathcal{N}(\lambda)$, $V_x \in \mathcal{N}(x)$ and $V_\lambda \times V_x \in \mathcal{N}(\lambda, x)$. As in the previous part, $\pi \circ \Lambda = \bar{\Lambda} \circ (\text{id}_{\mathbb{C}} \times \pi)$ and $\text{id}_{\mathbb{C}}$ is open. Hence $\text{id}_{\mathbb{C}} \times \pi$ is open and $\text{id}_{\mathbb{C}} \times \pi(V_\lambda \times V_x) \in \mathcal{N}(\lambda, \bar{x})$. As in the previous part we have that

$$\begin{aligned}\bar{\Lambda} \circ (\text{id}_{\mathbb{C}} \times \pi)(V_\lambda \times V_x) &\subset \bar{\Lambda} \circ (\text{id}_{\mathbb{C}} \times \pi)((\pi \circ \Lambda)^{-1}(U)) \\ &= \bar{\Lambda} \circ (\text{id}_{\mathbb{C}} \times \pi)((\bar{\Lambda} \circ (\text{id}_{\mathbb{C}} \times \pi))^{-1}(U)) \\ &\subset U\end{aligned}$$

So for each $U \in \mathcal{N}(\lambda\bar{x})$, there exists $\text{id}_{\mathbb{C}} \times \pi(V_\lambda \times V_x) \in \mathcal{N}(\lambda, \bar{x})$ such that $\bar{\Lambda}(\text{id}_{\mathbb{C}} \times \pi(V_\lambda \times V_x)) \subset U$. Hence $\bar{\Lambda}$ is continuous at $(\lambda, \bar{x})$. Since $\lambda \in \mathbb{C}$ and $\bar{x} \in X/M$ are arbitrary, $\bar{\Lambda}$ is continuous. □

Exercise 8.10.0.3. Let X be a topological vector space and $M \subset X$ a subspace. Then

1. $\pi_M \in L(X, X/M)$
2. π_M is open

Proof.

1. **FINISH!!!**

2. Define the action $\phi : M \times X \rightarrow X$ by $m \cdot x = x + m$. Then $M \cdot x = x + M$. Since for each $m \in M$, the map $x \mapsto x + m$ is continuous, Exercise 4.8.2.5 implies that $\pi_M : X \rightarrow X/M$ is open. □

Exercise 8.10.0.4. Let X be a topological vector space and $M \subset X$ a subspace. Then M is closed iff X/M is Hausdorff.

Proof. Suppose that M is closed. Define the action $\phi : M \times X \rightarrow X$ by $m \cdot x = m + x$. Denote by $\sim$, the equivalence relation induced by ϕ (i.e. $x \sim y$ iff $x - y \in M$). A previous exercise implies that $\pi : X \rightarrow X/M$ is open. Let $(x_\alpha, y_\alpha)_{\alpha \in A} \subset \sim$ be a net and $(x, y) \in X \times X$. Suppose that $(x_\alpha, y_\alpha) \rightarrow (x, y)$. Then $x_\alpha \rightarrow x$ and $y_\alpha \rightarrow y$. Therefore $x_\alpha - y_\alpha \rightarrow x - y$. Since for each $\alpha \in A$, $x_\alpha - y_\alpha \in M$ and M is closed, we have that $x - y \in M$. Hence $(x, y) \in \sim$ and $\sim$ is closed. Since π is open, a previous exercise in the section on separation and countability implies that X/M is Hausdorff.

Conversely, suppose that X/M is Hausdorff. Then $\{0 + M\}$ is closed in X/M . Since $\pi : X \rightarrow X/M$ is continuous, we have that $M = \pi^{-1}(0 + M)$ is closed in X . □

Exercise 8.10.0.5. Let X be a topological vector spaces and $\phi : X \rightarrow \mathbb{C}$ linear. Then $\ker \phi$ is closed iff ϕ is continuous.

Note: need to show that if $T : X \rightarrow Y$ is linear, then T is continuous iff $\bar{T} : X/\ker T \rightarrow Y$ is continuous

Proof. Suppose that ϕ is continuous. Since $\{0\} \subset \mathbb{C}$ is closed, $\ker \phi = \phi^{-1}(\{0\})$ is closed.

Conversely, suppose that $\ker \phi$ is closed. Then $X/\ker \phi$ is Hausdorff. Hence **FINISH!!!** □

Exercise 8.10.0.6. Let X be a topological vector space and $(\phi_j)_{j \in [n]} \subset X^*$ and $\phi \in X^*$. If $\bigcap_{j \in [n]} \ker \phi_j \subset \ker \phi$, then there exists

$(\lambda_j)_{j \in [n]} \in \mathbb{K}$ such that $\phi = \sum_{j \in [n]} \lambda_j \phi_j$.

Proof. Suppose that $\bigcap_{j \in [n]} \ker \phi_j \subset \ker \phi$. Define $T \in L(X, \mathbb{R}^n)$ by $T := (\phi_j)_{j \in [n]}$. Then $\ker T = \bigcap_{j \in [n]} \ker \phi_j$. We note that each $x_1, x_2 \in X$,

$$\begin{aligned}
 x_1 \sim_T x_2 &\iff T(x_1) = T(x_2) \\
 &\iff T(x_1 - x_2) = 0 \\
 &\iff x_1 - x_2 \in \ker T \\
 &\iff x_1 - x_2 \in \bigcap_{j \in [n]} \ker \phi_j \\
 &\implies x_1 - x_2 \in \ker \phi \\
 &\iff \phi(x_1 - x_2) = 0 \\
 &\iff \phi(x_1) = \phi(x_2).
 \end{aligned}$$

Thus ϕ is $\sim_T$ -invariant. Then there exists a unique $\bar{\phi}_0 \in L(\text{Im } T, \mathbb{R})$ such that $\bar{\phi}_0 \circ T = \phi$. Define $\bar{\phi} \in (\mathbb{R}^n)^*$ by $\bar{\phi} := \bar{\phi}_0 \oplus 0_{(\text{Im } T)^\perp}$. Since $(\mathbb{R}^n)^* = \text{span}(\epsilon_j : j \in [n])$, there exists $(\lambda_j)_{j \in [n]}$ such that $\bar{\phi} = \sum_{j \in [n]} \lambda_j \epsilon_j$. Then

$$\begin{aligned}
 \phi &= \bar{\phi} \circ T \\
 &= \left(\sum_{j \in [n]} \lambda_j \epsilon_j \right) \circ T \\
 &= \sum_{j \in [n]} \lambda_j \epsilon_j \circ T \\
 &= \sum_{j \in [n]} \lambda_j \phi_j.
 \end{aligned}$$

FINISH!!! (Clean up) □

8.11 Compactness

Note 8.11.0.1. pulling from [5, 6]

Exercise 8.11.0.2. Let X be a topological vector space. Suppose that X is Hausdorff. Let $(K_j)_{j \in [n]} \subset \mathcal{K}(X)$. If for each $j \in [n]$, K_j is convex, then $\text{conv} \left(\bigcup_{j \in [n]} K_j \right) \in \mathcal{K}(X)$.

Proof. Suppose that for each $j \in [n]$, K_j is convex.

- Suppose that for each $j \in [n]$, $K_j \neq \emptyset$. **An ex in convex sets section** implies that

$$\text{conv} \left(\bigcup_{j \in [n]} K_j \right) = \left\{ \sum_{j \in [n]} \lambda_j a_j : (\lambda_j)_{j \in [n]} \in \Delta^{n-1} \text{ and } (a_j)_{j \in [n]} \in \prod_{j \in [n]} K_j \right\}.$$

Define $T : \mathbb{R}^n \times X^n \rightarrow X$ by

$$T((\lambda_j)_{j \in [n]}, (a_j)_{j \in [n]}) := \sum_{j \in [n]} \lambda_j a_j.$$

make ex somewhere implies that $\Delta^{n-1} \in \mathcal{K}(\mathbb{R}^n)$. Since $(K_j)_{j \in [n]} \subset \mathcal{K}(X)$, **Tychonoff's theorem (ref)** implies that $\prod_{j \in [n]} K_j \in \mathcal{K}(X^n)$. **Tychonoff's theorem (ref)** then implies that $\Delta^{n-1} \times \prod_{j \in [n]} K_j \in \mathcal{K}(\mathbb{R}^n \times X^n)$. Since X is a locally convex space, **(ref ex)**

implies that $T \in L(\mathbb{R}^n \times X^n; X)$. Thus

$$\begin{aligned} \operatorname{conv} \left(\bigcup_{j \in [n]} K_j \right) &= \left\{ \sum_{j \in [n]} \lambda_j a_j : (\lambda_j)_{j \in [n]} \in \Delta^{n-1} \text{ and } (a_j)_{j \in [n]} \in \prod_{j \in [n]} K_j \right\} \\ &= T(\Delta^{n-1} \times \prod_{j \in [n]} K_j) \\ &\in \mathcal{K}(X). \end{aligned}$$

- Set $J := \{j \in [n] : K_j = \emptyset\}$. Then **the prev part** implies that

$$\begin{aligned} \operatorname{conv} \left(\bigcup_{j \in [n]} K_j \right) &= \operatorname{conv} \left(\bigcup_{j \in J^c} K_j \right) \\ &\in \mathcal{K}(X). \end{aligned}$$

□

Exercise 8.11.0.3. Let X be a locally convex space and $E \subset X$. If E is precompact, then $\operatorname{conv} E$ is precompact.

Proof. Suppose that E is precompact. Then **get neighborhood U around 0, use compactness and translation to get finitely many $(x_j)_{j \in [n]} \subset E$ such that $(U + x_j)_{j \in [n]}$ is open cover of $\operatorname{cl} E$.**

see 4.3 pg 50 in [5]

□

8.12 Duality

8.12.1 Weak and Weak-* Topology

Definition 8.12.1.1. Let X, Y and Z be vector spaces over $\mathbb{K}$ and $b \in \mathcal{L}^2(X, Y; Z)$. Then (X, Y, Z, b) is said to be a $(X, Y; Z)$ -pairing.

Definition 8.12.1.2. Let X, Y and Z be vector spaces and $b \in \mathcal{L}^2(X, Y; Z)$. We define the **dual pairing** of b , denoted $b^* \in \mathcal{L}^2(Y, X; Z)$, by $b^*(y, x) := b(x, y)$.

Exercise 8.12.1.3. Let X, Y and Z be vector spaces and $b \in \mathcal{L}^2(X, Y; Z)$. Then b^* is a $(Y, X; Z)$ -pairing.

Proof. Clear. □

Definition 8.12.1.4. Let X, Y and Z be topological vector spaces and $b \in \mathcal{L}^2(X, Y; Z)$. We define the **weak topology on X induced by b** , denoted $\sigma_b(X, Y)$ by

$$\sigma_b(X, Y) := \tau_X(b(\cdot, y) : y \in Y)$$

We define the **weak topology on Y induced by b** , denoted $\sigma_b(Y, X)$, by $\sigma_b(Y, X) := \sigma_{b^*}(Y, X)$.

Exercise 8.12.1.5. Let X, Y and Z be topological vector spaces, $b \in \mathcal{L}^2(X, Y; Z)$, $(x_\alpha)_{\alpha \in A} \subset X$ a net, $(y_\alpha)_{\alpha \in A} \subset Y$ a net and $x \in X, y \in Y$. Then

1. $x_\alpha \rightarrow x$ in $(X, \sigma_b(X, Y))$ iff for each $y \in Y$, $b(x_\alpha, y) \rightarrow b(x, y)$
2. $y_\alpha \rightarrow y$ in $(Y, \sigma_b(Y, X))$ iff for each $x \in X$, $b(x, y_\alpha) \rightarrow b(x, y)$.

Proof. Immediate by Exercise 4.3.2.15. □

Exercise 8.12.1.6. Let X, Y and Z be topological vector spaces and $b \in \mathcal{L}^2(X, Y; Z)$. Then

1. $(X, \sigma_b(X, Y))$ is a topological vector space
2. $(Y, \sigma_b(Y, X))$ is a topological vector space

Proof.

1. Let $(u_\alpha)_{\alpha \in A}, (v_\alpha)_{\alpha \in A} \subset X$ and $(\lambda_\alpha)_{\alpha \in A} \subset \mathbb{K}$ be nets and $u, v \in X$ and $\lambda \in \mathbb{K}$. Suppose that $u_\alpha \rightarrow u$ and $v_\alpha \rightarrow v$ in $(X, \sigma_b(X, Y))$ and $\lambda_\alpha \rightarrow \lambda$ in $(\mathbb{K}, \mathcal{T}_{\mathbb{K}})$. Let $y \in Y$. Then $b(u_\alpha, y) \rightarrow b(u, y)$ and $b(v_\alpha, y) \rightarrow b(v, y)$. Since Z is a topological vector space,

$$\begin{aligned} b(u_\alpha + v_\alpha, y) &= b(u_\alpha, y) + b(v_\alpha, y) \\ &\rightarrow b(u, y) + b(v, y) \\ &= b(u + v, y) \end{aligned}$$

and

$$\begin{aligned} b(\lambda_\alpha u_\alpha, y) &= \lambda_\alpha b(u_\alpha, y) \\ &\rightarrow \lambda b(u, y) \\ &= b(\lambda u, y) \end{aligned}$$

Since $y \in Y$ is arbitrary, $u_\alpha + v_\alpha \rightarrow u + v$ and $\lambda_\alpha u_\alpha \rightarrow \lambda u$. Hence addition $X \times X \rightarrow X$ and scalar multiplication $\mathbb{K} \times X \rightarrow X$ are continuous.

2. Since $\sigma_b(X, Y) = \sigma_{b^*}(Y, X)$, (1) implies (2). □

Definition 8.12.1.7. Let X, Y, Z be vector spaces and $b \in \mathcal{L}^2(X, Y; Z)$. Then

- Y is said to **separate the points of X via b** if for each $x \in X$, $x \neq 0$ implies that there exists $y \in Y$ such that $b(x, y) \neq 0$

- X is said to **separate the points of Y via b** if X separates the points of Y via b^*

Exercise 8.12.1.8. Let X, Y and Z be topological vector spaces and $b \in \mathcal{L}^2(X, Y; Z)$. Then Y separates the points of X via b iff for each $x \in X$, $x \neq 0$ implies that there exists $y \in Y$ such that $b(x, y) \neq 0$.

Proof. **FINISH!!! FIX!!!** □

Exercise 8.12.1.9. **just cite ex in topology separation axioms section exercise about weak (initial topology)**

Let X, Y and Z be topological vector spaces and $b \in \mathcal{L}^2(X, Y; Z)$. Suppose that Z is Hausdorff.

1. if Y separates the points of X via b , then $(X, \sigma_b(X, Y))$ is Hausdorff
- 2.

Proof.

1. Since $\sigma_b(X, Y) = \tau_X(b(\cdot, y) : y \in Y)$ and $(b(\cdot, y))_{y \in Y}$ separates the points of X , Exercise 4.12.1.9 implies that $(X, \sigma_b(X, Y))$ is Hausdorff.
2. **FINISH!!!** □

Definition 8.12.1.10. Let X be a topological vector space.

- We define the **canonical pairing of $(X, \mathcal{T})$ with X^*** , denoted $\langle \cdot, \cdot \rangle : X^* \times X \rightarrow \mathbb{K}$, by $\langle \phi, x \rangle := \phi(x)$.
- For each $x \in X$, we define $\hat{x} : X^* \rightarrow \mathbb{K}$ by $\hat{x} := \langle \cdot, x \rangle$.
- We define $\hat{X} \subset \mathbb{K}^{X^*}$ by $\hat{X} = \{\hat{x} : x \in X\}$.

Note 8.12.1.11. When the

Definition 8.12.1.12. Let $(X, \mathcal{T})$ be a topological vector space. We define the **weak topology on X** , denoted $\mathcal{T}_w$, by $\mathcal{T}_w := \sigma_{\langle \cdot, \cdot \rangle}(X, (X, \mathcal{T})^*)$.

Exercise 8.12.1.13. Let $(X, \mathcal{T})$ be a topological vector space. Then $\mathcal{T}_w = \tau_X((X, \mathcal{T})^*)$.

Proof. Clear. □

Definition 8.12.1.14. Let $(X, \mathcal{T})$ be a topological vector space, $(x_\alpha)_{\alpha \in A} \subset X$ and $x \in X$. Then $(x_\alpha)_{\alpha \in A}$ is said to **converge weakly to x** , denoted $x_\alpha \xrightarrow{w} x$, if $x_\alpha \rightarrow x$ in $(X, \mathcal{T}_w)$.

Exercise 8.12.1.15. Let X be a topological vector, $(x_\alpha)_{\alpha \in A} \subset X$ a net and $x \in X$. Then $x_\alpha \xrightarrow{w} x$ iff for each $\lambda \in X^*$, $\lambda(x_\alpha) \rightarrow \lambda(x)$.

Proof. Immediate by Exercise 8.12.1.5. □

Definition 8.12.1.16. Let $(X, \mathcal{T})$ be a topological vector space. We define the **weak-* topology on $(X, \mathcal{T})^*$** , denoted $\mathcal{T}_{w*}$, by $\mathcal{T}_{w*} := \sigma_{\langle \cdot, \cdot \rangle}((X, \mathcal{T})^*, X)$.

Exercise 8.12.1.17. Let X be a topological vector space. Then $\mathcal{T}_{w*} = \tau_{X^*}(\hat{X})$.

Proof. Clear. □

Definition 8.12.1.18. Let $(X, \mathcal{T})$ be a topological vector space, $(\lambda_\alpha)_{\alpha \in A} \subset (X, \mathcal{T})^*$ and $\lambda \in (X, \mathcal{T})^*$. Then $(\lambda_\alpha)_{\alpha \in A}$ is said to **converge in weak-* to λ** , denoted $\lambda_\alpha \xrightarrow{w*} \lambda$, if $\lambda_\alpha \rightarrow \lambda$ in $((X, \mathcal{T})^*, \mathcal{T}_{w*})$.

Exercise 8.12.1.19. Let X be a topological vector, $(\lambda_\alpha)_{\alpha \in A} \subset X^*$ a net and $\lambda \in X^*$. Then $\lambda_\alpha \xrightarrow{w*} \lambda$ iff for each $x \in X$, $\lambda_\alpha(x) \rightarrow \lambda(x)$.

Proof. Immediate by Exercise 8.12.1.5. □

Exercise 8.12.1.20. Let X be a topological vector space.

1. If X^* separates the points of X , then $(X, \mathcal{T}_w)$ is a locally convex space
2. $(X^*, \mathcal{T}_{w^*})$ is a locally convex space

Proof.

1. Suppose that X^* separates the points of X . For $\lambda \in X^*$, define $p_\lambda : X \rightarrow [0, \infty)$ by $p_\lambda = |\lambda|$. Set $\mathcal{P}_w = \{p_\lambda : \lambda \in X^*\}$. Then $\mathcal{P}_w$ separates the points of X . Let $(x_\alpha)_{\alpha \in A} \subset X$ be a net and $x \in X$. Suppose that $x_\alpha \xrightarrow{w} x$. Let $\lambda \in X^*$. Then

$$\begin{aligned} p_\lambda(x_\alpha - x) &= |\lambda(x_\alpha - x)| \\ &= |\lambda(x_\alpha) - \lambda(x)| \\ &\rightarrow 0 \end{aligned}$$

So $x_\alpha \rightarrow x$ in $\tau_X(\pi_p : p \in \mathcal{P}_w)$.

Conversely, suppose that $x_\alpha \rightarrow x$ in $\tau_X(\pi_p : p \in \mathcal{P}_w)$. Then for each $x \in X$,

$$\begin{aligned} |\lambda(x_\alpha) - \lambda(x)| &= p_\lambda(x_\alpha - x) \\ &\rightarrow 0 \end{aligned}$$

So that $\lambda(x_\alpha) \rightarrow \lambda(x)$ and $x_\alpha \xrightarrow{w} x$. [ref prev ex in locally convex section](#) implies that $\mathcal{T}_w = \mathcal{T}_{\mathcal{P}_w}$ and $(X, \mathcal{T}_w)$ is a locally convex space.

2. For $x \in X$, define $p_x : X^* \rightarrow [0, \infty)$ by $p_x = |\hat{x}|$. Set $\mathcal{P}_{w^*} = \{p_x : x \in X\}$. Let $\phi \in X^*$. Suppose that $\phi \neq 0$. Then there exists $x \in X$ such that

$$\begin{aligned} \hat{x}(\phi) &= \phi(x) \\ &\neq 0 \end{aligned}$$

So $\mathcal{P}_{w^*}$ separates the points of X^* . Let $(\lambda_\alpha)_{\alpha \in A} \subset X^*$ be a net and $\lambda \in X^*$. Suppose that $\lambda_\alpha \xrightarrow{w^*} \lambda$. Let $x \in X$. Then

$$\begin{aligned} p_x(\lambda_\alpha - \lambda) &= |\hat{x}(\lambda_\alpha - \lambda)| \\ &= |\hat{x}(\lambda_\alpha) - \hat{x}(\lambda)| \\ &\rightarrow 0 \end{aligned}$$

So $\lambda_\alpha \rightarrow \lambda$ in $\tau_{X^*}(\pi_p : p \in \mathcal{P}_{w^*})$.

Conversely, suppose that $\lambda_\alpha \rightarrow \lambda$ in $\tau_{X^*}(\pi_p : p \in \mathcal{P}_{w^*})$. Then for each $x \in X$,

$$\begin{aligned} |\hat{x}(\lambda_\alpha) - \hat{x}(\lambda)| &= p_x(\lambda_\alpha - \lambda) \\ &\rightarrow 0 \end{aligned}$$

So that $\hat{x}(\lambda_\alpha) \rightarrow \hat{x}(\lambda)$ and $\lambda_\alpha \xrightarrow{w^*} \lambda$. [ref prev ex in locally convex section](#) implies that $\mathcal{T}_{w^*} = \mathcal{T}_{\mathcal{P}_{w^*}}$ and $(X^*, \mathcal{T}_{w^*})$ is a locally convex space. □

Note 8.12.1.21. ([change notation or reword](#)) Let X be a topological vector space. When we equip X^* with the weak-* topology, we write X^{**} in place of $(X^*)^*$.

Exercise 8.12.1.22. Let X be a topological vector space. Then $X^{**} = \hat{X}$.

Hint: If $f \in X^{**}$, $(x_j)_{j \in [n]} \subset X$ and $\bigcap_{j \in [n]} B_{r^{-1}|\hat{x}_j|}(0, 1) \subset B_{|f|}(0, 1)$, then $\bigcap_{j \in [n]} \ker \hat{x}_j \subset \ker f$.

Proof. Set $\mathcal{P}_{w^*} := \{|\hat{x}| : x \in X\}$. By definition $\mathcal{T}_{w^*} = \mathcal{T}_{\mathcal{P}_{w^*}}$. Let $f \in X^{**}$. Define $p_f \in \text{SemiNorm}(X^*, \mathcal{T}_{w^*})$ by $p_f := |f|$. By continuity, $B_{p_f}(0, 1)$ is open. [A previous exercise in locally convex section](#) implies that there exists $n \in \mathbb{N}$, $(p_j)_{j \in [n]} \subset \mathcal{P}_{w^*}$ and $r > 0$ such that

$$\begin{aligned} \bigcap_{j \in [n]} B_{r^{-1}p_j}(0, 1) &= \bigcap_{j \in [n]} B_{p_j}(0, r) \\ &\subset B_{p_f}(0, 1). \end{aligned}$$

A previous exercise implies that

$$\begin{aligned} p_f &\leq \max_{j \in [n]} [r^{-1} p_j] \\ &= r^{-1} \max_{j \in [n]} p_j. \end{aligned}$$

By definition, there exists $(x_j)_{j \in [n]} \subset X$ such that for each $j \in [n]$, $p_j = |\hat{x}_j|$. A prev ex then implies that

$$\begin{aligned} \bigcap_{j \in [n]} \ker \hat{x}_j &= \bigcap_{j \in [n]} \ker p_j \\ &= \ker \max_{j \in [n]} p_j \\ &= \subset \ker p_f \\ &= \ker f. \end{aligned}$$

A prev ex in quotient space section then implies that there exists $(\lambda_j)_{j \in [n]} \subset \mathbb{K}$ such that

$$\begin{aligned} f &= \sum_{j \in [n]} \lambda_j \hat{x}_j \\ &\in \hat{X}. \end{aligned}$$

Since $f \in X^{**}$ is arbitrary, we have that for each $f \in X^{**}$, $f \in \hat{X}$. Thus

$$\begin{aligned} X^{**} &\subset \hat{X} \\ &\subset X^{**}. \end{aligned}$$

Hence $X^{**} = \hat{X}$. □

Exercise 8.12.1.23. Let X be a topological vector space. For each net $(x_\alpha)_{\alpha \in \Gamma} \subset X$ and $x \in X$, $x_\alpha \rightarrow x$, implies that $x \xrightarrow{w} x$.

Proof. Suppose that $x_\alpha \rightarrow x$. Let $\phi \in X^*$. Since ϕ is continuous, $\phi(x_\alpha) \rightarrow \phi(x)$. Since $\phi \in X^*$ is arbitrary, we have that for each $\phi \in X^*$, $\phi(x_\alpha) \rightarrow \phi(x)$. Thus $x_\alpha \xrightarrow{w} x$. □

Definition 8.12.1.24. Let $(X, \mathcal{T}_X), (Y, \mathcal{T}_Y)$ be topological vector spaces and $f : X \rightarrow Y$. Then f is said to be $(\mathcal{T}_X, \mathcal{T}_Y)$ -**weakly continuous** if f is $(\mathcal{T}_{X,w}, \mathcal{T}_{Y,w})$ -continuous.

Note 8.12.1.25. When the context is clear, we may say “weakly continuous” in place of “ $(\mathcal{T}_X, \mathcal{T}_Y)$ -weakly continuous”.

Exercise 8.12.1.26. Let X be topological vector spaces and $f : X \rightarrow Y$. Then f is weakly continuous iff for each net $(x_\alpha)_{\alpha \in \Gamma} \subset X$ and $x \in X$, $x_\alpha \xrightarrow{w} x$ implies that $f(x_\alpha) \xrightarrow{w} f(x)$.

Proof. Clear by Exercise 4.3.2.13. □

Exercise 8.12.1.27. Let $(X, \mathcal{T}_X), (Y, \mathcal{T}_Y)$ be topological vector spaces. Then

1. For each $f : X \rightarrow Y$, f is $(\mathcal{T}_X, \mathcal{T}_Y)$ -continuous implies that f is $(\mathcal{T}_X, \mathcal{T}_{Y,w})$ -continuous.
2. $L((X, \mathcal{T}_X), (Y, \mathcal{T}_{Y,w})) \subset L((X, \mathcal{T}_{X,w}), (Y, \mathcal{T}_{Y,w}))$.

Proof.

1. Let $f : X \rightarrow Y$. Suppose that f is $(\mathcal{T}_X, \mathcal{T}_Y)$ -continuous. Then for each $\phi \in (Y, \mathcal{T}_Y)^*$, we have that $\phi \circ f$ is $(\mathcal{T}_X, \mathcal{T}_{\mathbb{K}})$ -continuous. Since $\mathcal{T}_{Y,w} = \tau_Y((Y, \mathcal{T}_Y)^*)$, Exercise 4.2.0.14 implies that f is $(\mathcal{T}_X, \mathcal{T}_{Y,w})$ -continuous.

2. Let $T \in L((X, \mathcal{T}_X), (Y, \mathcal{T}_Y))$, $(x_\alpha)_{\alpha \in \Gamma} \subset X$ a net and $x \in X$. Suppose that $x_\alpha \xrightarrow{w} x$. Let $\phi \in Y^*$. Then $\phi \circ T \in X^*$. Thus

$$\begin{aligned}\phi(T(x_\alpha)) &= \phi \circ T(x_\alpha) \\ &\rightarrow \phi \circ T(x) \\ &= \phi(T(x)).\end{aligned}$$

Since $\phi \in Y^*$ is arbitrary, we have that for each $\phi \in Y^*$, $\phi(T(x_\alpha)) \rightarrow \phi(T(x))$. Hence $T(x_\alpha) \xrightarrow{w} T(x)$. Since $(x_\alpha)_{\alpha \in \Gamma} \subset X$ a net and $x \in X$ with $x_\alpha \rightarrow x$ are arbitrary, we have that for each net $(x_\alpha)_{\alpha \in \Gamma} \subset X$ and $x \in X$, $x_\alpha \rightarrow x$ implies that $T(x_\alpha) \xrightarrow{w} T(x)$. Thus $T \in L((X, \mathcal{T}_{X,w}), (Y, \mathcal{T}_{Y,w}))$. Since $T \in L((X, \mathcal{T}_X), (Y, \mathcal{T}_Y))$ is arbitrary, we have that for each $T \in L((X, \mathcal{T}_X), (Y, \mathcal{T}_Y))$, $T \in L((X, \mathcal{T}_{X,w}), (Y, \mathcal{T}_{Y,w}))$. Hence $L((X, \mathcal{T}_X), (Y, \mathcal{T}_Y)) \subset L((X, \mathcal{T}_{X,w}), (Y, \mathcal{T}_{Y,w}))$

□

Exercise 8.12.1.28. Let X be a topological vector space, $E \in \mathcal{K}(X)$. If X^* separates the points of X , then

1. $\mathcal{T}_w \cap E = \mathcal{T} \cap E$,
2. for each net $(x_\alpha)_{\alpha \in \Gamma} \subset E$ and $x \in E$, $x_\alpha \rightarrow x$ iff $x_\alpha \xrightarrow{w} x$.

Proof. Suppose that X^* separates the points of X . Exercise 17.3.0.5 implies that $(X, \mathcal{T}_w)$ is Hausdorff. Exercise 4.14.7.4 then implies that

1. $\mathcal{T}_w \cap E = \mathcal{T} \cap E$,
2. for each net $(x_\alpha)_{\alpha \in \Gamma} \subset E$ and $x \in E$, $x_\alpha \rightarrow x$ iff $x_\alpha \xrightarrow{w} x$.

□

Exercise 8.12.1.29. Let X be a locally convex space and $E \in \mathcal{K}(X)$. Then

1. $\mathcal{T}_w \cap E = \mathcal{T} \cap E$,
2. for each net $(x_\alpha)_{\alpha \in \Gamma} \subset E$ and $x \in E$, $x_\alpha \rightarrow x$ iff $x_\alpha \xrightarrow{w} x$.

Proof. Clear by Exercise 8.7.0.5 and [prev ex](#).

□

8.12.2 Annihilators

Definition 8.12.2.1. Let X, Y and Z be topological vector spaces, $b \in \mathcal{L}^2(X, Y; Z)$ and $A \subset X$. We define the **annihilator of A with respect to b** , denoted $A^{\perp_b}$, by $A^{\perp_b} := \{y \in Y : \text{for each } x \in A, b(x, y) = 0\}$.

Note 8.12.2.2. Let $A \subset X$ and $B \subset Y$. When the context is clear, we write $A^\perp$ and $B^\perp$ in place of $A^{\perp_b}$ and $B^{\perp_{b^*}}$ respectively.

Exercise 8.12.2.3. Let X, Y and Z be topological vector spaces, $b \in \mathcal{L}^2(X, Y; Z)$, $A \subset X$ and $B \subset Y$. Then

1. $A^\perp$ is a subspace of Y .
- 2.
- 3.
- 4.
- 5.
- 6.

Proof. 1. Let $y_1, y_2 \in A^\perp$, $\lambda \in \mathbb{K}$ and $x \in A$. Then

$$b()$$

- 2.

- 3.
- 4.
- 5.
- 6.

□

8.12.3 Adjoints

Definition 8.12.3.1.

Chapter 9

Banach Spaces

Note 9.0.0.1. For the remainder of the notes, we assume familiarity with [section on summation ??](#) ([ref appendix on summation of vector functions](#))

9.1 Introduction

need to define norms before seminorm section, so probably need to define pseudometrics somewhere in metric space chapter and then show that seminorms induce pseudometrics and norms induce metrics $d_{\|\cdot\|}$, before this chapter

Definition 9.1.0.1. Let $(X, \|\cdot\|)$ be a normed vector space. Then $(X, \|\cdot\|)$ is said to be a **Banach space** if $(X, d_{\|\cdot\|})$ is complete.

Exercise 9.1.0.2. We have that $(C^1([0, 1]), \|\cdot\|_\infty)$ is not a Banach space.

Proof. ([make ex in chapter on metric spaces and show \$C\(X\)\$ is complete when \$X\$ is compact, then ref exercise](#)). Set $X := C^1([0, 1])$ and $Y := C([0, 1])$. Then $X \subset Y$. Recall that for each $a, b \geq 0$ and $p \in \mathbb{N}$,

$$(a^{\frac{1}{p}} + b^{\frac{1}{p}})^p = \sum_{n=0}^p \binom{p}{n} a^{\frac{n}{p}} b^{\frac{p-n}{p}} \geq a + b$$

Thus $(a + b)^{\frac{1}{p}} \leq a^{\frac{1}{p}} + b^{\frac{1}{p}}$.

Define $(f_n)_{n \in \mathbb{N}} \subset X$ and $f \in Y$ by $f_n(x) := \sqrt{(x - \frac{1}{2})^2 + \frac{1}{n^2}}$ and $f(x) := |x - \frac{1}{2}|$. Then $f \notin X$. Note that for each $n \in \mathbb{N}$, $f \leq f_n$. Our observation above ([ref observation](#)) implies that for each $x \in X$,

$$\begin{aligned} f_n(x) &= \left[(x - \frac{1}{2})^2 + \frac{1}{n^2} \right]^{\frac{1}{2}} \\ &\leq |x - \frac{1}{2}| + \frac{1}{n} \end{aligned}$$

Thus for each $n \in \mathbb{N}$, $0 \leq f_n - f \leq \frac{1}{n}$. This implies that

$$\begin{aligned} \|f_n - f\|_\infty &= \sup_{x \in [0, 1]} |f_n(x) - f(x)| \\ &\leq \frac{1}{n} \\ &\rightarrow 0. \end{aligned}$$

Therefore $f_n \xrightarrow{u} f$. Thus there exists $(f_n)_{n \in \mathbb{N}} \subset X$ and $f \in Y$ such that $f_n \rightarrow f$ and $f \notin X$. Exercise 4.13.1.4 then implies that X is not closed. Exercise 6.7.2.2 implies that, X is not complete. $\square$

Definition 9.1.0.3. maybe get rid, of, just write out $\sum \|x_n\| < \infty$ instead of saying 'converges absolutely' or maybe discuss l^1 for Banach space valued functions. Let X be a normed vector space and $(x_i)_{i=1}^n \subset X$. The series $\sum_{i=1}^{\infty} x_i$ is said to **converge** if the sequence $s_n := \sum_{i=1}^n x_i$ converges. The series $\sum_{i=1}^{\infty} x_i$ is said to **converge absolutely** if $\sum_{i \in \mathbb{N}} \|x_i\| < \infty$.

Exercise 9.1.0.4. Let X be a normed vector space. Then X is complete iff for each $(x_i)_{i \in \mathbb{N}} \subset X$, $\sum_{i=1}^{\infty} x_i$ converges absolutely implies that $\sum_{i=1}^{\infty} x_i$ converges.

Hint: Given a Cauchy sequence $(x_n)_{n \in \mathbb{N}}$, obtain a subsequence $(x_{n_j})_{j \in \mathbb{N}} \subset (x_n)_{n \in \mathbb{N}}$ such that for each $j \in \mathbb{N}$, $\|x_{n_{j+1}} - x_{n_j}\| < 2^{-j}$. Define a new sequence $(y_j)_{j \in \mathbb{N}} \subset X$ by

$$y_j = \begin{cases} x_{n_1} & j = 1 \\ x_{n_j} - x_{n_{j-1}} & j \geq 2 \end{cases}$$

Proof. Suppose that X is complete. Let $(x_i)_{i \in \mathbb{N}} \subset X$. Suppose that $\sum_{i=1}^{\infty} x_i$ converges absolutely. Let $\epsilon > 0$. Choose $N \in \mathbb{N}$ such that for each $m, n \in \mathbb{N}$, if $m, n \geq N$ and $m < n$, then $\sum_{m+1}^n \|x_i\| < \epsilon$. Let $m, n \in \mathbb{N}$. Suppose that $m < n$. Then

$$\begin{aligned} \|s_n - s_m\| &= \left\| \sum_{i=1}^n x_i - \sum_{i=1}^m x_i \right\| \\ &= \left\| \sum_{i=m+1}^n x_i \right\| \\ &\leq \sum_{i=m+1}^n \|x_i\| \\ &< \epsilon \end{aligned}$$

Thus $(s_n)_{n \in \mathbb{N}}$ is Cauchy. Since X is complete, $\sum_{i=1}^{\infty} x_i$ converges.

Conversely, Suppose that for each $(x_i)_{i \in \mathbb{N}} \subset X$, $\sum_{i=1}^{\infty} x_i$ converges absolutely implies that $\sum_{i=1}^{\infty} x_i$ converges. Let $(x_i)_{i \in \mathbb{N}} \subset X$ be Cauchy. Proceed inductively to create a strictly increasing sequence $(n_i)_{i \in \mathbb{N}} \subset \mathbb{N}$ such that for each $m, n \in \mathbb{N}$, if $m, n \geq n_i$, then $\|x_m - x_n\| < 2^{-i}$. Define $(y_i)_{i \in \mathbb{N}} \subset X$ by

$$y_i = \begin{cases} x_{n_1} & i = 1 \\ x_{n_i} - x_{n_{i-1}} & i \geq 2 \end{cases}$$

Then $\sum_{i=1}^k y_i = x_{n_k}$ and

$$\begin{aligned} \sum_{i \in \mathbb{N}} \|y_i\| &= \|x_{n_1}\| + \sum_{i \in \mathbb{N}} \|x_{n_i} - x_{n_{i-1}}\| \\ &\leq \|x_{n_1}\| + 2 \sum_{i \in \mathbb{N}} 2^{-i} \\ &= \|x_{n_1}\| + 2 \end{aligned}$$

Hence $(x_{n_k})_{k \in \mathbb{N}} = (\sum_{i=1}^k y_i)_{i \in \mathbb{N}}$ converges. Since $(x_i)_{i \in \mathbb{N}}$ is cauchy and has a convergent subsequence, it converges. So X is complete. $\square$

Exercise 9.1.0.5. Let X be a normed vector space. Then addition $X \times X \rightarrow X$ and scalar multiplication $\mathbb{C} \times X \rightarrow X$ are continuous and $\|\cdot\| : X \rightarrow [0, \infty)$ is continuous.

Proof. Let $\epsilon > 0$. Choose $\delta = \frac{\epsilon}{2}$. Let $(x_1, y_1), (x_2, y_2) \in X \times X$. Suppose that

$$\max\{\|x_1 - x_2\|, \|y_1 - y_2\|\} < \delta$$

Then

$$\begin{aligned} \|(x_1 + y_1) - (x_2 + y_2)\| &= \|(x_1 - x_2) + (y_1 - y_2)\| \\ &\leq \|x_1 - x_2\| + \|y_1 - y_2\| \\ &< 2\delta \\ &= \epsilon \end{aligned}$$

Hence addition is uniformly continuous.

Let $(\lambda_1, x_1) \in \mathbb{C} \times X$ and $\epsilon > 0$. Choose $\delta = \min\{\frac{\epsilon}{2(|\lambda_1| + \|x_1\| + 1)}, \frac{\sqrt{\epsilon}}{\sqrt{2}}\}$. Let $(\lambda_2, x_2) \in \mathbb{C} \times X$. Suppose that

$$\max\{|\lambda_1 - \lambda_2|, \|x_1 - x_2\|\} < \delta$$

Then

$$\begin{aligned} \|\lambda_1 x_1 - \lambda_2 x_2\| &= \|\lambda_1 x_1 - \lambda_1 x_2 + \lambda_1 x_2 - \lambda_2 x_2\| \\ &= \|\lambda_1(x_1 - x_2) + (\lambda_1 - \lambda_2)x_2\| \\ &\leq |\lambda_1|\|x_1 - x_2\| + |\lambda_1 - \lambda_2|\|x_2\| \\ &\leq |\lambda_1|\|x_1 - x_2\| + |\lambda_1 - \lambda_2|(\|x_1 - x_2\| + \|x_1\|) \\ &< |\lambda_1|\delta + \delta(\delta + \|x_1\|) \\ &= (|\lambda_1| + \|x_1\|)\delta + \delta^2 \\ &< \frac{\epsilon}{2} + \frac{\epsilon}{2} \\ &= \epsilon \end{aligned}$$

Since $(\lambda_1, x_1) \in \mathbb{C} \times X$ is arbitrary, scalar multiplication is continuous.

Let $\epsilon > 0$. Choose $\delta = \epsilon$. Let $x, y \in X$. Suppose that $\|x - y\| < \delta$. Then

$$\begin{aligned} \left| \|x\| - \|y\| \right| &\leq \|x - y\| \\ &< \delta \\ &= \epsilon \end{aligned}$$

So $\|\cdot\| : X \rightarrow [0, \infty)$ is uniformly continuous. □

9.2 Bounded Operators

Definition 9.2.0.1. Let X, Y be a normed vector spaces and $T : X \rightarrow Y$ linear. Then T is said to be **bounded** if $T(\bar{B}(0, 1))$ is bounded. We define

$$L(X; Y) = \{T : X \rightarrow Y : T \text{ is linear and bounded}\}$$

When $X = Y$, we write $L(X)$.

Exercise 9.2.0.2. show $L(X; Y)$ is a vector space

Proof. **FINISH!!!** □

Exercise 9.2.0.3. Let X, Y be normed vector spaces and $T : X \rightarrow Y$ linear. Then T is bounded iff there exists $C \geq 0$ such that for each $x \in X$,

$$\|Tx\| \leq C\|x\|$$

Proof. can just cite exercise in metric space definition of bounded subset with $\|x\| = d(x, 0)$ Suppose that T is bounded. If $T = 0$, choose $C = 0$. Suppose that $T \neq 0$. Set $A = \{\|Tx\| : \|x\| = 1\}$. Since $T \neq 0$, there exists $x_0 \in X$ such that $\|x_0\| = 1$ so that $A \neq \emptyset$. Boundedness of T implies that A is bounded. Set $C = \sup A$. Let $x \in X$. If $x = 0$, then $Tx = 0$ and $\|Tx\| \leq C\|x\|$. Suppose that $x \neq 0$. Then $Tx = \|x\|T(\|x\|^{-1}x)$. Since $\|\|x\|^{-1}x\| = 1$, we have that

$$\begin{aligned} \|Tx\| &= \|T(\|x\|^{-1}x)\|\|x\| \\ &\leq C\|x\| \end{aligned}$$

Conversely, suppose that there exists $C \geq 0$ such that for each $x \in X$, $\|Tx\| \leq C\|x\|$. Let $x \in \bar{B}(0, 1)$. Then

$$\begin{aligned} \|Tx\| &\leq C\|x\| \\ &\leq C \end{aligned}$$

So that $T(\bar{B}(0, 1))$ is bounded. □

Exercise 9.2.0.4. Set $X = C^1([0, 1])$ and $Y = C([0, 1])$. Equip both X and Y with the sup norm. Define $T : X \rightarrow Y$ by $Tf = f'$. Then T is not bounded.

Proof. For the sake of contradiction, suppose that T is bounded. Then there exists $C \geq 0$ such that for each $f \in X$, $\|Tf\| \leq C\|f\|$. Choose $n \in \mathbb{N}$ such that $n > C$. Define $f \in X$ by $f(x) = x^n$. Then

$$\begin{aligned} n &= \|Tf\| \\ &\leq C\|f\| \\ &= C \end{aligned}$$

which is a contradiction. Hence T is not bounded. □

Exercise 9.2.0.5. Let X, Y be a normed vector spaces and $T : X \rightarrow Y$ a linear map. Then T is bounded iff there exists $r, s > 0$ such that $T(B(0, r)) \subset B(0, s)$

Proof. Suppose that T is bounded. Then there exists $C \geq 0$ such that for each $x \in X$, $\|Tx\| \leq C\|x\|$. Thus $T(B(0, 1)) \subset B(0, C+1)$. Conversely. Suppose that there exists $r, s > 0$ such that $T(B(0, r)) \subset B(0, s)$. Define $C = \frac{2s}{r}$. Let $x \in X$. Put $\alpha = \frac{r}{2\|x\|}$. Then $\alpha x \in B(0, r)$. So $T(\alpha x) = \alpha T(x) \in B(0, s)$. Hence

$$\begin{aligned} \|T(\alpha x)\| &= \|\alpha T(x)\| \\ &= |\alpha| \|T(x)\| \\ &= \frac{r}{2\|x\|} \|T(x)\| \\ &< s. \end{aligned}$$

Thus

$$\|Tx\| < \frac{2s}{r} \|x\| = C\|x\|$$

So T is bounded. □

Exercise 9.2.0.6. Let X, Y be normed vector spaces and $T : X \rightarrow Y$ a linear map. Then the following are equivalent:

1. T is continuous
2. there exists $x_0 \in X$ such that T is continuous at x_0
3. T is continuous at 0
4. T is bounded

Proof.

- (1) $\implies$ (2):

Suppose that T is continuous. Then for each $x \in X$, T is continuous at x . In particular, T is continuous at 0.

- (2) $\implies$ (3):

Suppose that there exists $x_0 \in X$ such that T is continuous at x_0 . Since T is linear, $T(0) = 0$. Let $(x_n)_{n \in \mathbb{N}} \subset X$. Suppose that $x_n \rightarrow 0$. Then $x_n + x_0 \rightarrow x_0$. Since T is continuous at x_0 , we have that

$$\begin{aligned} T(x_n) + T(x_0) &= T(x_n + x_0) \\ &\rightarrow T(x_0) \end{aligned}$$

Therefore

$$\begin{aligned} T(x_n) &= [T(x_n) + T(x_0)] - T(x_0) \\ &\rightarrow T(x_0) - T(x_0) \\ &= 0 \\ &= T(0). \end{aligned}$$

Since $(x_n)_{n \in \mathbb{N}} \subset X$ with $x_n \rightarrow 0$ is arbitrary, we have that for each $(x_n)_{n \in \mathbb{N}} \subset X$, $x_n \rightarrow 0$ implies that $T(x_n) \rightarrow T(0)$. Therefore T is continuous at 0.

- (3) $\implies$ (4):

Suppose that T is continuous at 0. Then there exists $\delta > 0$ such that for each $x \in X$, if $\|x\| < \delta$, then $\|Tx\| < 1$. Choose $C = \frac{2}{\delta}$. If $x = 0$, then $\|Tx\| \leq C\|x\|$. Suppose that $\|x\| \neq 0$. Define $y = \frac{\delta}{2\|x\|}x$. Then $\|y\| < \delta$. So

$$\begin{aligned} 1 &> \|Ty\| \\ &= \frac{\delta}{2\|x\|} \|Tx\| \end{aligned}$$

Thus

$$\begin{aligned} \|Tx\| &< \frac{2}{\delta} \|x\| \\ &= C\|x\| \end{aligned}$$

Hence T is bounded.

- (4) $\implies$ (1)

Suppose that T is bounded. Then there exists $C \geq 0$ such that for each $x \in X$, $\|Tx\| \leq C\|x\|$. Let $\epsilon > 0$. Choose $\delta = \frac{\epsilon}{C+1}$. Let $x, y \in X$. Suppose that $\|x - y\| < \delta$. Then

$$\begin{aligned} \|Tx - Ty\| &= \|T(x - y)\| \\ &\leq C\|x - y\| \\ &< (C + 1)\delta \\ &= \epsilon \end{aligned}$$

Thus T is continuous.

□

Definition 9.2.0.7. Let X, Y be normed vector spaces. We define the **operator norm on $L(X; Y)$** , denoted $\|\cdot\| : L(X; Y) \rightarrow [0, \infty)$, by

$$\|T\| = \inf\{C \geq 0 : \text{for each } x \in X, \|Tx\| \leq C\|x\|\}.$$

Exercise 9.2.0.8. Let X, Y be normed vector spaces. If $X \neq \{0\}$, then the operator norm on $L(X; Y)$ is given by:

1. $\|T\| = \sup_{\|x\|=1} \|Tx\|$
2. $\|T\| = \sup_{x \neq 0} \|x\|^{-1} \|Tx\|$
3. $\|T\| = \inf\{C \geq 0 : \text{for each } x \in X, \|Tx\| \leq C\|x\|\}$

Proof. Since $X \neq \{0\}$, the supremums in (1) and (2) are well defined. Let $T \in L(X; Y)$. By linearity of T , the sets over which the supremums are taken in (1) and (2) are the same. So (1) and (2) are equal.

Now, set $M = \sup_{\|x\|=1} \|Tx\|$ and $m = \inf\{C \geq 0 : \text{for each } x \in X, \|Tx\| \leq C\|x\|\}$. Let $x \in X$. If $\|x\| = 0$, then $\|Tx\| \leq M\|x\|$.

Suppose that $\|x\| \neq 0$. Then

$$\begin{aligned} \|Tx\| &= \left(\|T(\|x\|^{-1}x)\| \right) \|x\| \\ &\leq M\|x\| \end{aligned}$$

Hence $M \in \{C \geq 0 : \text{for each } x \in X, \|Tx\| \leq C\|x\|\}$ and $m \leq M$. Let $C \in \{C \geq 0 : \text{for each } x \in X, \|Tx\| \leq C\|x\|\}$. Suppose that $\|x\| = 1$. Then $\|Tx\| \leq C\|x\| = C$. So $M \leq C$. Therefore $M \leq m$. So $M = m$ and the supremum in (1) is the same as the infimum in (3). □

Note 9.2.0.9. From here on, unless stated otherwise, we assume $X \neq \{0\}$. (check if this is needed or where it is used, well it is used anywhere the norm is described by the sup over nonzero elements, otherwise the norm of a linear map $T : \{0\} \rightarrow Y$ is $\|T\| = 0$)

Exercise 9.2.0.10. Let X, Y be normed vector spaces and $T \in L(X; Y)$. Then for each $x \in X$, $\|Tx\| \leq \|T\|\|x\|$

Proof. Let $x \in X$. If $x = 0$, then $\|Tx\| \leq \|T\|\|x\|$. Suppose that $x \neq 0$. The previous exercise implies that

$$\begin{aligned} \|Tx\| &= \|T(\|x\|^{-1}x)\| \|x\| \\ &\leq \left(\sup_{\|x\|=1} \|Tx\| \right) \|x\| \\ &= \|T\| \|x\| \end{aligned}$$

□

Exercise 9.2.0.11. Let X, Y be normed vector spaces. Then $\|\cdot\| : L(X; Y) \rightarrow [0, \infty)$ is a norm on $L(X; Y)$.

Proof. Let $S, T \in L(X; Y)$ and $\alpha \in \mathbb{C}$.

- It is clear that $S + T$ is linear. For each $x \in X$,

$$\begin{aligned} \|(S + T)x\| &= \|Sx + Tx\| \\ &\leq \|Sx\| + \|Tx\| \\ &\leq \|S\|\|x\| + \|T\|\|x\| \\ &= (\|S\| + \|T\|)\|x\| \end{aligned}$$

So $S + T \in L(X; Y)$ and $\|S + T\| \leq \|S\| + \|T\|$

- It is clear that αT is linear. For each $x \in X$,

$$\begin{aligned}
 \|\alpha T\| &= \sup_{\|x\|=1} \|(\alpha T)x\| \\
 &= \sup_{\|x\|=1} |\alpha| \|Tx\| \\
 &= |\alpha| \sup_{\|x\|=1} \|Tx\| \\
 &= |\alpha| \|T\|
 \end{aligned}$$

So $\alpha T \in L(X; Y)$ and $\|\alpha T\| \leq |\alpha| \|T\|$.

- Suppose that $\|T\| = 0$. Let $x \in X$. Then

$$\begin{aligned}
 \|Tx\| &\leq \|T\| \|x\| \\
 &= 0
 \end{aligned}$$

So $Tx = 0$. Since $x \in X$ is arbitrary, we have that $T = 0$.

Therefore $L(X; Y)$ is a vector space and $\|\cdot\| : L(X; Y) \rightarrow [0, \infty)$ is a norm. □

Exercise 9.2.0.12. Let $(X, \|\cdot\|_X), (Y, \|\cdot\|_Y)$ be normed vector spaces and $T \in L(X; Y)$. Then

1. T is an isometry iff for each $x \in X$,

$$\|Tx\|_Y = \|x\|_X$$

2. If $X \neq \{0\}$ and T is an isometry, then $\|T\| = 1$.

Proof. Set $d_X := d_{\|\cdot\|_X}$ and $d_Y := d_{\|\cdot\|_Y}$.

1. • $(\implies)$:

Suppose that T is an isometry. Then for each $x \in X$,

$$\begin{aligned}
 \|Tx\| &= d_Y(Tx, 0) \\
 &= d_Y(Tx, T0) \\
 &= d_X(x, 0) \\
 &= \|x\|.
 \end{aligned}$$

- $(\impliedby)$:

Suppose that for each $x \in X$, $\|Tx\| = \|x\|$. Then for each $x_1, x_2 \in X$,

$$\begin{aligned}
 d_Y(Tx_1, Tx_2) &= \|Tx_1 - Tx_2\| \\
 &= \|T(x_1 - x_2)\| \\
 &= \|x_1 - x_2\| \\
 &= d_X(x_1, x_2).
 \end{aligned}$$

Thus T is an isometry.

2. Suppose that $X \neq \{0\}$ and T is an isometry. **the prev part (ref)** implies that $\|T\| \leq 1$. Since $X \neq \{0\}$, there exists $x_0 \in X$ such that $\|x_0\|_X = 1$. **A prev ex** then implies that

$$\begin{aligned}
 \|T\| &= \sup_{x: \|x\|_X=1} \|T(x)\|_Y \\
 &\geq \|T(x_0)\|_Y \\
 &= \|x_0\|_X \quad (\text{by ref prev part}) \\
 &= 1.
 \end{aligned}$$

Since

$$\begin{aligned} 1 &\leq \|T\| \\ &\leq 1, \end{aligned}$$

we have that $\|T\| = 1$.

□

Exercise 9.2.0.13. Let X, Y, Z be normed vector spaces, $T \in L(X; Y)$ and $S \in L(Y, Z)$. Define $ST : X \rightarrow Z$ by $STx = S(Tx)$. Then $ST \in L(X, Z)$ and $\|ST\| \leq \|S\|\|T\|$.

Proof. Clearly ST is linear. Let $x \in X$. Then

$$\begin{aligned} \|STx\| &= \|S(Tx)\| \\ &\leq \|S\|\|Tx\| \\ &\leq \|S\|\|T\|\|x\| \end{aligned}$$

So $\|ST\| \leq \|S\|\|T\|$.

□

Definition 9.2.0.14. Let X, Y be normed vector spaces and $T \in L(X; Y)$. Then T is said to be **invertible** or an **isomorphism** if T is a bijection and $T^{-1} \in L(Y, X)$.

Definition 9.2.0.15. Let X be a normed vector space. Define $GL(X) := \{T \in L(X, X) : T \text{ is invertible}\}$.

Exercise 9.2.0.16. Let X, Y be normed vector spaces. If Y is complete, then $L(X; Y)$ is complete.

Proof. Suppose that Y is complete. Let $(T_n)_{n \in \mathbb{N}} \subset L(X; Y)$. Suppose that $(T_n)_{n \in \mathbb{N}}$ is Cauchy.

- Since for each $m, n \in \mathbb{N}$, $\|T_m\| - \|T_n\| \leq \|T_m - T_n\|$, we have that $(\|T_n\|)_{n \in \mathbb{N}} \subset [0, \infty)$ is Cauchy. Hence $\lim_{n \rightarrow \infty} \|T_n\|$ exists.

Let $x \in X$ and $m, n \in \mathbb{N}$. Then

$$\begin{aligned} \|T_mx - T_nx\| &= \|(T_m - T_n)x\| \\ &\leq \|T_m - T_n\|\|x\| \end{aligned}$$

So $(T_nx)_{n \in \mathbb{N}} \subset Y$ is Cauchy and hence converges. Since Y is Hausdorff, there exists a unique $y \in Y$ such that $T_nx \rightarrow y$. Since $x \in X$ is arbitrary, we have that for each $x \in X$, $\#\{y \in Y : T_nx \rightarrow y\} = 1$. **ref ex on axiom of choice in product section or set theory section** then implies that there exists a unique $T : X \rightarrow Y$ such that $Tx = \lim_{n \rightarrow \infty} T_nx$. Since addition and scalar multiplication are continuous, for each $x_1, x_2 \in X$ and $\lambda \in \mathbb{K}$,

$$\begin{aligned} T(x_1 + \lambda x_2) &= \lim_{n \rightarrow \infty} T_n(x_1 + \lambda x_2) \\ &= \lim_{n \rightarrow \infty} [T_n(x_1) + \lambda T_n(x_2)] \\ &= \lim_{n \rightarrow \infty} T_n(x_1) + \lambda \lim_{n \rightarrow \infty} T_n(x_2) \quad (\text{by ref continuity of addition and scalar multiplication ex's}) \\ &= T(x_1) + \lambda T(x_2). \end{aligned}$$

Hence T is linear. Let $x \in X$ and $\epsilon > 0$. Choose $N \in \mathbb{N}$ such that for each $n \in \mathbb{N}$, if $n \geq N$, then $\|Tx - T_nx\| < \epsilon$. Then for each $n \in \mathbb{N}$, if $n \geq N$ we have that

$$\begin{aligned} \|Tx\| &\leq \|Tx - T_nx\| + \|T_nx\| \\ &< \epsilon + \|T_nx\| \\ &\leq \epsilon + \|T_n\|\|x\| \end{aligned}$$

Since $\lim_{n \rightarrow \infty} \|T_n\|$ exists, **continuity of addition and multiplication in $\mathbb{K}$** implies that $\|Tx\| \leq \epsilon + (\lim_{n \rightarrow \infty} \|T_n\|)\|x\|$. Since $\epsilon > 0$ is arbitrary, $\|Tx\| \leq (\lim_{n \rightarrow \infty} \|T_n\|)\|x\|$. Since $x \in X$ is arbitrary, we have that for each $x \in X$, $\|Tx\| \leq (\lim_{n \rightarrow \infty} \|T_n\|)\|x\|$. Thus $T \in L(X; Y)$ and $\|T\| \leq \lim_{n \rightarrow \infty} \|T_n\|$.

- Let $\epsilon > 0$. Since $(T_n)_{n \in \mathbb{N}}$ is Cauchy, there exists $N \in \mathbb{N}$ such that for each $m, n \in \mathbb{N}$, $n, m \geq N$ implies that $\|T_n - T_m\| < \epsilon$. Let $n \in \mathbb{N}$. Suppose that $n \geq N$. Since addition $\text{add} : X \times X \rightarrow X$, scalar multiplication $\text{scalar_mult} : \mathbb{K} \times X \rightarrow X$ and the norm $\|\cdot\| : X \rightarrow [0, \infty)$ are continuous (ref ex), we have that for each $x \in X$,

$$\begin{aligned}
 \|(T_n - T)(x)\| &= \|T_n(x) - T(x)\| \\
 &= \|T_n(x) - \lim_{m \rightarrow \infty} T_m(x)\| \\
 &= \lim_{m \rightarrow \infty} \|T_n(x) - T_m(x)\| \\
 &= \lim_{m \rightarrow \infty} \|(T_n - T_m)(x)\| \\
 &= \limsup_{m \rightarrow \infty} \|(T_n - T_m)(x)\| \\
 &\leq \limsup_{m \rightarrow \infty} \|T_n - T_m\| \|x\| \\
 &\leq \epsilon \|x\|.
 \end{aligned}$$

Hence $\|T_n - T\| \leq \epsilon$. Since $n \in \mathbb{N}$ with $n \geq N$ is arbitrary, we have that for each $n \in \mathbb{N}$, $n \geq N$ implies that $\|T_n - T\| \leq \epsilon$. Therefore $T_n \rightarrow T$.

Since $(T_n)_{n \in \mathbb{N}} \subset L(X; Y)$ with $(T_n)_{n \in \mathbb{N}}$ Cauchy is arbitrary, we have that for each $(T_n)_{n \in \mathbb{N}} \subset L(X; Y)$, $(T_n)_{n \in \mathbb{N}}$ is Cauchy implies that there exists $T \in L(X; Y)$ such that $T_n \rightarrow T$. Hence $L(X; Y)$ is complete. $\square$

9.3 Direct Sums

Definition 9.3.0.1. Let X, Y be normed vector spaces and $p \in [1, \infty]$. Let $\|\cdot\|'_p : \mathbb{R}^2 \rightarrow [0, \infty)$ denote the usual l^p norm. We define $\|\cdot\|_p : X \oplus Y \rightarrow [0, \infty)$ by

$$\|(x, y)\|_p = \|(\|x\|, \|y\|)\|'_p$$

Exercise 9.3.0.2. Let X, Y be normed vector spaces. Then

1. for each $p \in [1, \infty]$, $\|\cdot\|_p : X \oplus Y \rightarrow [0, \infty)$ is a norm on $X \oplus Y$
2. $\{\|\cdot\|_p : p \in [1, \infty]\}$ are equivalent.

Proof.

1. Let $p \in [1, \infty]$, $(x_1, y_1), (x_2, y_2) \in X \oplus Y$ and $\lambda \in \mathbb{C}$.

- Clearly if $(x_1, y_1) = (0, 0)$, then $\|S\|_p = 0$. Conversely, suppose that $\|(x_1, y_1)\|_p = 0$. Then $\|x_1\| = 0$ and $\|y_1\| = 0$. So $x_1 = 0$ and $y_1 = 0$. Therefore $S = 0$.

•

$$\begin{aligned} \|\lambda(x_1, y_1)\|_p &= \|(\|\lambda x_1\|, \|\lambda y_1\|)\|'_p \\ &= \|(|\lambda|\|x_1\|, |\lambda|\|y_1\|)\|'_p \\ &= \| |\lambda|(\|x_1\|, \|y_1\|) \|'_p \\ &= |\lambda| \|(\|x_1\|, \|y_1\|)\|'_p \\ &= |\lambda| \|(x_1, y_1)\|_p \end{aligned}$$

•

$$\begin{aligned} \|(x_1, y_1) + (x_2, y_2)\|_p &= \|(\|x_1 + x_2\|, \|y_1 + y_2\|)\|'_p \\ &\leq \|(\|x_1\| + \|x_2\|, \|y_1\| + \|y_2\|)\|'_p \\ &= \|(\|x_1\|, \|y_1\|) + (\|x_2\|, \|y_2\|)\|'_p \\ &\leq \|(\|x_1\|, \|y_1\|)\|'_p + \|(\|x_2\|, \|y_2\|)\|'_p \\ &= \|(x_1, y_1)\|_p + \|(x_2, y_2)\|_p \end{aligned}$$

2. All norms on $\mathbb{R}^2$ are equivalent.

□

Exercise 9.3.0.3. Let X, Y be Banach spaces. Then $X \oplus Y$ equipped with $\|\cdot\|_p : X \oplus Y \rightarrow [0, \infty)$ is a Banach space.

Proof.

□

Exercise 9.3.0.4. Let X, Y and Z be Banach spaces and $p \in [0, \infty]$. Equip $Y \oplus Z$ with $\|\cdot\|_p$. Let $T \in L(X, Y \oplus Z)$ with $T = (T_Y, T_Z)$. Then $T_Y \in L(X; Y)$ and $T_Z \in L(X, Z)$.

Proof. Let $x \in X$. Then $\|T_Y(x)\|, \|T_Z(x)\| \leq$

FINISH!!!

□

Definition 9.3.0.5. Let X, Y and Z be Banach spaces and $p \in [0, \infty]$. Let $\|\cdot\|'_p : \mathbb{R}^2 \rightarrow [0, \infty)$ denote the usual l^p norm. Equip $Y \oplus Z$ with $\|\cdot\|_p$. Let $T \in L(X, Y \oplus Z)$ with $T = (T_Y, T_Z)$. Define $\|T\|_p : L(X, Y \oplus Z) \rightarrow [0, \infty)$ by

$$\|T\|_p = \|(\|T_Y\|, \|T_Z\|)\|'_p$$

Exercise 9.3.0.6. Let X, Y and Z be Banach spaces and $p \in [0, \infty]$. Then $\|T\|_p : L(X, Y \oplus Z) \rightarrow [0, \infty)$ is a norm on $L(X, Y \oplus Z)$.

Proof. Let $\lambda \in \mathbb{C}$ and $S, T \in L(X, Y \oplus Z)$ with $S = (S_Y, S_Z)$ and $T = (T_Y, T_Z)$.

- Clearly if $S = 0$, then $\|S\|_p = 0$. Conversely, suppose that $\|S\|_p = 0$. Then $\|S_Y\| = 0$ and $\|S_Z\| = 0$. So $S_Y = 0$ and $S_Z = 0$. Therefore $S = 0$.

•

$$\begin{aligned}
\|\lambda S\|_p &= \|(\|\lambda S_Y\|, \|\lambda S_Z\|)\|'_p \\
&= \|(|\lambda|\|S_Y\|, |\lambda|\|S_Z\|)\|'_p \\
&= \|\lambda\|(\|S_Y\|, \|S_Z\|)\|'_p \\
&= |\lambda|(\|S_Y\|, \|S_Z\|)\|'_p \\
&= |\lambda|\|S\|_p
\end{aligned}$$

•

$$\begin{aligned}
\|S + T\|_p &= \|(\|S_Y + T_Y\|, \|S_Z + T_Z\|)\|'_p \\
&\leq \|(\|S_Y\| + \|T_Y\|, \|S_Z\| + \|T_Z\|)\|'_p \\
&= \|(\|S_Y\|, \|S_Z\|) + (\|T_Y\|, \|T_Z\|)\|'_p \\
&\leq \|(\|S_Y\|, \|S_Z\|)\|'_p + \|(\|T_Y\|, \|T_Z\|)\|'_p \\
&= \|S\|_p + \|T\|_p
\end{aligned}$$

So $\|\cdot\|_p : L(X, Y \oplus Z) \rightarrow [0, \infty)$ is a norm on $L(X, Y \oplus Z)$. □

Exercise 9.3.0.7. Let X, Y and Z be Banach spaces and $p \in [0, \infty]$. Equip $Y \oplus Z$ with $\|\cdot\|_p$. Let $T \in L(X, Y \oplus Z)$ with $T = (T_Y, T_Z)$. Then $\|T\| \leq 2^{1/p}\|T\|_p$.

Proof. Let $x \in X$. If $p < \infty$, then

$$\begin{aligned}
\|T(x)\|_p &= \|(\|T_Y(x)\|, \|T_Z(x)\|)\|_p \\
&= \|(\|T_Y(x)\|, \|T_Z(x)\|)\|'_p \\
&= \left(\|T_Y(x)\|^p + \|T_Z(x)\|^p \right)^{1/p} \\
&\leq \left(\|T_Y\|^p \|x\|^p + \|T_Z\|^p \|x\|^p \right)^{1/p} \\
&\leq \left[(\|T_Y\|^p + \|T_Z\|^p) \|x\|^p + (\|T_Y\|^p + \|T_Z\|^p) \|x\|^p \right]^{1/p} \\
&= \left[2(\|T_Y\|^p + \|T_Z\|^p) \|x\|^p \right]^{1/p} \\
&= 2^{1/p} \|T\|_p \|x\|
\end{aligned}$$

Hence $\|T\| \leq 2^{1/p}\|T\|_p$. If $p = \infty$, then

$$\begin{aligned}
\|T(x)\|_\infty &= \max(\|T_Y(x)\|, \|T_Z(x)\|) \\
&\leq \max(\|T_Y\| \|x\|, \|T_Z\| \|x\|) \\
&\leq \max \left[\max(\|T_Y\|, \|T_Z\|) \|x\|, \max(\|T_Y\|, \|T_Z\|) \|x\| \right] \\
&= \max(\|T_Y\|, \|T_Z\|) \|x\| \\
&= \|T\|_\infty \|x\|
\end{aligned}$$

Hence

$$\begin{aligned}\|T\| &\leq \|T\|_\infty \\ &= 2^{1/\infty} \|T\|_\infty\end{aligned}$$

□

Exercise 9.3.0.8. Let X and $X_1, \dots, X_n$ be Banach spaces and $p \in [0, \infty]$. Equip $\bigoplus_{j=1}^n X_j$ with $\|\cdot\|_p$. Let $T \in L(X, \bigoplus_{j=1}^n X_j)$. Then $\|T\| \leq n^{1/p} \|T\|_p$.

Proof. Similar to the previous exercise.

□

9.4 Quotient Spaces

Definition 9.4.0.1. Let X be a normed vector space and $M \subset X$ a closed subspace. Define $\|\cdot\| : X/M \rightarrow [0, \infty)$ by

$$\|x + M\|_{X/M} := \inf_{y \in M} \|x + y\|$$

We call $\|\cdot\|_{X/M}$ the **subspace norm on X/M**

Note 9.4.0.2. When the context is clear, we write $\|\cdot\|$ in place of $\|\cdot\|_{X/M}$.

Exercise 9.4.0.3. Let X be a normed vector space and $M \subsetneq X$ a proper, closed subspace of M . Then

1. The previously defined subspace norm on X/M is well defined and is a norm.
2. For each $\epsilon > 0$, there exists $x \in X$ such that $\|x\| = 1$ and $\|x + M\| \geq 1 - \epsilon$.
3. The projection map $\pi : X \rightarrow X/M$ defined by $\pi(x) = x + M$ is continuous and $\|\pi\| = 1$.
4. If X is complete, then X/M is complete.

Proof.

1. Let $x, y \in X$ and $\alpha \in \mathbb{C}$. Suppose that $x + M = y + M$. Then there exists $m \in M$ such that $x = y + m$. Since M is a subspace, the map $T : M \rightarrow M$ given by $Tx = x + m$ is a bijection. So

$$\inf_{z \in M} \|y + m + z\| = \inf_{z \in M} \|y + z\|$$

which implies that

$$\begin{aligned} \|x + M\| &= \inf_{z \in M} \|x + z\| \\ &= \inf_{z \in M} \|y + m + z\| \\ &= \inf_{z \in M} \|y + z\| \\ &= \|y + M\| \end{aligned}$$

So $\|\cdot\| : X/M \rightarrow [0, \infty)$ is well defined.

We observe that for each $z, w \in M$,

$$\|x + y + z\| \leq \|x + w\| + \|y + w + z\|$$

Taking infimums over M with respect to z in this inequality implies that for each $w \in M$,

$$\begin{aligned} \inf_{z \in M} \|x + y + z\| &\leq \inf_{z \in M} (\|x + w\| + \|y + w + z\|) \\ &= \|x + w\| + \inf_{z \in M} \|y + w + z\| \end{aligned}$$

Again we use the fact that for each $w \in M$,

$$\inf_{z \in M} \|y + w + z\| = \inf_{z \in M} \|y + z\|$$

This implies that for each $w \in M$,

$$\inf_{z \in M} \|x + y + z\| \leq \|x + w\| + \inf_{z \in M} \|y + z\|$$

Therefore, taking infimums over M with respect to w in this inequality yields

$$\begin{aligned} \|x + y + M\| &= \inf_{z \in M} \|x + y + z\| \\ &\leq \inf_{w \in M} \left(\|x + w\| + \inf_{z \in M} \|y + z\| \right) \\ &= \inf_{w \in M} \|x + w\| + \inf_{z \in M} \|y + z\| \\ &= \|x + M\| + \|y + M\| \end{aligned}$$

If $\alpha = 0$, then $\alpha x = 0$. Choosing $z = 0 \in M$ gives $\|\alpha x + M\| = 0 = |\alpha| \|x + M\|$. Suppose that $\alpha \neq 0$. Then the map $T : M \rightarrow M$ given by $Tx = \alpha^{-1}x$ is a bijection and thus $\inf_{z \in M} \|x + \alpha^{-1}z\| = \inf_{z \in M} \|x + z\|$. Hence we have that

$$\begin{aligned} \|\alpha x + M\| &= \inf_{z \in M} \|\alpha x + z\| \\ &= \inf_{z \in M} |\alpha| \|x + \alpha^{-1}z\| \\ &= |\alpha| \inf_{z \in M} \|x + \alpha^{-1}z\| \\ &= |\alpha| \inf_{z \in M} \|x + z\| \\ &= |\alpha| \|x + M\| \end{aligned}$$

Suppose that $\|x\| = 0$. Choose a sequence $(z_n)_{n \in \mathbb{N}} \subset M$ such that

$$\begin{aligned} \lim_{n \rightarrow \infty} \|x - z_n\| &= \inf_{z \in M} \|x + z\| \\ &= 0 \end{aligned}$$

Then $\lim_{n \rightarrow \infty} z_n = x$. Since M is closed, $x \in M$. Hence $x + M = 0 + M$.

2. Since M is a proper subspace, there exists $v \in X$ such that $v \notin M$. Then $\|v + M\| \neq 0$. Let $\epsilon > 0$. Then

$$\begin{aligned} (1 - \epsilon)^{-1} \|v + M\| &> \|v + M\| \\ &= \inf_{y \in M} \|v + y\| \end{aligned}$$

So there exists $z \in M$ such that $\|v + z\| < (1 - \epsilon)^{-1} \|v + M\|$. Since $v + M \neq 0 + M$, we have that $v + z \neq 0$. Choose $x = \|v + z\|^{-1}(v + z)$. Then $\|x\| = 1$ and

$$\begin{aligned} \|x + M\| &= \|v + z\|^{-1} \|v + z + M\| \\ &= \|v + z\|^{-1} \|v + M\| \\ &> 1 - \epsilon \end{aligned}$$

3. Let $x \in X$. Taking $z = 0$, we see that $\|\pi(x)\| = \|x + M\| \leq \|x + z\| = \|x\|$. So π is bounded and in particular,

$$\sup_{\|x\|=1} \|\pi(x)\| \leq 1$$

From (2) we see that

$$\sup_{\|x\|=1} \|\pi(x)\| \geq 1$$

Hence $\|\pi\| = 1$.

4. Suppose that X is complete. Let $(x_i + M)_{i \in \mathbb{N}} \subset X/M$. Suppose that $\sum_{i \in \mathbb{N}} \|x_i + M\| < \infty$. Let $\epsilon > 0$. Then for each $i \in \mathbb{N}$, there exists $z_i \in M$ such that $\|x_i + z_i\| < \|x_i + M\| + \epsilon 2^{-i}$. Define the sequence $(a_i)_{i \in \mathbb{N}} \subset X$ by $a_i = x_i + z_i$. Then we have

$$\begin{aligned} \sum_{i \in \mathbb{N}} \|a_i\| &= \sum_{i \in \mathbb{N}} \|x_i + z_i\| \\ &\leq \sum_{i \in \mathbb{N}} \left(\|x_i + M\| + \epsilon 2^{-i} \right) \\ &= \sum_{i \in \mathbb{N}} \|x_i + M\| + \epsilon \end{aligned}$$

Since $\epsilon > 0$ is arbitrary, it follows that

$$\sum_{i \in \mathbb{N}} \|a_i\| \leq \sum_{i \in \mathbb{N}} \|x_i + M\| < \infty$$

Since X is complete, $\sum_{i=1}^{\infty} a_i$ converges in X . Define $(s_n)_{n \in \mathbb{N}} \subset X$ and $s \in X$ by $s_n = \sum_{i=1}^n a_i$ and $s = \sum_{i=1}^{\infty} a_i$. Since $\lim_{n \rightarrow \infty} s_n = s$, and $\pi : X \rightarrow X/M$ is continuous, it follows that $\lim_{n \rightarrow \infty} \pi(s_n) = \pi(s)$. Since

$$\begin{aligned} \pi(s_n) &= \sum_{i=1}^n a_i + M \\ &= \sum_{i=1}^n x_i + M \end{aligned}$$

We have that $\sum_{i=1}^{\infty} x_i + M$ converges which implies that X/M is complete.

□

Exercise 9.4.0.4. Let X, Y be normed vector spaces and $T \in L(X; Y)$. Then

1. $\ker T$ is closed
2. there exists a unique map $S : X/\ker T \rightarrow T(X)$ such that $T = S \circ \pi$. Furthermore S is a bounded linear bijection and $\|S\| = \|T\|$.

Proof.

1. Since T is continuous and $\ker T = T^{-1}(\{0\})$, we have that $\ker T$ is closed.
2. Suppose that there exists $S_1, S_2 \in L(X/\ker T, T(X))$ such that $T = S_1 \circ \pi$ and $T = S_2 \circ \pi$. Let $x \in X$. Then

$$S_1(x + \ker T) = S_1(\pi(x)) = T(x) = S_2(\pi(x)) = S_2(x + \ker T)$$

So $S_1 = S_2$. Therefore such a map is unique.

Define $S : X/\ker T \rightarrow T(X)$ by $S(x + \ker T) = T(x)$. Then S is clearly a linear bijection that satisfies $T = S \circ \pi$. Let $x \in X$ and $z \in \ker T$. Then

$$\begin{aligned} \|S(x + \ker T)\| &= \|T(x)\| \\ &= \|T(x + z)\| \\ &\leq \|T\| \|x + z\| \end{aligned}$$

Thus

$$\|S(x + \ker T)\| \leq \|T\| \inf_{z \in \ker T} \|x + z\| = \|T\| \|x + \ker T\|$$

So S is bounded and $\|S\| \leq \|T\|$. This implies that

$$\|T\| = \|S \circ \pi\| \leq \|S\| \|\pi\| = \|S\|$$

Thus $\|S\| = \|T\|$.

□

Exercise 9.4.0.5. Let X, Y be normed vector spaces. Define $\phi : L(X; Y) \times X \rightarrow Y$ by $\phi(T, x) = Tx$. Then ϕ is continuous.

Proof. Let $(T_1, x_1) \in L(X; Y) \times X$ and $\epsilon > 0$. Choose $\delta = \min\{\frac{\epsilon}{2(\|x_1\| + \|T_1\| + 1)}, \frac{\sqrt{\epsilon}}{\sqrt{2}}\}$. Let $(t_2, x_2) \in L(X; Y) \times X$. Suppose that

$$\|(T_1, x_1) - (T_2, x_2)\| = \max\{\|T_1 - T_2\|, \|x_1 - x_2\|\} < \delta$$

Then

$$\begin{aligned} \|\phi(T_1, x_1) - \phi(T_2, x_2)\| &= \|T_1x_1 - T_2x_2\| \\ &= \|T_1x_1 - T_2x_1 + T_2x_1 - T_2x_2\| \\ &\leq \|(T_1 - T_2)x_1\| + \|T_2(x_1 - x_2)\| \\ &\leq \|T_1 - T_2\|\|x_1\| + \|T_2\|\|x_1 - x_2\| \\ &\leq \|T_1 - T_2\|\|x_1\| + (\|T_1 - T_2\| + \|T_1\|)\|x_1 - x_2\| \\ &< \delta\|x_1\| + (\delta + \|T_1\|)\delta \\ &= \delta(\|T_1\| + \|x_1\|) + \delta^2 \\ &< \frac{\epsilon}{2} + \frac{\epsilon}{2} \\ &= \epsilon \end{aligned}$$

So ϕ is continuous. □

Exercise 9.4.0.6. Let X be a normed vector space and $M \subset X$ a subspace. Then $\overline{M}$ is a subspace.

Proof. Let $x, y \in \overline{M}$ and $\alpha \in \mathbb{C}$. Then there exist sequences $(x_n)_{n \in \mathbb{N}} \subset M$ and $(y_n)_{n \in \mathbb{N}} \subset M$ such that $x_n \rightarrow x$ and $y_n \rightarrow y$. Since M is a subspace, $(x_n + y_n)_{n \in \mathbb{N}} \subset M$ and $(\alpha x_n)_{n \in \mathbb{N}} \subset M$. Since addition and scalar multiplication are continuous, we have that $x_n + y_n \rightarrow x + y$ and $\alpha x_n \rightarrow \alpha x$. Thus $x + y \in \overline{M}$ and $\alpha x \in \overline{M}$ and hence $\overline{M}$ is a subspace. □

9.5 Applications of the Hahn-Banach Theorem

Definition 9.5.0.1. Let X be a normed vector space over $\mathbb{C}$, and $T : X \rightarrow \mathbb{C}$. Then T is said to be a **bounded linear functional** on X if $T \in L(X, \mathbb{C})$. We define the **dual space** of X , denoted X^* , by $X^* = L(X, \mathbb{C})$.

Note 9.5.0.2. We define X^* similarly when X is a normed vector space over $\mathbb{R}$.

Definition 9.5.0.3. Let X be a normed vector space and $p : X \rightarrow \mathbb{R}$ a sublinear functional. Then p is said to be **bounded** if there exists $M > 0$ such that for each $x \in X$, $p(x) \leq M\|x\|$.

Exercise 9.5.0.4. Let X be a normed vector space and $p : X \rightarrow \mathbb{R}$ a sublinear functional. Then p is bounded iff p is Lipschitz.

Proof. Suppose that p is bounded. Then there exists $M > 0$ such that for each $x \in X$, $p(x) \leq M\|x\|$. Let $x, y \in X$. Then the previous exercise implies that

$$\begin{aligned} -M\|x - y\| &= -M\|y - x\| \\ &\leq -p(y - x) \\ &\leq p(x) - p(y) \\ &\leq p(x - y) \\ &\leq M\|x - y\| \end{aligned}$$

So that

$$|p(x) - p(y)| \leq M\|x - y\|$$

and p is Lipschitz. Conversely, suppose that p is Lipschitz. Then there exists $M > 0$ such that for each $x, y \in X$, $|p(x) - p(y)| \leq M\|x - y\|$. Let $x \in X$. Then

$$\begin{aligned} p(x) &\leq |p(x)| \\ &= |p(x) - p(0)| \\ &\leq M\|x - 0\| \\ &\leq M\|x\| \end{aligned}$$

So p is bounded. □

Exercise 9.5.0.5. Let X be a normed vector space, $p : X \rightarrow \mathbb{R}$ a bounded sublinear functional and $\phi : X \rightarrow \mathbb{R}$ a linear functional. If $\phi \leq p$, then $\phi \in X^*$.

Proof. Since p is Lipschitz, there exists $M > 0$ such that for each $x \in X$,

$$\begin{aligned} p(x) &\leq |p(x)| \\ &\leq M\|x\| \end{aligned}$$

Let $x \in X$. Then

$$\begin{aligned} \phi(x) &\leq p(x) \\ &\leq |p(x)| \\ &\leq M\|x\| \end{aligned}$$

and therefore

$$\begin{aligned} -M\|x\| &= -M\| -x \| \\ &\leq -p(-x) \\ &\leq -\phi(-x) \\ &= \phi(x) \end{aligned}$$

So that $|\phi(x)| \leq M\|x\|$ and $\phi \in X^*$. □

Exercise 9.5.0.6. Let X be a normed vector space and $p : X \rightarrow \mathbb{R}$ a bounded sublinear functional. Then there exists $\phi \in X^*$ such that for each $x \in X$, $\phi(x) \leq p(x)$.

Proof. A previous exercise in the section on sublinear functionals in the topological vector space chapter implies there exists $\phi : X \rightarrow \mathbb{R}$ such that ϕ is linear and $\phi \leq p$. The previous exercise implies that $\phi \in X^*$. $\square$

Exercise 9.5.0.7. Equivalency of linearity (Bounded Case)

Let X be a normed vector space and $p : X \rightarrow \mathbb{R}$ a bounded sublinear functional. Then the following are equivalent:

1. there exists a unique $\phi \in X^*$ such that $\phi \leq p$
2. for each $x \in X$, $-p(-x) = p(x)$
3. p is linear

Proof. Basically the same as last time. $\square$

Exercise 9.5.0.8. Let X be a normed vector space, $M \subset X$ a subspace and $f \in M^*$. Then there exists $F \in X^*$ such that $\|F\| = \|f\|$ and $F|_M = f$.

Proof. If $f = 0$, Choose $F = 0$. Suppose $f \neq 0$. Then $\|f\| \neq 0$ and there exists $x_0 \in M$ such that $x_0 \neq 0$. Thus $\|f\| \neq 0$. Define $p : X \rightarrow [0, \infty)$ by $p(x) = \|f\|\|x\|$. Then p is a sublinear functional on X and for each $x \in M$, $|f(x)| \leq p(x)$. So there exists a linear functional $F : X \rightarrow \mathbb{C}$ such that for each $x \in X$, $|F(x)| \leq p(x) = \|f\|\|x\|$ and $F|_M = f$. Thus $F \in X^*$ with $\|F\| \leq \|f\|$. Also

$$\|F\| = \sup_{\substack{x \in X \\ \|x\|=1}} |F(x)| \geq \sup_{\substack{x \in M \\ \|x\|=1}} |F(x)| = \sup_{\substack{x \in M \\ \|x\|=1}} |f(x)| = \|f\|$$

So $\|F\| = \|f\|$. $\square$

Exercise 9.5.0.9. Let X be a normed vector space, $M \subsetneq X$ a proper closed subspace and $x \in X \setminus M$. Then there exists $F \in X^*$ such that $F|_M = 0$, $\|F\| = 1$ and $F(x) = \|x + M\| \neq 0$.

Hint: Consider $f : M + \mathbb{C}x \rightarrow \mathbb{C}$ defined by $f(m + \lambda x) = \lambda\|x + M\|$.

Proof. Define $f : M + \mathbb{C}x \rightarrow \mathbb{C}$ as above. Clearly f is linear and $f|_M = 0$. Let $m \in M$ and $\lambda \in \mathbb{C}$. If $\lambda = 0$, then $|f(m + \lambda x)| = 0 \leq \|m + \lambda x\|$. Suppose that $\lambda \neq 0$. Then

$$\begin{aligned} |f(m + \lambda x)| &= |\lambda|\|x + M\| \\ &= \|\lambda x + M\| \\ &= \inf_{z \in M} \|z + \lambda x\| \\ &\leq \|m + \lambda x\| \end{aligned}$$

So $f \in (M + \mathbb{C}x)^*$ and $\|f\| \leq 1$. Let $\epsilon > 0$. A previous exercise tells us that there exist $m \in M, \lambda \in \mathbb{C}$ such that $\|m + \lambda x\| = 1$ and $\|m + \lambda x + M\| > 1 - \epsilon$. Then

$$\begin{aligned} |f(m + \lambda x)| &= |\lambda|\|x + M\| \\ &= \|\lambda x + M\| \\ &= \|m + \lambda x + M\| \\ &> 1 - \epsilon \end{aligned}$$

So

$$\|f\| = \sup_{\substack{z \in M + \mathbb{C}x \\ \|z\|=1}} |f(z)| \geq 1$$

Hence $\|f\| = 1$. The same exercise also tells us that $f(x) = \|x + M\| \neq 0$. Using the previous exercise, there exists $F \in X^*$ such that $\|F\| = \|f\| = 1$ and $F|_{M + \mathbb{C}x} = f$. $\square$

Exercise 9.5.0.10. Let X be a normed vector space and $x \in X$. If $x \neq 0$, then there exists $F \in X^*$ such that $\|F\| = 1$ and $F(x) = \|x\|$.

Proof. Define $f : \mathbb{C}x \rightarrow \mathbb{C}$ by $f(\lambda x) = \lambda\|x\|$. Then f is linear and $f(x) = \|x\|$. Clearly

$$\sup_{\substack{z \in \mathbb{C}x \\ \|z\|=1}} |f(z)| = 1$$

So $f \in (\mathbb{C}x)^*$ and $\|f\| = 1$. By a previous exercise, there exists $F \in X^*$ such that $\|F\| = \|f\| = 1$ and $F|_{\mathbb{C}x} = f$. □

Exercise 9.5.0.11. Let X be a normed vector space and $x \in X$. Then $x = 0$ iff for each $\phi \in X^*$, $\phi(x) = 0$.

Proof. Clear by previous exercise. □

Exercise 9.5.0.12. Let X be a normed vector space. Then X^* separates the points of X .

Proof. Let $x, y \in X$. Suppose that $x \neq y$. Then $x - y \neq 0$. The previous exercise implies that there exists $F \in X^*$ such that $\|F\| = 1$ and

$$F(x) - F(y) = F(x - y) = \|x - y\| \neq 0$$

Thus $F(x) \neq F(y)$ and X^* separates the points of X . □

Exercise 9.5.0.13. Let X be a normed vector space and $f : X \rightarrow \mathbb{C}$ a linear functional on X . Then f is bounded iff $\ker f$ is closed.

Proof. Suppose that f is continuous. Since $\{0\}$ is closed, we have that $\ker f = f^{-1}(\{0\})$ is closed. Conversely, suppose that $\ker f$ is closed. If $\ker f = X$, then $f = 0$ and f is continuous. Suppose that $\ker f \neq X$. Then $\ker f$ is a proper, closed subspace of X . A previous exercise tells us that there exists $x \in X$ such that $\|x\| = 1$ and $\|x + \ker f\| > \frac{1}{2}$. Let $y \in X$. Suppose that $\|y\| < \frac{1}{2}$. Then for each $z \in \ker f$,

$$\begin{aligned} \|z - (x + y)\| &= \|(z - x) - y\| \\ &\geq \|z - x\| - \|y\| \\ &> \frac{1}{2} - \frac{1}{2} \\ &= 0 \end{aligned}$$

So $x + y \notin \ker f$. Therefore $f(B(x, \frac{1}{2})) \cap \{0\} = \emptyset$. If $f(B(x, \frac{1}{2}))$ is unbounded, then $f(B(x, \frac{1}{2})) = \mathbb{C}$ by linearity. This is a contradiction since $0 \notin f(B(x, \frac{1}{2}))$. So There exists $s > 0$ such that $f(B(x, \frac{1}{2})) \subset B(0, s)$ and thus f is bounded. □

Exercise 9.5.0.14. Let X be a normed vector space.

1. Let $M \subset X$ be a proper closed subspace of X and $x \in X \setminus M$. Then $M + \mathbb{C}x$ is closed.
2. Let $M \subset X$ be a finite dimensional subspace of X . Then M is closed.

Proof. 1. Let $y \in X$ and $(y_n)_{n \in \mathbb{N}} \subset M + \mathbb{C}x$. Suppose that $y_n \rightarrow y$. If $y \in M$, then $y \in M + \mathbb{C}x$. Suppose that $y \notin M$. For each $n \in \mathbb{N}$, there exists $m_n \in M$ and $\lambda_n \in \mathbb{C}$ such that $y_n = m_n + \lambda_n x$. A previous exercise tells us that there exists $F \in X^*$ such that $\|F\| = 1$, $F|_M = 0$ and $F(x) = \|x + M\| \neq 0$. Since F is continuous, $F(y_n) \rightarrow F(y)$. Since for each $n \in \mathbb{N}$,

$$F(y_n) = F(m_n + \lambda_n x) = F(m_n) + \lambda_n F(x) = \lambda_n F(x)$$

we have that $\lambda_n F(x) \rightarrow F(y)$. Since $F(x) \neq 0$, this implies that $\lambda_n \rightarrow F(y)F(x)^{-1}$. It follows that $\lambda_n x \rightarrow F(y)F(x)^{-1}x$. Since for each $n \in \mathbb{N}$, $m_n = y_n - \lambda_n x$, we know that $m_n \rightarrow y - F(y)F(x)^{-1}x$. Since $(m_n)_{n \in \mathbb{N}} \subset M$ and M is closed, we have that $y - F(y)F(x)^{-1}x \in M$ and therefore $y \in M + \mathbb{C}x$. Hence $M + \mathbb{C}x$ is closed.

2. If $M = X$, then M is closed. Suppose that $M \neq X$. Let $(x_i)_{i=1}^n$ be a basis for M . Define $N_0 = \{0\}$ and for each $i = 1, 2, \dots, n$, define $N_i = N_{i-1} + \mathbb{C}x_i$. Since N_0 is a proper closed subspace of X and $x_1 \in X \setminus N_0$, (1) implies that N_1 is closed. Proceed inductively to obtain that $M = N_n$ is closed. □

Exercise 9.5.0.15. Let X be an infinite-dimensional normed vector space.

1. There exists a sequence $(x_n)_{n \in \mathbb{N}} \subset X$ such that for each $m, n \in \mathbb{N}$, $\|x_n\| = 1$ and if $m \neq n$, then $\|x_m - x_n\| > \frac{1}{2}$.
2. X is not locally compact.

Proof.

1. Define $N_0 = \{0\}$. Then N_0 is a closed proper subspace of X . Choose $x_1 \in X$ such that $\|x_1\| = 1$. Using the results of previous exercises, we proceed inductively. For each $n \geq 2$ we define $N_{n-1} = \text{span}(x_1, x_2, \dots, x_{n-1})$. Then N_{n-1} is a closed proper subspace of X . Thus we may choose $x_n \in X$ such that $\|x_n\| = 1$ and $\|x_n + N_{n-1}\| > \frac{1}{2}$. Let $m, n \in \mathbb{N}$. Suppose that $m < n$. Then $x_m \in N_{n-1}$. Thus $\|x_n - x_m\| \geq \|x_n + N_{n-1}\| > \frac{1}{2}$.
2. Suppose that X is locally compact. Then $\bar{B}(0, 1)$ is compact and therefore sequentially compact. Using $(x_n)_{n \in \mathbb{N}} \subset \bar{B}(0, 1)$ defined in (1), we see that there exists a subsequence $(x_{n_k})_{k \in \mathbb{N}}$, $x \in \bar{B}(0, 1)$ such that $x_{n_k} \rightarrow x$. Then $(x_{n_k})_{k \in \mathbb{N}}$ is Cauchy. So there exists $N \in \mathbb{N}$ such that for each $j, k \in \mathbb{N}$, if $j, k \geq N$, then $\|x_{n_j} - x_{n_k}\| < \frac{1}{2}$. Then $\|x_{n_N} - x_{n_{N+1}}\| < \frac{1}{2}$. This is a contradiction since by construction, $\|x_{n_N} - x_{n_{N+1}}\| > \frac{1}{2}$. Thus X is not locally compact.

□

9.6 Applications of the Baire Category Theorem

Theorem 9.6.0.1. Open Mapping Theorem:

Let X, Y be Banach spaces and $T \in L(X; Y)$. If T is surjective, then T is open.

Corollary 9.6.0.2. Let X, Y be Banach spaces and $T \in L(X; Y)$. If T is a bijection, then $T^{-1} \in L(X; Y)$.

Definition 9.6.0.3. [move to set theory/intro to math notes](#) Let X, Y be sets and $f : X \rightarrow Y$. We define the **graph of f** , denoted $\Gamma(f) \subset X \times Y$, by $\Gamma(f) = \{(x, y) \in X \times Y : f(x) = y\}$.

Theorem 9.6.0.4. Closed Graph Theorem:

Let X, Y be Banach spaces and $T : X \rightarrow Y$ a linear map. If $\Gamma(T)$ is closed, then $T \in L(X; Y)$.

Note 9.6.0.5. We recall that $\Gamma(T)$ is closed iff for each $(x_n)_{n \in \mathbb{N}} \subset X$, $x \in X$ and $y \in Y$, $x_n \rightarrow x$ and $T(x_n) \rightarrow y$ implies that $T(x) = y$.

Exercise 9.6.0.6. Uniform Boundedness Principle:

Let X, Y be Banach spaces and $S \subset L(X; Y)$. If for each $x \in X$,

$$\sup_{T \in S} \|Tx\| < \infty$$

then

$$\sup_{T \in S} \|T\| < \infty$$

Proof. [Finish!!!](#) □

Exercise 9.6.0.7. Let μ be counting measure on $(\mathbb{N}, \mathcal{P}(\mathbb{N}))$. Define $h : \mathbb{N} \rightarrow \mathbb{N}$ and ν on $(\mathbb{N}, \mathcal{P}(\mathbb{N}))$ by $h(n) = n$ and $d\nu = h d\mu$. Define $X = L^1(\nu)$ and $Y = L^1(\mu)$. Equip both X and Y with the L^1 norm with respect to μ .

1. We have that X is a proper subspace of Y and therefore X is not complete.
2. Define $T : X \rightarrow Y$ by $Tf(n) = nf(n)$. Then T is linear, $\Gamma(T)$ is closed, and T is unbounded.
3. Define $S : Y \rightarrow X$ by $Sg(n) = \frac{1}{n}g(n)$. Then $S \in L(Y, X)$, S is surjective and S is not open.

Proof.

1. Note that for each $f : \mathbb{N} \rightarrow \mathbb{C}$,

$$\begin{aligned} \|f\|_{\mu,1} &= \sum_{n=1}^{\infty} |f(n)| \\ &\leq \sum_{n=1}^{\infty} n|f(n)| \\ &= \|f\|_{\nu,1} \end{aligned}$$

Hence X is a subspace of Y . Define $f : \mathbb{N} \rightarrow \mathbb{C}$ by $f(n) = \frac{1}{n^2}$. Then

$$\|f\|_{\mu,1} = \sum_{n=1}^{\infty} \frac{1}{n^2} < \infty$$

So $f \in Y$. However

$$\|f\|_{\nu,1} = \sum_{n=1}^{\infty} \frac{1}{n} = \infty$$

So $f \notin X$. Thus X is a proper subspace of Y . Let $g \in Y$ and $\epsilon > 0$. Since the simple functions are dense in $L^1(\mu)$, there exists $\phi \in L^1(\mu)$ such that $\|g - \phi\|_{\mu,1} < \epsilon$. Then there exist $(c_i)_{i=1}^k \subset \mathbb{C}$ and $(E_i)_{i=1}^k \subset \mathcal{P}(\mathbb{N})$ such that for each $i, j \in \{1, 2, \dots, k\}$, E_i is finite, $i \neq j$ implies that $E_i \cap E_j = \emptyset$ and

$$\phi = \sum_{i=1}^k c_i \chi_{E_i}$$

Define $c = \max\{|c_i| : i = 1, 2, \dots, k\}$ and $m = \max \left[\bigcup_{i=1}^k E_i \right]$. Then

$$\begin{aligned} \|\phi\|_{\nu,1} &= \sum_{n=1}^m n|\phi(n)| \\ &\leq \sum_{n=1}^m mc \\ &= cm^2 \\ &< \infty \end{aligned}$$

Hence $\phi \in X$ and X is dense in Y . Since X is a dense, proper subspace, it is not closed. Since Y is complete and $X \subset Y$ is not closed, we have that X is not complete.

2. Clearly T is linear. Let $(f_j)_{j \in \mathbb{N}} \subset X$, $f \in X$ and $g \in Y$. Suppose that $f_j \xrightarrow{L^1(\mu)} f$ and $Tf_j \xrightarrow{L^1(\mu)} g$. Note that for each $j \in \mathbb{N}$ and $n \in \mathbb{N}$,

$$|f_j(n) - f(n)| \leq \sum_{n=1}^{\infty} |f_j(n) - f(n)| = \|f_j - f\|_{\mu,1}$$

and

$$|nf_j(n) - g(n)| \leq \sum_{n=1}^{\infty} |nf_j(n) - g(n)| = \|Tf_j - g\|_{\mu,1}$$

Thus for each $n \in \mathbb{N}$, $f_j(n) \xrightarrow{j} f(n)$ and $nf_j(n) \xrightarrow{j} g(n)$. This implies that for each $n \in \mathbb{N}$, $nf(n) = g(n)$. Thus $Tf = g$ which implies that $\Gamma(T)$ is closed. Suppose, for the sake of contradiction, that T is bounded. Then there exists $C \geq 0$ such that for each $f \in X$, $\|Tf\|_{\mu,1} \leq C\|f\|_{\mu,1}$. Choose $n \in \mathbb{N}$ such that $n > C$. Define $f : \mathbb{N} \rightarrow \mathbb{C}$ by $f = \chi_{\{n\}}$. As established above, $S^+ \subset L^1(\mu)$. Then $\|f\|_{\mu,1} = 1$ and

$$\begin{aligned} \|Tf\|_{\mu,1} &= n \\ &> C \\ &= C\|f\|_{\mu,1} \end{aligned}$$

which is a contradiction. So T is unbounded.

3. Clearly S is linear. Let $g \in Y$. Then

$$\begin{aligned} \|Sg\|_{\mu,1} &= \sum_{n=1}^{\infty} \frac{1}{n} |g(n)| \\ &\leq \sum_{n=1}^{\infty} |g(n)| \\ &= \|g\|_{\mu,1} \end{aligned}$$

So S is bounded and $\|S\| \leq 1$. Thus $S \in L(Y, X)$. Let $f \in X$. Define $g : \mathbb{N} \rightarrow \mathbb{C}$ by $g(n) = nf(n)$. By definition, $g \in Y$ and we have that

$$\begin{aligned} Sg(n) &= \frac{1}{n} g(n) \\ &= f(n) \end{aligned}$$

Hence $Sg = f$ and thus S is surjective. Let $g \in Y$. Suppose that $Sg = 0$. Then

$$\sum_{n=1}^{\infty} \frac{1}{n} |g(n)| = \|Sg\| = 0$$

Thus for each $n \in \mathbb{N}$, $g(n) = 0$. Hence $\ker S = \{0\}$ and S is injective. Note that for each $A \subset Y$, $S(A) = T^{-1}(A)$. If S is open, then T is continuous which as shown above is a contradiction. So g is not open.

□

Exercise 9.6.0.8. Let $X = C^1([0, 1])$ and $Y = C([0, 1])$. Equip both X and Y with the uniform norm.

1. Then X is not complete [move this to introduction section before showing differentiation is not bounded and reference here](#)
2. Define $T : X \rightarrow Y$ by $Tf = f'$. Then $\Gamma(T)$ is closed and T is not bounded.

Proof.

1. Recall that for each $a, b \geq 0$ and $p \in \mathbb{N}$,

$$(a^{\frac{1}{p}} + b^{\frac{1}{p}})^p = \sum_{n=0}^p \binom{p}{n} a^{\frac{n}{p}} b^{\frac{p-n}{p}} \geq a + b$$

Thus $(a + b)^{\frac{1}{p}} \leq a^{\frac{1}{p}} + b^{\frac{1}{p}}$.

For each $n \in \mathbb{N}$, define $f_n : [0, 1] \rightarrow \mathbb{C}$ by $f_n(x) = \sqrt{(x - \frac{1}{2})^2 + \frac{1}{n^2}}$. Then $(f_n)_{n \in \mathbb{N}} \subset X$. Define $f : [0, 1] \rightarrow \mathbb{C}$ by $f(x) = |x - \frac{1}{2}|$. Then $f \in Y \cap X^c$. Note that for each $n \in \mathbb{N}$, $f \leq f_n$. Our observation above implies that for each $x \in X$,

$$\begin{aligned} f_n(x) &= \left[(x - \frac{1}{2})^2 + \frac{1}{n^2} \right]^{\frac{1}{2}} \\ &\leq |x - \frac{1}{2}| + \frac{1}{n} \end{aligned}$$

Thus $0 \leq f_n - f \leq \frac{1}{n}$. This implies that $f_n \xrightarrow{u} f$. Since $f \notin X$, X is not complete.

2. Let $(f_n)_{n \in \mathbb{N}} \subset X$, $f \in X$ and $g \in Y$. Suppose that $f_n \xrightarrow{u} f$ and $Tf_n \xrightarrow{u} g$. Let $x \in [0, 1]$. Then $f_n(x) \rightarrow f(x)$ and $f_n(0) \rightarrow f(0)$ and $f'_n \xrightarrow{u} g$. Applying the DCT to this sequence of integrable functions that converges uniformly to an integrable function on a finite measure space, [ref ex in analysis notes](#) implies that

$$\begin{aligned} f_n(x) - f_n(0) &= \int_{[0, x]} f'_n dm \\ &\rightarrow \int_{[0, x]} g dm \end{aligned}$$

Since $f_n(x) - f_n(0) \rightarrow f(x) - f(0)$, we know that

$$f(x) - f(0) = \int_{[0, x]} g dm$$

Thus $Tf = g$ and $\Gamma(T)$ is closed.

By Exercise 9.2.0.4, T is not bounded.

□

Exercise 9.6.0.9. Let X, Y be Banach spaces and $T \in L(X; Y)$. Then $X/\ker T \cong T(X)$ iff $T(X)$ is closed.

Proof. Since X is a Banach space and T is continuous, we have that $\ker T$ is closed and $X/\ker T$ is a Banach space. Suppose that $X/\ker T \cong T(X)$. Then $T(X)$ is complete. Since Y is complete, this implies that $T(X)$ is closed.

Conversely Suppose that $T(X)$ is closed. Then $T(X)$ is complete. Define $S : X/\ker T \rightarrow T(X)$ by $S(x + \ker T) = T(x)$. A previous exercise tells us that the map $S : X/\ker T \rightarrow T(X)$ defined by $S(x + \ker T) = T(x)$ is a bounded linear bijection. Since $T(X)$ is complete and S is surjective, S^{-1} is bounded and thus S is an isomorphism. □

Exercise 9.6.0.10. Let X be a separable Banach space. Define $B_X = \{x \in X : \|x\| < 1\}$. Let $(x_n)_{n \in \mathbb{N}} \subset B_X$ a dense subset of the unit ball and μ the counting measure on $(\mathbb{N}, \mathcal{P}(\mathbb{N}))$. Define $T : L^1(\mu) \rightarrow X$ by

$$Tf = \sum_{n=1}^{\infty} f(n)x_n$$

Then

1. T is well defined and $T \in L(L^1(\mu), X)$
2. T is surjective
3. There exists a closed subspace $K \subset L^1(\mu)$ such that $L^1(\mu)/K \cong X$

Proof. 1. Let $f \in L^1(\mu)$. Since X is complete and

$$\begin{aligned} \sum_{n=1}^{\infty} \|f(n)x_n\| &= \sum_{n=1}^{\infty} |f(n)| \|x_n\| \\ &\leq \sum_{n=1}^{\infty} |f(n)| \\ &< \infty \end{aligned}$$

we have that $\sum_{n=1}^{\infty} f(n)x_n$ converges and thus $Tf \in X$. Hence T is well defined.

Clearly T is linear. Let $f \in L^1(\mu)$. Then

$$\begin{aligned} \|Tf\| &= \left\| \sum_{n=1}^{\infty} f(n)x_n \right\| \\ &\leq \sum_{n=1}^{\infty} \|f(n)x_n\| \\ &\leq \sum_{n=1}^{\infty} |f(n)| \\ &= \|f\|_1 \end{aligned}$$

So T is bounded with $\|T\| \leq 1$.

2. Let $x \in X$. Suppose that $\|x\| < 1$. Then $x \in B_X$. So there exists $n_1 \in \mathbb{N}$ such that $\|x - x_{n_1}\| < \frac{1}{2}$. Then $2(x - x_{n_1}) \in B_X$. Since for each $j \in \mathbb{N}$, $B_X \setminus (x_n)_{n=1}^j$ is dense in B_X , there exists $n_2 \in \mathbb{N}$ such that $x_{n_2} \notin (x_n)_{n=1}^{n_1}$ and $\|2(x - x_{n_1}) - x_{n_2}\| < \frac{1}{2}$ which implies that $\|x - (x_{n_1} - \frac{1}{2}x_{n_2})\| < \frac{1}{4}$.

Proceed inductively to obtain a subsequence $(x_{n_k})_{k \in \mathbb{N}}$ such that for each $k \geq 2$, $x_{n_k} \notin (x_n)_{n=1}^{n_{k-1}}$ and $\|x - \sum_{j=1}^k 2^{1-j}x_{n_j}\| < \frac{1}{2^k}$.

Then $x = \sum_{k=1}^{\infty} 2^{1-k}x_{n_k}$.

Define $f : \mathbb{N} \rightarrow \mathbb{C}$ by $f = \sum_{k=1}^{\infty} 2^{1-k}\chi_{\{n_k\}}$. Then $\|f\|_1 = \sum_{k=1}^{\infty} 2^{1-k} < \infty$, so $f \in L^1(\mu)$ and $Tf = \sum_{k=1}^{\infty} 2^{1-k}x_{n_k} = x$. Now, suppose that $\|x\| \geq 1$, then $\frac{1}{2\|x\|}x \in B_X$. The above argument shows that there exists $f \in L^1(\mu)$ such that $Tf = \frac{1}{2\|x\|}x$. Then $2\|x\|f \in L^1(\mu)$ and $T(2\|x\|f) = 2\|x\|Tf = x$.

So for each $x \in X$, there exists $f \in L^1(\mu)$ such that $Tf = x$ and thus T is surjective.

3. Since X is a Banach space and T is surjective, the previous exercise implies that $L^1(\mu)/\ker T \cong X$.

□

9.7 Duality

Note 9.7.0.1. Let X be a normed vector space. Then X^* is a normed vector space. In general the weak-* topology on X^* is not necessarily the same as the norm topology on X^* . In the context of normed vector spaces, we will write X^{**} to denote $(X^*)^*$ when X^* is equipped with the norm topology and $\widehat{X}$ to denote $(X^*)^*$ when X^* is equipped with the weak-* topology. **write** $X^{**} = ((X, \mathcal{T}_{\|\cdot\|_X})^*, \mathcal{T}_{\|\cdot\|_{X^*}})^*$ and $\widehat{X} = ((X, \mathcal{T}_{\|\cdot\|_X})^*, \mathcal{T}_{w^*})^*$?

Exercise 9.7.0.2. Let X be a normed vector space and $x \in X$. Define $\hat{x} : X^* \rightarrow \mathbb{C}$ by $\hat{x}(f) = f(x)$. Then $\hat{x} \in X^{**}$ and $\|\hat{x}\| = \|x\|$.

Hint: Hahn-Banach theorem

Proof. Let $f, g \in X^*$ and $\lambda \in \mathbb{C}$. Then

$$\hat{x}(f + \lambda g) = (f + \lambda g)(x) = f(x) + \lambda g(x) = \hat{x}(f) + \lambda \hat{x}(g)$$

So $\hat{x}$ is linear. For each $f \in X^*$,

$$|\hat{x}(f)| = |f(x)| \leq \|x\| \|f\|$$

Hence $\hat{x} \in X^{**}$ with $\|\hat{x}\| \leq \|x\|$. If $x = 0$, then $\hat{x} = 0$ and $\|\hat{x}\| = \|x\|$. Suppose that $x \neq 0$. Then a previous exercise implies that there exists $F \in X^*$ such that $\|F\| = 1$ and $F(x) = \|x\|$. Then we have that

$$\sup_{\substack{f \in X^* \\ \|f\|=1}} |\hat{x}(f)| = \sup_{\substack{f \in X^* \\ \|f\|=1}} |f(x)| \geq |F(x)| = \|x\|$$

Hence $\|\hat{x}\| = \|x\|$. □

Exercise 9.7.0.3. Let X be a normed vector space. If X is separable, then there exist $(\phi_n)_{n \in \mathbb{N}} \subset X^*$ such that for each $n \in \mathbb{N}$, $\|\phi_n\| = 1$ and for each $x \in X$,

$$\|x\| = \sup_{n \in \mathbb{N}} |\phi_n(x)|$$

Hint: Let $(x_n)_{n \in \mathbb{N}} \subset X$ be a dense subset. A previous exercise on the Hahn-Banach theorem implies that for each n , there exists $\phi_n \in X^*$ such that $\|\phi_n\| = 1$ and $\phi_n(x_n) = \|x_n\|$. Then for each $x \in X$,

$$\|x\| = \sup_{n \in \mathbb{N}} |\phi_n(x)|$$

Proof. Suppose that X is separable. Then there exists $(x_n)_{n \in \mathbb{N}} \subset X$ such that $(x_n)_{n \in \mathbb{N}}$ is dense in X . A previous exercise on the Hahn-Banach theorem implies that for each n , there exists $\phi_n \in X^*$ such that $\|\phi_n\| = 1$ and $\phi_n(x_n) = \|x_n\|$. Let $x \in X$. Then

$$\begin{aligned} \|x\| &= \|\hat{x}\| \\ &= \sup_{\substack{\phi \in X^* \\ \|\phi\|=1}} \|\hat{x}(\phi)\| \\ &= \sup_{\substack{\phi \in X^* \\ \|\phi\|=1}} \|\phi(x)\| \\ &\geq \sup_{n \in \mathbb{N}} \|\phi_n(x)\| \end{aligned}$$

Let $\epsilon > 0$. Choose $N \in \mathbb{N}$ such that $\|x - x_N\| < \epsilon/2$. Then

$$\begin{aligned} \|x\| &\leq \|x - x_N\| + \|x_N\| \\ &= \|x - x_N\| + |\phi_N(x_N)| \\ &\leq \|x - x_N\| + |\phi_N(x_N - x)| + |\phi_N(x)| \\ &\leq \|x - x_N\| + \|\phi_N\| \|x_N - x\| + |\phi_N(x)| \\ &\leq 2\|x - x_N\| + |\phi_N(x)| \\ &< 2\frac{\epsilon}{2} + |\phi_N(x)| \\ &\leq \epsilon + \sup_{n \in \mathbb{N}} |\phi_n(x)| \end{aligned}$$

Since $\epsilon > 0$ is arbitrary, $\|x\| \leq \sup_{n \in \mathbb{N}} |\phi_n(x)|$. So $\|x\| = \sup_{n \in \mathbb{N}} |\phi_n(x)|$. □

Exercise 9.7.0.4. Let X be a normed vector space. Define $\phi : X \rightarrow X^{**}$ by $\phi(x) = \hat{x}$. Then ϕ is a linear isometry.

Proof. Let $x, y \in X$ and $\lambda \in \mathbb{C}$. Then for each $f \in X^*$, we have that

$$\begin{aligned} \phi(x + \lambda y)(f) &= \widehat{x + \lambda y}(f) \\ &= f(x + \lambda y) \\ &= f(x) + \lambda f(y) \\ &= \hat{x}(f) + \lambda \hat{y}(f) \\ &= \phi(x)(f) + \lambda \phi(y)(f) \end{aligned}$$

So $\phi(x + \lambda y) = \phi(x) + \lambda \phi(y)$ and ϕ is linear. The previous exercise tells us that

$$\begin{aligned} \|\phi(x) - \phi(y)\| &= \|\phi(x - y)\| \\ &= \|\widehat{x - y}\| = \|x - y\| \end{aligned}$$

So ϕ is an isometry. □

Definition 9.7.0.5. Let X be a normed vector space and define $\phi : X \rightarrow X^{**}$ as above. We define $\hat{X} = \phi(X) \subset X^{**}$. Since $\hat{X}$ and X are isomorphic, we may identify X as a subset of X^{**} .

Definition 9.7.0.6. Let X be a normed vector space and define $\phi : X \rightarrow X^{**}$ as above. Then X is said to be **reflexive** if ϕ is surjective. In this case ϕ is then an isomorphism

Definition 9.7.0.7. Let X, Y be normed vector spaces and $T \in L(X; Y)$. Define the **adjoint of T** , denoted $T^* : Y^* \rightarrow X^*$, by $T^*(f) = f \circ T$.

Exercise 9.7.0.8. Let X, Y be normed vector spaces and $T \in L(X; Y)$.

1. Then $T^* \in L(Y^*, X^*)$.
2. Applying the result from (1) twice, we have that $T^{**} \in L(X^{**}, Y^{**})$. We have that for each $x \in X$, $T^{**}(\hat{x}) = \widehat{T(x)}$.
3. T^* is injective iff $T(X)$ is dense in Y .
4. If $T^*(Y^*)$ is dense in X^* , then T is injective. The converse is true if X is reflexive.

Proof.

1. Let $f \in Y^*$. Then $\|T^*(f)\| = \|f \circ T\| \leq \|T\| \|f\|$. So $T^* \in L(Y^*, X^*)$ with $\|T^*\| \leq \|T\|$.

2. Let $x \in X$. Let $f \in Y^*$. Then

$$\begin{aligned} T^{**}(\hat{x})(f) &= \hat{x} \circ T^*(f) \\ &= \hat{x}(T^*(f)) \\ &= \hat{x}(f \circ T) \\ &= f \circ T(x) \\ &= f(T(x)) \\ &= \widehat{T(x)}(f) \end{aligned}$$

Hence $T^{**}(\hat{x}) = \widehat{T(x)}$.

3. Suppose that $T(X)$ is not dense in Y . Then $\text{cl } T(X) \neq Y$. So $T(X)$ is a proper closed subspace of Y and there exists $y \in Y$ such that $y \notin \text{cl } T(X)$. By a previous exercise, there exists $f \in Y^*$ such that $f(y) = \|y + \text{cl } T(X)\| \neq 0$, $\|f\| = 1$ and $f|_{\text{cl } T(X)} = 0$. Let $x \in X$. Then $T^*(f)(x) = f \circ T(x) = 0$. Hence $T^*(f) = 0 = T^*(0)$. Since $f \neq 0$, T^* is not injective. Now suppose that $T(X)$ is dense in Y . Let $f, g \in Y^*$. Define $h \in Y^*$ by $h = f - g$. Suppose that $T^*(f) = T^*(g)$. Then $T^*(h) = 0$. So for each $x \in X$, $h(T(x)) = 0$. Let $y \in Y$ and $\epsilon > 0$. By continuity, there exists $\delta > 0$ such that for each $y' \in Y$, if $\|y - y'\| < \delta$, then $\|h(y) - h(y')\| < \epsilon$. Since $T(X)$ is dense in Y , there exists $x \in X$ such that $\|y - T(x)\| < \delta$. Thus

$$\begin{aligned} \|h(y)\| &\leq \|h(y) - h(T(x))\| + \|h(T(x))\| \\ &= \|h(y) - h(T(x))\| \\ &< \epsilon \end{aligned}$$

Since $\epsilon > 0$ is arbitrary, $\|h(y)\| = 0$. This implies that $h(y) = 0$ and therefore $f(y) = g(y)$. Since $y \in Y$ is arbitrary, $f = g$ and T^* is injective.

4. For the sake of contradiction, suppose that $T^*(Y^*)$ is dense in X^* and T is not injective. Then there exist $x_1, x_2 \in X$ such that $x_1 \neq x_2$ and $T(x_1) = T(x_2)$. Define $x = x_1 - x_2$. Then $x \neq 0$ and $T(x) = 0$. A previous exercise implies that there exists $F \in X^*$ such that $F(x) = \|x\| \neq 0$ and $\|F\| = 1$. Let $\epsilon > 0$. Choose $g \in Y^*$ such that $\|F - T^*(g)\| < \epsilon$. Then

$$\begin{aligned} \|x\| &= |F(x)| \\ &\leq |F(x) - T^*(g)(x)| + |T^*(g)(x)| \\ &< \epsilon\|x\| + |g(T(x))| \\ &= \epsilon\|x\| \end{aligned}$$

Since $\epsilon > 0$ is arbitrary, we have that $\|x\| = 0$ which is a contradiction. Hence if $T^*(Y^*)$ is dense in X^* , then T is injective.

Now, suppose that X is reflexive and T is injective. Let $\phi_1, \phi_2 \in X^{**}$. Suppose that $T^{**}(\phi_1) = T^{**}(\phi_2)$. Then $T^{**}(\phi_1 - \phi_2) = 0$. Since X is reflexive, there exist $x_1, x_2 \in X$ such that $\phi_1 = \hat{x}_1$ and $\phi_2 = \hat{x}_2$. Define $x = x_1 - x_2$. Then $T^{**}(\hat{x}) = 0$. So for each $f \in Y^*$,

$$\begin{aligned} T^{**}(\hat{x})(f) &= \hat{x} \circ T^*(f) \\ &= \hat{x}(T^*(f)) \\ &= \hat{x}(f \circ T) \\ &= f \circ T(x) \\ &= f(T(x)) \\ &= 0 \end{aligned}$$

Suppose that $T(x) \neq 0$. Then a previous exercise implies that there exists $g \in Y^*$ such that $g(T(x)) = \|T(x)\| \neq 0$ and $\|g\| = 1$. This is a contradiction since $g(T(x)) = 0$. So $T(x) = 0$. Since T is injective, this implies that $x = 0$. Hence $\hat{x} = 0$ and thus $\phi_1 = \phi_2$. Thus T^{**} is injective. By (3), we have that $T^*(Y^*)$ is dense in X^* .

□

Exercise 9.7.0.9. Let X be a normed vector space. Then X is reflexive iff X^* is reflexive.

Proof. Suppose that X is reflexive. Let $\alpha \in X^{***}$. Define $f : X \rightarrow \mathbb{C}$ by $f(x) = \alpha(\hat{x})$. Clearly f is linear and a previous exercise tells us that for each $x \in X$,

$$\begin{aligned} |f(x)| &\leq \|\alpha\| \|\hat{x}\| \\ &= \|\alpha\| \|x\| \end{aligned}$$

So $f \in X^*$. Let $\phi \in X^{**}$. Since X is reflexive, there exists $x \in X$ such that $\phi = \hat{x}$. Then

$$\begin{aligned}\alpha(\phi) &= \alpha(\hat{x}) \\ &= f(x) \\ &= \hat{x}(f) \\ &= \hat{f}(\hat{x}) \\ &= \hat{f}(\phi)\end{aligned}$$

Hence $\alpha = \hat{f}$. Thus the map $X^* \rightarrow X^{***}$ given by $f \mapsto \hat{f}$ is surjective and so X^* is reflexive.

Conversely, suppose that X^* is reflexive. Since $\phi : X \rightarrow X^{**}$ given by $\phi(x) = \hat{x}$ is an isometry, $\widehat{X} \subset X^{**}$ is closed. For the sake of contradiction, suppose that $\widehat{X} \neq X^{**}$. Then there exists $\alpha \in X^{**}$ such that $\alpha \notin \widehat{X}$. Thus there exists $F \in X^{***}$ such that $\|F\| = 1$, $F(\alpha) = \|\alpha + \widehat{X}\| \neq 0$ and $F|_{\widehat{X}} = 0$. Since X^* is reflexive, there exists $f \in X^*$ such that $F = \hat{f}$. A previous exercise tells us that $\|f\| = \|\hat{f}\| = \|F\| = 1$. Since for each $x \in X$, $f(x) = \hat{x}(f) = \hat{f}(\hat{x}) = F(\hat{x}) = 0$, we have that $f = 0$. Thus $\|f\| = 0$, a contradiction. So $\widehat{X} = X^{**}$ and X is reflexive. $\square$

Exercise 9.7.0.10. Let X be a normed vector space. Then

1. $\mathcal{T}_{w^*} \subset \mathcal{T}_{X^*}$
2. For each $E \subset X^*$, if E is weak-* closed, then E is norm closed

Proof.

1. Since $\widehat{X} \subset X^{**}$, we have that

$$\begin{aligned}\mathcal{T}_{w^*} &= \tau_{X^*}(\widehat{X}) \\ &\subset \tau_{X^*}(X^{**}) \\ &= \mathcal{T}_{X^*} \quad (\text{by ref ex that } \mathcal{T}_{X^*} = \tau_{X^*}(X^{**}))\end{aligned}$$

2. Let $E \subset X^*$. Suppose that E is weak-* closed. Then

$$\begin{aligned}E^c &\in \mathcal{T}_{w^*} \\ &\subset \mathcal{T}_{X^*}\end{aligned}$$

So E is norm closed. $\square$

Exercise 9.7.0.11. Let X be a Banach space.

1. Let $(x_n)_{n \in \mathbb{N}} \subset X$ and $x \in X$. If $x_n \xrightarrow{w} x$, then (x_n) is $d_{\|\cdot\|_X}$ -bounded.
2. Let $(\phi_n)_{n \in \mathbb{N}} \subset X^*$ and $\phi \in X^*$. If $\phi_n \xrightarrow{w^*} \phi$, then (ϕ_n) is $d_{\|\cdot\|_{X^*}}$ -bounded.

Proof.

1. **FINISH!!!**
2. We note that

$$\begin{aligned}\phi_n \xrightarrow{w^*} \phi &\iff \text{for each } x \in X, \phi_n(x) \rightarrow \phi(x) \\ &\implies \text{for each } x \in X, (\phi_n(x))_{n \in \mathbb{N}} \subset \mathbb{R} \text{ is } d_{|\cdot|}\text{-bounded} \\ &\iff \text{for each } x \in X, \sup_{n \in \mathbb{N}} |\phi_n(x)| < \infty. \\ &\implies \sup_{n \in \mathbb{N}} \|\phi_n\|_{X^*} < \infty \quad (\text{by ref ex, unif boundedness principle}) \\ &\implies (\phi_n)_{n \in \mathbb{N}} \subset X^* \text{ is } d_{\|\cdot\|_{X^*}}\text{-bounded.}\end{aligned}$$

□

Exercise 9.7.0.12. Let X be a normed vector space and $E \subset X$. Suppose that E is $\mathcal{T}_{\|\cdot\|}$ -dense in X . Let $(\phi_n)_{n \in \mathbb{N}} \subset X^*$ and $\phi \in X^*$. Suppose that $(\phi_n)_{n \in \mathbb{N}}$ is $d_{\|\cdot\|, X^*}$ -bounded. Then $\phi_n \xrightarrow{w^*} \phi$ iff for each $x \in E$, $\phi_n(x) \rightarrow \phi(x)$.

Proof.

• ($\Rightarrow$):

Suppose that $\phi_n \xrightarrow{w^*} \phi$. Then for each $x \in X$, $\phi_n(x) \rightarrow \phi(x)$. Since $E \subset X$, we have that for each $x \in E$, $\phi_n(x) \rightarrow \phi(x)$.

• ($\Leftarrow$):

Suppose that for each $x \in E$, $\phi_n(x) \rightarrow \phi(x)$. Since $(\phi_n)_{n \in \mathbb{N}}$ is $d_{\|\cdot\|, X^*}$ -bounded, $\sup_{n \in \mathbb{N}} \|\phi_n\|_{X^*} < \infty$. Define $M \in [0, \infty)$ by

$M := \sup_{n \in \mathbb{N}} \|\phi_n\|_{X^*}$. Let $\epsilon > 0$. Since E is $\mathcal{T}_{\|\cdot\|}$ -dense in X , there exists $x_0 \in E$ such that $\|x_0 - x\| \leq \epsilon 3^{-1} (M + 1)^{-1} (\|\phi\| + 1)^{-1}$.

Since $x_0 \in E$, by assumption, we have that $\phi_n(x_0) \rightarrow \phi(x_0)$. Therefore there exists $N \in \mathbb{N}$ such that for each $n \in \mathbb{N}$, $n \geq N$ implies that $|\phi_n(x_0) - \phi(x_0)| < \epsilon/3$. Then for each $n \in \mathbb{N}$, $n \geq N$ implies that

$$\begin{aligned} |\phi_n(x) - \phi(x)| &\leq |\phi_n(x) - \phi_n(x_0)| + |\phi_n(x_0) - \phi(x_0)| + |\phi(x_0) - \phi(x)| \\ &\leq \|\phi_n\| \|x - x_0\| + |\phi_n(x_0) - \phi(x_0)| + \|\phi\| \|x - x_0\| \\ &< M \frac{\epsilon}{3(M+1)} + \frac{\epsilon}{3} + \|\phi\| \frac{\epsilon}{3(\|\phi\| + 1)} \\ &< \frac{\epsilon}{3} + \frac{\epsilon}{3} + \frac{\epsilon}{3} \\ &= \epsilon. \end{aligned}$$

Since $\epsilon > 0$ is arbitrary, we have that for each $\epsilon > 0$, there exists $N \in \mathbb{N}$ such that for each $n \in \mathbb{N}$, $n \geq N$ implies that $|\phi_n(x) - \phi(x)| < \epsilon$. Hence $\phi_n(x) \rightarrow \phi(x)$. Since $x \in X$ is arbitrary, we have that for each $x \in X$, $\phi_n(x) \rightarrow \phi(x)$. Hence $\phi_n \xrightarrow{w^*} \phi$.

□

Definition 9.7.0.13. Let X be a normed vector space, $M \subset X$ and $N \subset X^*$. We define the **annihilator** of M and the annihilator of N , denoted by $M^\perp \subset X^*$ and ${}^\perp N \subset X$ respectively, by

$$\begin{aligned} M^\perp &= \{\phi \in X^* : \text{for each } x \in M, \phi(x) = 0\} \\ {}^\perp N &= \{x \in X : \text{for each } \phi \in N, \phi(x) = 0\} \end{aligned}$$

Exercise 9.7.0.14. Let X be a normed vector space, $M \subset X$ and $N \subset X^*$. Then

1.

$$M^\perp = \bigcap_{x \in M} \ker \hat{x}$$

2.

$${}^\perp N = \bigcap_{\phi \in N} \ker \phi$$

Proof.

1.

$$\begin{aligned} M^\perp &= \{\phi \in X^* : \text{for each } x \in M, \phi(x) = 0\} \\ &= \bigcap_{x \in M} \{\phi \in X^* : \phi(x) = 0\} \\ &= \bigcap_{x \in M} \{\phi \in X^* : \hat{x}(\phi) = 0\} \\ &= \bigcap_{x \in M} \ker \hat{x} \end{aligned}$$

2.

$$\begin{aligned}
{}^\perp N &= \{x \in X : \text{for each } \phi \in N, \phi(x) = 0\} \\
&= \bigcap_{\phi \in N} \{x \in X : \phi(x) = 0\} \\
&= \bigcap_{\phi \in N} \ker \phi
\end{aligned}$$

□

Exercise 9.7.0.15. Let X be a normed vector space, $M \subset X$ and $N \subset X^*$. Then

1. $M^\perp$ is weak-* closed
2. ${}^\perp N$ is closed

Proof.

1. Let $(\phi_n)_{n \in \mathbb{N}} \subset M^\perp$ and $\phi \in X^*$. Suppose that $\phi_n \xrightarrow{w^*} \phi$. Then for each $x \in X$, $\phi_n(x) \rightarrow \phi(x)$. Let $x \in M$. By definition, for each $n \in \mathbb{N}$, $\phi_n(x) = 0$. Thus $\phi_n(x) \rightarrow 0$ which implies that $\phi(x) = 0$ and $\phi \in \ker \hat{x}$. Since $x \in M$ is arbitrary,

$$\begin{aligned}
\phi &\in \bigcap_{x \in M} \ker \hat{x} \\
&= M^\perp
\end{aligned}$$

2. Let $(x_n)_{n \in \mathbb{N}} \subset {}^\perp N$ and $x \in X$. Suppose that $x_n \rightarrow x$. Let $\phi \in N$. Continuity implies that $\phi(x_n) \rightarrow \phi(x)$. By definition, for each $n \in \mathbb{N}$, $\phi(x_n) = 0$. Thus $\phi(x_n) \rightarrow 0$ which implies that $\phi(x) = 0$. So $x \in \ker \phi$. Since $\phi \in N$ is arbitrary,

$$\begin{aligned}
x &\in \bigcap_{\phi \in N} \ker \phi \\
&= {}^\perp N
\end{aligned}$$

□

Exercise 9.7.0.16. Let X be a normed vector space, $M \subset X$ and $N \subset X^*$. Then

1. ${}^\perp(M^\perp) = \text{cl } M$, i.e. the norm closure of M
2. $({}^\perp N)^\perp = \text{cl}_{w^*}(N)$, i.e. the weak-* closure of N .

Proof.

1. Let $x \in M$, then by definition, for each $\phi \in M^\perp$, $\phi(x) = 0$. Again by definition, $x \in {}^\perp(M^\perp)$. So $M \subset {}^\perp(M^\perp)$. Since ${}^\perp(M^\perp)$ is closed, $\text{cl } M \subset {}^\perp(M^\perp)$. For the sake of contradiction, suppose that ${}^\perp(M^\perp) \not\subset \text{cl } M$. Then there exists $x \in {}^\perp(M^\perp)$ such that $x \notin \text{cl } M$. Exercise 9.5.0.9 implies that there exists $\phi \in X^*$ such that $\phi|_{\text{cl } M} = 0$, $\|\phi\| = 1$ and $\phi(x) = \|x + \text{cl } M\| > 0$. By definition, $\phi \in M^\perp$. Since $\phi(x) \neq 0$, we have that $x \notin {}^\perp(M^\perp)$. This is a contradiction and so ${}^\perp(M^\perp) \subset \text{cl } M$.

2.

□

Exercise 9.7.0.17. Banach-Alaoglu Theorem:

Let X be a normed vector space. Then $\bar{B}(0, 1)$ is w^* -compact.

Proof. For $x \in X$, define $D_x \subset \mathbb{C}$ by $D_x := \bar{B}_{\mathbb{C}}(0, \|x\|)$. Then for each $x \in X$, D_x is compact. Define $D \subset \mathbb{C}^X$ by $D := \prod_{x \in X} D_x$. Tychonoff's theorem Exercise 4.14.3.1 implies that D is compact. Let $\phi \in \bar{B}_{X^*}(0, 1)$. Then for each $x \in X$, $|\phi(x)| \leq \|x\|$. Hence $\phi \in D$. Since $\phi \in \bar{B}_{X^*}(0, 1)$ is arbitrary, we have that $\bar{B}_{X^*}(0, 1) \subset D$. Let $(\phi_\alpha)_{\alpha \in A} \subset \bar{B}_{X^*}(0, 1)$ and $\phi \in D$. Suppose that $\phi_\alpha \xrightarrow{\text{p.w.}} \phi$. Then for each $x, y \in X$ and $\lambda \in \mathbb{C}$,

$$\begin{aligned} \phi(x + \lambda y) &= \lim_{\alpha \in A} \phi_\alpha(x + \lambda y) \\ &= \lim_{\alpha \in A} [\phi_\alpha(x) + \lambda \phi_\alpha(y)] \\ &= \lim_{\alpha \in A} \phi_\alpha(x) + \lambda \lim_{\alpha \in A} \phi_\alpha(y) \\ &= \phi(x) + \lambda \phi(y). \end{aligned}$$

So ϕ is linear. Let $x \in X$. Suppose that $\|x\| = 1$. Since $|\cdot| : \mathbb{C} \rightarrow \mathbb{C}$ is continuous, we have that $|\phi_\alpha(x)| \rightarrow |\phi(x)|$. Since $\|x\| = 1$, we have that

$$\begin{aligned} |\phi_\alpha(x)| &\leq \|\phi_\alpha\| \|x\| \\ &\leq \|x\| \\ &= 1. \end{aligned}$$

Since $x \in X$ with $\|x\| = 1$ is arbitrary, we have that

$$\begin{aligned} \|\phi\| &= \sup_{\|x\|=1} |\phi(x)| \\ &\leq 1. \end{aligned}$$

Thus $\phi \in \bar{B}_{X^*}(0, 1)$. Since $(\phi_\alpha)_{\alpha \in A} \subset \bar{B}_{X^*}(0, 1)$ and $\phi \in D$ with $\phi_\alpha \xrightarrow{\text{p.w.}} \phi$ are arbitrary, we have that for each $(\phi_\alpha)_{\alpha \in A} \subset \bar{B}_{X^*}(0, 1)$ and $\phi \in D$, $\phi_\alpha \xrightarrow{\text{p.w.}} \phi$ implies that $\phi \in \bar{B}_{X^*}(0, 1)$. Hence $\bar{B}_{X^*}(0, 1)$ is closed in D . Since D is compact, $\bar{B}_{X^*}(0, 1)$ is compact. Since for each $(\phi_\alpha)_{\alpha \in A} \subset \bar{B}_{X^*}(0, 1)$ and $\phi \in \bar{B}_{X^*}(0, 1)$, $\phi_\alpha \xrightarrow{w^*} \phi$ iff $\phi_\alpha \xrightarrow{\text{p.w.}} \phi$, we have that $\bar{B}_{X^*}(0, 1)$ in weak-* is compact. $\square$

Exercise 9.7.0.18. Let $(X, \|\cdot\|_X)$ be a normed vector space. Define $B^* \subset X^*$ and $\mathcal{T}_{w^*, B^*} \subset \mathcal{P}(B^*)$ by $B^* := \bar{B}_{\|\cdot\|_{X^*}}(0, 1)$ and $\mathcal{T}_{w^*, B^*} := \mathcal{T}_{w^*} \cap B^*$. Then $(X, \mathcal{T}_{\|\cdot\|_X})$ is separable iff $(B^*, \mathcal{T}_{w^*, B^*})$ is a Polish space.

Proof. Define $\mathcal{T}_{\mathcal{P}, B^*} \subset \mathcal{P}(B^*)$ by $\mathcal{T}_{\mathcal{P}, B^*} := \mathcal{T}_{\mathcal{P}} \cap B^*$.

• ($\implies$):

Suppose that X is separable. Then there exists $E \subset X$ such that E is countable and $\mathcal{T}_{\|\cdot\|}$ -dense.

1. Define $\mathcal{P} \subset \text{SemiNorm}(X^*, \mathcal{T}_{w^*})$ by $\mathcal{P} := \{|\hat{x}| : x \in E\}$. We note that $\mathcal{P}$ is countable. Let $(\phi_\alpha)_{\alpha \in A} \subset B^*$ a net and $\phi \in B^*$.

- Suppose that $\phi_\alpha \xrightarrow{\mathcal{T}_{w^*, B^*}} \phi$. **ref ex in subspace topology section** implies that $\phi_\alpha \xrightarrow{w^*} \phi$. Then for each $x \in E$, $|\hat{x}|(\phi_\alpha - \phi) \rightarrow 0$. Thus $\phi_\alpha \xrightarrow{\mathcal{T}_{\mathcal{P}}} \phi$. **ref subspace top section again** implies that $\phi_\alpha \xrightarrow{\mathcal{T}_{\mathcal{P}} \cap B^*} \phi$. **ref ex in subspace topology section** again implies that $\phi_\alpha \xrightarrow{\mathcal{T}_{w^*, B^*}} \phi$.
- Conversely, suppose that $\phi_\alpha \xrightarrow{\mathcal{T}_{\mathcal{P}, B^*}} \phi$. **subspace top ex** again implies that $\phi_\alpha \xrightarrow{\mathcal{T}_{\mathcal{P}}} \phi$. Let $x \in X$ and $\epsilon > 0$. Then there exists $x_0 \in E$ such that $\|x - x_0\| < \min\{3^{-1}(\|\phi\| + 1)^{-1}\epsilon, 3^{-1}\epsilon\}$. Since $\phi_\alpha \xrightarrow{\mathcal{T}_{\mathcal{P}}} \phi$, we have that $|\hat{x}_0|(\phi_\alpha - \phi) \rightarrow 0$. Thus there exist $\alpha_0 \in A$ such that for each $\alpha \in A$, $\alpha \geq \alpha_0$ implies that $|\hat{x}_0|(\phi_\alpha - \phi) < \epsilon/3$. Then for each $\alpha \in A$,

$\alpha \geq \alpha_0$ implies that

$$\begin{aligned}
 |\hat{x}|(\phi_\alpha - \phi) &= |\hat{x}(\phi_\alpha - \phi)| \\
 &= |\hat{x}(\phi_\alpha) - \hat{x}(\phi)| \\
 &= |\phi_\alpha(x) - \phi(x)| \\
 &\leq |\phi_\alpha(x) - \phi_\alpha(x_0)| + |\phi_\alpha(x_0) - \phi(x_0)| + |\phi(x_0) - \phi(x)| \\
 &= |\phi_\alpha(x - x_0)| + |(\phi_\alpha - \phi)(x_0)| + |\phi(x_0 - x)| \\
 &\leq \|\phi\| \|x - x_0\| + |\hat{x}_0(\phi_\alpha - \phi)| + \|\phi\| \|x_0 - x\| \\
 &\leq \|x - x_0\| + |\hat{x}_0|(\phi_\alpha - \phi) + \|\phi\| \|x_0 - x\| \\
 &< \frac{\epsilon}{3} + \frac{\epsilon}{3} + \frac{\|\phi\|\epsilon}{3(\|\phi\| + 1)} \\
 &< \frac{\epsilon}{3} + \frac{\epsilon}{3} + \frac{\epsilon}{3} \\
 &= \epsilon.
 \end{aligned}$$

Since $\epsilon > 0$ is arbitrary, we have that for each $\epsilon > 0$, there exists $\alpha_0 \in A$ such that for each $\alpha \in A$, $\alpha \geq \alpha_0$ implies that $|\hat{x}|(\phi_\alpha - \phi) < \epsilon$. Thus $|\hat{x}|(\phi_\alpha - \phi) \rightarrow 0$. Since $x \in X$ is arbitrary, we have that for each $x \in X$, $|\hat{x}|(\phi_\alpha - \phi) \rightarrow 0$.

Then $\phi_\alpha \xrightarrow{w^*} \phi$. **ref subspace top section again** implies that $\phi_\alpha \xrightarrow{\mathcal{T}_{w^*, B^*}} \phi$.

Since $\phi_\alpha \xrightarrow{\mathcal{T}_{w^*, B^*}} \phi$ iff $\phi_\alpha \xrightarrow{\mathcal{T}_{\mathcal{P}, B^*}} \phi$, we have that $\mathcal{T}_{w^*, B^*} = \mathcal{T}_{\mathcal{P}, B^*}$. Define $d : B^* \times B^* \rightarrow [0, \infty)$ by $d := d_{\mathcal{P}}|_{B^*}$. **ref ex in lcs section** implies that **(define $d_{\mathcal{P}}$)** $d_{\mathcal{P}}|_{B^*} : B^* \times B^* \rightarrow [0, \infty)$ is a metric on B^* and **ref possibly other ex** implies that

$$\begin{aligned}
 \mathcal{T}_d &= \mathcal{T}_{d_{\mathcal{P}}|_{B^*}} \\
 &= \mathcal{T}_{\mathcal{P}, B^*} \\
 &= \mathcal{T}_{w^*, B^*}.
 \end{aligned}$$

2. **Banach-Alaoglu theorem** implies that $(B^*, \mathcal{T}_{w^*, B^*})$ is compact. **ref ex in compact metric space section** implies that (B^*, d) is complete.
3. **ref/make ex in metric space section about compact metric spaces being separable** implies that $(B^*, \mathcal{T}_{w^*, B^*})$ is separable.

Thus $(B^*, \mathcal{T}_{w^*, B^*})$ is a Polish space.

• ($\Leftarrow$):

Suppose that $(B^*, \mathcal{T}_{w^*, B^*})$ is a Polish space. **FINISH!!!**

□

9.8 Compact Operators

Definition 9.8.0.1.

9.9 Multilinear Maps

Definition 9.9.0.1. Let $X_1, \dots, X_n, Y$ be normed vector spaces and $T : \prod_{j=1}^n X_j \rightarrow Y$. Suppose that T is multilinear. Then T is said to be **bounded** if there exists $C \geq 0$ such that for each $x_1, \dots, x_n \in X$,

$$\|T(x_1, \dots, x_n)\| \leq C\|x_1\| \cdots \|x_n\|$$

We define

$$L^n(X_1, \dots, X_n; Y) = \left\{ T : \prod_{j=1}^n X_j \rightarrow Y : T \text{ is multilinear and bounded} \right\}$$

If $X_1 = \dots = X_n = X$, we write $L^n(X; Y)$ in place of $L^n(X_1, \dots, X_n; Y)$.

Note 9.9.0.2. For the remainder of this section we will primarily consider $L^2(X_1, X_2; Y)$ to avoid notational clutter, but all results immediately generalize to $L^n(X_1, \dots, X_n; Y)$

Exercise 9.9.0.3. Let X_1, X_2 and Y be normed vector spaces and $T : X_1 \times X_2 \rightarrow Y$ bilinear. Then the following are equivalent:

1. T is continuous
2. T is continuous at $(0, 0)$
3. T is bounded

Proof.

- (1) $\implies$ (2):
Trivial

- (2) $\implies$ (3):
Suppose that T is continuous at $(0, 0)$. For the sake of contradiction, suppose that T is not bounded. Then for each $C \geq 0$, there exist $(x_1, x_2) \in X_1 \times X_2$ such that $\|T(x_1, x_2)\| > C\|x_1\|\|x_2\|$. Hence there exist $(a_n)_{n \in \mathbb{N}} \subset X_1$ and $(b_n)_{n \in \mathbb{N}} \subset X_2$ such that for each $n \in \mathbb{N}$, $\|T(a_n, b_n)\| > n^2\|a_n\|\|b_n\|$. Hence for each $n \in \mathbb{N}$, $\|a_n\|, \|b_n\| > 0$. Define

$$(a'_n)_{n \in \mathbb{N}} \subset X_1$$

and $(b'_n)_{n \in \mathbb{N}} \subset X_2$ by $a'_n = \frac{a_n}{n\|a_n\|}$ and $b'_n = \frac{b_n}{n\|b_n\|}$. Then $(a'_n, b'_n) \rightarrow (0, 0)$. Continuity implies that $T(a'_n, b'_n) \rightarrow 0$. By construction, for each $n \in \mathbb{N}$,

$$\begin{aligned} \|T(a'_n, b'_n)\| &= \frac{1}{n^2\|a_n\|\|b_n\|} \|T(a_n, b_n)\| \\ &> \frac{n^2\|a_n\|\|b_n\|}{n^2\|a_n\|\|b_n\|} \\ &= 1 \end{aligned}$$

which is a contradiction. So T is bounded.

- (3) $\implies$ (1):
Suppose that T is bounded. Then there exists $C > 0$ such that for each $(x_1, x_2) \in X_1 \times X_2$, $\|T(x_1, x_2)\| \leq C\|x_1\|\|x_2\|$. Let $(a, b) \in X_1 \times X_2$ and $(a_n, b_n)_{n \in \mathbb{N}} \subset X_1 \times X_2$. Suppose that $(a_n, b_n) \rightarrow (a, b)$. Then $a_n \rightarrow a$, $b_n \rightarrow b$ and $(a_n)_{n \in \mathbb{N}}, (b_n)_{n \in \mathbb{N}}$ are bounded. So there exists $B \geq 0$ such that for each $n \in \mathbb{N}$ $\|b_n\| \leq B$. Hence

$$\begin{aligned} \|T(a_n, b_n) - T(a, b)\| &= \|T(a_n, b_n) - T(a, b_n) + T(a, b_n) - T(a, b)\| \\ &\leq \|T(a_n, b_n) - T(a, b_n)\| + \|T(a, b_n) - T(a, b)\| \\ &= \|T(a_n - a, b_n)\| + \|T(a, b_n - b)\| \\ &\leq C(\|a_n - a\|\|b_n\| + \|a\|\|b_n - b\|) \\ &\leq C(\|a_n - a\|B + \|a\|\|b_n - b\|) \\ &\rightarrow 0 \end{aligned}$$

Thus T is continuous.

□

Definition 9.9.0.4. Let X_1, X_2 and Y be normed vector spaces and $T \in L^2(X_1, X_2; Y)$. We define the **operator norm** on $L^2(X_1, X_2; Y)$, denoted $\|\cdot\| : L^2(X_1, X_2; Y) \rightarrow [0, \infty)$, by

$$\|T\| = \inf\{C \geq 0 : \text{for each } (x_1, x_2) \in X_1 \times X_2, \|T(x_1, x_2)\| \leq C\|x_1\|\|x_2\|\}$$

Exercise 9.9.0.5. Let X_1, X_2 and Y be normed vector spaces. If $X_1 \neq \{0\}$ and $X_2 \neq \{0\}$, then the operator norm on $L^2(X_1, X_2; Y)$ is given by:

1. $\|T\| = \sup_{\|x_1\|=1, \|x_2\|=1} \|T(x_1, x_2)\|$
2. $\|T\| = \sup_{x_1 \neq 0, x_2 \neq 0} \|x_1\|^{-1}\|x_2\|^{-1}\|T(x_1, x_2)\|$
3. $\|T\| = \inf\{C \geq 0 : \text{for each } (x_1, x_2) \in X_1 \times X_2, \|T(x_1, x_2)\| \leq C\|x_1\|\|x_2\|\}$

Proof. Since $X_1 \neq \{0\}$ and $X_2 \neq \{0\}$, the supremums in (1) and (2) are well defined. Let $T \in L^2(X_1, X_2; Y)$. Bilinearity of T implies that the sets over which the supremums are taken in (1) and (2) are the same. So (1) and (2) are equal.

Now, set

$$M = \sup_{\|x_1\|=1, \|x_2\|=1} \|T(x_1, x_2)\|$$

and

$$m = \inf\{C \geq 0 : \text{for each } (x_1, x_2) \in X_1 \times X_2, \|T(x_1, x_2)\| \leq C\|x_1\|\|x_2\|\}$$

Let $(x_1, x_2) \in X_1 \times X_2$. If $\|x_1\| = 0$ or $\|x_2\| = 0$, then $T(x_1, x_2) = 0$ and $\|T(x_1, x_2)\| \leq M\|x_1\|\|x_2\|$. Suppose that $\|x_1\| \neq 0$ and $\|x_2\| \neq 0$. Then

$$\begin{aligned} \|T(x_1, x_2)\| &= \left(\|T(\|x_1\|^{-1}x_1, \|x_2\|^{-1}x_2)\| \right) \|x_1\|\|x_2\| \\ &\leq M\|x_1\|\|x_2\| \end{aligned}$$

Hence $M \in \{C \geq 0 : \text{for each } x \in X, \|Tx\| \leq C\|x\|\}$ and $m \leq M$. Let $C \in \{C \geq 0 : \text{for each } (x_1, x_2) \in X_1 \times X_2, \|T(x_1, x_2)\| \leq C\|x_1\|\|x_2\|\}$. Suppose that $\|x_1\| = 1$ and $\|x_2\| = 1$. Then $\|T(x_1, x_2)\| \leq C\|x_1\|\|x_2\| = C$. So $M \leq C$. Therefore $M \leq m$. So $M = m$ and the supremum in (1) is the same as the infimum in (3). □

Exercise 9.9.0.6. Let X_1, X_2 and Y be normed vector spaces and $T \in L(X_1, X_2; Y)$. Then for each $(x_1, x_2) \in X_1 \times X_2$, $\|T(x_1, x_2)\| \leq \|T\|\|x_1\|\|x_2\|$.

Proof. Let $(x_1, x_2) \in X_1 \times X_2$. If $x_1 = 0$ or $x_2 = 0$, then

$$\begin{aligned} \|T(x_1, x_2)\| &= \|0\| \\ &= 0 \\ &= \|T\|\|x_1\|\|x_2\| \end{aligned}$$

Suppose that $x_1 \neq 0$ and $x_2 \neq 0$. The previous exercise implies that

$$\begin{aligned} \|T(x_1, x_2)\| &= \|T(\|x_1\|^{-1}\|x_1\|, \|x_2\|^{-1}\|x_2\|)\| \|x_1\|\|x_2\| \\ &\leq \left(\sup_{\|x_1\|=1, \|x_2\|=1} \|T(x_1, x_2)\| \right) \|x_1\|\|x_2\| \\ &= \|T\|\|x_1\|\|x_2\| \end{aligned}$$

□

Exercise 9.9.0.7. Let X_1, X_2 and Y be normed vector spaces. Then $L(X_1, X_2; Y)$ is a vector space and $\|\cdot\| : L^2(X_1, X_2; Y) \rightarrow [0, \infty)$ is a norm.

Proof. Let $S, T \in L(X_1, X_2; Y)$ and $\lambda \in \mathbb{C}$.

- It is clear that $S + T : X_1 \times X_2 \rightarrow Y$ is multilinear. For each $(x_1, x_2) \in X_1 \times X_2$,

$$\begin{aligned} \|(S + T)(x_1, x_2)\| &= \|S(x_1, x_2) + T(x_1, x_2)\| \\ &\leq \|S(x_1, x_2)\| + \|T(x_1, x_2)\| \\ &\leq \|S\|\|x_1\|\|x_2\| + \|T\|\|x_1\|\|x_2\| \\ &= (\|S\| + \|T\|)\|x_1\|\|x_2\| \end{aligned}$$

So $S + T \in L(X_1, X_2; Y)$ and $\|S + T\| \leq \|S\| + \|T\|$.

- It is clear that $\lambda T : X_1 \times X_2 \rightarrow Y$ is multilinear. For each $(x_1, x_2) \in X_1 \times X_2$,

$$\begin{aligned} \|(\lambda T)(x_1, x_2)\| &= \|\lambda T(x_1, x_2)\| \\ &= |\lambda| \|T(x_1, x_2)\| \\ &\leq |\lambda| \|T\| \|x_1\| \|x_2\| \end{aligned}$$

So $\lambda T \in L(X_1, X_2; Y)$ and $\|\lambda T\| \leq |\lambda| \|T\|$.

- Suppose that $\|T\| = 0$. Let $(x_1, x_2) \in X_1 \times X_2$. If $x_1 = 0$ or $x_2 = 0$, then $T(x_1, x_2) = 0$. Suppose that $x_1 \neq 0$ and $x_2 \neq 0$. Then

$$\begin{aligned} \|T(x_1, x_2)\| &\leq \|T\| \|x_1\| \|x_2\| \\ &= 0 \end{aligned}$$

So $T(x_1, x_2) = 0$. Since $(x_1, x_2) \in X_1 \times X_2$ is arbitrary, $T = 0$.

Therefore $L(X_1, X_2; Y)$ is a vector space and $\|\cdot\| : L(X_1, X_2; Y) \rightarrow [0, \infty)$ is a norm. □

Note 9.9.0.8. Let X, Y, Z be sets. We recall the definition of $\text{cur} : Y^{X \times Y} \rightarrow (Z^Y)^X$ from Definition 1.5.0.16.

Exercise 9.9.0.9. Let X_1, X_2, Y be normed vector spaces. Then

1. $\text{cur} : (X_1 \times X_2)^Y \rightarrow (Y^{X_2})^{X_1}$ is linear
2. $\text{cur}|_{L(X_1, X_2; Y)} : L(X_1, X_2; Y) \rightarrow L(X_1; L(X_2; Y))$
3. $\text{cur}|_{L(X_1, X_2; Y)}$ is an isometry
4. $\text{cur}|_{L(X_1, X_2; Y)}$ is surjective
5. $\text{cur}|_{L(X_1, X_2; Y)} : L(X_1, X_2; Y) \rightarrow L(X_1; L(X_2; Y))$ is an isometric isomorphism.

Proof.

1. Let $S, T \in (X_1 \times X_2)^Y$, $\lambda \in \mathbb{C}$ and $(x_1, x_2) \in X_1 \times X_2$. Then

$$\begin{aligned} \text{cur}(S + \lambda T)(x_1)(x_2) &= (S + \lambda T)(x_1, x_2) \\ &= S(x_1, x_2) + \lambda T(x_1, x_2) \\ &= \text{cur}(S)(x_1)(x_2) + \lambda \text{cur}(T)(x_1)(x_2) \\ &= \text{cur}(S)(x_1)(x_2) + \lambda \text{cur}(T)(x_1)(x_2) \\ &= [\text{cur}(S) + \lambda \text{cur}(T)](x_1)(x_2) \end{aligned}$$

Since $(x_1, x_2) \in X_1 \times X_2$ is arbitrary, $\text{cur}(S + \lambda T) = \text{cur}(S) + \lambda \text{cur}(T)$. Since $S, T \in (X_1 \times X_2)^Y$ and $\lambda \in \mathbb{C}$ are arbitrary, $\text{cur} : (X_1 \times X_2)^Y \rightarrow (Y^{X_2})^{X_1}$ is linear.

2. Let $T \in L(X_1, X_2; Y)$ and $x_1 \in X_1$. Since T is bilinear, for each $u, v \in X_2$ and $\lambda \in \mathbb{C}$,

$$\begin{aligned}\operatorname{cur}(T)(x_1)(u + \lambda v) &= T(x_1, u + \lambda v) \\ &= T(x_1, u) + \lambda T(x_1, v) \\ &= \operatorname{cur}(T)(x_1)(u) + \lambda \operatorname{cur}(T)(x_1)(v)\end{aligned}$$

So for each $x_1 \in X_1$, $\operatorname{cur}(T)(x_1)$ is linear. Let $a, b \in X_1$, $\alpha \in \mathbb{C}$ and $x_2 \in X_2$. Then

$$\begin{aligned}\operatorname{cur}(T)(a + \alpha b)(x_2) &= T(a + \alpha b, x_2) \\ &= T(a, x_2) + \alpha T(b, x_2) \\ &= \operatorname{cur}(T)(a)(x_2) + \alpha \operatorname{cur}(T)(b)(x_2) \\ &= [\operatorname{cur}(T)(a) + \alpha \operatorname{cur}(T)(b)](x_2)\end{aligned}$$

Since $x_2 \in X_2$ is arbitrary, $\operatorname{cur}(T)(a + \alpha b) = \operatorname{cur}(T)(a) + \alpha \operatorname{cur}(T)(b)$. Since $a, b \in X_1$ and $\alpha \in \mathbb{C}$ are arbitrary, $\operatorname{cur}(T)$ is linear. Let $(x_1, x_2) \in X_1 \times X_2$. Then

$$\begin{aligned}\|\operatorname{cur}(T)(x_1)(x_2)\| &= \|T(x_1, x_2)\| \\ &\leq (\|T\| \|x_1\|) \|x_2\|\end{aligned}$$

So $\operatorname{cur}(T)(x_1) \in L(X_2, Y)$ and $\|\operatorname{cur}(T)(x_1)\| \leq \|T\| \|x_1\|$. Since $x_1 \in X_1$ is arbitrary, $\operatorname{cur}(T) \in L(X_1; L(X_2; Y))$ and $\|\operatorname{cur} T\| \leq \|T\|$. Since $T \in L(X_1, X_2; Y)$ is arbitrary, $\operatorname{cur}(L(X_1, X_2; Y)) \subset L(X_1; L(X_2; Y))$. Therefore $\operatorname{cur}|_{L(X_1, X_2; Y)} : L(X_1, X_2; Y) \rightarrow L(X_1; L(X_2; Y))$.

3. Let $T \in L(X_1, X_2; Y)$. A previous exercise and an exercise in the section on real numbers imply that

$$\begin{aligned}\|\operatorname{cur}(T)\| &= \sup_{\|x_1\|=1} \|T(x_1)\| \\ &= \sup_{\|x_1\|=1} \left[\sup_{\|x_2\|=1} \|T(x_1)(x_2)\| \right] \\ &= \sup_{\|x_1\|=1, \|x_2\|=1} \|T(x_1)(x_2)\| \\ &= \|T\|\end{aligned}$$

So $\operatorname{cur}|_{L(X_1, X_2; Y)} : L(X_1, X_2; Y) \rightarrow L(X_1; L(X_2; Y))$ is an isometry.

4. Let $T \in L(X_1; L(X_2; Y))$. Define $S : X_1 \times X_2 \rightarrow Y$ by $S(x_1, x_2) = T(x_1)(x_2)$. It is straightforward to show that S is bilinear and for each $(x_1, x_2) \in X_1 \times X_2$,

$$\begin{aligned}\|S(x_1, x_2)\| &= \|T(x_1)(x_2)\| \\ &\leq \|T(x_1)\| \|x_2\| \\ &\leq \|T\| \|x_1\| \|x_2\|\end{aligned}$$

So $S \in L(X_1, X_2; Y)$ and $\|S\| \leq \|T\|$. By construction $\operatorname{cur}(S) = T$. Since $T \in L(X_1; L(X_2; Y))$ is arbitrary, we have that for each $T \in L(X_1; L(X_2; Y))$, there exists $S \in L(X_1, X_2; Y)$ such that $T = \operatorname{cur}(S)$. Hence $\operatorname{cur}|_{L(X_1, X_2; Y)}$ is surjective.

5. Since $\operatorname{cur}|_{L(X_1, X_2; Y)}$ is an isometry, $\operatorname{cur}|_{L(X_1, X_2; Y)}$ is injective. From the previous part, we know that $\operatorname{cur}|_{L(X_1, X_2; Y)}$ is surjective. Hence $\operatorname{cur}|_{L(X_1, X_2; Y)}$ is a bijection. The first part implies that $\operatorname{cur}|_{L(X_1, X_2; Y)}$ is linear. Hence $\operatorname{cur}|_{L(X_1, X_2; Y)} : L(X_1, X_2; Y) \rightarrow L(X_1; L(X_2; Y))$ is an isometric isomorphism. □

Exercise 9.9.0.10. Let X_1, X_2, Y be normed vector spaces. If Y is complete, then $L(X_1, X_2; Y)$ is complete.

Proof. Suppose that Y is complete. Then $L(X_1; L(X_2; Y))$ is complete. Since $L(X_1; L(X_2; Y))$ is isometrically isomorphic to $L(X_1, X_2; Y)$, we have that $L(X_1, X_2; Y)$ is complete. □

Definition 9.9.0.11. Let X_1, X_2 be normed vector spaces, $\phi_1 \in X_1^*$ and $\phi_2 \in X_2^*$. Define $\phi_1 \otimes \phi_2 : X_1 \times X_2$ by $\phi_1 \otimes \phi_2(x_1, x_2) = \phi_1(x_1)\phi_2(x_2)$.

Exercise 9.9.0.12. Let X_1, X_2 be normed vector spaces, $\phi_1 \in X_1^*$ and $\phi_2 \in X_2^*$. Then $\phi_1 \otimes \phi_2 \in L^2(X_1, X_2; \mathbb{C})$.

Proof. Clear. □

Exercise 9.9.0.13. Let X_1, X_2 be normed vector spaces and $(x_1, x_2) \in X_1 \times X_2$. If for each $\phi_1 \in X_1^*$ and $\phi_2 \in X_2^*$, $\phi_1 \otimes \phi_2(x_1, x_2) = 0$, then $x_1 = 0$ or $x_2 = 0$.

Proof. Suppose that $x_1 \neq 0$ and $x_2 \neq 0$. The previous section implies that there exist $\phi_1 \in X_1^*$ and $\phi_2 \in X_2^*$ such that $\phi_1(x_1) = \|x_1\| \neq 0$ and $\phi_2(x_2) = \|x_2\| \neq 0$. Then

$$\begin{aligned}\phi_1 \otimes \phi_2(x_1, x_2) &= \phi_1(x_1)\phi_2(x_2) \\ &\neq 0\end{aligned}$$

□

9.10 Tensor Products of Banach Spaces

(cite "intro to tensor products of banach spaces by Ryan")

9.11 Injective Tensor Product

Definition 9.11.0.1. Let X, Y be vector spaces, $x \in X$ and $y \in Y$. We define $x \otimes_\epsilon y : X^* \times Y^* \rightarrow \mathbb{K}$ by $x \otimes_\epsilon y(\phi, \psi) := \phi(x)\psi(y)$.

Exercise 9.11.0.2. Let X, Y be vector spaces, $x \in X$ and $y \in Y$. Then $x \otimes_\epsilon y \in L^2(X^*, Y^*; \mathbb{K})$.

Proof. Let $\phi_1, \phi_2 \in X^*$, $\psi \in Y^*$ and $\lambda \in \mathbb{K}$. Then

$$\begin{aligned} x \otimes_\epsilon y(\phi_1 + \lambda\phi_2, \psi) &= [\phi_1 + \lambda\phi_2](x)\psi(y) \\ &= \phi_1(x)\psi(y) + \lambda\phi_2(x)\psi(y) \\ &= x \otimes_\epsilon y(\phi_1, \psi) + \lambda x \otimes_\epsilon y(\phi_2, \psi) \end{aligned}$$

Since $\phi_1, \phi_2 \in X^*$, $\psi \in Y^*$ and $\lambda \in \mathbb{K}$ are arbitrary, we have that for each $\psi \in Y^*$, $x \otimes_\epsilon y(\cdot, \psi)$ is linear. Similarly for each $\phi \in X^*$, $x \otimes_\epsilon y(\phi, \cdot)$ is linear. Hence $x \otimes_\epsilon y$ is bilinear and $x \otimes_\epsilon y \in L^2(X^*, Y^*; \mathbb{K})$. $\square$

Definition 9.11.0.3. Let X, Y be vector spaces. We define

- the **injective tensor product of X and Y** , denoted $X \otimes Y \subset L^2(X^*, Y^*; \mathbb{K})$, by

$$X \otimes_\epsilon Y := \text{span}(x \otimes_\epsilon y : x \in X \text{ and } y \in Y),$$

- the **injective tensor map**, denoted $\otimes_\epsilon : X \times Y \rightarrow X \otimes Y$, by $\otimes_\epsilon(x, y) := x \otimes_\epsilon y$.

Exercise 9.11.0.4. Let X, Y be vector spaces, $(x_j)_{j=1}^n \subset X$ and $(y_j)_{j=1}^n \subset Y$. The following are equivalent:

1. $\sum_{j=1}^n x_j \otimes y_j = 0$
2. for each $\phi \in X^*$ and $\psi \in Y^*$, $\sum_{j=1}^n \phi(x_j)\psi(y_j) = 0$
3. for each $\phi \in X^*$, $\sum_{j=1}^n \phi(x_j)y_j = 0$
4. for each $\psi \in Y^*$, $\sum_{j=1}^n \psi(y_j)x_j = 0$

Proof.

1. (1) $\implies$ (2) :

Suppose that $\sum_{j=1}^n x_j \otimes y_j = 0$. Let $\phi \in X^*$ and $\psi \in Y^*$. Then

$$\begin{aligned} \sum_{j=1}^n \phi(x_j)\psi(y_j) &= \phi\left(\sum_{j=1}^n \psi(y_j)x_j\right) \\ &= \end{aligned}$$

2.

3.

$\square$

9.12 Projective Tensor Product

Definition 9.12.0.1. Let X, Y be vector spaces, $x \in X$ and $y \in Y$. We define $x \otimes_\pi y : L^2(X, Y; \mathbb{K}) \rightarrow \mathbb{K}$ by $x \otimes_\pi y(\phi, \psi) := \phi(x)\psi(y)$.

Exercise 9.12.0.2. Let X, Y be vector spaces, $x \in X$ and $y \in Y$. Then $x \otimes_\pi y \in L^2(X, Y; \mathbb{K})^*$.

Proof. Let $A, B \in L^2(X, Y; \mathbb{K})$ and $\lambda \in \mathbb{K}$. Then

$$\begin{aligned} x \otimes_\pi y(A + \lambda B) &= (A + \lambda B)(x, y) \\ &= A(x, y) + \lambda B(x, y) \\ &= x \otimes_\pi y(A) + \lambda x \otimes_\pi y(B) \end{aligned}$$

and

$$\begin{aligned} |x \otimes_\pi y(A)| &= |A(x, y)| \\ &\leq \|x\| \|y\| \|A\| \end{aligned}$$

Since $A, B \in L^2(X, Y; \mathbb{K})$ and $\lambda \in \mathbb{K}$ are arbitrary, we have that $x \otimes_\pi y$ is linear and $\|x \otimes_\pi y\| \leq \|x\| \|y\|$. Hence $x \otimes_\pi y \in L^2(X, Y; \mathbb{K})^*$ $\square$

Definition 9.12.0.3. Let X, Y be vector spaces. We define

- the **projective tensor product of X and Y** , denoted $X \otimes Y \subset L^2(X^*, Y^*; \mathbb{K})$, by

$$X \otimes_\pi Y := \text{span}(x \otimes_\pi y : x \in X \text{ and } y \in Y),$$

- the **projective tensor map**, denoted $\otimes_\pi : X \times Y \rightarrow X \otimes Y$, by $\otimes_\pi(x, y) := x \otimes_\pi y$.

Exercise 9.12.0.4. Let X, Y be vector spaces, $(x_j)_{j=1}^n \subset X$ and $(y_j)_{j=1}^n \subset Y$. The following are equivalent:

1. $\sum_{j=1}^n x_j \otimes y_j = 0$
2. for each $\phi \in X^*$ and $\psi \in Y^*$, $\sum_{j=1}^n \phi(x_j)\psi(y_j) = 0$
3. for each $\phi \in X^*$, $\sum_{j=1}^n \phi(x_j)y_j = 0$
4. for each $\psi \in Y^*$, $\sum_{j=1}^n \psi(y_j)x_j = 0$

Hint: For (4) $\implies$ (1), set $E := \text{span}(x_j)_{j=1}^n$, $F := \text{span}(y_j)_{j=1}^n$ and define $B \in L^2(E, F; \mathbb{K})^*$ by $B := A|_{E \times F}$. Use the fact that $L^2(E, F; \mathbb{K}) = E^* \otimes_\epsilon F^*$ **make this an exercise in the section on multilinear maps.**

Proof.

1. (1) $\implies$ (2) :

Suppose that $\sum_{j=1}^n x_j \otimes_\pi y_j = 0$. Let $\phi \in X^*$ and $\psi \in Y^*$. Define $B_{\phi, \psi} \in L^2(X, Y; \mathbb{K})$ by $B_{\phi, \psi}(x, y) := \phi(x)\psi(y)$. Then

$$\begin{aligned} \sum_{j=1}^n \phi(x_j)\psi(y_j) &= \sum_{j=1}^n B_{\phi, \psi}(x_j, y_j) \\ &= \sum_{j=1}^n x_j \otimes_\pi y_j(B_{\phi, \psi}) \\ &= \left[\sum_{j=1}^n x_j \otimes_\pi y_j \right](B_{\phi, \psi}) \\ &= u(B_{\phi, \psi}) \\ &= 0 \end{aligned}$$

Since $\phi \in X^*$ and $\psi \in Y^*$ are arbitrary, for each $\phi \in X^*$ and $\psi \in Y^*$, $\sum_{j=1}^n \phi(x_j)\psi(y_j) = 0$.

2. (2) $\implies$ (3) :

Suppose that for each $\phi \in X^*$ and $\psi \in Y^*$, $\sum_{j=1}^n \phi(x_j)\psi(y_j) = 0$. Let $\phi \in X^*$ and $\psi \in Y^*$. Then

$$\begin{aligned} \psi\left(\sum_{j=1}^n \phi(x_j)y_j\right) &= \sum_{j=1}^n \phi(x_j)\psi(y_j) \\ &= 0. \end{aligned}$$

Since $\psi \in Y^*$ is arbitrary, Exercise 9.5.0.11 implies that $\sum_{j=1}^n \phi(x_j)y_j = 0$. Since $\phi \in X^*$ is arbitrary, we have that for each $\phi \in X^*$, $\sum_{j=1}^n \phi(x_j)y_j = 0$.

3. (3) $\implies$ (4):

Suppose that for each $\phi \in X^*$, $\sum_{j=1}^n \phi(x_j)y_j = 0$. Let $\psi \in Y^*$. Then

$$\begin{aligned} \phi\left(\sum_{j=1}^n \psi(y_j)x_j\right) &= \sum_{j=1}^n \phi(x_j)\psi(y_j) \\ &= \psi\left(\sum_{j=1}^n \phi(x_j)y_j\right) \\ &= \psi(0) \\ &= 0. \end{aligned}$$

Since $\phi \in X^*$ is arbitrary, we have that $\sum_{j=1}^n \psi(y_j)x_j = 0$. Since $\psi \in Y^*$ is arbitrary, we have that for each $\psi \in Y^*$, $\sum_{j=1}^n \psi(y_j)x_j = 0$.

4. (4) $\implies$ (1):

Suppose that for each $\psi \in Y^*$, $\sum_{j=1}^n \psi(y_j)x_j = 0$. Set $E := \text{span}(x_j : j \in [n])$, $F := \text{span}(y_j : j \in [n])$ and define $B \in L^2(E, F; \mathbb{K})$ by $B := A|_{E \times F}$. **make exercise about $\phi_{0,k} \otimes_\epsilon \psi_{0,l} : k, l$ is a basis when $(\phi_{0,k})_{k=1}^n$ and $(\psi_{0,l})_{l=1}^n$ are bases for E^* , which can be obtained from a basis for E and F , there exist $(\phi_{0,j})_{j=1}^n \subset E^*$ and $(\psi_{0,j})_{j=1}^n \subset F^*$ such that $B = \sum_{k=1}^n \phi_{0,k} \otimes_\epsilon \psi_{0,k}$. Let $j \in [n]$. Exercise 9.5.0.8 implies that there exist $(\phi_j)_{j=1}^n \subset X^*$ and $(\psi_j)_{j=1}^n \subset Y^*$ such that for each $j \in [n]$, $\|\phi_j\| = \|\psi_{0,j}\|$,**

$\|\psi_j\| = \|\psi_{0,j}\|$, $\phi_j|_E = \phi_{0,j}$ and $\psi_j|_F = \psi_{0,j}$. Then

$$\begin{aligned}
 u(A) &= \sum_{j=1}^n x_j \otimes y_j(A) \\
 &= \sum_{j=1}^n A(x_j, y_j) \\
 &= \sum_{j=1}^n B(x_j, y_j) \\
 &= \sum_{j=1}^n \left[\sum_{k=1}^n \phi_{0,k} \otimes \psi_{0,k}(x_j, y_j) \right] \\
 &= \sum_{j=1}^n \left[\sum_{k=1}^n \phi_{0,k}(x_j) \psi_{0,k}(y_j) \right] \\
 &= \sum_{j=1}^n \left[\sum_{k=1}^n \phi_k(x_j) \psi_k(y_j) \right] \\
 &= \sum_{k=1}^n \left[\sum_{j=1}^n \phi_k(x_j) \psi_k(y_j) \right] \\
 &= \sum_{k=1}^n \left[\phi_k \left(\sum_{j=1}^n \psi_k(y_j) x_j \right) \right] \\
 &= \sum_{k=1}^n \phi_k(0) \\
 &= 0
 \end{aligned}$$

Since $A \in L^2(X, Y; \mathbb{K})$ is arbitrary, we have that for each $A \in L^2(X, Y; \mathbb{K})$, $u(A) = 0$. Hence $u = 0$.

□

Definition 9.12.0.5. Let $(X, \|\cdot\|_X), (Y, \|\cdot\|_Y)$ be Banach spaces. We define the **projective norm on $X \otimes Y$** , denoted $\|\cdot\|_\pi : X \otimes Y \rightarrow [0, \infty)$, by

$$\|u\|_\pi := \inf \left\{ \sum_{j=1}^n \|x_j\|_X \|y_j\|_Y : (x_j)_{j=1}^n \subset X, (y_j)_{j=1}^n \subset Y \text{ and } u = \sum_{j=1}^n x_j \otimes y_j \right\}$$

Exercise 9.12.0.6. Let X, Y be Banach spaces. Then

1. $\|\cdot\|_\pi$ is a norm on $X \otimes Y$
2. for each $x \in X$ and $y \in Y$, $\|x \otimes_\pi y\|_\pi = \|x\| \|y\|$.

Proof.

1. (a) Let $u \in X \otimes Y$. Suppose that $\|u\|_\pi = 0$. Set

$$V_u := \left\{ \sum_{j=1}^n \|x_j\| \|y_j\| : (x_j)_{j=1}^n \subset X, (y_j)_{j=1}^n \subset Y \text{ and } u = \sum_{j=1}^n x_j \otimes y_j \right\}$$

Let $\phi \in X^*$, $\psi \in Y^*$ and $\epsilon > 0$. Then there exist $(x_j)_{j=1}^n \subset X$ and $(y_j)_{j=1}^n \subset Y$ such that $u = \sum_{j=1}^n x_j \otimes y_j$ and

$$\begin{aligned} \sum_{j=1}^n \|x_j\| \|y_j\| &< \inf V_u + \epsilon / (\|\phi\| \|\psi\| + 1) \\ &= \|u\|_\pi + \epsilon / (\|\phi\| \|\psi\| + 1) \\ &= \epsilon / (\|\phi\| \|\psi\| + 1). \end{aligned}$$

$$\begin{aligned} |u(\phi, \psi)| &= \sum_{j=1}^n |\phi(x_j) \psi(y_j)| \\ &\leq \sum_{j=1}^n \|\phi\| \|\psi\| \|x_j\| \|y_j\| \\ &= \|\phi\| \|\psi\| \sum_{j=1}^n \|x_j\| \|y_j\| \\ &< \|\phi\| \|\psi\| \frac{\epsilon}{\|\phi\| \|\psi\| + 1} \\ &\leq \epsilon \end{aligned}$$

Since $\epsilon > 0$ is arbitrary, we have that $u(\phi, \psi) = 0$. Since $\phi \in X^*$ and $\psi \in Y^*$ are arbitrary, we have that for each $\phi \in X^*$ and $\psi \in Y^*$, $u(\phi, \psi) = 0$. Therefore $u = 0$.

(b) Let $u \in X \otimes Y$ and $\lambda \in \mathbb{K}$. Set

$$V_u := \left\{ \sum_{j=1}^n \|x_j\| \|y_j\| : (x_j)_{j=1}^n \subset X, (y_j)_{j=1}^n \subset Y \text{ and } u = \sum_{j=1}^n x_j \otimes y_j \right\}$$

and

$$V_{\lambda u} := \left\{ \sum_{j=1}^n \|x_j\| \|y_j\| : (x_j)_{j=1}^n \subset X, (y_j)_{j=1}^n \subset Y \text{ and } \lambda u = \sum_{j=1}^n x_j \otimes y_j \right\}.$$

Let $\epsilon > 0$. Since $\|u\|_\pi = \inf V_u$, there exists $a \in V_u$ such that $a < \|u\|_\pi + \epsilon / (|\lambda| + 1)$. Therefore, there exists $(x_j)_{j=1}^n \subset X$ and $(y_j)_{j=1}^n \subset Y$ such that $u = \sum_{j=1}^n x_j \otimes y_j$ and $a = \sum_{j=1}^n \|x_j\| \|y_j\|$. Then $\lambda u = \sum_{j=1}^n (\lambda x_j) \otimes y_j$ and

$$\begin{aligned} \|\lambda u\|_\pi &= \inf V_{\lambda u} \\ &\leq \sum_{j=1}^n \|\lambda x_j\| \|y_j\| \\ &= |\lambda| \sum_{j=1}^n \|x_j\| \|y_j\| \\ &= |\lambda| a \\ &< |\lambda| \left(\|u\|_\pi + \frac{\epsilon}{|\lambda| + 1} \right) \\ &= |\lambda| \|u\|_\pi + \epsilon \frac{|\lambda|}{|\lambda| + 1} \\ &< |\lambda| \|u\|_\pi + \epsilon \end{aligned}$$

Since $\epsilon > 0$ is arbitrary, we have that $\|\lambda u\|_\pi \leq |\lambda| \|u\|_\pi$.

- Suppose that $\lambda = 0$. Then $\lambda u = 0 = 0 \otimes 0$. Hence

$$\begin{aligned} 0 &\leq \|\lambda u\|_\pi \\ &= \inf V_{\lambda u} \\ &\leq \|0\| \|0\| \\ &= 0 \end{aligned}$$

Thus

$$\begin{aligned} \|\lambda u\|_\pi &= 0 \\ &= |\lambda| \|u\|_\pi \end{aligned}$$

- Suppose that $\lambda \neq 0$. Then

$$\begin{aligned} \|u\|_\pi &= \|\lambda^{-1}(\lambda u)\|_\pi \\ &\leq |\lambda^{-1}| \|\lambda u\|_\pi \\ &= |\lambda|^{-1} \|\lambda u\|_\pi \end{aligned}$$

Hence $|\lambda| \|u\|_\pi \leq \|\lambda u\|_\pi$. Thus $\|\lambda u\|_\pi = |\lambda| \|u\|_\pi$.

(c)

2.

□

9.13 Unbounded Operators

Definition 9.13.0.1. Let X, Y be normed vector spaces, $D \subset X$ a subspace and $T : D \rightarrow Y$ linear. Then T is said to be an **operator from X to Y** . We define

$$\text{Op}(X, Y) := \{T : D \rightarrow Y : D \subset X \text{ is a subspace and } T \text{ is an operator from } X \text{ to } Y\}.$$

Note 9.13.0.2. By definition, for each $T \in \text{Op}(X, Y)$, $0 \in \text{dom}(T)$.

Definition 9.13.0.3. Let X, Y, Z be normed vector spaces. We define

- $\text{add}_{\text{unb}} : \text{Op}(X, Y) \times \text{Op}(X, Y) \rightarrow \text{Op}(X, Y)$ by

$$\text{add}_{\text{unb}}(S, T) := S|_{\text{dom}(T)} + T|_{\text{dom}(S)},$$

- $\text{scalar_mult}_{\text{unb}} : \mathbb{K} \times \text{Op}(X, Y) \rightarrow \text{Op}(X, Y)$ by

$$\text{scalar_mult}_{\text{unb}}(\lambda, T) := \lambda T,$$

- $\circ_{\text{unb}} : \text{Op}(Y, Z) \times \text{Op}(X, Y) \rightarrow \text{Op}(X, Z)$ by

$$S \circ_{\text{unb}} T := S \circ (T|_{T^{-1}(\text{dom}(S))}).$$

Note 9.13.0.4. When the context is clear, we write $S + T$, λT and $S \circ T$ in place of $\text{add}_{\text{unb}}(S, T)$, $\text{scalar_mult}_{\text{unb}}(\lambda, T)$ and $S \circ_{\text{unb}} T$ respectively.

Definition 9.13.0.5. Let X, Y be normed vector spaces and $S, T \in \text{Op}(X, Y)$.

- Then S is said to be an **linear extension of T** if $S \in \text{Ext}(T)$.
- We define $\text{OpExt}(T, X) := \{S \in \text{Op}(X, Y) : S \text{ is a linear extension of } T\}$.

Note 9.13.0.6. When the context is clear, we may write $\text{OpExt}(T)$ in place of $\text{OpExt}(T, X)$.

Exercise 9.13.0.7. For each $T \in \text{Op}(X, Y)$, if $T \in L(\text{dom}(T), Y)$, then there exists $S \in \text{Ext}(T) \cap L(X; Y)$ such that $\|S\| = \|T\|$.
maybe move this as corollary after hahn banach after rewrite

Proof. Hahn Banach or continuity if $\text{dom}(T)$ is dense in X . **FINISH!!!**

□

9.13.1 Closed Operators

Definition 9.13.1.1. Let X, Y be normed vector spaces and $T \in \text{Op}(X, Y)$. Then T is said to be

- **closed** if $\Gamma(T) \subset X \times Y$ is closed.
- **closable** if there exists $S \in \text{Ext}(T)$ such that S is closed.

Exercise 9.13.1.2. Let X, Y be normed vector spaces and $T \in \text{Op}(X, Y)$. If T is closable, then there exists a unique $S \in \text{Ext}(T)$ such that S is closed and for each $S' \in \text{Ext}(T)$, S' is closed implies that $S' \in \text{Ext}(S)$.

Proof. Suppose that T is closeable. Then there exists $S_0 \in \text{Ext}(T)$ such that S_0 is closed. Exercise 17.3.0.5 (ref ex in set theory section about restrictions/extensions of functions) implies that $\Gamma(T) \subset \Gamma(S_0)$. Since S_0 is closed, we have that $\Gamma(S_0) \subset X \times Y$ is closed. Thus

$$\begin{aligned}\Gamma(T) &\subset \text{cl}\Gamma(T) \\ &\subset \text{cl}\Gamma(S_0) \\ &= \Gamma(S_0).\end{aligned}$$

Exercise 17.3.0.5 (ref ex in top section that π_1, π_2 are closed maps) implies that $\pi_1(\text{cl}\Gamma(T)) \subset X$ and $\pi_2(\text{cl}\Gamma(T)) \subset Y$ are closed. Define $D \subset X$ and $S : D \rightarrow Y$ by $D := \pi_1(\text{cl}\Gamma(T))$ and $S(x) :=$ by **FINISH!!!** □

Chapter 10

Hilbert Spaces

10.1 TODO

- Express $V^* \cong \bar{V}$ where $\bar{V}$ is just V , but with $\lambda * v = \lambda^* v$. so Rieze rep theorem reads $V \cong \bar{V}^*$ or $V \cong \bar{V}^*$
- discuss projection maps
- show internal direct sum isomorphic to external
- discuss quotient hilbert space?
- discus subspaces

10.2 Introduction

10.2.1 Inner product Spaces

Definition 10.2.1.1. Let H be a vector space and $\langle \cdot, \cdot \rangle : H \times H \rightarrow \mathbb{C}$. Then

- $\langle \cdot, \cdot \rangle$ is said to be an **inner product** on H if for each $x, y, z \in H$ and $\lambda \in \mathbb{C}$
 1. $\langle x, y + \lambda z \rangle = \langle x, y \rangle + \lambda \langle x, z \rangle$
 2. $\langle x, y \rangle = \langle y, x \rangle^*$
 3. $\langle x, x \rangle \geq 0$
 4. if $\langle x, x \rangle = 0$, then $x = 0$.
- $(H, \langle \cdot, \cdot \rangle)$ is said to be a **inner product space** if $\langle \cdot, \cdot \rangle$ is an inner product on H .

Note 10.2.1.2. When the context is clear, we suppress the inner product $\langle \cdot, \cdot \rangle : H \times H \rightarrow \mathbb{C}$.

Note 10.2.1.3. In mathematics, inner products are conventionally linear in the first argument. The convention in physics is linearity in the second argument. The physics convention notationally generalizes the dot product as matrix multiplication when identifying $\mathbb{C}^n$ with $\mathbb{C}^{n \times 1}$ as is done in an introductory linear algebra class. For example, for $x, y \in \mathbb{C}^n$ $\langle x, y \rangle = \bar{x}^\top y$.

Definition 10.2.1.4. Let $(H, \langle \cdot, \cdot \rangle)$ be an inner product space. We define $\langle \cdot, \cdot \rangle_{\bar{H}} : \bar{H} \times \bar{H} \rightarrow \mathbb{C}$ by $\langle x, y \rangle_{\bar{H}} := \langle x, y \rangle^*$.

Exercise 10.2.1.5. Let $(H, \langle \cdot, \cdot \rangle)$ be an inner product space. Then $(\bar{H}, \langle \cdot, \cdot \rangle_{\bar{H}})$ is an inner product space.

Proof. Let $x, y, z \in \bar{H}$ and $\lambda \in \mathbb{C}$. Then

1.

$$\begin{aligned}
\langle x, y + \lambda^* z \rangle_{\bar{H}} &= \langle x, y + \lambda^* z \rangle^* \\
&= \langle x, y \rangle^* + \langle x, \lambda^* z \rangle^* \\
&= \langle x, y \rangle^* + \langle x, \lambda^* z \rangle^* \\
&= \langle x, y \rangle^* + \langle \lambda^* z, x \rangle \\
&= \langle x, y \rangle^* + \lambda \langle z, x \rangle \\
&= \langle x, y \rangle^* + \lambda \langle x, z \rangle^* \\
&= \langle x, y \rangle_{\bar{H}} + \lambda \langle x, z \rangle_{\bar{H}}
\end{aligned}$$

2.

$$\begin{aligned}
\langle x, y \rangle_{\bar{H}} &= \langle x, y \rangle^* \\
&= (\langle y, x \rangle^*)^* \\
&= \langle y, x \rangle_{\bar{H}}^*
\end{aligned}$$

3.

$$\begin{aligned}
\langle x, x \rangle_{\bar{H}} &= \langle x, x \rangle^* \\
&= \langle x, x \rangle \\
&\geq 0
\end{aligned}$$

4. Suppose that $\langle x, x \rangle_{\bar{H}} = 0$. Then

$$\begin{aligned}
\langle x, x \rangle &= \langle x, x \rangle^* \\
&= \langle x, x \rangle_{\bar{H}} \\
&= 0.
\end{aligned}$$

Therefore $x = 0$.Thus $\langle \cdot, \cdot \rangle_{\bar{H}}$ is an inner product on $\bar{H}$ and $(\bar{H}, \langle \cdot, \cdot \rangle_{\bar{H}})$ is an inner product space. □**Exercise 10.2.1.6.** Let H be an inner product space, $(x_j)_{j=1}^n, (y_j)_{j=1}^n \subset H$ and $(\alpha_j)_{j=1}^n, (\beta_j)_{j=1}^n \subset \mathbb{C}$. Then

$$\left\langle \sum_{i=1}^n \alpha_i x_i, \sum_{j=1}^n \beta_j y_j \right\rangle = \sum_{i=1}^n \sum_{j=1}^n \alpha_i^* \beta_j \langle x_i, y_j \rangle$$

Proof. Clear. □**Definition 10.2.1.7.** Let $(H, \langle \cdot, \cdot \rangle)$ be an inner product space. Define the **induced norm**, denoted $\| \cdot \| : \rightarrow \mathbb{C}$, by

$$\|x\| = \langle x, x \rangle^{1/2}$$

Note 10.2.1.8. Unless otherwise specified, we only consider the induced norm on any given Hilbert space.**Exercise 10.2.1.9. Cauchy-Schwarz Inequality**Let H be an inner product space. Then for each $x, y \in H$, $|\langle x, y \rangle| \leq \|x\| \|y\|$ and $|\langle x, y \rangle| = \|x\| \|y\|$ iff $x \in \text{span}(y)$.**Hint:** For $x, y \in H$, put $z = \text{sgn} \langle x, y \rangle^* y$ and Consider $f : \mathbb{R} \rightarrow [0, \infty)$ defined by $f(t) = \|x - tz\|^2$ *Proof.* Let $x, y \in H$. If $y = 0$, then the claim holds trivially. Suppose that $y \neq 0$. Put $z = \text{sgn} \langle x, y \rangle^* y$. So $\langle x, z \rangle = |\langle x, y \rangle|$ and $\|z\| = \|y\|$. Define $f : \mathbb{R} \rightarrow [0, \infty)$ by

$$f(t) = \|x - tz\|^2$$

Then for each $t \in \mathbb{R}$,

$$\begin{aligned} 0 &\leq f(t) \\ &= \|x - tz\|^2 \\ &= \|x\|^2 + |t|^2 \|z\|^2 - 2\operatorname{Re}(t\langle x, z \rangle) \\ &= \|x\|^2 + t^2 \|y\|^2 - 2t|\langle x, y \rangle| \end{aligned}$$

Thus f is a quadratic with a minimum at $t_0 = \frac{|\langle x, y \rangle|}{\|y\|^2}$. Hence

$$\begin{aligned} 0 &\leq f(t_0) \\ &= \|x\|^2 + \frac{|\langle x, y \rangle|^2}{\|y\|^4} - 2\frac{|\langle x, y \rangle|^2}{\|y\|^2} \\ &= \|x\|^2 - \frac{|\langle x, y \rangle|^2}{\|y\|^2} \end{aligned}$$

Which implies that

$$|\langle x, y \rangle|^2 \leq \|x\|^2 \|y\|^2$$

and hence the claim holds. Clearly if $x \in \operatorname{span}(y)$, then equality holds. Conversely, if equality holds, then $x - z = 0$ which implies that $x \in \operatorname{span}(y)$. $\square$

Exercise 10.2.1.10. Let H be an inner product space. Then the induced norm, $\|\cdot\| : H \rightarrow \mathbb{C}$, is a norm.

Proof. Let $x, y \in H$ and $\lambda \in \mathbb{C}$. Then

1. By definition, if $\|x\| = 0$, then $\langle x, x \rangle = 0$, which implies that $x = 0$.
2. Note that

$$\begin{aligned} \|\lambda x\|^2 &= \langle \lambda x, \lambda x \rangle \\ &= \lambda * \lambda \langle x, x \rangle \\ &= |\lambda|^2 \|x\|^2 \end{aligned}$$

$$\text{So } \|\lambda x\| = |\lambda| \|x\|$$

3. The Cauchy-Schwarz inequality implies that

$$\begin{aligned} \|x + y\|^2 &= \|x\|^2 + \|y\|^2 + 2\operatorname{Re}(\langle x, y \rangle) \\ &\leq \|x\|^2 + \|y\|^2 + 2|\langle x, y \rangle| \\ &\leq \|x\|^2 + \|y\|^2 + 2\|x\|\|y\| \\ &= (\|x\| + \|y\|)^2 \end{aligned}$$

$$\text{Hence } \|x + y\| \leq \|x\| + \|y\|.$$

$\square$

Exercise 10.2.1.11. Let H be an inner-product space and $y \in H$. Then $y = 0$ iff for each $x \in H$, $\langle y, x \rangle = 0$.

Proof.

• ($\implies$):

Suppose that $y = 0$. Let $x \in H$. Then

$$\begin{aligned} \langle y, x \rangle &= \langle 0, x \rangle \\ &= \langle 0 + 0, x \rangle \\ &= \langle 0, x \rangle + \langle 0, x \rangle \\ &= \langle y, x \rangle + \langle y, x \rangle \end{aligned}$$

Hence $\langle y, x \rangle = 0$. Since $x \in H$ is arbitrary, we have that for each $x \in H$, $\langle y, x \rangle = 0$.

- ($\Leftarrow$):

Suppose that $y \neq 0$. Then

$$\begin{aligned} 0 &\neq \|y\| \\ &= \langle y, y \rangle \end{aligned}$$

Define $x \in H$ by $x := y$. Then $\langle y, x \rangle \neq 0$. Hence $y \neq 0$ implies that there exists $x \in H$ such that $\langle y, x \rangle \neq 0$. By contrapositive, if for each $x \in H$, $\langle y, x \rangle = 0$, then $y = 0$. □

Exercise 10.2.1.12. Let $H_1, H_2 \in \text{Obj}(\mathbf{Hilb})$ and $A \in L(H_1, H_2)$. Then $A = 0$ iff for each $x \in H_1$ and $y \in H_2$, $\langle y, Ax \rangle = 0$.

Proof.

- ($\Rightarrow$):

Suppose that $A = 0$. Then for each $x \in H_1$ and $y \in H_2$,

$$\begin{aligned} \langle y, Ax \rangle &= \langle y, 0 \rangle \\ &= 0. \end{aligned}$$

- ($\Leftarrow$):

Suppose that for each $x \in H_1$ and $y \in H_2$, $\langle y, Ax \rangle = 0$. Let $x \in H_1$. Then for each $y \in H_2$,

$$\begin{aligned} \langle Ax, y \rangle &= \langle y, Ax \rangle^* \\ &= 0. \end{aligned}$$

Exercise 10.2.1.11 implies that $Ax = 0$. Since $x \in H_1$ is arbitrary, we have that for each $x \in H_1$, $Ax = 0$. Hence $A = 0$. □

Definition 10.2.1.13. Let X be a normed vector space. Then X is said to satisfy the **parallelogram law** if for each $x, y \in X$,

$$\|x + y\|^2 + \|x - y\|^2 = 2(\|x\|^2 + \|y\|^2).$$

Exercise 10.2.1.14. Polarization Identity:

Let H be an inner product space. For each $x, y \in H$,

$$\langle x, y \rangle = \frac{1}{4} \left[\|x + y\|^2 - \|x - y\|^2 - i\|x + iy\|^2 + \|x - iy\|^2 \right].$$

Proof. Let $x, y \in H$. Then

$$\begin{aligned} \|x + y\|^2 - \|x - y\|^2 - i\|x + iy\|^2 + i\|x - iy\|^2 &= 4\text{Re} \langle x, y \rangle - 4i\text{Re} \langle x, iy \rangle \\ &= 4\text{Re} \langle x, y \rangle - 4i\text{Re} i\langle x, y \rangle \\ &= 4\text{Re} \langle x, y \rangle + 4\text{Im} \langle x, y \rangle \\ &= 4\langle x, y \rangle. \end{aligned}$$

Hence

$$\langle x, y \rangle = \frac{1}{4} \left[\|x + y\|^2 - \|x - y\|^2 - i\|x + iy\|^2 + i\|x - iy\|^2 \right].$$

Since $x, y \in H$ are arbitrary, we have that for each $x, y \in H$,

$$\langle x, y \rangle = \frac{1}{4} \left[\|x + y\|^2 - \|x - y\|^2 - i\|x + iy\|^2 + i\|x - iy\|^2 \right].$$

□

Exercise 10.2.1.15. Parallelogram Law Equivalence:

Let X be a normed vector space. There exists $\langle \cdot, \cdot \rangle : X \times X \rightarrow \mathbb{C}$ such that $(X, \langle \cdot, \cdot \rangle)$ is an inner product space and for each $x \in X$, $\|x\|^2 = \langle x, x \rangle$ iff X satisfies the parallelogram law.

Proof.

• ($\Rightarrow$):

Suppose that there exists $\langle \cdot, \cdot \rangle : X \times X \rightarrow \mathbb{C}$ such that $(X, \langle \cdot, \cdot \rangle)$ is an inner product space and for each $x \in X$, $\|x\|^2 = \langle x, x \rangle$. Let $x, y \in X$. Then

$$\begin{aligned} \|x + y\|^2 + \|x - y\|^2 &= (\|x\|^2 + \|y\|^2 + 2\operatorname{Re}(\langle x, y \rangle)) + (\|x\|^2 + \|y\|^2 - 2\operatorname{Re}(\langle x, y \rangle)) \\ &= 2(\|x\|^2 + \|y\|^2) \end{aligned}$$

Since $x, y \in H$ are arbitrary, we have that for each $x, y \in X$, $\|x + y\|^2 + \|x - y\|^2 = 2(\|x\|^2 + \|y\|^2)$. Hence X satisfies the parallelogram law.

• ($\Leftarrow$):

Suppose that X satisfies the parallelogram law. Define $\langle \cdot, \cdot \rangle : X \times X \rightarrow \mathbb{C}$ by

$$\langle x, y \rangle := \frac{1}{4} \left[\|x + y\|^2 - \|x - y\|^2 - i\|x + iy\|^2 + i\|x - iy\|^2 \right].$$

Let $x, y, z \in X$ and $\lambda \in \mathbb{C}$.

1. We note that

$$\begin{aligned} \langle y, x \rangle &= \frac{1}{4} \left[\|y + x\|^2 - \|y - x\|^2 - i\|y + ix\|^2 + i\|y - ix\|^2 \right] \\ &= \frac{1}{4} \left[\|x + y\|^2 - \|x - y\|^2 - i\|iy - x\|^2 + i\|iy + x\|^2 \right] \\ &= \frac{1}{4} \left[\|x + y\|^2 - \|x - y\|^2 - i\|x - iy\|^2 + i\|x + iy\|^2 \right] \\ &= \frac{1}{4} \left[\|x + y\|^2 - \|x - y\|^2 - i\|x + iy\|^2 + i\|x - iy\|^2 \right]^* \\ &= \langle x, y \rangle^*. \end{aligned}$$

2. • By repeated application of the parallelogram law, we have that

$$\begin{aligned} 4\langle x, y + z \rangle &= \|x + (y + z)\|^2 - \|x - (y + z)\|^2 - i\|x + i(y + z)\|^2 + i\|x - i(y + z)\|^2 \\ &= \|(x + y) + z\|^2 - \|(x - y) - z\|^2 - i\|(x + iy) + iz\|^2 + i\|(x - iy) - iz\|^2 \\ &= \left[2(\|x + y\|^2 + \|z\|^2) - \|(x + y) - z\|^2 \right] \\ &\quad - \left[2(\|x - y\|^2 + \|z\|^2) - \|(x - y) + z\|^2 \right] \\ &\quad - i \left[2(\|x + iy\|^2 + \|z\|^2) - \|(x + iy) - iz\|^2 \right] \\ &\quad + i \left[2(\|x - iy\|^2 + \|z\|^2) - \|(x - iy) + iz\|^2 \right] \\ &= 2(\|x + y\|^2 - \|x - y\|^2 - i\|x + iy\|^2 + i\|x - iy\|^2) \\ &\quad + (\|(x - y) + z\|^2 - \|(x + y) - z\|^2 - i\|(x - iy) + iz\|^2 + i\|(x + iy) - iz\|^2) \\ &= 8\langle x, y \rangle + \|(x - y) + z\|^2 - \|(x + y) - z\|^2 - i\|(x - iy) + iz\|^2 + i\|(x + iy) - iz\|^2. \end{aligned}$$

Applying the parallelogram, we obtain

$$\begin{aligned}
 \|(x - y) + z\|^2 - \|(x + y) - z\|^2 &= \|(x + z) - y\|^2 - \|(x - z) + y\|^2 \\
 &= \left[2(\|x + z\|^2 + \|y\|^2) - \|(x + z) + y\|^2 \right] \\
 &\quad - \left[2(\|x - z\|^2 + \|y\|^2) - \|(x - z) - y\|^2 \right] \\
 &= 2(\|x + z\|^2 - \|x - z\|^2) + \|(x - z) - y\|^2 - \|(x + z) + y\|^2
 \end{aligned}$$

and

$$\begin{aligned}
 \|(x + iy) - iz\|^2 - \|(x - iy) + iz\|^2 &= \|(x - iz) + iy\|^2 - \|(x + iz) - iy\|^2 \\
 &= \left[2(\|x - iz\|^2 + \|y\|^2) - \|(x - iz) - iy\|^2 \right] \\
 &\quad - \left[2(\|x + iz\|^2 + \|y\|^2) - \|(x + iz) + iy\|^2 \right] \\
 &= 2(\|x - iz\|^2 - \|x + iz\|^2) + \|(x + iz) + iy\|^2 - \|(x - iz) - iy\|^2.
 \end{aligned}$$

Therefore

$$\begin{aligned}
 &\|(x - y) + z\|^2 - \|(x + y) - z\|^2 \\
 &- i\|(x - iy) + iz\|^2 + i\|(x + iy) - iz\|^2 = 2(\|x + z\|^2 - \|x - z\|^2) + \|(x - z) - y\|^2 - \|(x + z) + y\|^2 \\
 &\quad + 2(i\|x - iz\|^2 - i\|x + iz\|^2) + i\|(x + iz) + iy\|^2 - i\|(x - iz) - iy\|^2 \\
 &= 2(\|x + z\|^2 - \|x - z\|^2 - i\|x + iz\|^2 + i\|x - iz\|^2) \\
 &\quad + \left[\|(x - z) - y\|^2 - \|(x + z) + y\|^2 \right. \\
 &\quad \left. + i\|(x + iz) + iy\|^2 - i\|(x - iz) - iy\|^2 \right] \\
 &= 8\langle x, z \rangle + \left[\|x - (y + z)\|^2 - \|x + (y + z)\|^2 \right. \\
 &\quad \left. + i\|x + i(y + z)\|^2 - i\|x - i(y + z)\|^2 \right] \\
 &= 8\langle x, z \rangle - \left[\|x + (y + z)\|^2 - \|x - (y + z)\|^2 \right. \\
 &\quad \left. - i\|x + i(y + z)\|^2 + i\|x - i(y + z)\|^2 \right] \\
 &= 8\langle x, z \rangle - 4\langle x, y + z \rangle.
 \end{aligned}$$

Combining the previous equations, we have that

$$\begin{aligned}
 4\langle x, y + z \rangle &= 8\langle x, y \rangle + 8\langle x, z \rangle - 4\langle x, y + z \rangle \\
 &= 8(\langle x, y \rangle + \langle x, z \rangle) - 4\langle x, y + z \rangle.
 \end{aligned}$$

Thus $8\langle x, y + z \rangle = 8(\langle x, y \rangle + \langle x, z \rangle)$ and finally $\langle x, y + z \rangle = \langle x, y \rangle + \langle x, z \rangle$.

- By induction and additivity, for each $m \in \mathbb{Z}$, $\langle x, mz \rangle = m\langle x, z \rangle$. Let $q \in \mathbb{Q}$. Then there exist $m \in \mathbb{Z}$ and $n \in \mathbb{N}$ such that

$q = m/n$. Then

$$\begin{aligned}
\langle x, qz \rangle &= m\langle x, n^{-1}z \rangle \\
&= m\frac{1}{4}\left[\|x + n^{-1}z\|^2 - \|x - n^{-1}z\|^2 - i\|x + in^{-1}z\|^2 + i\|x - in^{-1}z\|^2\right] \\
&= m\frac{1}{4}\left[n^2\|nx + z\|^2 - n^2\|x - n^{-1}z\|^2 - n^2i\|x + in^{-1}z\|^2 + n^2i\|x - in^{-1}z\|^2\right] \\
&= m\frac{1}{4}\left[n^{-2}\|nx + z\|^2 - n^{-2}\|nx - z\|^2 - n^{-2}i\|nx + iz\|^2 + n^{-2}i\|nx - iz\|^2\right] \\
&= mn^{-2}\langle nx, z \rangle \\
&= mn^{-2}\langle z, nx \rangle^* \\
&= mn^{-2}[n\langle z, x \rangle]^* \\
&= mn^{-2}n\langle z, x \rangle^* \\
&= q\langle x, z \rangle.
\end{aligned}$$

Since $q \in \mathbb{Q}$ is arbitrary, we have that for each $q \in \mathbb{Q}$, $\langle x, qz \rangle = q\langle x, z \rangle$. We note that

$$\begin{aligned}
\langle x, iz \rangle &= \frac{1}{4}\left[\|x + (iz)\|^2 - \|x - (iz)\|^2 - i\|x + i(iz)\|^2 + i\|x - i(iz)\|^2\right] \\
&= \frac{1}{4}\left[\|x + iz\|^2 - \|x - iz\|^2 - i\|x - z\|^2 + i\|x + z\|^2\right] \\
&= \frac{1}{4}\left[i\|x + z\|^2 - i\|x - z\|^2 + \|x + iz\|^2 - \|x - iz\|^2\right] \\
&= i\frac{1}{4}\left[\|x + z\|^2 - \|x - z\|^2 - i\|x + iz\|^2 + i\|x - iz\|^2\right] \\
&= i\langle x, z \rangle.
\end{aligned}$$

Let $\lambda_0 \in \mathbb{Q}[i]$. Then there exist $a, b \in \mathbb{Q}$ such that $\lambda_0 = a + ib$. Then from before, we have that

$$\begin{aligned}
\langle x, \lambda_0 z \rangle &= \langle x, (a + ib)z \rangle \\
&= \langle x, az + ibz \rangle \\
&= \langle x, az + ibz \rangle \\
&= \langle x, az \rangle + \langle x, ibz \rangle \\
&= a\langle x, z \rangle + i\langle x, bz \rangle \\
&= a\langle x, z \rangle + ib\langle x, z \rangle \\
&= (a + ib)\langle x, z \rangle \\
&= \lambda_0\langle x, z \rangle.
\end{aligned}$$

Since $\lambda_0 \in \mathbb{Q}[i]$ is arbitrary, we have that for each $\lambda_0 \in \mathbb{Q}[i]$, $\langle x, \lambda_0 z \rangle = \lambda_0\langle x, z \rangle$. Since $\mathbb{Q}[i]$ is dense in $\mathbb{C}$, there exists $(\lambda_n)_{n \in \mathbb{N}} \subset \mathbb{Q}[i]$ such that $\lambda_n \rightarrow \lambda$. Since $\langle \cdot, \cdot \rangle$ is continuous, we have that

$$\begin{aligned}
\langle x, \lambda z \rangle &= \lim_{n \rightarrow \infty} \langle x, \lambda_n z \rangle \\
&= \lim_{n \rightarrow \infty} [\lambda_n \lim_{n \rightarrow \infty} \langle x, z \rangle] \\
&= \lambda\langle x, z \rangle.
\end{aligned}$$

Hence $\langle x, y + \lambda z \rangle = \langle x, y \rangle + \lambda\langle x, z \rangle$.

3. The parallelogram law implies that

$$\begin{aligned}
 \langle x, x \rangle &= \frac{1}{4} \left[\|x + x\|^2 - \|x - x\|^2 - i\|x + ix\|^2 + i\|x - ix\|^2 \right] \\
 &= \frac{1}{4} \left[4\|x\|^2 - i\|(1+i)x\|^2 + i\|(1-i)x\|^2 \right] \\
 &= \frac{1}{4} \left[4\|x\|^2 - i|1+i|\|x\|^2 + i|1-i|\|x\|^2 \right] \\
 &= \frac{1}{4} \left[4\|x\|^2 + i(|1-i| - |1+i|)\|x\|^2 \right] \\
 &= \frac{1}{4} \left[4\|x\|^2 + i(|1-i| - |1+i|)\|x\|^2 \right] \\
 &= \frac{1}{4} \left[4\|x\|^2 + i(\sqrt{2} - \sqrt{2})\|x\|^2 \right] \\
 &= \|x\|^2 \\
 &\geq 0.
 \end{aligned}$$

4. if $\langle x, x \rangle = 0$, then from before, we have that

$$\begin{aligned}
 \|x\|^2 &= \langle x, x \rangle \\
 &= 0.
 \end{aligned}$$

Since $\|\cdot\|$ is a norm on X , $x = 0$.

Thus $\langle \cdot, \cdot \rangle$ is an inner product on X and $\langle x, x \rangle = \|x\|^2$. □

Definition 10.2.1.16. Let H be an inner product space, $x, y \in H$ and $S \subset H$. Then

1. x and y are said to be **orthogonal**, denoted $x \perp y$, if $\langle x, y \rangle = 0$.
2. S is said to be **orthogonal** if for each $x, y \in S$, $x \neq y$ implies that $x \perp y$.

Definition 10.2.1.17. Let H be an inner product space and $E \subset H$ a closed subspace. We define the **orthogonal complement of E** , denoted $E^\perp$, by

$$E^\perp = \{x \in H : \text{for each } y \in E, x \perp y\}$$

Exercise 10.2.1.18. Let H be an inner product space and $E \subset H$. Then $E^\perp$ is a closed subspace of H .

Proof.

- Let $x, y \in E^\perp$ and $\lambda \in \mathbb{C}$. Then for each $z \in E$,

$$\begin{aligned}
 \langle x + \lambda y, z \rangle &= \langle x, z \rangle + \lambda \langle y, z \rangle \\
 &= 0
 \end{aligned}$$

Hence $x + \lambda y \in E^\perp$. Thus $E^\perp$ is a subspace of H .

- Let $(x_n)_{n \in \mathbb{N}} \subset E^\perp$ and $x \in H$. Suppose that $x_n \rightarrow x$. Then for each $z \in E$, continuity implies that

$$\begin{aligned}
 \langle x, z \rangle &= \lim_{n \rightarrow \infty} \langle x_n, z \rangle \\
 &= \lim_{n \rightarrow \infty} 0 \\
 &= 0
 \end{aligned}$$

Hence $x \in E^\perp$. Since $(x_n)_{n \in \mathbb{N}} \subset E^\perp$ and $x \in H$ with $x_n \rightarrow x$ are arbitrary, we have that $E^\perp$ is closed.

□

Exercise 10.2.1.19. Let H be an inner product space. Then

1. $H^\perp = \{0\}$.
2. $\{0\}^\perp = H$

Proof.

1. Let $z \in H^\perp$. By definition, for each $x \in H$, $\langle z, x \rangle = 0$. A previous exercise implies that $z = 0$. Since $z \in H^\perp$ is arbitrary, $H^\perp = \{0\}$.
2. Let $z \in H$. Trivially, for each $x \in \{0\}$, $\langle z, x \rangle = 0$. Thus $z \in \{0\}^\perp$. Since $z \in H$ is arbitrary, $H \subset \{0\}^\perp$. Since trivially, $\{0\}^\perp \subset H$, we have that $\{0\}^\perp = H$.

□

Exercise 10.2.1.20. Let H be an inner product space and $E, F \subset H$. If $E \subset F$, then $F^\perp \subset E^\perp$.

Proof. Suppose that $E \subset F$. Let $x \in F^\perp$ and $z \in E$. By definition, for each $y \in F$, $\langle x, y \rangle = 0$. Since $E \subset F$, $z \in F$. Hence $\langle x, z \rangle = 0$. Since $x \in F^\perp$ is arbitrary, we have that for each $z \in E$, $\langle x, z \rangle = 0$. Hence $x \in E^\perp$. Since $x \in F^\perp$ is arbitrary, we have that $F^\perp \subset E^\perp$. □

Exercise 10.2.1.21. Pythagorean Theorem:

Let H be an inner product space and $(x_j)_{j=1}^n \subset H$. If $(x_j)_{j=1}^n$ is orthogonal, then

$$\left\| \sum_{j=1}^n x_j \right\|^2 = \sum_{j=1}^n \|x_j\|^2$$

Proof. We have that

$$\begin{aligned} \left\| \sum_{j=1}^n x_j \right\|^2 &= \left\langle \sum_{i=1}^n x_i, \sum_{j=1}^n x_j \right\rangle \\ &= \sum_{i=1}^n \sum_{j=1}^n \langle x_i, x_j \rangle \\ &= \sum_{j=1}^n \langle x_j, x_j \rangle \\ &= \sum_{j=1}^n \|x_j\|^2 \end{aligned}$$

□

Exercise 10.2.1.22. Let H be an inner product space and $S \subset H$. Suppose that $0 \notin S$. If S is orthogonal, then S is linearly independent.

Proof. Let $x_1, \dots, x_n \in S$ and $c_1, \dots, c_n \in \mathbb{C}$. Suppose that $\sum_{j=1}^n c_j x_j = 0$. Since $(c_j x_j)_{j=1}^n$ is orthogonal, the Pythagorean theorem implies that

$$\begin{aligned} 0 &= \left\| \sum_{i=1}^n c_i x_i \right\|^2 \\ &= \sum_{j=1}^n |c_j|^2 \|x_j\|^2 \end{aligned}$$

So for each $j \in \{1, \dots, n\}$, $c_j = 0$ and S is linearly independent. □

Definition 10.2.1.23. Let H be an inner product space and $S \subset H$. Then S is said to be **orthonormal** if S is orthogonal and for each $x \in S$, $\|x\| = 1$.

Note 10.2.1.24. If S is orthonormal, then for each $u, v \in S$, $\langle u, v \rangle = \delta_{u,v}$.

Exercise 10.2.1.25. Bessel's Inequality:

Let H be an inner product space and $S \subset H$. If S is orthonormal, then for each $x \in H$,

1. $\sum_{u \in S} |\langle u, x \rangle|^2 \leq \|x\|^2$
2. $\{u \in S : \langle u, x \rangle \neq 0\}$ is countable.

Proof.

1. Suppose that S is orthonormal. Let $x \in H$. We consider the measure space $(S, \mathcal{P}(S), \#)$. Define $f_x : S \rightarrow [0, \infty)$ by $f_x(u) = \langle u, x \rangle$. Basic results about counting measure imply that

$$\begin{aligned} \sum_{u \in S} |\langle u, x \rangle|^2 &= \int |f_x|^2 d\# \\ &= \sup \left\{ \sum_{u \in F} |f_x(u)|^2 : F \subset S \text{ and } \#(F) < \infty \right\} \end{aligned}$$

Let $F \subset S$ finite. Then the Pythagorean theorem (Exercise 10.2.1.21) implies that

$$\begin{aligned} 0 &\leq \left\| x - \sum_{u \in F} \langle u, x \rangle u \right\|^2 \\ &= \|x\|^2 + \left\| \sum_{u \in F} \langle u, x \rangle u \right\|^2 - 2 \operatorname{Re} \left\langle x, \sum_{u \in F} \langle u, x \rangle u \right\rangle \\ &= \|x\|^2 + \sum_{u \in F} |\langle u, x \rangle|^2 \|u\|^2 - 2 \sum_{u \in F} |\langle u, x \rangle|^2 \\ &= \|x\|^2 - \sum_{u \in F} |\langle u, x \rangle|^2 \end{aligned}$$

Thus

$$\sum_{u \in F} |\langle u, x \rangle|^2 \leq \|x\|^2$$

Since $F \subset S$ such that $\#(F) < \infty$ was arbitrary, we have that

$$\begin{aligned} \sum_{u \in S} |\langle u, x \rangle|^2 &= \sup \left\{ \sum_{u \in F} |\langle u, x \rangle|^2 : F \subset S \text{ and } \#F < \infty \right\} \\ &\leq \|x\|^2. \end{aligned}$$

2. Since

$$\int |f_x|^2 d\# < \infty$$

basic results about counting measure imply that $\{u \in S : \langle u, x \rangle \neq 0\}$ is countable.

□

Definition 10.2.1.26. Let $(H, \langle \cdot, \cdot \rangle)$ be an inner product space. Then $(H, \langle \cdot, \cdot \rangle)$ is said to be a **Hilbert space** if $(H, \|\cdot\|)$ is a Banach space.

Note 10.2.1.27. We recall from Exercise 9.1.0.4 that if H is a Hilbert space, then for each $(x_j)_{j \in \mathbb{N}} \subset H$, $\sum_{j \in \mathbb{N}} \|x_j\| < \infty$ implies that

$\sum_{j \in \mathbb{N}} x_j$ exists.

Exercise 10.2.1.28. Let H be a Hilbert space and $S \subset H$. Suppose that S is orthonormal and S is countable.

1. Let $(a_u)_{u \in S} \subset \mathbb{K}$. If $\sum_{u \in S} |a_u|^2 < \infty$, then

(a) $\sum_{u \in S} a_u u$ exists,

(b) $\left\| \sum_{u \in S} a_u u \right\|^2 = \sum_{u \in S} |a_u|^2$.

2. Let $(a_u)_{u \in S}, (b_u)_{u \in S} \subset \mathbb{K}$. If $\sum_{u \in S} |a_u|^2 < \infty$ and $\sum_{u \in S} |b_u|^2 < \infty$, then

(a) $\sum_{u \in S} \sum_{v \in S} a_u^* b_v$ exists,

(b) $\langle \sum_{u \in S} a_u u, \sum_{v \in S} b_v v \rangle = \sum_{u \in S} \sum_{v \in S} a_u^* b_v$.

Proof.

1. Suppose that $\sum_{u \in S} |a_u|^2 < \infty$.

(a) • If $\#S < \infty$, then clearly $\sum_{u \in S} a_u u$ exists.

• Suppose that $\#S = \infty$. Then there exists $(u_j)_{j \in \mathbb{N}} \subset H$ such that $S = (u_j)_{j \in \mathbb{N}}$. Define $(y_n)_{n \in \mathbb{N}} \subset H$ by

$$y_n := \sum_{j \in [n]} a_j u_j.$$

Let $\epsilon > 0$. Since $\sum_{j \in \mathbb{N}} |a_j|^2 < \infty$, $\left(\sum_{j \in [n]} |a_j|^2 \right)_{n \in \mathbb{N}}$ is Cauchy. Thus for each $\epsilon > 0$, there exists $N \in \mathbb{N}$ such that for each $m, n \in \mathbb{N}$, $m, n \geq N$ implies that if $m < n$, then

$$\sum_{m+1}^n |a_j|^2 < \epsilon^2$$

Then for each $m, n \in \mathbb{N}$, $m, n \geq N$ implies that if $m < n$, then

$$\begin{aligned} \|y_n - y_m\|^2 &= \left\| \sum_{j \in [n]} a_j u_j - \sum_{j \in [m]} a_j u_j \right\|^2 \\ &= \left\| \sum_{m+1}^n a_j u_j \right\|^2 \\ &= \sum_{m+1}^n |a_j|^2 \\ &< \epsilon^2 \end{aligned}$$

and $\|y_n - y_m\| < \epsilon$. Since $\epsilon > 0$ is arbitrary, we have that for each $\epsilon > 0$, there exists $N \in \mathbb{N}$ such that for each $m, n \in \mathbb{N}$, $m, n \geq N$ implies that $\|y_n - y_m\| < \epsilon$. Hence $(y_n)_{n \in \mathbb{N}}$ is Cauchy. Since H is complete, there exists $y \in H$ such that $y_n \rightarrow y$. By definition,

$$y = \sum_{j \in \mathbb{N}} a_j u_j \quad (\text{maybe refer to summations section on summing of vector functions}).$$

(b) • If $\#S < \infty$, then the Pythagorean theorem (Exercise 10.2.1.21) implies that

$$\begin{aligned} \left\| \sum_{u \in S} a_u u \right\|^2 &= \sum_{u \in S} \|a_u u\|^2 \quad (\text{since } (a_u u)_{u \in S} \text{ is orthogonal}) \\ &= \sum_{u \in S} |a_u|^2. \end{aligned}$$

- Suppose that $\#S = \infty$. Write $S = (u_j)_{j \in \mathbb{N}}$ and define $(y_n)_{n \in \mathbb{N}} \subset H$ and $y \in H$ as in the prev part. Continuity of $\langle \cdot, \cdot \rangle : H \times H \rightarrow \mathbb{C}$ implies that

$$\begin{aligned}
\left\| \sum_{u \in S} a_u u \right\|^2 &= \left\| \sum_{j \in \mathbb{N}} a_j u_j \right\|^2 \\
&= \|y\|^2 \\
&= \langle y, y \rangle \\
&= \lim_{m \rightarrow \infty} \langle y_m, y \rangle \\
&= \lim_{m \rightarrow \infty} \left[\lim_{n \rightarrow \infty} \langle y_m, y_n \rangle \right] \\
&= \lim_{m \rightarrow \infty} \left[\lim_{n \rightarrow \infty} \left\langle \sum_{j \in [m]} a_j u_j, \sum_{k \in [n]} a_k u_k \right\rangle \right] \\
&= \lim_{m \rightarrow \infty} \left[\lim_{n \rightarrow \infty} \left(\sum_{j \in [m]} \sum_{k \in [n]} a_j^* a_k \langle u_j, u_k \rangle \right) \right] \\
&= \lim_{m \rightarrow \infty} \left[\lim_{n \rightarrow \infty} \left(\sum_{j \in [m]} \sum_{k \in [n]} a_j^* a_k \delta_{j,k} \right) \right] \\
&= \lim_{m \rightarrow \infty} \left[\lim_{n \rightarrow \infty} \left(\sum_{j \in [m]} |a_j|^2 \right) \right] \\
&= \lim_{m \rightarrow \infty} \left[\sum_{j \in [m]} |a_j|^2 \right] \\
&= \sum_{j \in \mathbb{N}} |a_j|^2 \\
&= \sum_{u \in S} |a_u|^2
\end{aligned}$$

2. Suppose that $\sum_{u \in S} |a_u|^2 < \infty$ and $\sum_{u \in S} |b_u|^2 < \infty$.

- (a) make ex on $\mathbb{K}^{\mathbb{N}}$ being a Hilbert space, use cauchy-schwarz or sums or something
- (b) use continuity of $\langle \cdot, \cdot \rangle$

□

Exercise 10.2.1.29. Let H be a Hilbert space and $S \subset H$. Suppose that S is orthonormal. Then the following are equivalent:

1. For each $x \in H$, if for each $u \in S$, $\langle u, x \rangle = 0$, then $x = 0$.
2. For each $x \in H$, there exists $S_* \subset S$ such that S_* is countable, $x = \sum_{u \in S_*} \langle u, x \rangle u$, for each $u \in S_*$, $\langle u, x \rangle \neq 0$ and for each $u \in S \setminus S_*$, $\langle u, x \rangle = 0$.
3. For each $x \in H$, $\|x\|^2 = \sum_{u \in S} |\langle u, x \rangle|^2$.

Proof.

- (1) $\implies$ (2):

(maybe make a portion of this part, like the finite assumption of S_* , into another exercise and just ref it here) Suppose that for each $x \in H$, if for each $u \in S$, $\langle u, x \rangle = 0$, then $x = 0$. Let $x \in H$. Set $S_* := \{u \in S : \langle u, x \rangle \neq 0\}$. The previous exercise implies that S_* is countable.

– Suppose that $\#S_* < \infty$.

* Suppose that $\#S_* = 0$, then for each $u \in S$, $\langle u, x \rangle = 0$. By assumption,

$$\begin{aligned} x &= 0 \\ &= \sum_{u \in S_*} \langle u, x \rangle u. \end{aligned}$$

* Suppose that $\#S_* \neq 0$. Then there exists $n \in \mathbb{N}$ and $(u_j)_{j \in [n]} \subset S$ such that $S_* = (u_j)_{j \in [n]}$. Define $x' \in H$ by $x' := \sum_{j \in [n]} \langle u_j, x \rangle u_j$. Let $u \in S$. Then

$$\begin{aligned} \langle u, x - x' \rangle &= \left\langle u, x - \sum_{j \in [n]} \langle u_j, x \rangle u_j \right\rangle \\ &= \langle u, x \rangle - \left\langle u, \sum_{j \in [n]} \langle u_j, x \rangle u_j \right\rangle \\ &= \langle u, x \rangle - \sum_{j \in [n]} \langle u_j, x \rangle \langle u, u_j \rangle \end{aligned}$$

· If $u \in S_*$, then there exists $j_0 \in [n]$ such that $u = u_{j_0}$ and

$$\begin{aligned} \langle u, x - x' \rangle &= \langle u, x \rangle - \sum_{j \in [n]} \langle u_j, x \rangle \langle u, u_j \rangle \\ &= \langle u_{j_0}, x \rangle - \sum_{j \in [n]} \langle u_{j_0}, x \rangle \langle u_{j_0}, u_j \rangle \\ &= \langle u_{j_0}, x \rangle - \sum_{j \in [n]} \langle u_{j_0}, x \rangle \delta_{j_0, j} \\ &= \langle u_{j_0}, x \rangle - \langle u_{j_0}, x \rangle \\ &= 0. \end{aligned}$$

· If $u \in S \setminus S_*$, then

$$\begin{aligned} \langle u, x - x' \rangle &= \langle u, x \rangle - \sum_{j \in [n]} \langle u_j, x \rangle \langle u, u_j \rangle \\ &= 0 - \sum_{j \in [n]} \langle u_j, x \rangle 0 \\ &= 0. \end{aligned}$$

Since $u \in S$ is arbitrary, we have that for each $u \in S$, $\langle u, x - x' \rangle = 0$. By assumption, $x - x' = 0$. Hence

$$\begin{aligned} x &= x' \\ &= \sum_{j \in [n]} \langle u_j, x \rangle u_j \\ &= \sum_{u \in S_*} \langle u, x \rangle u. \end{aligned}$$

– Suppose that $\#S_* = \infty$. Then there exist $(u_j)_{j \in \mathbb{N}} \subset S$ such that $S_* = (u_j)_{j \in \mathbb{N}}$. Exercise 10.2.1.25 (Bessel's inequality) implies that $\sum_{j \in \mathbb{N}} |\langle u_j, x \rangle|^2 \leq \|x\|^2$. Exercise 10.2.1.28 implies that $\sum_{j \in \mathbb{N}} \langle u_j, x \rangle u_j$ exists. Define $(y_n)_{n \in \mathbb{N}} \subset H$ and $y \in H$ by $y_n := y_n \sum_{j \in [n]} \langle u_j, x \rangle u_j$ and $y := \sum_{j \in \mathbb{N}} \langle u_j, x \rangle u_j$. Since $y_n \rightarrow y$, y is Cauchy. Continuity of $\langle \cdot, \cdot \rangle : H \times H \rightarrow \mathbb{C}$ implies that

* for each $u \in S \setminus S_*$,

$$\begin{aligned}
 \langle u, x - y \rangle &= \langle u, x \rangle - \langle u, y \rangle \\
 &= \langle u, x \rangle - \lim_{n \rightarrow \infty} \langle u, y_n \rangle \\
 &= \langle u, x \rangle - \lim_{n \rightarrow \infty} \sum_{j \in [n]} \langle u_j, x \rangle \langle u, u_j \rangle \\
 &= 0 - 0 \\
 &= 0
 \end{aligned}$$

* for each $k \in \mathbb{N}$,

$$\begin{aligned}
 \langle u_k, x - y \rangle &= \langle u_k, x \rangle - \langle u_k, y \rangle \\
 &= \langle u_k, x \rangle - \lim_{n \rightarrow \infty} \langle u_k, y_n \rangle \\
 &= \langle u_k, x \rangle - \lim_{n \rightarrow \infty} \sum_{j=1}^n \langle u_j, x \rangle \langle u_k, u_j \rangle \\
 &= \langle u_k, x \rangle - \langle u_k, x \rangle \\
 &= 0
 \end{aligned}$$

So for each $u \in S$, $\langle u, x - y \rangle = 0$. By assumption, $x - y = 0$ and hence

$$\begin{aligned}
 x &= y \\
 &= \sum_{j \in \mathbb{N}} \langle u_j, x \rangle u_j \\
 &= \sum_{u \in S_*} \langle u, x \rangle u.
 \end{aligned}$$

• (2) $\implies$ (3):

Suppose that for each $x \in H$, there exists $S_* \subset S$ such that S_* is countable, $x = \sum_{u \in S_*} \langle u, x \rangle u$, for each $u \in S_*$, $\langle u, x \rangle \neq 0$ and for each $u \in S \setminus S_*$, $\langle u, x \rangle = 0$. Let $x \in H$. By assumption there exist $S_* \subset S$ such that S_* is countable, $x = \sum_{u \in S_*} \langle u, x \rangle u$, for each $u \in S_*$, $\langle u, x \rangle \neq 0$ and for each $u \in S \setminus S_*$, $\langle u, x \rangle = 0$. Exercise 10.2.1.21 then implies that

$$\begin{aligned}
 \|x\|^2 &= \left\| \sum_{u \in S_*} \langle u, x \rangle u \right\|^2 \\
 &= \sum_{u \in S_*} \|\langle u, x \rangle u\|^2 \\
 &= \sum_{u \in S_*} |\langle u, x \rangle|^2 \|u\|^2 \\
 &= \sum_{u \in S_*} |\langle u, x \rangle|^2 \\
 &= \sum_{u \in S_*} |\langle u, x \rangle|^2 + 0 \\
 &= \sum_{u \in S_*} |\langle u, x \rangle|^2 + \sum_{u \in S \setminus S_*} |\langle u, x \rangle|^2 \\
 &= \sum_{u \in S} |\langle u, x \rangle|^2.
 \end{aligned}$$

• (3) $\implies$ (1):

Suppose that for each $x \in H$, $\|x\|^2 = \sum_{u \in S} |\langle u, x \rangle|^2$. Let $x \in H$. Suppose that for each $u \in S$, $\langle u, x \rangle = 0$. Then

$$\begin{aligned}\|x\|^2 &= \sum_{u \in S} |\langle u, x \rangle|^2 \\ &= 0\end{aligned}$$

Thus $x = 0$.

□

Definition 10.2.1.30. Let H be a Hilbert space and $S \subset H$. Then S is said to be an **orthonormal basis for H** if

1. S is orthonormal
2. for each $x \in H$, if for each $u \in S$, $\langle u, x \rangle = 0$, then $x = 0$

Exercise 10.2.1.31. Let H be a Hilbert space and $(u_j)_{j \in \mathbb{N}} \subset H$. If $(u_j)_{j \in \mathbb{N}}$ is an orthonormal basis for H , then for each $x \in H$,

$$x = \sum_{j \in \mathbb{N}} \langle u_j, x \rangle u_j.$$

Proof. Suppose that $(u_j)_{j \in \mathbb{N}}$ is an orthonormal basis for H . Let $x \in H$. Exercise 10.2.1.29 implies that there exists $S_* \subset S$ such that S_* is countable, $x = \sum_{u \in S_*} \langle u, x \rangle u$, for each $u \in S_*$, $\langle u, x \rangle \neq 0$ and for each $u \in S \setminus S_*$, $\langle u, x \rangle = 0$. Thus there $J \subset \mathbb{N}$ such that $S_* = (u_j)_{j \in J}$. Therefore

$$\begin{aligned}x &= \sum_{u \in S_*} \langle u, x \rangle u \\ &= \sum_{j \in J} \langle u_j, x \rangle u_j \\ &= \sum_{j \in J} \langle u_j, x \rangle u_j + 0 \\ &= \sum_{j \in J} \langle u_j, x \rangle u_j + \sum_{j \in \mathbb{N} \setminus J} \langle u_j, x \rangle u_j \\ &= \sum_{j \in \mathbb{N}} \langle u_j, x \rangle u_j.\end{aligned}$$

□

Exercise 10.2.1.32. Let H be a Hilbert space. Then there exists $S \subset H$ such that S is an orthonormal basis for H .

Proof. **ZORN's Lemma**

FINSH!!!!

□

Exercise 10.2.1.33. Let H be a Hilbert space. Then the following are equivalent:

1. for each $S \subset H$, S is an orthonormal basis for H implies that S is countable,
2. there exists $S \subset H$ such that S is an orthonormal basis for H and S is countable,
3. H is separable.

Proof.

- (1) $\implies$ (2):

Suppose that for each $S \subset H$, S is an orthonormal basis for H implies that S is countable. Exercise 10.2.1.32 implies that there exists $S \subset H$ such that S is an orthonormal basis for H . By assumption, S is countable. Thus there exists $S \subset H$ such that S is an orthonormal basis for H and S is countable.

• (2) $\implies$ (3):

Suppose that there exists $S \subset H$ such that S is an orthonormal basis for H and S is countable.

- Since $\mathbb{K}$ is separable, there exists $S_{\mathbb{K}} \subset \mathbb{K}$ such that $S_{\mathbb{K}}$ is dense in $\mathbb{K}$ and $S_{\mathbb{K}}$ is countable. Define $S_H \subset H$ by

$$S_H := \left\{ \sum_{\lambda \in F} \sum_{u \in E} \lambda u : F \subset S_{\mathbb{K}}, E \subset S, \#F < \infty \text{ and } \#E < \infty \right\}.$$

Then S_H is countable.

- Let $x \in H$ and $\epsilon > 0$. Exercise 10.2.1.29 implies that there exists $S_* \subset S$ such that $x = \sum_{u \in S_*} \langle u, x \rangle u$. Then there exists

$N \in \mathbb{N}$ and $(u_j)_{j \in [N]} \subset S_*$ such that $\left\| x - \sum_{j \in [N]} \langle u_j, x \rangle u_j \right\| < \epsilon/2$. Define $x_0 \in H$ by $x_0 := \sum_{j \in [N]} \langle u_j, x \rangle u_j$. Since $S_{\mathbb{K}}$ is dense in $\mathbb{K}$, for each $j \in [N]$, there exists $\lambda_j \in S_{\mathbb{K}}$ such that $|\lambda_j - \langle u_j, x \rangle| < \epsilon 2^{-(j+1)}$.

- Define $x' \in S_H$ by $x' := \sum_{j \in [N]} \lambda_j u_j$. Then

$$\begin{aligned} \|x - x'\| &\leq \|x - x_0\| + \|x_0 - x'\| \\ &= \|x - x_0\| + \left\| \sum_{j \in [N]} \langle u_j, x \rangle u_j - \sum_{j \in [N]} \lambda_j u_j \right\| \\ &< \|x - x_0\| + \left\| \sum_{j \in [N]} (\langle u_j, x \rangle - \lambda_j) u_j \right\| \\ &\leq \|x - x_0\| + \sum_{j \in [N]} |\langle u_j, x \rangle - \lambda_j| \|u_j\| \\ &= \|x - x_0\| + \sum_{j \in [N]} |\langle u_j, x \rangle - \lambda_j| \\ &< \frac{\epsilon}{2} + \sum_{j \in [N]} \frac{\epsilon}{2^{j+1}} \\ &< \frac{\epsilon}{2} + \frac{\epsilon}{2} \\ &= \epsilon. \end{aligned}$$

Since $x \in H$ and $\epsilon > 0$ are arbitrary, we have that for each $x \in H$ and $\epsilon > 0$, there exists $x' \in S_H$ such that $\|x - x'\| < \epsilon$. Thus S_H is dense in H .

- Since S_H is dense in H and S_H is countable, H is separable.

• (3) $\implies$ (1):

- Suppose that H is separable. Let $S \subset H$. Suppose that S is an orthonormal basis for H . Since H is separable, there exists $(x_n)_{n \in \mathbb{N}} \subset H$ such that $(x_n)_{n \in \mathbb{N}}$ is dense in H . Exercise 10.2.1.29 implies that for each $n \in \mathbb{N}$, there exists $S_n \subset S$ such that S_n is countable, $x = \sum_{u \in S_n} \langle u, x \rangle u$, for each $u \in S_n$, $\langle u, x \rangle \neq 0$ and for each $u \in S \setminus S_n$, $\langle u, x \rangle = 0$. Define $S' \subset S$

by $S' := \bigcup_{n \in \mathbb{N}} S_n$. Then S' is countable. We note that for each $n \in \mathbb{N}$,

$$x_n = \sum_{u \in S_n} \langle u, x_n \rangle u \tag{10.1}$$

$$= \sum_{u \in S_n} \langle u, x_n \rangle u + 0 \tag{10.2}$$

$$= \sum_{u \in S_n} \langle u, x_n \rangle u + \sum_{u \in S' \setminus S_n} \langle u, x_n \rangle u \tag{10.3}$$

$$= \sum_{u \in S'} \langle u, x_n \rangle u. \tag{10.4}$$

- Let $x \in X$. Exercise 10.2.1.29 implies that there exists $S_* \subset S$ such that S_* is countable, $x = \sum_{u \in S_*} \langle u, x \rangle u$, for each $u \in S_*$, $\langle u, x \rangle \neq 0$ and for each $u \in S \setminus S_*$, $\langle u, x \rangle = 0$. Define $x' \in H$ by $x' := \sum_{u \in S'} \langle u, x \rangle u$. For the sake of contradiction, suppose that $S_* \not\subset S'$. Then there exists $u_0 \in S_* \setminus S'$. Define $\epsilon > 0$ by $\epsilon := |\langle u_0, x \rangle|$. Since

$$\begin{aligned} \|x - x'\| &= \left\| \sum_{u \in S_*} \langle u, x \rangle u - \sum_{u \in S'} \langle u, x \rangle u \right\| \\ &= \left\| \left(\sum_{u \in S_* \setminus S'} \langle u, x \rangle u + \sum_{u \in S_* \cap S'} \langle u, x \rangle u \right) - \left(\sum_{u \in S' \setminus S_*} \langle u, x \rangle u + \sum_{u \in S_* \cap S'} \langle u, x \rangle u \right) \right\| \\ &= \left\| \sum_{u \in S_* \setminus S'} \langle u, x \rangle u - \sum_{u \in S' \setminus S_*} \langle u, x \rangle u \right\|, \end{aligned}$$

we have that

$$\begin{aligned} \|x - x'\|^2 &= \left\| \sum_{u \in S_* \setminus S'} \langle u, x \rangle u - \sum_{u \in S' \setminus S_*} \langle u, x \rangle u \right\|^2 \\ &= \sum_{u \in S_* \setminus S'} |\langle u, x \rangle|^2 + \sum_{u \in S' \setminus S_*} |\langle u, x \rangle|^2 \\ &\geq \sum_{u \in S_* \setminus S'} |\langle u, x \rangle|^2 \\ &\geq |\langle u_0, x \rangle|^2. \end{aligned}$$

Thus

$$\begin{aligned} \|x - x'\| &\geq |\langle u_0, x \rangle| \\ &= \epsilon. \end{aligned}$$

Since $(x_n)_{n \in \mathbb{N}}$ is dense in H , there exists $N \in \mathbb{N}$ such that $\|x - x_N\| < \epsilon/2$. Since

$$\begin{aligned} \sum_{u \in S'} \|\langle u, x_N - x \rangle u\|^2 &= \sum_{u \in S'} |\langle u, x_N - x \rangle|^2 \\ &\leq \|x_N - x\|^2 \quad (\text{by Bessel's inequality Exercise 10.2.1.25}) \\ &< \infty, \end{aligned}$$

Exercise 10.2.1.28 implies that $\sum_{u \in S'} \langle u, x_N - x \rangle u$ exists and

$$\begin{aligned} \left\| \sum_{u \in S'} \langle u, x_N - x \rangle u \right\|^2 &= \sum_{u \in S'} |\langle u, x_N - x \rangle|^2 \\ &\leq \|x_N - x\|^2. \end{aligned}$$

Therefore

$$\begin{aligned}
\|x - x'\| &\leq \|x - x_N\| + \|x_N - x'\| \\
&= \|x - x_N\| + \left\| \sum_{u \in S_N} \langle u, x_N \rangle u - \sum_{u \in S'} \langle u, x \rangle u \right\| \\
&= \|x - x_N\| + \left\| \sum_{u \in S'} \langle u, x_N \rangle u - \sum_{u \in S'} \langle u, x \rangle u \right\| \quad (\text{by the note above}) \\
&= \|x - x_N\| + \left\| \sum_{u \in S'} (\langle u, x_N \rangle - \langle u, x \rangle) u \right\| \\
&= \|x - x_N\| + \left\| \sum_{u \in S'} \langle u, x_N - x \rangle u \right\| \\
&\leq \|x - x_N\| + \left\| \sum_{u \in S'} \langle u, x_N - x \rangle u \right\| \\
&\leq \|x - x_N\| + \|x_N - x\| \\
&< \frac{\epsilon}{2} + \frac{\epsilon}{2} \\
&= \epsilon.
\end{aligned}$$

Hence

$$\begin{aligned}
\epsilon &\leq \|x - x'\| \\
&< \epsilon,
\end{aligned}$$

which is a contradiction. Therefore $S_* \subset S'$. Since $x \in X$ is arbitrary, we have that for each $x \in H$, there exists $S_* \subset S'$ such that S_* is countable, $x = \sum_{u \in S_*} \langle u, x \rangle u$, for each $u \in S_*$, $\langle u, x \rangle \neq 0$ and for each $u \in S \setminus S_*$, $\langle u, x \rangle = 0$. Exercise

10.2.1.29 then implies that S' is an orthonormal basis for H .

- For the sake of contradiction, suppose that $S \not\subset S'$. Then there exists $u_0 \in S \setminus S'$. Since S' is an orthonormal basis for H , Exercise 10.2.1.29 implies that

$$\begin{aligned}
1 &= \|u_0\|^2 \\
&= \sum_{u \in S'} |\langle u_0, u \rangle|^2 \\
&= \sum_{u \in S'} \delta_{u, u_0} \quad (\text{since } S' \subset S) \\
&= 0 \quad (\text{since } u_0 \in S \setminus S'),
\end{aligned}$$

which is a contradiction. Hence

$$\begin{aligned}
S' &\subset S \\
&\subset S'.
\end{aligned}$$

Thus $S = S'$ and S is countable. Since $S \subset H$ with S an orthonormal basis for H is arbitrary, we have that for each $S \subset H$, S is an orthonormal basis for H implies that S is countable. □

Exercise 10.2.1.34. Let H be a Hilbert space. Suppose that H is separable. Let $(u_j)_{j \in \mathbb{N}}$ be an orthonormal basis for H . Then for each $x_1, x_2 \in H$,

$$\langle x_1, x_2 \rangle = \sum_{j \in \mathbb{N}} \langle x_1, u_j \rangle \langle u_j, x_2 \rangle.$$

Proof. Let $x_1, x_2 \in H$. Since $(u_j)_{j \in \mathbb{N}}$ is an orthonormal basis for H , Exercise 10.2.1.29 implies that $x_1 = \sum_{j \in \mathbb{N}} \langle u_j, x_1 \rangle u_j$ and $x_2 = \sum_{j \in \mathbb{N}} \langle u_j, x_2 \rangle u_j$. Then □

10.2.2 Riesz Representation Theorem

Exercise 10.2.2.1. Let H be a Hilbert space, $E \subset H$ a closed subspace of H , $x \in H$, $y \in E$ and $z \in E^\perp$. If $x = y + z$, then $\|x + E\|_{H/E} = \|z\|$.

Proof. Suppose that $x = y + z$. Let $y' \in E$. The Pythagorean theorem implies that

$$\begin{aligned}\|x - y'\|^2 &= \|y + z - y'\|^2 \\ &= \|y - y'\|^2 + \|z\|^2 \\ &\geq \|z\|^2\end{aligned}$$

Thus

$$\begin{aligned}\|x + E\| &= \inf_{y' \in E} \|x - y'\| \\ &\geq \|z\|\end{aligned}$$

Also,

$$\begin{aligned}\|x + E\| &\leq \|x - y\| \\ &= \|z\|\end{aligned}$$

Hence $\|x + E\| = \|z\|$. □

Exercise 10.2.2.2. Let H be a Hilbert space, $E \subset H$ a closed subspace of H and $x \in H$. Then

1. there exists a unique $y_0 \in E$ such that $\|x - y_0\| = \|x + E\|$

Hint: Suppose $(y_n)_{n \in \mathbb{N}} \subset E$ satisfies $\|x - y_n\| \rightarrow \|x + E\|$. Show that $(y_n)_{n \in \mathbb{N}}$ is Cauchy using the parallelogram law.

2. there exist unique $y_0 \in E$ and $z_0 \in E^\perp$ such that $x = y_0 + z_0$ and $\|z_0\| = \|x + E\|$

Hint: Set $z_0 := x - y_0$ and for $u \in E$, choose $\lambda \in \mathbb{C}$ such that $\langle x, \lambda u \rangle \in \mathbb{R}$. Consider $f(t) = \|z - t\lambda u\|^2$.

Proof. Set $s := \|x + E\|$.

1. • **(Existence):**

Choose $(y_n)_{n \in \mathbb{N}} \subset E$ such that for each $n \in \mathbb{N}$, $\|x - y_n\| < s + 1/n$. Define $(a_n)_{n \in \mathbb{N}} \subset H$ by $a_n = x - y_n$. Let $m, n \in \mathbb{N}$. The parallelogram law implies that

$$\begin{aligned}2(\|x - y_n\|^2 + \|x - y_m\|^2) &= 2(\|a_n\|^2 + \|a_m\|^2) \\ &= \|a_n + a_m\|^2 + \|a_n - a_m\|^2 \\ &= \|2x - (y_n + y_m)\|^2 + \|y_m - y_n\|^2 \\ &= 4\|x - 2^{-1}(y_n + y_m)\|^2 + \|y_m - y_n\|^2\end{aligned}$$

Since $y_n, y_m \in E$, $2^{-1}(y_n + y_m) \in E$ and therefore $\|x - 2^{-1}(y_n + y_m)\| \geq s$. Since $m, n \in \mathbb{N}$ are arbitrary, we have that for each $m, n \in \mathbb{N}$,

$$\begin{aligned}\|y_m - y_n\|^2 &= 2(\|x - y_n\|^2 + \|x - y_m\|^2) - 4\|x - 2^{-1}(y_n + y_m)\|^2 \\ &\leq 2(\|x - y_n\|^2 + \|x - y_m\|^2) - 4s^2\end{aligned}$$

Let $\epsilon > 0$. Choose $N \in \mathbb{N}$ such that $1/N < (s^2 + \epsilon^2/4)^{1/2} - s$. Let $m, n \in \mathbb{N}$. Suppose that $m, n \geq N$. Then

$$\begin{aligned}\|y_m - y_n\|^2 &= 2\|x - y_n\|^2 + 2\|x - y_m\|^2 - 4s^2 \\ &< 2(s + 1/n)^2 + 2(s + 1/m)^2 - 4s^2 \\ &< 2(s + 1/N)^2 + 2(s + 1/N)^2 - 4s^2 \\ &= 4(s + 1/N)^2 - 4s^2 \\ &< 4(s^2 + \epsilon^2/4) - 4s^2 \\ &= \epsilon^2\end{aligned}$$

Thus $\|y_m - y_n\| < \epsilon$. Since $\epsilon > 0$ is arbitrary, we have that for each $\epsilon > 0$, there exists $N \in \mathbb{N}$ such that for each $m, n \in \mathbb{N}$, $m, n \geq N$ implies that $\|y_m - y_n\| < \epsilon$. So $(y_n)_{n \in \mathbb{N}}$ is Cauchy and since H is complete, there exists $y_0 \in H$ such that $y_n \rightarrow y_0$. Continuity of the inner product, addition and scalar multiplication implies that

$$\begin{aligned}\|x - y_0\| &= \lim_{n \rightarrow \infty} \|x - y_n\| \\ &= s \\ &= \|x + E\|\end{aligned}$$

• **(Uniqueness):**

Let $y_1 \in E$. Suppose that $\|x - y_1\| = \|x + E\|$. Similarly to part (1), the parallelogram law implies that

$$\begin{aligned}\|y_1 - y_0\|^2 &\leq 2(\|x - y_1\|^2 + \|x - y_0\|^2) - 4s^2 \\ &= 2(s^2 + s^2) - 4s^2 \\ &= 0\end{aligned}$$

Hence $\|y_1 - y_0\| = 0$ and $y_1 = y_0$.

2. • **(Existence):**

Set $z_0 := x - y_0$. Let $u \in E$. Set

$$\lambda := \begin{cases} (\operatorname{sgn}\langle z_0, u \rangle)^{-1} & \langle z_0, u \rangle \neq 0 \\ 1 & \langle z_0, u \rangle = 0 \end{cases}$$

and $v := \lambda u$. So $\langle z_0, v \rangle \in \mathbb{R}$. Define $f : \mathbb{R} \rightarrow \mathbb{R}$ by $f(t) = \|z_0 - tv\|^2$. By construction, since $y_0, v \in E$, we have that for each $t \in \mathbb{R}$, $y_0 - tv \in E$ and therefore

$$\begin{aligned}f(t) &= \|z_0 - tv\|^2 \\ &= \|x - (y_0 - tv)\|^2 \\ &\geq \inf_{y \in E} \|x - y\|^2 \\ &= \|x - y_0\|^2 \\ &= f(0)\end{aligned}$$

Since f is smooth and has a local minimum at $t = 0$, $f'(0) = 0$. Furthermore, for each $t \in \mathbb{R}$,

$$\begin{aligned}f(t) &= \|z_0\|^2 - 2t \operatorname{Re}(\langle z_0, v \rangle) + t^2 \|v\|^2 \\ &= \|z_0\|^2 - 2t \langle z_0, v \rangle + t^2 \|v\|^2\end{aligned}$$

so that $f'(0) = 2\langle z_0, v \rangle$. Thus

$$\begin{aligned}\langle z_0, u \rangle &= \lambda^{-1} \langle z_0, \lambda u \rangle \\ &= \lambda^{-1} \langle z_0, v \rangle \\ &= 0\end{aligned}$$

Since $u \in E$ is arbitrary, we have that $z_0 \in E^\perp$. **(Uniqueness):**

Suppose that there exist $y_1 \in E$ and $z_1 \in E^\perp$ such that $x = y_1 + z_1$ and $\|z_1\| = \|x + E\|$. Since $z_1 = x - y_1$, by assumption,

$$\begin{aligned}\|x - y_1\| &= \|z_1\| \\ &= \|x + E\|\end{aligned}$$

Since $y_1 \in E$, uniqueness in part (1) implies that $y_1 = y_0$. Hence

$$\begin{aligned}z_1 &= x - y_1 \\ &= x - y_0 \\ &= z_0\end{aligned}$$

□

Exercise 10.2.2.3. Let H be a Hilbert space and $E \subset H$ a closed subspace of H . Then $(E^\perp)^\perp = E$.

Proof.

- Let $x \in (E^\perp)^\perp$ and $x_0 \in H$. **The previous exercise** implies that there exist unique $y, y_0 \in E$ and $z, z_0 \in E^\perp$ such that $x = y + z$, $x_0 = y_0 + z_0$, $\|x\| = \|y + E\|$ and $\|z_0\| = \|x + E\|$. Since $x \in (E^\perp)^\perp$, $z_0 \in E^\perp$ and $y_0 \in E$, we have that

$$\begin{aligned}\langle x, x_0 \rangle &= \langle x, y_0 + z_0 \rangle \\ &= \langle x, y_0 \rangle + \langle x, z_0 \rangle \\ &= \langle x, y_0 \rangle \\ &= \langle y + z, y_0 \rangle \\ &= \langle y, y_0 \rangle + \langle z, y_0 \rangle \\ &= \langle y, y_0 \rangle\end{aligned}$$

Similarly, since $y \in E$ and $z_0 \in E^\perp$, we have that

$$\begin{aligned}\langle y, x_0 \rangle &= \langle y, y_0 + z_0 \rangle \\ &= \langle y, y_0 \rangle + \langle y, z_0 \rangle \\ &= \langle y, y_0 \rangle\end{aligned}$$

Therefore

$$\begin{aligned}\langle z, x_0 \rangle &= \langle x - y, x_0 \rangle \\ &= \langle x, x_0 \rangle - \langle y, x_0 \rangle \\ &= \langle y, y_0 \rangle - \langle y, y_0 \rangle \\ &= 0\end{aligned}$$

Since $x_0 \in H$ is arbitrary, **a previous exercise** implies that $z = 0$. Hence

$$\begin{aligned}x &= y + z \\ &= y \\ &\in E\end{aligned}$$

Since $x \in (E^\perp)^\perp$ is arbitrary, we have that $(E^\perp)^\perp \subset E$.

- Let $y \in E$ and $z \in E^\perp$. By definition of $E^\perp$, $\langle y, z \rangle = 0$. Since $z \in E^\perp$ is arbitrary, we have that for each $z \in E^\perp$, $\langle y, z \rangle = 0$. Hence $y \in (E^\perp)^\perp$. Since $y \in E$ is arbitrary, we have that $E \subset (E^\perp)^\perp$.

Since $(E^\perp)^\perp \subset E$ and $E \subset (E^\perp)^\perp$, we have that $(E^\perp)^\perp = E$. □

Exercise 10.2.2.4. Let H be a Hilbert space and $E, F \subset H$ closed subspaces of H . Then $E = F$ iff $E^\perp = F^\perp$.

Proof. If $E = F$, then clearly $E^\perp = F^\perp$.

Conversely, suppose that $E^\perp = F^\perp$. Then **the previous exercise** and part (1) imply that

$$\begin{aligned}E &= (E^\perp)^\perp \\ &= (F^\perp)^\perp \\ &= F\end{aligned}$$

□

Exercise 10.2.2.5. Riesz Representation Theorem:

Let H be a Hilbert space. For each $\phi \in H^*$, there exists a unique $y \in H$ such that for each $x \in H$, $\phi(x) = \langle y, x \rangle$.

Hint: If $x \notin \ker \phi$, then there exists $z \in (\ker \phi)^\perp$ such that $\|z\| = 1$. Consider $u := \phi(x)z - \phi(z)x$. Then $u \in \ker \phi$ and consider $\langle z, u \rangle$.

Proof. Let $\phi \in H^*$.

• **(Existence):**

- Suppose that $\phi = 0$. Set $y := 0$. [A previous exercise](#) implies that for each $x \in H$, $\phi(x) = \langle y, x \rangle$.
- Suppose that $\phi \neq 0$. Set $E := \ker \phi$. Since ϕ is continuous, E is a closed subspace of H . Since $\phi \neq 0$, $E \neq H$. [The previous exercise](#) then implies that $E^\perp \neq \{0\}$. Thus there exists $z \in E^\perp$ such that $\|z\| = 1$. Define $y \in H$ by $y := \phi(z)^* z$. Let $x \in H$. Define $u \in H$ by $u := \phi(x)z - \phi(z)x$. Then

$$\begin{aligned}\phi(u) &= \phi(x)\phi(z) - \phi(z)\phi(x) \\ &= 0\end{aligned}$$

Therefore $u \in E$. Since $z \in E^\perp$, we have that

$$\begin{aligned}0 &= \langle z, u \rangle \\ &= \langle z, \phi(x)z - \phi(z)x \rangle \\ &= \langle z, \phi(x)z \rangle - \langle z, \phi(z)x \rangle \\ &= \phi(x)\langle z, z \rangle - \phi(z)\langle z, x \rangle \\ &= \phi(x)\|z\|^2 - \phi(z)\langle z, x \rangle \\ &= \phi(x) - \langle \phi(z)^* z, x \rangle \\ &= \phi(x) - \langle y, x \rangle\end{aligned}$$

Since $x \in H$ is arbitrary, we have that for each $x \in H$, $\phi(x) = \langle y, x \rangle$.

• **(Uniqueness):**

Let $y' \in H$. Suppose that for each $x \in H$, $\phi(x) = \langle y', x \rangle$. Then for each $x \in H$, $\langle y - y', x \rangle = 0$. [A previous exercise](#) implies that $y - y' = 0$. Thus $y' = y$. □

Definition 10.2.2.6. Let H be a Hilbert space and $y \in H$. We define the **adjoint of y** , denoted $y^* \in H^*$, by $y^*(x) := \langle y, x \rangle$.

Exercise 10.2.2.7. Let H be a Hilbert Space. We define $\flat : \bar{H} \rightarrow H^*$ by $\flat(y) := y^*$. Then $\flat \in L(\bar{H}, H^*)$ and $\flat$ is an isometric isomorphism.

Proof.

- Let $y, z \in \bar{H}$ and $\lambda \in \mathbb{C}$. Then for each $x \in H$,

$$\begin{aligned}\flat(y + \lambda^* z)(x) &= (y + \lambda^* z)^*(x) \\ &= \langle y + \lambda^* z, x \rangle \\ &= \langle y + \lambda^* z, x \rangle \\ &= \langle y, x \rangle + \langle \lambda^* z, x \rangle \\ &= \langle y, x \rangle + \lambda \langle z, x \rangle \\ &= y^*(x) + \lambda z^*(x) \\ &= (y^* + \lambda z^*)(x) \\ &= (\flat(y) + \lambda \flat(z))(x).\end{aligned}$$

Thus $\flat(y + \lambda^* z) = \flat(y) + \lambda \flat(z)$. Since $y, z \in \bar{H}$ and $\lambda \in \mathbb{C}$ are arbitrary, we have that $\flat$ is linear.

- Let $y \in \bar{H}$. The Cauchy-Schwarz inequality implies that for each $x \in H$,

$$\begin{aligned}|\flat(y)(x)| &= |y^*(x)| \\ &= |\langle y, x \rangle| \\ &\leq \|y\| \|x\|.\end{aligned}$$

Thus $\|\flat(y)\| \leq \|y\|$. Since $y \in \bar{H}$ is arbitrary, we have that for each $y \in \bar{H}$, $\|\flat(y)\| \leq \|y\|$. Hence $\flat \in L(\bar{H}, H^*)$ and $\|\flat\| \leq 1$. We note that

$$\begin{aligned}
 \|\flat\| &= \sup_{\|y\|=1} \|\flat(y)\| \\
 &= \sup_{\|y\|=1} \|y^*\| \\
 &= \sup_{\|y\|=1} \left[\sup_{\|x\|=1} |y^*(x)| \right] \\
 &= \sup_{\|y\|=1} \left[\sup_{\|x\|=1} |\langle y, x \rangle| \right] \\
 &\geq \sup_{\|y\|=1} |\langle y, y \rangle| \\
 &= \sup_{\|y\|=1} \|y\|^2 \\
 &= 1,
 \end{aligned}$$

Therefore $\|\flat\| = 1$. Exercise 9.2.0.12 implies that $\flat$ is an isometry.

- Since $\flat$ is an isometry, $\flat$ is injective. Exercise 10.2.2.5 implies that $\flat$ is surjective. Hence $\flat$ is a bijection. Since $\flat$ is an isometry, $\|\flat^{-1}\| = 1$. Hence $\flat$ is an isomorphism. **reword this an make exercise in previous chapter about surjective isometries being isomorphisms**

□

Definition 10.2.2.8. Let H be a Hilbert Space. We define $\sharp : H^* \rightarrow \bar{H}$ by $\sharp := \flat^{-1}$.

10.3 Operators on Hilbert Spaces

10.3.1 Adjoint Operators

Note 10.3.1.1. We recall the definition of the adjoint of a bounded linear map from Exercise ??.

Exercise 10.3.1.2. Let H_1, H_2 be Hilbert spaces and $A \in L(H_1, H_2)$. Then there exists a unique $B \in L(H_2, H_1)$ such that for each $x_1 \in H_1$ and $x_2 \in H_2$,

$$\langle x_1, Bx_2 \rangle = \langle Ax_1, x_2 \rangle$$

Hint: B is essentially A^* where we identify H^* with $\bar{H}$.

Proof. Define $B := \sharp \circ A^* \circ \flat$. □

Definition 10.3.1.3. Adjoint of an Operator:

Let H_1, H_2 be a Hilbert space and $A \in L(H_1, H_2)$. We define the **adjoint of A** , denoted A^* , to be the unique $B \in L(H_2, H_1)$ such that for each $x_1 \in H_1$ and $x_2 \in H_2$,

$$\langle x_1, Bx_2 \rangle = \langle Ax_1, x_2 \rangle$$

Exercise 10.3.1.4. Let $H_1, H_2, H_3 \in \text{Obj}(\mathbf{Hilb})$.

1. For each $A, B \in L(H_1, H_2)$ and $\lambda \in \mathbb{C}$,

- (a) $(A^*)^* = A$
- (b) $(A + B)^* = A^* + B^*$
- (c) $(\lambda A)^* = \lambda^* A^*$

2. For each $A \in L(H_1, H_2)$ and $B \in L(H_2, H_3)$,

- (a) $(BA)^* = A^* B^*$
- (b) $H_1 = H_3$ implies that A and B commute iff A^* and B^* commute.

Proof.

1. Let $A, B \in L(H_1, H_2)$, $\lambda \in \mathbb{C}$.

(a) Let $x_1 \in H_1$ and $x_2 \in H_2$. Then

$$\begin{aligned} \langle x_2, Ax_1 \rangle^* &= \langle Ax_1, x_2 \rangle \\ &= \langle x_1, A^* x_2 \rangle^* \quad (\text{by definition}) \\ &= \langle A^* x_2, x_1 \rangle^*. \end{aligned}$$

Thus $\langle x_2, Ax_1 \rangle = \langle A^* x_2, x_1 \rangle$. Since $x_1 \in H_1$ and $x_2 \in H_2$ are arbitrary, we have that for each $x_1 \in H_1$ and $x_2 \in H_2$, $\langle x_2, Ax_1 \rangle = \langle A^* x_2, x_1 \rangle$. Uniqueness of $(A^*)^*$ implies that $(A^*)^* = A$.

(b) For each $x_1 \in H_1$, $x_2 \in H_2$,

$$\begin{aligned} \langle (A + B)x_1, x_2 \rangle &= \langle Ax_1 + Bx_1, x_2 \rangle \\ &= \langle Ax_1, x_2 \rangle + \langle Bx_1, x_2 \rangle \\ &= \langle x_1, A^* x_2 \rangle + \langle x_1, B^* x_2 \rangle \\ &= \langle x_1, A^* x_2 + B^* x_2 \rangle \\ &= \langle x_1, (A^* + B^*)x_2 \rangle. \end{aligned}$$

Uniqueness of $(A + B)^*$ implies that $(A + B)^* = A^* + B^*$.

(c) For each $x_1 \in H_1$, $x_2 \in H_2$,

$$\begin{aligned}\langle (\lambda A)x_1, x_2 \rangle &= \langle \lambda(Ax_1), x_2 \rangle \\ &= \lambda^* \langle Ax_1, x_2 \rangle \\ &= \lambda^* \langle x_1, A^*x_2 \rangle \\ &= \langle x_1, \lambda^*(A^*x_2) \rangle \\ &= \langle x_1, (\lambda^*A^*)x_2 \rangle.\end{aligned}$$

Uniqueness of $(\lambda A)^*$ implies that $(\lambda A)^* = \lambda^*A^*$.

2. Let $A \in L(H_1, H_2)$ and $B \in L(H_2, H_3)$.

(a) For each $x_1 \in H_1$, $x_2 \in H_2$,

$$\begin{aligned}\langle BAx_1, x_2 \rangle &= \langle Ax_1, B^*x_2 \rangle \\ &= \langle x_1, A^*B^*x_2 \rangle\end{aligned}$$

Uniqueness of $(BA)^*$ implies that $(BA)^* = A^*B^*$.

(b) Suppose that $H_1 = H_3$. If A and B commute, then

$$\begin{aligned}A^*B^* &= (BA)^* \\ &= (AB)^* \\ &= B^*A^*.\end{aligned}$$

Conversely, if A^* and B^* commute then

$$\begin{aligned}AB &= (B^*A^*)^* \\ &= (A^*B^*)^* \\ &= BA.\end{aligned}$$

□

10.3.2 Unitary Maps

Definition 10.3.2.1. Unitary Maps: Let $H_1, H_2 \in \text{Obj}(\mathbf{Hilb})$ and $U \in L(H_1, H_2)$. Then U is said to be **unitary** if U is invertible and for each $x, y \in H_1$, $\langle Ux, Uy \rangle = \langle x, y \rangle$.

Exercise 10.3.2.2. Let $H_1, H_2 \in \text{Obj}(\mathbf{Hilb})$ and $U \in L(H_1, H_2)$. Then the following are equivalent:

1. U is unitary
2. U is invertible and $U^{-1} = U^*$
3. U is surjective and U is an isometry

Proof.

- (1) $\implies$ (2):

Suppose that U is unitary. Then U is invertible and for each $x_1, x_2 \in H_1$, $\langle Ux_1, Ux_2 \rangle = \langle x_1, x_2 \rangle$. Let $x, y \in H$. Then

$$\begin{aligned}\langle y, x \rangle &= \langle Uy, Ux \rangle \\ &= \langle U^*Uy, x \rangle.\end{aligned}$$

Therefore

$$\begin{aligned} 0 &= \langle U^*Uy, x \rangle - \langle y, x \rangle \\ &= \langle U^*Uy - y, x \rangle \\ &= \langle (U^*U - I)y, x \rangle. \end{aligned}$$

Since $x \in H_1$ and $y \in H_2$ are arbitrary, for each $x \in H_1$ and $y \in H_2$, $\langle (U^*U - I)y, x \rangle = 0$. Exercise 10.2.1.12 implies that $(U^*U - I) = 0$. Hence $U^*U = I$. Therefore

$$\begin{aligned} U &= UI \\ &= U(U^*U) \\ &= (UU^*)U. \end{aligned}$$

Since U is invertible, we have that

$$\begin{aligned} I &= UU^{-1} \\ &= [(UU^*)U]U^{-1} \\ &= (UU^*)(UU^{-1}) \\ &= (UU^*)I \\ &= UU^*. \end{aligned}$$

Thus $U^*U = I$ and $UU^* = I$. Hence $U^{-1} = U^*$.

• (2) $\implies$ (3):

Suppose that U is invertible and $U^{-1} = U^*$. Since U is invertible, U is surjective. Let $x \in H_1$. Then

$$\begin{aligned} \|Ux\|^2 &= \langle Ux, Ux \rangle \\ &= \langle x, U^*Ux \rangle \\ &= \langle x, x \rangle \\ &= \|x\|^2. \end{aligned}$$

Hence $\|Ux\| = \|x\|$. Since $x \in X$ is arbitrary, we have that for each $x \in X$, $\|Ux\| = \|x\|$. Exercise 9.2.0.12 implies that U is an isometry.

• (3) $\implies$ (1):

Suppose that U is surjective and U is an isometry. Since U is an isometry, U is injective. Hence U is a bijection. Since U is an isometry, U^{-1} is an isometry. Thus $U^{-1} \in L(H_2, H_1)$ and U is invertible. Let $x, y \in H_1$. The polarization identity Exercise 10.2.1.14 and Exercise 9.2.0.12 imply that

$$\begin{aligned} \langle Ux, Uy \rangle &= \frac{1}{4} \left[\|Ux + Uy\|^2 - \|Ux - Uy\|^2 - i\|Ux + iUy\|^2 + i\|Ux - iUy\|^2 \right] \\ &= \frac{1}{4} \left[\|U(x + y)\|^2 - \|U(x - y)\|^2 - i\|U(x + iy)\|^2 + i\|U(x - iy)\|^2 \right] \\ &= \frac{1}{4} \left[\|x + y\|^2 - \|x - y\|^2 - i\|x + iy\|^2 + i\|x - iy\|^2 \right] \\ &= \langle x, y \rangle. \end{aligned}$$

Since $x, y \in H_1$ are arbitrary, we have that for each $x, y \in H_1$, $\langle Ux, Uy \rangle = \langle x, y \rangle$. Hence U is unitary. □

Definition 10.3.2.3. Let $H_1, H_2 \in \text{Obj}(\mathbf{Hilb})$. We define $U(H_1, H_2) := \{U \in L(H_1, H_2) : U \text{ is unitary}\}$. When $H_1 = H_2 = H$, we write $U(H)$ in place of $U(H, H)$.

Exercise 10.3.2.4. Let $H \in \text{Obj}(\mathbf{Hilb})$. Then $U(H)$ is a group.

Proof. **FINISH!!!**

□

Exercise 10.3.2.5. Let $H \in \text{Obj}(\mathbf{Hilb})$. Then $\mathfrak{b} \in U(\bar{H}, H^*)$.

Proof. **FINISH!!!**

□

Definition 10.3.2.6. Let H be a Hilbert space and $Q \in L(H)$. Then Q is said to be **self-adjoint** if

$$Q = Q^*$$

Exercise 10.3.2.7. Let $H \in \text{Obj}(\mathbf{Hilb})$. Then I is unitary and self-adjoint.

Proof. For each $x, y \in H$, $\langle Ix, Iy \rangle = \langle x, y \rangle$. So I is unitary. Exercise 10.3.2.2 implies that

$$\begin{aligned} I^* &= I^{-1} \\ &= I. \end{aligned}$$

Thus I is self-adjoint.

□

10.3.3 Spectral Theory (move to section on compact operators on hilbert space or banach algebra)

Exercise 10.3.3.1. Let H be a Hilbert space and $Q \in L(H)$. If Q is a self-adjoint then

1. the eigenvalues of Q are real. **reword**
2. the eigenvectors of Q corresponding to distinct eigenvalues are orthogonal.

Proof. Suppose that Q is self-adjoint.

1. Let λ be an eigenvalue of Q with corresponding eigenvector x . Then

$$\begin{aligned}\lambda\langle x, x \rangle &= \langle x, Qx \rangle \\ &= \langle Qx, x \rangle \\ &= \lambda^* \langle x, x \rangle\end{aligned}$$

Thus $\lambda = \lambda^*$ and is real

2. Let λ_1 and λ_2 be eigenvalues of Q with corresponding eigenvectors x_1 and x_2 . Suppose that $\lambda_1 \neq \lambda_2$. Then

$$\begin{aligned}\lambda_2 \langle x_1, x_2 \rangle &= \langle x_1, Qx_2 \rangle \\ &= \langle Qx_1, x_2 \rangle \\ &= \lambda_1 \langle x_1, x_2 \rangle\end{aligned}$$

So $(\lambda_2 - \lambda_1)\langle x_1, x_2 \rangle = 0$. Which implies that $\langle x_1, x_2 \rangle = 0$

□

Exercise 10.3.3.2. Let H be a Hilbert space, $A, B \in L(H)$ and $\lambda \in \mathbb{R}$. Suppose that A, B are self-adjoint. If A and B commute and then λAB is self-adjoint.

Proof.

$$\begin{aligned}(\lambda AB)^* &= \lambda^* (AB)^* \\ &= \lambda B^* A^* \\ &= \lambda BA \\ &= \lambda AB\end{aligned}$$

□

Definition 10.3.3.3. Adjoint of a Vector:

Let H be a Hilbert space and $x \in H$. We define the **adjoint** of x , denoted $x^* \in H^*$, by $x^*y = \langle x, y \rangle$.

Note 10.3.3.4. In mathematics, where linearity of the inner product is in the first argument, x^* is typically referred to by $u_x \in H^*$ where $u_x(y) = \langle y, x \rangle$. In physics, where the inner product with linearity in the second argument, $x^*\phi$ is usually written in the so-called “bra-ket” notation as $\langle x|\phi \rangle$ which works smoothly since it aligns with the linearity of $u_x(\phi_1 + \lambda\phi_2)$ and the conjugate-linearity of $u_{x_1+\lambda x_2}(\phi)$. In this way, it generalizes the notation for $\langle x, y \rangle = x^T y$ for $\mathbb{R}^n$ to $\langle x, y \rangle = x^* y$ for $\mathbb{C}^n$.

Exercise 10.3.3.5. Let H be a Hilbert space, $x, y \in H$ and $\lambda \in \mathbb{C}$. Then

1. $(x + y)^* = x^* + y^*$
2. $(\lambda x)^* = \lambda^* x^*$

Proof. Clear.

□

Definition 10.3.3.6. Let H be a Hilbert space, $x, y \in H$ and $A \in L(H)$. We define

1. $x^* A \in H^*$ by $(x^* A)y = x^*(Ay)$
2. $xy^* \in L(H)$ by $(xy^*)z = (y^*z)x$

Exercise 10.3.3.7. Let H be a Hilbert space, $A \in L(H)$ and $x \in H$. Then

$$(Ax)^* = x^* A^*$$

Proof. Let $y \in H$. Then

$$\begin{aligned}(Ax)^*y &= \langle Ax, y \rangle \\ &= \langle x, A^*y \rangle \\ &= x^*A^*y\end{aligned}$$

□

Definition 10.3.3.8. Commutator:

Let H be a Hilbert space and $A, B \in L(H)$. The **commutator** of A and B , denoted $[A, B]$, is defined by

$$[A, B] = AB - BA$$

Exercise 10.3.3.9. Let H be a Hilbert space and $A, B, C \in L(H)$. Then

1. $[AB, C] = A[B, C] + [A, C]B$
2. $[A, BC] = B[A, C] + [A, B]C$

Proof.

1.

$$\begin{aligned}[AB, C] &= ABC - CAB \\ &= ABC - ACB + ACB - CAB \\ &= A(BC - CB) + (AC - CA)B \\ &= A[B, C] + [A, C]B\end{aligned}$$

2. Similar to (1).

□

Exercise 10.3.3.10. Let $H_1, H_2, H_3, H_4 \in \text{Obj}(\mathbf{Hilb})$ and $A \in L(H_3, H_4)$, $B \in L(H_2, H_3)$ and $C \in L(H_1, H_2)$. Suppose that H, K, L are separable. Let $(e_j^2)_{j \in \mathbb{N}}$ and $(e_j^3)_{j \in \mathbb{N}}$ be orthonormal bases for H_2, H_3 and H_4 respectively. Then

$$ABC = \sum_{j \in \mathbb{N}} \sum_{k \in \mathbb{N}} \langle e_j^3, Be_k^2 \rangle (Ae_j^3)^* (e_k^2 C)$$

FINISH!!!

Proof. content...

□

10.4 Subspaces of Hilbert Spaces

10.4.1 Introduction

Exercise 10.4.1.1. Let $(H, \langle \cdot, \cdot \rangle) \in \text{Obj}(\mathbf{Hilb})$ and $E \subset H$ a closed subspace of H . Then $(E, \langle \cdot, \cdot \rangle|_{E \times E})$ is a Hilbert space.

Proof. Clearly $(E, \langle \cdot, \cdot \rangle|_{E \times E})$ is an inner product space. Since E is closed and H is complete, E is complete. Hence $(E, \langle \cdot, \cdot \rangle|_{E \times E})$ is a Hilbert space. $\square$

10.4.2 Orthogonal Projections

Definition 10.4.2.1. Let $H \in \text{Obj}(\mathbf{Hilb})$ and $P \in L(H)$. Then P is said to be **idempotent** if $P^2 = P$.

Exercise 10.4.2.2. Let $H \in \text{Obj}(\mathbf{Hilb})$ and $P \in L(H)$. Then P is idempotent iff $I - P$ is idempotent.

Proof.

- $(\implies)$:
Suppose that P is idempotent. Then

$$\begin{aligned} (I - P)^2 &= (I - P)(I - P) \\ &= I^2 - IP - PI + P^2 \\ &= I - 2P + P \\ &= I - P \end{aligned}$$

So $I - P$ is idempotent.

- $(\impliedby)$:
Suppose that $I - P$ is idempotent. Part (1) implies that $I - (I - P) = P$ is idempotent.

$\square$

Exercise 10.4.2.3. Let $H \in \text{Obj}(\mathbf{Hilb})$, $E \subset H$ a closed subspace of H and $P \in L(H)$. If $P|_E = I_E$ and $\text{Img } P \subset E$, then P is idempotent

Proof. Suppose that $P|_E = I_E$ and $\text{Img } P \subset E$. Let $x \in H$ and define $y \in E$ by $y := P(x)$. Then

$$\begin{aligned} P^2(x) &= P(y) \\ &= P|_E(y) \\ &= I_E(y) \\ &= y \\ &= P(x) \end{aligned}$$

Since $x \in H$ is arbitrary, we have that $P^2 = P$ and P is idempotent. $\square$

Definition 10.4.2.4. Let $H \in \text{Obj}(\mathbf{Hilb})$ and $E \subset H$ a closed subspace of H . We define the **orthogonal projection onto E** , denoted $P_E : H \rightarrow E$ by

$$P_E(x) = \arg \min_{y \in E^\perp} \|x - y\|$$

Note 10.4.2.5. Exercise 10.2.2.2 implies that P_E is well-defined.

Exercise 10.4.2.6. Let $H \in \text{Obj}(\mathbf{Hilb})$ and $E \subset H$ a closed subspace of H . Then

1. $P_E|_E = I_E$
2. $\text{Img } P_E = E$
3. P_E is linear

$$4. P_E \in L(H, E) \text{ and } \|P_E\| = \begin{cases} 0, & E = \{0\} \\ 1 & E \neq \{0\} \end{cases}$$

5. P_E is self-adjoint

6. P_E is idempotent

7. $\ker P_E = E^\perp$

Proof.

1. Let $x \in E$. Then

$$\begin{aligned} P_E(x) &= \arg \min_{y \in E^\perp} \|x - y\| \\ &= x \end{aligned}$$

Since $x \in E$ is arbitrary, $P_E|_E = I_E$.

2. By definition, $\text{Img } P_E \subset E$. The previous part implies that $E \subset \text{Img } P_E$. Thus $\text{Img } P_E = E$.

3. Let $x_1, x_2 \in H$ and $\lambda \in \mathbb{C}$. A previous exercise in the introduction section implies that there exist unique $y_1, y_2 \in E$ and $z_1, z_2 \in E^\perp$ such that $x_1 = y_1 + z_1$, $x_2 = y_2 + z_2$, $\|x_1 + E\| = \|z_1\|$ and $\|x_2 + E\| = \|z_2\|$. Then $y_1 + \lambda y_2 \in E$, $z_1 + \lambda z_2 \in E^\perp$ and $x_1 + \lambda x_2 = (y_1 + \lambda y_2) + (z_1 + \lambda z_2)$. An exercise in the introduction section implies that $\|x_1 + \lambda x_2 + E\| = \|z_1 + \lambda z_2\|$. Uniqueness implies that

$$\begin{aligned} P_E(x_1 + \lambda x_2) &= y_1 + \lambda y_2 \\ &= P_E(x_1) + \lambda P_E(x_2) \end{aligned}$$

Since $x_1, x_2 \in H$ and $\lambda \in \mathbb{C}$ are arbitrary, P_E is linear.

4. Let $x \in H$. Then there exist unique $y \in E$ and $z \in E^\perp$ such that $x = y + z$ and $\|x + E\| = \|z\|$. The Pythagorean theorem implies that

$$\begin{aligned} \|P_E(x)\|^2 &= \|y\|^2 \\ &\leq \|y\|^2 + \|z\|^2 \\ &= \|y + z\|^2 \\ &= \|x\|^2 \end{aligned}$$

So $\|P_E(x)\| \leq \|x\|$. Since $x \in H$ is arbitrary, $P_E \in L(H, E)$ and $\|P_E\| \leq 1$. If $E = \{0\}$, then $P_E = 0$ and therefore $\|P_E\| = 0$. Suppose that $E \neq \{0\}$. Then there exists $y \in E$ such that $\|y\| = 1$. Hence

$$\begin{aligned} \|P_E\| &= \sup_{y' \neq 0} \|y'\|^{-1} \|P_E(y')\| \\ &\geq \|P_E(y)\| \\ &= \|y\| \\ &= 1 \end{aligned}$$

So $\|P_E\| = 1$.

5. **FINISH!!!**

6. Since $P \in L(H)$, $P|_E = I_E$ and $\text{Img } P \subset E$, a previous exercise implies that P_E is idempotent.

7. • Let $x \in \ker P_E$. Exercise 10.2.2.2 implies that there exists unique $z_0 \in E^\perp$ such that $x = P_E(x) + z_0$ and $\|z_0\| = \|x + E\|$. Since $x \in \ker P_E$, we have that

$$\begin{aligned} x &= \\ &= P_E(x) + z_0 \\ &= 0 + z_0 \\ &= z_0. \end{aligned}$$

Hence $x \in E^\perp$. Since $x \in \ker P$ is arbitrary, we have that $\ker P \subset E^\perp$.

- Let $x \in E^\perp$. Then Exercise 10.2.2.2 implies that $P_E(x) = 0$. Thus $x \in \ker P_E$. Since $x \in E^\perp$ is arbitrary, we have that $E^\perp \subset \ker P_E$.

Since $E^\perp \subset \ker P_E$ and $P_E \subset \ker E^\perp$, we have that $\ker P_E = E^\perp$.

□

Exercise 10.4.2.7. **FINISH!!!**

Proof.

$$\begin{aligned} \|A\|^2 &= \left(\sup_{\|x\|=1} \|Ax\| \right)^2 \\ &= \sup_{\|x\|=1} \|Ax\|^2 \\ &= \sup_{\|x\|=1} \langle Ax, Ax \rangle \\ &= \sup_{\|x\|=1} \langle x, A^*Ax \rangle \\ &= \end{aligned}$$

FINISH!!!

□

Exercise 10.4.2.8. Let $H \in \text{Obj}(\mathbf{Hilb})$, $E \subset H$ a closed subspace of H . Then there exists a unique $P \in L(H)$ such that

1. P is self-adjoint
2. P is idempotent
3. $\text{Img } P = E$

Hint: for uniqueness, if P and Q satisfy (1) – (3), consider $(P - Q)^*(P - Q)$

Proof.

- **(Existence):**

P_E satisfies (1) – (3).

- **(Uniqueness):**

Let $Q \in L(H, E)$. Suppose that Q is self-adjoint, Q is idempotent and $\text{Img } Q = E$. The previous exercise implies that $P|_E, Q|_E = I_E$. Then

$$\begin{aligned} (P - Q)^*(P - Q) &= P^*P - P^*Q - Q^*P + Q^*Q \\ &= P^2 - PQ - QP + Q^2 \\ &= P^2 - P|_E Q - Q|_E P + Q^2 \\ &= P - Q - P + Q \\ &= 0 \end{aligned}$$

make exercise about $\|T\| = \|T^*\|$ and $\|T\| = \max\langle x, Ty \rangle : \|x\|, \|y\| = 1$

$$\begin{aligned}\|(P - Q)\|^2 &= \|(P - Q)^*(P - Q)\| \\ &= \|0\| \\ &= 0.\end{aligned}$$

Thus $\|P - Q\| = 0$ and therefore $P - Q = 0$. Hence $P = Q$ Thus $\|(P - Q)^*(P - Q)\|$ **FINISH!!!**

□

10.5 Direct Sums of Hilbert spaces

need to find universal property for direct sum

Definition 10.5.0.1. Let $(H_\alpha, \langle \cdot, \cdot \rangle_\alpha)_{\alpha \in A} \subset \text{Obj}(\mathbf{Hilb})$. We define $s_A : \prod_{\alpha \in A} H_\alpha \rightarrow [0, \infty)^A$ by $s_A(x) = (\|x_\alpha\|_\alpha)_{\alpha \in A}$.

Exercise 10.5.0.2. Let $(H_\alpha, \langle \cdot, \cdot \rangle_\alpha)_{\alpha \in A} \subset \text{Obj}(\mathbf{Hilb})$. Then

1. for each $x, y \in s_A^{-1}(l^2(A))$, $(\langle x_\alpha, y_\alpha \rangle_\alpha)_{\alpha \in A} \in l^1(A)$
2. for each $x, y \in \prod_{\alpha \in A} H_\alpha$, $\|s_A(x) - s_A(y)\|_2 \leq \|s_A(x - y)\|_2$

Proof.

1. Let $x, y \in s_A^{-1}(l^2(A))$. Then $s_A(x), s_A(y) \in l^2(A)$. Therefore $s_A(x)s_A(y) \in l^1(A)$ and

$$\begin{aligned} \|\langle x_\alpha, y_\alpha \rangle_\alpha\|_1 &= \sum_{\alpha \in A} |\langle x_\alpha, y_\alpha \rangle_\alpha| \\ &\leq \sum_{\alpha \in A} \|x_\alpha\|_\alpha \|y_\alpha\|_\alpha \\ &= \|s_A(x)s_A(y)\|_1 \\ &< \infty \end{aligned}$$

2. Let $x, y \in \prod_{\alpha \in A} H_\alpha$. The reverse triangle inequality implies that

$$\begin{aligned} \|s_A(x) - s_A(y)\|_2^2 &= \|(\|x_\alpha\|_\alpha)_{\alpha \in A} - (\|y_\alpha\|_\alpha)_{\alpha \in A}\|_2^2 \\ &= \|(\|x_\alpha\|_\alpha - \|y_\alpha\|_\alpha)_{\alpha \in A}\|_2^2 \\ &= \sum_{\alpha \in A} |\|x_\alpha\|_\alpha - \|y_\alpha\|_\alpha|^2 \\ &\leq \sum_{\alpha \in A} \|x_\alpha - y_\alpha\|_\alpha^2 \\ &= \|(\|x - y\|_\alpha)_{\alpha \in A}\|_2^2 \\ &= \|s_A(x - y)\|_2^2 \end{aligned}$$

Thus $\|s_A(x) - s_A(y)\|_2 \leq \|s_A(x - y)\|_2$.

□

Definition 10.5.0.3. Let $(H_\alpha, \langle \cdot, \cdot \rangle_\alpha)_{\alpha \in A} \subset \text{Obj}(\mathbf{Hilb})$. We define $\bigoplus_{\alpha \in A} H_\alpha \subset \prod_{\alpha \in A} H_\alpha$ and $\langle \cdot, \cdot \rangle : \left[\bigoplus_{\alpha \in A} H_\alpha \right]^2 \rightarrow \mathbb{C}$ by

$$\bigoplus_{\alpha \in A} H_\alpha = s_A^{-1}(l^2(A))$$

and

$$\langle x, y \rangle = \sum_{\alpha \in A} \langle x_\alpha, y_\alpha \rangle_\alpha$$

We define the **direct sum of** $(H_\alpha, \langle \cdot, \cdot \rangle_\alpha)_{\alpha \in A}$ to be $(\bigoplus_{\alpha \in A} H_\alpha, \langle \cdot, \cdot \rangle)$.

Exercise 10.5.0.4. Let $(H_\alpha, \langle \cdot, \cdot \rangle_\alpha)_{\alpha \in A} \subset \text{Obj}(\mathbf{Hilb})$. Then

1. for each $x \in \prod_{\alpha \in A} H_\alpha$, $x \in \bigoplus_{\alpha \in A} H_\alpha$ iff $s_A(x) \in l^2(A)$

$$2. s_A|_{\bigoplus_{\alpha \in A} H_\alpha} : \bigoplus_{\alpha \in A} H_\alpha \rightarrow l^2(A)$$

Proof. Immediate by definition □

Exercise 10.5.0.5. Let $(H_\alpha, \langle \cdot, \cdot \rangle_\alpha)_{\alpha \in A} \subset \text{Obj}(\mathbf{Hilb})$. Set $H := \bigoplus_{\alpha \in A} H_\alpha$. Then

1. H is a vector space
2. $\langle \cdot, \cdot \rangle : H \times H \rightarrow \mathbb{C}$ is an inner product
3. $(H, \langle \cdot, \cdot \rangle)$ is a Hilbert space

Proof.

1. Clearly $\prod_{\alpha \in A} H_\alpha$ is a vector space. Let $x, y \in H$ and $\lambda \in \mathbb{C}$. The previous exercise implies that $s_A(x), s_A(y) \in l^2(A)$. Therefore $s_A(x)s_A(y) \in L^1(A)$ and

$$\begin{aligned} \sum_{\alpha \in A} \|(x + \lambda y)_\alpha\|_\alpha^2 &= \sum_{\alpha \in A} \|x_\alpha + \lambda y_\alpha\|_\alpha^2 \\ &= \sum_{\alpha \in A} \left[\|x_\alpha\|_\alpha^2 + |\lambda|^2 \|y_\alpha\|_\alpha^2 + 2 \operatorname{Re}(\langle x_\alpha, y_\alpha \rangle_\alpha) \right] \\ &\leq \sum_{\alpha \in A} \|x_\alpha\|_\alpha^2 + |\lambda|^2 \sum_{\alpha \in A} \|y_\alpha\|_\alpha^2 + 2 \sum_{\alpha \in A} \|x_\alpha\|_\alpha \|y_\alpha\|_\alpha \\ &= \|s_A(x)\|_2^2 + |\lambda|^2 \|s_A(y)\|_2^2 + 2 \|s_A(x)s_A(y)\|_1 \\ &< \infty \end{aligned}$$

Thus H is a vector space.

2. Let $x, y, z \in H$ and $\lambda \in \mathbb{C}$. Then

(a)

$$\begin{aligned} \langle x, y + \lambda z \rangle &= \sum_{\alpha \in A} \langle x_\alpha, (y + \lambda z)_\alpha \rangle_\alpha \\ &= \sum_{\alpha \in A} \langle x_\alpha, y_\alpha + \lambda z_\alpha \rangle_\alpha \\ &= \sum_{\alpha \in A} \left[\langle x_\alpha, y_\alpha \rangle_\alpha + \lambda \langle x_\alpha, z_\alpha \rangle_\alpha \right] \\ &= \sum_{\alpha \in A} \langle x_\alpha, y_\alpha \rangle_\alpha + \lambda \sum_{\alpha \in A} \langle x_\alpha, z_\alpha \rangle_\alpha \\ &= \langle x, y \rangle + \lambda \langle x, z \rangle \end{aligned}$$

(b)

$$\begin{aligned} \langle x, y \rangle &= \sum_{\alpha \in A} \langle x_\alpha, y_\alpha \rangle_\alpha \\ &= \sum_{\alpha \in A} \langle y_\alpha, x_\alpha \rangle_\alpha^* \\ &= \left(\sum_{\alpha \in A} \langle y_\alpha, x_\alpha \rangle_\alpha \right)^* \\ &= \langle y, x \rangle \end{aligned}$$

(c)

$$\begin{aligned}\langle x, x \rangle &= \sum_{\alpha \in A} \langle x_\alpha, x_\alpha \rangle_\alpha \\ &\geq 0\end{aligned}$$

(d) Suppose that $\langle x, x \rangle = 0$. Then

$$\begin{aligned}0 &= \langle x, x \rangle \\ &= \sum_{\alpha \in A} \langle x_\alpha, x_\alpha \rangle_\alpha \\ &= \sum_{\alpha \in A} \|x_\alpha\|_\alpha^2\end{aligned}$$

Thus for each $\alpha \in A$, $\|x_\alpha\|_\alpha = 0$. Therefore for each $\alpha \in A$, $x_\alpha = 0$. Hence $x = 0$.

So $\langle \cdot, \cdot \rangle : H \times H \rightarrow \mathbb{C}$ is an inner product on H .

3. Let $(x_j)_{j \in \mathbb{N}} \subset H$. Suppose that $(x_j)_{j \in \mathbb{N}}$ Cauchy. Let $\alpha \in A$ and $\epsilon > 0$. Since $(x_j)_{j \in \mathbb{N}}$ Cauchy, there exists $N \in \mathbb{N}$ such that for each $m, n \in \mathbb{N}$, $m, n \geq N$ implies that $\|x_m - x_n\| < \epsilon$. Let $m, n \in \mathbb{N}$. Suppose that $m, n \geq N$. Then

$$\begin{aligned}\|x_{m,\alpha} - x_{n,\alpha}\|_\alpha^2 &\leq \sum_{\beta \in A} \|x_{m,\beta} - x_{n,\beta}\|_\beta^2 \\ &= \|x_m - x_n\|^2 \\ &< \epsilon^2\end{aligned}$$

Thus $\|x_{m,\alpha} - x_{n,\alpha}\|_\alpha \leq \epsilon$. Since $\epsilon > 0$ is arbitrary, we have that for each $\epsilon > 0$, there exists $N \in \mathbb{N}$ such that for each $m, n \in \mathbb{N}$, $m, n \geq N$ implies that $\|x_{m,\alpha} - x_{n,\alpha}\|_\alpha < \epsilon$. Hence $(x_{j,\alpha})_{j \in \mathbb{N}}$ is Cauchy. Since H_α is complete, there exists $x_\alpha \in H_\alpha$ such that $x_{j,\alpha} \rightarrow x_\alpha$. Since $\alpha \in A$ is arbitrary, we have that for each $\alpha \in A$, there exists $x_\alpha \in H_\alpha$ such that $x_{j,\alpha} \rightarrow x_\alpha$. Define $x \in \prod_{\alpha \in A} H_\alpha$

by $x = (x_\alpha)_{\alpha \in A}$.

Let $\epsilon > 0$. Since $(x_n)_{n \in \mathbb{N}}$ is Cauchy, there exists $N \in \mathbb{N}$ such that for each $m, n \in \mathbb{N}$, $m, n \geq N$ implies that $\|x_m - x_n\| \leq \epsilon$. Let $n \in \mathbb{N}$. Suppose that $n \geq N$. Fatou's lemma imply that

$$\begin{aligned}\|s_A(x - x_n)\|_2^2 &= \sum_{\alpha \in A} \|x_\alpha - x_{n,\alpha}\|_\alpha^2 \\ &= \sum_{\alpha \in A} \lim_{m \rightarrow \infty} \|x_{m,\alpha} - x_{n,\alpha}\|_\alpha^2 \\ &\leq \liminf_{m \rightarrow \infty} \sum_{\alpha \in A} \|x_{m,\alpha} - x_{n,\alpha}\|_\alpha^2 \\ &= \liminf_{m \rightarrow \infty} \|x_m - x_n\|^2 \\ &\leq \epsilon^2\end{aligned}$$

Thus $\|s_A(x - x_n)\|_2 \leq \epsilon$. Since $\epsilon > 0$ is arbitrary, we have that for each $\epsilon > 0$, there exists $N_\epsilon \in \mathbb{N}$ such that for each $n \in \mathbb{N}$, $n \geq N_\epsilon$ implies that $\|s_A(x - x_n)\|_2 \leq \epsilon$.

In particular, setting $\epsilon = 1$ and $n = N_1$, [A previous exercise](#) implies that

$$\begin{aligned}\|s_A(x)\|_2 &\leq \|s_A(x) - s_A(x_{N_1})\|_2 + \|s_A(x_{N_1})\|_2 \\ &= \|s_A(x) - s_A(x_{N_1})\|_2 + \|x_{N_1}\| \\ &\leq \|s_A(x - x_{N_1})\|_2 + \|x_{N_1}\| \\ &\leq 1 + \|x_{N_1}\| \\ &< \infty\end{aligned}$$

so that $s_A(x) \in l^2(A)$ and therefore $x \in \bigoplus_{\alpha \in A} H_\alpha$.

Since $x \in H$, we have that for each $n \in \mathbb{N}$, $\|x - x_n\| = \|s_A(x - x_n)\|_2$. Thus from before, for each $\epsilon > 0$, there exists $N_\epsilon \in \mathbb{N}$ such that for each $n \in \mathbb{N}$, $n \geq N_\epsilon$ implies that $\|x - x_n\| \leq \epsilon$. Hence $x_n \rightarrow x$. Since $(x_n)_{n \in \mathbb{N}} \subset H$ with $(x_n)_{n \in \mathbb{N}}$ Cauchy is arbitrary, we have that for each $(x_n)_{n \in \mathbb{N}} \subset H$, $(x_n)_{n \in \mathbb{N}}$ is Cauchy implies that there exists $x \in H$ such that $x_n \rightarrow x$. Hence H is complete. Thus $(H, \langle \cdot, \cdot \rangle)$ is a Hilbert space. □

Note 10.5.0.6. This construction might work for Banach spaces with norms satisfying $\|x_\alpha + y_\alpha\|_\alpha^2 \leq \|x_\alpha\|_\alpha^2 + \|y_\alpha\|_\alpha^2 + 2\|x_\alpha\|_\alpha\|y_\alpha\|_\alpha$.

Exercise 10.5.0.7. Let $H \in \text{Obj}(\mathbf{Hilb})$, A an index set and for each $\alpha \in A$, E_α a closed subspace of H . Then $H \cong \bigoplus_{\alpha \in A} E_\alpha$ iff

1. for each $\alpha, \beta \in A$, $\alpha \neq \beta$ implies that $E_\alpha \cap E_\beta = \{0\}$
2. for each $x \in H$, there exist $(x_\alpha)_{\alpha \in A} \in \prod_{\alpha \in A} E_\alpha$ such that $x = \sum_{\alpha \in A} x_\alpha$

Proof.

- ($\Rightarrow$):
Suppose that $H \cong \bigoplus_{\alpha \in A} E_\alpha$. Let $x \in H$.
- ($\Leftarrow$):

□

Exercise 10.5.0.8. Let $H \in \text{Obj}(\mathbf{Hilb})$ and $E \subset H$ a closed subspace of H . Then $E^\perp \oplus E$ something about $E^\perp \cong H/E$ and adding up to H . **FINISH!!!**

Proof.

□

10.6 Tensor Products

Exercise 10.6.0.1. Let $H_1, \dots, H_n \in \text{Obj}(\mathbf{Hilb})$, $\phi \in L^n(H_1, \dots, H_n; \mathbb{C})$. Suppose that for each $j \in [n]$, H_j is separable. For each $j \in [n]$, let $B_j \subset H_j$ be an orthonormal basis for H_j . Then

$$\sum_{e_1 \in B_1} \cdots \sum_{e_n \in B_n} |\phi(e_1, \dots, e_n)|^2 = \sum_{e'_1 \in B'_1} \cdots \sum_{e'_n \in B'_n} |\phi(e'_1, \dots, e'_n)|^2.$$

Proof. **Induction on n** □

Definition 10.6.0.2. Let $H_1, \dots, H_n \in \text{Obj}(\mathbf{Hilb})$, $\phi \in L^n(H_1, \dots, H_n; \mathbb{C})$. Suppose that for each $j \in [n]$, H_j is separable. For each $j \in [n]$, let $B_j \subset H_j$ be an orthonormal basis for H_j . We define $\|\cdot\|_{HS} : L^n(H_1, \dots, H_n; \mathbb{C}) \rightarrow [0, \infty]$ by

$$\|\phi\|_{HS} := \left[\sum_{e_1 \in B_1} \sum_{e_1 \in B_1} \cdots \sum_{e_n \in B_n} |\phi(e_1, \dots, e_n)|^2 \right]^{1/2}.$$

We define the **Hilbert-Schmidt functionals on $H_1, \dots, H_n$** , denoted $HS^n(H_1, \dots, H_n; \mathbb{C})$, by

$$HS^n(H_1, \dots, H_n; \mathbb{C}) := \{\phi \in L^n(H_1, \dots, H_n; \mathbb{C}) : \|\phi\|_{HS} < \infty\}.$$

We define $\langle \cdot, \cdot \rangle_{HS} : HS^n(H_1, \dots, H_n; \mathbb{C}) \rightarrow \mathbb{C}$ by

$$\langle \phi, \psi \rangle_{HS} := \sum_{e_1 \in B_1} \cdots \sum_{e_n \in B_n} \phi(e_1, \dots, e_n)^* \psi(e_1, \dots, e_n).$$

When $H_1 = \cdots = H_n = H$, we write $HS^n(H; \mathbb{C})$ in place of $HS^n(H_1, \dots, H_n; \mathbb{C})$.

Exercise 10.6.0.3. Let $H_1, \dots, H_n \in \text{Obj}(\mathbf{Hilb})$, $\phi \in L^n(H_1, \dots, H_n; \mathbb{C})$. Suppose that for each $j \in [n]$, H_j is separable. For each $j \in [n]$, let $B_j \subset H_j$ be an orthonormal basis for H_j .

1. $\langle \cdot, \cdot \rangle_{HS}$ is an inner product on $HS^n(H_1, \dots, H_n; \mathbb{C})$
2. for each $\phi \in HS^n(H_1, \dots, H_n; \mathbb{C})$, $\langle \phi, \phi \rangle_{HS} = \|\phi\|_{HS}^2$.

Proof.

1. Let $\phi, \psi, \chi \in HS^n(H_1, \dots, H_n; \mathbb{C})$ and $\lambda \in \mathbb{C}$. Then

$$\begin{aligned} \langle \phi, \psi + \lambda \chi \rangle_{HS} &= \sum_{e_1 \in B_1} \cdots \sum_{e_n \in B_n} \phi(e_1, \dots, e_n)^* \left[\psi(e_1, \dots, e_n) + \lambda \chi(e_1, \dots, e_n) \right] \\ &= \sum_{e_1 \in B_1} \cdots \sum_{e_n \in B_n} \phi(e_1, \dots, e_n)^* \psi(e_1, \dots, e_n) + \lambda \sum_{e_1 \in B_1} \cdots \sum_{e_n \in B_n} \phi(e_1, \dots, e_n)^* \chi(e_1, \dots, e_n) \\ &= \langle \phi, \psi \rangle_{HS} + \lambda \langle \phi, \chi \rangle_{HS}. \end{aligned}$$

(b)

$$\begin{aligned} \langle \phi, \psi \rangle_{HS} &= \sum_{e_1 \in B_1} \cdots \sum_{e_n \in B_n} \phi(e_1, \dots, e_n)^* \psi(e_1, \dots, e_n) \\ &= \left[\sum_{e_1 \in B_1} \cdots \sum_{e_n \in B_n} \phi(e_1, \dots, e_n) \psi(e_1, \dots, e_n)^* \right]^* \\ &= \left[\sum_{e_1 \in B_1} \cdots \sum_{e_n \in B_n} \psi(e_1, \dots, e_n)^* \phi(e_1, \dots, e_n) \right]^* \\ &= \langle \psi, \phi \rangle_{HS}^*. \end{aligned}$$

(c)

$$\begin{aligned}
\langle \phi, \psi \rangle_{HS} &= \sum_{e_1 \in B_1} \cdots \sum_{e_n \in B_n} \phi(e_1, \dots, e_n)^* \phi(e_1, \dots, e_n) \\
&= \sum_{e_1 \in B_1} \cdots \sum_{e_n \in B_n} |\phi(e_1, \dots, e_n)|^2 \\
&= \|\phi\|_{HS}^2 \\
&\geq 0
\end{aligned}$$

(d) Suppose that $\langle \phi, \phi \rangle_{HS} = 0$. Then

$$\begin{aligned}
\sum_{e_1 \in B_1} \cdots \sum_{e_n \in B_n} |\phi(e_1, \dots, e_n)|^2 &= \langle \phi, \phi \rangle_{HS} \\
&= 0.
\end{aligned}$$

Thus for each $e_{1,j_1} \in B_1, \dots, e_{n,j_n} \in B_n$, $\phi(e_1, \dots, e_n) = 0$. **make exercise about how multilinear maps are determined by value on tuples of orthonormal basis elements** Exercise 17.3.0.5 implies that $\phi = 0$.

2. Clear.

FINISH!!!

□

Exercise 10.6.0.4. Let $H \in \text{Obj}(\mathbf{Hilb})$. Suppose that H is separable. Then

1. for each $\phi \in H^*$, $\|\phi\|_{HS} = \|\phi\|$
2. $HS(H; \mathbb{C}) = \mathbb{H}^*$.

Proof. Since H is separable, there exists $(e_j)_{j \in \mathbb{N}} \subset H$ such that $(e_j)_{j \in \mathbb{N}}$ is an orthonormal basis for H .

1. Let $\phi \in H^*$. Then

$$\begin{aligned}
\|\phi\| &= \sup_{\|x\|=1} |\phi(x)| \\
&\leq
\end{aligned}$$

2.

□

Exercise 10.6.0.5. content...**Exercise 10.6.0.6.** associativity of tensor product

Definition 10.6.0.7. Let $H_1, H_2, H \in \text{Obj}(\mathbf{Hilb})$, $\alpha \in L^2(H_1, H_2; H)$. Suppose that for each $j \in [n]$, H_j is separable. For $j \in [n]$, let $B_j \subset H_j$ be an orthonormal basis for H_j . Then α is said to be **Hilbert-Schmidt** if

$$\sum_{e^1 \in B_1} \cdots \sum_{e^n \in B_n} |\phi(e^1, \dots, e^n)|^2 < \infty$$

Definition 10.6.0.8. Let $H_1, H_2, H \in \text{Obj}(\mathbf{Hilb})$, $\alpha \in L^2(H_1, H_2; H)$, $(e_j)_{j \in \mathbb{N}}$ and $(f_k)_{k \in \mathbb{N}}$ orthonormal bases for H_1 and H_2 respectively. Then α is said to be **weakly Hilbert-Schmidt** if there exists $M \geq 0$ such that for each $x \in H$,

Definition 10.6.0.9. Let $H_1, H_2, K \in \text{Obj}(\mathbf{Hilb})$ and $\alpha \in L^2(H_1, H_2; K)$. Then (K, α) is said to be a **tensor product of H_1 and H_2** if for each vector space Z and $\beta \in L^2(X, Y; Z)$, there exists a unique $\phi \in L^1(T; Z)$ such that $\phi \circ \alpha = \beta$, i.e. the following diagram commutes:

$$\begin{array}{ccc}
X \times Y & \xrightarrow{\alpha} & T \\
& \searrow \beta & \downarrow \phi \\
& & Z
\end{array}$$

Exercise 10.6.0.10. Let H be a Hilbert space and $a_1, a_2 \in H$. Then for each $x_1, x_2 \in H$,

$$\langle x_1, (a_1 \otimes a_2^*)x_2 \rangle = \langle x_1, a_1 \rangle \langle a_2, x_2 \rangle$$

Proof. For each $x_1, x_2 \in H$,

$$\begin{aligned} \langle x_1, (a_1 \otimes a_2^*)x_2 \rangle &= \langle x_1, (a_2^*x_2)a_1 \rangle \\ &= (a_2^*x_2) \langle x_1, a_1 \rangle \\ &= \langle a_2, x_2 \rangle \langle x_1, a_1 \rangle \\ &= \langle x_1, a_1 \rangle \langle a_2, x_2 \rangle. \end{aligned}$$

□

Exercise 10.6.0.11. Let H be a Hilbert space. Suppose that H is separable. Let $(e_j)_{j \in \mathbb{N}}$ be an orthonormal basis of H . Then

$$I = \sum_{j \in \mathbb{N}} e_j \otimes e_j^*$$

Proof. For each $x_1, x_2 \in H$,

$$\begin{aligned} \left\langle x_1, \left(\sum_{j \in \mathbb{N}} e_j \otimes e_j^* \right) x_2 \right\rangle &= \left\langle x_1, \sum_{j \in \mathbb{N}} (e_j \otimes e_j^*) x_2 \right\rangle \\ &= \sum_{j \in \mathbb{N}} \langle x_1, (e_j \otimes e_j^*) x_2 \rangle \\ &= \sum_{j \in \mathbb{N}} \langle x_1, e_j \rangle \langle e_j, x_2 \rangle \quad (\text{by the prev ex}) \\ &= \left\langle \sum_{j \in \mathbb{N}} \langle e_j, x_1 \rangle e_j, \sum_{k \in \mathbb{N}} \langle e_k, x_2 \rangle e_k \right\rangle \quad (\text{by Exercise 10.2.1.28}) \\ &= \langle x_1, x_2 \rangle \\ &= \langle x_1, Ix_2 \rangle. \end{aligned}$$

A prev ex then implies that

$$I = \sum_{j \in \mathbb{N}} e_j \otimes e_j^*.$$

□

Exercise 10.6.0.12. Let H_1, H_2 be Hilbert spaces. Suppose that H_1 is separable. Let $(e_j)_{j \in \mathbb{N}}$ be an orthonormal basis of H_1 . Then for each $A \in L(H_1, H_2)$,

$$AA^* = \sum_{j \in \mathbb{N}} (Ae_j) \otimes (Ae_j)^*.$$

Proof. For each $y_1, y_2 \in H_2$,

$$\begin{aligned}
 \left\langle y_1, \left[\sum_{j \in \mathbb{N}} (Ae_j) \otimes (Ae_j)^* \right] y_2 \right\rangle &= \left\langle y_1, \sum_{j \in \mathbb{N}} [(Ae_j) \otimes (Ae_j)^*] y_2 \right\rangle \\
 &= \sum_{j \in \mathbb{N}} \langle y_1, [(Ae_j) \otimes (Ae_j)^*] y_2 \rangle \\
 &= \sum_{j \in \mathbb{N}} \langle y_1, Ae_j \rangle \langle Ae_j, y_2 \rangle \quad (\text{by the prev ex}) \\
 &= \sum_{j \in \mathbb{N}} \langle A^* y_1, e_j \rangle \langle e_j, A^* y_2 \rangle \\
 &= \sum_{j \in \mathbb{N}} \langle A^* y_1, (e_j \otimes e_j^*) A^* y_2 \rangle \quad (\text{by the prev ex}) \\
 &= \left\langle A^* y_1, \left[\sum_{j \in \mathbb{N}} e_j \otimes e_j^* \right] A^* y_2 \right\rangle \\
 &= \langle A^* y_1, I_{H_1} A^* y_2 \rangle \quad (\text{by the prev part}) \\
 &= \langle A^* y_1, A^* y_2 \rangle \\
 &= \langle y_1, AA^* y_2 \rangle.
 \end{aligned}$$

A prev ex then implies that $AA^* = \sum_{j \in \mathbb{N}} (Ae_j) \otimes (Ae_j)^*$.

□

OLD STUFF

Note 10.6.0.13. This section assumes familiarity with the algebraic tensor product of two vector spaces. See section ??? of [?algebra] for details.

Definition 10.6.0.14. Let X, Y and Z be Banach spaces and $\phi \in L^2(X, Y; Z)$. Then (Z, ϕ) is said to be a **tensor product** of X with Y if

1. $\text{span } \phi(X \times Y)$ is dense in Z
2. for each Banach space W and $\psi \in L^2(X, Y; W)$, there exists a unique $\psi' \in L(Z, W)$ such that $\psi' \circ \phi = \psi$, i.e. such that the following diagram commutes:

$$\begin{array}{ccc} X \times Y & \xrightarrow{\phi} & Z \\ & \searrow \psi & \downarrow \psi' \\ & & W \end{array}$$

If (Z, ϕ) is a tensor product of X with Y . We often write $Z = X \otimes Y$ and for each $x \in X, y \in Y$, we often write $\phi(x, y) = x \otimes y$.

Exercise 10.6.0.15. Let X and Y be Banach spaces, $U \subset X$ and $V \subset Y$. Set $W = \{u \otimes v : u \in U \text{ and } v \in V\} \subset X \otimes Y$. If U and V are linearly independent, then W is linearly independent.

Hint: For $\phi \in X^*, \psi \in Y^*$, define $T \in L^2(X, Y; \mathbb{C})$ by $T(x, y) = \phi(x)\psi(y)$.

Proof. Let $w = \sum_{u \in U} \sum_{v \in V} \lambda_{u,v} u \otimes v$. Suppose that $w = 0$. Let $\phi \in X^*$ and $\psi \in Y^*$. Define $T \in L^2(X, Y; \mathbb{C})$ by $T(x, y) = \phi(x)\psi(y)$. By definition of the tensor product, there exists a unique $T' \in L(X \otimes Y, \mathbb{C})$ such that for each $x \in X$ and $y \in Y$, $T'(x \otimes y) = T(x, y)$. Then

$$\begin{aligned} 0 &= T'(w) \\ &= T'\left(\sum_{u \in U} \sum_{v \in V} \lambda_{u,v} u \otimes v\right) \\ &= \sum_{u \in U} \sum_{v \in V} \lambda_{u,v} T'(u \otimes v) \\ &= \sum_{u \in U} \sum_{v \in V} \lambda_{u,v} T(u, v) \\ &= \sum_{u \in U} \sum_{v \in V} \lambda_{u,v} \phi(u) \psi(v) \\ &= \phi\left(\sum_{u \in U} \sum_{v \in V} \lambda_{u,v} \psi(v) u\right) \end{aligned}$$

Since $\phi \in X^*$ is arbitrary, a previous exercise in the section on linear functionals implies that

$$\begin{aligned} 0 &= \sum_{u \in U} \sum_{v \in V} \lambda_{u,v} \psi(v) u \\ &= \sum_{u \in U} \left(\sum_{v \in V} \lambda_{u,v} \psi(v) \right) u \end{aligned}$$

Linear independence of U implies that for each $u \in U$,

$$\begin{aligned} 0 &= \sum_{v \in V} \lambda_{u,v} \psi(v) \\ &= \psi\left(\sum_{v \in V} \lambda_{u,v} v\right) \end{aligned}$$

Since $\psi \in Y^*$ is arbitrary, for each $u \in U$,

$$\sum_{v \in V} \lambda_{u,v} v = 0$$

Linear independence of V implies that for each $u \in U, v \in V, \lambda_{u,v} = 0$. Hence W is linearly independent. □

Exercise 10.6.0.16. Uniqueness:

Let X, Y and Z be Banach spaces and $\phi \in L^2(X, Y; Z)$. Suppose that (Z, ϕ) is a tensor product of X with Y . Then (Z, ϕ) is unique up to isomorphism.

Proof. Let W be a Banach space and $\psi \in L^2(X, Y; W)$. Suppose that (W, ψ) is a tensor product of X with Y . Since (Z, ϕ) is a tensor product of X with Y , there exists a unique $\psi' \in L(Z, W)$ such that $\psi' \circ \phi = \psi$. Since (W, ψ) is a tensor product of X with Y , there exists a unique $\phi' \in L(W, Z)$ such that $\phi' \circ \psi = \phi$. Thus the following diagram commutes:

$$\begin{array}{ccc} & & W \\ & \nearrow \psi & \downarrow \phi' \\ X \times Y & \xrightarrow{\phi} & Z \\ & \searrow \psi & \downarrow \psi' \\ & & W \end{array}$$

On the other hand, since (W, ψ) is a tensor product of X with Y , there exists a unique $\Psi \in L(W)$ such that $\Psi \circ \psi = \psi$. Thus the following diagram commutes:

$$\begin{array}{ccc} X \times Y & \xrightarrow{\psi} & W \\ & \searrow \psi & \downarrow \Psi \\ & & W \end{array}$$

Since $I_W \in L(W)$ and $I_W \circ \psi = \psi$, uniqueness of Ψ implies that $\Psi = I_W$. From the first diagram, we see that $\psi' \circ \phi'$ satisfies $(\psi' \circ \phi') \circ \psi = \psi$. Since $\psi' \circ \phi' \in L(W)$, uniqueness of Ψ implies that $\Psi = \psi' \circ \phi'$. Thus $\psi' \circ \phi' = I_W$.

Similarly, we could have initially considered the following diagram:

$$\begin{array}{ccc} & & Z \\ & \nearrow \phi & \downarrow \psi' \\ X \times Y & \xrightarrow{\psi} & W \\ & \searrow \phi & \downarrow \phi' \\ & & Z \end{array}$$

Playing a similar game, we could use the fact that there exists a unique $\Phi \in L(Z)$ such that $\Phi \circ \phi = \phi$ to obtain the following diagram:

$$\begin{array}{ccc} X \times Y & \xrightarrow{\phi} & Z \\ & \searrow \phi & \downarrow \Phi \\ & & Z \end{array}$$

As before, uniqueness enables us to conclude that $\phi' \circ \psi' = I_Z$. Thus ψ' and ϕ' are isomorphisms and $Z \cong W$. □

Note 10.6.0.17. The following definitions and exercises will cover the explicit construction of a tensor product of Banach spaces.

Definition 10.6.0.18. Let X and Y be Banach spaces. Define $X \otimes^{\text{alg}} Y = \text{span}\{x \otimes y : x \in X \text{ and } y \in Y\}$ to be the algebraic tensor product of X with Y (see section ??? of [?algebra] for details).

Exercise 10.6.0.19. Let X and Y be Banach spaces and $x \otimes y \in X \otimes^{\text{alg}} Y$. If for each $\phi \in X^*$ and $\psi \in Y^*$, $\phi \otimes \psi(x, y) = 0$, then $x \otimes y = 0$.

Proof. The previous section tells us that for each $\phi \in X^*$ and $\psi \in Y^*$, $\phi \otimes \psi(x, y) = 0$, then $x = 0$ or $y = 0$. This implies that $x \otimes y = 0$. □

Definition 10.6.0.20. The Projective Norm:

Define $\|\cdot\|_\pi : X \otimes^{\text{alg}} Y \rightarrow [0, \infty)$ by

$$\|u\|_\pi = \inf \left\{ \sum_{j=1}^n \|x_j\| \|y_j\| : (x_j)_{j=1}^n \subset X, (y_j)_{j=1}^n \subset Y \text{ and } u = \sum_{j=1}^n x_j \otimes y_j \right\}$$

Exercise 10.6.0.21. Let X and Y be Banach spaces. Then $\|\cdot\|_\pi : X \otimes^{\text{alg}} Y \rightarrow [0, \infty)$ is a norm on $X \otimes^{\text{alg}} Y$.

Proof.

- Let $\lambda \in \mathbb{C}$, $u \in X \otimes^{\text{alg}} Y$. If $\lambda = 0$, then $\lambda u = 0u = 0 \otimes 0$ and clearly $\|\lambda u\|_\pi = 0 = |\lambda| \|u\|_\pi$. Suppose that $\lambda \neq 0$. Let $\epsilon > 0$. Then there exist $(x_j)_{j=1}^n \subset X$ and $(y_j)_{j=1}^n \subset Y$ such that $u = \sum_{j=1}^n x_j \otimes y_j$ and $\sum_{j=1}^n \|x_j\| \|y_j\| < \|u\|_\pi + \epsilon/|\lambda|$. Then $\lambda u = \sum_{j=1}^n (\lambda x_j) \otimes y_j$. Therefore

$$\begin{aligned} \|\lambda u\|_\pi &\leq \sum_{j=1}^n \|\lambda x_j\| \|y_j\| \\ &\leq |\lambda| \sum_{j=1}^n \|x_j\| \|y_j\| \\ &< |\lambda| \left(\|u\|_\pi + \frac{\epsilon}{|\lambda|} \right) \\ &= |\lambda| \|u\|_\pi + \epsilon \end{aligned}$$

Since $\epsilon > 0$ is arbitrary, $\|\lambda u\|_\pi \leq |\lambda| \|u\|_\pi$. For the sake of contradiction, suppose that $\|\lambda u\|_\pi < |\lambda| \|u\|_\pi$. Then there exists $(x_j)_{j=1}^n \subset X$ and $(y_j)_{j=1}^n \subset Y$ such that $\lambda u = \sum_{j=1}^n x_j \otimes y_j$ and $\sum_{j=1}^n \|x_j\| \|y_j\| < |\lambda| \|u\|_\pi$. Hence $u = \sum_{j=1}^n (\lambda^{-1} x_j) \otimes y_j$. This implies that

$$\begin{aligned} \|u\|_\pi &\leq \sum_{j=1}^n \|\lambda^{-1} x_j\| \|y_j\| \\ &= |\lambda|^{-1} \sum_{j=1}^n \|x_j\| \|y_j\| \\ &< |\lambda|^{-1} |\lambda| \|u\|_\pi \\ &= \|u\|_\pi \end{aligned}$$

which is a contradiction. Therefore $\|\lambda u\|_\pi \geq |\lambda| \|u\|_\pi$ which implies that $\|\lambda u\|_\pi = |\lambda| \|u\|_\pi$

- Let $u, v \in X \otimes^{\text{alg}} Y$ and $\epsilon > 0$. Then there exist $(x_j)_{j=1}^n, (a_k)_{k=1}^m \subset X$ and $(y_j)_{j=1}^n, (b_k)_{k=1}^m \subset Y$ such that $u = \sum_{j=1}^n x_j \otimes y_j$, $v = \sum_{k=1}^m a_k \otimes b_k$, $\sum_{j=1}^n \|x_j\| \|y_j\| < \|u\|_\pi + \epsilon/2$ and $\sum_{k=1}^m \|a_k\| \|b_k\| < \|v\|_\pi + \epsilon/2$. Then $u + v = \sum_{j=1}^n x_j \otimes y_j + \sum_{k=1}^m a_k \otimes b_k$ which implies that

$$\begin{aligned} \|u + v\|_\pi &\leq \sum_{j=1}^n \|x_j\| \|y_j\| + \sum_{k=1}^m \|a_k\| \|b_k\| \\ &< \|u\|_\pi + \epsilon/2 + \|v\|_\pi + \epsilon/2 \\ &= \|u\|_\pi + \|v\|_\pi + \epsilon \end{aligned}$$

Since $\epsilon > 0$ is arbitrary, $\|u + v\|_\pi \leq \|u\|_\pi + \|v\|_\pi$.

- Let $u \in X \otimes^{\text{alg}} Y$. Suppose that $\|u\| = 0$. Let $\phi \in X^*$ and $\psi \in Y^*$ and $\epsilon > 0$. Then there exist $(x_j)_{j=1}^n \subset X$ and $(y_j)_{j=1}^n \subset Y$ such that $u = \sum_{j=1}^n x_j \otimes y_j$ and

$$\sum_{j=1}^n \|x_j\| \|y_j\| < \frac{\epsilon}{\|\phi\| \|\psi\| + 1}$$

Then

$$\begin{aligned}
 \sum_{j=1}^n |\phi \otimes \psi(x_j, y_j)| &= \sum_{j=1}^n |\phi(x_j)\psi(y_j)| \\
 &\leq \sum_{j=1}^n \|\phi\| \|x_j\| \|\psi\| \|y_j\| \\
 &= \|\phi\| \|\psi\| \sum_{j=1}^n \|x_j\| \|y_j\| \\
 &< \|\phi\| \|\psi\| \frac{\epsilon}{\|\phi\| \|\psi\| + 1} \\
 &< \epsilon
 \end{aligned}$$

Then for each $j \in \{1, \dots, n\}$, $|\phi \otimes \psi(x_j, y_j)| < \epsilon$. **FINISH!!!** Try using sequences and continuity and a common refinement of representation and averaging

□

Exercise 10.6.0.22. Existence:

Proof.

□

10.7 Rigged Hilbert Spaces

Introduce rigged hilbert spaces, talk about how schwarz space is a rigged hilbert space, then introduce gateaux derivative $\frac{dL}{d\phi(x)} := dL(\phi)(\delta(x))$ of L at ϕ in the direction of $\delta(x)$ (the dirac delta function), for functionals L on schwarz space, then talk about equations of motion and qft, see "quantum theory for gifted amateur" book section on functionals for details, try to formalize.

10.8 MISC, unitary transformations

Definition 10.8.0.1. Let H be a Hilbert space, $T \in GL(H)$ and $E \subset H$ a closed subspace. Then E is said to be **invariant under** T if $T(E) = E$.

Exercise 10.8.0.2. Let H be a Hilbert space, $U \in U(H)$ and $E \subset H$ a closed subspace. If E is invariant under U , then $U(E^\perp) = U(E)^\perp$.

Proof. Suppose that E is invariant under U . Let $y \in E$ and $x_0 \in E^\perp$. Since E is invariant under U , $U(E) = E$. Hence there exists $x \in E$ such that $Ux = y$. Since $x_0 \in E^\perp$ and $x \in E$, $\langle x_0, x \rangle = 0$. Since $U \in U(H)$,

$$\begin{aligned} \langle U(x_0), y \rangle &= \langle U(x_0), U(x) \rangle \\ &= \langle U(x_0), U(x) \rangle \\ &= \langle x_0, x \rangle \\ &= 0 \end{aligned}$$

Since $y \in E$ is arbitrary, we have that for each $y \in E$, $\langle U(x_0), y \rangle = 0$. Therefore $Ux_0 \in E^\perp$. Since $x_0 \in E^\perp$ is arbitrary, we have that for each $x_0 \in E^\perp$, $Ux_0 \in E^\perp$. Thus $U(E^\perp) \subset E^\perp$. For the sake of contradiction, suppose that $E^\perp \not\subset U(E^\perp)$. Then there exists $y \in E^\perp$ such that $y \notin U(E^\perp)$. Since $H = E \oplus E^\perp$ and **FINISH!!!** show $U(E \oplus E^\perp) = U(E) \oplus U(E^\perp)$. $\square$

Part III

Differentiation

Chapter 11

Differentiation

11.1 TODO

- Finish implicit and inverse function theorems

Note 11.1.0.1. Much of the material in this chapter discusses maps $f : X \rightarrow Y$ where X and Y are Banach spaces. It is often the case that a discussion requires the base fields of X and Y to agree or to both be real. We note that in these cases, every complex vector space is also a real vector space. In particular, if X is a finite dimensional complex vector space with dimension n , then X is a finite dimensional real vector space of dimension $2n$.

11.2 The Gateaux Derivative

Definition 11.2.0.1. Let X, Y be Banach spaces, $A \subset X$ open, $f : A \rightarrow Y$, $x_0 \in A$ and $x \in X$. Then f is said to be

1. **right-hand-differentiable at x_0 in the direction x** if the limit

$$\lim_{t \rightarrow 0^+} \frac{f(x_0 + tx) - f(x_0)}{t}$$

exists. If f is right-hand-differentiable at x_0 in the direction x , we define the **right-hand derivative** of f at x_0 in the direction x , denoted by $d^+f(x_0; x)$, to be the above limit.

2. **left-hand-differentiable at x_0 in the direction x** if the limit

$$\lim_{t \rightarrow 0^-} \frac{f(x_0 + tx) - f(x_0)}{t}$$

exists. If f is right-hand-differentiable at x_0 in the direction x , we define the **left-hand derivative** of f at x_0 in the direction x , denoted by $d^-f(x_0; x)$, to be the above limit.

3. **differentiable at x_0 in the direction x** if the limit

$$\lim_{t \rightarrow 0} \frac{f(x_0 + tx) - f(x_0)}{t}$$

exists. If f is differentiable at x_0 in the direction x , we define the **derivative** of f at x_0 in the direction x , denoted by $df(x_0; x)$, to be the above limit.

Exercise 11.2.0.2. Let X, Y be Banach spaces, $A \subset X$ open, $f : A \rightarrow \mathbb{R}$ and $x_0 \in A$. Then $df(x_0; 0) = 0$.

Proof. Clear. □

Definition 11.2.0.3. The Gateaux Derivative:

Let X, Y be Banach spaces, $A \subset X$ open, $f : A \rightarrow Y$ and $x_0 \in A$. Then f is said to be

1. **right-hand Gateaux differentiable at x_0** if for each $x \in X$, $d^+f(x_0; x)$ exists. We define the **right-hand Gateaux derivative** of f at x_0 , denoted $d^+f(x_0) : X \rightarrow Y$, to be

$$d^+f(x_0)(x) = d^+f(x_0; x)$$

2. **left-hand Gateaux differentiable at x_0** if for each $x \in X$, $d^-f(x_0; x)$ exists. We define the **left-hand Gateaux derivative** of f at x_0 , denoted $d^-f(x_0) : X \rightarrow Y$, to be

$$d^-f(x_0)(x) = d^-f(x_0; x)$$

3. **Gateaux differentiable at x_0** if for each $x \in X$, $df(x_0; x)$ exists. We define the **Gateaux derivative** of f at x_0 , denoted $df(x_0) : X \rightarrow Y$, to be

$$df(x_0)(x) = df(x_0; x)$$

Definition 11.2.0.4. Let Y be a Banach space, $A \subset \mathbb{R}$ open and $f : A \rightarrow Y$. Then f is said to be **Gateaux differentiable** if for each $x \in A$, f is Gateaux differentiable at x . If f is Gateaux differentiable, we define $df : A \rightarrow Y^X$ by $x_0 \mapsto df(x_0)$.

Exercise 11.2.0.5. Let X, Y be Banach spaces, $A \subset X$ open, $f, g : A \rightarrow Y$, $\lambda \in \mathbb{R}$ and $x_0 \in A$. If f, g are Gateaux differentiable at x_0 , then $f + \lambda g$ is Gateaux differentiable at x_0 and $d[f + \lambda g](x_0) = df(x_0) + \lambda dg(x_0)$.

Proof. Similar to the case of the derivative from Calc I. □

Exercise 11.2.0.6. Let X, Y be Banach spaces, $A \subset X$ open, $f : A \rightarrow Y$ and $x_0 \in A$. Suppose that f is Gateaux differentiable at x_0 . Then for each $\lambda \in \mathbb{R}$ and $x \in X$,

$$df(x_0)(\lambda x) = \lambda df(x_0)(x)$$

Proof. Let $\lambda \in \mathbb{R}$ and $x \in X$. Then

$$\begin{aligned} df(x_0)(\lambda x) &= \lim_{t \rightarrow 0} \frac{f(x_0 + t\lambda x) - f(x_0)}{t} \\ &= \lim_{t \rightarrow 0} \lambda \frac{f(x_0 + t\lambda x) - f(x_0)}{\lambda t} \\ &= \lambda \lim_{t \rightarrow 0} \frac{f(x_0 + t\lambda x) - f(x_0)}{\lambda t} \\ &= \lambda \lim_{t \rightarrow 0} \frac{f(x_0 + tx) - f(x_0)}{t} \\ &= \lambda df(x_0)(x) \end{aligned}$$

□

Exercise 11.2.0.7. Let X, Y be Banach spaces, $A \subset X$ open, $f : A \rightarrow Y$. If f is constant, then f is Gateaux differentiable and for each $x_0 \in A, x \in X$,

$$df(x_0)(x) = 0$$

Proof. Suppose that f is constant. Then there exists $c \in Y$ such that for each $x \in A, f(x) = c$. Let $x_0 \in A, x \in X$. Then

$$\begin{aligned} df(x_0)(x) &= \lim_{t \rightarrow 0} \frac{f(x_0 + tx) - f(x_0)}{t} \\ &= \lim_{t \rightarrow 0} \frac{c - c}{t} \\ &= 0 \end{aligned}$$

□

Exercise 11.2.0.8. Let X, Y be Banach spaces, $A \subset X$ open, $f : A \rightarrow Y$. If f is linear, then f is Gateaux differentiable and for each $x_0 \in A, x \in X$,

$$df(x_0)(x) = f(x)$$

Proof. Suppose that f is linear. Let $x_0 \in A, x \in X$. Then

$$\begin{aligned} df(x_0)(x) &= \lim_{t \rightarrow 0} \frac{f(x_0 + tx) - f(x_0)}{t} \\ &= \lim_{t \rightarrow 0} \frac{f(x_0) + tf(x) - f(x_0)}{t} \\ &= f(x) \end{aligned}$$

□

Exercise 11.2.0.9. There exist Banach spaces X, Y , and $f : X \rightarrow Y$ such that f is Gateaux differentiable and for each $x \in X, f$ is not continuous at x .

Hint: use Exercise 11.2.0.8

Proof. Set $X = C^1([0, 1])$ and $Y = C([0, 1])$. Equip both X and Y with the sup norm. Define $T : X \rightarrow Y$ by $Tf = f'$. Then Exercise 9.2.0.4 implies that T is not bounded. Since T is linear, Exercise 11.2.0.8 implies that T is Gateaux differentiable. Since T is not bounded, Exercise 9.2.0.6 implies that T is not continuous at 0. Then Exercise 9.2.0.6 tells us that for each $x \in X, T$ is not continuous at x . □

Exercise 11.2.0.10. Set $A = \{(x, y) \in \mathbb{R}^2 : y = -x^2 \text{ and } x \neq 0\}$. Define $f : \mathbb{R}^2 \setminus A \rightarrow \mathbb{R}$ by

$$f(x, y) = \begin{cases} 0 & (x, y) = (0, 0) \\ \frac{x^4 y}{x^6 + y^3} & \text{otherwise} \end{cases}$$

Then f is Gateaux differentiable at $(0, 0)$ and f is not continuous at $(0, 0)$.

Hint: Consider the set $B = \{(x, x^2 : x \in \mathbb{R})\} \subset \mathbb{R}^2 \setminus A$.

Proof. □

Exercise 11.2.0.11. Let Y be a Banach space, $A \subset \mathbb{R}$ open, $f : A \rightarrow Y$ and $x_0 \in A$. Suppose that f is Gateaux differentiable at x_0 . Then $df(x_0) \in L(\mathbb{R}, Y)$.

Proof.

- Exercise 11.2.0.6 implies that for each $x, y, \lambda \in \mathbb{R}$,

$$\begin{aligned} df(x_0)(x + \lambda y) &= df(x_0)((x + \lambda y)1) \\ &= (x + \lambda y)df(x_0)(1) \\ &= xdf(x_0)(1) + \lambda ydf(x_0)(1) \\ &= df(x_0)(x) + \lambda df(x_0)(y) \end{aligned}$$

So $df(x_0) \in \mathcal{L}(\mathbb{R}, Y)$.

- Since for each $x \in \mathbb{R}$,

$$\begin{aligned} \|df(x_0)(x)\| &= \|xdf(x_0)(1)\| \\ &= |x|\|df(x_0)(1)\| \end{aligned}$$

We have that $df(x_0) : \mathbb{R} \rightarrow Y$ is bounded with $\|df(x_0)\| \leq \|df(x_0)(1)\|$. So $df(x_0)$ is continuous.

Thus $df(x_0) \in L(\mathbb{R}, Y)$. □

Exercise 11.2.0.12. Let X be a Banach space, $A \subset X$ open, $f : A \rightarrow \mathbb{R}$ and $x_0 \in A$. If f is Gateaux differentiable at x_0 and f has a local extremum at x_0 , then $df(x_0) = 0$.

Proof. Suppose that f is Gateaux differentiable at x_0 and f has a local minimum point at x_0 . Then there exists $\delta > 0$ such that $B(x_0, \delta) \subset A$ and for each $y \in B(x_0, \delta)$, $f(x_0) \leq f(y)$.

For the sake of contradiction, suppose that $df(x_0) \neq 0$. Then there exists $x \in X$ such that $x \neq 0$ and $df(x_0)(x) \neq 0$.

First, suppose that $df(x_0)(x) < 0$. Choose $\epsilon = -df(x_0)(x) > 0$. Then there exists $t_0 > 0$ such that for each $t \in B^*(0, t_0)$, $x_0 + tx \in B(x_0, \delta)$ and

$$\left| \frac{f(x_0 + tx) - f(x_0)}{t} - df(x_0)(x) \right| < \epsilon$$

This implies that for each $t \in B^*(0, t_0)$,

$$\begin{aligned} \frac{f(x_0 + tx) - f(x_0)}{t} &< \epsilon + df(x_0)(x) \\ &= 0 \end{aligned}$$

and hence $f(x_0 + tx) < f(x_0)$, which is a contradiction.

Now, suppose that $df(x_0)(x) > 0$. Then

$$\begin{aligned} df(x_0)(-x) &= -df(x_0)(x) \\ &< 0 \end{aligned}$$

Similarly to above, this implies that there exists $t_0 > 0$ such that for each $t \in B^*(0, t_0)$, $x_0 - tx \in B(x_0, \delta)$ and $f(x_0 - tx) < f(x_0)$ which is a contradiction. So $df(x_0)(x) = 0$ and $df(x_0) = 0$.

If f has a local maximum at x_0 , then $-f$ has a local minimum point at x_0 . Then

$$\begin{aligned} df(x_0) &= -d[-f](x_0) \\ &= -0 \\ &= 0 \end{aligned}$$

□

Exercise 11.2.0.13. Let X, Y, Z be Banach spaces, $A \subset X$ open, $B \subset Y$ open, $f : A \rightarrow Y$, $g : B \rightarrow Z$ and $x_0 \in A$. Suppose that f is affine. If g is Gateaux differentiable at $f(x_0)$, then $g \circ f$ is Gateaux differentiable at x_0 and

$$d(g \circ f)(x_0)(x) = dg(f(x_0))(df(x_0)(x))$$

(DEFINE AFFINE SOMEWHERE)

Proof. Suppose that g is Gateaux differentiable at $f(x_0)$. Since f is affine, there exists $h : A \rightarrow Y$ and $c \in Y$ such that h is linear and $f = h + c$. Then

$$\begin{aligned} df(x_0) &= dh(x_0) \\ &= h \end{aligned}$$

Let $x \in X$. Choose $\delta > 0$ such that for each $t \in B(0, \delta) \subset \mathbb{R}$, $f(x_0) + th(x) \in B$. Then for each $t \in B^*(0, \delta)$,

$$\begin{aligned} g \circ f(x_0 + tx) &= g\left(f(x_0) + t \frac{f(x_0 + tx) - f(x_0)}{t}\right) \\ &= g(f(x_0) + th(x)) \end{aligned}$$

This implies that

$$\begin{aligned} d(g \circ f)(x_0) &= \lim_{t \rightarrow 0} \frac{g \circ f(x_0 + tx) - g(f(x_0))}{t} \\ &= \lim_{t \rightarrow 0} \frac{g(f(x_0) + th(x)) - g(f(x_0))}{t} \\ &= dg(f(x_0))(h(x)) \\ &= dg(f(x_0))(df(x_0)(x)) \end{aligned}$$

□

11.3 The Frechet Derivative

Exercise 11.3.0.1. Let X, Y be a normed vector spaces and $\phi : X \rightarrow Y$ linear. If $\phi(h) = o(\|h\|)$ as $h \rightarrow 0$, then $\phi = 0$.

Proof. Let $h_0 \in X$. If $h_0 = 0$, then $\phi(h_0) = 0$. Suppose that $h_0 \neq 0$. Define $(h_n)_{n \in \mathbb{N}} \subset X$ by

$$h_n = \frac{h_0}{n}$$

Then $h_n \rightarrow 0$. By continuity of ϕ and our initial assumption we have that

$$\begin{aligned} \|h_0\|^{-1}\phi(h_0) &= \phi\left(\frac{h_0}{\|h_0\|}\right) \\ &= \phi\left(\frac{h_n}{\|h_n\|}\right) \\ &= \frac{\phi(h_n)}{\|h_n\|} \\ &\rightarrow 0 \end{aligned}$$

which implies that $\|h_0\|^{-1}\phi(h_0) = 0$. So $\phi(h_0) = 0$ and hence $\phi = 0$. □

Exercise 11.3.0.2. Let X, Y be a normed vector spaces, $A \subset X$ open, $f : A \rightarrow Y$ and $x_0 \in A$. Suppose that there exists $\phi : X \rightarrow Y$ such that ϕ is linear and

$$f(x_0 + h) = f(x_0) + \phi(h) + o(\|h\|) \quad \text{as } h \rightarrow 0$$

then ϕ is unique.

Proof. Suppose that there exists $\psi : X \rightarrow Y$ such that ψ is linear and such that

$$f(x_0 + h) = f(x_0) + \psi(h) + o(\|h\|) \quad \text{as } h \rightarrow 0$$

Then $\phi(h) - \psi(h) = o(h)$. Since $\phi - \psi$ is linear, the previous exercise implies that $\phi = \psi$. □

Note 11.3.0.3. Recall that for Banach spaces X and Y ,

$$\text{cur} : L^n(X; Y) \rightarrow L(X; L(X; \cdots; L(X; Y)) \cdots)$$

is an isometric isomorphism and we may identify $L(X; L(X; \cdots; L(X; Y)) \cdots)$ as $L^n(X; Y)$.

Definition 11.3.0.4. Frechet Derivative:

Let X, Y be a Banach spaces, $A \subset X$ open, $f : A \rightarrow Y$ and $x_0 \in A$.

1. • Then f is said to be **Frechet differentiable at x_0** if there exists $\phi \in L(X; Y)$ such that,

$$f(x_0 + h) = f(x_0) + \phi(h) + o(\|h\|) \quad \text{as } h \rightarrow 0$$

- If f is Frechet differentiable at x_0 , we define the **Frechet derivative of f at x_0** , denoted $Df(x_0)$, to be the unique $\phi \in L(X; Y)$ such that,

$$f(x_0 + h) = f(x_0) + \phi(h) + o(\|h\|) \quad \text{as } h \rightarrow 0.$$

- We say that f is **Frechet differentiable** if for each $x \in A$, f is Frechet differentiable at x .
- If f is Frechet differentiable, we define the **Frechet derivative of f** , denoted $Df : A \rightarrow L(X; Y)$, by $x \mapsto D^{(1)}f(x)$.

2. Continuing inductively, we set $D^0 f = f$ and for $n \geq 2$,

- f is said to be **n -th order Frechet differentiable at x_0** if f is $(n - 1)$ -th order Frechet differentiable and $D^{n-1}f$ is Frechet differentiable at x_0 .
- If f is n -th order Frechet differentiable at x_0 , we define $D^n f(x_0) \in L^n(X; Y)$ by

$$D^n f(x_0) = D[D^{n-1}f](x_0)$$

- We say that f is **n -th order Frechet differentiable** if f is $(n-1)$ -th order Frechet differentiable and for each $x \in A$, $D^{n-1}f$ is Frechet differentiable at x .
- If f is n -th order Frechet differentiable, we define the **n -th order Frechet derivative of f** , denoted $D^n f : A \rightarrow L^n(X; Y)$, by $x \mapsto D^n f(x)$

3. If f is n -th order differentiable, then f is said to be **continuously n -th order differentiable** if $D^n f$ is continuous. We define

$$C^n(A, Y) = \{f : A \rightarrow Y : f \text{ is continuously } n\text{-th order differentiable}\}$$

Exercise 11.3.0.5. Let X, Y be a banach spaces, $A \subset X$ open, $f, g : A \rightarrow Y$, $\lambda \in \mathbb{R}$ and $x_0 \in A$. If f and g are Frechet differentiable at x_0 , then $f + \lambda g$ is Frechet differentiable at x_0 and $D(f + \lambda g)(x_0) = Df(x_0) + \lambda Dg(x_0)$.

Proof. Suppose that f and g are Frechet differentiable at x_0 . Then

$$f(x_0 + h) = f(x_0) + Df(x_0)(h) + o(\|h\|) \quad \text{as } h \rightarrow 0$$

and

$$g(x_0 + h) = g(x_0) + Dg(x_0)(h) + o(\|h\|) \quad \text{as } h \rightarrow 0$$

This implies that

$$\begin{aligned} (f + \lambda g)(x_0 + h) &= f(x_0 + h) + \lambda g(x_0 + h) \\ &= f(x_0) + Df(x_0)(h) + o(\|h\|) + \lambda g(x_0) + \lambda Dg(x_0)(h) + o(\|h\|) \\ &= (f + \lambda g)(x_0) + [Df(x_0) + \lambda Dg(x_0)](h) + o(\|h\|) \quad \text{as } h \rightarrow 0 \end{aligned}$$

Since $Df(x_0) + \lambda Dg(x_0) \in L(X; Y)$, $f + \lambda g$ is Frechet differentiable at x_0 and $D(f + \lambda g)(x_0) = Df(x_0) + \lambda Dg(x_0)$. $\square$

Exercise 11.3.0.6. Let X, Y be a banach spaces, $A \subset X$ open, $f : A \rightarrow Y$ and $x_0 \in A$. If f is Frechet differentiable at x_0 , then f is continuous at x_0 .

Proof. Suppose that f is Frechet differentiable at x_0 . Then $f(x) - f(x_0) = Df(x_0)(x - x_0) + o(\|x - x_0\|)$ as $x \rightarrow x_0$. Hence $\|f(x) - f(x_0)\| \leq \|Df(x_0)\| \|x - x_0\| + o(\|x - x_0\|)$ as $x \rightarrow x_0$. This implies that $f(x) \rightarrow f(x_0)$ as $x \rightarrow x_0$ and therefore f is continuous at x_0 . $\square$

Exercise 11.3.0.7. Let X, Y be a banach spaces, $A \subset X$ open, $f : A \rightarrow Y$ and $x_0 \in A$. If f is Frechet differentiable at x_0 , then f is Gateaux differentiable at x_0 and $df(x_0) = Df(x_0)$.

Proof. Suppose that f is Frechet differentiable at x_0 . Then $f(x_0 + h) = f(x_0) + Df(x_0)(h) + o(\|h\|)$ as $h \rightarrow 0$. Let $x \in X$. Exercise 17.3.0.5 **ref little o ex about ϕ linear and $\phi(h) = o(h)$ implying $\phi(tx) = o(t)$** implies that $f(x_0 + tx) - f(x_0) = tDf(x_0)(x) + o(t)$ as $t \rightarrow 0$. **?? ref ex in asymptotic notation appendices about $g > 0$ around x_0 implies that $f = o(g)$ iff $\lim \|f\|/\|g\| = 0$.**

$$\begin{aligned} df(x_0; x) &= \lim_{t \rightarrow 0} \frac{f(x_0 + tx) - f(x_0)}{t} \\ &= \lim_{t \rightarrow 0} \frac{f(x_0) + tDf(x_0)(x) + o(t) - f(x_0)}{t} \quad (\text{by Exercise .2.0.7}) \\ &= \lim_{t \rightarrow 0} \left[Df(x_0)(x) + \frac{o(t)}{t} \right] \\ &= Df(x_0)(x) + \lim_{t \rightarrow 0} \left[\frac{o(t)}{t} \right] \\ &= Df(x_0)(x) \quad (\text{by Exercise .2.0.5}) \end{aligned}$$

Hence f is differentiable at x_0 in the direction x . Since $x \in X$ is arbitrary, f is Gateaux differentiable at x_0 and $df(x_0) = Df(x_0)$. $\square$

Exercise 11.3.0.8. Let X be a Banach space, $A \subset X$ open, $f : A \rightarrow \mathbb{R}$ and $x_0 \in A$. If f is Frechet differentiable at x_0 and f has a local extremum at x_0 , then $Df(x_0) = 0$.

Proof. Suppose that f is Frechet differentiable at x_0 and f has a local extremum at x_0 . Two previous exercises imply that f is Gateaux differentiable at x_0 and

$$\begin{aligned} Df(x_0) &= df(x_0) \\ &= 0 \end{aligned}$$

□

Definition 11.3.0.9. Let X, Y be a Banach spaces, $A \subset X$ open, $f : A \rightarrow Y$ and $x_0 \in A$. Suppose that f is Frechet differentiable at x_0 . Define $R_f(x_0) : A - x_0 \rightarrow Y$ by

$$R_f(x_0)(h) = f(x_0 + h) - f(x_0) - Df(x_0)(h)$$

move somewhere else

Exercise 11.3.0.10. Let X, Y be a Banach spaces, $A \subset X$ open, $f : A \rightarrow Y$ and $x_0 \in A$. If f is Frechet differentiable at x_0 , then

$$f(x_0 + h) - f(x_0) = O(\|h\|) \text{ as } h \rightarrow 0$$

Proof. Suppose that f is Frechet differentiable at x_0 . Then $R_f(h) = o(\|h\|)$ as $h \rightarrow 0$. Hence there exists $\delta > 0$ such that $B(0, \delta) \subset A - x_0$ and for each $h \in B(0, \delta)$, $\|R_f(h)\| \leq \|h\|$. Hence for each $h \in B(0, \delta)$

$$\begin{aligned} \|f(x_0 + h) - f(x_0)\| &= \|Df(x_0)(h) + R_f(x_0)(h)\| \\ &\leq \|Df(x_0)(h)\| + \|R_f(x_0)(h)\| \\ &\leq \|Df(x_0)\| \|h\| + \|h\| \\ &= (\|Df(x_0)\| + 1) \|h\| \end{aligned}$$

□

Exercise 11.3.0.11. Chain Rule:

Let X, Y, Z be a Banach spaces, $A \subset X$ open, $B \subset Y$ open, $f : A \rightarrow Y$, $g : B \rightarrow Z$ and $x_0 \in A$. Suppose that $f(x_0) \in B$. If f is Frechet differentiable at x_0 and g is Frechet differentiable at $f(x_0)$, then $g \circ f$ is Frechet differentiable at x_0 and

$$D(g \circ f)(x_0) = Dg(f(x_0)) \circ Df(x_0)$$

Proof. Suppose that f is Frechet differentiable at x_0 and g is Frechet differentiable at $f(x_0)$.

- The previous exercise implies that there exists $\delta^* > 0$ and $K > 0$ such that for each $h \in B(0, \delta^*)$, $\|f(x_0 + h) - f(x_0)\| \leq K\|h\|$. Let $\epsilon > 0$. Since $R_g(f(x_0))(h') = o(\|h'\|)$ as $h' \rightarrow 0$, there exists $\delta' > 0$ such that for each $h' \in B(0, \delta')$, $\|R_g(f(x_0))(h')\| \leq \frac{\epsilon}{K}\|h'\|$. Choose $\delta = \min(\delta'/K, \delta^*)$. Let $h \in B(0, \delta)$. Then

$$\begin{aligned} \|f(x_0 + h) - f(x_0)\| &\leq K\|h\| \\ &< \delta' \end{aligned}$$

This implies that

$$\begin{aligned} \|R_g(f(x_0))(f(x_0 + h) - f(x_0))\| &\leq \frac{\epsilon}{K} \|f(x_0 + h) - f(x_0)\| \\ &\leq \frac{\epsilon}{K} K \|h\| \\ &\leq \epsilon \|h\| \end{aligned}$$

So $R_g(f(x_0))(f(x_0 + h) - f(x_0)) = o(\|h\|)$ as $h \rightarrow 0$.

- Since $\|Dg(f(x_0))(R_f(x_0)(h))\| \leq \|Dg(f(x_0))\| \|R_f(x_0)(h)\|$ and $R_f(x_0)(h) = o(h)$ as $h \rightarrow 0$, we have that $Dg(f(x_0))(R_f(x_0)(h)) = o(h)$ as $h \rightarrow 0$.

- Combining the previous two observations, we have that $Dg(f(x_0))(R_f(x_0)(h)) + R_g(f(x_0))(f(x_0+h) - f(x_0)) = o(\|h\|)$ as $h \rightarrow 0$.
- All together, we obtain

$$\begin{aligned}
g \circ f(x_0 + h) &= g(f(x_0)) + f(x_0 + h) - f(x_0) \\
&= g(f(x_0)) + Dg(f(x_0))(f(x_0 + h) - f(x_0)) + R_g(f(x_0))(f(x_0 + h) - f(x_0)) \\
&= g(f(x_0)) + Dg(f(x_0))(Df(x_0)(h) + R_f(x_0)(h)) \\
&\quad + R_g(f(x_0))(f(x_0 + h) - f(x_0)) \\
&= g(f(x_0)) + Dg(f(x_0))(Df(x_0)(h)) + Dg(f(x_0))(R_f(x_0)(h)) \\
&\quad + R_g(f(x_0))(f(x_0 + h) - f(x_0)) \\
&= g \circ f(x_0) + Dg(f(x_0)) \circ Df(x_0)(h) + o(\|h\|) \text{ as } h \rightarrow 0
\end{aligned}$$

So $g \circ f$ is Frechet differentiable at x_0 and $D(g \circ f)(x_0) = Dg(f(x_0)) \circ Df(x_0)$. □

Exercise 11.3.0.12. Let Y be a Banach space, $A \subset \mathbb{R}$ open and $f : A \rightarrow Y$. Then f is Gateaux differentiable iff f is Frechet differentiable.

Proof. Suppose that f is Gateaux differentiable. Let $x_0 \in A$. Exercise 11.2.0.11 implies that $df(x_0) \in L(\mathbb{R}, Y)$. By defintion,

$$\lim_{h \rightarrow 0} \left\| \frac{f(x_0 + h) - f(x_0)}{h} - df(x_0)(1) \right\| = 0$$

This is equivalent to saying that

$$f(x_0 + h) = f(x_0) + df(x_0)(h) + o(|h|) \quad \text{as } h \rightarrow 0$$

So f is Frechet differentiable at x_0 and $Df(x_0) = df(x_0)$. □

Exercise 11.3.0.13. Let X be a Banach spaces, $U_F, U_G \in \mathcal{T}_X$, $F : U_F \rightarrow Y$,

Exercise 11.3.0.14. Let X, Y be Banach spaces, $U_F, U_G \in \mathcal{T}_X$, $F : U_F \rightarrow Y$, $G : U_G \rightarrow Y$ and $a \in U_1 \cap U_2$. If there exists $U \in \mathcal{T}_X$ such that $a \in U$, $U \subset U_1 \cap U_2$ and $F|_U = G|_U$, then $DF(a) = DG(a)$.

Proof. Suppose that there exists $U \in \mathcal{T}_X$ such that $a \in U$, $U \subset U_1 \cap U_2$ and $F|_U = G|_U$. Then (make a prev ex that for each $x \in X$, $D\iota_U(x) = \text{id}_X$)

$$\begin{aligned}
DF(a) &= DF(a) \circ \text{id}_X &= DF(a) \circ D\iota_U(a) \\
&= D(F \circ \iota_U)(a) \\
&= DF|_U(a) \\
&= DG|_U(a) \\
&= D(G \circ \iota_U)(a) \\
&= DG(a) \circ D\iota_U(a) \\
&= DF(a) \circ \text{id}_X \\
&= DG(a).
\end{aligned}$$
□

11.4 The Calc I Derivative

Definition 11.4.0.1. Calc I Derivative:

Let Y be a Banach space, $A \subset \mathbb{R}$ or $\mathbb{C}$ open, $f : A \rightarrow Y$ and $x_0 \in A$.

1. • If f is Frechet differentiable at x_0 , we define the **calc I derivative of f at x_0** , denoted

$$f'(x_0) \text{ or } \frac{df}{dt}(x_0)$$

by

$$\begin{aligned} f'(x_0) &:= \lim_{t \rightarrow 0} \frac{f(x_0 + t) - f(x_0)}{t} \\ &= df(x_0)(1) \\ &= Df(x_0)(1) \end{aligned}$$

- If f is Frechet differentiable, we define $f' : A \rightarrow Y$ by $x \mapsto f'(x)$.
2. Continuing inductively, we set $f^{(0)} = f$ and for $n \geq 1$,
 - if $f^{(n-1)}$ is Frechet differentiable at x_0 , we define the **(n) -th order calc I derivative of f at x_0** , denoted $f^{(n)}(x_0)$, by

$$f^{(n)}(x_0) = [f^{(n-1)}]'(x_0)$$

- if $f^{(n-1)}$ is Frechet differentiable, we define $f^{(n)} : A \rightarrow Y$ by

$$f^{(n)} = [f^{(n-1)}]'$$

Exercise 11.4.0.2. Let Y be a Banach space, $A \subset \mathbb{R}$ open and $f : A \rightarrow Y$. If f is n -th order Frechet differentiable, then for each $x_0 \in A$ and $k \in \{1, \dots, n\}$,

$$f^{(k)}(x_0) = D^k f(x_0)(1^{\oplus k})$$

Proof. Let $x_0 \in A$. We proceed by induction. The base case is true by definition. Let $k \in \{1, \dots, n\}$. Suppose the claim is true for $k-1$. Then

$$f^{(k-1)}(x_0) = D^{k-1} f(x_0)(1^{\oplus(k-1)})$$

Since f is n -th order Frechet differentiable,

$$D^{k-1} f(x_0 + h) = D^{k-1} f(x_0) + D^k f(x_0)(h) + o(\|h\|) \quad \text{as } h \rightarrow 0$$

This implies that

$$\begin{aligned} f^{(k-1)}(x_0 + h) &= D^{k-1} f(x_0 + h)(1^{\oplus(k-1)}) \\ &= D^{k-1} f(x_0)(1^{\oplus(k-1)}) + D^k f(x_0)(h)(1^{\oplus(k-1)}) + o(\|h\|) \quad \text{as } h \rightarrow 0 \end{aligned}$$

Therefore for each $h \in \mathbb{R}$,

$$Df^{(k-1)}(x_0)(h) = D^k f(x_0)(h)(1^{\oplus(k-1)})$$

and by definition,

$$\begin{aligned} f^{(k)}(x_0) &= [f^{(k-1)}]'(x_0) \\ &= Df^{(k-1)}(x_0)(1) \\ &= D^k f(x_0)(1^{\oplus k}) \end{aligned}$$

□

Exercise 11.4.0.3. Let X, Y be Banach spaces, $A \subset X$ open, $f \in C^n(A, Y)$, $x_0 \in A$, and $h \in X$. Suppose that $\{x_0 + th : t \in [0, 1]\} \subset A$. Define and $g : (0, 1) \rightarrow Y$ by

$$g(t) = f(x_0 + th)$$

Then for each $k \in \{1, \dots, n\}$ and $t \in (0, 1)$,

$$g^{(k)}(t) = D^k f(x_0 + th)(h^{\oplus k})$$

Proof. We proceed by induction. It is straightforward to show that the claim is true for $k = 1$.

Let $k \in \{1, \dots, n\}$. Suppose that $g^{(k-1)}(t) = D^{k-1} f(x_0 + th)(h^{\oplus(k-1)})$. Since $f \in C^k(A, Y)$,

$$D^{k-1} f(x_0 + s_0 h + th) = D^{k-1} f(x_0 + s_0 h) + D^k f(x_0 + s_0 h)(th) + o(\|t\|) \quad \text{as } t \rightarrow 0$$

The previous exercise implies that

$$\begin{aligned} g^{(k-1)}(s_0 + t) &= D^{k-1} g(s_0 + t)(1^{\oplus(k-1)}) \\ &= D^{k-1} f(x_0 + s_0 h + th)(h^{\oplus(k-1)}) \\ &= D^{k-1} f(x_0 + s_0 h)(h^{\oplus(k-1)}) + D^k f(x_0 + s_0 h)(th)(h^{\oplus(k-1)}) + o(\|t\|) \quad \text{as } t \rightarrow 0 \end{aligned}$$

Hence

$$Dg^{(k-1)}(s_0)(t) = D^k f(x_0 + s_0 h)(th)(h^{\oplus(k-1)})$$

and

$$\begin{aligned} g^{(k)}(t) &= Dg^{(k-1)}(t)(1) \\ &= D^k f(x_0 + th)(h^{\oplus k}) \end{aligned}$$

□

11.5 Mean Value Theorem

Exercise 11.5.0.1. Let X be a Banach space, $A \subset X$ open and convex, and $f : A \rightarrow \mathbb{R}$. If f is continuous and Gateaux differentiable, then for each $x, y \in A$, there exists $t^* \in (0, 1)$ such that $f(x) - f(y) = df(t^*x + (1 - t^*)y)(x - y)$.

Proof. Suppose that f is continuous and Gateaux differentiable. Let $x, y \in A$. Define $h : [0, 1] \rightarrow X$ by $h(t) = tx + (1 - t)y$. Set $g = f \circ h : [0, 1] \rightarrow \mathbb{R}$. Then g is continuous on $[0, 1]$ and Exercise 11.2.0.13 implies that g is Gateaux differentiable on $(0, 1)$. Then Exercise 11.3.0.12 Exercise 11.2.0.13 and the mean value theorem implies that there exists $t^* \in (0, 1)$ such that

$$\begin{aligned} f(x) - f(y) &= g(1) - g(0) \\ &= g'(t^*) \\ &= dg(t^*)(1) \\ &= df(h(t^*))(dh(t^*)(1)) \\ &= df(h(t^*))(h'(t^*)) \\ &= df(t^*x + (1 - t^*)y)(x - y) \end{aligned}$$

□

Exercise 11.5.0.2. Let X be a Banach space, $A \subset X$ open and convex, and $f : A \rightarrow \mathbb{R}$. If f is Frechet differentiable, then for each $x, y \in A$, there exists $t^* \in (0, 1)$ such that $f(x) - f(y) = Df(t^*x + (1 - t^*)y)(x - y)$.

Proof. Suppose that f is Frechet differentiable. Then f is continuous and Gateaux differentiable. Now apply the previous exercise. □

Exercise 11.5.0.3. Mean Value Theorem:

Let X, Y be Banach spaces, $A \subset X$ open and convex and $f : A \rightarrow Y$. If f is Frechet differentiable, then for each $x, y \in A$, there exists $t^* \in (0, 1)$ such that

$$\|f(x) - f(y)\| \leq \|Df(t^*x + (1 - t^*)y)\| \|x - y\|$$

Hint: For $x, y \in A$ with $f(x) \neq f(y)$, using a Hahn-Banach argument, find $\lambda \in Y^*$ such that $\|\lambda\| = 1$ and $\lambda(f(x) - f(y)) = \|f(x) - f(y)\|$.

Proof. Suppose that f is Frechet differentiable. Let $x, y \in A$. The claim is clearly true when $f(x) = f(y)$. Suppose that $f(x) \neq f(y)$. An exercise in the section on linear functionals implies that there exists $\lambda \in Y^*$ such that $\lambda(f(x) - f(y)) = \|f(x) - f(y)\|$ and $\|\lambda\| = 1$. Define $g : [0, 1] \rightarrow \mathbb{R}$ by

$$g(t) = \lambda(f(tx + (1 - t)y))$$

Then g is continuous and (Frechet) differentiable on $(0, 1)$ with

$$Dg(t)(h) = \lambda \circ Df(tx + (1 - t)y)((x - y)h)$$

which implies that

$$\begin{aligned} g'(t) &= Dg(t)(1) \\ &= \lambda \circ Df(tx + (1 - t)y)((x - y)) \end{aligned}$$

The mean value theorem implies that there exists $t^* \in (0, 1)$ such that

$$\begin{aligned} \|f(x) - f(y)\| &= \lambda(f(x) - f(y)) \\ &= g(1) - g(0) \\ &= g'(t^*) \\ &= \lambda \circ Df(t^*x + (1 - t^*)y)((x - y)) \end{aligned}$$

Taking absolute values, we see that

$$\begin{aligned} \|f(x) - f(y)\| &= |\lambda \circ Df(t^*x + (1 - t^*)y)((x - y))| \\ &\leq \|\lambda\| \|Df(t^*x + (1 - t^*)y)\| \|x - y\| \\ &\leq \|Df(t^*x + (1 - t^*)y)\| \|x - y\| \end{aligned}$$

□

Exercise 11.5.0.4. Let X, Y be Banach spaces, $A \subset X$ open and convex and $f : A \rightarrow Y$. Suppose that f is Frechet differentiable. If for each $x \in A$, $Df(x) = 0$, then f is constant.

Proof. Suppose that for each $x \in A$, $Df(x) = 0$. Let $x, y \in A$. Then the mean value theorem implies that there exists $t \in (0, 1)$ such that

$$\begin{aligned}\|f(x) - f(y)\| &\leq \|Df(tx + (1-t)y)\| \|x - y\| \\ &= 0\end{aligned}$$

So $f(x) = f(y)$. □

Exercise 11.5.0.5. Let X, Y be Banach spaces, $A \subset X$ open and convex and $f, g : A \rightarrow Y$. Suppose that f and g are Frechet differentiable. If $Df = Dg$, then there exists $c \in Y$ such that $f = g + c$.

Proof. Suppose that $Df = Dg$. Then $D(f - g) = 0$ and the previous exercise implies that $f - g$ is constant. □

Exercise 11.5.0.6. Let X, Y be a Banach spaces, $A \subset \mathbb{R}$ open and $f : A \rightarrow Y$. Suppose that f is Frechet differentiable. Then $f' \in C(A, Y)$ iff $f \in C^1(A, Y)$.

Proof. Suppose that $f' \in C(A, Y)$. Let $x, y \in A$ and $h \in \mathbb{R}$. Then

$$\begin{aligned}\|(Df(x) - Df(y))(h)\| &= \|Df(x)(h) - Df(y)(h)\| \\ &= \|hf'(x) - hf'(y)\| \\ &= \|h(f'(x) - f'(y))\| \\ &= \|f'(x) - f'(y)\| |h|\end{aligned}$$

So $\|Df(x) - Df(y)\| \leq \|f'(x) - f'(y)\|$. Hence continuity of f' implies continuity of Df and $f \in C^1(A, Y)$. Conversely, suppose that $f \in C^1(A, Y)$. Let $x, y \in A$. Then

$$\begin{aligned}\|f'(x) - f'(y)\| &= \|Df(x)(1) - Df(y)(1)\| \\ &= \|(Df(x) - Df(y))(1)\| \\ &\leq \|Df(x) - Df(y)\|\end{aligned}$$

Hence continuity of Df implies continuity of f' and $f' \in C(A, Y)$. □

Exercise 11.5.0.7. Let X, Y be Banach spaces, $A \subset X$ open and convex and $f : A \rightarrow Y$. Suppose that f is Frechet differentiable. Then f is Lipschitz iff Df is bounded.

Proof. Suppose that f is Lipschitz. Then there exists $M > 0$ such that for each $x, y \in A$, $\|f(y) - f(x)\| \leq M\|y - x\|$. Let $x \in A$ and $h \in X$. Suppose that $\|h\| = 1$. Since $f(x + th) = f(x) + Df(x)(th) + o(|t|)$ as $t \rightarrow 0$, we have that

$$\begin{aligned}|t|\|Df(x)(h)\| &= \|Df(x)(th)\| \\ &\leq \|f(x + th) - f(x)\| + o(|t|) \text{ as } t \rightarrow 0 \\ &\leq M\|th\| + o(|t|) \text{ as } t \rightarrow 0 \\ &= M|t| + o(|t|) \text{ as } t \rightarrow 0\end{aligned}$$

Hence $\|Df(x)(h)\| \leq M + o(1)$ as $t \rightarrow 0$ which implies that $\|Df(x)(h)\| \leq M$. Thus

$$\begin{aligned}\|Df(x)\| &= \sup\{\|Df(x)(h)\| : h \in X \text{ and } \|h\| = 1\} \\ &\leq M\end{aligned}$$

Since $x \in A$ is arbitrary, Df is bounded.

Conversely, suppose that Df is bounded. Then there exists $M > 0$ such that for each $x \in A$, $\|Df(x)\| \leq M$. Let $x, y \in A$. The mean value theorem implies that there exists $t^* \in (0, 1)$ such that

$$\begin{aligned}\|f(x) - f(y)\| &\leq \|Df(t^*x + (1 - t^*)y)\| \|x - y\| \\ &\leq M\|x - y\|\end{aligned}$$

Therefore f is Lipschitz. □

11.6 Taylor's Theorem

Drop assumption of separable Banach spaces, only need separably valued maps

Exercise 11.6.0.1. Let Y be a separable Banach space and $f \in C([a, b], Y)$ so that $f \in L_Y^1([a, b], \mathcal{B}([a, b]), m)$. Define $F : (a, b) \rightarrow Y$ by

$$F(x) := \int_{(a, x]} f \, dm$$

Then $F \in C^1((a, b), Y)$ and for each $x_0 \in (a, b)$ and $F'(x_0) = f(x_0)$.

Proof. Let $x_0 \in (a, b)$ and $h \in (0, b - x_0)$. Then continuity implies that

$$\begin{aligned} \frac{1}{\|h\|} \left| \int_{(x_0, x_0+h]} f - f(x_0) \, dm \right| &\leq \frac{1}{\|h\|} \max_{x \in (x_0, x_0+h]} |f(x) - f(x_0)| \|h\| \\ &= \max_{x \in [x_0, x_0+h]} |f(x) - f(x_0)| \\ &\rightarrow 0 \text{ as } h \rightarrow 0 \end{aligned}$$

So

$$\int_{(x_0, x_0+h]} f - f(x_0) \, dm = o(\|h\|) \quad \text{as } h \rightarrow 0$$

Therefore

$$\begin{aligned} F(x_0 + h) &= \int_{(a, x_0+h]} f \, dm \\ &= \int_{(a, x_0]} f \, dm + \int_{(x_0, x_0+h]} f \, dm \\ &= \int_{(a, x_0]} f \, dm + hf(x_0) + \int_{(x_0, x_0+h]} f - f(x_0) \, dm \\ &= F(x_0) + hf(x_0) + o(\|h\|) \quad \text{as } h \rightarrow 0 \end{aligned}$$

The case is similar for $h \in (x_0 - b, 0)$. Since the map $h \mapsto f(x_0)h$ is bounded, F is Frechet differentiable at x_0 and $DF(x_0)(h) = f(x_0)h$. This implies that $F'(x_0) = f(x_0)$ and a previous exercise implies tells us that continuity of f implies continuity of DF . So $F \in C^1(A, Y)$. $\square$

Exercise 11.6.0.2. Fundamental Theorem of Calculus: Let Y be a separable Banach space and $f \in C^1((a, b), Y)$. Then for each $x, x_0 \in (a, b)$, $x_0 < x$ implies that

1. f' is Bochner integrable on $(x_0, x]$
- 2.

$$f(x) - f(x_0) = \int_{(x_0, x]} f' \, dm$$

Proof.

1. Since $f \in C^1((a, b), Y)$, a previous exercise tells us that $f' \in C_Y(a, b)$. Let $x, x_0 \in (a, b)$. Suppose that $x_0 < x$. Choose $c, d \in (a, b)$ such that $a < c < x_0 < x < d < b$. Then f' is continuous on $[c, d]$ and hence Bochner-integrable on $(c, d]$ and $(x_0, x]$.
2. Define $g : (c, d) \rightarrow Y$ by

$$g(\xi) = \int_{(c, \xi]} f' \, dm$$

Then the previous exercise implies that $g \in C_Y^1(c, d)$ and for each $t \in (c, d)$, $g'(t) = f'(t)$. Let $t \in (c, d)$ and $h \in \mathbb{R}$. Then

$$\begin{aligned} Dg(t)(h) &= hg'(t) \\ &= hf'(t) \\ &= Df(t)(h) \end{aligned}$$

So $Dg = Df$ on (c, d) . A previous exercise implies that there exists $c \in Y$ such that $f = g + c$ on (c, d) . Then

$$\begin{aligned} f(x) - f(x_0) &= g(x) + c - (g(x_0) + c) \\ &= g(x) - g(x_0) \\ &= \int_{(c, x]} f' dm - \int_{(c, x_0]} f' dm \\ &= \int_{(x_0, x]} f' dm \end{aligned}$$

□

Exercise 11.6.0.3. Let Y be a Banach space, $A \subset \mathbb{R}$ open and $g : A \rightarrow Y$. If g is n -th order Frechet differentiable, then

$$\frac{d}{dt} \sum_{k=0}^{n-1} \frac{(1-t)^k}{k!} g^{(k)}(t) = \frac{(1-t)^{n-1}}{(n-1)!} g^{(n)}(t)$$

Proof. Taking the derivative yields a telescoping series. □

Exercise 11.6.0.4. Taylor's Theorem I:

Let X be a Banach space, Y a separable Banach space, $A \subset X$ open and convex, $f \in C^{n+1}(A, Y)$, $x_0 \in A$, and $h \in A - x_0$. Then

$$f(x_0 + h) = \sum_{k=0}^n \frac{1}{k!} D^k f(x_0)(h^{\oplus k}) + R(x_0)(h)$$

where $R(x_0) : A - x_0 \rightarrow Y$ is defined by

$$R(x_0)(h) = \frac{1}{n!} \int_{(0,1)} (1-t)^n D^{n+1} f(x_0 + th)(h^{\oplus(n+1)}) dm(t)$$

and $R(x_0)(h) = o(\|h\|^n)$ as $h \rightarrow 0$.

Hint: Define $g : (0, 1) \rightarrow Y$ by

$$g(t) = f(x_0 + th)$$

Then use the previous exercise and the fundamental theorem of calculus.

Proof. Let $h \in X$. Suppose that $x_0 + h \in A$. Define $g : (0, 1) \rightarrow Y$ by

$$g(t) = f(x_0 + th)$$

For each $k \in \{1, \dots, n+1\}$, a previous exercise implies that $g^{(k)}(t) = D^k f(x_0 + th)(h^{\oplus k})$, so $g^{(k)}(0) = D^k f(x_0)(h^{\oplus k})$. The previous exercise and the fundamental theorem of calculus tell us that

$$\begin{aligned} f(x_0 + h) - \sum_{k=0}^n \frac{1}{k!} D^k f(x_0)(h^{\oplus k}) &= g(1) - \sum_{k=0}^n \frac{1}{k!} g^{(k)}(0) \\ &= \int_{(0,1)} \left[\frac{d}{dt} \sum_{k=0}^n \frac{(1-t)^k}{k!} g^{(k)}(t) \right] dm(t) \\ &= \int_{(0,1)} \frac{(1-t)^n}{n!} g^{(n+1)}(t) dm(t) \\ &= \frac{1}{n!} \int_{(0,1)} (1-t)^n D^{n+1} f(x_0 + th)(h^{\oplus(n+1)}) dm(t) \\ &= R(x_0)(h) \end{aligned}$$

Note that

$$\frac{1}{n+1} = \frac{1}{n!} \int_{(0,1)} (1-t)^n dm(t)$$

Since $D^{n+1}f$ is continuous at x_0 , there exists $\delta_1 > 0$ such that for each $h \in B(0, \delta_1)$, $x_0 + h \in A$ and

$$\|D^{n+1}f(x_0 + h) - D^{n+1}f(x_0)\| < 1$$

Let $\epsilon > 0$. Choose $\delta_2 > 0$ such that

$$\frac{1}{n+1} \left(\|D^{n+1}f(x_0)\| + 1 \right) \delta_2 < \epsilon$$

Set $\delta = \min(\delta_1, \delta_2)$. Let $h \in B(0, \delta)$. Then

$$\begin{aligned} \|R(x_0)(h)\| &= \left\| \int_{(0,1)} \frac{1}{n!} \int_{(0,1)} (1-t)^n D^{n+1}f(x_0 + th)(h^{\oplus(n+1)}) dm(t) \right\| \\ &\leq \frac{1}{n!} \int_{(0,1)} \|(1-t)^n D^{n+1}f(x_0 + th)(h^{\oplus(n+1)})\| dm(t) \\ &\leq \max_{t \in [0,1]} \|D^{n+1}f(x_0 + th)\| \|h\|^{n+1} \frac{1}{n!} \int_{(0,1)} (1-t)^n dm(t) \\ &\leq \frac{1}{n+1} \left(\|D^{n+1}f(x_0)\| + \max_{t \in [0,1]} \|D^{n+1}f(x_0 + th) - D^{n+1}f(x_0)\| \right) \|h\|^{n+1} \\ &< \frac{1}{n+1} \left(\|D^{n+1}f(x_0)\| + 1 \right) \|h\|^{n+1} \\ &< \epsilon \|h\|^n \end{aligned}$$

So $R(x_0)(h) = o(\|h\|^n)$ as $h \rightarrow 0$. □

Exercise 11.6.0.5. Taylor's Theorem II:

Let X be a Banach space, Y a separable Banach space, $A \subset X$ open and convex, $f \in C^n(A, Y)$, $x_0 \in A$, and $h \in A - x_0$. Then there exists $R(x_0) : A - x_0 \rightarrow Y$ such that

$$f(x_0 + h) = \sum_{k=0}^n \frac{1}{k!} D^k f(x_0)(h^{\oplus k}) + R(x_0)(h)$$

and $R(x_0)(h) = o(\|h\|^n)$ as $h \rightarrow 0$.

Hint: use Taylor's theorem and expand the derivative inside the integral.

Proof. This is clear by definition for $n = 1$. Suppose that $n \geq 2$. Taylor's theorem implies that

$$f(x_0 + h) = \sum_{k=0}^{n-2} \frac{1}{k!} D^k f(x_0)(h^{\oplus k}) + S(x_0)(h)$$

where $S(x_0) : A - x_0 \rightarrow Y$ is defined by

$$S(x_0)(h) = \frac{1}{(n-2)!} \int_{(0,1)} (1-t)^{n-2} D^{n-1}f(x_0 + th)(h^{\oplus(n-1)}) dm(t)$$

and $S(x_0; h) = o(\|h\|^{n-2})$ as $h \rightarrow 0$. Define $T^{n-1}(x_0) : A - x_0 \rightarrow L^{n-1}(X; Y)$ by

$$T^{n-1}(x_0)(h) = D^{n-1}f(x_0 + h) - D^{n-1}f(x_0) - D^n f(x_0)(h)$$

so that

$$D^{n-1}f(x_0 + h) = D^{n-1}f(x_0) + D^n f(x_0)(h) + T^{n-1}(x_0)(h)$$

and $T^{n-1}(x_0)(h) = o(\|h\|)$ as $h \rightarrow 0$.

Define $R(x_0) : A - x_0 \rightarrow Y$ by

$$R(x_0)(h) = \frac{1}{(n-2)!} \int_{(0,1)} (1-t)^{n-2} T^{n-1}(x_0)(th)(h^{\oplus(n-1)}) dm(t)$$

Note that

$$\int_0^1 (1-t)^{n-2} dt = \frac{1}{n-1}$$

$$\int_0^1 (1-t)^{n-2} t dt = \frac{1}{n(n-1)}$$

Let $\epsilon > 0$. Choose $\delta > 0$ such that for each $h \in B(0, \delta)$, $h \in A - x_0$ and

$$\|T^{n-1}(x_0)(h)\| \leq \epsilon n! \|h\|$$

Let $h \in B(0, \delta)$. Then

$$\begin{aligned} \|R(x_0)(h)\| &= \left\| \frac{1}{(n-2)!} \int_{(0,1)} (1-t)^{n-2} T^{n-1}(x_0)(th)(h^{\oplus(n-1)}) dm(t) \right\| \\ &\leq \frac{1}{(n-2)!} \int_{(0,1)} (1-t)^{n-2} \|T^{n-1}(x_0)(th)(h^{\oplus(n-1)})\| dm(t) \\ &\leq \frac{1}{(n-2)!} \int_{(0,1)} (1-t)^{n-2} \|T^{n-1}(x_0)(th)\| \|h\|^{n-1} dm(t) \\ &\leq \frac{\epsilon}{(n-2)!} n! \|h\|^n \int_{(0,1)} (1-t)^{n-2} t dm(t) \\ &= \epsilon \|h\|^n \end{aligned}$$

So that $R(x_0)(h) = o(\|h\|^n)$ as $h \rightarrow 0$.

Then

$$\begin{aligned} S(x_0)(h) &= \frac{1}{(n-2)!} \int_{(0,1)} (1-t)^{n-2} D^{n-1}f(x_0 + th)(h^{\oplus(n-1)}) dm(t) \\ &= \frac{1}{(n-2)!} \int_{(0,1)} (1-t)^{n-2} D^{n-1}f(x_0)(h^{\oplus(n-1)}) dm(t) \\ &\quad + \frac{1}{(n-2)!} \int_{(0,1)} (1-t)^{n-2} t D^n f(x_0)(h)(h^{\oplus(n-1)}) dm(t) \\ &\quad + \frac{1}{(n-2)!} \int_{(0,1)} (1-t)^{n-2} T^{n-1}(x_0)(th)(h^{\oplus(n-1)}) dm(t) \\ &= \frac{1}{(n-1)!} D^{n-1}f(x_0)(h^{\oplus(n-1)}) + \frac{1}{n!} D^n f(x_0)(h^{\oplus n}) + R_f(x_0)(h) \end{aligned}$$

Hence

$$\begin{aligned} f(x_0 + h) &= \sum_{k=0}^{n-2} \frac{1}{k!} D^k f(x_0)(h^{\oplus k}) + S(x_0)(h) \\ &= \sum_{k=0}^n \frac{1}{k!} D^k f(x_0)(h^{\oplus k}) + R(x_0)(h) \end{aligned}$$

□

Exercise 11.6.0.6. Taylor's Theorem III:

Let X be a Banach space, $A \subset X$ open and convex, $f \in C^n(A)$, $x_0 \in A$, and $h \in A - x_0$. Then there exists $t^* \in (0, 1)$ such that

$$f(x_0 + h) = \sum_{k=0}^{n-1} \frac{1}{k!} D^k f(x_0)(h^{\oplus k}) + \frac{1}{(n-1)!} (1-t^*)^{n-1} D^n f(x_0 + t^*h)(h^{\oplus n})$$

Hint: use Taylor's theorem and the mean value theorem.

Proof. Taylors Theorem implies that

$$f(x_0 + h) = \sum_{k=0}^{n-1} \frac{1}{k!} D^k f(x_0)(h^{\oplus k}) + R(x_0)(h)$$

where

$$R(x_0)(h) = \frac{1}{(n-1)!} \int_{(0,1)} (1-t)^{n-1} D^n f(x_0 + th)(h^{\oplus n}) dm(t)$$

Define $F \in C^1([0, 1])$ by

$$F(t) = \int_{(0,t]} \frac{1}{(n-1)!} (1-s)^{n-1} D^n f(x_0 + sh)(h^{\oplus n}) dm(s)$$

Then the fundamental theorem of calculus implies that

$$F'(t) = \frac{1}{(n-1)!} (1-t)^{n-1} D^n f(x_0 + th)(h^{\oplus n})$$

The mean value theorem implies that there exists $t^* \in (0, 1)$ such that

$$\begin{aligned} R(x_0)(h) &= F(1) - F(0) \\ &= F'(t^*) \\ &= \frac{1}{(n-1)!} (1-t^*)^{n-1} D^n f(x_0 + t^*h)(h^{\oplus n}) \end{aligned}$$

□

Exercise 11.6.0.7. Let X be a Banach space, $A \subset X$ open and convex and $f \in C^2(A)$, $x_0 \in A$. If f has a local minimum at x_0 , then $D^2f(x_0)$ is positive semidefinite.

Proof. Suppose that f has a local minimum at x_0 , then $Df(x_0) = 0$. Let $x \in X$. Then

$$\begin{aligned} 0 &\leq f(x+h) - f(x_0) \\ &= \frac{1}{2} D^2f(x_0)(h, h) + o(\|h\|^2) \quad \text{as } h \rightarrow 0 \end{aligned}$$

Let $h \in X$. Then

$$0 \leq \frac{1}{2} t^2 D^2f(x_0)(h, h) + o(t^2) \quad \text{as } t \rightarrow 0$$

This implies that $D^2f(x_0)(h, h) \geq 0$. So $D^2f(x_0)$ is positive semidefinite.

□

11.7 Implicit and Inverse Function Theorems

Definition 11.7.0.1. Let $(x_0, y_0) \in U$. Then f is said to be **partial Frechet differentiable with respect to X at (x_0, y_0)** if f^{y_0} is Frechet differentiable at x_0 .

Suppose that f is partial Frechet differentiable with respect to X at (x_0, y_0) . We define the **partial Frechet derivative of f with respect to X at (x_0, y_0)** , denoted $D_X f(x_0, y_0) \in L(X, Z)$, by

$$D_X f(x_0, y_0) = Df^{y_0}(x_0)$$

Suppose that for each $y \in Y$, f^y is Frechet differentiable. We define the **partial Frechet derivative of f with respect to X** , denoted $D_X f : X \times Y \rightarrow L(X, Z)$, by

$$D_X f(x, y) = Df^y(x)$$

We define partial Frechet differentiability with respect to Y similarly.

Exercise 11.7.0.2. Let X, Y and Z be Banach spaces, $f : X \times Y \rightarrow Z$ and $(x_0, y_0) \in X \times Y$. If f is Frechet differentiable at (x_0, y_0) , then f is partial Frechet differentiable at (x_0, y_0) with respect to X and Y and for each $h_X \in X$, $h_Y \in Y$,

$$Df(x_0, y_0)(h_X, h_Y) = D_X f(x_0, y_0)(h_X) + D_Y f(x_0, y_0)(h_Y)$$

Proof. Suppose that f is Frechet differentiable at (x_0, y_0) . Then

$$f[(x_0, y_0) + (h_X, h_Y)] = f(x_0, y_0) + Df(x_0, y_0)(h_X, h_Y) + o(\|(h_X, h_Y)\|_{X \oplus Y}) \text{ as } (h_X, h_Y) \rightarrow (0, 0)$$

Since there exist $C_1, C_2 > 0$ such that for each $h_X \in X$ and $h_Y \in Y$,

$$C_1(\|x\| + \|y\|) \leq \|(h_X, h_Y)\|_{X \oplus Y} \leq C_2(\|x\| + \|y\|)$$

we have that

$$f^{y_0}(x_0 + h_X) = f^{y_0}(x_0) + Df^{y_0}(x_0)(h_X) + o(\|h_X\|) \text{ as } h_X \rightarrow 0$$

Therefore $f^{y_0} : X \rightarrow Z$ is Frechet differentiable at x_0 and $Df^{y_0}(x_0) = Df(x_0, y_0)(h_X, 0)$. Hence f is partial Frechet differentiable at (x_0, y_0) with respect to X and for each $h_X \in X$, $D_X f(x_0, y_0)(h_X) = Df(x_0, y_0)(h_X, 0)$. Similarly, f is partial Frechet differentiable at (x_0, y_0) with respect to Y and for each $h_Y \in Y$, $D_Y f(x_0, y_0)(h_Y) = Df(x_0, y_0)(0, h_Y)$. Let $h_X \in X$ and $h_Y \in Y$. Then

$$\begin{aligned} Df(x_0, y_0)(h_X, h_Y) &= Df(x_0, y_0)[(h_X, 0) + (0, h_Y)] \\ &= Df(x_0, y_0)(h_X, 0) + Df(x_0, y_0)(0, h_Y) \\ &= D_X f(x_0, y_0)(h_X) + D_Y f(x_0, y_0)(h_Y) \end{aligned}$$

□

Exercise 11.7.0.3. Let X, Y and Z be Banach spaces, $U \subset X \times Y$ open, $f : U \rightarrow Z$ and $n \in \mathbb{N}$. If f is $C^1(U, Z)$, then $D_X f, D_Y f \in C(U, Z)$.

Proof. Suppose that f is $C^1(U, Z)$. Then $Df \in C(U, Z)$. Define $\phi_X : X \rightarrow X \times Y$ and $\phi_Y : Y \rightarrow X \times Y$ by $\phi_X(x) = (x, 0)$ and $\phi_Y(y) = (0, y)$. Then $\phi_X \in L(X, X \times Y)$ and $\phi_Y \in L(Y, X \times Y)$. The previous exercise implies that for each $(x, y) \in U$, $D_X f(x, y) = Df(x, y) \circ \phi_X$. Let $(x, y), (x_0, y_0) \in U$. Then

$$\begin{aligned} \|D_X f(x, y) - D_X f(x_0, y_0)\| &= \|Df(x, y) \circ \phi_X - Df(x_0, y_0) \circ \phi_X\| \\ &= \|(Df(x, y) - Df(x_0, y_0)) \circ \phi_X\| \\ &\leq \|Df(x, y) - Df(x_0, y_0)\| \|\phi_X\| \end{aligned}$$

□

Exercise 11.7.0.4. Let X, Y and Z be Banach spaces, $U \subset X \times Y$ open, $F : U \rightarrow Z$, $(x_0, y_0) \in U$. Suppose that F is partial Frechet differentiable with respect to Y on U and F and $D_Y F$ continuous at (x_0, y_0) . Then there

Proof. Set $L = D_Y F(x_0, y_0)$. Define $G : U \rightarrow Z$ by $G(x, y) = y - L^{-1}F(x, y)$. Then $G(x_0, y_0) = y_0$ and since $F \in C^1(U, Z)$, $G \in C^1(U, Z)$. The previous exercise implies that $D_Y G \in C(U, Z)$. Note that for each $(x, y) \in U$,

$$\begin{aligned} D_Y G(x, y) &= \text{id}_Y - L^{-1} D_Y F(x, y) \\ &= L^{-1}(L - D_Y F(x, y)) \end{aligned}$$

which implies that $D_Y G(x_0, y_0) = 0$. Set $\epsilon = 1/2$. Since U is open and $D_Y G$ is continuous at (x_0, y_0) there exist $\delta_X, \delta_Y > 0$ such that for each $x \in B(x_0, \delta_X)$ and $y \in B(y_0, \delta_Y)$, $(x, y) \in U$ and

$$\begin{aligned} \|D_Y G(x, y)\| &= \|D_Y G(x, y) - D_Y G(x_0, y_0)\| \\ &< \epsilon \end{aligned}$$

Set $A = B(x_0, \delta_X)$ and $B = B(y_0, \delta_Y)$. Let $x \in A$ and $y_1, y_2 \in B$. Define $l : [0, 1] \rightarrow B$ by $l(t) = ty_1 + (1 - t)y_2$. The mean value theorem implies that

$$\begin{aligned} \|G(x, y_1) - G(x, y_2)\| &\leq \sup_{t \in [0, 1]} \|D_Y G(x, l(t))\| \|y_1 - y_2\| \\ &\leq \epsilon \|y_1 - y_2\| \\ &= \frac{1}{2} \|y_1 - y_2\| \end{aligned}$$

Hence, for each $x \in X$ and $y \in Y$, $\|G(x, y)\| \leq \frac{1}{2} \|y_1 - y_2\|$. For $x \in A$, define $T_x : B \rightarrow B$ by $T_x(y) = G(x, y)$. □

11.8 The Gradient

Definition 11.8.0.1. Let H be a Hilbert space, $f : H \rightarrow \mathbb{C}$ and $x_0 \in H$. Suppose that f is Frechet differentiable at x_0 . Then $Df(x_0) \in H^*$. We define the **gradient of f at x_0** , denoted $\nabla f(x_0) \in H$, via the Riesz representation theorem to be the unique element of H satisfying

$$\langle \nabla f(x_0), y \rangle = Df(x_0)(y) \quad \text{for each } y \in H$$

Definition 11.8.0.2. Let $H \subset Y^X$ be a Hilbert space and $F : H \rightarrow \mathbb{R}$. We define the **functional derivative** F with respect to $\phi(x)$ of $f : X \rightarrow Y$, denoted $\frac{\delta F}{\delta \phi} : X \rightarrow \mathbb{R}$, by . **functional derivative relate this to rigged hilbert spaces and $\frac{\delta F}{\delta \phi(x_0)} = DF(\phi)(\delta_{x_0})$**

Chapter 12

Banach Lattices

Note 12.0.0.1. This chapter assumes familiarity with cones [ref section on convex sets subsection on cones](#).

To Do:

- cover Freudenthal spectral theorem [wikipedia link here](#) and its special cases. For example,
 - in measure theory notes, cover radon nikodym theorem
 - in banach algebra chapter later, cover spectral theorem for normal operators
 - in some place, cover validity of the poisson formula
 - in probability notes, cover Doob's almost sure convergence theorem in probability notes [Doob's almost sure convergence theorem](#)
 - in measure theory notes, try to show MCT for bochner integrable functions on Banach lattices
-

12.1 Introduction

12.1.1 Ordered Vector Spaces

Definition 12.1.1.1. Let X be a vector space and $\leq \subset X \times X$ a partial order on X . Then $(X, \leq)$ is said to be an **ordered vector space** if for each $x, y \in X$, $x \leq y$ implies that

1. for each $z \in X$, $x + z \leq y + z$,
2. for each $\lambda \in \mathbb{R}_+$, $\lambda x \leq \lambda y$,

Note 12.1.1.2. When the context is clear we may write X in place of $(X, \leq)$.

Exercise 12.1.1.3. Let $(X, \leq)$ be an ordered vector space. Then

1. for each $x \in X$, $0_X \leq x$ implies that $-x \leq 0_X$,
2. for each $x, y \in X$, $x \leq y$ implies that $-y \leq -x$.

Proof. 1. Let $x \in X$. Suppose that $0_X \leq x$. Since $(X, \leq)$ is an ordered vector space,

$$\begin{aligned} -x &= 0_X - x \\ &\leq x - x \\ &= 0_X. \end{aligned}$$

Since $x \in X$ with $0_X \leq x$ is arbitrary, we have that for each $x \in X$, $0_X \leq x$ implies that $-x \leq 0_X$.

2. For each $x, y \in X$,

$$\begin{aligned}
 x \leq y &\implies x - x \leq y - x \\
 &\implies 0_X \leq y - x \\
 &\implies -(y - x) \leq 0_X \quad (\text{by prev part}) \\
 &\implies x - y \leq 0_X \\
 &\implies (x - y) - x \leq 0_X - x \\
 &\implies -y \leq -x.
 \end{aligned}$$

□

Definition 12.1.1.4. Let $(X, \leq)$ be a Riesz space and $x \in X$.

- Then x is said to be **positive** if $0_X \leq x$.
- We define the **positive $\leq$ -cone of X** , denoted $(X, \leq)_+$, by $(X, \leq)_+ := \{x \in X : x \text{ is positive}\}$.

Note 12.1.1.5. When the context is clear,

- we may say “positive cone of X ” instead of “positive $\leq$ -cone of X ”,
- we may write X_+ in place of $(X, \leq)_+$.

Exercise 12.1.1.6. Let X be an ordered vector space. Then $X_+ \in \text{PtConvCone}(X)$.

Proof.

- Let $y \in \mathbb{R}_+ X_+$. Then there exists $a \in \mathbb{R}_+$ and $x \in X_+$ such that $y = ax$. Since $x \in X_+$, we have that $0_X \leq x$. Since X is an ordered vector space,

$$\begin{aligned}
 0_X &= a0_X \\
 &\leq ax \\
 &= y.
 \end{aligned}$$

Hence $y \in X_+$. Since $y \in \mathbb{R}_+ X_+$ is arbitrary, we have that for each $y \in \mathbb{R}_+ X_+$, $y \in X_+$. Hence $\mathbb{R}_+ X_+ \subset X_+$. Thus $X_+ \in \text{Cone}(X)$.

- Let $x \in X_+ + X_+$. Then there exists $x_1, x_2 \in X_+$ such that $x = x_1 + x_2$. Then $0_X \leq x_1$ and $0_X \leq x_2$. Since X is an ordered vector space, we have that

$$\begin{aligned}
 0_X &= 0_X + 0_X \\
 &\leq x_1 + 0_X \\
 &\leq x_1 + x_2.
 \end{aligned}$$

Hence

$$\begin{aligned}
 x &= x_1 + x_2 \\
 &\in X_+.
 \end{aligned}$$

Since $x \in X_+ + X_+$ is arbitrary, we have that for each $x \in X_+ + X_+$, $x \in X_+$. Thus $X_+ + X_+ \subset X_+$. Since $X_+ \in \text{Cone}(X)$ and $X_+ + X_+ \subset X_+$, **an ex in subsection on cones** implies that $X_+ \in \text{ConvCone}(X)$.

- Let $x \in X_+ \cap -X_+$. Then $x \in X_+$ and $x \in -X_+$. Thus there exists $y \in X_+$ such that $x = -y$. Since $x, y \in X_+$, $0_X \leq x$ and $0_X \leq y$. Therefore

$$\begin{aligned} 0_X &\leq y \\ &= -x \\ &\leq 0_X \quad (\text{by prev ex}) \\ &\leq 0_X. \end{aligned}$$

Thus $0_X \leq -x$ and $-x \leq 0_X$. Since $\leq$ is a partial order, $-x = 0_X$. Hence

$$\begin{aligned} x &= 0_X \\ &\in \{0_X\}. \end{aligned}$$

Since $x \in X_+ \cap -X_+$ is arbitrary, we have that for each $x \in X_+ \cap -X_+$, $x \in \{0_X\}$. Thus

$$\begin{aligned} X_+ \cap -X_+ &\subset \{0_X\} \\ &\subset X_+ \cap -X_+. \end{aligned}$$

Thus $X_+ \cap -X_+ = \{0_X\}$. Therefore $X_+ \in \text{PtConvCone}(X)$.

□

Definition 12.1.1.7. Let X be a vector space and $C \in \text{PtConvCone}(X)$. We define the **partial order on X induced by C** , denoted $\leq_C \subset X \times X$, by $\leq_C := \{(x, y) \in X \times X : y - x \in C\}$.

Exercise 12.1.1.8. Let X be a vector space and $C \in \text{PtConvCone}(X)$. Then

1. $\leq_C$ is a partial order
2. $(X, \leq_C)$ is an ordered vector space.

Proof.

1. • – Let $x \in X$. Since

$$\begin{aligned} x - x &= 0_X \\ &\in C, \end{aligned}$$

- we have that $x \leq_C x$. Since $x \in X$ is arbitrary, we have that for each $x \in X$, $x \leq_C x$. Therefore $\leq_C$ is reflexive.
- Let $x, y, z \in X$. Suppose that $x \leq_C y$ and $y \leq_C z$. Then $y - x, z - y \in C$. Thus

$$\begin{aligned} z - x &= (z - y) + (y - x) \\ &\in C + C \\ &\subset C \quad (\text{since } C \in \text{ConvCone}(X)). \end{aligned}$$

Hence $x \leq_C z$. Since $x, y, z \in X$ with $x \leq_C y$ and $y \leq_C z$ are arbitrary, we have that for each $x, y, z \in X$, $x \leq_C y$ and $y \leq_C z$ implies that $x \leq_C z$. Therefore $\leq_C$ is transitive.

Since $\leq_C$ is reflexive and transitive, $\leq_C$ is a preorder on X .

- Let $x, y \in X$. Suppose that $x \leq_C y$ and $y \leq_C x$. Then $y - x, x - y \in C$. Hence

$$\begin{aligned} y - x &\in C \cap -C \\ &= \{0_X\} \quad (\text{since } C \text{ is pointed}). \end{aligned}$$

Thus $y - x = 0_X$ and $x = y$. Since $x, y \in X$ with $x \leq_C y$ and $y \leq_C x$ are arbitrary, we have that for each $x, y \in X$, $x \leq_C y$ and $y \leq_C x$ implies that $x = y$. Therefore $\leq_C$ is a symmetric. Since $\leq_C$ is a preorder on X and $\leq_C$ is symmetric, $\leq_C$ is a partial order on X .

2. Let $x, y \in X$. Suppose that $x \leq_C y$. Then $y - x \in C$.

(a) Let $z \in X$. Since $y - x \in C$, we have that

$$\begin{aligned} (y + z) - (x + z) &= y - x \\ &\in C. \end{aligned}$$

Thus $x + z \leq y + z$. Since $z \in X$ is arbitrary, we have that for each $z \in X$, $x + z \leq y + z$.

(b) Let $\lambda \in \mathbb{R}_+$. Since $y - x \in C$ and $C \in \text{Cone}(X)$, we have that

$$\begin{aligned} \lambda y - \lambda x &= \lambda(y - x) \\ &\in C. \end{aligned}$$

Thus $\lambda x \leq_C \lambda y$. Since $\lambda \in \mathbb{R}_+$ is arbitrary, we have that for each $\lambda \in \mathbb{R}_+$, $\lambda x \leq \lambda y$.

Since $x, y \in X$ with $x \leq_C y$ are arbitrary, we have that for each $x, y \in X$, $x \leq_C y$ implies that

(a) for each $z \in X$, $x + z \leq y + z$,

(b) for each $\lambda \in \mathbb{R}_+$, $\lambda x \leq \lambda y$.

Thus $(X, \leq_C)$ is an ordered vector space. *reorganize this and the definition*

□

Exercise 12.1.1.9. There is a bijection between $\text{PtConvCone}(X)$ and partial orders $\leq_C$ on X such that $(X, \leq_C)$ is an ordered vector space. Is this a natural transformation?

Proof. **FINISH!!!**

□

12.1.2 Riesz Spaces

Definition 12.1.2.1. Let $(X, \leq)$ be an ordered vector space. Then $(X, \leq)$ is said to be a **Riesz space** or **vector lattice** if

- $(X, \leq)$ is an ordered vector space,
- $(X, \leq)$ is a lattice.

Exercise 12.1.2.2.

Chapter 13

Banach Algebras

13.1 Introduction

Definition 13.1.0.1. Let X be a Banach space and $\mu : X \times X \rightarrow X$. Then (X, μ) is said to be a **Banach algebra** if

1. (X, μ) is an associative algebra
2. $\mu \in L^2(X)$ and $\|\mu\| \leq 1$

Note 13.1.0.2. By definition in the section on multilinear maps, condition (2) is equivalent to the assumption that for each $x, y \in X$, $\|xy\| \leq \|x\|\|y\|$.

Definition 13.1.0.3. Let X be a Banach algebra and $e \in X$. Then e is said to be an **identity** if for each $x \in X$, $ex = xe = x$.

Definition 13.1.0.4. Let X be a Banach algebra. Then X is said to be **unital** if there exists $e \in X$ such that e is an identity.

Exercise 13.1.0.5. Let X be a unital Banach algebra. Then there exists a unique $e \in X$ such that e is an identity.

Proof.

• **Existence:**

By definition, there exists $e \in X$ such that e is an identity.

• **Uniqueness:**

Let $e' \in X$. Suppose that e' is an identity. Then

$$\begin{aligned} e' &= e'e \\ &= e' \end{aligned}$$

□

Exercise 13.1.0.6. Let X be a unital Banach algebra. If $X \neq \{0\}$, then $1 \leq \|e\|$.

Proof. Suppose that $X \neq \{0\}$. Then $e \neq 0$ which implies that $\|e\| > 0$. Since

$$\|e\| = \|ee\| \leq \|e\|\|e\|$$

we have that $1 \leq \|e\|$.

□

Exercise 13.1.0.7. Fundamental Example:

Let X be a Banach space. Then $GL(X)$ is a unital Banach algebra.

Proof. Clear.

□

Definition 13.1.0.8. Let X be a unital Banach algebra and $x, y \in X$. Then y is said to be an **inverse** of x if $xy = yx = e$.

Definition 13.1.0.9. Let X be a unital Banach algebra and $x \in X$. Then y is said to be **invertible** if there exists $y \in X$ such that y is an inverse of x .

Exercise 13.1.0.10. Let X be a unital Banach algebra and $x \in X$. If x is invertible, then there exists a unique $y \in X$ such that y is an inverse of x .

Proof. Suppose that x is invertible.

• **Existence:**

By definition, there exists $y \in X$ such that y is an inverse of x .

• **Uniqueness:**

Let $y' \in X$. Suppose that y' is an inverse of x . Then

$$\begin{aligned} y' &= y'e \\ &= y'(xy) \\ &= (y'x)y \\ &= ey \\ &= y \end{aligned}$$

□

Definition 13.1.0.11. Let X be a unital Banach algebra. We define $G(X) = \{x \in X : x \text{ is invertible}\}$.

Exercise 13.1.0.12. Let X be a unital Banach algebra. Then $G(X)$ is a group.

Proof. Clear.

□

Definition 13.1.0.13. Let X be a unital Banach algebra and $x \in G(X)$. We define the **inverse of x** , denoted x^{-1} , to be the unique $y \in X$ such that $yx = xy = e$.

Exercise 13.1.0.14. Let X be a unital Banach algebra, $x \in G(X)$ and $\lambda \in \mathbb{C}^\times$. Then $\lambda x \in G(X)$ is and $(\lambda x)^{-1} = \lambda^{-1}x^{-1}$.

Proof. We have that

$$\begin{aligned} (\lambda^{-1}x^{-1})(\lambda x) &= ((\lambda^{-1}\lambda)x^{-1})x \\ &= (1x^{-1})x \\ &= x^{-1}x \\ &= e \end{aligned}$$

Similarly, $(\lambda x)(\lambda^{-1}x^{-1}) = e$. Hence $\lambda x \in G(X)$ and $(\lambda x)^{-1} = \lambda^{-1}x^{-1}$.

□

Exercise 13.1.0.15. Let X be a unital Banach algebra and $x, y \in G(X)$. Then $xy \in G(X)$ is and $(xy)^{-1} = y^{-1}x^{-1}$.

Proof. We have that

$$\begin{aligned} (y^{-1}x^{-1})(xy) &= y^{-1}(x^{-1}(xy)) \\ &= y^{-1}((x^{-1}x)y) \\ &= y^{-1}(ey) \\ &= y^{-1}y \\ &= e \end{aligned}$$

Similarly, $(xy)(y^{-1}x^{-1}) = e$. Hence $xy \in G(X)$ and $(xy)^{-1} = y^{-1}x^{-1}$.

□

Exercise 13.1.0.16. Let X be a unital Banach algebra and $x, y \in X$.

1. If $xy \in G(X)$ and $y \in G(X)$, then $x \in G(X)$.
2. If $xy \in G(X)$ and $yx \in G(X)$, then $x \in G(X)$ and $y \in G(X)$.
3. If $xy = yx$ and $x \notin G(X)$, then $xy \notin G(X)$.

Proof.

1. Suppose that $xy \in G(X)$ and $y \in G(X)$. Since $xy \in G(X)$, there exists $z \in G(X)$ such that $z(xy) = (xy)z = e$. Since $z, y \in G(X)$, we have that $yz \in G(X)$ and $(yz)^{-1} = z^{-1}y^{-1}$. Therefore

$$\begin{aligned} z(xy) = e &\implies xy = z^{-1} \\ &\implies x = z^{-1}y^{-1} \\ &\implies x = (yz)^{-1} \end{aligned}$$

Hence $x \in G(X)$.

2. Suppose that $xy, yx \in G(X)$. Then there exists $z \in G(X)$ such that $z(xy) = (xy)z = e$. Then $x(yz) = e$ and since $yx \in G(X)$, we have that

$$\begin{aligned} z(xy) = e &\implies (zx)y = e \\ &\implies (zx)yx = x \\ &\implies zx = x(yx)^{-1} \\ &\implies y(zx) = y(x(yx)^{-1}) \\ &\implies (yz)x = (yx)(yx)^{-1} \\ &\implies (yz)x = e \end{aligned}$$

Since $(yz)x = x(yz) = e$, we have that $x \in G(X)$. Similarly, $y \in G(X)$.

3. Suppose that $xy = yx$ and $x \notin G(X)$. Part (2) implies that $xy \notin G(X)$ or $yx \notin G(X)$. Since $xy = yx$, we have that $xy \notin G(X)$. □

Exercise 13.1.0.17. Let X be a unital Banach algebra.

1. For each $x \in X$, if $\|x\| < 1$, then $e - x \in G(X)$ and

$$(e - x)^{-1} = \sum_{n=0}^{\infty} x^n$$

2. For each $x \in X$ and $\lambda \in \mathbb{C}$, if $\|x\| < |\lambda|$, then $\lambda e - x \in G(X)$ and

$$(\lambda e - x)^{-1} = \sum_{n=0}^{\infty} \lambda^{-(n+1)} x^n$$

3. For each $x, y \in X$, if $x \in G(X)$ and $\|y\| < \|x^{-1}\|^{-1}$, then $x - y \in G(X)$ and

$$(x - y)^{-1} = x^{-1} \sum_{n=0}^{\infty} (yx^{-1})^n$$

4. For each $x, y \in X$, if $x \in G(X)$ and $\|x - y\| < \|x^{-1}\|^{-1}$, then $y \in G(X)$ and

$$y^{-1} = x^{-1} \sum_{n=0}^{\infty} (e - yx^{-1})^n$$

5. $G(X)$ is open

Proof.

1. Let $x \in X$. Suppose that $\|x\| < 1$. Then

$$\sum_{n=0}^{\infty} \|x^n\| \leq \sum_{n=0}^{\infty} \|x\|^n < \infty$$

Since X is a complete, $\sum_{n=0}^{\infty} x^n$ converges in X .

Define $(s_k)_{k=0}^{\infty} \subset X$ and $s \in X$ by $s_k = \sum_{n=0}^k x^n$ and $s = \sum_{n=0}^{\infty} x^n$. Then for each $k \in \mathbb{N}$,

$$\begin{aligned} (e - x)s_k &= s_k - xs_k \\ &= e - x^{k+1} \end{aligned}$$

Since $x^k \rightarrow 0$ as $k \rightarrow \infty$, we have that $(e - x)s_k \rightarrow e$ as $k \rightarrow \infty$. Since multiplication on Banach algebras is continuous, we have that $(e - x)s_k \rightarrow (e - x)s$ as $k \rightarrow \infty$. Uniqueness of limits implies that $(e - x)s = e$. A similar argument implies that $s(e - x) = e$. Thus $e - x \in G(X)$ and $(e - x)^{-1} = \sum_{n=0}^{\infty} x^n$.

2. Let $x \in X$ and $\lambda \in \mathbb{C}$. Suppose that $\|x\| < |\lambda|$. Then $\lambda \neq 0$ and

$$\begin{aligned} \|\lambda^{-1}x\| &= |\lambda^{-1}|\|x\| \\ &= |\lambda|^{-1}\|x\| \\ &< |\lambda|^{-1}|\lambda| \\ &= 1 \end{aligned}$$

By (1), we have that $e - \lambda^{-1}x \in G(X)$ and

$$\begin{aligned} (e - \lambda^{-1}x)^{-1} &= \sum_{n=0}^{\infty} (\lambda^{-1}x)^n \\ &= \sum_{n=0}^{\infty} \lambda^{-n}x^n \end{aligned}$$

Therefore,

$$\begin{aligned} \lambda e - x &= \lambda(e - \lambda^{-1}x) \\ &\in G(X) \end{aligned}$$

and

$$\begin{aligned} (\lambda e - x)^{-1} &= (\lambda(e - \lambda^{-1}x))^{-1} \\ &= \lambda^{-1}(e - \lambda^{-1}x)^{-1} \\ &= \lambda^{-1} \sum_{n=0}^{\infty} \lambda^{-n}x^n \\ &= \sum_{n=0}^{\infty} \lambda^{-(n+1)}x^n \end{aligned}$$

3. Let $x, y \in X$. Suppose that $x \in G(X)$ and $\|y\| < \|x^{-1}\|^{-1}$. Then

$$\begin{aligned}\|yx^{-1}\| &\leq \|y\|\|x^{-1}\| \\ &< \|x^{-1}\|^{-1}\|x^{-1}\| \\ &= 1\end{aligned}$$

Hence $e - yx^{-1} \in G(X)$ and

$$(e - yx^{-1}) = \sum_{n=0}^{\infty} (yx^{-1})^n$$

This implies that

$$\begin{aligned}x - y &= (e - yx^{-1})x \\ &\in G(X)\end{aligned}$$

and

$$\begin{aligned}(x - y)^{-1} &= ((e - yx^{-1})x)^{-1} \\ &= x^{-1}(e - yx^{-1})^{-1} \\ &= x^{-1} \sum_{n=0}^{\infty} (yx^{-1})^n\end{aligned}$$

4. Let $x, y \in X$. Suppose that $x \in G(X)$ and $\|x - y\| < \|x^{-1}\|^{-1}$. Then (2) implies that

$$\begin{aligned}y &= x - (x - y) \\ &\in G(X)\end{aligned}$$

and

$$\begin{aligned}y^{-1} &= (x - (x - y))^{-1} \\ &= x^{-1} \sum_{n=0}^{\infty} ((x - y)x^{-1})^n \\ &= x^{-1} \sum_{n=0}^{\infty} (e - yx^{-1})^n\end{aligned}$$

5. Let $x \in G(X)$. Choose $\delta = \|x^{-1}\|^{-1}$. By (3), $B(x, \delta) \subset G(X)$. Since $x \in G(X)$ is arbitrary, $G(X)$ is open.

□

Definition 13.1.0.18. Let X be a unital Banach algebra. We define $\text{inv} : G(X) \rightarrow G(X)$ by $\text{inv}(x) = x^{-1}$.

Exercise 13.1.0.19. Let X be a unital Banach algebra. Then

1. for each $x, y \in X$, if $x \in G(X)$ and $\|y\| \leq \frac{1}{2}\|x^{-1}\|^{-1}$ so that $x - y \in G(X)$, then

$$\|(x - y)^{-1} - x^{-1}\| \leq 2\|x^{-1}\|^2\|y\|$$

2. $\text{inv} : G(X) \rightarrow G(X)$ is continuous

3. $G(X)$ is a topological group

Proof.

1. Let $x, y \in X$. Suppose that $x \in G(X)$ and $\|y\| \leq 2^{-1}\|x^{-1}\|^{-1}$. The previous exercise implies that

$$\begin{aligned} \|(x - y)^{-1} - x^{-1}\| &= \left\| x^{-1} \sum_{n=0}^{\infty} (yx^{-1})^n - x^{-1} \right\| \\ &= \left\| x^{-1} \sum_{n=1}^{\infty} (yx^{-1})^n \right\| \\ &\leq \|x^{-1}\| \sum_{n=1}^{\infty} (\|y\| \|x^{-1}\|)^n \\ &= \|x^{-1}\|^2 \|y\| \sum_{n=0}^{\infty} (\|y\| \|x^{-1}\|)^n \\ &= \|x^{-1}\|^2 \|y\| \sum_{n=0}^{\infty} \frac{1}{2^n} \\ &= 2\|x^{-1}\|^2 \|y\| \end{aligned}$$

2. Let $(x_n)_{n \in \mathbb{N}} \subset G(X)$ and $x \in G(X)$. Suppose that $x_n \rightarrow x$ in $G(X)$. Then $x_n \rightarrow x$ in X . Define $(y_n)_{n \in \mathbb{N}} \subset X$ by $y_n = x - x_n$. Then $y_n \rightarrow 0$ in X . Let $\epsilon > 0$. Then there exists $N \in \mathbb{N}$ such that for each $n \in \mathbb{N}$, $n \geq N$ implies that

$$\|y_n\| < \max \left(\frac{\epsilon}{2\|x^{-1}\|^2}, \frac{1}{2}\|x^{-1}\|^{-1} \right)$$

Let $n \in \mathbb{N}$. Suppose that $n \geq N$. By (1), we have that

$$\begin{aligned} \|\text{inv}(x_n) - \text{inv}(x)\| &= \|x_n^{-1} - x^{-1}\| \\ &= \|(x - y_n)^{-1} - x^{-1}\| \\ &\leq 2\|x^{-1}\|^2 \|y_n\| \\ &< \epsilon \end{aligned}$$

Hence $\text{inv}(x_n) \rightarrow \text{inv}(x)$ in X . Thus $\text{inv}(x_n) \rightarrow \text{inv}(x)$ in $G(X)$. So $\text{inv} : G(X) \rightarrow G(X)$ is continuous.

3. Since multiplication $G(X) \times G(X) \rightarrow G(X)$ and multiplicative inversion $\text{inv} : G(X) \rightarrow G(X)$ are continuous, $G(X)$ is a topological group. □

Exercise 13.1.0.20. Let X be a unital Banach algebra. Then $\text{inv} : G(X) \rightarrow G(X)$ is Frechet differentiable and for each $x \in G(X)$, $h \in X$, $D\text{inv}(x)(h) = -x^{-1}hx^{-1}$.

Proof. Let $x \in G(X)$ and $h \in B(x, \|x^{-1}\|^{-1})$. A previous exercise implies that $x + h \in G(X)$ and

$$\begin{aligned} \text{inv}(x + h) &= (x + h)^{-1} \\ &= x^{-1} \sum_{n=0}^{\infty} [(-h)x^{-1}]^n \\ &= x^{-1} - x^{-1}hx^{-1} + x^{-1} \sum_{n=2}^{\infty} [(-h)x^{-1}]^n \\ &= x^{-1} - x^{-1}hx^{-1} + o(\|h\|) \text{ as } h \rightarrow 0 \end{aligned}$$

Since the map $X \rightarrow X$ given by $h \mapsto -x^{-1}hx^{-1}$ is a bounded linear operator and $x^{-1} \sum_{n=2}^{\infty} [(-h)x^{-1}]^n = o(\|h\|)$ as $h \rightarrow 0$, we have that inv is differentiable at x and for each $h \in X$, $D\text{inv}(x)(h) = -x^{-1}hx^{-1}$. Since $x \in G(X)$ is arbitrary, we have that $\text{inv}(x)$ is differentiable. □

do all the other derivatives like power rule, product rule, etc

Exercise 13.1.0.21.

Proof. content...

**Exercise 13.1.0.22.**

Proof. content...

**Exercise 13.1.0.23.**

Proof. content...



13.2 Spectral Theory

Definition 13.2.0.1. Let X be a unital Banach algebra and $x \in X$. We define the

- **resolvent** of x , denoted $\rho(x)$, by

$$\rho(x) = \{\lambda \in \mathbb{C} : \lambda e - x \in G(X)\}$$

- **spectrum** of x , denoted $\sigma(x)$, by

$$\sigma(x) = \rho(x)^c$$

Exercise 13.2.0.2. Let X be a unital Banach algebra and $x \in X$. Then

1. $\rho(x)$ is open
2. $\sigma(x)$ is closed
3. $\sigma(x) \subset \bar{B}(0, \|x\|)$
4. $\sigma(x)$ is compact

Proof.

1. Let $\lambda \in \rho(x)$. Set $\delta = (\|(\lambda e - x)^{-1}\| \|e\|)^{-1} > 0$. Let $\lambda' \in B(\lambda, \delta)$. Then

$$\begin{aligned} \|(\lambda e - x) - (\lambda' e - x)\| &= |\lambda - \lambda'| \|e\| \\ &< \delta \|e\| \\ &= \|(\lambda e - x)^{-1}\|^{-1} \end{aligned}$$

A previous exercise implies that $\lambda' e - x \in G(X)$. Hence $\lambda' \in \rho(x)$. Since $\lambda' \in B(\lambda, \delta)$ is arbitrary, we have that $B(\lambda, \delta) \subset \rho(x)$. So for each $\lambda \in \rho(x)$, there exists $\delta > 0$ such that $B(\lambda, \delta) \subset \rho(x)$. Hence $\rho(x)$ is open.

2. Since $\sigma(x) = \rho(x)^c$ and $\rho(x)$ is open, we have that $\sigma(x)$ is closed.
3. Let $\lambda \in \sigma(x)$. For the sake of contradiction, suppose that $\|x\| < |\lambda|$. A previous exercise implies that $\lambda e - x \in G(X)$. Hence

$$\begin{aligned} \lambda &\in \rho(x) \\ &= \sigma(x)^c \end{aligned}$$

which is a contradiction. So $|\lambda| \leq \|x\|$ and thus $\lambda \in \bar{B}(0, \|x\|)$. Since $\lambda \in \sigma(x)$ is arbitrary, $\sigma(x) \subset \bar{B}(0, \|x\|)$.

4. Since $\sigma(x) \subset \mathbb{C}$ is closed and bounded, $\sigma(x)$ is compact.

□

Exercise 13.2.0.3. Let X be a unital Banach algebra and $x \in X$ and $p \in \mathbb{C}[t]$. Suppose that $\deg p \geq 1$. Then $\sigma(p(x)) = \{p(\lambda) : \lambda \in \sigma(x)\}$.

Hint: Consider the roots of $p(x) - p(\lambda)e$.

Proof. Let $\lambda \in \sigma(x)$. Then λe is a root of $p(t) - p(\lambda)e$. Therefore, there exists $q \in \mathbb{C}[t]$ such that $\deg q = \deg p - 1$ and $p(x) - p(\lambda)e = (x - \lambda e)q(x)$. Since $q(x)(x - \lambda e) = (x - \lambda e)q(x)$ and $(x - \lambda e) \notin G(X)$, a previous exercise implies that $p(x) - p(\lambda)e \notin G(X)$. Thus $p(\lambda) \in \sigma(p(x))$. Since $\lambda \in \sigma(x)$ is arbitrary, we have that $\{p(\lambda) : \lambda \in \sigma(x)\} \subset \sigma(p(x))$.

Conversely, let $\mu \in \sigma(p(x))$. Set $n = \deg p$. Then there exist $(a_j)_{j=1}^n \subset \mathbb{C}$ and $a \in \mathbb{C}$ such that

$$p(x) - \mu e = a \prod_{j=1}^n (x - a_j e)$$

Since $p(x) - \mu e \notin G(X)$, there exists $j \in \{1, \dots, n\}$ such that $(x - a_j e) \notin G(X)$. Thus $a_j \in \sigma(x)$. By construction

$$\begin{aligned} (p(a_j) - \mu)e &= p(a_j)e - \mu e \\ &= p(a_j e) - \mu e \\ &= 0 \end{aligned}$$

Thus

$$\begin{aligned} \mu &= p(a_j) \\ &\in \{p(\lambda) : \lambda \in \sigma(x)\} \end{aligned}$$

Since $\mu \in \sigma(p(x))$ is arbitrary, we have that $\sigma(p(x)) \subset \{p(\lambda) : \lambda \in \sigma(x)\}$. Hence $\sigma(p(x)) = \{p(\lambda) : \lambda \in \sigma(x)\}$. $\square$

Definition 13.2.0.4. Let X be a unital Banach algebra and $x \in X$. We define the **resolvent function** of x , denoted $R_x : \rho(x) \rightarrow G(X)$, by

$$R_x(\lambda) = (\lambda e - x)^{-1}$$

Exercise 13.2.0.5. Let X be a unital Banach algebra and $x \in X$. Then

1. $R_x : \rho(x) \rightarrow G(X)$ is Frechet differentiable and for each $\lambda \in \rho(x)$,

$$R'_x = -R_x^2$$

2. $R_x \in C^\infty(\rho(x))$ and for each $n \in \mathbb{N}$, $R_x^{(n)} = (-1)^n n! R_x^{n+1}$

Proof.

1. Define $S_x : \rho(x) \rightarrow G(X)$ by $S_x(\lambda) = \lambda e - x$. Then $R_x = \text{inv} \circ S_x$. Since S_x and inv are differentiable, $R_x = \text{inv} \circ S_x$ is differentiable. Previous exercises imply that for each $\lambda \in \rho(x)$, we have that

$$\begin{aligned} R'_x(\lambda) &= DR_x(\lambda)(1) \\ &= [D\text{inv}(S_x(\lambda)) \circ DS_x(\lambda)](1) \\ &= D\text{inv}(S_x(\lambda))(DS_x(\lambda)(1)) \\ &= D\text{inv}(S_x(\lambda))(e) \\ &= -S_x(\lambda)^{-1} e S_x(\lambda)^{-1} \\ &= -S_x(\lambda)^{-2} \\ &= -R_x(\lambda)^2 \end{aligned}$$

2. Let $n \in \mathbb{N}$. Suppose that $R_x \in C^{n-1}(\rho(x))$ and $R_x^{(n-1)} = (-1)^{n-1} (n-1)! R_x^n$. Then

$$\begin{aligned} R_x^{(n)} &= (R_x^{(n-1)})' \\ &= [(-1)^{n-1} (n-1)! R_x^n]' \\ &= (-1)^{n-1} (n-1)! (n R_x^{n-1}) (-R_x^2) \\ &= (-1)^n n! R_x^{n+1} \end{aligned}$$

By induction, for each $n \in \mathbb{N}$, $R_x \in C^n(\rho(x))$ and $R_x^{(n)} = (-1)^n n! R_x^{n+1}$. A previous exercise in the section of differentiability implies that for each $n \in \mathbb{N}$, $R_x \in C^n(\rho(x))$ iff $R_x^{(n)} \in C^0(\rho(x))$. Hence for each $n \in \mathbb{N}$, $R_x \in C^n(\rho(x))$ and therefore $R_x \in C^\infty(\rho(x))$. $\square$

Exercise 13.2.0.6. Let X be a unital Banach algebra and $x \in X$. Then $\sigma(x) \neq \emptyset$.

Hint: R_x is bounded and apply Liouville's theorem

Proof. Suppose that $\sigma(x) = \emptyset$. Then $\rho(x) = \mathbb{C}$ and the previous exercise implies that $R_x : \mathbb{C} \rightarrow G(X)$ is differentiable. We observe that for each $\lambda \in \mathbb{C}^\times$,

$$\begin{aligned} R_x(\lambda) &= (\lambda e - x)^{-1} \\ &= \lambda^{-1}(e - \lambda^{-1}x)^{-1} \end{aligned}$$

Since $\lambda^{-1} \rightarrow 0$ as $\lambda \rightarrow \infty$ and $\text{inv} : G(X) \rightarrow G(X)$ is continuous, we have that

$$\begin{aligned} (e - \lambda^{-1}x)^{-1} &\rightarrow e^{-1} \\ &= e \end{aligned}$$

as $\lambda \rightarrow \infty$. Hence $R_x(\lambda) \rightarrow 0$ as $\lambda \rightarrow \infty$. Thus $R_x : \mathbb{C} \rightarrow G(X)$ is bounded. Louville's theorem implies that $R_x = 0$. This is a contradiction since $0 \notin G(X)$. $\square$

Exercise 13.2.0.7. Gelfand-Mazur Theorem:

Let X be a unital Banach algebra. Suppose that $G(X) = X \setminus \{0\}$. Then $X = \mathbb{C}e$.

Proof. For the sake of contradiction, suppose that $X \neq \mathbb{C}e$. Then there exists $x \in X$ such that for each $\lambda \in \mathbb{C}$, $\lambda e - x \neq 0$. Let $\lambda \in \mathbb{C}$. Since $G(X) = X \setminus \{0\}$, $\lambda e - x \in G(X)$ and $\lambda \in \rho(x)$. Since $\lambda \in \mathbb{C}$ is arbitrary, we have that $\rho(x) = \mathbb{C}$. Therefore $\sigma(x) = \emptyset$. This is a contradiction since Exercise 17.3.0.5 **the previous exercise** implies that $\sigma(x) \neq \emptyset$. Thus $X = \mathbb{C}e$. $\square$

Note 13.2.0.8. In particular, if $X \setminus \{0\} = G(X)$, then $X \cong \mathbb{C}$.

Definition 13.2.0.9. Let X be a unital Banach algebra and $x \in X$. We define the **spectral radius of x** , denoted by $r(x)$, by

$$r(x) = \sup\{|\lambda| : \lambda \in \sigma(x)\}$$

Exercise 13.2.0.10. Let X be a unital Banach algebra and $x \in X$. Then

1. $r(x) \leq \liminf \|x^n\|^{1/n}$
2. for each $n \in \mathbb{N}$, $\lambda \in \bar{B}(0, r(x))^c$ and $\phi \in X^*$, $|\phi((\lambda x)^n)| < \infty$ and $\phi \in X^*$, $\phi \circ R_x(\lambda) = \sum_{n=0}^{\infty} \lambda^{-(n+1)} \phi(x)^n$ is **Hint:** uniform boundedness principle

Proof.

1. Let $\lambda \in \sigma(x)$ and $n \in \mathbb{N}$. The previous exercise implies that $\lambda^n \in \sigma(x^n)$. Since $\lambda^n e - x^n \notin G(X)$, we have that

$$\begin{aligned} |\lambda|^n &= |\lambda^n| \\ &\leq \|x^n\| \end{aligned}$$

Therefore $|\lambda| \leq \|x^n\|^{1/n}$. Since $n \in \mathbb{N}$ is arbitrary, $|\lambda| \leq \liminf \|x^n\|^{1/n}$. Since $\lambda \in \sigma(x)$ is arbitrary, we have that

$$\begin{aligned} r(x) &= \sup_{\lambda \in \sigma(x)} |\lambda| \\ &\leq \liminf \|x^n\|^{1/n} \end{aligned}$$

2. Let $\lambda \in \mathbb{C}$. Suppose that $|\lambda| > r(x)$. Then $\lambda \in \rho(x)$. **A previous exercise** implies that $|\lambda| > \|x\|$. Let $\phi \in X^*$. Then

$$\begin{aligned} \phi \circ R_x(\lambda) &= \phi \left(\sum_{n=0}^{\infty} \lambda^{-(n+1)} x^n \right) \\ &= \lambda^{-1} \sum_{n=0}^{\infty} \phi((\lambda^{-1}x)^n). \end{aligned}$$

Hence

$$\begin{aligned}\|\phi \circ R_x(\lambda)\| &= \sum_{n=0}^{\infty} \|\phi\| \|\lambda\|^{-(n+1)} \|x\|^n \\ &< \infty\end{aligned}$$

and therefore $|\phi \circ R_x(\lambda)| \leq \sum_{n=0}^{\infty} r(x)^{-(n+1)} \|\phi\|^n \|x\|^n$. Thus for each $n \in \mathbb{N}$ and $\lambda \in \bar{B}(0, r(x))^c$, **FINISH!!!, need to use holomorphicity, make section on complex analysis and also for Banach algebra valued functions, show that for $D \subset \mathbb{C}$ open, $f : D \rightarrow X$ is holomorphic iff for each $\phi \in X^*$, $\phi \circ f$ is holomorphic.**

□

Exercise 13.2.0.11. Let X be a unital Banach algebra and $x \in X$. Then

$$r(x) = \lim_{n \rightarrow \infty} \|x^n\|^{1/n}$$

Definition 13.2.0.12. Let X, Y be Banach algebras and $\phi : X \rightarrow Y$. Then ϕ is said to be a **Banach algebra homomorphism** if

1. $\phi \in L(X, Y)$,
2. for each $a, b \in X$, $\phi(ab) = \phi(a)\phi(b)$.

Exercise 13.2.0.13. Let X, Y be Banach algebras and $\phi : X \rightarrow Y$. Suppose that X is unital and ϕ is a Banach algebra homomorphism. Then

1. $\phi(e) = 0$ iff $\phi = 0$,
2. $Y = \mathbb{C}$ implies that $\phi(e) \neq 0$ iff $\phi(e) = 1$.

Proof.

1. • ($\implies$) :
Suppose that $\phi(e) = 0$. Then for each $x \in X$,

$$\begin{aligned}\phi(x) &= \phi(xe) \\ &= \phi(x)\phi(e) \\ &= \phi(x)0 \\ &= 0.\end{aligned}$$

Thus $\phi = 0$

- ($\impliedby$) :
Suppose that $\phi = 0$. Then $\phi(e) = 0$.

2. Suppose that $Y = \mathbb{C}$.

- ($\implies$) :
Suppose that $\phi(e) \neq 0$. Then

$$\begin{aligned}\phi(e) &= \phi(ee) \\ &= \phi(e)\phi(e).\end{aligned}$$

Thus $\phi(e)(\phi(e) - 1) = 0$. Since $\phi(e) \neq 0$, $\phi(e) = 1$.

- ($\impliedby$) :
Suppose that $\phi(e) = 1$. Then clearly $\phi(e) \neq 0$.

□

Definition 13.2.0.14. Category of Banach algebras We define the **category of Banach algebras**, denoted **BanAlg**

Definition 13.2.0.15. Let X be a unital Banach algebra. We define the **spectrum of X** , denoted $\sigma(X)$, by $\sigma(X) := \text{Hom}_{\text{BanAlg}}(X, \mathbb{C}) \setminus \{0\}$.

Exercise 13.2.0.16. Let X be a unital Banach algebra and $\phi \in \sigma(X)$. Then

1. $\phi(e) \neq 0$,
2. $\phi(e) = 1$,
3. for each $x \in G(X)$, $\phi(x) \neq 0$,
4. $\|\phi\| \leq 1$.

Proof.

1. Since $\phi \in \sigma(X)$, $\phi \neq 0$. **The previous exercise** implies $\phi(e) \neq 0$.
2. **the previous exercise** implies that $\phi(e) = 1$ iff $\phi(e) \neq 0$. Since the previous part implies that $\phi(e) \neq 0$, we have that $\phi(e) = 1$.
3. Let $x \in G(X)$. Then

$$\begin{aligned}\phi(x)\phi(x^{-1}) &= \phi(xx^{-1}) \\ &= \phi(e) \\ &= 1\end{aligned}$$

and similarly, $\phi(x^{-1})\phi(x) = 1$. Hence $\phi(x) \neq 0$ and $\phi(x)^{-1} = \phi(x^{-1})$. $\phi(x) \neq 0$.

4. Let $x \in X$. If $\phi(x) = 0$, then

$$\begin{aligned}|\phi(x)| &= |0| \\ &= 0 \\ &\leq \|x\|.\end{aligned}$$

Suppose that $\phi(x) \neq 0$. For the sake of contradiction, suppose that $|\phi(x)| > \|x\|$. Define $\lambda \in \mathbb{C}$ by $\lambda := \phi(x)$. Since $|\lambda| > \|x\|$, **A previous exercise** then implies that $\lambda e - x \in G(X)$. **The previous part** implies that

$$\begin{aligned}\phi(x) - \phi(x) &= \lambda - \phi(x) \\ &= \lambda\phi(e) - \phi(x) \\ &= \phi(\lambda e - x) \\ &\neq 0.\end{aligned}$$

This is a contradiction. Hence $|\phi(x)| \leq \|x\|$. Since $x \in X$ is arbitrary, we have that for each $x \in X$, $|\phi(x)| \leq \|x\|$. Thus $\|\phi\| \leq 1$. □

Exercise 13.2.0.17. Let X be a unital Banach algebra. Then $\sigma(X)$ is weak-* compact.

Proof. The Banach-Alaoglu theorem Exercise ?? (**reference here**) implies that $\bar{B}_{X^*}(0, 1)$ is weak-* compact. Exercise 17.3.0.5 **the previous exercise** implies that $\sigma(X) \subset \bar{B}_{X^*}(0, 1)$. Let $(\phi_\alpha)_{\alpha \in A} \subset \sigma(X)$ and $\phi \in \bar{B}_{X^*}(0, 1)$. Suppose that $\phi_\alpha \xrightarrow{w^*} \phi$. Then

- Let $x, y \in X$. Since multiplication $\mu : X \times X \rightarrow X$ is norm continuous, μ is weak-* continuous. Thus

$$\begin{aligned}\phi(xy) &= \lim_{\alpha \in A} \phi_\alpha(xy) \\ &= \lim_{\alpha \in A} [\phi_\alpha(x)\phi_\alpha(y)] \\ &= [\lim_{\alpha \in A} \phi_\alpha(x)][\lim_{\alpha \in A} \phi_\alpha(y)] \\ &= \phi(x)\phi(y).\end{aligned}$$

So $\phi \in \text{Hom}_{\text{BanAlg}}(X, \mathbb{C})$.

- Since for each $\alpha \in A$, $\phi_\alpha \in \sigma(X)$, a **previous exercise** Exercise ?? implies that for each $\alpha \in A$, $\phi_\alpha(e) = 1$. Thus

$$\begin{aligned}\phi(e) &= \lim_{\alpha \in A} \phi_\alpha(e) \\ &= 1.\end{aligned}$$

Hence $\phi \neq 0$.

Thus $\phi \in \sigma(X)$. Since $(\phi_\alpha)_{\alpha \in A} \subset \sigma(X)$ and $\phi \in \bar{B}_{X^*}(0, 1)$ with $\phi_\alpha \xrightarrow{w^*} \phi$ are arbitrary, we have that for each $(\phi_\alpha)_{\alpha \in A} \subset \sigma(X)$ and $\phi \in \bar{B}_{X^*}(0, 1)$, $\phi_\alpha \xrightarrow{w^*} \phi$ implies that $\phi \in \sigma(X)$. Hence $\sigma(X)$ is weak-* closed in $\bar{B}_{X^*}(0, 1)$. Thus $\sigma(X)$ is weak-* compact. $\square$

Definition 13.2.0.18. Let X be a Banach algebra and $J \subset X$ a subalgebra. Then J is said to be a

- **left ideal of X** if for each $x \in X$, $y \in J$, $xy \in J$,
- **right ideal of X** if for each $x \in J$, $y \in X$, $xy \in J$,
- **ideal of X** if J is a left ideal of X and J is a right ideal of X .

Definition 13.2.0.19. Let X be a Banach algebra. We define $\mathcal{I}(X) := \{J \subset X : J \text{ is an ideal of } X\}$

Exercise 13.2.0.20. Let X be a Banach algebra and $J \subset X$ an ideal of X . Then $\text{cl } J$ is a ideal of X .

Proof. Let $x \in X$ and $y \in \text{cl } J$. Then there exists $(y_n)_{n \in \mathbb{N}} \subset J$ such that $y_n \rightarrow y$. Since multiplication $\mu : X \times X \rightarrow X$ is continuous, we have that $xy_n \rightarrow xy$. Since J is a left ideal of X , we have that $(xy_n)_{n \in \mathbb{N}} \subset J$. Since there exists $(z_n)_{n \in \mathbb{N}} \subset J$ such that $z_n \rightarrow xy$, we have that $xy \in \text{cl } J$. Since $x \in X$ and $y \in \text{cl } J$ are arbitrary, we have that for each $x \in X$ and $y \in \text{cl } J$, $xy \in \text{cl } J$. So $\text{cl } J$ is a left ideal of X . Similarly, $\text{cl } J$ is a right ideal of X . Hence $\text{cl } J$ is an ideal of X . $\square$

Definition 13.2.0.21. Let X be a Banach algebra and $J \in \mathcal{I}(X)$. Then J is said to be

- **proper** if $J \neq X$,
- **maximal** if J is proper and for each $J' \in \mathcal{I}(X)$, $J \subset J'$ implies that $J = J'$ or $J' = X$.

Exercise 13.2.0.22. Let X be a unital Banach algebra and $J \in \mathcal{I}(X)$. Then the following are equivalent:

1. J is proper
2. $e \notin J$
3. $J \cap G(X) = \emptyset$

Proof.

1. (1) $\implies$ (2) :
Suppose that $e \in J$. Then for each $x \in X$,

$$\begin{aligned}x &= xe \\ &\in J.\end{aligned}$$

So $X \subset J$ and $J = X$. Hence J is not proper.

2. (2) $\implies$ (3) :
Suppose that $J \cap G(X) \neq \emptyset$. Then there exists $x \in G(X)$ such that $x \in J$. Then

$$\begin{aligned}e &= x^{-1}x \\ &\in J.\end{aligned}$$

3. (3) $\implies$ (1) :
Suppose that $J \cap G(X) = \emptyset$. Since $e \in G(X)$, we have that $e \notin J$. Hence $J \neq X$ and J is proper.

□

Exercise 13.2.0.23. Let X be a unital Banach algebra and $J, K \in \mathcal{T}(X)$. Then

1. $J + K \in \mathcal{I}(X)$
2. $JK \in \mathcal{I}(X)$

FINISH!!!

Proof. content...

□

Exercise 13.2.0.24. Let X be a unital Banach algebra and $J \in \mathcal{I}(X)$. Suppose that J is a proper. Then

1. $\text{cl } J$ is proper,
2. there exists $J' \in \mathcal{I}(X)$ such that J' is a maximal and $J \subset J'$,
3. if J is maximal, then J is closed.

Proof.

1. Exercise 13.2.0.20 implies that $\text{cl } J$ is an ideal of X . Since J is proper, Exercise 13.2.0.22 implies that $J \subset G(X)^c$. Exercise 13.1.0.17 implies that $G(X)^c$ is closed. Hence $\text{cl } J \subset G(X)^c$. Another application of Exercise 13.2.0.22 then implies that $\text{cl } J$ is proper.
2. **zorn's lemma FINISH!!!**
3. Suppose that J is maximal. Since $J \subset \text{cl } J$ and J is maximal, we have that $J = \text{cl } J$ or $\text{cl } J = X$. Since J is maximal, J is proper. The previous part implies that $\text{cl } J$ is proper. Thus $\text{cl } J \neq X$. Hence $J = \text{cl } J$.

□

Exercise 13.2.0.25. Let X be a unital Banach algebra and $J \in \mathcal{I}(X)$. If J is maximal, then X/J is a Banach algebra.

Proof. Suppose that J is maximal. **A previous exercise in section on Banach spaces** implies that X/J is a Banach space. Let $a, b \in X/J$ and $\epsilon > 0$. Then there exist $x, y \in X$ such that $a = x + J$ and $b = y + J$. Set

$$\epsilon_0 := \min \left(\frac{\epsilon}{3(\|x + J\| + 1)}, \frac{\epsilon}{3(\|y + J\| + 1)}, \sqrt{\frac{\epsilon}{3}} \right)$$

Then $\epsilon_0 > 0$. Thus there exist $z, w \in J$ such that $\|x + z\| < \|x + J\| + \epsilon_0$ and $\|y + w\| < \|y + J\| + \epsilon_0$. Since J is an ideal of X , we have that $xw, zy, zw \in J$. Therefore $xw + zy + zw \in J$ and

$$\begin{aligned} \|ab\| &= \|xy + J\| \\ &= \inf_{q \in J} \|xy + q\| \\ &\leq \|xy + xw + zy + zw\| \\ &= \|(x + z)(y + w)\| \\ &\leq \|x + z\| \|y + w\| \\ &< (\|x + J\| + \epsilon_0)(\|y + J\| + \epsilon_0) \\ &= \|x + J\| \|y + J\| + \epsilon_0 \|x + J\| + \epsilon_0 \|y + J\| + \epsilon_0^2 \\ &< \|x + J\| \|y + J\| + \frac{\epsilon}{3} + \frac{\epsilon}{3} + \frac{\epsilon}{3} \\ &= \|x + J\| \|y + J\| + \epsilon \\ &= \|a\| \|b\| + \epsilon. \end{aligned}$$

Since $\epsilon > 0$ is arbitrary, we have that for each $\epsilon > 0$, $\|ab\| < \|a\| \|b\| + \epsilon$. Therefore $\|ab\| \leq \|a\| \|b\|$. Since $a, b \in X/J$ are arbitrary, we have that for each $a, b \in X/J$, $\|ab\| \leq \|a\| \|b\|$. Hence X/J is a Banach algebra. □

Note 13.2.0.26. We denote the canonical projection of X onto X/J by $\pi : X \rightarrow X/J$.

Exercise 13.2.0.27. Let X be a unital Banach algebra, $J \in \mathcal{I}(X)$ and $K \in \mathcal{I}(X/J)$. Then

1. $\pi^{-1}(K)$ is an ideal of X and $J \subset \pi^{-1}(K)$,
2. X is commutative implies that J is maximal iff that $X/J = \mathbb{C}(e + J)$.

Proof.

1. **The preimage of an ideal is an ideal.** Let $x \in X$ and $y \in \pi^{-1}(K)$. Then $\pi(y) \in K$. Since $K \in \mathcal{I}(X/J)$, we have that $\pi(x)\pi(y) \in K$. Therefore

$$\begin{aligned}\pi(xy) &= \pi(x)\pi(y) \\ &\in K.\end{aligned}$$

Hence $xy \in \pi^{-1}(K)$. Since $x \in X$ and $y \in \pi^{-1}(K)$ is arbitrary, we have that for each $x \in X$ and $y \in \pi^{-1}(K)$, $xy \in \pi^{-1}(K)$.

2. Suppose that X is commutative. **A previous exercise** implies that X/J is a commutative unital Banach algebra. Since J is maximal, **a previous exercise, or exercise in appendices** implies that X/J is a field. The Gelfand-Mazur theorem Exercise 17.3.0.5 (reference) implies that $X/J = \mathbb{C}(e + J)$.

□

Exercise 13.2.0.28.

1. $\mathcal{I}(M_2(\mathbb{C})) = \{(0)\}$.
2. $M_2(\mathbb{C})/(0)$ is not a division ring.

Proof.

- 1.
- 2.

□

Definition 13.2.0.29. Let X be a Banach algebra. We define $\mathcal{I}_{\max}(X) := \{J \in \mathcal{I}(X) : J \text{ is a maximal}\}$.

Exercise 13.2.0.30. Let X be a unital Banach algebra.

1. For each $\phi \in \sigma(X)$, $\ker \phi \in \mathcal{I}_{\max}(X)$.
Hint: Let $J' \in \mathcal{I}(X)$. Suppose that $\ker \phi \subset J'$ and $\ker \phi \neq J'$. Write $e = a + b$ with $a \in \ker \phi$ and $b \in J' \setminus \ker \phi$.
2. Define $\alpha : \sigma(X) \rightarrow \mathcal{I}_{\max}(X)$ by $\alpha(\phi) = \ker \phi$. Then α is a bijection.

Proof.

1. Let $\phi \in \sigma(X)$. Let $x \in X$ and $y \in \ker \phi$. Then

$$\begin{aligned}\phi(xy) &= \phi(x)\phi(y) \\ &= \phi(x)0 \\ &= 0.\end{aligned}$$

Thus $xy \in \ker \phi$. Since $x \in X$ and $y \in \ker \phi$ are arbitrary, we have that for each $x \in X$ and $y \in \ker \phi$, $xy \in \ker \phi$. Hence $\ker \phi$ is a left ideal of X . Similarly, $\ker \phi$ is a right ideal of X . Hence $\ker \phi \in \mathcal{I}(X)$. Since $\phi \in \sigma(X)$, $\phi \neq 0$ and therefore $\ker \phi \neq X$. So $\ker \phi$ is proper. Let $J' \in \mathcal{I}(X)$. Suppose $\ker \phi \subset J'$ and $\ker \phi \neq J'$. Then there exists $x \in J'$ such that $x \notin \ker \phi$. Hence $\phi(x) \neq 0$. Set $x' := \phi(x)^{-1}x$. Then

$$\begin{aligned}\phi(e - x') &= \phi(e) - \phi(x') \\ &= 1 - \phi(x)^{-1}\phi(x) \\ &= 1 - 1 \\ &= 0.\end{aligned}$$

Thus

$$\begin{aligned} e - x' &\in \ker \phi \\ &\subset J'. \end{aligned}$$

Since J' is an ideal of X and $x \in J'$, we have that $[\phi(x)^{-1}e]x \in J'$. Therefore

$$\begin{aligned} e &= e - x' + x' \\ &= (e - x') + [\phi(x)^{-1}e]x \\ &\in J' \end{aligned}$$

A previous exercise implies that $J' = X$. Since $J' \in \mathcal{I}(X)$ with $\ker \phi \subset J'$ and $\ker \phi \neq J'$ is arbitrary, we have that for each $J' \in \mathcal{I}(X)$, $\ker \phi \subset J'$ and $\ker \phi \neq J'$ implies that $J' = X$. Thus $\ker \phi$ is maximal.

2. Gelfand-Mazur Theorem

□

Exercise 13.2.0.31. If $\sigma(X) \neq \emptyset$ and $\dim X \geq 2$ (i.e. $X \neq \mathbb{C}e$), then $\mathcal{I}(X) \neq \{(0)\}$. If X is commutative, then for each $x \in G(X)$, there exists $J \in \mathcal{I}(X)$ such that $x \in J$ (so that if $x \neq 0$ and $x \in G(X)$, then $\mathcal{I}(X) \neq \{(0)\}$). Thus there exists $\phi \in \sigma(X)$ since $\sigma(X)$ is bijective to $\mathcal{I}(X)$, so commutativity implies that.

note: For example $\mathcal{I}(\mathbb{C}^{n \times n}) = \{(0)\}$. why isnt then $\mathbb{C}^{n \times n}/(0)$ a division ring? so it must not be the case that division by a maximal ideal always results in a division ring see first stackexchange answer

Proof. Let $\phi \in \sigma(X)$. Then $\ker \phi$ is a maximal ideal of X . Since $\dim X \geq 2$, $\dim \ker \phi \geq 1$. Hence $\ker \phi \neq (0)$. □

• $\implies$:

• $\Longleftarrow$:

• $\implies$:

• $\Longleftarrow$:

13.2.1 *-Algebras

Definition 13.2.1.1. Let X be an algebra and $*$: $X \rightarrow X$.

- Then $*$ is said to be an **involution on X** if for each $x, y \in X$ and $\lambda \in \mathbb{C}$,

1. $(x + y)^* = x^* + y^*$,
2. $(\lambda x)^* = \lambda^* x^*$,
3. $(xy)^* = y^* x^*$,
4. $(x^*)^* = x$.

- If X is a Banach algebra, then $(X, *)$ is said to be a **C^* -algebra** if for each $x \in X$, $\|x^* x\| = \|x\|^2$.

Note 13.2.1.2. We note that $*$ $\in \text{Hom}_{\mathbf{Vect}_{\mathbb{C}}}(X, \text{conj}(X))$ and therefore $0^* = 0$. (need to define the conjugate of a vector space $\text{conj}(X)$)

Exercise 13.2.1.3. Let X be a C^* -algebra. Then

1. for each $x \in X$, $\|x\| = \|x^*\|$
2. for each $x, y \in X$, $\|x^* y\| = \|y^* x\|$
3. $*$ $\in L(X, \text{conj}(X))$.

Proof. 1. Let $x \in X$. If $x = 0$, then $x^* = 0$ and therefore $\|x\| = \|x^*\|$. Suppose that $\|x\| \neq 0$. Then

$$\begin{aligned} \|x\|^2 &= \|x^* x\| \\ &\leq \|x^*\| \|x\|. \end{aligned}$$

Thus $\|x\| \leq \|x^*\|$. Similarly, we have that $\|x^*\| \leq \|(x^*)^*\|$. Therefore

$$\begin{aligned} \|x\| &\leq \|x^*\| \\ &\leq \|(x^*)^*\| \\ &= \|x\|. \end{aligned}$$

Hence $\|x^*\| = \|x\|$.

2. Let $x, y \in X$. We have that

$$\begin{aligned} \|x^* y\| &= \|(x^* y)^*\| \\ &= \|y^* (x^*)^*\| \\ &= \|y^* x\|. \end{aligned}$$

3. As noted above, $*$ $\in \text{Hom}_{\mathbf{Vect}_{\mathbb{C}}}(X, \text{conj}(X))$. The previous part implies that for each $x \in X$, $\|x^*\| = \|x\|$. Hence $\|*\| \leq 1$. Similarly,

$$\begin{aligned} \|*\| &= \sup_{\|x\|=1} \|x^*\| \\ &= \end{aligned}$$

□

Exercise 13.2.1.4. Let X be a C^* -algebra. Then $*$: $X \rightarrow X$ is continuous.

Proof.

□

13.3 Exponential Map

Note 13.3.0.1. ref summation appendix Exercise 9.1.0.4 exercise about $(x_n) \in l_X^1(\mathbb{N})$ implies $\sum_{n \in \mathbb{N}} x_n \in X$.

Exercise 13.3.0.2. Let X be a Banach space. Then for each $x \in X$, $(x^n/n!) \in l_X^1(\mathbb{N})$.

Proof. For each $x \in X$,

$$\begin{aligned} \sum_{j \in \mathbb{N}_0} \left\| \frac{x^n}{n!} \right\| &\leq \sum_{n \in \mathbb{N}_0} \frac{\|x\|^n}{n!} \\ &= e^{\|x\|} \\ &< \infty \end{aligned}$$

and therefore $(x^n/n!) \in l_X^1(\mathbb{N})$. □

Definition 13.3.0.3. Let X be a Banach algebra. We define the **exponential map on** X , denoted $\exp : X \rightarrow X$, by

$$\exp(x) := \sum_{j \in \mathbb{N}_0} \frac{x^j}{j!}$$

Exercise 13.3.0.4.

Chapter 14

Differential Equations

14.1 Introduction

- note: need to define differentiation $A \subset X$ $f : A \rightarrow Y$ and A is not open, i.e. f is differentiable at $x_0 \in \text{int } A$ if there exists $U \in \mathcal{N}^{\text{opn}}(x, \mathcal{T}_X)$ and $g : U \rightarrow Y$ such that g is differentiable at x_0 and $g|_{U \cap A} = f|_{U \cap A}$, then show that this is independent of g .
- need to define

$$\int_{t_0}^t f \, dm$$

as

$$\int_{(t_0, t]} f \, dm$$

if $t \geq t_0$ and

$$\int_{(t, t_0]} f \, dm$$

if $t < t_0$, maybe have to connect this to the stieltjes integral or otherwise justify it, see intro to math notes section on FTC for similar reason to motivate, basically want to extend $F : [t_0, b] \rightarrow \mathbb{R}/X$ to $G : [a, b] \rightarrow \mathbb{R}/X$.

Definition 14.1.0.1. Let X be a Banach space, $\Omega \subset \mathcal{T}_{\mathbb{R} \times X}$, $f : \Omega \rightarrow X$, $(t_0, x_0) \in \Omega$. Let $I \subset \text{int } \pi_1(\Omega)$ and $u : I \rightarrow \pi_2(\Omega)$. Suppose that u is differentiable. Then u is said to **solve the initial value problem associated to f and (t_0, x_0)** if

1. $u' = f(\cdot, u)$
2. $u(t_0) = x_0$.

Exercise 14.1.0.2. Let X be a Banach space, $\Omega \subset \mathbb{R} \times X$, $f : \Omega \rightarrow X$, $(t_0, x_0) \in \text{int } \Omega$, $I \subset \text{int } \pi_1(\Omega)$ and $u : I \rightarrow \pi_2(\Omega)$. Suppose that u is differentiable and u solves the initial value problem associated to f and (t_0, x_0) . Then for each $n \in \mathbb{N}_0$, $f \in C^n(\Omega, X)$ implies that $u \in C^{n+1}(I, X)$.

Proof. Let $n \in \mathbb{N}_0$. Suppose that $f \in C^n(\Omega, X)$. We proceed by induction on n .

- **Base Case:**

Suppose that $n = 0$. By assumption, $u' = f(\cdot, u)$. Since u' is differentiable, u' is continuous. Since $f \in C(\Omega, X)$,

$$\begin{aligned} u' &= f(\cdot, u) \\ &\in C(\Omega, X). \end{aligned}$$

Thus $u \in C^1(I, X)$.

- **Induction Step:**

Let $k \in \mathbb{N}$. Suppose that **the claim** is true for $n = k - 1$. Suppose that $f \in C^k(\Omega, X)$. Then $f \in C^{k-1}(\Omega, X)$. By **the induction**

hypothesis, $u \in C^k(I, X)$. (ref ex here) then implies that $(u') \in C^{k-1}(I, X)$. Since $f \in C^k(\Omega, X)$, (make/ref generalized chain rule : $f \in C^k, g \in C^k$ implies $g \circ f \in C^k$) implies that

$$\begin{aligned} u^{(k)} &= (u')^{(k-1)} \\ &= [f(\cdot, u)]^{(k-1)} \\ &\in C^1(I, X). \end{aligned}$$

(ref ex) then implies that $u \in C^{k+1}(I, X)$. Thus the claim is true for $n = k$. Since $k \in \mathbb{N}$ is arbitrary, we have that for each $k \in \mathbb{N}$, the claim is true for $n = k - 1$ implies that the claim is true for $n = k$.

By induction, we have that for each $n \in \mathbb{N}_0$, $f \in C^n(\Omega, X)$ implies that $u \in C^{n+1}(I, X)$. **FINISH!!!** $\square$

If we are given a solution $u \in C(I, X)$ to the initial value problem above, then intuitively **taylor theorem section (FTC)** implies that

$$\begin{aligned} u(t) &= u(t_0) + \int_{t_0}^t f(s, u(s)) dm(s) \\ &= x_0 + \int_{t_0}^t f(s, u(s)) dm(s). \end{aligned}$$

Thus u is a fixed point of the map $T : C^1(I, X) \rightarrow C^1(I, X)$ given by

$$[Tu](t) := x_0 + \int_{t_0}^t f(s, u(s)) ds.$$

If we were able to deduce under which assumptions, if any, the map T is a contraction mapping, then under such assumptions, the Banach fixed-point theorem Exercise 17.3.0.5 (rex ex) would guarantee the existence and uniqueness of a fixed point of T (i.e. a solution to the initial value problem) which could be approximated by starting with any $\phi_0 \in C_1(I, X)$, (e.g. $\phi_0(t) := x_0$), and then iterating to obtain the sequence $(\phi_n)_{n \in \mathbb{N}} \subset C(I, X)$ defined by $\phi_n := T^n \phi_0$, which converge exponentially (**CHECK THIS MOTIVATION!!!**)

Exercise 14.1.0.3. Picard-Lindelöf Theorem:

Let X be a separable (maybe just f separably valued) Banach space, $\Omega \subset \mathbb{R} \times X$, $(t_0, x_0) \in \text{Int } \Omega$ and $f : \Omega \rightarrow X$. Suppose that f is continuous, $\sup_{t \in \pi_1(\Omega)} \text{Lip}(f(t, \cdot)) < \infty$ and for each $A \subset \Omega$, A is bounded implies $f|_A$ is bounded. Then there exists $\epsilon > 0$, $B \subset \pi_2(\Omega)$ and a unique $u : [t_0 - \epsilon, t_0 + \epsilon] \rightarrow B$ such that u is differentiable and u solves the initial value problem associated to f and (t_0, x_0) .

Proof.

• Existence:

- Define $C \in \mathbb{R}_+$ by $C := \sup_{t \in \pi_1(\Omega)} \text{Lip}(f(t, \cdot))$. Define $\rho \in (0, 1)$ by $\rho := C(C + 1)^{-1}$. Since $(t_0, x_0) \in \text{Int } \Omega$, (ref ex in product topology section) implies that there exist $U \in \mathcal{N}^{\text{opn}}(t_0, \mathcal{T}_{\mathbb{R}})$ and $V \in \mathcal{N}^{\text{opn}}(x_0, \mathcal{T}_X)$ such that

$$\begin{aligned} (t_0, x_0) &\in U \times V \\ &\subset \text{Int } \Omega. \end{aligned}$$

Since $V \in \mathcal{N}^{\text{opn}}(x_0, \mathcal{T}_X)$, there exists $\delta_0 > 0$ such that $B_X(x_0, \delta_0) \subset V$. Define $\delta > 0$ and $B \subset X$ by $\delta := \delta_0/2$ and $B := \bar{B}_X(x_0, \delta)$. Since $U \in \mathcal{N}^{\text{opn}}(t_0, \mathcal{T}_{\mathbb{R}})$, there exists $\epsilon_0 > 0$ such that $(t_0 - \epsilon_0, t_0 + \epsilon_0) \subset U$. Define $\epsilon > 0$, $I \subset \mathbb{R}$ by $\epsilon := \min\{\epsilon_0/2, \delta(M + 1)^{-1}, (C + 1)^{-1}\}$ and $I := [t_0 - \epsilon, t_0 + \epsilon]$. Since $I \times B$ is bounded, $f|_{I \times B}$ is bounded. Define $M \in \mathbb{R}_+$ by $M := \sup_{(t, x) \in I \times B} \|f\|$

- Since X is separable (or f separably valued) and f is continuous, ex from taylor theorem section and the FTC imply that for each $u \in C(I, \pi_2(\Omega))$,

$$\begin{aligned} f(\cdot, u) &\in C(I, X) \\ &\subset L_X^1(I, \mathcal{B}(I), m). \end{aligned}$$

Define $T : C(I, B) \rightarrow X^I$ by

$$[Tu](t) := x_0 + \int_{t_0}^t f(s, u(s)) dm(s).$$

(ref first ex in taylor theorem section) then implies that

$$\begin{aligned} T(C(I, B)) &\subset C^1(I, X) \\ &\subset C(I, X). \end{aligned}$$

Let $u \in C(I, B)$. Then for each $t \in I$,

$$\begin{aligned} \|[Tu](t) - x_0\| &= \left\| \left(x_0 + \int_{t_0}^t f(s, u(s)) dm(s) \right) - x_0 \right\| \\ &= \left\| \int_{t_0}^t f(s, u(s)) dm(s) \right\| \\ &\leq \int_{t_0}^t \|f(s, u(s))\| dm(s) \\ &\leq \int_{t_0}^t M dm(s) \\ &\leq M|t - t_0| \\ &\leq M\epsilon \\ &\leq M \frac{\delta}{M+1} \\ &< \delta. \end{aligned}$$

Thus $T(u) \in C(I, B)$. Since $u \in C(I, B)$ is arbitrary, we have that $T(C(I, B)) \subset C(I, B)$.

– Let $u, v \in C(I, B)$. Since for each $t \in I$,

$$\begin{aligned} \|Tu(t) - Tv(t)\| &= \left\| \left[x_0 + \int_{(t_0, t]} f(s, u(s)) dm(s) \right] - \left[x_0 + \int_{(t_0, t]} f(s, v(s)) dm(s) \right] \right\| \\ &= \left\| \int_{(t_0, t]} f(s, u(s)) dm(s) - \int_{(t_0, t]} f(s, v(s)) dm(s) \right\| \\ &= \left\| \int_{(t_0, t]} [f(s, u(s)) - f(s, v(s))] dm(s) \right\| \\ &\leq \int_{(t_0, t]} \|f(s, u(s)) - f(s, v(s))\| dm(s) \\ &\leq \int_{(t_0, t]} C\|u(s) - v(s)\| dm(s) \\ &= \int_{(t_0, t]} C\|u - v\|_\infty dm(s) \\ &= C\|u - v\|_\infty |t - t_0| \\ &< C\epsilon\|u - v\|_\infty \\ &\leq \frac{C}{C+1}\|u - v\|_\infty \\ &= \rho\|u - v\|_\infty, \end{aligned}$$

we have that $\|Tu - Tv\| \leq \rho\|u - v\|_\infty$. Since $u, v \in C(I, B)$ are arbitrary, we have that for each $u, v \in C(I, B)$, $\|Tu - Tv\| \leq \rho\|u - v\|_\infty$. Hence $T \in C_{\text{Lip}}(C(I, B), C(I, B))$ and T is a contraction mapping. Since B is closed, X is complete, and $B \subset X$, Exercise 17.3.0.5 (ref ex in completeness section on metric spaces) implies that B is complete. Since I is compact and B is complete, Exercise 17.3.0.5 (ref ex in completeness section on function spaces in metric spaces

chapter) implies that $C(I, B)$ is complete. Define $u_0 \in C(I, B)$ by $u_0(t) := x_0$. Exercise 6.7.1.13 then implies that there exists a unique $u_* \in C(I, B)$ such that $(T^n)(u_0) \xrightarrow{\text{unif}} u_*$ and

$$\begin{aligned} x_0 + \int_{t_0}^t f(s, u_*(s)) dm(s) &= T(u_*)(t) \\ &= u_*(t). \end{aligned}$$

1. (ref first ex in taylor theorem section) then implies that $u_* \in C^1(I, B)$ and

$$u'_* = f(\cdot, u_*).$$

2. We have that

$$\begin{aligned} u_*(t_0) &= x_0 + \int_{t_0}^{t_0} f(s, u_*(s)) dm(s) \\ &= x_0 + \int_{(t_0, t_0]} f(s, u_*(s)) dm(s) \\ &= x_0 + \int_{\emptyset} f(s, u_*(s)) dm(s) \\ &= x_0. \end{aligned}$$

Thus u_* solves the initial value problem associated to f and (t_0, x_0) .

• **Uniqueness:**

Let $u : I \rightarrow B$. Suppose that u is differentiable and u solves the initial value problem associated to f and (t_0, x_0) . Since f is continuous, the prev ex implies that $u' \in C(I, B) \subset L^1(I, \mathcal{B}(I), m)$ and an ex from Taylor theorem section implies that for each $t \in I$,

$$\begin{aligned} u(t) &= u(t_0) + \int_{t_0}^t u'(s) dm(s) \\ &= x_0 + \int_{t_0}^t f(s, u(s)) dm(s) \\ &= [Tu](t). \end{aligned}$$

Thus $Tu = u$. Uniqueness of u_* implies that $u = u_*$.

FINISH!!! (clean up)

□

14.2 Semigroup Theory

Chapter 15

Convex Analysis

15.1 Subadditive Functions

make ex about sup of subadditive functions being subadditive, etc, see [7]

Pulling this section from [1] and [7]

Definition 15.1.0.1. Let $(G, +)$ be a commutative semigroup, $A \subset G$ a subsemigroup and $f : A \rightarrow H$. Then f is said to be

- **subadditive** if for each $x, y \in A$, $f(x + y) \leq f(x) + f(y)$
- **superadditive** if $-f$ is subadditive.

Exercise 15.1.0.2. Fekete's Lemma I:

Let $(a_n)_{n \in \mathbb{N}} \subset \mathbb{R}$. Suppose that $(a_n)_{n \in \mathbb{N}}$ is subadditive. Then

1. $(a_n/n)_{n \in \mathbb{N}}$ is bounded,
2. $\frac{a_n}{n} \rightarrow \inf_{n \in \mathbb{N}} \frac{a_n}{n}$.

Proof. 1. Let $k \in \mathbb{N}$. Define $M_k \in \mathbb{R}$ by $M_k := \max_{k \in [k]_0} a_j$. Let $n \in \mathbb{N}$. Suppose that $n \geq k + 1$. The **division algorithm** (add to intro to math notes) implies that there exists $q \in \mathbb{N}_0$ and $r \in [k]_0$ such that $n = kq + r$. Then

$$\begin{aligned} \frac{a_n}{n} &= \frac{a_{kq+r}}{n} \\ &\leq \frac{a_{kq+r}}{n} \\ &\leq \frac{a_{kq} + a_r}{n} \quad (\text{by subadditivity}) \\ &= \frac{a_{kq}}{n} + \frac{a_r}{n} \\ &\leq \frac{qa_k}{n} + \frac{a_r}{n} \quad (\text{by subadditivity}) \\ &\leq \frac{qa_k}{kq} + \frac{a_r}{n} \\ &= \frac{a_k}{k} + \frac{a_r}{n} \\ &\leq \frac{a_k}{k} + \frac{M_k}{n}. \end{aligned}$$

Since $n \in \mathbb{N}$ with $n \geq k + 1$ is arbitrary, we have that for each $n \in \mathbb{N}$, $n \geq k + 1$ implies that $\frac{a_n}{n} \leq \frac{a_k}{k} + \frac{M_k}{n}$. In particular, we have that $(a_n/n)_{n \in \mathbb{N}}$ is bounded.

2. From above (ref above), we have that

$$\begin{aligned}\limsup_{n \rightarrow \infty} \frac{a_n}{n} &\leq \limsup_{n \rightarrow \infty} \left(\frac{a_k}{k} + \frac{M_k}{n} \right) \\ &\leq \limsup_{n \rightarrow \infty} \frac{a_k}{k} + \limsup_{n \rightarrow \infty} \frac{M_k}{n} \\ &= \frac{a_k}{k}.\end{aligned}$$

Since $k \in \mathbb{N}$ is arbitrary, we have that

$$\begin{aligned}\limsup_{n \rightarrow \infty} \frac{a_n}{n} &\leq \inf_{k \in \mathbb{N}} \frac{a_k}{k} \\ &= \inf_{n \in \mathbb{N}} \frac{a_n}{n} \\ &\leq \liminf_{n \rightarrow \infty} \frac{a_n}{n}.\end{aligned}$$

Hence $a_n/n \rightarrow \inf_{n \in \mathbb{N}} \frac{a_n}{n}$. □

Exercise 15.1.0.3. Let $A \subset \mathbb{R}$ be a subsemigroup. Suppose that for each $x, y \in A$, $x \leq y$ implies that $y - x \in A$. Let $f : A \rightarrow \mathbb{R}$. Suppose that f subadditive. Then for each $x, y \in A$, $x \leq y$ implies that $f(y) - f(x) \leq f(y - x)$.

Proof. Let $x, y \in A$. Suppose that $x \leq y$. Then

$$\begin{aligned}f(y) &= f(x + (y - x)) \\ &\leq f(x) + f(y - x).\end{aligned}$$

Therefore $f(y) - f(x) \leq f(y - x)$. Since $x, y \in A$ with $x \leq y$ are arbitrary, we have that for each $x, y \in A$, $x \leq y$ implies that $f(y) - f(x) \leq f(y - x)$. □

Exercise 15.1.0.4. Let $A \subset \mathbb{R}$ be a submonoid of $\mathbb{R}$ and $f : A \rightarrow \mathbb{R}$. If f is subadditive, then $f(0) \geq 0$.

Proof. Suppose that f is subadditive. Then

$$\begin{aligned}f(0) &= f(0 + 0) \\ &\leq f(0) + f(0).\end{aligned}$$

Therefore $0 \leq f(0)$. □

next ex pulling from [7]

Exercise 15.1.0.5. Fekete's Lemma II:

Let $f : [0, \infty) \rightarrow \mathbb{R}$. If f is subadditive, then

$$\lim_{t \rightarrow \infty} \frac{f(t)}{t} = \inf_{t > 0} \frac{f(t)}{t}.$$

FINISH!!! make bounded exercise

Proof. Set $\alpha := \inf_{t > 0} f(t)/t$.

- Suppose that $\alpha > -\infty$. Let $\epsilon > 0$. Then there exists $t_0 > 0$ such that $f(t_0)/t_0 < \alpha + \epsilon$. We note that for each $t \geq 2t_0$, $t/t_0 \geq 2$. For each $t \geq 2t_0$, define $n_t \in \mathbb{N}$ by $n_t := \lfloor t/t_0 - 1 \rfloor$. Let $t \geq 2t_0$. Then $(n_t + 1)t_0 \leq t \leq (n_t + 2)t_0$ and $t_0 \leq t - n_t t_0 \leq 2t_0$. (make prev exercise about boundedness of subadditive functions on compact sets) implies that there exist $m_{t_0}, M_{t_0} \in \mathbb{R}$ such that for each $s \in [t_0, 2t_0]$, $f(s) \in [m_{t_0}, M_{t_0}]$. Therefore

$$\begin{aligned}\frac{m_{t_0}}{t} &\leq \frac{f(t - n_t t_0)}{t} \\ &\leq \frac{M_{t_0}}{t}.\end{aligned}$$

Since $t \geq 2t_0$ is arbitrary, we have that for each $t \geq 2t_0$,

$$\frac{m_{t_0}}{t} \leq \frac{f(t - n_t t_0)}{t} \leq \frac{M_{t_0}}{t}.$$

Therefore

$$\lim_{t \rightarrow \infty} \frac{f(t - n_t t_0)}{t} = 0.$$

Since for each $t \geq 2t_0$,

$$\begin{aligned} \frac{n_t}{t} &= \frac{\lfloor t/t_0 - 1 \rfloor}{t} \\ &\leq \frac{\lfloor t/t_0 \rfloor}{t} \\ &\leq \frac{t/t_0}{t} \\ &= \frac{1}{t_0}, \end{aligned}$$

we have that

$$\limsup_{t \rightarrow \infty} \frac{n_t t_0}{t} \leq 1.$$

Since

$$\begin{aligned} \alpha &< \frac{f(t)}{t} \\ &= \frac{f(n_t t_0 + [t - n_t t_0])}{t} \\ &\leq \frac{f(n_t t_0)}{t} + \frac{f(t - n_t t_0)}{t} \\ &\leq \frac{n_t f(t_0)}{t} + \frac{f(t - n_t t_0)}{t} \\ &= \frac{n_t t_0}{t} \frac{f(t_0)}{t_0} + \frac{f(t - n_t t_0)}{t} \\ &< \frac{n_t t_0}{t} (\alpha + \epsilon) + \frac{M_{t_0}}{t}, \end{aligned}$$

we have that

$$\begin{aligned} \alpha &\leq \limsup_{t \rightarrow \infty} \frac{f(t)}{t} \\ &\leq \limsup_{t \rightarrow \infty} \left(\frac{n_t t_0}{t} (\alpha + \epsilon) + \frac{M_{t_0}}{t} \right) \\ &\leq \limsup_{t \rightarrow \infty} \frac{n_t t_0}{t} (\alpha + \epsilon) + \limsup_{t \rightarrow \infty} \frac{M_{t_0}}{t} \\ &= (\alpha + \epsilon) \limsup_{t \rightarrow \infty} \frac{n_t t_0}{t} (\alpha + \epsilon) + \lim_{t \rightarrow \infty} \frac{n_t t_0}{t} (\alpha + \epsilon) \\ &= (\alpha + \epsilon) 1 + 0 \\ &= \alpha + \epsilon. \end{aligned}$$

Since $\epsilon > 0$ is arbitrary, we have that for each $\epsilon > 0$,

$$\alpha \leq \limsup_{t \rightarrow \infty} \frac{f(t)}{t} \leq \alpha + \epsilon.$$

Hence

$$\begin{aligned}\limsup_{t \rightarrow \infty} \frac{f(t)}{t} &= \alpha \\ &= \inf_{t > 0} \frac{f(t)}{t} \\ &\leq \liminf_{t \rightarrow \infty} \frac{f(t)}{t}.\end{aligned}$$

Thus

$$\lim_{t \rightarrow \infty} \frac{f(t)}{t} = \inf_{t > 0} \frac{f(t)}{t}.$$

- Suppose that $\alpha = -\infty$ **FINISH!!!!** I think if $f : [0, \infty) \rightarrow \mathbb{R}$, then this cant happen since $f(0) \geq 0$, well maybe f needs to be continuous at 0 for this to be the case, idk

□

15.2 Convex Sets

15.2.1 Introduction

Definition 15.2.1.1. Let X be a vector space.

- Let $A \subset X$. Then A is said to be **convex** if for each $x, y \in A$, and $t \in [0, 1]$, $tx + (1 - t)y \in A$.
- We define

$$\text{Conv}(X) := \{A \subset X : A \text{ is convex}\}.$$

Exercise 15.2.1.2. Let X be a vector space. Then $\emptyset \in \text{Conv}(X)$.

Proof. Clear.

□

Exercise 15.2.1.3. Let X, Y be vector spaces, $f : X \rightarrow Y$ and $A \in \text{Conv}(X)$. If f is linear, then $f(A) \in \text{Conv}(Y)$.

Proof. Suppose that f is linear. Let $y_1, y_2 \in f(A)$ and $t \in [0, 1]$. Then there exist $x_1, x_2 \in A$ such that $y_1 = f(x_1)$ and $y_2 = f(x_2)$. Since A is convex, $tx_1 + (1 - t)x_2 \in A$. Then

$$\begin{aligned}ty_1 + (1 - t)y_2 &= tf(x_1) + (1 - t)f(x_2) \\ &= f(tx_1 + (1 - t)x_2) \\ &\in f(A).\end{aligned}$$

Since $y_1, y_2 \in f(A)$ and $t \in [0, 1]$ are arbitrary, we have that for each $y_1, y_2 \in f(A)$ and $t \in [0, 1]$, $ty_1 + (1 - t)y_2 \in f(A)$. Hence $f(A)$ is convex. □

Exercise 15.2.1.4. Let X be a vector space. Then for each $\mathcal{A} \subset \text{Conv}(X)$, $\bigcap_{A \in \mathcal{A}} A \in \text{Conv}(X)$.

Proof. Let $\mathcal{A} \subset \text{Conv}(X)$, $x, y \in \bigcap_{A \in \mathcal{A}} A$ and $t \in [0, 1]$. Then for each $A \in \mathcal{A}$, $x, y \in A$. Let $A \in \mathcal{A}$. Since A is convex, $tx + (1 - t)y \in A$. Since $A \in \mathcal{A}$ is arbitrary, we have that for each $A \in \mathcal{A}$, $tx + (1 - t)y \in A$. Hence $tx + (1 - t)y \in \bigcap_{A \in \mathcal{A}} A$. Since $x, y \in \bigcap_{A \in \mathcal{A}} A$ and $t \in [0, 1]$ are arbitrary, we have that for each $x, y \in \bigcap_{A \in \mathcal{A}} A$ and $t \in [0, 1]$, $tx + (1 - t)y \in \bigcap_{A \in \mathcal{A}} A$. Hence $\bigcap_{A \in \mathcal{A}} A \in \text{Conv}(X)$. □

Definition 15.2.1.5. Let X be a vector space and $A \subset X$. Define $\mathcal{S} \subset \text{Conv}(X)$ by

$$\mathcal{S} := \{S \subset X : S \in \text{Conv}(X) \text{ and } A \subset S\}.$$

We define the **convex hull** of A , denoted $\text{conv } A \in \text{Conv}(X)$, by

$$\text{conv } A = \bigcap_{S \in \mathcal{S}} S.$$

Note 15.2.1.6. We may think of $\text{conv } A$ as the smallest convex set containing A .

Definition 15.2.1.7. For each $n \in \mathbb{N}_0$, We define the **standard n -simplex**, denoted $\Delta^n \subset \mathbb{R}^{n+1}$, by

$$\Delta^n := \left\{ (\lambda_j)_{j \in [n+1]} : \sum_{j \in [n+1]} \lambda_j = 1 \right\}.$$

Exercise 15.2.1.8. Let $n \in \mathbb{N}_0$. Then Δ^n is convex.

Proof. Let $(\lambda_j)_{j \in [n+1]}, (\mu_j)_{j \in [n+1]} \in \Delta^n$ and $t \in [0, 1]$. We note that

$$\begin{aligned} \sum_{j \in [n+1]} [t\lambda_j + (1-t)\mu_j] &= t \sum_{j \in [n+1]} \lambda_j + (1-t) \sum_{j \in [n+1]} \mu_j \\ &= t + (1-t) \\ &= 1. \end{aligned}$$

Therefore

$$\begin{aligned} t((\lambda_j)_{j \in [n+1]}) + (1-t)(\lambda_j)_{j \in [n+1]} &= (t\lambda_j + (1-t)\mu_j)_{j \in [n+1]} \\ &\in \Delta^n. \end{aligned}$$

Since $(\lambda_j)_{j \in [n+1]}, (\mu_j)_{j \in [n+1]} \in \Delta^n$ and $t \in [0, 1]$ are arbitrary, we have that for each $(\lambda_j)_{j \in [n+1]}, (\mu_j)_{j \in [n+1]} \in \Delta^n$ and $t \in [0, 1]$, $t((\lambda_j)_{j \in [n+1]}) + (1-t)(\lambda_j)_{j \in [n+1]} \in \Delta^n$. $\square$

Definition 15.2.1.9. Let X be a vector space, $A \subset X$.

- Let $x \in X$. Then x is said to be a **convex combinations of elements of A** if there exist $n \in \mathbb{N}$ and $(a_j)_{j \in [n]} \subset A$ and $(\lambda_j)_{j \in [n]} \in \Delta^{n-1}$ such that $x = \sum_{j \in [n]} \lambda_j a_j$ and $\sum_{j \in [n]} \lambda_j = 1$.
- We define the **convex span of A** , denoted $\text{convspan } A \subset X$, by

$$\text{convspan } A := \{x \in X : x \text{ is a convex combination of elements of } A\}$$

Exercise 15.2.1.10. Let X be a vector space and $A \subset X$. Then

1. $A \subset \text{convspan } A$
2. $\text{convspan } A$ is convex

Proof.

1. Let $x \in A$, then

$$\begin{aligned} x &= 1x \\ &\in \text{convspan } A \end{aligned}$$

So $A \subset \text{convspan } A$.

2. Let $x, y \in \text{convspan } A$ and $t \in [0, 1]$. Then there exist $(a_i)_{i \in [n]}, (b_j)_{j \in [m]} \subset A$ and $(\lambda_i)_{i \in [n]} \in \Delta^{n-1}, (\mu_j)_{j \in [m]} \in \Delta^{m-1}$ such that $x = \sum_{i \in [n]} \lambda_i a_i$ and $y = \sum_{j \in [m]} \mu_j b_j$. Then

$$\begin{aligned} tx + (1-t)y &= t \left[\sum_{i \in [n]} \lambda_i a_i \right] + (1-t) \left[\sum_{j \in [m]} \mu_j b_j \right] \\ &= \sum_{i \in [n]} t\lambda_i a_i + \sum_{j \in [m]} (1-t)\mu_j b_j \end{aligned}$$

Since

$$\begin{aligned} \sum_{i \in [n]} t\lambda_i + \sum_{j \in [m]} (1-t)\mu_j &= t \sum_{i \in [n]} \lambda_i + (1-t) \sum_{j \in [m]} \mu_j \\ &= t + (1-t) \\ &= 1, \end{aligned}$$

we have that $tx + (1-t)y \in \text{convspan } A$. Since $x, y \in \text{convspan } A$ and $t \in [0, 1]$ are arbitrary, we have that for each $x, y \in \text{convspan } A$ and $t \in [0, 1]$, $tx + (1-t)y \in \text{convspan } A$. Hence $\text{convspan } A$ is convex. □

Exercise 15.2.1.11. Let X be a vector space and $A \subset X$. If A is convex, then

1. for each $n \in \mathbb{N}$, $(a_j)_{j \in [n]} \subset A$ and $(\lambda_j)_{j \in [n]} \in \Delta^{n-1}$, $\sum_{j \in [n]} \lambda_j a_j \in A$.

Hint: proceed by induction on n

2. $\text{conv } A = \text{convspan } A$

Proof. Suppose that A is convex.

1. Let $n \in \mathbb{N}$, $(a_j)_{j \in [n]} \subset A$ and $(\lambda_j)_{j \in [n]} \in \Delta^{n-1}$.
 - If $n = 2$, then by definition, $\sum_{j \in [n]} \lambda_j a_j \in A$.
 - Suppose that the claim is true for $n-1$. Since $\sum_{j \in [n]} \lambda_j = 1$, then there $k \in \{1, \dots, n\}$ such that $\lambda_k > 0$. Choose $l \in \{1, \dots, n\}$ such that $l \neq k$. Set $S = \{1, \dots, n\} \setminus \{l\}$. Then $1 - \lambda_l > 0$ and

$$\begin{aligned} x &= \sum_{j \in [n]} \lambda_j a_j \\ &= \lambda_l a_l + \sum_{j \in S} \lambda_j a_j \\ &= \lambda_l a_l + (1 - \lambda_l) \sum_{j \in S} \frac{\lambda_j}{1 - \lambda_l} a_j \end{aligned}$$

Since

$$\begin{aligned} \sum_{j \in S} \frac{\lambda_j}{1 - \lambda_l} &= \frac{1 - \lambda_l}{1 - \lambda_l} \\ &= 1 \end{aligned}$$

our induction hypothesis implies that

$$\sum_{j \in S} \frac{\lambda_j}{1 - \lambda_l} a_j \in A$$

Since A is convex, by definition we have that

$$\begin{aligned} x &= \lambda_l a_l + (1 - \lambda_l) \left[\sum_{j \in S} \frac{\lambda_j}{1 - \lambda_l} a_j \right] \\ &\in A \end{aligned}$$

By induction, for each $n \in \mathbb{N}$, $(a_j)_{j \in [n]} \subset A$ and $(\lambda_j)_{j \in [n]} \in \Delta^{n-1}$, $\sum_{j \in [n]} \lambda_j a_j \in A$.

2. • Since $A \subset \text{convspan } A$ and $\text{convspan } A$ is convex, Definition .2.0.11 (ref definition of convex hull) implies $\text{conv } A \subset \text{convspan } A$.

- The previous part implies that $\text{convspan } A \subset A$.

Since $\text{conv } A \subset \text{convspan } A$ and $\text{convspan } A \subset A$, we have that $\text{conv } A = \text{convspan } A$.

□

Exercise 15.2.1.12. Let X be a vector space and $(A_j)_{j \in [n]} \subset X$. Suppose that for each $j \in [n]$, A_j is convex. Then

$$\text{conv} \left(\bigcup_{j \in [n]} A_j \right) = \left\{ \sum_{j \in [n]} \lambda_j a_j : (\lambda_j)_{j \in [n]} \in \Delta^{n-1} \text{ and } (a_j)_{j \in [n]} \in \prod_{j \in [n]} A_j \right\}.$$

Proof. Define $A, B \subset X$ by $A := \bigcup_{j \in [n]} A_j$ and $B := \left\{ \sum_{j \in [n]} \lambda_j a_j : (\lambda_j)_{j \in [n]} \in \Delta^{n-1} \text{ and } (a_j)_{j \in [n]} \in \prod_{j \in [n]} A_j \right\}$.

- Clearly $B \subset \text{conv } A$.
- Let $x \in \text{conv } A$. **a prev ex about conv A = convspan A** implies that there exist $(\lambda_j)_{j \in [n]} \in \Delta^{n-1}$ and $(x_j)_{j \in [n]} \subset A$ such that $x = \sum_{j \in [n]} \lambda_j x_j$.
 - For each $j \in [n]$, define $K_j \subset [n]$ by $K_j := \{k \in [n] : x_j \in A_k\}$. Let $j \in [n]$. Since $x_j \in A$ and $A = \bigcup_{j \in [n]} A_j$, there exists $k \in [n]$ such that $x_j \in A_k$. Thus $K_j \neq \emptyset$. Since $j \in [n]$ is arbitrary, we have that for each $j \in [n]$, $K_j \neq \emptyset$. Define $F_K : [n] \rightarrow \mathcal{P}([n])$ by $F_K(j) := K_j$. **(make and ref ex about finite choice)** then implies that there exists $K : [n] \rightarrow [n]$ such that K is a selection of F_K .
 - By construction, for each $k \in [n]$, $K^{-1}(k) \neq \emptyset$. Define $F_J : [n] \rightarrow \mathcal{P}([n])$ by $F_J(k) := K^{-1}(k)$. finite choice implies that there exists $J : [n] \rightarrow [n]$ such that J is a selection of F_J .

Define $(\mu_k)_{k \in [n]} [0, 1]^n$ and $(y_k)_{k \in [n]} \subset X$ by $\mu_k := \sum_{j \in K^{-1}(k)} \lambda_j$ and

$$y_k := \begin{cases} x_{J(k)}, & \mu_k = 0 \\ \mu_k^{-1} \sum_{j \in K^{-1}(k)} \lambda_j x_j, & \mu_k \neq 0. \end{cases}$$

We note that

$$\begin{aligned} \sum_{k \in [n]} \mu_k &= \sum_{k \in [n]} \sum_{j \in K^{-1}(k)} \lambda_j \\ &= \sum_{j \in [n]} \lambda_j \\ &= 1 \end{aligned}$$

Thus $(\mu_k)_{k \in [n]} \in \Delta^{n-1}$. Let $k \in [n]$.

- If $\mu_k = 0$, then

$$\begin{aligned} y_k &= x_{J(k)} \\ &\in A_k \end{aligned}$$

- Assume that $\mu_k \neq 0$. Since

$$\begin{aligned} \sum_{j \in K^{-1}(k)} \frac{\lambda_j}{\mu_k} &= \frac{1}{\mu_k} \sum_{j \in K^{-1}(k)} \lambda_j \\ &= \frac{\mu_k}{\mu_k} \\ &= 1, \end{aligned}$$

we have that $(\lambda_j/\mu_k)_{j \in K^{-1}(k)} \in \Delta^{\#K^{-1}(k)-1}$. Since $(x_j)_{j \in K^{-1}(k)} \subset A_k$ and A_k is convex, we have that

$$\begin{aligned} y_k &= \sum_{j \in K^{-1}(k)} \frac{\lambda_j}{\mu_k} x_j \\ &\in A_k. \end{aligned}$$

Since $k \in [n]$ is arbitrary, we have that for each $k \in [n]$, $y_k \in A_k$. Thus $(y_k)_{k \in [n]} \in \prod_{k \in [n]} A_k$. Define $L \subset [n]$ by $L := \{k \in [n] : \mu_k \neq 0\}$. We note that if $k \in L^c$, then for each $j \in K^{-1}(k)$,

$$\begin{aligned} 0 &\leq \lambda_j \\ &\leq \sum_{j' \in K^{-1}(k)} \lambda_{j'} \\ &= \mu_k \\ &= 0 \end{aligned}$$

and $\lambda_j = 0$. Therefore

$$\begin{aligned} x &= \sum_{j \in [n]} \lambda_j x_j \\ &= \sum_{k \in [n]} \left[\sum_{j \in K^{-1}(k)} \lambda_j x_j \right] \\ &= \sum_{k \in L} \left[\sum_{j \in K^{-1}(k)} \lambda_j x_j \right] + \sum_{k \in L^c} \left[\sum_{j \in K^{-1}(k)} \lambda_j x_j \right] \\ &= \sum_{k \in L} \left[\mu_k \sum_{j \in K^{-1}(k)} \frac{\lambda_j}{\mu_k} x_j \right] + \sum_{k \in L^c} \left[\sum_{j \in K^{-1}(k)} 0 x_j \right] \\ &= \sum_{k \in L} \mu_k y_k + \sum_{k \in L^c} \mu_k y_k \\ &= \sum_{k \in [n]} \mu_k y_k \\ &\in B. \end{aligned}$$

Since $x \in \text{conv } A$ is arbitrary, we have that for each $x \in \text{conv } A$, $x \in B$. Thus $\text{conv } A \subset B$.

Since $B \subset \text{conv } A$ and $\text{conv } A \subset B$, we have that $\text{conv } A = B$. □

Exercise 15.2.1.13. Let X be a vector space and $A, B \subset X$ convex and $\lambda \in \mathbb{K}$. Then

1. $A + B$ is convex
2. λA is convex

Proof.

1. Let $x, y \in A + B$ and $t \in [0, 1]$. Then there exist $a_x, a_y \in A$, $b_x, b_y \in B$ such that $x = a_x + b_x$ and $y = a_y + b_y$. Since A and B are convex, $ta_x + (1-t)a_y \in A$ and $tb_x + (1-t)b_y \in B$. Hence

$$\begin{aligned} tx + (1-t)y &= ta_x + tb_x + (1-t)a_y + (1-t)b_y \\ &= [ta_x + (1-t)a_y] + [tb_x + (1-t)b_y] \\ &\in A + B \end{aligned}$$

So $A + B$ is convex.

2. Let $x, y \in \lambda A$ and $t \in [0, 1]$. Then there exist $a_x, a_y \in A$ such that $x = \lambda a_x$ and $y = \lambda a_y$. Since A is convex, $ta_x + (1-t)a_y \in A$. Therefore

$$\begin{aligned} tx + (1-t)y &= t\lambda a_x + (1-t)\lambda a_y \\ &= \lambda[ta_x + (1-t)a_y] \\ &\in \lambda A \end{aligned}$$

So λA is convex. □

Definition 15.2.1.14. Let X be a vector space and $A \in \text{Conv}(X)$.

- Let $x \in A$. Then x is said to be an **extreme point** of A if for each $y, z \in A$, $y \neq z$ implies that $x \notin [y, z]$.
- We define $\text{extrm } A := \{x \in A : x \text{ is an extreme point of } A\}$.

Exercise 15.2.1.15. [extreme point equiv wiki](#)

Proof. **FINISH!!!** □

Exercise 15.2.1.16. Let X be a vector space and $A \in \text{Conv}(X)$. Then $\text{conv}(\text{extrm } A) = \text{conv } A$.

Proof. **FINISH!!!** □

15.2.2 Cones

Definition 15.2.2.1. Let X be a vector space and $C \subset X$. Then C is said to be a

- **cone** of X if $\mathbb{R}_+ C \subset C$.
- We define $\text{Cone}(X) \subset \mathcal{P}(X)$ by $\text{Cone}(X) := \{C \subset X : C \text{ is a cone of } X\}$.

Definition 15.2.2.2. Let X be a vector space. We define $\text{ConvCone}(X) \subset \mathcal{P}(X)$ by $\text{ConvCone}(X) := \text{Conv}(X) \cap \text{Cone}(X)$.

Exercise 15.2.2.3. Let X be a vector space and $C \in \text{Cone}(X)$. Then $C \in \text{ConvCone}(X)$ iff $C + C \subset C$.

Proof.

- ($\implies$):
Suppose that $C \in \text{ConvCone}(X)$. Let $x \in C + C$. Then there exists $x_1, x_2 \in C$ such that $x = x_1 + x_2$. Since $C \in \text{Cone}(X)$, $2x_1, 2x_2 \in C$. Define $y_1, y_2 \in C$ by $y_1 := 2x_1$ and $y_2 := 2x_2$. Since $C \in \text{Conv}(X)$, we have that

$$\begin{aligned} x &= x_1 + x_2 \\ &= 2^{-1}y_1 + 2^{-1}y_2 \\ &\in C. \end{aligned}$$

Since $x \in C + C$ is arbitrary, we have that for each $x \in C + C$, $x \in C$. Hence $C + C \subset C$.

- ($\impliedby$):
Suppose that $C + C \subset C$. Let $x, y \in C$ and $t \in [0, 1]$. Since $C \in \text{Cone}(X)$, $tx \in C$ and $(1-t)y \in C$. By assumption,

$$\begin{aligned} tx + (1-t)y &\in C + C \\ &\subset C. \end{aligned}$$

Since $x, y \in C$ and $t \in [0, 1]$ are arbitrary, we have that for each $x, y \in C$ and $t \in [0, 1]$, $tx + (1-t)y \in C$. Hence $C \in \text{Conv}(X)$. Thus $C \in \text{ConvCone}(X)$. □

discus cone generated by subset, faces of cones, extreme points of cones, etc

Definition 15.2.2.4. Let X be a vector space.

- Let $C \in \text{Cone}(X)$. Then C is said to be **pointed** if $C \cap -C = \{0\}$,
- We define $\text{PtConvCone}(X) \subset \mathcal{P}(X)$ by $\text{PtConvCone}(X) := \{C \in \text{ConvCone}(X) : C \text{ is pointed}\}$.

15.3 Convex Functions

Note 15.3.0.1. In the remainder of this chapter, we assume all vector spaces are real.

Definition 15.3.0.2. Let X be a vector space and $f : A \rightarrow \mathbb{R}$. Then f is said to be **convex** if for each $x, y \in A$, $t \in [0, 1]$,

$$f(tx + (1 - t)y) \leq tf(x) + (1 - t)f(y)$$

Definition 15.3.0.3. Let X be a vector space, $A \in \text{Conv}(X)$ and $f : A \rightarrow \mathbb{R}$. Then f is said to be **strictly convex** if for each $x, y \in A$, $t \in (0, 1)$, $x \neq y$ implies that

$$f(tx + (1 - t)y) < tf(x) + (1 - t)f(y)$$

Exercise 15.3.0.4. Let X be a vector space, $f \in X^*$ and $g : X \rightarrow \mathbb{R}$ constant. Then f and g are convex.

Proof. Let $x, y \in X$ and $t \in [0, 1]$. Put $c = g(0)$. Then

$$f(tx + (1 - t)y) = tf(x) + (1 - t)f(y)$$

and

$$\begin{aligned} g(tx + (1 - t)y) &= c \\ &= tc + (1 - t)c \\ &= tg(x) + (1 - t)g(y) \end{aligned}$$

So f and g are convex. □

Exercise 15.3.0.5. Star-shapedness: Let $f : [0, \infty) \rightarrow \mathbb{R}$ be convex. If $f(0) \leq 0$, then for each $x \in [0, \infty)$, $t \in [0, 1]$, $f(tx) \leq tf(x)$.

Proof. Suppose that $f(0) \leq 0$. Let $x \in [0, \infty)$ and $t \in [0, 1]$. Then

$$\begin{aligned} f(tx) &= f(tx + (1 - t)0) \\ &\leq tf(x) + (1 - t)f(0) \\ &\leq tf(x) \end{aligned}$$

□

Exercise 15.3.0.6. Superadditivity:

Let $f : [0, \infty) \rightarrow \mathbb{R}$ be convex. If $f(0) = 0$, then for each $x, y \in [0, \infty)$,

$$f(x) + f(y) \leq f(x + y)$$

Hint: $f(x) = f\left(\frac{x}{x+y}(x+y)\right)$

Proof. Suppose that $f(0) = 0$. Let $x, y \in [0, \infty)$. If $x + y = 0$, then $x = y = 0$ and $f(x) + f(y) = 0 = f(x + y)$. Suppose that $x + y \neq 0$. Then the previous exercise implies that

$$\begin{aligned} f(x) + f(y) &= f\left(\frac{x}{x+y}(x+y)\right) + f\left(\frac{y}{x+y}(x+y)\right) \\ &\leq \frac{x}{x+y}f(x+y) + \frac{y}{x+y}f(x+y) \\ &= f(x+y) \end{aligned}$$

□

Exercise 15.3.0.7. Let X be a vector space, $A \in \text{Conv}(X)$, $f, g : A \rightarrow \mathbb{R}$ and $\lambda \geq 0$. If f, g are convex, then

1. $f + g$ is convex

2. λf is convex

Proof. Suppose that f and g are convex. Let $x, y \in A$ and $t \in [0, 1]$. Then

$$\begin{aligned} (f + \lambda g)(tx + (1-t)y) &= f(tx + (1-t)y) + \lambda g(tx + (1-t)y) \\ &\leq tf(x) + (1-t)f(y) + t\lambda g(x) + (1-t)\lambda g(y) \\ &= t(f(x) + \lambda g(x)) + (1-t)(f(y) + \lambda g(y)) \\ &= t(f + \lambda g)(x) + (1-t)(f + \lambda g)(y) \end{aligned}$$

□

Definition 15.3.0.8. Let X be a vector space and $A \in \text{Conv}(X)$.

- Let $f : A \rightarrow \mathbb{R}$. Then f is said to be **affine** if there exists $\phi \in X^*$, $a \in \mathbb{R}$ constant such that $f = \phi|_A + a$.

- We define $\text{Aff}(A) \subset \mathbb{R}^A$ by $\text{Aff}(A) := \{f \in \mathbb{R}^A : f \text{ is affine}\}$.

Exercise 15.3.0.9. Let X be a vector space, $A \in \text{Conv}(X)$ and $f \in \text{Aff}(A)$. Then f is convex.

Proof. Since $f \in \text{Aff}(A)$, there exists $\phi \in X^*$, $a \in \mathbb{R}$ constant such that $f = \phi|_A + a$. Define $g : A \rightarrow \mathbb{R}$ by $g(x) = a$. **Some previous exercises** imply that $\phi|_A$, g are convex and $f = \phi|_A + g$ is convex. □

Exercise 15.3.0.10. Let X be a vector space, $A \in \text{Conv}(X)$, $f : \mathbb{R} \rightarrow \mathbb{R}$ and $g : A \rightarrow \mathbb{R}$. If f is convex and increasing and g is convex, then $f \circ g$ is convex.

Proof. Let $t \in [0, 1]$ and $x, y \in A$. Then convexity of g implies that

$$g(tx + (1-t)y) \leq tg(x) + (1-t)g(y)$$

and we have

$$\begin{aligned} f \circ g(tx + (1-t)y) &= f(g(tx + (1-t)y)) \\ &\leq f(tg(x) + (1-t)g(y)) && (f \text{ increasing}) \\ &\leq tf(g(x)) + (1-t)f(g(y)) && (f \text{ convex}) \\ &= tf \circ g(x) + (1-t)f \circ g(y) \end{aligned}$$

So $f \circ g$ is convex. □

Exercise 15.3.0.11. Let X be a vector space, $A \in \text{Conv}(X)$, $f : A \rightarrow \mathbb{R}$ convex and $x_0 \in A$. Then f has a local minimum point at x_0 iff f has a global minimum point at x_0 .

Proof. If f has a global minimum point at x_0 , then f has a local minimum point at x_0 . Conversely, suppose that f has a local minimum point at x_0 . Then there exists $\delta > 0$ such that for each $x \in B(x_0, \delta) \cap A$, $f(x_0) \leq f(x)$. For the sake of contradiction, suppose that f does not have a global minimum point at x_0 . Then there exists $x' \in A$ such that $f(x') < f(x_0)$. Put $t_0 = \min(\frac{\delta}{\|x' - x_0\| + 1}, 1) > 0$. Let $t \in (0, t_0)$, then

$$\begin{aligned} \|(tx' + (1-t)x_0) - x_0\| &= t\|x' - x_0\| \\ &< \frac{\|x' - x_0\|\delta}{\|x' - x_0\| + 1} \\ &< \delta \end{aligned}$$

so that $tx' + (1-t)x_0 \in B(x_0, \delta) \cap A$ and hence $f(x_0) \leq f(tx' + (1-t)x_0)$. Therefore

$$\begin{aligned} f(x_0) &\leq f(tx' + (1-t)x_0) \\ &\leq tf(x') + (1-t)f(x_0) && (\text{convexity of } f) \\ &< tf(x_0) + (1-t)f(x_0) \\ &= f(x_0) \end{aligned}$$

which is a contradiction. Hence f has a global minimum point at x_0 . □

Exercise 15.3.0.12. Let X be a vector space, $A \in \text{Conv}(X)$, $f : A \rightarrow \mathbb{R}$ strictly convex and $x_0 \in X$. If f has a local minimum point at x_0 , then f has a unique global minimum point at x_0 .

Proof. Suppose that f has a local minimum point at x_0 . The previous exercise implies that f has a global minimum point at x_0 . For the sake of contradiction suppose that there exists $x_1 \in X$ such that f has a global minimum point at x_1 and $x_0 \neq x_1$. This implies $f(x_0) = f(x_1)$. Set $t = 1/2$. Strict convexity implies that

$$\begin{aligned} f(tx_0 + (1-t)x_1) &< tf(x_0) + (1-t)f(x_1) \\ &= f(x_0) \end{aligned}$$

which is a contradiction since f has a global minimum point at x_0 . □

Definition 15.3.0.13. (rewrite in terms of slice maps) Let X, Y be vector spaces, $A \subset X \oplus Y$. For $y \in Y$, define

$$A^y = \{x \in X : (x, y) \in A\}$$

and $f^y : A^y \rightarrow \mathbb{R}$ by

$$f^y(x) = f(x, y)$$

Exercise 15.3.0.14. (rewrite in terms of slice maps) Let X, Y be vector spaces, $A \subset X \oplus Y$ convex and $f : A \rightarrow \mathbb{R}$ convex. Then for each $y \in \pi_2(A)$,

1. A^y is convex
2. f^y is convex

where $\pi_2 : X \times Y \rightarrow Y$, the canonical projection of $X \times Y$ onto Y given by $\pi_2(x, y) = y$.

Proof. Let $y \in \pi_2(A)$, $x_1, x_2 \in A^y$ and $t \in [0, 1]$. Then by definition, $(x_1, y), (x_2, y) \in A$.

1. Convexity of A implies that $(tx_1 + (1-t)x_2, y) \in A$. Hence $tx_1 + (1-t)x_2 \in A^y$ and A^y is convex.
2. Convexity of f implies that

$$\begin{aligned} f^y(tx_1 + (1-t)x_2) &= f(tx_1 + (1-t)x_2, y) \\ &= f(t(x_1, y) + (1-t)(x_2, y)) \\ &\leq tf(x_1, y) + (1-t)f(x_2, y) \\ &= tf^y(x_1) + (1-t)f^y(x_2) \end{aligned}$$

and so f^y is convex. □

Exercise 15.3.0.15. Let X, Y be vector spaces and $A \subset X, B \subset Y$. If A and B are convex, then $A \times B \subset X \oplus Y$ is convex.

Proof. Suppose that A and B are convex. Let $(x_1, y_1), (x_2, y_2) \in A \times B$ and $t \in [0, 1]$. Convexity of A and B implies that $tx_1 + (1-t)x_2 \in A$ and $ty_1 + (1-t)y_2 \in B$. Therefore

$$\begin{aligned} t(x_1, y_1) + (1-t)(x_2, y_2) &= (tx_1 + (1-t)x_2, ty_1 + (1-t)y_2) \\ &\in A \times B \end{aligned}$$

□

Exercise 15.3.0.16. (rewrite in terms of slice maps) Let X, Y be vector spaces and $A \subset X, B \subset Y$ convex (implying that $A \times B$ is convex) and $f : A \times B \rightarrow \mathbb{R}$ convex. Suppose that for each $y \in B$, $\{f(x, y) : x \in A\}$ is bounded below. Then $\inf_{y \in B} f^y$ is convex

Proof. Put $g = \inf_{y \in B} f^y$. Let $x_1, x_2 \in A$, $y_1, y_2 \in B$ and $t \in [0, 1]$. Put $y' = ty_1 + (1-t)y_2$. Then convexity of f implies that

$$\begin{aligned} g(tx_1 + (1-t)x_2) &\leq f^{y'}(tx_1 + (1-t)x_2) \\ &= f(tx_1 + (1-t)x_2, ty_1 + (1-t)y_2) \\ &= f(t(x_1, y_1) + (1-t)(x_2, y_2)) \\ &\leq tf(x_1, y_1) + (1-t)f(x_2, y_2) \\ &= tf^{y_1}(x_1) + (1-t)f^{y_2}(x_2) \end{aligned}$$

Since $y_1 \in B$ is arbitrary, we have that

$$g(tx_1 + (1-t)x_2) \leq tg(x_1) + (1-t)f^{y_2}(x_2)$$

Similarly, since $y_2 \in B$ is arbitrary, we have that

$$g(tx_1 + (1-t)x_2) \leq tg(x_1) + (1-t)g(x_2)$$

and f is convex. □

Exercise 15.3.0.17. Let X be a vector space, $A \in \text{Conv}(X)$ and $(f_\lambda)_{\lambda \in \Lambda} \subset \mathbb{R}^A$. Suppose that for each $\lambda \in \Lambda$, f_λ is convex. Define

- $A^* = \{x \in A : \sup_{\lambda \in \Lambda} f_\lambda(x) < \infty\}$
- $f^* : A^* \rightarrow \mathbb{R}$ by $f^*(x) = \sup_{\lambda \in \Lambda} f_\lambda(x)$

Then

1. A^* is convex
2. f^* is convex

Proof. 1. Let $x, y \in A$ and $t \in [0, 1]$. By definition, $\sup_{\lambda \in \Lambda} f_\lambda(x), \sup_{\lambda \in \Lambda} f_\lambda(y) < \infty$. Therefore

$$\begin{aligned} \sup_{\lambda \in \Lambda} f_\lambda(tx + (1-t)y) &\leq \sup_{\lambda \in \Lambda} [tf_\lambda(x) + (1-t)f_\lambda(y)] \\ &\leq t \sup_{\lambda \in \Lambda} f_\lambda(x) + (1-t) \sup_{\lambda \in \Lambda} f_\lambda(y) \\ &< \infty \end{aligned}$$

So $tx + (1-t)y \in A$.

2. By definition, the previous part implies that for each $x, y \in A^*$, $f^*(tx + (1-t)y) \leq tf^*(x) + (1-t)f^*(y)$. So $f^* : A^* \rightarrow \mathbb{R}$ is convex. □

Exercise 15.3.0.18. Let X be a normed vector space, $A \subset X$ open and convex, $f : A \rightarrow \mathbb{R}$ convex and $x_0 \in A$. If f is continuous at x_0 , then f is locally Lipschitz at x_0 .

Hint: Given x_1, x_2 near x_0 Choose a z near x_0 s.t. x_1 is a convex combination of x_2 and z . Then repeat but with x_2 as a convex combination of x_1 and z

Proof. By continuity, f is locally bounded at x_0 . So there exist $M, \delta > 0$ such that $B(x_0, \delta) \subset A$ and for each $x \in B(x_0, \delta)$, $|f(x)| \leq M$. Put $\delta' = \frac{\delta}{2}$ and choose $U = B(x_0, \delta')$. Then $U \subset A$ and $U \in \mathcal{N}(x_0)$.

Let $x_1, x_2 \in U$. Suppose that $x_1 \neq x_2$. Define $\alpha = \|x_1 - x_2\| > 0$, $p = \frac{\alpha}{\alpha + \delta'}$, $q = 1 - p$ and $z = p^{-1}(x_1 - qx_2)$. Then $x_1 = pz + qx_2$ and

$$\begin{aligned} \|z - x_1\| &= \|(p^{-1} - 1)x_1 - p^{-1}qx_2\| \\ &= \frac{1-p}{p}\alpha \\ &= \frac{\delta'}{\alpha}\alpha \\ &= \delta' \end{aligned}$$

Therefore

$$\begin{aligned}\|z - x_0\| &\leq \|z - x_1\| + \|x_1 - x_0\| \\ &< \delta' + \delta' \\ &= \delta\end{aligned}$$

So $z \in B(x_0, \delta)$, which implies that

$$\begin{aligned}f(z) - f(x_2) &\leq |f(z) - f(x_2)| \\ &\leq |f(z)| + |f(x_2)| \\ &\leq 2M\end{aligned}$$

Since $x_1 = pz + qx_2$, convexity of f implies that $f(x_1) \leq pf(z) + qf(x_2)$. Hence

$$\begin{aligned}f(x_1) - f(x_2) &\leq pf(z) - pf(x_2) \\ &= p(f(z) - f(x_2)) \\ &\leq p2M \\ &= \frac{\alpha}{\alpha + \delta'} 2M \\ &\leq \alpha 2M \\ &= 2M\|x_1 - x_2\|\end{aligned}$$

Similarly, choosing $z = p^{-1}(x_2 - qx_1)$, yields $f(x_2) - f(x_1) \leq 2M\|x_1 - x_2\|$ which implies that

$$|f(x_1) - f(x_2)| \leq 2M\|x_1 - x_2\|$$

and f is Lipschitz on U . □

15.4 The Subdifferential

Exercise 15.4.0.1. Let X be a Banach space, $A \subset X$ open and convex, $f : A \rightarrow \mathbb{R}$ convex, $x_0 \in A$ and $x \in X$. Define $T = \{t \in \mathbb{R} : x_0 + tx \in A\}$. Then there exist $a, b \in (0, \infty]$ such that $T = (-a, b)$.

Proof. Continuity of scalar multiplication and addition implies that T is an open neighborhood of 0. Let $t > 0$ and $s \in [0, t]$. Then $\frac{s}{t} \in [0, 1]$ and by convexity of A , $x_0 + tx \in A$ implies that

$$\begin{aligned} x_0 + sx &= \frac{s}{t}(x_0 + tx) + \left(1 - \frac{s}{t}\right)x_0 \\ &\in A \end{aligned}$$

Thus $[0, t] \subset T$. Similarly, $x_0 - tx \in A$ implies that $[-t, 0] \subset T$.

Define $a, b \in (0, \infty]$ by $a = \sup\{t > 0 : x_0 - tx \in A\}$ and $b = \sup\{t > 0 : x_0 + tx \in A\}$. Then $(-a, b) = T$. $\square$

Definition 15.4.0.2. Let X be a Banach space, $A \subset X$ open and convex, $f : A \rightarrow \mathbb{R}$ convex, $x_0 \in A$ and $x \in X$. Define T as in the previous exercise and choose $t_0 > 0$ such that $(-t_0, t_0) \subset T$. For $t \in (0, t_0)$, define the difference quotient $q : (-t_0, t_0) \setminus \{0\} \rightarrow \mathbb{R}$ by

$$q(t) = \frac{f(x_0 + tx) - f(x_0)}{t}$$

Exercise 15.4.0.3. Let X be a Banach space, $A \subset X$ open and convex, $f : A \rightarrow \mathbb{R}$ convex, $x_0 \in A$ and $x \in X$. Define t_0 as above. Then

1. $q(t)$ is increasing on $(0, t_0)$
2. $q(-t)$ decreasing on $(0, t_0)$

Hint: As an example, look at the graph of $f(x) = x^2$. For the algebra, start at the desired end inequality and work backwards

Proof.

1. Let $s, t \in (0, t_0)$ and suppose that $s \leq t$. Then $x_0 + sx, x_0 + tx \in A$. Note that since $0 < s \leq t$, $\frac{s}{t} \in (0, 1]$ and $1 - \frac{s}{t} = \frac{t-s}{t} \in (0, 1]$. Also, since A is convex, we have that

$$\left(\frac{t-s}{t}\right)x_0 + \left(\frac{s}{t}\right)(x_0 + tx) \in A$$

Convexity of f implies that

$$\begin{aligned} f(x_0 + sx) &= f\left(\left(\frac{t-s}{t}\right)x_0 + \left(\frac{s}{t}\right)(x_0 + tx)\right) \\ &\leq \left(\frac{t-s}{t}\right)f(x_0) + \left(\frac{s}{t}\right)f(x_0 + tx) \end{aligned}$$

This implies that

$$tf(x_0 + sx) \leq (t-s)f(x_0) + sf(x_0 + tx)$$

and after rearranging, we get

$$tf(x_0 + sx) - tf(x_0) \leq sf(x_0 + tx) - sf(x_0)$$

and so finally, dividing both sides by st , we obtain

$$\begin{aligned} q(s) &= \frac{f(x_0 + sx) - f(x_0)}{s} \\ &\leq \frac{f(x_0 + tx) - f(x_0)}{t} \\ &= q(t) \end{aligned}$$

as desired.

2. Similar to (1). □

Exercise 15.4.0.4. Let X be a Banach space, $A \subset X$ open and convex, $f : A \rightarrow \mathbb{R}$ convex, $x_0 \in A$ and $x \in X$. Define t_0 as before. Then for each $t \in (0, t_0)$,

$$q(-t) \leq q(t)$$

Hint: for sufficiently small t , convexity of f implies that $f(x_0) \leq \frac{1}{2}f(x_0 - 2tx) + \frac{1}{2}f(x_0 + 2tx)$

Proof. Choose t_0 as in the previous exercise. Since convexity of f implies that for each $t \in (0, t_0/2)$,

$$f(x_0) \leq \frac{1}{2}f(x_0 - 2tx) + \frac{1}{2}f(x_0 + 2tx)$$

we have that for each $t \in (0, t_0/2)$,

$$\begin{aligned} q(-2t) &= \frac{f(x_0 - 2tx) - f(x_0)}{-2t} \\ &\leq \frac{f(x_0 + 2tx) - f(x_0)}{2t} \\ &= q(2t) \end{aligned}$$

So for each $t \in (0, t_0)$, $q(-t) \leq q(t)$. □

Exercise 15.4.0.5. Let X be a Banach space, $A \subset X$ open and convex, $f : A \rightarrow \mathbb{R}$ convex and $x_0 \in A$. Then

1. f is left-hand and right-hand Gateaux differentiable at x_0 with $d^-f(x_0) \leq d^+f(x_0)$
2. for each $x \in X$, $d^-f(x_0)(x) = -d^+f(x_0)(-x)$

Proof.

1. Let $x \in X$. Choose $t_0 > 0$ as in the previous two exercises. Let $t, u \in (0, t_0)$. Choose $s \in (0, \min(u, t))$. The previous two exercises imply that

$$\begin{aligned} q(-u) &\leq q(-s) \\ &\leq q(s) \\ &\leq q(t) \end{aligned}$$

and therefore $q(t)$ is an upper bound for $\{q(-u) : u \in (0, t_0)\}$ and $d^-f(x_0)(x) = \sup_{u \in (0, t_0)} q(-u)$ exists with $d^-f(x_0)(x) \leq q(t)$.

Since $t \in (0, t_0)$ is arbitrary, $d^-f(x_0)(x)$ is a lower bound for $\{q(t) : t \in (0, t_0)\}$. Therefore

$$d^+f(x_0)(x) = \inf_{t \in (0, t_0)} q(t)$$

exists with $d^+f(x_0)(x) \geq d^-f(x_0)(x)$.

2. By definition, we have

$$\begin{aligned} d^-f(x_0)(x) &= \lim_{t \rightarrow 0^+} \frac{f(x_0 - tx) - f(x_0)}{-t} \\ &= - \lim_{t \rightarrow 0^+} \frac{f(x_0 + tx) - f(x_0)}{t} \\ &= -d^+f(x_0)(-x) \end{aligned}$$

□

Exercise 15.4.0.6. Let X be a Banach space, $A \subset X$ open and convex, $f : A \rightarrow \mathbb{R}$ convex and $x_0 \in A$. Then $d^+f(x_0) : X \rightarrow \mathbb{R}$ is a sublinear functional.

Proof. Let $x, y \in X$ and $k \geq 0$. If $k = 0$, then clearly

$$d^+f(x_0)(kx) = kd^+(x_0)(x)$$

If $k > 0$. Then

$$\begin{aligned} d^+f(x_0)(kx) &= \lim_{t \rightarrow 0^+} \frac{f(x_0 + tkx) - f(x_0)}{t} \\ &= k \lim_{t \rightarrow 0^+} \frac{f(x_0 + tkx) - f(x_0)}{tk} \\ &= kd^+f(x_0)(x) \end{aligned}$$

Define $t_0 > 0$ as before and let $t \in (0, \frac{t_0}{2})$. Note that

$$x_0 + tx + ty = \frac{1}{2}(x_0 + 2tx) + \frac{1}{2}(x_0 + 2ty)$$

Convexity of f implies that

$$f(x_0 + tx + ty) \leq \frac{1}{2}f(x_0 + 2tx) + \frac{1}{2}f(x_0 + 2ty)$$

which implies that

$$\frac{f(x_0 + tx + ty) - f(x_0)}{t} \leq \frac{f(x_0 + 2tx) - f(x_0)}{2t} + \frac{f(x_0 + 2ty) - f(x_0)}{2t}$$

Therefore

$$\begin{aligned} d^+f(x_0)(x + y) &= \lim_{t \rightarrow 0^+} \frac{f(x_0 + t(x + y)) - f(x_0)}{t} \\ &= \lim_{t \rightarrow 0^+} \frac{f(x_0 + tx + ty) - f(x_0)}{t} \\ &\leq \lim_{t \rightarrow 0^+} \left[\frac{f(x_0 + 2tx) - f(x_0)}{2t} + \frac{f(x_0 + 2ty) - f(x_0)}{2t} \right] \\ &= \lim_{t \rightarrow 0^+} \frac{f(x_0 + 2tx) - f(x_0)}{2t} + \lim_{t \rightarrow 0^+} \frac{f(x_0 + 2ty) - f(x_0)}{2t} \\ &= d^+f(x_0)(x) + d^+f(x_0)(y) \end{aligned}$$

□

Exercise 15.4.0.7. Let X be a Banach space, $A \subset X$ open and convex, $f : A \rightarrow \mathbb{R}$ convex and $x_0 \in A$. Then for each $x \in A$,

$$d^+f(x_0)(x - x_0) \leq f(x) - f(x_0)$$

Proof. Let $x \in A$. Define $T = \{t \in \mathbb{R} : x_0 + t(x - x_0) \in A\}$ similarly to earlier. Clearly $1 \in T$ and

$$\begin{aligned} d^+f(x_0)(x - x_0) &= \inf_{t \in (0, 1]} \frac{f(x_0 + t(x - x_0)) - f(x_0)}{t} \\ &\leq f(x) - f(x_0) \end{aligned}$$

□

Exercise 15.4.0.8. Let X be a Banach space, $A \subset X$ open and convex, $f : A \rightarrow \mathbb{R}$ convex and $x_0 \in A$. If f is continuous at x_0 , then $d^+f(x_0)$ is Lipschitz (equivalently bounded).

Proof. Suppose that f is continuous at x_0 . A previous exercise about convex functions tells us that f is locally Lipschitz at x_0 , so there exists $\delta, M > 0$ such that for each $x_1, x_2 \in B(x_0, \delta)$, $|f(x_1) - f(x_2)| \leq M\|x_1 - x_2\|$. Let $x \in X$ and define $t_0 = \frac{\delta}{\|x\|+1}$ so that for each $t \in (0, t_0)$,

$$\begin{aligned} \|(x_0 + tx) - x_0\| &= t\|x\| \\ &\leq t_0\|x\| \\ &= \frac{\delta\|x\|}{\|x\|+1} \\ &< \delta \end{aligned}$$

and $x_0 + tx \in B(x_0, \delta)$. Then for each $t \in (0, t_0)$,

$$\begin{aligned} d^+f(x_0)(x) &\leq \frac{f(x_0 + tx) - f(x_0)}{t} \\ &\leq \frac{|f(x_0 + tx) - f(x_0)|}{t} \\ &\leq t^{-1}M\|(x_0 + tx) - x_0\| \\ &= M\|x\| \end{aligned}$$

Thus $d^+f(x_0)$ is a bounded sublinear functional and a previous exercise in the section on sublinear functionals implies this is equivalent to $d^+f(x_0)$ being Lipschitz. $\square$

Exercise 15.4.0.9. Let X be a Banach space, $A \subset X$ open and convex, $f : A \rightarrow \mathbb{R}$ convex and $x_0 \in A$. If f is continuous at x_0 , then there exists $\phi \in X^*$ such that $\phi \leq d^+f(x_0)$.

Proof. Suppose that f is continuous at x_0 . The previous exercise implies that $d^+f(x_0)$ is Lipschitz (equivalently bounded). A previous exercise in the section discussing sublinear functionals tells us that boundedness of $d^+f(x_0)$ implies that there exists $\phi \in X^*$ such that $\phi \leq d^+f(x_0)$. $\square$

Definition 15.4.0.10. Subdifferential:

Let X be a Banach space, $A \subset X$ open and convex, $f : A \rightarrow \mathbb{R}$ convex and $x_0 \in A$. We define the **subdifferential of f at x_0** , denoted $\partial f(x_0)$, to be

$$\partial f(x_0) = \{\phi \in X^* : \text{for each } x \in A, f(x_0) + \phi(x - x_0) \leq f(x)\}$$

Exercise 15.4.0.11. Let X be a Banach space, $A \subset X$ open and convex, $f : A \rightarrow \mathbb{R}$ convex and $x_0 \in A$. If f is continuous at x_0 , then $\partial f(x_0) \neq \emptyset$.

Proof. Suppose that f is continuous at x_0 . The previous exercise tells us that there exists $\phi \in X^*$ such that $\phi \leq d^+f(x_0)$. Let $x \in A$. A previous exercise implies that

$$\begin{aligned} \phi(x - x_0) &\leq d^+f(x_0)(x - x_0) \\ &\leq f(x) - f(x_0) \end{aligned}$$

Then $f(x_0) + \phi(x - x_0) \leq f(x)$. $\square$

Exercise 15.4.0.12. Let X be a Banach space, $A \subset X$ open and convex, $f : A \rightarrow \mathbb{R}$ convex, $\phi \in X^*$ and $x_0 \in A$. Then

1. for each $x \in A$,

$$\phi(x - x_0) \leq f(x) - f(x_0)$$

iff

$$\phi \leq d^+f(x_0)$$

2. $\partial f(x_0) = \{\phi \in X^* : \phi \leq d^+f(x_0)\}$

Proof.

1. Suppose that for each $x \in A$, $\phi(x - x_0) \leq f(x) - f(x_0)$. Let $x \in X$. Define t_0 as before. Then for each $t \in (0, t_0)$,

$$\begin{aligned} t\phi(x) &= \phi((x_0 + tx) - x_0) \\ &\leq f(x_0 + tx) - f(x_0) \end{aligned}$$

This implies that $\phi(x) \leq d^+f(x_0)(x)$.

Conversely, suppose that $\phi \leq d^+f(x_0)$. Let $x \in A$. A previous exercise implies that,

$$\begin{aligned} \phi(x - x_0) &\leq d^+f(x_0)(x - x_0) \\ &\leq f(x) - f(x_0) \end{aligned}$$

2. Clear. □

Exercise 15.4.0.13. Let X be a Banach space, $A \subset X$ open and convex, $f : A \rightarrow \mathbb{R}$ convex and $x_0 \in A$. If f is continuous at x_0 , then the following are equivalent:

1. f is Gateaux differentiable at x_0
2. $d^+f(x_0)$ is linear
3. $\#\partial f(x_0) = 1$

Proof. Suppose that f is continuous at x_0 . Then $d^+f(x_0)$ is Lipschitz and bounded.

- (1) $\implies$ (2):

Suppose that f is Gateaux differentiable at x_0 . Let $x \in X$. Then a previous exercise implies that

$$\begin{aligned} -df^+(x_0)(-x) &= df^-f(x_0)(x) \\ &= df^+f(x_0)(x) \end{aligned}$$

An exercise in the section on sublinear functionals implies that $df^+f(x_0)$ is linear.

- (2) $\implies$ (3):

Suppose that $df^+f(x_0)$ is linear. Let $\phi \in \partial f(x_0)$. The previous exercise implies that $\phi \leq df^+f(x_0)$. Equivalence of linearity in the section on sublinear functionals implies that $d^+f(x_0) = \phi$.

- (3) $\implies$ (1):

Suppose that $\#\partial f(x_0) = 1$. Since $\partial f(x_0) = \{\phi \in X^* : \phi \leq d^+f(x_0)\}$, equivalence of linearity in the section on sublinear functionals implies that $d^+f(x_0)$ is linear. This implies that $d^+f(x_0) = d^-f(x_0)$ and which implies that f is Gateaux differentiable at x_0 . □

Exercise 15.4.0.14. Let X be a Banach space, $A \subset X$ open and convex, $f, g : A \rightarrow \mathbb{R}$ convex, $\lambda \geq 0$ and $x_0 \in A$. Then

$$\partial f(x_0) + \lambda \partial g(x_0) \subset \partial[f + \lambda g](x_0)$$

Proof. Let $\zeta \in \partial f(x_0) + \lambda \partial g(x_0)$. Then there exist $\phi \in \partial f(x_0)$ and $\psi \in \partial g(x_0)$ such that $\zeta = \phi + \lambda\psi$. A previous exercise implies that $\phi \leq d^+f(x_0)$ and $\lambda\psi \leq \lambda d^+g(x_0) = d^+[\lambda g](x_0)$. Hence

$$\begin{aligned} \zeta &= \phi + \lambda\psi \\ &\leq d^+f(x_0) + d^+[\lambda g](x_0) \\ &= d^+[f + \lambda g](x_0) \end{aligned}$$

So $\zeta \in \partial[f + \lambda g](x_0)$ □

Exercise 15.4.0.15. Let X be a Banach space, $A \subset X$ open and convex, $f : A \rightarrow \mathbb{R}$ convex and $x_0 \in A$. If f is continuous at x_0 , then f has a global minimum point at x_0 iff $0 \in \partial f(x_0)$.

Proof. Suppose that f has a global minimum point at x_0 . Let $x \in X$. Then

$$\begin{aligned} d^+f(x_0)(x) &= \lim_{t \rightarrow 0^+} \frac{f(x_0 + tx) - f(x_0)}{t} \\ &\geq 0 \end{aligned}$$

So $0 \leq df^+(x_0)$ and $0 \in \partial f(x_0)$.

Conversely, suppose that $0 \in \partial f(x_0)$. Let $x \in A$. Then

$$\begin{aligned} 0 &= 0(x - x_0) \\ &\leq f(x) - f(x_0) \end{aligned}$$

So that $f(x_0) \leq f(x)$ which implies that f has a global minimum point at x_0 . □

Exercise 15.4.0.16. Let X be a Banach space, $A \subset X$ open and convex, $f : A \rightarrow \mathbb{R}$ convex and $x_0 \in A$. If f is Frechet differentiable at x_0 , then $\partial f(x_0) = \{Df(x_0)\}$.

Proof. Clear. □

Exercise 15.4.0.17. Let X be a Banach space, $A \subset X$ open and convex, $f : A \rightarrow \mathbb{R}$ convex and $x_0 \in A$. Suppose that f is Frechet differentiable at x_0 . If $Df(x_0) = 0$, then f has a global minimum point at x_0 .

Proof. Suppose that $Df(x_0) = 0$. Since $\partial f(x_0) = \{Df(x_0)\}$, a previous exercise implies that f has a global minimum point at x_0 . □

Exercise 15.4.0.18. Let X be a Banach space, $A \subset X$ open and convex, $f : A \rightarrow \mathbb{R}$ convex and $x_0 \in A$. Suppose that f is Frechet differentiable at x_0 . Then for each $x \in A$, $f(x) \geq f(x_0) + Df(x_0)(x - x_0)$

Proof. Since $Df(x_0) \in \partial f(x_0)$, for each $x \in A$, $Df(x_0)(x - x_0) \leq f(x) - f(x_0)$. □

Exercise 15.4.0.19. Let X be a Banach space, $A \subset X$ open and convex, $f : A \rightarrow \mathbb{R}$. Suppose that f is Frechet differentiable. Then f is convex iff for each $x_0, x \in A$, $f(x) \geq f(x_0) + Df(x_0)(x - x_0)$.

Proof. Suppose that f is convex. Then the previous exercise implies that for each $x_0, x \in A$, $f(x) \geq f(x_0) + Df(x_0)(x - x_0)$. Conversely, suppose that for each $x_0, x \in A$, $f(x) \geq f(x_0) + Df(x_0)(x - x_0)$. Let $x_0, x, y \in A$. Then $f(x) \geq f(x_0) + Df(x_0)(x - x_0)$ and $f(y) \geq f(x_0) + Df(x_0)(y - x_0)$.

FINISH!!! □

Exercise 15.4.0.20. Let X be a Banach space, $A \subset X$ open and convex, and $f \in C^2(A)$. Then f is convex iff for each $x_0 \in A$, $D^2f(x_0)$ is positive semidefinite.

Hint: Define $g : A \rightarrow \mathbb{R}$ by $g(x) = f(x) - Df(x_0)(x - x_0)$ and show g is convex and use Taylor's Theorem

Proof. Suppose that f is convex. Let $x_0 \in X$. Define $g : A \rightarrow \mathbb{R}$ by $g(x) = f(x) - Df(x_0)(x - x_0)$. Since g is the sum of a convex function and an affine function, g is convex. Since $f \in C^2(A)$, we have that $g \in C^2(A)$ and it is straightforward to show that for each $x \in A$, $Dg(x) = Df(x) - Df(x_0)$ and $D^2g(x) = D^2f(x)$. In particular, $Dg(x_0) = 0$. Hence g has a global minimum point at x_0 . This implies that $D^2f(x_0)$ is positive semidefinite. Conversely, suppose that for each $x_0 \in A$, $D^2f(x_0)$ is positive semidefinite. Let

FINISH!!! □

15.5 Conjugacy

15.5.1 Convex Conjugate

Definition 15.5.1.1. Let X be a topological vector space, $A \in \text{Conv}(X)$ and $f : A \rightarrow \mathbb{R}$. We define the

- **convex conjugate domain of f** , denoted $\text{dom}_{\text{conj}} f \subset X^*$, by

$$\text{dom}_{\text{conj}} f := \left\{ \phi \in X^* : \sup_{x \in A} [\phi(x) - f(x)] < \infty \right\}$$

- **convex conjugate of f** , denoted $\text{conj } f : \text{dom}_{\text{conj}} f \rightarrow \mathbb{R}$, by

$$[\text{conj } f](\phi) := \sup_{x \in A} [\phi(x) - f(x)]$$

Note 15.5.1.2.

- When the context is clear, we may write A^* and f^* in place of $\text{dom}_{\text{conj}} f$ and $\text{conj } f$ respectively so that

$$A^* := \left\{ \phi \in X^* : \sup_{x \in A} [\phi(x) - f(x)] < \infty \right\}, \quad f^*(\phi) := \sup_{x \in A} [\phi(x) - f(x)].$$

- If X is a Hilbert space, we may identify $A^* \subset X$ and $f^* : A^* \rightarrow \mathbb{R}$ via the Riesz representation theorem by

$$A^* := \left\{ y \in X : \sup_{x \in A} [\langle y, x \rangle - f(x)] < \infty \right\}$$

and

$$f^*(y) := \sup_{x \in A} [\langle y, x \rangle - f(x)]$$

Exercise 15.5.1.3. Let X be a topological vector space, $A \in \text{Conv}(X)$ and $f : A \rightarrow \mathbb{R}$. Then $A^{**} = \left\{ \hat{x} : x \in X \text{ and } \sup_{\phi \in A^*} [\hat{x}(\phi) - f^*(\phi)] < \infty \right\}$.

Proof. By definition,

$$\begin{aligned} A^{**} &= \left\{ \mu \in X^{**} : \sup_{\phi \in A^*} [\mu(\phi) - f^*(\phi)] < \infty \right\} \\ &= \left\{ \mu \in \widehat{X} : \sup_{\phi \in A^*} [\mu(\phi) - f^*(\phi)] < \infty \right\} \quad (\text{by ex in duality section on tvs chapter}) \\ &= \left\{ \hat{x} : x \in X \text{ and } \sup_{\phi \in A^*} [\hat{x}(\phi) - f^*(\phi)] < \infty \right\}. \end{aligned}$$

□

Note 15.5.1.4. Let $\iota : X \rightarrow \widehat{X}$ be the canonical embedding. When the context is clear, we identify

- A^{**} with $\iota^{-1}(A^{**})$,
- f^{**} with $f^{**} \circ \iota$.

Exercise 15.5.1.5. Let X be a vector space, $A \in \text{Conv}(X)$ and $f : A \rightarrow \mathbb{R}$. Then

1. for each $x \in A$ and $\phi \in A^*$, $f^*(\phi) \geq \phi(x) - f(x)$,
2. $A \subset A^{**}$.

Proof.

1. For each $x \in A$ and $\phi \in A^*$,

$$\begin{aligned} f^*(\phi) &= \sup_{x' \in A} [\phi(x') - f(x')] \\ &\geq \phi(x) - f(x). \end{aligned}$$

2. Let $x \in A$. Then

$$\begin{aligned} f^{**}(x) &= \sup_{\phi \in A^*} [\hat{x}(\phi) - f^*(\phi)] \\ &= \sup_{\phi \in A^*} [\phi(x) - f^*(\phi)] \\ &\leq f(x) \quad (\text{by the previous part}) \\ &< \infty. \end{aligned}$$

Thus $x \in A^{**}$. Since $x \in A$ is arbitrary, we have that for each $x \in A$, $x \in A^{**}$. Hence $A \subset A^{**}$. □

Exercise 15.5.1.6. Let X be a locally convex space, $A \in \text{Conv}(X)$ and $f : A \rightarrow \mathbb{R}$. Then

1. A^* is convex, $f^* : A^* \rightarrow \mathbb{R}$ is convex and weak* lower semicontinuous.
2. A^{**} is convex and $f^{**} : A^{**} \rightarrow \mathbb{R}$ is convex and weakly lower semicontinuous.

Proof. **FINISH!!!, clean up a bit and expand.**

1. For $x \in A$, define $g_x : X^* \rightarrow \mathbb{R}$ by $g_x(\phi) = \hat{x}(\phi) - f(x)$. Then for each $x \in A$, g_x is convex and weak* lower semicontinuous since it is affine and w^* -continuous. Exercise 15.3.0.17 implies that $A^* = \{\phi \in X^* : \sup_{x \in A} g_x(\phi) < \infty\}$ is convex and $f^* = \sup_{x \in A} g_x$ is convex. **ex in topology section on semicontinuity** implies that f^* is w^* -lower semicontinuous.
2. For $\phi \in A^*$, define $h_\phi : X \rightarrow \mathbb{R}$ by $h_\phi(x) = \phi(x) - f^*(\phi)$. Then for each $\phi \in A^*$, h_ϕ is convex and weakly lower semicontinuous since it is affine and weakly continuous. Exercise 15.3.0.17 implies that $A^{**} = \{x \in X : \sup_{\phi \in A^*} h_\phi(x) < \infty\}$ is convex and $f^{**} = \sup_{\phi \in A^*} h_\phi$ is convex. **ex in topology section on semicontinuity** implies that f^* is weakly lower semicontinuous. □

Exercise 15.5.1.7. Let X be a vector space, $A \in \text{Conv}(X)$ and $f : A \rightarrow \mathbb{R}$. Then $f^{**}|_A \leq f$.

Proof. Let $x \in A$ and $\epsilon > 0$. **A prev ex** implies that $x \in A^{**}$. Since

$$\begin{aligned} f^{**}(x) - \epsilon &< f^{**}(x) \\ &= \sup_{\phi \in A^*} [\hat{x}(\phi) - f^*(\phi)], \end{aligned}$$

there exists $\phi \in A^*$ such that $\hat{x}(\phi) - f^*(\phi) > f^{**}(x) - \epsilon$. Hence

$$\begin{aligned} f^{**}(x) - \epsilon &< \hat{x}(\phi) - f^*(\phi) \\ &= \phi(x) - f^*(\phi) \\ &\leq \phi(x) - [\phi(x) - f(x)] \quad (\text{by prev ex}) \\ &= f(x). \end{aligned}$$

Since $\epsilon > 0$ is arbitrary, we have that for each $\epsilon > 0$, $f^{**}(x) - \epsilon < f(x)$. Hence $f^{**}(x) \leq f(x)$. Since $x \in A$ is arbitrary, we have that for each $x \in A$, $f^{**}(x) \leq f(x)$. Thus $f^{**}|_A \leq f$. □

Exercise 15.5.1.8. Let X be a locally convex space, $A \in \text{Conv}(X)$, $f : A \rightarrow \mathbb{R}$ and $x_0 \in X$. Suppose that f is convex and f is lower semicontinuous.

1. if $x_0 \in A$, then for each $\epsilon > 0$, there exists $\phi \in A^*$ such that for each $x \in A$, $f(x) > f(x_0) + \phi(x - x_0) - \epsilon$
2. if $x_0 \notin A$, then for each $M \in \mathbb{R}$, there exists $\phi \in A^*$ such that for each $x \in A$, $f(x) > M + \phi(x - x_0)$

Hint: Apply second Hahn-Banach separation theorem to $\{(x_0, f(x_0) - \epsilon)\}$ and $\text{epi } f$.

Proof.

1. Suppose that $x_0 \in A$. Let $\epsilon > 0$. Since f is convex and lower semicontinuous, $\text{epi } f \subset X \times \mathbb{R}$ is convex and closed, $\{(x_0, f(x_0) - \epsilon)\} \subset X \times \mathbb{R}$ is convex and compact and $\{(x_0, f(x_0) - \epsilon)\} \cap \text{epi } f = \emptyset$. Thus, there exists $\lambda \in \mathbb{R}$, $\psi \in X^*$ and $k \in \mathbb{R}$ such that for each $x \in A$ and $r \geq f(x)$,

$$\psi(x) + \lambda r < k < \psi(x_0) + \lambda(f(x_0) - \epsilon)$$

Taking $(x, r) = (x_0, f(x_0))$ implies that $0 < -\lambda\epsilon$ and hence that $\lambda < 0$. Set $\phi = |\lambda|^{-1}\psi$. For $x \in A$, set $r = f(x)$. Then

$$\begin{aligned} \psi(x) - |\lambda|f(x) &< \psi(x_0) - |\lambda|(f(x_0) - \epsilon) \\ \iff |\lambda|^{-1}\psi(x) - f(x) &< |\lambda|^{-1}\psi(x_0) - (f(x_0) - \epsilon) \\ \iff \phi(x) - f(x) &< \phi(x_0) - (f(x_0) - \epsilon) \\ \iff f(x) &> f(x_0) + \phi(x - x_0) - \epsilon \end{aligned}$$

Since for each $x \in A$, $\phi(x) - f(x) < \phi(x_0) - f(x_0) + \epsilon$, we have that

$$\begin{aligned} \sup_{x \in A} [\phi(x) - f(x)] &\leq \phi(x_0) - f(x_0) + \epsilon \\ &< \infty \end{aligned}$$

So $\phi \in A^*$.

2. Suppose that $x_0 \notin A$. Let $M \in \mathbb{R}$. Repeat the previous argument for (x_0, M) and $\text{epi } f$.

□

Exercise 15.5.1.9. Let X be a locally convex space, $A \in \text{Conv}(X)$ and $f : A \rightarrow \mathbb{R}$. Suppose that f is convex and f is lower semicontinuous. Then

1. $A = A^{**}$,
2. $f = f^{**}$.

Proof.

1. A previous exercise implies that $A \subset A^{**}$. Let $x_0 \in X$. Suppose that $x_0 \notin A$. Let $M \in \mathbb{R}$. The previous exercise implies that there exists $\phi_0 \in A^*$ such that for each $x \in A$, $f(x) > M + \phi_0(x - x_0)$. Then

$$\begin{aligned} \phi_0(x_0) - f^*(\phi_0) &= \phi_0(x_0) - \sup_{x \in A} [\phi_0(x) - f(x)] \\ &= \phi_0(x_0) + \inf_{x \in A} [f(x) - \phi_0(x)] \\ &\geq \phi_0(x_0) + (M - \phi_0(x_0)) \\ &= M \end{aligned}$$

Therefore

$$\begin{aligned} \sup_{\phi \in A^*} [\phi(x_0) - f^*(\phi)] &\geq \phi_0(x_0) - f^*(\phi_0) \\ &\geq M \end{aligned}$$

Since $M \in \mathbb{R}$ is arbitrary,

$$\sup_{\phi \in A^*} [\phi(x_0) - f^*(\phi)] = \infty$$

and $x_0 \notin A^{**}$. So $A^c \subset (A^{**})^c$, which implies that $A^{**} \subset A$. Thus $A^{**} = A$.

2. A previous exercise and the prev part imply that

$$\begin{aligned} f^{**} &= f^{**}|_A \\ &\leq f. \end{aligned}$$

Suppose that $f \not\leq f^{**}$. Then there exists $x_0 \in A$ such that $f(x_0) > f^{**}(x_0)$. Choose $\epsilon > 0$ such that $f(x_0) > f^{**}(x_0) + 2\epsilon$. A previous exercise implies that there exists $\phi \in A^*$ such that for each $x \in A$, $f(x) > f(x_0) + \phi(x - x_0) - \epsilon$. Choose $a \in A$ such that $f^*(\phi) - \epsilon < \phi(a) - f(a)$. Then

$$\begin{aligned} f(x_0) &> f^{**}(x_0) + 2\epsilon \\ &\geq \phi(x_0) - f^*(\phi) + 2\epsilon \\ &> \phi(x_0 - a) + f(a) + \epsilon \\ &> \phi(x_0 - a) + f(x_0) + \phi(a - x_0) - \epsilon + \epsilon \\ &= f(x_0) \end{aligned}$$

which is a contradiction. So $f \leq f^{**}$ and hence $f = f^{**}$.

□

15.5.2 Convex Envelope

Definition 15.5.2.1. Let X be a locally convex space, $A \in \text{Conv}(X)$ and $f : A \rightarrow \mathbb{R}$. We define the **convex envelope of f** , denoted $\text{env } f : A \rightarrow \mathbb{R}$, by

$$\text{env } f(x) := \sup\{h(x) : h \in \text{Aff}(A) \text{ and } h \leq f\}.$$

Exercise 15.5.2.2.

Chapter 16

Topological Groups

16.1 Introduction

Definition 16.1.0.1. Let G be a group, we define $\text{mult} : G \times G \rightarrow G$ and $\text{inv} : G \rightarrow G$ by $\text{mult}(g, h) = gh$ and $\text{inv}(g) = g^{-1}$ respectively.

Definition 16.1.0.2. Let G be a group and $\mathcal{T}$ a topology on G . Then $(G, \mathcal{T})$ is said to be a **topological group** if $\text{mult} : G \times G \rightarrow G$ and $\text{inv} : G \rightarrow G$ are continuous.

Note 16.1.0.3. For the remainder of this chapter, measurability is in reference to $(G, \mathcal{B}(\mathcal{T}))$. That is, the measurable sets are the Borel sets.

Definition 16.1.0.4. Let G be a topological group. We define

$$\text{Homeo}(G) = \{\phi : G \rightarrow G : \phi \text{ is a homeomorphism}\}$$

Note 16.1.0.5. Let G be a topological group. Then $\text{Homeo}(G)$ is a group.

Exercise 16.1.0.6. Let G be a topological group. Then $\text{inv} \in \text{Homeo}(G)$.

Proof. By assumption inv is continuous. We know from basic group theory that inv is a bijection with $\text{inv}^{-1} = \text{inv}$. □

Definition 16.1.0.7. Let G be a group and $S \subset G$, then S is said to be **symmetric** if $\text{inv}(S) = S$, (i.e. $S^{-1} = S$).

Definition 16.1.0.8. Let G be a topological group and $\phi : G \rightarrow G$. Then ϕ is said to be an **automorphism** of G if ϕ is a homomorphism and a homeomorphism. We define

$$\text{Aut}(G) = \{\phi : G \rightarrow G : \phi \text{ is an automorphism}\}$$

Exercise 16.1.0.9. Let G be a topological group. Then $\text{inv} \in \text{Aut}(G)$ iff G is abelian.

Proof. Basic group theory tells us that inv is a homomorphism iff G is abelian. □

Definition 16.1.0.10. Let G be a group and $g \in G$. We define the **left and right translation maps**, denoted $l_g : G \rightarrow G$ and $r_g : G \rightarrow G$ respectively, by $l_g(x) = gx$ and $r_g(x) = xg^{-1}$.

Exercise 16.1.0.11. Let G be a topological group and $g \in G$. Then $l_g, r_g \in \text{Homeo}(G)$.

Proof. By assumption l_g and r_g are continuous. We know from basic group theory that l_g and r_g are bijections with $l_g^{-1} = l_{g^{-1}}$ and $r_g^{-1} = r_{g^{-1}}$ so l_g and r_g are homeomorphisms. □

Exercise 16.1.0.12. Let G be a topological group. Define $\phi, \psi : G \rightarrow \text{Homeo}(G)$ by $\phi(g) = l_g$ and $\psi(g) = r_g$. Then ϕ, ψ are homomorphisms.

Proof. Let $g_1, g_2 \in G$. Then

$$l_{g_1} \circ l_{g_2}(x) = l_{g_1}(g_2x) = g_1g_2x = l_{g_1g_2}(x)$$

and

$$r_{g_1} \circ r_{g_2}(x) = r_{g_1}(xg_2^{-1}) = xg_2^{-1}g_1^{-1} = x(g_1g_2)^{-1} = r_{g_1g_2}(x)$$

□

Exercise 16.1.0.13. Let G be a topological group. Then for each $U \subset G$ and $g \in G$, if U is open, then gU , Ug and U^{-1} are open.

Proof. Let $U \subset G$ and $g \in G$. Suppose that U is open. Since l_g, r_g and inv are homeomorphisms, $l_g(U) = gU$, $r_g(U) = Ug$ and $\text{inv}(U) = U^{-1}$ are open. □

Definition 16.1.0.14. Let G be a topological group, $y \in G$ and $f \in L^0$. Define $L_y, R_y : L^0(G) \rightarrow L^0(G)$ by $L_yf = f \circ l_y^{-1}$ and $R_yf = f \circ r_y^{-1}$, that is, $L_yf(x) = f(y^{-1}x)$ and $R_yf(x) = f(xy)$.

Exercise 16.1.0.15. Let G be a topological group and $y \in G$. Then $L_y, R_y \in \text{Sym}(L^0(G))$.

Proof. It is straight forward to show that $L_y^{-1} = L_{y^{-1}}$ and $R_y^{-1} = R_{y^{-1}}$. □

Exercise 16.1.0.16. Let G be a topological group. Define $\phi, \psi : G \rightarrow \text{Sym}(L^0(G))$ by $\phi(y) = L_y$ and $\psi(y) = R_y$. Then ϕ and ψ are homomorphisms.

Proof. Let $y, z \in G$ and $f \in L^0(G)$. Then

$$\begin{aligned} L_y \circ L_z(f) &= L_y(L_z(f)) \\ &= L_y(f \circ l_z^{-1}) \\ &= (f \circ l_z^{-1}) \circ l_y^{-1} \\ &= f \circ (l_z^{-1} \circ l_y^{-1}) \\ &= f \circ (l_y \circ l_z)^{-1} \\ &= f \circ l_{yz}^{-1} \\ &= L_{yz}(f) \end{aligned}$$

The case is similar for R_y and R_z . □

Exercise 16.1.0.17. Let G be a topological group, $U \in \mathcal{B}(G)$ and $y \in G$. Then $L_y\chi_U = \chi_{yU}$ and $R_y\chi_U = \chi_{Uy^{-1}}$.

Proof. Let $x \in G$. Then

$$\begin{aligned} L_y\chi_U(x) = 1 &\iff y^{-1}x \in U \\ &\iff x \in yU \\ &\iff \chi_{yU}(x) = 1 \end{aligned}$$

The case is similar for R_y . □

Exercise 16.1.0.18. Let G be a topological group, $y \in G$ and $f \in L^0(G)$. Then $\text{supp}(L_yf) = y \text{supp}(f)$ and $\text{supp}(R_yf) = \text{supp}(f)y^{-1}$.

Proof. Put $A = \{x \in G : L_yf(x) \neq 0\}$ and $B = \{x \in G : f(x) \neq 0\}$. Then

$$\begin{aligned} x \in A &\iff L_yf(x) \neq 0 \\ &\iff f(y^{-1}x) \neq 0 \\ &\iff y^{-1}x \in B \\ &\iff x \in yB \end{aligned}$$

Thus $A = yB$ which implies that $\text{cl } A = y \text{cl } B$. Therefore $\text{supp}(L_yf) = y \text{supp}(f)$. □

Exercise 16.1.0.19. Let G be a topological group and $y \in G$. Then L_y, R_y are linear and if we restrict to the bounded measurable functions, then $L_y, R_y \in L(B(G))$ and $\|L_y\|, \|R_y\| = 1$.

Proof. Let $f, g \in L^0(G)$ and $\lambda \in \mathbb{C}$. Then

$$\begin{aligned} L_y(\lambda f + g)(x) &= (\lambda f + g)(y^{-1}x) \\ &= \lambda f(y^{-1}x) + g(y^{-1}x) \\ &= \lambda L_y f(x) + L_y g(x) \end{aligned}$$

So L_y is linear. Next, we restrict to $B(G) \cap L^0$. We note that

$$\{|f(y^{-1}x)| : x \in y \operatorname{supp}(f)\} = \{|f(x)| : x \in \operatorname{supp}(f)\}$$

This implies that

$$\begin{aligned} \|L_y f\|_u &= \sup_{x \in \operatorname{supp}(L_y f)} |L_y f(x)| \\ &= \sup_{x \in y \operatorname{supp}(f)} |f(y^{-1}x)| \\ &= \sup_{x \in \operatorname{supp}(f)} |f(x)| \\ &= \|f\|_u \end{aligned}$$

So L_y is bounded. Hence $L_y \in L(L^0)$. The case is similar for R_y . □

Definition 16.1.0.20. Let G be a topological group. We say that G is a **locally compact group** if G is locally compact and Hausdorff.

16.2 Group Actions

16.2.1 Introduction

Note 16.2.1.1. Let X, Y, Z be sets. We recall that for $f : X \times Y \rightarrow Z$, $a \in X$ and $b \in Y$, the maps $f_a : Y \rightarrow Z$ and $f^b : X \rightarrow Z$ are defined by

$$f_a(y) = f(a, y) \quad f^b(x) = f(x, b)$$

Definition 16.2.1.2. Let $\mathcal{C}$ a concrete category with products, $G, X \in \text{Obj}(\mathcal{C})$ and $\phi \in \text{Hom}_{\mathcal{C}}(G \times X, X)$. Suppose that G is a group. Then ϕ is said to be a **group action** of G on X if

1. for each $x \in X$, $\phi_e = \text{id}_X$
2. for each $g, h \in G$, $\phi_{gh} = \phi_g \circ \phi_h$

Note 16.2.1.3. When the context is clear, we will write $g \cdot x$ in place of $\phi(g, x)$.

Exercise 16.2.1.4. Let $\mathcal{C}$ a category with products, $G, X \in \text{Obj}(\mathcal{C})$ and $\phi \in \text{Hom}_{\mathcal{C}}(G \times X, X)$. Suppose that G is a group and ϕ group action. Then for each $g \in G$, $\phi_g \in \text{Aut}(X)$.

Proof. Let $g \in G$. Then

$$\begin{aligned} \phi_g \circ \phi_{g^{-1}}(x) &= \phi_g(\phi_{g^{-1}}(x)) \\ &= g \cdot (g^{-1} \cdot x) \\ &= (gg^{-1}) \cdot x \\ &= e \cdot x \\ &= x \end{aligned}$$

Since $x \in X$ is arbitrary, $\phi_g \circ \phi_{g^{-1}} = \text{id}_X$. Similarly, $\phi_{g^{-1}} \circ \phi_g = \text{id}_X$. Hence $\phi_g \in \text{Aut}(X)$. □

Definition 16.2.1.5. Let $\mathcal{C}$ a category with products, $G, X \in \text{Obj}(\mathcal{C})$ and $\phi \in \text{Hom}_{\mathcal{C}}(G \times X, X)$. Suppose that G is a group and ϕ group action. We define $\hat{\phi} : G \rightarrow \text{Aut}(X)$ by $\hat{\phi}(g) = \phi_g$.

Exercise 16.2.1.6. Let $\mathcal{C}$ a category with products, $G, X \in \text{Obj}(\mathcal{C})$ and $\phi \in \text{Hom}_{\mathcal{C}}(G \times X, X)$. Suppose that G is a group and ϕ group action. Then $\hat{\phi} : G \rightarrow \text{Aut}(X)$ is a group homomorphism.

Proof. Clear by definition. □

16.2.2 Homogeneous Spaces

Definition 16.2.2.1. Let G be a topological group, X a topological space and $\phi : G \times X \rightarrow X$ a continuous group action. Then (X, ϕ) is said to be a **homogeneous G -space** if

- ϕ is transitive
- for each $x \in X$, $\phi_x : G \rightarrow X$ is open

Definition 16.2.2.2. Let G be a topological group, $H < G$. We define $\phi_H : H \times G \rightarrow G$ by $\phi(h, g) = gh^{-1}$.

Exercise 16.2.2.3.

Exercise 16.2.2.4. Let G be a topological group, $H < G$ a closed subgroup of G . Then $(G/H, \phi_H)$

Exercise 16.2.2.5. Let G be a topological group, $H < G$ a closed subgroup of G and (X, ϕ) a homogeneous G -space.

16.2.3 Common Examples

Exercise 16.2.3.1. Let H be a Hilbert space and $x, y \in H$. Then $\|x\| = \|y\|$ iff there exists $U \in U(H)$ such that $x = Uy$.

Proof.

- ($\implies$) : Suppose that $\|x\| = \|y\|$. An exercise
- ($\impliedby$) :

□

Exercise 16.2.3.2. Let H be a Hilbert space. Then

1. $\|\cdot\| : H \rightarrow [0, \infty)$ is a quotient map
2. $H/U(H)$ is homeomorphic to $[0, \infty)$

Proof. content...

□

Definition 16.2.3.3. Let $n, k \in \mathbb{N}$. Suppose that $n \geq k$. We define the **Stiefel manifold**, denoted $V_k(\mathbb{R}^n)$, by

$$V_k(\mathbb{R}^n) = \{A \in \mathbb{R}^{n \times k} : A^*A = I\}$$

We define the **orthogonal matrices**, denoted by $O(n)$, by

$$O(n) = V_n(\mathbb{R}^n)$$

Note 16.2.3.4. We note that for each $X \in V_k(\mathbb{R}^n)$, $\text{rank } X = k$ and for each $U \in O(n)$, $UU^* = I$.

Exercise 16.2.3.5. Let $X, Y \in \mathbb{R}^{n \times k}$. Suppose that $\text{rank } X = k$ and $\text{rank } Y = k$. Then $XX^* = YY^*$ iff there exists $U \in O(k)$ such that $X = YU$.

Hint: $\text{rank } X = \text{rank } X^*X$.

Proof.

- ($\implies$) :
Suppose that $XX^* = YY^*$. Since $\text{rank } X = k$, we have that

$$\begin{aligned} \text{rank } XX^* &= \text{rank } X \\ &= k \end{aligned}$$

Since $X^*X \in \mathbb{R}^{k \times k}$, X^*X is invertible. Hence

$$\begin{aligned} X &= XI \\ &= X(X^*X)(X^*X)^{-1} \\ &= (XX^*)X(X^*X)^{-1} \\ &= (YY^*)X(X^*X)^{-1} \\ &= Y(Y^*X)(X^*X)^{-1} \end{aligned}$$

Set $U = (Y^*X)(X^*X)^{-1}$. Then $X = YU$ and

$$\begin{aligned} U^*U &= \left((Y^*X)(X^*X)^{-1} \right)^* (Y^*X)(X^*X)^{-1} \\ &= (X^*X)^{-1} (X^*Y)(Y^*X)(X^*X)^{-1} \\ &= (X^*X)^{-1} X^*(YY^*)X(X^*X)^{-1} \\ &= (X^*X)^{-1} X^*(XX^*)X(X^*X)^{-1} \\ &= (X^*X)^{-1} (X^*X)(X^*X)(X^*X)^{-1} \\ &= I \end{aligned}$$

Thus $U \in O(k)$.

- ($\Leftarrow$):

Suppose that there exists $U \in O(k)$ such that $X = YU$. Then

$$\begin{aligned}
 XX^* &= (YU)(YU)^* \\
 &= (YU)(U^*Y^*) \\
 &= Y(UU^*)Y^* \\
 &= YIY^* \\
 &= YY^*
 \end{aligned}$$

□

Exercise 16.2.3.6. Define $f : V$

16.3 Quotient Groups

Definition 16.3.0.1. Let

16.4 Automorphism Groups of Metric Spaces

Definition 16.4.0.1. Let (X, τ) be a topological space. Define

$$\text{Aut}(X) = \{\sigma : X \rightarrow X : \sigma \text{ is a homeomorphism}\}$$

Exercise 16.4.0.2. Let (X, d) be a compact metric space. Then $(\text{Aut}(X), d_u)$ is a topological group.

Proof. Let $(\sigma_n)_{n \in \mathbb{N}}, (\tau_n)_{n \in \mathbb{N}} \subset \text{Aut}(X)$ and $\sigma, \tau \in \text{Aut}(X)$. Suppose that $\sigma_n \xrightarrow{u} \sigma$ and $\tau_n \xrightarrow{u} \tau$.

1. Let $\epsilon > 0$. Since X is compact and σ is continuous, σ is uniformly continuous. Then there exists $\delta > 0$ such that for each $x, y \in X$, $d(x, y) < \delta$ implies that $d(\sigma(x), \sigma(y)) \leq \epsilon/2$. Choose $N_\sigma \in \mathbb{N}$ such that for each $n \in \mathbb{N}$, $n \geq N_\sigma$ implies that $d_u(\sigma_n, \sigma) < \epsilon/2$. Choose $N_\tau \in \mathbb{N}$ such that for each $n \in \mathbb{N}$, $n \geq N_\tau$ implies that $d_u(\tau_n, \tau) < \delta$. Put $N = \max(N_\sigma, N_\tau)$. Let $n \in \mathbb{N}$ and $x \in X$. Suppose that $n \geq N$. Then

$$\begin{aligned} d(\sigma_n \circ \tau_n(x), \sigma \circ \tau(x)) &\leq d(\sigma_n(\tau_n(x)), \sigma(\tau_n(x))) + d(\sigma(\tau_n(x)), \sigma(\tau(x))) \\ &< \epsilon/2 + \epsilon/2 \\ &= \epsilon \end{aligned}$$

So $d_u(\sigma_n \circ \tau_n, \sigma \circ \tau) \leq \epsilon$ and $\circ : \text{Aut}(X)^2 \rightarrow \text{Aut}(X)$ is continuous.

2. Suppose that $\sigma = \text{id}_X$. Let $\epsilon > 0$. Then there exists $N \in \mathbb{N}$ such that for each $n \in \mathbb{N}$, $n \geq N$ implies that $d_u(\sigma_n, \text{id}_X) < \epsilon$. Let $n \in \mathbb{N}$. Suppose that $n \geq N$. Then

$$\begin{aligned} \sup_{x \in X} d(\sigma_n^{-1}(x), x) &= \sup_{x \in \sigma_n(X)} d(\sigma_n^{-1}(x), x) \\ &= \sup_{x \in X} d(\sigma_n^{-1}(\sigma_n(x)), \sigma_n(x)) \\ &= \sup_{x \in X} d(x, \sigma_n(x)) \\ &< \epsilon \end{aligned}$$

So $\sigma_n^{-1} \xrightarrow{u} \text{id}_X$. Now suppose that $\sigma \neq \text{id}_X$. Since $\sigma_n \xrightarrow{u} \sigma$, part (1) implies that $\sigma^{-1} \circ \sigma_n \xrightarrow{u} \text{id}_X$. Applying the result from above, we get that $\sigma_n^{-1} \circ \sigma \xrightarrow{u} \text{id}_X$. Applying part (1) again implies that $\sigma_n^{-1} \xrightarrow{u} \sigma^{-1}$. So the map $\sigma \mapsto \sigma^{-1}$ is continuous.

Hence $\text{Aut}(X)$ is a topological group. □

Definition 16.4.0.3. Let (X, d) be a metric space. Define

$$\text{Aut}(X, d) = \{\sigma : X \rightarrow X : \sigma \text{ is an isometric isomorphism}\}$$

Exercise 16.4.0.4. Let (X, d) be a compact metric space. Then $(\text{Aut}(X, d), d_u)$ is a compact subgroup of $(\text{Aut}(X), d_u)$.

Proof. Clearly, $(\text{Aut}(X, d), d_u)$ is a topological subgroup. To show compactness, use the Arzela Ascoli theorem. □

Definition 16.4.0.5. Let (X, τ) be a topological space and $\mu : \mathcal{B}(X) \rightarrow \mathbb{R}$ a Borel measure. Define

$$\text{Aut}(X, \mu) = \{\sigma \in \text{Aut}(X) : \sigma_*\mu = \mu\}$$

Exercise 16.4.0.6. Let (X, d) be a compact metric space and $\mu : \mathcal{B}(X) \rightarrow \mathbb{R}$ an outer-regular Borel measure. Then $\text{Aut}(X, \mu)$ is a closed subgroup of $\text{Aut}(X)$.

Proof. It is clear that $\text{Aut}(X, \mu)$ is a subgroup of $\text{Aut}(X)$. Let $(\sigma_n)_{n \in \mathbb{N}} \subset \text{Aut}(X, \mu)$ and $\sigma \in \text{Aut}(X)$. Suppose that $\sigma_n \xrightarrow{u} \sigma$. Let $E \subset X$ be closed, $U \subset X$ open and suppose that $E \subset U$. An exercise in the section on metric spaces tells us that there exists $N \in \mathbb{N}$ such that for each $n \in \mathbb{N}$, $n \geq N$ implies that $\sigma(E) \subset \sigma_n(U)$. Then

$$\begin{aligned} \mu(\sigma(E)) &\leq \mu(\sigma_n(U)) \\ &= \mu(U) \end{aligned}$$

Therefore, since μ is outer regular, $\mu(\sigma(E)) \leq \mu(E)$. Since $\sigma_n^{-1} \xrightarrow{u} \sigma^{-1}$, we may apply the above argument to obtain that

$$\begin{aligned}\mu(E) &= \mu(\sigma^{-1}(\sigma(E))) \\ &\leq \mu(\sigma(E))\end{aligned}$$

Hence $\mu(E) = \mu(\sigma(E))$. Applying the whole argument above thus far to σ^{-1} , we see that $\mu(E) = \mu(\sigma^{-1}(E))$. Since $E \subset X$ is an arbitrary closed set and $\mathcal{B}(X) = \sigma(E \subset X : E \text{ is closed})$, we have that $\mu = \sigma_*\mu$. Thus $\sigma \in \text{Aut}(X, \mu)$ which implies that $\text{Aut}(X, \mu)$ is closed. $\square$

Definition 16.4.0.7. Let (X, d) be a compact metric space and $\mu : \mathcal{B}(X) \rightarrow \mathbb{R}$ an outer-regular Borel measure. Define $\text{Aut}(X, d, \mu) = \text{Aut}(X, d) \cap \text{Aut}(X, \mu)$.

Exercise 16.4.0.8. Let (X, d) be a compact metric space and $\mu : \mathcal{B}(X) \rightarrow \mathbb{R}$ an outer-regular Borel measure. Then $\text{Aut}(X, d, \mu)$ is compact.

Proof. Since $\text{Aut}(X, d)$ is compact and $\text{Aut}(X, \mu)$ is closed, $\text{Aut}(X, d, \mu)$ is compact. $\square$

Chapter 17

Group Actions

17.1 Introduction

Note 17.1.0.1. For a set X , a group G and a (left) group action $\phi : G \times X \rightarrow X$, we will write $\phi(g, x)$ as $g \cdot x$.

Definition 17.1.0.2. Let X be a set, G a group, $\phi : G \times X \rightarrow X$ a group action and $g \in G$. Define $l_g : X \rightarrow X$ by

$$l_g(x) = g \cdot x$$

Definition 17.1.0.3. Let X be a topological space, G a group and $\phi : G \times X \rightarrow X$ a group action. Then ϕ is said to be X -continuous if for each $g \in G$, ϕ_g is continuous.

Exercise 17.1.0.4. Let X be a topological space, G a group and $\phi : G \times X \rightarrow X$ an X -continuous group action. Then for each $g \in G$, $\phi_g \in \text{Homeo}(X)$.

Proof. Let $g \in G$, then ϕ_g and $\phi_g^{-1} = \phi_{g^{-1}}$ are continuous, so $\phi_g \in \text{Homeo}(X)$. □

Definition 17.1.0.5. Let (X, d) be a metric space, G a group, and $\phi : G \times X \rightarrow X$ a group action. Then ϕ is said to be an **isometric group action** if for each $g \in G$, $\phi_g : X \rightarrow X$ is an isometry.

Exercise 17.1.0.6. Let (X, d) be a metric space, G a group, and $\phi : G \times X \rightarrow X$ an isometric group action. Then ϕ is X -continuous.

Proof. Clear since isometries are continuous. □

Definition 17.1.0.7. Let X be a set, G a group, $\phi : G \times X \rightarrow X$ a group action. We define the relation $\sim_\phi \subset X \times X$ by

$$\sim_\phi = \{(a, b) \in X \times X : \text{there exists } g \in G : a = g \cdot b\}$$

Exercise 17.1.0.8. Let X be a set, G a group, $\phi : G \times X \rightarrow X$ a group action. Then $\sim_\phi$ is an equivalence relation on X .

Proof. Let $a, b, c \in X$.

- **(reflexivity):** Then $a = e \cdot a$. Hence $a \sim_\phi a$.
- **(symmetry):** Suppose that $a \sim_\phi b$. Then there exists $g \in G$ such that $a = g \cdot b$. Hence $b = g^{-1} \cdot a$. Thus $b \sim_\phi a$.
- **(transitivity):** Suppose that $a \sim_\phi b$ and $b \sim_\phi c$. Then there exist $g, h \in G$ such that $a = g \cdot b$ and $b = h \cdot c$. Then

$$\begin{aligned} a &= g \cdot b \\ &= g \cdot (h \cdot c) \\ &= (gh) \cdot c \end{aligned}$$

Hence $a \sim_\phi c$. □

Definition 17.1.0.9. Let X be a set, G a group and $\phi : G \times X \rightarrow X$ a group action. We define the **quotient of X by G** , denoted X/G , by

$$X/G = X / \sim_\phi$$

We denote the projection from X onto X/G by $\pi : X \rightarrow X/G$.

Definition 17.1.0.10. Let X be a set, G a group, $\phi : G \times X \rightarrow X$ a group action and $f : X \rightarrow \mathbb{C}$. Then f is said to be **ϕ -invariant** if for each $g \in G$ and $x \in X$, $f(g \cdot x) = f(x)$.

Exercise 17.1.0.11. Let X be a set, G a group, $\phi : G \times X \rightarrow X$ a group action and $f : X \rightarrow \mathbb{C}$. Then f is ϕ -invariant iff f is $\sim_\phi$ -invariant.

Proof.

• ($\implies$) :

Suppose that f is ϕ -invariant. Let $a, b \in X$. Suppose that $a \sim_\phi b$. Then there exists $g \in G$ such that $a = g \cdot b$. Since f is ϕ -invariant,

$$\begin{aligned} f(a) &= f(g \cdot b) \\ &= f(b) \end{aligned}$$

Since $a, b \in X$ such that $a \sim_\phi b$ are arbitrary, we have that f is $\sim_\phi$ -invariant.

• ($\impliedby$) :

Suppose that f is $\sim_\phi$ -invariant. Let $g \in G$ and $x \in X$. By definition, $x \sim_\phi g \cdot x$. Since f is $\sim_\phi$ -invariant, $f(g \cdot x) = f(x)$. Since $g \in G$ and $x \in X$ are arbitrary, f is ϕ -invariant.

□

Exercise 17.1.0.12. Let X, Y be a topological spaces, G a topological group, $\phi : G \times X \rightarrow X$ a continuous group action and $f : X \rightarrow Y$ a homeomorphism.

maybe introduce the unitary matrices, the stiefel manifold, the grassman manifold here, maybe introduce in geometry notes

17.2 Group Actions on Metric Spaces

Note 17.2.0.1. This section establishes the criteria for the existence of a metric on the orbit space of a metric space under a group action.

Definition 17.2.0.2. Let (X, d) be a metric space, G a group, and $\phi : G \times X \rightarrow X$ a group action. We define $\bar{d} : X/G \times X/G \rightarrow [0, \infty)$ by

$$\bar{d}(\bar{x}, \bar{y}) = \inf_{\substack{a \in \bar{x} \\ b \in \bar{y}}} d(a, b)$$

Exercise 17.2.0.3. Let (X, d) be a metric space, G a group, and $\phi : G \times X \rightarrow X$ an isometric group action. Then for each $x, y \in X$,

$$\bar{d}(\bar{x}, \bar{y}) = \inf_{g \in G} d(g \cdot x, y)$$

Proof. Let $x, y \in X$, $a \in \bar{x}$ and $b \in \bar{y}$. Then there exists there exists $g_a, g_b \in G$ such that $a = g_a \cdot x$ and $b = g_b \cdot y$. Set $g = g_b^{-1}g_a$. Since the map $z \mapsto g_b^{-1} \cdot z$ is an isometry,

$$\begin{aligned} d(a, b) &= d(g_a \cdot x, g_b \cdot y) \\ &= d(g_b^{-1}g_a \cdot x, y) \\ &= d(g \cdot x, y) \end{aligned}$$

Let $\epsilon > 0$. Then there exist $a^* \in \bar{x}$ and $b^* \in \bar{y}$ such that $d(a^*, b^*) < \bar{d}(\bar{x}, \bar{y}) + \epsilon$. The above argument implies that that there exists $g^* \in G$ such that

$$\begin{aligned} \inf_{g \in G} d(g \cdot x, y) &\leq d(g^* \cdot x, y) \\ &= d(a^*, b^*) \\ &< \bar{d}(\bar{x}, \bar{y}) + \epsilon \end{aligned}$$

Since $\epsilon > 0$ is arbitrary,

$$\inf_{g \in G} d(g \cdot x, y) \leq \bar{d}(\bar{x}, \bar{y})$$

Conversely, since $\{(g \cdot x, y) : g \in G\} \subset \{(a, b) : a \in \bar{x}, b \in \bar{y}\}$, we have that

$$\inf_{g \in G} d(g \cdot x, y) \geq \bar{d}(\bar{x}, \bar{y})$$

□

Exercise 17.2.0.4. Let (X, d) be a metric space, G a group, and $\phi : G \times X \rightarrow X$ an isometric group action. Then for each $x, y, z \in X$,

$$\bar{d}(\bar{x}, \bar{y}) \leq \bar{d}(\bar{x}, \bar{z}) + \bar{d}(\bar{z}, \bar{y})$$

Proof. Let $x, y, z \in X$. An exercise in section (2.1) implies that $d(\bar{x}, \bar{y}) \leq d(\bar{x}, z) + d(z, \bar{y})$. The previous exercise implies that

$$\begin{aligned} d(\bar{x}, z) &= \inf_{a \in \bar{x}} d(a, z) \\ &= \inf_{g \in G} d(g \cdot x, z) \\ &= \bar{d}(\bar{x}, \bar{z}) \end{aligned}$$

Similarly, $d(z, \bar{y}) = \bar{d}(\bar{z}, \bar{y})$. Then

$$\begin{aligned} d(\bar{x}, \bar{y}) &\leq d(\bar{x}, z) + d(z, \bar{y}) \\ &= \bar{d}(\bar{x}, \bar{z}) + \bar{d}(\bar{z}, \bar{y}) \end{aligned}$$

□

Exercise 17.2.0.5. Let (X, d) be a metric space, G a group, and $\phi : G \times X \rightarrow X$ an isometric group action. If for each $x \in X$, $\bar{x}$ is closed, then for each $x, y \in X$, $\bar{d}(\bar{x}, \bar{y}) = 0$ implies that $\bar{x} = \bar{y}$.

Proof. Suppose that for each $x \in X$, $\bar{x}$ is closed. Let $x, y \in X$. Suppose that $\bar{d}(\bar{x}, \bar{y}) = 0$. Then $\inf_{g \in G} d(g \cdot x, y) = 0$. Hence there exists $(g_n)_{n \in \mathbb{N}} \subset G$ such that $g_n \cdot x \rightarrow y$. Since $(g_n \cdot x)_{n \in \mathbb{N}} \subset \bar{x}$ and $\bar{x}$ is closed, $y \in \bar{x}$. Thus $\bar{x} = \bar{y}$. $\square$

Exercise 17.2.0.6. Let (X, d) be a metric space, G a group, and $\phi : G \times X \rightarrow X$ an isometric group action. If for each $x \in X$, $\bar{x}$ is closed, then $\bar{d}$ is a metric on X/G .

Proof. Clear by preceeding exercises. $\square$

Exercise 17.2.0.7. Let (X, d) be a metric space, $(G, \mathcal{T}_G)$ a topological group, and $\phi : G \times X \rightarrow X$ an isometric group action. Suppose that G is compact and for each $x \in X$, the map $g \mapsto g \cdot x$ is $(\mathcal{T}_G, \mathcal{T}_d)$ -continuous. Then $\bar{d}$ is a metric on X/G .

Proof. Let $x \in X$. Since G is compact and the map $g \mapsto g \cdot x$ is $(\mathcal{T}_G, \mathcal{T}_d)$ -continuous, $\bar{x} = G \cdot x$ is compact and therefore closed. The previous exercise implies that $\bar{d}$ is a metric. $\square$

Exercise 17.2.0.8. Let (X, d) be a metric space, G a group, and $\phi : G \times X \rightarrow X$ an isometric group action. Suppose that $\bar{d}$ is a metric on X/G . Then the projection map $\pi : X \rightarrow X/G$, is $(d, \bar{d})$ -Lipschitz and therefore $(\mathcal{T}_d, \mathcal{T}_{\bar{d}})$ -continuous.

Proof. Let $x, y \in X$. Then

$$\begin{aligned} \bar{d}(\pi(x), \pi(y)) &= \bar{d}(\bar{x}, \bar{y}) \\ &= \inf_{g \in G} d(g \cdot x, y) \\ &\leq d(x, y) \end{aligned}$$

$\square$

Exercise 17.2.0.9. Let (X, d) be a metric space, G a group, and $\phi : G \times X \rightarrow X$ an isometric group action. Suppose that $\bar{d}$ is a metric on X/G . Let $(x_n)_{n \in \mathbb{N}} \subset X$ and $x \in X$. Then $\bar{x}_n \xrightarrow{\bar{d}} \bar{x}$ iff there exists a sequence $(g_n)_{n \in \mathbb{N}}$ such that $g_n \cdot x_n \xrightarrow{d} x$.

Proof. Suppose that $\bar{x}_n \xrightarrow{\bar{d}} \bar{x}$. For $n \in \mathbb{N}$, choose $g_n \in G$ such that $d(g_n \cdot x_n, x) < \bar{d}(\bar{x}_n, \bar{x}) + 2^{-n}$. Then $d(g_n \cdot x_n, x) \rightarrow 0$ and $g_n \cdot x_n \xrightarrow{d} x$.

Conversely, suppose that there exists a sequence $(g_n)_{n \in \mathbb{N}}$ such that $g_n \cdot x_n \xrightarrow{d} x$. Since $\pi : X \rightarrow X/G$ is $(\mathcal{T}_d, \mathcal{T}_{\bar{d}})$ -continuous, we have that

$$\begin{aligned} g_n \cdot x_n \xrightarrow{d} x &\implies \pi(g_n \cdot x_n) \xrightarrow{\bar{d}} \pi(x) \\ &\implies \bar{x}_n \xrightarrow{\bar{d}} \bar{x} \end{aligned}$$

$\square$

Exercise 17.2.0.10. Let X be a set, $d_1, d_2 : X^2 \rightarrow [0, \infty)$ metrics, G a group and $\phi : G \times X \rightarrow X$ an isometric group action. Suppose that d_1 and d_2 are **Top**-equivalent.

1. Then $\bar{d}_1$ is a metric on X/G iff $\bar{d}_2$ is a metric on X/G
2. If $\bar{d}_1$ and $\bar{d}_2$ are metrics, then $\bar{d}_1$ and $\bar{d}_2$ are **Top**-equivalent.

Proof.

1. $\implies$ Suppose that $\bar{d}_1$ is a metric. Let $x, y \in X$. Suppose that $\bar{d}_2(\bar{x}, \bar{y}) = 0$. Then there exist $(g_n)_{n \in \mathbb{N}} \subset G$ such that $d_2(g_n \cdot x, y) \rightarrow 0$. Since d_1 and d_2 are **Top**-equivalent, $d_1(g_n \cdot x, y) \rightarrow 0$. Thus $\bar{d}_1(\bar{x}, \bar{y}) = 0$. Since $\bar{d}_1$ is a metric, $\bar{x} = \bar{y}$. Hence $\bar{d}_2$ is a metric.
- $\longleftarrow$ Similar.

2. Suppose that $\bar{d}_1$ and $\bar{d}_2$ are metrics. Let $(\bar{x}_n)_{n \in \mathbb{N}} \subset X/G$ and $\bar{x} \in X/G$.

- Suppose that $\bar{x}_n \xrightarrow{\bar{d}_1} \bar{x}$. Then there exists a sequence $(g_n)_{n \in \mathbb{N}}$ such that $g_n \cdot x_n \xrightarrow{d_1} x$. Since d_1 and d_2 are **Top**-equivalent, $g_n \cdot x_n \xrightarrow{d_2} x$. This implies that $\bar{x}_n \xrightarrow{\bar{d}_2} \bar{x}$.
- Suppose that $\bar{x}_n \xrightarrow{\bar{d}_2} \bar{x}$. Then similarly to above, $\bar{x}_n \xrightarrow{\bar{d}_1} \bar{x}$.

□

Exercise 17.2.0.11. Let X be a set, $d_1, d_2 : X^2 \rightarrow [0, \infty)$ metrics on X , G a group and $\phi : G \times X \rightarrow X$ an isometric group action. Suppose that d_1 and d_2 are equivalent. If $\bar{d}_1$ and $\bar{d}_2$ are metrics, then $\bar{d}_1$ and $\bar{d}_2$ are equivalent.

Proof. Suppose that $\bar{d}_1$ and $\bar{d}_2$ are metrics. Since d_1, d_2 are equivalent, there exist $C_1, C_2 > 0$ such that for each $x, y \in X$, $C_1 d_1(x, y) \leq d_2(x, y) \leq C_2 d_1(x, y)$. Let $x, y \in X$. Then

$$\begin{aligned} C_1 \bar{d}_1(\bar{x}, \bar{y}) &= C_1 \inf_{g \in G} d_1(g \cdot x, y) \\ &= \inf_{g \in G} C_1 d_1(g \cdot x, y) \\ &\leq \inf_{g \in G} d_2(g \cdot x, y) \\ &= \bar{d}_2(\bar{x}, \bar{y}) \end{aligned}$$

and

$$\begin{aligned} \bar{d}_2(\bar{x}, \bar{y}) &= \inf_{g \in G} d_2(g \cdot x, y) \\ &\leq \inf_{g \in G} C_2 d_1(g \cdot x, y) \\ &= C_2 \inf_{g \in G} d_1(g \cdot x, y) \\ &= C_2 \bar{d}_1(\bar{x}, \bar{y}) \end{aligned}$$

So that $C_1 \bar{d}_1 \leq \bar{d}_2 \leq C_2 \bar{d}_1$

□

Exercise 17.2.0.12. Let (X, d) be a metric space, G a group and $\phi : G \times X \rightarrow X$ an isometric group action. Suppose that $\bar{d}$ is a metric. Then $\pi : X \rightarrow X/G$ is a $(\mathcal{T}_d, \mathcal{T}_{\bar{d}})$ -quotient map.

Proof.

- Clearly π is surjective.
- Let $C \subset X/G$. Suppose that C is closed. Since π is $(\mathcal{T}_d, \mathcal{T}_{\bar{d}})$ -continuous, if $\pi^{-1}(C)$ is closed. Conversely, suppose that $\pi^{-1}(C)$ is closed. Let $(\bar{x}_n)_{n \in \mathbb{N}} \subset C$ and $\bar{x} \in X/G$. Suppose that $\bar{x}_n \xrightarrow{\bar{d}} \bar{x}$. Then there exists $(g_n)_{n \in \mathbb{N}} \subset G$ such that $g_n \cdot x_n \xrightarrow{d} x$. Since $(g_n \cdot x_n)_{n \in \mathbb{N}} \subset \pi^{-1}(C)$, $x \in \pi^{-1}(C)$. Hence $\bar{x} \in C$ and C is closed. Then Exercise 4.8.1.3 implies that π is a $(\mathcal{T}_d, \mathcal{T}_{\bar{d}})$ -quotient map.

□

Exercise 17.2.0.13. Let (X, d) be a metric space, G a group and $\phi : G \times X \rightarrow X$ an isometric group action. Suppose that $\bar{d}$ is a metric. Then $\pi : X \rightarrow X/G$ is $(\mathcal{T}_d, \mathcal{T}_{\bar{d}})$ -open.

Proof. Let $U \subset X$. Suppose that U is open. Then

$$\pi^{-1}(\pi(U)) = \bigcup_{g \in G} g \cdot U$$

Since for each $g \in G$, ϕ_g is an isometry and thus a homeomorphism, we have that for each $g \in G$, $g \cdot U$ is open. Therefore

$$\pi^{-1}(\pi(U)) = \bigcup_{g \in G} g \cdot U$$

is open. Exercise 4.8.1.9 implies that π is open.

□

Exercise 17.2.0.14. Let (X, d) be a metric space, G a group and $\phi : G \times X \rightarrow X$ an isometric group action. Suppose that $\bar{d}$ is a metric. Then $\bar{\pi} : X/G \rightarrow X/G$ is a $(\mathcal{T}_{X/G}, \mathcal{T}_{\bar{d}})$ -homeomorphism.

Proof. The previous exercises imply that $\pi : X \rightarrow X/G$ is a $(\mathcal{T}_d, \mathcal{T}_{\bar{d}})$ -quotient map and $(\mathcal{T}_d, \mathcal{T}_{\bar{d}})$ -open. Since for each $a, b \in X$, $a \sim b$ iff $\pi(a) = \pi(b)$, Exercise ?? implies that $\bar{\pi} : X/G \rightarrow X/G$ is a $(\mathcal{T}_{X/G}, \mathcal{T}_{\bar{d}})$ -homeomorphism. $\square$

Exercise 17.2.0.15. Let (X, d) be a metric space, G a group and $\phi : G \times X \rightarrow X$ an isometric group action. Suppose that $\bar{d}$ is a metric. Then $\bar{d}$ metrizes the quotient topology $\pi_*\mathcal{T}_d$ on X/G .

Proof. Immediate by the previous exercise. $\square$

17.3 Fundamental Examples

Note 17.3.0.1. This section uses results from the previous two sections to establish metrics on some fundamental orbit spaces of metric spaces under a group action.

Exercise 17.3.0.2. Procrustes Distance:

Consider the metric space $(\mathbb{C}^{n \times d}, \|\cdot\|_F)$, topological group $(U_d, \|\cdot\|_F)$ and the (right) action $\phi : X \times U_d \rightarrow X$ by $X \cdot U = XU$. Then

1. ϕ is a continuous isometric group action
2. U_d is compact
3. $\bar{d}$ is a metric on $\mathbb{C}^{n \times d}/U_d$

Proof. Clear. □

Exercise 17.3.0.3. Let X be a compact metric space and $\mu : \mathcal{B}(X) \rightarrow [0, \infty]$ a Borel measure. Define the (right) group action $\phi : L^1(\mu) \times \text{Aut}(X, \mu) \rightarrow L^1(\mu)$ by

$$f \cdot \sigma = f \circ \sigma$$

Then ϕ is an isometric group action.

Proof. Let $\sigma \in \text{Aut}(X, \mu)$ and $f \in L^1(\mu)$. Then

$$\begin{aligned} \|f \cdot \sigma\|_1 &= \int_X |f \circ \sigma| d\mu \\ &= \int_X |f| \circ \sigma d\mu \\ &= \int_{\sigma(X)} |f| d\sigma_*\mu \\ &= \int_{\sigma(X)} |f| d\mu \\ &= \int_X |f| d\mu \\ &= \|f\|_1 \end{aligned}$$

□

Exercise 17.3.0.4. Let X be a compact metric space and $\mu : \mathcal{B}(X) \rightarrow [0, \infty]$ a Radon measure. Define the (right) group action $\phi : L^1(\mu) \times \text{Aut}(X, \mu) \rightarrow L^1(\mu)$ by

$$f \cdot \sigma = f \circ \sigma$$

Then for each $f \in L^1(\mu)$, the map $\sigma \mapsto f \cdot \sigma$ is continuous.

Proof. Let $f \in L^1(\mu)$, $(\sigma_n)_{n \in \mathbb{N}} \subset \text{Aut}(X, \mu)$ and $\sigma \in \text{Aut}(X, \mu)$. Suppose that $\sigma_n \xrightarrow{u} \sigma$. Since μ is Radon, $C_c(X)$ is dense in $L^1(\mu)$ and therefore, there exists $\phi \in C_c(X)$ such that $\|\phi - f\| < \epsilon/3$. Since X is compact and μ is Radon, $\mu(X) < \infty$. Since ϕ is uniformly continuous, Exercise 7.1.0.13 implies that $\phi \circ \sigma_n \xrightarrow{u} \phi \circ \sigma$. So there exists $N \in \mathbb{N}$ such that for each $n \in \mathbb{N}$, $n \geq N$ implies that $\|\phi \circ \sigma_n - \phi \circ \sigma\|_u < \frac{\epsilon}{3(\mu(X)+1)}$. Let $n \in \mathbb{N}$. Suppose that $n \geq N$. Then

$$\begin{aligned} \|f \circ \sigma_n - f \circ \sigma\|_1 &\leq \|f \circ \sigma_n - \phi \circ \sigma_n\|_1 + \|\phi \circ \sigma_n - \phi \circ \sigma\|_1 + \|\phi \circ \sigma - f \circ \sigma\|_1 \\ &= \|(f - \phi) \circ \sigma_n\|_1 + \|\phi \circ \sigma_n - \phi \circ \sigma\|_1 + \|(\phi - f) \circ \sigma\|_1 \\ &= \|f - \phi\|_1 + \|\phi \circ \sigma_n - \phi \circ \sigma\|_1 + \|\phi - f\|_1 \\ &= \|f - \phi\|_1 + \|\phi \circ \sigma_n - \phi \circ \sigma\|_u \mu(X) + \|\phi - f\|_1 \\ &< \frac{\epsilon}{3} + \frac{\epsilon}{3} + \frac{\epsilon}{3} \\ &= \epsilon \end{aligned}$$

So that $f \circ \sigma_n \xrightarrow{u} f \circ \sigma$ which implies that the map $\sigma \mapsto f \cdot \sigma$ is continuous. □

Exercise 17.3.0.5. Cut Distance:

Let X be a compact metric space and $\mu : \mathcal{B}(X) \rightarrow [0, \infty]$ a Radon measure. Define the (right) group action $\phi : L^1(\mu) \times \text{Aut}(X, \mu) \rightarrow L^1(\mu)$ by

$$f \cdot \sigma = f \circ \sigma$$

Then

1. ϕ is an isometric group action
2. $\text{Aut}(X, d, \mu)$ is compact
3. for each $f \in L^1(\mu)$, the map $\sigma \mapsto f \cdot \sigma$ is continuous.
4. $\bar{d}$ is a metric on $L^1(\mu) / \text{Aut}(X, d, \mu)$

Proof. Clear by the preceeding exercises. □

Note 17.3.0.6. The preceeding distance is not quite the Cut distance, as the Cut norm only considers a subset of measurable sets for a function of two variables, but with some work, maybe I can show it is a distance.

17.4 Optimization over Compact Groups

Definition 17.4.0.1. We define $W \cdot U := WU$ and $W \triangleleft \Sigma := W^* \Sigma W$. Define $S(n) := \{\Sigma \in \mathbb{C}^{n \times n} : \Sigma^* = \Sigma\}$. Define $U(n, k) := \{U \in \mathbb{C}^{n \times k} : U^* U = I\}$.

Exercise 17.4.0.2. Let $n, k \in \mathbb{N}$. Suppose that $k \leq n$. Define $f : U(n, k) \times S(n) \rightarrow \mathbb{R}$ and $F : S(n) \rightarrow \mathbb{R}$ by $f(U, \Sigma) := \text{tr}(UU^* \Sigma)$ and $F(\Sigma) := \sup_{U \in U(n, k)} f(U, \Sigma)$. Let $\Sigma \in S(n)$. Let $\Lambda \in S(n)$. Suppose that there exists $V \in U(n)$ and $\Lambda \in S(n)$ such that Λ is diagonal and $\Sigma = V \Lambda V^*$. Then

1. for each $W \in U(n)$ and $U \in U(n, k)$, $f(W \cdot U, \Sigma) = f(U, W \triangleleft \Sigma)$,
2. $F(\Sigma) = F(\Lambda)$,
3. Suppose that $\Lambda = \text{diag}(\lambda_1, \dots, \lambda_n)$ and $\lambda_1 \geq \dots \geq \lambda_n \geq 0$. Then $\arg \min_{U \in U(n, k)} f(U, \Sigma) = V I_{n, k}$.

Proof.

1. Let $W \in U(n)$ and $U \in U(n, k)$. Then

$$\begin{aligned} f(W \cdot U, \Sigma) &= \text{tr}(WU U^* W^* \Sigma) \\ &= \text{tr}(U^* W^* \Sigma W U) \\ &= \text{tr}(U^* (W \triangleleft \Sigma) U) \\ &= \text{tr}(U^* U (W \triangleleft \Sigma)) \\ &= f(U, W \triangleleft \Sigma). \end{aligned}$$

2. The previous part implies that for each $W \in U(n)$ and $U \in U(n, k)$,

$$\begin{aligned} f(U, \Sigma) &= f(U, V^* \triangleleft \Lambda) \\ &= f(V^* \cdot U, \Lambda). \end{aligned}$$

Therefore

$$\begin{aligned} F(\Sigma) &= \sup_{U \in U(n, k)} f(U, \Sigma) \\ &= \sup_{U \in U(n, k)} f(V^* \cdot U, \Lambda) \\ &= \sup_{U \in U(n, k)} f(U, \Lambda) \\ &= F(\Lambda). \end{aligned}$$

3. Let $U \in U(n, k)$. For each $j \in [k]$, we define $u_j := U e_j$. We know that

$$\begin{aligned} \text{tr}(UU^* \Lambda) &= \text{tr}(U^* \Lambda U) \\ &= \sum_{j=1}^k \langle u_j, \Lambda u_j \rangle \\ &= \sum_{j=1}^k \lambda_j \langle u_j, u_j \rangle \\ &= \sum_{j=1}^k \lambda_j \|u_j\|^2 \\ &= \sum_{j=1}^k \lambda_j. \end{aligned}$$

Therefore

$$\begin{aligned}
 F(\Sigma) &= F(\Lambda) \\
 &= \sup_{U \in U(n,k)} \operatorname{tr}(U^* U \Lambda) \\
 &= \sup_{U \in U(n,k)} \sum_{j=1}^k \lambda_j \\
 &= \sum_{j=1}^k \lambda_j.
 \end{aligned}$$

Since

$$\begin{aligned}
 f(VI_{n,k}, \Sigma) &= \operatorname{tr}(VI_{n,k}(VI_{n,k})^* \Sigma) \\
 &= \operatorname{tr}((VI_{n,k})^* \Sigma VI_{n,k}) \\
 &= \operatorname{tr}(I_{k,n} V^* \Sigma VI_{n,k}) \\
 &= \operatorname{tr}(I_{k,n} V^* V \Lambda V^* VI_{n,k}) \\
 &= \operatorname{tr}(I_{k,n} \Lambda I_{n,k}) \\
 &= \sum_{j=1}^k \lambda_j,
 \end{aligned}$$

we have that $\arg \min_{U \in U(n,k)} f(U, \Sigma) = VI_{n,k}$.

□

Summation

discuss summation of vector valued functions, related summation to integration and the lebesgue integral and bochner integral

.1 Summation of Positive Functions

Definition .1.0.1. Let X be a set, $A \subset X$ and $f : X \rightarrow \overline{\mathbb{R}}_+$. We define **the sum of f over A** , denoted $\sum_{x \in A} f(x) \in \overline{\mathbb{R}}_+$, by

$$\sum_{x \in A} f(x) := \sup \left\{ \sum_{x \in F} f(x) : F \subset A \text{ and } \#F < \infty \right\}$$

Exercise .1.0.2. Let X be a set, $A, B \subset X$ and $f : X \rightarrow \overline{\mathbb{R}}_+$. If $A \subset B$, then $\sum_{x \in A} f(x) \leq \sum_{x \in B} f(x)$.

Proof. Suppose that $A \subset B$. Define $\mathcal{S}_A, \mathcal{S}_B \subset \overline{\mathbb{R}}_+$ by $\mathcal{S}_A := \left\{ \sum_{x \in F} f(x) : F \subset A \text{ and } \#F < \infty \right\}$ and $\mathcal{S}_B := \left\{ \sum_{x \in F} f(x) : F \subset B \text{ and } \#F < \infty \right\}$. Let $s \in \mathcal{S}_A$. Then there exists $F \subset A$ such that $\#F < \infty$ and $s = \sum_{x \in F} f(x)$. Since

$$\begin{aligned} F &\subset A \\ &\subset B, \end{aligned}$$

we have that $s \in \mathcal{S}_B$. Since $s \in \mathcal{S}_A$ is arbitrary, we have that for each $s \in \mathcal{S}_A$, $s \in \mathcal{S}_B$. Thus $\mathcal{S}_A \subset \mathcal{S}_B$. Hence

$$\begin{aligned} \sum_{x \in A} f(x) &= \sup \mathcal{S}_A \\ &\leq \sup \mathcal{S}_B \\ &= \sum_{x \in B} f(x). \end{aligned}$$

□

Exercise .1.0.3. Let X be a set, $(A_j)_{j \in \mathbb{N}} \subset X$ and $f : X \rightarrow \overline{\mathbb{R}}_+$. If $(A_j)_{j \in \mathbb{N}}$ are disjoint, then

$$\sum_{x \in \bigcup_{j \in \mathbb{N}} A_j} f(x)$$

Exercise .1.0.4. Change of Variable

Let X, Y be sets, $f : Y \rightarrow \mathbb{R}_+$ and $\sigma : X \rightarrow Y$. Then for each $B \subset Y$,

$$\sum_{x \in \sigma^{-1}(B)} [f \circ \sigma](x) = \sum_{y \in B} \# \sigma^{-1}(\{y\}) f(y)$$

Proof. Let $B \subset Y$. Then for each $F \subset B$, $\#F < \infty$ implies that

$$\begin{aligned}
 \sum_{x \in \sigma^{-1}(F)} [f \circ \sigma](x) &= \sum_{x \in \bigcup_{y \in F} \sigma^{-1}(\{y\})} [f \circ \sigma](x) \\
 &= \sum_{y \in F} \left[\sum_{x \in \sigma^{-1}(\{y\})} [f \circ \sigma](x) \right] \quad (\text{by (make prev ex) about sum over disjoint union is sum of sums}) \\
 &= \sum_{y \in F} \left[\sum_{x \in \sigma^{-1}(\{y\})} f(\sigma(x)) \right] \\
 &= \sum_{y \in F} \left[\sum_{x \in \sigma^{-1}(\{y\})} f(y) \right] \\
 &= \sum_{y \in F} \# \sigma^{-1}(\{y\}) f(y).
 \end{aligned}$$

FINISH!!! □

Exercise .1.0.5. Let X be a set, $A \subset X$ and $f : X \rightarrow \mathbb{R}_+$. Then for each $\sigma \in \text{Sym}(X)$,

$$\sum_{x \in X} [f \circ \sigma](x) = \sum_{x \in X} f(x)$$

Proof. Let $\sigma \in \text{Sym}(X)$. Then $X = \sigma(X)$ and for each $y \in X$, $\# \sigma^{-1}(\{y\}) = 1$. The prev ex then implies that

$$\begin{aligned}
 \sum_{x \in X} [f \circ \sigma](x) &= \sum_{x \in \sigma^{-1}(X)} [f \circ \sigma](x) \\
 &= \sum_{y \in X} \# \sigma^{-1}(\{y\}) f(y) \\
 &= \sum_{y \in X} f(y).
 \end{aligned}$$
□

Exercise .1.0.6. This definition coincides with the usual notion of summation when X is countable.

Proof. □

.2 Summation of Complex-Valued Functions

For $f : X \rightarrow \mathbb{C}$, we can write $f = g + ih$ where $g, h : X \rightarrow \mathbb{R}$. If

$$\sum_{x \in X} |f(x)| < \infty,$$

then the same is true for g^+, g^-, h^+, h^- . In this case, we may define

$$\sum_{x \in X} f(x)$$

in the obvious way.

The following note justifies the notation $\sum_{x \in X} f(x)$ where $f : X \rightarrow \mathbb{C}$.

Note .2.0.1. Let $f : X \rightarrow \mathbb{C}$ and $\alpha : X \rightarrow X$ a bijection. If $\sum_{x \in X} |f(x)| < \infty$, then $\sum_{x \in X} f(\alpha(x)) = \sum_{x \in X} f(x)$.

Asymptotic Notation

Definition .2.0.2. Let $X \in \text{Obj}(\mathbf{Top})$ and $x_0 \in X$.

- We define $\mathcal{F}((X, x_0), \mathbf{Norm}) := \{f \in Y^U : Y \in \text{Obj}(\mathbf{Norm}) \text{ and } U \in \mathcal{N}(x_0)\}$. (can we view this as a coslice category?)
- Let $Y \in \text{Obj}(\mathbf{Norm})$. We define $\mathcal{F}((X, x_0), Y) := \{f \in Y^U : U \in \mathcal{N}(x_0)\}$.

(change notation and find out what this object is and what its used for, its like a coslice category, need to learn more category theory and ask people)

Note .2.0.3. When the context is clear, we write

- $\mathcal{F}(x_0)$ in place of $\mathcal{F}((X, x_0), \mathbf{Norm})$
- $\mathcal{F}(x_0, Y)$ in place of $\mathcal{F}((X, x_0), Y)$

(change notation)

Definition .2.0.4. Let $X \in \text{Obj}(\mathbf{Top})$, $x_0 \in X$ and $f, g \in \mathcal{F}(x_0)$. Write $f : U_f \rightarrow Y_f$ and $g : U_g \rightarrow Y_g$. Then f is said to be **little-o of g near x_0** , denoted $f = o(g)$ as $x \rightarrow x_0$, if for each $\epsilon > 0$, there exists $U \in \mathcal{N}(x_0)$ such that $U \subset U_f \cap U_g$ and $\|f|_U\|_{Y_f} \leq \epsilon \|g|_U\|_{Y_g}$.

Exercise .2.0.5. Let $X \in \text{Obj}(\mathbf{Top})$, $x_0 \in X$ and $f, g \in \mathcal{F}(x_0, \mathbf{Norm})$. Write $f : U_f \rightarrow Y_f$ and $g : U_g \rightarrow Y_g$. If there exists $U \in \mathcal{N}(x_0)$ such that for each $x \in U \setminus \{x_0\}$, $\|g(x)\|_{Y_g} > 0$, then

$$f = o(g) \text{ as } x \rightarrow x_0 \text{ iff } \lim_{\substack{x \rightarrow x_0 \\ x \neq x_0}} \frac{\|f(x)\|_{Y_f}}{\|g(x)\|_{Y_g}} = 0.$$

Proof. Suppose that there exists $U \in \mathcal{N}(x_0)$ such that for each $x \in U \setminus \{x_0\}$, $\|g(x)\| > 0$. Define $V \subset X$ by $V := U \cap U_f \cap U_g$. Then $V \in \mathcal{N}(x_0)$, $V \subset U_f \cap U_g$ and for each $x \in V \setminus \{x_0\}$, $g(x) > 0$. We note that $x_0 \in \text{cl } V$. Define $h : V \setminus \{x_0\} \rightarrow \mathbb{R}$ by $h(x) := \|f(x)\|_{Y_f} / \|g(x)\|_{Y_g}$.

- ($\implies$):

Suppose that $f = o(g)$ as $x \rightarrow x_0$. Let $\epsilon > 0$. Then there exists $U_0 \in \mathcal{N}(x_0)$ such that $U_0 \subset U_f \cap U_g$ and $\|f|_{U_0}\|_{Y_f} \leq 2^{-1}\epsilon \|g|_{U_0}\|_{Y_g}$. Define $V_0 \subset X$ by $V_0 := V \cap U_0$. Then $V_0 \in \mathcal{N}(x_0, \mathcal{T}_X \cap V)$, and for each $x \in V_0 \setminus \{x_0\}$, we have that $\|g(x)\|_{Y_g} > 0$ and $\|f(x)\|_{Y_f} \leq 2^{-1}\epsilon \|g(x)\|_{Y_g}$. Then for each $x \in V_0 \setminus \{x_0\}$,

$$\begin{aligned} \left| h(x) - 0 \right| &= \frac{\|f(x)\|}{\|g(x)\|} \\ &\leq \frac{\epsilon}{2} \\ &< \epsilon. \end{aligned}$$

Since $\epsilon > 0$ is arbitrary, we have that for each $\epsilon > 0$, there exists $V_0 \in \mathcal{N}(x_0, \mathcal{T}_X \cap V)$ such that for each $x \in V_0 \setminus \{x_0\}$, $|h(x) - 0| < \epsilon$. Thus $\lim_{\substack{x \rightarrow x_0 \\ x \neq x_0}} h = 0$.

In Topology section, should probably define $\lim_{x \rightarrow x_0} f$ to be defined for $f : U \setminus \{x_0\} \rightarrow Y$ where $U \in \mathcal{N}(x_0)$, i.e. $\lim_{x \rightarrow x_0}$ is an operator on $\{f : U \setminus \{x_0\} \rightarrow Y : U \in \mathcal{N}(x_0)\}$, then for $g : X \rightarrow Y$, define $\lim_{x \rightarrow x_0} g := \lim_{x \rightarrow x_0} g|_{U \setminus \{x_0\}}$ for some $U \in \mathcal{N}(x_0, \mathcal{T}_X)$ and show that this definition is independent of U , so we should be thinking of equivalence classes of functions, like germs, except only on punctured neighborhood, need to explicitly state what $\lim_{x \rightarrow x_0} f(x)$ means when $f(x)$ is not defined outside a punctured neighborhood so that things like $\lim_{t \rightarrow 3} \frac{2t}{(3-t)(t-2)}$ make sense even though it doesn't make sense as a function on $\mathbb{R}$

• ($\Leftarrow$):

might want to specifically use the notation $\lim_{\substack{x \rightarrow x_0 \\ x \neq x_0}}$ for traditional limits when we don't care about the value at x_0 , this will be in line with the wikipedia page on filters and limits [wikipeage on limits of functions in terms of filters](#)

FINISH!!!!

FINISH!!!!

FINISH!!!!

FINISH!!!!

□

Exercise .2.0.6. Let X, Y, Z be normed vector spaces, $U \in \mathcal{T}_X$, $f : U \rightarrow Y$, $g : U \rightarrow Z$ and $x_0 \in U$. For each $x \in U$, define f_x : If $f = o(g)$ as $x \rightarrow x_0$, then for each $x \in X$, $f(tx) = o(g(t))$ (**maybe delete**)

Proof. Suppose that $f(h) = o(\|h\|)$ as $h \rightarrow 0$. Let $h \in X$ and $\epsilon > 0$. Choose $\delta' > 0$ such that for each $h' \in B(0, \delta')$, $h' \in A$ and

$$\|f(h')\| \leq \frac{\epsilon}{\|h\| + 1} \|h'\|$$

Choose $\delta > 0$ such that for each $t \in B_{\mathbb{R}}(0, \delta)$, $th \in B_X(0, \delta')$. Let $t \in B(0, \delta)$. Then

$$\begin{aligned} \|f(th)\| &\leq \frac{\epsilon}{\|h\| + 1} |t| \|h\| \\ &< \epsilon |t| \end{aligned}$$

So $f(th) = o(t)$ as $t \rightarrow 0$. □

Exercise .2.0.7. Let X and Y be normed vector spaces, $A \subset X$ open and $f : A \rightarrow Y$. Suppose that $0 \in A$. If $f(h) = o(\|h\|)$ as $h \rightarrow 0$, then for each $h \in X$, $f(th) = o(t)$ as $t \rightarrow 0$.

Proof. Suppose that $f(h) = o(\|h\|)$ as $h \rightarrow 0$. Let $h \in X$ and $\epsilon > 0$. Choose $\delta' > 0$ such that for each $h' \in B(0, \delta')$, $h' \in A$ and

$$\|f(h')\| \leq \frac{\epsilon}{\|h\| + 1} \|h'\|$$

Choose $\delta > 0$ such that for each $t \in B_{\mathbb{R}}(0, \delta)$, $th \in B_X(0, \delta')$. Let $t \in B(0, \delta)$. Then

$$\begin{aligned} \|f(th)\| &\leq \frac{\epsilon}{\|h\| + 1} |t| \|h\| \\ &< \epsilon |t| \end{aligned}$$

So $f(th) = o(t)$ as $t \rightarrow 0$. □

Exercise .2.0.8. Let X be a topological space, Y, Z be normed vector spaces, $f : X \rightarrow Y$, $g : X \rightarrow Z$ and $x_0 \in X$. If $f = o(g)$ as $x \rightarrow x_0$, then $\|f\|$

Definition .2.0.9. Let X be a topological space, $x_0 \in X$ and $h \in \mathcal{F}(x_0)$. Then there exist $Y_h \in \text{Obj}(\mathbf{Norm})$ and $U_h \in \mathcal{N}(x_0)$ such that $h \in Y_h^{U_h}$.

- Let $Y \in \text{Obj}(\mathbf{Norm})$ and $f, g \in \mathcal{F}(x_0, Y)$. Then there exist $U_f, U_g \in \mathcal{N}(x_0)$ such that $f \in Y^{U_f}$ and $g \in Y^{U_g}$. Then f is said to be **$o(h)$ -equivalent to g near x_0** , denoted $f \sim_{x_0}^o(h)g$, if there exists $U \in \mathcal{N}(x_0)$ such that $U \subset U_f \cap U_g$ and $f|_U - g|_U = o(h)$ as $x \rightarrow x_0$.

•

Then f is said to be **little-o equivalent to g near x_0** , denoted $f \sim_{x_0}^o g$, if for each $\epsilon > 0$, there exists $U \in \mathcal{N}(x_0)$ such that $U \subset U_f \cap U_g$ and $\|f|_U\|_{Y_f} \leq \epsilon \|g|_U\|_{Y_g}$.

Exercise .2.0.10. Let X be a topological space and $x_0 \in X$. Then $\sim_{x_0}^o \in \text{Equiv}(\mathcal{F}(x_0, o))$.

Proof.

1. Let $f \in \mathcal{F}(x_0)$. Then there exist $Y \in \text{Obj}(\mathbf{Norm})$, $U \in \mathcal{N}(x_0)$ such that $f \in Y^U$. Let $\epsilon > 0$. Define $U_0 \in \mathcal{N}(x_0)$ by $U_0 := U$. Then $U_0 \subset U \cap U$ and

$$\|f|_{U_0}\| = \|f|_U\|_Y \leq \epsilon \|g|_U\|_Y.$$

2. Let $f, g \in \mathcal{F}(x_0)$. Then there exist $Y, Z \in \text{Obj}(\mathbf{Norm})$, $U_f, U_g \in \mathcal{N}(x_0)$ such that $f \in Y^{U_f}$ and $g \in Z^{U_g}$.
3. Let $f, g, h \in \mathcal{F}(x_0)$. Then there exist $Y_f, Y_g, Y_h \in \text{Obj}(\mathbf{Norm})$ and $U_f, U_g, U_h \in \mathcal{N}(x_0)$ such that $f \in Y_f^{U_f}$, $g \in Y_g^{U_g}$ and $h \in Y_h^{U_h}$. Suppose that

□

Definition .2.0.11. Let X be a topological space, Y, Z be normed vector spaces, $f : X \rightarrow Y$, $g : X \rightarrow Z$ and $x_0 \in X$. Then we write

$$f = O(g) \quad \text{as } x \rightarrow x_0$$

if there exists $U \in \mathcal{N}(x_0)$ and $M \geq 0$ such that for each $x \in U$,

$$\|f(x)\| \leq M \|g(x)\|$$

Categories

move to notation?

Definition .2.0.12. We define the category of topological measure spaces, denoted $\mathbf{TopMsr}_+$, by

- $\text{Obj}(\mathbf{TopMsr}_+) := \{(X, \mu) : X \in \text{Obj}(\mathbf{Top}) \text{ and } \mu \in M(X)\}$
- $\text{Hom}_{\mathbf{TopMsr}_+}((X, \mu), (Y, \nu)) := \text{Hom}_{\mathbf{Top}}(X, Y) \cap \text{Hom}_{\mathbf{Msr}_+}((X, \mathcal{B}(X), \mu), (Y, \mathcal{B}(Y), \nu))$

Rings

.3 Introduction

Definition .3.0.1. Let R be a ring and $I, J \subset R$ ideals. We define

- $I + J \subset R$ by $I + J := \{i + j : i \in I \text{ and } j \in J\}$,
- $IJ \subset R$ by $IJ := \{i_1j_1 + i_2j_2 : i_1, i_2 \in I \text{ and } j_1, j_2 \in J\}$.

Exercise .3.0.2. Let R be a ring and $I, J \subset R$ ideals. Then $I + J$ and IJ are ideals.

Proof. **FINISH!!!** □

.4 Quotient Rings

Exercise .4.0.1. Let R be a commutative unital ring and $I \subset R$ and ideal. Then I is maximal iff R/I is a field.

Proof. **A previous exercise** implies that R/I is a commutative unital ring.

- $(\implies)$:

Suppose that I is maximal. Let $k + I \in R/I$. Suppose that $k + I \neq 0 + I$. Then $k \notin I$. Define $J \subset R$ by $J := (k) + I$. **A previous exercise** implies that J is an ideal. By construction $I \subset J$ and $k \in J$. Hence $I \neq J$. Since I is maximal, $J = R$. Since $1 \in J$, there exists $r \in R$ and $i \in I$ such that $1 = rk + i$. Hence

$$\begin{aligned} 1 + I &= rk + i + I \\ &= rk + I \\ &= (r + I)(k + I). \end{aligned}$$

Thus $(k + I)$ is invertible. Since $k + I \in R/I$ such that $k + I \neq 0 + I$ is arbitrary, we have that for each $k + I \in R/I$, $k + I \neq 0 + I$ implies that $k + I$ is invertible. Thus R/I is a field.

- $(\impliedby)$:

Suppose that R/I is a field. Let $J \subset R$. Suppose that J is an ideal, $I \subset J$ and $I \neq J$. Then there exists $j \in J$ such that $j \notin I$. Thus $j + I \neq 0 + I$. Since R/I is a field, $j + I$ is invertible. Hence there exists $k \in R$ such that $jk + I = 1 + I$. Hence

$$\begin{aligned} 1 - jk &\in I \\ &\subset J. \end{aligned}$$

Since J is an ideal, $jk \in J$ and therefore

$$\begin{aligned} 1 &= (1 - jk) + jk \\ &\in J. \end{aligned}$$

Since $1 \in J$, $J = R$. Since $J \subset R$ with J an ideal and $I \subset J$ is arbitrary, we have that for each $J \subset R$, J an ideal and $I \subset J$ implies that $J = I$ or $J = R$. Hence I is maximal. □

Bibliography

- [1] Angshuman R Goswami, *Revisiting Fekete's Lemma, Subadditive and Periodic Sequences: AR Goswami*, Results in Mathematics **80** (2025), no. 5, 131.
- [2] Gerald B Folland, *Real analysis: modern techniques and their applications*, John Wiley & Sons, 1999.
- [3] Robert R Phelps, *Lectures on Choquet's theorem*, Springer, 2002.
- [4] Gunnar F Stefánsson, *Pettis integrability*, Transactions of the American Mathematical Society **330** (1992), no. 1, 401–418.
- [5] H.H. and Wolff Schaefer M.P.H., *Topological Vector Spaces*, Graduate Texts in Mathematics, Springer New York, 1999.
- [6] L. and Beckenstein Narici E., *Topological Vector Spaces*, Chapman & Hall/CRC Pure and Applied Mathematics, CRC Press, 2010.
- [7] Einar Hille, *Functional Analysis and Semi-Groups*, Dutt Press, 2007.
- [8] ———, *Methods in classical and functional analysis*, Addison Wesley Publishing, 1972.
- [9] E Zeidler, *Nonlinear Functional Analysis and its Applications I.(Fixed Point Theorems) 1986*, Springer-Verlag New York, Inc.
- [10] Nicholas and Tarieladze Vakhania Vazha and Chobanyan, *Probability distributions on Banach spaces*, Springer Science & Business Media, 2012.
- [11] David Simmons, *Conditional measures and conditional expectation; Rohlin's disintegration theorem*, Vol. 32, Discrete and Continuous Dynamical Systems, 2012.
- [12] Edwin and Savage Hewitt Leonard J, *Symmetric measures on Cartesian products*, Transactions of the American Mathematical Society **80** (1955), no. 2, 470–501.