

Introduction to Category Theory

Carson James

Contents

Notation	vii
Preface	1
1 Preliminary Material	3
1.1 First Order Logic	3
1.2 von Neumann–Bernays–Gödel (NBG) Set Theory	3
1.2.1 TO DO	4
1.2.2 Relations	4
1.2.3 Functions	5
1.2.4 Digraphs	5
I Fundamental Concepts	7
2 Basic Concepts	9
3 Categories	11
3.1 Introduction	11
3.2 Fundamental Examples	13
3.3 Opposite Categories	14
4 Functors	17
4.1 Introduction	17
4.2 Fundamental Examples	18
4.3 Category of Small Categories	21
4.4 Comma Categories	22
5 Natural Transformations	27
5.1 Introduction	27
5.2 Fundamental Examples	31
5.3 Category of Functors	33
5.4 Diagonal Functor	34
6 Isomorphisms	37
6.1 Introduction	37
6.2 Isomorphisms between Categories	39
6.3 Natural Isomorphisms	40
II Fundamental Structures	43
7 Introduction	45
7.1 Initial and Terminal Objects	45

7.2	Products and Coproducts	50
7.3	Equalizers and Coequalizers	57
7.4	Pullbacks and Pushouts	60
7.5	Exponentials	61
7.6	Monomorphisms and Epimorphisms	63
7.6.1	Introduction	63
7.6.2	Extremal Monomorphisms and Epimorphisms	67
7.6.3	Regular Monomorphisms and Epimorphisms	68
7.7	Subobjects and Quotient Objects	74
8	Constructions on Categories	75
8.1	Subcategories	75
8.2	Product and Coproduct Categories	78
8.2.1	Introduction	78
8.2.2	Slicing	84
8.3	Functor Categories	88
8.4	Opposition	93
8.5	Currying	97
8.6	Equalizer and Coequalizer Categories	102
9	Limits	103
9.1	Cones and Cocones	103
9.2	Limits and Colimits	109
9.2.1	Universal Morphisms	109
9.2.2	Limits and Colimits	110
9.3	Fundamental Examples	110
9.3.1	Products and Coproducts	110
9.3.2	Equalizers and Coequalizers	111
9.3.3	Projective and Inductive Limits	111
9.4	Completeness and Cocompleteness	112
III	Basic Results	117
10	Yoneda Lemma	119
10.1	Representable Functors	119
10.2	Fundamental Examples	123
11	Adjunctions	125
12	Monads	127
12.1	Introduction	127
12.2	TO DO	127
13	Monoidal Categories	129
14	To Do	131
A	App	133
A.1	Reading Diagrams and associated digraphs of diagrams	133

Notation

$\mathcal{M}_+(X, \mathcal{A})$	finite measures on $(X, \mathcal{A})$
v	velocity

Preface

cc-by-nc-sa

Chapter 1

Preliminary Material

1.1 First Order Logic

maybe reorganize and move to logic notes?

1.2 von Neumann–Bernays–Gödel (NBG) Set Theory

maybe just use iterated extensions to not worry about if stuff is well defined or small, etc, see [iterated NBG math overflow](#)

Definition 1.2.0.1. Let x be a class. Then x is said to be a set iff there exists a class A such that $x \in A$.

Definition 1.2.0.2. Let x and y be classes. Then x is said to be a **subclass** of y , denoted $x \subset y$, if for each set a , $a \in x$ implies that $a \in y$.

Definition 1.2.0.3. Let x and y be classes. Then x is said to be **equal** to y if $x \subset y$ and $y \subset x$.

Axiom 1.2.0.4. Axiom of Extensionality:

Let x and y be classes. If for each set a , $a \in x$ iff $a \in y$, then $x = y$.

Axiom 1.2.0.5. Axiom of Pairing:

Let a, b be sets. Then there exists a set p such that for each for each set x , $x \in p$ iff $x = a$ or $x = b$.

Definition 1.2.0.6. product of two classes

Definition 1.2.0.7. Let A, B be classes and $R \subset A \times B$. elation from A to B .

Note 1.2.0.8. We can define cartesian products, relations, and functions for classes just like for sets.

Exercise 1.2.0.9. Let a, b be sets. Then there exists a unique set p such that for each for each set x , $x \in p$ iff $x = a$ or $x = b$.

Proof. By Axiom 1.2.0.5 implies that there exists a set p such that for each for each set x , $x \in p$ iff $x = a$ or $x = b$. Let q be a set. Suppose that for each for each set x , $x \in q$ iff $x = a$ or $x = b$. Then $\square$

Definition 1.2.0.10. Let x and y be sets. We define $(x, y) = \{\}$, denoted

Axiom 1.2.0.11. Axiom of Replacement:

Let A, B be classes and $f : A \rightarrow B$. If A is a set, then $f(A)$ is a set.

Axiom 1.2.0.12. Schema of Specification:

Let ϕ a propositional function on sets. Then there exists a class A such that for each set x , $x \in A$ iff $\phi(x)$.

Exercise 1.2.0.13. There exists a class A such that for each class x , $x \in A$ iff x is a set.

Proof. Define ϕ by

$$\phi(x) : x = x$$

Axiom 1.2.0.12 implies that there exists a class A such that for each set x , $x \in A$ iff $x = x$. Let x be a class. If $x \in A$, then by definition, x is a set.

Conversely, if x is a set, then by construction, $x \in A$. □

Exercise 1.2.0.14. There exists a class A such that for each class G and $*$: $G \times G \rightarrow G$, $(G, *) \in A$ iff $(G, *)$ is a group.

Proof. Define ϕ_1 , ϕ_2 and ϕ_3 by

- $\phi_1(G, *) : * : G \times G \rightarrow G$ is associative
- $\phi_2(G, *) : \text{there exists } e \in G \text{ such that for each } g \in G, e * g = g * e = g$
- $\phi_3(G, *) : \text{for each } g \in G \text{ there exists } h \in G \text{ such that } g * h = h * g = e$

Define ϕ by

$$\phi(G, *) : \phi_1(G, *) \text{ and } \phi_2(G, *) \text{ and } \phi_3(G, *)$$

Then there exists a class A such that for each set G and $*$: $G \times G \rightarrow G$, $(G, *) \in A$ iff $\phi(G, *)$ “is a group”. Therefore, for each group $(G, *)$, $(G, *) \in A$. **FINISH!!!** □

1.2.1 TO DO

1. [cover existence of subclasses, products of classes to be able to define class relations and subsequently class functions](#)
2. need to merge this with set theory notes and into to math notes, can first have naive set theory and then the NBG set theory or something

1.2.2 Relations

can I define relations on classes instead of just sets? do i need to move this stuff before to prelim material since we need class functions which I guess are class relations?

apparently yes [class, set theory notes \(link, but already downloaded\)](#)

Definition 1.2.2.1. Let A, B be classes and $R \subset A \times B$. Then R is said to be a **relation between A and B** . [write out 1st order logic definition like in notes class, set theory notes \(link, but already downloaded\)](#)

Definition 1.2.2.2. Let A, B, C be classes, R a relation between A and B , S a relation between B and C . We define the **composition relation of R with S between A and C** , denoted $R \circ S$, by $R \circ S := \{(a, c) \in A \times C : \text{there exists } b \in B \text{ such that } (a, b) \in R \text{ and } (b, c) \in S\}$.

Exercise 1.2.2.3. [show composition of class relations exists](#)

Exercise 1.2.2.4. Let A, B, C, D be classes, R a relation between A and B , S relation between B and C and T relation between C and D . Then $(T \circ S) \circ R = T \circ (S \circ R)$.

Proof.

- Let $(a, d) \in (T \circ S) \circ R$. Then there exists $b \in B$ such that $(a, b) \in R$ and $(b, d) \in T \circ S$. Since $(b, d) \in T \circ S$, there exists $c \in C$ such that $(b, c) \in S$ and $(c, d) \in T$. Since $(a, b) \in R$ and $(b, c) \in S$, $(a, c) \in S \circ R$. Since $(a, c) \in S \circ R$ and $(c, d) \in T$, we have that $(a, d) \in T \circ (S \circ R)$. Since $(a, d) \in (T \circ S) \circ R$ is arbitrary, we have that for each $(a, d) \in (T \circ S) \circ R$, $(a, d) \in T \circ (S \circ R)$. Hence $(T \circ S) \circ R \subset T \circ (S \circ R)$.
- Similarly, $T \circ (S \circ R) \subset (T \circ S) \circ R$.

Since $(T \circ S) \circ R \subset T \circ (S \circ R)$ and $T \circ (S \circ R) \subset (T \circ S) \circ R$, we have that $(T \circ S) \circ R = T \circ (S \circ R)$. □

[review, input any necessary axioms for classes](#)

Definition 1.2.2.5. Let A be a class. We define the **diagonal relation on A** , denoted Δ_A , by $\Delta_A := \{(a, a) : a \in A\}$. [make and show exists in a previous ex](#)

Exercise 1.2.2.6. Let A, B be classes and R a relation between A and B . Then

1. $R \circ \Delta_A = R$
2. $\Delta_B \circ R = R$

Proof.

1.
 - Let $(a, b) \in R \circ \Delta_A$. Then there exists $a' \in A$ such that $(a, a') \in \Delta_A$ and $(a', b) \in R$. By definition of Δ_A , $a = a'$. Therefore $(a, b) \in R$. Since $(a, b) \in R \circ \Delta_A$ is arbitrary, we have that for each $(a, b) \in R \circ \Delta_A$, $(a, b) \in R$. Hence $R \circ \Delta_A \subset R$.
 - Let $(a, b) \in R$. Then $a \in A$. Therefore $(a, a) \in \Delta_A$. Since $(a, a) \in \Delta_A$ and $(a, b) \in R$, we have that $(a, b) \in R \circ \Delta_A$. Since $(a, b) \in R$ is arbitrary, we have that for each $(a, b) \in R$, $(a, b) \in R \circ \Delta_A$. Hence $R \circ \Delta_A \subset R$.

Since $R \circ \Delta_A \subset R$ and $R \circ \Delta_A \subset R$, we have that $R \circ \Delta_A = R$.

2. Similar to (1).

□

1.2.3 Functions

1.2.4 Digraphs

Definition 1.2.4.1. talk about what it means for two diagrams to commute, talk about parallel morphisms being the only exception

Definition 1.2.4.2.

- We say a digraph (G, E) **commutes** if for each $x, y \in G$ and paths p, q , if p, q start at x and end at y , then $p = q$ **need to fix this definition**

Note 1.2.4.3. We make the following exception about parallel arrows. if two ore more edges are equal, we draw them as curved arrows, not parallel.

We define parallel arrows **FINISH!!!**

Part I

Fundamental Concepts

Chapter 2

Basic Concepts

Chapter 3

Categories

3.1 Introduction

Definition 3.1.0.1. Let $\mathcal{C}_0, \mathcal{C}_1$ be classes and $\text{dom}, \text{cod} : \mathcal{C}_1 \rightarrow \mathcal{C}_0$. Define $\mathcal{C}_1^{(2)} \subset \mathcal{C}_1 \times \mathcal{C}_1$ by $\mathcal{C}_1^{(2)} := \{(g, f) \in \mathcal{C}_1 \times \mathcal{C}_1 : \text{dom}(g) = \text{cod}(f)\}$. Let $\circ : \mathcal{C}_1^{(2)} \rightarrow \mathcal{C}_1$. Then $(\mathcal{C}_0, \mathcal{C}_1, \text{dom}, \text{cod}, \circ)$ is said to be a **category** if

1. **(Composition):**

for each $(g, f) \in \mathcal{C}_1^{(2)}$, $\text{dom}(g \circ f) = \text{dom}(f)$ and $\text{cod}(g \circ f) = \text{cod}(g)$

2. **(Associativity):**

for each $f, g, h \in \mathcal{C}_1$, if $(g, f), (h, g) \in \mathcal{C}_1^{(2)}$, then

$$(h \circ g) \circ f = h \circ (g \circ f)$$

3. **(Identity):**

for each $X \in \mathcal{C}_0$, there exists $e \in \mathcal{C}_1$ such that $\text{dom}(e) = X$, $\text{cod}(e) = X$ and for each $f, g \in \mathcal{C}_1$, if $\text{dom}(f) = X$ and $\text{cod}(g) = X$, then

$$f \circ e = f \text{ and } e \circ g = g$$

Definition 3.1.0.2. Let $\mathcal{C}$ be a category. Write $\mathcal{C} = (\mathcal{C}_0, \mathcal{C}_1, \text{dom}, \text{cod}, \circ)$.

• We define the

- **objects of $\mathcal{C}$** , denoted $\text{Obj}(\mathcal{C})$, by $\text{Obj}(\mathcal{C}) := \mathcal{C}_0$
- **morphisms of $\mathcal{C}$** , denoted Hom , by $\text{Hom}_{\mathcal{C}} := \mathcal{C}_1$
- **domain function of $\mathcal{C}$** , denoted $\text{dom}_{\mathcal{C}}$, by $\text{dom}_{\mathcal{C}} := \text{dom}$
- **codomain function of $\mathcal{C}$** , denoted $\text{cod}_{\mathcal{C}}$, by $\text{cod}_{\mathcal{C}} := \text{cod}$
- **composition operator of $\mathcal{C}$** , denoted $\circ_{\mathcal{C}}$, by $\circ_{\mathcal{C}} := \circ$

- For $X, Y \in \text{Obj}(\mathcal{C})$, we define the **Morphisms of $\mathcal{C}$ from X to Y** , denoted $\text{Hom}_{\mathcal{C}}(X, Y)$, by $\text{Hom}_{\mathcal{C}}(X, Y) := \{f \in \text{Hom}(\mathcal{C}) : \text{dom}_{\mathcal{C}}(f) = X \text{ and } \text{cod}_{\mathcal{C}}(f) = Y\}$.

Note 3.1.0.3. When the context is clear, we write dom , cod and $g \circ f$ in place of $\text{dom}_{\mathcal{C}}$, $\text{cod}_{\mathcal{C}}$ and $g \circ_{\mathcal{C}} f$ respectively.

Note 3.1.0.4. We typically define a category $\mathcal{C}$ by specifying

- $\text{Obj}(\mathcal{C})$
- for $X, Y \in \text{Obj}(\mathcal{C})$, the class $\text{Hom}_{\mathcal{C}}(X, Y)$
- for $X, Y, Z \in \text{Obj}(\mathcal{C})$, $f \in \text{Hom}_{\mathcal{C}}(X, Y)$ and $g \in \text{Hom}_{\mathcal{C}}(Y, Z)$, the composite morphism $g \circ f \in \text{Hom}_{\mathcal{C}}(X, Z)$.

and then show

- well-definedness of composition
- associativity of composition
- existence of identities

Exercise 3.1.0.5. Uniqueness of Identities:

Let $\mathcal{C}$ be a category. Then for each $A \in \text{Obj}(\mathcal{C})$, there exists a unique $e \in \text{End}_{\mathcal{C}}(A)$ such that for each $B \in \text{Obj}(\mathcal{C})$, $f \in \text{Hom}_{\mathcal{C}}(A, B)$ and $g \in \text{Hom}_{\mathcal{C}}(B, A)$, $f \circ e = f$ and $e \circ g = g$.

Proof. Let $A \in \text{Obj}(\mathcal{C})$.

- **Existence:**

Since $\mathcal{C}$ is a category, by definition there exists $e \in \text{Hom}_{\mathcal{C}}(A, A)$ such that for each $B \in \text{Obj}(\mathcal{C})$, $f \in \text{Hom}_{\mathcal{C}}(A, B)$ and $g \in \text{Hom}_{\mathcal{C}}(B, A)$, $f \circ e = f$ and $e \circ g = g$.

- **Uniqueness:**

Let $e_1, e_2 \in \text{Hom}_{\mathcal{C}}(A, A)$. Suppose that for each $B \in \text{Obj}(\mathcal{C})$, $f \in \text{Hom}_{\mathcal{C}}(A, B)$ and $g \in \text{Hom}_{\mathcal{C}}(B, A)$, $f \circ e_1 = f$, $f \circ e_2 = f$, $e_1 \circ g = g$ and $e_2 \circ g = g$. Then

$$\begin{aligned} e_1 &= e_2 \circ e_1 \\ &= e_2. \end{aligned}$$

□

Definition 3.1.0.6. Let $\mathcal{C}$ be a category and $A \in \text{Obj}(\mathcal{C})$. We define the **identity morphism on A** , denoted $\text{id}_A^{\mathcal{C}} \in \text{Hom}_{\mathcal{C}}(A, A)$, to be the unique $e \in \text{Hom}_{\mathcal{C}}(A, A)$ such that for each $B \in \text{Obj}(\mathcal{C})$, $f \in \text{Hom}_{\mathcal{C}}(A, B)$ and $g \in \text{Hom}_{\mathcal{C}}(B, A)$, $f \circ e = f$ and $e \circ g = g$.

Note 3.1.0.7. When the context is clear, we write id_A in place of $\text{id}_A^{\mathcal{C}}$.

Definition 3.1.0.8. Let $\mathcal{C}$ be a category. Then $\mathcal{C}$ is said to be

- **small** if $\text{Obj}(\mathcal{C})$ and $\text{Hom}_{\mathcal{C}}$ are sets
- **locally small** if for each $A, B \in \text{Obj}(\mathcal{C})$, $\text{Hom}_{\mathcal{C}}(A, B)$ is a set

Exercise 3.1.0.9. Let $\mathcal{C}$ be a category. If $\mathcal{C}$ is small, then $\mathcal{C}$ is a set.

Proof. Suppose that $\mathcal{C}$ is small. Then $\text{Obj}(\mathcal{C})$ and $\text{Hom}_{\mathcal{C}}$ are sets. Then $\mathcal{P}(\text{Obj}(\mathcal{C}))$, $\mathcal{P}(\text{Hom}_{\mathcal{C}})$ and $\text{Obj}(\mathcal{C})^{\text{Hom}_{\mathcal{C}}}$ are sets. Hence $\text{Obj}(\mathcal{C}) \times \text{Hom}_{\mathcal{C}} \times \text{Obj}(\mathcal{C})^{\text{Hom}_{\mathcal{C}}} \times \text{Obj}(\mathcal{C})^{\text{Hom}_{\mathcal{C}}}$ is a set. By definition, $\mathcal{C} = (\text{Obj}(\mathcal{C}), \text{Hom}_{\mathcal{C}}, \text{dom}, \text{cod}) \in \text{Obj}(\mathcal{C}) \times \text{Hom}_{\mathcal{C}} \times \text{Obj}(\mathcal{C})^{\text{Hom}_{\mathcal{C}}} \times \text{Obj}(\mathcal{C})^{\text{Hom}_{\mathcal{C}}}$. By definition, $\mathcal{C}$ is a set. □

Exercise 3.1.0.10. There exists a class A such that $\mathcal{C} \in A$ iff $\mathcal{C}$ is a small category.

Proof. Exercise 3.1.0.9 implies that for each category $\mathcal{C}$, $\mathcal{C}$ is small implies that $\mathcal{C}$ is a set. Define ϕ by

$$\phi(\mathcal{C}) : \mathcal{C} \text{ is a small category}$$

Then Axiom 1.2.0.12 implies that there exists a class A such that $\mathcal{C} \in A$ iff $\mathcal{C}$ is a small category. □

prosets

Definition 3.1.0.11. Let $\mathcal{P}$ be a category. Then $\mathcal{P}$ is said to be a

1. **proset category** if for each $a, b \in \text{Obj}(\mathcal{P})$, $\# \text{Hom}_{\mathcal{P}}(a, b) \leq 1$
2. **poset category** if $\mathcal{P}$ is a proset category and for each $a, b \in \text{Obj}(\mathcal{P})$, $\# \text{Hom}_{\mathcal{P}}(a, b) = 1$ and $\# \text{Hom}_{\mathcal{P}}(b, a) = 1$ implies that $a = b$.

Definition 3.1.0.12. Let $\mathcal{P}$ be a proset. Set $P := \text{Obj}(\mathcal{P})$. We define the **less-than-or-equal-to relation** on P , denoted $\leq \subset P \times P$, by

$$\leq := \{(a, b) \in P \times P : \text{Hom}_{\mathcal{P}}(a, b) \neq \emptyset\}.$$

Definition 3.1.0.13. Let $\mathcal{C}$ be a category. Then $\mathcal{C}$ is said to be **discrete** if $\text{Hom}_{\mathcal{C}} = \{\text{id}_X : X \in \text{Obj}(\mathcal{C})\}$.

3.2 Fundamental Examples

Definition 3.2.0.1. We define the **empty category**, denoted $\mathbf{0}$, by

- $\text{Obj}(\mathbf{0}) = \emptyset$
- $\text{Hom}_{\mathbf{0}} = \emptyset$

Exercise 3.2.0.2. We have that $\mathbf{0}$ is a category.

Proof. Vacuously true. □

Definition 3.2.0.3. We define the **trivial category**, denoted $\mathbf{1}$, by

- $\text{Obj}(\mathbf{1}) = \{*\}$
- $\text{Hom}_{\mathbf{1}} = \{\text{id}_*\}$

Exercise 3.2.0.4. We have that $\mathbf{1}$ is a category.

Proof. Clear. □

Definition 3.2.0.5. We define **Set** by

- $\text{Obj}(\mathbf{Set}) = \{A : A \text{ is a set}\}$
- for each $A, B \in \text{Obj}(\mathbf{Set})$, $\text{Hom}_{\mathbf{Set}}(A, B) = \{f : f : A \rightarrow B\}$
- for $A, B, C \in \mathbf{Set}$, $f \in \text{Hom}_{\mathbf{Set}}(A, B)$ and $g \in \text{Hom}_{\mathbf{Set}}(B, C)$, $g \circ_{\mathbf{Set}} f = g \circ f$.

Exercise 3.2.0.6. We have that **Set** is a category.

Proof.

- **well-definedness of composition:**
- **associativity of composition:**
- **existence of identities:**

FINISH!!! □

Exercise 3.2.0.7. Let $A, B \in \text{Obj}(\mathbf{Set})$, show $\mathcal{R}(A, B)$ is a category, (category of relations)

Exercise 3.2.0.8. **preordered set is a category**

FIX!!!! need to decide what to do about relations, on the one hand, we can define relations as classes like in the previous section, on another hand, we can define the category of relations based on **Set**, on the third hand, looking at ‘Elementary categories, elementary toposes’ by McLarty who shows you can define relations in terms of subobjects, i.e. morphisms, does this more general idea also form a category, of which the category of relations based on set can be considered a subobject(i.e. a subcategory)?

Definition 3.2.0.9. We define the **category of relations**, denoted **Rel**, by

- $\text{Obj}(\mathbf{Rel}) := \text{Obj}(\mathbf{Set})$
- for each $A, B \in \text{Obj}(\mathbf{Rel})$, $\text{Hom}_{\mathbf{Rel}}(A, B) := \mathcal{P}(A \times B)$
- for each $A, B, C \in \text{Obj}(\mathbf{Rel})$, $R \in \text{Hom}_{\mathbf{Rel}}(A, B)$ and $S \in \text{Hom}_{\mathbf{Rel}}(B, C)$, we define $S \circ_{\mathbf{Rel}} R := \{(a, c) \in A \times C : \text{there exists } b \in B \text{ such that } (a, b) \in R \text{ and } (b, c) \in S\}$

Proof.

- **well-definedness of composition:**

By definition, for each $A, B, C \in \text{Obj}(\mathbf{Rel})$, $R \in \text{Hom}_{\mathbf{Rel}}(A, B)$ and $S \in \text{Hom}_{\mathbf{Rel}}(B, C)$, we have that

$$\begin{aligned} S \circ_{\mathbf{Rel}} R &\in \mathcal{P}(A \times C) \\ &= \text{Hom}_{\mathbf{Rel}}(A, C). \end{aligned}$$

So $\circ_{\mathbf{Rel}}$ is well-defined.

- **associativity of composition:**

ref ex in basic concepts chapter

- **existence of identities:**

ref ex in basic concepts chapter

FINISH!!!

□

maybe discuss a quiver as a category whos objects are vertices and morphisms are and composition is given by concatenation (i.e. I guess a quiver equipped with the path algebra), maybe this is not appropriate [quiver wikipedia](#)

Definition 3.2.0.10. We define the **category of quivers**, denoted $\mathbf{Quiv}$, by

- $\text{Obj}(\mathbf{Quiv})$
- for $\mathcal{Q}, \mathcal{L} \in \text{Obj}(\mathbf{Quiv})$,

$$\text{Hom}_{\mathbf{Quiv}}(\mathcal{Q}, \mathcal{L}) := \{\phi : \phi \text{ is a quiver}\}$$

3.3 Opposite Categories

Definition 3.3.0.1. Let $\mathcal{C}$ be a category, we define the dual of $\mathcal{C}$ or the **opposite of $\mathcal{C}$** , denoted $\mathcal{C}^{\text{op}}$, by

- $\text{Obj}(\mathcal{C}^{\text{op}}) := \text{Obj}(\mathcal{C})$
- for $X, Y \in \text{Obj}(\mathcal{C}^{\text{op}})$, $\text{Hom}_{\mathcal{C}^{\text{op}}}(X, Y) := \text{Hom}_{\mathcal{C}}(Y, X)$
- for $X, Y, Z \in \text{Obj}(\mathcal{C}^{\text{op}})$ and $f \in \text{Hom}_{\mathcal{C}^{\text{op}}}(X, Y)$, $g \in \text{Hom}_{\mathcal{C}^{\text{op}}}(Y, Z)$, $g \circ_{\mathcal{C}^{\text{op}}} f := f \circ_{\mathcal{C}} g$

Exercise 3.3.0.2. Let $\mathcal{C}$ be a category. Then $\mathcal{C}^{\text{op}}$ is a category.

Proof.

- for $W, X, Y, Z \in \text{Obj}(\mathcal{C})$, $f \in \text{Hom}_{\mathcal{C}^{\text{op}}}(W, X)$ and $g \in \text{Hom}_{\mathcal{C}^{\text{op}}}(X, Y)$ and $h \in \text{Hom}_{\mathcal{C}^{\text{op}}}(Y, Z)$. Then

$$\begin{aligned} (h \circ_{\mathcal{C}^{\text{op}}} g) \circ_{\mathcal{C}^{\text{op}}} f &= f \circ_{\mathcal{C}} (h \circ_{\mathcal{C}^{\text{op}}} g) \\ &= f \circ_{\mathcal{C}} (g \circ_{\mathcal{C}} h) \\ &= (f \circ_{\mathcal{C}} g) \circ_{\mathcal{C}} h \\ &= h \circ_{\mathcal{C}^{\text{op}}} (f \circ_{\mathcal{C}} g) \\ &= h \circ_{\mathcal{C}^{\text{op}}} (g \circ_{\mathcal{C}^{\text{op}}} f) \end{aligned}$$

So composition is associative.

- Let $X \in \text{Obj}(\mathcal{C})$ and $f, g \in \text{Hom}_{\mathcal{C}^{\text{op}}}$. Suppose that $\text{dom}(f) = X$ and $\text{cod}(g) = X$ Then

$$\begin{aligned} f \circ_{\mathcal{C}^{\text{op}}} \text{id}_X &= \text{id}_X \circ_{\mathcal{C}} f \\ &= f \end{aligned}$$

and

$$\begin{aligned} \text{id}_X \circ_{\mathcal{C}^{\text{op}}} g &= g \circ_{\mathcal{C}} \text{id}_X \\ &= g \end{aligned}$$

So $(\text{id}_X)_{\mathcal{C}^{\text{op}}} = (\text{id}_X)_{\mathcal{C}}$.

□

Exercise 3.3.0.3. Let $\mathcal{C}$ be a category. Then $(\mathcal{C}^{\text{op}})^{\text{op}} = \mathcal{C}$.

Proof. Set $\mathcal{D} := \mathcal{C}^{\text{op}}$.

- We have that

$$\begin{aligned} \text{Obj}((\mathcal{C}^{\text{op}})^{\text{op}}) &= \text{Obj}(\mathcal{D}^{\text{op}}) \\ &= \text{Obj}(\mathcal{D}), \quad (\text{by definition}) \\ &= \text{Obj}(\mathcal{C}^{\text{op}}) \\ &= \text{Obj}(\mathcal{C}), \quad (\text{by definition}). \end{aligned}$$

- Let $X, Y \in \text{Obj}((\mathcal{C}^{\text{op}})^{\text{op}})$. Then

$$\begin{aligned} \text{Hom}_{(\mathcal{C}^{\text{op}})^{\text{op}}}(X, Y) &= \text{Hom}_{\mathcal{D}^{\text{op}}}(X, Y) \\ &= \text{Hom}_{\mathcal{D}}(Y, X), \quad (\text{by definition}) \\ &= \text{Hom}_{\mathcal{C}^{\text{op}}}(Y, X) \\ &= \text{Hom}_{\mathcal{C}}(X, Y), \quad (\text{by definition}). \end{aligned}$$

□

Exercise 3.3.0.4. Let $\mathcal{P}$ be a category. Then

1. $\mathcal{P}$ is a proset iff $\mathcal{P}^{\text{op}}$ is a proset
2. $\mathcal{P}$ is a poset iff $\mathcal{P}^{\text{op}}$ is a poset

Proof. **FINISH!!!**

□

Definition 3.3.0.5. Let $\mathcal{P}$ be a proset. Set $P := \text{Obj}(\mathcal{P})$. We define the **less-than-or-equal-to relation** on P , denoted $\leq \subset P \times P$, by

$$\leq := \{(a, b) \in P \times P : \text{Hom}_{\mathcal{P}^{\text{op}}}(a, b) \neq \emptyset\}.$$

Chapter 4

Functors

4.1 Introduction

Definition 4.1.0.1. Let $\mathcal{C}$ and $\mathcal{D}$ be categories and $F_0 : \text{Obj}(\mathcal{C}) \rightarrow \text{Obj}(\mathcal{D})$, $F_1 : \text{Hom}_{\mathcal{C}} \rightarrow \text{Hom}_{\mathcal{D}}$ class functions. Set $F = (F_0, F_1)$. Then F is said to be a $(\mathcal{C}, \mathcal{D})$ -functor, denoted $F : \mathcal{C} \rightarrow \mathcal{D}$, if

1. for each $A, B \in \text{Obj}(\mathcal{C})$ and $f \in \text{Hom}_{\mathcal{C}}(A, B)$, $F_1(f) \in \text{Hom}_{\mathcal{D}}(F_0(A), F_0(B))$
2. for each $A, B, C \in \text{Obj}(\mathcal{C})$, $f \in \text{Hom}_{\mathcal{C}}(A, B)$ and $g \in \text{Hom}_{\mathcal{C}}(B, C)$, $F_1(g \circ f) = F_1(g) \circ F_1(f)$
3. for each $A \in \text{Obj}(\mathcal{C})$, $F_1(\text{id}_A) = \text{id}_{F_0(A)}$

Note 4.1.0.2. When the context is clear, we may say “functor” instead of “ $(\mathcal{C}, \mathcal{D})$ -functor”.

Note 4.1.0.3. For $A \in \text{Obj}(\mathcal{C})$ and $f \in \text{Hom}_{\mathcal{C}}$, we typically write $F(A)$ and $F(f)$ instead of $F_0(A)$ and $F_1(f)$ respectively. **maybe discuss how this is just a quiver homomorphism**

Definition 4.1.0.4. Let $\mathcal{C}$, $\mathcal{D}$ and $\mathcal{E}$ be categories and $F : \mathcal{C} \rightarrow \mathcal{D}$, $G : \mathcal{D} \rightarrow \mathcal{E}$ functors. We define the **composition of G with F** , denoted $G \circ F : \mathcal{C} \rightarrow \mathcal{E}$, by

- $G \circ F(A) := G(F(A))$
- $G \circ F(f) := G(F(f))$

Note 4.1.0.5. Writing $G \circ F = (H_0, H_1)$, we have that $H_0 = G_0 \circ F_0$ and $H_1 = G_1 \circ F_1$.

Exercise 4.1.0.6. Let $\mathcal{C}$, $\mathcal{D}$ and $\mathcal{E}$ be categories and $F : \mathcal{C} \rightarrow \mathcal{D}$, $G : \mathcal{D} \rightarrow \mathcal{E}$ functors. Then $G \circ F : \mathcal{C} \rightarrow \mathcal{E}$ is a functor.

Proof.

1. Let $A, B \in \text{Obj}(\mathcal{C})$ and $f \in \text{Hom}_{\mathcal{C}}(A, B)$. Since $F(f) \in \text{Hom}_{\mathcal{D}}(F(A), F(B))$, we have that $G(F(f)) \in \text{Hom}_{\mathcal{E}}(G(F(A)), G(F(B)))$. Then

$$\begin{aligned} G \circ F(f) &= G(F(f)) \\ &\in \text{Hom}_{\mathcal{E}}(G(F(A)), G(F(B))) \\ &= \text{Hom}_{\mathcal{E}}(G \circ F(A), G \circ F(B)) \end{aligned}$$

2. Let $A, B, C \in \text{Obj}(\mathcal{C})$, $f \in \text{Hom}_{\mathcal{C}}(A, B)$ and $g \in \text{Hom}_{\mathcal{C}}(B, C)$. Then

$$\begin{aligned} G \circ F(g \circ f) &= G(F(g \circ f)) \\ &= G(F(g) \circ F(f)) \\ &= G(F(g)) \circ G(F(f)) \\ &= G \circ F(g) \circ G \circ F(f) \end{aligned}$$

3. Let $A \in \text{Obj}(\mathcal{C})$. Then

$$\begin{aligned} G \circ F(\text{id}_A) &= G(F(\text{id}_A)) \\ &= G(\text{id}_{F(A)}) \\ &= \text{id}_{G(F(A))} \\ &= \text{id}_{G \circ F(A)} \end{aligned}$$

So $G \circ F : \mathcal{C} \rightarrow \mathcal{E}$ is a functor. □

Exercise 4.1.0.7. Let $\mathcal{C}, \mathcal{D}, \mathcal{E}, \mathcal{F}$ be categories and $F : \mathcal{C} \rightarrow \mathcal{D}$, $G : \mathcal{D} \rightarrow \mathcal{E}$, $H : \mathcal{E} \rightarrow \mathcal{F}$ functors. Then $(H \circ G) \circ F = H \circ (G \circ F)$.

Proof. Let $A, B \in \text{Obj}(\mathcal{C})$ and $f \in \text{Hom}_{\mathcal{C}}(A, B)$. Then

•

$$\begin{aligned} (H \circ G) \circ F(A) &= H \circ G(F(A)) \\ &= H(G(F(A))) \\ &= H(G \circ F(A)) \\ &= H \circ (G \circ F)(A) \end{aligned}$$

•

$$\begin{aligned} (H \circ G) \circ F(f) &= H \circ G(F(f)) \\ &= H(G(F(f))) \\ &= H(G \circ F(f)) \\ &= H \circ (G \circ F)(f) \end{aligned}$$

Hence $(H \circ G) \circ F = H \circ (G \circ F)$. □

4.2 Fundamental Examples

discuss a quiver as a functor from the free quiver (a category) to **Set** or other category [quiver wikipedia](#)

Definition 4.2.0.1. Let $\mathcal{C}$ be a category. We define the **identity functor from $\mathcal{C}$ to $\mathcal{C}$** , denoted $\text{id}_{\mathcal{C}} : \mathcal{C} \rightarrow \mathcal{C}$, by

- $\text{id}_{\mathcal{C}}(A) := A, \quad (A \in \text{Obj}(\mathcal{C}))$
- $\text{id}_{\mathcal{C}}(f) := f, \quad (f \in \text{Hom}_{\mathcal{C}})$

Exercise 4.2.0.2. Let $\mathcal{C}$ be a category. Then $\text{id}_{\mathcal{C}} : \mathcal{C} \rightarrow \mathcal{C}$ is a functor.

Proof.

1. Let $A, B \in \text{Obj}(\mathcal{C})$ and $f \in \text{Hom}_{\mathcal{C}}(A, B)$. Then

$$\begin{aligned} \text{id}_{\mathcal{C}}(f) &= f \\ &\in \text{Hom}_{\mathcal{C}}(A, B) \\ &= \text{Hom}_{\mathcal{C}}(\text{id}_{\mathcal{C}}(A), \text{id}_{\mathcal{C}}(B)) \end{aligned}$$

2. Let $A, B, C \in \text{Obj}(\mathcal{C})$, $f \in \text{Hom}_{\mathcal{C}}(A, B)$ and $g \in \text{Hom}_{\mathcal{C}}(B, C)$. Then

$$\begin{aligned} \text{id}_{\mathcal{C}}(g \circ f) &= g \circ f \\ &= \text{id}_{\mathcal{C}}(g) \circ \text{id}_{\mathcal{C}}(f) \end{aligned}$$

3. Let $A \in \text{Obj}(\mathcal{C})$. Then

$$\begin{aligned}\text{id}_{\mathcal{C}}(\text{id}_A) &= \text{id}_A \\ &= \text{id}_{\text{id}_{\mathcal{C}}(A)}\end{aligned}$$

□

Exercise 4.2.0.3. Let $\mathcal{C}$ and $\mathcal{D}$ be categories and $F : \mathcal{C} \rightarrow \mathcal{D}$. Then

$$1. \text{id}_{\mathcal{D}} \circ F = F$$

$$2. F \circ \text{id}_{\mathcal{C}} = F$$

Proof.

1. Let $A, B \in \text{Obj}(\mathcal{C})$ and $f \in \text{Hom}_{\mathcal{C}}(A, B)$. Then

$$\begin{aligned}\text{id}_{\mathcal{D}} \circ F(A) &= \text{id}_{\mathcal{D}}(F(A)) \\ &= F(A)\end{aligned}$$

and

$$\begin{aligned}\text{id}_{\mathcal{D}} \circ F(f) &= \text{id}_{\mathcal{D}}(F(f)) \\ &= F(f)\end{aligned}$$

Since $A, B \in \text{Obj}(\mathcal{C})$ and $f \in \text{Hom}_{\mathcal{C}}(A, B)$ are arbitrary, $\text{id}_{\mathcal{D}} \circ F = F$.

2. Let $A, B \in \text{Obj}(\mathcal{C})$ and $f \in \text{Hom}_{\mathcal{C}}(A, B)$. Then

$$\begin{aligned}F \circ \text{id}_{\mathcal{C}}(A) &= F(\text{id}_{\mathcal{C}}(A)) \\ &= F(A)\end{aligned}$$

and

$$\begin{aligned}F \circ \text{id}_{\mathcal{C}}(f) &= F(\text{id}_{\mathcal{C}}(f)) \\ &= F(f)\end{aligned}$$

Since $A, B \in \text{Obj}(\mathcal{C})$ and $f \in \text{Hom}_{\mathcal{C}}(A, B)$ are arbitrary, $F \circ \text{id}_{\mathcal{C}} = F$.

□

Definition 4.2.0.4. Let $\mathcal{C}$ be a category. We define the **empty functor** from $\mathbf{0}$ to $\mathcal{C}$, denoted $E_{\mathcal{C}} : \mathbf{0} \rightarrow \mathcal{C}$ by $(E_{\mathcal{C}})_0 := \emptyset$ and $(E_{\mathcal{C}})_1 := \emptyset$.

Exercise 4.2.0.5. Let $\mathcal{C}$ be a category. Then $E_{\mathcal{C}} : \mathbf{0} \rightarrow \mathcal{C}$ is a functor.

Proof. Since $\text{Obj}(\mathbf{0}) = \emptyset$ and $\text{Hom}_{\mathbf{0}} = \emptyset$, this is vacuously true. □

Definition 4.2.0.6. Let $\mathcal{C}, \mathcal{D}$ be categories and $X \in \text{Obj}(\mathcal{D})$. We define the **constant functor** from $\mathcal{C}$ onto X , denoted $\Delta_X^{\mathcal{C}} : \mathcal{C} \rightarrow \mathcal{D}$ by

- $\Delta_X^{\mathcal{C}}(A) = X$
- $\Delta_X^{\mathcal{C}}(f) = \text{id}_X$

Exercise 4.2.0.7. Let $\mathcal{C}, \mathcal{D}$ be categories and $X \in \text{Obj}(\mathcal{D})$. Then $\Delta_X^{\mathcal{C}} : \mathcal{C} \rightarrow \mathcal{D}$ is a functor.

Proof.

1. Let $A, B \in \text{Obj}(\mathcal{C})$ and $f \in \text{Hom}_{\mathcal{C}}(A, B)$. Then

$$\begin{aligned}\Delta_X^{\mathcal{C}}(f) &= \text{id}_X \\ &\in \text{Hom}_{\mathcal{D}}(X, X) \\ &= \text{Hom}_{\mathcal{D}}(\Delta_X^{\mathcal{C}}(A), \Delta_X^{\mathcal{C}}(B))\end{aligned}$$

2. Let $A, B, C \in \text{Obj}(\mathcal{C})$, $f \in \text{Hom}_{\mathcal{C}}(A, B)$ and $g \in \text{Hom}_{\mathcal{C}}(B, C)$. Then

$$\begin{aligned}\Delta_X^{\mathcal{C}}(g \circ f) &= \text{id}_X \\ &= \text{id}_X \circ \text{id}_X \\ &= \Delta_X^{\mathcal{C}}(g) \circ \Delta_X^{\mathcal{C}}(f)\end{aligned}$$

3. Let $A \in \text{Obj}(\mathcal{C})$. Then

$$\begin{aligned}\Delta_X^{\mathcal{C}}(\text{id}_A) &= \text{id}_X \\ &= \text{id}_{\Delta_X^{\mathcal{C}}(A)}\end{aligned}$$

So $\Delta_X^{\mathcal{C}} : \mathcal{C} \rightarrow \mathcal{D}$ is a functor. □

Note 4.2.0.8. We recall for $X, Y \in \text{Obj}(\mathbf{Set})$, $f \in \text{Hom}_{\mathbf{Set}}(X, Y)$, the image and preimage maps induced by f , denoted $f_* \in \text{Hom}_{\mathbf{Set}}(\mathcal{P}(X), \mathcal{P}(Y))$ and $f^* \in \text{Hom}_{\mathbf{Set}}(\mathcal{P}(Y), \mathcal{P}(X))$ are defined by $f_*(A) := f(A)$ and $f^*(B) := f^{-1}(B)$. make exercise in set theory notes showing $(g \circ f)_* = g_* \circ f_*$ and similar for contravariant version

Definition 4.2.0.9. We define the

- **image power set functor on $\mathbf{Set}$** , denoted $\mathcal{P}_* : \mathbf{Set} \rightarrow \mathbf{Set}$ by
 - $\mathcal{P}_*(X) := \mathcal{P}(X)$, $X \in \text{Obj}(\mathbf{Set})$
 - $\mathcal{P}_*(f) := f_*$, $f \in \text{Hom}_{\mathbf{Set}}$
- **preimage power set functor on $\mathbf{Set}$** , denoted $\mathcal{P}^* : \mathbf{Set}^{\text{op}} \rightarrow \mathbf{Set}$ by
 - $\mathcal{P}^*(X) := \mathcal{P}(X)$, $X \in \text{Obj}(\mathbf{Set})$
 - $\mathcal{P}^*(f) := f^*$, $f \in \text{Hom}_{\mathbf{Set}}$

Exercise 4.2.0.10. We have that

1. $\mathcal{P}_* : \mathbf{Set} \rightarrow \mathbf{Set}$ is a functor.
2. $\mathcal{P}^* : \mathbf{Set}^{\text{op}} \rightarrow \mathbf{Set}$ is a functor.

Proof.

1. (a) For each $X, Y \in \text{Obj}(\mathbf{Set})$ and $f \in \text{Hom}_{\mathbf{Set}}(X, Y)$,

$$\begin{aligned}\mathcal{P}_*(f) &= f_* \\ &\in \text{Hom}_{\mathbf{Set}}(\mathcal{P}(X), \mathcal{P}(Y)) \quad (\text{ref ex in set theory section intro to math notes}) \\ &= \text{Hom}_{\mathbf{Set}}(\mathcal{P}_*(X), \mathcal{P}_*(Y)).\end{aligned}$$

- (b) For each $X, Y, Z \in \text{Obj}(\mathbf{Set})$, $f \in \text{Hom}_{\mathbf{Set}}(X, Y)$ and $g \in \text{Hom}_{\mathbf{Set}}(Y, Z)$,

$$\begin{aligned}\mathcal{P}_*(g \circ_{\mathbf{Set}} f) &= (g \circ_{\mathbf{Set}} f)_* \\ &= g_* \circ_{\mathbf{Set}} f_* \quad (\text{ref ex in set theory section intro to math notes}) \\ &= \mathcal{P}_*(g) \circ_{\mathbf{Set}} \mathcal{P}_*(f).\end{aligned}$$

(c) For each $X \in \text{Obj}(\mathbf{Set})$,

$$\begin{aligned}\mathcal{P}_*(\text{id}_X^{\mathbf{Set}}) &= \text{id}_{\mathcal{P}(X)}^{\mathbf{Set}} \quad (\text{ref ex in set theory section intro to math notes}) \\ &= \text{id}_{\mathcal{P}_*(X)}^{\mathbf{Set}}.\end{aligned}$$

Thus $\mathcal{P}_* : \mathbf{Set} \rightarrow \mathbf{Set}$ is a functor.

2. (a) Let $X, Y \in \text{Obj}(\mathbf{Set}^{\text{op}})$ and $f \in \text{Hom}_{\mathbf{Set}^{\text{op}}}(X, Y)$. Then $X \in \text{Obj}(\mathbf{Set})$ and $f \in \text{Hom}_{\mathbf{Set}}(Y, X)$. Therefore

$$\begin{aligned}\mathcal{P}^*(f) &= f^* \\ &\in \text{Hom}_{\mathbf{Set}}(\mathcal{P}(X), \mathcal{P}(Y)) \quad (\text{ref ex in set theory section intro to math notes}) \\ &= \text{Hom}_{\mathbf{Set}}(\mathcal{P}^*(X), \mathcal{P}^*(Y)).\end{aligned}$$

Since $X, Y \in \text{Obj}(\mathbf{Set}^{\text{op}})$ and $f \in \text{Hom}_{\mathbf{Set}^{\text{op}}}(X, Y)$ are arbitrary, we have that for each $X, Y \in \text{Obj}(\mathbf{Set}^{\text{op}})$ and $f \in \text{Hom}_{\mathbf{Set}^{\text{op}}}(X, Y)$, $\mathcal{P}^*(f) \in \text{Hom}_{\mathbf{Set}^{\text{op}}}(\mathcal{P}^*(X), \mathcal{P}^*(Y))$.

(b) Let $X, Y, Z \in \text{Obj}(\mathbf{Set}^{\text{op}})$, $f \in \text{Hom}_{\mathbf{Set}^{\text{op}}}(X, Y)$ and $g \in \text{Hom}_{\mathbf{Set}^{\text{op}}}(Y, Z)$. Then $X, Y, Z \in \text{Obj}(\mathbf{Set})$, $f \in \text{Hom}_{\mathbf{Set}}(Y, X)$ and $g \in \text{Hom}_{\mathbf{Set}}(Z, Y)$. Therefore

$$\begin{aligned}\mathcal{P}^*(g \circ_{\mathbf{Set}^{\text{op}}} f) &= (g \circ_{\mathbf{Set}^{\text{op}}} f)^* \\ &= (f \circ_{\mathbf{Set}} g)^* \\ &= g^* \circ_{\mathbf{Set}} f^* \quad (\text{ref ex in set theory section intro to math notes}) \\ &= \mathcal{P}^*(g) \circ_{\mathbf{Set}} \mathcal{P}^*(f).\end{aligned}$$

(c) For each $X \in \text{Obj}(\mathbf{Set}^{\text{op}})$,

$$\begin{aligned}\mathcal{P}^*(\text{id}_X^{\mathbf{Set}^{\text{op}}}) &= (\text{id}_X^{\mathbf{Set}^{\text{op}}})^* \\ &= (\text{id}_X^{\mathbf{Set}})^* \\ &= \text{id}_{\mathcal{P}(X)}^{\mathbf{Set}^{\text{op}}} \quad (\text{ref ex in set theory section intro to math notes}) \\ &= \text{id}_{\mathcal{P}^*(X)}^{\mathbf{Set}^{\text{op}}}.\end{aligned}$$

Thus $\mathcal{P}^* : \mathbf{Set}^{\text{op}} \rightarrow \mathbf{Set}$ is a functor. □

4.3 Category of Small Categories

Exercise 4.3.0.1. Let $\mathcal{C}$ and $\mathcal{D}$ be categories and $F : \mathcal{C} \rightarrow \mathcal{D}$. If $\mathcal{C}$ is small, then F is a set.

Proof. Suppose that $\mathcal{C}$ is small. Then $\text{Obj}(\mathcal{C})$ and $\text{Hom}_{\mathcal{C}}$ are sets. By definition, there exist $F_0 : \text{Obj}(\mathcal{C}) \rightarrow \text{Obj}(\mathcal{D})$ and $F_1 : \text{Hom}_{\mathcal{C}} \rightarrow \text{Hom}_{\mathcal{D}}$ such that $F = (F_0, F_1)$. Axiom 1.2.0.11 implies that $F_0(\text{Obj}(\mathcal{C}))$ and $F_1(\text{Hom}_{\mathcal{C}})$ are sets. Therefore, $\text{Obj}(\mathcal{C}) \times F_0(\text{Obj}(\mathcal{C}))$ and $\text{Hom}_{\mathcal{C}} \times F_1(\text{Hom}_{\mathcal{C}})$ are sets. Hence $\mathcal{P}(\text{Obj}(\mathcal{C}) \times F_0(\text{Obj}(\mathcal{C})))$ and $\mathcal{P}(\text{Hom}_{\mathcal{C}} \times F_1(\text{Hom}_{\mathcal{C}}))$ are sets. Since $F_0 \subset \text{Obj}(\mathcal{C}) \times F_0(\text{Obj}(\mathcal{C}))$ and $F_1 \subset \text{Hom}_{\mathcal{C}} \times F_1(\text{Hom}_{\mathcal{C}})$, we have that $F_0 \in \mathcal{P}(\text{Obj}(\mathcal{C}) \times F_0(\text{Obj}(\mathcal{C})))$ and $F_1 \in \mathcal{P}(\text{Hom}_{\mathcal{C}} \times F_1(\text{Hom}_{\mathcal{C}}))$. Hence F_0 and F_1 are sets. Thus $F = (F_0, F_1)$ is a set. □

Exercise 4.3.0.2. Let $\mathcal{C}$ and $\mathcal{D}$ be categories. Suppose that $\mathcal{C}$ is small. Then there exists a class A such that for each class F , $F \in A$ iff $F : \mathcal{C} \rightarrow \mathcal{D}$.

Proof. Let $\mathcal{C}$ and $\mathcal{D}$ be categories. Suppose that $\mathcal{C}$ is small. Define ϕ by

$$\phi(F) : F : \mathcal{C} \rightarrow \mathcal{D}$$

Then there exists a class A such that for each set F , $F \in A$ iff $\phi(F)$. Let F be a class. Suppose that $F \in A$. By Definition 1.2.0.1, F is a set. Since F is a set and $F \in A$, we have that $\phi(F)$. Hence $F : \mathcal{C} \rightarrow \mathcal{D}$.

Conversely, suppose that $F : \mathcal{C} \rightarrow \mathcal{D}$. Exercise 4.3.0.1 implies that F is a set. Since F is a set and $\phi(F)$ is true, we have that $F \in A$. □

Definition 4.3.0.3. We define **Cat** by

- $\text{Obj}(\mathbf{Cat}) = \{\mathcal{C} : \mathcal{C} \text{ is a small category}\}.$
- for $\mathcal{C}, \mathcal{D} \in \text{Obj}(\mathbf{Cat})$,

$$\text{Hom}_{\mathbf{Cat}}(\mathcal{C}, \mathcal{D}) = \{F : \mathcal{C} \rightarrow \mathcal{D}\}$$

- for $\mathcal{C}, \mathcal{D}, \mathcal{E} \in \text{Obj}(\mathbf{Cat})$, $F \in \text{Hom}_{\mathbf{Cat}}(\mathcal{C}, \mathcal{D})$ and $G \in \text{Hom}_{\mathbf{Cat}}(\mathcal{D}, \mathcal{E})$,

$$G \circ_{\mathbf{Cat}} F = G \circ F$$

Exercise 4.3.0.4. We have that **Cat** is

1. a category
2. locally small

Proof.

1. Exercise 4.1.0.6 implies that composition is well defined. Exercise 4.1.0.7 implies that composition is associative. Exercise 4.2.0.2 and Exercise 4.2.0.3 imply the existence of identities.
2. Let $\mathcal{C}, \mathcal{D} \in \text{Obj}(\mathbf{Cat})$ and $F \in \text{Hom}_{\mathbf{Cat}}(\mathcal{C}, \mathcal{D})$. Definition 3.1.0.8 implies that $\text{Obj}(\mathcal{C})$, $\text{Obj}(\mathcal{D})$, $\text{Hom}_{\mathcal{C}}$ and $\text{Hom}_{\mathcal{D}}$ are sets. Then $\text{Obj}(\mathcal{D})^{\text{Obj}(\mathcal{C})}$ and $\text{Hom}_{\mathcal{D}}^{\text{Hom}_{\mathcal{C}}}$ are sets. Hence $\text{Obj}(\mathcal{D})^{\text{Obj}(\mathcal{C})} \times \text{Hom}_{\mathcal{D}}^{\text{Hom}_{\mathcal{C}}}$ is a set. Let $F \in \text{Hom}_{\mathbf{Cat}}(\mathcal{C}, \mathcal{D})$. Then there exist $F_0 \in \text{Obj}(\mathcal{D})^{\text{Obj}(\mathcal{C})}$ and $F_1 \in \text{Hom}_{\mathcal{D}}^{\text{Hom}_{\mathcal{C}}}$ such that $F = (F_0, F_1)$. Therefore $F \in \text{Obj}(\mathcal{D})^{\text{Obj}(\mathcal{C})} \times \text{Hom}_{\mathcal{D}}^{\text{Hom}_{\mathcal{C}}}$. Since $F \in \text{Hom}_{\mathbf{Cat}}(\mathcal{C}, \mathcal{D})$ is arbitrary,

$$\text{Hom}_{\mathbf{Cat}}(\mathcal{C}, \mathcal{D}) \subset \text{Obj}(\mathcal{D})^{\text{Obj}(\mathcal{C})} \times \text{Hom}_{\mathcal{D}}^{\text{Hom}_{\mathcal{C}}}$$

which implies that $\text{Hom}_{\mathbf{Cat}}(\mathcal{C}, \mathcal{D})$ is a set. Therefore, **Cat** is locally small. □

4.4 Comma Categories

Definition 4.4.0.1. Let $\mathcal{A}, \mathcal{B}, \mathcal{C}$ be categories and $S : \mathcal{A} \rightarrow \mathcal{C}$, $T : \mathcal{B} \rightarrow \mathcal{C}$ functors. We define the **comma category of S to T** , denoted $(S \downarrow T)$, by

- $\text{Obj}(S \downarrow T) = \coprod_{(A, B) \in \text{Obj}(\mathcal{A}) \times \text{Obj}(\mathcal{B})} \text{Hom}_{\mathcal{C}}(S(A), T(B))$
- For $(A_1, B_1, h_1), (A_2, B_2, h_2) \in \text{Obj}(S \downarrow T)$,

$$\begin{aligned} \text{Hom}_{(S \downarrow T)}((A_1, B_1, h_1), (A_2, B_2, h_2)) = \\ \{(\alpha, \beta) \in \text{Hom}_{\mathcal{A}}(A_1, A_2) \times \text{Hom}_{\mathcal{B}}(B_1, B_2) : T(\beta) \circ_{\mathcal{C}} h_1 = h_2 \circ_{\mathcal{C}} S(\alpha)\} \end{aligned}$$

i.e. for $(A_1, B_1, h_1), (A_2, B_2, h_2) \in \text{Obj}(S \downarrow T)$, $\alpha \in \text{Hom}_{\mathcal{A}}(A_1, A_2)$ and $\beta \in \text{Hom}_{\mathcal{B}}(B_1, B_2)$, $(\alpha, \beta) \in \text{Hom}_{(S \downarrow T)}((A_1, B_1, h_1), (A_2, B_2, h_2))$ iff the following diagram commutes:

$$\begin{array}{ccc} S(A_1) & \xrightarrow{S(\alpha)} & S(A_2) \\ h_1 \downarrow & & \downarrow h_2 \\ T(B_1) & \xrightarrow{T(\beta)} & T(B_2) \end{array}$$

- For
 - $(A_1, B_1, h_1), (A_2, B_2, h_2), (A_3, B_3, h_3) \in \text{Obj}(S \downarrow T)$
 - $(\alpha_{12}, \beta_{12}) \in \text{Hom}_{(S \downarrow T)}((A_1, B_1, h_1), (A_2, B_2, h_2))$
 - $(\alpha_{23}, \beta_{23}) \in \text{Hom}_{(S \downarrow T)}((A_2, B_2, h_2), (A_3, B_3, h_3))$

we define

$$(\alpha_{23}, \beta_{23}) \circ_{(S \downarrow T)} (\alpha_{12}, \beta_{12}) = (\alpha_{23} \circ_{\mathcal{A}} \alpha_{12}, \beta_{23} \circ_{\mathcal{B}} \beta_{12})$$

Exercise 4.4.0.2. Let $\mathcal{A}, \mathcal{B}, \mathcal{C}$ be categories and $S : \mathcal{A} \rightarrow \mathcal{C}, T : \mathcal{B} \rightarrow \mathcal{C}$ functors. Then $(S \downarrow T)$ is a category.

Proof.

• **well-definedness of composition:**

Let

- $(A_1, B_1, h_1), (A_2, B_2, h_2), (A_3, B_3, h_3) \in \text{Obj}(S \downarrow T)$
- $(\alpha_{12}, \beta_{12}) \in \text{Hom}_{(S \downarrow T)}((A_1, B_1, h_1), (A_2, B_2, h_2))$
- $(\alpha_{23}, \beta_{23}) \in \text{Hom}_{(S \downarrow T)}((A_2, B_2, h_2), (A_3, B_3, h_3))$

By definition, $\alpha_{12} \in \text{Hom}_{\mathcal{A}}(A_1, A_2), \alpha_{23} \in \text{Hom}_{\mathcal{A}}(A_2, A_3), \beta_{12} \in \text{Hom}_{\mathcal{B}}(B_1, B_2), \beta_{23} \in \text{Hom}_{\mathcal{B}}(B_2, B_3), T(\beta_{12}) \circ_{\mathcal{C}} h_1 = h_2 \circ S(\alpha_{12})$ and $T(\beta_{23}) \circ_{\mathcal{C}} h_2 = h_3 \circ S(\alpha_{23})$,

i.e. the following diagram commutes:

$$\begin{array}{ccccc} S(A_1) & \xrightarrow{S(\alpha_{12})} & S(A_2) & \xrightarrow{S(\alpha_{23})} & S(A_3) \\ h_1 \downarrow & & \downarrow h_2 & & \downarrow h_3 \\ T(B_1) & \xrightarrow{T(\beta_{12})} & T(B_2) & \xrightarrow{T(\beta_{23})} & T(B_3) \end{array}$$

Then $\alpha_{23} \circ_{\mathcal{A}} \alpha_{12} \in \text{Hom}_{\mathcal{A}}(A_1, A_3), \beta_{23} \circ_{\mathcal{B}} \beta_{12} \in \text{Hom}_{\mathcal{B}}(B_1, B_3)$ and

$$\begin{aligned} T(\beta_{23} \circ_{\mathcal{B}} \beta_{12}) \circ_{\mathcal{C}} h_1 &= (T(\beta_{23}) \circ_{\mathcal{C}} T(\beta_{12})) \circ_{\mathcal{C}} h_1 \\ &= T(\beta_{23}) \circ_{\mathcal{C}} (T(\beta_{12}) \circ_{\mathcal{C}} h_1) \\ &= T(\beta_{23}) \circ_{\mathcal{C}} (h_2 \circ_{\mathcal{C}} S(\alpha_{12})) \\ &= (T(\beta_{23}) \circ_{\mathcal{C}} h_2) \circ_{\mathcal{C}} S(\alpha_{12}) \\ &= (h_3 \circ_{\mathcal{C}} S(\alpha_{23})) \circ_{\mathcal{C}} S(\alpha_{12}) \\ &= h_3 \circ_{\mathcal{C}} (S(\alpha_{23}) \circ_{\mathcal{C}} S(\alpha_{12})) \\ &= h_3 \circ_{\mathcal{C}} S(\alpha_{23} \circ_{\mathcal{A}} \alpha_{12}) \end{aligned}$$

i.e. the following diagram commutes:

$$\begin{array}{ccc} S(A_1) & \xrightarrow{S(\alpha_{23} \circ_{\mathcal{A}} \alpha_{12})} & S(A_3) \\ h_1 \downarrow & & \downarrow h_3 \\ T(B_1) & \xrightarrow{T(\beta_{23} \circ_{\mathcal{B}} \beta_{12})} & T(B_3) \end{array}$$

Hence $(\alpha_{23} \circ_{\mathcal{A}} \alpha_{12}, \beta_{23} \circ_{\mathcal{B}} \beta_{12}) \in \text{Hom}_{(S \downarrow T)}((A_1, B_1, h_1), (A_3, B_3, h_3))$ and composition is well defined.

• **associativity of composition:**

Let

- $(A_1, B_1, h_1), (A_2, B_2, h_2), (A_3, B_3, h_3), (A_4, B_4, h_4) \in \text{Obj}(S \downarrow T)$
- $(\alpha_{12}, \beta_{12}) \in \text{Hom}_{(S \downarrow T)}((A_1, B_1, h_1), (A_2, B_2, h_2))$
- $(\alpha_{23}, \beta_{23}) \in \text{Hom}_{(S \downarrow T)}((A_2, B_2, h_2), (A_3, B_3, h_3))$
- $(\alpha_{34}, \beta_{34}) \in \text{Hom}_{(S \downarrow T)}((A_3, B_3, h_3), (A_4, B_4, h_4))$

Then

$$\begin{aligned} [(\alpha_{34}, \beta_{34}) \circ_{(S \downarrow T)} (\alpha_{23}, \beta_{23})] \circ_{(S \downarrow T)} (\alpha_{12}, \beta_{12}) &= (\alpha_{34} \circ_{\mathcal{A}} \alpha_{23}, \beta_{34} \circ_{\mathcal{B}} \beta_{23}) \circ_{(S \downarrow T)} (\alpha_{12}, \beta_{12}) \\ &= ([\alpha_{34} \circ_{\mathcal{A}} \alpha_{23}] \circ_{\mathcal{A}} \alpha_{12}, [\beta_{34} \circ_{\mathcal{B}} \beta_{23}] \circ_{\mathcal{B}} \beta_{12}) \\ &= (\alpha_{34} \circ_{\mathcal{A}} [\alpha_{23} \circ_{\mathcal{A}} \alpha_{12}], \beta_{34} \circ_{\mathcal{B}} [\beta_{23} \circ_{\mathcal{B}} \beta_{12}]) \\ &= (\alpha_{34}, \beta_{34}) \circ_{(S \downarrow T)} (\alpha_{23} \circ_{\mathcal{A}} \alpha_{12}, \beta_{23} \circ_{\mathcal{B}} \beta_{12}) \\ &= (\alpha_{34}, \beta_{34}) \circ_{(S \downarrow T)} [(\alpha_{23}, \beta_{23}) \circ_{(S \downarrow T)} (\alpha_{12}, \beta_{12})] \end{aligned}$$

So composition is associative.

• **existence of identities:**

Let

- $(A_1, B_1, h_1), (A_2, B_2, h_2) \in \text{Obj}(S \downarrow T)$
- $(\alpha, \beta) \in \text{Hom}_{(S \downarrow T)}((A_1, B_1, h_1), (A_2, B_2, h_2))$

By definition,

- $\alpha \in \text{Hom}_{\mathcal{A}}(A_1, A_2), \beta \in \text{Hom}_{\mathcal{B}}(B_1, B_2)$
- $h_1 \in \text{Hom}_{\mathcal{C}}(S(A_1), T(B_1)), h_2 \in \text{Hom}_{\mathcal{C}}(S(A_2), T(B_2))$
- $T(\beta) \circ h_1 = h_2 \circ S(\alpha)$

Since $\text{id}_{A_1} \in \text{Hom}_{\mathcal{A}}(A_1, A_1), \text{id}_{B_1} \in \text{Hom}_{\mathcal{B}}(B_1, B_1)$, and

$$\begin{aligned} T(\text{id}_{B_1}) \circ_{\mathcal{C}} h_1 &= \text{id}_{T(B_1)} \circ_{\mathcal{C}} h_1 \\ &= h_1 \\ &= h_1 \circ_{\mathcal{C}} \text{id}_{S(A_1)} \\ &= h_1 \circ_{\mathcal{C}} S(\text{id}_{A_1}) \end{aligned}$$

i.e. the following diagram commutes:

$$\begin{array}{ccc} S(A_1) & \xrightarrow{S(\text{id}_{A_1})} & S(A_1) \\ h_1 \downarrow & & \downarrow h_1 \\ T(B_1) & \xrightarrow{T(\text{id}_{B_1})} & T(B_1) \end{array}$$

we have that $(\text{id}_{A_1}, \text{id}_{B_1}) \in \text{Hom}_{(S \downarrow T)}((A_1, B_1, h_1), (A_1, B_1, h_1))$. Similarly $(\text{id}_{A_2}, \text{id}_{B_2}) \in \text{Hom}_{(S \downarrow T)}((A_2, B_2, h_2), (A_2, B_2, h_2))$. Therefore

$$\begin{aligned} (\alpha, \beta) \circ_{(S \downarrow T)} (\text{id}_{A_1}, \text{id}_{B_1}) &= (\alpha \circ_{\mathcal{A}} \text{id}_{A_1}, \beta \circ_{\mathcal{B}} \text{id}_{B_1}) \\ &= (\alpha, \beta) \end{aligned}$$

and

$$\begin{aligned} (\text{id}_{A_2}, \text{id}_{B_2}) \circ_{(S \downarrow T)} (\alpha, \beta) &= (\text{id}_{A_2} \circ_{\mathcal{A}} \alpha, \text{id}_{B_2} \circ_{\mathcal{B}} \beta) \\ &= (\alpha, \beta) \end{aligned}$$

Since $(A_1, B_1, h_1), (A_2, B_2, h_2) \in \text{Obj}(S \downarrow T)$ and

$(\alpha, \beta) \in \text{Hom}_{(S \downarrow T)}((A_1, B_1, h_1), (A_2, B_2, h_2))$ are arbitrary, we have that for each $(A, B, h) \in \text{Obj}(S \downarrow T)$, $\text{id}_{(A, B, h)} = (\text{id}_A, \text{id}_B)$. □

Definition 4.4.0.3. Let $\mathcal{C}$ be a category. We define the **arrow category of $\mathcal{C}$** , denoted $\mathbf{Arr}(\mathcal{C})$, by $\mathbf{Arr}(\mathcal{C}) := (\text{id}_{\mathcal{C}} \downarrow \text{id}_{\mathcal{C}})$.

Note 4.4.0.4. explain with diagram how in the case $\alpha = \Delta_X^1$ and how we can contract one edge of the rectangle diagram to get a triangle

Definition 4.4.0.5. Let $\mathcal{C}, \mathcal{D}$ be categories, $X \in \text{Obj}(\mathcal{D})$ and $F : \mathcal{C} \rightarrow \mathcal{D}$. We define the **comma category from F to X** , denoted $(F \downarrow X)$, by $(F \downarrow X) := (F \downarrow \Delta_X^1)$.

We may make the following identification:

$$\bullet \text{Obj}(F \downarrow X) = \coprod_{A \in \text{Obj}(\mathcal{C})} \text{Hom}_{\mathcal{D}}(F(A), X)$$

- For $(A_1, f_1), (A_2, f_2) \in \text{Obj}(F \downarrow X)$,

$$\text{Hom}_{(X \downarrow F)}((A_1, f_1), (A_2, f_2)) = \{\alpha \in \text{Hom}_{\mathcal{C}}(A_1, A_2) : f_2 \circ F(\alpha) = f_1\},$$

i.e. for $(A_1, f_1), (A_2, f_2) \in \text{Obj}(F \downarrow X)$ and $\alpha \in \text{Hom}_{A_1, A_2}$, $\alpha \in \text{Hom}_{(F \downarrow X)}((A_1, f_1), (A_2, f_2))$ iff the following diagram commutes:

$$\begin{array}{ccc} F(A_1) & \xrightarrow{F(\alpha)} & F(A_2) \\ & \searrow f_1 \quad \swarrow f_2 & \\ & X & \end{array}$$

- For
 - $(A_1, f_1), (A_2, f_2), (A_3, f_3) \in \text{Obj}(F \downarrow X)$
 - $\alpha \in \text{Hom}_{(F \downarrow X)}((A_1, f_1), (A_2, f_2))$
 - $\beta \in \text{Hom}_{(F \downarrow X)}((A_2, f_2), (A_3, f_3))$

we define

$$\beta \circ_{(F \downarrow X)} \alpha = \beta \circ_{\mathcal{C}} \alpha$$

Definition 4.4.0.6. Let $\mathcal{C}, \mathcal{D}$ be categories, $X \in \text{Obj}(\mathcal{D})$ and $F : \mathcal{C} \rightarrow \mathcal{D}$. We define the **comma category from X to F** , denoted $(X \downarrow F)$, by $(X \downarrow F) := (\Delta_X^1 \downarrow F)$.

We may make the following identification:

- $\text{Obj}(X \downarrow F) = \coprod_{A \in \text{Obj}(\mathcal{C})} \text{Hom}_{\mathcal{D}}(X, F(A))$
- For $(A_1, f_1), (A_2, f_2) \in \text{Obj}(X \downarrow F)$,

$$\text{Hom}_{(X \downarrow F)}((A_1, f_1), (A_2, f_2)) = \{\alpha \in \text{Hom}_{\mathcal{C}}(A_1, A_2) : F(\alpha) \circ f_1 = f_2\},$$

i.e. for $(A_1, f_1), (A_2, f_2) \in \text{Obj}(X \downarrow F)$ and $\alpha \in \text{Hom}_{A_1, A_2}$, $\alpha \in \text{Hom}_{(X \downarrow F)}((A_1, f_1), (A_2, f_2))$ iff the following diagram commutes:

$$\begin{array}{ccc} & X & \\ f_1 \swarrow & & \searrow f_2 \\ F(A_1) & \xrightarrow{F(\alpha)} & F(A_2) \end{array}$$

- For
 - $(A_1, f_1), (A_2, f_2), (A_3, f_3) \in \text{Obj}(X \downarrow F)$
 - $\alpha \in \text{Hom}_{(X \downarrow F)}((A_1, f_1), (A_2, f_2))$
 - $\beta \in \text{Hom}_{(X \downarrow F)}((A_2, f_2), (A_3, f_3))$

we define

$$\beta \circ_{(X \downarrow F)} \alpha = \beta \circ_{\mathcal{C}} \alpha$$

Definition 4.4.0.7. Let $\mathcal{C}$ be a category and $X \in \text{Obj}(\mathcal{C})$. We define the

- **slice category of $\mathcal{C}$ over X** or the **over category of $\mathcal{C}$ over X** , denoted $\mathcal{C} \downarrow X$, by $\mathcal{C} \downarrow X := (\text{id}_{\mathcal{C}} \downarrow X)$.
- **coslice category of $\mathcal{C}$ over X** or the **under category of $\mathcal{C}$ under X** , denoted $X \downarrow \mathcal{C}$, by $X \downarrow \mathcal{C} := (X \downarrow \text{id}_{\mathcal{C}})$.

give examples in posets, and other

Chapter 5

Natural Transformations

5.1 Introduction

Definition 5.1.0.1. Let $\mathcal{C}$ and $\mathcal{D}$ be categories, $F, G : \mathcal{C} \rightarrow \mathcal{D}$ and $\alpha : \text{Obj}(\mathcal{C}) \rightarrow \text{Hom}_{\mathcal{D}}$. Then α is said to be a **natural transformation from F to G** , denoted $\alpha : F \Rightarrow G$, if

1. for each $A \in \text{Obj}(\mathcal{C})$, $\alpha_A \in \text{Hom}_{\mathcal{D}}(F(A), G(A))$
2. for each $A, B \in \text{Obj}(\mathcal{C})$ and $f \in \text{Hom}_{\mathcal{C}}(A, B)$, $G(f) \circ \alpha_A = \alpha_B \circ F(f)$, i.e. the following diagram commutes:

$$\begin{array}{ccc} F(A) & \xrightarrow{\alpha_A} & G(A) \\ F(f) \downarrow & & \downarrow G(f) \\ F(B) & \xrightarrow{\alpha_B} & G(B) \end{array}$$

Definition 5.1.0.2. Vertical Composition:

Let $\mathcal{C}, \mathcal{D}$ be categories, $F, G, H : \mathcal{C} \rightarrow \mathcal{D}$ functors and $\alpha : F \Rightarrow G$, $\beta : G \Rightarrow H$ natural transformations. We define the **vertical composition of β with α** , denoted $\beta \circ \alpha : F \Rightarrow H$, by

$$(\beta \circ \alpha)_A = \beta_A \circ \alpha_A$$

Exercise 5.1.0.3. Let $\mathcal{C}, \mathcal{D}$ be categories, $F, G, H : \mathcal{C} \rightarrow \mathcal{D}$ functors and $\alpha : F \Rightarrow G$, $\beta : G \Rightarrow H$ natural transformations. Then $\beta \circ \alpha : F \Rightarrow H$ is a natural transformation.

Proof.

1. Let $A \in \text{Obj}(\mathcal{C})$. Since $\alpha_A \in \text{Hom}_{\mathcal{D}}(F(A), G(A))$ and $\beta_A \in \text{Hom}_{\mathcal{D}}(G(A), H(A))$, we have that

$$\begin{aligned} (\beta \circ \alpha)_A &= \beta_A \circ \alpha_A \\ &\in \text{Hom}_{\mathcal{D}}(F(A), H(A)) \end{aligned}$$

2. Let $A, B \in \text{Obj}(\mathcal{C})$ and $f \in \text{Hom}_{\mathcal{C}}(A, B)$. Since $\alpha : F \Rightarrow G$ and $\beta : G \Rightarrow H$, $G(f) \circ \alpha_A = \alpha_B \circ F(f)$ and $H(f) \circ \beta_A = \beta_B \circ G(f)$. Therefore

$$\begin{aligned} H(f) \circ (\beta \circ \alpha)_A &= H(f) \circ (\beta_A \circ \alpha_A) \\ &= (H(f) \circ \beta_A) \circ \alpha_A \\ &= (\beta_B \circ G(f)) \circ \alpha_A \\ &= \beta_B \circ (G(f) \circ \alpha_A) \\ &= \beta_B \circ (\alpha_B \circ F(f)) \\ &= (\beta_B \circ \alpha_B) \circ F(f) \\ &= (\beta \circ \alpha)_B \circ F(f) \end{aligned}$$

So $\beta \circ \alpha : F \Rightarrow H$ is a natural transformation. $\square$

Exercise 5.1.0.4. Let $\mathcal{C}, \mathcal{D}$ be categories, $F, G, H, K : \mathcal{C} \rightarrow \mathcal{D}$ functors and $\alpha : F \Rightarrow G$, $\beta : G \Rightarrow H$ and $\gamma : H \Rightarrow K$ natural transformations. Then

$$(\gamma \circ \beta) \circ \alpha = \gamma \circ (\beta \circ \alpha)$$

Proof. Let $A \in \text{Obj}(\mathcal{C})$. By definition,

$$\begin{aligned} [(\gamma \circ \beta) \circ \alpha]_A &= (\gamma \circ \beta)_A \circ \alpha_A \\ &= (\gamma_A \circ \beta_A) \circ \alpha_A \\ &= \gamma_A \circ (\beta_A \circ \alpha_A) \\ &= \gamma_A \circ (\beta \circ \alpha)_A \\ &= [\gamma \circ (\beta \circ \alpha)]_A \end{aligned}$$

Since $A \in \text{Obj}(\mathcal{C})$ is arbitrary,

$$(\gamma \circ \beta) \circ \alpha = \gamma \circ (\beta \circ \alpha)$$

$\square$

Definition 5.1.0.5. Horizontal Composition:

Let $\mathcal{A}, \mathcal{B}, \mathcal{C}$ be categories, $F, G : \mathcal{A} \rightarrow \mathcal{B}$, $F', G' : \mathcal{B} \rightarrow \mathcal{C}$ functors and $\alpha : F \Rightarrow G$, $\alpha' : F' \Rightarrow G'$ natural transformations. We define the **horizontal composition of β with α** , denoted $\beta * \alpha : F' \circ F \Rightarrow G' \circ G$, by

$$(\alpha' * \alpha)_A = \alpha'_{G(A)} \circ F'(\alpha_A)$$

Exercise 5.1.0.6. Let $\mathcal{A}, \mathcal{B}, \mathcal{C}$ be categories, $F, G : \mathcal{A} \rightarrow \mathcal{B}$, $F', G' : \mathcal{B} \rightarrow \mathcal{C}$ functors and $\alpha : F \Rightarrow G$, $\alpha' : F' \Rightarrow G'$ natural transformations. Then $\alpha' \circ \alpha : F' \circ F \Rightarrow G' \circ G$ is a natural transformation.

Proof.

1. Let $A \in \text{Obj}(\mathcal{A})$. Since $\alpha : F \Rightarrow G$, we have that $\alpha_A \in \text{Hom}_{\mathcal{B}}(F(A), G(A))$. Since $F' : \mathcal{B} \rightarrow \mathcal{C}$, we have that

$$\begin{aligned} F'(\alpha_A) &\in \text{Hom}_{\mathcal{C}}(F'(F(A)), F'(G(A))) \\ &= \text{Hom}_{\mathcal{C}}(F' \circ F(A), F' \circ G(A)). \end{aligned}$$

Since $G : \mathcal{A} \rightarrow \mathcal{B}$, we have that $G(A) \in \text{Obj}(\mathcal{B})$. Since $\alpha' : F' \Rightarrow G'$, we have that

$$\begin{aligned} \alpha'_{G(A)} &\in \text{Hom}_{\mathcal{C}}(F'(G(A)), G'(G(A))) \\ &= \text{Hom}_{\mathcal{C}}(F' \circ G(A), G' \circ G(A)). \end{aligned}$$

Therefore

$$\begin{aligned} (\alpha' * \alpha)_A &= \alpha'_{G(A)} \circ F'(\alpha_A) \\ &\in \text{Hom}_{\mathcal{C}}(F' \circ F(A), G' \circ G(A)). \end{aligned}$$

Since $A \in \text{Obj}(\mathcal{A})$ is arbitrary, we have that for each $A \in \text{Obj}(\mathcal{A})$, $(\alpha' * \alpha)_A \in \text{Hom}_{\mathcal{C}}(F' \circ F(A), G' \circ G(A))$.

2. Let $A_1, A_2 \in \text{Obj}(\mathcal{A})$ and $f \in \text{Hom}_{\mathcal{A}}(A_1, A_2)$. Since $F, G : \mathcal{A} \rightarrow \mathcal{B}$, we have that $F(f) \in \text{Hom}_{\mathcal{B}}(F(A_1), F(A_2))$ and $G(f) \in \text{Hom}_{\mathcal{B}}(G(A_1), G(A_2))$. Since $\alpha : F \Rightarrow G$, we have that for each $k \in [2]$, $\alpha_{A_k} \in \text{Hom}_{\mathcal{B}}(F(A_k), G(A_k))$ and $G(f) \circ \alpha_{A_1} = \alpha_{A_2} \circ F(f)$, i.e. the following diagram commutes:

$$\begin{array}{ccc} F(A_1) & \xrightarrow{\alpha_{A_1}} & G(A_1) \\ F(f) \downarrow & & \downarrow G(f) \\ F(A_2) & \xrightarrow{\alpha_{A_2}} & G(A_2) \end{array}$$

Since $F', G' : \mathcal{B} \rightarrow \mathcal{C}$, we have that

$$\begin{aligned} F' \circ F(f) &= F'(F(f)) \\ &\in \text{Hom}_{\mathcal{C}}(F'(F(A_1)), F'(F(A_2))) \\ &= \text{Hom}_{\mathcal{C}}(F' \circ F(A_1), F' \circ F(A_2)), \end{aligned}$$

$$\begin{aligned} F' \circ G(f) &= F'(G(f)) \\ &\in \text{Hom}_{\mathcal{C}}(F'(G(A_1)), F'(G(A_2))) \\ &= \text{Hom}_{\mathcal{C}}(F' \circ G(A_1), F' \circ G(A_2)), \end{aligned}$$

$$\begin{aligned} G' \circ G(f) &= G'(G(f)) \\ &\in \text{Hom}_{\mathcal{C}}(G'(G(A_1)), G'(G(A_2))) \\ &= \text{Hom}_{\mathcal{C}}(G' \circ G(A_1), G' \circ G(A_2)), \end{aligned}$$

and for each $k \in [2]$,

$$\begin{aligned} F'(\alpha_{A_k}) &\in \text{Hom}_{\mathcal{C}}(F'(F(A_k)), F'(G(A_k))) \\ &= \text{Hom}_{\mathcal{C}}(F' \circ F(A_k), F' \circ G(A_k)). \end{aligned}$$

Since $\alpha' : F' \Rightarrow G'$, we have that for each $k \in [2]$,

$$\begin{aligned} \alpha'_{G(A_k)} &\in \text{Hom}_{\mathcal{C}}(F'(G(A_k)), G'(G(A_k))) \\ &= \text{Hom}_{\mathcal{C}}(F' \circ G(A_k), G' \circ G(A_k)). \end{aligned}$$

Therefore

$$\begin{aligned} [F' \circ G(f)] \circ F'(\alpha_{A_1}) &= F'(G(f)) \circ F'(\alpha_{A_1}) \\ &= F'(G(f) \circ \alpha_{A_1}) \\ &= F'(\alpha_{A_2} \circ F(f)) \\ &= F'(\alpha_{A_2}) \circ F'(F(f)) \\ &= F'(\alpha_{A_2}) \circ [F' \circ F(f)], \end{aligned}$$

and

$$\begin{aligned} [G' \circ G(f)] \circ \alpha'_{G(A_1)} &= G'(G(f)) \circ \alpha'_{G(A_1)} \\ &= \alpha'_{G(A_2)} \circ F'(G(f)) \\ &= \alpha'_{G(A_2)} \circ [F' \circ G(f)], \end{aligned}$$

i.e. the following diagram commutes:

$$\begin{array}{ccccc} F' \circ F(A_1) & \xrightarrow{F'(\alpha_{A_1})} & F' \circ G(A_1) & \xrightarrow{\alpha'_{G(A_1)}} & G' \circ G(A_1) \\ \downarrow F' \circ F(f) & & \downarrow F' \circ G(f) & & \downarrow G' \circ G(f) \\ F' \circ F(A_2) & \xrightarrow{F'(\alpha_{A_2})} & F' \circ G(A_2) & \xrightarrow{\alpha'_{G(A_2)}} & G' \circ G(A_2) \end{array}$$

Hence

$$\begin{aligned} [G' \circ G(f)] \circ (\alpha' * \alpha)_{A_1} &= [G' \circ G(f)] \circ [\alpha'_{G(A_1)} \circ F'(\alpha_{A_1})] \\ &= ([G' \circ G(f)] \circ \alpha'_{G(A_1)}) \circ F'(\alpha_{A_1}) \\ &= (\alpha'_{G(A_2)} \circ [F' \circ G(f)]) \circ F'(\alpha_{A_1}) \\ &= \alpha'_{G(A_2)} \circ ([F' \circ G(f)] \circ F'(\alpha_{A_1})) \\ &= \alpha'_{G(A_2)} \circ (F'(\alpha_{A_2}) \circ [F' \circ F(f)]) \\ &= [\alpha'_{G(A_2)} \circ F'(\alpha_{A_2})] \circ [F' \circ F(f)] \\ &= (\alpha' * \alpha)_{A_2} \circ [F' \circ F(f)], \end{aligned}$$

i.e. the following diagram commutes:

$$\begin{array}{ccc} F' \circ F(A_1) & \xrightarrow{(\alpha' * \alpha)_{A_1}} & G' \circ G(A_1) \\ F' \circ F(f) \downarrow & & \downarrow G' \circ G(f) \\ F' \circ F(A_2) & \xrightarrow{(\alpha' * \alpha)_{A_2}} & G' \circ G(A_2) \end{array}$$

Since $A_1, A_2 \in \text{Obj}(\mathcal{A})$ and $f \in \text{Hom}_{\mathcal{A}}(A_1, A_2)$ are arbitrary, we have that for each $A_1, A_2 \in \text{Obj}(\mathcal{A})$ and $f \in \text{Hom}_{\mathcal{A}}(A_1, A_2)$, $[G' \circ G(f)] \circ (\alpha' * \alpha)_{A_1} = (\alpha' * \alpha)_{A_2} \circ [F' \circ F(f)]$.

Thus $\alpha' * \alpha : F' \circ F \Rightarrow G' \circ G$. □

Exercise 5.1.0.7. Let

- $\mathcal{C}_1, \mathcal{C}_2, \mathcal{C}_3, \mathcal{C}_4$ be categories,
- $F_{1,2}, G_{1,2} : \mathcal{C}_1 \rightarrow \mathcal{C}_2$, $F_{2,3}, G_{2,3} : \mathcal{C}_2 \rightarrow \mathcal{C}_3$, $F_{3,4}, G_{3,4} : \mathcal{C}_3 \rightarrow \mathcal{C}_4$
- $\alpha^{1,2} : F_{1,2} \Rightarrow G_{1,2}$, $\alpha^{2,3} : F_{2,3} \Rightarrow G_{2,3}$, $\alpha^{3,4} : F_{3,4} \Rightarrow G_{3,4}$.

Then $\alpha^{3,4} * (\alpha^{2,3} * \alpha^{1,2}) = (\alpha^{3,4} * \alpha^{2,3}) * \alpha^{1,2}$.

Proof. Let $A \in \text{Obj}(\mathcal{C}_1)$. By definition and [ref prev ex](#),

- $\alpha^{2,3} * \alpha^{1,2} : F_{2,3} \circ F_{1,2} \Rightarrow G_{2,3} \circ G_{1,2}$ and $(\alpha^{2,3} * \alpha^{1,2})_A = \alpha_{G_{1,2}(A)}^{2,3} \circ F_{2,3}(\alpha_A^{1,2})$
- $\alpha^{3,4} * \alpha^{2,3} : F_{3,4} \circ F_{2,3} \Rightarrow G_{3,4} \circ G_{2,3}$ and $(\alpha^{3,4} * \alpha^{2,3})_{G_{1,2}(A)} = \alpha_{G_{2,3}(G_{1,2}(A))}^{3,4} \circ F_{3,4}(\alpha_{G_{1,2}(A)}^{2,3})$.

Therefore

$$\begin{aligned} [\alpha^{3,4} * (\alpha^{2,3} * \alpha^{1,2})]_A &= \alpha_{G_{2,3} \circ G_{1,2}(A)}^{3,4} \circ F_{3,4}([\alpha^{2,3} * \alpha^{1,2}]_A) \\ &= \alpha_{G_{2,3}(G_{1,2}(A))}^{3,4} \circ F_{3,4}(\alpha_{G_{1,2}(A)}^{2,3} \circ F_{2,3}(\alpha_A^{1,2})) \\ &= \alpha_{G_{2,3}(G_{1,2}(A))}^{3,4} \circ [F_{3,4}(\alpha_{G_{1,2}(A)}^{2,3}) \circ F_{3,4}(F_{2,3}(\alpha_A^{1,2}))] \\ &= \alpha_{G_{2,3}(G_{1,2}(A))}^{3,4} \circ (F_{3,4}(\alpha_{G_{1,2}(A)}^{2,3}) \circ [F_{3,4} \circ F_{2,3}(\alpha_A^{1,2})]) \\ &= [\alpha_{G_{2,3}(G_{1,2}(A))}^{3,4} \circ F_{3,4}(\alpha_{G_{1,2}(A)}^{2,3})] \circ [F_{3,4} \circ F_{2,3}(\alpha_A^{1,2})] \\ &= [(\alpha^{3,4} * \alpha^{2,3})_{G_{1,2}(A)}] \circ [F_{3,4} \circ F_{2,3}(\alpha_A^{1,2})] \\ &= [(\alpha^{3,4} * \alpha^{2,3}) * \alpha^{1,2}]_A \end{aligned}$$

Since $A \in \text{Obj}(\mathcal{C}_1)$ is arbitrary, we have that for each $A \in \text{Obj}(\mathcal{C}_1)$, $[\alpha^{3,4} * (\alpha^{2,3} * \alpha^{1,2})]_A = [(\alpha^{3,4} * \alpha^{2,3}) * \alpha^{1,2}]_A$. Thus $\alpha^{3,4} * (\alpha^{2,3} * \alpha^{1,2}) = (\alpha^{3,4} * \alpha^{2,3}) * \alpha^{1,2}$. □

Exercise 5.1.0.8. Interchange law:

[wikipedia interchange law](#)

FINISH!!!

Exercise 5.1.0.9. $F : \mathcal{A} \rightarrow \mathcal{B}$ and $G : \mathcal{B} \rightarrow \mathcal{C}$. Then $\text{id}_G * \text{id}_F = \text{id}_{G \circ F}$

Proof. **FINISH!!!** □

Definition 5.1.0.10. Whiskering: Let $\mathcal{A}, \mathcal{B}, \mathcal{C}, \mathcal{D}$ be categories, $F, G : \mathcal{B} \rightarrow \mathcal{C}$, $H : \mathcal{A} \rightarrow \mathcal{B}$, $K : \mathcal{C} \rightarrow \mathcal{D}$ and $\alpha : F \Rightarrow G$. We define

- $\alpha H : F \circ H \Rightarrow G \circ H$ by $\alpha H := \alpha * \text{id}_H$.
- $K \alpha : K \circ F \Rightarrow K \circ G$ by $K \alpha := \text{id}_K * \alpha$

Exercise 5.1.0.11. Let $\mathcal{A}, \mathcal{B}, \mathcal{C}, \mathcal{D}$ be categories, $F, G : \mathcal{B} \rightarrow \mathcal{C}$, $H : \mathcal{A} \rightarrow \mathcal{B}$, $K : \mathcal{C} \rightarrow \mathcal{D}$ and $\alpha : F \Rightarrow G$. Then

1. for each $A \in \mathcal{A}$, $(\alpha H)_A = \alpha_{H(A)}$

2. for each $B \in \mathcal{B}$, $(K\alpha)_B = K(\alpha_B)$.

Proof.

1. For each $A \in \mathcal{A}$, we have that

$$\begin{aligned} (\alpha H)_A &= (\alpha * \text{id}_H)_A \\ &= \alpha_{H(A)} \circ F((\text{id}_H)_A) \\ &= \alpha_{H(A)} \circ F(\text{id}_{H(A)}) \\ &= \alpha_{H(A)} \circ \text{id}_{F(H(A))} \\ &= \alpha_{H(A)} \circ \text{id}_{F \circ H(A)} \\ &= \alpha_{H(A)}. \end{aligned}$$

2. For each $B \in \mathcal{B}$, we have that

$$\begin{aligned} (K\alpha)_B &= (\text{id}_K * \alpha)_B \\ &= ((\text{id}_K)_{G(B)}) \circ K(\alpha_B) \\ &= \text{id}_{K(G(B))} \circ K(\alpha_B) \\ &= \text{id}_{K \circ G(B)} \circ K(\alpha_B) \\ &= K(\alpha_B). \end{aligned}$$

□

Exercise 5.1.0.12. Let $\mathcal{A}_1, \mathcal{A}_2, \mathcal{A}_3$ be categories, $F_{1,2} : \mathcal{A}_1 \rightarrow \mathcal{A}_2$, $F_{2,3} : \mathcal{A}_2 \rightarrow \mathcal{A}_3$, $\alpha^{1,2} : F_{1,2} \rightarrow G_{1,2}$, $\alpha^{2,3} : F_{2,3} \Rightarrow G_{2,3}$. Then

1. $\alpha_2 * \alpha_1 = \alpha_2 * G_2 \circ F_2 \alpha_1$
2. $\alpha_2 * \alpha_1 = G_2 \alpha_2 \circ \alpha_1 F_1$

Proof.

- 1.
- 2.

FINISH!!!

□

Definition 5.1.0.13. We define $\text{Nat}(F, G) \times \text{Funct}(\mathcal{A}, \mathcal{B})$ $(\alpha, H) \mapsto \alpha H$ and similar for the other whisker map. **FINISH!!!**

5.2 Fundamental Examples

Definition 5.2.0.1. Let $\mathcal{C}, \mathcal{D}$ be categories and $F : \mathcal{C} \rightarrow \mathcal{D}$. We define the **identity natural transformation from F to F** , denoted $\text{id}_F : F \Rightarrow F$, by

$$(\text{id}_F)_A = \text{id}_{F(A)}^{\mathcal{D}}$$

Exercise 5.2.0.2. Let $\mathcal{C}, \mathcal{D}$ be categories and $F : \mathcal{C} \rightarrow \mathcal{D}$. Then $\text{id}_F : F \Rightarrow F$ is a natural transformation.

Proof.

1. Let $A \in \text{Obj}(\mathcal{C})$. Then

$$\begin{aligned} (\text{id}_F)_A &= \text{id}_{F(A)}^{\mathcal{D}} \\ &\in \text{Hom}_{\mathcal{D}}(F(A), F(A)) \end{aligned}$$

2. Let $A, B \in \text{Obj}(\mathcal{C})$, $f \in \text{Hom}_{\mathcal{C}}(A, B)$. Then

$$\begin{aligned} F(f) \circ_{\mathcal{D}} (\text{id}_F)_A &= F(f) \circ_{\mathcal{D}} \text{id}_{F(A)}^{\mathcal{D}} \\ &= F(f) \\ &= \text{id}_{F(B)}^{\mathcal{D}} \circ_{\mathcal{D}} F(f) \\ &= (\text{id}_F)_B \circ_{\mathcal{D}} F(f), \end{aligned}$$

i.e. the following diagram commutes:

$$\begin{array}{ccc} F(A) & \xrightarrow{(\text{id}_F)_A} & F(A) \\ F(f) \downarrow & & \downarrow F(f) \\ F(B) & \xrightarrow{(\text{id}_F)_B} & F(B). \end{array}$$

□

Exercise 5.2.0.3. Let $\mathcal{C}, \mathcal{D}$ be categories, $F, G : \mathcal{C} \rightarrow \mathcal{D}$ and $\alpha : F \Rightarrow G$. Then

1. $\text{id}_G \circ \alpha = \alpha$
2. $\alpha \circ \text{id}_F = \alpha$

Proof.

1. Let $A \in \text{Obj}(\mathcal{C})$. Then

$$\begin{aligned} (\text{id}_G \circ \alpha)_A &= (\text{id}_G)_A \circ \alpha_A \\ &= \text{id}_{G(A)} \circ \alpha_A \\ &= \alpha_A \end{aligned}$$

Since $A \in \text{Obj}(\mathcal{C})$ is arbitrary, $\text{id}_G \circ \alpha = \alpha$

2. Let $A \in \text{Obj}(\mathcal{C})$. Then

$$\begin{aligned} (\alpha \circ \text{id}_F)_A &= \alpha_A \circ (\text{id}_F)_A \\ &= \alpha_A \circ \text{id}_{F(A)} \\ &= \alpha_A \end{aligned}$$

Since $A \in \text{Obj}(\mathcal{C})$ is arbitrary, $\alpha \circ \text{id}_F = \alpha$.

□

talk about following example preimages and set comprehension vs post composition and predicates will need to talk about contravariant functors from $\text{preim}, \text{pred}$, denoted $F, G : \mathbf{Set} \rightarrow \mathbf{Set}$ respectively given by $F(X) := \mathcal{P}(X)$, $F(f) := f^{-1}$ and $G(X) := \{0, 1\}^X$, $G(f) := - \circ f$. This post/functor is interesting because it highlights the associativity of $(\text{NOT} \circ \phi) \circ f = \text{NOT} \circ (\phi \circ f)$ as a commutativity $f^{-1}(A^c) = (f^{-1}(A))^c$. That is if $X = f^{-1}$ and $Y = -^c$, then $XYA = YXA$ so that $XY = YX$. It begs the question whether or not commutativity can always be recast as associativity in some other context.

below is wild speculation and off topic question The author writes that associativity shows you that for any appropriate f, ϕ , post composing with NOT first and then pre composing with f vs precomposing with f first then poste composing with NOT are the same thing and in a sense orthogonal and independent, as in they do not affect the outcome. So is there a general view in which associativity (or via natural transformation commutativity) is a measure of independence?

Can we connect this with probability/physics in an information theoretic sense using the motivation/intuition that the kl divergence (or other notion of info loss) is a functor? i.e. $XY = YX$ or $x \circ (y \circ a) = (x \circ y) \circ a$ iff $\text{info_loss}[\text{joint}(X, Y), \otimes(X, Y)] = 0$ or some generalization? We would need a notion of joint for X and Y and a (not product because category of prob dists doesnt have products) tensor product, so maybe we need to consider monoidal cats)

what X, Y would correspond to classical/quantum probability? In the classical/quantum case, X, Y are marginal systems and $\text{joint}(X, Y)$ is the given bipartite system from which we isolate the marginals. In the classical case, $\text{info_loss}[\text{joint}(X, Y), \otimes(X, Y)] = 0$ iff $\text{joint}(X, Y) = \otimes(X, Y)$, that is, there is the two marginal states of the bipartite system contain as much information as the bipartite system itself, i.e. there is no interaction. Im not sure what the associated commuting/associative operations would be, but maybe we can get a clue from the above example or by looking further into similar examples, maybe toposes or other tools cover this?

If it were possible to find commutative/associative operations that capture two subsystems being independent and not having an interaction in the classical/quantum probabilistic sense, then this would seem to suggest that we would be able to recast interaction between subsystems in terms of some associative/commutative operations. How would this compare/contrast with the characterization of interaction via homological algebra using exactness?

5.3 Category of Functors

Exercise 5.3.0.1. Let $\mathcal{C}$ and $\mathcal{D}$ be categories, $F, G : \mathcal{C} \rightarrow \mathcal{D}$ and $\alpha : F \Rightarrow G$. If $\mathcal{C}$ is small, then α is a set.

Proof. Suppose that $\mathcal{C}$ is small. Then $\text{Obj}(\mathcal{C})$ is a set. Since $\alpha : \text{Obj}(\mathcal{C}) \rightarrow \text{Hom}_{\mathcal{D}}$, Axiom 1.2.0.11 implies that $\alpha(\text{Obj}(\mathcal{C}))$ is a set. Then $\text{Obj}(\mathcal{C}) \times \alpha(\text{Obj}(\mathcal{C}))$ is a set. Therefore $\mathcal{P}(\text{Obj}(\mathcal{C}) \times \alpha(\text{Obj}(\mathcal{C})))$ is a set. Since $\alpha \subset \text{Obj}(\mathcal{C}) \times \alpha(\text{Obj}(\mathcal{C}))$, we have that $\alpha \in \mathcal{P}(\text{Obj}(\mathcal{C}) \times \alpha(\text{Obj}(\mathcal{C})))$ which implies that α is a set. $\square$

Exercise 5.3.0.2. Let $\mathcal{C}$ and $\mathcal{D}$ be categories and $F, G : \mathcal{C} \rightarrow \mathcal{D}$. If $\mathcal{C}$ is small, then there exists a class A such that for each class α , $\alpha \in A$ iff $\alpha : F \Rightarrow G$.

Proof. Suppose that $\mathcal{C}$ is small. Define ϕ by

$$\phi(\alpha) : \alpha : F \Rightarrow G$$

Axiom 1.2.0.12 implies that there exists a class A such that for each set α , $\alpha \in A$ iff $\phi(\alpha)$. Let α be a class. Suppose that $\alpha \in A$. By Definition 1.2.0.1, α is a set. Since α is a set and $\alpha \in A$, we have that $\phi(\alpha)$. Hence $\alpha : F \Rightarrow G$. Conversely, suppose that $\alpha : F \Rightarrow G$. Since $\mathcal{C}$ is small, Exercise 5.3.0.1 implies that α is a set. Since $\phi(\alpha)$, we have that $\alpha \in A$. $\square$

Definition 5.3.0.3. Let $\mathcal{C}$ and $\mathcal{D}$ be categories. Suppose that $\mathcal{C}$ is small. We define the **functor category from $\mathcal{C}$ to $\mathcal{D}$** , denoted $\mathcal{D}^{\mathcal{C}}$, by

- $\text{Obj}(\mathcal{D}^{\mathcal{C}}) = \{F : \mathcal{C} \rightarrow \mathcal{D}\}$
- For $F, G \in \text{Obj}(\mathcal{D}^{\mathcal{C}})$, $\text{Hom}_{\mathcal{D}^{\mathcal{C}}}(F, G) = \{\alpha : \alpha : F \Rightarrow G\}$
- For $F, G, H \in \text{Obj}(\mathcal{D}^{\mathcal{C}})$, $\alpha \in \text{Hom}_{\mathcal{D}^{\mathcal{C}}}(F, G)$ and $\beta \in \text{Hom}_{\mathcal{D}^{\mathcal{C}}}(G, H)$, $\beta \circ_{\mathcal{D}^{\mathcal{C}}} \alpha = \beta \circ \alpha$

Exercise 5.3.0.4. Let $\mathcal{C}$ and $\mathcal{D}$ be categories. Suppose that $\mathcal{C}$ is small. Then $\mathcal{D}^{\mathcal{C}}$ is a category.

Proof. Exercise 5.1.0.3 implies that composition is well-defined. Exercise 5.1.0.6 implies that composition is associative. Exercise 5.2.0.2 and Exercise 5.2.0.3 imply the existence of identities. $\square$

Exercise 5.3.0.5. Let $\mathcal{C}$ and $\mathcal{D}$ be categories. Suppose that $\mathcal{C}$ and $\mathcal{D}$ are small. Then $\mathcal{D}^{\mathcal{C}}$ is small.

Proof. We have that

$$\begin{aligned} \mathcal{C} \text{ and } \mathcal{D} \text{ are small} &\iff \text{Obj}(\mathcal{C}), \text{Obj}(\mathcal{D}), \text{Hom}_{\mathcal{C}}, \text{Hom}_{\mathcal{D}} \in \text{Obj}(\mathbf{Set}) \\ &\iff \text{Obj}(\mathcal{D})^{\text{Obj}(\mathcal{C})}, \text{Hom}_{\mathcal{D}}^{\text{Hom}_{\mathcal{C}}} \in \text{Obj}(\mathbf{Set}) \\ &\iff \text{Obj}(\mathcal{D})^{\text{Obj}(\mathcal{C})} \times \text{Hom}_{\mathcal{D}}^{\text{Hom}_{\mathcal{C}}} \in \text{Obj}(\mathbf{Set}). \end{aligned}$$

Since $\mathcal{D}^{\mathcal{C}} \subset \text{Obj}(\mathcal{D})^{\text{Obj}(\mathcal{C})} \times \text{Hom}_{\mathcal{D}}^{\text{Hom}_{\mathcal{C}}}$ and $\mathcal{C}, \mathcal{D}$ are small, we have that $\mathcal{D}^{\mathcal{C}} \in \text{Obj}(\mathbf{Set})$. Thus $\mathcal{D}^{\mathcal{C}}$ is small. $\square$

5.4 Diagonal Functor

Definition 5.4.0.1. Let $\mathcal{C}, \mathcal{D}$ be categories, $X, Y \in \text{Obj}(\mathcal{D})$ and $f \in \text{Hom}_{\mathcal{D}}(X, Y)$. We define the **constant natural transformation on $\mathcal{C}$ at f** , denoted $\delta_f^{\mathcal{C}} : \Delta_X^{\mathcal{C}} \Rightarrow \Delta_Y^{\mathcal{C}}$, by

$$(\delta_f^{\mathcal{C}})_A = f$$

Exercise 5.4.0.2. Let $\mathcal{C}, \mathcal{D}$ be categories, $X, Y \in \text{Obj}(\mathcal{D})$ and $f \in \text{Hom}_{\mathcal{D}}(X, Y)$. Then $\delta_f^{\mathcal{C}} : \Delta_X^{\mathcal{C}} \Rightarrow \Delta_Y^{\mathcal{C}}$ is a natural transformation.

Proof.

1. By definition, for each $A \in \text{Obj}(\mathcal{C})$ $(\delta_f^{\mathcal{C}})_A \in \text{Hom}_{\mathcal{D}}(\Delta_X^{\mathcal{C}}(A), \Delta_Y^{\mathcal{C}}(A))$.
2. Let $A, B \in \text{Obj}(\mathcal{C})$ and $g \in \text{Hom}_{\mathcal{C}}(A, B)$. Then

$$\begin{aligned} \Delta_Y^{\mathcal{C}}(g) \circ (\delta_f^{\mathcal{C}})_A &= \text{id}_Y \circ f \\ &= f \\ &= f \circ \text{id}_X \\ &= (\delta_f^{\mathcal{C}})_B \circ \Delta_X^{\mathcal{C}}(g) \end{aligned}$$

i.e. the following diagrams commute:

$$\begin{array}{ccc} \Delta_X^{\mathcal{C}}(A) & \xrightarrow{(\delta_f^{\mathcal{C}})_A} & \Delta_Y^{\mathcal{C}}(A) \\ \Delta_X^{\mathcal{C}}(g) \downarrow & & \downarrow \Delta_Y^{\mathcal{C}}(g) \\ \Delta_X^{\mathcal{C}}(B) & \xrightarrow{(\delta_f^{\mathcal{C}})_B} & \Delta_Y^{\mathcal{C}}(B) \end{array} = \begin{array}{ccc} X & \xrightarrow{f} & Y \\ \text{id}_X \downarrow & & \downarrow \text{id}_Y \\ X & \xrightarrow{f} & Y \end{array}$$

So $\delta_f^{\mathcal{C}} : \Delta_X^{\mathcal{C}} \Rightarrow \Delta_Y^{\mathcal{C}}$ is a natural transformation. □

Exercise 5.4.0.3. Let $\mathcal{C}, \mathcal{D}$ be categories, $X, Y, Z \in \text{Obj}(\mathcal{D})$, $f \in \text{Hom}_{\mathcal{D}}(X, Y)$ and $g \in \text{Hom}_{\mathcal{D}}(Y, Z)$. Then $\delta_{g \circ f}^{\mathcal{C}} = \delta_g^{\mathcal{C}} \circ \delta_f^{\mathcal{C}}$.

Proof. Let $A \in \text{Obj}(\mathcal{C})$. Then

$$\begin{aligned} (\delta_{g \circ f}^{\mathcal{C}})_A &= g \circ f \\ &= (\delta_g^{\mathcal{C}})_A \circ (\delta_f^{\mathcal{C}})_A \\ &= (\delta_g^{\mathcal{C}} \circ \delta_f^{\mathcal{C}})_A \end{aligned}$$

Since $A \in \text{Obj}(\mathcal{C})$ is arbitrary, $\delta_{g \circ f}^{\mathcal{C}} = \delta_g^{\mathcal{C}} \circ \delta_f^{\mathcal{C}}$. □

Exercise 5.4.0.4. Let $\mathcal{C}, \mathcal{D}$ be categories and $X \in \text{Obj}(\mathcal{D})$. Then $\delta_{\text{id}_X}^{\mathcal{C}} = \text{id}_{\Delta_X^{\mathcal{C}}}$.

Proof. Let $A \in \text{Obj}(\mathcal{C})$. Then

$$\begin{aligned} (\delta_{\text{id}_X}^{\mathcal{C}})_A &= \text{id}_X \\ &= \text{id}_{\Delta_X^{\mathcal{C}}(A)} \\ &= (\text{id}_{\Delta_X^{\mathcal{C}}})_A \end{aligned}$$

Since $A \in \text{Obj}(\mathcal{C})$ is arbitrary, $\delta_{\text{id}_X}^{\mathcal{C}} = \text{id}_{\Delta_X^{\mathcal{C}}}$. □

Definition 5.4.0.5. Let $\mathcal{C}, \mathcal{D}$ be categories. Suppose that $\mathcal{C}$ is small. We define the **$\mathcal{C}$ -ary diagonal functor** on $\mathcal{D}$, denoted by $\Delta^{\mathcal{C}} : \mathcal{D} \rightarrow \mathcal{D}^{\mathcal{C}}$, by

- $\Delta^{\mathcal{C}}(X) = \Delta_X^{\mathcal{C}}$
- $\Delta^{\mathcal{C}}(f) = \delta_f^{\mathcal{C}}$

Exercise 5.4.0.6. Let $\mathcal{C}, \mathcal{D}$ be categories. Suppose that $\mathcal{C}$ is small. Then $\Delta^{\mathcal{C}} : \mathcal{D} \rightarrow \mathcal{D}^{\mathcal{C}}$ is a functor.

Proof.

1. Exercise 5.4.0.2 implies that for each $X, Y \in \text{Obj}(\mathcal{D})$ and $f \in \text{Hom}_{\mathcal{D}}(X, Y)$, $\Delta^{\mathcal{C}}(f) \in \text{Hom}_{\mathcal{D}^{\mathcal{C}}}(\Delta^{\mathcal{C}}(X), \Delta^{\mathcal{C}}(Y))$
2. Exercise 5.4.0.3 implies that for each $X, Y, Z \in \text{Obj}(\mathcal{D})$, $f \in \text{Hom}_{\mathcal{D}}(X, Y)$ and $g \in \text{Hom}_{\mathcal{D}}(Y, Z)$, $\Delta^{\mathcal{C}}(g \circ f) = \Delta^{\mathcal{C}}(g) \circ \Delta^{\mathcal{C}}(f)$
3. Exercise 5.4.0.4 implies that for each $X \in \text{Obj}(\mathcal{D})$, $\Delta^{\mathcal{C}}(\text{id}_X) = \text{id}_{\Delta^{\mathcal{C}}(X)}$

So $\Delta^{\mathcal{C}} : \mathcal{D} \rightarrow \mathcal{D}^{\mathcal{C}}$ is a functor. □

Exercise 5.4.0.7. Let $\mathcal{C}, \mathcal{D}$ be categories. Suppose that $\mathcal{C} \neq \mathbf{0}$ and $\mathcal{C}$ is small. Then $\Delta^{\mathcal{C}} : \mathcal{D} \rightarrow \mathcal{D}^{\mathcal{C}}$ is full and faithful **define full/faithful**.

Proof. Since $\mathcal{C} \neq \mathbf{0}$, $\text{Obj}(\mathcal{C}) \neq \emptyset$. Therefore there exists $C \in \text{Obj}(\mathcal{C})$.

• **faithful:**

Let $X, Y \in \text{Obj}(\mathcal{D})$ and $f, g \in \text{Hom}_{\mathcal{D}}(X, Y)$. Suppose that $\Delta^{\mathcal{C}}(f) = \Delta^{\mathcal{C}}(g)$. Then

$$\begin{aligned} \delta_f^{\mathcal{C}} &= \Delta^{\mathcal{C}}(f) \\ &= \Delta^{\mathcal{C}}(g) \\ &= \delta_g^{\mathcal{C}}. \end{aligned}$$

In particular,

$$\begin{aligned} f &= (\delta_f^{\mathcal{C}})_C \\ &= (\delta_g^{\mathcal{C}})_C \\ &= g. \end{aligned}$$

Since $f, g \in \text{Hom}_{\mathcal{D}}(X, Y)$ with $\Delta^{\mathcal{C}}(f) = \Delta^{\mathcal{C}}(g)$ are arbitrary, we have that for each $f, g \in \text{Hom}_{\mathcal{D}}(X, Y)$, $\Delta^{\mathcal{C}}(f) = \Delta^{\mathcal{C}}(g)$ implies that $f = g$. Thus $(\Delta^{\mathcal{C}})_{X,Y} : \text{Hom}_{\mathcal{D}}(X, Y) \rightarrow \text{Hom}_{\mathcal{D}^{\mathcal{C}}}(\Delta^{\mathcal{C}}(X), \Delta^{\mathcal{C}}(Y))$ is injective. Since $X, Y \in \text{Obj}(\mathcal{D})$ are arbitrary, we have that for each $X, Y \in \text{Obj}(\mathcal{D})$, $(\Delta^{\mathcal{C}})_{X,Y} : \text{Hom}_{\mathcal{D}}(X, Y) \rightarrow \text{Hom}_{\mathcal{D}^{\mathcal{C}}}(\Delta^{\mathcal{C}}(X), \Delta^{\mathcal{C}}(Y))$ is injective. Thus $\Delta^{\mathcal{C}}$ is faithful.

• **full:**

Let $X, Y \in \text{Obj}(\mathcal{D})$ and $\alpha \in \text{Hom}_{\mathcal{D}^{\mathcal{C}}}(\Delta^{\mathcal{C}}(X), \Delta^{\mathcal{C}}(Y))$. Then $\alpha : \Delta_X^{\mathcal{C}} \Rightarrow \Delta_Y^{\mathcal{C}}$. Therefore

– for each $A \in \text{Obj}(\mathcal{C})$,

$$\begin{aligned} \alpha_A &\in \text{Hom}_{\mathcal{D}}(\Delta_X^{\mathcal{C}}(A), \Delta_Y^{\mathcal{C}}(A)) \\ &= \text{Hom}_{\mathcal{D}}(X, Y), \end{aligned}$$

– for each $A, B \in \text{Obj}(\mathcal{C})$, and $f \in \text{Hom}_{\mathcal{C}}(A, B)$,

$$\begin{aligned} \alpha_A &= \text{id}_Y \circ \alpha_A \\ &= \Delta_Y^{\mathcal{C}}(f) \circ \alpha_Y \\ &= \alpha_B \circ \Delta_X^{\mathcal{C}}(f) \\ &= \alpha_B \circ \text{id}_X \\ &= \alpha_B, \end{aligned}$$

i.e. the following diagrams commute:

$$\begin{array}{ccc} \Delta_X^{\mathcal{C}}(A) & \xrightarrow{\alpha_A} & \Delta_Y^{\mathcal{C}}(A) \\ \Delta_X^{\mathcal{C}}(f) \downarrow & & \downarrow \Delta_Y^{\mathcal{C}}(f) \\ \Delta_X^{\mathcal{C}}(B) & \xrightarrow{\alpha_B} & \Delta_Y^{\mathcal{C}}(B) \end{array} = \begin{array}{ccc} X & \xrightarrow{\alpha_A} & Y \\ \text{id}_X \downarrow & & \downarrow \text{id}_Y \\ X & \xrightarrow{\alpha_B} & Y \end{array} = \begin{array}{ccc} & \alpha_A & \\ X & \curvearrowright & Y \\ & \alpha_B & \end{array}$$

Thus $\alpha_C \in \text{Hom}_{\mathcal{D}}(X, Y)$ and for each $A \in \text{Obj}(\mathcal{C})$,

$$\begin{aligned}\alpha_A &= \alpha_C \\ &= (\delta_{\alpha_C}^{\mathcal{C}})_A.\end{aligned}$$

Hence

$$\begin{aligned}\alpha &= \delta_{\alpha_C}^{\mathcal{C}} \\ &= \Delta^{\mathcal{C}}(\alpha_C).\end{aligned}$$

Since $\alpha \in \text{Hom}_{\mathcal{D}^{\mathcal{C}}}(\Delta^{\mathcal{C}}(X), \Delta^{\mathcal{C}}(Y))$ is arbitrary, we have that for each $\alpha \in \text{Hom}_{\mathcal{D}^{\mathcal{C}}}(\Delta^{\mathcal{C}}(X), \Delta^{\mathcal{C}}(Y))$, there exists $f \in \text{Hom}_{\mathcal{D}}(X, Y)$ such that $\Delta^{\mathcal{C}}(f) = \alpha$. Therefore $(\Delta^{\mathcal{C}})_{X,Y} : \text{Hom}_{\mathcal{D}}(X, Y) \rightarrow \text{Hom}_{\mathcal{D}^{\mathcal{C}}}(\Delta^{\mathcal{C}}(X), \Delta^{\mathcal{C}}(Y))$ is surjective. Since $X, Y \in \text{Obj}(\mathcal{D})$ are arbitrary, we have that for each $X, Y \in \text{Obj}(\mathcal{D})$, $(\Delta^{\mathcal{C}})_{X,Y} : \text{Hom}_{\mathcal{D}}(X, Y) \rightarrow \text{Hom}_{\mathcal{D}^{\mathcal{C}}}(\Delta^{\mathcal{C}}(X), \Delta^{\mathcal{C}}(Y))$ is surjective. Thus $\Delta^{\mathcal{C}}$ is full.

□

Chapter 6

Isomorphisms

6.1 Introduction

Definition 6.1.0.1. Let $\mathcal{C}$ be a category.

- Let $A \in \text{Obj}(\mathcal{C})$
 - Let $f \in \text{Hom}_{\mathcal{C}}(A, A)$. Then f is said to be an **endomorphism**,
 - We define $\text{End}_{\mathcal{C}}(A) := \text{Hom}_{\mathcal{C}}(A, A)$.
- We define $\text{End}_{\mathcal{C}} := \bigcup_{A \in \text{Obj}(\mathcal{C})} \text{End}_{\mathcal{C}}(A)$.

Definition 6.1.0.2. Let $\mathcal{C}$ be a category

- Let $A, B \in \text{Obj}(\mathcal{C})$.
 - Let $f \in \text{Hom}_{\mathcal{C}}(A, B)$. Then f is said to be an **isomorphism** if there exists $g \in \text{Hom}_{\mathcal{C}}(B, A)$ such that $g \circ f = \text{id}_A$ and $f \circ g = \text{id}_B$, i.e. the following diagram commutes:

$$\begin{array}{ccc} & g & \\ \text{id}_A \hookrightarrow & A & \xleftarrow{\quad} B \xrightarrow{\quad} \text{id}_B \\ & f & \end{array}$$

- We define $\text{Iso}_{\mathcal{C}}(A, B) := \{f \in \text{Hom}_{\mathcal{C}}(A, B) : f \text{ is an isomorphism}\}$
- We define $\text{Iso}_{\mathcal{C}} := \bigcup_{(A, B) \in \text{Obj}(\mathcal{C})^2} \text{Iso}_{\mathcal{C}}(A, B)$.

Definition 6.1.0.3. Let $\mathcal{C}$ be a category.

- Let $A \in \text{Obj}(\mathcal{C})$
 - Let $f \in \text{Iso}_{\mathcal{C}}(A, A)$. Then f is said to be an **automorphism**,
 - We define $\text{Aut}_{\mathcal{C}}(A) := \text{Iso}_{\mathcal{C}}(A, A)$.
- We define $\text{Aut}_{\mathcal{C}} := \bigcup_{A \in \text{Obj}(\mathcal{C})} \text{Aut}_{\mathcal{C}}(A)$.

Exercise 6.1.0.4. Uniqueness of Inverses:

Let $\mathcal{C}$ be a category, $A, B \in \text{Obj}(\mathcal{C})$ and $f \in \text{Iso}_{\mathcal{C}}(A, B)$. Then there exists a unique $g \in \text{Iso}_{\mathcal{C}}(B, A)$ such that $g \circ f = \text{id}_A$ and $f \circ g = \text{id}_B$.

Proof.

- **Existence:**

By definition, since f is an isomorphism, there exists $g \in \text{Hom}_{\mathcal{C}}(B, A)$ such that $g \circ f = \text{id}_A$ and $f \circ g = \text{id}_B$. By definition, $g \in \text{Iso}_{\mathcal{C}}(B, A)$.

- **Uniqueness:**

Let $g' \in \text{Iso}_{\mathcal{C}}(B, A)$. Suppose that $g' \circ f = \text{id}_A$, $f \circ g' = \text{id}_B$. Then

$$\begin{aligned} g' &= g' \circ \text{id}_B \\ &= g' \circ (f \circ g) \\ &= (g' \circ f) \circ g \\ &= \text{id}_A \circ g \\ &= g \end{aligned}$$

□

Definition 6.1.0.5. Let $\mathcal{C}$ be a category, $A, B \in \text{Obj}(\mathcal{C})$ and $f \in \text{Hom}_{\mathcal{C}}(A, B)$. Suppose that f is an isomorphism. We define the **inverse of f** , denoted $f^{-1} \in \text{Iso}_{\mathcal{C}}(B, A)$, to be the unique $g \in \text{Iso}_{\mathcal{C}}(B, A)$ such that $g \circ f = \text{id}_A$ and $f \circ g = \text{id}_B$.

Exercise 6.1.0.6. Let $\mathcal{C}$ be a category and $A \in \text{Obj}(\mathcal{C})$. Then $\text{id}_A \in \text{Aut}_{\mathcal{C}}(A)$ and $(\text{id}_A)^{-1} = \text{id}_A$.

Proof. Since $\text{id}_A \circ \text{id}_A = \text{id}_A$, we have that $\text{id}_A \in \text{Aut}_{\mathcal{C}}(A)$ and $(\text{id}_A)^{-1} = \text{id}_A$. □

Exercise 6.1.0.7. Let $\mathcal{C}$ be a category and $A, B \in \text{Obj}(\mathcal{C})$. Then for each $f \in \text{Iso}_{\mathcal{C}}(A, B)$, $(f^{-1})^{-1} = f$.

Proof. Let $f \in \text{Iso}_{\mathcal{C}}(A, B)$. By definition, $f \circ f^{-1} = \text{id}_B$ and $f^{-1} \circ f = \text{id}_A$. Hence $(f^{-1})^{-1} = f$. □

Exercise 6.1.0.8. Let $\mathcal{C}$ be a category and $A, B, C \in \text{Obj}(\mathcal{C})$. Then for each $f \in \text{Iso}_{\mathcal{C}}(A, B)$ and $g \in \text{Iso}_{\mathcal{C}}(B, C)$, $g \circ f \in \text{Iso}_{\mathcal{C}}(A, C)$ and $(g \circ f)^{-1} = f^{-1} \circ g^{-1}$.

Proof. Let $f \in \text{Iso}_{\mathcal{C}}(A, B)$ and $g \in \text{Iso}_{\mathcal{C}}(B, C)$. Then

$$\begin{aligned} (f^{-1} \circ g^{-1}) \circ (g \circ f) &= ((f^{-1} \circ g^{-1}) \circ g) \circ f \\ &= (f^{-1} \circ (g^{-1} \circ g)) \circ f \\ &= (f^{-1} \circ \text{id}_B) \circ f \\ &= f^{-1} \circ f \\ &= \text{id}_A \end{aligned}$$

and

$$\begin{aligned} (g \circ f) \circ (f^{-1} \circ g^{-1}) &= ((g \circ f) \circ f^{-1}) \circ g^{-1} \\ &= (g \circ (f \circ f^{-1})) \circ g^{-1} \\ &= (g \circ \text{id}_B) \circ g^{-1} \\ &= g \circ g^{-1} \\ &= \text{id}_C. \end{aligned}$$

Hence $g \circ f \in \text{Iso}_{\mathcal{C}}(A, C)$ and $(g \circ f)^{-1} = f^{-1} \circ g^{-1}$. □

Definition 6.1.0.9. Let $\mathcal{C}$ be a category.

- Let $A, B \in \text{Obj}(\mathcal{C})$. Then A is said to be **isomorphic** to B if $\text{Iso}_{\mathcal{C}}(A, B) \neq \emptyset$.
- We define the **isomorphism class relation on $\text{Obj}(\mathcal{C})$** , denoted $\cong_{\mathcal{C}} \in \mathcal{R}(A, B)$, by $A \cong_{\mathcal{C}} B$ iff A is isomorphic to B .

Note 6.1.0.10. When the context is clear, we may write $\cong$ in place of $\cong_{\mathcal{C}}$.

Exercise 6.1.0.11. Let $\mathcal{C}$ be a category. Then $\cong \in \text{Equiv}(\text{Obj}(\mathcal{C}))$ **does $\mathcal{C}$ need to be small or just locally small?**

Proof.

1. **reflexivity:**

Let $A \in \text{Obj}(\mathcal{C})$. Exercise 6.1.0.6 implies that $\text{id}_A \in \text{Aut}(A)$. Thus $A \cong A$. Since $A \in \text{Obj}(\mathcal{C})$ is arbitrary, we have that for each $A \in \text{Obj}(\mathcal{C})$, $A \cong A$ and thus $\cong$ is reflexive.

2. **symmetry:**

Let $A, B \in \text{Obj}(\mathcal{C})$. Suppose that $A \cong B$. Then there exists $f \in \text{Iso}_{\mathcal{C}}(A, B)$. Exercise 6.1.0.7 implies that $f^{-1} \in \text{Iso}_{\mathcal{C}}(B, A)$. Hence $B \cong A$. Since $A, B \in \text{Obj}(\mathcal{C})$ with $A \cong B$ are arbitrary, we have that for each $A, B \in \text{Obj}(\mathcal{C})$, $A \cong B$ implies that $B \cong A$ and thus $\cong$ is reflexive.

3. **transitivity:**

Let $A, B, C \in \text{Obj}(\mathcal{C})$. Suppose that $A \cong B$ and $B \cong C$. Then there exist $f \in \text{Iso}_{\mathcal{C}}(A, B)$ and $g \in \text{Iso}_{\mathcal{C}}(B, C)$. Exercise 6.1.0.8 implies that $g \circ f \in \text{Iso}_{\mathcal{C}}(A, C)$. Thus $A \cong C$. Since $A, B, C \in \text{Obj}(\mathcal{C})$ with $A \cong B$ and $B \cong C$ are arbitrary, we have that for each $A, B, C \in \text{Obj}(\mathcal{C})$, $A \cong B$ and $B \cong C$ implies that $A \cong C$ and thus $\cong$ is transitive.

Thus $\cong \in \text{Equiv}(\text{Obj}(\mathcal{C}))$. □

Exercise 6.1.0.12. Let $\mathcal{C}$ be a category and $\mathcal{E} \subset \text{Obj}(\mathcal{C})$. Then for each $A, B \in \mathcal{E}$, $\# \text{Hom}_{\mathcal{C}}(A, B) = 1$ iff for each $A, B \in \mathcal{E}$, there exists $f \in \text{Iso}_{\mathcal{C}}(A, B)$ such that $\text{Hom}_{\mathcal{C}}(A, B) = \{f\}$.

Proof.

• ($\implies$):

Suppose that for each $A, B \in \mathcal{E}$, $\# \text{Hom}_{\mathcal{C}}(A, B) = 1$. Let $A, B \in \mathcal{E}$. By assumption, $\text{Hom}_{\mathcal{C}}(A, A) = \{\text{id}_A\}$, $\text{Hom}_{\mathcal{C}}(B, B) = \{\text{id}_B\}$ and there exist $f \in \text{Hom}_{\mathcal{C}}(A, B)$ and $g \in \text{Hom}_{\mathcal{C}}(B, A)$ such that $\text{Hom}_{\mathcal{C}}(A, B) = \{f\}$ and $\text{Hom}_{\mathcal{C}}(B, A) = \{g\}$. Since

$$\begin{aligned} g \circ f &\in \text{Hom}_{\mathcal{C}}(A, A) \\ &= \{\text{id}_A\}, \end{aligned}$$

we have that $g \circ f = \text{id}_A$. Similarly, $f \circ g = \text{id}_B$, i.e. the following diagram commutes:

$$\begin{array}{ccc} & & g \\ & \curvearrowright & \\ \text{id}_A \hookrightarrow & A & \xrightarrow{f} B \\ & \curvearrowleft & \\ & & \text{id}_B \end{array}$$

Hence $f \in \text{Iso}_{\mathcal{C}}(A, B)$. Since $A, B \in \mathcal{E}$ are arbitrary, we have that for each $A, B \in \mathcal{E}$, there exists $f \in \text{Iso}_{\mathcal{C}}(A, B)$ such that $\text{Hom}_{\mathcal{C}}(A, B) = \{f\}$.

• ($\impliedby$):

Clearly if for each $A, B \in \mathcal{E}$, there exists $f \in \text{Iso}_{\mathcal{C}}(A, B)$ such that $\text{Hom}_{\mathcal{C}}(A, B) = \{f\}$, then for each $A, B \in \mathcal{E}$, $\# \text{Hom}_{\mathcal{C}}(A, B) = 1$. □

6.2 Isomorphisms between Categories

Exercise 6.2.0.1. Let $\mathcal{C}, \mathcal{D} \in \text{Obj}(\mathbf{Cat})$ and $F \in \text{Hom}_{\mathbf{Cat}}(\mathcal{C}, \mathcal{D})$. Write $F = (F_0, F_1)$. Then $F \in \text{Iso}_{\mathbf{Cat}}$ iff $F_0, F_1 \in \text{Iso}_{\mathbf{Set}}$

Proof.

• ($\implies$):

Suppose that $F \in \text{Iso}_{\mathbf{Cat}}$. Then there exists $G \in \text{Hom}_{\mathbf{Cat}}(\mathcal{D}, \mathcal{C})$ such that $G \circ_{\mathbf{Cat}} F = \text{id}_{\mathcal{C}}$ and $F \circ_{\mathbf{Cat}} G = \text{id}_{\mathcal{D}}$. Write $G = (G_0, G_1)$. By Definition 4.1.0.4, we have that $G_0 \circ_{\mathbf{Set}} F_0 = \text{id}_{\text{Obj}(\mathcal{C})}^{\mathbf{Set}}$, $F_0 \circ_{\mathbf{Set}} G_0 = \text{id}_{\text{Obj}(\mathcal{D})}^{\mathbf{Set}}$, $G_1 \circ_{\mathbf{Set}} F_1 = \text{id}_{\text{Hom}_{\mathcal{C}}}^{\mathbf{Set}}$ and $F_1 \circ_{\mathbf{Set}} G_1 = \text{id}_{\text{Hom}_{\mathcal{D}}}^{\mathbf{Set}}$. Hence $F_0, G_0 \in \text{Iso}_{\mathbf{Set}}$.

• ($\impliedby$):

Suppose that $F_0, F_1 \in \text{Iso}_{\mathbf{Set}}$. Define $G_0 := F_0^{-1}$ and $G_1 := F_1^{-1}$.

1. Let $A', B' \in \text{Obj}(\mathcal{D})$ and $f' \in \text{Hom}_{\mathcal{D}}(A', B')$. Define $f \in \text{Hom}_{\mathcal{C}}$ and $A, B \in \text{Obj}(\mathcal{C})$ by $f := G_1(f')$, $A := \text{dom}(f)$ and $B := \text{cod}(f)$. Since $F \in \text{Hom}_{\mathbf{Cat}}(\mathcal{C}, \mathcal{D})$, we have that

$$\begin{aligned} f' &= F_1(f) \\ &\in \text{Hom}_{\mathcal{D}}(F_0(A), F_0(B)). \end{aligned}$$

Since $f' \in \text{Hom}_{\mathcal{D}}(A', B')$, we have that

$$\begin{aligned} A' &= \text{dom}(f') \\ &= F_0(A). \end{aligned}$$

Similarly, $B' = F_0(B)$. Hence $G_0(A') = A$, $G_0(B') = B$ and

$$\begin{aligned} G_1(f') &= G_1(F_1(f)) \\ &= f \\ &\in \text{Hom}_{\mathcal{C}}(A, B) \\ &= \text{Hom}_{\mathcal{C}}(A, B) \\ &= \text{Hom}_{\mathcal{C}}(G_0(A'), G_0(B')). \end{aligned}$$

2. Let $A', B', C' \in \text{Obj}(\mathcal{D})$, $f' \in \text{Hom}_{\mathcal{D}}(A', B')$ and $g' \in \text{Hom}_{\mathcal{D}}(B', C')$. Define $f, g \in \text{Hom}_{\mathcal{C}}$ by $f := G_1(f')$ and $g := G_1(g')$. Then $F_1(f) = f'$ and $F_1(g) = g'$. Since $F \in \text{Hom}_{\mathbf{Cat}}(\mathcal{C}, \mathcal{D})$, we have that

$$\begin{aligned} G_1(g' \circ_{\mathcal{D}} f') &= G_1(F_1(g) \circ_{\mathcal{D}} F_1(f)) \\ &= G_1(F_1(g \circ_{\mathcal{C}} f)) \\ &= g \circ_{\mathcal{C}} f \\ &= G_1(g') \circ_{\mathcal{C}} G_1(f'). \end{aligned}$$

3. Let $A' \in \text{Obj}(\mathcal{D})$. Define $A \in \text{Obj}(\mathcal{C})$ by $A := G_0(A')$. Then $F_0(A) = A'$ and since $F \in \text{Hom}_{\mathbf{Cat}}(\mathcal{C}, \mathcal{D})$, we have that

$$\begin{aligned} G_1(\text{id}_{A'}^{\mathcal{D}}) &= G_1(\text{id}_{F_0(A)}^{\mathcal{D}}) \\ &= G_1(F_1(\text{id}_A^{\mathcal{C}})) \\ &= \text{id}_A^{\mathcal{C}} \\ &= \text{id}_{G_0(A')}^{\mathcal{C}}. \end{aligned}$$

Hence $G \in \text{Hom}_{\mathbf{Cat}}(\mathcal{D}, \mathcal{C})$. By construction, $G \circ_{\mathbf{Cat}} F = \text{id}_{\mathcal{C}}$ and $F \circ_{\mathbf{Cat}} G = \text{id}_{\mathcal{D}}$. Thus $F \in \text{Iso}_{\mathbf{Cat}}$. □

6.3 Natural Isomorphisms

Definition 6.3.0.1. Let $\mathcal{C}$ and $\mathcal{D}$ be categories, $F, G : \mathcal{C} \rightarrow \mathcal{D}$ and $\alpha : F \Rightarrow G$. Then α is said to be a **natural isomorphism** if for each $A \in \text{Obj}(\mathcal{C})$, $\alpha_A \in \text{Iso}_{\mathcal{D}}(F(A), G(A))$.

Definition 6.3.0.2. Let $\mathcal{C}$ and $\mathcal{D}$ be categories, $F, G : \mathcal{C} \rightarrow \mathcal{D}$ and $\alpha : F \Rightarrow G$. Suppose that α is a natural isomorphism. We define $\alpha^{-1} : G \Rightarrow F$ by $(\alpha^{-1})_A = \alpha_A^{-1}$.

Exercise 6.3.0.3. Let $\mathcal{C}$ and $\mathcal{D}$ be categories, $F, G : \mathcal{C} \rightarrow \mathcal{D}$ and $\alpha : F \Rightarrow G$. Suppose that α is a natural isomorphism. Then $\alpha^{-1} : G \Rightarrow F$ is a natural transformation

Proof.

1. Let $A \in \text{Obj}(\mathcal{C})$. Since $\alpha_A \in \text{Hom}_{\mathcal{D}}(F(A), G(A))$, we have that

$$\begin{aligned} (\alpha^{-1})_A &= \alpha_A^{-1} \\ &\in \text{Hom}_{\mathcal{D}}(G(A), F(A)) \end{aligned}$$

2. Let $A, B \in \text{Obj}(\mathcal{C})$ and $f \in \text{Hom}_{\mathcal{C}}(A, B)$. Since $G(f) \circ \alpha_A = \alpha_B \circ F(f)$, i.e. the following diagram commutes:

$$\begin{array}{ccc} F(A) & \xrightarrow{\alpha_A} & G(A) \\ F(f) \downarrow & & \downarrow G(f) \\ F(B) & \xrightarrow{\alpha_B} & G(B) \end{array}$$

we have that

$$\begin{aligned} F(f) \circ (\alpha^{-1})_A &= F(f) \circ \alpha_A^{-1} \\ &= \text{id}_{F(B)} \circ (F(f) \circ \alpha_A^{-1}) \\ &= (\alpha_B^{-1} \circ \alpha_B) \circ (F(f) \circ \alpha_A^{-1}) \\ &= \alpha_B^{-1} \circ (\alpha_B \circ (F(f) \circ \alpha_A^{-1})) \\ &= \alpha_B^{-1} \circ ((\alpha_B \circ F(f)) \circ \alpha_A^{-1}) \\ &= \alpha_B^{-1} \circ ((G(f) \circ \alpha_A) \circ \alpha_A^{-1}) \\ &= \alpha_B^{-1} \circ (G(f) \circ (\alpha_A \circ \alpha_A^{-1})) \\ &= \alpha_B^{-1} \circ (G(f) \circ \text{id}_{G(A)}) \\ &= \alpha_B^{-1} \circ G(f) \\ &= (\alpha^{-1})_B \circ G(f) \end{aligned}$$

i.e. the following diagram commutes:

$$\begin{array}{ccc} G(A) & \xrightarrow{(\alpha^{-1})_A} & F(A) \\ G(f) \downarrow & & \downarrow F(f) \\ G(B) & \xrightarrow{(\alpha^{-1})_B} & F(B) \end{array}$$

So $\alpha^{-1} : G \Rightarrow F$.

□

Exercise 6.3.0.4. Let $\mathcal{C}$ and $\mathcal{D}$ be categories, $F, G : \mathcal{C} \rightarrow \mathcal{D}$ and $\alpha : F \Rightarrow G$. Suppose that α is a natural isomorphism. Then $\alpha^{-1} \circ \alpha = \text{id}_F$ and $\alpha \circ \alpha^{-1} = \text{id}_G$.

Proof. Let $A \in \text{Obj}(\mathcal{C})$. Then

$$\begin{aligned} (\alpha^{-1} \circ \alpha)_A &= (\alpha^{-1})_A \circ \alpha_A \\ &= \alpha_A^{-1} \circ \alpha_A \\ &= \text{id}_{F(A)} \\ &= (\text{id}_F)_A \end{aligned}$$

and

$$\begin{aligned} (\alpha \circ \alpha^{-1})_A &= \alpha_A \circ (\alpha^{-1})_A \\ &= \alpha_A \circ \alpha_A^{-1} \\ &= \text{id}_{G(A)} \\ &= (\text{id}_G)_A \end{aligned}$$

Since $A \in \text{Obj}(\mathcal{C})$ is arbitrary, $\alpha^{-1} \circ \alpha = \text{id}_F$ and $\alpha \circ \alpha^{-1} = \text{id}_G$.

□

Exercise 6.3.0.5. Let $\mathcal{C}$ and $\mathcal{D}$ be categories. Suppose that $\mathcal{C}$ is small. Let $F, G \in \text{Obj}(\mathcal{D}^{\mathcal{C}})$ and $\alpha \in \text{Hom}_{\mathcal{D}^{\mathcal{C}}}(F, G)$. Then α is a natural isomorphism iff $\alpha \in \text{Iso}_{\mathcal{D}^{\mathcal{C}}}(F, G)$.

Proof.

- ($\implies$):

Suppose that α is a natural isomorphism. Exercise 6.3.0.4 implies that $\alpha \in \text{Iso}_{\mathcal{D}^{\mathcal{C}}}(F, G)$.

- ($\impliedby$):

Suppose that $\alpha \in \text{Iso}_{\mathcal{D}^{\mathcal{C}}}(F, G)$. Let $A \in \text{Obj}(\mathcal{C})$. Then

$$\begin{aligned}\alpha_A \circ (\alpha^{-1})_A &= (\alpha \circ \alpha^{-1})_A \\ &= (\text{id}_G)_A \\ &= \text{id}_{G(A)}\end{aligned}$$

and similarly, $\alpha_A^{-1} \circ \alpha_A = \text{id}_{F(A)}$. Thus $\alpha_A \in \text{Iso}_{\mathcal{D}}(F(A), G(A))$. Since $A \in \text{Obj}(\mathcal{C})$ is arbitrary, we have that for each $A \in \text{Obj}(\mathcal{C})$, $\alpha_A \in \text{Iso}_{\mathcal{D}}(F(A), G(A))$. By definition, α is a natural isomorphism.

□

Part II

Fundamental Structures

Chapter 7

Introduction

7.1 Initial and Terminal Objects

Definition 7.1.0.1. Let $\mathcal{C}$ be a category.

- Let $X \in \text{Obj}(\mathcal{C})$. Then
 - X is said to be **$\mathcal{C}$ -terminal** if for each $A \in \text{Obj}(\mathcal{C})$, there exists $f \in \text{Hom}_{\mathcal{C}}(A, X)$ such that $\text{Hom}_{\mathcal{C}}(A, X) = \{f\}$.
 - X is said to be **$\mathcal{C}$ -coterminial** or **$\mathcal{C}$ -initial** if for each $A \in \text{Obj}(\mathcal{C})$, there exists $f \in \text{Hom}_{\mathcal{C}}(X, A)$ such that $\text{Hom}_{\mathcal{C}}(X, A) = \{f\}$.
- We define
 - $\text{Term}_{\mathcal{C}} := \{X \in \text{Obj}(\mathcal{C}) : X \text{ is } \mathcal{C}\text{-terminal}\}$.
 - $\text{Coterm}_{\mathcal{C}} := \{X \in \text{Obj}(\mathcal{C}) : X \text{ is } \mathcal{C}\text{-coterminial}\}$.

Note 7.1.0.2. When the context is clear, we say “terminal” in place of “ $\mathcal{C}$ -terminal” and “coterminial” (or “initial”) in place of “ $\mathcal{C}$ -coterminial” (or “ $\mathcal{C}$ -initial”) respectively.

Exercise 7.1.0.3. Let $\mathcal{C}$ be a category. Then

1. $\text{Term}_{\mathcal{C}} = \text{Coterm}_{\mathcal{C}^{\text{op}}}$,
2. $\text{Coterm}_{\mathcal{C}} = \text{Term}_{\mathcal{C}^{\text{op}}}$.

Proof.

1. • Let $X \in \text{Term}_{\mathcal{C}}$ and $A \in \text{Obj}(\mathcal{C}^{\text{op}})$. Then $A \in \text{Obj}(\mathcal{C})$. Since $X \in \text{Term}_{\mathcal{C}}$, there exists $f \in \text{Hom}_{\mathcal{C}}(A, X)$ such that $\text{Hom}_{\mathcal{C}}(A, X) = \{f\}$. Then

$$\begin{aligned}\text{Hom}_{\mathcal{C}^{\text{op}}}(X, A) &= \text{Hom}_{\mathcal{C}}(A, X) \\ &= \{f\}.\end{aligned}$$

Since $A \in \text{Obj}(\mathcal{C}^{\text{op}})$ is arbitrary, we have that for each $A \in \text{Obj}(\mathcal{C}^{\text{op}})$, there exists $f \in \text{Hom}_{\mathcal{C}^{\text{op}}}(X, A)$ such that $\text{Hom}_{\mathcal{C}^{\text{op}}}(X, A) = \{f\}$. Thus $X \in \text{Coterm}_{\mathcal{C}^{\text{op}}}$. Since $X \in \text{Term}_{\mathcal{C}}$ is arbitrary, we have that for each $X \in \text{Term}_{\mathcal{C}}$, $X \in \text{Coterm}_{\mathcal{C}^{\text{op}}}$. Thus $\text{Term}_{\mathcal{C}} \subset \text{Coterm}_{\mathcal{C}^{\text{op}}}$.

- ($\Leftarrow$):
Suppose that $X \in \text{Coterm}_{\mathcal{C}^{\text{op}}}$. Let $A \in \text{Obj}(\mathcal{C})$. Then $A \in \text{Obj}(\mathcal{C}^{\text{op}})$. Since $X \in \text{Coterm}_{\mathcal{C}^{\text{op}}}$, there exists $f \in \text{Hom}_{\mathcal{C}^{\text{op}}}(X, A)$ such that $\text{Hom}_{\mathcal{C}^{\text{op}}}(X, A) = \{f\}$. Then

$$\begin{aligned}\text{Hom}_{\mathcal{C}}(A, X) &= \text{Hom}_{\mathcal{C}^{\text{op}}}(X, A) \\ &= \{f\}.\end{aligned}$$

Since $A \in \text{Obj}(\mathcal{C})$ is arbitrary, we have that for each $A \in \text{Obj}(\mathcal{C})$, there exists $f \in \text{Hom}_{\mathcal{C}}(A, X)$ such that $\text{Hom}_{\mathcal{C}}(A, X) = \{f\}$. Thus $X \in \text{Term}_{\mathcal{C}}$. Since $X \in \text{Coterm}_{\mathcal{C}^{\text{op}}}$ is arbitrary, we have that for each $X \in \text{Coterm}_{\mathcal{C}^{\text{op}}}$, $X \in \text{Term}_{\mathcal{C}}$. Thus $\text{Coterm}_{\mathcal{C}^{\text{op}}} \subset \text{Term}_{\mathcal{C}}$.

Since $\text{Term}_{\mathcal{C}} \subset \text{Coterm}_{\mathcal{C}^{\text{op}}}$ and $\text{Coterm}_{\mathcal{C}^{\text{op}}} \subset \text{Term}_{\mathcal{C}}$, we have that $\text{Term}_{\mathcal{C}} = \text{Coterm}_{\mathcal{C}^{\text{op}}}$.

2. **similar to prev part.**

□

Exercise 7.1.0.4. Let $\mathcal{C}$ be a category. Then

1. for each $X, Y \in \text{Term}_{\mathcal{C}}$, there exists $f \in \text{Iso}_{\mathcal{C}}(X, Y)$ such that $\text{Hom}_{\mathcal{C}}(X, Y) = \{f\}$.
2. for each $X, Y \in \text{Coterm}_{\mathcal{C}}$, there exists $f \in \text{Iso}_{\mathcal{C}}(X, Y)$ such that $\text{Hom}_{\mathcal{C}}(X, Y) = \{f\}$.

Proof.

1. By Definition 7.1.0.1, for each $X, Y \in \text{Term}_{\mathcal{C}}$, $\#\text{Hom}_{\mathcal{C}}(X, Y) = 1$. Exercise 6.1.0.12 then implies that for each $X, Y \in \text{Term}_{\mathcal{C}}$, there exists $f \in \text{Iso}_{\mathcal{C}}(X, Y)$ such that $\text{Hom}_{\mathcal{C}}(X, Y) = \{f\}$.

2. **Similar to prev part**

□

Note 7.1.0.5. In particular,

- for each $1, 1' \in \text{Term}_{\mathcal{C}}$ are terminal, then $1 \cong_{\mathcal{C}} 1'$, $\text{Hom}_{\mathcal{C}}(1, 1') = \text{Iso}_{\mathcal{C}}(1, 1')$ and $\#\text{Hom}_{\mathcal{C}}(0, 0') = 1$.
- for each $0, 0' \in \text{Coterm}_{\mathcal{C}}$, then $0 \cong_{\mathcal{C}} 0'$, $\text{Hom}_{\mathcal{C}}(0, 0') = \text{Iso}_{\mathcal{C}}(0, 0')$ and $\#\text{Hom}_{\mathcal{C}}(0, 0') = 1$

FINISH!!! maybe rework ordering terminal first, then coterminal

Exercise 7.1.0.6. Let $X, Y \in \text{Obj}(\mathbf{Set})$. Then

1. X is initial iff $X = \emptyset$.
2. Y is terminal iff $Y \cong_{\mathbf{Set}} \{\emptyset\}$.

Proof.

1. • ($\implies$):
Suppose that X is initial. Since $\text{Hom}_{\mathbf{Set}}(X, \{0, 1\}) \cong_{\mathbf{Set}} \mathcal{P}(X)$, we have that

$$\begin{aligned} \#\text{Hom}_{\mathbf{Set}}(X, \{0, 1\}) = 1 &\iff \#\mathcal{P}(X) = 1 \\ &\iff \mathcal{P}(X) = \{\emptyset\} \\ &\iff X = \emptyset. \end{aligned}$$

Since X is initial, there exists $f \in \text{Hom}_{\mathbf{Set}}(X, \{0, 1\})$ such that $\text{Hom}_{\mathbf{Set}}(X, \{0, 1\}) = \{f\}$. Thus $X = \emptyset$.

- ($\impliedby$):
Suppose that $X = \emptyset$. Let $A \in \text{Obj}(\mathbf{Set})$.
– We note that for each $f \in \text{Hom}_{\mathbf{Set}}$,

$$\begin{aligned} f \in \text{Hom}_{\mathbf{Set}}(X, A) &\iff f \in \text{Hom}_{\mathbf{Set}}(\emptyset, A) \\ &\implies f \subset \emptyset \times A \\ &= \iff f \subset \emptyset \\ &\iff f = \emptyset. \end{aligned}$$

Thus for each $f \in \text{Hom}_{\mathbf{Set}}(X, A)$, $f \in \{\emptyset\}$. Hence $\text{Hom}_{\mathbf{Set}}(X, A) \subset \{\emptyset\}$.

– Since

$$\begin{aligned}\emptyset &\in \text{Hom}_{\mathbf{Set}}(\emptyset, A) \\ &= \text{Hom}_{\mathbf{Set}}(X, A),\end{aligned}$$

we have that $\{\emptyset\} \subset \text{Hom}_{\mathbf{Set}}(X, A)$.

Thus $\text{Hom}_{\mathbf{Set}}(X, A) = \{\emptyset\}$. Therefore X is initial.

2. • ($\implies$):

Suppose that Y is terminal. Since $\#\text{Hom}_{\mathbf{Set}}(\{\emptyset\}, Y) = \#Y$, we have that

$$\begin{aligned}\#\text{Hom}_{\mathbf{Set}}(\{\emptyset\}, Y) = 1 &\iff \#Y = 1 \\ &\iff Y \cong_{\mathbf{Set}} \{\emptyset\}.\end{aligned}$$

Since Y is terminal, there exists $f \in \text{Hom}_{\mathbf{Set}}(\{\emptyset\}, Y)$ such that $\text{Hom}_{\mathbf{Set}}(\{\emptyset\}, Y) = \{f\}$. Thus $Y \cong_{\mathbf{Set}} \{\emptyset\}$.

• ($\impliedby$):

Suppose that $Y \cong_{\mathbf{Set}} \{\emptyset\}$. Then there exists $y \in Y$ such that $Y = \{y\}$. Let $A \in \text{Obj}(\mathbf{Set})$. Define $f_0 \in \text{Hom}_{\mathbf{Set}}(A, Y)$ by $f_0(a) := y$. Let $f \in \text{Hom}_{\mathbf{Set}}(A, Y)$. Let $a \in A$. Since

$$\begin{aligned}f(a) &\in Y \\ &= \{y\},\end{aligned}$$

we have that $f(a) = y$. Since $a \in A$ is arbitrary, we have that for each $a \in A$,

$$\begin{aligned}f(a) &= y \\ &= f_0(a).\end{aligned}$$

Thus $f = f_0$. Since $f \in \text{Hom}_{\mathbf{Set}}(A, Y)$ is arbitrary, we have that for each $f \in \text{Hom}_{\mathbf{Set}}(A, Y)$, $f = f_0$. Hence $\text{Hom}_{\mathbf{Set}}(A, Y) = \{f_0\}$. Since $A \in \text{Obj}(\mathbf{Set})$ is arbitrary, we have that for each $A \in \text{Obj}(\mathbf{Set})$, there exists $f_0 \in \text{Hom}_{\mathbf{Set}}(A, Y)$ such that $\text{Hom}_{\mathbf{Set}}(A, Y) = \{f_0\}$. Thus Y is terminal. □

Exercise 7.1.0.7. Let $\mathcal{C}$ be a category. Suppose that $\mathcal{C}$ is locally small. Let $X \in \text{Obj}(\mathcal{C})$. Then

1. X is $\mathcal{C}$ -initial iff for each $A \in \text{Obj}(\mathcal{C})$, $\text{Hom}_{\mathcal{C}}(X, A)$ is **Set**-terminal.
2. X is $\mathcal{C}$ -terminal iff for each $A \in \text{Obj}(\mathcal{C})$, $\text{Hom}_{\mathcal{C}}(A, X)$ is **Set**-terminal.

Proof.

1. We have that

$$\begin{aligned}X \text{ is } \mathcal{C}\text{-initial} &\iff \text{for each } A \in \text{Obj}(\mathcal{C}), \#\text{Hom}_{\mathcal{C}}(X, A) = 1 \\ &\iff \text{for each } A \in \text{Obj}(\mathcal{C}), \#\text{Hom}_{\mathcal{C}}(X, A) \cong_{\mathbf{Set}} \{\emptyset\} \\ &\iff \text{for each } A \in \text{Obj}(\mathcal{C}), \#\text{Hom}_{\mathcal{C}}(X, A) \text{ is } \mathbf{Set}\text{-terminal,} \quad (\text{by Exercise 7.1.0.6}).\end{aligned}$$

2. We have that

$$\begin{aligned}X \text{ is } \mathcal{C}\text{-terminal} &\iff X \text{ is } \mathcal{C}^{\text{op}}\text{-initial,} \quad (\text{by Exercise 7.1.0.3}) \\ &\iff \text{for each } A \in \text{Obj}(\mathcal{C}^{\text{op}}), \text{Hom}_{\mathcal{C}^{\text{op}}}(X, A) \text{ is } \mathbf{Set}\text{-terminal,} \quad (\text{by ref prev part}) \\ &\iff \text{for each } A \in \text{Obj}(\mathcal{C}), \text{Hom}_{\mathcal{C}}(A, X) \text{ is } \mathbf{Set}\text{-terminal}\end{aligned}$$

□

Exercise 7.1.0.8. We have that

1. $\mathbf{0}$ is **Cat**-initial.
2. $\mathbf{1}$ is **Cat**-terminal.

Proof.

1. Let $\mathcal{C} \in \text{Obj}(\mathbf{Cat})$. It is clear that $\text{Hom}_{\mathbf{Cat}}(\mathbf{0}, \mathcal{C}) = \{E_{\mathcal{C}}\}$. Hence $\mathbf{0}$ is **Cat**-initial. **Fill out like below ex**
2. Let $\mathcal{C} \in \text{Obj}(\mathbf{Cat})$. Let $F : \mathcal{C} \rightarrow \mathbf{1}$. Since $\text{Obj}(\mathbf{1}) = \{*\}$ and $\text{Hom}_{\mathbf{1}}(*, *) = \{\text{id}_*^{\mathbf{1}}\}$, we have for each $A, B \in \text{Obj}(\mathcal{C})$ and $f \in \text{Hom}_{\mathcal{C}}(A, B)$, $F(A) = *$ and $F(f) = \text{id}_*^{\mathbf{1}}$. Thus $F = \Delta_*^{\mathcal{C}}$. Since $F : \mathcal{C} \rightarrow \mathbf{1}$ is arbitrary, we have that for each $F : \mathcal{C} \rightarrow \mathbf{1}$, $F = \Delta_*^{\mathcal{C}}$. Thus $\text{Hom}_{\mathbf{Cat}}(\mathcal{C}, \mathbf{1}) = \{\Delta_*^{\mathcal{C}}\}$. Since $\mathcal{C} \in \text{Obj}(\mathbf{Cat})$ is arbitrary, we have that for each $\mathcal{C} \in \text{Obj}(\mathbf{Cat})$, $\text{Hom}_{\mathbf{Cat}}(\mathcal{C}, \mathbf{1}) = \{\Delta_*^{\mathcal{C}}\}$. Hence $\mathbf{1}$ is **Cat**-terminal. □

Definition 7.1.0.9. Let $\mathcal{C}, \mathcal{D}$ be categories and $0, 1 \in \text{Obj}(\mathcal{D})$ and $F : \mathcal{C} \rightarrow \mathcal{D}$. Suppose that 0 is $\mathcal{D}$ -initial and 1 is $\mathcal{D}$ -terminal. Then for each $A \in \text{Obj}(\mathcal{C})$, there exists $f_A \in \text{Hom}_{\mathcal{D}}(0, F(A))$ and $g_A \in \text{Hom}_{\mathcal{D}}(F(A), 1)$ such that $\text{Hom}_{\mathcal{D}}(0, F(A)) = \{f_A\}$ and $\text{Hom}_{\mathcal{D}}(F(A), 1) = \{g_A\}$. We define the

- **initial natural transformation induced by 0 from $\Delta_0^{\mathcal{C}}$ to F** , denoted $\eta^{(0,F)} : \Delta_0^{\mathcal{C}} \Rightarrow F$, by $\eta_A^{(0,F)} := f_A$.
- **terminal natural transformation induced by 1 from F to $\Delta_1^{\mathcal{C}}$** , denoted $\phi^{(F,1)} : F \Rightarrow \Delta_1^{\mathcal{C}}$, by $\phi_A^{(F,1)} := g_A$.

Exercise 7.1.0.10. Let $\mathcal{C}, \mathcal{D}$ be categories and $0 \in \text{Obj}(\mathcal{D})$ and $F : \mathcal{C} \rightarrow \mathcal{D}$. Suppose that 0 is $\mathcal{D}$ -initial and 1 is $\mathcal{D}$ -terminal. Then

1. $\eta^{(0,F)} : \Delta_0^{\mathcal{C}} \Rightarrow F$ is a natural transformation.
2. $\phi^{(F,1)} : F \Rightarrow \Delta_1^{\mathcal{C}}$ is a natural transformation.

Proof.

1. (a) By definition, for each $A \in \text{Obj}(\mathcal{C})$, $\eta_A^{(0,F)} \in \text{Hom}_{\mathcal{D}}(\Delta_0^{\mathcal{C}}(A), F(A))$
- (b) Let $A, B \in \text{Obj}(\mathcal{C})$ and $f \in \text{Hom}_{\mathcal{C}}(A, B)$. Since

$$\begin{aligned} F(f) \circ \eta_A^{(0,F)} &\in \text{Hom}_{\mathcal{D}}(0, F(B)) \\ &= \{\eta_B^{(0,F)}\} \end{aligned}$$

we have that

$$\begin{aligned} F(f) \circ \eta_A^{(0,F)} &= \eta_B^{(0,F)} \\ &= \eta_B^{(0,F)} \circ \text{id}_0^{\mathcal{C}} \end{aligned}$$

i.e. the following diagram commutes:

$$\begin{array}{ccc} \Delta_0^{\mathcal{C}}(A) & \xrightarrow{\eta_A^{(0,F)}} & F(A) \\ \Delta_0^{\mathcal{C}}(f) \downarrow & & \downarrow F(f) \\ \Delta_0^{\mathcal{C}}(B) & \xrightarrow{\eta_B^{(0,F)}} & F(B) \end{array} = \begin{array}{ccc} 0 & \xrightarrow{\eta_A^{(0,F)}} & F(A) \\ \text{id}_0^{\mathcal{C}} \downarrow & & \downarrow F(f) \\ 0 & \xrightarrow{\eta_B^{(0,F)}} & F(B) \end{array}$$

So $\eta^{(0,F)} : \Delta_0^{\mathcal{C}} \Rightarrow F$ is a natural transformation.

2. **FINISH!!!** □

Exercise 7.1.0.11. Let $\mathcal{C}, \mathcal{D}$ be categories and $0, 1 \in \text{Obj}(\mathcal{D})$. Suppose that $\mathcal{C}$ is small.

1. If 0 is $\mathcal{D}$ -initial, then $\Delta_0^{\mathcal{C}}$ is $\mathcal{D}^{\mathcal{C}}$ -initial.
2. If 1 is $\mathcal{D}$ -terminal, then $\Delta_1^{\mathcal{C}}$ is $\mathcal{D}^{\mathcal{C}}$ -terminal.

Proof.

1. Suppose that 0 is $\mathcal{D}$ -initial. Let $F \in \text{Obj}(\mathcal{D}^{\mathcal{C}})$, $\alpha \in \text{Hom}_{\mathcal{D}^{\mathcal{C}}}(\Delta_0^{\mathcal{C}}, F)$ and $A \in \text{Obj}(\mathcal{C})$. Then

$$\begin{aligned}\alpha_A &\in \text{Hom}_{\mathcal{D}}(\Delta_0^{\mathcal{C}}(A), F(A)) \\ &= \text{Hom}_{\mathcal{D}}(0, F(A)) \\ &= \{\eta_A^{(0,F)}\}\end{aligned}$$

Hence $\alpha_A = \eta_A^{(0,F)}$. Since $A \in \text{Obj}(\mathcal{C})$ is arbitrary, we have that for each $A \in \text{Obj}(\mathcal{C})$, $\alpha_A = \eta_A^{(0,F)}$. Thus $\alpha = \eta^{(0,F)}$. Since $\alpha \in \text{Hom}_{\mathcal{D}^{\mathcal{C}}}(\Delta_0^{\mathcal{C}}, F)$ is arbitrary, we have that for each $\alpha \in \text{Hom}_{\mathcal{D}^{\mathcal{C}}}(\Delta_0^{\mathcal{C}}, F)$, $\alpha = \eta^{(0,F)}$. This $\text{Hom}_{\mathcal{D}^{\mathcal{C}}}(\Delta_0^{\mathcal{C}}, F) = \{\eta^{(0,F)}\}$. Therefore $\Delta_0^{\mathcal{C}}$ is $\mathcal{D}^{\mathcal{C}}$ -initial.


2. use duality

□

7.2 Products and Coproducts

Definition 7.2.0.1. Let $\mathcal{C}$ be a category, A a set (class?), $(X_\alpha)_{\alpha \in A} \in \text{Obj}(\mathcal{C})^A$ and $X \in \text{Obj}(\mathcal{C})$.


- Let $(\pi_\alpha)_{\alpha \in A} \in \prod_{\alpha \in A} \text{Hom}_{\mathcal{C}}(X, X_\alpha)$. Then $(X, (\pi_\alpha)_{\alpha \in A})$ is said to be a $\mathcal{C}$ -**product** of $(X_\alpha)_{\alpha \in A}$ if for each $Y \in \text{Obj}(\mathcal{C})$ and $(p_\alpha)_{\alpha \in A} \in \prod_{\alpha \in A} \text{Hom}_{\mathcal{C}}(Y, X_\alpha)$, there exists a unique $\phi \in \text{Hom}_{\mathcal{C}}(Y, X)$ such that for each $\alpha \in A$, from $\pi_\alpha \circ_{\mathcal{C}} \phi = p_\alpha$, i.e. the following diagram commutes:



- We define

$$\text{Prod}_{\mathcal{C}}((X_\alpha)_{\alpha \in A}) := \left\{ (X, (\pi_\alpha)_{\alpha \in A}) \in \text{Obj}(\mathcal{C}) \times \prod_{\alpha \in A} \text{Hom}_{\mathcal{C}}(X, X_\alpha) : (X, (\pi_\alpha)_{\alpha \in A}) \text{ is a } \mathcal{C}\text{-product of } (X_\alpha)_{\alpha \in A} \right\}.$$

- Let $(\iota_\alpha)_{\alpha \in A} \in \prod_{\alpha \in A} \text{Hom}_{\mathcal{C}}(X_\alpha, X)$. Then $(X, (\iota_\alpha)_{\alpha \in A})$ is said to be a $\mathcal{C}$ -**coproduct** of $(X_\alpha)_{\alpha \in A}$ if for each $Y \in \text{Obj}(\mathcal{C})$ and $(i_\alpha)_{\alpha \in A} \in \prod_{\alpha \in A} \text{Hom}_{\mathcal{C}}(X_\alpha, Y)$, there exists a unique $\phi \in \text{Hom}_{\mathcal{C}}(X, Y)$ such that for each $\alpha \in A$, from $\phi \circ_{\mathcal{C}} \iota_\alpha = i_\alpha$, i.e. the following diagram commutes:



- We define

$$\text{Coproduct}_{\mathcal{C}}((X_\alpha)_{\alpha \in A}) := \left\{ (X, (\iota_\alpha)_{\alpha \in A}) \in \text{Obj}(\mathcal{C}) \times \prod_{\alpha \in A} \text{Hom}_{\mathcal{C}}(X_\alpha, X) : (X, (\iota_\alpha)_{\alpha \in A}) \text{ is a } \mathcal{C}\text{-coproduct of } (X_\alpha)_{\alpha \in A} \right\}.$$

Exercise 7.2.0.2. Let $\mathcal{C}$ be a category, A a set (class?) and $(X_\alpha)_{\alpha \in A} \in \text{Obj}(\mathcal{C})^A$. Then

1. $\text{Prod}_{\mathcal{C}^{\text{op}}}((X_\alpha)_{\alpha \in A}) = \text{Coproduct}_{\mathcal{C}}((X_\alpha)_{\alpha \in A})$.
2. $\text{Coproduct}_{\mathcal{C}^{\text{op}}}((X_\alpha)_{\alpha \in A}) = \text{Prod}_{\mathcal{C}}((X_\alpha)_{\alpha \in A})$.

Proof.

1. • Let $(X, (\pi_\alpha)_{\alpha \in A}) \in \text{Prod}_{\mathcal{C}^{\text{op}}}((X_\alpha)_{\alpha \in A})$. We note that $(X_\alpha)_{\alpha \in A} \in \text{Obj}(\mathcal{C})^A$, $X \in \text{Obj}(\mathcal{C})$ and $(\pi_\alpha)_{\alpha \in A} \in \prod_{\alpha \in A} \text{Hom}_{\mathcal{C}}(X_\alpha, X)$. Let $Y \in \text{Obj}(\mathcal{C})$ and $(i_\alpha)_{\alpha \in A} \in \prod_{\alpha \in A} \text{Hom}_{\mathcal{C}}(X_\alpha, Y)$.
 - **Existence:**
Then $Y \in \text{Obj}(\mathcal{C}^{\text{op}})$ and $(i_\alpha)_{\alpha \in A} \in \prod_{\alpha \in A} \text{Hom}_{\mathcal{C}^{\text{op}}}(Y, X_\alpha)$. Since $(X, (\pi_\alpha)_{\alpha \in A}) \in \text{Prod}_{\mathcal{C}^{\text{op}}}((X_\alpha)_{\alpha \in A})$, there exists a unique $\phi \in \text{Hom}_{\mathcal{C}^{\text{op}}}(Y, X)$ such that for each $\alpha \in A$, $\pi_\alpha \circ_{\mathcal{C}^{\text{op}}} \phi = i_\alpha$. Then $\phi \in \text{Hom}_{\mathcal{C}}(X, Y)$ and for each $\alpha \in A$, $\phi \circ_{\mathcal{C}} \pi_\alpha = i_\alpha$.
 - **Uniqueness:**
Let $\phi' \in \text{Hom}_{\mathcal{C}}(X, Y)$. Suppose that for each $\alpha \in A$, $\phi' \circ_{\mathcal{C}} \pi_\alpha = i_\alpha$. Then $\phi' \in \text{Hom}_{\mathcal{C}^{\text{op}}}(Y, X)$ and for each $\alpha \in A$, $\pi_\alpha \circ_{\mathcal{C}^{\text{op}}} \phi' = i_\alpha$. Uniqueness of ϕ implies that $\phi' = \phi$.

Since $Y \in \text{Obj}(\mathcal{C})$ and $(i_\alpha)_{\alpha \in A} \in \prod_{\alpha \in A} \text{Hom}_{\mathcal{C}}(X_\alpha, Y)$ are arbitrary, we have that for each $Y \in \text{Obj}(\mathcal{C})$ and $(i_\alpha)_{\alpha \in A} \in \prod_{\alpha \in A} \text{Hom}_{\mathcal{C}}(X_\alpha, Y)$, there exists a unique $\phi \in \text{Hom}_{\mathcal{C}}(X, Y)$ such that for each $\alpha \in A$, $\phi \circ_C \pi_\alpha = i_\alpha$. Thus $(X, (\pi_\alpha)_{\alpha \in A}) \in \text{Coproduct}_{\mathcal{C}}((X_\alpha)_{\alpha \in A})$. Since $(X, (\pi_\alpha)_{\alpha \in A}) \in \text{Prod}_{\mathcal{C}^{\text{op}}}((X_\alpha)_{\alpha \in A})$ is arbitrary, we have that for each $(X, (\pi_\alpha)_{\alpha \in A}) \in \text{Prod}_{\mathcal{C}^{\text{op}}}((X_\alpha)_{\alpha \in A})$, $(X, (\pi_\alpha)_{\alpha \in A}) \in \text{Coproduct}_{\mathcal{C}}((X_\alpha)_{\alpha \in A})$. Hence $\text{Prod}_{\mathcal{C}^{\text{op}}}((X_\alpha)_{\alpha \in A}) \subset \text{Coproduct}_{\mathcal{C}}((X_\alpha)_{\alpha \in A})$.

- **Similar, fill out!!!**

Since $\text{Prod}_{\mathcal{C}^{\text{op}}}((X_\alpha)_{\alpha \in A}) \subset \text{Coproduct}_{\mathcal{C}}((X_\alpha)_{\alpha \in A})$ and $\text{Coproduct}_{\mathcal{C}}((X_\alpha)_{\alpha \in A}) \subset \text{Prod}_{\mathcal{C}^{\text{op}}}((X_\alpha)_{\alpha \in A})$, we have that $\text{Prod}_{\mathcal{C}^{\text{op}}}((X_\alpha)_{\alpha \in A}) = \text{Coproduct}_{\mathcal{C}}((X_\alpha)_{\alpha \in A})$.

2. **The prev part (ref)** implies that

$$\begin{aligned} \text{Coproduct}_{\mathcal{C}^{\text{op}}}((X_\alpha)_{\alpha \in A}) &= \text{Prod}_{(\mathcal{C}^{\text{op}})^{\text{op}}}((X_\alpha)_{\alpha \in A}) \\ &= \text{Prod}_{\mathcal{C}}((X_\alpha)_{\alpha \in A}). \end{aligned}$$

□

Exercise 7.2.0.3. Let $\mathcal{C}$ be a category, A a set (**class?**) and $(X_\alpha)_{\alpha \in A} \in \text{Obj}(\mathcal{C})^A$. Then

1. for each $(X, (\pi_\alpha^X)_{\alpha \in A}), (Y, (\pi_\alpha^Y)_{\alpha \in A}) \in \text{Prod}_{\mathcal{C}}((X_\alpha)_{\alpha \in A})$, $X \cong_{\mathcal{C}} Y$.
2. for each $(X, (\iota_\alpha^X)_{\alpha \in A}), (Y, (\iota_\alpha^Y)_{\alpha \in A}) \in \text{Coproduct}_{\mathcal{C}}((X_\alpha)_{\alpha \in A})$, $X \cong_{\mathcal{C}} Y$.

Proof.

1. Let $(X, (\pi_\alpha^X)_{\alpha \in A}), (Y, (\pi_\alpha^Y)_{\alpha \in A}) \in \text{Prod}_{\mathcal{C}}((X_\alpha)_{\alpha \in A})$.

- – By definition, there exists a unique $\phi_{Y,X} \in \text{Hom}_{\mathcal{C}}(Y, X)$ such that for each $\alpha \in A$, $\pi_\alpha^X \circ_C \phi_{Y,X} = \pi_\alpha^Y$. Similarly, there exists a unique $\phi_{X,Y} \in \text{Hom}_{\mathcal{C}}(X, Y)$ such that for each $\alpha \in A$, $\pi_\alpha^Y \circ_C \phi_{X,Y} = \pi_\alpha^X$.

$$\begin{aligned} \pi_\alpha^Y \circ_C (\phi_{X,Y} \circ_C \phi_{Y,X}) &= (\pi_\alpha^Y \circ_C \phi_{X,Y}) \circ_C \phi_{Y,X} \\ &= \pi_\alpha^X \circ_C \phi_{Y,X} \\ &= \pi_\alpha^X. \end{aligned}$$

- By definition, there exists a unique $\phi \in \text{Hom}_{\mathcal{C}}(Y, Y)$ such that $\pi_\alpha^Y \circ \phi = \pi_\alpha^Y$.
- We note that $\pi_\alpha^Y \circ \text{id}_Y^{\mathcal{C}} = \pi_\alpha^Y$.

Thus the following diagrams commutes:

$$\begin{array}{ccc} \begin{array}{ccc} Y & & Y \\ \phi_{X,Y} \circ_C \phi_{Y,X} \downarrow & \searrow \pi_\alpha & \downarrow \phi \\ Y & & Y \\ \pi_\alpha \searrow & & \searrow \pi_\alpha \\ & X_\alpha & \end{array} & \begin{array}{ccc} Y & & Y \\ \phi \downarrow & \searrow \pi_\alpha & \downarrow \phi \\ Y & & Y \\ \pi_\alpha \searrow & & \searrow \pi_\alpha \\ & X_\alpha & \end{array} & \begin{array}{ccc} Y & & Y \\ \text{id}_Y^{\mathcal{C}} \downarrow & \searrow \pi_\alpha & \downarrow \text{id}_Y^{\mathcal{C}} \\ Y & & Y \\ \pi_\alpha \searrow & & \searrow \pi_\alpha \\ & X_\alpha & \end{array} \end{array}$$

Uniqueness of ϕ then implies that

$$\begin{aligned} \phi_{X,Y} \circ_C \phi_{Y,X} &= \phi \\ &= \text{id}_Y^{\mathcal{C}}, \end{aligned}$$

- Similarly, $\phi_{Y,X} \circ_C \phi_{X,Y} = \text{id}_X^{\mathcal{C}}$.

Hence $\phi_{Y,X} \in \text{Iso}_{\mathcal{C}}(Y, X)$ and $X \cong_{\mathcal{C}} Y$. Since $(X, (\pi_\alpha^X)_{\alpha \in A}), (Y, (\pi_\alpha^Y)_{\alpha \in A}) \in \text{Prod}_{\mathcal{C}}((X_\alpha)_{\alpha \in A})$ are arbitrary, we have that for each $(X, (\pi_\alpha^X)_{\alpha \in A}), (Y, (\pi_\alpha^Y)_{\alpha \in A}) \in \text{Prod}_{\mathcal{C}}((X_\alpha)_{\alpha \in A})$, $X \cong_{\mathcal{C}} Y$.

2. Let $(X, (\iota_\alpha^X)_{\alpha \in A}), (Y, (\iota_\alpha^Y)_{\alpha \in A}) \in \text{Coproduct}_{\mathcal{C}}((X_\alpha)_{\alpha \in A})$. **The prev ex (ref)** implies that $(X, (\iota_\alpha^X)_{\alpha \in A}), (Y, (\iota_\alpha^Y)_{\alpha \in A}) \in \text{Prod}_{\mathcal{C}^{\text{op}}}((X_\alpha)_{\alpha \in A})$. **The prev part (ref)** then implies that $X \cong_{\mathcal{C}^{\text{op}}} Y$. **An exercise in isomorphism section (ref)** implies that $X \cong_{\mathcal{C}} Y$.

□

Note 7.2.0.4.

- If $(X, (\pi_\alpha)_{\alpha \in A}) \in \text{Prod}_{\mathcal{C}}((X_\alpha)_{\alpha \in A})$, we may write $\prod_{\alpha \in A} X_\alpha$ in place of X .
- If $(X, (\iota_\alpha)_{\alpha \in A}) \in \text{Coproduct}_{\mathcal{C}}((X_\alpha)_{\alpha \in A})$, we may write $\coprod_{\alpha \in A} X_\alpha$ in place of X .
- If $A = \{1, 2\}$, we may write
 - $\text{Prod}_{\mathcal{C}}(X_1, X_2)$ and $\text{Coproduct}_{\mathcal{C}}(X_1, X_2)$ in place of $\text{Prod}_{\mathcal{C}}((X_\alpha)_{\alpha \in A})$ and $\text{Coproduct}_{\mathcal{C}}((X_\alpha)_{\alpha \in A})$ respectively.
 - $X_1 \times X_2$ in place of $\prod_{\alpha \in A} X_\alpha$ and $X_1 + X_2$ in place of $\coprod_{\alpha \in A} X_\alpha$

Note 7.2.0.5. Other common notations for coproduct include $\bigoplus_{\alpha \in A} X_\alpha$ and $X_1 \oplus X_2$.

Exercise 7.2.0.6. Let $\mathcal{C}$ be a category, A a set (class?), $X \in \text{Obj}(\mathcal{C})$, $(X_\alpha)_{\alpha \in A} \in \text{Obj}(\mathcal{C})^A$, $(\prod_{\alpha \in A} X_\alpha, (\pi_\alpha)_{\alpha \in A}) \in \text{Prod}_{\mathcal{C}}((X_\alpha)_{\alpha \in A})$ and $f, g \in \text{Hom}_{\mathcal{C}}(X, \prod_{\alpha \in A} X_\alpha)$. If for each $\alpha \in A$, $\pi_\alpha \circ_{\mathcal{C}} f = \pi_\alpha \circ_{\mathcal{C}} g$, then $f = g$.

Proof. Suppose that for each $\alpha \in A$, $\pi_\alpha \circ_{\mathcal{C}} f = \pi_\alpha \circ_{\mathcal{C}} g$. Since $(\pi_\alpha \circ_{\mathcal{C}} g)_{\alpha \in A} \in \prod_{\alpha \in A} \text{Hom}_{\mathcal{C}}(X, X_\alpha)$ and $(\prod_{\alpha \in A} X_\alpha, (\pi_\alpha)_{\alpha \in A}) \in \text{Prod}_{\mathcal{C}}((X_\alpha)_{\alpha \in A})$, there exists a unique $\phi \in \text{Hom}_{\mathcal{C}}(X, \prod_{\alpha \in A} X_\alpha)$ such that for each $\alpha \in A$, $\pi_\alpha \circ_{\mathcal{C}} \phi = \pi_\alpha \circ_{\mathcal{C}} g$, i.e. the following diagram commutes:

$$\begin{array}{ccc}
 X & & \\
 \phi \downarrow & \searrow \pi_\alpha \circ_{\mathcal{C}} g & \\
 \prod_{\alpha \in A} X_\alpha & & \\
 \pi_\alpha \searrow & & \\
 & X_\alpha &
 \end{array}$$

Since for each $\alpha \in A$, $\pi_\alpha \circ_{\mathcal{C}} f = \pi_\alpha \circ_{\mathcal{C}} g$ and $\pi_\alpha \circ_{\mathcal{C}} g = \pi_\alpha \circ_{\mathcal{C}} \phi$, uniqueness of ϕ implies that

$$\begin{aligned}
 f &= \phi \\
 &= g.
 \end{aligned}$$

□

Definition 7.2.0.7. Let $\mathcal{C}$ be a category, A a set (class?), $(X_\alpha)_{\alpha \in A}, (Y_\alpha)_{\alpha \in A} \in \text{Obj}(\mathcal{C})^A$, $(f_\alpha)_{\alpha \in A} \in \prod_{\alpha \in A} \text{Hom}_{\mathcal{C}}(X_\alpha, Y_\alpha)$, $(\prod_{\alpha \in A} X_\alpha, (\pi_\alpha^X)_{\alpha \in A}) \in \text{Prod}_{\mathcal{C}}((X_\alpha)_{\alpha \in A})$ and $(\prod_{\alpha \in A} Y_\alpha, (\pi_\alpha^Y)_{\alpha \in A}) \in \text{Prod}_{\mathcal{C}}((Y_\alpha)_{\alpha \in A})$. We define the $\mathcal{C}$ -**product of** $(f_\alpha)_{\alpha \in A}$, denoted $\prod_{\alpha \in A} f_\alpha \in \text{Hom}_{\mathcal{C}}(\prod_{\alpha \in A} X_\alpha, \prod_{\alpha \in A} Y_\alpha)$ to be the unique $f \in \text{Hom}_{\mathcal{C}}(\prod_{\alpha \in A} X_\alpha, \prod_{\alpha \in A} Y_\alpha)$ such that for each $\alpha \in A$, $\pi_\alpha^Y \circ_{\mathcal{C}} f = f_\alpha \circ_{\mathcal{C}} \pi_\alpha^X$, i.e the following diagram commutes:

$$\begin{array}{ccc}
 \prod_{\alpha \in A} X_\alpha & & \\
 f \downarrow & \searrow f_\alpha \circ_{\mathcal{C}} \pi_\alpha^X & \\
 \prod_{\alpha \in A} Y_\alpha & & \\
 \pi_\alpha^Y \searrow & & \\
 & Y_\alpha &
 \end{array}$$

Exercise 7.2.0.8. Let $\mathcal{C}$ be a category, A a set (class?), $(X_\alpha)_{\alpha \in A}, (Y_\alpha)_{\alpha \in A} \in \text{Obj}(\mathcal{C})^A$, $(f_\alpha)_{\alpha \in A} \in \prod_{\alpha \in A} \text{Hom}_{\mathcal{C}}(X_\alpha, Y_\alpha)$, $(\prod_{\alpha \in A} X_\alpha, (\pi_\alpha^X)_{\alpha \in A}) \in \text{Prod}_{\mathcal{C}}((X_\alpha)_{\alpha \in A})$ and $(\prod_{\alpha \in A} Y_\alpha, (\pi_\alpha^Y)_{\alpha \in A}) \in \text{Prod}_{\mathcal{C}}((Y_\alpha)_{\alpha \in A})$. Then $(\pi_\alpha^X, \pi_\alpha^Y)_{\alpha \in A} \in \prod_{\alpha \in A} \text{Hom}_{\mathbf{Arr}(\mathcal{C})}(\prod_{\alpha \in A} X_\alpha, \prod_{\alpha \in A} Y_\alpha)$.

Proof. By construction, for each $\alpha \in A$,

$$f_\alpha \circ_{\mathcal{C}} \pi_\alpha^X = \pi_\alpha^Y \circ_{\mathcal{C}} \prod_{\alpha \in A} f_\alpha,$$

i.e. the following diagram commutes:

$$\begin{array}{ccc} \prod_{\alpha \in A} X_\alpha & \xrightarrow{\pi_\alpha^X} & X_\alpha \\ \prod_{\alpha \in A} f_\alpha \downarrow & & \downarrow f_\alpha \\ \prod_{\alpha \in A} Y_\alpha & \xrightarrow{\pi_\alpha^Y} & Y_\alpha \end{array}$$

Thus $(\pi_\alpha^X, \pi_\alpha^Y)_{\alpha \in A} \in \prod_{\alpha \in A} \text{Hom}_{\mathbf{Arr}(\mathcal{C})}(\prod_{\alpha \in A} f_\alpha, f_\alpha)$. □

Next exercise shows that the product of morphisms is in fact a product object in the arrow category

Exercise 7.2.0.9. Let $\mathcal{C}$ be a category, A a set (class?), $(X_\alpha)_{\alpha \in A}, (Y_\alpha)_{\alpha \in A} \in \text{Obj}(\mathcal{C})^A$, $(f_\alpha)_{\alpha \in A} \in \prod_{\alpha \in A} \text{Hom}_{\mathcal{C}}(X_\alpha, Y_\alpha)$, $(\prod_{\alpha \in A} X_\alpha, (\pi_\alpha^X)_{\alpha \in A}) \in \text{Prod}_{\mathcal{C}}((X_\alpha)_{\alpha \in A})$ and $(\prod_{\alpha \in A} Y_\alpha, (\pi_\alpha^Y)_{\alpha \in A}) \in \text{Prod}_{\mathcal{C}}((Y_\alpha)_{\alpha \in A})$. Define $(\pi_\alpha)_{\alpha \in A} \in \prod_{\alpha \in A} \text{Hom}_{\mathbf{Arr}(\mathcal{C})}(\prod_{\alpha \in A} f_\alpha, f_\alpha)$ by $\pi_\alpha := (\pi_\alpha^X, \pi_\alpha^Y)$. Then $(\prod_{\alpha \in A} f_\alpha, \pi_\alpha) \in \text{Prod}_{\mathbf{Arr}(\mathcal{C})}((f_\alpha)_{\alpha \in A})$.

Proof. Let $g \in \text{Obj}(\mathbf{Arr}(\mathcal{C}))$ and $(p_\alpha)_{\alpha \in A} \in \prod_{\alpha \in A} \text{Hom}_{\mathbf{Arr}(\mathcal{C})}(g, f_\alpha)$. Define $A, B \in \text{Obj}(\mathcal{C})$ by $A := \text{dom}_{\mathcal{C}}(g)$ and $B := \text{cod}_{\mathcal{C}}(g)$. Write $p_\alpha = (p_\alpha^A, p_\alpha^B)$. We note that

- By definition, for each $\alpha \in A$, $f_\alpha \circ_{\mathcal{C}} p_\alpha^A = p_\alpha^B \circ_{\mathcal{C}} g$, i.e. the following diagram commutes:

$$\begin{array}{ccc} A & \xrightarrow{p_\alpha^A} & X_\alpha \\ g \downarrow & & \downarrow f_\alpha \\ B & \xrightarrow{p_\alpha^B} & Y_\alpha \end{array}$$

- By construction, for each $\alpha \in A$, $f_\alpha \circ_{\mathcal{C}} \pi_\alpha^X = \pi_\alpha^Y \circ_{\mathcal{C}} \prod_{\alpha \in A} f_\alpha$, i.e. the following diagram commutes:

$$\begin{array}{ccc} \prod_{\alpha \in A} X_\alpha & \xrightarrow{\pi_\alpha^X} & X_\alpha \\ \prod_{\alpha \in A} f_\alpha \downarrow & & \downarrow f_\alpha \\ \prod_{\alpha \in A} Y_\alpha & \xrightarrow{\pi_\alpha^Y} & Y_\alpha \end{array}$$

• **Existence:**

By definition, there exists a unique $\phi_X \in \text{Hom}_{\mathcal{C}}(A, \prod_{\alpha \in A} X_\alpha)$ such that for each $\alpha \in A$, $\pi_\alpha^X \circ_{\mathcal{C}} \phi_X = p_\alpha^A$, i.e. the following diagram commutes:

$$\begin{array}{ccc} A & & \\ \phi_X \downarrow & \searrow p_\alpha^A & \\ \prod_{\alpha \in A} X_\alpha & & \\ & \searrow \pi_\alpha^X & \\ & & X_\alpha \end{array}$$

Similarly, there exists a unique $\phi_Y \in \text{Hom}_{\mathcal{C}}(B, \prod_{\alpha \in A} Y_{\alpha})$ such that for each $\alpha \in A$, $\pi_{\alpha}^Y \circ_{\mathcal{C}} \phi_Y = p_{\alpha}^B$, i.e. the following diagram commutes:

$$\begin{array}{ccc}
 B & & \\
 \phi_Y \downarrow & \searrow p_{\alpha}^B & \\
 \prod_{\alpha \in A} Y_{\alpha} & & \\
 \pi_{\alpha}^Y \searrow & & \downarrow \\
 & & Y_{\alpha}
 \end{array}$$

Then for each $\alpha \in A$,

$$\begin{aligned}
 \pi_{\alpha}^Y \circ_{\mathcal{C}} \left(\prod_{\alpha \in A} f_{\alpha} \circ_{\mathcal{C}} \phi_X \right) &= (\pi_{\alpha}^Y \circ_{\mathcal{C}} \prod_{\alpha \in A} f_{\alpha}) \circ_{\mathcal{C}} \phi_X \\
 &= (f_{\alpha} \circ_{\mathcal{C}} \pi_{\alpha}^X) \circ_{\mathcal{C}} \phi_X \\
 &= f_{\alpha} \circ_{\mathcal{C}} (\pi_{\alpha}^X \circ_{\mathcal{C}} \phi_X) \\
 &= f_{\alpha} \circ_{\mathcal{C}} p_{\alpha}^A \\
 &= p_{\alpha}^B \circ_{\mathcal{C}} g \\
 &= (\pi_{\alpha}^Y \circ_{\mathcal{C}} \phi_Y) \circ_{\mathcal{C}} g \\
 &= \pi_{\alpha}^Y \circ_{\mathcal{C}} (\phi_Y \circ_{\mathcal{C}} g).
 \end{aligned}$$

the prev ex (ref) then implies that $\prod_{\alpha \in A} f_{\alpha} \circ_{\mathcal{C}} \phi_X = \phi_Y \circ_{\mathcal{C}} g$, i.e. the following diagram commutes:

$$\begin{array}{ccc}
 A & \xrightarrow{\phi_X} & \prod_{\alpha \in A} X_{\alpha} \\
 g \downarrow & & \downarrow \prod_{\alpha \in A} f_{\alpha} \\
 B & \xrightarrow{\phi_Y} & \prod_{\alpha \in A} Y_{\alpha}
 \end{array}$$

Thus $(\phi_X, \phi_Y) \in \text{Hom}_{\mathbf{Arr}(\mathcal{C})}(g, \prod_{\alpha \in A} f_{\alpha})$. Define $\phi \in \text{Hom}_{\mathbf{Arr}(\mathcal{C})}(g, \prod_{\alpha \in A} f_{\alpha})$ by $\phi := (\phi_X, \phi_Y)$. Let $\alpha \in A$. Since $\pi_{\alpha}^X \circ_{\mathcal{C}} \phi_X = p_{\alpha}^A$ and $\pi_{\alpha}^Y \circ_{\mathcal{C}} \phi_Y = p_{\alpha}^B$, we have that

$$\begin{aligned}
 \pi_{\alpha} \circ_{\mathbf{Arr}(\mathcal{C})} \phi &= (\pi_{\alpha}^X, \pi_{\alpha}^Y) \circ_{\mathbf{Arr}(\mathcal{C})} (\phi_X, \phi_Y) \\
 &= (\pi_{\alpha}^X \circ_{\mathcal{C}} \phi_X, \pi_{\alpha}^Y \circ_{\mathcal{C}} \phi_Y) \\
 &= (p_{\alpha}^A, p_{\alpha}^B) \\
 &= p_{\alpha}.
 \end{aligned}$$

Since $\alpha \in A$ is arbitrary, we have that for each $\alpha \in A$, $\pi_{\alpha} \circ_{\mathbf{Arr}(\mathcal{C})} \phi = p_{\alpha}$, i.e. the following diagram commutes:

$$\begin{array}{ccc}
 g & & \\
 \phi \downarrow & \searrow p_{\alpha} & \\
 \prod_{\alpha \in A} f_{\alpha} & & \\
 \pi_{\alpha} \searrow & & \downarrow \\
 & & f_{\alpha}
 \end{array}$$

• **Uniqueness:**

Let $\phi' \in \text{Hom}_{\mathbf{Arr}(\mathcal{C})}(g, \prod_{\alpha \in A} f_{\alpha})$. Write $\phi' = (\phi'_X, \phi'_Y)$. Suppose that for each $\alpha \in A$, $\pi_{\alpha} \circ_{\mathbf{Arr}(\mathcal{C})} \phi' = p_{\alpha}$, i.e. the following

diagram commutes:

$$\begin{array}{ccc}
 g & & \\
 \phi' \downarrow & \searrow p_\alpha & \\
 \prod_{\alpha \in A} f_\alpha & & \\
 \pi_\alpha \searrow & & \downarrow \\
 & & f_\alpha
 \end{array}$$

Let $\alpha \in A$. Then

$$\begin{aligned}
 (\pi_\alpha^X \circ_{\mathcal{C}} \phi'_X, \pi_\alpha^Y \circ_{\mathcal{C}} \phi'_Y) &= (\pi_\alpha^X, \pi_\alpha^Y) \circ_{\mathbf{Arr}(\mathcal{C})} (\phi'_X, \phi'_Y) \\
 &= \pi_\alpha \circ_{\mathbf{Arr}(\mathcal{C})} \phi' \\
 &= p_\alpha \\
 &= (p_\alpha^A, p_\alpha^B).
 \end{aligned}$$

Therefore $\pi_\alpha^X \circ_{\mathcal{C}} \phi'_X = p_\alpha^A$ and $\pi_\alpha^Y \circ_{\mathcal{C}} \phi'_Y = p_\alpha^B$. Since $\alpha \in A$ is arbitrary, we have that for each $\alpha \in A$, $\pi_\alpha^X \circ_{\mathcal{C}} \phi'_X = p_\alpha^A$ and $\pi_\alpha^Y \circ_{\mathcal{C}} \phi'_Y = p_\alpha^B$, i.e. the following diagrams commute:

$$\begin{array}{ccc}
 A & & B \\
 \phi'_X \downarrow & \searrow p_\alpha^A & \phi'_Y \downarrow \\
 \prod_{\alpha \in A} X_\alpha & & \prod_{\alpha \in A} Y_\alpha \\
 \pi_\alpha^X \searrow & & \pi_\alpha^Y \searrow \\
 & & X_\alpha \quad Y_\alpha
 \end{array}$$

Uniqueness of ϕ_X and ϕ_Y imply that $\phi'_X = \phi_X$ and $\phi'_Y = \phi_Y$. Thus

$$\begin{aligned}
 \phi' &= (\phi'_X, \phi'_Y) \\
 &= (\phi_X, \phi_Y) \\
 &= \phi.
 \end{aligned}$$

Since $\phi' \in \text{Hom}_{\mathbf{Arr}(\mathcal{C})}(g, \prod_{\alpha \in A} f_\alpha)$ is arbitrary, we have that for each $\phi' \in \text{Hom}_{\mathbf{Arr}(\mathcal{C})}(g, \prod_{\alpha \in A} f_\alpha)$, $\phi' = \phi$.

Thus there exists a unique $\phi \in \text{Hom}_{\mathbf{Arr}(\mathcal{C})}(g, \prod_{\alpha \in A} f_\alpha)$ such that for each $\alpha \in A$, $\pi_\alpha \circ_{\mathbf{Arr}(\mathcal{C})} \phi = p_\alpha$. Since $g \in \text{Obj}(\mathbf{Arr}(\mathcal{C}))$ and $(p_\alpha)_{\alpha \in A} \in \prod_{\alpha \in A} \text{Hom}_{\mathbf{Arr}(\mathcal{C})}(g, f_\alpha)$ are arbitrary, we have that for each $g \in \text{Obj}(\mathbf{Arr}(\mathcal{C}))$ and $(p_\alpha)_{\alpha \in A} \in \prod_{\alpha \in A} \text{Hom}_{\mathbf{Arr}(\mathcal{C})}(g, f_\alpha)$, there exists a unique $\phi \in \text{Hom}_{\mathbf{Arr}(\mathcal{C})}(g, \prod_{\alpha \in A} f_\alpha)$ such that for each $\alpha \in A$, $\pi_\alpha \circ_{\mathbf{Arr}(\mathcal{C})} \phi = p_\alpha$, i.e. the following diagram commutes:

$$\begin{array}{ccc}
 g & & \\
 \phi \downarrow & \searrow p_\alpha & \\
 \prod_{\alpha \in A} f_\alpha & & \\
 \pi_\alpha \searrow & & \downarrow \\
 & & f_\alpha
 \end{array}$$

Hence $(\prod_{\alpha \in A} f_\alpha, (\pi_\alpha)_{\alpha \in A}) \in \text{Prod}_{\mathbf{Arr}(\mathcal{C})}((f_\alpha)_{\alpha \in A})$. □

Definition 7.2.0.10. Let $\mathcal{C}$ be a category, A a set (class?), $X \in \text{Obj}(\mathcal{C})$, $(Y_\alpha)_{\alpha \in A} \in \text{Obj}(\mathcal{C})^A$, $(f_\alpha)_{\alpha \in A} \in \prod_{\alpha \in A} \text{Hom}_{\mathcal{C}}(X, Y_\alpha)$ and $(\prod_{\alpha \in A} Y_\alpha, (\pi_\alpha^Y)_{\alpha \in A}) \in \text{Prod}_{\mathcal{C}}((Y_\alpha)_{\alpha \in A})$. We define the **tupling of** $(f_\alpha)_{\alpha \in A}$, denoted $(f_\alpha)_\alpha \in \text{Hom}_{\mathcal{C}}(X, \prod_{\alpha \in A} Y_\alpha)$ to be the unique

$\phi \in \text{Hom}_{\mathcal{C}}(X, \prod_{\alpha \in A} Y_{\alpha})$ such that for each $\alpha \in A$, $\pi_{\alpha} \circ \phi = f_{\alpha}$, i.e. the following diagram commutes:

$$\begin{array}{ccc}
 g & & \\
 \downarrow \phi & \searrow p_{\alpha} & \\
 \prod_{\alpha \in A} f_{\alpha} & & \\
 \searrow \pi_{\alpha} & & \downarrow \\
 & & f_{\alpha}
 \end{array}$$

Definition 7.2.0.11. Let $\mathcal{C}$ be a category. Then $\mathcal{C}$ is said to


- **have all products** if for each $A \in \text{Obj}(\mathbf{Set})$ and $(X_{\alpha})_{\alpha \in A} \in \text{Obj}(\mathcal{C})^A$, $\text{Prod}_{\mathcal{C}}((A_{\alpha})_{\alpha \in A}) \neq \emptyset$,
- **have all coproducts** if for each $A \in \text{Obj}(\mathbf{Set})$ and $(X_{\alpha})_{\alpha \in A} \in \text{Obj}(\mathcal{C})^A$, $\text{Prod}_{\mathcal{C}}((A_{\alpha})_{\alpha \in A}) \neq \emptyset$,
- **have all finite products** if for each $A, B \in \text{Obj}(\mathcal{C})$, $\text{Prod}_{\mathcal{C}}(A, B) \neq \emptyset$,
- **have all finite coproducts** if for each $A, B \in \text{Obj}(\mathcal{C})$, $\text{Coproduct}_{\mathcal{C}}(A, B) \neq \emptyset$,

7.3 Equalizers and Coequalizers


need to redo diagrams either with parallel arrows and describe some precise way to define them so that 'commuting' makes sense, or use commutative squares

Definition 7.3.0.1. Let $\mathcal{C}$ be a category. Let $A, B \in \text{Obj}(\mathcal{C})$ and $g, h \in \text{Hom}_{\mathcal{C}}(A, B)$.


- Let $C \in \text{Obj}(\mathcal{C})$.
 - Let $f \in \text{Hom}_{\mathcal{C}}(C, A)$. Then (C, f) is said to be a **$\mathcal{C}$ -equalizer of g and h** if
 1. $g \circ_C f = h \circ_C f$, i.e. the following diagram commutes:




2. for each $C' \in \text{Obj}(\mathcal{C})$ and $f' \in \text{Hom}_{\mathcal{C}}(C', A)$, if $g \circ_C f' = h \circ_C f'$, then there exists a unique $\phi \in \text{Hom}_{\mathcal{C}}(C', C)$ such that $f \circ_C \phi = f'$, i.e. the following diagram commutes:



- Let $f \in \text{Hom}_{\mathcal{C}}(B, C)$. Then (C, f) is said to be a **$\mathcal{C}$ -coequalizer of g and h** if
 1. $f \circ_C g = f \circ_C h$, i.e. the following diagram commutes:



2. for each $C' \in \text{Obj}(\mathcal{C})$ and $f' \in \text{Hom}_{\mathcal{C}}(B, C')$, if $f' \circ_C g = f' \circ_C h$, then there exists a unique $\phi \in \text{Hom}_{\mathcal{C}}(C, C')$ such that $\phi \circ_C f = f'$, i.e. the following diagram commutes:



- We define

- $\text{Equi}_{\mathcal{C}}(g, h) := \{(C, f) \in \text{Obj}(\mathcal{C}) \times \text{Hom}_{\mathcal{C}} : \text{cod}_{\mathcal{C}}(f) = A \text{ and } f \text{ is a } \mathcal{C}\text{-equalizer of } g \text{ and } h\}$
- $\text{Coequi}_{\mathcal{C}}(g, h) := \{(C, f) \in \text{Obj}(\mathcal{C}) \times \text{Hom}_{\mathcal{C}} : \text{dom}_{\mathcal{C}}(f) = B \text{ and } f \text{ is a } \mathcal{C}\text{-coequalizer of } g \text{ and } h\}$

Definition 7.3.0.2. Let $\mathcal{C}$ be a category.

- Let $C \in \text{Obj}(\mathcal{C})$ and $f \in \text{Hom}_{\mathcal{C}}$. Then (C, f) is said to be a
 1. $\mathcal{C}$ -equalizer if there exist $A, B \in \text{Obj}(\mathcal{C})$ and $g, h \in \text{Hom}_{\mathcal{C}}(A, B)$ such that $f \in \text{Hom}_{\mathcal{C}}(C, A)$ and f is a $\mathcal{C}$ -equalizer of g and h .
 2. $\mathcal{C}$ -coequalizer if there exist $A, B \in \text{Obj}(\mathcal{C})$ and $g, h \in \text{Hom}_{\mathcal{C}}(A, B)$ such that $f \in \text{Hom}_{\mathcal{C}}(B, C)$ and f is a $\mathcal{C}$ -coequalizer of g and h .
- We define
 - $\text{Equi}_{\mathcal{C}} := \{(C, f) \in \text{Obj}(\mathcal{C}) \times \text{Hom}_{\mathcal{C}} : (C, f) \text{ is a } \mathcal{C}\text{-equalizer}\}$
 - $\text{Coequi}_{\mathcal{C}} := \{(C, f) \in \text{Obj}(\mathcal{C}) \times \text{Hom}_{\mathcal{C}} : (C, f) \text{ is a } \mathcal{C}\text{-coequalizer}\}$

Exercise 7.3.0.3. Let $\mathcal{C}$ be a category. Then

1. for each $A, B \in \text{Obj}(\mathcal{C})$ and $g, h \in \text{Hom}_{\mathcal{C}^{\text{op}}}(A, B)$,
 - (a) $\text{Equi}_{\mathcal{C}^{\text{op}}}(g, h) = \text{Coequi}_{\mathcal{C}}(g, h)$
 - (b) $\text{Coequi}_{\mathcal{C}^{\text{op}}}(g, h) = \text{Equi}_{\mathcal{C}}(g, h)$
2. (a) $\text{Equi}_{\mathcal{C}^{\text{op}}} = \text{Coequi}_{\mathcal{C}}$
- (b) $\text{Coequi}_{\mathcal{C}^{\text{op}}} = \text{Equi}_{\mathcal{C}}$

Proof.

1. (a) • Let $(C, f) \in \text{Equi}_{\mathcal{C}^{\text{op}}}(f, g)$. By definition (ref def), $C \in \text{Obj}(\mathcal{C}^{\text{op}})$, $f \in \text{Hom}_{\mathcal{C}^{\text{op}}}(C, A)$ and
 - $g \circ_{\mathcal{C}^{\text{op}}} f = h \circ_{\mathcal{C}^{\text{op}}} f$,
 - for each $C' \in \text{Obj}(\mathcal{C}^{\text{op}})$ and $f' \in \text{Hom}_{\mathcal{C}^{\text{op}}}(C', A)$, if $g \circ_{\mathcal{C}^{\text{op}}} f' = h \circ_{\mathcal{C}^{\text{op}}} f'$, then there exists a unique $\phi \in \text{Hom}_{\mathcal{C}^{\text{op}}}(C', C)$ such that $f \circ_{\mathcal{C}^{\text{op}}} \phi = f'$.

We note that $g, h \in \text{Hom}_{\mathcal{C}}(B, A)$ and $f \in \text{Hom}_{\mathcal{C}}(B, C)$.

- i. Thus

$$\begin{aligned} f \circ_{\mathcal{C}} g &= g \circ_{\mathcal{C}^{\text{op}}} f \\ &= h \circ_{\mathcal{C}^{\text{op}}} f \\ &= f \circ_{\mathcal{C}} h. \end{aligned}$$

- ii. Let $C' \in \text{Obj}(\mathcal{C})$ and $f' \in \text{Hom}_{\mathcal{C}}(B, C')$. Suppose that $f' \circ_{\mathcal{C}} g = f' \circ_{\mathcal{C}} h$. Then $f' \in \text{Hom}_{\mathcal{C}^{\text{op}}}(C', B)$ and

$$\begin{aligned} g \circ_{\mathcal{C}^{\text{op}}} f' &= f' \circ_{\mathcal{C}} g \\ &= f' \circ_{\mathcal{C}} h \\ &= h \circ_{\mathcal{C}^{\text{op}}} f'. \end{aligned}$$

Thus there exists a unique $\phi \in \text{Hom}_{\mathcal{C}^{\text{op}}}(C', C)$ such that $f \circ_{\mathcal{C}^{\text{op}}} \phi = f'$. Therefore there exists a unique $\phi \in \text{Hom}_{\mathcal{C}}(C, C')$ such that

$$\begin{aligned} \phi \circ_{\mathcal{C}} f &= f \circ_{\mathcal{C}^{\text{op}}} \phi \\ &= f'. \end{aligned}$$

Since $C' \in \text{Obj}(\mathcal{C})$ and $f' \in \text{Hom}_{\mathcal{C}}(B, C')$ are arbitrary, we have that for each $C' \in \text{Obj}(\mathcal{C})$ and $f' \in \text{Hom}_{\mathcal{C}}(B, C')$, there exists a unique $\phi \in \text{Hom}_{\mathcal{C}}(C, C')$ such that $\phi \circ_{\mathcal{C}} f = f'$.

Thus $(C, f) \in \text{Coequi}_{\mathcal{C}}(f, g)$. Since $(C, f) \in \text{Equi}_{\mathcal{C}^{\text{op}}}(f, g)$ is arbitrary, we have that for each $(C, f) \in \text{Equi}_{\mathcal{C}^{\text{op}}}(f, g)$, $(C, f) \in \text{Coequi}_{\mathcal{C}}(f, g)$. Thus $\text{Equi}_{\mathcal{C}^{\text{op}}}(f, g) \subset \text{Coequi}_{\mathcal{C}}(f, g)$.

- Similarly, $\text{Coequi}_{\mathcal{C}}(f, g) \subset \text{Equi}_{\mathcal{C}^{\text{op}}}(f, g)$. **show details**

Since $\text{Equi}_{\mathcal{C}^{\text{op}}}(f, g) \subset \text{Coequi}_{\mathcal{C}}(f, g)$ and $\text{Coequi}_{\mathcal{C}}(f, g) \subset \text{Equi}_{\mathcal{C}^{\text{op}}}(f, g)$, we have that $\text{Equi}_{\mathcal{C}^{\text{op}}}(f, g) = \text{Coequi}_{\mathcal{C}}(f, g)$

(b) **Similar**

2. **FINISH!!!**

FINISH!!!

□

7.4 Pullbacks and Pushouts

7.5 Exponentials

Definition 7.5.0.1. Let $\mathcal{C}$ be a category and $B, C, D \in \text{Obj}(\mathcal{C})$.

- Suppose that $\mathcal{C}$ has finite products.

- Let $\text{eval} : D \times B \rightarrow C$. Then (D, eval) is said to be a **$\mathcal{C}$ -exponential object of B and C** if for each $A \in \text{Obj}(\mathcal{C})$ and $f \in \text{Hom}_{\mathcal{C}}(A \times B, C)$, there exists a unique $\phi \in \text{Hom}_{\mathcal{C}}(A, D)$ such that $\text{eval} \circ_C (\phi \times \text{id}_B^C) = f$, i.e. the following diagram commutes:

$$\begin{array}{ccc} A & A \times B & \\ \phi \downarrow & \phi \times \text{id}_B^C \downarrow & \searrow f \\ D & D \times B & \xrightarrow{\text{eval}} C \end{array}$$

- We define

$$\text{Exp}_{\mathcal{C}}(B, C) := \left\{ (D, \text{eval}) \in \coprod_{D \in \text{Obj}(\mathcal{C})} \text{Hom}_{\mathcal{C}}(D \times B, C) : (D, \text{eval}) \text{ is a } \mathcal{C}\text{-exponential of } B \text{ and } C \right\}.$$

- Suppose that $\mathcal{C}$ has finite coproducts.

- Let $\text{coeval} : C \rightarrow D + B$. Then (D, coeval) is said to be a **$\mathcal{C}$ -coexponential object of B and C** if for each $A \in \text{Obj}(\mathcal{C})$ and $f \in \text{Hom}_{\mathcal{C}}(C, A + B)$, there exists a unique $\phi \in \text{Hom}_{\mathcal{C}}(D, A)$ such that $(\phi + \text{id}_B^C) \circ_C \text{coeval} = f$, i.e. the following diagram commutes:

$$\begin{array}{ccc} D & D + B & \\ \phi \downarrow & \phi + \text{id}_B^C \downarrow & \nwarrow \text{coeval} \\ A & A + B & \xleftarrow{f} C \end{array}$$

- We define

$$\text{Coexp}_{\mathcal{C}}(B, C) := \left\{ (D, \text{coeval}) \in \coprod_{D \in \text{Obj}(\mathcal{C})} \text{Hom}_{\mathcal{C}}(C, D + B) : (D, \text{coeval}) \text{ is a } \mathcal{C}\text{-coexponential of } B \text{ and } C \right\}.$$

Exercise 7.5.0.2. Let $\mathcal{C}$ be a category, A a set (class?) and $(X_{\alpha})_{\alpha \in A} \in \text{Obj}(\mathcal{C})^A$. Then

1. $\text{Exp}_{\mathcal{C}^{\text{op}}}(B, C) = \text{Coexp}_{\mathcal{C}}(B, C)$.
2. $\text{Coexp}_{\mathcal{C}^{\text{op}}}(B, C) = \text{Exp}_{\mathcal{C}}(B, C)$.

Proof.

1. • Let $(D, \text{eval}) \in \text{Exp}_{\mathcal{C}^{\text{op}}}(B, C)$. We note that $(X_{\alpha})_{\alpha \in A} \in \text{Obj}(\mathcal{C})^A$, $X \in \text{Obj}(\mathcal{C})$ and $(\pi_{\alpha})_{\alpha \in A} \in \prod_{\alpha \in A} \text{Hom}_{\mathcal{C}}(X_{\alpha}, X)$. Let $Y \in \text{Obj}(\mathcal{C})$ and $(i_{\alpha})_{\alpha \in A} \in \prod_{\alpha \in A} \text{Hom}_{\mathcal{C}}(X_{\alpha}, Y)$.

– **Existence:**

Then $Y \in \text{Obj}(\mathcal{C}^{\text{op}})$ and $(i_{\alpha})_{\alpha \in A} \in \prod_{\alpha \in A} \text{Hom}_{\mathcal{C}^{\text{op}}}(Y, X_{\alpha})$. Since $(X, (\pi_{\alpha})_{\alpha \in A}) \in \text{Prod}_{\mathcal{C}^{\text{op}}}((X_{\alpha})_{\alpha \in A})$, there exists a unique $\phi \in \text{Hom}_{\mathcal{C}^{\text{op}}}(Y, X)$ such that for each $\alpha \in A$, $\pi_{\alpha} \circ_{\mathcal{C}^{\text{op}}} \phi = i_{\alpha}$. Then $\phi \in \text{Hom}_{\mathcal{C}}(X, Y)$ and for each $\alpha \in A$, $\phi \circ_C \pi_{\alpha} = i_{\alpha}$.

– **Uniqueness:**

Let $\phi' \in \text{Hom}_{\mathcal{C}}(X, Y)$. Suppose that for each $\alpha \in A$, $\phi' \circ_C \pi_{\alpha} = i_{\alpha}$. Then $\phi' \in \text{Hom}_{\mathcal{C}^{\text{op}}}(Y, X)$ and for each $\alpha \in A$, $\pi_{\alpha} \circ_{\mathcal{C}^{\text{op}}} \phi' = i_{\alpha}$. Uniqueness of ϕ implies that $\phi' = \phi$.

Since $Y \in \text{Obj}(\mathcal{C})$ and $(i_{\alpha})_{\alpha \in A} \in \prod_{\alpha \in A} \text{Hom}_{\mathcal{C}}(X_{\alpha}, Y)$ are arbitrary, we have that for each $Y \in \text{Obj}(\mathcal{C})$ and $(i_{\alpha})_{\alpha \in A} \in \prod_{\alpha \in A} \text{Hom}_{\mathcal{C}}(X_{\alpha}, Y)$, there exists a unique $\phi \in \text{Hom}_{\mathcal{C}}(X, Y)$ such that for each $\alpha \in A$, $\phi \circ_C \pi_{\alpha} = i_{\alpha}$. Thus $(X, (\pi_{\alpha})_{\alpha \in A}) \in \text{Coproduct}_{\mathcal{C}}((X_{\alpha})_{\alpha \in A})$. Since $(X, (\pi_{\alpha})_{\alpha \in A}) \in \text{Prod}_{\mathcal{C}^{\text{op}}}((X_{\alpha})_{\alpha \in A})$ is arbitrary, we have that for each $(X, (\pi_{\alpha})_{\alpha \in A}) \in \text{Prod}_{\mathcal{C}^{\text{op}}}((X_{\alpha})_{\alpha \in A})$, $(X, (\pi_{\alpha})_{\alpha \in A}) \in \text{Coproduct}_{\mathcal{C}}((X_{\alpha})_{\alpha \in A})$. Hence $\text{Prod}_{\mathcal{C}^{\text{op}}}((X_{\alpha})_{\alpha \in A}) \subset \text{Coproduct}_{\mathcal{C}}((X_{\alpha})_{\alpha \in A})$.

- Similar, fill out!!!

Since $\text{Prod}_{\mathcal{C}^{\text{op}}}((X_\alpha)_{\alpha \in A}) \subset \text{Coproduct}_{\mathcal{C}}((X_\alpha)_{\alpha \in A})$ and $\text{Coproduct}_{\mathcal{C}}((X_\alpha)_{\alpha \in A}) \subset \text{Prod}_{\mathcal{C}^{\text{op}}}((X_\alpha)_{\alpha \in A})$, we have that $\text{Prod}_{\mathcal{C}^{\text{op}}}((X_\alpha)_{\alpha \in A}) = \text{Coproduct}_{\mathcal{C}}((X_\alpha)_{\alpha \in A})$.

2. The prev part (ref) implies that

$$\begin{aligned} \text{Coproduct}_{\mathcal{C}^{\text{op}}}((X_\alpha)_{\alpha \in A}) &= \text{Prod}_{(\mathcal{C}^{\text{op}})^{\text{op}}}((X_\alpha)_{\alpha \in A}) \\ &= \text{Prod}_{\mathcal{C}}((X_\alpha)_{\alpha \in A}). \end{aligned}$$

□

Exercise 7.5.0.3. Let $\mathcal{C}$ be a category, A a set (class?) and $(X_\alpha)_{\alpha \in A} \in \text{Obj}(\mathcal{C})^A$. Then

1. for each $(X, (\pi_\alpha^X)_{\alpha \in A}), (Y, (\pi_\alpha^Y)_{\alpha \in A}) \in \text{Prod}_{\mathcal{C}}((X_\alpha)_{\alpha \in A})$, $X \cong_{\mathcal{C}} Y$.
2. for each $(X, (\iota_\alpha^X)_{\alpha \in A}), (Y, (\iota_\alpha^Y)_{\alpha \in A}) \in \text{Coproduct}_{\mathcal{C}}((X_\alpha)_{\alpha \in A})$, $X \cong_{\mathcal{C}} Y$.

Exercise 7.5.0.4. content...

7.6 Monomorphisms and Epimorphisms

7.6.1 Introduction

Definition 7.6.1.1. Let $\mathcal{C}$ be a category and $A, B \in \text{Obj}(\mathcal{C})$ and $f \in \text{Hom}_{\mathcal{C}}(A, B)$. Then f is said to be

- a **monomorphism** if for each $C \in \text{Obj}(\mathcal{C})$ and $g, h \in \text{Hom}_{\mathcal{C}}(C, A)$, $f \circ g = f \circ h$ implies that $g = h$, i.e. we have the following implication of commutative diagrams:

$$\begin{array}{c} C \\ \searrow g \\ A \xrightarrow{f} B \\ \nearrow h \\ C \end{array} \implies \begin{array}{ccc} & g & \\ C & \curvearrowright & A \\ & h & \end{array}$$

- an **epimorphism** if for each $C \in \text{Obj}(\mathcal{C})$ and $g, h \in \text{Hom}_{\mathcal{C}}(B, C)$, $g \circ f = h \circ f$ implies that $g = h$, i.e. we have the following implication of commutative diagrams:

$$\begin{array}{ccc} & & C \\ & g \nearrow & \\ A \xrightarrow{f} B & & \\ & h \searrow & \\ & & C \end{array} \implies \begin{array}{ccc} & g & \\ B & \curvearrowright & C \\ & h & \end{array}$$

make note about common short-hand using parallel arrows

Definition 7.6.1.2. Let $\mathcal{C}$ be a category.

- We define
 - $\text{Mono}_{\mathcal{C}} := \{f \in \text{Hom}_{\mathcal{C}} : f \text{ is a monomorphism}\}$
 - $\text{Epi}_{\mathcal{C}} := \{f \in \text{Hom}_{\mathcal{C}} : f \text{ is an epimorphism}\}$
- Let $A, B \in \text{Obj}(\mathcal{C})$. We define
 - $\text{Mono}_{\mathcal{C}}(A, B) := \text{Mono}_{\mathcal{C}} \cap \text{Hom}_{\mathcal{C}}(A, B)$.
 - $\text{Epi}_{\mathcal{C}}(A, B) := \text{Epi}_{\mathcal{C}} \cap \text{Hom}_{\mathcal{C}}(A, B)$.

Exercise 7.6.1.3. Let $\mathcal{C}$ be a category. Then

- for each $A, B \in \text{Obj}(\mathcal{C})$,
 - $\text{Mono}_{\mathcal{C}^{\text{op}}}(A, B) = \text{Epi}_{\mathcal{C}}(B, A)$
 - $\text{Epi}_{\mathcal{C}^{\text{op}}}(A, B) = \text{Mono}_{\mathcal{C}}(B, A)$
- $\text{Mono}_{\mathcal{C}^{\text{op}}} = \text{Epi}_{\mathcal{C}}$
 - $\text{Epi}_{\mathcal{C}^{\text{op}}} = \text{Mono}_{\mathcal{C}}$

Proof. **FIX!!!**

- Let $A, B \in \text{Obj}(\mathcal{C})$.

- Let $m \in \text{Mono}_{\mathcal{C}^{\text{op}}}(A, B)$. Then $m \in \text{Hom}_{\mathcal{C}}(B, A)$. Let $C \in \text{Obj}(\mathcal{C})$ and $f, g \in \text{Hom}_{\mathcal{C}}(A, C)$. Suppose that $f \circ_{\mathcal{C}} m = g \circ_{\mathcal{C}} m$. Then $m \circ_{\mathcal{C}^{\text{op}}} f = m \circ_{\mathcal{C}^{\text{op}}} g$. Since $m \in \text{Mono}_{\mathcal{C}^{\text{op}}}$, we have that $f = g$. Since $C \in \text{Obj}(\mathcal{C})$ and $f, g \in \text{Hom}_{\mathcal{C}}(A, C)$ are arbitrary, we have that for each $C \in \text{Obj}(\mathcal{C})$ and $f, g \in \text{Hom}_{\mathcal{C}}(A, C)$, $f \circ_{\mathcal{C}} m = g \circ_{\mathcal{C}} m$ implies that $f = g$. Thus $m \in \text{Epi}_{\mathcal{C}}(B, A)$. Since $m \in \text{Mono}_{\mathcal{C}^{\text{op}}}(A, B)$ is arbitrary, we have that for each $m \in \text{Mono}_{\mathcal{C}^{\text{op}}}(A, B)$, $m \in \text{Epi}_{\mathcal{C}}(B, A)$. Hence $\text{Mono}_{\mathcal{C}^{\text{op}}}(A, B) \subset \text{Epi}_{\mathcal{C}}(B, A)$.
- Let $e \in \text{Epi}_{\mathcal{C}}(B, A)$. Then $e \in \text{Hom}_{\mathcal{C}^{\text{op}}}(A, B)$. Let $C \in \text{Obj}(\mathcal{C}^{\text{op}})$ and $f, g \in \text{Hom}_{\mathcal{C}^{\text{op}}}(C, A)$. Suppose that $e \circ_{\mathcal{C}^{\text{op}}} f = e \circ_{\mathcal{C}^{\text{op}}} g$. Then $f \circ_{\mathcal{C}} e = g \circ_{\mathcal{C}} e$. Since $e \in \text{Epi}_{\mathcal{C}}$, we have that $f = g$. Since $C \in \text{Obj}(\mathcal{C}^{\text{op}})$ and $f, g \in \text{Hom}_{\mathcal{C}^{\text{op}}}(C, A)$ are arbitrary, we have that for each $C \in \text{Obj}(\mathcal{C}^{\text{op}})$ and $f, g \in \text{Hom}_{\mathcal{C}^{\text{op}}}(C, A)$, $e \circ_{\mathcal{C}^{\text{op}}} f = e \circ_{\mathcal{C}^{\text{op}}} g$ implies that $f = g$. Thus $e \in \text{Mono}_{\mathcal{C}^{\text{op}}}(A, B)$. Since $e \in \text{Epi}_{\mathcal{C}}(B, A)$ is arbitrary, we have that for each $e \in \text{Epi}_{\mathcal{C}}(B, A)$, $e \in \text{Mono}_{\mathcal{C}^{\text{op}}}(A, B)$. Hence $\text{Epi}_{\mathcal{C}}(B, A) \subset \text{Mono}_{\mathcal{C}^{\text{op}}}(A, B)$.

Since $\text{Mono}_{\mathcal{C}^{\text{op}}}(A, B) \subset \text{Epi}_{\mathcal{C}}(B, A)$ and $\text{Epi}_{\mathcal{C}}(B, A) \subset \text{Mono}_{\mathcal{C}^{\text{op}}}(A, B)$, we have that $\text{Mono}_{\mathcal{C}^{\text{op}}}(A, B) = \text{Epi}_{\mathcal{C}}(B, A)$.

(b)

2. (a) Since $A, B \in \text{Obj}(\mathcal{C})$ are arbitrary, we have that for each $A, B \in \text{Obj}(\mathcal{C})$, $\text{Mono}_{\mathcal{C}^{\text{op}}}(A, B) = \text{Epi}_{\mathcal{C}}(B, A)$. Hence

$$\begin{aligned}
 \text{Mono}_{\mathcal{C}^{\text{op}}} &= \bigcup_{(A, B) \in \text{Obj}(\mathcal{C}^{\text{op}})^2} \text{Mono}_{\mathcal{C}^{\text{op}}}(A, B) \\
 &= \bigcup_{(A, B) \in \text{Obj}(\mathcal{C})^2} \text{Mono}_{\mathcal{C}^{\text{op}}}(A, B) \\
 &= \bigcup_{(B, A) \in \text{Obj}(\mathcal{C})^2} \text{Epi}_{\mathcal{C}}(B, A) \\
 &= \text{Epi}_{\mathcal{C}}.
 \end{aligned}$$

- (b) The previous part (ref needed) implies that

$$\begin{aligned}
 \text{Mono}_{\mathcal{C}} &= \text{Mono}_{(\mathcal{C}^{\text{op}})^{\text{op}}} \\
 &= \text{Epi}_{\mathcal{C}^{\text{op}}}.
 \end{aligned}$$

□

Exercise 7.6.1.4. Let $\mathcal{C}$ be a category and $A, B, C \in \text{Obj}(\mathcal{C})$. Then

1. for each $f \in \text{Mono}_{\mathcal{C}}(A, B)$ and $g \in \text{Mono}_{\mathcal{C}}(B, C)$, $g \circ_{\mathcal{C}} f \in \text{Mono}_{\mathcal{C}}(A, C)$,
2. for each $f \in \text{Epi}_{\mathcal{C}}(A, B)$ and $g \in \text{Epi}_{\mathcal{C}}(B, C)$, $g \circ_{\mathcal{C}} f \in \text{Epi}_{\mathcal{C}}(A, C)$.

Proof.

1. Let $f \in \text{Mono}_{\mathcal{C}}(A, B)$, $g \in \text{Mono}_{\mathcal{C}}(B, C)$, $X \in \text{Obj}(\mathcal{C})$ and $\alpha, \beta \in \text{Hom}_{\mathcal{C}}(X, A)$. We note that since

$$\begin{aligned}
 (g \circ_{\mathcal{C}} f) \circ_{\mathcal{C}} \alpha &= (g \circ_{\mathcal{C}} f) \circ_{\mathcal{C}} \beta \iff g \circ_{\mathcal{C}} (f \circ_{\mathcal{C}} \alpha) = g \circ_{\mathcal{C}} (f \circ_{\mathcal{C}} \beta) \\
 &\implies f \circ_{\mathcal{C}} \alpha = f \circ_{\mathcal{C}} \beta \quad (g \in \text{Mono}_{\mathcal{C}}) \\
 &\implies \alpha = \beta \quad (f \in \text{Mono}_{\mathcal{C}}).
 \end{aligned}$$

Since $X \in \text{Obj}(\mathcal{C})$ and $\alpha, \beta \in \text{Hom}_{\mathcal{C}}(X, A)$ with $(g \circ_{\mathcal{C}} f) \circ_{\mathcal{C}} \alpha = (g \circ_{\mathcal{C}} f) \circ_{\mathcal{C}} \beta$ are arbitrary, we have that for each $X \in \text{Obj}(\mathcal{C})$ and $\alpha, \beta \in \text{Hom}_{\mathcal{C}}(X, A)$, $(g \circ_{\mathcal{C}} f) \circ_{\mathcal{C}} \alpha = (g \circ_{\mathcal{C}} f) \circ_{\mathcal{C}} \beta$ implies that $\alpha = \beta$. Hence $g \circ_{\mathcal{C}} f \in \text{Mono}_{\mathcal{C}}(A, C)$. Since $f \in \text{Mono}_{\mathcal{C}}(A, B)$ and $g \in \text{Mono}_{\mathcal{C}}(B, C)$ are arbitrary, we have that for each $f \in \text{Mono}_{\mathcal{C}}(A, B)$, $g \in \text{Mono}_{\mathcal{C}}(B, C)$, $g \circ_{\mathcal{C}} f \in \text{Mono}_{\mathcal{C}}(A, C)$.

2. Let $f \in \text{Epi}_{\mathcal{C}}(A, B)$ and $g \in \text{Epi}_{\mathcal{C}}(B, C)$. Exercise 10.1.0.10 (ref prev ex) implies that $f \in \text{Mono}_{\mathcal{C}^{\text{op}}}(B, A)$ and $g \in \text{Mono}_{\mathcal{C}^{\text{op}}}(C, B)$, (ref prev part) then implies that

$$\begin{aligned}
 g \circ_{\mathcal{C}} f &= f \circ_{\mathcal{C}^{\text{op}}} g \\
 &\in \text{Mono}_{\mathcal{C}^{\text{op}}} \\
 &= \text{Epi}_{\mathcal{C}}(A, C).
 \end{aligned}$$

Since $f \in \text{Epi}_{\mathcal{C}}(A, B)$ and $g \in \text{Epi}_{\mathcal{C}}(B, C)$ are arbitrary, we have that for each $f \in \text{Epi}_{\mathcal{C}}(A, B)$ and $g \in \text{Epi}_{\mathcal{C}}(B, C)$, $g \circ_{\mathcal{C}} f \in \text{Epi}_{\mathcal{C}}(A, C)$.

□

Exercise 7.6.1.5. Let $\mathcal{C}$ be a category, $A, B, C \in \text{Obj}(\mathcal{C})$ and $f \in \text{Hom}_{\mathcal{C}}(A, B)$ and $g \in \text{Hom}_{\mathcal{C}}(B, C)$. Then

1. $g \in \text{Mono}_{\mathcal{C}}$ implies that $g \circ_{\mathcal{C}} f \in \text{Mono}_{\mathcal{C}}$ iff $f \in \text{Mono}_{\mathcal{C}}$.
2. $f \in \text{Epi}_{\mathcal{C}}$ implies that $g \circ_{\mathcal{C}} f \in \text{Epi}_{\mathcal{C}}$ iff $g \in \text{Epi}_{\mathcal{C}}$.

Proof.

1. Suppose that $g \in \text{Mono}_{\mathcal{C}}$.

- ($\implies$):
Suppose that $g \circ_{\mathcal{C}} f \in \text{Mono}_{\mathcal{C}}$. Let $D \in \text{Obj}(\mathcal{C})$ and $\alpha, \beta \in \text{Hom}_{\mathcal{C}}(D, A)$. Suppose that $f \circ_{\mathcal{C}} \alpha = f \circ_{\mathcal{C}} \beta$. Then

$$\begin{aligned} (g \circ_{\mathcal{C}} f) \circ_{\mathcal{C}} \alpha &= g \circ_{\mathcal{C}} (f \circ_{\mathcal{C}} \alpha) \\ &= g \circ_{\mathcal{C}} (f \circ_{\mathcal{C}} \beta) \\ &= (g \circ_{\mathcal{C}} f) \circ_{\mathcal{C}} \beta. \end{aligned}$$

Since $g \circ_{\mathcal{C}} f \in \text{Mono}_{\mathcal{C}}$, $\alpha = \beta$. Since $D \in \text{Obj}(\mathcal{C})$ and $\alpha, \beta \in \text{Hom}_{\mathcal{C}}(D, A)$ with $f \circ_{\mathcal{C}} \alpha = f \circ_{\mathcal{C}} \beta$ are arbitrary, we have that for each $D \in \text{Obj}(\mathcal{C})$ and $\alpha, \beta \in \text{Hom}_{\mathcal{C}}(D, A)$, $f \circ_{\mathcal{C}} \alpha = f \circ_{\mathcal{C}} \beta$ implies that $\alpha = \beta$. Thus $f \in \text{Mono}_{\mathcal{C}}$.

- ($\impliedby$):
Suppose that $f \in \text{Mono}_{\mathcal{C}}$. Since $g \in \text{Mono}_{\mathcal{C}}$, Exercise 10.1.0.10 (ref prev ex) implies that $g \circ_{\mathcal{C}} f \in \text{Mono}_{\mathcal{C}}$.

2. Suppose that $f \in \text{Epi}_{\mathcal{C}}$. Exercise 10.1.0.10 (ref prev ex about duality) implies that $f \in \text{Mono}_{\mathcal{C}^{\text{op}}}$. Then

$$\begin{aligned} g \circ_{\mathcal{C}} f \in \text{Epi}_{\mathcal{C}} &\iff g \circ_{\mathcal{C}} f \in \text{Mono}_{\mathcal{C}^{\text{op}}} \quad (\text{ref prev ex about duality}) \\ &\iff f \circ_{\mathcal{C}^{\text{op}}} g \in \text{Mono}_{\mathcal{C}^{\text{op}}} \\ &\iff g \in \text{Mono}_{\mathcal{C}^{\text{op}}} \quad (\text{ref prev part}) \\ &\iff g \in \text{Epi}_{\mathcal{C}} \quad (\text{ref prev ex about duality}). \end{aligned}$$

□

Exercise 7.6.1.6. Let $\mathcal{C}$ be a category and $A, B \in \text{Obj}(\mathcal{C})$. Then $\text{Iso}_{\mathcal{C}}(A, B) \subset \text{Mono}_{\mathcal{C}}(A, B) \cap \text{Epi}_{\mathcal{C}}(A, B)$.

Proof. Let $f \in \text{Iso}_{\mathcal{C}}(A, B)$.

- (monomorphism):
Let $C \in \text{Obj}(\mathcal{C})$ and $g, h \in \text{Hom}_{\mathcal{C}}(C, A)$. Suppose that $f \circ_{\mathcal{C}} g = f \circ_{\mathcal{C}} h$. Then

$$\begin{aligned} g &= \text{id}_A \circ_{\mathcal{C}} g \\ &= (f^{-1} \circ_{\mathcal{C}} f) \circ_{\mathcal{C}} g \\ &= f^{-1} \circ_{\mathcal{C}} (f \circ_{\mathcal{C}} g) \\ &= f^{-1} \circ_{\mathcal{C}} (f \circ_{\mathcal{C}} h) \\ &= (f^{-1} \circ_{\mathcal{C}} f) \circ_{\mathcal{C}} h \\ &= \text{id}_A \circ_{\mathcal{C}} h \\ &= h \end{aligned}$$

Since $g, h \in \text{Hom}_{\mathcal{C}}(C, A)$ with $f \circ_{\mathcal{C}} g = f \circ_{\mathcal{C}} h$ are arbitrary, we have that for each $g, h \in \text{Hom}_{\mathcal{C}}(C, A)$, $f \circ_{\mathcal{C}} g = f \circ_{\mathcal{C}} h$ implies that $g = h$. Therefore $f \in \text{Mono}_{\mathcal{C}}(A, B)$.

- (epimorphism):

Let $C \in \text{Obj}(\mathcal{C})$ and $g, h \in \text{Hom}_{\mathcal{C}}(B, C)$. Suppose that $g \circ_C f = h \circ_C f$. Then

$$\begin{aligned}
 g &= g \circ_C \text{id}_B \\
 &= g \circ_C (f \circ_C f^{-1}) \\
 &= (g \circ_C f) \circ_C f^{-1} \\
 &= (h \circ_C f) \circ_C f^{-1} \\
 &= h \circ_C (f \circ_C f^{-1}) \\
 &= h \circ_C \text{id}_B \\
 &= h
 \end{aligned}$$

Since $g, h \in \text{Hom}_{\mathcal{C}}(B, C)$ with $g \circ_C f = h \circ_C f$ are arbitrary, we have that for each $g, h \in \text{Hom}_{\mathcal{C}}(B, C)$, $g \circ_C f = h \circ_C f$ implies that $g = h$. Therefore $f \in \text{Epi}_{\mathcal{C}}(A, B)$.

Thus $f \in \text{Mono}_{\mathcal{C}}(A, B) \cap \text{Epi}_{\mathcal{C}}(A, B)$. Since $f \in \text{Iso}_{\mathcal{C}}(A, B)$ is arbitrary, we have that for each $f \in \text{Iso}_{\mathcal{C}}(A, B)$, $f \in \text{Mono}_{\mathcal{C}}(A, B) \cap \text{Epi}_{\mathcal{C}}(A, B)$. Hence $\text{Iso}_{\mathcal{C}}(A, B) \subset \text{Mono}_{\mathcal{C}}(A, B) \cap \text{Epi}_{\mathcal{C}}(A, B)$. $\square$

Exercise 7.6.1.7. Let $\mathcal{C}$ be a category and $X \in \text{Obj}(\mathcal{C})$. Then $\text{id}_X \in \text{Mono}_{\mathcal{C}} \cap \text{Epi}_{\mathcal{C}}$.

Proof. [ref prev ex in isomorphism section](#) implies that $\text{id}_X \in \text{Aut}_{\mathcal{C}}(X)$. [ref prev ex](#) implies that $\text{id}_X \in \text{Mono}_{\mathcal{C}} \cap \text{Epi}_{\mathcal{C}}$. $\square$

Exercise 7.6.1.8. [mono/epi in Set](#) Let $A, B \in \text{Obj}(\mathbf{Set})$ and $f \in \text{Hom}_{\mathbf{Set}}(A, B)$. Then

1. f is a monomorphism iff f is injective
2. f is an epimorphism iff f is surjective

Hint: consider $C := \{0\}$ and $C := \{0, 1\}$.

Proof.

1. • ($\implies$):
Suppose that f is injective. Let $C \in \text{Obj}(\mathbf{Set})$ and $g, h \in \text{Hom}_{\mathbf{Set}}(C, A)$. Suppose that $f \circ_{\mathbf{Set}} g = f \circ_{\mathbf{Set}} h$. Let $x \in C$. Then $f(g(x)) = f(h(x))$. Injectivity of f implies that $g(x) = h(x)$. Since $x \in C$ is arbitrary, $g = h$. Hence $f \in \text{Mono}_{\mathbf{Set}}$.
- ($\impliedby$):
Conversely, suppose that $f \in \text{Mono}_{\mathbf{Set}}$. Let $a, b \in A$. Suppose that $f(a) = f(b)$. Set $C = \{0\}$ and define $g, h : C \rightarrow A$ by $g(0) = a$ and $h(0) = b$. Then

$$\begin{aligned}
 f \circ_{\mathbf{Set}} g(0) &= f(g(0)) \\
 &= f(a) \\
 &= f(b) \\
 &= f(h(0)) \\
 &= f \circ_{\mathbf{Set}} h(0)
 \end{aligned}$$

Therefore $f \circ_{\mathbf{Set}} g = f \circ_{\mathbf{Set}} h$. Since $f \in \text{Mono}_{\mathbf{Set}}$, we have that $g = h$. Hence

$$\begin{aligned}
 a &= g(0) \\
 &= h(0) \\
 &= b
 \end{aligned}$$

Since $a, b \in A$ with $f(a) = f(b)$ are arbitrary, we have that for each $a, b \in A$, $f(a) = f(b)$ implies that $a = b$. Thus f is injective.

2. • ($\Rightarrow$):

Suppose that f is surjective. Let $C \in \text{Obj}(\mathcal{C})$ and $g, h \in \text{Hom}_{\mathbf{Set}}(B, C)$. Suppose that $g \circ_{\mathbf{Set}} f = h \circ_{\mathbf{Set}} f$. Let $y \in B$. Surjective of f implies that there exists $x \in A$ such that $y = f(x)$. Then

$$\begin{aligned} g(y) &= g(f(x)) \\ &= g \circ_{\mathbf{Set}} f(x) \\ &= h \circ_{\mathbf{Set}} f(x) \\ &= h(f(x)) \\ &= h(y) \end{aligned}$$

Since $y \in B$ is arbitrary, we have that for each $y \in B$, $g(y) = h(y)$. Hence $g = h$. Since $C \in \text{Obj}(\mathcal{C})$ and $g, h \in \text{Hom}_{\mathbf{Set}}(B, C)$ with $g \circ_{\mathbf{Set}} f = h \circ_{\mathbf{Set}} f$ are arbitrary, we have that for each $C \in \text{Obj}(\mathcal{C})$ and $g, h \in \text{Hom}_{\mathbf{Set}}(B, C)$, $g \circ_{\mathbf{Set}} f = h \circ_{\mathbf{Set}} f$ implies that $g = h$. Hence f is an epimorphism.

• ($\Leftarrow$):

Conversely, suppose that f is an epimorphism. Define $C \in \text{Obj}(\mathbf{Set})$ and $g, h : B \rightarrow C$ by $C := \{0, 1\}$, $g := \chi_{f(A)}$ and $h := \chi_B$. Then $g \circ_{\mathbf{Set}} f = h \circ_{\mathbf{Set}} f$. Since $f \in \text{Epi}_{\mathcal{C}}$, $g = h$ and $f(A) = B$. Hence f is surjective.

□

7.6.2 Extremal Monomorphisms and Epimorphisms

Definition 7.6.2.1. Let $\mathcal{C}$ be a category, $A, B \in \text{Obj}(\mathcal{C})$ and $f \in \text{Hom}_{\mathcal{C}}(A, B)$. Then f is said to be an

• **extremal monomorphism** if

- $f \in \text{Mono}_{\mathcal{C}}$
- for each $C \in \text{Obj}(\mathcal{C})$, $e \in \text{Epi}_{\mathcal{C}}(A, C)$ and $g \in \text{Hom}_{\mathcal{C}}(C, B)$, $f = g \circ_{\mathcal{C}} e$ implies that $e \in \text{Iso}_{\mathcal{C}}(A, C)$.

• **extremal epimorphism** if

- $f \in \text{Epi}_{\mathcal{C}}$
- for each $C \in \text{Obj}(\mathcal{C})$, $g \in \text{Hom}_{\mathcal{C}}(A, C)$ and $m \in \text{Mono}_{\mathcal{C}}(C, B)$, $f = m \circ_{\mathcal{C}} g$ implies that $m \in \text{Iso}_{\mathcal{C}}(C, B)$.

Definition 7.6.2.2. Let $\mathcal{C}$ be a category.

• We define

- $\text{ExtMono}_{\mathcal{C}} := \{f \in \text{Hom}_{\mathcal{C}} : f \text{ is an extremal monomorphism}\},$
- $\text{ExtEpi}_{\mathcal{C}} := \{f \in \text{Hom}_{\mathcal{C}} : f \text{ is an extremal epimorphism}\}.$

• Let $A, B \in \text{Obj}(\mathcal{C})$. We define

- $\text{ExtMono}_{\mathcal{C}}(A, B) := \text{ExtMono}_{\mathcal{C}} \cap \text{Hom}_{\mathcal{C}}(A, B),$
- $\text{ExtEpi}_{\mathcal{C}}(A, B) := \text{ExtEpi}_{\mathcal{C}} \cap \text{Hom}_{\mathcal{C}}(A, B).$

Exercise 7.6.2.3. Let $\mathcal{C}$ be a category. Then

1. for each $A, B \in \text{Obj}(\mathcal{C})$,
 - (a) $\text{ExtMono}_{\mathcal{C}^{\text{op}}}(A, B) = \text{ExtEpi}_{\mathcal{C}}(B, A)$
 - (b) $\text{ExtEpi}_{\mathcal{C}^{\text{op}}}(A, B) = \text{ExtMono}_{\mathcal{C}}(B, A)$
2.
 - (a) $\text{ExtMono}_{\mathcal{C}^{\text{op}}} = \text{ExtEpi}_{\mathcal{C}}$
 - (b) $\text{ExtEpi}_{\mathcal{C}^{\text{op}}} = \text{ExtMono}_{\mathcal{C}}$

Proof. **FIX!!!**

1. Let $A, B \in \text{Obj}(\mathcal{C})$.

- Let $f \in \text{ExtMono}_{\mathcal{C}^{\text{op}}}(A, B)$.
 - Then $f \in \text{Mono}_{\mathcal{C}^{\text{op}}}(A, B)$. [ref prev ex in section on monomorphisms](#) implies that $f \in \text{Epi}_{\mathcal{C}}(B, A)$.
 - Let $C \in \text{Obj}(\mathcal{C})$, $g \in \text{Hom}_{\mathcal{C}}(B, C)$ and $m \in \text{Mono}_{\mathcal{C}}(C, A)$. Suppose that $f = m \circ_{\mathcal{C}} g$. Then $g \in \text{Hom}_{\mathcal{C}^{\text{op}}}(C, B)$, [a prev ex in section on monomorphisms](#) implies that $m \in \text{Epi}_{\mathcal{C}^{\text{op}}}(A, C)$ and $f = g \circ_{\mathcal{C}^{\text{op}}} m$. Since $f \in \text{ExtMono}_{\mathcal{C}^{\text{op}}}(A, B)$, we have that $m \in \text{Iso}_{\mathcal{C}^{\text{op}}}(B, C)$. Exercise 10.1.0.10 ([make and ref prev ex that \$\text{Iso}_{\mathcal{C}} = \text{Iso}_{\mathcal{C}^{\text{op}}}\$](#)) implies that $m \in \text{Iso}_{\mathcal{C}}(C, B)$. Since $C \in \text{Obj}(\mathcal{C})$, $g \in \text{Hom}_{\mathcal{C}}(B, C)$ and $m \in \text{Mono}_{\mathcal{C}}(C, A)$ are arbitrary, we have that for each $C \in \text{Obj}(\mathcal{C})$, $g \in \text{Hom}_{\mathcal{C}}(B, C)$ and $m \in \text{Mono}_{\mathcal{C}}(C, A)$, $f = m \circ_{\mathcal{C}} g$ implies that $m \in \text{Iso}_{\mathcal{C}}(C, A)$.

Thus $f \in \text{ExtEpi}_{\mathcal{C}}(B, A)$. Since $f \in \text{ExtMono}_{\mathcal{C}^{\text{op}}}(A, B)$ is arbitrary, we have that for each $f \in \text{ExtMono}_{\mathcal{C}^{\text{op}}}(A, B)$, $f \in \text{ExtEpi}_{\mathcal{C}}(B, A)$. Hence $\text{ExtMono}_{\mathcal{C}^{\text{op}}}(A, B) \subset \text{ExtEpi}_{\mathcal{C}}(B, A)$.

- Let $f \in \text{ExtEpi}_{\mathcal{C}}(B, A)$.
 - Then $f \in \text{Epi}_{\mathcal{C}}(B, A)$. [ref prev ex in section on monomorphisms](#) implies that $f \in \text{Mono}_{\mathcal{C}^{\text{op}}}(A, B)$.
 - Let $C \in \text{Obj}(\mathcal{C}^{\text{op}})$, $e \in \text{Epi}_{\mathcal{C}^{\text{op}}}(A, C)$ and $g \in \text{Hom}_{\mathcal{C}^{\text{op}}}(C, B)$. Suppose that $f = g \circ_{\mathcal{C}^{\text{op}}} e$. Then $g \in \text{Hom}_{\mathcal{C}}(B, C)$, [a prev ex in section on monomorphisms](#) implies that $e \in \text{Mono}_{\mathcal{C}}(C, A)$ and $f = e \circ_{\mathcal{C}} g$. Since $f \in \text{ExtEpi}_{\mathcal{C}}(B, A)$, we have that $e \in \text{Iso}_{\mathcal{C}}(C, A)$. Exercise 10.1.0.10 ([make and ref prev ex that \$\text{Iso}_{\mathcal{C}} = \text{Iso}_{\mathcal{C}^{\text{op}}}\$](#)) implies that $e \in \text{Iso}_{\mathcal{C}^{\text{op}}}(A, C)$. Since $C \in \text{Obj}(\mathcal{C}^{\text{op}})$, $e \in \text{Epi}_{\mathcal{C}^{\text{op}}}(A, C)$ and $g \in \text{Hom}_{\mathcal{C}^{\text{op}}}(C, B)$ are arbitrary, we have that for each $C \in \text{Obj}(\mathcal{C}^{\text{op}})$, $e \in \text{Epi}_{\mathcal{C}^{\text{op}}}(A, C)$ and $g \in \text{Hom}_{\mathcal{C}^{\text{op}}}(C, B)$, $f = g \circ_{\mathcal{C}^{\text{op}}} e$ implies that $e \in \text{Iso}_{\mathcal{C}^{\text{op}}}(A, C)$.

Thus $f \in \text{ExtMono}_{\mathcal{C}^{\text{op}}}(A, B)$. Since $f \in \text{ExtEpi}_{\mathcal{C}}(B, A)$ is arbitrary, we have that for each $f \in \text{ExtMono}_{\mathcal{C}^{\text{op}}}(A, B)$, $f \in \text{ExtEpi}_{\mathcal{C}}(B, A)$. Hence $\text{ExtMono}_{\mathcal{C}^{\text{op}}}(A, B) \subset \text{ExtEpi}_{\mathcal{C}}(B, A)$.

Since $\text{ExtMono}_{\mathcal{C}^{\text{op}}}(A, B) \subset \text{ExtEpi}_{\mathcal{C}}(B, A)$ and $\text{ExtMono}_{\mathcal{C}^{\text{op}}}(A, B) \subset \text{ExtEpi}_{\mathcal{C}}(B, A)$, we have that $\text{ExtMono}_{\mathcal{C}^{\text{op}}}(A, B) = \text{ExtEpi}_{\mathcal{C}}(B, A)$. Since $A, B \in \text{Obj}(\mathcal{C})$ are arbitrary, we have that for each $A, B \in \text{Obj}(\mathcal{C})$, $\text{ExtMono}_{\mathcal{C}^{\text{op}}}(A, B) = \text{ExtEpi}_{\mathcal{C}}(B, A)$. Therefore

$$\begin{aligned}
 \text{ExtMono}_{\mathcal{C}^{\text{op}}} &= \bigcup_{(A, B) \in \text{Obj}(\mathcal{C}^{\text{op}})^2} \text{ExtMono}_{\mathcal{C}^{\text{op}}}(A, B) \\
 &= \bigcup_{(A, B) \in \text{Obj}(\mathcal{C})^2} \text{ExtMono}_{\mathcal{C}^{\text{op}}}(A, B) \\
 &= \bigcup_{(B, A) \in \text{Obj}(\mathcal{C})^2} \text{ExtEpi}_{\mathcal{C}}(B, A) \\
 &= \text{ExtEpi}_{\mathcal{C}}.
 \end{aligned}$$

2. [The previous part \(ref needed\)](#) implies that

$$\begin{aligned}
 \text{ExtMono}_{\mathcal{C}} &= \text{ExtMono}_{(\mathcal{C}^{\text{op}})^{\text{op}}} \\
 &= \text{ExtEpi}_{\mathcal{C}^{\text{op}}}.
 \end{aligned}$$

□

7.6.3 Regular Monomorphisms and Epimorphisms

Exercise 7.6.3.1. Let $\mathcal{C}$ be a category, $A, B \in \text{Obj}(\mathcal{C})$ and $f, g \in \text{Hom}_{\mathcal{C}}(A, B)$. [Rewrite using notation \$\text{Equi}_{\mathcal{C}}\$](#)

1. Let $M \in \text{Obj}(\mathcal{C})$ and $m \in \text{Hom}_{\mathcal{C}}(M, A)$. If (M, m) is an equalizer of f and g , then $m \in \text{Mono}_{\mathcal{C}}(M, A)$.
2. Let $E \in \text{Obj}(\mathcal{C})$ and $e \in \text{Hom}_{\mathcal{C}}(B, E)$. If (E, e) is a coequalizer of f and g , then $e \in \text{Epi}_{\mathcal{C}}(B, E)$.

Proof.

1. Suppose that (M, m) is an equalizer of f and g . Let $C \in \text{Obj}(\mathcal{C})$ and $\alpha, \beta \in \text{Hom}_{\mathcal{C}}(C, M)$. Suppose that $m \circ_C \alpha = m \circ_C \beta$. Since (M, m) is an equalizer of f and g , we have that $f \circ_C m = g \circ_C m$, i.e. the following diagram commutes:

$$\begin{array}{ccc} M & \xrightarrow{m} & A \\ & & \swarrow f \\ & & B \\ & & \searrow g \\ & & B \end{array}$$

Define $M' \in \text{Obj}(\mathcal{C})$ and $m' \in \text{Hom}_{\mathcal{C}}(M', A)$ by $M' := C$ and $m' := m \circ_C \alpha$. Then

$$\begin{aligned} f \circ_C m' &= f \circ_C (m \circ_C \alpha) \\ &= (f \circ_C m) \circ_C \alpha \\ &= (g \circ_C m) \circ_C \alpha \\ &= g \circ_C (m \circ_C \alpha) \\ &= g \circ_C m'. \end{aligned}$$

Since (E, m) is an equalizer of f and g , there exists a unique $\phi \in \text{Hom}_{\mathcal{C}}(E', E)$ such that $m \circ_C \phi = m'$, i.e. the following diagram commutes:

$$\begin{array}{ccc} E' & \xrightarrow{m'} & A \\ \downarrow \phi & & \swarrow f \\ E & \xrightarrow{m} & B \\ & & \searrow g \\ & & B \end{array}$$

Since $m \circ_C \alpha = m'$, uniqueness of ϕ implies that $\phi = \alpha$. Since

$$\begin{aligned} m \circ_C \beta &= m \circ_C \alpha \\ &= m', \end{aligned}$$

uniqueness of ϕ implies that $\phi = \beta$. Hence $\alpha = \beta$. Since $C \in \text{Obj}(\mathcal{C})$ and $\alpha, \beta \in \text{Hom}_{\mathcal{C}}(C, E)$ with $m \circ_C \alpha = m \circ_C \beta$ are arbitrary, we have that for each $C \in \text{Obj}(\mathcal{C})$ and $\alpha, \beta \in \text{Hom}_{\mathcal{C}}(C, E)$, $m \circ_C \alpha = m \circ_C \beta$ implies that $\alpha = \beta$, i.e. we have the following implication of commutative diagrams:

$$\begin{array}{ccc} C & & \\ \searrow \alpha & & \\ & E & \xrightarrow{m} A \\ \nearrow \beta & & \\ C & & \end{array} \implies \begin{array}{ccc} & \alpha & \\ C & \xrightarrow{\quad} & E \\ & \beta & \end{array}$$

Thus $m \in \text{Mono}_{\mathcal{C}}(E, A)$.

2. **FINISH!!!**

□

Definition 7.6.3.2. Let $\mathcal{C}$ be a category, $A, B \in \text{Obj}(\mathcal{C})$ and $f \in \text{Hom}_{\mathcal{C}}(A, B)$. Then f is said to be a

- **regular monomorphism** if there exists $C \in \text{Obj}(\mathcal{C})$ and $g, h \in \text{Hom}_{\mathcal{C}}(B, C)$ such that $(A, f) \in \text{Equi}_{\mathcal{C}}(g, h)$.

- **regular epimorphism** if there exists $C \in \text{Obj}(\mathcal{C})$ and $g, h \in \text{Hom}_{\mathcal{C}}(C, A)$ such that $(B, f) \in \text{Coequi}_{\mathcal{C}}(g, h)$.

Definition 7.6.3.3. Let $\mathcal{C}$ be a category.

- We define
 - $\text{RegMono}_{\mathcal{C}} := \{f \in \text{Hom}_{\mathcal{C}} : f \text{ is a regular monomorphism}\},$
 - $\text{RegEpi}_{\mathcal{C}} := \{f \in \text{Hom}_{\mathcal{C}} : f \text{ is a regular epimorphism}\}.$
- Let $A, B \in \text{Obj}(\mathcal{C})$. We define
 - $\text{RegMono}_{\mathcal{C}}(A, B) := \text{RegMono}_{\mathcal{C}} \cap \text{Hom}_{\mathcal{C}}(A, B),$
 - $\text{RegEpi}_{\mathcal{C}}(A, B) := \text{RegEpi}_{\mathcal{C}} \cap \text{Hom}_{\mathcal{C}}(A, B).$

Exercise 7.6.3.4. Let $\mathcal{C}$ be a category. Then

1. for each $A, B \in \text{Obj}(\mathcal{C})$,
 - (a) $\text{RegMono}_{\mathcal{C}^{\text{op}}}(A, B) = \text{RegEpi}_{\mathcal{C}}(B, A)$
 - (b) $\text{RegEpi}_{\mathcal{C}^{\text{op}}}(A, B) = \text{RegMono}_{\mathcal{C}}(B, A)$
2.
 - (a) $\text{RegMono}_{\mathcal{C}^{\text{op}}} = \text{RegEpi}_{\mathcal{C}}$
 - (b) $\text{RegEpi}_{\mathcal{C}^{\text{op}}} = \text{RegMono}_{\mathcal{C}}$

Proof. **FINISH!!!** □

Exercise 7.6.3.5. Let $\mathcal{C}$ be a category. Then for each $A, B \in \text{Obj}(\mathcal{C})$,

1. $\text{RegMono}_{\mathcal{C}}(A, B) \subset \text{ExtMono}_{\mathcal{C}}(A, B),$
2. $\text{RegEpi}_{\mathcal{C}}(A, B) \subset \text{ExtEpi}_{\mathcal{C}}(A, B).$

Proof. Let $A, B \in \text{Obj}(\mathcal{C})$. **FINISH!!!**

- 1.
 - 2.
-

Exercise 7.6.3.6. Let $A, B \in \text{Obj}(\mathbf{Set})$ and $f, g \in \text{Hom}_{\mathbf{Set}}(A, B)$. Then $(\text{Equi}(f, g), \iota_{\text{Equi}(f, g)}) \in \text{Equi}_{\mathbf{Set}}(f, g)$.

Proof. Define $E \in \text{Obj}(\mathbf{Set})$ by $E := \text{Equi}(f, g)$.

1. By definition, $E = \{x \in A : f(x) = g(x)\}$. Therefore for each $x \in E$,

$$\begin{aligned}
 f \circ_{\mathbf{Set}} \iota_E(x) &= f(\iota_E(x)) \\
 &= f(x) \\
 &= g(x) \\
 &= g(\iota_E(x)) \\
 &= g \circ_{\mathbf{Set}} \iota_E(x).
 \end{aligned}$$

Thus $f \circ_{\mathbf{Set}} \iota_E = g \circ_{\mathbf{Set}} \iota_E$.

2. Let $E' \in \text{Obj}(\mathbf{Set})$ and $e' \in \text{Hom}_{\mathbf{Set}}(E', A)$. Suppose that $f \circ_{\mathbf{Set}} e' = g \circ_{\mathbf{Set}} e'$. Let $x \in E'$. Define $a \in A$ by $a := e'(x)$. Then

$$\begin{aligned} f(a) &= f(e'(x)) \\ &= f \circ_{\mathbf{Set}} e'(x) \\ &= f \circ_{\mathbf{Set}} e'(x) \\ &= g(e'(x)) \\ &= g(a). \end{aligned}$$

Thus $a \in \text{Equi}(f, g)$. Therefore

$$\begin{aligned} e'(x) &= a \\ &\in \text{Equi}(f, g) \\ &= E. \end{aligned}$$

Since $x \in E'$ is arbitrary, we have that for each $x \in E'$, $e'(x) \in E$. Define $\phi \in \text{Hom}_{\mathbf{Set}}(E', E)$ by $\phi(x) := e'(x)$.

- For each $x \in E'$,

$$\begin{aligned} \iota_E \circ_{\mathbf{Set}} \phi(x) &= \iota_E(\phi(x)) \\ &= \iota_E(e'(x)) \\ &= e'(x). \end{aligned}$$

Thus $\iota_E \circ_{\mathbf{Set}} \phi = e'$.

- Let $\phi' \in \text{Hom}_{\mathbf{Set}}(E', E)$. Suppose that $\iota_E \circ_{\mathbf{Set}} \phi' = e'$. Then for each $x \in E'$,

$$\begin{aligned} \phi'(x) &= \iota_E(\phi'(x)) \\ &= \iota_E \circ_{\mathbf{Set}} \phi'(x) \\ &= e'(x) \\ &= \iota_E \circ_{\mathbf{Set}} \phi(x) \\ &= \iota_E(\phi(x)) &= \phi(x). \end{aligned}$$

Hence $\phi' = \phi$. Since $\phi' \in \text{Hom}_{\mathbf{Set}}(E', E)$ with $\iota_E \circ_{\mathbf{Set}} \phi' = e'$ is arbitrary, we have that for each $\phi' \in \text{Hom}_{\mathbf{Set}}(E', E)$, $\iota_E \circ_{\mathbf{Set}} \phi' = e'$ implies that $\phi' = \phi$.

Thus there exists a unique $\phi \in \text{Hom}_{\mathbf{Set}}(E', E)$ such that $\iota_E \circ_{\mathbf{Set}} \phi = e'$. Since $E' \in \text{Obj}(\mathbf{Set})$ and $e' \in \text{Hom}_{\mathbf{Set}}(E', A)$ with $f \circ_{\mathbf{Set}} e' = g \circ_{\mathbf{Set}} e'$ are arbitrary, we have that for each $E' \in \text{Obj}(\mathbf{Set})$ and $e' \in \text{Hom}_{\mathbf{Set}}(E', A)$, $f \circ_{\mathbf{Set}} e' = g \circ_{\mathbf{Set}} e'$ implies that there exists a unique $\phi \in \text{Hom}_{\mathbf{Set}}(E', E)$ such that $\iota_E \circ_{\mathbf{Set}} \phi = e'$.

Thus $(E, \iota_E) \in \text{Equi}_{\mathbf{Set}}(f, g)$. □

Exercise 7.6.3.7. We have that $\text{RegMono}_{\mathbf{Set}} = \text{Mono}_{\mathbf{Set}}$.

Proof.

- **a prev ex** implies that $\text{RegMono}_{\mathbf{Set}} \subset \text{Mono}_{\mathbf{Set}}$.
- Let $m \in \text{Mono}_{\mathbf{Set}}$. Define $M, A \in \text{Obj}(\mathbf{Set})$ by $M := \text{dom}_{\mathbf{Set}}(m)$ and $A := \text{cod}_{\mathbf{Set}}(m)$. **a prev ex** implies that m is injective. Define $B \in \text{Obj}(\mathbf{Set})$ and $f, g \in \text{Hom}_{\mathbf{Set}}(A, B)$ by $B := m(M) \cup \{0, 1\}$ and

$$f(x) := \begin{cases} x, & x \in m(M) \\ 0, & x \in m(M)^c \end{cases}, \quad g(x) := \begin{cases} x, & x \in m(M) \\ 1, & x \in m(M)^c \end{cases}.$$


Define $E \in \text{Obj}(\mathbf{Set})$ by $E := \text{Equi}(f, g)$. Then

$$\begin{aligned} E &= \text{Equi}(f, g) \\ &= \{x \in A : f(x) = g(x)\} \\ &= m(M). \end{aligned}$$

1. By construction, for each $x \in M$,


$$\begin{aligned} f \circ_{\mathbf{Set}} m(x) &= f(m(x)) \\ &= g(m(x)) \quad (\text{since } m(x) \in E) \\ &= g \circ_{\mathbf{Set}} m(x). \end{aligned}$$

Thus $f \circ_{\mathbf{Set}} m = g \circ_{\mathbf{Set}} m$, i.e. the following diagram commutes:



2. Let $M' \in \mathbf{Obj}(\mathbf{Set})$ and $m' \in \mathbf{Hom}_{\mathbf{Set}}(M', A)$. Suppose that $f \circ_{\mathbf{Set}} m' = g \circ_{\mathbf{Set}} m'$. Exercise 10.1.0.10 [ref prev ex](#) implies that $(E, \iota_E) \in \mathbf{Equi}_{\mathbf{Set}}(f, g)$. Thus
- there exists a unique $\alpha \in \mathbf{Hom}_{\mathbf{Set}}(M, E)$ such that $\iota_E \circ_{\mathbf{Set}} \alpha = m$,
 - there exists a unique $\alpha' \in \mathbf{Hom}_{\mathbf{Set}}(M', E)$ such that $\iota_E \circ_{\mathbf{Set}} \alpha' = m'$,

i.e. the following diagram commutes:



Since $\iota_E \circ m|_E = m$, uniqueness of α implies that $\alpha = m|_E$. Since m is injective, $m|_E \in \mathbf{Iso}_{\mathbf{Set}}(M, E)$.

- Define $\phi \in \mathbf{Hom}_{\mathbf{Set}}(M', M)$ by $\phi' := \alpha^{-1} \circ_{\mathbf{Set}} \alpha'$. Then

$$\begin{aligned} m \circ_{\mathbf{Set}} \phi &= (\iota_E \circ_{\mathbf{Set}} m|_E) \circ_{\mathbf{Set}} (\alpha^{-1} \circ_{\mathbf{Set}} \alpha') \\ &= \iota_E \circ_{\mathbf{Set}} (m|_E \circ_{\mathbf{Set}} \alpha^{-1}) \circ_{\mathbf{Set}} \alpha' \\ &= \iota_E \circ_{\mathbf{Set}} (\alpha \circ_{\mathbf{Set}} \alpha^{-1}) \circ_{\mathbf{Set}} \alpha' \\ &= \iota_E \circ_{\mathbf{Set}} \text{id}_E \circ_{\mathbf{Set}} \alpha' \\ &= \iota_E \circ_{\mathbf{Set}} \alpha' \\ &= m'. \end{aligned}$$

- * Let $\phi_0 \in \mathbf{Hom}_{\mathbf{Set}}(M', M)$. Suppose that $m \circ \phi_0 = m'$. Then

$$\begin{aligned} \iota_E \circ_{\mathbf{Set}} (\alpha \circ_{\mathbf{Set}} \phi_0) &= (\iota_E \circ_{\mathbf{Set}} \alpha) \circ_{\mathbf{Set}} \phi_0 \\ &= (\iota_E \circ_{\mathbf{Set}} m|_E) \circ_{\mathbf{Set}} \phi_0 \\ &= m \circ_{\mathbf{Set}} \phi_0 \\ &= m'. \end{aligned}$$

Uniqueness of α' implies that $\alpha \circ_{\mathbf{Set}} \phi_0 = \alpha'$. Since $\phi_0 \in \mathbf{Hom}_{\mathbf{Set}}(M', M)$ such that $m \circ \phi_0 = m'$ is arbitrary, we have that for each $\phi_0 \in \mathbf{Hom}_{\mathbf{Set}}(M', M)$, $m \circ \phi_0 = m'$ implies that $\alpha \circ_{\mathbf{Set}} \phi_0 = \alpha'$.

- * In particular, [ref prev part \(above\)](#) implies that $\alpha \circ_{\mathbf{Set}} \phi = \alpha'$.

* Let $\phi' \in \text{Hom}_{\mathbf{Set}}(M', M)$. Suppose that $m \circ \phi' = m'$. **ref above prev parts** implies that

$$\begin{aligned}\alpha \circ_{\mathbf{Set}} \phi' &= \alpha' \\ &= \alpha \circ_{\mathbf{Set}} \phi.\end{aligned}$$

Since

$$\begin{aligned}\alpha &= m|_E \\ &\in \text{Iso}_{\mathbf{Set}}(M, E) \\ &\subset \text{Mono}_{\mathbf{Set}}(M, E),\end{aligned}$$

we have that $\phi' = \phi$. Since $\phi' \in \text{Hom}_{\mathbf{Set}}(M', M)$ with $m \circ \phi' = m'$ is arbitrary, we have that for each $\phi' \in \text{Hom}_{\mathbf{Set}}(M', M)$, $m \circ \phi' = m'$ implies that $\phi' = \phi$.

Thus there exists a unique $\phi \in \text{Hom}_{\mathbf{Set}}(M', M)$ such that $m \circ_{\mathbf{Set}} \phi = m'$. Since $M' \in \text{Obj}(\mathbf{Set})$ and $m' \in \text{Hom}_{\mathbf{Set}}(M', A)$ with $f \circ_{\mathbf{Set}} m' = g \circ_{\mathbf{Set}} m'$ are arbitrary, we have that for each $M' \in \text{Obj}(\mathbf{Set})$ and $m' \in \text{Hom}_{\mathbf{Set}}(M', A)$, $f \circ_{\mathbf{Set}} m' = g \circ_{\mathbf{Set}} m'$ implies that there exists a unique $\phi \in \text{Hom}_{\mathbf{Set}}(M', M)$ such that $m \circ_{\mathbf{Set}} \phi = m'$, i.e. the following diagram commutes:

$$\begin{array}{ccccc} M' & & & & B \\ & \searrow m' & & \nearrow f & \\ & & A & & \\ & \nearrow m & & \searrow g & \\ M & & & & B \end{array}$$

ϕ (vertical arrow from M' to M)

Therefore $(M, m) \in \text{Equi}_{\mathbf{Set}}(f, g)$. Hence $m \in \text{RegMono}_{\mathbf{Set}}$. Since $m \in \text{Mono}_{\mathbf{Set}}$ is arbitrary, we have that for each $m \in \text{Mono}_{\mathbf{Set}}$, $m \in \text{RegMono}_{\mathbf{Set}}$. Thus $\text{Mono}_{\mathbf{Set}} \subset \text{RegMono}_{\mathbf{Set}}$.

Since $\text{RegMono}_{\mathbf{Set}} \subset \text{Mono}_{\mathbf{Set}}$ and $\text{Mono}_{\mathbf{Set}} \subset \text{RegMono}_{\mathbf{Set}}$, we have that $\text{RegMono}_{\mathbf{Set}} = \text{Mono}_{\mathbf{Set}}$. □

7.7 Subobjects and Quotient Objects

Definition 7.7.0.1. maybe move this to subsection on subcategories, define wide category generated by hom closed morphisms Let $\mathcal{C}$ be a category and $H \subset \text{Hom}_{\mathcal{C}}$. Then H is said to be **hom closed** if

- 1.
- 2.

Definition 7.7.0.2. define wide category with all morphisms mono, resp epi Let $\mathcal{C}$ be a category. We define

- the restriction of $\mathcal{C}$ to $\text{Mono}_{\mathcal{C}}$, denoted $\mathbf{Mono}(\mathcal{C}, M)$, by
 - $\text{Obj}(\mathbf{Mono}(\mathcal{C}, M)) := \text{Obj}(\mathcal{C})$
 - For $A, B \in \text{Obj}(\mathcal{C})$, $\text{Hom}_{\mathbf{Mono}(\mathcal{C}, M)}(A, B) := \text{Mono}_{\mathcal{C}}(A, B)$
 - For $A, B, C \in \text{Obj}(\mathcal{C})$, $f \in \text{Hom}_{\mathbf{Mono}(\mathcal{C}, M)}(A, B)$ and $g \in \text{Hom}_{\mathbf{Mono}(\mathcal{C}, M)}(B, C)$, $g \circ_{\mathbf{Mono}(\mathcal{C}, M)} f := g \circ_{\mathcal{C}} f$.
- the restriction of $\mathcal{C}$ to $\text{Epi}_{\mathcal{C}}$, denoted $\mathbf{Epi}(\mathcal{C})$, by
 - $\text{Obj}(\mathbf{Epi}(\mathcal{C})) := \text{Obj}(\mathcal{C})$
 - For $A, B \in \text{Obj}(\mathcal{C})$, $\text{Hom}_{\mathbf{Epi}(\mathcal{C})}(A, B) := \text{Epi}_{\mathcal{C}}(A, B)$
 - For $A, B, C \in \text{Obj}(\mathcal{C})$, $f \in \text{Hom}_{\mathbf{Epi}(\mathcal{C})}(A, B)$ and $g \in \text{Hom}_{\mathbf{Epi}(\mathcal{C})}(B, C)$, $g \circ_{\mathbf{Epi}(\mathcal{C})} f := g \circ_{\mathcal{C}} f$.

Exercise 7.7.0.3. Let $\mathcal{C}$ be a category. Then

1. (a) $\mathbf{Mono}(\mathcal{C}, M)$ is a category ,
 (b) $\mathbf{Mono}(\mathcal{C}, M)$ is a wide subcategory of $\mathcal{C}$,
2. (a) $\mathbf{Epi}(\mathcal{C})$ is a category,
 (b) $\mathbf{Epi}(\mathcal{C})$ is a wide subcategory of $\mathcal{C}$.

Proof. clear. fill out, ref exercices showing that composition of mono is mono, id_X is mono □

Exercise 7.7.0.4. Let $\mathcal{C}$ be a category and $X \in \text{Obj}(\mathcal{C})$. Then

1. $\mathbf{Mono}(\mathcal{C}, M)/X$ is a proset category.
- 2.

Proof. 1. Let $(A, f), (B, g) \in \mathbf{Mono}(\mathcal{C})/X$ and $\phi \in \text{Hom}_{\mathbf{Mono}(\mathcal{C})/X}((A, f), (B, g))$. Then

- 2.

□

Definition 7.7.0.5. Let $\mathcal{C}$ be a category, $M \subset \text{Mono}_{\mathcal{C}}$, $A, B \in \text{Obj}(\mathcal{C})$ and $f \in \text{Mono}_{\mathcal{C}}(A, B)$. Then (A, f) is said to be an M -subobject of B if $f \in M$.

Definition 7.7.0.6. Let $\mathcal{C}$ be a category, $M \subset \text{Mono}_{\mathcal{C}}$ and $X \in \text{Obj}(\mathcal{C})$. We define the **category of M -subobjects of X** , denoted $\text{Sub}_{\mathcal{C}}^M(X)$, by

- $\text{Obj}(\text{Sub}_{\mathcal{C}}^M(X)) := \{(A, \alpha) \in \text{Obj}(\mathcal{C}/X) : \alpha \text{ is an } M\text{-subobject of } X\}$
- for $(A, f), (B, g) \in \text{Obj}(\text{Sub}_{\mathcal{C}}^M(X))$, $\text{Hom}_{(\text{Sub}_{\mathcal{C}}^M(X))}((A, f), (B, g)) := \text{Hom}_{\mathcal{C}/X}((A, f), (B, g))$
-

is there a notion of full subcategory generated by a set of objects? can we say that $\text{Sub}_{\mathcal{C}}^M(X)$ is the full subcategory of $\mathcal{C}/X$ generated by the M -subobjects of X ?

Exercise 7.7.0.7. show $\text{Sub}_{\mathcal{C}}^M(X)$ is a preordered set/category, i.e. if (A, f) and (B, g) are M -subobjects of X , then $\# \text{Hom}_{\text{Sub}_{\mathcal{C}}^M(X)}((A, f), (B, g)) = 1$.

Constructions on Categories

are subcategories subobjects in **Cat** where subobjects are defined as in ‘Elementary categories, elementary toposes’ by mcLarty?, how to place this subsection?

The subcategories generated by a graph as an adjunction should also be applicable to describing the topology generated by a subset I think, not sure if there is a lattice formulation of this topic as well and if the adjunct description generalizes this lattice description. Maybe there is just a straight lattice based characterization of generated subcategories when we order by inclusion functors.

So: cats (and their graphs) functors natty transf subcats (with gal conn between posets of subcats and graphs and then subcats generated by subgraphs) iso/mono/epi init/terminal prod(including) coprod equal/coequal pull/push-subobj? END!!!!!!!!!!!!!!!!!!!!!!END!!!!!!!!!!!!!!!!!!!!!!

- We define $\text{Subcat}(\mathcal{C}) := \{\mathcal{D} \in \mathbf{Cat} : \mathcal{L} \text{ is a subcategory of } \mathcal{C}\}$. need to give more justification for why we restrict to $\mathbf{Cat}$, with universes, this should not be an issue, but need to formalize it

- We define the **subcategory relation on** $\text{Subcat}(\mathcal{C})$, denoted $\leq_{\text{Cat}} \subset \text{Subcat}(\mathcal{C}) \times \text{Subcat}(\mathcal{C})$, by $\mathcal{D} \leq_{\text{Cat}} \mathcal{C}$ iff $\mathcal{D}$ is a subcat of $\mathcal{C}$.

Definition 8.1.0.2. Let $\mathcal{C}$ be a category and $\mathcal{D} \in \text{Subcat}(\mathcal{C})$. We define the **inclusion functor** from $\mathcal{D}$ to $\mathcal{C}$, denoted $\iota_{\mathcal{D}}^{\mathcal{C}} : \mathcal{D} \rightarrow \mathcal{C}$, by

- $\iota_{\mathcal{D}}^{\mathcal{C}}(X) := X, \quad X \in \text{Obj}(\mathcal{D})$
- $\iota_{\mathcal{D}}^{\mathcal{C}}(f) := f \in \text{Hom}_{\mathcal{D}}$

Note 8.1.0.3. When the context is clear, we may write $\iota_{\mathcal{D}}$ in place of $\iota_{\mathcal{D}}^{\mathcal{C}}$.

Exercise 8.1.0.4. Let $\mathcal{C}$ be a category and $\mathcal{D} \in \text{Subcat}(\mathcal{C})$. Then $\iota_{\mathcal{D}} : \mathcal{D} \rightarrow \mathcal{C}$ is a functor.

Proof.

- Let $X, Y \in \text{Obj}(\mathcal{D})$ and $f \in \text{Hom}_{\mathcal{D}}(X, Y)$. By Definition 8.1.0.1,

$$\begin{aligned} \iota_{\mathcal{D}}(f) &= f \\ &\in \text{Hom}_{\mathcal{D}}(X, Y) \\ &\subset \text{Hom}_{\mathcal{C}}(X, Y) \\ &= \text{Hom}_{\mathcal{C}}(\iota_{\mathcal{D}}(X), \iota_{\mathcal{D}}(Y)). \end{aligned}$$

- Let $X, Y, Z \in \text{Obj}(\mathcal{D})$, $g \in \text{Hom}_{\mathcal{D}}(X, Y)$ and $h \in \text{Hom}_{\mathcal{D}}(Y, Z)$. By Definition 8.1.0.1,

$$\begin{aligned} \iota_{\mathcal{D}}(g \circ_{\mathcal{D}} f) &= g \circ_{\mathcal{D}} f \\ &= g \circ_{\mathcal{C}} f \\ &= \iota_{\mathcal{D}}(g) \circ_{\mathcal{C}} \iota_{\mathcal{D}}(f). \end{aligned}$$

- Let $X \in \text{Obj}(\mathcal{D})$. By Definition 8.1.0.1,

$$\begin{aligned} \iota_{\mathcal{D}}(\text{id}_X^{\mathcal{D}}) &= \text{id}_X^{\mathcal{D}} \\ &= \text{id}_X^{\mathcal{C}} \\ &= \text{id}_{\iota_{\mathcal{D}}(X)}^{\mathcal{C}}. \end{aligned}$$

Hence $\iota_{\mathcal{D}} : \mathcal{D} \rightarrow \mathcal{C}$ is a functor. □

Definition 8.1.0.5. We define the **forgetful functor from Cat to Quiv**, denoted $\text{Forget}_{\text{Quiv}}^{\text{Cat}} : \text{Cat} \rightarrow \text{Quiv}$ **finish**

Definition 8.1.0.6. Let $\mathcal{C}$ be a category and $H \subset \text{Hom}_{\mathcal{C}}$. We define the **subcategory of $\mathcal{C}$ generated by H** , denoted $\text{SubcatGen}(H, \mathcal{C})$, by $\text{SubcatGen}(H, \mathcal{C}) :=$

Exercise 8.1.0.7. content...

Definition 8.1.0.8.

Definition 8.1.0.9. Let $\mathcal{C}$ and $\mathcal{D}$ be categories. Suppose that $\mathcal{D}$ is a subcategory of $\mathcal{C}$. Then $\mathcal{C}$ is said to be

- **full** if for each $A, B \in \text{Obj}(\mathcal{D})$, $\text{Hom}_{\mathcal{D}}(A, B) = \text{Hom}_{\mathcal{C}}(A, B)$,
- **wide** or **lluf** if for each $\text{Obj}(\mathcal{D}) = \text{Obj}(\mathcal{C})$.

Note 8.1.0.10. The motivation for the adjective “lluf” seems to be that it is just “full” backwards.

Exercise 8.1.0.11. content...

Exercise 8.1.0.12. **erase after making iff** Let $\mathcal{C}$ be a category, $\mathcal{D}_0 \subset \text{Obj}(\mathcal{C})$ and $\mathcal{D}_1 \subset \text{Hom}_{\mathcal{C}}$. Define $\mathcal{D}_1^{(2)} \subset \mathcal{D}_1 \times \mathcal{D}_1$ by $\mathcal{D}_1^{(2)} := \{(g, f) \in \mathcal{D}_1 \times \mathcal{D}_1 : \text{dom}_{\mathcal{C}}(g) = \text{cod}_{\mathcal{C}}(f)\}$. If

- $\text{dom}_{\mathcal{C}}(\mathcal{D}_1), \text{cod}_{\mathcal{C}}(\mathcal{D}_1) \subset \mathcal{D}_0$,
- $\circ_{\mathcal{C}}(\mathcal{D}_1^{(2)}) \subset \mathcal{D}_1$,
- for each $X \in \mathcal{D}_0$, $\text{id}_X^{\mathcal{C}} \in \mathcal{D}_1$,

then

1. $(\mathcal{D}_0, \mathcal{D}_1, \text{dom}_{\mathcal{C}}|_{\mathcal{D}_1}, \text{cod}_{\mathcal{C}}|_{\mathcal{D}_1}, \circ_{\mathcal{C}}|_{\mathcal{D}_1^{(2)}})$ is a category,
2. $(\mathcal{D}_0, \mathcal{D}_1, \text{dom}_{\mathcal{C}}|_{\mathcal{D}_1}, \text{cod}_{\mathcal{C}}|_{\mathcal{D}_1}, \circ_{\mathcal{C}}|_{\mathcal{D}_1^{(2)}})$ is a subcategory of $\mathcal{C}$.

Proof. Set $\text{dom}_{\mathcal{D}} := \text{dom}_{\mathcal{C}}|_{\mathcal{D}_1}$, $\text{cod}_{\mathcal{D}} := \text{cod}_{\mathcal{C}}|_{\mathcal{D}_1}$, $\circ_{\mathcal{D}} := \circ_{\mathcal{C}}|_{\mathcal{D}_1^{(2)}}$ and $\mathcal{D} := (\mathcal{D}_0, \mathcal{D}_1, \text{dom}_{\mathcal{C}}|_{\mathcal{D}_1}, \text{cod}_{\mathcal{C}}|_{\mathcal{D}_1}, \circ_{\mathcal{C}}|_{\mathcal{D}_1^{(2)}})$.

1. (a) **(Composition:)**

For each $(g, f) \in \mathcal{D}_1^{(2)}$,

$$\begin{aligned} \text{dom}_{\mathcal{D}}(g \circ_{\mathcal{D}} f) &= \text{dom}_{\mathcal{C}}(g \circ_{\mathcal{C}} f) \\ &= \text{dom}_{\mathcal{C}}(f) \\ &= \text{dom}_{\mathcal{D}}(f). \end{aligned}$$

and

$$\begin{aligned} \text{cod}_{\mathcal{D}}(g \circ_{\mathcal{D}} f) &= \text{cod}_{\mathcal{C}}(g \circ_{\mathcal{C}} f) \\ &= \text{cod}_{\mathcal{C}}(g) \\ &= \text{cod}_{\mathcal{D}}(g). \end{aligned}$$

- (b) **(Associativity:)**

For each $f, g, h \in \mathcal{D}_1$, if $(g, f), (h, g) \in \mathcal{D}_1^{(2)}$, then

$$\begin{aligned} (h \circ_{\mathcal{D}} g) \circ_{\mathcal{D}} f &= (h \circ_{\mathcal{C}} g) \circ_{\mathcal{C}} f \\ &= h \circ_{\mathcal{C}} (g \circ_{\mathcal{C}} f) \\ &= h \circ_{\mathcal{D}} (g \circ_{\mathcal{D}} f). \end{aligned}$$

- (c) **(Identity:)**

Let $X \in \mathcal{D}_0$. Since $\text{id}_X^{\mathcal{C}} \in \mathcal{D}_1$,

$$\begin{aligned} \text{dom}_{\mathcal{D}}(\text{id}_X^{\mathcal{C}}) &= \text{dom}_{\mathcal{C}}(\text{id}_X^{\mathcal{C}}) \\ &= X, \end{aligned}$$

$$\begin{aligned} \text{cod}_{\mathcal{D}}(\text{id}_X^{\mathcal{C}}) &= \text{cod}_{\mathcal{C}}(\text{id}_X^{\mathcal{C}}) \\ &= X, \end{aligned}$$

and for each $f, g \in \mathcal{D}_1$, if $\text{dom}_{\mathcal{D}}(f) = X$ and $\text{cod}_{\mathcal{D}}(g) = X$, then

$$\begin{aligned} f \circ_{\mathcal{D}} \text{id}_X^{\mathcal{C}} &= f \circ_{\mathcal{C}} \text{id}_X^{\mathcal{C}} \\ &= f \end{aligned}$$

and

$$\begin{aligned} \text{id}_X^{\mathcal{C}} \circ_{\mathcal{D}} g &= \text{id}_X^{\mathcal{C}} \circ_{\mathcal{C}} g \\ &= g. \end{aligned}$$

Hence $\mathcal{D}$ is a category.

2. Clear by construction.

□

8.2 Product and Coproduct Categories

8.2.1 Introduction

Note 8.2.1.1. there may be an issue with the definition of infinite products/coproducts of categories. The idea is that if for some $\alpha_0 \in A$, $\text{Obj}(\mathcal{C}_{\alpha_0})$ is not a set, then $\bigcup \text{Obj}(\mathcal{C}_\alpha)$ is not a set. So it might be fine to define a single (or finitely many) object(s) of the product category: $(X_\alpha)_{\alpha \in A} : A \rightarrow \bigcup_{\alpha \in A} \text{Obj}(\mathcal{C}_\alpha)$ such that for each $\alpha_0 \in A$, $(X_\alpha)_{\alpha \in A}(\alpha_0) \in \text{Obj}(\mathcal{C}_{\alpha_0})$, but this single object may be in fact not a set. In which case, we can not make a 'class' of these objects since classes contained as members in other classes are sets. So we either iterate the von Neumann-Bernays-Godel set theory to higher level classes, or we use universes or something, but this needs to be made precise and needs to be applied to all constructions going forward. FINISH!!!

Definition 8.2.1.2. Let $(\mathcal{C})_{\alpha \in A}$ be a collection of categories. We define the **product category of** $(\mathcal{C})_{\alpha \in A}$, denoted $\prod_{\alpha \in A} \mathcal{C}_\alpha$, by

- $\text{Obj}\left(\prod_{\alpha \in A} \mathcal{C}_\alpha\right) := \prod_{\alpha \in A} \text{Obj}(\mathcal{C}_\alpha)$

- for each $(X_\alpha)_{\alpha \in A}, (Y_\alpha)_{\alpha \in A} \in \text{Obj}\left(\prod_{\alpha \in A} \mathcal{C}_\alpha\right)$,

$$\text{Hom}_{\prod_{\alpha \in A} \mathcal{C}_\alpha}((X_\alpha)_{\alpha \in A}, (Y_\alpha)_{\alpha \in A}) := \prod_{\alpha \in A} \text{Hom}_{\mathcal{C}_\alpha}(X_\alpha, Y_\alpha)$$

- $(X_\alpha)_{\alpha \in A}, (Y_\alpha)_{\alpha \in A}, (Z_\alpha)_{\alpha \in A} \in \text{Obj}\left(\prod_{\alpha \in A} \mathcal{C}_\alpha\right)$, $(f_\alpha)_{\alpha \in A} \in \text{Hom}_{\prod_{\alpha \in A} \mathcal{C}_\alpha}((X_\alpha)_{\alpha \in A}, (Y_\alpha)_{\alpha \in A})$ and $(g_\alpha)_{\alpha \in A} \in \text{Hom}_{\prod_{\alpha \in A} \mathcal{C}_\alpha}((Y_\alpha)_{\alpha \in A}, (Z_\alpha)_{\alpha \in A})$

$$(g_\alpha)_{\alpha \in A} \circ \prod_{\alpha \in A} \mathcal{C}_\alpha (f_\alpha)_{\alpha \in A} := (g_\alpha \circ_{\mathcal{C}_\alpha} f_\alpha)_{\alpha \in A}$$

Note 8.2.1.3. When $A = \{1, 2\}$, we may write $\mathcal{C}_1 \times \mathcal{C}_2$ in place of $\prod_{\alpha \in A} \mathcal{C}_\alpha$.

Exercise 8.2.1.4. Let $(\mathcal{C})_{\alpha \in A}$ be a collection of categories. Then $\prod_{\alpha \in A} \mathcal{C}_\alpha$ is a category.

Proof.

- **well-definedness of composition:**

Let $(X_\alpha^1)_{\alpha \in A}, (X_\alpha^2)_{\alpha \in A}, (X_\alpha^3)_{\alpha \in A} \in \text{Obj}\left(\prod_{\alpha \in A} \mathcal{C}_\alpha\right)$, $(f_\alpha^{12})_{\alpha \in A} \in \text{Hom}_{\prod_{\alpha \in A} \mathcal{C}_\alpha}((X_\alpha^1)_{\alpha \in A}, (X_\alpha^2)_{\alpha \in A})$ and $(f_\alpha^{23})_{\alpha \in A} \in \text{Hom}_{\prod_{\alpha \in A} \mathcal{C}_\alpha}((X_\alpha^2)_{\alpha \in A}, (X_\alpha^3)_{\alpha \in A})$

Then

$$\begin{aligned} (f_\alpha^{23})_{\alpha \in A} \circ \prod_{\alpha \in A} \mathcal{C}_\alpha (f_\alpha^{12})_{\alpha \in A} &= (f_\alpha^{23} \circ_{\mathcal{C}_\alpha} f_\alpha^{12})_{\alpha \in A} \\ &\in \text{Hom}_{\prod_{\alpha \in A} \mathcal{C}_\alpha}((X_\alpha^1)_{\alpha \in A}, (X_\alpha^3)_{\alpha \in A}). \end{aligned}$$

Thus, composition is well defined.

- **associativity of composition:**

Let $(X_\alpha^1)_{\alpha \in A}, (X_\alpha^2)_{\alpha \in A}, (X_\alpha^3)_{\alpha \in A}, (X_\alpha^4)_{\alpha \in A} \in \text{Obj}\left(\prod_{\alpha \in A} \mathcal{C}_\alpha\right)$, $(f_\alpha^{12})_{\alpha \in A} \in \text{Hom}_{\prod_{\alpha \in A} \mathcal{C}_\alpha}((X_\alpha^1)_{\alpha \in A}, (X_\alpha^2)_{\alpha \in A})$, $(f_\alpha^{23})_{\alpha \in A} \in \text{Hom}_{\prod_{\alpha \in A} \mathcal{C}_\alpha}((X_\alpha^2)_{\alpha \in A}, (X_\alpha^3)_{\alpha \in A})$

and $(f_\alpha^{34})_{\alpha \in A} \in \text{Hom}_{\prod_{\alpha \in A} \mathcal{C}_\alpha}((X_\alpha^3)_{\alpha \in A}, (X_\alpha^4)_{\alpha \in A})$. Then

$$\begin{aligned} [(f_\alpha^{34})_{\alpha \in A} \circ \prod_{\alpha \in A} \mathcal{C}_\alpha (f_\alpha^{23})_{\alpha \in A}] \circ \prod_{\alpha \in A} \mathcal{C}_\alpha (f_\alpha^{12})_{\alpha \in A} &= (f_\alpha^{34} \circ_{\mathcal{C}_\alpha} f_\alpha^{12})_{\alpha \in A} \circ \prod_{\alpha \in A} \mathcal{C}_\alpha (f_\alpha^{12})_{\alpha \in A} \\ &= ([f_\alpha^{34} \circ_{\mathcal{C}_\alpha} f_\alpha^{23}] \circ_{\mathcal{C}_\alpha} f_\alpha^{12})_{\alpha \in A} \\ &= (f_\alpha^{34} \circ_{\mathcal{C}_\alpha} [f_\alpha^{23} \circ_{\mathcal{C}_\alpha} f_\alpha^{12}])_{\alpha \in A} \\ &= (f_\alpha^{34})_{\alpha \in A} \circ \prod_{\alpha \in A} \mathcal{C}_\alpha (f_\alpha^{23} \circ_{\mathcal{C}_\alpha} f_\alpha^{12})_{\alpha \in A} \\ &= (f_\alpha^{34})_{\alpha \in A} \circ \prod_{\alpha \in A} \mathcal{C}_\alpha [(f_\alpha^{23})_{\alpha \in A} \circ \prod_{\alpha \in A} \mathcal{C}_\alpha (f_\alpha^{12})_{\alpha \in A}]. \end{aligned}$$

Thus composition is associative.

• **existence of identities:**

Let $(X_\alpha)_{\alpha \in A} \in \text{Obj}\left(\prod_{\alpha \in A} \mathcal{C}_\alpha\right)$, $(f_\alpha)_{\alpha \in A}, (g_\alpha)_{\alpha \in A} \in \text{Hom}_{\prod_{\alpha \in A} \mathcal{C}_\alpha}$. Suppose that $\text{dom}_{\prod_{\alpha \in A} \mathcal{C}_\alpha}((f_\alpha)_{\alpha \in A}) = (X_\alpha)_{\alpha \in A}$ and $\text{cod}_{\prod_{\alpha \in A} \mathcal{C}_\alpha}((g_\alpha)_{\alpha \in A}) = (X_\alpha)_{\alpha \in A}$. Then

$$\begin{aligned} (f_\alpha)_{\alpha \in A} \circ \prod_{\alpha \in A} \mathcal{C}_\alpha (\text{id}_{X_\alpha}^{\mathcal{C}_\alpha})_{\alpha \in A} &= (f_\alpha \circ_{\mathcal{C}_\alpha} \text{id}_{X_\alpha}^{\mathcal{C}_\alpha})_{\alpha \in A} \\ &= (f_\alpha)_{\alpha \in A} \end{aligned}$$

and

$$\begin{aligned} (\text{id}_{X_\alpha}^{\mathcal{C}_\alpha})_{\alpha \in A} \circ \prod_{\alpha \in A} \mathcal{C}_\alpha (g_\alpha)_{\alpha \in A} &= (\text{id}_{X_\alpha}^{\mathcal{C}_\alpha} \circ_{\mathcal{C}_\alpha} g_\alpha)_{\alpha \in A} \\ &= (g_\alpha)_{\alpha \in A}. \end{aligned}$$

Therefore $\text{id}_{(X_\alpha)_{\alpha \in A}}^{\prod_{\alpha \in A} \mathcal{C}_\alpha} = (\text{id}_{X_\alpha}^{\mathcal{C}_\alpha})_{\alpha \in A}$.

□

Exercise 8.2.1.5. Let $(\mathcal{C}_\alpha)_{\alpha \in A}$ be a collection of categories. Then for each $\alpha \in A$, $\mathcal{C}_\alpha$ is small iff $\prod_{\alpha \in A} \mathcal{C}_\alpha$ is small.

Proof. We have that

$$\begin{aligned} \text{for each } \alpha \in A, \mathcal{C}_\alpha \text{ is small} &\iff \text{for each } \alpha \in A, \text{Obj}(\mathcal{C}_\alpha) \in \text{Obj}(\text{Set}) \text{ and } \text{Hom}_{\mathcal{C}_\alpha} \in \text{Obj}(\text{Set}) \\ &\iff \prod_{\alpha \in A} \text{Obj}(\mathcal{C}_\alpha) \in \text{Obj}(\mathbf{Set}) \text{ and } \prod_{\alpha \in A} \text{Hom}_{\mathcal{C}_\alpha} \in \text{Obj}(\mathbf{Set}) \\ &\iff \text{Obj}\left(\prod_{\alpha \in A} \mathcal{C}_\alpha\right) \in \text{Obj}(\mathbf{Set}) \text{ and } \text{Hom}_{\prod_{\alpha \in A} \mathcal{C}_\alpha} \in \text{Obj}(\mathbf{Set}) \\ &\iff \prod_{\alpha \in A} \mathcal{C}_\alpha \text{ is small.} \end{aligned}$$

□

Definition 8.2.1.6. Let $(\mathcal{C})_{\alpha \in A}$ be a collection of categories. We define the **coproduct category of $(\mathcal{C})_{\alpha \in A}$** , denoted $\coprod_{\alpha \in A} \mathcal{C}_\alpha$, by

- $\text{Obj}\left(\coprod_{\alpha \in A} \mathcal{C}_\alpha\right) := \coprod_{\alpha \in A} \text{Obj}(\mathcal{C}_\alpha)$
- for each $(\alpha_X, X), (\alpha_Y, Y) \in \text{Obj}\left(\coprod_{\alpha \in A} \mathcal{C}_\alpha\right)$,

$$\text{Hom}_{\prod_{\alpha \in A} \mathcal{C}_\alpha}((\alpha_X, X), (\alpha_Y, Y)) := \begin{cases} \{\alpha\} \times \text{Hom}_{\mathcal{C}_\alpha}(X, Y), & \alpha_X = \alpha_Y \\ \emptyset, & \alpha_X \neq \alpha_Y \end{cases}$$

- for each $\alpha \in A$, $X, Y, Z \in \text{Obj}(\mathcal{C}_\alpha)$, $f \in \text{Hom}_{\mathcal{C}_\alpha}(X, Y)$ and $g \in \text{Hom}_{\mathcal{C}_\alpha}(Y, Z)$,

$$(\alpha, g) \circ \coprod_{\alpha' \in A} c_{\alpha'} (\alpha, f) := (\alpha, g \circ c_\alpha f)$$

Note 8.2.1.7.

- For each $(\alpha_X, X), (\alpha_Y, Y) \in \text{Obj}\left(\coprod_{\alpha \in A} \mathcal{C}_\alpha\right)$ and $(\alpha_f, f) \in \text{Hom}_{\coprod_{\alpha \in A} \mathcal{C}_\alpha}$, we have that

$$\begin{aligned} (\alpha_f, f) \in \text{Hom}_{\coprod_{\alpha \in A} \mathcal{C}_\alpha}((\alpha_X, X), (\alpha_Y, Y)) &\implies \text{Hom}_{\coprod_{\alpha \in A} \mathcal{C}_\alpha}((\alpha_X, X), (\alpha_Y, Y)) \neq \emptyset \\ &\implies \alpha_X = \alpha_Y = \alpha_f. \end{aligned}$$

- By definition, $\text{Hom}_{\coprod_{\alpha \in A} \mathcal{C}_\alpha} = \coprod_{\alpha \in A} \text{Hom}_{\mathcal{C}_\alpha}$.

Exercise 8.2.1.8. Let $(\mathcal{C})_{\alpha \in A}$ be a collection of categories. Then $\coprod_{\alpha \in A} \mathcal{C}_\alpha$ is a category.

Proof.

- **well-definedness of composition:**

Let $(\alpha_1, X_1), (\alpha_2, X_2), (\alpha_3, X_3) \in \text{Obj}\left(\coprod_{\alpha \in A} \mathcal{C}_\alpha\right)$, $(\alpha_{12}, f_{12}) \in \text{Hom}_{\coprod_{\alpha \in A} \mathcal{C}_\alpha}((\alpha_1, X_1), (\alpha_2, X_2))$ and $(\alpha_{23}, f_{23}) \in \text{Hom}_{\coprod_{\alpha \in A} \mathcal{C}_\alpha}((\alpha_2, X_2), (\alpha_3, X_3))$. Then $\alpha_1 = \alpha_2 = \alpha_3 = \alpha_{12} = \alpha_{23}$. Set $\alpha_0 := \alpha_1$. Since $f_{12} \in \text{Hom}_{\mathcal{C}_{\alpha_0}}(X_1, X_2)$ and $f_{23} \in \text{Hom}_{\mathcal{C}_{\alpha_0}}(X_2, X_3)$, we have that $f_{23} \circ c_{\alpha_0} f_{12} \in \text{Hom}_{\mathcal{C}_{\alpha_0}}(X_1, X_3)$ and

$$\begin{aligned} (\alpha_{23}, f_{23}) \circ \coprod_{\alpha \in A} c_\alpha (\alpha_{12}, f_{12}) &= (\alpha_0, f_{23}) \circ \coprod_{\alpha \in A} c_\alpha (\alpha_0, f_{12}) \\ &= (\alpha_0, f_{23} \circ c_{\alpha_0} f_{12}) \\ &\in \text{Hom}_{\coprod_{\alpha \in A} \mathcal{C}_\alpha}((\alpha_0, X_1), (\alpha_0, X_3)) \\ &= \text{Hom}_{\coprod_{\alpha \in A} \mathcal{C}_\alpha}((\alpha_1, X_1), (\alpha_3, X_3)). \end{aligned}$$

Thus, composition is well defined.

- **associativity of composition:**

Let $(\alpha_1, X_1), (\alpha_2, X_2), (\alpha_3, X_3), (\alpha_4, X_4) \in \text{Obj}\left(\coprod_{\alpha \in A} \mathcal{C}_\alpha\right)$, $(\alpha_{12}, f_{12}) \in \text{Hom}_{\coprod_{\alpha \in A} \mathcal{C}_\alpha}((\alpha_1, X_1), (\alpha_2, X_2))$, $(\alpha_{23}, f_{23}) \in \text{Hom}_{\coprod_{\alpha \in A} \mathcal{C}_\alpha}((\alpha_2, X_2), (\alpha_3, X_3))$ and $(\alpha_{34}, f_{34}) \in \text{Hom}_{\coprod_{\alpha \in A} \mathcal{C}_\alpha}((\alpha_3, X_3), (\alpha_4, X_4))$. Then $\alpha_1 = \alpha_2 = \alpha_3 = \alpha_4 = \alpha_{12} = \alpha_{23} = \alpha_{34}$. Set $\alpha_0 := \alpha_1$. Since $f_{12} \in \text{Hom}_{\mathcal{C}_{\alpha_0}}(X_1, X_2)$, $f_{23} \in \text{Hom}_{\mathcal{C}_{\alpha_0}}(X_2, X_3)$ and $f_{34} \in \text{Hom}_{\mathcal{C}_{\alpha_0}}(X_3, X_4)$, we have that

$$\begin{aligned} (\alpha_{34}, f_{34}) \circ \coprod_{\alpha \in A} c_\alpha [(\alpha_{23}, f_{23}) \circ \coprod_{\alpha \in A} c_\alpha (\alpha_{12}, f_{12})] &= (\alpha_0, f_{34}) \circ \coprod_{\alpha \in A} c_\alpha [(\alpha_0, f_{23}) \circ \coprod_{\alpha \in A} c_\alpha (\alpha_0, f_{12})] \\ &= (\alpha_0, f_{34}) \circ \coprod_{\alpha \in A} c_\alpha (\alpha_0, f_{23} \circ c_{\alpha_0} f_{12}) \\ &= (\alpha_0, f_{34} \circ c_{\alpha_0} [f_{23} \circ c_{\alpha_0} f_{12}]) \\ &= (\alpha_0, [f_{34} \circ c_{\alpha_0} f_{23}] \circ c_{\alpha_0} f_{12}) \\ &= (\alpha_0, f_{34} \circ c_{\alpha_0} f_{23}) \circ \coprod_{\alpha \in A} c_\alpha (\alpha_0, f_{12}) \\ &= [(\alpha_0, f_{34}) \circ \coprod_{\alpha \in A} c_\alpha (\alpha_0, f_{23})] \circ \coprod_{\alpha \in A} c_\alpha (\alpha_0, f_{12}) \\ &= [(\alpha_{34}, f_{34}) \circ \coprod_{\alpha \in A} c_\alpha (\alpha_{23}, f_{23})] \circ \coprod_{\alpha \in A} c_\alpha (\alpha_{12}, f_{12}). \end{aligned}$$

Thus composition is associative.

• **existence of identities:**

Let $(\alpha_X, X) \in \text{Obj} \left(\coprod_{\alpha \in A} \mathcal{C}_\alpha \right)$, $(\alpha_f, f), (\alpha_g, g) \in \text{Hom}_{\coprod_{\alpha \in A} \mathcal{C}_\alpha}$. Suppose that $\text{dom}_{\coprod_{\alpha \in A} \mathcal{C}_\alpha}(\alpha_f, f) = (\alpha_X, X)$ and $\text{cod}_{\coprod_{\alpha \in A} \mathcal{C}_\alpha}(\alpha_g, g) = (\alpha_X, X)$. Then $\alpha_X = \alpha_f = \alpha_g$, $f, g \in \text{Hom}_{\mathcal{C}_{\alpha_X}}$, $\text{dom}_{\mathcal{C}_{\alpha_X}}(f) = X$ and $\text{cod}_{\mathcal{C}_{\alpha_X}}(g) = X$. Hence

$$\begin{aligned} (\alpha_f, f) \circ \coprod_{\alpha \in A} \mathcal{C}_\alpha (\alpha_X, \text{id}_X^{\mathcal{C}_{\alpha_X}}) &= (\alpha_X, f) \circ \coprod_{\alpha \in A} \mathcal{C}_\alpha (\alpha_X, \text{id}_X^{\mathcal{C}_{\alpha_X}}) \\ &= (\alpha_X, f \circ_{\mathcal{C}_{\alpha_X}} \text{id}_X^{\mathcal{C}_{\alpha_X}}) \\ &= (\alpha_X, f) \\ &= (\alpha_f, f) \end{aligned}$$

and

$$\begin{aligned} (\alpha_X, \text{id}_X^{\mathcal{C}_{\alpha_X}}) \circ \coprod_{\alpha \in A} \mathcal{C}_\alpha (\alpha_g, g) &= (\alpha_X, \text{id}_X^{\mathcal{C}_{\alpha_X}}) \circ \coprod_{\alpha \in A} \mathcal{C}_\alpha (\alpha_X, g) \\ &= (\alpha_X, \text{id}_X^{\mathcal{C}_{\alpha_X}} \circ_{\mathcal{C}_{\alpha_X}} g) \\ &= (\alpha_X, g) \\ &= (\alpha_g, g). \end{aligned}$$

Therefore $\text{id}_{(\alpha_X, X)}^{\coprod_{\alpha \in A} \mathcal{C}_\alpha} = (\alpha_X, \text{id}_X^{\mathcal{C}_{\alpha_X}})$.

□

Exercise 8.2.1.9. Let $(\mathcal{C}_\alpha)_{\alpha \in A}$ be a collection of categories. Then for each $\alpha \in A$, $\mathcal{C}_\alpha$ is small iff $\coprod_{\alpha \in A} \mathcal{C}_\alpha$ is small.

Proof. We have that

$$\begin{aligned} \text{for each } \alpha \in A, \mathcal{C}_\alpha \text{ is small} &\iff \text{for each } \alpha \in A, \text{Obj}(\mathcal{C}_\alpha) \in \text{Obj}(\text{Set}) \text{ and } \text{Hom}_{\mathcal{C}_\alpha} \in \text{Obj}(\text{Set}) \\ &\iff \coprod_{\alpha \in A} \text{Obj}(\mathcal{C}_\alpha) \in \text{Obj}(\mathbf{Set}) \text{ and } \coprod_{\alpha \in A} \text{Hom}_{\mathcal{C}_\alpha} \in \text{Obj}(\mathbf{Set}) \\ &\iff \text{Obj} \left(\coprod_{\alpha \in A} \mathcal{C}_\alpha \right) \in \text{Obj}(\mathbf{Set}) \text{ and } \text{Hom}_{\coprod_{\alpha \in A} \mathcal{C}_\alpha} \in \text{Obj}(\mathbf{Set}) \\ &\iff \coprod_{\alpha \in A} \mathcal{C}_\alpha \text{ is small.} \end{aligned}$$

□

Definition 8.2.1.10. Let $\mathcal{A}, \mathcal{B}, \mathcal{C}, \mathcal{D}$ be categories and $F : \mathcal{A} \rightarrow \mathcal{C}$, $G : \mathcal{B} \rightarrow \mathcal{D}$. We define the **product functor of F and G** , denoted $F \times G : \mathcal{A} \times \mathcal{B} \rightarrow \mathcal{C} \times \mathcal{D}$, by

- $F \times G(A, B) := (F(A), G(B)), \quad (A, B) \in \text{Obj}(\mathcal{A} \times \mathcal{B}),$
- $F \times G(f, g) := (F(f), G(g)), \quad (f, g) \in \text{Hom}_{\mathcal{A} \times \mathcal{B}}$

Exercise 8.2.1.11. Let $\mathcal{A}, \mathcal{B}, \mathcal{C}, \mathcal{D}$ be categories and $F : \mathcal{A} \rightarrow \mathcal{C}$, $G : \mathcal{B} \rightarrow \mathcal{D}$. Then $F \times G$ is a functor.

Proof.

1. Let $X, Y \in \text{Obj}(\mathcal{A} \times \mathcal{B})$ and $h \in \text{Hom}_{\mathcal{A} \times \mathcal{B}}(X, Y)$. Then there exist $A_1, A_2 \in \text{Obj}(\mathcal{A})$, $B_1, B_2 \in \text{Obj}(\mathcal{B})$, $f \in \text{Hom}_{\mathcal{A}}(A_1, A_2)$ and $g \in \text{Hom}_{\mathcal{B}}(B_1, B_2)$ such that $X = (A_1, B_1)$, $Y = (A_2, B_2)$ and $h = (f, g)$. Since $F(f) \in \text{Hom}_{\mathcal{C}}(F(A_1), F(A_2))$ and $G(g) \in \text{Hom}_{\mathcal{D}}(G(B_1), G(B_2))$, we have that

$$\begin{aligned} F \times G(h) &= F \times G(f, g) \\ &= (F(f), G(g)) \\ &\in \text{Hom}_{\mathcal{A} \times \mathcal{B}}((F(A_1), G(B_1)), (F(A_2), G(B_2))) \\ &= \text{Hom}_{\mathcal{A} \times \mathcal{B}}(F \times G(A_1, B_1), F \times G(A_2, B_2)) \\ &= \text{Hom}_{\mathcal{A} \times \mathcal{B}}(F \times G(X), F \times G(Y)). \end{aligned}$$

2. Let $X_1, X_2, X_3 \in \text{Obj}(\mathcal{A} \times \mathcal{B})$, $h_{1,2} \in \text{Hom}_{\mathcal{A} \times \mathcal{B}}(X_1, X_2)$ and $h_{2,3} \in \text{Hom}_{\mathcal{A} \times \mathcal{B}}(X_2, X_3)$. Then there exist $A_1, A_2, A_3 \in \text{Obj}(\mathcal{A})$, $B_1, B_2, B_3 \in \text{Obj}(\mathcal{B})$, $f_{1,2} \in \text{Hom}_{\mathcal{A}}(A_1, A_2)$, $g_{1,2} \in \text{Hom}_{\mathcal{B}}(B_1, B_2)$, $f_{2,3} \in \text{Hom}_{\mathcal{A}}(A_2, A_3)$ and $g_{2,3} \in \text{Hom}_{\mathcal{B}}(B_2, B_3)$, such that $X_1 = (A_1, B_1)$, $X_2 = (A_2, B_2)$, $X_3 = (A_3, B_3)$, $h_{1,2} = (f_{1,2}, g_{1,2})$ and $h_{2,3} = (f_{2,3}, g_{2,3})$. Then

$$\begin{aligned}
 F \times G(h_{2,3} \circ_{\mathcal{A} \times \mathcal{B}} h_{1,2}) &= F \times G((f_{2,3}, g_{2,3}) \circ_{\mathcal{A} \times \mathcal{B}} (f_{1,2}, g_{1,2})) \\
 &= F \times G(f_{2,3} \circ_{\mathcal{A}} f_{1,2}, g_{2,3} \circ_{\mathcal{B}} g_{1,2}) \\
 &= (F(f_{2,3} \circ_{\mathcal{A}} f_{1,2}), G(g_{2,3} \circ_{\mathcal{B}} g_{1,2})) \\
 &= (F(f_{2,3}) \circ_{\mathcal{C}} F(f_{1,2}), G(g_{2,3}) \circ_{\mathcal{D}} G(g_{1,2})) \\
 &= (F(f_{2,3}), G(g_{2,3})) \circ_{\mathcal{C} \times \mathcal{D}} (F(f_{1,2}), G(g_{1,2})) \\
 &= F \times G(f_{2,3}, g_{2,3}) \circ_{\mathcal{C} \times \mathcal{D}} F \times G(f_{1,2}, g_{1,2}) \\
 &= F \times G(h_{2,3}) \circ_{\mathcal{C} \times \mathcal{D}} F \times G(h_{1,2}).
 \end{aligned}$$

3. Let $X \in \text{Obj}(\mathcal{A} \times \mathcal{B})$. Then there exists $A \in \text{Obj}(\mathcal{A})$ and $B \in \text{Obj}(\mathcal{B})$ such that $X = (A, B)$. Then

$$\begin{aligned}
 F \times G(\text{id}_X^{A \times B}) &= F \times G(\text{id}_{(A,B)}^{A \times B}) \\
 &= F \times G(\text{id}_A^A, \text{id}_B^B) \\
 &= (F(\text{id}_A^A), G(\text{id}_B^B)) \\
 &= (\text{id}_{F(A)}^{\mathcal{C}}, \text{id}_{G(B)}^{\mathcal{D}}) \\
 &= \text{id}_{(F(A), G(B))}^{\mathcal{C} \times \mathcal{D}} \\
 &= \text{id}_{F \times G(A,B)}^{\mathcal{C} \times \mathcal{D}} \\
 &= \text{id}_{F \times G(X)}^{\mathcal{C} \times \mathcal{D}}.
 \end{aligned}$$

Hence $F \times G$ is a functor. □

Exercise 8.2.1.12. Let $\mathcal{C}_1, \mathcal{C}_2, \mathcal{C}_3, \mathcal{D}_1, \mathcal{D}_2, \mathcal{D}_3 \in \text{Obj}(\mathbf{Cat})$, $F_{1,2} \in \text{Hom}_{\mathbf{Cat}}(\mathcal{C}_1, \mathcal{C}_2)$, $F_{2,3} \in \text{Hom}_{\mathbf{Cat}}(\mathcal{C}_2, \mathcal{C}_3)$, $G_{1,2} \in \text{Hom}_{\mathbf{Cat}}(\mathcal{D}_1, \mathcal{D}_2)$ and $G_{2,3} \in \text{Hom}_{\mathbf{Cat}}(\mathcal{D}_2, \mathcal{D}_3)$. Then

$$(F_{2,3} \circ_{\mathbf{Cat}} F_{1,2}) \times (G_{2,3} \circ_{\mathbf{Cat}} G_{1,2}) = (F_{2,3} \times G_{2,3}) \circ_{\mathbf{Cat}} (F_{1,2} \times G_{1,2}).$$

Proof.

- For each $A \in \text{Obj}(\mathcal{C}_1)$ and $B \in \text{Obj}(\mathcal{D}_1)$, we have that

$$\begin{aligned}
 [(F_{2,3} \circ_{\mathbf{Cat}} F_{1,2}) \times (G_{2,3} \circ_{\mathbf{Cat}} G_{1,2})](A, B) &= (F_{2,3} \circ_{\mathbf{Cat}} F_{1,2}(A), (G_{2,3} \circ_{\mathbf{Cat}} G_{1,2}(B))) \\
 &= (F_{2,3}(F_{1,2}(A)), G_{2,3}(G_{1,2}(B))) \\
 &= (F_{2,3} \times G_{2,3})(F_{1,2}(A), G_{1,2}(B)) \\
 &= (F_{2,3} \times G_{2,3})((F_{1,2} \times G_{1,2})(A, B)) \\
 &= [(F_{2,3} \times G_{2,3}) \circ_{\mathbf{Cat}} (F_{1,2} \times G_{1,2})](A, B).
 \end{aligned}$$

- For each $A_1, A_2 \in \text{Obj}(\mathcal{C}_1)$, $B_1, B_2 \in \text{Obj}(\mathcal{D}_1)$ Then

$$\begin{aligned}
 (F_{2,3} \circ_{\mathbf{Cat}} F_{1,2}) \times (G_{2,3} \circ_{\mathbf{Cat}} G_{1,2})(f, g) &= (F_{2,3} \circ_{\mathbf{Cat}} F_{1,2}(f), G_{2,3} \circ_{\mathbf{Cat}} G_{1,2}(g)) \\
 &= (F_{2,3}(F_{1,2}(f)), G_{2,3}(G_{1,2}(g))) \\
 &= (F_{2,3} \times G_{2,3})(F_{1,2}(f), G_{1,2}(g)) \\
 &= (F_{2,3} \times G_{2,3})((F_{1,2} \times G_{1,2})(f, g)) \\
 &= (F_{2,3} \times G_{2,3}) \circ_{\mathbf{Cat}} (f, g).
 \end{aligned}$$

Hence $(F_{2,3} \circ_{\mathbf{Cat}} F_{1,2}) \times (G_{2,3} \circ_{\mathbf{Cat}} G_{1,2}) = (F_{2,3} \times G_{2,3}) \circ_{\mathbf{Cat}} (F_{1,2} \times G_{1,2})$. □

Exercise 8.2.1.13. Let $\mathcal{C}, \mathcal{D} \in \text{Obj}(\mathbf{Cat})$. Then $\text{id}_{\mathcal{C}}^{\mathbf{Cat}} \times \text{id}_{\mathcal{D}}^{\mathbf{Cat}} = \text{id}_{\mathcal{C} \times \mathcal{D}}^{\mathbf{Cat}}$.

Proof.

- For each $(A, B) \in \text{Obj}(\mathcal{C} \times \mathcal{D})$,

$$\begin{aligned} \text{id}_{\mathcal{C}}^{\text{Cat}} \times \text{id}_{\mathcal{D}}^{\text{Cat}}(A, B) &= (\text{id}_{\mathcal{C}}^{\text{Cat}}(A), \text{id}_{\mathcal{D}}^{\text{Cat}}(B)) \\ &= (A, B) \\ &= \text{id}_{\mathcal{C} \times \mathcal{D}}^{\text{Cat}}(A, B). \end{aligned}$$

- For each $(f, g) \in \text{Hom}_{\mathcal{C} \times \mathcal{D}}$,

$$\begin{aligned} \text{id}_{\mathcal{C}}^{\text{Cat}} \times \text{id}_{\mathcal{D}}^{\text{Cat}}(f, g) &= (\text{id}_{\mathcal{C}}^{\text{Cat}}(f), \text{id}_{\mathcal{D}}^{\text{Cat}}(g)) \\ &= (f, g) \\ &= \text{id}_{\mathcal{C} \times \mathcal{D}}^{\text{Cat}}(f, g). \end{aligned}$$

Hence $\text{id}_{\mathcal{C}}^{\text{Cat}} \times \text{id}_{\mathcal{D}}^{\text{Cat}} = \text{id}_{\mathcal{C} \times \mathcal{D}}^{\text{Cat}}$. □

Definition 8.2.1.14. We define the **product functor on $\mathbf{Cat}$** , denoted $(- \times -) : \mathbf{Cat} \times \mathbf{Cat} \rightarrow \mathbf{Cat}$, by

- $(- \times -)(\mathcal{A}, \mathcal{B}) := \mathcal{A} \times \mathcal{B}$, $(\mathcal{A}, \mathcal{B}) \in \text{Obj}(\mathbf{Cat} \times \mathbf{Cat})$,
- $(- \times -)(F, G) := F \times G$, $(\mathcal{C}_1, \mathcal{D}_1), (\mathcal{C}_2, \mathcal{D}_2) \in \text{Obj}(\mathbf{Cat} \times \mathbf{Cat})$ and $(F, G) \in \text{Hom}_{\mathbf{Cat} \times \mathbf{Cat}}((\mathcal{C}_1, \mathcal{D}_1), (\mathcal{C}_2, \mathcal{D}_2))$.

Exercise 8.2.1.15. We have that $(- \times -) : \mathbf{Cat} \times \mathbf{Cat} \rightarrow \mathbf{Cat}$ is a functor.

Proof.

1. Let $(\mathcal{C}_1, \mathcal{D}_1), (\mathcal{C}_2, \mathcal{D}_2) \in \text{Obj}(\mathbf{Cat} \times \mathbf{Cat})$ and $(F, G) \in \text{Hom}_{\mathbf{Cat} \times \mathbf{Cat}}((\mathcal{C}_1, \mathcal{D}_1), (\mathcal{C}_2, \mathcal{D}_2))$. Exercise 8.2.1.11 implies that

$$\begin{aligned} (- \times -)(F, G) &= F \times G \\ &\in \text{Hom}_{\mathbf{Cat}}(\mathcal{C}_1 \times \mathcal{D}_1, \mathcal{C}_2 \times \mathcal{D}_2) \\ &= \text{Hom}_{\mathbf{Cat}}((- \times -)(\mathcal{C}_1, \mathcal{D}_1), (- \times -)(\mathcal{C}_2, \mathcal{D}_2)). \end{aligned}$$

2. Let $(\mathcal{C}_1, \mathcal{D}_1), (\mathcal{C}_2, \mathcal{D}_2), (\mathcal{C}_3, \mathcal{D}_3) \in \text{Obj}(\mathbf{Cat} \times \mathbf{Cat})$, $(F_{1,2}, G_{1,2}) \in \text{Hom}_{\mathbf{Cat} \times \mathbf{Cat}}((\mathcal{C}_1, \mathcal{D}_1), (\mathcal{C}_2, \mathcal{D}_2))$ and $(F_{2,3}, G_{2,3}) \in \text{Hom}_{\mathbf{Cat} \times \mathbf{Cat}}((\mathcal{C}_2, \mathcal{D}_2), (\mathcal{C}_3, \mathcal{D}_3))$. Exercise 8.2.1.12 implies that

$$\begin{aligned} (- \times -)((F_{2,3}, G_{2,3}) \circ_{\mathbf{Cat} \times \mathbf{Cat}} (F_{1,2}, G_{1,2})) &= (- \times -)(F_{2,3} \circ_{\mathbf{Cat}} F_{1,2}, G_{2,3} \circ_{\mathbf{Cat}} G_{1,2}) \\ &= (F_{2,3} \circ_{\mathbf{Cat}} F_{1,2}) \times (G_{2,3} \circ_{\mathbf{Cat}} G_{1,2}) \\ &= (F_{2,3} \times G_{2,3}) \circ_{\mathbf{Cat}} (F_{1,2} \times G_{1,2}) \\ &= (- \times -)(F_{2,3}, G_{2,3}) \circ_{\mathbf{Cat}} (- \times -)(F_{1,2}, G_{1,2}). \end{aligned}$$

3. Exercise 8.2.1.13 implies that for each $(\mathcal{C}, \mathcal{D}) \in \text{Obj}(\mathbf{Cat} \times \mathbf{Cat})$,

$$\begin{aligned} (- \times -)(\text{id}_{(\mathcal{C}, \mathcal{D})}^{\mathbf{Cat} \times \mathbf{Cat}}) &= (- \times -)(\text{id}_{\mathcal{C}}^{\mathbf{Cat}}, \text{id}_{\mathcal{D}}^{\mathbf{Cat}}) \\ &= \text{id}_{\mathcal{C}}^{\mathbf{Cat}} \times \text{id}_{\mathcal{D}}^{\mathbf{Cat}} \\ &= \text{id}_{\mathcal{C} \times \mathcal{D}}^{\mathbf{Cat}} \\ &= \text{id}_{(- \times -)(\mathcal{C}, \mathcal{D})}^{\mathbf{Cat}}. \end{aligned}$$

Hence $(- \times -) : \mathbf{Cat} \times \mathbf{Cat} \rightarrow \mathbf{Cat}$ is a functor. □

Exercise 8.2.1.16. maybe switch order of 'basic construction and operations' and 'fundamental structures' and redo the previous exercises by showing that if $\mathcal{C}$ has finite products, then $\times : \mathcal{C} \times \mathcal{C} \rightarrow \mathcal{C}$ given by $\times(A, B) := A \times B$ and $\times(f, g) := f \times g$ is a functor and then the previous exercise would be a corollary with $\mathcal{C} = \mathbf{Cat}$ (if we show that $(\mathcal{C} \times \mathcal{D}, (\pi_{\mathcal{C}}, \pi_{\mathcal{D}})) \in \text{Prod}_{\mathbf{Cat}}(\mathcal{C}, \mathcal{D})$ and $F \times G$ is the product of morphisms in $\mathbf{Cat}$ as defined in the section on products in the basic).

8.2.2 Slicing

Definition 8.2.2.1. Let $\mathcal{A}, \mathcal{B}, \mathcal{D}$ be categories, $F : \mathcal{A} \times \mathcal{B} \rightarrow \mathcal{D}$ and $A_0 \in \text{Obj}(\mathcal{A})$, $B_0 \in \text{Obj}(\mathcal{B})$. We define the

- **slice functor of F at B_0** , denoted $F(\cdot, B_0) : \mathcal{A} \rightarrow \mathcal{D}$ by
 - $F(\cdot, B_0)(A) := F(A, B_0)$, $A \in \text{Obj}(\mathcal{A})$
 - $F(\cdot, B_0)(f) := F(f, \text{id}_{B_0}^{\mathcal{B}})$, $f \in \text{Hom}_{\mathcal{A}}$
- **slice functor of F at A_0** , denoted $F(A_0, \cdot) : \mathcal{B} \rightarrow \mathcal{D}$
 - $F(A_0, \cdot)(B) := F(A_0, B)$, $B \in \mathcal{B}$
 - $F(A_0, \cdot)(g) := F(\text{id}_{A_0}^{\mathcal{A}}, g)$, $g \in \text{Hom}_{\mathcal{B}}$

Exercise 8.2.2.2. Let $\mathcal{A}, \mathcal{B}, \mathcal{D}$ be categories, $F : \mathcal{A} \times \mathcal{B} \rightarrow \mathcal{D}$. Then

1. for each $B_0 \in \text{Obj}(\mathcal{B})$, $F(\cdot, B_0) : \mathcal{A} \rightarrow \mathcal{D}$ is a functor,
2. for each $A_0 \in \text{Obj}(\mathcal{A})$, $F(A_0, \cdot) : \mathcal{B} \rightarrow \mathcal{D}$ is a functor.

Proof.

1. Let $B_0 \in \text{Obj}(\mathcal{B})$.
 - (a) Let $A_1, A_2 \in \text{Obj}(\mathcal{A})$ and $f_{12} \in \text{Hom}_{\mathcal{A}}(A_1, A_2)$. Then $(f_{12}, \text{id}_{B_0}^{\mathcal{B}}) \in \text{Hom}_{\mathcal{A} \times \mathcal{B}}((A_1, B_0), (A_2, B_0))$. Since $F : \mathcal{A} \times \mathcal{B} \rightarrow \mathcal{D}$, we have that

$$\begin{aligned} F(\cdot, B_0)(f_{12}) &= F(f_{12}, \text{id}_{B_0}^{\mathcal{B}}) \\ &\in \text{Hom}_{\mathcal{D}}(F(A_1, B_0), F(A_2, B_0)) \\ &= \text{Hom}_{\mathcal{D}}(F(\cdot, B_0)(A_1), F(\cdot, B_0)(A_2)) \end{aligned}$$

- (b) Let $A_1, A_2, A_3 \in \text{Obj}(\mathcal{A})$, $f_{12} \in \text{Hom}_{\mathcal{A}}(A_1, A_2)$ and $f_{23} \in \text{Hom}_{\mathcal{A}}(A_2, A_3)$. Since $F : \mathcal{A} \times \mathcal{B} \rightarrow \mathcal{D}$, we have that

$$\begin{aligned} F(\cdot, B_0)(f_{23} \circ_{\mathcal{A}} f_{12}) &= F(f_{23} \circ_{\mathcal{A}} f_{12}, \text{id}_{B_0}^{\mathcal{B}}) \\ &= F(f_{23} \circ_{\mathcal{A}} f_{12}, \text{id}_{B_0}^{\mathcal{B}} \circ_{\mathcal{B}} \text{id}_{B_0}^{\mathcal{B}}) \\ &= F[(f_{23}, \text{id}_{B_0}^{\mathcal{B}}) \circ_{\mathcal{A} \times \mathcal{B}} (f_{12}, \text{id}_{B_0}^{\mathcal{B}})] \\ &= F(f_{23}, \text{id}_{B_0}^{\mathcal{B}}) \circ_{\mathcal{D}} F(f_{12}, \text{id}_{B_0}^{\mathcal{B}}) \\ &= F(\cdot, B_0)(f_{23}) \circ_{\mathcal{D}} F(\cdot, B_0)(f_{12}). \end{aligned}$$

- (c) Let $A \in \text{Obj}(\mathcal{A})$. Then

$$\begin{aligned} F(\cdot, B_0)(\text{id}_A^{\mathcal{A}}) &= F(\text{id}_A^{\mathcal{A}}, \text{id}_{B_0}^{\mathcal{B}}) \\ &= F(\text{id}_{(A, B_0)}^{A \times B}) \\ &= \text{id}_{F(A, B_0)}^{\mathcal{D}} \\ &= \text{id}_{F(\cdot, B_0)(A)}^{\mathcal{D}}. \end{aligned}$$

Thus $F(\cdot, B_0) : \mathcal{A} \rightarrow \mathcal{D}$ is a functor.

2. Similar to previous part [ref prev part](#).

□

Definition 8.2.2.3. Let $\mathcal{A}, \mathcal{B}$ be categories, $A_0 \in \text{Obj}(\mathcal{A})$ and $B_0 \in \text{Obj}(\mathcal{B})$. We define the

- **embedding functor of $\mathcal{A}$ into $\mathcal{A} \times \mathcal{B}$ at B_0** , denoted $\iota_{\mathcal{A}}^{B_0} : \mathcal{A} \rightarrow \mathcal{A} \times \mathcal{B}$, by $\iota_{\mathcal{A}}^{B_0} := \text{id}_{\mathcal{A}} \times \text{id}_{\mathcal{B}}(\cdot, B_0)$

- **embedding functor of $\mathcal{B}$ into $\mathcal{A} \times \mathcal{B}$ at A_0** , denoted $\iota_{\mathcal{B}}^{A_0} : \mathcal{B} \rightarrow \mathcal{A} \times \mathcal{B}$, by $\iota_{\mathcal{B}}^{A_0} := \text{id}_{\mathcal{A}} \times \text{id}_{\mathcal{B}}(A_0, \cdot)$.

Exercise 8.2.2.4. Let $\mathcal{A}, \mathcal{B}$ be categories. Then

1. for each $B_0 \in \text{Obj}(\mathcal{B})$, $\iota_{\mathcal{A}}^{B_0} : \mathcal{A} \rightarrow \mathcal{A} \times \mathcal{B}$ is a functor and

- $\iota_{\mathcal{A}}^{B_0}(A) := (A, B_0)$, $A \in \text{Obj}(\mathcal{A})$,
- $\iota_{\mathcal{A}}^{B_0}(f) := (f, \text{id}_{B_0}^{\mathcal{B}})$, $f \in \text{Hom}_{\mathcal{A}}$,

2. for each $A_0 \in \text{Obj}(\mathcal{B})$, $\iota_{\mathcal{B}}^{A_0} : \mathcal{B} \rightarrow \mathcal{A} \times \mathcal{B}$ is a functor and

- $\iota_{\mathcal{B}}^{A_0}(B) := (A_0, B)$, $B \in \text{Obj}(\mathcal{B})$,
- $\iota_{\mathcal{B}}^{A_0}(g) := (\text{id}_{A_0}^{\mathcal{A}}, g)$, $g \in \text{Hom}_{\mathcal{B}}$.

Proof.

1. Let $B_0 \in \text{Obj}(\mathcal{B})$. By **prev exercises**, $\iota_{\mathcal{A}}^{B_0}$ is a functor. By definition, we have that

- for each $A \in \text{Obj}(\mathcal{A})$,

$$\begin{aligned} \iota_{\mathcal{A}}^{B_0}(A) &= \text{id}_{\mathcal{A}} \times \text{id}_{\mathcal{B}}(A, B_0) \\ &= (\text{id}_{\mathcal{A}}(A), \text{id}_{\mathcal{B}}(B_0)) \\ &= (A, B_0), \end{aligned}$$

- for each $f \in \text{Hom}_{\mathcal{A}}$,

$$\begin{aligned} \iota_{\mathcal{A}}^{B_0}(f) &= \text{id}_{\mathcal{A}} \times \text{id}_{\mathcal{B}}(f, \text{id}_{B_0}^{\mathcal{B}}) \\ &= (\text{id}_{\mathcal{A}}(f), \text{id}_{\mathcal{B}}(\text{id}_{B_0}^{\mathcal{B}})) \\ &= (f, \text{id}_{B_0}^{\mathcal{B}}). \end{aligned}$$

2. Similar to prev part (ref prev part)

□

Definition 8.2.2.5. Let $\mathcal{A}, \mathcal{B}, \mathcal{C}$ be categories, $F, G : \mathcal{A} \times \mathcal{B} \rightarrow \mathcal{C}$, $\alpha : F \Rightarrow G$, $A_0 \in \text{Obj}(\mathcal{A})$ and $B_0 \in \text{Obj}(\mathcal{B})$. We define the

- **slice transformation of α at A_0** , denoted $\alpha_{(A_0, \cdot)} : F(A_0, \cdot) \Rightarrow G(A_0, \cdot)$, by $(\alpha_{(A_0, \cdot)})_B := \alpha_{(A_0, B)}$
- **slice transformation of α at B_0** , denoted $\alpha_{(\cdot, B_0)} : F(\cdot, B_0) \Rightarrow G(\cdot, B_0)$, by $(\alpha_{(\cdot, B_0)})_A := \alpha_{(A, B_0)}$.

Exercise 8.2.2.6. Let $\mathcal{A}, \mathcal{B}, \mathcal{C}$ be categories, $F, G : \mathcal{A} \times \mathcal{B} \rightarrow \mathcal{C}$ and $\alpha : F \Rightarrow G$. Then

1. for each $A_0 \in \text{Obj}(\mathcal{A})$, $\alpha_{(A_0, \cdot)} : F(A_0, \cdot) \Rightarrow G(A_0, \cdot)$ is a natural transformation,
2. for each $B_0 \in \text{Obj}(\mathcal{B})$, $\alpha_{(\cdot, B_0)} : F(\cdot, B_0) \Rightarrow G(\cdot, B_0)$ is a natural transformation.

Proof.

1. Let $A_0 \in \text{Obj}(\mathcal{A})$.

- By construction, for each $B \in \text{Obj}(\mathcal{B})$,

$$\begin{aligned} (\alpha_{(A_0, \cdot)})_B &= \alpha_{(A_0, B)} \\ &\in \text{Hom}_{\mathcal{C}}(F(A_0, B), G(A_0, B)) \\ &= \text{Hom}_{\mathcal{C}}(F(A_0, \cdot)(B), G(A_0, \cdot)(B)). \end{aligned}$$

- Let $B_1, B_2 \in \text{Obj}(\mathcal{B})$ and $g \in \text{Hom}_{\mathcal{B}}(B_1, B_2)$. Since $\alpha : F \Rightarrow G$, we have that for each $g \in \text{Hom}_{\mathcal{B}}(B_1, B_2)$, $G(\text{id}_{A_0}^A, g) \circ_C \alpha_{(A_0, B_1)} = \alpha_{(A_0, B_2)} \circ_C F(\text{id}_{A_0}^A, g)$, i.e. the following diagram commutes:

$$\begin{array}{ccc} F(A_0, B_1) & \xrightarrow{\alpha_{(A_0, B_1)}} & G(A_0, B_1) \\ F(\text{id}_{A_0}^A, g) \downarrow & & \downarrow G(\text{id}_{A_0}^A, g) \\ F(A_0, B_2) & \xrightarrow{\alpha_{(A_0, B_2)}} & G(A_0, B_2). \end{array}$$

Therefore

$$\begin{aligned} G(A_0, \cdot)(g) \circ_C (\alpha_{(A_0, \cdot)})_{B_1} &= G(\text{id}_{A_0}^A, g) \circ_C \alpha_{(A_0, B_1)} \\ &= \alpha_{(A_0, B_2)} \circ_C F(\text{id}_{A_0}^A, g) \\ &= (\alpha_{(A_0, \cdot)})_{B_2} \circ_C F(A_0, \cdot)(g), \end{aligned}$$

i.e. the following diagram commutes:

$$\begin{array}{ccc} F(A_0, \cdot)(B_1) & \xrightarrow{(\alpha_{(A_0, \cdot)})_{B_1}} & G(A_0, \cdot)(B_1) \\ F(A_0, \cdot)(g) \downarrow & & \downarrow G(A_0, \cdot)(g) \\ F(A_0, \cdot)(B_2) & \xrightarrow{(\alpha_{(A_0, \cdot)})_{B_2}} & G(A_0, \cdot)(B_2). \end{array}$$

Since $B_1, B_2 \in \text{Obj}(\mathcal{B})$ and $g \in \text{Hom}_{\mathcal{B}}(B_1, B_2)$ are arbitrary, we have that for each $B_1, B_2 \in \text{Obj}(\mathcal{B})$, $G(A_0, \cdot)(g) \circ_C (\alpha_{(A_0, \cdot)})_{B_1} = (\alpha_{(A_0, \cdot)})_{B_2} \circ_C F(A_0, \cdot)(g)$.

Thus $\alpha_{(A_0, \cdot)} : F(A_0, \cdot) \Rightarrow G(A_0, \cdot)$ is a natural transformation. Since $A_0 \in \text{Obj}(\mathcal{A})$ is arbitrary, we have that for each $A_0 \in \text{Obj}(\mathcal{A})$, $\alpha_{(A_0, \cdot)} : F(A_0, \cdot) \Rightarrow G(A_0, \cdot)$ is a natural transformation.

2. **Similar to previous part (ref prev part)**

□

Exercise 8.2.2.7. Let $\mathcal{A}, \mathcal{B}, \mathcal{C}$ be categories, $F_1, F_2, F_3 : \mathcal{A} \times \mathcal{B} \rightarrow \mathcal{C}$, $\alpha_{1,2} : F_1 \Rightarrow F_2$ and $\alpha_{2,3} : F_2 \Rightarrow F_3$. Then

1. for each $A_0 \in \text{Obj}(\mathcal{A})$, $(\alpha_{2,3} \circ \alpha_{1,2})_{(A_0, \cdot)} = (\alpha_{2,3})_{(A_0, \cdot)} \circ (\alpha_{1,2})_{(A_0, \cdot)}$.
2. for each $B_0 \in \text{Obj}(\mathcal{B})$, $(\alpha_{2,3} \circ \alpha_{1,2})_{(\cdot, B_0)} = (\alpha_{2,3})_{(\cdot, B_0)} \circ (\alpha_{1,2})_{(\cdot, B_0)}$.

Proof.

1. Let $A_0 \in \text{Obj}(\mathcal{A})$ and $B \in \text{Obj}(\mathcal{B})$. Then

$$\begin{aligned} [(\alpha_{2,3} \circ \alpha_{1,2})_{(A_0, \cdot)}]_B &= (\alpha_{2,3} \circ \alpha_{1,2})_{(A_0, B)} \\ &= (\alpha_{2,3})_{(A_0, B)} \circ (\alpha_{1,2})_{(A_0, B)} \\ &= [(\alpha_{2,3})_{(A_0, \cdot)}]_B \circ [(\alpha_{1,2})_{(A_0, \cdot)}]_B \\ &= [(\alpha_{2,3})_{(A_0, \cdot)} \circ (\alpha_{1,2})_{(A_0, \cdot)}]_B. \end{aligned}$$

Since $B \in \text{Obj}(\mathcal{B})$ is arbitrary, we have that for each $B \in \text{Obj}(\mathcal{B})$, $[(\alpha_{2,3} \circ \alpha_{1,2})_{(A_0, \cdot)}]_B = [(\alpha_{2,3})_{(A_0, \cdot)} \circ (\alpha_{1,2})_{(A_0, \cdot)}]_B$. Thus $(\alpha_{2,3} \circ \alpha_{1,2})_{(A_0, \cdot)} = (\alpha_{2,3})_{(A_0, \cdot)} \circ (\alpha_{1,2})_{(A_0, \cdot)}$. Since $A_0 \in \text{Obj}$ is arbitrary, we have that for each $A_0 \in \text{Obj}(\mathcal{A})$, $(\alpha_{2,3} \circ \alpha_{1,2})_{(A_0, \cdot)} = (\alpha_{2,3})_{(A_0, \cdot)} \circ (\alpha_{1,2})_{(A_0, \cdot)}$.

2. **Similar to prev part (ref prev part)**

□

Exercise 8.2.2.8. Let $\mathcal{A}, \mathcal{B}, \mathcal{C}$ be categories and $F \in \mathcal{A} \times \mathcal{B} \rightarrow \mathcal{C}$. Then

1. for each $A_0 \in \text{Obj}(\mathcal{A})$, $(\text{id}_F)_{(A_0, \cdot)} = \text{id}_{F(A_0, \cdot)}$,

2. for each $B_0 \in \text{Obj}(\mathcal{A})$, $(\text{id}_F)_{(\cdot, B_0)} = \text{id}_{F(\cdot, B_0)}$.

Proof.

1. Let $A_0 \in \text{Obj}(\mathcal{A})$. Then for each $B \in \text{Obj}(\mathcal{B})$,

$$\begin{aligned} [(\text{id}_F)_{(A_0, \cdot)}]_B &= (\text{id}_F)_{(A_0, B)} \\ &= \text{id}_{F(A_0, B)} \\ &= \text{id}_{F(A_0, \cdot)(B)} \\ &= (\text{id}_{F(A_0, \cdot)})_B. \end{aligned}$$

Hence $(\text{id}_F)_{(A_0, \cdot)} = \text{id}_{F(A_0, \cdot)}$. Since $A_0 \in \text{Obj}(\mathcal{A})$ is arbitrary, we have that for each $A_0 \in \text{Obj}(\mathcal{A})$, $(\text{id}_F)_{(A_0, \cdot)} = \text{id}_{F(A_0, \cdot)}$.

2. **Similar to (ref) prev part**

□

8.3 Functor Categories

Definition 8.3.0.1. Let $\mathcal{C}, \mathcal{D} \in \text{Obj}(\mathbf{Cat})$. We define the **evaluation functor from $\mathcal{C} \times \mathcal{D}^c$ to $\mathcal{D}$** , denoted $\text{eval}_{(\mathcal{C}, \mathcal{D})} : \mathcal{C} \times \mathcal{D}^c \rightarrow \mathcal{D}$, by

- $\text{eval}_{(\mathcal{C}, \mathcal{D})}(A, F) := F(A), \quad (A, F) \in \text{Obj}(\mathcal{C} \times \mathcal{D}^c)$
- $\text{eval}_{(\mathcal{C}, \mathcal{D})}(f, \alpha) := G(f) \circ_{\mathcal{D}} \alpha_A, \quad (f, \alpha) \in \text{Hom}_{\mathcal{C} \times \mathcal{D}^c}$

Exercise 8.3.0.2. Let $\mathcal{C}, \mathcal{D} \in \text{Obj}(\mathbf{Cat})$. Then $\text{eval}_{(\mathcal{C}, \mathcal{D})} : \text{eval} : \mathcal{C} \times \mathcal{D}^c \rightarrow \mathcal{D}$ is a functor.

Proof.

1. Let $(A, F), (B, G) \in \text{Obj}(\mathcal{C} \times \mathcal{D}^c)$ and $(f, \alpha) \in \text{Hom}_{\mathcal{C} \times \mathcal{D}^c}((A, F), (B, G))$. Then $G(f) \in \text{Hom}_{\mathcal{D}}(G(A), G(B))$ and $\alpha_A \in \text{Hom}_{\mathcal{D}}(F(A), G(A))$. Hence

$$\begin{aligned} \text{eval}_{(\mathcal{C}, \mathcal{D})}(f, \alpha) &= G(f) \circ_{\mathcal{D}} \alpha_A \\ &\in \text{Hom}_{\mathcal{D}}(F(A), G(B)) \\ &= \text{Hom}_{\mathcal{D}}(\text{eval}_{(\mathcal{C}, \mathcal{D})}(A, F), \text{eval}_{(\mathcal{C}, \mathcal{D})}(B, G)). \end{aligned}$$

2. Let $(A, F), (B, G), (C, H) \in \text{Obj}(\mathcal{C} \times \mathcal{D}^c)$, $(f, \alpha) \in \text{Hom}_{\mathcal{C} \times \mathcal{D}^c}((A, F), (B, G))$ and $(g, \beta) \in \text{Hom}_{\mathcal{C} \times \mathcal{D}^c}((B, G), (C, H))$. Since $\beta \in \text{Hom}_{\mathcal{D}^c}(G, H)$, we have that $H(f) \circ_{\mathcal{D}} \beta_A = \beta_B \circ_{\mathcal{D}} G(f)$, i.e. the following diagram commutes:

$$\begin{array}{ccc} G(A) & \xrightarrow{\beta_A} & H(A) \\ G(f) \downarrow & & \downarrow H(f) \\ G(B) & \xrightarrow{\beta_B} & H(B). \end{array}$$

$$\begin{aligned} \text{eval}_{(\mathcal{C}, \mathcal{D})}((g, \beta) \circ_{\mathcal{C} \times \mathcal{D}^c} (f, \alpha)) &= \text{eval}_{(\mathcal{C}, \mathcal{D})}(g \circ_{\mathcal{C}} f, \beta \circ_{\mathcal{D}^c} \alpha) \\ &= H(g \circ_{\mathcal{C}} f) \circ_{\mathcal{D}} (\beta \circ_{\mathcal{D}^c} \alpha)_A \\ &= H(g) \circ_{\mathcal{D}} (H(f) \circ_{\mathcal{D}} \beta_A) \circ_{\mathcal{D}} \alpha_A \\ &= H(g) \circ_{\mathcal{D}} (\beta_B \circ_{\mathcal{D}} G(f)) \circ_{\mathcal{D}} \alpha_A \\ &= (H(g) \circ_{\mathcal{D}} \beta_B) \circ_{\mathcal{D}} (G(f) \circ_{\mathcal{D}} \alpha_A) \\ &= \text{eval}_{(\mathcal{C}, \mathcal{D})}(g, \beta) \circ_{\mathcal{D}} \text{eval}_{(\mathcal{C}, \mathcal{D})}(f, \alpha). \end{aligned}$$

3. For each $(A, F) \in \text{Obj}(\mathcal{C} \times \mathcal{D}^c)$, we have that

$$\begin{aligned} \text{eval}_{(\mathcal{C}, \mathcal{D})}(\text{id}_{(A, F)}^{\mathcal{C} \times \mathcal{D}^c}) &= \text{eval}_{(\mathcal{C}, \mathcal{D})}(\text{id}_A^{\mathcal{C}}, \text{id}_F^{\mathcal{D}^c}) \\ &= \text{eval}_{(\mathcal{C}, \mathcal{D})}(\text{id}_A^{\mathcal{C}}, \text{id}_F^{\mathcal{D}^c}) \\ &= F(\text{id}_A^{\mathcal{C}}) \circ_{\mathcal{D}} (\text{id}_F^{\mathcal{D}^c})_A \\ &= \text{id}_{F(A)}^{\mathcal{D}} \circ_{\mathcal{D}} \text{id}_{F(A)}^{\mathcal{D}} \\ &= \text{id}_{F(A)}^{\mathcal{D}} \\ &= \text{id}_{\text{eval}_{(\mathcal{C}, \mathcal{D})}(A, F)}^{\mathcal{D}} \end{aligned}$$

Hence $\text{eval}_{(\mathcal{C}, \mathcal{D})} : \mathcal{C} \times \mathcal{D}^c \rightarrow \mathcal{D}$ is a functor. □

Exercise 8.3.0.3. Find two functors $F, G : \mathbf{Cat} \times \mathbf{Cat} \rightarrow \mathbf{Cat}$ such that $\text{eval} \mapsto \text{eval}_{(\mathcal{C}, \mathcal{D})}$ is a natural isomorphism $F \Rightarrow G$.

Definition 8.3.0.4. Let $(\mathcal{A}_1, \mathcal{B}_1), (\mathcal{A}_2, \mathcal{B}_2) \in \text{Obj}(\mathbf{Cat}^{\text{op}} \times \mathbf{Cat})$, $(F, G) \in \text{Hom}_{\mathbf{Cat}^{\text{op}} \times \mathbf{Cat}}((\mathcal{A}_1, \mathcal{B}_1), (\mathcal{A}_2, \mathcal{B}_2))$, $H, J \in \text{Obj}(\mathcal{B}_1^{\mathcal{A}_1})$ and $\alpha \in \text{Hom}_{\mathcal{B}_1^{\mathcal{A}_1}}(H, J)$.

- For each $H \in \text{Obj}(\mathcal{B}_1^{\mathcal{A}_1})$, we define the **internal Hom functor object induced by F and G at H** , denoted $G^F(H) \in \text{Obj}(\mathcal{B}_2^{\mathcal{A}_2})$, by $G^F(H) := G \circ_{\mathbf{Cat}} H \circ_{\mathbf{Cat}} F$
- **internal Hom functor natural transformation induced by F and G at α** , denoted $G^F(\alpha) : G^F(H) \Rightarrow G^F(J)$, by $G^F(\alpha)_A := G(\alpha_{F(A)})$.

Exercise 8.3.0.5. Let $(\mathcal{A}_1, \mathcal{B}_1), (\mathcal{A}_2, \mathcal{B}_2) \in \text{Obj}(\mathbf{Cat}^{\text{op}} \times \mathbf{Cat})$, $(F, G) \in \text{Hom}_{\mathbf{Cat}^{\text{op}} \times \mathbf{Cat}}((\mathcal{A}_1, \mathcal{B}_1), (\mathcal{A}_2, \mathcal{B}_2))$, $H, J \in \text{Obj}(\mathcal{B}_1^{\mathcal{A}_1})$ and $\alpha \in \text{Hom}_{\mathcal{B}_1^{\mathcal{A}_1}}(H, J)$. Then $G^F(\alpha) : G^F(H) \Rightarrow G^F(J)$ is a natural transformation.

Proof.

- Let $A \in \text{Obj}(\mathcal{A}_2)$. Since $\alpha \in \text{Hom}_{\mathcal{B}_1^{\mathcal{A}_1}}(H, J)$, we have that $\alpha_{F(A)} \in \text{Hom}_{\mathcal{B}_1}(H(F(A)), J(F(A)))$. Then

$$\begin{aligned} G^F(\alpha)_A &= G(\alpha_{F(A)}) \\ &\in \text{Hom}_{\mathcal{B}_2}(G(H(F(A))), G(J(F(A)))) \\ &= \text{Hom}_{\mathcal{B}_2}(G^F(H)(A), G^F(J)(A)). \end{aligned}$$

Since $A \in \text{Obj}(\mathcal{A}_2)$ is arbitrary, we have that for each $A \in \text{Obj}(\mathcal{A}_2)$, $G^F(\alpha)_A \in \text{Hom}_{\mathcal{B}_2}(G^F(H)(A), G^F(J)(A))$.

- Let $A_1, A_2 \in \text{Obj}(\mathcal{A}_2)$ and $f \in \text{Hom}_{\mathcal{A}_2}(A_1, A_2)$. Define $A'_1, A'_2 \in \text{Obj}(\mathcal{A}_1)$ and $f' \in \text{Hom}_{\mathcal{A}_1}(A'_1, A'_2)$ by $A'_k := F(A_k)$ and $f' := F(f)$. Since $\alpha \in \text{Hom}_{\mathcal{B}_1^{\mathcal{A}_1}}(H, J)$, we have that

$$J(f') \circ_{\mathcal{B}_1} \alpha_{A'_1} = \alpha_{A'_2} \circ_{\mathcal{B}_1} H(f'),$$

i.e. the following diagram commutes:

$$\begin{array}{ccc} H(A'_1) & \xrightarrow{\alpha_{A'_1}} & J(A'_1) \\ H(f') \downarrow & & \downarrow J(f') \\ H(A'_2) & \xrightarrow{\alpha_{A'_2}} & J(A'_2). \end{array}$$

Therefore,

$$\begin{aligned} G^F(J)(f) \circ_{\mathcal{B}_2} G^F(\alpha)_{A_1} &= [G \circ_{\mathbf{Cat}} J \circ_{\mathbf{Cat}} F](f) \circ_{\mathcal{B}_2} G(\alpha_{F(A_1)}) \\ &= G(J(F(f))) \circ_{\mathcal{B}_2} G(\alpha_{F(A_1)}) \\ &= G(J(f')) \circ_{\mathcal{B}_1} G(\alpha_{A'_2}) \\ &= G(J(f') \circ_{\mathcal{B}_1} \alpha_{A'_1}) \\ &= G(\alpha_{A'_2} \circ_{\mathcal{B}_1} H(f')) \\ &= G(\alpha_{A'_2}) \circ_{\mathcal{B}_2} G(H(f')) \\ &= G(\alpha_{F(A_2)}) \circ_{\mathcal{B}_2} G(H(F(f))) \\ &= G(\alpha_{F(A_2)}) \circ_{\mathcal{B}_2} [G \circ_{\mathbf{Cat}} H \circ_{\mathbf{Cat}} F](f) \\ &= G^F(\alpha)_{A_2} \circ_{\mathcal{B}_2} G^F(H)(f), \end{aligned}$$

i.e. the following diagram commutes:

$$\begin{array}{ccc} G^F(H)(A_1) & \xrightarrow{(G^F_\alpha)_{A_1}} & G^F(J)(A_1) \\ G^F(H)(f) \downarrow & & \downarrow G^F(J)(f) \\ G^F(H)(A_2) & \xrightarrow{(G^F_\alpha)_{A_2}} & G^F(J)(A_2). \end{array}$$

Since $A_1, A_2 \in \text{Obj}(\mathcal{A}_2)$ and $f \in \text{Hom}_{\mathcal{A}_2}(A_1, A_2)$ are arbitrary, we have that for each $A_1, A_2 \in \text{Obj}(\mathcal{A}_2)$ and $f \in \text{Hom}_{\mathcal{A}_2}(A_1, A_2)$, $G^F(J)(f) \circ_{\mathcal{B}_2} G^F(\alpha)_{A_1} = G^F(\alpha)_{A_2} \circ_{\mathcal{B}_2} G^F(H)(f)$.

Thus $G^F(\alpha) : G^F(H) \Rightarrow G^F(J)$ is a natural transformation. □

Exercise 8.3.0.6. Let $(\mathcal{A}_1, \mathcal{B}_1), (\mathcal{A}_2, \mathcal{B}_2) \in \text{Obj}(\mathbf{Cat}^{\text{op}} \times \mathbf{Cat})$, $(F, G) \in \text{Hom}_{\mathbf{Cat}^{\text{op}} \times \mathbf{Cat}}((\mathcal{A}_1, \mathcal{B}_1), (\mathcal{A}_2, \mathcal{B}_2))$, $H, J, K \in \text{Obj}(\mathcal{B}_1^{\mathcal{A}_1})$, $\alpha \in \text{Hom}_{\mathcal{B}_1^{\mathcal{A}_1}}(H, J)$ and $\beta \in \text{Hom}_{\mathcal{B}_1^{\mathcal{A}_1}}(J, K)$. Then $G^F(\beta \circ_{\mathcal{B}_1^{\mathcal{A}_1}} \alpha) = G^F(\beta) \circ_{\mathcal{B}_2^{\mathcal{A}_2}} G^F(\alpha)$.

Proof. For each $A \in \text{Obj}(\mathcal{A}_2)$,

$$\begin{aligned} G^F(\beta \circ_{\mathcal{B}_1^{\mathcal{A}_1}} \alpha)_A &= G([\beta \circ_{\mathcal{B}_1^{\mathcal{A}_1}} \alpha]_{F(A)}) \\ &= G(\beta_{F(A)} \circ_{\mathcal{B}_1} \alpha_{F(A)}) \\ &= G(\beta_{F(A)}) \circ_{\mathcal{B}_2} G(\alpha_{F(A)}) \\ &= G^F(\beta)_A \circ_{\mathcal{B}_2} G^F(\alpha)_A \\ &= (G^F(\beta) \circ_{\mathcal{B}_2^{\mathcal{A}_2}} G^F(\alpha))_A. \end{aligned}$$

Hence $G^F(\beta \circ_{\mathcal{B}_1^{\mathcal{A}_1}} \alpha) = G^F(\beta) \circ_{\mathcal{B}_2^{\mathcal{A}_2}} G^F(\alpha)$. □

Exercise 8.3.0.7. Let $(\mathcal{A}_1, \mathcal{B}_1), (\mathcal{A}_2, \mathcal{B}_2) \in \text{Obj}(\mathbf{Cat}^{\text{op}} \times \mathbf{Cat})$, $(F, G) \in \text{Hom}_{\mathbf{Cat}^{\text{op}} \times \mathbf{Cat}}((\mathcal{A}_1, \mathcal{B}_1), (\mathcal{A}_2, \mathcal{B}_2))$, $H \in \text{Obj}(\mathcal{B}_1^{\mathcal{A}_1})$, Then $G^F(\text{id}_H^{\mathcal{B}_1^{\mathcal{A}_1}}) = \text{id}_{G^F(H)}^{\mathcal{B}_2^{\mathcal{A}_2}}$.

Proof. For each $A \in \mathcal{A}_2$,

$$\begin{aligned} G^F(\text{id}_H^{\mathcal{B}_1^{\mathcal{A}_1}})_A &= G((\text{id}_H^{\mathcal{B}_1^{\mathcal{A}_1}})_{F(A)}) \\ &= G(\text{id}_{H(F(A))}^{\mathcal{B}_1}) \\ &= \text{id}_{G(H(F(A)))}^{\mathcal{B}_1} \\ &= \text{id}_{G \circ_{\mathbf{Cat}} H \circ_{\mathbf{Cat}} F(A)}^{\mathcal{B}_1} \\ &= \text{id}_{G^F(H)(A)}^{\mathcal{B}_1} \\ &= (\text{id}_{G^F(H)}^{\mathcal{B}_2^{\mathcal{A}_2}})_A. \end{aligned}$$

Hence $G^F(\text{id}_H^{\mathcal{B}_1^{\mathcal{A}_1}}) = \text{id}_{G^F(H)}^{\mathcal{B}_2^{\mathcal{A}_2}}$. □

Definition 8.3.0.8. Let $(\mathcal{A}_1, \mathcal{B}_1), (\mathcal{A}_2, \mathcal{B}_2) \in \text{Obj}(\mathbf{Cat}^{\text{op}} \times \mathbf{Cat})$ and $(F, G) \in \text{Hom}_{\mathbf{Cat}^{\text{op}} \times \mathbf{Cat}}((\mathcal{A}_1, \mathcal{B}_1), (\mathcal{A}_2, \mathcal{B}_2))$. We define the **internal Hom functor of Cat at (F, G)** , denoted $[F, G]_{\mathbf{Cat}} : \mathcal{B}_1^{\mathcal{A}_1} \rightarrow \mathcal{B}_2^{\mathcal{A}_2}$ or $G^F : \mathcal{B}_1^{\mathcal{A}_1} \rightarrow \mathcal{B}_2^{\mathcal{A}_2}$, by

- $[F, G]_{\mathbf{Cat}}(H) := G^F(H)$, $H \in \text{Obj}(\mathcal{B}_1^{\mathcal{A}_1})$
- $[F, G]_{\mathbf{Cat}}(\alpha) := G^F(\alpha)$, $\alpha \in \text{Hom}_{\mathcal{B}_1^{\mathcal{A}_1}}$

Exercise 8.3.0.9. Let $(\mathcal{A}_1, \mathcal{B}_1), (\mathcal{A}_2, \mathcal{B}_2) \in \text{Obj}(\mathbf{Cat}^{\text{op}} \times \mathbf{Cat})$ and $(F, G) \in \text{Hom}_{\mathbf{Cat}^{\text{op}} \times \mathbf{Cat}}((\mathcal{A}_1, \mathcal{B}_1), (\mathcal{A}_2, \mathcal{B}_2))$. Then $G^F : \mathcal{B}_1^{\mathcal{A}_1} \rightarrow \mathcal{B}_2^{\mathcal{A}_2}$ is a functor.

Proof.

1. Exercise 8.3.0.5 implies that for each $H, J \in \text{Obj}(\mathcal{B}_1^{\mathcal{A}_1})$ and $\alpha \in \text{Hom}_{\mathcal{B}_1^{\mathcal{A}_1}}(H, J)$, $G^F(\alpha) \in \text{Hom}_{\mathcal{B}_2^{\mathcal{A}_2}}(G^F(H), G^F(J))$.
2. Exercise 8.3.0.6 implies that for each $H, J, K \in \text{Obj}(\mathcal{B}_1^{\mathcal{A}_1})$, $\alpha \in \text{Hom}_{\mathcal{B}_1^{\mathcal{A}_1}}(H, J)$ and $\beta \in \text{Hom}_{\mathcal{B}_1^{\mathcal{A}_1}}(J, K)$, $G^F(\beta \circ_{\mathcal{B}_1^{\mathcal{A}_1}} \alpha) = G^F(\beta) \circ_{\mathcal{B}_2^{\mathcal{A}_2}} G^F(\alpha)$.
3. Exercise 8.3.0.7 implies that for each $A \in \text{Obj}(\mathcal{A}_2)$, $G^F(\text{id}_H^{\mathcal{B}_1^{\mathcal{A}_1}}) = \text{id}_{G^F(H)}^{\mathcal{B}_2^{\mathcal{A}_2}}$.

Hence $G^F : \mathcal{B}_1^{\mathcal{A}_1} \rightarrow \mathcal{B}_2^{\mathcal{A}_2}$ is a functor. □

Exercise 8.3.0.10. Let $(\mathcal{A}_1, \mathcal{B}_1), (\mathcal{A}_2, \mathcal{B}_2), (\mathcal{A}_3, \mathcal{B}_3) \in \text{Obj}(\mathbf{Cat}^{\text{op}} \times \mathbf{Cat})$, $(F_{1,2}, G_{1,2}) \in \text{Hom}_{\mathbf{Cat}^{\text{op}} \times \mathbf{Cat}}((\mathcal{A}_1, \mathcal{B}_1), (\mathcal{A}_2, \mathcal{B}_2))$ and $(F_{2,3}, G_{2,3}) \in \text{Hom}_{\mathbf{Cat}^{\text{op}} \times \mathbf{Cat}}((\mathcal{A}_2, \mathcal{B}_2), (\mathcal{A}_3, \mathcal{B}_3))$. Then

$$(G_{2,3} \circ_{\mathbf{Cat}} G_{1,2})^{F_{2,3} \circ_{\mathbf{Cat}^{\text{op}}} F_{1,2}} = G_{2,3}^{F_{2,3}} \circ_{\mathbf{Cat}} G_{1,2}^{F_{1,2}}.$$

Proof.

- For each $H \in \text{Obj}(\mathcal{B}_1^{\mathcal{A}_1})$,

$$\begin{aligned}
 (G_{2,3} \circ_{\mathbf{Cat}} G_{1,2})^{F_{2,3} \circ_{\mathbf{Cat}^{\text{op}}} F_{1,2}}(H) &= G_{2,3} \circ_{\mathbf{Cat}} G_{1,2} \circ_{\mathbf{Cat}} H \circ_{\mathbf{Cat}} (F_{2,3} \circ_{\mathbf{Cat}^{\text{op}}} F_{1,2}) \\
 &= G_{2,3} \circ_{\mathbf{Cat}} G_{1,2} \circ_{\mathbf{Cat}} H \circ_{\mathbf{Cat}} (F_{1,2} \circ_{\mathbf{Cat}} F_{2,3}) \\
 &= G_{2,3} \circ_{\mathbf{Cat}} (G_{1,2} \circ_{\mathbf{Cat}} H \circ_{\mathbf{Cat}} F_{1,2}) \circ_{\mathbf{Cat}} F_{2,3} \\
 &= G_{2,3} \circ_{\mathbf{Cat}} (G_{1,2}^{F_{1,2}}(H)) \circ_{\mathbf{Cat}} F_{2,3} \\
 &= G_{2,3}^{F_{2,3}}(G_{1,2}^{F_{1,2}}(H)) \\
 &= G_{2,3}^{F_{2,3}} \circ_{\mathbf{Cat}} G_{1,2}^{F_{1,2}}(H).
 \end{aligned}$$

- Let $H, K \in \text{Obj}(\mathcal{B}_1^{\mathcal{A}_1})$ and $\alpha \in \text{Hom}_{\mathcal{B}_1^{\mathcal{A}_1}}(H, K)$. Then for each $A \in \text{Obj}(\mathcal{A}_3)$,

$$\begin{aligned}
 [(G_{2,3} \circ_{\mathbf{Cat}} G_{1,2})^{F_{2,3} \circ_{\mathbf{Cat}^{\text{op}}} F_{1,2}}(\alpha)]_A &= [(G_{2,3} \circ_{\mathbf{Cat}} G_{1,2})^{F_{1,2} \circ_{\mathbf{Cat}} F_{2,3}}(\alpha)]_A \\
 &= G_{2,3} \circ_{\mathbf{Cat}} G_{1,2}(\alpha_{F_{1,2} \circ_{\mathbf{Cat}} F_{2,3}}(A)) \\
 &= G_{2,3}(G_{1,2}(\alpha_{F_{1,2}(F_{2,3}(A))})) \\
 &= G_{2,3}(G_{1,2}^{F_{1,2}}(\alpha)_{F_{2,3}(A)}) \\
 &= [G_{2,3}^{F_{2,3}}(G_{1,2}^{F_{1,2}}(\alpha))]_A \\
 &= [G_{2,3}^{F_{2,3}} \circ_{\mathbf{Cat}} G_{1,2}^{F_{1,2}}(\alpha)]_A
 \end{aligned}$$

Thus $(G_{2,3} \circ_{\mathbf{Cat}} G_{1,2})^{F_{2,3} \circ_{\mathbf{Cat}^{\text{op}}} F_{1,2}}(\alpha) = G_{2,3}^{F_{2,3}} \circ_{\mathbf{Cat}} G_{1,2}^{F_{1,2}}(\alpha)$. Since $H, K \in \text{Obj}(\mathcal{B}_1^{\mathcal{A}_1})$ and $\alpha \in \text{Hom}_{\mathcal{B}_1^{\mathcal{A}_1}}(H, K)$ are arbitrary, we have that for each $H, K \in \text{Obj}(\mathcal{B}_1^{\mathcal{A}_1})$ and $\alpha \in \text{Hom}_{\mathcal{B}_1^{\mathcal{A}_1}}(H, K)$, $(G_{2,3} \circ_{\mathbf{Cat}} G_{1,2})^{F_{2,3} \circ_{\mathbf{Cat}^{\text{op}}} F_{1,2}}(\alpha) = G_{2,3}^{F_{2,3}} \circ_{\mathbf{Cat}} G_{1,2}^{F_{1,2}}(\alpha)$.

Therefore $(G_{2,3} \circ_{\mathbf{Cat}} G_{1,2})^{F_{2,3} \circ_{\mathbf{Cat}^{\text{op}}} F_{1,2}} = G_{2,3}^{F_{2,3}} \circ_{\mathbf{Cat}} G_{1,2}^{F_{1,2}}$. □

Exercise 8.3.0.11. Let $\mathcal{A}, \mathcal{B} \in \text{Obj}(\mathbf{Cat})$. Then

$$(\text{id}_{\mathcal{B}}^{\mathbf{Cat}})^{\text{id}_{\mathcal{A}}^{\mathbf{Cat}}} = \text{id}_{\mathcal{B}^{\mathcal{A}}}^{\mathbf{Cat}}.$$

Proof. Let $\mathcal{A}, \mathcal{B} \in \text{Obj}(\mathbf{Cat})$.

- For each $H \in \text{Obj}(\mathcal{B}^{\mathcal{A}})$,

$$\begin{aligned}
 (\text{id}_{\mathcal{B}}^{\mathbf{Cat}})^{\text{id}_{\mathcal{A}}^{\mathbf{Cat}}}(H) &= \text{id}_{\mathcal{B}}^{\mathbf{Cat}} \circ_{\mathbf{Cat}} H \circ_{\mathbf{Cat}} \text{id}_{\mathcal{A}}^{\mathbf{Cat}} \\
 &= H \\
 &= \text{id}_{\mathcal{B}^{\mathcal{A}}}^{\mathbf{Cat}}(H).
 \end{aligned}$$

- Let $H, K \in \text{Obj}(\mathcal{B}^{\mathcal{A}})$ and $\alpha \in \text{Hom}_{\mathcal{B}^{\mathcal{A}}}(H, K)$. Then for each $A \in \text{Obj}(\mathcal{A})$,

$$\begin{aligned}
 [(\text{id}_{\mathcal{B}}^{\mathbf{Cat}})^{\text{id}_{\mathcal{A}}^{\mathbf{Cat}}}(\alpha)]_A &= \text{id}_{\mathcal{B}}^{\mathbf{Cat}}(\alpha_{\text{id}_{\mathcal{A}}^{\mathbf{Cat}}(A)}) \\
 &= \text{id}_{\mathcal{B}}^{\mathbf{Cat}}(\alpha_A) \\
 &= \alpha_A \\
 &= [\text{id}_{\mathcal{B}^{\mathcal{A}}}^{\mathbf{Cat}}(\alpha)]_A
 \end{aligned}$$

Hence $(\text{id}_{\mathcal{B}}^{\mathbf{Cat}})^{\text{id}_{\mathcal{A}}^{\mathbf{Cat}}}(\alpha) = \text{id}_{\mathcal{B}^{\mathcal{A}}}^{\mathbf{Cat}}(\alpha)$. □

Definition 8.3.0.12. We define the **internal Hom functor of Cat**, denoted $[\cdot, \cdot]_{\mathbf{Cat}} : \mathbf{Cat}^{\text{op}} \times \mathbf{Cat} \rightarrow \mathbf{Cat}$, by

- $[\mathcal{C}, \mathcal{D}]_{\mathbf{Cat}} := \mathcal{D}^{\mathcal{C}}$, $(\mathcal{C}, \mathcal{D}) \in \text{Obj}(\mathbf{Cat}^{\text{op}} \times \mathbf{Cat})$,

- $[F, G]_{\mathbf{Cat}} := G^F, \quad (F, G) \in \text{Hom}_{\mathbf{Cat}^{\text{op}} \times \mathbf{Cat}}.$

Exercise 8.3.0.13. We have that $[\cdot, \cdot]_{\mathbf{Cat}} : \mathbf{Cat}^{\text{op}} \times \mathbf{Cat} \rightarrow \mathbf{Cat}$ is a functor.

Note 8.3.0.14. When the context is clear, we may write $[\cdot, \cdot]$ in place of $[\cdot, \cdot]_{\mathbf{Cat}}$.

Proof.

1. Let $(\mathcal{A}_1, \mathcal{B}_1), (\mathcal{A}_2, \mathcal{B}_2) \in \text{Obj}(\mathbf{Cat}^{\text{op}} \times \mathbf{Cat})$ $(F, G) \in \text{Hom}_{\mathbf{Cat}^{\text{op}} \times \mathbf{Cat}}((\mathcal{A}_1, \mathcal{B}_1), (\mathcal{A}_2, \mathcal{B}_2))$. Exercise 8.3.0.9 implies that

$$\begin{aligned} [\cdot, \cdot](F, G) &= G^F \\ &\in \text{Hom}_{\mathbf{Cat}}(\mathcal{B}_1^{\mathcal{A}_1}, \mathcal{B}_2^{\mathcal{A}_2}) \\ &= \text{Hom}_{\mathbf{Cat}}([\mathcal{A}_1, \mathcal{B}_1], [\mathcal{A}_2, \mathcal{B}_2]) \\ &= \text{Hom}_{\mathbf{Cat}}([\cdot, \cdot](\mathcal{A}_1, \mathcal{B}_1), [\cdot, \cdot](\mathcal{A}_2, \mathcal{B}_2)). \end{aligned}$$

2. $(\mathcal{A}_1, \mathcal{B}_1), (\mathcal{A}_2, \mathcal{B}_2), (\mathcal{A}_2, \mathcal{B}_3) \in \text{Obj}(\mathbf{Cat}^{\text{op}} \times \mathbf{Cat})$ $(F_{1,2}, G_{1,2}) \in \text{Hom}_{\mathbf{Cat}^{\text{op}} \times \mathbf{Cat}}((\mathcal{A}_1, \mathcal{B}_1), (\mathcal{A}_2, \mathcal{B}_2))$ and $(F_{2,3}, G_{2,3}) \in \text{Hom}_{\mathbf{Cat}^{\text{op}} \times \mathbf{Cat}}((\mathcal{A}_2, \mathcal{B}_2), (\mathcal{A}_2, \mathcal{B}_3))$. Exercise 8.3.0.10 then implies that

$$\begin{aligned} [\cdot, \cdot]((F_{2,3}, G_{2,3}) \circ_{\mathbf{Cat}^{\text{op}} \times \mathbf{Cat}} (F_{1,2}, G_{1,2})) &= [\cdot, \cdot](F_{2,3} \circ_{\mathbf{Cat}^{\text{op}}} F_{1,2}, G_{2,3} \circ_{\mathbf{Cat}} G_{1,2}) \\ &= (G_{2,3} \circ_{\mathbf{Cat}} G_{1,2})^{F_{2,3} \circ_{\mathbf{Cat}^{\text{op}}} F_{1,2}} \\ &= G_{2,3}^{F_{2,3}} \circ_{\mathbf{Cat}} G_{1,2}^{F_{1,2}} \\ &= [\cdot, \cdot](F_{2,3}, G_{2,3}) \circ_{\mathbf{Cat}} [\cdot, \cdot](F_{1,2}, G_{1,2}). \end{aligned}$$

3. Exercise 8.3.0.11 implies that for each $\mathcal{A}, \mathcal{B} \in \text{Obj}(\mathbf{Cat})$,

$$\begin{aligned} [\cdot, \cdot](\text{id}_{(\mathcal{A}, \mathcal{B})}^{\mathbf{Cat}^{\text{op}} \times \mathbf{Cat}}) &= [\cdot, \cdot](\text{id}_{\mathcal{A}}^{\mathbf{Cat}}, \text{id}_{\mathcal{B}}^{\mathbf{Cat}}) \\ &= (\text{id}_{\mathcal{B}}^{\mathbf{Cat}})^{\text{id}_{\mathcal{A}}^{\mathbf{Cat}}} \\ &= \text{id}_{\mathcal{B}^{\mathcal{A}}}^{\mathbf{Cat}} \\ &= \text{id}_{[\cdot, \cdot](\mathcal{A}, \mathcal{B})}^{\mathbf{Cat}}. \end{aligned}$$

Hence $[\cdot, \cdot] : \mathbf{Cat}^{\text{op}} \times \mathbf{Cat} \rightarrow \mathbf{Cat}$ is a functor. □

8.4 Opposition

relabel and organize

Definition 8.4.0.1. Let $\mathcal{C}, \mathcal{D}$ be categories and $F : \mathcal{C} \rightarrow \mathcal{D}$. We define the **opposite of F** , denoted $F^{\text{op}} : \mathcal{C}^{\text{op}} \rightarrow \mathcal{D}^{\text{op}}$, by

- $F^{\text{op}}(A) := F(A), \quad A \in \text{Obj}(\mathcal{C}^{\text{op}})$
- $F^{\text{op}}(f) := F(f), \quad f \in \text{Hom}_{\mathcal{C}^{\text{op}}}$

Exercise 8.4.0.2. Let $\mathcal{C}, \mathcal{D}$ be categories and $F : \mathcal{C} \rightarrow \mathcal{D}$. Then $F^{\text{op}} : \mathcal{C}^{\text{op}} \rightarrow \mathcal{D}^{\text{op}}$ is a functor.

Proof.

1. For each $A, B \in \text{Obj}(\mathcal{C}^{\text{op}})$ and $f \in \text{Hom}_{\mathcal{C}^{\text{op}}}(A, B)$,

$$\begin{aligned} F^{\text{op}}(f) &= f \\ &\in \text{Hom}_{\mathcal{C}^{\text{op}}}(A, B) \\ &= \text{Hom}_{\mathcal{C}^{\text{op}}}(F^{\text{op}}(A), F^{\text{op}}(B)). \end{aligned}$$

2. For each $A, B, C \in \text{Obj}(\mathcal{C}^{\text{op}})$, $f \in \text{Hom}_{\mathcal{C}^{\text{op}}}(A, B)$ and $g \in \text{Hom}_{\mathcal{C}^{\text{op}}}(B, C)$,

$$\begin{aligned} F^{\text{op}}(g \circ_{\mathcal{C}^{\text{op}}} f) &= g \circ_{\mathcal{C}^{\text{op}}} f \\ &= F^{\text{op}}(g) \circ_{\mathcal{C}^{\text{op}}} F^{\text{op}}(f). \end{aligned}$$

3. For each $A \in \text{Obj}(\mathcal{C}^{\text{op}})$,

$$\begin{aligned} F^{\text{op}}(\text{id}_A^{\mathcal{C}^{\text{op}}}) &= \text{id}_A^{\mathcal{C}^{\text{op}}} \\ &= \text{id}_{F^{\text{op}}(A)}^{\mathcal{C}^{\text{op}}}. \end{aligned}$$

Hence $F^{\text{op}} : \mathcal{C}^{\text{op}} \rightarrow \mathcal{D}^{\text{op}}$ is a functor. □

Exercise 8.4.0.3. Let $\mathcal{A}, \mathcal{B}, \mathcal{C} \in \text{Obj}(\mathbf{Cat})$, $F \in \text{Hom}_{\mathbf{Cat}}(\mathcal{A}, \mathcal{B})$ and $G \in \text{Hom}_{\mathbf{Cat}}(\mathcal{B}, \mathcal{C})$. Then $G \circ_{\mathbf{Cat}} F^{\text{op}} = G^{\text{op}} \circ_{\mathbf{Cat}} F^{\text{op}}$.

Proof.

- For each $A \in \text{Obj}(\mathcal{A}^{\text{op}})$,

$$\begin{aligned} G \circ_{\mathbf{Cat}} F^{\text{op}}(A) &= G \circ_{\mathbf{Cat}} F(A) \\ &= G(F(A)) \\ &= G^{\text{op}}(F^{\text{op}}(A)) \\ &= G^{\text{op}} \circ_{\mathbf{Cat}} F^{\text{op}}(A). \end{aligned}$$

- For each $A, B, C \in \text{Obj}(\mathcal{A}^{\text{op}})$ and $f \in \text{Hom}_{\mathcal{A}^{\text{op}}}(A, B)$,

$$\begin{aligned} G \circ_{\mathbf{Cat}} F^{\text{op}}(f) &= G \circ_{\mathbf{Cat}} F(f) \\ &= G(F(f)) \\ &= G^{\text{op}}(F^{\text{op}}(f)) \\ &= G^{\text{op}} \circ_{\mathbf{Cat}} F^{\text{op}}(f). \end{aligned}$$

Thus $G \circ_{\mathbf{Cat}} F^{\text{op}} = G^{\text{op}} \circ_{\mathbf{Cat}} F^{\text{op}}$. □

Exercise 8.4.0.4. Let $\mathcal{C} \in \text{Obj}(\mathbf{Cat})$. Then $\text{id}_{\mathcal{C}}^{\mathbf{Cat}^{\text{op}}} = \text{id}_{\mathcal{C}^{\text{op}}}^{\mathbf{Cat}}$.

Proof. • For each $A \in \text{Obj}(\mathcal{C}^{\text{op}})$,

$$\begin{aligned} \text{id}_{\mathcal{C}}^{\text{Cat}^{\text{op}}}(A) &= \text{id}_{\mathcal{C}}^{\text{Cat}}(A) \\ &= A \\ &= \text{id}_{\mathcal{C}^{\text{op}}}^{\text{Cat}}(A). \end{aligned}$$

• For each $A, B \in \text{Obj}(\mathcal{C}^{\text{op}})$ and $f \in \text{Hom}_{\mathcal{C}^{\text{op}}}(A, B)$,

$$\begin{aligned} \text{id}_{\mathcal{C}}^{\text{Cat}^{\text{op}}}(f) &= \text{id}_{\mathcal{C}}^{\text{Cat}}(f) \\ &= f \\ &= \text{id}_{\mathcal{C}^{\text{op}}}^{\text{Cat}}(f). \end{aligned}$$

Thus $\text{id}_{\mathcal{C}}^{\text{Cat}^{\text{op}}} = \text{id}_{\mathcal{C}^{\text{op}}}^{\text{Cat}}$. □

Definition 8.4.0.5. We define the **opposite functor on Cat**, denoted $\text{op}^{\text{Cat}} : \mathbf{Cat} \rightarrow \mathbf{Cat}$, by

- $\text{op}^{\text{Cat}}(\mathcal{C}) := \mathcal{C}^{\text{op}}$
- $\text{op}^{\text{Cat}}(F) := F^{\text{op}}$

Note 8.4.0.6. When the context is clear, we may write op in place of op^{Cat} .

Exercise 8.4.0.7. We have that $\text{op}^{\text{Cat}} : \mathbf{Cat} \rightarrow \mathbf{Cat}$ is a functor.

Proof.

1. Exercise 8.4.0.2 implies that for each $\mathcal{C}, \mathcal{D} \in \text{Obj}(\mathbf{Cat})$ and $F \in \text{Hom}_{\mathbf{Cat}}(\mathcal{C}, \mathcal{D})$,

$$\begin{aligned} \text{op}(F) &= F^{\text{op}} \\ &\in \text{Hom}_{\mathbf{Cat}}(\mathcal{C}^{\text{op}}, \mathcal{D}^{\text{op}}) \\ &= \text{Hom}_{\mathbf{Cat}}(\text{op}(\mathcal{C}), \text{op}(\mathcal{D})). \end{aligned}$$

2. Exercise 8.4.0.3 implies that for each $\mathcal{A}, \mathcal{B}, \mathcal{C} \in \text{Obj}(\mathbf{Cat})$, $F \in \text{Hom}_{\mathbf{Cat}}(\mathcal{A}, \mathcal{B})$ and $G \in \text{Hom}_{\mathbf{Cat}}(\mathcal{B}, \mathcal{C})$,

$$\begin{aligned} \text{op}(G \circ_{\mathbf{Cat}} F) &= G \circ_{\mathbf{Cat}} F^{\text{op}} \\ &= G^{\text{op}} \circ_{\mathbf{Cat}} F^{\text{op}} \\ &= \text{op}((G) \circ_{\mathbf{Cat}} \text{op}(F)). \end{aligned}$$

3. Exercise 8.4.0.4 implies that for each $\mathcal{C} \in \text{Obj}(\mathbf{Cat})$,

$$\begin{aligned} \text{op}(\text{id}_{\mathcal{C}}^{\text{Cat}}) &= \text{id}_{\mathcal{C}}^{\text{Cat}^{\text{op}}} &= \text{id}_{\mathcal{C}^{\text{op}}}^{\text{Cat}} \\ &= \text{id}_{\text{op}(\mathcal{C})}^{\text{Cat}}. \end{aligned}$$

Thus $\text{op} : \mathbf{Cat} \rightarrow \mathbf{Cat}$ is a functor. □

Definition 8.4.0.8. Let $\mathcal{C}, \mathcal{D}$ be categories, $F, G : \mathcal{C} \rightarrow \mathcal{D}$ functors and $\alpha : F \Rightarrow G$ a natural transformation. We define the **opposite** of α , denoted $\alpha^{\text{op}} : G^{\text{op}} \Rightarrow F^{\text{op}}$, by

$$\alpha_A^{\text{op}} := \alpha_A, \quad A \in \text{Obj}(\mathcal{C}^{\text{op}})$$

Exercise 8.4.0.9. Let $\mathcal{C}, \mathcal{D}$ be categories, $F, G : \mathcal{C} \rightarrow \mathcal{D}$ functors and $\alpha : F \Rightarrow G$ a natural transformation. Then $\alpha^{\text{op}} : G^{\text{op}} \Rightarrow F^{\text{op}}$ is a natural transformation.

Proof.

- Let $A \in \text{Obj}(\mathcal{C}^{\text{op}})$. Then $A \in \text{Obj}(\mathcal{C})$ and

$$\begin{aligned}\alpha_A^{\text{op}} &= \alpha_A \\ &\in \text{Hom}_{\mathcal{D}}(F(A), G(A)) \\ &= \text{Hom}_{\mathcal{D}^{\text{op}}}(G(A), F(A)) \\ &= \text{Hom}_{\mathcal{D}^{\text{op}}}(G^{\text{op}}(A), F^{\text{op}}(A)).\end{aligned}$$

Since $A \in \text{Obj}(\mathcal{C}^{\text{op}})$ is arbitrary, we have that for each $A \in \text{Obj}(\mathcal{C}^{\text{op}})$, $\alpha_A^{\text{op}} \in \text{Hom}_{\mathcal{D}^{\text{op}}}(G^{\text{op}}(A), F^{\text{op}}(A))$.

- Let $A, B \in \text{Obj}(\mathcal{C}^{\text{op}})$ and $f \in \text{Hom}_{\mathcal{C}^{\text{op}}}(A, B)$. Then $A, B \in \text{Obj}(\mathcal{C})$ and $f \in \text{Hom}_{\mathcal{C}}(B, A)$. Since $\alpha : F \Rightarrow G$ is a natural transformation, $G(f) \circ_{\mathcal{D}} \alpha_B = \alpha_A \circ_{\mathcal{D}} F(f)$, i.e. the following diagram commutes:

$$\begin{array}{ccc} F(B) & \xrightarrow{\alpha_B} & G(B) \\ F(f) \downarrow & & \downarrow G(f) \\ F(A) & \xrightarrow{\alpha_A} & G(A). \end{array}$$

Therefore

$$\begin{aligned}F^{\text{op}}(f) \circ_{\mathcal{D}^{\text{op}}} \alpha_A^{\text{op}} &= F(f) \circ_{\mathcal{D}^{\text{op}}} \alpha_A \\ &= \alpha_A \circ_{\mathcal{D}} F(f) \\ &= G(f) \circ_{\mathcal{D}} \alpha_B \\ &= \alpha_B \circ_{\mathcal{D}^{\text{op}}} G^{\text{op}}(f) \\ &= \alpha_B^{\text{op}} \circ_{\mathcal{D}^{\text{op}}} G^{\text{op}}(f),\end{aligned}$$

i.e. the following diagram commutes:

$$\begin{array}{ccc} G^{\text{op}}(A) & \xrightarrow{\alpha_A^{\text{op}}} & F^{\text{op}}(A) \\ G^{\text{op}}(f) \downarrow & & \downarrow F^{\text{op}}(f) \\ G^{\text{op}}(B) & \xrightarrow{\alpha_B^{\text{op}}} & F^{\text{op}}(B). \end{array}$$

Since $A, B \in \text{Obj}(\mathcal{C}^{\text{op}})$ and $f \in \text{Hom}_{\mathcal{C}^{\text{op}}}(A, B)$ are arbitrary, we have that for each $A, B \in \text{Obj}(\mathcal{C}^{\text{op}})$ and $f \in \text{Hom}_{\mathcal{C}^{\text{op}}}(A, B)$, $F^{\text{op}}(f) \circ_{\mathcal{D}^{\text{op}}} \alpha_A^{\text{op}} = \alpha_B^{\text{op}} \circ_{\mathcal{D}^{\text{op}}} G^{\text{op}}(f)$.

Thus $\alpha^{\text{op}} : G^{\text{op}} \Rightarrow F^{\text{op}}$ is a natural transformation. □

Exercise 8.4.0.10. Let $\mathcal{C}, \mathcal{D} \in \text{Obj}(\mathbf{Cat})$, $F, G, H \in \text{Obj}(\mathcal{D}^{\mathcal{C}})$, $\alpha \in \text{Hom}_{\mathcal{D}^{\mathcal{C}}}(F, G)$ and $\beta \in \text{Hom}_{\mathcal{D}^{\mathcal{C}}}(G, H)$. Then $(\beta \circ_{\mathcal{D}^{\mathcal{C}}} \alpha)^{\text{op}} = \alpha^{\text{op}} \circ_{(\mathcal{D}^{\text{op}})^{\mathcal{C}^{\text{op}}}} \beta^{\text{op}}$.

Proof. For each $A \in \text{Obj}(\mathcal{C}^{\text{op}})$,

$$\begin{aligned}(\beta \circ_{\mathcal{D}^{\mathcal{C}}} \alpha)_A^{\text{op}} &= (\beta \circ_{\mathcal{D}^{\mathcal{C}}} \alpha)_A \\ &= \beta_A \circ_{\mathcal{D}} \alpha_A \\ &= \alpha_A \circ_{\mathcal{D}^{\text{op}}} \beta_A \\ &= \alpha_A^{\text{op}} \circ_{\mathcal{D}^{\text{op}}} \beta_A^{\text{op}} \\ &= (\alpha^{\text{op}} \circ_{(\mathcal{D}^{\text{op}})^{\mathcal{C}^{\text{op}}}} \beta^{\text{op}})_A.\end{aligned}$$

Thus $(\beta \circ_{\mathcal{D}^{\mathcal{C}}} \alpha)^{\text{op}} = \alpha^{\text{op}} \circ_{(\mathcal{D}^{\text{op}})^{\mathcal{C}^{\text{op}}}} \beta^{\text{op}}$. □

Exercise 8.4.0.11. Let $\mathcal{C}, \mathcal{D} \in \text{Obj}(\mathbf{Cat})$ and $F \in \text{Obj}(\mathcal{D}^{\mathcal{C}})$. Then $(\text{id}_F^{\mathcal{D}^{\mathcal{C}}})^{\text{op}} = \text{id}_{F^{\text{op}}}^{(\mathcal{D}^{\text{op}})^{\mathcal{C}^{\text{op}}}}$.

Proof. For each $A \in \text{Obj}(\mathcal{C}^{\text{op}})$,

$$\begin{aligned} (\text{id}_F^{\mathcal{D}^{\text{c op}}})_A &= (\text{id}_F^{\mathcal{D}^{\text{c}}})_A \\ &= \text{id}_{F(A)}^{\mathcal{D}} \\ &= \text{id}_{F(A)}^{\mathcal{D}^{\text{op}}} \\ &= \text{id}_{F^{\text{op}}(A)}^{\mathcal{D}^{\text{op}}} \\ &= (\text{id}_{F^{\text{op}}}^{(\mathcal{D}^{\text{op}})^{\mathcal{C}^{\text{op}}}})_A. \end{aligned}$$

Thus $(\text{id}_F^{\mathcal{D}^{\text{c}}})^{\text{op}} = \text{id}_{F^{\text{op}}}^{(\mathcal{D}^{\text{op}})^{\mathcal{C}^{\text{op}}}}$. □

Definition 8.4.0.12. Let $\mathcal{C}, \mathcal{D} \in \text{Obj}(\mathbf{Cat})$. We define the **opposite functor on $\mathcal{D}^{\text{c}}$** , denoted $\text{op}_{(\mathcal{C}, \mathcal{D})} : \mathcal{D}^{\text{c}} \rightarrow [(\mathcal{D}^{\text{op}})^{\mathcal{C}^{\text{op}}}]^{\text{op}}$, by

- $\text{op}_{(\mathcal{C}, \mathcal{D})}(F) := F^{\text{op}}$
- $\text{op}_{(\mathcal{C}, \mathcal{D})}(\alpha) := \alpha^{\text{op}}$

Exercise 8.4.0.13. Let $\mathcal{C}, \mathcal{D} \in \text{Obj}(\mathbf{Cat})$. Then $\text{op}_{(\mathcal{C}, \mathcal{D})} : \mathcal{D}^{\text{c}} \rightarrow [(\mathcal{D}^{\text{op}})^{\mathcal{C}^{\text{op}}}]^{\text{op}}$ is a functor.

Proof.

1. Let $F, G \in \text{Obj}(\mathcal{D}^{\text{c}})$ and $\alpha \in \text{Hom}_{\mathcal{D}^{\text{c}}}(F, G)$. Exercise 8.4.0.2 implies that

$$\begin{aligned} \text{op}_{(\mathcal{C}, \mathcal{D})}(F), \text{op}_{(\mathcal{C}, \mathcal{D})}(G) &\in \text{Obj}((\mathcal{D}^{\text{op}})^{\mathcal{C}^{\text{op}}}) \\ &= \text{Obj}([(\mathcal{D}^{\text{op}})^{\mathcal{C}^{\text{op}}}]^{\text{op}}) \end{aligned}$$

and Exercise 8.4.0.9 implies that

$$\begin{aligned} \text{op}_{(\mathcal{C}, \mathcal{D})}(\alpha) &\in \text{Hom}_{(\mathcal{D}^{\text{op}})^{\mathcal{C}^{\text{op}}}}(G^{\text{op}}, F^{\text{op}}) \\ &= \text{Hom}_{[(\mathcal{D}^{\text{op}})^{\mathcal{C}^{\text{op}}}]^{\text{op}}}(F^{\text{op}}, G^{\text{op}}) \\ &= \text{Hom}_{[(\mathcal{D}^{\text{op}})^{\mathcal{C}^{\text{op}}}]^{\text{op}}}(\text{op}_{(\mathcal{C}, \mathcal{D})}(F), \text{op}_{(\mathcal{C}, \mathcal{D})}(G)). \end{aligned}$$

Since $F, G \in \text{Obj}(\mathcal{D}^{\text{c}})$ and $\alpha \in \text{Hom}_{\mathcal{D}^{\text{c}}}(F, G)$ are arbitrary, we have that for each $F, G \in \text{Obj}(\mathcal{D}^{\text{c}})$ and $\alpha \in \text{Hom}_{\mathcal{D}^{\text{c}}}(F, G)$, $\text{op}_{(\mathcal{C}, \mathcal{D})}(\alpha) \in \text{Hom}_{[(\mathcal{D}^{\text{op}})^{\mathcal{C}^{\text{op}}}]^{\text{op}}}(\text{op}_{(\mathcal{C}, \mathcal{D})}(F), \text{op}_{(\mathcal{C}, \mathcal{D})}(G))$.

2. Let $F, G, H \in \text{Obj}(\mathcal{D}^{\text{c}})$, $\alpha \in \text{Hom}_{\mathcal{D}^{\text{c}}}(F, G)$ and $\beta \in \text{Hom}_{\mathcal{D}^{\text{c}}}(G, H)$. Exercise 8.4.0.10 implies that

$$\begin{aligned} \text{op}_{(\mathcal{C}, \mathcal{D})}(\beta \circ_{\mathcal{D}^{\text{c}}} \alpha) &= \beta \circ_{\mathcal{D}^{\text{c}}} \alpha^{\text{op}} \\ &= \alpha^{\text{op}} \circ_{(\mathcal{D}^{\text{op}})^{\mathcal{C}^{\text{op}}}} \beta^{\text{op}} \\ &= \beta^{\text{op}} \circ_{[(\mathcal{D}^{\text{op}})^{\mathcal{C}^{\text{op}}}]^{\text{op}}} \alpha^{\text{op}} \\ &= \text{op}_{(\mathcal{C}, \mathcal{D})}(\beta) \circ_{[(\mathcal{D}^{\text{op}})^{\mathcal{C}^{\text{op}}}]^{\text{op}}} \text{op}_{(\mathcal{C}, \mathcal{D})}(\alpha). \end{aligned}$$

3. Exercise 8.4.0.11 implies that for each $F \in \text{Obj}(\mathcal{D}^{\text{c}})$,

$$\begin{aligned} \text{op}_{(\mathcal{C}, \mathcal{D})}(\text{id}_F^{\mathcal{D}^{\text{c}}}) &= (\text{id}_F^{\mathcal{D}^{\text{c}}})^{\text{op}} \\ &= \text{id}_{F^{\text{op}}}^{(\mathcal{D}^{\text{op}})^{\mathcal{C}^{\text{op}}}} \\ &= \text{id}_{\text{op}_{(\mathcal{C}, \mathcal{D})}(F)}^{(\mathcal{D}^{\text{op}})^{\mathcal{C}^{\text{op}}}}. \end{aligned}$$

Thus $\text{op}_{(\mathcal{C}, \mathcal{D})} : \mathcal{D}^{\text{c}} \rightarrow [(\mathcal{D}^{\text{op}})^{\mathcal{C}^{\text{op}}}]^{\text{op}}$ is a functor. □

Definition 8.4.0.14. We define $\text{op} : [\cdot, \cdot]_{\mathbf{Cat}} \Rightarrow \text{op}^{\mathbf{Cat}} \circ_{\mathbf{Cat}} [\cdot, \cdot] \circ (\text{op}^{\mathbf{Cat}}, \text{op}^{\mathbf{Cat}})$ by $\text{op}_{(\mathcal{C}, \mathcal{D})} := \text{op}_{(\mathcal{C}, \mathcal{D})}$ **rename stuff and reorganize the opposite stuff**

Exercise 8.4.0.15. $\emptyset$

8.5 Currying

Definition 8.5.0.1. Let $\mathcal{A}, \mathcal{B}, \mathcal{C}$ be categories, $F : \mathcal{A} \times \mathcal{B} \rightarrow \mathcal{C}$, $A_1, A_2 \in \text{Obj}(\mathcal{A})$, $B_1, B_2 \in \text{Obj}(\mathcal{B})$, $f \in \text{Hom}_{\mathcal{A}}(A_1, A_2)$ and $g \in \text{Hom}_{\mathcal{B}}(B_1, B_2)$. We define the

- **natural transformation from $F(A_1, \cdot)$ to $F(A_2, \cdot)$ induced by f** , denoted $\alpha(F)^f : F(A_1, \cdot) \Rightarrow F(A_2, \cdot)$, by $\alpha(F)_B^f := F(f, \text{id}_B^{\mathcal{B}})$,
- **natural transformation from $F(\cdot, B_1)$ to $F(\cdot, B_2)$ induced by g** , denoted $\beta(F)^g : F(\cdot, B_1) \Rightarrow F(\cdot, B_2)$, by $\beta(F)_A^g := F(\text{id}_A^{\mathcal{A}}, g)$,

Exercise 8.5.0.2. Let $\mathcal{A}, \mathcal{B}, \mathcal{C}$ be categories and $F : \mathcal{A} \times \mathcal{B} \rightarrow \mathcal{C}$. Then

1. for each $A_1, A_2 \in \text{Obj}(\mathcal{A})$ and $f \in \text{Hom}_{\mathcal{A}}(A_1, A_2)$, $\alpha(F)^f : F(A_1, \cdot) \Rightarrow F(A_2, \cdot)$ is a natural transformation,
2. for each $B_1, B_2 \in \text{Obj}(\mathcal{B})$ and $g \in \text{Hom}_{\mathcal{B}}(B_1, B_2)$, $\beta(F)^g : F(\cdot, B_1) \Rightarrow F(\cdot, B_2)$ is a natural transformation.

Proof.

1. Let $A_1, A_2 \in \text{Obj}(\mathcal{A})$.

- Let $B \in \text{Obj}(\mathcal{B})$. Then $(f, \text{id}_B^{\mathcal{B}}) \in \text{Hom}_{\mathcal{A} \times \mathcal{B}}((A_1, B), (A_2, B))$. Therefore

$$\begin{aligned} \alpha(F)_B^f &= F(f, \text{id}_B^{\mathcal{B}}) \\ &\in \text{Hom}_{\mathcal{A} \times \mathcal{B}}(F(A_1, B), F(A_2, B)) \\ &= \text{Hom}_{\mathcal{A} \times \mathcal{B}}(F(A_1, \cdot)(B), F(A_2, \cdot)(B)). \end{aligned}$$

Since $B \in \text{Obj}(\mathcal{B})$ is arbitrary, we have that for each $B \in \text{Obj}(\mathcal{B})$, $\alpha(F)_B^f \in \text{Hom}_{\mathcal{A} \times \mathcal{B}}(F(A_1, \cdot)(B), F(A_2, \cdot)(B))$.

- Let $B_1, B_2 \in \text{Obj}(\mathcal{B})$, $f \in \text{Hom}_{\mathcal{A}}(A_1, A_2)$ and $g \in \text{Hom}_{\mathcal{B}}(B_1, B_2)$. Then

$$\begin{aligned} F(A_2, \cdot)(g) \circ_{\mathcal{C}} \alpha(F)_{B_1}^f &= F(\text{id}_{A_2}^{\mathcal{A}}, g) \circ_{\mathcal{C}} F(f, \text{id}_{B_1}^{\mathcal{B}}) \\ &= F((\text{id}_{A_2}^{\mathcal{A}}, g) \circ_{\mathcal{A} \times \mathcal{B}} (f, \text{id}_{B_1}^{\mathcal{B}})) \\ &= F(\text{id}_{A_2}^{\mathcal{A}} \circ_{\mathcal{A}} f, g \circ_{\mathcal{B}} \text{id}_{B_1}^{\mathcal{B}}) \\ &= F(f, g) \\ &= F(f \circ_{\mathcal{A}} \text{id}_{A_1}^{\mathcal{A}}, \text{id}_{B_2}^{\mathcal{B}} \circ_{\mathcal{B}} g) \\ &= F((f, \text{id}_{B_2}^{\mathcal{B}}) \circ_{\mathcal{A} \times \mathcal{B}} (\text{id}_{A_1}^{\mathcal{A}}, g)) \\ &= F(f, \text{id}_{B_2}^{\mathcal{B}}) \circ_{\mathcal{C}} F(\text{id}_{A_1}^{\mathcal{A}}, g) \\ &= \alpha(F)_{B_2}^f \circ_{\mathcal{C}} F(A_1, \cdot)(g), \end{aligned}$$

i.e. the following diagram commutes:

$$\begin{array}{ccc} F(A_1, \cdot)(B_1) & \xrightarrow{\alpha(F)_{B_1}^f} & F(A_2, \cdot)(B_1) \\ \downarrow F(A_1, \cdot)(g) & & \downarrow F(A_2, \cdot)(g) \\ F(A_1, \cdot)(B_2) & \xrightarrow{\alpha(F)_{B_2}^f} & F(A_2, \cdot)(B_2). \end{array}$$

Since $B_1, B_2 \in \text{Obj}(\mathcal{B})$ and $g \in \text{Hom}_{\mathcal{B}}(B_1, B_2)$ are arbitrary, we have that for each $B_1, B_2 \in \text{Obj}(\mathcal{B})$ and $g \in \text{Hom}_{\mathcal{B}}(B_1, B_2)$, $F(A_2, \cdot)(g) \circ_{\mathcal{C}} \alpha(F)_{B_1}^f = \alpha(F)_{B_2}^f \circ_{\mathcal{C}} F(A_1, \cdot)(g)$.

Thus $\alpha(F)^f : F(A_1, \cdot) \Rightarrow F(A_2, \cdot)$ is a natural transformation. Since $A_1, A_2 \in \text{Obj}(\mathcal{A})$ and $f \in \text{Hom}_{\mathcal{A}}(A_1, A_2)$ are arbitrary, we have that for each $A_1, A_2 \in \text{Obj}(\mathcal{A})$ and $f \in \text{Hom}_{\mathcal{A}}(A_1, A_2)$, $\alpha(F)^f : F(A_1, \cdot) \Rightarrow F(A_2, \cdot)$ is a natural transformation.

2. Similar to **prev part (ref part)**.

□

Exercise 8.5.0.3. Let $\mathcal{A}, \mathcal{B}, \mathcal{C}$ be categories and $F : \mathcal{A} \times \mathcal{B} \rightarrow \mathcal{C}$. Then

1. for each $A_1, A_2, A_3 \in \text{Obj}(\mathcal{A})$, $f_{1,2} \in \text{Hom}_{\mathcal{A}}(A_1, A_2)$ and $f_{2,3} \in \text{Hom}_{\mathcal{A}}(A_2, A_3)$, $\alpha(F)^{f_{2,3} \circ_{\mathcal{A}} f_{1,2}} = \alpha(F)^{f_{2,3}} \circ_{\mathcal{C}^{\mathcal{B}}} \alpha(F)^{f_{1,2}}$
2. for each $B_1, B_2, B_3 \in \text{Obj}(\mathcal{B})$, $g_{1,2} \in \text{Hom}_{\mathcal{B}}(B_1, B_2)$ and $g_{2,3} \in \text{Hom}_{\mathcal{B}}(B_2, B_3)$, $\beta(F)^{g_{2,3} \circ_{\mathcal{B}} g_{1,2}} = \beta(F)^{g_{2,3}} \circ_{\mathcal{C}^{\mathcal{A}}} \beta(F)^{g_{1,2}}$

Proof.

1. Let $A_1, A_2, A_3 \in \text{Obj}(\mathcal{A})$, $f_{1,2} \in \text{Hom}_{\mathcal{A}}(A_1, A_2)$ and $f_{2,3} \in \text{Hom}_{\mathcal{A}}(A_2, A_3)$. Then for each $B \in \text{Obj}(\mathcal{B})$,

$$\begin{aligned}
 (\alpha(F)^{f_{2,3} \circ_{\mathcal{A}} f_{1,2}})_B &= F(f_{2,3} \circ_{\mathcal{A}} f_{1,2}, \text{id}_B^{\mathcal{B}}) \quad (\text{by Definition 8.5.0.1}) \\
 &= F(f_{2,3} \circ_{\mathcal{A}} f_{1,2}, \text{id}_B^{\mathcal{B}} \circ_{\mathcal{B}} \text{id}_B^{\mathcal{B}}) \\
 &= F((f_{2,3}, \text{id}_B^{\mathcal{B}}) \circ_{\mathcal{A} \times \mathcal{B}} (f_{1,2}, \text{id}_B^{\mathcal{B}})) \\
 &= F(f_{2,3}, \text{id}_B^{\mathcal{B}}) \circ_{\mathcal{C}} F(f_{1,2}, \text{id}_B^{\mathcal{B}}) \\
 &= \alpha(F)_B^{f_{2,3}} \circ_{\mathcal{C}} \alpha(F)_B^{f_{1,2}} \\
 &= (\alpha(F)^{f_{2,3}} \circ_{\mathcal{C}^{\mathcal{B}}} \alpha(F)^{f_{1,2}})_B.
 \end{aligned}$$

Thus $\alpha(F)^{f_{2,3} \circ_{\mathcal{A}} f_{1,2}} = \alpha(F)^{f_{2,3}} \circ_{\mathcal{C}^{\mathcal{B}}} \alpha(F)^{f_{1,2}}$.

2. similar to prev part (ref prev part)

□

Exercise 8.5.0.4. Let $\mathcal{A}, \mathcal{B}, \mathcal{C}$ be categories and $F : \mathcal{A} \times \mathcal{B} \rightarrow \mathcal{C}$. Then

1. For each $A \in \text{Obj}(\mathcal{A})$, $\alpha(F)^{\text{id}_A^{\mathcal{A}}} = \text{id}_{F(A, \cdot)}^{\mathcal{C}^{\mathcal{B}}}$.
2. For each $B \in \text{Obj}(\mathcal{B})$, $\beta(F)^{\text{id}_B^{\mathcal{B}}} = \text{id}_{F(\cdot, B)}^{\mathcal{C}^{\mathcal{A}}}$.

Proof.

1. For each $B \in \text{Obj}(\mathcal{B})$,

$$\begin{aligned}
 \alpha(F)_B^{\text{id}_A^{\mathcal{A}}} &= F(\text{id}_A^{\mathcal{A}}, \text{id}_B^{\mathcal{B}}) \\
 &= F(\text{id}_{A \times B}^{\mathcal{A} \times \mathcal{B}}) \\
 &= \text{id}_{F(A, B)}^{\mathcal{C}} \\
 &= \text{id}_{F(A, \cdot)(B)}^{\mathcal{C}} \\
 &= (\text{id}_{F(A, \cdot)})_B \quad (\text{by Definition 5.2.0.1}).
 \end{aligned}$$

Since $B \in \text{Obj}(\mathcal{B})$ is arbitrary, we have that for each $B \in \text{Obj}(\mathcal{B})$, $\alpha(F)_B^{\text{id}_A^{\mathcal{A}}} = (\text{id}_{F(A, \cdot)})_B$. Thus $\alpha(F)^{\text{id}_A^{\mathcal{A}}} = \text{id}_{F(A, \cdot)}$.

2. Similar to prev part (ref prev part).

□

Definition 8.5.0.5. Let $\mathcal{A}, \mathcal{B}, \mathcal{C} \in \text{Obj}(\mathbf{Cat})$ and $F \in \text{Obj}(\mathcal{C}^{\mathcal{A} \times \mathcal{B}})$. We define the **Curry functor induced by F** , denoted $\text{cur}(F) : \mathcal{A} \rightarrow \mathcal{C}^{\mathcal{B}}$ by

- $\text{cur}(F)(A) := F(A, \cdot)$, ($A \in \text{Obj}(\mathcal{A})$),
- $\text{cur}(F)(f) := \alpha(F)^f$, ($f \in \text{Hom}_{\mathcal{A}}$),

Exercise 8.5.0.6. Let $\mathcal{A}, \mathcal{B}, \mathcal{C} \in \text{Obj}(\mathbf{Cat})$ and $F \in \text{Obj}(\mathcal{C}^{\mathcal{A} \times \mathcal{B}})$. Then $\text{cur}(F) : \mathcal{A} \rightarrow \mathcal{C}^{\mathcal{B}}$ is a functor.

Proof.

1. Let $A_1, A_2 \in \text{Obj}(\mathcal{A})$ and $f \in \text{Hom}_{\mathcal{A}}(A_1, A_2)$. Exercise 8.5.0.2 implies that

$$\begin{aligned} \text{cur}(F)(f) &= \alpha(F)^f \\ &\in \text{Hom}_{\mathcal{C}^{\mathcal{B}}}(F(A_1, \cdot), F(A_2, \cdot)) \\ &= \text{Hom}_{\mathcal{C}^{\mathcal{B}}}(\text{cur}(F)(A_1), \text{cur}(F)(A_2)). \end{aligned}$$

2. Let $A_1, A_2, A_3 \in \text{Obj}(\mathcal{A})$, $f_{1,2} \in \text{Hom}_{\mathcal{A}}(A_1, A_2)$ and $f_{2,3} \in \text{Hom}_{\mathcal{A}}(A_2, A_3)$. Exercise 8.5.0.3 implies that

$$\begin{aligned} \text{cur}(F)(f_{2,3} \circ_{\mathcal{A}} f_{1,2}) &= \alpha(F)^{f_{2,3} \circ_{\mathcal{A}} f_{1,2}} \\ &= \alpha(F)^{f_{2,3}} \circ_{\mathcal{C}^{\mathcal{B}}} \alpha(F)^{f_{1,2}} \\ &= \text{cur}(F)(f_{2,3}) \circ_{\mathcal{C}^{\mathcal{B}}} \text{cur}(F)(f_{1,2}). \end{aligned}$$

3. Let $A \in \text{Obj}(\mathcal{A})$. Exercise 8.5.0.4 implies that,

$$\begin{aligned} \text{cur}(F)(\text{id}_A^{\mathcal{A}}) &= \alpha(F)^{\text{id}_A^{\mathcal{A}}} \\ &= \text{id}_{F(A, \cdot)}^{\mathcal{C}^{\mathcal{B}}} \\ &= \text{id}_{\text{cur}(F)(A)}^{\mathcal{C}^{\mathcal{B}}}. \end{aligned}$$

Hence $\text{cur}(F) : \mathcal{A} \rightarrow \mathcal{C}^{\mathcal{B}}$ is a functor. □

Definition 8.5.0.7. Let $\mathcal{A}, \mathcal{B}, \mathcal{C} \in \text{Obj}(\mathbf{Cat})$ and $F, G \in \text{Obj}(\mathcal{C}^{\mathcal{A} \times \mathcal{B}})$ and $\alpha \in \text{Hom}_{\mathcal{C}^{\mathcal{A} \times \mathcal{B}}}(F, G)$. We define the **Curry transformation induced by α** , denoted $\text{cur}(\alpha) : \text{cur}(F) \Rightarrow \text{cur}(G)$ by

$$\text{cur}(\alpha)_A := \alpha_{(A, \cdot)}, \quad A \in \text{Obj}(\mathcal{A}).$$

Exercise 8.5.0.8. Let $\mathcal{A}, \mathcal{B}, \mathcal{C} \in \text{Obj}(\mathbf{Cat})$, $F, G \in \text{Obj}(\mathcal{C}^{\mathcal{A} \times \mathcal{B}})$ and $\alpha \in \text{Hom}_{\mathcal{C}^{\mathcal{A} \times \mathcal{B}}}(F, G)$. Then $\text{cur}(\alpha) : \text{cur}(F) \Rightarrow \text{cur}(G)$ is a natural transformation.

Proof.

- Exercise 8.2.2.6 implies that for each $A \in \text{Obj}(\mathcal{A})$,

$$\begin{aligned} \text{cur}(\alpha)_A &= \alpha_{(A, \cdot)} \\ &\in \text{Hom}_{\mathcal{C}^{\mathcal{B}}}(F(A, \cdot), G(A, \cdot)) \\ &= \text{Hom}_{\mathcal{C}^{\mathcal{B}}}(\text{cur}(F)(A), \text{cur}(G)(A)). \end{aligned}$$

- Let $A_1, A_2 \in \text{Obj}(\mathcal{A})$ and $f \in \text{Hom}_{\mathcal{A}}(A_1, A_2)$. Since $\alpha \in \text{Hom}_{\mathcal{C}^{\mathcal{A} \times \mathcal{B}}}(F, G)$, we have that

$$G(f, \text{id}_B^{\mathcal{B}}) \circ_{\mathcal{C}} \alpha_{(A_1, B)} = \alpha_{(A_2, B)} \circ_{\mathcal{C}} F(f, \text{id}_B^{\mathcal{B}}),$$

i.e. the following diagram commutes:

$$\begin{array}{ccc} F(A_1, B) & \xrightarrow{\alpha_{(A_1, B)}} & G(A_1, B) \\ F(f, \text{id}_B^{\mathcal{B}}) \downarrow & & \downarrow G(f, \text{id}_B^{\mathcal{B}}) \\ F(A_2, B) & \xrightarrow{\alpha_{(A_2, B)}} & G(A_2, B). \end{array}$$

Therefore for each $B \in \text{Obj}(\mathcal{B})$,

$$\begin{aligned} (\text{cur}(G)(f) \circ_{\mathcal{C}^{\mathcal{B}}} \text{cur}(\alpha)_{A_1})_B &= \text{cur}(G)(f)_B \circ_{\mathcal{C}} (\text{cur}(\alpha)_{A_1})_B \\ &= G(f, \text{id}_B^{\mathcal{B}}) \circ_{\mathcal{C}} \alpha_{(A_1, B)} \\ &= \alpha_{(A_2, B)} \circ_{\mathcal{C}} F(f, \text{id}_B^{\mathcal{B}}) \\ &= \text{cur}(\alpha)_{A_1}_B \circ_{\mathcal{C}} \text{cur}(F)(f)_B \\ &= (\text{cur}(\alpha)_{A_1} \circ_{\mathcal{C}} \text{cur}(F)(f))_B. \end{aligned}$$

Hence $\text{cur}(G)(f) \circ_{\mathcal{C}^{\mathcal{B}}} \text{cur}(\alpha)_{A_1} = \text{cur}(\alpha)_{A_2} \circ_{\mathcal{C}} \text{cur}(F)(f)$, i.e. the following diagram commutes:

$$\begin{array}{ccc} \text{cur}(F)(A_1) & \xrightarrow{\text{cur}(\alpha)_{A_1}} & \text{cur}(G)(A_1) \\ \text{cur}(F)(f) \downarrow & & \downarrow \text{cur}(G)(f) \\ \text{cur}(F)(A_2) & \xrightarrow{\text{cur}(\alpha)_{A_2}} & \text{cur}(G)(A_2). \end{array}$$

Since $A_1, A_2 \in \text{Obj}(\mathcal{A})$ and $f \in \text{Hom}_{\mathcal{A}}(A_1, A_2)$ are arbitrary, we have that for each $A_1, A_2 \in \text{Obj}(\mathcal{A})$ and $f \in \text{Hom}_{\mathcal{A}}(A_1, A_2)$, $\text{cur}(G)(f) \circ_{\mathcal{C}^{\mathcal{B}}} \text{cur}(\alpha)_{A_1} = \text{cur}(\alpha)_{A_2} \circ_{\mathcal{C}} \text{cur}(F)(f)$.

Thus $\text{cur}(\alpha) : \text{cur}(F) \Rightarrow \text{cur}(G)$ is a natural transformation. □

Exercise 8.5.0.9. Let $\mathcal{A}, \mathcal{B}, \mathcal{C} \in \text{Obj}(\mathbf{Cat})$, $F_1, F_2, F_3 \in \text{Obj}(\mathcal{C}^{\mathcal{A} \times \mathcal{B}})$, $\alpha_{1,2} \in \text{Hom}_{\mathcal{C}^{\mathcal{A} \times \mathcal{B}}}(F_1, F_2)$ and $\alpha_{2,3} \in \text{Hom}_{\mathcal{C}^{\mathcal{A} \times \mathcal{B}}}(F_2, F_3)$. Then $\text{cur}(\alpha_{2,3} \circ_{\mathcal{C}^{\mathcal{A} \times \mathcal{B}}} \alpha_{1,2}) = \text{cur}(\alpha_{2,3}) \circ_{(\mathcal{C}^{\mathcal{B}})^{\mathcal{A}}} \text{cur}(\alpha_{1,2})$.

Proof. Exercise 8.2.2.7 implies that for each $A \in \text{Obj}(\mathcal{A})$,

$$\begin{aligned} \text{cur}(\alpha_{2,3} \circ_{\mathcal{C}^{\mathcal{A} \times \mathcal{B}}} \alpha_{1,2})_A &= (\alpha_{2,3} \circ_{\mathcal{C}^{\mathcal{A} \times \mathcal{B}}} \alpha_{1,2})_{(A, \cdot)} \\ &= (\alpha_{2,3})_{(A, \cdot)} \circ_{\mathcal{C}^{\mathcal{B}}} (\alpha_{1,2})_{(A, \cdot)} \\ &= \text{cur}(\alpha_{2,3})_A \circ_{\mathcal{C}^{\mathcal{B}}} \text{cur}(\alpha_{1,2})_A \\ &= [\text{cur}(\alpha_{2,3}) \circ_{(\mathcal{C}^{\mathcal{B}})^{\mathcal{A}}} \text{cur}(\alpha_{1,2})]_A. \end{aligned}$$

Therefore $\text{cur}(\alpha_{2,3} \circ_{\mathcal{C}^{\mathcal{A} \times \mathcal{B}}} \alpha_{1,2}) = \text{cur}(\alpha_{2,3}) \circ_{(\mathcal{C}^{\mathcal{B}})^{\mathcal{A}}} \text{cur}(\alpha_{1,2})$. □

Exercise 8.5.0.10. Let $\mathcal{A}, \mathcal{B}, \mathcal{C} \in \text{Obj}(\mathbf{Cat})$ and $F \in \text{Obj}(\mathcal{C}^{\mathcal{A} \times \mathcal{B}})$. Then $\text{cur}(\text{id}_F^{\mathcal{C}^{\mathcal{A} \times \mathcal{B}}}) = \text{id}_{\text{cur}(F)^{(\mathcal{C}^{\mathcal{B}})^{\mathcal{A}}}}$.

Proof. Exercise 8.2.2.8 implies that for each $A \in \text{Obj}(\mathcal{A})$,

$$\begin{aligned} \text{cur}(\text{id}_F^{\mathcal{C}^{\mathcal{A} \times \mathcal{B}}})_A &= (\text{id}_F^{\mathcal{C}^{\mathcal{A} \times \mathcal{B}}})_{(A, \cdot)} \\ &= \text{id}_{F(A, \cdot)}^{\mathcal{C}^{\mathcal{B}}} \\ &= \text{id}_{\text{cur}(F)(A)}^{\mathcal{C}^{\mathcal{B}}} \\ &= (\text{id}_{\text{cur}(F)}^{(\mathcal{C}^{\mathcal{B}})^{\mathcal{A}}})_A. \end{aligned}$$

Hence $\text{cur}(\text{id}_F^{\mathcal{C}^{\mathcal{A} \times \mathcal{B}}}) = \text{id}_{\text{cur}(F)^{(\mathcal{C}^{\mathcal{B}})^{\mathcal{A}}}}$. □

Definition 8.5.0.11. Let $\mathcal{A}, \mathcal{B}, \mathcal{C} \in \text{Obj}(\mathbf{Cat})$. We define the **Curry functor from $\mathcal{C}^{\mathcal{A} \times \mathcal{B}}$ to $(\mathcal{C}^{\mathcal{B}})^{\mathcal{A}}$** , denoted $\text{cur}_{(\mathcal{A}, \mathcal{B}, \mathcal{C})}^{\mathbf{Cat}} : \mathcal{C}^{\mathcal{A} \times \mathcal{B}} \rightarrow (\mathcal{C}^{\mathcal{B}})^{\mathcal{A}}$ by

- $\text{cur}_{(\mathcal{A}, \mathcal{B}, \mathcal{C})}^{\mathbf{Cat}}(F) := \text{cur}(F)$
- $\text{cur}_{(\mathcal{A}, \mathcal{B}, \mathcal{C})}^{\mathbf{Cat}}(\alpha) := \text{cur}(\alpha)$.

Exercise 8.5.0.12. Let $\mathcal{A}, \mathcal{B}, \mathcal{C} \in \text{Obj}(\mathbf{Cat})$. Then $\text{cur}_{(\mathcal{A}, \mathcal{B}, \mathcal{C})}^{\mathbf{Cat}} : \mathcal{C}^{\mathcal{A} \times \mathcal{B}} \rightarrow (\mathcal{C}^{\mathcal{B}})^{\mathcal{A}}$ is a functor.

Proof.

1. Let $F, G \in \text{Obj}(\mathcal{C}^{\mathcal{A} \times \mathcal{B}})$ and $\alpha \in \text{Hom}_{\mathcal{C}^{\mathcal{A} \times \mathcal{B}}}(F, G)$. Exercise 8.5.0.8 implies that

$$\begin{aligned} \text{cur}_{(\mathcal{A}, \mathcal{B}, \mathcal{C})}^{\mathbf{Cat}}(\alpha) &= \text{cur}(\alpha) \\ &\in \text{Hom}_{\mathcal{C}^{\mathcal{B}}}(\text{cur}(F), \text{cur}(G)) \\ &= \text{Hom}_{\mathcal{C}^{\mathcal{B}}}(\text{cur}_{(\mathcal{A}, \mathcal{B}, \mathcal{C})}^{\mathbf{Cat}}(F), \text{cur}_{(\mathcal{A}, \mathcal{B}, \mathcal{C})}^{\mathbf{Cat}}(G)). \end{aligned}$$

2. Let $F_1, F_2, F_3 \in \text{Obj}(\mathcal{C}^{\mathcal{A} \times \mathcal{B}})$, $\alpha_{1,2} \in \text{Hom}_{\mathcal{C}^{\mathcal{A} \times \mathcal{B}}}(F_1, F_2)$ and $\alpha_{2,3} \in \text{Hom}_{\mathcal{C}^{\mathcal{A} \times \mathcal{B}}}(F_2, F_3)$. Exercise 8.5.0.9 implies that

$$\begin{aligned} \text{cur}_{(\mathcal{A}, \mathcal{B}, \mathcal{C})}^{\text{Cat}}(\alpha_{2,3} \circ_{\mathcal{C}^{\mathcal{A} \times \mathcal{B}}} \alpha_{1,2}) &= \text{cur}((\alpha_{2,3} \circ_{\mathcal{C}^{\mathcal{A} \times \mathcal{B}}} \alpha_{1,2})) \\ &= \text{cur}(\alpha_{2,3}) \circ_{(\mathcal{C}^{\mathcal{B}})^{\mathcal{A}}} \text{cur}(\alpha_{1,2}) \\ &= \text{cur}_{(\mathcal{A}, \mathcal{B}, \mathcal{C})}^{\text{Cat}}(\alpha_{2,3}) \circ_{(\mathcal{C}^{\mathcal{B}})^{\mathcal{A}}} \text{cur}_{(\mathcal{A}, \mathcal{B}, \mathcal{C})}^{\text{Cat}}(\alpha_{1,2}). \end{aligned}$$

3. Exercise 8.5.0.10 implies that

$$\begin{aligned} \text{cur}_{(\mathcal{A}, \mathcal{B}, \mathcal{C})}^{\text{Cat}}(\text{id}_F^{\mathcal{C}^{\mathcal{A} \times \mathcal{B}}}) &= \text{cur}(\text{id}_F^{\mathcal{C}^{\mathcal{A} \times \mathcal{B}}}) \\ &= \text{id}_{\text{cur}(F)}^{(\mathcal{C}^{\mathcal{B}})^{\mathcal{A}}} \\ &= \text{id}_{\text{cur}_{(\mathcal{A}, \mathcal{B}, \mathcal{C})}^{\text{Cat}}(F)}^{(\mathcal{C}^{\mathcal{B}})^{\mathcal{A}}}. \end{aligned}$$

Thus $\text{cur}_{(\mathcal{A}, \mathcal{B}, \mathcal{C})}^{\text{Cat}} : \mathcal{C}^{\mathcal{A} \times \mathcal{B}} \rightarrow (\mathcal{C}^{\mathcal{B}})^{\mathcal{A}}$ is a functor. □

Definition 8.5.0.13. We define the **Currying natural transformation on Cat**, denoted $\text{cur}^{\text{Cat}} : [(- \times -)_{\text{Cat}}, -]_{\text{Cat}} \Rightarrow [-, [-, -]_{\text{Cat}}]_{\text{Cat}}$, by $(\text{cur}^{\text{Cat}})_{(\mathcal{A}, \mathcal{B}, \mathcal{C})} := \text{cur}_{(\mathcal{A}, \mathcal{B}, \mathcal{C})}^{\text{Cat}}$

Exercise 8.5.0.14. We have that $\text{cur}^{\text{Cat}} : [(- \times -)_{\text{Cat}}, -]_{\text{Cat}} \Rightarrow [-, [-, -]_{\text{Cat}}]_{\text{Cat}}$ is a natural transformation.

Proof.

show it is in correct Hom set

Show diagram commutes □

To Do:

- make a note that $[-, [-, -]] : \text{Cat}^{\text{op}} \times (\text{Cat}^{\text{op}} \times \text{Cat}) \rightarrow \text{Cat}$ means $[-, -] \circ \text{id}_{\text{Cat}} \times [-, -]$
- define opposite functor $OP : \mathcal{C} \rightarrow \mathcal{C}^{\text{op}}$
- show that $\text{cur}^{\text{Cat}} : F \Rightarrow G$ is a natural transformation (and natural isomorphism)

Definition 8.5.0.15.

Exercise 8.5.0.16. content...

Proof. content... □

Definition 8.5.0.17. Let $\mathcal{A}, \mathcal{B}, \mathcal{C} \in \text{Obj}(\text{Cat})$. We define the **Currying functor from $(\mathcal{C})^{\mathcal{A} \times \mathcal{B}}$ to $(\mathcal{C}^{\mathcal{B}})^{\mathcal{A}}$** , denoted $\text{cur}_{(\mathcal{A}, \mathcal{B}, \mathcal{C})}^{\text{Cat}} : (\mathcal{C})^{\mathcal{A} \times \mathcal{B}} \rightarrow (\mathcal{C}^{\mathcal{B}})^{\mathcal{A}}$, by

- $\text{cur}_{(\mathcal{A}, \mathcal{B}, \mathcal{C})}^{\text{Cat}}(F) := \text{cur}(F)$, ($A \in \text{Obj}(\mathcal{A})$)
- $\text{cur}_{(\mathcal{A}, \mathcal{B}, \mathcal{C})}^{\text{Cat}}(\alpha)(A) := \alpha_{(A, \cdot)}$.

(need exercise showing that if $\mathcal{A}, \mathcal{B}$ are small, then $\mathcal{A} \times \mathcal{B}$ is small)

Exercise 8.5.0.18. Let $\mathcal{A}, \mathcal{B}, \mathcal{C} \in \text{Obj}(\text{Cat})$. Then $\text{cur}_{(\mathcal{A}, \mathcal{B}, \mathcal{C})}^{\text{Cat}} : \mathcal{C}^{\mathcal{A} \times \mathcal{B}} \rightarrow (\mathcal{C}^{\mathcal{B}})^{\mathcal{A}}$ is a functor.

Proof.

1. Let $F, G \in \text{Obj}(\mathcal{C}^{\mathcal{A} \times \mathcal{B}})$ and $\alpha \in \text{Hom}_{\mathcal{C}^{\mathcal{A} \times \mathcal{B}}}(F, G)$. Then for each $A \in \text{Obj}(\mathcal{A})$,

$$\begin{aligned} \text{cur}_{(\mathcal{A}, \mathcal{B}, \mathcal{C})}^{\text{Cat}}(\alpha)(A) &= \alpha_{(A, \cdot)} \\ &\in \text{Hom}_{\mathcal{C}^{\mathcal{B}}}(F(A, \cdot), G(A, \cdot)) \quad (\text{by ref prev ex}) \\ &= \text{Hom}_{\mathcal{C}^{\mathcal{B}}}(\text{cur}_{(\mathcal{A}, \mathcal{B}, \mathcal{C})}^{\text{Cat}}(F)(A), \text{cur}_{(\mathcal{A}, \mathcal{B}, \mathcal{C})}^{\text{Cat}}(G)(A)) \end{aligned}$$

2.

3. □

8.6 Equalizer and Coequalizer Categories

Chapter 9

Limits

9.1 Cones and Cocones

Definition 9.1.0.1. Let $\mathcal{J}, \mathcal{C}$ be categories and $D : \mathcal{J} \rightarrow \mathcal{C}$. Then D is said to be a **diagram of type $\mathcal{J}$ in $\mathcal{C}$** .

Note 9.1.0.2. We are usually interested in the case that $\mathcal{J}$ is small. We will identify a diagram D with its image **and we visualize a diagram by the generating morphisms in $\mathcal{J}$** .

Example 9.1.0.3. **create example reference** Define $\mathcal{J}$ by

- $\text{Obj}(\mathcal{J}) = \{1, 2, 3, 4\}$,
- for $i, j \in \text{Obj}(\mathcal{J})$, $\text{Hom}_{\mathcal{J}}(i, j) = \{a_{ij}\}$.

Let $\mathcal{C}$ be a category and $D : \mathcal{J} \rightarrow \mathcal{C}$. Without including the identity morphisms or compositions, we can visualize D **by its set of generating morphisms (i.e. not including the identity morphisms)** as follows:

$$\begin{array}{ccc}
 & 1 \xrightarrow{a_{1,2}} 2 & \\
 a_{1,3} \swarrow & & \searrow a_{2,4} \\
 3 & \xrightarrow{a_{3,4}} 4 & \\
 & \xrightarrow{D} & \\
 & D_1 \xrightarrow{D_{1,2}} D_2 & \\
 D_{1,3} \swarrow & & \searrow D_{2,4} \\
 D_3 & \xrightarrow{D_{3,4}} D_4 &
 \end{array}$$

Definition 9.1.0.4. Let $\mathcal{J}, \mathcal{C}$ be categories. Suppose that $\mathcal{J}$ is small. Let $D \in \text{Obj}(\mathcal{C}^{\mathcal{J}})$.

- We define the **category of cones over D** , denoted $\mathbf{Cone}(D)$, by $\mathbf{Cone}(D) = (\Delta^{\mathcal{J}} \downarrow D)$.
- We define the **category of cocones under D** , denoted $\mathbf{Cocone}(D)$, by $\mathbf{Cocone}(D) = (D \downarrow \Delta^{\mathcal{J}})$.

Note 9.1.0.5.

- By Definition 4.4.0.5 and Definition 5.4.0.5,

$$\begin{aligned}
 \text{Obj}(\mathbf{Cone}(D)) &= \{(X, \phi) : X \in \text{Obj}(\mathcal{C}) \text{ and } \phi \in \text{Hom}_{\mathcal{C}^{\mathcal{J}}}(\Delta^{\mathcal{J}}(X), D)\} \\
 &= \{(X, \phi) : X \in \text{Obj}(\mathcal{C}) \text{ and } \phi : \Delta_X^{\mathcal{J}} \Rightarrow D\}
 \end{aligned}$$

and for $(X, \phi), (Y, \psi) \in \text{Obj}(\mathbf{Cone}(D))$,

$$\begin{aligned}
 \text{Hom}_{\mathbf{Cone}(D)}((X, \phi), (Y, \psi)) &= \{\alpha \in \text{Hom}_{\mathcal{C}}(X, Y) : \psi \circ_{\mathcal{C}^{\mathcal{J}}} \Delta^{\mathcal{J}}(\alpha) = \phi\} \\
 &= \{\alpha \in \text{Hom}_{\mathcal{C}}(X, Y) : \psi \circ_{\mathcal{C}^{\mathcal{J}}} \delta_{\alpha}^{\mathcal{J}} = \phi\} \\
 &= \{\alpha \in \text{Hom}_{\mathcal{C}}(X, Y) : \text{for each } j \in \mathcal{J}, \psi_j \circ_{\mathcal{C}} \alpha = \phi_j\}.
 \end{aligned}$$

make ex in subcategory section and rephrase this in terms of the commutative diagram on only the generating morphisms. This will enable us to more precisely cover limits like products and projective systems where the identity morphisms are not typically included in the definition of a diagram, but rather only the set of generating morphisms. Therefore, $(X, \phi) \in \text{Obj}(\mathbf{Cone}(D))$ iff for each $i, j \in \text{Obj}(\mathcal{J})$ and $f_{i,j} \in \text{Hom}_{\mathcal{J}}(i, j)$, the following diagram commutes:

$$\begin{array}{ccc} & X & \\ \phi_i \swarrow & & \searrow \phi_j \\ D_i & \xrightarrow{D_{f_{i,j}}} & D_j \end{array}$$

and $\alpha \in \text{Hom}_{\mathbf{Cone}(D)}((X, \phi), (Y, \psi))$ iff for each $i, j \in \text{Obj}(\mathcal{J})$ and $(i, j) \in \text{Hom}_{\mathcal{J}}(i, j)$ the following diagram commutes:

$$\begin{array}{ccc} & X & \\ \phi_i \swarrow & \alpha \downarrow & \searrow \phi_j \\ & Y & \\ \psi_i \swarrow & & \searrow \psi_j \\ D_i & \xrightarrow{D_{i,j}} & D_j \end{array}$$

- By Definition 4.4.0.6 and Definition 5.4.0.5,

$$\begin{aligned} \text{Obj}(\mathbf{Cocone}(D)) &= \{(X, \phi) : X \in \text{Obj}(\mathcal{C}) \text{ and } \phi \in \text{Hom}_{\mathcal{C}\mathcal{J}}(D, \Delta^{\mathcal{J}}(X))\} \\ &= \{(X, \phi) : X \in \text{Obj}(\mathcal{C}) \text{ and } \phi : D \Rightarrow \Delta_X^{\mathcal{J}}\} \end{aligned}$$

and for $(X, \phi), (Y, \psi) \in \text{Obj}(\mathbf{Cocone}(D))$,

$$\begin{aligned} \text{Hom}_{\mathbf{Cocone}(D)}((X, \phi), (Y, \psi)) &= \{\alpha \in \text{Hom}_{\mathcal{C}}(X, Y) : \Delta^{\mathcal{J}}(\alpha) \circ_{\mathcal{C}\mathcal{J}} \phi = \psi\} \\ &= \{\alpha \in \text{Hom}_{\mathcal{C}}(X, Y) : \delta_{\alpha}^{\mathcal{J}} \circ_{\mathcal{C}\mathcal{J}} \phi = \psi\} \\ &= \{\alpha \in \text{Hom}_{\mathcal{C}}(X, Y) : \text{for each } j \in \mathcal{J}, \alpha \circ_{\mathcal{C}} \phi_j = \psi_j\}. \end{aligned}$$

make ex in subcategory section and rephrase this in terms of the commutative diagram on only the generating morphisms. This will enable us to more precisely cover limits like products and projective systems where the identity morphisms are not typically included in the definition of a diagram, but rather only the set of generating morphisms. Therefore, $(X, \phi) \in \text{Obj}(\mathbf{Cone}(D))$ iff for each $i, j \in \text{Obj}(\mathcal{J})$ and $f_{i,j} \in \text{Hom}_{\mathcal{J}}(i, j)$, the following diagram commutes:

$$\begin{array}{ccc} D_i & \xrightarrow{D_{f_{i,j}}} & D_j \\ & \searrow \phi_i & \swarrow \phi_j \\ & X & \end{array}$$


and $\alpha \in \text{Hom}_{\mathbf{Cone}(D)}((X, \phi), (Y, \psi))$ iff for each $i, j \in \text{Obj}(\mathcal{J})$ and $(i, j) \in \text{Hom}_{\mathcal{J}}(i, j)$ the following diagram commutes:

$$\begin{array}{ccc} D_i & \xrightarrow{D_{f_{i,j}}} & D_j \\ & \searrow \phi_i & \swarrow \phi_j \\ & X & \\ \psi_i \swarrow & \alpha \downarrow & \searrow \psi_j \\ & Y & \end{array}$$


note that in the case of say a product of projective system, the category $\mathcal{J}$ does not include the identity morphisms in the Hom sets, to avoid having to specify the category with all morphisms, rephrase this definition in terms of the subcategory generated by a set of morphisms. This will need some exercises made in section on subcategories about natural transformations on a generated subcategory needing only to satisfy the composition rules in the def natl transf's on the generating morphisms of that category. maybe other similar exercises needed as well. For example in a product, when we write the diagram, we include the objects X, Y but we never include the morphisms id_X, id_Y in the diagram.

Example 9.1.0.6. ref example Define $\mathcal{J}$ and $\mathcal{D}$ as in previous example.

- Let $(X, \phi) \in \text{Obj}(\mathbf{Cone}(\mathcal{D}))$. We can visualize (X, ϕ) as follows:



- Let $(X, \phi) \in \text{Obj}(\mathbf{Cocone}(\mathcal{D}))$. We can visualize (X, ϕ) as follows:



FIX!!!!

Exercise 9.1.0.7. show $\mathbf{Cocone}(\mathcal{D}) \cong \mathbf{Cone}(\mathcal{D})^{\text{op}}$ (ref ex about comma categories and opposite categories)

Proof.

□

Exercise 9.1.0.8. Let $\mathcal{C}$ be a category, $\mathcal{J} \in \text{Obj}(\mathbf{Cat})$ and $D_1, D_2 \in \text{Obj}(\mathcal{C}^{\mathcal{J}})$.

1. Let $\alpha \in \text{Hom}_{\mathcal{C}^{\mathcal{J}}}(D_1, D_2)$.

- (a) For each $(X, \phi) \in \text{Obj}(\mathbf{Cone}(D_1))$,

$$(X, \alpha \circ_{\mathcal{C}^{\mathcal{J}}} \phi) \in \text{Obj}(\mathbf{Cone}(D_2)).$$

- (b) For each $(X, \phi), (X', \phi') \in \text{Obj}(\mathbf{Cone}(D_1))$ and $\eta \in \text{Hom}_{\mathbf{Cone}(D_1)}((X, \phi), (X', \phi'))$,

$$\eta \in \text{Hom}_{\mathbf{Cone}(D_2)}((X, \alpha \circ_{\mathcal{C}^{\mathcal{J}}} \phi), (X', \alpha \circ_{\mathcal{C}^{\mathcal{J}}} \phi')).$$

2. Let $\beta \in \text{Hom}_{\mathcal{C}^{\mathcal{J}}}(D_2, D_1)$.

- (a) FIX for cocones

(b)

Proof.

1. (a) Let $(X, \phi) \in \text{Obj}(\mathbf{Cone}(D_1))$. Then $X \in \text{Obj}(\mathcal{C})$ and $\phi \in \text{Hom}_{\mathcal{C}^{\mathcal{J}}}(\Delta_X^{\mathcal{J}}, D_1)$. Therefore $\alpha \circ_{\mathcal{C}^{\mathcal{J}}} \phi \in \text{Hom}_{\mathcal{C}^{\mathcal{J}}}(\Delta_X^{\mathcal{J}}, D_2)$. Hence $(X, \alpha \circ_{\mathcal{C}^{\mathcal{J}}} \phi) \in \text{Obj}(\mathbf{Cone}(D_2))$.
- (b) Let $(X, \phi), (X', \phi') \in \text{Obj}(\mathbf{Cone}(D_1))$ and $\eta \in \text{Hom}_{\mathbf{Cone}(D_1)}((X, \phi), (X', \phi'))$. Then $\eta \in \text{Hom}_{\mathcal{C}}(X, X')$ and $\phi' \circ_{\mathcal{C}^{\mathcal{J}}} \Delta^{\mathcal{J}}(\eta) = \phi$. Therefore $\alpha \circ_{\mathcal{C}^{\mathcal{J}}} \phi' \circ_{\mathcal{C}^{\mathcal{J}}} \Delta^{\mathcal{J}}(\eta) = \alpha \circ_{\mathcal{C}^{\mathcal{J}}} \phi$. Hence $\eta \in \text{Hom}_{\mathbf{Cone}(D_2)}((X, \alpha \circ_{\mathcal{C}^{\mathcal{J}}} \phi), (X', \alpha \circ_{\mathcal{C}^{\mathcal{J}}} \phi'))$.
2. use duality

□

Definition 9.1.0.9. Let $\mathcal{C}$ be a category, $\mathcal{J} \in \text{Obj}(\mathbf{Cat})$ and $D_1, D_2 \in \text{Obj}(\mathcal{C}^{\mathcal{J}})$.

1. Let $\alpha \in \text{Hom}_{\mathcal{C}^{\mathcal{J}}}(D_1, D_2)$. We define the **cone functor induced by α** , denoted $\text{Cn}_{\alpha} : \mathbf{Cone}(D_1) \rightarrow \mathbf{Cone}(D_2)$, by
 - $\text{Cn}_{\alpha}(X, \phi) := (X, \alpha \circ_{\mathcal{C}^{\mathcal{J}}} \phi), \quad (X, \phi) \in \text{Obj}(\mathbf{Cone}(D_1)),$
 - $\text{Cn}_{\alpha}(\eta) := \eta, \quad \eta \in \text{Hom}_{\mathbf{Cone}(D_1)}.$
2. similar for cocone functor

Exercise 9.1.0.10. Let $\mathcal{C}$ be a category, $\mathcal{J} \in \text{Obj}(\mathbf{Cat})$ and $D_1, D_2 \in \text{Obj}(\mathcal{C}^{\mathcal{J}})$.

1. Let $\alpha \in \text{Hom}_{\mathcal{C}^{\mathcal{J}}}(D_1, D_2)$. Then $\text{Cn}_{\alpha} : \mathbf{Cone}(D_1) \rightarrow \mathbf{Cone}(D_2)$ is a functor.
2. similar for cocone functor

Proof.

1. We note that ref prev ex implies that Cn_{α} is well-defined.
 - (a) ref prev ex implies that for each $(X, \phi), (X', \phi') \in \text{Obj}(\mathbf{Cone}(D_1))$, and $\eta \in \text{Hom}_{\mathbf{Cone}(D_1)}((X, \phi), (X', \phi'))$,

$$\begin{aligned} \text{Cn}_{\alpha}(\eta) &= \eta \\ &\in \text{Hom}_{\mathbf{Cone}(D_2)}((X, \alpha \circ_{\mathcal{C}^{\mathcal{J}}} \phi), (X', \alpha \circ_{\mathcal{C}^{\mathcal{J}}} \phi')) \\ &= \text{Hom}_{\mathbf{Cone}(D_2)}(\text{Cn}_{\alpha}(X, \phi), \text{Cn}_{\alpha}(X', \phi')). \end{aligned}$$
 - (b) Let $(X, \phi), (Y, \psi), (Z, \chi) \in \text{Obj}(\mathbf{Cone}(D_1))$ and $\mu \in \text{Hom}_{\mathbf{Cone}(D_1)}((X, \phi), (Y, \psi))$ and $\nu \in \text{Hom}_{\mathbf{Cone}(D_1)}((Y, \psi), (Z, \chi))$. Since $\mathbf{Cone}(D_1) = (\Delta^{\mathcal{J}} \downarrow D_1)$ and $\mathbf{Cone}(D_2) = (\Delta^{\mathcal{J}} \downarrow D_2)$, Definition 4.4.0.5 implies that

$$\begin{aligned} \text{Cn}_{\alpha}(\nu \circ_{\mathbf{Cone}(D_1)} \mu) &= \nu \circ_{\mathbf{Cone}(D_2)} \text{Cn}_{\alpha}(\mu) \\ &= \nu \circ_{(\Delta^{\mathcal{J}} \downarrow D_2)} \text{Cn}_{\alpha}(\mu) \\ &= \nu \circ_{\mathcal{C}} \text{Cn}_{\alpha}(\mu) \\ &= \nu \circ_{(\Delta^{\mathcal{J}} \downarrow D_2)} \text{Cn}_{\alpha}(\mu) \\ &= \nu \circ_{\mathbf{Cone}(D_2)} \text{Cn}_{\alpha}(\mu) \\ &= \text{Cn}_{\alpha}(\nu) \circ_{\mathbf{Cone}(D_2)} \text{Cn}_{\alpha}(\mu). \end{aligned}$$

- (c) By Definition 9.1.0.4,

$$\begin{aligned} \text{id}_{(X, \phi)}^{\mathbf{Cone}(D_1)} &= \text{id}_{(X, \phi)}^{(\Delta^{\mathcal{J}} \downarrow D_1)} \\ &= \text{id}_X^{\mathcal{C}} \\ &= \text{id}_{(X, \alpha \circ_{\mathcal{C}^{\mathcal{J}}} \phi)}^{(\Delta^{\mathcal{J}} \downarrow D_2)} \\ &= \text{id}_{(X, \alpha \circ_{\mathcal{C}^{\mathcal{J}}} \phi)}^{\mathbf{Cone}(D_2)}. \end{aligned}$$

Hence for each $(X, \phi) \in \text{Obj}(\mathbf{Cone}(D_1))$,

$$\begin{aligned} \text{Cn}_\alpha(\text{id}_{(X, \phi)}^{\mathbf{Cone}(D_1)}) &= \text{id}_{(X, \phi)}^{\mathbf{Cone}(D_1)} \\ &= \text{id}_{(X, \alpha \circ_{\mathcal{C}^{\mathcal{J}}} \phi)}^{\mathbf{Cone}(D_2)} \\ &= \text{id}_{\text{Cn}_\alpha(X, \phi)}^{\mathbf{Cone}(D_2)}. \end{aligned}$$

Hence $\text{Cn}_\alpha : \mathbf{Cone}(D_1) \rightarrow \mathbf{Cone}(D_2)$ is a functor.

2. use duality

□

Definition 9.1.0.11. Let $\mathcal{C}$ be a category, $\mathcal{J} \in \text{Obj}(\mathbf{Cat})$.

1. We define the **Cone functor**, denoted $\mathbf{Cone}(\cdot) : \mathcal{C}^{\mathcal{J}} \rightarrow \mathbf{Cat}$, by

- $\mathbf{Cone}(\cdot)(D) := \mathbf{Cone}(D)$, $D \in \text{Obj}(\mathcal{C}^{\mathcal{J}})$
- $\mathbf{Cone}(\cdot)(\alpha) := \text{Cn}_\alpha$, $\alpha \in \text{Hom}_{\mathcal{C}^{\mathcal{J}}}$.

2. define cocone functor

Exercise 9.1.0.12. Let $\mathcal{C}$ be a category, $\mathcal{J} \in \text{Obj}(\mathbf{Cat})$.

1. We have that $\mathbf{Cone}(\cdot) : \mathcal{C}^{\mathcal{J}} \rightarrow \mathbf{Cat}$ is a functor.

2. show cocone functor is functor

Proof.

1. Set $C := \mathbf{Cone}(\cdot)$.

(a) Let $D_1, D_2 \in \text{Obj}(\mathcal{C}^{\mathcal{J}})$ and $\alpha \in \text{Hom}_{\mathcal{C}^{\mathcal{J}}}(D_1, D_2)$. Then

$$\begin{aligned} C(\alpha) &= \text{Cn}_\alpha \\ &\in \text{Hom}_{\mathbf{Cat}}(\mathbf{Cone}(D_1), \mathbf{Cone}(D_2)) \\ &= \text{Hom}_{\mathbf{Cat}}(C(D_1), C(D_2)). \end{aligned}$$

(b) Let $D_1, D_2, D_3 \in \text{Obj}(\mathcal{C}^{\mathcal{J}})$, $\alpha \in \text{Hom}_{\mathcal{C}^{\mathcal{J}}}(D_1, D_2)$ and $\beta \in \text{Hom}_{\mathcal{C}^{\mathcal{J}}}(D_2, D_3)$. Then for each $(X, \phi) \in \text{Obj}(\mathbf{Cone}(D_1))$,

$$\begin{aligned} [C(\beta \circ_{\mathcal{C}^{\mathcal{J}}} \alpha)](X, \phi) &= \text{Cn}_{\beta \circ_{\mathcal{C}^{\mathcal{J}}} \alpha}(X, \phi) \\ &= (X, [\beta \circ_{\mathcal{C}^{\mathcal{J}}} \alpha] \circ_{\mathcal{C}^{\mathcal{J}}} \phi) \\ &= (X, \beta \circ_{\mathcal{C}^{\mathcal{J}}} [\alpha \circ_{\mathcal{C}^{\mathcal{J}}} \phi]) \\ &= \text{Cn}_\beta(X, \alpha \circ_{\mathcal{C}^{\mathcal{J}}} \phi) \\ &= \text{Cn}_\beta(\text{Cn}_\alpha(X, \phi)) \\ &= C(\beta)[C(\alpha)(X, \phi)] \\ &= [C(\beta) \circ_{\mathbf{Cat}} C(\alpha)](X, \phi) \end{aligned}$$

Therefore $C(\beta \circ_{\mathcal{C}^{\mathcal{J}}} \alpha) = C(\beta) \circ_{\mathbf{Cat}} C(\alpha)$.

(c) Let $D \in \text{Obj}(\mathcal{C}^{\mathcal{J}})$. Then for each $(X, \phi), (X', \phi') \in \text{Obj}(\mathbf{Cone}(D))$ and $\eta \in \text{Hom}_{\mathbf{Cone}(D)}((X, \phi), (X', \phi'))$,

$$\begin{aligned} C(\text{id}_D^{\mathcal{C}^{\mathcal{J}}})(X, \phi) &= \text{Cn}_{\text{id}_D^{\mathcal{C}^{\mathcal{J}}}}(X, \phi) \\ &= (X, \text{id}_D^{\mathcal{C}^{\mathcal{J}}} \circ_{\mathcal{C}^{\mathcal{J}}} \phi) \\ &= (X, \phi) \\ &= \text{id}_{\mathbf{Cone}(D)}^{\mathbf{Cat}}(X, \phi) \\ &= \text{id}_{C(D)}^{\mathbf{Cat}}(X, \phi) \end{aligned}$$

and

$$\begin{aligned}
 C(\mathrm{id}_D^{\mathcal{C}^{\mathcal{J}}})(\eta) &= \mathrm{Cn}_{\mathrm{id}_D^{\mathcal{C}^{\mathcal{J}}}}(\eta) \\
 &= \eta \\
 &= \mathrm{id}_{\mathbf{Cone}(D)}^{\mathbf{Cat}}(\eta) \\
 &= \mathrm{id}_{C(D)}^{\mathbf{Cat}}(\eta).
 \end{aligned}$$

Thus $C(\mathrm{id}_D^{\mathcal{C}^{\mathcal{J}}}) = \mathrm{id}_{C(D)}^{\mathbf{Cat}}$.

Hence $\mathbf{Cone}(\cdot) : \mathcal{C}^{\mathcal{J}} \rightarrow \mathbf{Cat}$ is a functor.

2. **FINISH for cocones!!!!**

□

9.2 Limits and Colimits

9.2.1 Universal Morphisms

make subsection instead of section and put it in section on limits after section on cones and cocones

Definition 9.2.1.1. Let $\mathcal{C}, \mathcal{D}$ be categories, $X \in \text{Obj}(\mathcal{D})$, $F : \mathcal{C} \rightarrow \mathcal{D}$ and $A \in \text{Obj}(\mathcal{C})$.

- Let $f \in \text{Hom}_{\mathcal{D}}(F(A), X)$. Then (A, f) is said to be a **universal morphism from F to X** if for each $A' \in \text{Obj}(\mathcal{C})$ and $f' \in \text{Hom}_{\mathcal{D}}(F(A'), X)$, there exists a unique $\alpha \in \text{Hom}_{\mathcal{C}}(A', A)$ such that $f \circ_{\mathcal{D}} F(\alpha) = f'$, i.e. the following diagram commutes:

$$\begin{array}{ccc}
 & F(A') & \\
 A' & \downarrow F(\alpha) & \searrow f' \\
 \downarrow \alpha & F(A) & \\
 A & \downarrow f & \\
 & X &
 \end{array}$$

- Let $f \in \text{Hom}_{\mathcal{D}}(X, F(A))$. Then (A, f) is said to be a **universal morphism from X to F** if for each $A' \in \text{Obj}(\mathcal{C})$ and $f' \in \text{Hom}_{\mathcal{D}}(X, F(A'))$, there exists a unique $\alpha \in \text{Hom}_{\mathcal{C}}(A, A')$ such that $F(\alpha) \circ_{\mathcal{D}} f = f'$, i.e. the following diagram commutes:

$$\begin{array}{ccc}
 & X & \\
 A & \downarrow f & \searrow f' \\
 \downarrow \alpha & F(A) & \\
 A' & \downarrow F(\alpha) & \\
 & F(A') &
 \end{array}$$

Exercise 9.2.1.2. Let $\mathcal{C}, \mathcal{D}$ be categories, $X \in \text{Obj}(\mathcal{D})$, $F : \mathcal{C} \rightarrow \mathcal{D}$ and $A \in \text{Obj}(\mathcal{C})$.

1. Let $f \in \text{Hom}_{\mathcal{D}}(F(A), X)$. Then (A, f) is a universal morphism from F to X iff $(A, f) \in \text{Term}_{(F \downarrow X)}$.
2. Let $f \in \text{Hom}_{\mathcal{D}}(X, F(A))$. Then (A, f) is a universal morphism from X to F iff $(A, f) \in \text{Coterm}_{(X \downarrow F)}$.

Proof.

1. By Definition 4.4.0.5, $\text{Obj}(F \downarrow X) = \coprod_{A' \in \text{Obj}(\mathcal{C})} \text{Hom}_{\mathcal{D}}(F(A'), X)$. In particular, $(A, f) \in \text{Obj}(F \downarrow X)$. By Definition 4.4.0.5, $\text{Hom}_{(F \downarrow X)}((A', f'), (A, f)) = \{\alpha \in \text{Hom}_{\mathcal{C}}(A', A) : f \circ_{\mathcal{D}} F(\alpha) = f'\}$. By Definition 9.2.1.1,

$$\begin{aligned}
 (A, f) \text{ is a universal morphism from } F \text{ to } X &\iff \text{for each } A' \in \text{Obj}(\mathcal{C}) \text{ and } f' \in \text{Hom}_{\mathcal{D}}(F(A'), X), \text{ there exists a unique } \alpha \in \text{Hom}_{\mathcal{C}}(A', A) \text{ such that } f \circ_{\mathcal{D}} F(\alpha) = f' \\
 &\iff \text{for each } (A', f') \in \text{Obj}(F \downarrow X) \text{ there exists a unique } \alpha \in \text{Hom}_{(F \downarrow X)}(A', A) \text{ such that } f \circ_{\mathcal{D}} F(\alpha) = f' \\
 &\iff (A, f) \in \text{Term}_{(F \downarrow X)}.
 \end{aligned}$$

2. Similar to prev part

FINISH!!!

□

Note 9.2.1.3. make a comment on how if (A, f) is universal from X to F , then for each (A', f') , f' is a post-processing of f and (A, f) is universal from F to X , then for each (A', f') , f' is a pre-processing of f

9.2.2 Limits and Colimits

Definition 9.2.2.1. Let $\mathcal{J}, \mathcal{C}$ be categories. Suppose that $\mathcal{J}$ is small. Let $D \in \text{Obj}(\mathcal{C}^{\mathcal{J}})$.

- Let $(X, \phi) \in \text{Obj}(\mathbf{Cone}(D))$. Then (X, ϕ) is said to be a **D -limit**, denoted $(X, \phi) = \lim D$, if (X, ϕ) is a universal morphism from $\Delta^{\mathcal{J}}$ to D .
- We define the **limit set of D** , denoted $\text{limSet}(D)$, by

$$\text{limSet } D := \{(X, \phi) \in \text{Obj}(\mathbf{Cone}(D)) : \text{Obj}(\mathbf{Cone}(D)) \text{ is a } D\text{-limit}\}.$$

- Let $(X, \phi) \in \text{Obj}(\mathbf{Cocone}(D))$. Then (X, ϕ) is said to be a **D -colimit**, denoted $(X, \phi) = \text{colim } D$, if (X, ϕ) is a universal morphism from D to $\Delta^{\mathcal{J}}$.
- We define the **colimit set of D** , denoted $\text{colimSet}(D)$, by

$$\text{colimSet } D := \{(X, \phi) \in \text{Obj}(\mathbf{Cocone}(D)) : \text{Obj}(\mathbf{Cocone}(D)) \text{ is a } D\text{-colimit}\}.$$

maybe define lim set, but maybe not, we want to be able to write $(X, \phi) = \lim D$, but we need some guarantee that anything we would possibly want to say is invariant under isomorphism, other wise different limits of D may always be interchangeable, ask someone if we need to be careful here and define limit sets. For instance [commutativity](#)

Note 9.2.2.2. Let $\mathcal{J}, \mathcal{C}$ be categories. Suppose that $\mathcal{J}$ is small. Let $D \in \text{Obj}(\mathcal{C}^{\mathcal{J}})$.

- Let $(X, \phi) \in \text{Obj}(\mathbf{Cone}(D))$. Then

$$\begin{aligned} (X, \phi) \in \text{limSet } D &\iff (X, \phi) \text{ is } \mathbf{Cone}(D)\text{-terminal} \\ &\iff \text{for each } (X', \phi') \in \text{Obj}(\mathbf{Cone}(D)), \text{ there exists a unique} \\ &\quad \alpha \in \text{Hom}_{\mathcal{C}}(X', X) \text{ such that for each } j \in \mathcal{J}, \phi_j \circ \alpha = \phi'_j \end{aligned}$$

- Let $(X, \phi) \in \text{Obj}(\mathbf{Cocone}(D))$. Then

$$\begin{aligned} (X, \phi) \in \text{colimSet } D &\iff (X, \phi) \text{ is } \mathbf{Cocone}(D)\text{-initial} \\ &\iff \text{for each } (X', \phi') \in \text{Obj}(\mathbf{Cocone}(D)), \text{ there exists a unique} \\ &\quad \alpha \in \text{Hom}_{\mathcal{C}}(X, X') \text{ such that for each } j \in \mathcal{J}, \alpha \circ \phi_j = \phi'_j \end{aligned}$$

Exercise 9.2.2.3. Let $\mathcal{J}, \mathcal{C}$ be categories. Suppose that $\mathcal{J}$ is small. Let $D \in \text{Obj}(\mathcal{C}^{\mathcal{J}})$.

1. For each $(X, \phi), (Y, \psi) \in \text{limSet } D$, there exists $f \in \text{Iso}_{\mathbf{Cone}(D)}((X, \phi), (Y, \psi))$ such that $\text{Hom}_{\mathbf{Cone}(D)}((X, \phi), (Y, \psi)) = \{f\}$.
make/ref ex in comma category section about inverses implies that $f \in \text{Iso}_{\mathcal{C}}(X, Y)$.

2. colimits

Proof.

1. Let $(X, \phi), (Y, \psi) \in \text{limSet } D$. Then $(X, \phi), (Y, \psi) \in \text{Cone}(D)$ and $(X, \phi), (Y, \psi)$ are $\text{Cone}(D)$ -terminal. Exercise 7.1.0.4 implies that there exists $f \in \text{Iso}_{\mathbf{Cone}(D)}((X, \phi), (Y, \psi))$ such that $\text{Hom}_{\mathbf{Cone}(D)}((X, \phi), (Y, \psi)) = \{f\}$.

2. colimits

□

9.3 Fundamental Examples

9.3.1 Products and Coproducts

Definition 9.3.1.1. Let $\mathcal{J}$ be a discrete category, $\mathcal{C}$ a category, $D \in \mathcal{C}^{\mathcal{J}}$ and $(X, \pi) \in \mathbf{Cone}(D)$. Then (X, π) is said to be a **product** of D .

Note 9.3.1.2. We may think of a product of D as a tuple $(X,)$

Exercise 9.3.1.3. show that if $\mathbf{1}$ is terminal, then $X \times \mathbf{1} \cong X$.

Exercise 9.3.1.4. show that if $\mathcal{C}$ has products, then $\mathcal{C}$ has a copy morphism (define a copy morphism) see page 32 in ‘elementary categories, elementary toposes’ by McLarty.

9.3.2 Equalizers and Coequalizers

9.3.3 Projective and Inductive Limits

Definition 9.3.3.1. Let $\mathcal{J}$ be a **directed** poset, $\mathcal{C}$ a category and $D \in \mathcal{C}^{\mathcal{J}}$. Then D is said to be a **$\mathcal{C}$ -projective system**.

Note 9.3.3.2. We may think of

- a $\mathcal{C}$ -projective system as a tuple $((X_j)_{j \in J}, (\pi_{j,k})_{(j,k) \in \leq})$ where $(J, \leq)$ is a **directed** poset, $(X_j)_{j \in J} \subset \text{Obj}(\mathcal{C})$ and $(\pi_{j,k})_{(j,k) \in \leq} \subset \text{Hom}_{\mathcal{C}}$ satisfy that for each $j, k, l \in J$,
 1. $j \leq k$ implies that $\pi_{j,k} \in \text{Hom}_{\mathcal{C}}(X_k, X_j)$,
 2. $\pi_{j,j} = \text{id}_{X_j}$,
 3. $j \leq k$ and $k \leq l$ implies that $\pi_{j,k} \circ_{\mathcal{C}} \pi_{k,l} = \pi_{j,l}$.
- a cone over D as a tuple $(X, (\pi_j)_{j \in J})$ where $X \in \text{Obj}(\mathcal{C})$ and $(\pi_j)_{j \in J} \subset \text{Hom}_{\mathcal{C}}$ satisfy that for each $j, k \in J$, $j \leq k$ implies that $\pi_{j,k} \circ_{\mathcal{C}} \pi_k = \pi_j$.

make some diagrams

Exercise 9.3.3.3. Let $\mathcal{C}$ be a category and $(X_j)_{j \in \mathbb{N}} \subset \text{Obj}(\mathcal{C})$. Define $(Y_n)_{n \in \mathbb{N}} \subset \text{Obj}(\mathcal{C})$ and $(\pi_{n,k})_{n \leq k} \in \prod_{n \leq k} \text{Hom}_{\mathcal{C}}(Y_k, Y_n)$ by $Y_n := \prod_{j=1}^n X_j$ and $\pi_{n,k}$. Suppose that for each $(\prod_{n \in \mathbb{N}}, (\pi_n)_{n \in \mathbb{N}})$ **arg1**.

Definition 9.3.3.4. Let $(\mathcal{J}, \mathcal{C}, D)$ be a $\mathcal{C}$ -projective system and $(X, \phi) \in \text{Obj}(\mathbf{Cone}(D))$. Then (X, ϕ) is said to be a $\mathcal{C}$ -projective limit of $(\mathcal{J}, \mathcal{C}, D)$ if (X, ϕ) is a limit of D .

Note 9.3.3.5. We may think of a projective limit if $(\mathcal{J}, \mathcal{C}, D)$ as a cone $(X, (\pi_j)_{j \in J})$ over D satisfying that for each cone $(Y, (\tau_j)_{j \in J})$ over D , there exists a unique $\alpha \in \text{Hom}_{\mathcal{C}}(Y, X)$ such that $\tau_j = \pi_j \circ_{\mathcal{C}} \alpha$.

make some diagrams

9.4 Completeness and Cocompleteness

Definition 9.4.0.1. Let $\mathcal{C}$ be a category and $\mathcal{J} \in \text{Obj}(\mathbf{Cat})$. Then $\mathcal{C}$ is said to

- **have all $\mathcal{J}$ -limits** if for each $D \in \text{Obj}(\mathcal{C}^{\mathcal{J}})$, $\lim_{\text{Set}} D \neq \emptyset$.
- **have all $\mathcal{J}$ -colimits** if for each $D \in \text{Obj}(\mathcal{C}^{\mathcal{J}})$, $\text{colim}_{\text{Set}} D \neq \emptyset$.

Definition 9.4.0.2. Let $\mathcal{C}$ be a category and $\mathcal{J} \in \text{Obj}(\mathbf{Cat})$.

- – Suppose that $\mathcal{C}$ has all $\mathcal{J}$ -limits. Let $L \in \prod_{D \in \text{Obj}(\mathcal{C}^{\mathcal{J}})} \lim_{\text{Set}} D$. Then L is said to be a $\mathcal{C}^{\mathcal{J}}$ -**limit choice function**.
– We define $\lim_{\text{Choice}}(\mathcal{C}^{\mathcal{J}}) := \prod_{D \in \text{Obj}(\mathcal{C}^{\mathcal{J}})} \lim_{\text{Set}} D$.
- – Suppose that $\mathcal{C}$ has all $\mathcal{J}$ -colimits. Let $C \in \prod_{D \in \text{Obj}(\mathcal{C}^{\mathcal{J}})} \text{colim}_{\text{Set}} D$. Then C is said to be a $\mathcal{C}^{\mathcal{J}}$ -**colimit choice function**.
– We define $\text{colim}_{\text{Choice}}(\mathcal{C}^{\mathcal{J}}) := \prod_{D \in \text{Obj}(\mathcal{C}^{\mathcal{J}})} \text{colim}_{\text{Set}} D$.

Note 9.4.0.3. Let $L \in \lim_{\text{Choice}}(\mathcal{C}^{\mathcal{J}})$, $D_1, D_2 \in \text{Obj}(\mathcal{C}^{\mathcal{J}})$ and $\alpha \in \text{Hom}_{\mathcal{C}^{\mathcal{J}}}(D_1, D_2)$. Write $\text{Cn}_{\alpha}(L_{D_1}) = (X, \phi)$ and $L_{D_2} = (Y, \psi)$. Since $L_{D_2} \in \lim_{\text{Set}} D_2$, L_{D_2} is $\text{Cone}(D_2)$ -terminal. Thus there exists $f \in \text{Hom}_{\mathbf{Cone}(D_2)}(\text{Cn}_{\alpha}(L_{D_1}), L_{D_2})$ such that $\text{Hom}_{\mathbf{Cone}(D_2)}(\text{Cn}_{\alpha}(L_{D_1}), \{f\})$. By Note 9.1.0.5, $f \in \text{Hom}_{\mathcal{C}}(X, Y)$.

Definition 9.4.0.4. Let $\mathcal{C}$ be a category and $\mathcal{J} \in \text{Obj}(\mathbf{Cat})$.

1. Suppose that $\mathcal{C}$ has all $\mathcal{J}$ -limits. Let $L \in \lim_{\text{Choice}}(\mathcal{C}^{\mathcal{J}})$, $D_1, D_2 \in \text{Obj}(\mathcal{C}^{\mathcal{J}})$ and $\alpha \in \text{Hom}_{\mathcal{C}^{\mathcal{J}}}(D_1, D_2)$. Write $\text{Cn}_{\alpha}(L_{D_1}) = (X, \phi)$ and $L_{D_2} = (Y, \psi)$. We define the **cone morphism induced by L and α** , denoted $\eta_{\alpha}^L \in \text{Hom}_{\mathcal{C}}$ to be the unique $f \in \text{Hom}_{\mathcal{C}}(X, Y)$ such that $\text{Hom}_{\mathbf{Cone}(D_2)}(\text{Cn}_{\alpha}(L_{D_1}), L_{D_2}) = \{f\}$.
2. **do something similar for colimits**

Definition 9.4.0.5. Let $\mathcal{C}$ be a category and $\mathcal{J} \in \text{Obj}(\mathbf{Cat})$.

- Suppose that $\mathcal{C}$ has all $\mathcal{J}$ -limits. Let $L \in \lim_{\text{Choice}}(\mathcal{C}^{\mathcal{J}})$.
– For $D_1, D_2 \in \text{Obj}(\mathcal{C}^{\mathcal{J}})$ and $\alpha \in \text{Hom}_{\mathcal{C}^{\mathcal{J}}}(D_1, D_2)$, we define $\eta_{\alpha}^L \in \text{Hom}_{\mathbf{Cone}(D_2)}(\text{Cn}_{\alpha}(L_{D_1}), L_{D_2})$ as in Exercise ??.
– We define the **L -limit functor**, denoted $\lim_L : \mathcal{C}^{\mathcal{J}} \rightarrow \mathcal{C}$, by
* $\lim_L(D) := \text{proj}_{\text{Obj}(\mathcal{C})}(L_D)$, $D \in \text{Obj}(\mathcal{C}^{\mathcal{J}})$
* $\lim_L(\alpha) := \eta_{\alpha}^L$, $\alpha \in \text{Hom}_{\mathcal{C}^{\mathcal{J}}}$.

- **define something similar for colimits, might need to change notation from η_{α}^L to something else**

Exercise 9.4.0.6. Let $\mathcal{C}$ be a category and $\mathcal{J} \in \text{Obj}(\mathbf{Cat})$.

1. Suppose that $\mathcal{C}$ has all $\mathcal{J}$ -limits. Let $L \in \lim_{\text{Choice}}(\mathcal{C}^{\mathcal{J}})$. Then $\lim_L : \mathcal{C}^{\mathcal{J}} \rightarrow \mathcal{C}$ is a functor.
2. **do something similar for colimits**

Proof.

1. (a) Let $D_1, D_2 \in \text{Obj}(\mathcal{C}^{\mathcal{J}})$ and $\alpha \in \text{Hom}_{\mathcal{C}^{\mathcal{J}}}(D_1, D_2)$. Write $L_{D_1} = (X_1, \phi_1)$ and $L_{D_2} = (X_2, \phi_2)$. Then $\lim_L(D_1) = X_1$ and $\lim_L(D_2) = X_2$. By Definition 9.1.0.4,

$$\begin{aligned} \text{Hom}_{\mathbf{Cone}(D_2)}(\text{Cn}_{\alpha}(L_{D_1}), L_{D_2}) &= \text{Hom}_{\mathbf{Cone}(D_2)}((X_1, \alpha \circ_{\mathcal{C}^{\mathcal{J}}} \phi_1), (X_2, \phi_2)) \\ &\subset \text{Hom}_{\mathcal{C}}(X_1, X_2). \end{aligned}$$

Therefore

$$\begin{aligned}
 \lim_L(\alpha) &= \eta_\alpha^L \\
 &\in \text{Hom}_{\mathbf{Cone}(D_2)}(\text{Cn}_\alpha(L_{D_1}), L_{D_2}) \\
 &\subset \text{Hom}_{\mathcal{C}}(X_1, X_2) \\
 &= \text{Hom}_{\mathcal{C}}(\lim_L(D_1), \lim_L(D_2)).
 \end{aligned}$$

(b) Let $D_1, D_2, D_3 \in \text{Obj}(\mathcal{C}^\mathcal{J})$, $\alpha \in \text{Hom}_{\mathcal{C}^\mathcal{J}}(D_1, D_2)$ and $\beta \in \text{Hom}_{\mathcal{C}^\mathcal{J}}(D_2, D_3)$. Write $L_{D_1} = (X_1, \phi_1)$, $L_{D_2} = (X_2, \phi_2)$ and $L_{D_3} = (X_3, \phi_3)$. Then $\lim_L(D_1) = X_1$, $\lim_L(D_2) = X_2$ and $\lim_L(D_3) = X_3$. By Definition 9.4.0.4 and Definition 9.1.0.4,

- $\text{Hom}_{\mathbf{Cone}(D_2)}(\text{Cn}_\alpha(L_{D_1}), L_{D_2}) = \{\eta_\alpha^L\}$,
- $\text{Hom}_{\mathbf{Cone}(D_3)}(\text{Cn}_\beta(L_{D_2}), L_{D_3}) = \{\eta_\beta^L\}$,
- $\text{Hom}_{\mathbf{Cone}(D_3)}(\text{Cn}_{\beta \circ_{\mathcal{C}^\mathcal{J}} \alpha}(L_{D_1}), L_{D_3}) = \{\eta_{\beta \circ_{\mathcal{C}^\mathcal{J}} \alpha}^L\}$

and

- $\eta_\alpha^L \in \text{Hom}_{\mathcal{C}}(X_1, X_2)$ and $\phi_2 \circ_{\mathcal{C}^\mathcal{J}} \Delta^\mathcal{J}(\eta_\alpha^L) = \alpha \circ_{\mathcal{C}^\mathcal{J}} \phi_1$
- $\eta_\beta^L \in \text{Hom}_{\mathcal{C}}(X_2, X_3)$ and $\phi_3 \circ_{\mathcal{C}^\mathcal{J}} \Delta^\mathcal{J}(\eta_\beta^L) = \beta \circ_{\mathcal{C}^\mathcal{J}} \phi_2$.

Therefore $\eta_\beta^L \circ_{\mathcal{C}} \eta_\alpha^L \in \text{Hom}_{\mathcal{C}}(X_1, X_3)$ and

$$\begin{aligned}
 \phi_3 \circ_{\mathcal{C}^\mathcal{J}} \Delta^\mathcal{J}(\eta_\beta^L \circ_{\mathcal{C}} \eta_\alpha^L) &= \phi_3 \circ_{\mathcal{C}^\mathcal{J}} [\Delta^\mathcal{J}(\eta_\beta^L) \circ_{\mathcal{C}^\mathcal{J}} \Delta^\mathcal{J}(\eta_\alpha^L)] \\
 &= [\phi_3 \circ_{\mathcal{C}^\mathcal{J}} \Delta^\mathcal{J}(\eta_\beta^L)] \circ_{\mathcal{C}^\mathcal{J}} \Delta^\mathcal{J}(\eta_\alpha^L) \\
 &= [\beta \circ_{\mathcal{C}^\mathcal{J}} \phi_2] \circ_{\mathcal{C}^\mathcal{J}} \Delta^\mathcal{J}(\eta_\alpha^L) \\
 &= \beta \circ_{\mathcal{C}^\mathcal{J}} [\phi_2 \circ_{\mathcal{C}^\mathcal{J}} \Delta^\mathcal{J}(\eta_\alpha^L)] \\
 &= \beta \circ_{\mathcal{C}^\mathcal{J}} [\alpha \circ_{\mathcal{C}^\mathcal{J}} \phi_1] \\
 &= [\beta \circ_{\mathcal{C}^\mathcal{J}} \alpha] \circ_{\mathcal{C}^\mathcal{J}} \phi_1.
 \end{aligned}$$

Hence

$$\begin{aligned}
 \eta_\beta^L \circ_{\mathcal{C}} \eta_\alpha^L &\in \text{Hom}_{\mathbf{Cone}(D_3)}(X_1, (\beta \circ_{\mathcal{C}^\mathcal{J}} \alpha) \circ_{\mathcal{C}^\mathcal{J}} \phi_1, (X_3, \phi_3)) \\
 &= \text{Hom}_{\mathbf{Cone}(D_3)}(\text{Cn}_{\beta \circ_{\mathcal{C}^\mathcal{J}} \alpha}(L_{D_1}), L_{D_3}) \\
 &= \{\eta_{\beta \circ_{\mathcal{C}^\mathcal{J}} \alpha}^L\}.
 \end{aligned}$$

Thus

$$\begin{aligned}
 \lim_L(\beta \circ_{\mathcal{C}^\mathcal{J}} \alpha) &= \eta_{\beta \circ_{\mathcal{C}^\mathcal{J}} \alpha}^L \\
 &= \eta_\beta^L \circ_{\mathcal{C}} \eta_\alpha^L \\
 &= \lim_L(\beta) \circ_{\mathcal{C}} \lim_L(\alpha).
 \end{aligned}$$

(c) Let $D \in \text{Obj}(\mathcal{C}^\mathcal{J})$. Write $L_D = (X, \phi)$. Exercise 10.1.0.10 **ref prev ex** implies that $\mathbf{Cone}(\cdot) : \mathcal{C}^\mathcal{J} \rightarrow \mathbf{Cat}$ and

$$\begin{aligned}
 \text{Cn}_{\text{id}_D^{\mathcal{C}^\mathcal{J}}} &= \mathbf{Cone}(\cdot)(\text{id}_D^{\mathcal{C}^\mathcal{J}}) \\
 &= \text{id}_{\mathbf{Cone}(D)}^{\mathbf{Cat}}.
 \end{aligned}$$

Then

$$\begin{aligned}
 \text{Cn}_{\text{id}_D^{\mathcal{C}^\mathcal{J}}}(L_D) &= \text{id}_{\mathbf{Cone}(D)}^{\mathbf{Cat}}(L_D) \\
 &= L_D.
 \end{aligned}$$

Since L_D is **Cone**(D)-terminal,

$$\begin{aligned} \lim_L(\mathrm{id}_D^{\mathcal{C}^{\mathcal{J}}}) &= \eta_{\mathrm{id}_D^{\mathcal{C}^{\mathcal{J}}}}^L \\ &\in \mathrm{Hom}_{\mathbf{Cone}(D)}(\mathrm{Cn}_{\mathrm{id}_D^{\mathcal{C}^{\mathcal{J}}}}(L_D), L_D) \\ &= \mathrm{Hom}_{\mathbf{Cone}(D)}(L_D, L_D) \\ &= \{\mathrm{id}_{L_D}^{\mathbf{Cone}(D)}\}. \end{aligned}$$

By Definition ?? **ref cone def**,

$$\begin{aligned} \mathrm{id}_{L_D}^{\mathbf{Cone}(D)} &= \mathrm{id}_{L_D}^{(\Delta^{\mathcal{J}} \downarrow D)} \\ &= \mathrm{id}_{(X, \phi)}^{(\Delta^{\mathcal{J}} \downarrow D)} \\ &= \mathrm{id}_X^{\mathcal{C}}. \end{aligned}$$

Therefore

$$\begin{aligned} \lim_L(\mathrm{id}_D^{\mathcal{C}^{\mathcal{J}}}) &= \mathrm{id}_{L_D}^{\mathbf{Cone}(D)} \\ &= \mathrm{id}_X^{\mathcal{C}} \\ &= \mathrm{id}_{\lim_L(D)}^{\mathcal{C}}. \end{aligned}$$

Hence $\lim_L : \mathcal{C}^{\mathcal{J}} \rightarrow \mathcal{C}$ is a functor.

2. colimit functor

□

Exercise 9.4.0.7. Let $\mathcal{C}$ be a category and $\mathcal{J} \in \mathrm{Obj}(\mathbf{Cat})$.

- Suppose that $\mathcal{C}$ has all $\mathcal{J}$ -limits. Let $L_1, L_2 \in \mathrm{limChoice}(\mathcal{C}^{\mathcal{J}})$. We have that $\prod_{D \in \mathrm{Obj}(\mathcal{C}^{\mathcal{J}})} \mathrm{Hom}_{\mathbf{Cone}(D)}(L_1(D), L_2(D)) = \prod_{D \in \mathrm{Obj}(\mathcal{C}^{\mathcal{J}})} \mathrm{Iso}_{\mathbf{Cone}(D)}$ and there exists $f : \mathrm{Obj}(\mathcal{C}^{\mathcal{J}}) \rightarrow \mathrm{Hom}_{\mathcal{C}}$ such that $\prod_{D \in \mathrm{Obj}(\mathcal{C}^{\mathcal{J}})} \mathrm{Hom}_{\mathbf{Cone}(D)}(L_1(D), L_2(D)) = \{f\}$.

2. something similar for colimits.

Proof.

- We note that $D \in \mathrm{Obj}(\mathcal{C}^{\mathcal{J}})$,

$$\begin{aligned} \{L_1(D), L_2(D)\} &\subset \mathrm{limSet } D \\ &\subset \mathrm{Term}_{\mathbf{Cone}(D)}, \end{aligned}$$

Exercise 7.1.0.4 implies that for each $D \in \mathrm{Obj}(\mathcal{C}^{\mathcal{J}})$, $\mathrm{Iso}_{\mathbf{Cone}(D)}(L_1(D), L_2(D)) = \mathrm{Hom}_{\mathbf{Cone}(D)}(L_1(D), L_2(D))$ and $\# \mathrm{Hom}_{\mathbf{Cone}(D)}$

- Then $\prod_{D \in \mathrm{Obj}(\mathcal{C}^{\mathcal{J}})} \mathrm{Hom}_{\mathbf{Cone}(D)}(L_1(D), L_2(D)) = \prod_{D \in \mathrm{Obj}(\mathcal{C}^{\mathcal{J}})} \mathrm{Iso}_{\mathbf{Cone}(D)}(L_1(D), L_2(D))$ and **The axiom of choice** implies that

there exists a $f \in \prod_{D \in \mathrm{Obj}(\mathcal{C}^{\mathcal{J}})} \mathrm{Hom}_{\mathbf{Cone}(D)}(L_1(D), L_2(D))$ such that $\prod_{D \in \mathrm{Obj}(\mathcal{C}^{\mathcal{J}})} \mathrm{Hom}_{\mathbf{Cone}(D)}(L_1(D), L_2(D)) = \{f\}$.

2. similar for colimits

□

Definition 9.4.0.8. Let $\mathcal{C}$ be a category and $\mathcal{J} \in \mathrm{Obj}(\mathbf{Cat})$.

- Suppose that $\mathcal{C}$ has all $\mathcal{J}$ -limits. Let $L_1, L_2 \in \mathrm{limChoice}(\mathcal{C}^{\mathcal{J}})$. We define the **natural transformation from $\lim_{L_1}$ to $\lim_{L_2}$** , denoted $f^{L_1, L_2} : \mathrm{Obj}(\mathcal{C}^{\mathcal{J}}) \rightarrow \mathrm{Hom}_{\mathcal{C}}$ to be the unique $f \in \prod_{D \in \mathrm{Obj}(\mathcal{C}^{\mathcal{J}})} \mathrm{Hom}_{\mathbf{Cone}(D)}(L_1(D), L_2(D))$.

- similar for colimits

Exercise 9.4.0.9. Let $\mathcal{C}$ be a category and $\mathcal{J} \in \text{Obj}(\mathbf{Cat})$.

1. Suppose that $\mathcal{C}$ has all $\mathcal{J}$ -limits. Let $L_1, L_2 \in \text{limChoice}(\mathcal{C}^{\mathcal{J}})$. Then $f^{L_1, L_2} \in \text{Iso}_{\mathcal{C}^{\mathcal{C}^{\mathcal{J}}}}(\lim_{L_1}, \lim_{L_2})$.

2. something similar for colimits.

Proof.

1. • Let $D \in \text{Obj}(\mathcal{C}^{\mathcal{J}})$. Write $L_1(D) = (X, \phi)$ and $L_2(D) = (Y, \psi)$. By Exercise 9.4.0.9 and Definition 9.4.0.12,

$$\begin{aligned} f_D^{L_1, L_2} &\in \text{Hom}_{\text{Cone}(D)}((L_1(D), L_2(D))) \\ &\subset \text{Hom}_{\mathcal{C}}(X, Y) \quad (\text{by ref ex cone def}) \\ &= \text{Hom}_{\mathcal{C}}(\lim_{L_1}(D), \lim_{L_2}(D)). \end{aligned}$$

Since $D \in \text{Obj}(\mathcal{C}^{\mathcal{J}})$ is arbitrary, we have that for each $D \in \text{Obj}(\mathcal{C}^{\mathcal{J}})$, $f_D^{L_1, L_2} \in \text{Hom}_{\mathcal{C}}(\lim_{L_1}(D), \lim_{L_2}(D))$.

• Let $D, E \in \text{Obj}(\mathcal{C}^{\mathcal{J}})$ and $\alpha \in \text{Hom}_{\mathcal{C}^{\mathcal{J}}}(D, E)$. Write $L_1(D) = (X^D, \phi^D)$, $L_1(E) = (X^E, \phi^E)$, $L_2(D) = (Y^D, \psi^D)$, $L_2(E) = (Y^E, \psi^E)$. Exercise 9.1.0.10 implies that $\text{Cn}_{\alpha} : \text{Cone}(D) \rightarrow \text{Cone}(E)$, $\text{Cn}_{\alpha}(L_1(D)), \text{Cn}_{\alpha}(L_2(D)) \in \text{Obj}(\text{Cone}(E))$ and

$$\begin{aligned} f_D^{L_1, L_2} &= \text{Cn}_{\alpha}(f_D^{L_1, L_2}) \\ &\in \text{Hom}_{\text{Cone}(E)}(\text{Cn}_{\alpha}(L_1(D)), \text{Cn}_{\alpha}(L_2(D))) \\ &= \text{Hom}_{\text{Cone}(E)}((X^D, \alpha \circ_{\mathcal{C}^{\mathcal{J}}} \phi^D), (Y^D, \alpha \circ_{\mathcal{C}^{\mathcal{J}}} \psi^D)) \\ &\subset \text{Hom}_{\mathcal{C}}(X^D, Y^D) \\ &= \text{Hom}_{\mathcal{C}}(\lim_{L_1}(D), \lim_{L_2}(D)). \end{aligned}$$

By Definition 9.4.0.5,

$$\begin{aligned} \eta_{\alpha}^{L_1} &= \lim_{L_1}(\alpha) \\ &\in \text{Hom}_{\text{Cone}(E)}(\text{Cn}_{\alpha}(L_1(D)), L_1(E)) \\ &\subset \text{Hom}_{\mathcal{C}}(X^D, X^E). \end{aligned}$$

and

$$\begin{aligned} \eta_{\alpha}^{L_2} &= \lim_{L_2}(\alpha) \\ &\in \text{Hom}_{\text{Cone}(E)}(\text{Cn}_{\alpha}(L_2(D)), L_2(E)) \\ &\subset \text{Hom}_{\mathcal{C}}(Y^D, Y^E). \end{aligned}$$

Since $L_2(E) \in \text{Term}_{\text{Cone}(E)}$, there exists a unique $g \in \text{Hom}_{\text{Cone}(E)}(\text{Cn}_{\alpha}(L_1(D)), L_2(E))$. Thus

$$\begin{aligned} \lim_{L_2}(\alpha) \circ_{\mathcal{C}} f_D^{L_1, L_2} &= \eta_{\alpha}^{L_2} \circ_{\mathcal{C}} f_D^{L_1, L_2} \\ &\in \text{Hom}_{\text{Cone}(E)}(\text{Cn}_{\alpha}(L_1(D)), L_2(E)) \\ &= \{g\} \end{aligned}$$

and similarly

$$\begin{aligned} f_E^{L_1, L_2} \circ_{\mathcal{C}} \lim_{L_1}(\alpha) &= f_E^{L_1, L_2} \circ_{\mathcal{C}} \eta_{\alpha}^{L_1} \\ &\in \text{Hom}_{\text{Cone}(E)}(\text{Cn}_{\alpha}(L_1(D)), L_2(E)) \\ &\in \{g\}. \end{aligned}$$

Thus

$$\begin{aligned} \lim_{L_2}(\alpha) \circ_{\mathcal{C}} f_D^{L_1, L_2} &= g \\ &= f_E^{L_1, L_2} \circ_{\mathcal{C}} \lim_{L_1}(\alpha), \end{aligned}$$

i.e. the following diagrams commute:

$$\begin{array}{ccc} \lim_{L_1}(D) & \xrightarrow{f_D^{L_1, L_2}} & \lim_{L_2}(D) \\ \lim_{L_2}(\alpha) \downarrow & & \downarrow \lim_{L_2}(\alpha) \\ \lim_{L_1}(E) & \xrightarrow{f_E^{L_1, L_2}} & \lim_{L_2}(E) \end{array} = \begin{array}{ccc} X^D & \xrightarrow{f_D^{L_1, L_2}} & Y^D \\ \eta_\alpha^{L_1} \downarrow & & \downarrow \eta_\alpha^{L_2} \\ X^E & \xrightarrow{f_E^{L_1, L_2}} & Y^E \end{array}$$

Since $D, E \in \text{Obj}(\mathcal{C}^{\mathcal{J}})$ and $\alpha \in \text{Hom}_{\mathcal{C}^{\mathcal{J}}}(D, E)$ are arbitrary, we have that for each $D, E \in \text{Obj}(\mathcal{C}^{\mathcal{J}})$ and $\alpha \in \text{Hom}_{\mathcal{C}^{\mathcal{J}}}(D, E)$, $\lim_{L_2}(\alpha) \circ_{\mathcal{C}} f_D^{L_1, L_2} = f_E^{L_1, L_2} \circ_{\mathcal{C}} \lim_{L_1}(\alpha)$. Thus $f^{L_1, L_2} \in \text{Hom}_{\mathcal{C}^{\mathcal{C}^{\mathcal{J}}}}(\lim_{L_1}, \lim_{L_2})$.

- Since for each $D \in \text{Obj}(\mathcal{C}^{\mathcal{J}})$,

$$\begin{aligned} f_D^{L_1, L_2} &\in \text{Hom}_{\text{Cone}(D)}((L_1(D), L_2(D))) \\ &= \text{Iso}_{\text{Cone}(D)}((L_1(D), L_2(D))) \\ &\subset \text{Iso}_{\mathcal{C}}(\lim_{L_1}(D), \lim_{L_2}(D)) \quad (\text{by make/ref ex in comma cat section}), \end{aligned}$$

we have that $f^{L_1, L_2} \in \text{Iso}_{\mathcal{C}^{\mathcal{C}^{\mathcal{J}}}}(\lim_{L_1}, \lim_{L_2})$.

2. colimits

□

Note 9.4.0.10. now can define $\times^L : \mathcal{C} \times \mathcal{C} \rightarrow \mathcal{C}$ and other functors as a restriction of $\lim$ to a subcategory of $\mathcal{C}^{\mathcal{J}}$ which are embeddings of $\mathcal{J}$, (like when $\mathcal{J}$ is just two dots, i.e. the product diagram) and explain people usually drop the L .

Definition 9.4.0.11. Let $\mathcal{C}$ be a category. Then $\mathcal{C}$ is said to be

- **complete** if for each $J \in \text{Obj}(\mathbf{Cat})$, $\mathcal{C}$ has all $\mathcal{J}$ -limits.
- **cocomplete** if for each $J \in \text{Obj}(\mathbf{Cat})$, $\mathcal{C}$ has all $\mathcal{J}$ -colimits.

Part III

Basic Results

Chapter 10

Yoneda Lemma

10.1 Representable Functors

Exercise 10.1.0.1. Let $\mathcal{C}$ be a category, $(A_1, A_2), (B_1, B_2) \in \text{Obj}(\mathcal{C}^{\text{op}} \times \mathcal{C})$ and $(f_1, f_2) \in \text{Hom}_{\mathcal{C}^{\text{op}} \times \mathcal{C}}((A_1, A_2), (B_1, B_2))$. Then for each $h \in \text{Hom}_{\mathcal{C}}(A_1, A_2)$, $f_2 \circ_{\mathcal{C}} h \circ_{\mathcal{C}} f_1 \in \text{Hom}_{\mathcal{C}}(B_1, B_2)$.

Proof. By definition of $\mathcal{C}^{\text{op}} \times \mathcal{C}$, we have that

•

$$\begin{aligned} A_1, B_1 &\in \text{Obj}(\mathcal{C}^{\text{op}}) \\ &= \text{Obj}(\mathcal{C}), \end{aligned}$$

$$\begin{aligned} f_1 &\in \text{Hom}_{\mathcal{C}^{\text{op}}}(A_1, B_1) \\ &= \text{Hom}_{\mathcal{C}}(B_1, A_1), \end{aligned}$$

- $A_2, B_2 \in \text{Obj}(\mathcal{C})$ and $f_2 \in \text{Hom}_{\mathcal{C}}(A_2, B_2)$.

Therefore, for each $h \in \text{Hom}_{\mathcal{C}}(A_1, A_2)$, $f_2 \circ_{\mathcal{C}} h \circ_{\mathcal{C}} f_1 \in \text{Hom}_{\mathcal{C}}(B_1, B_2)$. □

Note 10.1.0.2. If $\mathcal{C}$ is locally small, then for each $(A_1, A_2) \in \text{Obj}(\mathcal{C}^{\text{op}} \times \mathcal{C})$, $\text{Hom}_{\mathcal{C}}(A_1, A_2) \in \text{Obj}(\mathbf{Set})$.

Definition 10.1.0.3. Let $\mathcal{C}$ be a locally small category. We define the **Hom-functor of $\mathcal{C}$** , denoted $\text{Hom}_{\mathcal{C}}(\cdot, \cdot) : \mathcal{C}^{\text{op}} \times \mathcal{C} \rightarrow \mathbf{Set}$, by

- $\text{Hom}_{\mathcal{C}}(\cdot, \cdot)(A_1, A_2) := \text{Hom}_{\mathcal{C}}(A_1, A_2), \quad (A_1, A_2) \in \text{Obj}(\mathcal{C}^{\text{op}} \times \mathcal{C})$
- $\text{Hom}_{\mathcal{C}}(\cdot, \cdot)(f_1, f_2)(h) := f_2 \circ_{\mathcal{C}} h \circ_{\mathcal{C}} f_1, \quad (f_1, f_2) \in \text{Hom}_{\mathcal{C}^{\text{op}} \times \mathcal{C}}((A_1, A_2), (B_1, B_2)), h \in \text{Hom}_{\mathcal{C}}(A_1, A_2)$

Exercise 10.1.0.4. Let $\mathcal{C}$ be a locally small category. Then $\text{Hom}_{\mathcal{C}} : \mathcal{C}^{\text{op}} \times \mathcal{C} \rightarrow \mathbf{Set}$ is a functor.

Proof. Set $H := \text{Hom}_{\mathcal{C}}(\cdot, \cdot)$.

1. **The previous ex** implies that for each $(A_1, A_2), (B_1, B_2) \in \text{Obj}(\mathcal{C}^{\text{op}} \times \mathcal{C})$ and $(f_1, f_2) \in \text{Hom}_{\mathcal{C}^{\text{op}} \times \mathcal{C}}((A_1, A_2), (B_1, B_2))$,

$$\begin{aligned} H(f_1, f_2) &\in \text{Hom}_{\mathbf{Set}}(\text{Hom}_{\mathcal{C}}(A_1, A_2), \text{Hom}_{\mathcal{C}}(B_1, B_2)) \\ &= \text{Hom}_{\mathbf{Set}}(H(A_1, A_2), H(B_1, B_2)) \end{aligned}$$

2. Let $(A_1, A_2), (B_1, B_2), (C_1, C_2) \in \text{Obj}(\mathcal{C}^{\text{op}} \times \mathcal{C})$, $(f_1, f_2) \in \text{Hom}_{\mathcal{C}^{\text{op}} \times \mathcal{C}}((A_1, A_2), (B_1, B_2))$ and $(g_1, g_2) \in \text{Hom}_{\mathcal{C}^{\text{op}} \times \mathcal{C}}((B_1, B_2), (C_1, C_2))$. Then for each $h \in \text{Hom}_{\mathcal{C}}(A_1, A_2)$, we have that

$$\begin{aligned}
 H((g_1, g_2) \circ_{\mathcal{C}^{\text{op}} \times \mathcal{C}} (f_1, f_2))(h) &= H(g_1 \circ_{\mathcal{C}^{\text{op}}} f_1, g_2 \circ_{\mathcal{C}} f_2)(h) \\
 &= H(f_1 \circ_{\mathcal{C}} g_1, g_2 \circ_{\mathcal{C}} f_2)(h) \\
 &= (g_2 \circ_{\mathcal{C}} f_2) \circ_{\mathcal{C}} h \circ_{\mathcal{C}} (f_1 \circ_{\mathcal{C}} g_1) \\
 &= g_2 \circ_{\mathcal{C}} (f_2 \circ_{\mathcal{C}} h \circ_{\mathcal{C}} f_1) \circ_{\mathcal{C}} g_1 \\
 &= g_2 \circ_{\mathcal{C}} [H(f_1, f_2)(h)] \circ_{\mathcal{C}} g_1 \\
 &= H(g_1, g_2)(H(f_1, f_2)(h)) \\
 &= H(g_1, g_2) \circ_{\mathbf{Set}} H(f_1, f_2)(h).
 \end{aligned}$$

Thus $H((g_1, g_2) \circ_{\mathcal{C}^{\text{op}} \times \mathcal{C}} (f_1, f_2)) = H(g_1, g_2) \circ_{\mathbf{Set}} H(f_1, f_2)$. Since $(A_1, A_2), (B_1, B_2), (C_1, C_2) \in \text{Obj}(\mathcal{C}^{\text{op}} \times \mathcal{C})$, $(f_1, f_2) \in \text{Hom}_{\mathcal{C}^{\text{op}} \times \mathcal{C}}((A_1, A_2), (B_1, B_2))$ and $(g_1, g_2) \in \text{Hom}_{\mathcal{C}^{\text{op}} \times \mathcal{C}}((B_1, B_2), (C_1, C_2))$ are arbitrary, we have that for each $(A_1, A_2), (B_1, B_2), (C_1, C_2) \in \text{Obj}(\mathcal{C}^{\text{op}} \times \mathcal{C})$, $(f_1, f_2) \in \text{Hom}_{\mathcal{C}^{\text{op}} \times \mathcal{C}}((A_1, A_2), (B_1, B_2))$ and $(g_1, g_2) \in \text{Hom}_{\mathcal{C}^{\text{op}} \times \mathcal{C}}((B_1, B_2), (C_1, C_2))$, $H((g_1, g_2) \circ_{\mathcal{C}^{\text{op}} \times \mathcal{C}} (f_1, f_2)) = H(g_1, g_2) \circ_{\mathbf{Set}} H(f_1, f_2)$.

3. Let $(A_1, A_2) \in \text{Obj}(\mathcal{C}^{\text{op}} \times \mathcal{C})$. Then for each $h \in \text{Hom}_{\mathcal{C}}(A_1, A_2)$,

$$\begin{aligned}
 H(\text{id}_{(A_1, A_2)}^{\mathcal{C}^{\text{op}} \times \mathcal{C}})(h) &= H(\text{id}_{A_1}^{\mathcal{C}^{\text{op}}}, \text{id}_{A_2}^{\mathcal{C}})(h) \\
 &= H(\text{id}_{A_1}^{\mathcal{C}}, \text{id}_{A_2}^{\mathcal{C}})(h) \\
 &= \text{id}_{A_2}^{\mathcal{C}} \circ_{\mathcal{C}} h \circ_{\mathcal{C}} \text{id}_{A_1}^{\mathcal{C}} \\
 &= h \\
 &= \text{id}_{\text{Hom}_{\mathcal{C}}(A_1, A_2)}^{\mathbf{Set}}(h) \\
 &= \text{id}_{H(A_1, A_2)}^{\mathbf{Set}}(h).
 \end{aligned}$$

Thus $H(\text{id}_{(A_1, A_2)}^{\mathcal{C}^{\text{op}} \times \mathcal{C}}) = \text{id}_{H(A_1, A_2)}^{\mathbf{Set}}$. Since $(A_1, A_2) \in \text{Obj}(\mathcal{C}^{\text{op}} \times \mathcal{C})$ is arbitrary, we have that for each $(A_1, A_2) \in \text{Obj}(\mathcal{C}^{\text{op}} \times \mathcal{C})$, $\text{Hom}_{\mathcal{C}}(\text{id}_{(A_1, A_2)}^{\mathcal{C}^{\text{op}} \times \mathcal{C}}) = \text{id}_{H(A_1, A_2)}^{\mathbf{Set}}$.

Hence $\text{Hom}_{\mathcal{C}} : \mathcal{C}^{\text{op}} \times \mathcal{C} \rightarrow \mathbf{Set}$ is a functor. □

Note 10.1.0.5. If $\mathcal{C}$ is locally small, Exercise 8.2.2.2 implies that for each $X \in \text{Obj}(\mathcal{C})$, $\text{Hom}_{\mathcal{C}}(X, \cdot) : \mathcal{C} \rightarrow \mathbf{Set}$ and $\text{Hom}_{\mathcal{C}}(\cdot, X) : \mathcal{C}^{\text{op}} \rightarrow \mathbf{Set}$ are functors.

Exercise 10.1.0.6. Let $\mathcal{C}$ be a category. Suppose that $\mathcal{C}$ is locally small. Then

1. for each $X \in \text{Obj}(\mathcal{C}^{\text{op}})$, $\text{Hom}_{\mathcal{C}^{\text{op}}}(X, \cdot) = \text{Hom}_{\mathcal{C}}(\cdot, X)$
2. for each $X \in \text{Obj}(\mathcal{C})$, $\text{Hom}_{\mathcal{C}}(X, \cdot) = \text{Hom}_{\mathcal{C}^{\text{op}}}(\cdot, X)$

Proof. We note that Exercise 10.1.0.10 (ref ex) implies that $(\mathcal{C}^{\text{op}})^{\text{op}} = \mathcal{C}$. So $(\mathcal{C}^{\text{op}})^{\text{op}} \times \mathcal{C}^{\text{op}} = \mathcal{C} \times \mathcal{C}^{\text{op}}$.

1. Let $X \in \text{Obj}(\mathcal{C}^{\text{op}})$. Then $X \in \text{Obj}(\mathcal{C})$.

- Let $A \in \text{Obj}(\mathcal{C}^{\text{op}})$. Then $A \in \text{Obj}(\mathcal{C})$ and

$$\begin{aligned}
 \text{Hom}_{\mathcal{C}^{\text{op}}}(X, \cdot)(A) &= \text{Hom}_{\mathcal{C}^{\text{op}}}(\cdot, \cdot)(X, A) \\
 &= \text{Hom}_{\mathcal{C}^{\text{op}}}(X, A) \\
 &= \text{Hom}_{\mathcal{C}}(A, X) \\
 &= \text{Hom}_{\mathcal{C}}(\cdot, \cdot)(A, X) \\
 &= \text{Hom}_{\mathcal{C}}(\cdot, X)(A).
 \end{aligned}$$

- Let $A, B \in \text{Obj}(\mathcal{C}^{\text{op}})$ and $f \in \text{Hom}_{\mathcal{C}^{\text{op}}}(A, B)$ and $h \in \text{Hom}_{\mathcal{C}^{\text{op}}}(X, A)$. Then $A, B \in \text{Obj}(\mathcal{C})$, $f \in \text{Hom}_{\mathcal{C}}(B, A)$ and $h \in \text{Hom}_{\mathcal{C}}(A, X)$. Thus $(\text{id}_X^{\mathcal{C}}, f) \in \text{Hom}_{\mathcal{C} \times \mathcal{C}^{\text{op}}}((X, A), (X, B))$ and

$$\begin{aligned} \text{Hom}_{\mathcal{C}^{\text{op}}}(X, \cdot)(f)(h) &= \text{Hom}_{\mathcal{C}^{\text{op}}}(\cdot, \cdot)(\text{id}_X^{\mathcal{C}}, f)(h) \\ &= f \circ_{\mathcal{C}^{\text{op}}} h \circ_{\mathcal{C}^{\text{op}}} \text{id}_X^{\mathcal{C}} \\ &= \text{id}_X^{\mathcal{C}} \circ_{\mathcal{C}} h \circ_{\mathcal{C}} f \\ &= \text{Hom}_{\mathcal{C}}(\cdot, \cdot)(f, \text{id}_X^{\mathcal{C}})(h) \\ &= \text{Hom}_{\mathcal{C}}(\cdot, X)(f)(h) \end{aligned}$$

Since $h \in \text{Hom}_{\mathcal{C}^{\text{op}}}(X, A)$ is arbitrary, we have that for each $h \in \text{Hom}_{\mathcal{C}^{\text{op}}}(X, A)$, $\text{Hom}_{\mathcal{C}^{\text{op}}}(X, \cdot)(f)(h) = \text{Hom}_{\mathcal{C}}(\cdot, X)(f)(h)$. Thus $\text{Hom}_{\mathcal{C}^{\text{op}}}(X, \cdot)(f) = \text{Hom}_{\mathcal{C}}(\cdot, X)(f)$.

Hence $\text{Hom}_{\mathcal{C}^{\text{op}}}(X, \cdot) = \text{Hom}_{\mathcal{C}}(\cdot, X)$.

2. **FINISH!!!** (similar)

□

Definition 10.1.0.7. Let $\mathcal{C}$ be a category. Suppose that $\mathcal{C}$ is locally small. Let $F : \mathcal{C} \rightarrow \mathbf{Set}$.

- Let $X \in \text{Obj}(\mathcal{C})$ and $\phi : F \Rightarrow \text{Hom}_{\mathcal{C}}(X, \cdot)$. Then (X, ϕ) is said to be a **representation of F** if ϕ is a natural isomorphism.
- Then F is said to be **representable** if there exists $X \in \text{Obj}(\mathcal{C})$ and $\phi : F \Rightarrow \text{Hom}_{\mathcal{C}}(X, \cdot)$ such that (X, ϕ) is a representation of F .

Exercise 10.1.0.8. Let $\mathcal{C}$ be a category. Suppose that $\mathcal{C}$ is small. Let $X \in \text{Obj}(\mathcal{C})$. Let $1 \in \text{Obj}(\mathbf{Set})$. Suppose that 1 is **Set**-terminal. Then

1. X is $\mathcal{C}$ -initial iff $\text{Hom}_{\mathcal{C}}(X, \cdot) \cong_{\mathbf{Set}^{\mathcal{C}}} \Delta_1^{\mathcal{C}}$.
2. X is $\mathcal{C}$ -terminal iff $\text{Hom}_{\mathcal{C}^{\text{op}}}(\cdot, X) \cong_{\mathbf{Set}^{\mathcal{C}^{\text{op}}}} \Delta_1^{\mathcal{C}^{\text{op}}}$.

Proof. Define $F \in \mathbf{Set}^{\mathcal{C}}$ by $F := \text{Hom}_{\mathcal{C}}(X, \cdot)$.

1. Exercise 7.1.0.11 implies that $\phi^{(F,1)} \in \text{Hom}_{\mathbf{Set}^{\mathcal{C}}}(F, \Delta_1^{\mathcal{C}})$. Exercise 7.1.0.6 implies that there exists $x \in 1$ such that $1 = \{x\}$.

- ($\implies$):
Suppose that X is $\mathcal{C}$ -initial. Let $A \in \text{Obj}(\mathcal{C})$. Since $F(A) = \text{Hom}_{\mathcal{C}}(X, A)$, Exercise 7.1.0.7 implies that $F(A)$ is **Set**-terminal. Since 1 is **Set**-terminal, Exercise 7.1.0.4 implies that $F(A) \cong_{\mathbf{Set}} 1$ and $\text{Hom}_{\mathbf{Set}}(F(A), 1) = \text{Iso}_{\mathbf{Set}}(F(A), 1)$. Therefore

$$\begin{aligned} \phi_A^{(F,1)} &\in \text{Hom}_{\mathbf{Set}}(F(A), 1) \\ &= \text{Iso}_{\mathbf{Set}}(F(A), 1) \\ &= \text{Iso}_{\mathbf{Set}}(F(A), \Delta_1^{\mathcal{C}}(A)). \end{aligned}$$

Since $A \in \text{Obj}(\mathcal{C})$ is arbitrary, we have that for each $A \in \text{Obj}(\mathcal{C})$, $\phi_A^{(F,1)} \in \text{Iso}_{\mathbf{Set}}(F(A), \Delta_1^{\mathcal{C}}(A))$. Thus $\phi^{(F,1)} : F \Rightarrow \Delta_1^{\mathcal{C}}$ is a natural isomorphism. Exercise 6.3.0.5 then implies that $\phi^{(F,1)} \in \text{Iso}_{\mathbf{Set}^{\mathcal{C}}}(F, \Delta_1^{\mathcal{C}})$. Since $\text{Iso}_{\mathbf{Set}^{\mathcal{C}}}(F, \Delta_1^{\mathcal{C}}) \neq \emptyset$, we have that $F \cong_{\mathbf{Set}^{\mathcal{C}}} \Delta_1^{\mathcal{C}}$.

- ($\impliedby$):
Suppose that $F \cong_{\mathbf{Set}^{\mathcal{C}}} \Delta_1^{\mathcal{C}}$. Then there exists $\phi \in \text{Iso}_{\mathbf{Set}^{\mathcal{C}}}(F, \Delta_1^{\mathcal{C}})$. Exercise 6.3.0.5 then implies that $\phi : F \Rightarrow \Delta_1^{\mathcal{C}}$ is a natural isomorphism. Let $A \in \text{Obj}(\mathcal{C})$. Then

$$\begin{aligned} \phi_A &\in \text{Iso}_{\mathbf{Set}}(F(A), \Delta_1^{\mathcal{C}}(A)) \\ &= \text{Iso}_{\mathbf{Set}}(F(A), 1). \end{aligned}$$

Since $\text{Iso}_{\mathbf{Set}}(F(A), 1) \neq \emptyset$ and 1 is **Set**-terminal, we have that

$$\begin{aligned} \text{Hom}_{\mathcal{C}}(X, A) &= F(A) \\ &\cong_{\mathbf{Set}} 1 \\ &\cong_{\mathbf{Set}} \{\emptyset\}, \quad (\text{by Exercise 7.1.0.6}). \end{aligned}$$

Thus there exists $f \in \text{Hom}_{\mathcal{C}}(X, A)$ such that $\text{Hom}_{\mathcal{C}}(X, A) = \{f\}$. Since $A \in \text{Obj}(\mathcal{X})$ is arbitrary, we have that for each $A \in \text{Obj}(\mathcal{C})$, there exists $f \in \text{Hom}_{\mathcal{C}}(X, A)$ such that $\text{Hom}_{\mathcal{C}}(X, A) = \{f\}$. Hence X is $\mathcal{C}$ -initial.

2. We have that

$$\begin{aligned} X \text{ is } \mathcal{C}\text{-terminal} &\iff X \text{ is } \mathcal{C}^{\text{op}}\text{-initial}, \quad (\text{by Exercise 7.1.0.3}) \\ &\iff \text{Hom}_{\mathcal{C}^{\text{op}}}(X, \cdot) \cong_{\mathbf{Set}^{\mathcal{C}^{\text{op}}}} \Delta_1^{\mathcal{C}^{\text{op}}}, \quad (\text{by (ref) prev part}) \\ &\iff \text{Hom}_{\mathcal{C}}(\cdot, X) \cong_{\mathbf{Set}^{\mathcal{C}^{\text{op}}}} \Delta_1^{\mathcal{C}^{\text{op}}}, \quad (\text{by (ref) prev ex about } \text{Hom}_{\mathcal{C}}(\cdot, X) = \text{Hom}_{\mathcal{C}^{\text{op}}}(X, \cdot)) \end{aligned}$$

□

Exercise 10.1.0.9. Let $\mathcal{C}$ be a category, $F : \mathcal{C} \rightarrow \mathbf{Set}$, $X \in \text{Obj}(\mathbf{Set})$ and $(A, f) \in (X \downarrow F)$. Suppose that $\mathcal{C}$ is locally small, X is **Set**-terminal and (A, f) is a universal morphism from X to F . Then

relevant exercises

Exercise 8.3.0.2

Exercise 8.5.0.12

Exercise 10.1.0.10. Yoneda Lemma: Let $\mathcal{C}$ be a category. Suppose that $\mathcal{C}$ is locally small. Then

$$\Rightarrow \text{eval}_{(\mathcal{C}, \mathbf{Set})}$$

10.2 Fundamental Examples

look up applications of yoneda lemma to analysis. For example, see Tom LaGatta's answer on [stackexchange](#) and look up yoneda lemma construction of dedekind cuts for posets.

Chapter 11

Adjunctions

Exercise 11.0.0.1. Let

Definition 11.0.0.2. Let $\mathcal{C}, \mathcal{D}$ be categories, $F : \mathcal{C} \rightarrow \mathcal{D}$ and $G : \mathcal{D} \rightarrow \mathcal{C}$. Then (F, G) is said to be an **adjunction** if

Chapter 12

Monads

12.1 Introduction

Definition 12.1.0.1. Let $\mathcal{C} \in \text{Obj}(\mathbf{Cat})$, $T \in \text{End}_{\mathbf{Cat}}(X)$, $\eta : \text{id}_{\mathcal{C}} \Rightarrow T$ and $\mu : T^2 \Rightarrow T$. Then (T, η, μ) is said to be a **monad on $\mathcal{C}$** if

1. $\mu \circ T\mu = \mu \circ \mu T$
2. $\mu \circ T\eta = \text{id}_T$ and $\mu \circ \eta T = \text{id}_T$,

i.e. the following diagrams commute:

$$\begin{array}{ccc} T^3 & \xrightarrow{T\mu} & T^2 \\ \mu T \downarrow & & \downarrow \mu \\ T^2 & \xrightarrow{\mu} & T \end{array} \quad \begin{array}{ccc} T & \xrightarrow{\eta T} & T^2 \\ T\eta \downarrow & \searrow \text{id}_T & \downarrow \mu \\ T^2 & \xrightarrow{\mu} & T \end{array}$$

12.2 TO DO

- Define subcategories and full subcategories and show that if $\text{Obj}(D) \subset \text{Obj}(C)$ and for each $X, Y \in \text{Obj}(D)$, $\text{Hom}_D(X, Y) = \text{Hom}_C(X, Y)$, then D is a full subcategory of C . I used this in differential
- discuss projective/inductive systems and the projective/inductive limits and applications to topology and measure theory

Chapter 13

Monoidal Categories

Definition 13.0.0.1. Let $\mathcal{C}$ be a category, $\otimes : \mathcal{C} \times \mathcal{C} \rightarrow \mathcal{C}$, $\rho : (- \otimes -) \otimes \cdot \Rightarrow - \otimes (- \otimes -)$.

Chapter 14

To Do

- look into the schroder bernstein property [wikipedia](#)

Appendix A

App

A.1 Reading Diagrams and associated digraphs of diagrams

Definition A.1.0.1. Let

$$\begin{array}{ccc} C & \xrightarrow{g} & A \\ h \downarrow & & \downarrow f \\ A & \xrightarrow{f} & B \end{array} \implies \begin{array}{ccc} & \xrightarrow{g} & \\ C & & A \\ & \xleftarrow{h} & \end{array}$$

see an intro to the language of category theory by roman for description

Definition A.1.0.2. A diagram is said to be **commutative** if for each path of length ≥ 2 , in the associated digraph gives the same morphism.

