

Introduction to Order Theory

Carson James

Contents

Notation	ix
Preface	1
I Sets	3
1 Relations	5
1.1 Introduction	5
1.2 Homogeneous Relations	6
1.2.1 Subrelations	6
1.2.2 Product Relations	6
1.3 Binary Relations	7
1.3.1 Subrelations	10
1.3.2 Products of Relations	11
1.3.3 Subrelations	12
2 Operations	13
3 Equivalence Relations	15
3.1 Introduction	15
3.2 Products	20
3.3 Quotients	20
II Ordered Sets	27
3.4 To Do	29
4 Prosets	31
4.1 Introduction	31
4.2 Subprosets	32
4.3 Products	32
4.4 Quotients	33
5 Posets	35
5.1 Introduction	35
6 Directed Sets	37
6.1 Introduction	37
6.2 Subdirected Sets	38
6.3 Products of Directed Sets	39

7	Monotone Maps	41
7.1	Introduction	41
7.2	Galois Connections	44
8	Upper and Lower Sets	51
8.1	Introduction	51
8.2	Maximal Elements	62
8.3	Filters and Ideals	63
9	Suprema and Infima	65
9.1	Upper and Lower Bounds	65
9.2	SupSets and InfSets	70
9.3	Suprema and Infima	73
III	Lattices	79
10	Lattices	83
10.1	Semilattices	83
10.1.1	Skew Semilattices	83
10.1.2	Skew Semilattice Homomorphisms	85
10.1.3	Semilattices	86
10.1.4	Semilattice Homomorphisms	90
10.1.5	Semilattice Structures	93
10.2	Lattices	93
10.2.1	Skew Lattices	93
10.2.2	Lattices	95
10.2.3	Lattice Homomorphisms	98
10.3	Fundamental Examples	99
10.3.1	Bounded, Modular and Distributive Lattices	99
10.3.2	Heyting Algebras	103
10.3.3	Boolean Algebras	105
10.4	Lattice Structures	106
10.4.1	Irreducibility and Primality	106
10.4.2	Ideals and Filters	107
10.4.3	Properties of Ideals and Filters	114
10.5	Constructions with Lattices	116
10.5.1	Sublattices	116
10.5.2	Product Lattices	117
10.5.3	Coproduct Lattices	117
10.5.4	Quotient Lattices	117
11	Complete Lattices	119
11.1	Introduction	119
11.2	Complete Lattice Homomorphisms	120
11.3	Properties of Ideals and Filters	123
11.3.1	Prime Ideals and Filters in Complete Lattices	123
11.4	Constructions with Complete Lattices	125
11.4.1	Complete Sublattices	125
11.4.2	Products of Complete Lattices	125
11.4.3	Coproducts of Complete Lattices	125
11.4.4	Quotients of Complete Lattices	125
11.5	Fundamental Examples	126
11.5.1	Introduction	126
11.5.2	Sets	126

11.5.3	Equivalence Relations	126
11.5.4	Ideal and Filter Lattices	126
11.5.5	Topological Spaces	126
11.6	Least Upper Bound Property	126
11.7	Complete Lattice Homomorphisms	126

IV Continuous Lattices

127

12 Scott

131

Notation

$\mathcal{M}_+(X, \mathcal{A})$	finite measures on $(X, \mathcal{A})$
v	velocity

Preface

cc-by-nc-sa

Part I

Sets

Chapter 1

Relations

1.1 Introduction

Definition 1.1.0.1. Let $n \in \mathbb{N}$ and $(A_j)_{j \in [n]}$ a collection of sets.

- Let $R \subset \prod_{j \in [n]} A_j$. Then R is said to be a **n -ary relation over $(A_j)_{j \in [n]}$** .
- We define $\mathcal{R}(A_j : j \in [n]) := \mathcal{P}(\prod_{j \in [n]} A_j)$

1.2 Homogeneous Relations

Definition 1.2.0.1. Let $n \in \mathbb{N}$ and A a set.

- Let $R \subset A^n$. Then R is said to be a **homogeneous n -ary relation on A**
- We define $\mathcal{R}^n(A) := \mathcal{P}(A^n)$.

1.2.1 Subrelations

Definition 1.2.1.1. Let $n \in \mathbb{N}$, A, B sets, $R \in \mathcal{R}^n(A)$ and $S \in \mathcal{R}^n(B)$. Then

- S is said to be a **restriction of R** if $B \subset A$,
- R is said to be an **extension of S** if S is a restriction of R .

Definition 1.2.1.2. Let $n \in \mathbb{N}$, A a set, $B \subset A$ and $R \in \mathcal{R}^n(A)$. We define the **restriction of R to B** , denoted $R|_B$, by $R|_B := R \cap B^n$.

Exercise 1.2.1.3. Let $n \in \mathbb{N}$, A a set, $B \subset A$ and $R \in \mathcal{R}^n(A)$. Then $R|_B$ is a restriction of R .

Proof. Clear. **fill out later**

□

1.2.2 Product Relations

1.3 Binary Relations

Definition 1.3.0.1. Let A, B be sets.

- Let $R \subset A \times B$. Then R is said to be a **relation over A and B** .
- We define $\mathcal{R}(A, B) := \mathcal{P}(A \times B)$

Definition 1.3.0.2. Let A, B, C be sets, $R \in \mathcal{R}(A, B)$, $S \in \mathcal{R}(B, C)$. We define the **composition relation of S with R** , denoted $S \circ_{\text{Rel}} R \in \mathcal{R}(A, C)$, by

$$S \circ_{\text{Rel}} R := \{(a, c) \in A \times C : \text{there exists } b \in B \text{ such that } (a, b) \in R \text{ and } (b, c) \in S\}.$$

Note 1.3.0.3. When the context is clear, we write $S \circ R$ in place of $S \circ_{\text{Rel}} R$.

Exercise 1.3.0.4. Let A, B, C, D be sets, R a relation between A and B , S relation between B and C and T relation between C and D . Then $(T \circ S) \circ R = T \circ (S \circ R)$.

Proof.

- Let $(a, d) \in (T \circ S) \circ R$. Then there exists $b \in B$ such that $(a, b) \in R$ and $(b, d) \in T \circ S$. Since $(b, d) \in T \circ S$, there exists $c \in C$ such that $(b, c) \in S$ and $(c, d) \in T$. Since $(a, b) \in R$ and $(b, c) \in S$, $(a, c) \in S \circ R$. Since $(a, c) \in S \circ R$ and $(c, d) \in T$, we have that $(a, d) \in T \circ (S \circ R)$. Since $(a, d) \in (T \circ S) \circ R$ is arbitrary, we have that for each $(a, d) \in (T \circ S) \circ R$, $(a, d) \in T \circ (S \circ R)$. Hence $(T \circ S) \circ R \subset T \circ (S \circ R)$.
- Similarly, $T \circ (S \circ R) \subset (T \circ S) \circ R$.

Since $(T \circ S) \circ R \subset T \circ (S \circ R)$ and $T \circ (S \circ R) \subset (T \circ S) \circ R$, we have that $(T \circ S) \circ R = T \circ (S \circ R)$. □

Definition 1.3.0.5. Let A be a set. We define the **diagonal relation on A** , denoted $\Delta_A \in \mathcal{R}(A, A)$, by

$$\Delta_A := \{(a, a) : a \in A\}.$$

Exercise 1.3.0.6. Let A, B be sets and $R \in \mathcal{R}(A, B)$. Then

1. $R \circ \Delta_A = R$
2. $\Delta_B \circ R = R$

Proof.

1. • Let $(a, b) \in R \circ \Delta_A$. Then there exists $a' \in A$ such that $(a, a') \in \Delta_A$ and $(a', b) \in R$. By definition of Δ_A , $a = a'$. Therefore $(a, b) \in R$. Since $(a, b) \in R \circ \Delta_A$ is arbitrary, we have that for each $(a, b) \in R \circ \Delta_A$, $(a, b) \in R$. Hence $R \circ \Delta_A \subset R$.
 • Let $(a, b) \in R$. Then $a \in A$. Therefore $(a, a) \in \Delta_A$. Since $(a, a) \in \Delta_A$ and $(a, b) \in R$, we have that $(a, b) \in R \circ \Delta_A$. Since $(a, b) \in R$ is arbitrary, we have that for each $(a, b) \in R$, $(a, b) \in R \circ \Delta_A$. Hence $R \circ \Delta_A \supset R$.

Since $R \circ \Delta_A \subset R$ and $R \circ \Delta_A \supset R$, we have that $R \circ \Delta_A = R$.

2. Similar to (1). □

Exercise 1.3.0.7. Let A, B, C be sets.

1. Let $R \in \mathcal{R}(A, B)$ and $(S_\gamma)_{\gamma \in \Gamma} \subset \mathcal{R}(B, C)$. Then

$$(a) \left(\bigcup_{\gamma \in \Gamma} S_\gamma \right) \circ R = \bigcup_{\gamma \in \Gamma} (S_\gamma \circ R),$$

$$(b) \left(\bigcap_{\gamma \in \Gamma} S_\gamma \right) \circ R \subset \bigcap_{\gamma \in \Gamma} (S_\gamma \circ R)$$

2. Let $(R_\gamma)_{\gamma \in \Gamma} \subset \mathcal{R}(A, B)$ and $S \in \mathcal{R}(B, C)$. Then

$$(a) \quad S \circ \left(\bigcup_{\gamma \in \Gamma} R_\gamma \right) = \bigcup_{\gamma \in \Gamma} (S \circ R_\gamma).$$

$$(b) \quad S \circ \left(\bigcap_{\gamma \in \Gamma} R_\gamma \right) \subset \bigcap_{\gamma \in \Gamma} (S \circ R_\gamma).$$

Proof.

1. (a) We note that

$$\begin{aligned} (a, c) \in \left(\bigcup_{\gamma \in \Gamma} S_\gamma \right) \circ R &\iff \text{there exists } b \in B \text{ such that } (a, b) \in R \text{ and } (b, c) \in \bigcup_{\gamma \in \Gamma} S_\gamma \\ &\iff \text{there exists } b \in B \text{ and } \gamma_0 \in \Gamma \text{ such that } (a, b) \in R \text{ and } (b, c) \in S_{\gamma_0} \\ &\iff \text{there exists } \gamma_0 \in \Gamma \text{ such that } (a, c) \in S_{\gamma_0} \circ R \\ &\iff (a, c) \in \bigcup_{\gamma \in \Gamma} (S_\gamma \circ R). \end{aligned}$$

$$\text{Hence } \left(\bigcup_{\gamma \in \Gamma} S_\gamma \right) \circ R = \bigcup_{\gamma \in \Gamma} (S_\gamma \circ R).$$

(b) We note that

$$\begin{aligned} (a, c) \in \left(\bigcap_{\gamma \in \Gamma} S_\gamma \right) \circ R &\iff \text{there exists } b \in B \text{ such that } (a, b) \in R \text{ and } (b, c) \in \bigcap_{\gamma \in \Gamma} S_\gamma \\ &\iff \text{there exists } b \in B \text{ such that } (a, b) \in R \text{ and for each } \gamma \in \Gamma \text{ and } (b, c) \in S_\gamma \\ &\implies \text{for each } \gamma \in \Gamma, \text{ there exists } b \in B \text{ such that } (a, b) \in R \text{ and } (b, c) \in S_\gamma \\ &\iff \text{for each } \gamma \in \Gamma, (a, c) \in S_\gamma \circ R \\ &\iff (a, c) \in \bigcap_{\gamma \in \Gamma} (S_\gamma \circ R). \end{aligned}$$

$$\text{Hence } \left(\bigcap_{\gamma \in \Gamma} S_\gamma \right) \circ R = \bigcap_{\gamma \in \Gamma} (S_\gamma \circ R).$$

2. (a) We note that

$$\begin{aligned} (a, c) \in S \circ \left(\bigcup_{\gamma \in \Gamma} R_\gamma \right) &\iff \text{there exists } b \in B \text{ such that } (a, b) \in \bigcup_{\gamma \in \Gamma} R_\gamma \text{ and } (b, c) \in S \\ &\iff \text{there exists } b \in B \text{ and } \gamma_0 \in \Gamma \text{ such that } (a, b) \in R_{\gamma_0} \text{ and } (b, c) \in S \\ &\iff \text{there exists } \gamma_0 \in \Gamma \text{ such that } (a, c) \in S \circ R_{\gamma_0} \\ &\iff (a, c) \in \bigcup_{\gamma \in \Gamma} (S \circ R_\gamma). \end{aligned}$$

$$\text{Hence } S \circ \left(\bigcup_{\gamma \in \Gamma} R_\gamma \right) = \bigcup_{\gamma \in \Gamma} (S \circ R_\gamma).$$

(b) We note that

$$\begin{aligned} (a, c) \in S \circ \left(\bigcap_{\gamma \in \Gamma} R_\gamma \right) &\iff \text{there exists } b \in B \text{ such that } (a, b) \in \bigcap_{\gamma \in \Gamma} R_\gamma \text{ and } (b, c) \in S \\ &\iff \text{there exists } b \in B \text{ such that for each } \gamma \in \Gamma, (a, b) \in R_\gamma \text{ and } (b, c) \in S \\ &\implies \text{for each } \gamma \in \Gamma, \text{ there exists } b \in B \text{ such that } (a, b) \in R_\gamma \text{ and } (b, c) \in S \\ &\iff \text{for each } \gamma \in \Gamma, (a, c) \in S \circ R_\gamma \\ &\iff (a, c) \in \bigcap_{\gamma \in \Gamma} (S \circ R_\gamma). \end{aligned}$$

Hence $S \circ \left(\bigcap_{\gamma \in \Gamma} R_\gamma \right) \subset \bigcap_{\gamma \in \Gamma} (S_\gamma \circ R)$.

□

Exercise 1.3.0.8. Let A, B, C be sets.

1. Let $R_1, R_2 \in \mathcal{R}(A, B)$ and $S \in \mathcal{R}(B, C)$. If $R_1 \subset R_2$, then $S \circ R_1 \subset S \circ R_2$.
2. Let $R \in \mathcal{R}(A, B)$ and $S_1, S_2 \in \mathcal{R}(B, C)$. If $S_1 \subset S_2$, then $S_1 \circ R \subset S_2 \circ R$.
3. Let $R_1, R_2 \in \mathcal{R}(A, B)$ and $S_1, S_2 \in \mathcal{R}(B, C)$. If $R_1 \subset R_2$ and $S_1 \subset S_2$, then $S_1 \circ R_1 \subset S_2 \circ R_2$.

Proof.

1. Suppose that $R_1 \subset R_2$. Then

$$\begin{aligned} (a, c) \in S \circ R_1 &\iff \text{there exists } b \in B \text{ such that } (a, b) \in R_1 \text{ and } (b, c) \in S \\ &\implies \text{there exists } b \in B \text{ such that } (a, b) \in R_2 \text{ and } (b, c) \in S \\ &\iff (a, c) \in S \circ R_2. \end{aligned}$$

Hence $S \circ R_1 \subset S \circ R_2$.

2. Suppose that $S_1 \subset S_2$. Then

$$\begin{aligned} (a, c) \in S_1 \circ R &\iff \text{there exists } b \in B \text{ such that } (a, b) \in R \text{ and } (b, c) \in S_1 \\ &\implies \text{there exists } b \in B \text{ such that } (a, b) \in R \text{ and } (b, c) \in S_2 \\ &\iff (a, c) \in S_2 \circ R. \end{aligned}$$

Hence $S_2 \circ R \subset S_1 \circ R$.

3. Suppose that $R_1 \subset R_2$ and $S_1 \subset S_2$. Then **the previous parts** imply that

$$\begin{aligned} S_1 \circ R_1 &\subset S_2 \circ R_1 \\ &\subset S_2 \circ R_2. \end{aligned}$$

□

Definition 1.3.0.9. Let A, B be sets and $R \in \mathcal{R}(A, B)$. We define the **converse of** R , denoted $R^* \in \mathcal{R}(B, A)$, by

$$R^* := \{(b, a) \in B \times A : (a, b) \in R\}.$$

Exercise 1.3.0.10. Let A, B be sets and R a relation between A and B . Then $(R^*)^* = R$.

Proof.

$$\begin{aligned} (a, b) \in (R^*)^* &\iff (b, a) \in R^* \\ &\iff (a, b) \in R. \end{aligned}$$

Hence $(R^*)^* = R$.

□

Exercise 1.3.0.11. Let A, B, C be sets, $R \in \mathcal{R}(A, B)$ and $S \in \mathcal{R}(B, C)$. Then $(S \circ R)^* = R^* \circ S^*$.

Proof. We note that

$$\begin{aligned} (c, a) \in (S \circ R)^* &\iff (a, c) \in S \circ R \\ &\iff \text{there exists } b \in B \text{ such that } (a, b) \in R \text{ and } (b, c) \in S \\ &\iff \text{there exists } b \in B \text{ such that } (c, b) \in S^* \text{ and } (b, a) \in R^* \\ &\iff (c, a) \in R^* \circ S^*. \end{aligned}$$

Hence $(S \circ R)^* = R^* \circ S^*$.

□

Exercise 1.3.0.12. Let A and B be sets and $(R_\gamma)_{\gamma \in \Gamma} \subset \mathcal{R}(A, B)$. Then

1. $\left(\bigcup_{\gamma \in \Gamma} R_\gamma \right)^* = \bigcup_{\gamma \in \Gamma} R_\gamma^*$
2. $\left(\bigcap_{\gamma \in \Gamma} R_\gamma \right)^* = \bigcap_{\gamma \in \Gamma} R_\gamma^*$

Proof.

1. We note that

$$\begin{aligned}
 (b, a) \in \left(\bigcup_{\gamma \in \Gamma} R_\gamma \right)^* &\iff (a, b) \in \bigcup_{\gamma \in \Gamma} R_\gamma \\
 &\iff \text{there exists } \gamma_0 \in \Gamma \text{ such that } (a, b) \in R_{\gamma_0} \\
 &\iff \text{there exists } \gamma_0 \in \Gamma \text{ such that } (b, a) \in R_{\gamma_0}^* \\
 &\iff (b, a) \in \bigcup_{\gamma \in \Gamma} R_\gamma^*.
 \end{aligned}$$

$$\text{Hence } \left(\bigcup_{\gamma \in \Gamma} R_\gamma \right)^* = \bigcup_{\gamma \in \Gamma} R_\gamma^*.$$

2. We note that

$$\begin{aligned}
 (b, a) \in \left(\bigcap_{\gamma \in \Gamma} R_\gamma \right)^* &\iff (a, b) \in \bigcap_{\gamma \in \Gamma} R_\gamma \\
 &\iff \text{for each } \gamma \in \Gamma, (a, b) \in R_\gamma \\
 &\iff \text{for each } \gamma \in \Gamma, (b, a) \in R_\gamma^* \\
 &\iff (b, a) \in \bigcap_{\gamma \in \Gamma} R_\gamma^*.
 \end{aligned}$$

$$\text{Hence } \left(\bigcap_{\gamma \in \Gamma} R_\gamma \right)^* = \bigcap_{\gamma \in \Gamma} R_\gamma^*.$$

□

1.3.1 Subrelations

Definition 1.3.1.1. Let $n \in \mathbb{N}$, A, B, E, F sets, $R \in \mathcal{R}(A, B)$ and $S \in \mathcal{R}(E, F)$. Then

- S is said to be a **restriction of R** if $E \subset A$ and $F \subset B$,
- R is said to be an **extension of S** if S is a restriction of R .

Definition 1.3.1.2. Let A, B be sets, $E \subset A$, $F \subset B$ and $R \in \mathcal{R}(A, B)$. We define the **restriction of R over E and F** , denoted $R|_{E \times F} \in \mathcal{R}(E, F)$, by $R|_{E \times F} := R \cap (E \times F)$.

Exercise 1.3.1.3. Let A, B be sets, $E \subset A$, $F \subset B$ and $R \in \mathcal{R}(A, B)$. Then $R|_B$ is a restriction of R .

Proof. Clear. fill out later

□

maybe dont do this subsection, since relations are like morphisms, so unless you can find a categorical way to present restrictions on domain or codomain $f|_E$ or $f|_F$, just get rid of this subsection

Exercise 1.3.1.4. Let A, B be sets, $E \subset A$, $F \subset B$ and $R \in \mathcal{R}(A, B)$. Suppose that $E \subset A$ and $F \subset B$. Then $(R^*)|_{F \times E} = (R|_{E \times F})^*$.

Proof. We note that for each $b, a \in B \times A$,

$$\begin{aligned} (b, a) \in (R^*)|_{F \times E} &\iff (b, a) \in R^* \cap (F \times E) \\ &\iff (a, b) \in R \cap (E \times F) \\ &\iff (a, b) \in R|_{E \times F}. \end{aligned}$$

Hence $(R^*)|_{F \times E} = R|_{E \times F}$. □

1.3.2 Products of Relations

Definition 1.3.2.1. Let $(A_j)_{j \in [n]}, (B_j)_{j \in [n]}$ be sets and $(R_j)_{j \in [n]} \in \prod_{j \in [n]} \mathcal{R}(A_j, B_j)$. We define the **product relation of $(R_j)_{j \in [n]}$** ,

denoted $\prod_{j \in [n]}^{\text{Rel}} R_j \in \mathcal{R}\left(\prod_{j \in [n]} A_j, \prod_{j \in [n]} B_j\right)$, by

$$\prod_{j \in [n]}^{\text{Rel}} R_j := \left\{ ((a_j)_{j \in [n]}, (b_j)_{j \in [n]}) \in \left(\prod_{j \in [n]} A_j \right) \times \left(\prod_{j \in [n]} B_j \right) : \text{for each } j \in [n], (a_j, b_j) \in R_j \right\}.$$

Note 1.3.2.2.

- When $n = 2$, we may write $R_1 \times^{\text{Rel}} R_2$ in place of $\prod_{j \in [2]}^{\text{Rel}} R_j$,
- If for each $j \in [n]$, $R_j = R$, we may write $R^{\times^{\text{Rel}} n}$ in place of $\prod_{j \in [n]}^{\text{Rel}} R$,
- When the context is clear we may write $\prod_{j \in [n]} R_j$ and $R_1 \times R_2$ in place of $\prod_{j \in [n]}^{\text{Rel}} R_j$ and $R_1 \times^{\text{Rel}} R_2$.

Exercise 1.3.2.3. Let A, B, C, D be sets.

1. Let $R \in \mathcal{R}(A, B)$ and $(S_\gamma)_{\gamma \in \Gamma} \subset \mathcal{R}(C, D)$. Then

$$(a) \quad R \times \left(\bigcup_{\gamma \in \Gamma} S_\gamma \right) = \bigcup_{\gamma \in \Gamma} (R \times S_\gamma)$$

$$(b) \quad R \times \left(\bigcap_{\gamma \in \Gamma} S_\gamma \right) = \bigcap_{\gamma \in \Gamma} (R \times S_\gamma)$$

2. Let $(R_\gamma)_{\gamma \in \Gamma} \subset \mathcal{R}(A, B)$ and $S \in \mathcal{R}(C, D)$. Then

$$(a) \quad \left(\bigcup_{\gamma \in \Gamma} R_\gamma \right) \times S = \bigcup_{\gamma \in \Gamma} (R_\gamma \times S)$$

$$(b) \quad \left(\bigcap_{\gamma \in \Gamma} R_\gamma \right) \times S = \bigcap_{\gamma \in \Gamma} (R_\gamma \times S)$$

Proof.

1. (a) We note that

$$\begin{aligned} ((a, c), (b, d)) \in R \times \left(\bigcup_{\gamma \in \Gamma} S_\gamma \right) &\iff (a, b) \in R \text{ and } (c, d) \in \bigcup_{\gamma \in \Gamma} S_\gamma \\ &\iff (a, b) \in R \text{ and there exists } \gamma_0 \in \Gamma \text{ such that } (c, d) \in S_{\gamma_0} \\ &\iff \text{there exists } \gamma_0 \in \Gamma \text{ such that } (a, b) \in R \text{ and } (c, d) \in S_{\gamma_0} \\ &\iff \text{there exists } \gamma_0 \in \Gamma \text{ such that } ((a, c), (b, d)) \in R \times S_{\gamma_0} \\ &\iff ((a, c), (b, d)) \in \bigcup_{\gamma \in \Gamma} (R \times S_\gamma). \end{aligned}$$

$$\text{Hence } R \times \left(\bigcup_{\gamma \in \Gamma} S_\gamma \right) = \bigcup_{\gamma \in \Gamma} (R \times S_\gamma).$$

(b) We note that

$$\begin{aligned}
 ((a, c), (b, d)) \in R \times \left(\bigcap_{\gamma \in \Gamma} S_\gamma \right) &\iff (a, b) \in R \text{ and } (c, d) \in \bigcap_{\gamma \in \Gamma} S_\gamma \\
 &\iff (a, b) \in R \text{ for each } \gamma \in \Gamma, (c, d) \in S_\gamma \\
 &\iff \text{for each } \gamma \in \Gamma, (a, b) \in R \text{ and } (c, d) \in S_\gamma \\
 &\iff \text{for each } \gamma \in \Gamma, ((a, c), (b, d)) \in R \times S_\gamma \\
 &\iff ((a, c), (b, d)) \in \bigcap_{\gamma \in \Gamma} (R \times S_\gamma).
 \end{aligned}$$

$$\text{Hence } R \times \left(\bigcap_{\gamma \in \Gamma} S_\gamma \right) = \bigcap_{\gamma \in \Gamma} (R \times S_\gamma).$$

2. (a) We note that

$$\begin{aligned}
 ((a, c), (b, d)) \in \left(\bigcup_{\gamma \in \Gamma} R_\gamma \right) \times S &\iff (a, b) \in \left(\bigcup_{\gamma \in \Gamma} R_\gamma \right) \text{ and } (c, d) \in S \\
 &\iff \text{there exists } \gamma_0 \in \Gamma \text{ such that } (a, b) \in R_{\gamma_0} \text{ and } (c, d) \in S \\
 &\iff \text{there exists } \gamma_0 \in \Gamma \text{ such that } ((a, c), (b, d)) \in R_{\gamma_0} \times S \\
 &\iff ((a, c), (b, d)) \in \bigcup_{\gamma \in \Gamma} (R_\gamma \times S)
 \end{aligned}$$

$$\text{Hence } \left(\bigcup_{\gamma \in \Gamma} R_\gamma \right) \times S = \bigcup_{\gamma \in \Gamma} (R_\gamma \times S).$$

(b) We note that

$$\begin{aligned}
 ((a, c), (b, d)) \in \left(\bigcap_{\gamma \in \Gamma} R_\gamma \right) \times S &\iff (a, b) \in \left(\bigcap_{\gamma \in \Gamma} R_\gamma \right) \text{ and } (c, d) \in S \\
 &\iff \text{for each } \gamma \in \Gamma, (a, b) \in R_\gamma \text{ and } (c, d) \in S \\
 &\iff \text{for each } \gamma \in \Gamma, ((a, c), (b, d)) \in R_\gamma \times S \\
 &\iff ((a, c), (b, d)) \in \bigcap_{\gamma \in \Gamma} (R_\gamma \times S)
 \end{aligned}$$

$$\text{Hence } \left(\bigcap_{\gamma \in \Gamma} R_\gamma \right) \times S = \bigcap_{\gamma \in \Gamma} (R_\gamma \times S).$$

□

1.3.3 Subrelations

Exercise 1.3.3.1. Let A be a set, $B \subset A$ and $R \in \mathcal{R}^2(A)$. Then $(R^*)|_B = (R|_B)^*$

Proof. Clear by [prev ex in section on binary hetero relations](#).

□

Chapter 2

Operations

Chapter 3

Equivalence Relations

3.1 Introduction

Note 3.1.0.1. Let X be a set and $\nu \in \mathcal{R}^2(X)$. When the context is clear, we may write $a\nu b$ in place of $(a, b) \in \nu$.

Definition 3.1.0.2. Let X be a set. Suppose that $X \neq \emptyset$.

- Let $\theta \in \mathcal{R}^2(X)$. Then θ is said to be
 - **reflexive** if for each $a \in X$, $(a, a) \in \theta$
 - **symmetric** if for each $a, b \in X$, $(a, b) \in \theta$ implies that $(b, a) \in \theta$
 - **transitive** if for each $a, b, c \in X$, $(a, b) \in \theta$ and $(b, c) \in \theta$ implies that $(a, c) \in \theta$
 - an **equivalence relation on X** if
 - 1. θ is reflexive,
 - 2. θ is symmetric,
 - 3. θ is transitive.
- We define $\text{Equiv}(X) := \{\theta \in \mathcal{R}^2(X) : \theta \text{ is an equivalence relation on } X\}$

Exercise 3.1.0.3. Let X be a set and $\theta \in \mathcal{R}^2(X)$. Then

1. θ is reflexive iff $\Delta_X \subset \theta$
2. θ is symmetric iff $\theta^* \subset \theta$
3. θ is transitive iff $\theta \circ \theta \subset \theta$

Proof.

1.
 - ($\implies$):
Suppose that θ is reflexive. Let $(a, a) \in \Delta_X$. Then $a \in X$. Since θ is reflexive, $(a, a) \in \theta$. Since $(a, a) \in \Delta_X$ is arbitrary, we have that for each $(a, a) \in \Delta_X$, $(a, a) \in \theta$. Hence $\Delta_X \subset \theta$.
 - ($\impliedby$):
Suppose that $\Delta_X \subset \theta$. Let $a \in X$. By definition and assumption,

$$\begin{aligned} (a, a) &\in \Delta_X \\ &\subset \theta. \end{aligned}$$

Since $a \in X$ is arbitrary, we have that for each $a \in X$, $(a, a) \in \theta$. Hence θ is reflexive.

2.
 - ($\implies$):
Suppose that θ is symmetric. Let $(b, a) \in \theta^*$. By definition, $(a, b) \in \theta$. Since θ is symmetric, $(b, a) \in \theta$. Since $(b, a) \in \theta^*$ is arbitrary, we have that for each $(b, a) \in \theta^*$, $(b, a) \in \theta$. Hence $\theta^* \subset \theta$.

- ($\Leftarrow$):

Suppose that $\theta^* \subset \theta$. Let $a, b \in X$. Suppose that $(a, b) \in \theta$. By definition and assumption,

$$\begin{aligned} (b, a) &\in \theta^* \\ &\subset \theta. \end{aligned}$$

Since $a, b \in X$ with $(a, b) \in \theta$ are arbitrary, we have that for each $a, b \in X$, $(a, b) \in \theta$ implies that $(b, a) \in \theta$. Hence θ is symmetric.

3. • ($\Rightarrow$):

Suppose that θ is transitive. Let $(a, c) \in \theta \circ \theta$. By definition, there exists $b \in X$ such that $(a, b) \in \theta$ and $(b, c) \in \theta$. Since θ is transitive, $(a, c) \in \theta$. Since $(a, c) \in \theta \circ \theta$ is arbitrary, we have that for each $(a, c) \in \theta \circ \theta$, $(a, c) \in \theta$. Hence $\theta \circ \theta \subset \theta$.

- ($\Leftarrow$):

Suppose that $\theta \circ \theta \subset \theta$. Let $a, b, c \in X$. Suppose that $(a, b) \in \theta$ and $(b, c) \in \theta$. By definition and assumption,

$$\begin{aligned} (a, c) &\in \theta \circ \theta \\ &\subset \theta. \end{aligned}$$

Since $a, b, c \in X$ with $(a, b) \in \theta$ and $(b, c) \in \theta$ are arbitrary, we have that for each $a, b \in X$, $(a, b) \in \theta$ and $(b, c) \in \theta$ implies that $(a, c) \in \theta$. Hence θ is transitive.

□

Definition 3.1.0.4. Let X be a set. Suppose that $X \neq \emptyset$. We define the **universal relation on X** , denoted $\nabla_X \in \mathcal{R}^2(X)$, by $\nabla_X := X^2$.

Exercise 3.1.0.5. Let X be a set. Suppose that $X \neq \emptyset$. Then $\nabla_X \in \text{Equiv}(X)$.

Proof. Let $a, b, c \in X$.

1. Since $a \in X$, $(a, a) \in \nabla_X$.
2. Suppose that $(a, b) \in \nabla_X$. Since $a, b \in X$, $(b, a) \in X^2$. Therefore $(a, b) \in \nabla_X$ implies that $(b, a) \in \nabla_X$.
3. Suppose that $(a, b), (b, c) \in \nabla_X$. Since $a, c \in X$, $(a, c) \in X^2$. Therefore $(a, b), (b, c) \in X^2$ implies that $(a, c) \in \nabla_X$.

□

Exercise 3.1.0.6. Let X be a set. Suppose that $X \neq \emptyset$. Then $\Delta_X \in \text{Equiv}(X)$.

Proof. Let $a, b, c \in X$.

1. Since $a \in X$, $(a, a) \in \Delta_X$.
2. Suppose that $(a, b) \in \Delta_X$. Then $a = b$ and

$$\begin{aligned} (b, a) &= (a, a) \\ &\in \Delta_X. \end{aligned}$$

Therefore $(a, b) \in \Delta_X$ implies that $(b, a) \in \Delta_X$.

3. Suppose that $(a, b), (b, c) \in \Delta_X$. Then $a = b$ and $b = c$. Hence $a = c$ and

$$\begin{aligned} (a, c) &= (a, a) \\ &\in \Delta_X. \end{aligned}$$

Therefore $(a, b), (b, c) \in \Delta_X$ implies that $(a, c) \in \Delta_X$.

□

Exercise 3.1.0.7. Let X be a set. Suppose that $X \neq \emptyset$. Then for each $(\theta_\alpha)_{\alpha \in A} \subset \text{Equiv}(X)$, $\bigcap_{\alpha \in A} \theta_\alpha \in \text{Equiv}(X)$.

Proof. Let $(\theta_\alpha)_{\alpha \in A} \subset \text{Equiv}(X)$. Set $\theta := \bigcap_{\alpha \in A} \theta_\alpha$. Let $a, b, c \in X$.

1. Let $\alpha \in A$. Since $\theta_\alpha \in \text{Equiv}(X)$, $(a, a) \in \theta_\alpha$. Since $\alpha \in A$ is arbitrary, we have that for each $\alpha \in A$, $(a, a) \in \theta_\alpha$. Hence $(a, a) \in \theta$.
2. Suppose that $(a, b) \in \theta$. Let $\alpha \in A$. Then $(a, b) \in \theta_\alpha$. Since $\theta_\alpha \in \text{Equiv}(X)$, we have that $(b, a) \in \theta_\alpha$. Since $\alpha \in A$ is arbitrary, we have that for each $\alpha \in A$, $(b, a) \in \theta_\alpha$. Hence $(b, a) \in \theta$. Therefore $(a, b) \in \theta$ implies that $(b, a) \in \theta$.
3. Suppose that $(a, b) \in \theta$ and $(b, c) \in \theta$. Let $\alpha \in A$. Then $(a, b), (b, c) \in \theta_\alpha$. Since $\theta_\alpha \in \text{Equiv}(X)$, $(a, c) \in \theta_\alpha$. Since $\alpha \in A$ is arbitrary, we have that for each $\alpha \in A$, $(a, c) \in \theta_\alpha$. Hence $(a, c) \in \theta$. Therefore $(a, b) \in \theta$ and $(b, c) \in \theta$ implies that $(a, c) \in \theta$.

Hence $\theta \in \text{Equiv}(X)$. □

Definition 3.1.0.8. Let X be a set. Suppose that $X \neq \emptyset$. Let $\nu \in \mathcal{R}^2(X)$.

- We define the **equivalence relations containing ν** , denoted $\mathcal{E}(\nu)$, by $\mathcal{E}(\nu) := \{\theta \in \text{Equiv}(X) : \nu \subset \theta\}$.
- We define the **equivalence relation on X generated by ν** , denoted $\epsilon(\nu) \in \mathcal{R}^2(X)$ by $\epsilon(\nu) := \bigcap_{\theta \in \mathcal{E}(\nu)} \theta$.

Note 3.1.0.9.

- A previous ex implies that $\nabla_X \in \mathcal{E}(\nu)$.
- A previous ex (about intersections of equiv relats) implies that $\epsilon(\nu) \in \mathcal{E}(\nu)$.

Exercise 3.1.0.10. Let X be a set. Suppose that $X \neq \emptyset$. Let $\nu \in \mathcal{R}^2(X)$. Define $(\nu_n)_{n \in \mathbb{N}_0} \subset \mathcal{R}^2(X)$ by

$$\nu_n := \begin{cases} \Delta_X \cup \nu \cup \nu^*, & n = 0 \\ \nu_{n-1} \circ \nu_{n-1}, & n \geq 1. \end{cases}$$

Then

1. (a) for each $n \in \mathbb{N}$, ν_n is reflexive
 (b) for each $n \in \mathbb{N}$, ν_n is symmetric
 (c) for each $n \in \mathbb{N}_0$, $\nu_n \subset \nu_{n+1}$
2. $\epsilon(\nu) = \bigcup_{n \in \mathbb{N}_0} \nu_n$.

Proof.

1. (a) • **Base Case:**
 Since

$$\begin{aligned} \Delta_X &\subset \Delta_X \cup \nu \cup \nu^* \\ &= \nu_0, \end{aligned}$$

a previous ex implies that ν_0 is reflexive.

• **Induction Step:**

Let $n \in \mathbb{N}_0$. Suppose that ν_n is reflexive. Then that same prev ex implies that $\Delta_X \subset \nu_n$. two prev ex in section on relations about composition with identity and subsets then imply that

$$\begin{aligned} \Delta_X &= \Delta_X \circ \Delta_X \\ &\subset \nu_n \circ \nu_n \\ &= \nu_{n+1}. \end{aligned}$$

Another application of that prev ex implies that ν_{n+1} is reflexive.

By induction, for each $n \in \mathbb{N}_0$, ν_n is reflexive.

(b) • **Base Case:**

a previous ex about converse relations and unions, double converse and since $\Delta_X \in \text{Equiv}(X)$ and an ex about symmetric relations implies that

$$\begin{aligned}
 \nu_0^* &= (\Delta_X \cup \nu \cup \nu^*)^* \\
 &= \Delta_X^* \cup \nu^* \cup (\nu^*)^* \\
 &= \Delta_X^* \cup \nu^* \cup \nu \\
 &\subset \Delta_X \cup \nu^* \cup \nu \\
 &= \Delta_X \cup \nu \cup \nu^* \\
 &= \nu_0.
 \end{aligned}$$

that same previous ex about symmetric relations implies that ν_0 is symmetric.

• **Induction Step:**

Let $n \in \mathbb{N}_0$. Suppose that ν_n is symmetric. Then that same prev ex about symmetric and converse relations implies that $\nu_n^* \subset \nu_n$. Therefore prev ex imply

$$\begin{aligned}
 \nu_{n+1}^* &= (\nu_n \circ \nu_n)^* \\
 &= \nu_n^* \circ \nu_n^* \\
 &\subset \nu_n \circ \nu_n. &= \nu_{n+1}.
 \end{aligned}$$

Another application of that prev ex implies that ν_{n+1} is symmetric.

By induction, for each $n \in \mathbb{N}_0$, ν_n is symmetric.

(c) Let $n \in \mathbb{N}_0$. Since ν_n is reflexive, prev ex implies that $\Delta_X \subset \nu_n$. Therefore prev ex implies

$$\begin{aligned}
 \nu_n &= \Delta_X \circ \nu_n \\
 &\subset \nu_n \circ \nu_n \\
 &= \nu_{n+1}.
 \end{aligned}$$

Since $n \in \mathbb{N}_0$ is arbitrary, we have that for each $n \in \mathbb{N}_0$, $\nu_n \subset \nu_{n+1}$.

2. Set $\theta_0 := \bigcup_{n \in \mathbb{N}_0} \nu_n$.

• – **Base Case:**

Since $\epsilon(\nu) \in \mathcal{E}(\nu)$, we have that $\epsilon(\nu) \in \text{Equiv}(X)$ and $\nu \subset \epsilon(\nu)$. Since $\epsilon(\nu) \in \text{Equiv}(X)$, we have that $\epsilon(\nu)$ is reflexive and symmetric. A ref ex then implies that $\Delta_X \subset \epsilon(\nu)$ and $\epsilon(\nu)^* \subset \epsilon(\nu)$. Therefore a prev ex about converse relations and subsets

$$\begin{aligned}
 \nu_0 &= \Delta_X \cup \nu \cup \nu^* \\
 &\subset \epsilon(\nu) \cup \epsilon(\nu) \cup \epsilon(\nu)^* \\
 &\subset \epsilon(\nu) \cup \epsilon(\nu) \cup \epsilon(\nu) \\
 &= \epsilon(\nu).
 \end{aligned}$$

– **Induction Step:**

Let $n \in \mathbb{N}_0$. Suppose that $\nu_n \subset \epsilon(\nu)$. Then ref prev ex

$$\begin{aligned}
 \nu_{n+1} &= \nu_n \circ \nu_n \\
 &\subset \epsilon(\nu) \circ \epsilon(\nu) \\
 &= \epsilon(\nu).
 \end{aligned}$$

By induction, for each $n \in \mathbb{N}_0$, $\nu_n \subset \epsilon(\nu)$. Therefore

$$\begin{aligned}
 \theta_0 &= \bigcup_{n \in \mathbb{N}_0} \nu_n \\
 &\subset \epsilon(\nu).
 \end{aligned}$$

- – Since ν_0 is reflexive, we have that

$$\begin{aligned}\Delta_X &\subset \nu_0 \\ &\subset \bigcup_{n \in \mathbb{N}_0} \nu_n \\ &= \theta_0.\end{aligned}$$

Hence θ_0 is reflexive.

- Since for each $n \in \mathbb{N}_0$, ν_n is symmetric, we have that that **ref exs** for each $n \in \mathbb{N}_0$, $\nu_n^* \subset \nu_n$. Therefore

$$\begin{aligned}\theta_0^* &= \left(\bigcup_{n \in \mathbb{N}_0} \nu_n \right)^* \\ &= \bigcup_{n \in \mathbb{N}_0} \nu_n^* \\ &\subset \bigcup_{n \in \mathbb{N}_0} \nu_n \\ &= \theta_0.\end{aligned}$$

Hence θ_0 is symmetric.

- Let $a, b, c \in X$. Suppose that $(a, b) \in \theta_0$ and $(b, c) \in \theta_0$. Then there exist $n_1, n_2 \in \mathbb{N}_0$ such that $(a, b) \in \nu_{n_1}$ and $(b, c) \in \nu_{n_2}$. Set $n_0 := \max\{n_1, n_2\}$. Since for each $n \in \mathbb{N}$, $\nu_n \subset \nu_{n+1}$, we have that

$$\begin{aligned}(a, b) &\in \nu_{n_1} \\ &\subset \nu_{n_0}.\end{aligned}$$

and

$$\begin{aligned}(b, c) &\in \nu_{n_2} \\ &\subset \nu_{n_0}.\end{aligned}$$

Since $(a, b), (b, c) \in \nu_{n_0}$, we have that

$$\begin{aligned}(a, c) &\in \nu_{n_0} \circ \nu_{n_0} \\ &= \nu_{n_0+1} \\ &\subset \bigcup_{n \in \mathbb{N}_0} \nu_n \\ &= \theta_0.\end{aligned}$$

Since $a, b, c \in X$ with $(a, b) \in \theta_0$ and $(b, c) \in \theta_0$ are arbitrary, we have that for each $a, b, c \in X$, $(a, b) \in \theta_0$ and $(b, c) \in \theta_0$ implies that $(a, c) \in \theta_0$. Hence θ_0 is transitive.

Since θ_0 is reflexive, symmetric and transitive, $\theta_0 \in \text{Equiv}(X)$. Since

$$\begin{aligned}\nu &\subset \Delta_X \cup \nu \cup \nu^* \\ &= \nu_0 \\ &\subset \bigcup_{n \in \mathbb{N}_0} \nu_n \\ &= \theta_0,\end{aligned}$$

we have that $\theta_0 \in \mathcal{E}(\nu)$. Therefore

$$\begin{aligned}\epsilon(\nu) &= \bigcap_{\theta \in \mathcal{E}(\nu)} \theta \\ &\subset \theta_0.\end{aligned}$$

Since $\theta_0 \subset \epsilon(\nu)$ and $\epsilon(\nu) \subset \theta_0$, we have that $\theta_0 = \epsilon(\nu)$.

□

3.2 Products

Exercise 3.2.0.1. Let $(X_j)_{j \in [n]}$ be sets. Then for each $(\theta_j)_{j \in [n]} \in \prod_{j \in [n]} \text{Equiv}(X_j)$, $\Pi_{j \in [n]}^{\text{Rel}} \theta_j \in \text{Equiv}(\prod_{j \in [n]} X_j)$.

Proof. Set $X := \prod_{j \in [n]} X_j$ and define $\theta \in \mathcal{R}^2(X)$ by $\theta := \Pi_{j \in [n]}^{\text{Rel}} \theta_j$.

1. For each $(a_j)_{j \in [n]} \in X$,

$$\begin{aligned} \text{for each } j \in [n], a_j \in X_j &\implies \text{for each } j \in [n], (a_j, a_j) \in \theta_j \\ &\implies ((a_j)_{j \in [n]}, (a_j)_{j \in [n]}) \in \theta. \end{aligned}$$

Thus θ is reflexive.

2. For each $(a_j)_{j \in [n]}, (b_j)_{j \in [n]} \in X$,

$$\begin{aligned} ((a_j)_{j \in [n]}, (b_j)_{j \in [n]}) \in \theta &\iff \text{for each } j \in [n], (a_j, b_j) \in \theta_j \\ &\implies \text{for each } j \in [n], (b_j, a_j) \in \theta_j \\ &\iff ((b_j)_{j \in [n]}, (a_j)_{j \in [n]}) \in \theta. \end{aligned}$$

Thus θ is symmetric.

3. For each $(a_j)_{j \in [n]}, (b_j)_{j \in [n]}, (c_j)_{j \in [n]} \in X$,

$$\begin{aligned} ((a_j)_{j \in [n]}, (b_j)_{j \in [n]}) \in \theta \text{ and } ((b_j)_{j \in [n]}, (c_j)_{j \in [n]}) \in \theta &\iff \text{for each } j \in [n], (a_j, b_j) \in \theta_j \text{ and } (b_j, c_j) \in \theta_j \\ &\implies \text{for each } j \in [n], (a_j, c_j) \in \theta_j \\ &\iff ((a_j)_{j \in [n]}, (c_j)_{j \in [n]}) \in \theta. \end{aligned}$$

Thus θ is transitive.

Hence $\theta \in \text{Equiv}(X)$. □

Exercise 3.2.0.2.

Proof. □

3.3 Quotients

Exercise 3.3.0.1. Let $X, Y \in \text{Set}$, $\theta_X \in \text{Equiv}(X)$, $\theta_Y \in \text{Equiv}(Y)$ and $f : X \rightarrow Y$. If f is (θ_X, θ_Y) -invariant, then there exists a unique $\bar{f} : X/\theta_X \rightarrow Y/\theta_Y$ such that $\pi_{\theta_Y} \circ f = \bar{f} \circ \pi_{\theta_X}$, i.e. the following diagram commutes:

$$\begin{array}{ccc} X & \xrightarrow{f} & Y \\ \pi_{\theta_X} \downarrow & & \downarrow \pi_{\theta_Y} \\ X/\theta_X & \xrightarrow{\bar{f}} & Y/\theta_Y \end{array}$$

Proof. Suppose that f is (θ_X, θ_Y) -invariant.

• **Existence:**

Define $\bar{f} \subset (X/\theta_X) \times (Y/\theta_Y)$ by $\bar{f} := \{(\pi_{\theta_X}(a), \pi_{\theta_Y}(f(a))) : a \in X\}$.

- Let $c \in X/\theta_X$. Then there exists $a \in X$ such that $c := \pi_{\theta_X}(a)$. Define $d \in Y/\theta_Y$ by $d := \pi_{\theta_Y}(f(a))$. Then

$$\begin{aligned} (c, d) &= (\pi_{\theta_X}(a), \pi_{\theta_Y}(f(a))) \\ &\in \bar{f}. \end{aligned}$$

Since $c \in X/\theta_X$ is arbitrary, we have that for each $c \in X/\theta_X$, there exists $d \in Y/\theta_Y$ such that $(c, d) \in \bar{f}$.

- Let $(c_1, d_1), (c_2, d_2) \in \bar{f}$. Then there exists $a_1, a_2 \in X$ such that $(c_1, d_1) = (\pi_{\theta_X}(a_1), \pi_{\theta_Y}(f(a_1)))$, $(c_2, d_2) = (\pi_{\theta_X}(a_2), \pi_{\theta_Y}(f(a_2)))$. Suppose that $c_1 = c_2$. Then

$$\begin{aligned}\pi_{\theta_X}(a_1) &= c_1 \\ &= c_2 \\ &= \pi_{\theta_X}(a_2).\end{aligned}$$

Hence $a_1 \theta_X a_2$. Since f is (θ_X, θ_Y) -invariant, $f(a_1) \theta_Y f(a_2)$. Thus

$$\begin{aligned}d_1 &= \pi_{\theta_Y}(f(a_1)) \\ &= \pi_{\theta_Y}(f(a_2)) \\ &= d_2.\end{aligned}$$

Since $(c_1, d_1), (c_2, d_2) \in \bar{f}$ with $c_1 = c_2$ are arbitrary, we have that for each $(c_1, d_1), (c_2, d_2) \in \bar{f}$, $c_1 = c_2$ implies that $d_1 = d_2$.

Thus $\bar{f} : X/\theta_X \rightarrow Y/\theta_Y$. By construction, for each $a \in X$,

$$\begin{aligned}\pi_{\theta_Y} \circ f(a) &= \pi_{\theta_Y}(f(a)) \\ &= \bar{f}(\pi_{\theta_X}(a)) \\ &= \bar{f} \circ \pi_{\theta_X}(a).\end{aligned}$$

Thus $\pi_{\theta_Y} \circ f = \bar{f} \circ \pi_{\theta_X}$.

• **Uniqueness:**

Let $g : X/\theta_X \rightarrow Y/\theta_Y$. Suppose that $\pi_{\theta_Y} \circ f = g \circ \pi_{\theta_X}$. Let $c \in X/\theta_X$. Then there exists $a \in X$ such that $c = \pi_{\theta_X}(a)$. Then

$$\begin{aligned}g(c) &= g(\pi_{\theta_X}(a)) \\ &= g \circ \pi_{\theta_X}(a) \\ &= \pi_{\theta_Y} \circ f(a) \\ &= \bar{f} \circ \pi_{\theta_X}(a) \\ &= \bar{f}(\pi_{\theta_X}(a)) \\ &= \bar{f}(c).\end{aligned}$$

Since $c \in X/\theta_X$ is arbitrary, we have that for each $c \in X/\theta_X$, $g(c) = \bar{f}(c)$. Hence $g = \bar{f}$.

□

Definition 3.3.0.2. Let $X, Y \in \text{Set}$, $\theta_X \in \text{Equiv}(X)$, $\theta_Y \in \text{Equiv}(Y)$ and $f : X \rightarrow Y$. Suppose that f is (θ_X, θ_Y) -invariant. We define the **quotient of f by (θ_X, θ_Y)** , denoted $f/(\theta_X, \theta_Y) : X/\theta_X \rightarrow Y/\theta_Y$, to be the unique $\bar{f} : X/\theta_X \rightarrow Y/\theta_Y$ such that $\pi_{\theta_Y} \circ f = \bar{f} \circ \pi_{\theta_X}$.

Exercise 3.3.0.3. Let $X, Y \in \text{Set}$, $\theta \in \text{Equiv}(X)$ and $f : X \rightarrow Y$. If f is θ -invariant, then there exists a unique $\bar{f} : X/\theta \rightarrow Y$ such that $f = \bar{f} \circ \pi_\theta$, i.e. the following diagram commutes:

$$\begin{array}{ccc} X & & \\ \pi_\theta \downarrow & \searrow f & \\ X/\theta & \xrightarrow{\bar{f}} & Y \end{array}$$

Proof. Suppose that f is θ -invariant.

- **Existence:**

Since f is θ -invariant, we have that f is (θ, Δ_Y) -invariant. **make ref ex** implies that $\pi_{\Delta_Y} \in \text{Iso}_{\text{Set}}(Y, Y/\Delta_Y)$. Define $\bar{f} : X/\theta \rightarrow Y$ by $\bar{f} := \pi_{\Delta_Y}^{-1} \circ f/(\theta, \Delta_Y)$. Then

$$\begin{aligned} \bar{f} \circ \pi_\theta &= (\pi_{\Delta_Y}^{-1} \circ f/(\theta, \Delta_Y)) \circ \pi_\theta \\ &= \pi_{\Delta_Y}^{-1} \circ (f/(\theta, \Delta_Y) \circ \pi_\theta) \\ &= \pi_{\Delta_Y}^{-1} \circ (\pi_{\Delta_Y} \circ f) \\ &= (\pi_{\Delta_Y}^{-1} \circ \pi_{\Delta_Y}) \circ f \\ &= \text{id}_Y \circ f \\ &= f. \end{aligned}$$

- **Uniqueness:**

Let $g : X/\theta \rightarrow Y$. Suppose that $g \circ \pi_\theta = f$. Let $c \in X/\theta$. Then there exists $a \in X$ such that $c = \pi_\theta(a)$. Then

$$\begin{aligned} g(c) &= g(\pi_\theta(a)) \\ &= g \circ \pi_\theta(a) \\ &= f(a) \\ &= \bar{f} \circ \pi_\theta(a) \\ &= \bar{f}(\pi_\theta(a)) \\ &= \bar{f}(c). \end{aligned}$$

Since $c \in X/\theta_X$ is arbitrary, we have that for each $c \in X/\theta_X$, $g(c) = \bar{f}(c)$. Hence $g = \bar{f}$. □

Definition 3.3.0.4. Let $X, Y \in \text{Set}$, $\theta \in \text{Equiv}(X)$ and $f : X \rightarrow Y$. Suppose that f is (θ) -invariant. We define the **quotient of f by θ** , denoted $f/\theta : X/\theta \rightarrow Y$, to be the unique $\bar{f} : X/\theta \rightarrow Y$ such that $f = \bar{f} \circ \pi_\theta$.

Exercise 3.3.0.5. Let $n \in \mathbb{N}$, $(X_j)_{j \in [n]} \subset \text{Set}$ and $(\theta_j)_{j \in [n]} \in \prod_{j \in [n]} \text{Equiv}(X_j)$. Define $X, X' \in \text{Set}$, $\theta \in \text{Equiv}(X)$ and $\pi' : X \rightarrow X'$ by $X := \prod_{j \in [n]} X_j$, $X' := \prod_{j \in [n]} (X_j/\theta_j)$, $\theta := \prod_{j \in [n]}^{\text{Rel}} \theta_j$ and $\pi' := \prod_{j \in [n]} \pi_{\theta_j}$. Then π' is θ -invariant.

Proof. For each $(a_j)_{j \in [n]}, (b_j)_{j \in [n]} \in X$,

$$\begin{aligned} (a_j)_{j \in [n]} \sim_\theta (b_j)_{j \in [n]} &\iff \text{for each } j \in [n], a_j \sim_{\theta_j} b_j \\ &\iff \text{for each } j \in [n], \pi_{\theta_j}(a_j) = \pi_{\theta_j}(b_j) \\ &\iff (\pi_{\theta_j}(a_j))_{j \in [n]} = (\pi_{\theta_j}(b_j))_{j \in [n]} \\ &\iff \pi'((a_j)_{j \in [n]}) = \pi'((b_j)_{j \in [n]}). \end{aligned}$$

Hence π' is $\theta^{\times n}$ -invariant. □

Exercise 3.3.0.6. Let $n \in \mathbb{N}$, $(X_j)_{j \in [n]} \subset \text{Set}$ and $(\theta_j)_{j \in [n]} \in \prod_{j \in [n]} \text{Equiv}(X_j)$. Define $X, X' \in \text{Set}$, $\theta \in \text{Equiv}(X)$ and $\pi' : X \rightarrow X'$ by $X := \prod_{j \in [n]} X_j$, $X' := \prod_{j \in [n]} (X_j/\theta_j)$, $\theta := \prod_{j \in [n]}^{\text{Rel}} \theta_j$, $\pi := \pi_\theta$ and $\pi' := \prod_{j \in [n]} \pi_{\theta_j}$. Then $(\pi')/\theta \in \text{Iso}_{\text{Set}}(X/\theta, X')$.

Proof. Set $\alpha := (\pi')/\theta$. **A prev ex** implies that $\pi' = \alpha \circ \pi_\theta$

- **Injectivity:**

Let $c, d \in X/\theta$. Then there exist $(a_j)_{j \in [n]}, (b_j)_{j \in [n]} \in X$ such that $c = \pi_\theta((a_j)_{j \in [n]})$ and $d = \pi_\theta((b_j)_{j \in [n]})$. Suppose that $\alpha(c) = \alpha(d)$. Then

$$\begin{aligned} \pi'((a_j)_{j \in [n]}) &= \alpha(\pi_\theta((a_j)_{j \in [n]})) \\ &= \alpha(c) \\ &= \alpha(d) \\ &= \alpha(\pi_\theta((b_j)_{j \in [n]})) \\ &= \pi'((b_j)_{j \in [n]}). \end{aligned}$$

Thus for each $j \in [n]$, $\pi_{\theta_j}(a_j) = \pi_{\theta_j}(b_j)$. Hence for each $j \in [n]$, $a_j \sim_{\theta_j} b_j$. Therefore $(a_j)_{j \in [n]} \sim_{\theta} (b_j)_{j \in [n]}$. Thus

$$\begin{aligned} c &= \pi_{\theta}((a_j)_{j \in [n]}) \\ &= \pi_{\theta}((b_j)_{j \in [n]}) \\ &= d. \end{aligned}$$

Since $c, d \in X/\theta$ with $\alpha(c) = \alpha(d)$ are arbitrary, we have that for each $c, d \in X/\theta$, $\alpha(c) = \alpha(d)$ implies that $c = d$. Hence α is injective.

• **Surjectivity:**

Since for each $j \in [n]$, π_{θ_j} is surjective, **ref ex** implies that π' is surjective. Since $\pi' = \alpha \circ \pi_{\theta}$, **ref ex** implies that α is surjective.

FINISH!!! (ref exercises) □

Exercise 3.3.0.7. Let $n \in \mathbb{N}$, $(X_j)_{j \in [n]} \subset \text{Set}$, $Y \in \text{Set}$, $(\theta_{X_j})_{j \in [n]} \in \prod_{j \in [n]} \text{Equiv}(X_j)$, $\theta_Y \in \text{Equiv}(Y)$ and $f : \prod_{j \in [n]} X_j \rightarrow Y$. Define $X, X' \in \text{Set}$, $\theta \in \text{Equiv}(X)$ and $\pi' : X \rightarrow X'$ by $X := \prod_{j \in [n]} X_j$, $X' := \prod_{j \in [n]} (X_j/\theta_{X_j})$, $\theta := \prod_{j \in [n]}^{\text{Rel}} \theta_{X_j}$ and $\pi' := \prod_{j \in [n]} \pi_{\theta_{X_j}}$. If f is (θ, θ_Y) -invariant, then there exists a unique $\bar{f} : X' \rightarrow Y/\theta_Y$ such that $\pi_{\theta_Y} \circ f = \bar{f} \circ \pi'$, i.e. the following diagram commutes:

$$\begin{array}{ccc} X & \xrightarrow{f} & Y \\ \pi' \downarrow & & \downarrow \pi_{\theta_Y} \\ X' & \xrightarrow{\bar{f}} & Y/\theta_Y \end{array}$$

Proof. Suppose that f is (θ, θ_Y) -invariant.

• **Existence:**

A prev ex implies that π' is θ -invariant. **The prev ex** implies that $(\pi')/\theta \in \text{Iso}_{\text{Set}}(X/\theta, X')$. Define $\bar{f} : X' \rightarrow Y/\theta_Y$ by $\bar{f} := f/(\theta, \theta_Y) \circ [(\pi')/\theta]^{-1}$. Then

$$\begin{aligned} \bar{f} \circ \pi' &= (f/(\theta, \theta_Y) \circ [(\pi')/\theta]^{-1}) \circ ((\pi')/\theta \circ \pi_{\theta}) \\ &= f/(\theta, \theta_Y) \circ ([(\pi')/\theta]^{-1} \circ (\pi')/\theta) \circ \pi_{\theta} \\ &= f/(\theta, \theta_Y) \circ \text{id}_{X/\theta} \circ \pi_{\theta} \\ &= f/(\theta, \theta_Y) \circ \pi_{\theta} \\ &= \pi_{\theta_Y} \circ f. \end{aligned}$$

• **Uniqueness:**

Let $g : X' \rightarrow Y/\theta_Y$. Suppose that $\pi_{\theta_Y} \circ f = g \circ \pi'$. Let $c \in X'$. Then there exists $a \in X$ such that $c = \pi'(a)$. Then

$$\begin{aligned} g(c) &= g(\pi'(a)) \\ &= g \circ \pi'(a) \\ &= \pi_{\theta_Y} \circ f(a) \\ &= \bar{f} \circ \pi'(a) \\ &= \bar{f}(\pi'(a)) \\ &= \bar{f}(c). \end{aligned}$$

Since $c \in X'$ is arbitrary, we have that for each $c \in X'$, $g(c) = \bar{f}(c)$. Hence $g = \bar{f}$. □

Definition 3.3.0.8. Let $n \in \mathbb{N}$, $(X_j)_{j \in [n]} \subset \text{Set}$, $Y \in \text{Set}$, $(\theta_{X_j})_{j \in [n]} \in \prod_{j \in [n]} \text{Equiv}(X_j)$, $\theta_Y \in \text{Equiv}(Y)$ and $f : \prod_{j \in [n]} X_j \rightarrow Y$. Suppose that f is $\left(\prod_{j \in [n]}^{\text{Rel}} \theta_{X_j}, \theta_Y\right)$ -invariant. We define **multiquotient of f by $((\theta_{X_j})_{j \in [n]}, \theta_Y)$** , denoted $f / ((\theta_{X_j})_{j \in [n]}, \theta_Y) : \prod_{j \in [n]} X_j / \theta_{X_j} \rightarrow Y / \theta_Y$, to be the unique $\bar{f} : \prod_{j \in [n]} X_j / \theta_{X_j} \rightarrow Y / \theta_Y$ such that

$$\pi_{\theta_Y} \circ f = \bar{f} \circ \prod_{j \in [n]} \pi_{\theta_{X_j}}.$$

Note 3.3.0.9. If for each $j \in [n]$, $X_j = X$ and $\theta_{X_j} = \theta_X$, we may write $f / ((n : \theta_X), \theta_Y)$ in place of $f / ((\theta_{X_j})_{j \in [n]}, \theta_Y)$

Exercise 3.3.0.10. Let $n \in \mathbb{N}$, $(X_j)_{j \in [n]} \subset \text{Set}$, $Y \in \text{Set}$, $(\theta_j)_{j \in [n]} \in \prod_{j \in [n]} \text{Equiv}(X_j)$ and $f : \prod_{j \in [n]} X_j \rightarrow Y$. Define $X, X' \in \text{Set}$, $\theta \in \text{Equiv}(X)$ and $\pi' : X \rightarrow X'$ by $X := \prod_{j \in [n]} X_j$, $X' := \prod_{j \in [n]} (X_j / \theta_j)$, $\theta := \prod_{j \in [n]}^{\text{Rel}} \theta_j$ and $\pi' := \prod_{j \in [n]} \pi_{\theta_j}$. If f is θ -invariant, then there exists a unique $\bar{f} : X' \rightarrow Y$ such that $f = \bar{f} \circ \pi'$, i.e. the following diagram commutes:

$$\begin{array}{ccc} X & & \\ \pi' \downarrow & \searrow f & \\ X' & \xrightarrow{\bar{f}} & Y \end{array}$$

Proof. Suppose that f is θ -invariant.

• **Existence:**

Since f is θ -invariant, we have that f is (θ, Δ_Y) -invariant. Define $\bar{f}' : X' \rightarrow Y / \Delta_Y$ by $\bar{f}' := f / ((\theta_j)_{j \in [n]}, \Delta_Y)$. Then $\pi_{\Delta_Y} \circ f = \bar{f}' \circ \pi'$. **make ref ex** implies that $\pi_{\Delta_Y} \in \text{Iso}_{\text{Set}}(Y, Y / \Delta_Y)$. Define $\bar{f} : X' \rightarrow Y$ by $\bar{f} := \pi_{\Delta_Y}^{-1} \circ \bar{f}'$. Then

$$\begin{aligned} \bar{f} \circ \pi' &= (\pi_{\Delta_Y}^{-1} \circ \bar{f}') \circ \pi' \\ &= \pi_{\Delta_Y}^{-1} \circ (\bar{f}' \circ \pi') \\ &= \pi_{\Delta_Y}^{-1} \circ (\pi_{\Delta_Y} \circ f) \\ &= (\pi_{\Delta_Y}^{-1} \circ \pi_{\Delta_Y}) \circ f \\ &= \text{id}_Y \circ f \\ &= f. \end{aligned}$$

• **Uniqueness:**

Let $g : X' \rightarrow Y$. Suppose that $g \circ \pi' = f$. Let $c \in X'$. Then there exists $a \in X$ such that $c = \pi'(a)$. Then

$$\begin{aligned} g(c) &= g(\pi'(a)) \\ &= g \circ \pi'(a) \\ &= f(a) \\ &= \bar{f} \circ \pi'(a) \\ &= \bar{f}(\pi'(a)) \\ &= \bar{f}(c). \end{aligned}$$

Since $c \in X'$ is arbitrary, we have that for each $c \in X'$, $g(c) = \bar{f}(c)$. Hence $g = \bar{f}$. □

Definition 3.3.0.11. Let $n \in \mathbb{N}$, $(X_j)_{j \in [n]} \subset \text{Set}$, $Y \in \text{Set}$, $(\theta_j)_{j \in [n]} \in \prod_{j \in [n]} \text{Equiv}(X_j)$ and $f : \prod_{j \in [n]} X_j \rightarrow Y$. Suppose that f is $\prod_{j \in [n]}^{\text{Rel}} \theta_j$ -invariant. We define **multiquotient of f by $(\theta_j)_{j \in [n]}$** , denoted $f / (\theta_j)_{j \in [n]} : \prod_{j \in [n]} X_j / \theta_j \rightarrow Y$, to be the unique

$\bar{f} : \prod_{j \in [n]} X_j / \theta_j \rightarrow Y$ such that

$$f = \bar{f} \circ \prod_{j \in [n]} \pi_{\theta_j}.$$

Note 3.3.0.12. If for each $j \in [n]$, $X_j = X$ and $\theta_j = \theta$, we may write $f/(n : \theta)$ in place of $f/(\theta_j)_{j \in [n]}$

Part II

Ordered Sets

3.4 To Do

- Relabel exercises
- finish generated element subsection in galois connection section

at this point, there should be a simple structure capturing posets and $\downarrow S$, measurable spaces and $\sigma(\mathcal{A})$, topological spaces and $\tau(\mathcal{E})$, groups/rings/ $\dots$ /algebras and $\langle S \rangle / (E) / \dots / (F)$. It seems like we need a category $\mathcal{C}$ with some structure, another category $\mathcal{C}_F$ which is like $\mathcal{C}$, but less structured and some forgetful-like functor $F : \mathcal{C} \rightarrow \mathcal{C}_F$. eg, $F : \mathbf{Pro} \rightarrow \mathbf{Set}$ or $F : \mathbf{Top} \rightarrow \mathbf{Set}$ and we need an associated poset $\mathcal{P}$ containing the structure forgotten by F , e.g. the lower sets of X or the set of topologies on X and we need a “generating” or “contained” object $\mathcal{E}$ in $\mathcal{C}_F$ in the sense that there is at least one object A in $\mathcal{C}$ and monomorphism $\iota_A : \mathcal{E} \rightarrow F(A)$ in $\mathcal{C}_F$. We then need a minimal element in the structure poset $\mathcal{P}$ in some universal sense relating to these monomorphisms. **Ask people who know category theory**

Chapter 4

Prosets

4.1 Introduction

Definition 4.1.0.1. Proset:

Let X be a set and $\leq \subset X \times X$ a binary relation on X . Then

- $\leq$ is said to be a **preorder on X** if
 1. for each $a \in X$, $a \leq a$
 2. for each $a, b, c \in X$, $a \leq b$ and $b \leq c$ implies that $a \leq c$
- $(X, \leq)$ is said to be a **preordered set** or **proset** if $\leq$ is a preorder on X .

Definition 4.1.0.2. maybe just ref def about converse relation in set theory appendices section on heterogeneous binary relations

Let $(X, \leq)$ be a proset. We define the **dual order of $\leq$ on X** , denoted $\leq^*$, by $a \leq^* b$ iff $b \leq a$.

Exercise 4.1.0.3. Let $(X, \leq)$ be a proset. Then $\leq^*$ is a preorder on X .

Proof.

1. Let $a \in X$. Since $a \leq a$, we have that $a \leq^* a$.
2. Let $a, b, c \in X$. Suppose that $a \leq^* b$ and $b \leq^* c$. Then $b \leq a$ and $c \leq b$. Hence $c \leq a$. Thus $a \leq^* c$.

Therefore $\leq^*$ is a preorder on X . □

Definition 4.1.0.4. Let $(X, \leq)$ be a proset. We define $\sim_{\leq} \subset X \times X$ by $a \sim_{\leq} b$ iff $a \leq b$ and $b \leq a$.

Exercise 4.1.0.5. Let $(X, \leq)$ be a proset. Then $\sim_{\leq}$ is an equivalence relation on X .

Proof. Let $x, y, z \in X$.

1. Since $x \leq x$, $x \sim_{\leq} x$.
2. Suppose that $x \sim_{\leq} y$. Then $x \leq y$ and $y \leq x$. Thus $y \sim_{\leq} x$.
3. Suppose that $x \sim_{\leq} y$ and $y \sim_{\leq} z$. Then $x \leq y$, $y \leq x$, $y \leq z$ and $z \leq y$. Therefore $x \leq z$ and $z \leq x$. Hence $x \sim_{\leq} z$. □

Exercise 4.1.0.6. Let $(X, \leq)$ be a proset. Then for each $x, y \in X$, $x \sim_{\leq} y$ iff $x \sim_{\leq^*} y$.

Proof. We have that for each $x, y \in X$,

$$\begin{aligned} x \sim_{\leq} y &\iff x \leq y \text{ and } y \leq x \\ &\iff y \leq^* x \text{ and } x \leq^* y \\ &\iff x \sim_{\leq^*} y. \end{aligned}$$
□

Exercise 4.1.0.7. Let X be a set. Then $(\mathcal{P}(X), \subset)$ is a proset.

Proof.

1. For each $A \in \mathcal{P}(X)$, $A \subset A$.
2. For each $A, B, C \in \mathcal{P}(X)$, if $A \subset B$ and $B \subset C$, then $A \subset C$.

Thus $\subset$ is a preorder on $\mathcal{P}(X)$. Hence $(\mathcal{P}(X), \subset)$ is a proset. □

Definition 4.1.0.8. We define the

- **one point proset**, denoted $(\mathbf{1}, \leq_1)$, by
 - $\mathbf{1} := \{0\}$
 - $\leq_1 := \{(0, 0)\}$.
- **two point proset**, denoted $(\mathbf{2}, \leq_2)$, by
 - $\mathbf{2} := \{0, 1\}$
 - $\leq_2 := \{(0, 0), (0, 1), (1, 1)\}$.

Note 4.1.0.9. When the context is clear, we may write $\leq$ in place of $\leq_1$ and $\leq_2$.

Exercise 4.1.0.10. We have that

1. $(\mathbf{1}, \leq_1)$ is a proset
2. $(\mathbf{2}, \leq_2)$ is a proset

Proof. Clear, FINISH!!!

- 1.
- 2.

□

Talk about presets/posests where the order is induced by a map and another preset/poset, i.e. Let $(Y, \leq)$ be a preset/poset and $f : X \rightarrow Y$. Define $f^* \leq$ on X by $x_1 f^* \leq x_2$ iff $f(x_1) \leq f(x_2)$. Similarly, If $(X, \leq)$ is a preset/poset and $f : X \rightarrow Y$. Define $f_* \leq$ on Y by $y_1 f_* \leq y_2$ iff there exists $x_1 \in f^{-1}(y_1)$ and $x_2 \in f^{-1}(y_2)$ such that $x_1 \leq x_2$. When are these well defined? I think the posetal reflection is an example of the second case. An example of the first would be from a probability, define order on measurable sets $A \leq B$ iff $P(A) \leq P(B)$ so that one is larger than another if it is more likely.

4.2 Subprosets

Definition 4.2.0.1. ref defn in set theory appendices on restrictions of n -ary homogeneous relations

Note 4.2.0.2. Let $(X, \leq)$ be a proset and $A \subset X$. When the context is clear, we write $(A, \leq)$ in place of $(A, \leq|_A)$

4.3 Products

Definition 4.3.0.1. Let $(A, \leq_A)$ and $(B, \leq_B)$ be prosets. We define the

- **product preorder of $\leq_A$ and $\leq_B$ on $A \times B$** , denoted $\leq_A \otimes \leq_B$ by $(a_1, b_1) \leq_A \otimes \leq_B (a_2, b_2)$ iff $a_1 \leq_A a_2$ and $b_1 \leq_B b_2$.
- **product proset of $(A, \leq_A)$ and $(B, \leq_B)$** , denoted $(A, \leq_A) \otimes (B, \leq_B)$ by $(A, \leq_A) \otimes (B, \leq_B) := (A \times B, \leq_A \otimes \leq_B)$

Exercise 4.3.0.2. probably need to change notation since $\otimes$ might be reserved for something else. Let $(A, \leq_A)$ and $(B, \leq_B)$ be prosets. Then

1. $\leq_A \otimes \leq_B$ is a preorder on $A \times B$,
2. $(A, \leq_A) \otimes (B, \leq_B)$ is a proset.

Proof.

1. (a) Let $(a, b) \in A \times B$. Then $a \leq_A a$ and $b \leq_B b$. Therefore $(a, b) \leq_A \otimes \leq_B (a, b)$.
 - (b) Let $(a_1, b_1), (a_2, b_2), (a_3, b_3) \in A \times B$. Suppose that $(a_1, b_1) \leq_A \otimes \leq_B (a_2, b_2)$ and $(a_2, b_2) \leq_A \otimes \leq_B (a_3, b_3)$. Then $a_1 \leq_A a_2$, $a_2 \leq_A a_3$, $b_1 \leq_B b_2$ and $b_2 \leq_B b_3$. Therefore $a_1 \leq_A a_3$ and $b_1 \leq_B b_3$. Hence $(a_1, b_1) \leq_A \otimes \leq_B (a_3, b_3)$.
- Hence $\leq_A \otimes \leq_B$ is a preorder on $A \times B$.
2. Since $\leq_A \otimes \leq_B$ is a preorder on $A \times B$, $(X, \leq)$ is a proset.

□

4.4 Quotients

Chapter 5

Posets

5.1 Introduction

Definition 5.1.0.1. Poset:

Let $(X, \leq)$ be a poset. Then

- $\leq$ is said to be a **partial order on X** if for each $a, b \in X$, $a \leq b$ and $b \leq a$ implies that $a = b$
- $(X, \leq)$ is said to be a **partially ordered set** or **poset** if $\leq$ is a partial ordering on X .

Exercise 5.1.0.2. Let $(X, \leq)$ be a poset and $a, b \in X$. Then $a \sim_{\leq} b$ iff $a = b$.

Proof.

- $(\implies)$:
Suppose that $a \sim_{\leq} b$. Then $a \leq b$ and $b \leq a$. Since $\leq$ is a partial order, $a = b$.
- $(\impliedby)$:
Since $\sim_{\leq}$ is reflexive, $a = b$ implies that $a \sim_{\leq} b$.

□

Definition 5.1.0.3. Let $(X, \leq)$ be a poset. We define the

- **posetal reflection partial order on $X / \sim_{\leq}$** , denoted $\leq_{X / \sim_{\leq}} \subset X / \sim_{\leq} \times X / \sim_{\leq}$, by $y_1 \leq_{X / \sim_{\leq}} y_2$ iff there exists $x_1 \in y_1$ and $x_2 \in y_2$ such that $x_1 \leq x_2$.
- **the posetal reflection of $(X, \leq)$** , denoted $\text{PosRefl}(X, \leq)$, by $\text{PosRefl}(X, \leq) := (X / \sim_{\leq}, \leq_{X / \sim_{\leq}})$

Exercise 5.1.0.4. Let $(X, \leq)$ be a poset. Then $\text{PosRefl}(X, \leq)$ is a poset.

Proof. Set $Y := X / \sim_{\leq}$ and $\leq_Y := \leq_{X / \sim_{\leq}}$.

- (a) Let $y \in Y$. Then there exists $x \in y$. Since $x \leq x$, we have that $y \leq_Y y$.
(b) Let $y_1, y_2, y_3 \in Y$. Suppose that $y_1 \leq_Y y_2$ and $y_2 \leq_Y y_3$. Then there exist $x_1 \in y_1$, $x_2 \in y_2$ and $x_3 \in y_3$ such that $x_1 \leq x_2$ and $x_2 \leq x_3$. Therefore $x_1 \leq x_3$. Hence $y_1 \leq_Y y_3$.

Thus $\leq_Y$ is a preorder on Y .

- Let $y_1, y_2 \in Y$. Suppose that $y_1 \leq_Y y_2$ and $y_2 \leq_Y y_1$. Then there exist $a_1, b_1 \in y_1$, $a_2, b_2 \in y_2$ such that $a_1 \leq a_2$ and $b_2 \leq b_1$. Since $a_1, b_1 \in y_1$ and $a_2, b_2 \in y_2$, $a_1 \sim_{\leq} b_1$ and $a_2 \sim_{\leq} b_2$. Thus

$$\begin{aligned} a_1 &\leq a_2 \\ &\leq b_2. \end{aligned}$$

and

$$\begin{aligned} b_2 &\leq b_1 \\ &\leq a_1. \end{aligned}$$

Hence $a_1 \sim_{\leq} b_2$ and

$$\begin{aligned} y_1 &= \pi(a_1) \\ &= \pi(b_2) \\ &= y_2. \end{aligned}$$

Therefore $\leq_Y$ is a partial order on Y . □

Exercise 5.1.0.5. Let X be a set. Then $(\mathcal{P}(X), \subset)$ is a poset.

Proof. Exercise 4.1.0.7 implies that $(\mathcal{P}(X), \subset)$ is a poset. We note that for each $A, B \in \mathcal{P}(X)$, $A \subset B$ and $B \subset A$ implies that $A = B$ ref def of equality in set theory notes?. □

Exercise 5.1.0.6. We have that

1. $(\mathbf{1}, \leq_1)$ is a poset
2. $(\mathbf{2}, \leq_2)$ is a poset

Proof. Clear, FINISH!!! reference Exercise 4.1.0.10.

- 1.
- 2.

□

Chapter 6

Directed Sets

6.1 Introduction

Definition 6.1.0.1. Directed Set:

Let $(A, \leq)$ be a proset. Then

- $\leq$ is said to be a **direction on** A if for each $\alpha, \beta \in A$, $\text{ub}(\{\alpha, \beta\}, \leq) \neq \emptyset$.
- $(A, \leq)$ is said to be a **directed set** if
 1. $A \neq \emptyset$
 2. $\leq$ is a direction on A

6.2 Subdirected Sets

Definition 6.2.0.1. Let $(A, \leq)$ be a proset and $B \subset A$. Then B is said to be

- **$\leq$ -upward directed** if $(B, \leq|_B)$ is a directed set,
- **$\leq$ -downward directed** if $(B, \leq^*|_B)$ is a directed set.

Note 6.2.0.2. When the context is clear, we may say “upward directed” and “downward directed” instead of “ $\leq$ -upward directed” and “ $\leq$ -downward directed” respectively.

Exercise 6.2.0.3. Let $(A, \leq)$ be a proset and $B \subset A$. Then

1. B is $\leq$ -downward directed iff B is $\leq^*$ -upward directed.
2. B is $\leq$ -upward directed iff B is $\leq^*$ -downward directed.

Proof.

1. We note that

$$\begin{aligned} B \text{ is } \leq\text{-downward directed} &\iff (B, \leq^*|_B) \text{ is a directed set} \\ &\iff B \text{ is } \leq^*\text{-upward directed.} \end{aligned}$$

2. We note that

$$\begin{aligned} B \text{ is } \leq\text{-upward directed} &\iff B \text{ is } (\leq^*)^*\text{-upward directed} && \text{ref ex in set theory section on binary hetero relations about converse r} \\ &\iff B \text{ is } \leq^*\text{-downward directed} && \text{ref previous part.} \end{aligned}$$

□

6.3 Products of Directed Sets

Exercise 6.3.0.1. Let $(A, \leq_A)$ and $(B, \leq_B)$ be directed sets. Then

1. $\leq_A \otimes \leq_B$ is a direction on $A \times B$,
2. $(A, \leq_A) \otimes (B, \leq_B)$ is a directed set.

Proof.

1. (a) Exercise ?? implies that $\leq_A \otimes \leq_B$ is a preorder of $A \times B$.
 (b) Let $(a_1, b_1), (a_2, b_2) \in A \times B$. Then there exist $a \in A$ and $b \in B$ such that $a_1, a_2 \leq_A a$ and $b_1, b_2 \leq_B b$. Hence $(a_1, b_1), (a_2, b_2) \leq_A \otimes \leq_B (a, b)$.

Hence $\leq_A \otimes \leq_B$ is a direction on $A \times B$.

2. (a) Since $A \neq \emptyset$ and $B \neq \emptyset$, we have that $A \times B \neq \emptyset$.
 (b) From above, $\leq_A \otimes \leq_B$ is a direction on $A \times B$.

Hence $(A \times B, \leq_A \otimes \leq_B)$ is a directed set.

□

Chapter 7

Monotone Maps

7.1 Introduction

Definition 7.1.0.1. Let $(X, \leq_X)$ $(Y, \leq_Y)$ be posets.

- $f : X \rightarrow Y$. Then f is said to be
 - $(\leq_X, \leq_Y)$ -**order preserving** or $(\leq_X, \leq_Y)$ -**monotone** if for each $a, b \in X$, $a \leq_X b$ implies that $f(a) \leq_Y f(b)$.
 - $(\leq_X, \leq_Y)$ -**order reversing** or $(\leq_X, \leq_Y)$ -**antitone** if for each $a, b \in X$, $a \leq_X b$ implies that $f(b) \leq_Y f(a)$.
- We define
 - $\text{Mtn}((X, \leq_X), (Y, \leq_Y)) := \{f \in Y^X : f \text{ is } (\leq_X, \leq_Y)\text{-monotone}\},$
 - $\text{Antn}((X, \leq_X), (Y, \leq_Y)) := \{f \in Y^X : f \text{ is } (\leq_X, \leq_Y)\text{-antitone}\}.$

Note 7.1.0.2. When the context is clear

- we may say that f is “monotone” and “antitone” instead of “ $(\leq_X, \leq_Y)$ -monotone” and “ $(\leq_X, \leq_Y)$ -antitone” respectively.
- we may write $\text{Mtn}(X, Y)$ and $\text{Antn}(X, Y)$ in place of $\text{Mtn}((X, \leq_X), (Y, \leq_Y))$ and $\text{Antn}((X, \leq_X), (Y, \leq_Y))$ respectively.

need to go rewrite using notation Mtn and Antn

Exercise 7.1.0.3. Let $(X, \leq_X)$ $(Y, \leq_Y)$ be posets and $f : X \rightarrow Y$. Then

1. the following are equivalent:
 - (a) f is $(\leq_X, \leq_Y)$ -antitone
 - (b) f is $(\leq_X, \leq_Y^*)$ -monotone
 - (c) f is $(\leq_X^*, \leq_Y)$ -monotone
 - (d) f is $(\leq_X^*, \leq_Y^*)$ -antitone
2. the following are equivalent:
 - (a) f is $(\leq_X, \leq_Y)$ -monotone
 - (b) f is $(\leq_X, \leq_Y^*)$ -antitone
 - (c) f is $(\leq_X^*, \leq_Y)$ -antitone
 - (d) f is $(\leq_X^*, \leq_Y^*)$ -monotone

Proof.

1. • (a) $\implies$ (b):
Suppose that f is $(\leq_X, \leq_Y)$ -antitone. Let $a, b \in X$. Then

$$\begin{aligned} a \leq_X b &\implies f(b) \leq_Y f(a) \\ &\iff f(a) \leq_Y^* f(b). \end{aligned}$$

Since $a, b \in X$ are arbitrary, we have that for each $a, b \in X$, $a \leq_X b$ implies that $f(a) \leq_Y^* f(b)$. Hence f is $(\leq_X, \leq_Y^*)$ -monotone.

- (b) $\implies$ (c):
Suppose that f is $(\leq_X, \leq_Y^*)$ -monotone. Let $a, b \in X$. Then

$$\begin{aligned} a \leq_X^* b &\iff b \leq_X^* a \\ &\implies f(b) \leq_Y^* f(a) \\ &\iff f(a) \leq_Y f(b). \end{aligned}$$

Since $a, b \in X$ are arbitrary, we have that for each $a, b \in X$, $a \leq_X^* b$ implies that $f(a) \leq_Y f(b)$. Hence f is $(\leq_X^*, \leq_Y)$ -monotone.

- (c) $\implies$ (d):
Suppose that f is $(\leq_X^*, \leq_Y)$ -monotone. Let $a, b \in X$. Then

$$\begin{aligned} a \leq_X^* b &\implies f(a) \leq_Y f(b) \\ &\iff f(b) \leq_Y^* f(a). \end{aligned}$$

Since $a, b \in X$ are arbitrary, we have that for each $a, b \in X$, $a \leq_X^* b$ implies that $f(b) \leq_Y^* f(a)$. Hence f is $(\leq_X^*, \leq_Y^*)$ -antitone.

- (d) $\implies$ (a):
Suppose that f is $(\leq_X^*, \leq_Y^*)$ -antitone. Let $a, b \in X$. Then

$$\begin{aligned} a \leq_X b &\iff b \leq_X^* a \\ &\implies f(a) \leq_Y^* f(b) \\ &\iff f(b) \leq_Y f(a). \end{aligned}$$

Since $a, b \in X$ are arbitrary, we have that for each $a, b \in X$, $a \leq_X b$ implies that $f(b) \leq_Y f(a)$. Hence f is $(\leq_X, \leq_Y)$ -antitone.

2. Part (1) implies that

$$\begin{aligned} f \text{ is } (\leq_X, (\leq_Y^*)^*)\text{-monotone} &\iff f \text{ is } (\leq_X, \leq_Y^*)\text{-antitone} \quad (\text{by } 1(b) \iff 1(a)) \\ &\iff f \text{ is } (\leq_X^*, (\leq_Y^*)^*)\text{-antitone} \quad (\text{by } 1(a) \iff 1(d)) \\ &\iff f \text{ is } (\leq_X^*, \leq_Y^*)\text{-monotone} \quad (\text{by } 1(d) \iff 1(c)). \end{aligned}$$

Since $(\leq_Y^*)^* = \leq_Y$, we have that

$$\begin{aligned} f \text{ is } (\leq_X, \leq_Y)\text{-monotone} &\iff f \text{ is } (\leq_X, \leq_Y^*)\text{-antitone} \\ &\iff f \text{ is } (\leq_X^*, \leq_Y)\text{-antitone} \\ &\iff f \text{ is } (\leq_X^*, \leq_Y^*)\text{-monotone}. \end{aligned}$$

□

Exercise 7.1.0.4. Let $(X, \leq_X)$ $(Y, \leq_Y)$ be prosets and $f : X \rightarrow Y$. Then f is $(\leq_X, \leq_Y)$ -monotone iff f is $(\leq_X^*, \leq_Y^*)$ -monotone.

Proof.

- ($\Rightarrow$):

Suppose that f is $(\leq_X, \leq_Y)$ -monotone. Let $a, b \in X$. Suppose that $a \leq_X^* b$. Then $b \leq_X a$. Hence $f(b) \leq_Y f(a)$. Thus $f(a) \leq_Y^* f(b)$. Since $a, b \in X$ with $a \leq_X^* b$ are arbitrary, we have that for each $a, b \in X$, $a \leq_X^* b$ implies that $f(a) \leq_Y^* f(b)$. Therefore f is $(\leq_X^*, \leq_Y^*)$ -monotone.

- ($\Leftarrow$):

Suppose that f is $(\leq_X^*, \leq_Y^*)$ -monotone. Since $(\leq_X^*)^* = \leq_X$, the previous part implies that f is $(\leq_X, \leq_Y)$ -monotone.

□

Definition 7.1.0.5. Let $(X, \leq_X)$ and $(Y, \leq_Y)$ be posets and $f : X \rightarrow Y$. Then f is said to be a $(\leq_X, \leq_Y)$ -isomorphism if

1. f is a bijection,
2. f is $(\leq_X, \leq_Y)$ -monotone and f^{-1} is $(\leq_Y, \leq_X)$ -monotone.

Definition 7.1.0.6. Let X be a set and $(Y, \leq)$ a poset. We define the **preorder on Y^X induced by $\leq$** , denoted $\leq_{(Y, \leq)^X}$, by

$$f \leq_{(Y, \leq)^X} g \text{ iff for each } x \in X, f(x) \leq g(x)$$

Note 7.1.0.7. When the context is clear, we may write $\leq$ in place of $\leq_{(Y, \leq)^X}$.

Exercise 7.1.0.8. Let X be a set and $(Y, \leq)$ a poset. Then

1. $(Y^X, \leq_{(Y, \leq)^X})$ is a poset.
2. $(Y, \leq)$ a poset, implies that $(Y^X, \leq_{(Y, \leq)^X})$ is a poset.

Proof. Set $\leq' := \leq_{(Y, \leq)^X}$. Since $(Y, \leq)$ is a poset, $\leq$ is reflexive and transitive.

1.
 - Let $f \in Y^X$ and $x \in X$. Since $\leq$ is reflexive, we have that $f(x) \leq f(x)$. Since $x \in X$ is arbitrary, we have that for each $x \in X$, $f(x) \leq f(x)$. Hence $f \leq' f$. Since $f \in Y^X$ is arbitrary, we have that for each $f \in Y^X$, $f \leq' f$. Thus $\leq'$ is reflexive.
 - Let $f, g, h \in Y^X$ and $x \in X$. Suppose that $f \leq' g$ and $g \leq' h$. Then $f(x) \leq g(x)$ and $g(x) \leq h(x)$. Since $\leq$ is transitive, $f(x) \leq h(x)$. Since $x \in X$ is arbitrary, we have that for each $x \in X$, $f(x) \leq h(x)$. Thus $f \leq' h$. Since $f, g, h \in Y^X$ with $f \leq' g$ and $g \leq' h$ are arbitrary, we have that for each $f, g, h \in Y^X$, $f \leq' g$ and $g \leq' h$ implies that $f \leq' h$. Thus $\leq'$ is transitive.

Hence $\leq'$ is a preorder on Y^X and $(Y^X, \leq')$ is a poset.

2. Suppose that $(Y, \leq)$ is a poset. Then $\leq$ is antisymmetric. Let $f, g \in Y^X$ and $x \in X$. Suppose that $f \leq' g$ and $g \leq' f$. Then $f(x) \leq g(x)$ and $g(x) \leq f(x)$. Since $\leq$ is antisymmetric, $f(x) = g(x)$. Since $x \in X$ is arbitrary, we have that for each $x \in X$, $f(x) = g(x)$. Hence $f = g$. Since $f, g \in Y^X$ with $f \leq' g$ and $g \leq' f$ are arbitrary, we have that for each $f, g \in Y^X$, $f \leq' g$ and $g \leq' f$ implies that $f = g$. Thus $\leq'$ is antisymmetric. Thus $\leq'$ is a partial order on Y^X and $(Y^X, \leq')$ is a poset.

□

7.2 Galois Connections

ToDo: If this comes after groups rings, etc, give examples of defining the 'algebraic object generated by less structured objects' in terms of a galois connection. e.g. the subgroup/subring/upperset/sublattice/ideal/filter/sigma-alg/topology generated, subsets of set/powerset

Definition 7.2.0.1. Let $(X, \leq_X)$ and $(Y, \leq_Y)$ be prosets, $f : X \rightarrow Y$ and $g : Y \rightarrow X$. Then

- (f, g) is said to be a $(\leq_X, \leq_Y)$ -**monotone Galois connection** if for each $x \in X$ and $y \in Y$,

$$f(x) \leq_Y y \text{ iff } x \leq_X g(y),$$

- (f, g) is said to be a $(\leq_X, \leq_Y)$ -**antitone Galois connection** if for each $x \in X$ and $y \in Y$,

$$f(x) \leq_Y y \text{ iff } g(y) \leq_X x,$$

- f is said to be a $(\leq_X, \leq_Y)$ -**left adjoint of g** if (f, g) is a $(\leq_X, \leq_Y)$ -monotone Galois connection,
- g is said to be a $(\leq_X, \leq_Y)$ -**right adjoint of f** if (f, g) is a $(\leq_X, \leq_Y)$ -monotone Galois connection.
- f is said to **have a $(\leq_X, \leq_Y)$ -right adjoint** if there exists $f^* : Y \rightarrow X$ such that f^* is a $(\leq_X, \leq_Y)$ -right adjoint of f .
- g is said to **have a $(\leq_X, \leq_Y)$ -left adjoint** if there exists $g_* : X \rightarrow Y$ such that g_* is a $(\leq_X, \leq_Y)$ -left adjoint of g .

Note 7.2.0.2. When the context is clear, we say

- “monotone (resp. antitone) Galois connection” instead of “ $(\leq_X, \leq_Y)$ -monotone (resp. $(\leq_X, \leq_Y)$ -antitone) Galois connection”
- “left (resp. right) adjoint” instead of “ $(\leq_X, \leq_Y)$ -left (resp. $(\leq_X, \leq_Y)$ -right) adjoint”

Question:

There is an interpretation of prosets, monotonic maps and galois connections in terms of information.

- Prosets are spaces that encode informational direction: $x_1 \leq_X x_2$ iff x_1 contains less information than x_2 .
- Monotonic maps are those that preserve informational direction: if x_1 has less information than x_2 , then $f(x_1)$ has less information than $f(x_2)$.
- Galois connections encode informational direction in different contexts as well as minimal and maximal changes in information: $f(x)$ has less information than y in the context of Y . More specifically,

- * y contains at least as much information as x in the context of Y iff $f(x) \leq_Y y$.
- * $f(x)$ is the element y of Y such that

1. y contains at least as much information as x in the context of Y , i.e. $f(x) \leq_Y y$.

2. for each $y' \in Y$, if y contains at least as much information as x in the context of Y , then y' contains at least as much information as y in the context of Y , i.e. $f(x) \leq_Y y' \implies y \leq_X y'$.

So that $f(x)$ is the object in Y such that the increase in information (information gain) from x to $f(x)$ in the context of Y is minimal. Equivalently, the change in information $\Delta_Y I(x, f(x))$ is minimal

- * x has less information than y in the context of X iff $x \leq_X g(y)$.

- * $g(y)$ is the element x of X such that

1. y contains at least as much information as x in the context of X , i.e. $x \leq_X g(y)$.

2. for each $x' \in X$, if y contains at least as much information as x' in the context of X , then x contains at least as much information as x' in the context of X , i.e. $x' \leq_X g(y) \implies x' \leq_X x$.

So that $g(y)$ is the object in X such that the decrease in information (information loss) from y to $g(y)$ in the context of X is minimal. Equivalently, the change in information $\Delta_X I(y, g(y))$ is maximal (-1 is larger than -2)

This can be generalized to categories, functors and adjoint functors where

- $x_1 \leq_X x_2$ generalizes to $\text{Hom}_X(x_2, x_1) \neq \emptyset$, i.e. x_2 has at least as much information as x_1 generalizes to x_1 is a processed version of x_1 ,
- a monotone map $f : X \rightarrow Y$ generalizes to a functor $F : X \rightarrow Y$
- a Galois connection (f, g) generalizes to an adjunction (F, G) .

This categorical generalization also generalizes information theory and statistics. However, from most math sources, the typical language of adjunctions is in terms of efficiency instead of information change. For example $f(x)$ is the most 'efficient' way to obtain an object in Y from x which 'preserves x '. Similarly, $g(y)$ is the least wasteful way to obtain an object in X from y which ?????????? Could we rephrase this to describe dynamical systems, specifically, can we rephrase information or efficiency in terms of energy to capture describe physics? Something like the path in phase space from t_1 to t_2 is a right adjoint functor from the universe (a category) at a time t_1 to the universe at time t_2 so that the information loss is minimal. Maybe show that least action, or newtons equation is a manifestation of this.

Exercise 7.2.0.3. Let $(X, \leq_X)$ and $(Y, \leq_Y)$ be posets, $f : X \rightarrow Y$ and $g : Y \rightarrow X$. Then

1. (f, g) is a $(\leq_X, \leq_Y)$ -monotone Galois connection iff (g, f) is a $(\leq_Y^*, \leq_X^*)$ -monotone Galois connection
2. (f, g) is a $(\leq_X, \leq_Y)$ -antitone Galois connection iff (f, g) is a $(\leq_X^*, \leq_Y)$ -monotone Galois connection
3. (f, g) is a $(\leq_X, \leq_Y)$ -antitone Galois connection iff (g, f) is a $(\leq_Y, \leq_X)$ -antitone Galois connection

Proof.

1. • $(\implies)$:
Suppose that (f, g) is a $(\leq_X, \leq_Y)$ -monotone Galois connection. Let $y \in Y$ and $x \in X$. Then by assumption,

$$\begin{aligned} g(y) \leq_X^* x &\iff x \leq_X g(y) \\ &\iff f(x) \leq_Y y \\ &\iff y \leq_Y^* f(x). \end{aligned}$$

Since $y \in Y$ and $x \in X$ are arbitrary, we have that for each $y \in Y$ and $x \in X$, $g(y) \leq_X^* x$ iff $y \leq_Y^* f(x)$. Therefore (g, f) is a $(\leq_Y^*, \leq_X^*)$ -monotone Galois connection.

- $(\impliedby)$:
Suppose that (g, f) is a $(\leq_Y^*, \leq_X^*)$ -monotone Galois connection. Let $x \in X$ and $y \in Y$. Then by assumption,

$$\begin{aligned} f(x) \leq_Y y &\iff y \leq_Y^* f(x) \\ &\iff g(y) \leq_X^* x \\ &\iff x \leq_X g(y). \end{aligned}$$

Since $x \in X$ and $y \in Y$ are arbitrary, we have that for each $x \in X$ and $y \in Y$, $f(x) \leq_Y y$ iff $x \leq_X g(y)$. Therefore (f, g) is a $(\leq_X, \leq_Y)$ -monotone Galois connection.

2. • $(\implies)$:
Suppose that (f, g) is a $(\leq_X, \leq_Y)$ -antitone Galois connection. Let $x \in X$ and $y \in Y$. Then by assumption,

$$\begin{aligned} f(x) \leq_Y y &\iff g(y) \leq_X x \\ &\iff x \leq_X^* g(y). \end{aligned}$$

Since $x \in X$ and $y \in Y$ are arbitrary, we have that for each $x \in X$ and $y \in Y$, $f(x) \leq_Y y$ iff $x \leq_X^* g(y)$. Therefore (f, g) is a $(\leq_X^*, \leq_Y)$ -monotone Galois connection.

- $(\impliedby)$:
Suppose that (f, g) is a $(\leq_X^*, \leq_Y)$ -monotone Galois connection. Let $x \in X$ and $y \in Y$. Then by assumption,

$$\begin{aligned} f(x) \leq_Y y &\iff x \leq_X^* g(y) \\ &\iff g(y) \leq_X x. \end{aligned}$$

Since $x \in X$ and $y \in Y$ are arbitrary, we have that for each $x \in X$ and $y \in Y$, $f(x) \leq_Y y$ iff $g(y) \leq_X x$. Therefore (f, g) is a $(\leq_X, \leq_Y)$ -antitone Galois connection.

3. By (1) and (2),

$$\begin{aligned}
 (f, g) \text{ is a } (\leq_X, \leq_Y)\text{-antitone Galois connection} &\iff (f, g) \text{ is a } (\leq_X^*, \leq_Y)\text{-monotone Galois connection} \\
 &\iff (g, f) \text{ is a } (\leq_Y^*, (\leq_X^*)^*)\text{-monotone Galois connection} \\
 &\iff (g, f) \text{ is a } (\leq_Y^*, \leq_X)\text{-monotone Galois connection} \\
 &\iff (g, f) \text{ is a } (\leq_Y, \leq_X)\text{-antitone Galois connection.}
 \end{aligned}$$

□

Exercise 7.2.0.4. Let $(X, \leq_X)$ and $(Y, \leq_Y)$ be posets, $f : X \rightarrow Y$ and $g : Y \rightarrow X$. Then f is a $(\leq_X, \leq_Y)$ -left adjoint of g iff g is a $(\leq_X, \leq_Y)$ -right adjoint of f .

Proof. We have that

$$\begin{aligned}
 f \text{ is a } (\leq_X, \leq_Y)\text{-left adjoint of } g &\iff (f, g) \text{ is a } (\leq_X, \leq_Y)\text{-monotone Galois connection} \\
 &\iff g \text{ is a } (\leq_X, \leq_Y)\text{-right adjoint of } f.
 \end{aligned}$$

□

Exercise 7.2.0.5. Let $(X, \leq_X)$ and $(Y, \leq_Y)$ be posets, $f : X \rightarrow Y$ and $g : Y \rightarrow X$. Then

1. g is a $(\leq_X, \leq_Y)$ -right adjoint of f iff g is a $(\leq_Y^*, \leq_X^*)$ -left adjoint of f .
2. f is a $(\leq_X, \leq_Y)$ -left adjoint of g iff f is a $(\leq_Y^*, \leq_X^*)$ -right adjoint of g .

Proof.

1. Exercise 7.2.0.3 implies that that

$$\begin{aligned}
 g \text{ is a } (\leq_X, \leq_Y)\text{-right adjoint of } f &\iff (f, g) \text{ is a } (\leq_X, \leq_Y)\text{-monotone Galois connection} \\
 &\iff (g, f) \text{ is a } (\leq_Y^*, \leq_X^*)\text{-monotone Galois connection} \\
 &\iff g \text{ is a } (\leq_Y^*, \leq_X^*)\text{-left adjoint of } f.
 \end{aligned}$$

2. We have that

$$\begin{aligned}
 f \text{ is a } (\leq_X, \leq_Y)\text{-left adjoint of } g &\iff g \text{ is a } (\leq_X, \leq_Y)\text{-right adjoint of } f \quad (\text{by Exercise 7.2.0.4}) \\
 &\iff g \text{ is a } (\leq_Y^*, \leq_X^*)\text{-left adjoint of } f \quad (\text{by prev part}) \\
 &\iff f \text{ is a } (\leq_Y^*, \leq_X^*)\text{-right adjoint of } g \quad (\text{by Exercise 7.2.0.4}).
 \end{aligned}$$

□

Exercise 7.2.0.6. Let $(X, \leq_X)$ and $(Y, \leq_Y)$ be posets, $f : X \rightarrow Y$ and $g : Y \rightarrow X$. Then

1. (f, g) is a $(\leq_X, \leq_Y)$ -monotone Galois connection iff
 - (a) i. $\text{id}_X \leq_X g \circ f$
 - ii. $f \circ g \leq_Y \text{id}_Y$
 - (b) f is $(\leq_X, \leq_Y)$ -monotone and g is $(\leq_Y, \leq_X)$ -monotone
2. (f, g) is a $(\leq_X, \leq_Y)$ -antitone Galois connection iff
 - (a) i. $g \circ f \leq_X \text{id}_X$
 - ii. $f \circ g \leq_Y \text{id}_Y$
 - (b) f is $(\leq_X, \leq_Y)$ -antitone and g is $(\leq_Y, \leq_X)$ -antitone

Proof.

1. • ($\Rightarrow$):

Suppose that (f, g) is a $(\leq_X, \leq_Y)$ -monotone Galois connection.

- (a) i. Let $x \in X$. Define $y \in Y$ by $y := f(x)$. Since $f(x) \leq_Y f(x)$, we have that $f(x) \leq_Y y$. Therefore

$$\begin{aligned} x &\leq_X g(y) \\ &= g \circ f(x). \end{aligned}$$

Since $x \in X$ is arbitrary, we have that for each $x \in X$, $x \leq_X g \circ f(x)$. Hence $\text{id}_X \leq_X g \circ f$.

- ii. Let $y \in Y$. **the previous ex** implies that (g, f) is a $(\leq_Y^*, \leq_X^*)$ -monotone Galois connection. Part 1(a) then implies that for each $y \in Y$, $y \leq_Y^* f \circ g(y)$. Therefore for each $y \in Y$, $f \circ g(y) \leq_Y y$. Hence $f \circ g \leq_Y \text{id}_Y$.
- (b) – Let $a, b \in X$. Define $x \in X$ and $y \in Y$ by $x := a$ and $y := f(b)$. Suppose that $a \leq_X b$. Part 1(a) implies that

$$\begin{aligned} x &= a \\ &\leq_X b \\ &\leq_X g(f(x)) \\ &= g(y). \end{aligned}$$

Since (f, g) is a $(\leq_X, \leq_Y)$ -monotone Galois connection, we have that

$$\begin{aligned} f(a) &= f(x) \\ &\leq_Y y \\ &= f(b). \end{aligned}$$

Since $a, b \in X$ with $a \leq_X b$ are arbitrary, we have that for each $a, b \in X$, $a \leq_X b$ implies that $f(a) \leq_Y f(b)$. Thus f is $(\leq_X, \leq_Y)$ -monotone.

- Since (f, g) is a $(\leq_X, \leq_Y)$ -monotone Galois connection, **a previous ex** implies that (g, f) is a $(\leq_Y^*, \leq_X^*)$ -monotone Galois connection. The previous part implies that g is $(\leq_Y^*, \leq_X^*)$ -monotone. Exercise 7.1.0.3 then implies that g is $(\leq_Y, \leq_X)$ -monotone.

• ($\Leftarrow$):

Suppose that

- (a) i. for each $x \in X$, $x \leq_X g \circ f(x)$
 ii. for each $y \in Y$, $f \circ g(y) \leq_Y y$
 (b) f is $(\leq_X, \leq_Y)$ -monotone and g is $(\leq_Y, \leq_X)$ -monotone.

Let $x \in X$ and $y \in Y$.

– ($\Rightarrow$):

Suppose that $f(x) \leq_Y y$. Then by (a)i and (a)ii,

$$\begin{aligned} x &\leq_X g \circ f(x) \\ &\leq_X g(y). \end{aligned}$$

– ($\Leftarrow$):

Suppose that $x \leq_X g(y)$. Then by (a)i and (a)ii,

$$\begin{aligned} f(x) &\leq_Y f \circ g(y) \\ &\leq_Y y. \end{aligned}$$

Since $x \in X$ and $y \in Y$ are arbitrary, we have that for each $x \in X$ and $y \in Y$,

$$f(x) \leq_Y y \text{ iff } f(x) \leq_Y y.$$

Therefore (f, g) is a $(\leq_X, \leq_Y)$ -monotone Galois connection.

2. **the previous ex** implies that (f, g) is a $(\leq_X, \leq_Y)$ -antitone Galois connection iff (f, g) is a $(\leq_X^*, \leq_Y)$ -monotone Galois connection. Part (1) implies that (f, g) is a $(\leq_X^*, \leq_Y)$ -monotone Galois connection iff

- (a) i. for each $x \in X$, $x \leq_X^* g \circ f(x)$
 ii. for each $y \in Y$, $f \circ g(y) \leq_Y y$
- (b) f is $(\leq_X^*, \leq_Y)$ -monotone and g is $(\leq_Y, \leq_X^*)$ -monotone.

two previous exercise about monotonic functions and the dual order then imply that (f, g) is a $(\leq_X, \leq_Y)$ -antitone Galois connection iff

- (a) i. for each $x \in X$, $g \circ f(x) \leq_X x$
 ii. for each $y \in Y$, $f \circ g(y) \leq_Y y$
- (b) f is $(\leq_X, \leq_Y)$ -antitone and g is $(\leq_Y, \leq_X)$ -antitone.

□

Exercise 7.2.0.7. Let $(X, \leq_X)$ and $(Y, \leq_Y)$ be posets, $f : X \rightarrow Y$ and $g : Y \rightarrow X$. If (f, g) is a $(\leq_X, \leq_Y)$ -monotone Galois connection or (f, g) is a $(\leq_X, \leq_Y)$ -antitone Galois connection, then $f \sim_{\leq_Y} f \circ g \circ f$ and $g \sim_{\leq_X} g \circ f \circ g$.

Proof.

- Suppose that (f, g) a $(\leq_X, \leq_Y)$ -monotone Galois connection.

- Let $x \in X$.

- * Define $a \in X$ and $b \in Y$ by $a := g \circ f(x)$ and $b := f(x)$. Since

$$\begin{aligned} g \circ f(x) \leq g \circ f(x) &\iff a \leq_X g(b) \\ &\iff f(a) \leq_Y b \\ &\iff f \circ g \circ f(x) \leq_Y f(x), \end{aligned}$$

we have that $f \circ g \circ f(x) \leq_Y f(x)$.

- * **The prev ex** implies that $x \leq_X g \circ f(x)$ and f is $(\leq_X, \leq_Y)$ -monotone. Therefore $f(x) \leq_Y f \circ g \circ f(x)$.

Since $f \circ g \circ f(x) \leq_Y f(x)$ and $f(x) \leq_Y f \circ g \circ f(x)$, we have that $f \circ g \circ f(x) \sim_{\leq_Y} f(x)$. Since $x \in X$ is arbitrary, we have that for each $x \in X$, $f \circ g \circ f(x) \sim_{\leq_Y} f(x)$. Therefore $f \sim_{\leq_Y} f \circ g \circ f$.

- Let $y \in Y$.

- * Define $a \in X$ and $b \in Y$ by $a := g(y)$ and $b := f \circ g(y)$. Since

$$\begin{aligned} f \circ g(y) \leq_Y f \circ g(y) &\iff f(a) \leq_Y b \\ &\iff a \leq_X g(b) \\ &\iff g(y) \leq_X g \circ f \circ g(y), \end{aligned}$$

we have that $g(y) \leq_X g \circ f \circ g(y)$.

- * **The prev ex** implies that $f \circ g(y) \leq_Y y$ and g is $(\leq_Y, \leq_X)$ -monotone. Therefore $g \circ f \circ g(y) \leq_X g(y)$.

Since $g(y) \leq_X g \circ f \circ g(y)$ and $g \circ f \circ g(y) \leq_X g(y)$, we have that $g(y) \sim_{\leq_X} g \circ f \circ g(y)$. Since $y \in Y$ is arbitrary, we have that for each $y \in Y$, $g(y) \sim_{\leq_X} g \circ f \circ g(y)$. Therefore $g \sim_{\leq_X} g \circ f \circ g$.

- Suppose that (f, g) a $(\leq_X, \leq_Y)$ -antitone Galois connection. **The prev ex** then implies that (f, g) is a $(\leq_X^*, \leq_Y)$ -monotone Galois connection. Part (1) then implies that $f \sim_{\leq_Y} f \circ g \circ f$ and $g \sim_{\leq_X^*} g \circ f \circ g$. **Make previous ex that $x \sim_{\leq} y$ iff $x \sim_{\leq^*} y$** then implies that $g \sim_{\leq_X} g \circ f \circ g$.

FINSIH!!

□

Exercise 7.2.0.8. Let $(X, \leq_X)$ and $(Y, \leq_Y)$ be posets, $f : X \rightarrow Y$ and $g : Y \rightarrow X$. If (f, g) is a $(\leq_X, \leq_Y)$ -monotone Galois connection or (f, g) is a $(\leq_X, \leq_Y)$ -antitone Galois connection, then $f = f \circ g \circ f$ and $g = g \circ f \circ g$.

Proof. Suppose that (f, g) is a $(\leq_X, \leq_Y)$ -monotone Galois connection or (f, g) is a $(\leq_X, \leq_Y)$ -antitone Galois connection. Exercise 7.2.0.7 implies that $f \sim_{\leq_Y} f \circ g \circ f$ and $g \sim_{\leq_X} g \circ f \circ g$. Since $(X, \leq_X)$ and $(Y, \leq_Y)$ are posets, we have that $f = f \circ g \circ f$ and $g = g \circ f \circ g$. □

Exercise 7.2.0.9. Let $(X, \leq_X)$ and $(Y, \leq_Y)$ be posets, $f : X \rightarrow Y$ and $g : Y \rightarrow X$.

1. For each $g' : Y \rightarrow X$, if g and g' are right adjoints of f , then $g = g'$.
2. For each $f' : X \rightarrow Y$, if f and f' are left adjoints of g , then $f = f'$.

Proof.

1. Let $g' : Y \rightarrow X$. Suppose that g and g' are right adjoints of f . Then (f, g) and (f, g') are Galois connections. Exercise 7.2.0.8 implies that $f \circ g' \circ f = f$ and for each $x \in X$, $g' \circ f(x) \leq_Y g \circ f(g' \circ f(x))$. Therefore for each $x \in X$,

$$\begin{aligned} g' \circ f(x) &\leq g \circ f(g' \circ f(x)) \\ &= g(f \circ g' \circ f(x)) \\ &= g(f(x)) \\ &= g \circ f(x). \end{aligned}$$

Thus $g' \circ f \leq g \circ f$. Similarly, $g \circ f \leq g' \circ f$. Hence $g' \circ f = g \circ f$. Therefore

$$\begin{aligned} g &= g \circ f \circ g \\ &= g' \circ f \circ g. \end{aligned}$$

and

$$\begin{aligned} g' &= g' \circ f \circ g' \\ &= g \circ f \circ g'. \end{aligned}$$

Let $y \in Y$. Since $f \circ g'(y) \leq y$, we have that

$$\begin{aligned} g'(y) &= g \circ f \circ g'(y) \\ &= g(f \circ g'(y)) \\ &\leq g(y). \end{aligned}$$

Similarly, $g(y) \leq g'(y)$. Hence $g(y) = g'(y)$. Since $y \in Y$ is arbitrary, we have that for each $y \in Y$, $g(y) = g'(y)$. Thus $g = g'$.

2. Similar to (1).

TIDY UP!!!!

□

Definition 7.2.0.10. Let $(X, \leq_X)$ and $(Y, \leq_Y)$ be posets, $f : X \rightarrow Y$ and $g : Y \rightarrow X$.

- If f has a $(\leq_X, \leq_Y)$ -right adjoint, then we define the $(\leq_X, \leq_Y)$ -**right adjoint of f** , denoted $\text{RAdj}(f, (\leq_X, \leq_Y))$, to be the unique $g' : Y \rightarrow X$ such that g' is a $(\leq_X, \leq_Y)$ -right adjoint of f .
- If g has a $(\leq_X, \leq_Y)$ -left adjoint, then we define the $(\leq_X, \leq_Y)$ -**left adjoint of g** , denoted $\text{LAdj}(g, (\leq_X, \leq_Y))$, to be the unique $f' : X \rightarrow Y$ such that f' is a $(\leq_X, \leq_Y)$ -left adjoint of g .

Note 7.2.0.11. When the context is clear, we write f^* and g_* in place of $\text{RAdj}(f, (\leq_X, \leq_Y))$ and $\text{LAdj}(g, (\leq_X, \leq_Y))$ respectively.

Exercise 7.2.0.12. Let $(X, \leq_X)$, $(Y, \leq_Y)$, $(Z, \leq_Z)$ be posets, $f : X \rightarrow Y$ and $g : Y \rightarrow Z$.

1. If f has a right adjoint and g has a right adjoint, then $g \circ f$ has a right adjoint and $(g \circ f)^* = f^* \circ g^*$.
2. If f has a left adjoint and g has a left adjoint, then $g \circ f$ has a left adjoint and $(g \circ f)_* = f_* \circ g_*$.

Proof.

1. Suppose that f has a $(\leq_X, \leq_Y)$ -right adjoint and g has a $(\leq_X, \leq_Y)$ -right adjoint. Since (f, f^*) is a $(\leq_X, \leq_Y)$ -monotone Galois connection and (g, g^*) is a $(\leq_Y, \leq_Z)$ -monotone Galois connection, we have that for each $x \in X$, $y \in Y$ and $z \in Z$, $x \leq_X f^*(y)$ iff $f(x) \leq_Y y$ and $y \leq_Y g^*(z)$ iff $g(y) \leq_Z z$. Hence for each $x \in X$ and $z \in Z$,

$$\begin{aligned} x \leq_X f^* \circ g^*(z) &\iff x \leq_X f^*(g^*(z)) \\ &\iff f(x) \leq_Y g^*(z) \\ &\iff g(f(x)) \leq_Z z \\ &\iff g \circ f(x) \leq_Z z. \end{aligned}$$

Thus $(g \circ f, f^* \circ g^*)$ is a $(\leq_X, \leq_Z)$ -monotone Galois connection. Therefore $g \circ f$ has a $(\leq_X, \leq_Z)$ -right adjoint and Exercise 7.2.0.9 implies that $(g \circ f)^* = f^* \circ g^*$.

2. Suppose that f has a $(\leq_Y, \leq_X)$ -left adjoint and g has a $(\leq_Z, \leq_Y)$ -left adjoint. Exercise 7.2.0.5 implies that $\text{LAdj}(f, (\leq_Y, \leq_X)) = \text{RAdj}(f, (\leq_X^*, \leq_Y^*))$ and $\text{LAdj}(g, (\leq_Z, \leq_Y)) = \text{RAdj}(g, (\leq_Y^*, \leq_Z^*))$. The previous part implies that $g \circ f$ has a $(\leq_X^*, \leq_Z^*)$ -right adjoint and $\text{RAdj}(g \circ f, (\leq_X^*, \leq_Z^*)) = \text{LAdj}(f, (\leq_Y, \leq_X)) \circ \text{LAdj}(g, (\leq_Z, \leq_Y))$. Exercise 7.2.0.5 then implies that $g \circ f$ has a $(\leq_Z, \leq_X)$ -left adjoint and

$$\begin{aligned} \text{LAdj}(g \circ f, (\leq_Z, \leq_X)) &= \text{RAdj}(g \circ f, (\leq_X^*, \leq_Z^*)) \\ &= \text{LAdj}(f, (\leq_Y, \leq_X)) \circ \text{LAdj}(g, (\leq_Z, \leq_Y)). \end{aligned}$$

□

Exercise 7.2.0.13. Let X, Y be sets and $f : X \rightarrow Y$. We define $f_* : \mathcal{P}(X) \rightarrow \mathcal{P}(Y)$ and $f^* : \mathcal{P}(Y) \rightarrow \mathcal{P}(X)$ by $f_*(A) := f(A)$ and $f^*(B) := f^{-1}(B)$. Then (f_*, f^*) is a $(\subset_X, \subset_Y)$ -monotone Galois connection.

Proof. We note that for each $A \in \mathcal{P}(X)$ and $B \in \mathcal{P}(Y)$,

$$\begin{aligned} f_*(A) \subset_Y B &\iff f(A) \subset_Y B \\ &\iff f^{-1}(f(A)) \subset_X f^{-1}(B) \quad \text{by (ref ex in set theory notes)} \\ &\iff A \subset_X f^{-1}(B) \quad \text{by (ref ex in set theory notes)} \\ &\iff A \subset_X f^*(B). \end{aligned}$$

Thus (f_*, f^*) is a $(\subset_X, \subset_Y)$ -monotone Galois connection. □

Definition 7.2.0.14. maybe get rid of, ask if there is any point in rephrasing/renaming, possibly move right after definition of gallois connections, with discussion on 'information' Let $(X, \leq_X)$ and $(Y, \leq_Y)$ be posets, $f : X \rightarrow Y$ and $g : Y \rightarrow X$. Suppose that (f, g) is a $(\leq_X, \leq_Y)$ -monotone Galois connection.

- Let $x \in X$. We define the **right (f, g) -connection of x** , denoted $\text{RConn}_Y(x, (f, g)) \in Y$, by $\text{RConn}_Y(x, (f, g)) := f(x)$.
- Let $y \in Y$. We define the **left (f, g) -connection of y** , denoted $\text{LConn}_X(y, (f, g)) \in X$, by $\text{LConn}_X(y, (f, g)) := g(y)$.

Chapter 8

Upper and Lower Sets

8.1 Introduction

Definition 8.1.0.1. Let $(X, \leq)$ be a proset and $A \subset X$. Then

- A is said to be a **$\leq$ -upper set** if for each $a \in A$ and $x \in X$, $a \leq x$ implies that $x \in A$.
- A is said to be an **$\leq$ -lower set** if for each $a \in A$ and $x \in X$, $x \leq a$ implies that $x \in A$.

Note 8.1.0.2. When the context is clear, we say A is a

- “upper set” instead of “ $\leq$ -upper set”
- “lower set” instead of “ $\leq$ -lower set”

Exercise 8.1.0.3. Let $(X, \leq)$ be a proset. Then

1. X is a $\leq$ -upper set
2. X is a $\leq$ -lower set

Proof.

1. Let $a, x \in X$. Suppose that $a \leq x$. By assumption, $x \in X$. Since $a, x \in X$ with $a \leq x$ are arbitrary, we have that for each $a, x \in X$, $a \leq x$ implies that $x \in A$. Hence X is a $\leq$ -upper set.
2. Similar to (1).

□

Exercise 8.1.0.4. Let $(X, \leq)$ be a proset and $A \subset X$. Then

1. A is a $\leq$ -upper set iff A is a $\leq^*$ -lower set
2. A is a $\leq$ -lower set iff A is a $\leq^*$ -upper set

Proof.

1. • ($\implies$):
Suppose that A is a $\leq$ -upper set. Let $a \in A$ and $x \in X$. Suppose that $x \leq^* a$. Then $a \leq x$. Since A is a $\leq$ -upper set, $x \in A$. Since $a \in A$ and $x \in X$ with $x \leq^* a$ is arbitrary, we have that for each $a \in A$ and $x \in X$, $x \leq^* a$ implies that $x \in A$. Hence A is a $\leq^*$ -lower set.
- ($\impliedby$):
Suppose that A is a $\leq^*$ -lower set. Let $a \in A$ and $x \in X$. Suppose that $a \leq x$. Then $x \leq^* a$. Since A is a $\leq^*$ -lower set, $x \in A$. Since $a \in A$ and $x \in X$ with $a \leq x$ is arbitrary, we have that for each $a \in A$ and $x \in X$, $a \leq x$ implies that $x \in A$. Hence A is a $\leq$ -upper set.

2. Similar to (1). **USE DUALITY**

□

Exercise 8.1.0.5. Let $(X, \leq)$ be a proset and $A \subset X$. Then

1. A is an upper set iff A^c is a lower set
2. A is a lower set iff A^c is an upper set

Proof.

1. • ($\Rightarrow$):
Suppose that A is an upper set. Let $b \in A^c$ and $x \in X$. Suppose that $x \leq b$. For the sake of contradiction, suppose that $x \in A$. Since $x \leq b$ and A is an upper set, $b \in A$. This is a contradiction since $b \in A^c$. Hence $x \in A^c$. Since $b \in A^c$ and $x \in X$ with $x \leq b$ are arbitrary, we have that for each $b \in A^c$ and $x \in X$, $x \leq b$ implies that $x \in A^c$. Thus A^c is a lower set.
- ($\Leftarrow$):
Suppose that A^c is a $\leq$ -lower set. Exercise 8.1.0.4 implies that A^c is a $\leq^*$ -upper set. The previous part implies that A is a $\leq^*$ -lower set. Another application of Exercise 8.1.0.4 implies that A is a $\leq$ -upper set.
2. Similar to (1).

□

Exercise 8.1.0.6. Let $(X, \leq)$ be a proset, $(E_\alpha)_{\alpha \in A} \subset \mathcal{P}(X)$.

1. If for each $\alpha \in A$, E_α is an upper set, then

- (a) $\bigcup_{\alpha \in A} E_\alpha$ is an upper set
- (b) $\bigcap_{\alpha \in A} E_\alpha$ is an upper set

2. If for each $\alpha \in A$, E_α is a lower set, then

- (a) $\bigcup_{\alpha \in A} E_\alpha$ is a lower set
- (b) $\bigcap_{\alpha \in A} E_\alpha$ is a lower set

Proof.

1. Suppose that for each $\alpha \in A$, E_α is an upper set. Set $E := \bigcup_{\alpha \in A} E_\alpha$.

- (a) Let $e \in E$ and $x \in X$. Suppose that $e \leq x$. Since $e \in E$, there exists $\alpha \in A$ such that $e \in E_\alpha$. Since E_α is an upper set and $e \leq x$, we have that

$$\begin{aligned} x &\in E_\alpha \\ &\subset \bigcup_{\alpha \in A} E_\alpha \\ &= E. \end{aligned}$$

Since $e \in E$ and $x \in X$ with $e \leq x$ are arbitrary, we have that for each $e \in E$ and $x \in X$, $e \leq x$ implies that $x \in E$. Hence E is an upper set.

- (b) Let $\alpha \in A$. Since E_α is a $\leq$ -upper set, Exercise 8.1.0.5 implies that E_α^c is a $\leq$ -lower set. Exercise 8.1.0.4 then implies that E_α^c is a $\leq^*$ -upper set. Since $\alpha \in A$ is arbitrary, we have that for each $\alpha \in A$, E_α^c is a $\leq^*$ -upper set. The previous part implies that $\bigcup_{\alpha \in A} E_\alpha^c$ is a $\leq^*$ -upper set. Since $\bigcap_{\alpha \in A} E_\alpha = \left(\bigcup_{\alpha \in A} E_\alpha^c \right)^c$, Exercise 8.1.0.5 implies that $\bigcap_{\alpha \in A} E_\alpha$ is a $\leq^*$ -lower set. Exercise 8.1.0.4 then implies that $\bigcap_{\alpha \in A} E_\alpha$ is a $\leq$ -upper set.

2. Similar to (1).

□

Definition 8.1.0.7. Let $(X, \leq)$ be a poset and $S \subset X$. (define upper set generated by- as the generated element from a galois connection, then make exercise showing that $\uparrow(S, \leq) := \bigcap_{U \in \mathcal{U}(S, \leq)} U$) We define the

- $\leq$ -upper sets containing S , denoted $\mathcal{U}(S, \leq) \subset \mathcal{P}(X)$, by $\mathcal{U}(S, \leq) := \{U \subset X : U \text{ is a } \leq\text{-upper set and } S \subset U\}$
- $\leq$ -lower sets containing S , denoted $\mathcal{L}(S, \leq) \subset \mathcal{P}(X)$, by $\mathcal{L}(S, \leq) := \{L \subset X : L \text{ is a } \leq\text{-lower set and } S \subset L\}$.
- $\leq$ -upper set generated by S , denoted $\uparrow(S, \leq) \subset X$, by $\uparrow(S, \leq) := \bigcap_{U \in \mathcal{U}(S, \leq)} U$
- $\leq$ -lower set generated by S , denoted $\downarrow(S, \leq) \subset X$, by $\downarrow(S, \leq) := \bigcap_{L \in \mathcal{L}(S, \leq)} L$

Note 8.1.0.8.

- When the context is clear, we write
 - $\mathcal{U}(S), \mathcal{L}(S), \uparrow S$ and $\downarrow S$ in place of $\mathcal{U}(S, \leq), \mathcal{L}(S, \leq), \uparrow(S, \leq)$ and $\downarrow(S, \leq)$ respectively.
 - $\mathcal{U}(\leq)$ or $\mathcal{U}_X$ in place of $\mathcal{U}(\emptyset, \leq)$
 - $\mathcal{L}(\leq)$ or $\mathcal{L}_X$ in place of $\mathcal{L}(\emptyset, \leq)$.
- If $S = \{s\}$, we write $\uparrow s$ and $\downarrow s$ in place of $\uparrow S$ and $\downarrow S$ respectively.
- Exercise 8.1.0.3 implies that $X \in \mathcal{U}(S)$ and $X \in \mathcal{L}(S)$.
- Exercise 8.1.0.6 implies that $\uparrow S \in \mathcal{U}(S)$ and $\downarrow S \in \mathcal{L}(S)$.

Exercise 8.1.0.9. Let $(X, \leq)$ be a poset and $S \subset X$. Then

1. $\mathcal{U}(S, \leq^*) = \mathcal{L}(S, \leq)$,
2. $\mathcal{L}(S, \leq^*) = \mathcal{U}(S, \leq)$,
3. $\uparrow(S, \leq^*) = \downarrow(S, \leq)$,
4. $\downarrow(S, \leq^*) = \uparrow(S, \leq)$.

Proof.

1. • Let $L \in \mathcal{U}(S, \leq^*)$. Then $S \subset L$ and L is a $\leq^*$ -upper set. Exercise 8.1.0.4 then implies that L is a $\leq$ -lower set. Hence $L \in \mathcal{L}(S, \leq)$. Since $L \in \mathcal{U}(S, \leq^*)$ is arbitrary, we have that for each $L \in \mathcal{U}(S, \leq^*)$, $L \in \mathcal{L}(S, \leq)$. Thus $\mathcal{U}(S, \leq^*) \subset \mathcal{L}(S, \leq)$.
 • Let $L \in \mathcal{L}(S, \leq)$. Then $S \subset L$ and L is a $\leq$ -lower set. Another application of Exercise 8.1.0.4 implies that L is a $\leq^*$ -upper set. Hence $L \in \mathcal{U}(S, \leq^*)$. Since $L \in \mathcal{L}(S, \leq)$ is arbitrary, we have that for each $L \in \mathcal{L}(S, \leq)$, $L \in \mathcal{U}(S, \leq^*)$. Thus $\mathcal{L}(S, \leq) \subset \mathcal{U}(S, \leq^*)$.

Since $\mathcal{U}(S, \leq^*) \subset \mathcal{L}(S, \leq)$ and $\mathcal{L}(S, \leq) \subset \mathcal{U}(S, \leq^*)$, we have that $\mathcal{U}(S, \leq^*) = \mathcal{L}(S, \leq)$.

2. By (1), we have that

$$\begin{aligned} \mathcal{L}(S, \leq^*) &= \mathcal{U}(S, (\leq^*)^*) \\ &= \mathcal{U}(S, \leq). \end{aligned}$$

3. Part (1) implies that

$$\begin{aligned} \uparrow(S, \leq^*) &= \bigcap_{U \in \mathcal{U}(S, \leq^*)} U \\ &= \bigcap_{U \in \mathcal{L}(S, \leq)} U \\ &= \downarrow(S, \leq^*). \end{aligned}$$

4. Similar to (3).

□

Exercise 8.1.0.10. Let $(X, \leq)$ be a proset and $S \subset X$. Then

1. $S \in \mathcal{U}(S, \leq)$ iff $\uparrow S = S$
2. $S \in \mathcal{L}(S, \leq)$ iff $\downarrow S = S$

Proof.

1. • ($\implies$):
Suppose that $S \in \mathcal{U}(S, \leq)$. Then

$$\begin{aligned}\uparrow S &= \bigcap_{U \in \mathcal{U}(S, \leq)} U \\ &\subset S \\ &\subset \uparrow S.\end{aligned}$$

Hence $\uparrow S = S$.

- ($\impliedby$):
Suppose that $\uparrow S = S$. Then

$$\begin{aligned}S &= \uparrow S \\ &\in \mathcal{U}(S, \leq).\end{aligned}$$

2. Similar to (1).

□

Exercise 8.1.0.11. Let $(X, \leq)$ be a proset and $a \in X$. Then

1. $\uparrow a = \{x \in X : a \leq x\}$
2. $\downarrow a = \{x \in X : x \leq a\}$

Proof.

1. Define $U \subset X$ by $U := \{x \in X : a \leq x\}$.
• Let $u \in U$ and $x \in X$. Suppose that $u \leq x$. Then

$$\begin{aligned}a &\leq u \\ &\leq x.\end{aligned}$$

Hence $x \in U$. Since $u \in U$ and $x \in X$ with $u \leq x$ are arbitrary, we have that for each $u \in U$ and $x \in X$, $u \leq x$ implies that $x \in U$. Hence U is an upper set. Since $a \leq a$, $a \in U$. By definition, $U \in \mathcal{U}(a)$. Therefore

$$\begin{aligned}\uparrow a &= \bigcap_{U' \in \mathcal{U}(a)} U' \\ &\subset U.\end{aligned}$$

- Let $x \in U$ and $U' \in \mathcal{U}(a)$. Since $x \in U$, $a \leq x$. Since $U' \in \mathcal{U}(a)$, U' is an upper set and $a \in U'$. Thus $x \in U'$. Since $U' \in \mathcal{U}(a)$ is arbitrary, we have that for each $U' \in \mathcal{U}(a)$, $x \in U'$. Thus

$$\begin{aligned}x &\in \bigcap_{U' \in \mathcal{U}(a)} U' \\ &= \uparrow a.\end{aligned}$$

Since $x \in U$ is arbitrary, we have that for each $x \in U$, $x \in \uparrow a$. Hence $U \subset \uparrow a$.

Since $\uparrow a \subset U$ and $U \subset \uparrow a$, we have that $\uparrow a = U$.

2. Exercise 8.1.0.4 and (1) imply that

$$\begin{aligned}\downarrow(a, \leq) &= \uparrow(a, \leq^*) \\ &= \{x \in X : a \leq^* x\} \\ &= \{x \in X : x \leq a\}.\end{aligned}$$

□

Exercise 8.1.0.12. Let $(X, \leq)$ be a proset and $a, b \in X$.

1. If $\uparrow a = \uparrow b$, then $a \sim_{\leq} b$
2. If $\downarrow a = \downarrow b$, then $a \sim_{\leq} b$

follow up later with one about posets

Proof.

1. Suppose that $\uparrow a = \uparrow b$. Then

$$\begin{aligned}a &\in \uparrow a \\ &= \uparrow b.\end{aligned}$$

Hence $a \leq b$. Similarly, $b \leq a$. Hence $a \sim_{\leq} b$.

- 2.

□

Exercise 8.1.0.13. Let $(X, \leq)$ be a poset and $a, b \in X$.

1. If $\uparrow a = \uparrow b$, then $a = b$
2. If $\downarrow a = \downarrow b$, then $a = b$

Proof.

1. Suppose that $\uparrow a = \uparrow b$. Exercise 8.1.0.12 then implies that $a \sim_{\leq} b$. Since $(X, \leq)$ is a poset, we have that $a = b$.
2. Similar to (1).

□

Exercise 8.1.0.14. Let $(X, \leq)$ be a proset and $S_1, S_2 \subset X$. If $S_1 \subset S_2$, then

1. $\uparrow S_1 \subset \uparrow S_2$
2. $\downarrow S_1 \subset \downarrow S_2$

Proof. Suppose that $S_1 \subset S_2$.

1. Since $\uparrow S_2 \in \mathcal{U}(S_2)$, $S_2 \subset \uparrow S_2$ and $\uparrow S_2$ is an upper set. Since

$$\begin{aligned}S_1 &\subset S_2 \\ &\subset \uparrow S_2,\end{aligned}$$

we have that $\uparrow S_2 \in \mathcal{U}(S_1)$. Therefore

$$\begin{aligned}\uparrow S_1 &= \bigcap_{U \in \mathcal{U}(S_1)} U \\ &\subset \uparrow S_2.\end{aligned}$$

2. Part (1) and Exercise 8.1.0.9 implies that

$$\begin{aligned}\downarrow(S_1, \leq) &= \uparrow(S_1, \leq^*) \\ &\subset \uparrow(S_2, \leq^*) \\ &= \downarrow(S_2, \leq).\end{aligned}$$

□

Exercise 8.1.0.15. Let $(X, \leq)$ be a proset and $a, b \in X$. Then the following are equivalent:

1. $a \leq b$,
2. $\uparrow b \subset \uparrow a$,
3. $\downarrow a \subset \downarrow b$.

Proof.

1. (1) $\implies$ (2):

Suppose that $a \leq b$. Since $\uparrow a \in \mathcal{U}(a)$, we have that $a \in \uparrow a$ and $\uparrow a$ is an upper set. Since $a \leq b$, we have that $b \in \uparrow a$. Hence $\uparrow a \in \mathcal{U}(b)$. Therefore

$$\begin{aligned}\uparrow b &= \bigcap_{U \in \mathcal{U}(b)} U \\ &\subset \uparrow a.\end{aligned}$$

2. (2) $\implies$ (3):

Suppose that $\uparrow(b, \leq) \subset \uparrow(a, \leq)$. Exercise 8.1.0.11 then implies that

$$\begin{aligned}b &\in \uparrow(b, \leq) \\ &\subset \uparrow(a, \leq) \\ &= \{x \in X : a \leq x\}.\end{aligned}$$

Hence $a \leq b$. Thus $b \leq^* a$. Exercise 8.1.0.9 and part (1) $\implies$ (2) then imply that

$$\begin{aligned}\downarrow(a, \leq) &= \uparrow(a, \leq^*) \\ &\subset \uparrow(b, \leq^*) \\ &= \downarrow(b, \leq).\end{aligned}$$

3. (3) $\implies$ (1):

Suppose that $\downarrow a \subset \downarrow b$. Exercise 8.1.0.11 then implies that

$$\begin{aligned}a &\in \downarrow a \\ &\subset \downarrow b \\ &= \{x \in X : x \leq b\}.\end{aligned}$$

Hence $a \leq b$.

□

Exercise 8.1.0.16. Let $(X, \leq)$ be a proset and $S \subset X$. Then

1. $\uparrow S = \bigcup_{s \in S} \uparrow s$
2. $\downarrow S = \bigcup_{s \in S} \downarrow s$

Proof.

1. Define $U \subset X$ by $U := \bigcup_{s \in S} \uparrow s$.

- Since for each $s \in S$, $\uparrow s$ is an upper set, Exercise 8.1.0.6 implies that U is an upper set. Let $s \in S$. Then

$$\begin{aligned} s &\in \uparrow s \\ &\subset \bigcup_{s \in S} \uparrow s \\ &= U. \end{aligned}$$

Since $s \in S$ is arbitrary, we have that for each $s \in S$, $s \in U$. Hence $S \subset U$. Therefore $U \in \mathcal{U}(S)$ and

$$\begin{aligned} \uparrow S &= \bigcap_{U' \in \mathcal{U}(S)} U' \\ &\subset U. \end{aligned}$$

- Let $x \in U$ and $U' \in \mathcal{U}(S)$. Since $x \in U$, there exists $s \in S$ such that $x \in \uparrow s$. Exercise 8.1.0.11 then implies that $s \leq x$. Since $U' \in \mathcal{U}(S)$, U' is an upper set and $S \subset U'$. Then

$$\begin{aligned} s &\in S \\ &\subset U'. \end{aligned}$$

Since U' is an upper set and $s \leq x$, $x \in U'$. Since $U' \in \mathcal{U}(S)$ is arbitrary, we have that for each $U' \in \mathcal{U}(S)$, $x \in U'$. Thus

$$\begin{aligned} x &\in \bigcap_{U' \in \mathcal{U}(S)} U' \\ &= \uparrow S. \end{aligned}$$

Since $x \in U$ is arbitrary, we have that for each $x \in U$, $x \in \uparrow S$. Hence $U \subset \uparrow S$.

Since $\uparrow S \subset U$ and $U \subset \uparrow S$, we have that $\uparrow S = U$.

2. Similar to (1).

□

Exercise 8.1.0.17. Let $(X, \leq)$ be a proset and $(E_\alpha)_{\alpha \in A} \subset \mathcal{P}(X)$. Then

1. $\uparrow \bigcup_{\alpha \in A} E_\alpha = \bigcup_{\alpha \in A} \uparrow E_\alpha$
2. $\downarrow \bigcup_{\alpha \in A} E_\alpha = \bigcup_{\alpha \in A} \downarrow E_\alpha$

Proof.

1. • Let $x \in \uparrow \bigcup_{\alpha \in A} E_\alpha$. Exercise 8.1.0.16 implies that there exists $y \in \bigcup_{\alpha \in A} E_\alpha$ such that $x \in \uparrow y$. Then there exists $\alpha_0 \in A$ such that $y \in E_{\alpha_0}$. Since $\{y\} \subset E_{\alpha_0}$, Exercise 8.1.0.14 implies that $\uparrow y \subset \uparrow E_{\alpha_0}$. Therefore

$$\begin{aligned} x &\in \uparrow y \\ &\subset \uparrow E_{\alpha_0} \\ &\subset \bigcup_{\alpha \in A} \uparrow E_\alpha. \end{aligned}$$

Since $x \in \uparrow \bigcup_{\alpha \in A} E_\alpha$ is arbitrary, we have that for each $x \in \uparrow \bigcup_{\alpha \in A} E_\alpha$, $x \in \bigcup_{\alpha \in A} \uparrow E_\alpha$. Hence $\uparrow \bigcup_{\alpha \in A} E_\alpha \subset \bigcup_{\alpha \in A} \uparrow E_\alpha$.

- Let $x \in \bigcup_{\alpha \in A} \uparrow E_\alpha$. Then there exists $\alpha_0 \in A$ such that $x \in \uparrow E_{\alpha_0}$. Exercise 8.1.0.16 implies that there exists $y \in E_{\alpha_0}$ such that $x \in \uparrow y$. Since

$$\begin{aligned} \{y\} &\subset E_{\alpha_0} \\ &\subset \bigcup_{\alpha \in A} E_\alpha, \end{aligned}$$

Exercise 8.1.0.14 implies that

$$\begin{aligned} x &\in \uparrow y \\ &\subset \uparrow \bigcup_{\alpha \in A} E_\alpha. \end{aligned}$$

Since $x \in \bigcup_{\alpha \in A} \uparrow E_\alpha$ is arbitrary, we have that for each $x \in \bigcup_{\alpha \in A} \uparrow E_\alpha$, $x \in \uparrow \bigcup_{\alpha \in A} E_\alpha$. Hence $\bigcup_{\alpha \in A} \uparrow E_\alpha \subset \uparrow \bigcup_{\alpha \in A} E_\alpha$.

Since $\uparrow \bigcup_{\alpha \in A} E_\alpha \subset \bigcup_{\alpha \in A} \uparrow E_\alpha$ and $\bigcup_{\alpha \in A} \uparrow E_\alpha \subset \uparrow \bigcup_{\alpha \in A} E_\alpha$, we have that $\uparrow \bigcup_{\alpha \in A} E_\alpha = \bigcup_{\alpha \in A} \uparrow E_\alpha$.

2. Similar to (1).

□

Exercise 8.1.0.18. Let $(A, \leq)$ be a proset and $B \subset A$.

1. If B is upward directed, then $\downarrow B$ is upward directed
2. If B is downward directed, then $\uparrow B$ is downward directed

Proof.

1. Suppose that B is upward directed.

- Since B is upward directed, $(B, \leq|_B)$ is a directed set. Thus $B \neq \emptyset$. Thus there exists $x \in B$. Since $B \subset \downarrow B$, we have that $x \in \downarrow B$. Hence $\downarrow B \neq \emptyset$.
- Let $a, b \in \downarrow B$. Exercise 8.1.0.16 implies that there exists $x_a, x_b \in B$ such that $a \in \downarrow x_a$ and $b \in \downarrow x_b$. Since B is upward directed, $(B, \leq|_B)$ is a directed set. Thus $\text{ub}(\{x_a, x_b\}, \leq|_B) \neq \emptyset$. Therefore there exists $x_0 \in B$ such that $x_0 \geq x_a$ and $x_0 \geq x_b$. Hence

$$\begin{aligned} x_0 &\geq x_a \\ &\geq a. \end{aligned}$$

Similarly, $x_0 \geq b$. Since $B \subset \downarrow B$, $x_0 \in \downarrow B$. Thus $\text{ub}(\{a, b\}, \leq|_{\downarrow B}) \neq \emptyset$. Since $a, b \in \downarrow B$ are arbitrary, we have that for each $a, b \in \downarrow B$, $\text{ub}(\{a, b\}, \leq|_{\downarrow B}) \neq \emptyset$. Then $\leq|_{\downarrow B}$ is a direction on $\downarrow B$.

Hence B is upward directed.

2. Suppose that B is $\leq$ -downward directed. **The prev ex** implies that B is $\leq^*$ -upward directed. **The prev part** implies that $\downarrow(B, \leq^*)$ is $\leq^*$ -upward directed. **A prev ex** implies that $\downarrow(B, \leq^*) = \uparrow(B, \leq)$. Since $\uparrow(B, \leq)$ is $\leq^*$ -upward directed, **the prev ex** implies that $\uparrow(B, \leq)$ is $\leq$ -downward directed.

□

Exercise 8.1.0.19. Let $(X, \leq_X)$ $(Y, \leq_Y)$ be prosets and $f : X \rightarrow Y$. Then for each $U \subset Y$, U is a $\leq_Y$ -upper set implies that $f^{-1}(U)$ is a $\leq_X$ -upper set iff for each $L \subset Y$, L is a $\leq_Y$ -lower set implies that $f^{-1}(L)$ is a $\leq_X$ -lower set.

Proof.

- ($\Rightarrow$):

Suppose that for each $U \subset Y$, U is a $\leq_Y$ -upper set implies that $f^{-1}(U)$ is a $\leq_X$ -upper set. Let $L \subset Y$. Suppose that L is a $\leq_Y$ -lower set. Exercise 8.1.0.5 then implies that L^c is a $\leq_Y$ -upper set. By assumption, $f^{-1}(L^c)$ is a $\leq_X$ -upper set. Another application of Exercise 8.1.0.5 implies that $f^{-1}(L^c)^c$ is a $\leq_X$ -lower set. Since

$$\begin{aligned} f^{-1}(L) &= [f^{-1}(L^c)^c]^c \\ &= f^{-1}(L^c)^c, \end{aligned}$$

we have that $f^{-1}(L)$ is a $\leq_X$ -lower set. Since $L \subset Y$ such that L is a $\leq_Y$ -lower set is arbitrary, we have that for each $L \subset Y$, L is a $\leq_Y$ -lower set implies that $f^{-1}(L)$ is a $\leq_X$ -lower set.

- ($\Leftarrow$):

Similar to ($\Rightarrow$).

□

Exercise 8.1.0.20. Let $(X, \leq_X)$ $(Y, \leq_Y)$ be posets and $f : X \rightarrow Y$. If f is $(\leq_X, \leq_Y)$ -monotone, then for each $U \subset Y$, U is a $\leq_Y$ -upper set implies that $f^{-1}(U)$ is a $\leq_X$ -upper set.

Proof. Suppose that f is $(\leq_X, \leq_Y)$ -monotone. Let $U \subset Y$. Suppose that U is a $\leq_Y$ -upper set. Let $a \in f^{-1}(U)$ and $x \in X$. Suppose that $a \leq x$. Since $a \in f^{-1}(U)$, $f(a) \in U$. Since f is monotone, $f(a) \leq f(x)$. Since U is a $\leq_Y$ -upper set and $f(a) \in U$, we have that $f(x) \in U$. Hence $x \in f^{-1}(U)$. Since $a \in f^{-1}(U)$ and $x \in X$ with $a \leq x$ are arbitrary, we have that for each $a \in f^{-1}(U)$ and $x \in X$, $a \leq x$ implies that $x \in f^{-1}(U)$. Thus $f^{-1}(U)$ is a $\leq_X$ -upper set. Since $U \subset Y$ such that U is a $\leq_Y$ -upper set is arbitrary, we have that for each $U \subset Y$, U is a $\leq_Y$ -upper set implies that $f^{-1}(U)$ is a $\leq_X$ -upper set. □

Exercise 8.1.0.21. Let $(X, \leq_X)$ and $(Y, \leq_Y)$ be posets and $f : X \rightarrow Y$. Then f has a right adjoint iff for each $y \in Y$, there exists $x \in X$ such that $f^{-1}(\downarrow y) = \downarrow x$.

Proof.

- ($\Rightarrow$):

Suppose that f has a right adjoint. Then there exists $g : Y \rightarrow X$ such that (f, g) is a monotone Galois connection. [a prev ex in galois connections on posets section](#) Exercise ?? then implies that f and g are monotone, $\text{id}_X \leq_X g \circ f$ and $f \circ g \leq_Y \text{id}_Y$. Let $y \in Y$. Define $x \in X$ by $x := g(y)$.

– We note that for each $a \in X$,

$$\begin{aligned} a \in f^{-1}(\downarrow y) &\iff f(a) \in \downarrow y \\ &\iff f(a) \leq_Y y. \end{aligned}$$

Let $a \in f^{-1}(\downarrow y)$. Then

$$\begin{aligned} a &\leq_X g \circ f(a) \\ &= g(f(a)) \\ &\leq_X g(y) \\ &= x. \end{aligned}$$

Therefore $a \in \downarrow x$. Since $a \in f^{-1}(\downarrow y)$ is arbitrary, we have that for each $a \in f^{-1}(\downarrow y)$, $a \in \downarrow x$. Hence $a \in f^{-1}(\downarrow y) \subset \downarrow x$. Then $f(a) \in \downarrow$.

– Let $a \in \downarrow x$. Then $a \leq_X x$. Hence

$$\begin{aligned} f(a) &\leq_Y f(x) \\ &= f(g(y)) \\ &= f \circ g(y) \\ &\leq_Y y. \end{aligned}$$

Thus $f(a) \in \downarrow y$ and $a \in f^{-1}(\downarrow y)$. Since $a \in \downarrow x$ is arbitrary, we have that for each $a \in \downarrow x$, $a \in f^{-1}(\downarrow y)$. Hence $\downarrow x \subset f^{-1}(\downarrow y)$.

Since $f^{-1}(\downarrow y) \subset \downarrow x$ and $\downarrow x \subset f^{-1}(\downarrow y)$, we have that $f^{-1}(\downarrow y) = \downarrow x$. Since $y \in Y$ is arbitrary, we have that for each $y \in Y$, there exists $x \in X$ such that $f^{-1}(\downarrow y) = \downarrow x$.

• ($\Leftarrow$):

Suppose that for each $y \in Y$, there exists $x \in X$ such that $f^{-1}(\downarrow y) = \downarrow x$. The axiom of choice implies that there exists $g : Y \rightarrow X$ such that for each $y \in Y$, $f^{-1}(\downarrow y) = \downarrow g(y)$. Then for each $x \in X$ and $y \in Y$,

$$\begin{aligned} f(x) \leq_X y &\iff f(x) \in \downarrow y \\ &\iff x \in f^{-1}(\downarrow y) \\ &\iff x \in \downarrow g(y) \\ &\iff x \leq_Y g(y). \end{aligned}$$

Thus (f, g) is a monotone Galois connection. Hence f has a right adjoint. □

Definition 8.1.0.22. Let $(X, \leq_X)$ be a poset.

- Let $U \in \mathcal{U}(\emptyset, \leq_X)$. We define the $(\leq_X, \leq_2)$ -monotonic function induced by U , denoted $m_U : X \rightarrow \mathbf{2}$, by

$$m_U(x) := \begin{cases} 1, & x \in U \\ 0, & x \in U^c. \end{cases}$$

- Let $f : X \rightarrow \mathbf{2}$. Suppose that f is $(\leq_X, \leq_2)$ -monotonic. We define the $\leq_X$ -upper set induced by f , denoted $U_f \subset \mathcal{P}(X)$ by $U_f := f^{-1}(\{1\})$.

Exercise 8.1.0.23. Let $(X, \leq_X)$ be a poset.

1. Let $U \in \mathcal{U}(\leq_X)$. Then $m_U \in \text{Mtn}((X, \leq_X), (\mathbf{2}, \leq_2))$.
2. Let $f : X \rightarrow \mathbf{2}$. Suppose that f is $(\leq_X, \leq_2)$ -monotonic. Then $U_f \in \mathcal{U}(\leq_X)$.

Proof.

1. Let $x, y \in X$. Suppose that $x \leq_X y$.
 - Suppose that $x \in U$. Since $U \in \mathcal{U}(\leq_X)$ and $x \leq_X y$, we have that $y \in U$. Thus

$$\begin{aligned} m_U(x) &= 1 \\ &= m_U(y). \end{aligned}$$

Hence $m_U(x) \leq_2 m_U(y)$.

- Suppose that $x \notin U$. Then

$$\begin{aligned} m_U(x) &= 0 \\ &\leq m_U(y). \end{aligned}$$

Since $x, y \in X$ with $x \leq_X y$ are arbitrary, we have that for each $x, y \in X$, $x \leq_X y$ implies that $m_U(x) \leq_2 m_U(y)$. Thus m_U is monotone.

2. Let $x \in U_f$ and $y \in X$. Suppose that $x \leq_X y$. Since f is $(\leq_X, \leq_2)$ -monotonic, we have that

$$\begin{aligned} 1 &= f(x) \\ &\leq_2 f(y) \\ &\leq_2 1. \end{aligned}$$

Thus $f(y) = 1$ and

$$\begin{aligned} y &\in f^{-1}(\{1\}) \\ &= U_f. \end{aligned}$$

Since $x \in U_f$ and $y \in X$ with $x \leq_X y$ are arbitrary, we have that for each $x \in U_f$ and $y \in X$, $x \leq_X y$ implies that $y \in U_f$. Therefore $U_f \in \mathcal{U}(\leq_X)$.

□

Definition 8.1.0.24. Let X be a poset. We define the **monotone-to-upper-set map**, denoted $\Phi : \text{Mtn}(X, \mathbf{2}) \rightarrow \mathcal{U}_X$, by $\Phi(f) := U_f$.

Exercise 8.1.0.25. Then

1. $\Phi \in \text{Mtn}(\text{Mtn}(X, \mathbf{2}), \mathcal{U}_X)$.
2. Φ is an isomorphism.

Proof.

1. Let $f, g \in \text{Mtn}(X, \mathbf{2})$. Suppose that $f \leq g$. Let $x \in U_f$. Then $f(x) = 1$. Since $f \leq g$, we have that

$$\begin{aligned} 1 &= f(x) \\ &\leq g(x) \\ &\leq 1. \end{aligned}$$

Hence $g(x) = 1$ and $x \in U_g$. Since $x \in U_f$ is arbitrary, we have that for each $x \in U_f$, $x \in U_g$. Thus

$$\begin{aligned} \Phi(f) &= U_f \\ &\subset U_g \\ &= \Phi(g). \end{aligned}$$

Since $f, g \in \text{Mtn}(X, \mathbf{2})$ with $f \leq g$ are arbitrary, we have that for each $f, g \in \text{Mtn}(X, \mathbf{2})$, $f \leq g$ implies that $\Phi(f) \subset \Phi(g)$. Hence $\Phi \in \text{Mtn}(\text{Mtn}(X, \mathbf{2}), \mathcal{U}_X)$.

2. • – **injectivity:**

Let $f, g \in \text{Mtn}(X, \mathbf{2})$. Suppose that $\Phi(f) = \Phi(g)$. Then $U_f = U_g$ and therefore for each $x \in X$,

- * $x \in U_f$ implies that $x \in U_g$ and $f(x) = 1 = g(x)$.
- * $x \in U_f^c$ implies that $x \in U_g^c$ and $f(x) = 0 = g(x)$.

Thus for each $x \in X$, $f(x) = g(x)$. Hence $f = g$. Since $f, g \in \text{Mtn}(X, \mathbf{2})$ with $\Phi(f) = \Phi(g)$ are arbitrary, we have that for each $f, g \in \text{Mtn}(X, \mathbf{2})$, $\Phi(f) = \Phi(g)$ implies that $f = g$. Thus Φ is injective.

- **surjectivity:**

Let $U \in \mathcal{U}_X$. Define $f \in \text{Mtn}(X, \mathbf{2})$ by $f := m_U$. Then

$$\begin{aligned} \Phi(f) &= U_f \\ &= f^{-1}(\{1\}) \\ &= m_U^{-1}(\{1\}) \\ &= U \quad (\text{by Definition 8.1.0.22}). \end{aligned}$$

Since $U \in \mathcal{U}_X$ is arbitrary, we have that for each $U \in \mathcal{U}_X$, there exists $f \in \text{Mtn}(X, \mathbf{2})$ such that $\Phi(f) = U$. Thus Φ is surjective.

Therefore Φ is a bijection.

- Clearly for each $U \in \mathcal{U}_X$, $\Phi^{-1}(U) = m_U$. Let $U_1, U_2 \in \mathcal{U}_X$. Suppose that $U_1 \subset U_2$. Then for each $x \in X$,
 - $x \in U_2$ implies that

$$\begin{aligned} \Phi^{-1}(U_1)(x) &= m_{U_1}(x) \\ &\leq 1 \\ &= m_{U_2}(x) \\ &= \Phi^{-1}(U_2)(x). \end{aligned}$$

– $x \in U_2^c$ implies that $x \in U_1^c$ and

$$\begin{aligned}\Phi^{-1}(U_1)(x) &= m_{U_1}(x) \\ &= 0 \\ &= m_{U_2}(x) \\ &= \Phi^{-1}(U_2)(x).\end{aligned}$$

Thus $\Phi^{-1}(U_1) \leq \Phi^{-1}(U_2)$. Since $U_1, U_2 \in \mathcal{U}_X$ with $U_1 \subset U_2$ are arbitrary, we have that for each $U_1, U_2 \in \mathcal{U}_X$, $U_1 \subset U_2$ implies that $\Phi^{-1}(U_1) \leq \Phi^{-1}(U_2)$. Therefore $\Phi^{-1} \in \text{Mtn}(\mathcal{U}_X, \text{Mtn}(X, \mathbf{2}))$.

Thus Φ is an isomorphism.

□

need to do similar for lower sets $\mathcal{L}_X$ and antitone? morphisms

8.2 Maximal Elements

RELABEL, MAYBE RETITLE

Definition 8.2.0.1. Let $(X, \leq)$ be a proset and $a \in X$. Then a is said to be

- **$\leq$ -maximal** if for each $x \in X$, $a \leq x$ implies that $x \sim_{\leq} a$.
- **$\leq$ -minimal** if for each $x \in X$, $x \leq a$ implies that $x \sim_{\leq} a$.

Note 8.2.0.2. When the context is clear, we write “maximal” and “minimal” instead of “ $\leq$ -maximal” and “ $\leq$ -minimal” respectively.

Exercise 8.2.0.3. Let $(X, \leq)$ be a proset and $a \in X$. Then

1. a is maximal iff $\uparrow a = \pi_{X/\sim_{\leq}}(a)$
2. a is minimal iff $\downarrow a = \pi_{X/\sim_{\leq}}(a)$

Proof.

1. • ($\implies$):

Suppose that a is maximal.

- Let $x \in \uparrow a$. Exercise 8.1.0.11 implies $a \leq x$. Since a is maximal, $x \sim_{\leq} a$. Thus $x \in \pi_{X/\sim_{\leq}}(a)$. Since $x \in \uparrow a$ is arbitrary, we have that for each $x \in \uparrow a$, $x \in \pi_{X/\sim_{\leq}}(a)$. Hence $\uparrow a \subset \pi_{X/\sim_{\leq}}(a)$.
- Let $x \in \pi_{X/\sim_{\leq}}(a)$. Then $a \sim_{\leq} x$. Hence $a \leq x$ and $x \leq a$. Since $a \leq x$, we have that $x \in \uparrow a$. Since $x \in \pi_{X/\sim_{\leq}}(a)$ is arbitrary, we have that for each $x \in \pi_{X/\sim_{\leq}}(a)$, $x \in \uparrow a$. Hence $\pi_{X/\sim_{\leq}}(a) \subset \uparrow a$.

Since $\uparrow a \subset \pi_{X/\sim_{\leq}}(a)$ and $\pi_{X/\sim_{\leq}}(a) \subset \uparrow a$, we have that $\uparrow a = \pi_{X/\sim_{\leq}}(a)$.

- ($\impliedby$):

Suppose that $\uparrow a = \pi_{X/\sim_{\leq}}(a)$. Let $x \in X$. Suppose that $a \leq x$. Then

$$\begin{aligned}x &\in \uparrow a \\ &= \pi_{X/\sim_{\leq}}(a).\end{aligned}$$

Thus $x \sim_{\leq} a$. Since $x \in X$ with $a \leq x$ is arbitrary, we have that for each $x \in X$, $a \leq x$ implies that $x \sim_{\leq} a$. Hence a is maximal.

2. Similar to (1).

□

8.3 Filters and Ideals

Definition 8.3.0.1. Let $(X, \leq)$ be a poset and $F, I \subset X$. Then

- F is said to be a $\leq$ -**filter** if
 1. $F \in \mathcal{U}(\leq)$
 2. F is downward directed.
- I is said to be a $\leq$ -**ideal** if
 1. $I \in \mathcal{L}(\leq)$
 2. I is upward directed.

Definition 8.3.0.2.

Chapter 9

Suprema and Infima

9.1 Upper and Lower Bounds

Definition 9.1.0.1. maybe move to suprema infima section, or make a section on upper and lower bounds Let $(X, \leq)$ be a proset and $A \subset X$.

- Let $x \in X$. Then x is said to be a
 - **$\leq$ -upper bound of A** if for each $a \in A$, $a \leq x$
 - **$\leq$ -lower bound of A** if for each $a \in A$, $x \leq a$
- We define
 - $\text{ub}(A, \leq) := \{x \in X : x \text{ is a } \leq\text{-upper bound of } A\}$
 - $\text{lb}(A, \leq) := \{x \in X : x \text{ is a } \leq\text{-lower bound of } A\}$
- Then A is said to be
 - **$\leq$ -bounded above** if $\text{ub}(A, \leq) \neq \emptyset$
 - **$\leq$ -bounded below** if $\text{lb}(A, \leq) \neq \emptyset$

Note 9.1.0.2. When the context is clear, we write

- “upper bound” and “lower bound” instead of “ $\leq$ -upper bound” and “ $\leq$ -lower bound” respectively
- $\text{ub } A$ and $\text{lb } A$ in place of $\text{ub}(A, \leq)$ and $\text{lb}(A, \leq)$ respectively
- “bounded above” and “bounded below” instead of “ $\leq$ -bounded above” and “ $\leq$ -bounded below” respectively

Exercise 9.1.0.3. Let $(X, \leq)$ be a proset and $A \subset X$. Then

1. $\text{ub}(A, \leq) = \text{lb}(A, \leq^*)$
2. $\text{ub}(A, \leq^*) = \text{lb}(A, \leq)$

Proof.

1. We note that for each $x \in X$,

$$\begin{aligned} x \in \text{ub}(A, \leq) &\iff \text{for each } a \in A, a \leq x \\ &\iff \text{for each } a \in A, x \leq^* a \\ &\iff x \in \text{lb}(A, \leq^*). \end{aligned}$$

Hence $\text{ub}(A, \leq) = \text{lb}(A, \leq^*)$.

2. Similar to (1) **ref**.

□

Exercise 9.1.0.4. Let $(X, \leq)$ be a poset and $A \subset X$. Then

1. $\text{ub } A = \bigcap_{x \in A} \uparrow x$
2. $\text{lb } A = \bigcap_{x \in A} \downarrow x$

Proof.

1.
 - Let $a \in \text{ub } A$ and $x \in A$. Then $x \leq a$. Hence $a \in \uparrow x$. Since $x \in A$ is arbitrary, we have that for each $x \in A$, $a \in \uparrow x$. Thus $a \in \bigcap_{x \in A} \uparrow x$. Since $a \in \text{ub } A$ is arbitrary, we have that for each $a \in \text{ub } A$, $a \in \bigcap_{x \in A} \uparrow x$. Hence $\text{ub } A \subset \bigcap_{x \in A} \uparrow x$.
 - Let $a \in \bigcap_{x \in A} \uparrow x$ and $x_0 \in A$. Then $a \in \uparrow x_0$. Hence $x_0 \leq a$. Since $x_0 \in A$ is arbitrary, we have that for each $x_0 \in A$, $x_0 \leq a$. Hence $a \in \text{ub } A$. Since $a \in \bigcap_{x \in A} \uparrow x$ is arbitrary, we have that for each $a \in \bigcap_{x \in A} \uparrow x$, $a \in \text{ub } A$. Thus $\bigcap_{x \in A} \uparrow x \subset \text{ub } A$.

Since $\text{ub } A \subset \bigcap_{x \in A} \uparrow x$ and $\bigcap_{x \in A} \uparrow x \subset \text{ub } A$, we have that $\text{ub } A = \bigcap_{x \in A} \uparrow x$.

2. Similar to (1).

□

Exercise 9.1.0.5. Let $(X, \leq)$ be a poset and $A, B \subset X$. If $A \subset B$, then

1. $\text{ub } B \subset \text{ub } A$
2. $\text{lb } B \subset \text{lb } A$

Proof. Suppose that $A \subset B$.

1. Exercise 9.1.0.4 implies that

$$\begin{aligned} \text{ub } B &= \bigcap_{x \in B} \uparrow x \\ &\subset \bigcap_{x \in A} \uparrow x \\ &= \text{ub } A. \end{aligned}$$

2. Similar to (1). **ref**

□

Exercise 9.1.0.6. Let $(X, \leq)$ be a poset and $(E_\alpha)_{\alpha \in A} \subset \mathcal{P}(X)$. Then $\text{ub } \bigcup_{\alpha \in A} E_\alpha = \bigcap_{\alpha \in A} \text{ub } E_\alpha$.

Proof. Define $E_\vee \subset X$ by $E_\vee := \bigcup_{\alpha \in A} E_\alpha$. Exercise 9.1.0.4 then implies that

$$\begin{aligned} \text{ub } \bigcup_{\alpha \in A} E_\alpha &= \bigcap_{x \in E_\vee} \uparrow x \\ &= \bigcap_{\alpha \in A} \left[\bigcap_{x \in E_\alpha} \uparrow x \right] \\ &= \bigcap_{\alpha \in A} \text{ub } E_\alpha. \end{aligned}$$

□

Exercise 9.1.0.7. Let $(X, \leq)$ be a poset. For each $A \subset X$,

1. $A \subset \text{lb}(\text{ub } A)$
2. $A \subset \text{ub}(\text{lb } A)$

Proof.

1. Let $a \in A$. Then for each $x \in \text{ub } A$, $a \leq x$. Hence $a \in \text{lb}(\text{ub } A)$. Since $a \in A$ is arbitrary, we have that for each $a \in A$, $a \in \text{lb}(\text{ub } A)$. Thus $A \subset \text{lb}(\text{ub } A)$.
2. Let $a \in A$. Then for each $x \in \text{lb } A$, $x \leq a$. Hence $a \in \text{ub}(\text{lb } A)$. Since $a \in A$ is arbitrary, we have that for each $a \in A$, $a \in \text{ub}(\text{lb } A)$. Thus $A \subset \text{ub}(\text{lb } A)$.

□

Exercise 9.1.0.8. Let $(X, \leq)$ be a poset and $A, B \subset X$. Suppose that $B \subset A$. Then

1. $\text{ub}(B, \leq|_A) = \text{ub}(B, \leq) \cap A$.
2. $\text{lb}(B, \leq|_A) = \text{lb}(B, \leq) \cap A$.

Proof.

1. We note that for each $x \in X$,

$$\begin{aligned} x \in \text{ub}(B, \leq|_A) &\iff x \in A \text{ and for each } y \in B, y \leq|_A x \\ &\iff x \in A \text{ and for each } y \in B, y \leq x \\ &\iff x \in \text{ub}(B, \leq) \cap A. \end{aligned}$$

2. We have that

$$\begin{aligned} \text{lb}(B, \leq|_A) &= \text{ub}(B, \leq|_A^*) \quad (\text{by Exercise 9.1.0.3}) \\ &= \text{ub}(B, \leq^*|_A) \quad (\text{by ref ex in set theory notes in binary hetero relations}) \\ &= \text{ub}(B, \leq^*) \cap A \quad (\text{by ref prev part}) \\ &= \text{lb}(B, \leq) \cap A \quad (\text{by Exercise 9.1.0.3}). \end{aligned}$$

□

Exercise 9.1.0.9. Let $(X, \leq_X)$ and $(Y, \leq_Y)$ be posets, $f : X \rightarrow Y$ and $g : Y \rightarrow X$.

1. If (f, g) is a $(\leq_X, \leq_Y)$ -monotone Galois connection, then
 - (a) for each $A \subset X$, $\text{ub}(f(A), \leq_Y) = g^{-1}(\text{ub}(A, \leq_X))$,
 - (b) for each $B \subset Y$, $\text{lb}(g(B), \leq_X) = f^{-1}(\text{lb}(B, \leq_Y))$.
2. If (f, g) is a $(\leq_X, \leq_Y)$ -antitone Galois connection, then
 - (a) for each $A \subset X$, $\text{ub}(f(A), \leq_Y) = g^{-1}(\text{lb}(A, \leq_X))$,
 - (b) for each $B \subset Y$, $\text{ub}(g(B), \leq_X) = f^{-1}(\text{lb}(B, \leq_Y))$.

Proof. Since (f, g) is a $(\leq_X, \leq_Y)$ -monotone Galois connection, a prev ex implies that f is $(\leq_X, \leq_Y)$ -monotone, g is $(\leq_Y, \leq_X)$ -monotone, $\text{id}_X \leq_X g \circ f$ and $f \circ g \leq_Y \text{id}_Y$.

1. Suppose that (f, g) is a $(\leq_X, \leq_Y)$ -monotone Galois connection.
 - (a) Let $A \subset X$.

- Let $y \in \text{ub}(f(A), \leq_Y)$ and $a \in A$. Then $f(a) \leq_Y y$. Hence

$$\begin{aligned} a &\leq_X g \circ f(a) \\ &= g(f(a)) \\ &\leq_X g(y). \end{aligned}$$

Since $a \in A$ is arbitrary, we have that for each $a \in A$, $a \leq_X g(y)$. Thus $g(y) \in \text{ub}(A, \leq_X)$. Hence $y \in g^{-1}(\text{ub}(A, \leq_X))$. Since $y \in \text{ub}(f(A), \leq_Y)$ is arbitrary, we have that for each $y \in \text{ub}(f(A), \leq_Y)$, $y \in g^{-1}(\text{ub}(A, \leq_X))$. Thus $\text{ub}(f(A), \leq_Y) \subset g^{-1}(\text{ub}(A, \leq_X))$.

- Let $y \in g^{-1}(\text{ub}(A, \leq_X))$. Then $g(y) \in \text{ub}(A, \leq_X)$. Let $b \in f(A)$. Then there exists $a \in A$ such that $b = f(a)$. Then $a \leq_X g(y)$. Therefore

$$\begin{aligned} b &= f(a) \\ &\leq_Y f(g(y)) \\ &= f \circ g(y) \\ &= \leq_Y y. \end{aligned}$$

Since $b \in f(A)$ is arbitrary, we have that for each $b \in f(A)$, $b \leq_Y y$. Thus $y \in \text{ub}(f(A), \leq_Y)$. Since $y \in g^{-1}(\text{ub}(A, \leq_X))$ is arbitrary, we have that for each $y \in g^{-1}(\text{ub}(A, \leq_X))$, $y \in \text{ub}(f(A), \leq_Y)$. Hence $g^{-1}(\text{ub}(A, \leq_X)) \subset \text{ub}(f(A), \leq_Y)$.

Since $\text{ub}(f(A), \leq_Y) \subset g^{-1}(\text{ub}(A, \leq_X))$ and $g^{-1}(\text{ub}(A, \leq_X)) \subset \text{ub}(f(A), \leq_Y)$, we have that

$$\text{ub}(f(A), \leq_Y) = g^{-1}(\text{ub}(A, \leq_X)).$$

- (b) Since (f, g) is a $(\leq_X, \leq_Y)$ -monotone Galois connection, [a prev rex](#) implies that (g, f) is a $(\leq_Y^*, \leq_X^*)$ -monotone Galois connection. [The previous part \(ref\)](#) and Exercise 9.1.0.3 then imply that for each $B \subset Y$,

$$\begin{aligned} \text{lb}(g(B), \leq_X) &= \text{ub}(g(B), \leq_X^*) \\ &= f^{-1}(\text{ub}(B, \leq_Y^*)) \\ &= f^{-1}(\text{lb}(B, \leq_Y)). \end{aligned}$$

2. Suppose that (f, g) is a $(\leq_X, \leq_Y)$ -antitone Galois connection. [A prev ex](#) then implies that (f, g) is a $(\leq_X^*, \leq_Y)$ -monotone Galois connection and (g, f) is a $(\leq_Y^*, \leq_X)$ -monotone Galois connection. [The prev part](#) implies that

- (a) for each $A \subset X$,

$$\begin{aligned} \text{ub}(f(A), \leq_Y) &= g^{-1}(\text{ub}(A, \leq_X^*)) \\ &= g^{-1}(\text{lb}(A, \leq_X)). \end{aligned}$$

- (b) for each $B \subset Y$,

$$\begin{aligned} \text{ub}(g(B), \leq_X) &= f^{-1}(\text{ub}(B, \leq_Y^*)) \\ &= f^{-1}(\text{lb}(B, \leq_Y)). \end{aligned}$$

□

Definition 9.1.0.10. Let $(X, \leq)$ be a proset. We define the

- **$\leq$ -upper bound map**, denoted $\text{ub}(\cdot, \leq) : \mathcal{P}(X) \rightarrow \mathcal{P}(X)$, by $\text{ub}(\cdot, \leq)(A) := \text{ub}(A, \leq)$.
- **$\leq$ -lower bound map**, denoted $\text{lb}(\cdot, \leq) : \mathcal{P}(X) \rightarrow \mathcal{P}(X)$, by $\text{lb}(\cdot, \leq)(A) := \text{lb}(A, \leq)$.

Note 9.1.0.11. When the context is clear, we may write $\text{ub} : \mathcal{P}(X) \rightarrow \mathcal{P}(X)$ and $\text{lb} : \mathcal{P}(X) \rightarrow \mathcal{P}(X)$ in place of $\text{ub}(\cdot, \leq) : \mathcal{P}(X) \rightarrow \mathcal{P}(X)$ and $\text{lb}(\cdot, \leq) : \mathcal{P}(X) \rightarrow \mathcal{P}(X)$.

Exercise 9.1.0.12. Let $(X, \leq)$ be a proset. Then $(\text{ub}(\cdot, \leq), \text{lb}(\cdot, \leq))$ is a $(\subset^*, \subset^*)$ -antitone Galois connection.

Proof. Define $f, g : \mathcal{P}(X) \rightarrow \mathcal{P}(X)$ by $f := \text{ub}(\cdot, \leq)$ and $g := \text{lb}(\cdot, \leq)$.

- Exercise 9.1.0.7 then implies that for each $A \in \mathcal{P}(X)$, $\text{id}_{\mathcal{P}(X)}(A) \subset g \circ f(A)$ and $\text{id}_{\mathcal{P}(X)}(A) \subset f \circ g(A)$. Therefore for each $A \in \mathcal{P}(X)$, $g \circ f(A) \subset^* \text{id}_{\mathcal{P}(X)}(A)$ and $f \circ g(A) \subset^* \text{id}_{\mathcal{P}(X)}(A)$.
- Exercise 9.1.0.5 implies that f, g are $(\subset, \subset)$ -antitone. Exercise 7.1.0.3 implies that f, g are $(\subset^*, \subset^*)$ -antitone.

Exercise 7.2.0.6 then implies that (f, g) is a $(\subset^*, \subset^*)$ -antitone Galois connection. □

Exercise 9.1.0.13. Let $(X, \leq_X)$ $(Y, \leq_Y)$ be posets, $f : X \rightarrow Y$ and $A \subset X$. If f is $(\leq_X, \leq_Y)$ -monotone, then

1. $f(\text{ub}(A, \leq_X)) \subset \text{ub}(f(A), \leq_Y)$,
2. $f(\text{lb}(A, \leq_X)) \subset \text{lb}(f(A), \leq_Y)$

Proof. Suppose that f is $(\leq_X, \leq_Y)$ -monotone.

1. Let $x \in \text{ub}(A, \leq_X)$ and $y_0 \in f(A)$. Then there exists $x_0 \in A$ such that $f(x_0) = y_0$. Since $x \in \text{ub}(A, \leq_X)$, $x_0 \leq x$. Since f is $(\leq_X, \leq_Y)$ -monotone,

$$\begin{aligned} y_0 &= f(x_0) \\ &\leq f(x). \end{aligned}$$

Since $y_0 \in f(A)$ is arbitrary, we have that for each $y_0 \in f(A)$, $y_0 \leq f(x)$. Therefore $f(x) \in \text{ub}(f(A), \leq_Y)$. Since $x \in \text{ub}(A, \leq_X)$ is arbitrary, we have that for each $x \in \text{ub}(A, \leq_X)$, $f(x) \in \text{ub}(f(A), \leq_Y)$. Hence $f(\text{ub}(A, \leq_X)) \subset \text{ub}(f(A), \leq_Y)$.

2. Exercise 9.1.0.3 and [the prev part](#) imply that

$$\begin{aligned} f(\text{lb}(A, \leq_X)) &= f(\text{ub}(A, \leq_X^*)) \\ &\subset \text{ub}(f(A), \leq_Y^*) \\ &= \text{lb}(f(A), \leq_Y). \end{aligned}$$

□

9.2 SupSets and InfSets

Definition 9.2.0.1. Let $(X, \leq)$ be a proset and $A \subset X$.

- Let $x \in X$. Then x is said to be a
 - **$\leq$ -supremum of A or $\leq$ -least upper bound of A** , if
 1. $x \in \text{ub}(A, \leq)$
 2. for each $y \in \text{ub}(A, \leq)$, $x \leq y$
 - **$\leq$ -infimum of A or $\leq$ -greatest lower bound of A** , if
 1. $x \in \text{lb}(A, \leq)$
 2. for each $y \in \text{lb}(A, \leq)$, $y \leq x$.
- – We define $\text{supSet}(A, \leq) \subset X$ by $\text{supSet}(A, \leq) := \{x \in X : x \text{ is a } \leq\text{-supremum of } A\}$
- We define $\text{infSet}(A, \leq) \subset X$ by $\text{infSet}(A, \leq) := \{x \in X : x \text{ is an } \leq\text{-infimum of } A\}$

Note 9.2.0.2. When the context is clear,

- we say “supremum” and “infimum” in place of “ $\leq$ -supremum” and “ $\leq$ -infimum” respectively,
- we write $\text{supSet } A$ and $\text{infSet } A$ in place of $\text{supSet}(A, \leq)$ and $\text{infSet}(A, \leq)$ respectively.

Exercise 9.2.0.3. Let $(X, \leq)$ be a proset and $A \subset X$. Then

1. (a) $\text{supSet } A = \text{ub } A \cap \text{lb}(\text{ub } A)$
 (b) $\text{supSet } A = \text{infSet}(\text{ub } A)$
2. (a) $\text{infSet } A = \text{lb } A \cap \text{ub}(\text{lb } A)$
 (b) $\text{infSet } A = \text{supSet}(\text{lb } A)$

Proof.

1. (a) We note that for each $x \in X$,

$$\begin{aligned}
 x \in \text{supSet } A &\iff x \in \text{ub } A \text{ and for each } y \in \text{ub } A, x \leq y \\
 &\iff x \in \text{ub } A \text{ and } x \in \text{lb}(\text{ub } A) \\
 &\iff x \in \text{ub } A \cap \text{lb}(\text{ub } A).
 \end{aligned}$$

Therefore $\text{supSet } A = \text{ub } A \cap \text{lb}(\text{ub } A)$.

- (b) Exercise 9.1.0.12 implies that $(\text{ub}(\cdot, \leq), \text{lb}(\cdot, \leq))$ is a $(\subset^*, \subset^*)$ -antitone Galois connection. Since $(\mathcal{P}(X), \subset)$ is a poset Exercise 7.2.0.8 implies that $\text{ub}(\text{lb}(\text{ub } A)) = \text{ub } A$. Therefore

$$\begin{aligned}
 \text{supSet } A &= \text{ub } A \cap \text{lb}(\text{ub } A) \quad (\text{by prev part (ref)}) \\
 &= \text{ub}(\text{lb}(\text{ub } A)) \cap \text{lb}(\text{ub } A) \quad (\text{by prev part (ref)}) \\
 &= \text{lb}(\text{ub } A) \cap \text{ub}(\text{lb}(\text{ub } A)) \\
 &= \text{infSet}(\text{ub } A).
 \end{aligned}$$

2. Similar to (1) **USE DUALITY???**.

□

Exercise 9.2.0.4. Let $(X, \leq)$ be a proset and $A \subset X$. Then

1. $\text{supSet}(A, \leq^*) = \text{infSet}(A, \leq)$
2. $\text{supSet}(A, \leq) = \text{infSet}(A, \leq^*)$

Proof.

1. We have that

$$\begin{aligned}\text{supSet}(A, \leq^*) &= \text{ub}(A, \leq^*) \cap \text{lb}(\text{ub}(A, \leq^*), \leq^*) \quad (\text{by Exercise 9.2.0.3}) \\ &= \text{lb}(A, \leq) \cap \text{ub}(\text{lb}(A, \leq), \leq) \quad (\text{by Exercise 9.1.0.3}) \\ &= \text{infSet}(A, \leq) \quad (\text{by Exercise 9.2.0.3}).\end{aligned}$$

2. We have that

$$\begin{aligned}\text{supSet}(A, \leq) &= \text{supSet}(A, (\leq^*)^*) \\ &= \text{infSet}(A, \leq^*) \quad (\text{by prev part (ref prev part)}).\end{aligned}$$

Exercise 9.2.0.3 □

Exercise 9.2.0.5. Let $(X, \leq)$ be a proset and $A \subset X$. Then

1. for each $x, y \in \text{supSet}(A, \leq)$, $x \sim_{\leq} y$.
2. for each $x, y \in \text{infSet}(A, \leq)$, $x \sim_{\leq} y$.

Proof.

1. Let $x, y \in \text{supSet}(A, \leq)$. Then $x, y \in \text{ub}(A, \leq)$. Since $x \in \text{supSet}(A, \leq)$ and $y \in \text{ub}(A, \leq)$, $x \leq y$. Since $y \in \text{supSet}(A, \leq)$ and $x \in \text{ub}(A, \leq)$, $y \leq x$. Hence $x \sim_{\leq} y$. Since $x, y \in \text{supSet}(A, \leq)$ are arbitrary, we have that for each $x, y \in \text{supSet}(A, \leq)$, $x \sim_{\leq} y$.
2. We note that for each $x, y \in X$,

$$\begin{aligned}x, y \in \text{infSet}(A, \leq) &\iff x, y \in \text{supSet}(A, \leq^*) \quad (\text{by Exercise 9.2.0.4}) \\ &\implies x \sim_{\leq^*} y \quad (\text{by prev part (ref)}) \\ &\iff x \sim_{\leq} y \quad (\text{by Exercise 4.1.0.6}).\end{aligned}$$

□

Exercise 9.2.0.6. Let $(X, \leq)$ be a poset and $A \subset X$. Then

1. for each $x, y \in \text{supSet } A$, $x = y$.
2. for each $x, y \in \text{infSet } A$, $x = y$.

Proof.

1. Let $x, y \in \text{supSet } A$. Exercise 9.2.0.5 implies that $x \sim_{\leq} y$. Exercise 5.1.0.2 then implies that $x = y$.
2. Similar to (1).

□

Exercise 9.2.0.7. Let $(X, \leq)$ be a proset and $A, B \subset X$. If $A \subset B$, then

1. for each $s_A \in \text{supSet } A$ and $s_B \in \text{supSet } B$, $s_A \leq s_B$,
2. for each $l_A \in \text{infSet } A$ and $l_B \in \text{infSet } B$, $l_B \leq l_A$.

Proof. Suppose that $A \subset B$.

1. Let $s_A \in \text{supSet } A$ and $s_B \in \text{supSet } B$. Then Exercise 9.1.0.5 implies that

$$\begin{aligned} s_B &\in \text{supSet } B \\ &\subset \text{ub } B \\ &\subset \text{ub } A. \end{aligned}$$

Since $s_A \in \text{lb}(\text{ub } A)$, we have that $s_A \leq s_B$. Since $s_A \in \text{supSet } A$ and $s_B \in \text{supSet } B$ are arbitrary, we have that for each $s_A \in \text{supSet } A$ and $s_B \in \text{supSet } B$, $s_A \leq s_B$.

2. **similar, use duality**

□

Exercise 9.2.0.8. Let $(X, \leq_X)$ and $(Y, \leq_Y)$ be posets, $f : X \rightarrow Y$, $g : Y \rightarrow X$ and $A \subset X$. If (f, g) is a $(\leq_X, \leq_Y)$ -monotone Galois connection, then

1. $f(\text{supSet}(A, \leq_X)) \subset \text{supSet}(f(A), \leq_Y)$.
2. **can we say something about $\text{infSet}(f(A), \leq_Y)$ or $\text{infSet}(g(B), \leq_X)$?**

Proof. Suppose that (f, g) is a $(\leq_X, \leq_Y)$ -monotone Galois connection.

1. **A prev ex in Galois connection subsection** implies that f is $(\leq_X, \leq_Y)$ -monotone, g is $(\leq_Y, \leq_X)$ -monotone, $g \circ f \leq_X \text{id}_X$ and $\text{id}_Y \leq_Y f \circ g$. Let $s \in \text{supSet}(A, \leq_X)$. Since $\text{supSet}(A, \leq_X) \subset \text{ub}(A, \leq_X)$, we have that $s \in \text{ub}(A, \leq_X)$. Exercise 9.1.0.13 then implies that

$$\begin{aligned} f(s) &\in f(\text{ub}(A, \leq_X)) \\ &\subset \text{ub}(f(A), \leq_Y). \end{aligned}$$

Let $y \in \text{ub}(f(A), \leq_Y)$. **A prev ex in galois connection subsection** imply that

$$\begin{aligned} y &\in \text{ub}(f(A), \leq_Y) \\ &= g^{-1}(\text{ub}(A, \leq_X)). \end{aligned}$$

Then $g(y) \in \text{ub}(A, \leq_X)$. Since $s \in \text{supSet}(A, \leq_X)$, we have that $s \leq_X g(y)$. Hence

$$\begin{aligned} f(s) &\leq_Y f(g(y)) \\ &= f \circ g(y) \\ &\leq_Y y. \end{aligned}$$

Since $y \in \text{ub}(f(A), \leq_Y)$ is arbitrary, we have that for each $y \in \text{ub}(f(A), \leq_Y)$, $f(s) \leq_Y y$. Thus $f(s) \in \text{supSet}(f(A), \leq_Y)$. Since $s \in \text{supSet}(A, \leq_X)$ is arbitrary, we have that for each $s \in \text{supSet}(A, \leq_X)$, $f(s) \in \text{supSet}(f(A), \leq_Y)$. Hence $f(\text{supSet}(A, \leq_X)) \subset \text{supSet}(f(A), \leq_Y)$.

2. **FINISH!!!**

□

9.3 Suprema and Infima

relabel ex/def references

Definition 9.3.0.1. Let $(X, \leq)$ be a poset and $A \subset X$.

- If $\text{supSet } A \neq \emptyset$, we define the **supremum of** A , denoted $\sup A$, to be the unique $x \in X$ such that $x \in \text{supSet } A$.
- If $\text{infSet } A \neq \emptyset$, we define the **infimum of** A , denoted $\inf A$, to be the unique $x \in X$ such that $x \in \text{infSet } A$.
- We say that $\sup A$ (**resp.** $\inf A$) **exists** if $\text{supSet } A \neq \emptyset$ (**resp.** $\text{infSet } A \neq \emptyset$).

Exercise 9.3.0.2. Let $(X, \leq)$ be a poset and $A \subset X$. Then

- (a) $\sup A$ exists iff $\inf(\text{ub } A)$ exists
(b) $\sup A$ exists implies that $\sup A = \inf(\text{ub } A)$
- (a) $\inf A$ exists iff $\sup(\text{lb } A)$ exists
(b) $\inf A$ exists implies that $\inf A = \sup(\text{lb } A)$

Proof.

- (a) Exercise 9.2.0.3 implies that

$$\begin{aligned} \sup A \text{ exists} &\iff \text{supSet } A \neq \emptyset \\ &\iff \text{infSet}(\text{ub } A) \neq \emptyset \\ &\iff \inf(\text{ub } A) \text{ exists.} \end{aligned}$$

- (b) Suppose that $\sup A$ exists. **the previous part (a)** then implies that $\inf(\text{ub } A)$ exists. **a previous ex** implies that $\text{supSet } A = \{\sup A\}$ and $\text{infSet}(\text{ub } A) = \{\inf(\text{ub } A)\}$. **The previous parts (a)** then implies that

$$\begin{aligned} \{\sup A\} &= \text{supSet } A \\ &= \text{infSet}(\text{ub } A) \\ &= \{\inf(\text{ub } A)\}. \end{aligned}$$

Hence $\sup A = \inf(\text{ub } A)$.

- Similar to (1).

□

Exercise 9.3.0.3. Let $(X, \leq)$ be a poset and $A \subset X$. Then

- (a) $\sup(A, \leq)$ exists iff $\inf(A, \leq^*)$ exists
(b) $\sup A$ exists implies that $\sup(A, \leq) = \inf(A, \leq^*)$
- (a) $\inf(A, \leq)$ exists iff $\sup(A, \leq^*)$ exists
(b) $\inf A$ exists implies that $\inf(A, \leq) = \sup(A, \leq^*)$

Proof.

- (a) We have that

$$\begin{aligned} \sup(A, \leq) \text{ exists} &\iff \text{supSet}(A, \leq) \neq \emptyset \\ &\iff \text{infSet}(A, \leq^*) \neq \emptyset \quad (\text{by Exercise 9.2.0.4}) \\ &\iff \inf(A, \leq^*) \text{ exists.} \end{aligned}$$

- (b) Suppose that $\sup(A, \leq)$ exists. **The prev part** implies that $\inf(A, \leq^*)$ exists. Since $\sup\text{Set}(A, \leq) = \{\sup(A, \leq)\}$ and $\inf\text{Set}(A, \leq^*) = \{\inf(A, \leq^*)\}$, Exercise 9.2.0.4 implies that

$$\begin{aligned}\{\sup(A, \leq)\} &= \sup\text{Set}(A, \leq) \\ &= \inf\text{Set}(A, \leq^*) \\ &= \{\inf(A, \leq^*)\}.\end{aligned}$$

Thus $\sup(A, \leq) = \inf(A, \leq^*)$.

2. **Similar to (1).**

□

Exercise 9.3.0.4. Let $(X, \leq)$ be a poset and $A, B \subset X$. If $A \subset B$, then

1. $\sup A$ and $\sup B$ exist implies that $\sup A \leq \sup B$.
2. $\inf A$ and $\inf B$ exist implies that $\inf B \leq \inf A$.

Proof. 1. Since $\sup A \in \sup\text{Set } A$ and $\sup B \in \sup\text{Set } B$, Exercise 9.2.0.7 implies that $\sup A \leq \sup B$.

2. **FINISH!!!**

□

Exercise 9.3.0.5. Associativity of Supremum and Infima:

Let $(X, \leq)$ be a poset and $(E_\alpha)_{\alpha \in A} \subset \mathcal{P}(X)$.

1. Suppose that

- for each $\alpha \in A$, $\sup E_\alpha$ exists
- $\sup_{\alpha \in A} \left[\sup E_\alpha \right]$ exists.

Then $\sup \bigcup_{\alpha \in A} E_\alpha$ exists and

$$\sup \bigcup_{\alpha \in A} E_\alpha = \sup_{\alpha \in A} \left[\sup E_\alpha \right]$$

2. Suppose that

- for each $\alpha \in A$, $\inf E_\alpha$ exists
- $\inf_{\alpha \in A} \left[\inf E_\alpha \right]$ exists.

Then $\inf \bigcup_{\alpha \in A} E_\alpha$ exists

$$\inf \bigcup_{\alpha \in A} E_\alpha = \inf_{\alpha \in A} \left[\inf E_\alpha \right]$$

Proof.

1. Define $s \in X$ by $s := \sup_{\alpha \in A} \left[\sup E_\alpha \right]$.

- Let $x \in \bigcup_{\alpha \in A} E_\alpha$. Then there exists $\alpha_0 \in A$ such that $x \in E_{\alpha_0}$. We note by definition, $s \in \text{ub}\{\sup E_\alpha : \alpha \in A\}$. Hence for each $\alpha \in A$, $s \geq \sup E_\alpha$. Since $\sup E_{\alpha_0} \in \text{ub } E_{\alpha_0}$, we have that

$$\begin{aligned}x &\leq \sup E_{\alpha_0} \\ &\leq s.\end{aligned}$$

Since $x \in \bigcup_{\alpha \in A} E_\alpha$ is arbitrary, we have that for each $x \in \bigcup_{\alpha \in A} E_\alpha$, $x \leq s$. Hence $s \in \text{ub} \bigcup_{\alpha \in A} E_\alpha$.

- Let $x \in \text{ub} \bigcup_{\alpha \in A} E_\alpha$ and $\alpha_0 \in A$. Since $E_{\alpha_0} \subset \bigcup_{\alpha \in A} E_\alpha$, Exercise 9.1.0.5 implies that

$$\begin{aligned} x &\in \text{ub} \bigcup_{\alpha \in A} E_\alpha \\ &\subset \text{ub} E_{\alpha_0}. \end{aligned}$$

Therefore $\sup E_{\alpha_0} \leq x$. Since $\alpha_0 \in A$ is arbitrary, we have that for each $\alpha_0 \in A$, $\sup E_{\alpha_0} \leq x$. Hence $x \in \text{ub}\{\sup E_\alpha : \alpha \in A\}$. Therefore $s \leq x$. Since $x \in \text{ub} \bigcup_{\alpha \in A} E_\alpha$ is arbitrary, we have that for each $x \in \text{ub} \bigcup_{\alpha \in A} E_\alpha$, $s \leq x$.

Therefore $s \in \text{supSet} \bigcup_{\alpha \in A} E_\alpha$ and $\sup \bigcup_{\alpha \in A} E_\alpha = s$.

2. Similar to (1). **Maybe fill out, use duality**

□

Exercise 9.3.0.6. Let $(X, \leq)$ be a poset and $a, b \in X$. Then

1. (a) $\sup\{a\}$ exists and $\sup\{a\} = a$,
(b) $a \leq b$ iff $\sup\{a, b\} = b$,
2. (a) $\inf\{a\}$ exists and $\inf\{a\} = a$,
(b) $a \leq b$ iff $\inf\{a, b\} = a$,
3. $\sup \downarrow a = a$,
4. $\inf \uparrow a = a$.

Proof.

1. (a) Since $a \leq a$, $a \in \text{ub}\{a\}$. By definition, for each $x \in \text{ub}\{a\}$, $a \leq x$. Hence $a \in \text{supSet}\{a\}$. Thus $\sup\{a\}$ exists and $a = \sup\{a\}$.
(b) • ($\implies$):
Suppose that $a \leq b$. Since $b \leq b$, we have that for each $c \in \{a, b\}$, $c \leq b$. Hence $b \in \text{lb}\{a, b\}$. Let $c \in \text{ub}\{a, b\}$. Then $b \leq c$. Since $c \in \text{ub}\{a, b\}$ is arbitrary, we have that for each $c \in \text{ub}\{a, b\}$, $b \leq c$. Hence $b \in \text{supSet}\{a, b\}$ and $b = \sup\{a, b\}$.
• ($\impliedby$):
Suppose that $\sup\{a, b\}$ exists and $b = \sup\{a, b\}$. Then $b \in \text{ub}\{a, b\}$. Therefore $a \leq b$.
2. (a) **FINISH!!!**
(b) **FINISH!!!**
3. Since for each $x \in \downarrow a$, $x \leq a$, we have that $a \in \text{ub} \downarrow a$. Since $a \in \downarrow a$, we have that for each $x \in \text{ub} \downarrow a$, $a \leq x$. Hence $a = \sup \downarrow a$.
4. Similar to (3).

□

Exercise 9.3.0.7. Let $(X, \leq)$ be a poset. Then for each $a, b \in X$, $\sup\{a, b\}$ exists iff for each $n \in \mathbb{N}$ and $(a_k)_{k=1}^n \in X^n$, $\sup\{a_1, \dots, a_n\}$ exists and

$$\sup\{a_1, \dots, a_n\} = \begin{cases} a_1, & n = 1 \\ \sup\left\{\sup\{a_1, \dots, a_{n-1}\}, a_n\right\}, & n \geq 2. \end{cases}$$

Proof.

- ($\implies$):
Suppose that for each $a, b \in X$, $\sup\{a, b\}$ exists. Let $n \in \mathbb{N}$ and $(a_k)_{k=1}^n \in X^n$.

– **Base Case:**

Suppose that $n = 1$. Exercise 9.3.0.6 implies that $\sup\{a_1\}$ exists and $\sup\{a_1\} = a_1$.

– **Induction Step:**

Suppose that $n \geq 2$ and $\sup\{a_1, \dots, a_{n-1}\}$ exists. Then by assumption, $\sup\left\{\sup\{a_1, \dots, a_{n-1}\}, a_n\right\}$ exists. Exercise 9.3.0.5 then implies that $\sup\{a_1, \dots, a_n\}$ exists and

$$\begin{aligned}\sup\{a_1, \dots, a_n\} &= \sup\left\{\{a_1, \dots, a_{n-1}\} \cup \{a_n\}\right\} \\ &= \sup\left\{\sup\{a_1, \dots, a_{n-1}\}, \sup\{a_n\}\right\} \\ &= \sup\left\{\sup\{a_1, \dots, a_{n-1}\}, a_n\right\}.\end{aligned}$$

By induction, for each $n \in \mathbb{N}$ and $(a_k)_{k=1}^n \in X^n$, $\sup\{a_1, \dots, a_n\}$ exists and

$$\sup\{a_1, \dots, a_n\} = \begin{cases} a_1, & n = 1 \\ \sup\left\{\sup\{a_1, \dots, a_{n-1}\}, a_n\right\}, & n \geq 2. \end{cases}$$

- ($\Leftarrow$):
Clear.

□

Exercise 9.3.0.8. Let $(X, \leq)$ be a poset. Then

1. for each $a, b \in \text{ub } X$, $a = b$,
2. for each $a \in \text{ub } X$, $\sup X = a$,
3. for each $a, b \in \text{lb } X$, $a = b$,
4. for each $a \in \text{lb } X$, $\inf X = a$

Proof.

1. Let $a, b \in \text{ub } X$. Since $a \in X$ and $b \in \text{ub } X$, $a \leq b$. Similarly, $b \leq a$. Hence $a = b$.
2. Let $a, b \in \text{ub } X$. **the previous part** implies that $a = b$. Since $b \in \text{ub } X$ is arbitrary, we have that for each $b \in \text{ub } X$, $a \leq b$. Thus $a = \sup X$.
3. Similar to (1) **ref.**
4. Similar to (2) **ref.**

□

Exercise 9.3.0.9. Let $(X, \leq_X)$ and $(Y, \leq_Y)$ be posets, $f : X \rightarrow Y$ and $g : Y \rightarrow X$. Suppose that (f, g) is a monotone Galois connection. Then

1. for each $A \subset X$, if $\sup(A, \leq_X)$ exists, then $\sup(f(A), \leq_Y)$ exists and $f(\sup(A, \leq_X)) = \sup(f(A), \leq_Y)$.
2. for each $B \subset Y$, if $\inf(B, \leq_Y)$ exists, then $g(\inf(B, \leq_Y)) = \inf(g(B), \leq_X)$. **FIX!!!**

Proof.

1. Suppose that $\sup(A, \leq_X)$ exists. Then $\sup\text{Set}(A, \leq_X) \neq \emptyset$. Exercise 9.2.0.8 then implies that

$$\begin{aligned} \emptyset &\neq f(\sup\text{Set}(A, \leq_X)) \\ &\subset \sup\text{Set}(f(A), \leq_Y). \end{aligned}$$

Hence $\sup(f(A), \leq_Y)$ exists and $f(\sup(A, \leq_X)) = \sup(f(A), \leq_Y)$.

2. ???

□

Exercise 9.3.0.10. Let $(X, \leq_X)$ and $(Y, \leq_Y)$ be posets, $f : X \rightarrow Y$ and $g : Y \rightarrow X$. Suppose that (f, g) is a monotone Galois connection. Then

1. for each $x \in X$, $f(x) = \inf(g^{-1}(\uparrow(x, \leq_X)), \leq_Y)$
2. for each $y \in Y$, $g(y) = \sup(f^{-1}(\downarrow(y, \leq_Y)), \leq_X)$

Proof.

1. Let $x \in X$. We note that for each $y \in Y$,

$$\begin{aligned} y \in g^{-1}(\uparrow(x, \leq_X)) &\iff g(y) \in \uparrow(x, \leq_X) \\ &\iff x \leq_X g(y) \\ &\iff f(x) \leq_Y y \quad (\text{since } (f, g) \text{ is a Galois connection}) \\ &\iff y \in \uparrow(f(x), \leq_Y). \end{aligned}$$

Thus $g^{-1}(\uparrow(x, \leq_X)) = \uparrow(f(x), \leq_Y)$. Exercise 9.3.0.6 then implies that

$$\begin{aligned} \inf(g^{-1}(\uparrow(x, \leq_X)), \leq_Y) &= \inf(\uparrow(f(x), \leq_Y), \leq_Y) \\ &= f(x). \end{aligned}$$

Since $x \in X$ is arbitrary, we have that for each $x \in X$, $f(x) = \inf(g^{-1}(\uparrow(x, \leq_X)), \leq_Y)$.

2. Let $y \in Y$. Since (f, g) is a $(\leq_X, \leq_Y)$ -monotone Galois connection, Exercise ?? ref prev ex in poset galois connection section implies that (g, f) is a $(\leq_Y^*, \leq_X^*)$ -monotone Galois connection. The previous part and exercises in previous section on poset upper bounds implies that

$$\begin{aligned} g(y) &= \inf(f^{-1}(\uparrow(y, \leq_Y^*)), \leq_X^*) \\ &= \sup(f^{-1}(\downarrow(y, \leq_Y)), \leq_X). \end{aligned}$$

Since $y \in Y$ is arbitrary, we have that for each $y \in Y$, $g(y) = \sup(f^{-1}(\downarrow(y, \leq_Y)), \leq_X)$.

□

Definition 9.3.0.11. Let $(X, \leq)$ be a poset.

- Then $(X, \leq)$ is said to
 - * **have a one** or **have a top** if $\sup(X, \leq)$ exists,
 - * **have a zero** or **have a bottom** if $\inf(X, \leq)$ exists.
- be a
 - * **poset with one** or **poset with top** if $(X, \leq)$ has a one,
 - * **poset with zero** or **poset with bottom** if $(X, \leq)$ has a zero.
- – If $(X, \leq)$ has a one, we define $1_\leq \in X$ by $1_\leq := \sup(X, \leq)$.
- If $(X, \leq)$ has a zero, we define $0_\leq \in X$ by $0_\leq := \inf(X, \leq)$.

Note 9.3.0.12.

- We the context is clear, we may write 1 and 0 in place of $1_{\leq}$ and $0_{\leq}$ respectively.
- Another common notation for 1 and 0 is $\top$ and $\perp$ respectively.

Exercise 9.3.0.13. Let X be a poset. Then

1. X has a top iff there exists $a \in X$ such that for each $x \in X$, $\inf\{x, a\} = x$.
2. X has a bottom iff there exists $a \in X$ such that for each $x \in X$, $\sup\{x, a\} = x$.

Proof.

1.
 - ($\implies$):
If X has a top, then there exists $1 \in X$ such that $1 = \sup X$. Since for each $x \in X$, $x \leq 1$, Exercise 9.3.0.6 implies that for each $x \in X$, $\inf\{x, 1\} = x$.
 - ($\impliedby$):
Suppose that there exists $a \in X$ such that for each $x \in X$, $\inf\{x, a\} = x$. Exercise 9.3.0.6 implies that for each $x \in X$, $x \leq a$. Therefore $a \in \text{ub}(X)$. Exercise 9.3.0.8 implies that $a = \sup X$. Hence X has a top.
2. **Similar to prev part FINISH!!!**

□

Part III

Lattices

pulling from [20]

Chapter 10

Lattices

10.1 Semilattices

10.1.1 Skew Semilattices

cite Leech, skew lattices in rings

Definition 10.1.1.1. Let X be a set and $\diamond : X \times X \rightarrow X$ a binary operator on X . Then

- $\diamond$ is said to be a **skew semilattice operator on X** if $\diamond$ is associative and idempotent.
- $(X, \diamond)$ is said to be a **skew semilattice** if
 1. $X \neq \emptyset$
 2. $\diamond$ is a skew semilattice operator on X

Definition 10.1.1.2. Let $(X, \diamond)$ be an skew semilattice. We define the

- **join preorder on X induced by $\diamond$** , denoted $\leq_{\diamond}^{\vee} \subset X \times X$, by $a \leq_{\diamond}^{\vee} b$ iff $(b \diamond a) \diamond b = b$
- **meet preorder on X induced by $\diamond$** , denoted $\leq_{\diamond}^{\wedge} \subset X \times X$, by $a \leq_{\diamond}^{\wedge} b$ iff $(a \diamond b) \diamond a = a$.

Exercise 10.1.1.3. Let $(X, \diamond)$ be an skew semilattice. Then

1. $\leq_{\diamond}^{\vee}$ is a preorder on X .
Hint: If $cbc = c$ and $bab = b$, then
 - (a) $c = (cb)(abc)(abc)$ and $c = cab$
 - (b) $cac = (ca)(ca)(bc)$
2. $\leq_{\diamond}^{\wedge}$ is a preorder on X .

Proof.

1. Let $a, b, c \in X$.
 - (a) Since $\diamond$ is idempotent, we have that

$$\begin{aligned}(a \diamond a) \diamond a &= a \diamond a \\ &= a\end{aligned}$$

and therefore $a \leq_{\diamond}^{\vee} a$.

(b) Suppose that $a \leq_{\diamond}^{\vee} b$ and $b \leq_{\diamond}^{\vee} c$. Then $(b \diamond a) \diamond b = b$ and $(c \diamond b) \diamond c = c$. Since $\diamond$ is associative and idempotent,

$$\begin{aligned}
 c &= cbc \\
 &= cbabc \\
 &= (cb)(abc) \\
 &= (cb)(abc)(abc) \\
 &= c(bab)cabc \\
 &= cbcabc \\
 &= (cbc)abc \\
 &= cabc.
 \end{aligned}$$

Therefore

$$\begin{aligned}
 cac &= (ca)(c) \\
 &= (ca)(cab) \\
 &= (ca)(ca)(bc) \\
 &= (ca)(bc) \\
 &= cabc \\
 &= c.
 \end{aligned}$$

Hence $a \leq_{\diamond}^{\vee} c$.

So $\leq_{\diamond}^{\vee}$ is a preorder on X .

2. Similar to (1).

□

Exercise 10.1.1.4. Let $(X, \diamond)$ be an skew semilattice. Then

1. $(\leq_{\diamond}^{\vee})^* = \leq_{\diamond}^{\wedge}$
2. $(\leq_{\diamond}^{\wedge})^* = \leq_{\diamond}^{\vee}$

Proof.

1. Let $a, b \in X$. Then

$$\begin{aligned}
 a(\leq_{\diamond}^{\vee})^*b &\iff b \leq_{\diamond}^{\vee} a \\
 &\iff a \diamond b \diamond a = a \\
 &\iff a \leq_{\diamond}^{\wedge} b.
 \end{aligned}$$

2. Similar to (1).

□

Exercise 10.1.1.5. Let $(X, \diamond)$ be an skew semilattice. Then for each $a, b \in X$,

1. $a \diamond b \in \text{ub}(\{a, b\}, \leq_{\diamond}^{\vee})$
2. $a \diamond b \in \text{lb}(\{a, b\}, \leq_{\diamond}^{\wedge})$

Proof. Let $a, b \in X$.

1. We note that

$$\begin{aligned}
 (a \diamond b) \diamond [a \diamond (a \diamond b)] &= (a \diamond b) \diamond [(a \diamond a) \diamond b] \\
 &= (a \diamond b) \diamond (a \diamond b) \\
 &= a \diamond b.
 \end{aligned}$$

Similarly, $[(a \diamond b) \diamond b] \diamond (a \diamond b) = a \diamond b$. Therefore $a \leq_{\diamond}^{\vee} a \diamond b$ and $b \leq_{\diamond}^{\vee} a \diamond b$. Hence $a \diamond b \in \text{ub}(\{a, b\}, \leq_{\diamond}^{\vee})$.

2. **FINISH!!!**

□

10.1.2 Skew Semilattice Homomorphisms

Definition 10.1.2.1. Let $(X, \diamond_X)$ and $(Y, \diamond_Y)$ be skew semilattices and $f : X \rightarrow Y$. Then f is said to be a $(\diamond_X, \diamond_Y)$ -**homomorphism** if for each $a, b \in X$, $f(a \diamond_X b) = f(a) \diamond_Y f(b)$.

Note 10.1.2.2. When the context is clear, we say homomorphism instead of $(\diamond_X, \diamond_Y)$ -homomorphism.

Exercise 10.1.2.3. Let $(X, \diamond_X)$ and $(Y, \diamond_Y)$ be skew semilattices and $f : X \rightarrow Y$. If f is a $(\diamond_X, \diamond_Y)$ -homomorphism, then

1. f is $(\leq_{\diamond_X}^{\vee}, \leq_{\diamond_Y}^{\vee})$ -monotone
2. f is $(\leq_{\diamond_X}^{\wedge}, \leq_{\diamond_Y}^{\wedge})$ -monotone

Proof. Suppose that f is a $(\diamond_X, \diamond_Y)$ -homomorphism.

1. Let $a, b \in X$. Suppose that $a \leq_{\diamond_X}^{\vee} b$. Then $(b \diamond_X a) \diamond_X b = b$. Therefore

$$\begin{aligned}
 f(b) &= (b \diamond_X a) \diamond_X b \\
 &= f(b \diamond_X a) \diamond_Y f(b) \\
 &= (f(b) \diamond_Y f(a)) \diamond_Y f(b)
 \end{aligned}$$

Thus $f(a) \leq_{\diamond_Y}^{\vee} f(b)$. Since $a, b \in X$ with $a \leq_{\diamond_X}^{\vee} b$ are arbitrary, we have that for each $a, b \in X$, $a \leq_{\diamond_X}^{\vee} b$ implies that $f(a) \leq_{\diamond_Y}^{\vee} f(b)$. Hence f is $(\leq_{\diamond_X}^{\vee}, \leq_{\diamond_Y}^{\vee})$ -monotone.

2. **use duality?, FINISH!!!**

□

10.1.3 Semilattices

Definition 10.1.3.1. Let X be a set and $\diamond : X \times X \rightarrow X$. Then

- $\diamond$ is said to be a **semilattice operator on X** if $\diamond$ associative, idempotent and commutative.
- $(X, \diamond)$ is said to be a **semilattice** if
 1. $X \neq \emptyset$
 2. $\diamond$ is a semilattice operator on X .

Exercise 10.1.3.2. Let X be a set and $\diamond : X \times X \rightarrow X$. Then $\diamond$ is a semilattice operator on X iff

1. $\diamond$ is a skew semilattice operator on X ,
2. $\diamond$ is commutative.

Proof. Clear by definition. □

Exercise 10.1.3.3. Let $(X, \diamond)$ be a semilattice. Then

1. for each $a, b \in X$,
 - (a) $a \leq_{\diamond}^{\vee} b$ iff $a \diamond b = b$
 - (b) $a \leq_{\diamond}^{\wedge} b$ iff $a \diamond b = a$
2. (a) $\leq_{\diamond}^{\vee}$ is a partial order on X
- (b) $\leq_{\diamond}^{\wedge}$ is a partial order on X

Proof.

1. Let $a, b \in X$.
 - (a) Since $\diamond$ is commutative, associative and idempotent,

$$\begin{aligned}
 a \leq_{\diamond}^{\vee} b &\iff (b \diamond a) \diamond b = b \\
 &\iff (a \diamond b) \diamond b = b \\
 &\iff a \diamond (b \diamond b) = b \\
 &\iff a \diamond b = b.
 \end{aligned}$$

- (b) Similar to 1(a).

2. (a) Exercise 10.1.1.3 implies that $\leq_{\diamond}^{\vee}$ is a preorder on X . Let $a, b \in X$. Suppose that $a \leq_{\diamond}^{\vee} b$ and $b \leq_{\diamond}^{\vee} a$. Part (1) then implies that $a \diamond b = b$ and $b \diamond a = a$. Since $\diamond$ is commutative,

$$\begin{aligned}
 a &= b \diamond a \\
 &= a \diamond b \\
 &= b.
 \end{aligned}$$

Since $a, b \in X$ are arbitrary, we have that for each $a, b \in X$, if $a \leq_{\diamond}^{\vee} b$ and $b \leq_{\diamond}^{\vee} a$, then $a = b$. Thus $\leq_{\diamond}^{\vee}$ is a partial order on X .

- (b) Similar to 2(a). □

Exercise 10.1.3.4. Let $(X, \diamond)$ be a semilattice. Then for each $a, b \in X$,

1. $a \diamond b = \sup(\{a, b\}, \leq_{\diamond}^{\vee})$
2. $a \diamond b = \inf(\{a, b\}, \leq_{\diamond}^{\wedge})$

Proof. Let $a, b \in X$.

- Exercise 10.1.1.5 implies that $a \diamond b \in \text{ub}(\{a, b\}, \leq_{\diamond}^{\vee})$. Let $c \in \text{ub}(\{a, b\}, \leq_{\diamond}^{\vee})$. Since $a \leq_{\diamond}^{\vee} c$ and $b \leq_{\diamond}^{\vee} c$, Exercise 10.1.3.3 implies that $a \diamond c = c$ and $b \diamond c = c$. Therefore

$$\begin{aligned} (a \diamond b) \diamond c &= a \diamond (b \diamond c) \\ &= a \diamond c \\ &= c. \end{aligned}$$

Another application of Exercise 10.1.3.3 implies that $a \diamond b \leq_{\diamond}^{\vee} c$. Since $c \in \text{ub}(\{a, b\}, \leq_{\diamond}^{\vee})$ is arbitrary, we have that for each $c \in \text{ub}(\{a, b\}, \leq_{\diamond}^{\vee})$, $a \diamond b \leq_{\diamond}^{\vee} c$. Therefore $a \diamond b = \sup(\{a, b\}, \leq_{\diamond}^{\vee})$.

- FINISH!!!**

□

Definition 10.1.3.5. Let $(X, \leq)$ be a poset. Then $(X, \leq)$ is said to be a

- **join semilattice** if

1. $X \neq \emptyset$
2. for each $a, b \in X$, $\sup\{a, b\}$ exists.

- **meet semilattice** if

1. $X \neq \emptyset$
2. for each $a, b \in X$, $\inf\{a, b\}$ exists.

Exercise 10.1.3.6. Let $(X, \leq)$ be a poset. Then

1. $(X, \leq)$ is a join semilattice iff $(X, \leq^*)$ is a meet semilattice
2. $(X, \leq^*)$ is a join semilattice iff $(X, \leq)$ is a meet semilattice

Proof.

- UPDATE ONCE SUPREMA/INFIMA EXERCISES ARE REORGANIZED**

$$\begin{aligned} (X, \leq) \text{ is a join semilattice} &\iff X \neq \emptyset \text{ and for each } a, b \in X, \sup(\{a, b\}, \leq) \text{ exists} \\ &\iff X \neq \emptyset \text{ and for each } a, b \in X, \inf(\{a, b\}, \leq^*) \text{ exists} \quad (\text{by Exercise ?? ADD EXERCISE REF}) \\ &\iff (X, \leq) \text{ is a meet semilattice} \end{aligned}$$

- 2.

□

Exercise 10.1.3.7. Let $(X, \diamond)$ be a semilattice. Then

1. $(X, \leq_{\diamond}^{\vee})$ is a join semilattice
2. $(X, \leq_{\diamond}^{\wedge})$ is a meet semilattice

Proof. Exercise 10.1.3.4 implies that

1. For each $a, b \in X$, $\sup(\{a, b\}, \leq_{\diamond}^{\vee}) = a \diamond b$. Thus $(X, \leq_{\diamond}^{\vee})$ is a join semilattice.
2. for each $a, b \in X$, $\inf(\{a, b\}, \leq_{\diamond}^{\wedge}) = a \diamond b$. Thus $(X, \leq_{\diamond}^{\wedge})$ is a meet semilattice.

□

Definition 10.1.3.8. Let $(X, \leq)$ be a poset.

- If $(X, \leq)$ is a join semilattice, we define the **join operator on X induced by $\leq$** , denoted $\vee_{\leq} : X \times X \rightarrow X$, by $a \vee_{\leq} b := \sup\{a, b\}$.
- If $(X, \leq)$ is a meet semilattice, we define the **meet operator on X induced by $\leq$** , denoted $\wedge_{\leq} : X \times X \rightarrow X$, by $a \wedge_{\leq} b := \inf\{a, b\}$.

Exercise 10.1.3.9. Let $(X, \leq)$ be a poset.

1. If $(X, \leq)$ is a join semilattice, then $\vee_{\leq}$ is a semilattice operator on X .
2. If $(X, \leq)$ is a meet semilattice, then $\wedge_{\leq}$ is a semilattice operator on X .

Proof.

1. Suppose that $(X, \leq)$ is a join semilattice.

- (a) • Let $a, b, c \in X$. Exercise 9.3.0.5 implies that

$$\begin{aligned}
 (a \vee_{\leq} b) \vee_{\leq} c &= \sup\{\sup\{a, b\}, c\} \\
 &= \sup\{\sup\{a, b\}, \sup\{c\}\} \\
 &= \sup\{a, b, c\} \\
 &= \sup\{\sup\{a\}, \sup\{b, c\}\} \\
 &= \sup\{a, \sup\{b, c\}\} \\
 &= a \vee_{\leq} (b \vee_{\leq} c).
 \end{aligned}$$

Hence $\vee_{\leq}$ is associative.

- Let $a \in X$. Then

$$\begin{aligned}
 a \vee_{\leq} a &= \sup\{a, a\} \\
 &= \sup\{a\} \\
 &= a.
 \end{aligned}$$

Hence $\vee_{\leq}$ is idempotent.

- Let $a, b \in X$. Then

$$\begin{aligned}
 a \vee_{\leq} b &= \sup\{a, b\} \\
 &= \sup\{b, a\} \\
 &= b \vee_{\leq} a.
 \end{aligned}$$

Hence $\vee_{\leq}$ is commutative.

Since $\vee_{\leq}$ is associative, idempotent and commutative, $\vee_{\leq}$ is a semilattice operator on X .

2. Similar to (1).

□

Exercise 10.1.3.10. Let $(X, \leq)$ be a poset.

1. If $(X, \leq)$ is a join semilattice, then

- (a) $\leq_{\vee_{\leq}}^{\vee} = \leq$,
 (b) $\leq_{\vee_{\leq}}^{\wedge} = \leq^*$.

2. If $(X, \leq)$ is a meet semilattice, then

- (a) $\leq_{\wedge_{\leq}}^{\wedge} = \leq$,
 (b) $\leq_{\wedge_{\leq}}^{\vee} = \leq^*$.

Proof.

1. Suppose that $(X, \leq)$ is a join semilattice.

(a) Let $a, b \in X$. Then

$$\begin{aligned} a \leq_{\vee_{\leq}}^{\vee} b &\iff a \vee_{\leq} b = b \quad (\text{Exercise 10.1.3.3}) \\ &\iff \sup(\{a, b\}, \leq) = b \quad (\text{Definition 10.1.3.8}) \\ &\iff a \leq b \quad (\text{Exercise 9.3.0.6}). \end{aligned}$$

Since $a, b \in X$ are arbitrary, we have that $\leq_{\vee_{\leq}}^{\vee} = \leq$.

(b) Exercise 10.1.1.4 implies that

$$\begin{aligned} \leq_{\vee_{\leq}}^{\wedge} &= (\leq_{\vee_{\leq}}^{\vee})^* \\ &= \leq^*. \end{aligned}$$

2. Similar to (1).

□

Exercise 10.1.3.11. Let $(X, \diamond)$ be a semilattice. Then

1. $\leq_{\diamond}^{\vee}$ is the unique partial order $\leq$ on X such that $(X, \leq)$ is a join semilattice and for each $a, b \in X$, $a \diamond b = \sup(\{a, b\}, \leq)$.
2. $\leq_{\diamond}^{\wedge}$ is the unique partial order $\leq$ on X such that $(X, \leq)$ is a meet semilattice and for each $a, b \in X$, $a \diamond b = \inf(\{a, b\}, \leq)$.

Proof.

1. Let $\leq$ be partial order on X . Suppose that $(X, \leq)$ is a join semilattice and for each $a, b \in X$, $a \diamond b = \sup(\{a, b\}, \leq)$. Let $a, b \in X$. Then

$$\begin{aligned} a \leq b &\iff b = \sup(\{a, b\}, \leq) \quad (\text{Exercise 9.3.0.6}) \\ &\iff a \diamond b = b \quad (\text{by assumption}) \\ &\iff a \leq_{\diamond}^{\vee} b \quad (\text{Exercise 10.1.3.3}). \end{aligned}$$

Since $a, b \in X$ are arbitrary, we have that for each $a, b \in X$, $a \leq b$ iff $a \leq_{\diamond}^{\vee} b$. Hence $\leq = \leq_{\diamond}^{\vee}$.

2. Similar to (1).

□

Exercise 10.1.3.12. Let $(X, \leq)$ be a poset.

1. If $(X, \leq)$ is a join semilattice, then $\vee_{\leq}$ is the unique semilattice operator $\diamond$ on X such that for each $a, b \in X$, $a \diamond b = \sup(\{a, b\}, \leq)$.
2. If $(X, \leq)$ is a meet semilattice, then $\wedge_{\leq}$ is the unique semilattice operator $\diamond$ on X such that for each $a, b \in X$, $a \diamond b = \inf(\{a, b\}, \leq)$.

Proof.

1. Let $\diamond : X \times X \rightarrow X$ be a semilattice operator on X . Suppose that $a, b \in X$, $a \diamond b = \sup(\{a, b\}, \leq)$. Then for each $a, b \in X$,

$$\begin{aligned} a \diamond b &= \sup(\{a, b\}, \leq) \quad (\text{by assumption}) \\ &= a \vee_{\leq} b \quad (\text{Definition 10.1.3.8}). \end{aligned}$$

Hence $\diamond = \vee_{\leq}$.

2. Similar to (1).

□

Exercise 10.1.3.13. Let $(L, \leq)$ be a poset.

1. Suppose that $(L, \leq)$ is a join semilattice. Then $(L, \leq)$ has a zero iff for each $A \subset L$, A is finite implies that $\sup A$ exists.
2. Suppose that $(L, \leq)$ is a meet semilattice. Then $(L, \leq)$ has a one iff for each $A \subset L$, A is finite implies that $\inf A$ exists.

Proof.

1. • $(\implies)$:

Suppose that $(L, \leq)$ has a zero. Let $A \subset L$. Suppose that A is finite.

- Suppose that $\#A = 0$. Then $A = \emptyset$. Exercise 9.3.0.2 implies that

$$\begin{aligned} \sup A \text{ exists} &\iff \inf(\text{ub } A) \text{ exists} \\ &\iff \inf(\text{ub } \emptyset) \text{ exists} \\ &\iff \inf L \text{ exists.} \end{aligned}$$

Since L has a zero, Definition 9.3.0.11 implies that $\inf L$ exists. Therefore $\sup A$ exists.

- Suppose that $\#A \geq 1$. Since $(L, \leq)$ is a join semilattice, we have that for each $a, b \in L$, $\sup\{a, b\}$ exists. Exercise 9.3.0.7 then implies that $\sup A$ exists.

- $(\impliedby)$:

Suppose that for each $A \subset L$, A is finite implies that $\sup A$ exists. Since $\emptyset$ is finite, $\sup \emptyset$ exists. By Definition 9.3.0.11, $(L, \leq)$ has a zero.

2. **FINISH!!!!**

□

10.1.4 Semilattice Homomorphisms

Definition 10.1.4.1. Let $(X, \leq_X), (Y, \leq_Y)$ be posets and $f : X \rightarrow Y$.

1. Suppose that $(X, \leq_X), (Y, \leq_Y)$ are join semilattices. Then f is said to be $(\leq_X, \leq_Y)$ -**binary join preserving** if for each $a, b \in X$, $f(\sup(\{a, b\}, \leq_X)) = \sup(\{f(a), f(b)\}, \leq_Y)$.
2. Suppose that $(X, \leq_X), (Y, \leq_Y)$ are meet semilattices. Then f is said to be $(\leq_X, \leq_Y)$ -**binary meet preserving** if for each $a, b \in X$, $f(\inf(\{a, b\}, \leq_X)) = \inf(\{f(a), f(b)\}, \leq_Y)$.

Exercise 10.1.4.2. Let $(X, \leq_X), (Y, \leq_Y)$ be posets and $f : X \rightarrow Y$.

1. Suppose that $(X, \leq_X), (Y, \leq_Y)$ are join semilattices. Then f is finite $(\leq_X, \leq_Y)$ -binary join preserving iff f is a $(\vee_{\leq_X}, \vee_{\leq_Y})$ -homomorphism.
2. Suppose that $(X, \leq_X), (Y, \leq_Y)$ are meet semilattices. Then f is finite $(\leq_X, \leq_Y)$ -binary meet preserving iff f is a $(\wedge_{\leq_X}, \wedge_{\leq_Y})$ -homomorphism.

Proof. Clear.

□

Exercise 10.1.4.3. Let $(X, \diamond_X)$ and $(Y, \diamond_Y)$ be semilattices and $f : X \rightarrow Y$. Then the following are equivalent:

1. f is a $(\diamond_X, \diamond_Y)$ -homomorphism
2. f is $(\leq_{\diamond_X}^\vee, \leq_{\diamond_Y}^\vee)$ -binary join preserving
3. f is $(\leq_{\diamond_X}^\wedge, \leq_{\diamond_Y}^\wedge)$ -binary meet preserving

Proof.

- $(1) \implies (2)$:

Suppose that f is a $(\diamond_X, \diamond_Y)$ -homomorphism. **make/ref previous ex here** implies that $\vee_{\leq_{\diamond_X}^\vee} = \diamond_X$ and $\vee_{\leq_{\diamond_Y}^\vee} = \diamond_Y$. Therefore f is a $(\vee_{\leq_{\diamond_X}^\vee}, \vee_{\leq_{\diamond_Y}^\vee})$ -homomorphism. **the previous ex** implies that f is $(\leq_{\diamond_X}^\vee, \leq_{\diamond_Y}^\vee)$ -binary join preserving.

- (2) $\implies$ (3):
Suppose that **FINISH!!!**
- (3) $\implies$ (1):

□

Exercise 10.1.4.4. Let $(X, \leq_X), (Y, \leq_Y)$ be posets and $f : X \rightarrow Y$.

1. Suppose that $(X, \leq_X)$ and $(Y, \leq_Y)$ are join semilattices. Then f is $(\leq_X, \leq_Y)$ -monotone iff for each $a, b \in X$, $f(a) \vee_{\leq_Y} f(b) \leq_Y f(a \vee_{\leq_X} b)$.
2. Suppose that $(X, \leq_X)$ and $(Y, \leq_Y)$ are meet semilattices. Then f is $(\leq_X, \leq_Y)$ -monotone iff for each $a, b \in X$, $f(a \vee_{\leq_X} b) \leq_Y f(a) \wedge_{\leq_Y} f(b)$.

Proof.

1. • ($\implies$):
Suppose that f is $(\leq_X, \leq_Y)$ -monotone. Let $a, b \in X$. Since $a \vee_{\leq_X} b \in \text{ub}(\{a, b\}, \leq_X)$, Exercise 9.1.0.13 implies that

$$\begin{aligned} f(a \vee_{\leq_X} b) &\in f(\text{ub}(\{a, b\}, \leq_X)) \\ &\subset \text{ub}(f(\{a, b\}), \leq_Y) \\ &= \text{ub}(\{f(a), f(b)\}, \leq_Y) \end{aligned}$$

Therefore

$$\begin{aligned} f(a) \vee_{\leq_Y} f(b) &= \sup(\{f(a), f(b)\}, \leq_Y) \\ &\leq f(a \vee_{\leq_X} b). \end{aligned}$$

Since $a, b \in X$ are arbitrary, we have that for each $a, b \in X$, $f(a) \vee_{\leq_Y} f(b) \leq_Y f(a \vee_{\leq_X} b)$.

- ($\impliedby$):
Suppose that for each $a, b \in X$, $f(a) \vee_{\leq_Y} f(b) \leq_Y f(a \vee_{\leq_X} b)$. Let $a, b \in X$. Suppose that $a \leq_X b$. Exercise 9.3.0.6 then implies that $a \vee_{\leq_X} b = b$. By assumption,

$$\begin{aligned} f(a) \vee_{\leq_Y} f(b) &\leq_Y f(a \vee_{\leq_X} b) \\ &= f(b) \\ &\leq f(a) \vee_{\leq_Y} f(b). \end{aligned}$$

Therefore $f(a) \vee_{\leq_Y} f(b) = f(b)$. Another application of Exercise 9.3.0.6 implies that $f(a) \leq_Y f(b)$. Since $a, b \in X$ with $a \leq b$ are arbitrary, we have that for each $a, b \in X$, $a \leq_X b$ implies that $f(a) \leq_Y f(b)$. Hence f is $(\leq_X, \leq_Y)$ -monotone.

2. **use duality FINISH!!!**

□

Exercise 10.1.4.5. Let $(X, \leq_X), (Y, \leq_Y)$ be posets and $f : X \rightarrow Y$.

1. Suppose that $(X, \leq_X)$ and $(Y, \leq_Y)$ are join semilattices. If f binary join preserving, then f is $(\leq_X, \leq_Y)$ -monotone.
2. Suppose that $(X, \leq_X)$ and $(Y, \leq_Y)$ are meet semilattices. If f binary meet preserving, then f is $(\leq_X, \leq_Y)$ -monotone.

Proof.

1. Suppose that f is $(\leq_X, \leq_Y)$ -binary join preserving. Then **the previous exercise** implies that

$$\begin{aligned} \text{for each } a, b \in X, f(a) \vee_{\leq_Y} f(b) &= f(a \vee_{\leq_X} b) \implies \text{for each } a, b \in X, f(a) \vee_{\leq_Y} f(b) \leq f(a \vee_{\leq_X} b) \\ &\implies f \text{ is } (\leq_X, \leq_Y)\text{-monotone} \end{aligned}$$

2. **FINISH!!!** , use duality or just similar

□

Definition 10.1.4.6. Let $(X, \leq_X), (Y, \leq_Y)$ be posets and $f : X \rightarrow Y$.

1. Suppose that $(X, \leq_X), (Y, \leq_Y)$ are join semilattices with zero. Then f is said to be $(\leq_X, \leq_Y)$ -**finite join preserving** if for each $A \subset X$, A is finite implies that $f(\sup(A, \leq_X)) = \sup(f(A), \leq_Y)$.
2. Suppose that $(X, \leq_X), (Y, \leq_Y)$ are meet semilattices with one. Then f is said to be $(\leq_X, \leq_Y)$ -**finite meet preserving** if for each $A \subset X$, A is finite implies that $f(\inf(A, \leq_X)) = \inf(f(A), \leq_Y)$.

Exercise 10.1.4.7. Let $(L, \leq_L)$ and $(M, \leq_M)$ be posets and $f : L \rightarrow M$.

1. Suppose that $(L, \leq_L)$ and $(M, \leq_M)$ are join semilattices with zero. Then f is $(\leq_L, \leq_M)$ -finite join preserving iff f is $(\leq_L, \leq_M)$ -binary join preserving and $f(0_L) = 0_M$.
2. Suppose that $(L, \leq_L)$ and $(M, \leq_M)$ are meet semilattices with one. Then f is $(\leq_L, \leq_M)$ -finite meet preserving iff f is $(\leq_L, \leq_M)$ -binary meet preserving and $f(1_L) = 1_M$.

Proof.

1. • $(\implies)$:
Suppose that f is $(\leq_L, \leq_M)$ -finite join preserving. Then clearly f is $(\leq_L, \leq_M)$ -binary join preserving. Since $\emptyset$ is finite, **a prev ex about $\sup \emptyset = 0$** implies that

$$\begin{aligned} f(0_L) &= f(\sup(\emptyset, \leq_L)) \\ &= \sup(f(\emptyset), \leq_M) \\ &= \sup(\emptyset, \leq_M) \\ &= 0_M. \end{aligned}$$

- $(\impliedby)$:
Suppose that f is $(\leq_L, \leq_M)$ -binary join preserving and $f(0_L) = 0_M$. Let $A \subset L$. Suppose that A is finite.
 - Suppose that $\#A = 0$. Then $A = \emptyset$ and

$$\begin{aligned} f(\sup(A, \leq_L)) &= f(\sup(\emptyset, \leq_L)) \\ &= f(0_L) \\ &= 0_M \\ &= \sup(\emptyset, \leq_M) \\ &= \sup(f(\emptyset), \leq_M) \\ &= \sup(f(A), \leq_M). \end{aligned}$$

- Suppose that $\#A = 1$. Then there exists $a \in L$ such that $A = \{a\}$. Hence

$$\begin{aligned} f(\sup(A, \leq_L)) &= f(\sup(\{a\}, \leq_L)) \\ &= f(a) \\ &= \sup(\{f(a)\}, \leq_M) \\ &= \sup(f(A), \leq_M). \end{aligned}$$

- Suppose that $\#A \geq 2$. Then there exists $n \in \mathbb{N}$ and $a_1, \dots, a_n \in L$ such that $n \geq 2$ and $A = \{a_1, \dots, a_n\}$. **ref prev ex** implies that $\sup(A, \leq_L) = (\dots((a_1 \vee_{\leq_L} a_2) \vee_{\leq_L} a_3) \dots a_n)$. Then

$$\begin{aligned} f(\sup(A, \leq_L)) &= f(\dots((a_1 \vee_{\leq_L} a_2) \vee_{\leq_L} a_3) \vee_{\leq_L} \dots \vee_{\leq_L} a_n) \\ &= \end{aligned}$$

do by induction on $\#A$ so it is less messy.

2.

FINISH!!!

□

10.1.5 Semilattice Structures

Definition 10.1.5.1. Let $(X, \leq)$ be a poset and $E \subset X$.

1. Suppose that $(X, \leq)$ is a join semilattice. Then $E \in \mathcal{U}_X$ iff for each $x \in X$ and $a \in E$, $x \vee a \in E$.
2. Suppose that $(X, \leq)$ is a meet semilattice. Then $E \in \mathcal{L}_X$ iff for each $x \in X$ and $a \in E$, $x \wedge a \in E$.

Proof.

1. • $(\implies)$:
Suppose that $E \in \mathcal{U}_X$. Then for each $x \in X$ and $a \in E$, $a \leq x$ implies that $x \in E$. Let $x \in X$ and $a \in E$. Since $a \leq x \vee a$, we have that $x \vee a \in E$. Since $x \in X$ and $a \in E$ are arbitrary, we have that for each $x \in X$ and $a \in E$, $x \vee a \in E$.

- $(\impliedby)$:
Suppose that for each $x \in X$ and $a \in E$, $x \vee a \in E$. Let $x \in X$ and $a \in E$. Suppose that $a \leq x$. Then

$$\begin{aligned} x &= x \vee a \\ &\in E. \end{aligned}$$

Since $x \in X$ and $a \in E$ with $a \leq x$ are arbitrary, we have that for each $x \in X$ and $a \in E$, $a \leq x$ implies that $x \in E$. Hence $E \in \mathcal{U}_X$.

2. • $(\implies)$:
Suppose that E is upward directed. Then for each $x, y \in E$, there exists $z \in E$ such that $x, y \leq z$. Let $x, y \in E$. Since $a \leq x \vee a$, we have that $x \vee a \in E$. Since $x \in X$ and $a \in E$ are arbitrary, we have that for each $x \in X$ and $a \in E$, $x \vee a \in E$.

- $(\impliedby)$:
Suppose that for each $x \in X$ and $a \in E$, $x \vee a \in E$. Let $x \in X$ and $a \in E$. Suppose that $a \leq x$. Then

$$\begin{aligned} x &= x \vee a \\ &\in E. \end{aligned}$$

Since $x \in X$ and $a \in E$ with $a \leq x$ are arbitrary, we have that for each $x \in X$ and $a \in E$, $a \leq x$ implies that $x \in E$. Hence $E \in \mathcal{U}_X$.

3. **FINISH!!! use duality**

□

Exercise 10.1.5.2. Let $(X, \leq)$ be a poset, $F \in \mathcal{U}_X$ and $I \in \mathcal{L}_X$.

1. Suppose that $(X, \leq)$ is a join semilattice. Then $I \in \mathcal{I}(X)$ iff $I \in \mathcal{L}_X$ and for each $x, y \in I$, $x \wedge y \in I$.
2. Suppose that $(X, \leq)$ is a meet semilattice. Then $F \in \mathcal{F}(X)$ iff $F \in \mathcal{U}_X$ and for each $x, y \in F$, $x \wedge y \in F$.

10.2 Lattices

10.2.1 Skew Lattices

Definition 10.2.1.1. Let X be a set and $\vee, \wedge$ skew semilattice operators on X . Then $\vee$ and $\wedge$ are said to **satisfy the skew lattice absorption identities** if for each $a, b \in X$,

1. $a \wedge (a \vee b) = a$
2. $a \vee (a \wedge b) = a$
3. $(b \vee a) \wedge a = a$
4. $(b \wedge a) \vee a = a$

Exercise 10.2.1.2. Let X be a set and $\vee, \wedge$ skew semilattice operators on X . Then $\vee$ and $\wedge$ satisfy the skew lattice absorption identities iff for each $a, b \in X$,

1. $a \vee b = b$ iff $a \wedge b = a$
2. $a \vee b = a$ iff $a \wedge b = b$

Proof.

• $(\implies)$:

Suppose that $\vee$ and $\wedge$ satisfy the skew lattice absorption identities. Let $a, b \in X$.

1. $-(\implies)$:

If $a \vee b = b$, then

$$\begin{aligned} a \wedge b &= a \wedge (a \vee b) \\ &= a. \end{aligned}$$

$-(\impliedby)$:

If $a \wedge b = a$, then

$$\begin{aligned} a \vee b &= (a \wedge b) \vee b \\ &= b. \end{aligned}$$

Thus $a \vee b = b$ iff $a \wedge b = a$.

2. Similar to (1)

• $(\impliedby)$:

Suppose that for each $a, b \in X$,

1. $a \vee b = b$ iff $a \wedge b = a$,
2. $a \vee b = a$ iff $a \wedge b = b$.

Let $a, b \in X$.

1. Set $c := a \vee b$. By assumption,

$$\begin{aligned} a \wedge (a \vee b) &= a \iff a \wedge c = a \\ &\iff a \vee c = c \\ &\iff a \vee (a \vee b) = c \\ &\iff (a \vee a) \vee b = c \\ &\iff a \vee b = c \\ &\iff c = c. \end{aligned}$$

Since $c = c$, we have that $a \wedge (a \vee b) = a$.

2. Similar to (1)
3. Similar to (1)
4. Similar to (1)

□

Exercise 10.2.1.3. Let X be a set and $\vee, \wedge$ skew semilattice operators on X . If $\vee$ and $\wedge$ satisfy the skew lattice absorption identities, then for each $a, b \in X$, $b \vee a \vee b = b$ iff $a \wedge b \wedge a = a$.

Hint: Exercise 10.2.1.2 implies that $b \vee (a \vee b) = b$ iff $b \wedge (a \vee b) = a \vee b$. Use, skew lattice absorption identities and associativity.

Proof. Suppose that $\vee$ and $\wedge$ satisfy the skew lattice absorption identities. Let $a, b \in X$.

- ($\implies$):

Suppose that $b \vee a \vee b = b$. Then Exercise 10.2.1.2 implies that $b \wedge (a \vee b) = (a \vee b)$. Therefore

$$\begin{aligned} a \wedge b &= (a \wedge b) \vee [(a \wedge b) \wedge (a \vee b)] \quad (\text{absorption}) \\ &= (a \wedge b) \vee [a \wedge (b \wedge [a \vee b])] \quad (\text{associativity}) \\ &= (a \wedge b) \vee [a \wedge (a \vee b)] \quad (\text{from earlier}) \\ &= (a \wedge b) \vee a \quad (\text{absorption}). \end{aligned}$$

Another application of Exercise 10.2.1.2 implies that $(a \wedge b) \wedge a = a$.

- ($\impliedby$):

Similar to ($\implies$).

□

Definition 10.2.1.4. Let X be a set and $\vee, \wedge$ skew semilattice operators on X . Then $(X, \vee, \wedge)$ is said to be a **skew lattice** if

1. $X \neq \emptyset$
2. $\vee$ and $\wedge$ satisfy the skew lattice absorption identities

Exercise 10.2.1.5. Let $(X, \vee, \wedge)$ be an skew lattice. Then

1. $\leq_{\vee}^{\vee} = \leq_{\wedge}^{\wedge}$.
2. $\leq_{\vee}^{\wedge} = \leq_{\wedge}^{\vee} = (\leq_{\vee}^{\vee})^*$.

Proof.

1. Let $a, b \in X$. Since $(X, \vee, \wedge)$ is an skew lattice, $\vee$ and $\wedge$ satisfy the skew lattice absorption identities. Exercise 10.2.1.3 then implies that

$$\begin{aligned} a \leq_{\vee}^{\vee} b &\iff b \vee a \vee b = b \\ &\iff a \wedge b \wedge a = a \\ &\iff a \leq_{\wedge}^{\wedge} b. \end{aligned}$$

Since $a, b \in X$ are arbitrary, we have that for each $a, b \in X$, $a \leq_{\vee}^{\vee} b$ iff $a \leq_{\wedge}^{\wedge} b$. Hence $\leq_{\vee}^{\vee} = \leq_{\wedge}^{\wedge}$.

2. Exercise 10.1.1.4 and part (1) imply that

$$\begin{aligned} \leq_{\vee}^{\wedge} &= (\leq_{\vee}^{\vee})^* \\ &= (\leq_{\wedge}^{\wedge})^* \\ &= \leq_{\wedge}^{\vee}. \end{aligned}$$

□

Note 10.2.1.6. Let $(X, \vee, \wedge)$ be an skew lattice. When the context is clear, we write $\leq$ in place of $\leq_{\vee}^{\vee}$.

10.2.2 Lattices

Definition 10.2.2.1. Let X be a set and $\vee, \wedge$ semilattice operators on X . Then $\vee$ and $\wedge$ are said to **satisfy the lattice absorption identities** if for each $a, b \in X$,

1. $a \wedge (a \vee b) = a$
2. $a \vee (a \wedge b) = a$

Exercise 10.2.2.2. Let X be a set and $\vee, \wedge$ semilattice operators on X . Then $\vee$ and $\wedge$ satisfy the lattice absorption identities iff $\vee$ and $\wedge$ satisfy the skew lattice absorption identities.

Proof.

- ($\implies$):
Clear by commutativity.
- ($\impliedby$):
Immediate.

□

Definition 10.2.2.3. Let X be a set and $\vee, \wedge$ semilattice operators on X . Then $(X, \vee, \wedge)$ is said to be a **lattice** if

1. $X \neq \emptyset$
2. $\vee$ and $\wedge$ satisfy the lattice absorption identities.

Definition 10.2.2.4. Let $(X, \leq)$ be a poset. Then $(X, \leq)$ is said to be an **ordered lattice** if $(X, \leq)$ is a join semilattice and $(X, \leq)$ is a meet semilattice.

Exercise 10.2.2.5. Let $(X, \vee, \wedge)$ be a lattice. Then $(X, \leq_{\vee}^{\vee})$ is an ordered lattice.

Proof. Exercise 10.1.3.7 implies that $(X, \leq_{\vee}^{\vee})$ is a join semilattice and $(X, \leq_{\wedge}^{\wedge})$ is a meet semilattice. Exercise 10.2.1.5 implies that $\leq_{\vee}^{\vee} = \leq_{\wedge}^{\wedge}$. Therefore $(X, \leq_{\vee}^{\vee})$ is a join semilattice and $(X, \leq_{\vee}^{\vee})$ is a meet semilattice. Hence $(X, \leq_{\vee}^{\vee})$ is an ordered lattice. □

Exercise 10.2.2.6. Let $(X, \leq)$ be an ordered lattice. Then $(X, \vee_{\leq}, \wedge_{\leq})$ is a lattice.

Proof. Let $a, b \in X$.

1. • Since $\sup\{a, b\} \in \text{ub}\{a, b\}$ is an upper bound of $\{a, b\}$, we have that $a \leq \sup\{a, b\}$. Since $a \leq a$, we have that $a \in \text{lb}\{a, \sup\{a, b\}\}$. Hence $a \leq \inf\{a, \sup\{a, b\}\}$.
- Since $\inf\{a, \sup\{a, b\}\} \in \text{lb}\{a, \sup\{a, b\}\}$, we have that $\inf\{a, \sup\{a, b\}\} \leq a$.

Since $a \leq \inf\{a, \sup\{a, b\}\}$ and $\inf\{a, \sup\{a, b\}\} \leq a$, we have that

$$\begin{aligned} a \wedge_{\leq} (a \vee_{\leq} b) &= \inf\{a, \sup\{a, b\}\} \\ &= a. \end{aligned}$$

2. Similarly, $a \vee_{\leq} (a \wedge_{\leq} b) = a$.

Since $a, b \in X$ are arbitrary, we have that for each $a, b \in X$,

1. $a \wedge_{\leq} (a \vee_{\leq} b) = a$
2. $a \vee_{\leq} (a \wedge_{\leq} b) = a$

Hence $\vee_{\leq}$ and $\wedge_{\leq}$ satisfy the lattice absorption identities. Therefore $(X, \vee_{\leq}, \wedge_{\leq})$ is a lattice. □

Exercise 10.2.2.7. Let $(X, \leq)$ be an ordered lattice. Then

1. $\leq_{\vee_{\leq}}^{\vee} = \leq_{\wedge_{\leq}}^{\wedge} = \leq$
2. $\leq_{\vee_{\leq}}^{\wedge} = \leq_{\wedge_{\leq}}^{\vee} = \leq^*$

Proof. Since $(X, \leq)$ is an ordered lattice, $(X, \leq)$ is a join semilattice $(X, \leq)$ is a meet semilattice. Exercise 10.1.3.10 then implies that

- 1.

$$\begin{aligned} \leq_{\vee_{\leq}}^{\vee} &= \leq \\ &= \leq_{\wedge_{\leq}}^{\wedge} \end{aligned}$$

2.

$$\begin{aligned}\leq_{\vee \leq}^{\wedge} &= \leq^* \\ &= \leq_{\wedge \leq}^{\vee}\end{aligned}$$

□

Exercise 10.2.2.8. Let $(X, \vee, \wedge)$ be a lattice. Then $\leq_{\vee}^{\vee}$ is the unique partial order $\leq$ on X such that $(X, \leq)$ is an ordered lattice and for each $a, b \in X$, $a \vee b = \sup(\{a, b\}, \leq)$.

Proof. Let $\leq$ be a partial order on X . Suppose that $(X, \leq)$ is an ordered lattice and for each $a, b \in X$, $a \vee b = \sup(\{a, b\}, \leq)$. Since $(X, \vee)$ is a semilattice, $(X, \leq_{\vee}^{\vee})$ and $(X, \leq)$ are join semilattices and for each $a, b \in X$, $a \vee b = \sup(\{a, b\}, \leq_{\vee}^{\vee})$ and $a \vee b = \sup(\{a, b\}, \leq)$, Exercise 10.1.3.11 implies that $\leq = \leq_{\vee}^{\vee}$. □

Exercise 10.2.2.9. Let $(X, \leq)$ be an ordered lattice. Then $\vee_{\leq}, \wedge_{\leq}$ are the unique semilattice operators $\vee, \wedge$ on X such that $(X, \vee, \wedge)$ is a lattice and for each $a, b \in X$, $a \vee b = \sup(\{a, b\}, \leq)$

Proof. Let $\vee, \wedge$ be semilattice operators on X . Suppose that $(X, \vee, \wedge)$ is a lattice and for each $a, b \in X$, $a \vee b = \sup(\{a, b\}, \leq)$. Then for each $a, b \in X$,

$$\begin{aligned}a \vee_{\leq} b &= \sup(\{a, b\}, \leq) \quad (\text{Definition 10.1.3.8}) \\ &= a \vee b \quad (\text{by assumption}).\end{aligned}$$

Thus $\vee = \vee_{\leq}$. Since $(X, \vee, \wedge)$ is a lattice and for each $a, b \in X$, $a \vee b = \sup(\{a, b\}, \leq)$, Exercise 10.2.2.8 implies that $\leq_{\vee}^{\vee} = \leq$. Exercise 10.2.1.5 then implies that

$$\begin{aligned}\leq_{\wedge}^{\wedge} &= \leq_{\vee}^{\vee} \\ &= \leq.\end{aligned}$$

Therefore for each $a, b \in X$,

$$\begin{aligned}a \wedge b &= \inf(\{a, b\}, \leq_{\wedge}^{\wedge}) \quad (\text{Exercise 10.1.3.4}) \\ &= \inf(\{a, b\}, \leq) \\ &= a \wedge_{\leq} b \quad (\text{Definition 10.1.3.8}).\end{aligned}$$

Hence $\wedge = \wedge_{\leq}$. □

Definition 10.2.2.10. Let $(X, \vee, \wedge)$ be a lattice. We define the

- **$(\vee, \wedge)$ -dual join operator on X** , denoted $\vee^* : X \times X \rightarrow X$, by $\vee^* := \wedge$
- **$(\vee, \wedge)$ -dual meet operator on X** , denoted $\wedge^* : X \times X \rightarrow X$, by $\wedge^* := \vee$
- **dual lattice of $(X, \vee, \wedge)$** , denoted $(X, \vee, \wedge)^*$, by $(X, \vee, \wedge)^* := (X, \vee^*, \wedge^*)$.

Note 10.2.2.11. When the context is clear, we may say “dual join operator on X ” and “dual meet operator on X ” in place of “ $(\vee, \wedge)$ -dual join operator on X ” and “ $(\vee, \wedge)$ -dual meet operator on X ” respectively.

Exercise 10.2.2.12. Let $(X, \leq)$ be an ordered lattice. Then

1. $\vee_{\leq}^* = \vee_{\leq^*}$ and $\wedge_{\leq}^* = \wedge_{\leq^*}$
2. $(\vee^*)^* = \vee$ and $(\wedge^*)^* = \wedge$

Proof.

1. For each $a, b \in X$,

$$\begin{aligned} a \vee_{\leq}^* b &= a \wedge_{\leq} b \\ &= \inf(\{a, b\}, \leq) \\ &= \sup(\{a, b\}, \leq^*) \quad (\text{by ref ex}) \\ &= a \vee_{\leq^*} b \end{aligned}$$

Hence $\vee_{\leq}^* = \vee_{\leq^*}$. Similarly, $\wedge_{\leq}^* = \wedge_{\leq^*}$.

2. By Definition 10.2.2.10,

$$\begin{aligned} (\vee^*)^* &= \wedge^* \\ &= \vee. \end{aligned}$$

Similarly, $(\wedge^*)^* = \vee$.

□

Exercise 10.2.2.13. Let $(X, \vee, \wedge)$ be a lattice. Then $\leq_{\vee^*}^{\vee} = (\leq_{\vee}^{\vee})^*$.

Proof. We have that

$$\begin{aligned} \leq_{\vee^*}^{\vee} &= \leq_{\wedge}^{\vee} \quad (\text{by Definition ?? ref prev definition}) \\ &= (\leq_{\vee}^{\vee})^* \quad (\text{by Exercise ?? ref prev ex}) \end{aligned}$$

□

Note 10.2.2.14. We the context is clear, we write $\leq_X$ and $\leq_{X^*}$ in place of $\leq_{\vee}^{\vee}$ and $\leq_{\vee^*}^{\vee}$ respectively.

10.2.3 Lattice Homomorphisms

???

10.3 Fundamental Examples

10.3.1 Bounded, Modular and Distributive Lattices

Definition 10.3.1.1. Let $(L, \leq)$ be a lattice. Then $(L, \leq, 0, 1)$ is said to be a **bounded lattice** if 0 is a $\leq$ -zero and 1 is a $\leq$ -one
 put some exercises?

Definition 10.3.1.2. Let L be a lattice. Then L is said to be a **distributive lattice** if for each $a, b, c \in L$, $a \vee (b \wedge c) = (a \vee b) \wedge (a \vee c)$.

Exercise 10.3.1.3. Let L be a lattice. Then for each $a, b, c \in L$,

1. $(a \wedge b) \vee (a \wedge c) \leq a \wedge (b \vee c)$.
2. $a \wedge (b \vee c) \leq (a \vee b) \wedge (a \vee c)$

Proof. Let $a, b, c \in L$.

1. • Since $a \wedge b \leq a$ and $a \wedge c \leq a$, we have that $a \in \text{ub}\{a \wedge b, a \wedge c\}$. Therefore

$$\begin{aligned} (a \wedge b) \vee (a \wedge c) &= \sup\{a \wedge b, a \wedge c\} \\ &\leq a. \end{aligned}$$

- Since

$$\begin{aligned} a \wedge b &\leq b \\ &\leq b \vee c \end{aligned}$$

and

$$\begin{aligned} a \wedge c &\leq c \\ &\leq b \vee c, \end{aligned}$$

we have that $b \vee c \in \text{ub}\{a \wedge b, a \wedge c\}$. Therefore

$$\begin{aligned} (a \wedge b) \vee (a \wedge c) &= \sup\{a \wedge b, a \wedge c\} \\ &\leq b \vee c. \end{aligned}$$

Then $(a \wedge b) \vee (a \wedge c) \in \text{lb}\{a, b \vee c\}$. Hence

$$\begin{aligned} (a \wedge b) \vee (a \wedge c) &\leq \inf\{a, b \vee c\} \\ &= a \wedge (b \vee c). \end{aligned}$$

2. **FINISH!!! (use duality)**

□

Exercise 10.3.1.4. Let L be a lattice. Then for each $a, b, c \in L$,

1. $c \leq a$ implies that $(a \wedge b) \vee c \leq a \wedge (b \vee c)$
2. $a \leq c$ implies that $a \vee (b \wedge c) \leq (a \vee b) \wedge c$

Proof. Let $a, b, c \in L$.

1. Suppose that $c \leq a$. Then $a \wedge c = c$. The previous exercise then implies that

$$\begin{aligned} (a \wedge b) \vee c &= (a \wedge b) \vee (a \wedge c) \\ &\leq a \wedge (b \vee c). \end{aligned}$$

2. **FINISH!!!** use duality

□

Exercise 10.3.1.5. Let L be a lattice. Then for each $a, b, c \in L$,

$$(a \wedge b) \vee (b \wedge c) \vee (c \wedge a) \leq (a \vee b) \wedge (b \vee c) \wedge (c \vee a).$$

Proof. Let $a, b, c \in L$.

- We first note that

$$a \wedge b \leq a \leq a \vee b,$$

$$b \wedge c \leq b \leq a \vee b,$$

$$c \wedge a \leq a \leq a \vee b.$$

Therefore $a \vee b \in \text{ub}\{a \wedge b, b \wedge c, c \wedge a\}$ and

$$\begin{aligned} (a \wedge b) \vee (b \wedge c) \vee (c \wedge a) &= \sup\{a \wedge b, b \wedge c, c \wedge a\} \\ &\leq a \vee b. \end{aligned}$$

- Similarly,

$$a \wedge b \leq b \leq b \vee c,$$

$$b \wedge c \leq c \leq b \vee c,$$

$$c \wedge a \leq c \leq b \vee c.$$

Therefore $b \vee c \in \text{ub}\{a \wedge b, b \wedge c, c \wedge a\}$ and

$$\begin{aligned} (a \wedge b) \vee (b \wedge c) \vee (c \wedge a) &= \sup\{a \wedge b, b \wedge c, c \wedge a\} \\ &\leq b \vee c. \end{aligned}$$

- Finally, we have that

$$a \wedge b \leq a \leq c \vee a,$$

$$b \wedge c \leq c \leq c \vee a,$$

$$c \wedge a \leq c \leq c \vee a.$$

Therefore $c \vee a \in \text{ub}\{a \wedge b, b \wedge c, c \wedge a\}$ and

$$\begin{aligned} (a \wedge b) \vee (b \wedge c) \vee (c \wedge a) &= \sup\{a \wedge b, b \wedge c, c \wedge a\} \\ &\leq c \vee a. \end{aligned}$$

Hence $(a \wedge b) \vee (b \wedge c) \vee (c \wedge a) \in \text{lb}\{a \vee b, b \vee c, c \vee a\}$ and

$$\begin{aligned} (a \wedge b) \vee (b \wedge c) \vee (c \wedge a) &\leq \inf\{a \vee b, b \vee c, c \vee a\} \\ &= (a \vee b) \wedge (b \vee c) \wedge (c \vee a). \end{aligned}$$

□

Exercise 10.3.1.6. Let L be a lattice. Then the following are equivalent:

1. For each $a, b, c \in L$, $c \leq a$ implies that $a \wedge (b \vee c) = (a \wedge b) \vee c$.
2. For each $a, b, c \in L$, $c \leq a$ implies that $a \wedge (b \vee c) = (a \wedge b) \vee (a \wedge c)$.
3. For each $p, q, r \in L$, $p \wedge (q \vee (p \wedge r)) = (p \wedge q) \vee (p \wedge r)$.

Proof.

1. (1) $\implies$ (2):

Suppose that for each $a, b, c \in L$, $c \leq a$ implies that $a \wedge (b \vee c) = (a \wedge b) \vee c$. Let $a, b, c \in L$. Suppose that $c \leq a$. Then $a \wedge c = c$. By assumption

$$\begin{aligned} a \wedge (b \vee c) &= (a \wedge b) \vee c \\ &= (a \wedge b) \vee (a \wedge c) \end{aligned}$$

Since $a, b, c \in L$ with $c \leq a$ are arbitrary, we have that for each $a, b, c \in L$, $c \leq a$ implies that $a \wedge (b \vee c) = (a \wedge b) \vee (a \wedge c)$.

2. (2) $\implies$ (3):

Suppose that for each $a, b, c \in L$, $c \leq a$ implies that $a \wedge (b \vee c) = (a \wedge b) \vee (a \wedge c)$. Let $p, q, r \in L$. Define $a, b, c \in L$ by $a := p$, $b := q$ and $c := p \wedge r$. Then

$$\begin{aligned} a \wedge c &= p \wedge (p \wedge r) \\ &= (p \wedge p) \wedge r \\ &= p \wedge r. \end{aligned}$$

By assumption,

$$\begin{aligned} p \wedge (q \vee (p \wedge r)) &= a \wedge (b \vee c) \\ &= (a \wedge b) \vee (a \wedge c) \\ &= (p \wedge q) \vee (p \wedge r). \end{aligned}$$

Since $p, q, r \in L$ are arbitrary, we have that for each $p, q, r \in L$, $p \wedge (q \vee (p \wedge r)) = (p \wedge q) \vee (p \wedge r)$.

3. (3) $\implies$ (1):

Suppose that for each $p, q, r \in L$, $p \wedge (q \vee (p \wedge r)) = (p \wedge q) \vee (p \wedge r)$. Let $a, b, c \in L$. Suppose that $c \leq a$. Define $p, q, r \in L$ by $p := a$, $q := b$ and $r := c$. Since $c \leq a$, we have that

$$\begin{aligned} p \wedge r &= a \wedge c \\ &= c. \end{aligned}$$

By assumption,

$$\begin{aligned} a \wedge (b \vee c) &= p \wedge (q \vee (p \wedge r)) \\ &= (p \wedge q) \vee (p \wedge r) \\ &= (a \wedge b) \vee c. \end{aligned}$$

Since $a, b, c \in L$ with $c \leq a$ are arbitrary, we have that for each $a, b, c \in L$, $c \leq a$ implies that $a \wedge (b \vee c) = (a \wedge b) \vee c$.

□

Definition 10.3.1.7. Let L be a lattice. Then L is said to be **modular** if for each $a, b, c \in L$,

$$c \leq a \text{ implies that } a \wedge (b \vee c) = (a \wedge b) \vee c.$$

Exercise 10.3.1.8. Let L be a lattice. Then the following are equivalent:

1. For each $a, b, c \in L$, $a \wedge (b \vee c) = (a \wedge b) \vee (a \wedge c)$.
2. For each $p, q, r \in L$, $p \vee (q \wedge r) = (p \vee q) \wedge (p \vee r)$.

Proof.

1. (1) $\implies$ (2):

Suppose that for each $a, b, c \in L$, $a \wedge (b \vee c) = (a \wedge b) \vee (a \wedge c)$. Let $p, q, r \in L$. Define $a, b, c \in L$ by $a := p \vee q$, $b := p$ and $c := r$. By absorption,

$$\begin{aligned} a \wedge b &= (p \vee q) \wedge p \\ &= p \wedge (p \vee q) \\ &= p. \end{aligned}$$

By assumption,

$$\begin{aligned} a \wedge c &= (p \vee q) \wedge r \\ &= r \wedge (p \vee q) \\ &= (r \wedge p) \vee (r \wedge q). \end{aligned}$$

Then by assumption and absorption,

$$\begin{aligned} (p \vee q) \wedge (p \vee r) &= a \wedge (b \vee c) \\ &= (a \wedge b) \vee (a \wedge c) \\ &= p \vee [(r \wedge p) \vee (r \wedge q)] \\ &= [p \vee (r \wedge p)] \vee (r \wedge q) \\ &= [p \vee (p \wedge r)] \vee (q \wedge r) \\ &= p \vee (q \wedge r). \end{aligned}$$

Since $p, q, r \in L$ are arbitrary, we have that for each $p, q, r \in L$, $p \vee (q \wedge r) = (p \vee q) \wedge (p \vee r)$.

2. (2) $\implies$ (1):

Suppose that for each $p, q, r \in L$, $p \vee (q \wedge r) = (p \vee q) \wedge (p \vee r)$. Let $a, b, c \in L$. Define $p, q, r \in L$ by $p := a \wedge b$, $q := a$ and $r := c$. By absorption,

$$\begin{aligned} p \vee q &= (a \wedge b) \vee a \\ &= a \vee (a \wedge b) \\ &= a. \end{aligned}$$

By assumption,

$$\begin{aligned} p \vee r &= (a \wedge b) \vee c \\ &= c \vee (a \wedge b) \\ &= (c \vee a) \wedge (c \vee b). \end{aligned}$$

Then by assumption and absorption,

$$\begin{aligned} (a \wedge b) \vee (a \wedge c) &= p \vee (q \wedge r) \\ &= (p \vee q) \wedge (p \vee r) \\ &= a \wedge [(c \vee a) \wedge (c \vee b)] \\ &= [a \wedge (c \vee a)] \wedge (c \vee b) \\ &= [a \wedge (a \vee c)] \wedge (b \vee c) \\ &= a \wedge (b \vee c). \end{aligned}$$

Since $a, b, c \in L$ are arbitrary, we have that for each $a, b, c \in L$, $a \wedge (b \vee c) = (a \wedge b) \vee (a \wedge c)$.

□

Definition 10.3.1.9. Let L be a lattice. Then L is said to be **distributive** if for each $a, b, c \in L$,

$$a \wedge (b \vee c) = (a \wedge b) \vee (a \wedge c).$$

Exercise 10.3.1.10. Let L be a lattice. If L is distributive, then L is modular.

Proof. Suppose that L is distributive. **FINISH!!!** □

Exercise 10.3.1.11. Let L be a lattice. Then for each $a, b, c \in L$, $a \vee (b \wedge c) = (a \vee b) \wedge (a \vee c)$ iff $a \wedge (b \vee c) = (a \wedge b) \vee (a \wedge c)$.

Proof. Let $a, b, c \in L$.

- $(\implies)$:
Suppose that $a \vee (b \wedge c) = (a \vee b) \wedge (a \vee c)$.
- $(\impliedby)$:

□

10.3.2 Heyting Algebras

Definition 10.3.2.1. Let $(L, \vee, \wedge)$ be a lattice and $\rightarrow: L \times L \rightarrow L$. Then

- $\rightarrow$ is said to be a $(\vee, \wedge)$ -**Heyting implication operator** if for each $a, x, y \in L$, $(x \wedge a) \leq_L y$ iff $x \leq_L (a \rightarrow y)$,
- $(L, \vee, \wedge, \rightarrow)$ is said to be a **Heyting algebra** if
 1. $(L, \vee, \wedge)$ is a lattice with zero,
 2. $\rightarrow: L \times L \rightarrow L$ is a $(L, \vee, \wedge)$ -Heyting implication operator.

(ask if there is any way to show that L has a zero instead of showing it. Maybe there is a result saying that this is impossible in general.)

Note 10.3.2.2. When the context is clear, we say “Heyting implication operator” instead of “ $(L, \vee, \wedge)$ -Heyting implication operator” and we write L in place of $(L, \vee, \wedge, \rightarrow)$.

Exercise 10.3.2.3. Let L be a Heyting algebra. Then

1. L is a lattice with one
2. for each $a \in L$, $a \rightarrow a = 1_L$.

(ask if there is any way to show that L has a zero instead of showing it. Maybe there is a result saying that this is impossible in general.)

Proof. Let $a \in L$. Define $y \in L$ by $y := a$. Then

$$\begin{aligned} (x \wedge a) &\leq a \\ &= y. \end{aligned}$$

Then by **definition**, we have that $x \leq a \rightarrow y$. Since $x \in L$ is arbitrary, we have that for each $x \in L$,

$$\begin{aligned} x &\leq (a \rightarrow y) \\ &= a \rightarrow a. \end{aligned}$$

Thus $a \rightarrow a$ is a one of L . Since $a \in L$ is arbitrary, we have that for each $a \in L$, $a \rightarrow a$ is a one of L .

1. Since L is a lattice, $L \neq \emptyset$. Thus there exists $a \in L$. Since $a \rightarrow a \in L$ and $a \rightarrow a$ is a one of L , L is a lattice with one.
2. **the part above** implies that for each $a \in L$, $a \rightarrow a = 1_L$.

□

Exercise 10.3.2.4. Let $(L, \vee, \wedge)$ be a lattice with zero and $\rightarrow: L \times L \rightarrow L$. Then $(L, \vee, \wedge, \rightarrow)$ is a Heyting algebra iff for each $a \in L$, $((\cdot \wedge a), (a \rightarrow \cdot))$ is a $(\leq_L, \leq_L)$ -monotone Galois connection.

Can we get rid of assumption about having 0?

Proof. For each $a \in L$, define $f_a, g_a: L \rightarrow L$ by $f_a := \cdot \wedge a$ and $g_a := a \rightarrow \cdot$.

• $(\implies)$:

Suppose that $(L, \vee, \wedge, \rightarrow)$ is a Heyting algebra. By definition, $\rightarrow: L \times L \rightarrow L$ is a $(L, \vee, \wedge)$ -Heyting implication operator. Therefore, for each $a, x, y \in L$,

$$\begin{aligned} f_a(x) \leq_L y &\iff (x \wedge a) \leq_L y \\ &\iff x \leq_L (a \rightarrow y) \\ &\iff x \leq_L g_a(y). \end{aligned}$$

Thus for each $a \in L$, (f_a, g_a) is a $(\leq_L, \leq_L)$ -monotone Galois connection.

• $(\impliedby)$:

Suppose that for each $a \in L$, (f_a, g_a) is a $(\leq_L, \leq_L)$ -monotone Galois connection.

1. By assumption, $(L, \vee, \wedge)$ is a lattice with zero.
2. We have that for each $a, x, y \in L$,

$$\begin{aligned} (x \wedge a) \leq_L y &\iff f_a(x) \leq_L y \\ &\iff x \leq_L g_a(y) \quad (\text{by assumption}) \\ &\iff x \leq_L (a \rightarrow y). \end{aligned}$$

Hence $\rightarrow: L \times L \rightarrow L$ is a $(\vee, \wedge)$ -Heyting implication operator.

Thus $(L, \vee, \wedge, \rightarrow)$ is a Heyting algebra. □

Exercise 10.3.2.5. Let L be a Heyting algebra. Then for each $a, b \in L$, $a \rightarrow b = \sup\{x \in L : x \wedge a \leq b\}$ (maybe write $\max A$? need to define $\max A$).

Proof. Let $a, b \in L$. Define $A \subset L$ by $A := \{x \in L : x \wedge a \leq b\}$. Define $y \in L$ by $y := b$. We note that

$$\begin{aligned} x \in A &\iff x \wedge a \leq b \\ &\iff x \wedge a \leq y \\ &\iff x \leq a \rightarrow y \quad (\text{by def}) \\ &\iff x \leq a \rightarrow b. \end{aligned}$$

1. Define $x_0 \in L$ by $x_0 := (a \rightarrow b)$. Since

$$\begin{aligned} x_0 &= a \rightarrow b \\ &\leq a \rightarrow b, \end{aligned}$$

we have that $x_0 \in A$.

2. Let $x \in A$. Then

$$\begin{aligned} x &\leq a \rightarrow b \\ &= x_0. \end{aligned}$$

Since $x \in A$ is arbitrary, we have that for each $x \in A$, $x \leq x_0$.

Exercise ?? ref prev ex implies that $x_0 = \sup A$ (maybe write $\max A$? need to define $\max A$). □

Exercise 10.3.2.6. Let L be a Heyting algebra. Then for each $x, y \in L$, $x \leq y$ iff $x \rightarrow y = 1$.

Proof. Let $x, y \in L$. We note that $1 \wedge x = x$ and for each $a \in L$, $a = 1$ iff $1 \leq a$. Therefore

$$\begin{aligned} x \rightarrow y = 1 &\iff 1 \leq x \rightarrow y \\ &\iff 1 \wedge x \leq y \\ &\iff x \leq y. \end{aligned}$$

Since $x, y \in L$ are arbitrary, we have that for each $x, y \in L$, $x \leq y$ iff $x \rightarrow y = 1$. □

Definition 10.3.2.7. Let $(L, \vee, \wedge, \rightarrow)$ be a Heyting algebra. We define the **$\rightarrow$ -negation operator**, denoted $\neg_{\rightarrow} : L \rightarrow L$, by $\neg_{\rightarrow} x := x \rightarrow 0$.

Note 10.3.2.8. When the context is clear, we write $\neg$ in place of $\neg_{\rightarrow}$.

Exercise 10.3.2.9. Let L be a Heyting algebra. Then $\neg : L \rightarrow L$ is a pseudocomplement.

Proof. **FINISH!!!!**, define pseudocomplement (and complement and orthocomplemented lattice, etc) □

10.3.3 Boolean Algebras

10.4 Lattice Structures

10.4.1 Irreducibility and Primality

Definition 10.4.1.1. Let $(L, \leq)$ be a poset.

- Suppose that $(L, \leq)$ is a join semilattice.
 - Let $x \in L$. Then x is said to be
 - * **$\leq$ -join irreducible** if
 1. $x \neq 0$,
 2. for each $a, b \in L$, $x = a \vee b$ implies that $x = a$ or $x = b$,
 - * **$\leq$ -join prime**
 1. $x \neq 0$,
 2. if for each $a, b \in L$, $x \leq a \vee b$ implies that $x \leq a$ or $x \leq b$.
 - We define
 - * $\text{irred}_{\vee}(L, \leq) := \{x \in L : x \text{ is } \leq\text{-join irreducible}\}$,
 - * $\text{prime}_{\vee}(L, \leq) := \{x \in L : x \text{ is } \leq\text{-join prime}\}$.
- Suppose that $(L, \leq)$ is a meet semilattice.
 - Let $x \in L$. Then x is said to be
 - * **$\leq$ -meet irreducible**
 1. $x \neq 1$,
 2. if for each $a, b \in L$, $x = a \wedge b$ implies that $x = a$ or $x = b$,
 - * **$\leq$ -meet prime** if
 1. $x \neq 1$,
 2. for each $a, b \in L$, $a \wedge b \leq x$ implies that $a \leq x$ or $b \leq x$.
 - We define
 - * $\text{irred}_{\wedge}(L, \leq) := \{x \in L : x \text{ is } \leq\text{-meet irreducible}\}$,
 - * $\text{prime}_{\wedge}(L, \leq) := \{x \in L : x \text{ is } \leq\text{-meet prime}\}$.

Note 10.4.1.2. When the context is clear,

- we say “join irreducible”, “join prime”, “meet irreducible” and “meet prime” in place of “ $\leq$ -join irreducible”, “ $\leq$ -join prime”, “ $\leq$ -meet irreducible” and “ $\leq$ -meet prime” respectively,
- we write $\text{irred}_{\vee}(L)$, $\text{prime}_{\vee}(L)$, $\text{irred}_{\wedge}(L)$ and $\text{prime}_{\wedge}(L)$ in place of $\text{irred}_{\vee}(L, \leq)$, $\text{prime}_{\vee}(L, \leq)$, $\text{irred}_{\wedge}(L, \leq)$ and $\text{prime}_{\wedge}(L, \leq)$ respectively

Exercise 10.4.1.3. Let $(L, \leq)$ be a poset.

1. Suppose that $(L, \leq)$ is a join semilattice. Then $\text{prime}_{\vee}(L) \subset \text{irred}_{\vee}(L)$.
2. Suppose that $(L, \leq)$ is a meet semilattice. Then $\text{prime}_{\wedge}(L) \subset \text{irred}_{\wedge}(L)$.

Proof.

1. Let $x \in \text{prime}_{\vee}(L)$.
 - (a) Since x is join prime, x is not at zero of L .
 - (b) Let $a, b \in L$. Suppose that $x = a \vee b$. Then $x \leq a \vee b$. Since x is join prime, $x \leq a$ or $x \leq b$.

- Suppose that $x \leq a$. Then

$$\begin{aligned} a \vee b &= x \\ &\leq a \\ &\leq a \vee b \end{aligned}$$

and therefore $x = a$. Thus $x \leq a$ implies that $x = a$.

- Similarly, $x \leq b$ implies that $x = b$.

Since $x \leq a$ or $x \leq b$, we have that $x = a$ or $x = b$. Since $a, b \in L$ with $x = a \vee b$ are arbitrary, we have that for each $a, b \in L$, $x = a \vee b$ implies that $x = a$ or $x = b$.

Thus $x \in \text{irred}_\vee(L)$. Since $x \in \text{prime}_\vee(L)$ is arbitrary, we have that for each $x \in \text{prime}_\vee(L)$, $x \in \text{irred}_\vee(L)$. Hence $\text{prime}_\vee(L) \subset \text{irred}_\vee(L)$.

2. Similar to (1) **use duality**.

□

10.4.2 Ideals and Filters

Definition 10.4.2.1. Let $(L, \vee, \wedge)$ be a lattice.

- Let $J \subset L$. Then
 - J is said to be an $(L, \vee, \wedge)$ -**ideal** if
 1. $J \neq \emptyset$,
 2. for each $a, b \in J$, $a \vee b \in J$,
 3. for each $x \in L$ and $x \in J$, $x \wedge a \in J$,
 - J is said to be an $(L, \vee, \wedge)$ -**filter** if
 1. $J \neq \emptyset$,
 2. for each $a, b \in J$, $a \wedge b \in J$,
 3. for each $x \in L$ and $a \in J$, $x \vee a \in J$.
- We define
 - $\mathcal{I}(L, \vee, \wedge) := \{I \subset L : I \text{ is an } (L, \vee, \wedge)\text{-ideal}\}$
 - $\mathcal{F}(L, \vee, \wedge) := \{F \subset L : F \text{ is a } (L, \vee, \wedge)\text{-filter}\}$

Note 10.4.2.2. When the context is clear,

- we may say “ideal” and “filter” instead of “ $(L, \vee, \wedge)$ -ideal” and “ $(L, \vee, \wedge)$ -filter” respectively,
- we may write $\mathcal{I}(L)$ and $\mathcal{F}(L)$ in place of $\mathcal{I}(L, \vee, \wedge)$ and $\mathcal{F}(L, \vee, \wedge)$ respectively.

Exercise 10.4.2.3. Let L be a lattice. If $L \neq \emptyset$, then

1. $L \in \mathcal{I}(L)$,
2. $L \in \mathcal{F}(L)$.

Proof.

1. (a) Since L is a lattice, $L \neq \emptyset$.
 (b) Clearly for each $a, b \in L$, $a \vee b \in L$
 (c) Clearly for each $x \in L$ and $b \in L$, $a \wedge b \in L$
 2. Similar to (1) **or use duality?**.

□

Exercise 10.4.2.4. Let L be a lattice. Then for each $x \in L$,

1. $\downarrow x \in \mathcal{I}(L)$
2. $\uparrow x \in \mathcal{F}(L)$

Proof. Let $x \in L$.

1. (a) Since $x \in \downarrow x$, $\downarrow x \neq \emptyset$.
- (b) Let $a, b \in \downarrow x$. Then $a \leq x$ and $b \leq x$. Hence $x \in \text{ub}(\{a, b\})$. Therefore

$$\begin{aligned} a \vee b &= \sup\{a, b\} \\ &\leq x. \end{aligned}$$

Thus $a \wedge b \in \downarrow x$. Since $a, b \in \downarrow x$ are arbitrary, we have that for each $a, b \in \downarrow x$, $a \vee b \in \downarrow x$.

- (c) Let $y \in L$ and $a \in \downarrow x$. Then $a \leq x$. Therefore

$$\begin{aligned} y \wedge a &= \inf(\{y, a\}) \\ &\leq a \\ &\leq x. \end{aligned}$$

Hence $y \wedge a \in \downarrow x$. Since $y \in L$ and $a \in \downarrow x$ are arbitrary, we have that for each $y \in L$ and $a \in \downarrow x$, $y \wedge a \in \downarrow x$.

2. use duality

□

Exercise 10.4.2.5. Let $(L, \vee, \wedge)$ be a lattice and $J \subset L$. Then

1. $J \in \mathcal{I}(L)$ iff $J \in \mathcal{F}(L^*)$,
2. $J \in \mathcal{F}(L)$ iff $J \in \mathcal{I}(L^*)$.

Proof.

1. • ($\implies$):
Suppose that $J \in \mathcal{I}(L)$.
 - Since $J \in \mathcal{I}(L, \vee, \wedge)$, $J \neq \emptyset$.
 - Since $J \in \mathcal{I}(L)$, for each $a, b \in J$,

$$\begin{aligned} a \wedge^* b &= a \vee b \\ &\in J. \end{aligned}$$

- Since $J \in \mathcal{I}(L)$, for each $x \in L$ and $a \in J$,

$$\begin{aligned} x \vee^* a &= x \wedge a \\ &\in J. \end{aligned}$$

Thus $J \in \mathcal{F}(L^*)$.

- ($\impliedby$) Suppose that $J \in \mathcal{F}(L^*)$. Then

2. Similar to (1) ref prev part

□

Exercise 10.4.2.6. Let L be a lattice and $J \subset L$. Then

1. (a) $J \in \mathcal{I}(L)$ iff J is a $\leq$ -lower set and J is $\leq$ -upward directed.
 (b) $J \in \mathcal{I}(L^*)$ iff J is a $\leq^*$ -lower set and J is $\leq^*$ -upward directed.
2. (a) $J \in \mathcal{F}(L)$ iff J is an $\leq$ -upper set and J is $\leq$ -downward directed.
 (b) $J \in \mathcal{F}(L^*)$ iff J is an $\leq^*$ -upper set and J is $\leq^*$ -downward directed.

FIX!!!
REWORK!!!!
THINK ABOUT WHAT STATEMENTS WE WANT AND SHORTEST WAY TO SHOW THEM

Proof.

1. (a) • ($\implies$):
 Suppose that $J \in \mathcal{I}(L)$.
 – Let $x \in L$ and $a \in J$. Suppose that $x \leq a$. Then $x = \inf(\{x, a\}, \leq)$ and therefore

$$\begin{aligned} x &= \inf(\{x, a\}, \leq) \\ &= x \wedge_{\leq} a \\ &\in J. \end{aligned}$$

Since $a \in J$ and $x \in L$ with $x \leq a$ are arbitrary, we have that J is a down set.

- * Since $J \in \mathcal{I}(L)$, we have that $J \neq \emptyset$.
 * Let $a, b \in J$. Since $J \in \mathcal{I}(L)$, we have that $a \vee b \in J$. Since $a \vee b = \sup(\{a, b\}, \leq)$ and $\{a, b\} \subset J$, we have that

$$\begin{aligned} a \vee b &\in \text{ub}(\{a, b\}, \leq) \cap J \\ &= \text{ub}(\{a, b\}, \leq |_J) \quad (\text{by Exercise 9.1.0.8}). \end{aligned}$$

Therefore $\text{ub}(\{a, b\}, \leq |_J) \neq \emptyset$. Since $a, b \in J$ are arbitrary, we have that for each $a, b \in J$, $\text{ub}(\{a, b\}, \leq |_J) \neq \emptyset$.
 Hence $\leq |_J$ is a direction on J .

Thus $(J, \leq |_J)$ is a directed set. Therefore J is $\leq$ -upward directed.

- ($\impliedby$):
 Suppose that J is a $\leq$ -lower set and J is $\leq$ -upward directed.
 i. Since J is $\leq$ -upward directed, $(J, \leq |_J)$ is a directed set. Hence $J \neq \emptyset$.
 ii. Let $a, b \in J$. Since $(J, \leq |_J)$ is a directed set and $\{a, b\} \subset J$, we have that

$$\begin{aligned} \text{ub}(\{a, b\}, \leq) \cap J &= \text{ub}(\{a, b\}, \leq |_J) \quad (\text{by Exercise 9.1.0.8}) \\ &\neq \emptyset. \end{aligned}$$

Thus there exists

$$c \in \text{ub}(\{a, b\}, \leq) \cap J \tag{10.1}$$

$$\subset \text{ub}(\{a, b\}, \leq). \tag{10.2}$$

Therefore $\sup(\{a, b\}, \leq) \leq c$. Since J is a $\leq$ -lower set, $c \in J$ and $\sup(\{a, b\}, \leq) \leq c$, we have that

$$\begin{aligned} a \vee b &= \sup(\{a, b\}, \leq) \\ &\in J. \end{aligned}$$

Since $a, b \in J$ are arbitrary, we have that for each $a, b \in J$, $a \vee b \in J$.

- iii. Let $x \in L$ and $a \in J$. Then $x \wedge a \leq a$. Since J is a $\leq$ -lower set, $x \wedge a \in J$. Since $x \in L$ and $a \in J$ are arbitrary, we have that for each $x \in L$ and $a \in J$, $x \wedge a \in J$.

Therefore $J \in \mathcal{I}(L)$.

- (b) Exercise 10.2.2.13 and the previous part imply that $J \in \mathcal{I}(L^*)$ iff J is an $\leq^*$ -lower set and J is $\leq^*$ -upward directed.

2. (a) We have that

$$\begin{aligned}
 J \in \mathcal{F}(L) &\iff J \in \mathcal{I}(L^*) \quad (\text{by prev part}) \\
 &\iff J \text{ is an } \leq^* \text{-lower set and } J \text{ is } \leq^* \text{-upward directed} \quad (\text{by prev part}) \\
 &\iff J \text{ is an } \leq \text{-upper set and } J \text{ is } \leq \text{-downward directed} \quad (\text{by prev ex in directed set section})
 \end{aligned}$$

(b) Exercise 10.2.2.13 and the previous part imply that $J \in \mathcal{F}(L^*)$ iff J is an $\leq^*$ -upper set and J is $\leq^*$ -downward directed.

$$\begin{aligned}
 J \in \mathcal{F}(L^*) &\iff J \text{ is an } \leq^* \text{-upper set and } J \text{ is } \leq \text{-downward directed} \quad (\text{by prev part}) \\
 &\iff J \text{ is a } \leq \text{-lower set and } J \text{ is } \leq \text{-upward directed} \\
 &\iff J \text{ is a } \leq \text{-lower set and } J \text{ is } \leq \text{-upward directed}
 \end{aligned}$$

□

Exercise 10.4.2.7. Let L be a lattice.

1. If L has a zero, then for each $I \in \mathcal{I}(L)$, then $0 \in I$.
2. If L has a one, then for each $F \in \mathcal{F}(L)$, then $1 \in F$.

Proof.

1. Suppose that L has a zero. Let $J \in \mathcal{I}(L)$. Since $I \in \mathcal{I}(L)$, $I \neq \emptyset$ and a previous ex implies I is a lower set. Hence there exists $a \in I$. a prev ex implies that $0 \in \text{lb } L$. Thus $0 \leq a$. Since I is a lower set, $0 \in I$.
2. Similar to (1).

□

Exercise 10.4.2.8. Let L be a lattice.

1. Let $(I_\alpha)_{\alpha \in A} \subset \mathcal{I}(L)$.
 - (a) If $\bigcap_{\alpha \in A} I_\alpha \neq \emptyset$, then $\bigcap_{\alpha \in A} I_\alpha \in \mathcal{I}(L)$.
 - (b) If $A = \{1, 2\}$, then $\bigcap_{\alpha \in A} I_\alpha \in \mathcal{I}(L)$.
 - (c) If L has a zero, then $\bigcap_{\alpha \in A} I_\alpha \in \mathcal{I}(L)$.
2. Let $(F_\alpha)_{\alpha \in A} \subset \mathcal{F}(L)$.
 - (a) If $\bigcap_{\alpha \in A} F_\alpha \neq \emptyset$, then $\bigcap_{\alpha \in A} F_\alpha \in \mathcal{F}(L)$.
 - (b) If $A = \{1, 2\}$, then $\bigcap_{\alpha \in A} F_\alpha \in \mathcal{F}(L)$.
 - (c) If L has a one, then $\bigcap_{\alpha \in A} F_\alpha \in \mathcal{F}(L)$.

Proof.

1. Set $I := \bigcap_{\alpha \in A} I_\alpha$.
 - (a) Suppose that $I \neq \emptyset$.
 - i. By assumption $I \neq \emptyset$.
 - ii. Let $a, b \in I$ and $\alpha \in A$. Then $a, b \in I_\alpha$. Since $I_\alpha \in \mathcal{I}(L)$, $a \vee b \in I_\alpha$. Since $\alpha \in A$ is arbitrary, we have that for each $\alpha \in A$, $a \vee b \in I_\alpha$. Hence $a \vee b \in I$.

- iii. Let $x \in L$, $a \in I$ and $\alpha \in A$. Then $a \in I_\alpha$. Since $I_\alpha \in \mathcal{I}(L)$, $x \in L$ and $a \in I_\alpha$, we have that $x \wedge a \in I_\alpha$. Since $\alpha \in A$ is arbitrary, we have that for each $\alpha \in A$, $x \wedge a \in I_\alpha$. Hence $x \wedge a \in I$. Since $x \in L$ and $a \in I$ are arbitrary, we have that for each $x \in L$ and $a \in I$, $x \wedge a \in I$.

Thus $I \in \mathcal{I}(L)$.

- (b) Suppose that $A = \{1, 2\}$.

- i. Since $I_1, I_2 \in \mathcal{I}(L)$, $I_1 \neq \emptyset$ and $I_2 \neq \emptyset$. Thus there exist $x_1 \in I_1$ and $x_2 \in I_2$. Since $I_1 \in \mathcal{I}(L)$, $x_2 \in L$ and $x_1 \in I_1$, we have that $x_1 \wedge x_2 \in I_1$. Similarly, $x_1 \wedge x_2 \in I_2$. Hence $x_1 \wedge x_2 \in I_1 \cap I_2$ and $I_1 \cap I_2 \neq \emptyset$. Part 1(a) implies that $I_1 \cap I_2 \in \mathcal{I}(L)$.

- (c) Suppose that L has a zero. Let $\alpha \in A$. Since $I_\alpha \in \mathcal{I}(L)$, a previous exercise implies that $0 \in I_\alpha$. Since $\alpha \in A$ is arbitrary, we have that for each $\alpha \in A$, $0 \in I_\alpha$. Hence $0 \in I$. Thus $I \neq \emptyset$. The previous part implies that $I \in \mathcal{I}(L)$.

2. Similar to (1).

□

Definition 10.4.2.9. (consider changing notation) Let L be a lattice and $A \subset L$.

- We define

- $\text{Ideals}(A, L) := \{I \in \mathcal{I}(L) : S \subset I\}$
- $\text{Filters}(A, L) := \{F \in \mathcal{F}(L) : S \subset F\}$

- We define

- the L -ideal generated by A , denoted $\text{IdealGen}(A, L)$, by $\text{IdealGen}(A, L) := \bigcap_{I \in \text{Ideals}(A, L)} I$
- the L -filter generated by A , denoted $\text{FilterGen}(A, L)$, by $\text{FilterGen}(A, L) := \bigcap_{F \in \text{Filters}(A, L)} F$

Note 10.4.2.10. the prev ex implies that $\text{IdealGen}(A, L) \in \text{Ideals}(A, L)$ and $\text{FilterGen}(A, L) \in \text{Filters}(A, L)$.

Exercise 10.4.2.11. Let L be a lattice and $B \subset L$.

1. If B is upward directed, then
 - (a) $\downarrow B \in \mathcal{I}(L)$
 - (b) $\text{IdealGen}(B) = \downarrow(B, \leq_L)$.
2. If B is downward directed, then
 - (a) $\uparrow B \in \mathcal{F}(L)$
 - (b) $\text{FilterGen}(B) = \uparrow(B, \leq_L)$.

Proof.

1. Suppose that B is upward directed.

- (a) Exercise ??(ref ex) implies that $\downarrow B$ is upward directed and Exercise ?? (ref ex) implies that $\downarrow B$ is an lower set. Exercise ?? then implies that $\downarrow B \in \mathcal{I}(L)$.
- (b) The prev part implies that $\downarrow B \in \text{Ideals}(B, L)$. Therefore

$$\begin{aligned} \text{IdealGen}(B, L) &= \bigcap_{I \in \text{Ideals}(B, L)} I \\ &\subset \downarrow B. \end{aligned}$$

We note that for each $I \in \mathcal{I}(L)$, I is a lower set. Thus $\text{Ideals}(B, L) \subset \mathcal{L}(B, \leq_L)$. Hence

$$\begin{aligned} \downarrow B &= \bigcap_{I \in \text{Ideals}(B, L)} I \\ &\subset \uparrow B. \end{aligned}$$

2. Suppose that B is $\leq$ -downward directed. Exercise ?? implies that B is $\leq^*$ -upward directed. the prev part implies that $\uparrow(B, \leq^*) \in \mathcal{I}(L)$. Exercise ?? then implies that $\uparrow(B, \leq^*) \in \mathcal{I}(L)$

make new notation $\leq_L :=$ the unique partial order on L compatible with $\vee, \wedge$. Show $\leq_L^* = \leq_L$, make note that we write $\leq$ in place

□

Definition 10.4.2.12. define prefilter or filter basis, preideal or ideal basis as a downward directed/upward directed set, maybe don't need to, but look into what we need for topology.

Exercise 10.4.2.13. Let L be a lattice. Then

1. for each $I_1, I_2 \in \mathcal{I}(L)$, $\{x \in L : \text{there exist } a \in I_1 \text{ and } b \in I_2 \text{ such that } x \leq a \vee b\} \in \mathcal{I}(L)$.
2. for each $F_1, F_2 \in \mathcal{F}(L)$, $\{x \in L : \text{there exist } a \in F_1 \text{ and } b \in F_2 \text{ such that } a \vee b \leq x\} \in \mathcal{F}(L)$ (maybe FIX!!!).

Proof.

1. Let $I_1, I_2 \in \mathcal{I}(L)$. Set $I := \{x \in L : \text{there exist } a \in I_1 \text{ and } b \in I_2 \text{ such that } x \leq a \vee b\}$.
 - (a) Since $I_1, I_2 \in \mathcal{I}(L)$, we have that $I_1 \neq \emptyset$ and $I_2 \neq \emptyset$. Hence there exist $x_1 \in I_1$ and $x_2 \in I_2$. Since $x_1 \leq x_1 \vee x_2$, we have that $x_1 \in I$. Thus $I \neq \emptyset$.
 - (b) Let $a, b \in I$. Then there exist $x_a, x_b \in I_1$ and $y_a, y_b \in I_2$ such that $a \leq x_a \vee y_a$ and $b \leq x_b \vee y_b$. Define $z_1, z_2 \in L$ by $z_1 := x_a \vee x_b$ and $z_2 := y_a \vee y_b$. Since $I_1, I_2 \in \mathcal{I}(L)$, we have that $z_1 \in I_1$ and $z_2 \in I_2$. We note that

$$\begin{aligned} a &\leq x_a \vee y_a \\ &\leq (x_a \vee y_a) \vee (x_b \vee y_b) \\ &= (x_a \vee x_b) \vee (y_a \vee y_b) \\ &= z_1 \vee z_2. \end{aligned}$$

and similarly $b \leq z_1 \vee z_2$. Therefore $z_1 \vee z_2 \in \text{ub}(\{a, b\}, \leq)$. Hence

$$\begin{aligned} a \vee b &= \sup(\{a, b\}, \leq) \\ &\leq z_1 \vee z_2. \end{aligned}$$

Thus $a \vee b \in I$. Since $a, b \in I$ are arbitrary, we have that for each $a, b \in I$, $a \vee b \in I$.

- (c) Let $x \in L$ and $y \in I$. Then there exists $a \in I_1$ and $b \in I_2$ such that $y \leq a \vee b$. Then

$$\begin{aligned} x \wedge y &\leq y \\ &\leq a \vee b. \end{aligned}$$

Therefore $x \wedge y \in I$. Since $x \in L$ and $y \in I$ are arbitrary, we have that for each $x \in L$ and $y \in I$, $x \wedge y \in I$.

Therefore $I \in \mathcal{I}(L)$.

2. Similar to (1). FINISH!!!

□

Definition 10.4.2.14. Let L be a lattice. Define $\leq_{\mathcal{I}(L)} := \subset |_{\mathcal{I}(L)}$ (maybe standardize notation).

Exercise 10.4.2.15. Let L be a lattice.

1. Let $I_1, I_2 \in \mathcal{I}(L)$.
 - (a) $\sup(\{I_1, I_2\}, \leq_{\mathcal{I}(L)}) = \{x \in L : \text{there exist } a \in I_1 \text{ and } b \in I_2 \text{ such that } x \leq a \vee b\}$.
 - (b) $\inf(\{I_1, I_2\}, \leq_{\mathcal{I}(L)}) = I_1 \cap I_2$
2. Set $\leq_{\mathcal{F}(L)} := \subset |_{\mathcal{F}(L)}$ (maybe standardize notation). Let $F_1, F_2 \in \mathcal{F}(L)$. FIX!!!

- (a) $\sup(\{F_1, F_2\}, \leq_{\mathcal{I}(L)}) = \{x \in L : \text{there exist } a \in F_1 \text{ and } b \in F_2 \text{ such that } x \leq a \vee b\}.$
 (b) $\inf(\{F_1, F_2\}, \leq_{\mathcal{F}(L)}) = F_1 \cap F_2$

Proof.

1. (a) Set $I := \{x \in L : \text{there exist } a \in I_1 \text{ and } b \in I_2 \text{ such that } x \leq a \vee b\}$. (ref previous ex) implies that $I \in \mathcal{I}(L)$.
 - i. • Let $a \in I_1$. Since $I_2 \in \mathcal{I}(L)$, $I_2 \neq \emptyset$. Hence there exists $b \in I_2$. Then $a \leq a \vee b$. Since $a \in I_1$, $b \in I_2$ and $a \leq a \vee b$, we have that $a \in I$. Since $a \in I_1$ is arbitrary, we have that for each $a \in I_1$, $a \in I$. Thus $I_1 \subset I$.
 • Similarly, $I_2 \subset I$.
 Since $I_1, I_2 \subset I$, we have that $I \in \text{ub}(\{I_1, I_2\}, \leq_{\mathcal{I}(L)})$.
 - ii. Let $K \in \text{ub}(\{I_1, I_2\}, \leq_{\mathcal{I}(L)})$ and $x \in I$. Then $I_1, I_2 \subset K$ and there exist $a \in I_1$ and $b \in I_2$ such that $x \leq a \vee b$. Since $a \in I_1$, $b \in I_2$ and $I_1, I_2 \subset K$, we have that $a, b \in K$. Since $K \in \mathcal{I}(L)$, $a \vee b \in K$. Since $K \in \mathcal{I}(L)$ (ref a prev ex) implies that K is a lower set. Since K is a lower set, $a \vee b \in K$ and $x \leq a \vee b$, we have that $x \in K$. Since $x \in I$ is arbitrary, we have that for each $x \in I$, $x \in K$. Hence $I \subset K$. Since $K \in \text{ub}(\{I_1, I_2\}, \leq_{\mathcal{I}(L)})$ is arbitrary, we have that for each $K \in \text{ub}(\{I_1, I_2\}, \leq_{\mathcal{I}(L)})$, $I \leq_{\mathcal{I}(L)} K$.

Since

- i. $I \in \text{ub}(\{I_1, I_2\}, \leq_{\mathcal{I}(L)})$,
- ii. for each $K \in \text{ub}(\{I_1, I_2\}, \leq_{\mathcal{I}(L)})$, $I \leq_{\mathcal{I}(L)} K$,

we have that $I = \sup(\{I_1, I_2\}, \leq_{\mathcal{I}(L)})$.

- (b) (ref previous ex) implies that $I_1 \cap I_2 \in \mathcal{I}(L)$. Since $I_1 \cap I_2 \subset I_1$ and $I_1 \cap I_2 \subset I_2$, we have that $I_1 \cap I_2 \in \text{lb}(\{I_1, I_2\}, \leq_{\mathcal{I}(L)})$. Since $(\mathcal{I}(L), \leq_{\mathcal{I}(L)})$ is a subposet of $(\mathcal{P}(L), \subset)$, (define subposet/subposet and make exercise for the following fact) implies that

$$\begin{aligned} \inf(\{I_1, I_2\}, \leq_{\mathcal{I}(L)}) &\leq \inf(\{I_1, I_2\}, \subset) \\ &= I_1 \cap I_2 \\ &\leq \inf(\{I_1, I_2\}, \leq_{\mathcal{I}(L)}). \end{aligned}$$

Hence $I_1 \cap I_2 = \inf(\{I_1, I_2\}, \leq_{\mathcal{I}(L)})$.

2. Similar to (1). **FINISH!!!**

□

Definition 10.4.2.16. Let L be a lattice. Define $\vee_{\mathcal{I}(L)}, \wedge_{\mathcal{I}(L)} : \mathcal{I}(L) \times \mathcal{I}(L) \rightarrow \mathcal{I}(L)$ and $\vee_{\mathcal{F}(L)}, \wedge_{\mathcal{F}(L)} : \mathcal{F}(L) \times \mathcal{F}(L) \rightarrow \mathcal{F}(L)$ by

1. • $I_1 \vee_{\mathcal{I}(L)} I_2 := \{x \in L : \text{there exist } a \in I_1 \text{ and } b \in I_2 \text{ such that } x \leq a \vee b\}$
 • $I_1 \wedge_{\mathcal{I}(L)} I_2 := I_1 \cap I_2$.
2. **FIX!!!**
 - $F_1 \vee_{\mathcal{F}(L)} F_2 := \{x \in L : \text{there exist } a \in F_1 \text{ and } b \in F_2 \text{ such that } x \leq a \vee b\}$
 - $F_1 \wedge_{\mathcal{F}(L)} F_2 := F_1 \cap F_2$.

Exercise 10.4.2.17. Let L be a lattice. Then

1. $(\mathcal{I}(L), \vee_{\mathcal{I}(L)}, \wedge_{\mathcal{I}(L)})$ is a lattice
2. $(\mathcal{F}(L), \vee_{\mathcal{F}(L)}, \wedge_{\mathcal{F}(L)})$ is a lattice

Proof.

1. the previous exercise implies that $(\mathcal{I}(L), \leq_{\mathcal{I}(L)})$ is an ordered lattice. Exercise 10.2.2.6 then implies that $(\mathcal{I}(L), \vee_{\mathcal{I}(L)}, \wedge_{\mathcal{I}(L)})$ is a lattice.
- 2.

□

10.4.3 Properties of Ideals and Filters

Definition 10.4.3.1. Let L be a lattice.

- Let $I \in \mathcal{I}(L)$. Then I is said to be
 - **proper** if $I \neq L$.
 - **maximal** if I is maximal in $(\mathcal{I}(L) \setminus \{L\}, \subset)$.
 - **prime** if
 1. I is proper
 2. for each $a, b \in L$, $a \wedge b \in I$ implies that $a \in I$ or $b \in I$.
 - **principal** if there exists $a \in L$ such that $I = \downarrow a$.
- Let $F \in \mathcal{F}(L)$. Then F is said to be
 - **proper** if $F \neq L$.
 - **maximal** if F is maximal in $(\mathcal{F}(L) \setminus \{L\}, \subset)$.
 - **prime** if
 1. F is proper
 2. for each $a, b \in L$, $a \vee b \in F$ implies that $a \in F$ or $b \in F$.
 - **principal** if there exists $a \in L$ such that $F = \uparrow a$.

Exercise 10.4.3.2. Let L be a lattice.

1. For each $I \in \mathcal{I}(L)$.
 - (a) if I is proper, then for each $x \in I$, x is not a one of L .
 - (b) if I is principal, then I is proper iff for each $x \in I$, x is not a one of L .
2. For each $F \in \mathcal{F}(L)$.
 - (a) If F is proper, then for each $x \in F$, x is not a zero of L .
 - (b) If F is principal, then F is proper iff for each $x \in F$, x is not a zero of L .

Proof.

1. Let $I \in \mathcal{I}(L)$.
 - (a) Suppose that I is proper. Let $x \in I$. For the sake of contradiction, suppose that x is a one of L . (previous ex) then implies that $(L, \leq) = \{x\}$. Let $y \in L$. Then $y \leq x$. Since I is a lower set, $x \in I$ and $y \leq x$, we have that $y \in I$. Since $y \in L$ is arbitrary, we have that for each $y \in L$, $y \in I$. Thus

$$\begin{aligned} L &\subset I \\ &\subset L. \end{aligned}$$

Thus $I = L$ and I is not proper. Therefore x is not a one of L . Since $x \in I$ is arbitrary, we have that for each $x \in I$, x is not a one of L .

- (b) Suppose that I is principal.
 - ($\implies$):
If I is proper, then 1(a) implies that for each $x \in I$, x is not a one of L .

- ($\Leftarrow$):

Suppose that I is not proper. Then $I = L$. Since I is principal, there exists $a \in I$ such that $I = \downarrow a$. Let $x \in L$. Since

$$\begin{aligned} x &\in L \\ &= I \\ &= \downarrow a, \end{aligned}$$

we have that $x \leq a$. Hence

$$\begin{aligned} x \wedge a &= \inf\{x, a\} \\ &= x. \end{aligned}$$

Since $x \in L$ is arbitrary, we have that for each $x \in L$, $x \wedge a = x$. Hence a is a one of L . Therefore I is not proper implies that there exists $a \in I$ such that a is a one of L . By contrapositive, we have that if for each $x \in I$, x is not a one of L , then I is proper.

2. FINISH!!

□

Exercise 10.4.3.3. Let L be a lattice and $x \in L$.

1. Then $\downarrow x \in \mathcal{I}_p(L)$ iff $x \in \text{prime}_\wedge(L)$.
2. Then $\uparrow x \in \mathcal{F}_p(L)$ iff $x \in \text{prime}_\vee(L)$.

Proof.

1. • ($\Rightarrow$):

Suppose that $\downarrow x \in \mathcal{I}_p(L)$. Then $\downarrow x$ is proper and for each $a, b \in L$, $a \wedge b \in \downarrow x$ implies that $a \in \downarrow x$ or $b \in \downarrow x$.

- Since $\downarrow x$ is proper, **previous exercise** implies that x is not a one of L .
- Let $a, b \in L$. Suppose that $a \wedge b \leq x$. Since $\downarrow x$ is a lower set, $x \in \downarrow x$ and $a \wedge b \leq x$, we have that $a \wedge b \in \downarrow x$. Therefore $a \in \downarrow x$ or $b \in \downarrow x$. Then $a \leq x$ or $b \leq x$. Since $a, b \in L$ with $a \wedge b \leq x$ are arbitrary, we have that for each $a, b \in L$, $a \wedge b \leq x$ implies that $a \leq x$ or $b \leq x$.

Thus $x \in \text{prime}_\wedge(L)$.

- ($\Leftarrow$):

Suppose that $x \in \text{prime}_\wedge(L)$.

- For the sake of contradiction, suppose that $\downarrow x$ is not proper. Then $\downarrow x = L$. Thus $\text{ub } L = \{x\}$. **A previous ex** implies that x is a one of L . This is a contradiction since $x \in \text{prime}_\wedge(L)$. Therefore $\downarrow x$ is proper.
- Let $a, b \in L$. Suppose that $a \wedge b \in \downarrow x$. Then $a \wedge b \leq x$. Since $x \in \text{prime}_\wedge(L)$, $a \leq x$ or $b \leq x$. Hence $a \in \downarrow x$ or $b \in \downarrow x$. Since $a, b \in L$ with $a \wedge b \in \downarrow x$ are arbitrary, we have that for each $a, b \in L$, $a \wedge b \in \downarrow x$ implies that $a \in \downarrow x$ or $b \in \downarrow x$.

Therefore $\downarrow x \in \mathcal{I}_p(L)$.

2. Similar to (1) **FIX/CHECK/FINISH!!!**

□

Exercise 10.4.3.4. Let L be a lattice.

1. For each $I \in \mathcal{I}(L)$, $I \in \mathcal{I}_p(L)$ iff $I^c \in \mathcal{F}(L)$.
2. For each $F \in \mathcal{F}(L)$, $F \in \mathcal{F}_p(L)$ iff $F^c \in \mathcal{I}(L)$.

Proof.

1. Let $I \in \mathcal{I}(L)$.

- ($\Rightarrow$):

Suppose that $I \in \mathcal{I}_p(L)$. Then I is proper and for each $a, b \in L$, $a \wedge b \in I$ implies that $a \in I$ or $b \in I$.

(a) Since I is proper, $I \neq L$. Therefore $I^c \neq \emptyset$

(b) Let $a, b \in I^c$. For the sake of contradiction, suppose that $a \wedge b \in I$. Since $I \in \mathcal{I}_p(L)$, $a \in I$ or $b \in I$. This is a contradiction since $a \in I^c$ and $b \in I^c$. Therefore $a \wedge b \notin I$. Hence $a \wedge b \in I^c$. Since $a, b \in I^c$ are arbitrary, we have that for each $a, b \in I^c$, $a \wedge b \in I^c$.

(c) Since $I \in \mathcal{I}(L)$, a previous exercise implies that I is a lower set. an exercise in proset section implies that I^c is an upper set. Let $x \in L$ and $a \in I^c$. Since

$$\begin{aligned} x \vee a &\geq a \\ &\in I^c \end{aligned}$$

and I^c is an upper set, we have that $x \vee a \in I^c$. Since $x \in L$ and $a \in I^c$ are arbitrary, we have that for each $x \in L$ and $a \in I^c$, $x \vee a \in I^c$.¹

Thus $I^c \in \mathcal{F}(L)$.

- ($\Leftarrow$):

Suppose that $I^c \in \mathcal{F}(L)$.

(a) Since $I^c \in \mathcal{F}(L)$, $I^c \neq \emptyset$. Then $I \neq L$ and therefore I is proper.

(b) Let $a, b \in L$. Suppose that $a \wedge b \in I$. For the sake of contradiction, suppose that $a \notin I$ and $b \notin I$. Then $a \in I^c$ and $b \in I^c$. Since $I^c \in \mathcal{F}(L)$, we have that $a \wedge b \in I^c$. This is a contradiction since $a \wedge b \in I$. Hence $a \in I$ or $b \in I$. Since $a, b \in L$ with $a \wedge b \in I$ are arbitrary, we have that for each $a, b \in L$, $a \wedge b \in I$ implies that $a \in I$ or $b \in I$.

Thus $I \in \mathcal{I}_p(L)$.

2. use duality.

□

10.5 Contructions with Lattices

10.5.1 Sublattices

Definition 10.5.1.1. Let $(L, \vee_L, \wedge_L)$ and $(M, \vee_M, \wedge_M)$ be lattices. Then $(M, \vee_M, \wedge_M)$ is said to be a **sublattice of** $(L, \vee_L, \wedge_L)$ if

1. $M \subset L$
2. ι_M is a $((\vee_L, \wedge_L), (\vee_M, \wedge_M))$ -homomorphism.

Exercise 10.5.1.2. Let $(L, \vee_L, \wedge_L)$ and $(M, \vee_M, \wedge_M)$ be lattices. Then $(M, \vee_M, \wedge_M)$ is a sublattice of L iff

1. $M \subset L$
2. $\vee_L|_{M^2} = \vee_M$ and $\wedge_L|_{M^2} = \wedge_M$.

Proof.

- ($\Rightarrow$):

Suppose that $(M, \vee_M, \wedge_M)$ is a sublattice of L .

1. By definition, $M \subset L$.
2. By definition, ι_M is a $((\vee_L, \wedge_L), (\vee_M, \wedge_M))$ -homomorphism. Then for each $a, b \in M$,

$$\begin{aligned} \vee_M(a, b) &= a \vee_M b \\ &= \iota_M(a \vee_M b) \\ &= \iota_M(a) \vee_L \iota_M(b) \\ &= a \vee_L b \\ &= \vee_L|_{M^2}(a, b). \end{aligned}$$

Hence $\vee_L|_{M^2} = \vee_M$. Similarly, $\wedge_L|_{M^2} = \wedge_M$.

- ($\Leftarrow$):

Suppose that

1. $M \subset L$
2. $\vee_L|_{M^2} = \vee_M$ and $\wedge_L|_{M^2} = \wedge_M$.

1. By assumption $M \subset L$.
2. Since $\vee_L|_{M^2} = \vee_M$, we have that for each $a, b \in M$,

$$\begin{aligned}\iota_M(a \vee_M b) &= a \vee_M b \\ &= \vee_M(a, b) \\ &= \vee_L|_{M^2}(a, b) \\ &= \vee_L(a, b) \\ &= a \vee_L b.\end{aligned}$$

Hence ι_M is a $(\vee_M, \vee_L)$ -homomorphism. Similarly, ι_M is a $(\wedge_M, \wedge_L)$ -homomorphism. Therefore ι_M is a $((\vee_L, \wedge_L), (\vee_M, \wedge_M))$ -homomorphism.

Thus $(M, \vee_M, \wedge_M)$ is a sublattice of L .

FINISH!!!!

□

need to make preliminary ex showing that if $M \subset L$, $M \neq \emptyset$ and $\vee(M^2), \wedge(M^2) \subset M$, then $(M, \vee|_{M^2}, \wedge|_{M^2})$ is a lattice

Exercise 10.5.1.3. Let $(L, \vee, \wedge)$ be a lattice and $M \subset L$. Then $(M, \vee|_{M^2}, \wedge|_{M^2})$ is a sublattice of $(L, \vee, \wedge)$ iff

1. $M \neq \emptyset$
2. $\vee(M^2), \wedge(M^2) \subset M$.

Proof. Set $\vee_M := \vee|_{M^2}$ and $\wedge_M := \wedge|_{M^2}$

- ($\Rightarrow$):

Suppose that $(M, \vee_M, \wedge_M)$ is a sublattice of $(L, \vee, \wedge)$. Since $(M, \vee_M, \wedge_M)$ is a sublattice of $(L, \vee, \wedge)$, we have that $(M, \vee_M, \wedge_M)$ is a lattice.

1. Since $(M, \vee_M, \wedge_M)$ is a lattice, $M \neq \emptyset$.
2. Since $(M, \vee_M, \wedge_M)$ is a lattice, $\vee_M(M^2), \wedge_M(M^2) \subset M$.

- ($\Leftarrow$):

Suppose that

1. $M \neq \emptyset$
2. $\vee(M^2), \wedge(M^2) \subset M$.

1. By assumption $M \neq \emptyset$,
2. Since $\vee, \wedge$ are semilattice operators, $\vee, \wedge$ satisfy the lattice absorption identities and $\vee(M^2), \wedge(M^2) \subset M$, we have that $\vee_M, \wedge_M$ are semilattice operators and $\vee|_M, \wedge|_M$ satisfy the lattice absorption identities.

Thus $(M, \vee_M, \wedge_M)$ is a lattice. By assumption, $M \subset L$ and $\vee_M = \vee|_{M^2}$, $\wedge_M = \wedge|_{M^2}$, we have that $(M, \vee_M, \wedge_M)$ is a sublattice of $(L, \vee, \wedge)$.

□

10.5.2 Product Lattices

10.5.3 Coproduct Lattices

10.5.4 Quotient Lattices

Chapter 11

Complete Lattices

maybe move complete lattices to order chapter? two choices: a) cover complete lattices after order and before other structures

(including lattices?) so that we can define the 'subgroup/ring/lattice generated by a subset' or a 'subquiver generated by a sub set of vertices/edges' using the ideas of galois connection between two posets, one of which has extra structure b) cover complete lattices

after algebraic structures with lattices and note that we can capture all the subobjects with the notion of galois connection between complete lattices and give groups, rings etc as fundamental examples of complete lattices.

idk which is best option, it might make more sense to keep complete lattices after lattices, since we can talk about completely

prime ideals and other lattice-type objects, and we can give fundamental examples of complete lattices that were discussed prior like groups/rings/lattice ideals/etc

11.1 Introduction

Exercise 11.1.0.1. Let $(X, \leq)$ be a poset. Then for each $A \subset X$, $\sup A$ exists iff for each $A \subset X$, $\inf A$ exists.

Proof.

- $(\implies)$:
Suppose that for each $A \subset X$, $\sup A$ exists. Let $A \subset X$. We note that

$$\sup(\text{lb } A) \text{ exists} \iff \inf A \text{ exists} \quad (\text{by Exercise 9.3.0.2}).$$

Since $\text{lb } A \subset X$, then by assumption, $\sup(\text{lb } A)$ exists. Therefore $\inf A$ exists. Since $A \subset X$ is arbitrary, we have that for each $A \subset X$, $\inf A$ exists.

- $(\impliedby)$:
Similar to $(\implies)$.

□

Definition 11.1.0.2. Let $(X, \leq)$ be a poset.

- Then $(X, \leq)$ is said to be a **complete lattice** if for each $A \subset X$, $\sup A$ exists.
- If $(X, \leq)$ is a complete lattice, we define the **arbitrary join and meet operators on X** , denoted $\bigvee_{\leq} : \mathcal{P}(X) \rightarrow X$ and $\bigwedge_{\leq} : \mathcal{P}(X) \rightarrow X$ respectively, by $\bigvee_{\leq} A := \sup(A, \leq)$ and $\bigwedge_{\leq} A := \inf(A, \leq)$.

Note 11.1.0.3. When the context is clear, we may write $\bigvee$ and $\bigwedge$ in place of $\bigvee_{\leq}$ and $\bigwedge_{\leq}$ respectively.

Exercise 11.1.0.4. Let $(X, \leq)$ be a complete lattice. Then $(X, \vee_{\leq}, \wedge_{\leq}, \inf(L, \leq), \sup(L, \leq),)$ is a bounded lattice.

Proof. **FIX!!!**

1. **make previous exercise** implies that $\text{lb}(\emptyset, \leq) = X$. Since $(X, \leq)$ is a complete lattice and $\emptyset \subset X$, there exists $x_1 \in X$ such that $x_1 := \inf \emptyset$. In particular, $X \neq \emptyset$.

2.
 - **make previous ex saying $\inf \emptyset = \sup X$ and $\sup \emptyset = \inf X$ with the following logic**

$$\begin{aligned} \inf \emptyset &\in \text{lb } \emptyset \cap \text{ub}(\text{lb } \emptyset) \\ &= X \cap \text{lb}(X) \\ &= \text{lb } X. \end{aligned}$$

ref an ex implying that $\text{lb } X = \inf X$. Hence X is a one of X .

- similar x_0

Clearly $\bigvee X$ is a one of X and $\bigwedge X$ is a zero of X . □

11.2 Complete Lattice Homomorphisms

Definition 11.2.0.1. Let $(X, \leq_X), (Y, \leq_Y)$ be complete lattices and $f : X \rightarrow Y$. Then f is said to

1. **$(\leq_X, \leq_Y)$ -arbitrary join preserving** if for each $A \subset X$, $f(\sup(A, \leq_X)) = \sup(f(A), \leq_Y)$
2. **$(\leq_X, \leq_Y)$ -arbitrary meet preserving** if for each $A \subset X$, $f(\inf(A, \leq_X)) = \inf(f(A), \leq_Y)$

Exercise 11.2.0.2. Let $(X, \leq_X), (Y, \leq_Y)$ be complete lattices and $f : X \rightarrow Y$. Then

1. f is $(\leq_X, \leq_Y)$ -arbitrary join preserving iff f is $(\leq_X^*, \leq_Y^*)$ -arbitrary meet preserving.
2. f is $(\leq_X, \leq_Y)$ -arbitrary meet preserving iff f is $(\leq_X^*, \leq_Y^*)$ -arbitrary join preserving.

Proof.

1.
 - **$(\implies)$:**
Suppose that f is $(\leq_X, \leq_Y)$ -arbitrary join preserving. Then **a prev ex in poset section on suprema** implies that for each $A \subset X$,

$$\begin{aligned} f(\inf(A, \leq_X^*)) &= f(\sup(A, \leq_X)) \\ &= \sup(f(A), \leq_Y) \\ &= \inf(f(A), \leq_Y^*). \end{aligned}$$

Hence f is $(\leq_X^*, \leq_Y^*)$ -arbitrary meet preserving.

- **$(\impliedby)$:**
Similar to $(\implies)$. **ref.**

2. Similar to (1). **ref.** □

Definition 11.2.0.3. RENAME FINISH!!!! , maybe call right/left subinverse? Let $(X, \leq_X), (Y, \leq_Y)$ be complete lattices, $f : X \rightarrow Y$ and $g : Y \rightarrow X$. We define the

- **right $(\leq_X, \leq_Y)$ -quasiadjoint of f** , denoted $\text{rad}_{\leq_X}^{\leq_Y} f : Y \rightarrow X$, by $[\text{rad}_{\leq_X}^{\leq_Y} f](y) := \sup(f^{-1}(\downarrow(y, \leq_Y))), \leq_X$.
- **left $(\leq_X, \leq_Y)$ -quasiadjoint of g** , denoted $\text{lad}_{\leq_X}^{\leq_Y} g : X \rightarrow Y$, by $[\text{lad}_{\leq_X}^{\leq_Y} g](x) := \inf(f^{-1}(\uparrow(x, \leq_X))), \leq_Y$.

Exercise 11.2.0.4. Let $(X, \leq_X), (Y, \leq_Y)$ be complete lattices, $f : X \rightarrow Y$ and $g : Y \rightarrow X$. Define $r_f : Y \rightarrow X$ and $l_g : X \rightarrow Y$, by $r_f(y) := \sup f^{-1}(\downarrow y, \leq_X)$

1. Then

- (a) r_f is monotonic
- (b) l_g is monotonic

- 2. (a) f is arbitrary join preserving iff f has a right adjoint and $\text{rad } f = r_f$.
- (b) g is arbitrary meet preserving iff g has a left adjoint $\text{lad } g = l_g$.

Proof.

- 1. (a) For each $y_1, y_2 \in Y$,

$$\begin{aligned}
 y_1 \leq_Y y_2 &\implies \downarrow y_1 \subset \downarrow y_2 \\
 &\implies f^{-1}(\downarrow y_1) \subset f^{-1}(\downarrow y_2) \\
 &\implies \sup f^{-1}(\downarrow y_1) \leq_X \sup f^{-1}(\downarrow y_2) \quad (\text{by prev ex}) \\
 &\implies r_f(y_1) \leq_X r_f(y_2).
 \end{aligned}$$

Hence r_f is $(\leq_Y, \leq_X)$ -monotone.

- (b) Similar to prev part.

- 2. (a) • ($\implies$):

Suppose that f is arbitrary join preserving. an ex in subsection on semilattice homomorphisms Exercise ?? implies that f is monotone. The prev ex implies that r_f is monotone.

- For each $x \in X$,

$$\begin{aligned}
 f(x) \leq_Y f(x) &\implies f(x) \in \downarrow f(x) \\
 &\implies x \in f^{-1}(\downarrow f(x)) \\
 &\implies x \leq_X \sup f^{-1}(\downarrow f(x)) \\
 &\implies x \leq_X [r_f \circ f](x) \\
 &\implies \text{id}_X(x) \leq_X [r_f \circ f](x).
 \end{aligned}$$

Hence $\text{id}_X \leq_X r_f \circ f$.

- For each $y \in Y$,

$$\begin{aligned}
 [f \circ r_f](y) &= f(\sup f^{-1}(\downarrow y)) \\
 &= \sup f(f^{-1}(\downarrow y)) \quad (\text{since } f \text{ is arbitrary join preserving}) \\
 &\leq_Y \sup \downarrow y \quad (\text{by ex on image of preimage and ex in section on suprema and subsets}) \\
 &= y \quad (\text{by ex in section on suprema}) \\
 &= \text{id}_Y(y).
 \end{aligned}$$

Hence $f \circ r_f \leq_Y \text{id}_Y$.

Exercise 7.2.0.6 then implies that (f, r_f) is a monotone Galois connection. Thus f has a right adjoint and $\text{rad } f = r_f$.

- ($\impliedby$):

Suppose that f has a right adjoint and $\text{rad } f = r_f$. Then $(f, \text{rad } f)$ is a monotone Galois connection. Exercise 9.3.0.9 implies that for each $A \subset X$, $f(\sup A) = \sup f(A)$. Hence f is arbitrary join preserving.

- (b) Similar to the prev part FINISH!!!, rewrite

□

Comment on how the prev ex connects to probability theory in that the cdf of a random variable is a right continuous increasing function and therefore preserves infima. Its associated quantile function is actually its left adjoint whose most general definition is given above.

Exercise 11.2.0.5. Let $(X, \leq_X), (Y, \leq_Y)$ be complete lattices, $f : X \rightarrow Y$ and $g : Y \rightarrow X$. Suppose that (f, g) is a monotone Galois connection.

1. (a)
(b)
2. (a) $g(1_Y) = 1_X$ iff
(b)

Exercise 11.2.0.6. show that if g is a right adjoint, then $g(\inf \uparrow x) = \inf g(\uparrow x) = \inf \uparrow(g(x)) = g(x)$

Exercise 11.2.0.7.

try to capture idea of a forgetful functor in terms of posets. If we go from groups to sets, we 'lose information' a group with a relations (i.e. a quotient group) is somehow 'more informative' than the free group. So if I send a set to a group in the least informative way (i.e. in a way which preserves as much entropy as possible) then we should get the free group, anything else would be more informative, with a loss of entropy. So quotient objects of a free object can maybe be characterized by some loss of entropy.

For example in **Set**, $A \subset B$ (i.e. A is a subobject of B) is equivalent to saying that A is a quotient of B in the opposite category, which is equivalent to the complete atomic boolean algebras, or the power set, so that if $\iota : A \rightarrow B$ is a monomorphism in **Set**, then $\iota^{-1} : \mathcal{P}(B) \rightarrow \mathcal{P}(A)$ is an epimorphism in complete atomic boolean lattices and we think of $\mathcal{P}(A)$ as a quotient of $\mathcal{P}(B)$. So in this sense $\mathcal{P}(A)$ has less information or is less informative than $\mathcal{P}(B)$ (i.e. A is less informative than B in **Set**^{op}.)

If (X, x) is a set X with some structure x , and (A, a) is in the same category $\mathcal{C}$ (which we order by information level) and we can think of (A, a) as a quotient of (X, x) , so it is less informative as a structured object $(A, a) \leq_{\text{info}\mathcal{C}} (X, x)$, then if we change the context (apply a covariant functor) to land in Set^{op} and think of A quotient of X , (i.e. A is subset of X in **Set**), so that $A \leq_{\text{info}\text{Set}^{\text{op}}} B$. Then we would like an adjoint of F in the sense of a poset so that we obtain the other half a monotone galois connection $G : \text{Set}^{\text{op}} \rightarrow \mathcal{C}$. The idea here is that the order $\leq_{\text{info}\text{Set}^{\text{op}}}$ on Set^{op} is a part of $\leq_{\text{info}\mathcal{C}}$, that is $\leq_{\text{info}\mathcal{C}}$ is $\leq_{\text{info}\text{Set}^{\text{op}}}$ plus some extra order information

i.e. a restriction in some sense so $(A, a) \leq_{\mathcal{C}} (B, b)$ (i.e. there is a quotient from B to A , like free group to abelianization) implies $A \leq_{\text{Set}^{\text{op}}} B$. So then that $\leq_{\mathcal{C}} \subset \leq_{\text{Set}^{\text{op}}}$ (in some sense), so that $\text{ub}((A, a), \leq_{\mathcal{C}}) \subset \text{ub}(A, \leq_{\text{Set}^{\text{op}}})$ in some sense as sets, so that $\inf(\text{ub}(A, \leq_{\text{Set}^{\text{op}}}) \leq_{\text{Set}^{\text{op}}} \inf(\text{ub}((A, a), \leq_{\mathcal{C}}))) = A$. Then if we have a right adjoint $G : (\text{Set}^{\text{op}}, \leq_{\text{Set}^{\text{op}}}) \rightarrow (\mathcal{C}, \leq_{\mathcal{C}})$, which preserves meets and is monotone, we would have that (considering $\leq_{\mathcal{C}}$ as a subset of $\leq_{\text{Set}^{\text{op}}}$)

$$\begin{aligned} G(A) &= G(\inf \text{ub}(A, \leq_{\mathcal{C}}), \leq_{\mathcal{C}}) \\ &= \inf(G(\text{ub}(A, \leq_{\mathcal{C}}), \leq_{\mathcal{C}})) \\ &\geq \inf(G(\text{ub}(A, \leq_{\text{Set}^{\text{op}}}), \leq_{\mathcal{C}})) \end{aligned}$$

idk, I think we can show this aligns with the (least informative object (B, b) in $\mathcal{C}$ which is "above" A , i.e. the structured object (B, b) generated by A), i.e. the structured object "above" A with the least entropy gained from the minimum or zero structured element in the poset.

Try to show that a disintegration of measure, i.e. a conditional probability is the kind of the opposite or dual of this, i.e. it is the highest structured object (markov kernel + data) sitting below an unstructured object (markov kernel of a measure), so that going from (kernel + data) to disintegration results in the least entropy lost, maybe this is related to kl divergence as a measure of entropy loss, see baez/leinster/fritz paper.

In the case of statistics and say maximum likelihood estimation, and kl divergence as loss of estimate $\hat{\theta}(x_j : j \in \mathbb{N}) : (X, P_{\theta}) \rightarrow (X, P_{\hat{\theta}})$, there should be a categorical way of rephrasing so that the observed data $(x_j)_{j \in \mathbb{N}} \subset X$ could perhaps be viewed as maps $x_j : (\Lambda, \mu) \rightarrow (X, P_{\theta})$, so that $\hat{\theta} : \left[\prod_{j \in \mathbb{N}} \text{Hom}((\Lambda, \mu), (X, P_{\theta})) \right] \times (X, P_{\theta}) \rightarrow (X, P_{\hat{\theta}})$

We should be able to do the same in a different context, say the true group $(G, \cdot)$ is unknown, but we observe the elements of G and the fact that for each $a, b \in G$, $a \cdot b = b \cdot a$ (i.e. we are given the proposition that G is commutative), then what would be the best estimate for G under this information in the sense of minimizing info loss (i.e. maybe occams razor?)? Well it would probably be the free abelian group. Any further processed abelian group, (i.e. a quotient of the free abelian group) would lose more info, so even though the trivial group also satisfies the observation, it is suboptimal because we lose basically all info about G . Need a way to quantify info loss in this way. What would the analogouse 'data' be? here it seems we need a way to view relations about elements as a morphism into G from some other object, like in the case of statistics.

Vague idea: in two different contexts, say two different posets, we want to relate two objects in the ordering induced by $a \leq b$ iff b is a quotient of a , i.e. b is more informative, i.e. there is an arrow from a to b which exhibits entropy loss.

talk about the idea of the element in a complete lattice with more structure generated another element in a complete lattice with less structure. i.e. subgroup generated subset where subgroups ordered by inclusion, power set ordered by inclusion. Or topology generated by collection of subsets (i.e. subset of powerset) where topologies ordered by inclusion and double power set ordered by inclusion. In these cases, both posets are complete lattices since arbitrary meets exist. If we consider the map g from the more structured space P_2 to the less structured space P_1 , by sending an object to itself minus the structure, so a group to its underlying set, or a topology to itself considered as a subset of the power set, then meets are preserved. Thus g is a right adjoint to some left adjoint f from the less structured space P_1 to the more structured space P_2 , so f preserves arbitrary joins. For an object 1 in the less structured poset, we define the object in the more structured poset generated by $\text{obj}1$ to be the left adjoint of $\text{obj}1$, i.e. $f(\text{obj}1)$. we can find this in many cases by noting that in P_1 , $\text{obj}1$ is the sup of its upper bounds. Since sups are preserved by f , the generated structured object in P_2 is $f(\text{obj}1)$ which (from a prev ex) is the sup of f (of the lower bounds of $\text{obj}1$) i.e. the sup of $f(\downarrow o1)$, which should be the sup of all upper bounds of $f(\text{obj}1)$, which is the inf

ok in P_1 is less structured, P_2 is more structured, $f : P_1 \rightarrow P_2$ preserves joins, adjoint g preserves meets, then the object in P_2 generated by $a \in P_1$ is $f(a) = f(\sup \downarrow a) = \sup f(\downarrow a)$, or something like this

11.3 Properties of Ideals and Filters

11.3.1 Prime Ideals and Filters in Complete Lattices

Definition 11.3.1.1. Let L be a complete lattice.

- Let $I \in \mathcal{I}(L)$. Then I is said to be
 - **completely prime** if
 1. I is proper
 2. for each $A \subset L$, $\bigwedge A \in I$ implies that there exists $a \in A$ such that $a \in I$.
- Let $F \in \mathcal{F}(L)$. Then F is said to be
 - **completely prime** if
 1. F is proper
 2. for each $A \subset L$, $\bigvee A \in F$ implies that there exists $a \in A$ such that $a \in F$.

Exercise 11.3.1.2. Let L be a complete lattice.

1. For each $I \in \mathcal{I}(L)$, if I is completely prime, then $I \in \mathcal{I}_p(L)$.
2. For each $F \in \mathcal{F}(L)$, if F is completely prime, then $F \in \mathcal{F}_p(L)$.

Proof. Clear. fill out

□

Exercise 11.3.1.3. Let L be a complete lattice.

1. For each $I \in \mathcal{I}_{cp}(L)$,
 - (a) $\bigwedge I^c \in I^c$,
 - (b) $\bigwedge I^c \in \text{prime}_\vee(L)$.
2. For each $F \in \mathcal{F}_{cp}(L)$,
 - (a) $\bigvee F^c \in F^c$,
 - (b) $\bigvee F^c \in \text{prime}_\wedge(L)$.

Proof.

1. Let $I \in \mathcal{I}_{cp}(L)$.

- (a) For the sake of contradiction, suppose that $\bigwedge I^c \in I$. Since I is completely prime, there exists $a \in I^c$ such that $a \in I$. This is a contradiction. Hence $\bigwedge I^c \notin I$. Thus $\bigwedge I^c \in I^c$.
- (b) i. Since $I \in \mathcal{I}(L)$, **a previous ex** implies that $0 \in I$. Since $\bigwedge I^c \in I^c$, we have that $\bigwedge I^c \neq 0$.
- ii. Define $x \in L$ by $x := \bigwedge I^c$. Since $I \in \mathcal{I}(L)$, **a prev ex** implies that I is a lower set. Let $a, b \in L$. Suppose that $x \leq a \vee b$. For the sake of contradiction, suppose that $a \in I$ and $b \in I$. Since $I \in \mathcal{I}(L)$, we have that $a \vee b \in I$. Since I is a lower set and

$$\begin{aligned} x &\leq a \vee b \\ &\in I, \end{aligned}$$

we have that $x \in I$. This is a contradiction since $x \in I^c$. Hence $a \notin I$ or $b \notin I$. Thus $a \in I^c$ or $b \in I^c$. Since $x = \bigwedge I^c$, we have that $x \leq a$ or $x \leq b$. Since $a, b \in L$ with $x \leq a \vee b$ are arbitrary, we have that for each $a, b \in L$, $x \leq a \vee b$ implies that $x \leq a$ or $x \leq b$.

Hence $x \in \text{prime}_\vee(L)$.

2. **use duality**

□

Exercise 11.3.1.4. Let L be a complete lattice. Then

1. for each $x \in \text{prime}_\vee(L)$, $(\uparrow x)^c \in \mathcal{I}_{cp}(L)$.
2. for each $x \in \text{prime}_\wedge(L)$, $(\downarrow x)^c \in \mathcal{F}_{cp}(L)$.

Proof.

1. Let $x \in \text{prime}_\vee(L)$. **a prev ex** implies that $\uparrow x \in \mathcal{F}(L)$.

- (a) Since $x \in \text{prime}_\vee(L)$, $x \neq 0$. Hence $0 \in (\uparrow x)^c$. Thus $(\uparrow x)^c \neq \emptyset$.
- (b) Let $a, b \in (\uparrow x)^c$. For the sake of contradiction, suppose that $a \vee b \in \uparrow x$. Then $x \leq a \vee b$. Since $x \in \text{prime}_\vee(L)$, $x \leq a$ or $x \leq b$. Hence $a \in \uparrow x$ or $b \in \uparrow x$. This is a contradiction since $a \in (\uparrow x)^c$ and $b \in (\uparrow x)^c$. Therefore $a \vee b \in (\uparrow x)^c$. Since $a, b \in (\uparrow x)^c$ are arbitrary, we have that for each $a, b \in (\uparrow x)^c$, $a \vee b \in (\uparrow x)^c$.
- (c) Let $y \in L$ and $a \in (\uparrow x)^c$. For the sake of contradiction, suppose that $y \wedge a \in \uparrow x$. Then

$$\begin{aligned} x &\leq y \wedge a \\ &\leq a. \end{aligned}$$

Thus $a \in \uparrow x$, which is a contradiction. Hence $y \wedge a \in (\uparrow x)^c$. Since $y \in L$ and $a \in (\uparrow x)^c$ are arbitrary, we have that for each $y \in L$ and $a \in (\uparrow x)^c$, $y \wedge a \in (\uparrow x)^c$.

Thus $(\uparrow x)^c \in \mathcal{I}(L)$.

- (a) Since $\uparrow x \neq \emptyset$, we have that $(\uparrow x)^c \neq L$. Thus $(\uparrow x)^c$ is proper.
- (b) Let $A \subset L$. Suppose that $\bigwedge A \in (\uparrow x)^c$. For the sake of contradiction, suppose that for each $a \in A$, $a \notin (\uparrow x)^c$. Then for each $a \in A$, $a \in \uparrow x$. Hence for each $a \in A$, $x \leq a$. Therefore $x \in \text{lb } A$. Thus

$$\begin{aligned} x &\leq \inf A \\ &= \bigwedge A. \end{aligned}$$

Then $\bigwedge A \in \uparrow x$, which is a contradiction. Hence there exists $a \in A$ such that $a \in (\uparrow x)^c$. Since $A \subset L$ with $\bigwedge A \in (\uparrow x)^c$ is arbitrary, we have that for each $A \subset L$, $\bigwedge A \in (\uparrow x)^c$ implies that there exists $a \in A$ such that $a \in (\uparrow x)^c$.

Thus $(\uparrow x)^c \in \mathcal{I}_{cp}(L)$.

2. **similar or duality**

□

maybe make ex in earlier section on ideals that $\downarrow x = \downarrow y$ iff $x = y$ and similar for $\uparrow x$. Then in this section, show that for complete lattice L , $\mathcal{F}_{cp}(L) \cong \text{prime}_\wedge(L)$ bijective.

11.4 Constructions with Complete Lattices

11.4.1 Complete Sublattices

Definition 11.4.1.1. **define** complete sublattice Let L, M be lattices

11.4.2 Products of Complete Lattices

11.4.3 Coproducts of Complete Lattices

11.4.4 Quotients of Complete Lattices

11.5 Fundamental Examples

11.5.1 Introduction

Exercise 11.5.1.1. Let L be a complete lattice.

11.5.2 Sets

11.5.3 Equivalence Relations

Proof. Let X be a set. Suppose that $X \neq \emptyset$. Then $(\text{Equiv}, \subset)$ is a complete lattice. □

11.5.4 Ideal and Filter Lattices

Exercise 11.5.4.1. Let L be a lattice. If L has a zero, then

1. $(\mathcal{I}(L), \leq_{\mathcal{I}(L)})$ is a complete lattice.
2. $(\mathcal{F}(L), \leq_{\mathcal{F}(L)})$ is a complete lattice.

11.5.5 Topological Spaces

11.6 Least Upper Bound Property

Definition 11.6.0.1. Let $(X, \leq)$ be a poset. Then $(X, \leq)$ is said to satisfy the

- **least upper bound (LUB) property** if for each $A \subset X$, if $A \neq \emptyset$ and A is bounded above, then there exists $x \in X$ such that $x = \sup A$
- **greatest lower bound (GLB) property** if for each $A \subset X$, if $A \neq \emptyset$ and A is bounded below, then there exists $x \in X$ such that $x = \inf A$.

Exercise 11.6.0.2. LUB iff GLB

Proof. **FINISH!!!!** □

Definition 11.6.0.3. Suplattice:

Let $(L, \leq)$ be a poset. Then $(L, \leq)$ is said to be a **suplattice** if for each $A \subset L$, there exists $x \in L$ such that $x = \sup A$.

11.7 Complete Lattice Homomorphisms

Part IV

Continuous Lattices

pulling from [21]

Chapter 12

Scott

Definition 12.0.0.1.

Bibliography

- [1] Angshuman R Goswami, *Revisiting Fekete's Lemma, Subadditive and Periodic Sequences: AR Goswami*, Results in Mathematics **80** (2025), no. 5, 131.
- [2] Gerald B Folland, *Real analysis: modern techniques and their applications*, John Wiley & Sons, 1999.
- [3] Robert R Phelps, *Lectures on Choquet's theorem*, Springer, 2002.
- [4] Gunnar F Stefánsson, *Pettis integrability*, Transactions of the American Mathematical Society **330** (1992), no. 1, 401–418.
- [5] H.H. and Wolff Schaefer M.P.H., *Topological Vector Spaces*, Graduate Texts in Mathematics, Springer New York, 1999.
- [6] L. and Beckenstein Narici E., *Topological Vector Spaces*, Chapman & Hall/CRC Pure and Applied Mathematics, CRC Press, 2010.
- [7] Einar Hille, *Functional Analysis and Semi-Groups*, Dutt Press, 2007.
- [8] ———, *Methods in classical and functional analysis*, Addison Wesley Publishing, 1972.
- [9] E Zeidler, *Nonlinear Functional Analysis and its Applications I.(Fixed Point Theorems) 1986*, Springer-Verlag New York, Inc.
- [10] Nicholas and Tarieladze Vakhania Vazha and Chobanyan, *Probability distributions on Banach spaces*, Springer Science & Business Media, 2012.
- [11] David Simmons, *Conditional measures and conditional expectation; Rohlin's disintegration theorem*, Vol. 32, Discrete and Continuous Dynamical Systems, 2012.
- [12] Edwin and Savage Hewitt Leonard J, *Symmetric measures on Cartesian products*, Transactions of the American Mathematical Society **80** (1955), no. 2, 470–501.
- [13] John M Lee, *Introduction to Smooth Manifolds*, Springer, 2012.
- [14] ———, *Introduction to Riemannian Manifolds*, Springer, 2018.
- [15] Mark JD Hamilton, *Mathematical Gauge Theory*, Springer, 2017.
- [16] Liviu I Nicolaescu, *Lectures on the Geometry of Manifolds*, World Scientific, 2020.
- [17] Steven Rosenberg, *The Laplacian on a Riemannian manifold: an introduction to analysis on manifolds*, Cambridge University Press, 1997.
- [18] ———, *The Laplacian on a Riemannian manifold: an introduction to analysis on manifolds*, Cambridge University Press, 1997.
- [19] Emmanuel Hebey, *Sobolev Spaces on Riemannian Manifolds*, Springer Berlin, Heidelberg, 1996.
- [20] Brian A and Priestley Davey Hilary A, *Introduction to lattices and order*, Cambridge university press, 2002.
- [21] Gerhard and Hofmann Gierz Karl Heinrich and Keimel, *Continuous lattices and domains*, Vol. 93, Cambridge university press, 2003.