1

Introduction to Algebra

Carson James

Contents

Notation **ix**

Preface **1**

I Sets and Order **3**

1 Relations **5**
1.1 Binary Heterogeneous Relations . . . 5
1.1.1 Introduction . . . 5
1.1.2 Subrelations . . . 9
1.1.3 Products of Relations . . . 9
1.2 n-ary Heterogeneous Relations . . . 11
1.2.1 Introduction . . . 11
1.2.2 Subrelations . . . 14
1.2.3 Products of Relations . . . 15
1.3 n-ary Homogeneous Relations . . . 17
1.3.1 Introduction . . . 17
1.3.2 Subrelations . . . 17
1.3.3 Product Relations . . . 17
1.4 Binary Homogeneous Relations . . . 18
1.4.1 Subrelations . . . 18
1.5 Equivalence Relations . . . 19
1.6 Products . . . 24

2 Ordered Sets **25**
2.1 To Do . . . 25
2.2 Prosets . . . 25
2.2.1 Introduction . . . 25
2.2.2 Subprosets . . . 26
2.2.3 Products . . . 26
2.2.4 Quotients . . . 27
2.3 Posets . . . 28
2.3.1 Introduction . . . 28
2.4 Directed Sets . . . 29
2.4.1 Introduction . . . 29
2.4.2 Subdirected Sets . . . 29
2.4.3 Products of Directed Sets . . . 29
2.5 Monotone Maps . . . 30
2.5.1 Introduction . . . 30
2.5.2 Galois Connections . . . 31
2.6 Upper and Lower Sets . . . 39
2.6.1 Introduction . . . 39

2.6.2 Maximal Elements . . . 48
2.7 Suprema and Infima . . . 48
2.7.1 Upper and Lower Bounds . . . 48
2.7.2 SupSets and InfSets . . . 52
2.7.3 Suprema and Infima . . . 55

II Algebraic Structures 61

3 Semigroups, Monoids and Groups 63
3.1 Semigroups . . . 63
3.2 Monoids . . . 64
3.3 Groups . . . 64
3.3.1 Direct Products . . . 64

4 Rings 67
4.1 Introduction . . . 67

5 Modules 69
5.1 Introduction . . . 69

6 Vector Spaces 73

7 Fields 75

8 Quivers 77
8.1 Introduction . . . 77
8.2 Subquivers . . . 77

9 Lattices 83
9.0.1 To Do . . . 83
9.1 Introduction . . . 83
9.1.1 Skew Semilattices . . . 83
9.1.2 Skew Semilattice Homomorphisms . . . 85
9.1.3 Skew Lattices . . . 85
9.1.4 Semilattices . . . 88
9.1.5 Semilattice Homomorphisms . . . 92
9.1.6 Lattices . . . 93
9.1.7 Lattice Homomorphisms . . . 96
9.2 Fundamental Examples . . . 97
9.2.1 Bounded Lattices . . . 97
9.2.2 Bounded Lattice Homomorphisms . . . 98
9.2.3 Modular and Distributive Lattices . . . 99
9.2.4 Introduction . . . 99
9.2.5 Heyting Algebras . . . 103
9.2.6 Boolean Algebras . . . 105
9.3 Lattice Structures . . . 106
9.3.1 Irreducibility and Primality . . . 106
9.3.2 Ideals and Filters . . . 107
9.3.3 Properties of Ideals and Filters . . . 114
9.4 Contructions with Lattices . . . 116
9.4.1 Sublattices . . . 116
9.4.2 Product Lattices . . . 117
9.4.3 Coproduct Lattices . . . 117
9.4.4 Quotient Lattices . . . 117

10 Complete Lattices **119**
10.1 Introduction . . . 119
10.2 Complete Lattice Homomorphisms . . . 120
10.3 Properties of Ideals and Filters . . . 123
10.3.1 Prime Ideals and Filters in Complete Lattices . . . 123
10.4 Constructions with Complete Lattices . . . 125
10.4.1 Complete Sublattices . . . 125
10.4.2 Products of Complete Lattices . . . 125
10.4.3 Coproducts of Complete Lattices . . . 125
10.4.4 Quotients of Complete Lattices . . . 125
10.5 Fundamental Examples . . . 126
10.5.1 Introduction . . . 126
10.5.2 Sets . . . 126
10.5.3 Equivalence Relations . . . 126
10.5.4 Ideal and Filter Lattices . . . 126
10.5.5 Topological Spaces . . . 126
10.6 Least Upper Bound Property . . . 126
10.6.1 Complete Lattice Homomorphisms . . . 126

11 Some Chapter **127**
11.1 Closure Operators . . . 127

12 Universal Algebra **129**
12.1 Introduction . . . 129
12.2 Subalgebras . . . 131
12.3 Congruences . . . 133

A Summation **135**

B Asymptotic Notation **137**

C Categories **139**

D Vector Spaces **141**
D.1 Introduction . . . 141
D.2 Bases . . . 142
D.3 Multilinear Maps . . . 143
D.4 Tensor Products . . . 144

Notation

$\mathcal{M}_+(X, \mathcal{A})$	finite measures on $(X, \mathcal{A})$
v	velocity

Preface

cc-by-nc-sa

Part I

Sets and Order

Chapter 1

Relations

1.1 Binary Heterogeneous Relations

keep this part here, after section on sets similar to analysis notes, so we can assume arbitrary unions, products, etc, not just unions/products/etc of two sets

1.1.1 Introduction

Definition 1.1.1.1. Let A, B be sets.

- Let $R \subset A \times B$. Then R is said to be a **relation over** A **and** B.
- We define $\mathcal{R}(A, B) := \mathcal{P}(A, B)$

Definition 1.1.1.2. Let A, B, C be sets, $R \in \mathcal{R}(A, B)$, $S \in \mathcal{R}(B, C)$. We define the **composition relation of** S **with** R, denoted $S \circ_{\text{Rel}} R \in \mathcal{R}(A, C)$, by

$$S \circ_{\text{Rel}} R := \{(a, c) \in A \times C : \text{there exists } b \in B \text{ such that } (a, b) \in R \text{ and } (b, c) \in S\}.$$

Note 1.1.1.3. When the context is clear, we write $S \circ R$ in place of $S \circ_{\text{Rel}} R$.

Exercise 1.1.1.4. Let A, B, C, D be sets, R a relation between A and B, S relation between B and C and T relation between C and D. Then $(T \circ S) \circ R = T \circ (S \circ R)$.

Proof.

- Let $(a, d) \in (T \circ S) \circ R$. Then there exists $b \in B$ such that $(a, b) \in R$ and $(b, d) \in T \circ S$. Since $(b, d) \in T \circ S$, there exists $c \in C$ such that $(b, c) \in S$ and $(c, d) \in T$. Since $(a, b) \in R$ and $(b, c) \in S$, $(a, c) \in S \circ R$. Since $(a, c) \in S \circ R$ and $(c, d) \in T$, we have that $(a, d) \in T \circ (S \circ R)$. Since $(a, d) \in (T \circ S) \circ R$ is arbitrary, we have that for each $(a, d) \in (T \circ S) \circ R$, $(a, d) \in T \circ (S \circ R)$. Hence $(T \circ S) \circ R \subset T \circ (S \circ R)$.
- Similarly, $T \circ (S \circ R) \subset (T \circ S) \circ R$.

Since $(T \circ S) \circ R \subset T \circ (S \circ R)$ and $T \circ (S \circ R) \subset (T \circ S) \circ R$, we have that $(T \circ S) \circ R = T \circ (S \circ R)$. □

Definition 1.1.1.5. Let A be a set. We define the **diagonal relation on** A, denoted $\Delta_A \in \mathcal{R}(A, A)$, by

$$\Delta_A := \{(a, a) : a \in A\}.$$

Exercise 1.1.1.6. Let A, B be sets and $R \in \mathcal{R}(A, B)$. Then

1. $R \circ \Delta_A = R$
2. $\Delta_B \circ R = R$

Proof.

1. - Let $(a,b) \in R \circ \Delta_A$. Then there exists $a' \in A$ such that $(a,a') \in \Delta_A$ and $(a',b) \in R$. By definition of Δ_A, $a = a'$. Therefore $(a,b) \in R$. Since $(a,b) \in R \circ \Delta_A$ is arbitrary, we have that for each $(a,b) \in R \circ \Delta_A$, $(a,b) \in R$. Hence $R \circ \Delta_A \subset R$.
   - Let $(a,b) \in R$. Then $a \in A$. Therefore $(a,a) \in \Delta_A$. Since $(a,a) \in \Delta_A$ and $(a,b) \in R$, we have that $(a,b) \in R \circ \Delta_A$. Since $(a,b) \in R$ is arbitrary, we have that for each $(a,b) \in R$, $(a,b) \in R \circ \Delta_A$. Hence $R \circ \Delta_A \subset R$.

   Since $R \circ \Delta_A \subset R$ and $R \circ \Delta_A \subset R$, we have that $R \circ \Delta_A = R$.
2. Similar to (1).

□

Exercise 1.1.1.7. Let A, B, C be sets.

1. Let $R \in \mathcal{R}(A,B)$ and $(S_\gamma)_{\gamma\in\Gamma} \subset \mathcal{R}(B,C)$. Then
   (a) $\left(\bigcup_{\gamma\in\Gamma} S_\gamma\right) \circ R = \bigcup_{\gamma\in\Gamma} (S_\gamma \circ R)$,
   (b) $\left(\bigcap_{\gamma\in\Gamma} S_\gamma\right) \circ R \subset \bigcap_{\gamma\in\Gamma} (S_\gamma \circ R)$
2. Let $(R_\gamma)_{\gamma\in\Gamma} \subset \mathcal{R}(A,B)$ and $S \in \mathcal{R}(B,C)$. Then
   (a) $S \circ \left(\bigcup_{\gamma\in\Gamma} R_\gamma\right) = \bigcup_{\gamma\in\Gamma} (S \circ R_\gamma)$.
   (b) $S \circ \left(\bigcap_{\gamma\in\Gamma} R_\gamma\right) \subset \bigcap_{\gamma\in\Gamma} (S \circ R_\gamma)$.

Proof.

1. (a) We note that

$$\begin{aligned}
(a,c) \in \left(\bigcup_{\gamma\in\Gamma} S_\gamma\right) \circ R &\iff \text{there exists } b \in B \text{ such that } (a,b) \in R \text{ and } (b,c) \in \bigcup_{\gamma\in\Gamma} S_\gamma \\
&\iff \text{there exists } b \in B \text{ and } \gamma_0 \in \Gamma \text{ such that } (a,b) \in R \text{ and } (b,c) \in S_{\gamma_0} \\
&\iff \text{there exists } \gamma_0 \in \Gamma \text{ such that } (a,c) \in S_{\gamma_0} \circ R \\
&\iff (a,c) \in \bigcup_{\gamma\in\Gamma} (S_\gamma \circ R).
\end{aligned}$$

   Hence $\left(\bigcup_{\gamma\in\Gamma} S_\gamma\right) \circ R = \bigcup_{\gamma\in\Gamma} (S_\gamma \circ R)$.

   (b) We note that

$$\begin{aligned}
(a,c) \in \left(\bigcap_{\gamma\in\Gamma} S_\gamma\right) \circ R &\iff \text{there exists } b \in B \text{ such that } (a,b) \in R \text{ and } (b,c) \in \bigcap_{\gamma\in\Gamma} S_\gamma \\
&\iff \text{there exists } b \in B \text{ such that } (a,b) \in R \text{ and for each } \gamma \in \Gamma \text{ and } (b,c) \in S_\gamma \\
&\implies \text{for each } \gamma \in \Gamma, \text{ there exists } b \in B \text{ such that } (a,b) \in R \text{ and } (b,c) \in S_\gamma \\
&\iff \text{for each } \gamma \in \Gamma,\ (a,c) \in S_\gamma \circ R \\
&\iff (a,c) \in \bigcap_{\gamma\in\Gamma} (S_\gamma \circ R).
\end{aligned}$$

   Hence $\left(\bigcap_{\gamma\in\Gamma} S_\gamma\right) \circ R = \bigcap_{\gamma\in\Gamma} (S_\gamma \circ R)$.

2. (a) We note that

$$\begin{aligned}(a,c) \in S \circ \left(\bigcup_{\gamma\in\Gamma} R_\gamma\right) &\iff \text{there exists } b \in B \text{ such that } (a,b) \in \bigcup_{\gamma\in\Gamma} R_\gamma \text{ and } (b,c) \in S \\ &\iff \text{there exists } b \in B \text{ and } \gamma_0 \in \Gamma \text{ such that } (a,b) \in R_{\gamma_0} \text{ and } (b,c) \in S \\ &\iff \text{there exists } \gamma_0 \in \Gamma \text{ such that } (a,c) \in S \circ R_{\gamma_0} \\ &\iff (a,c) \in \bigcup_{\gamma\in\Gamma} (S \circ R_\gamma).\end{aligned}$$

Hence $S \circ \left(\bigcup_{\gamma\in\Gamma} R_\gamma\right) = \bigcup_{\gamma\in\Gamma} (S_\gamma \circ R)$.

(b) We note that

$$\begin{aligned}(a,c) \in S \circ \left(\bigcap_{\gamma\in\Gamma} R_\gamma\right) &\iff \text{there exists } b \in B \text{ such that } (a,b) \in \bigcap_{\gamma\in\Gamma} R_\gamma \text{ and } (b,c) \in S \\ &\iff \text{there exists } b \in B \text{ such that for each } \gamma \in \Gamma,\ (a,b) \in R_\gamma \text{ and } (b,c) \in S \\ &\implies \text{for each } \gamma \in \Gamma, \text{ there exists } b \in B \text{ such that } (a,b) \in R_\gamma \text{ and } (b,c) \in S \\ &\iff \text{for each } \gamma \in \Gamma,\ (a,c) \in S \circ R_\gamma \\ &\iff (a,c) \in \bigcap_{\gamma\in\Gamma} (S \circ R_\gamma).\end{aligned}$$

Hence $S \circ \left(\bigcap_{\gamma\in\Gamma} R_\gamma\right) \subset \bigcap_{\gamma\in\Gamma} (S_\gamma \circ R)$.

□

Exercise 1.1.1.8. Let A, B, C be sets.

1. Let $R_1, R_2 \in \mathcal{R}(A,B)$ and $S \in \mathcal{R}(B,C)$. If $R_1 \subset R_2$, then $S \circ R_1 \subset S \circ R_2$.
2. Let $R \in \mathcal{R}(A,B)$ and $S_1, S_2 \in \mathcal{R}(B,C)$. If $S_1 \subset S_2$, then $S_1 \circ R \subset S_2 \circ R$.
3. Let $R_1, R_2 \in \mathcal{R}(A,B)$ and $S_1, S_2 \in \mathcal{R}(B,C)$. If $R_1 \subset R_2$ and $S_1 \subset S_2$, then $S_1 \circ R_1 \subset S_2 \circ R_2$.

Proof.

1. Suppose that $R_1 \subset R_2$. Then

$$\begin{aligned}(a,c) \in S \circ R_1 &\iff \text{there exists } b \in B \text{ such that } (a,b) \in R_1 \text{ and } (b,c) \in S \\ &\implies \text{there exists } b \in B \text{ such that } (a,b) \in R_2 \text{ and } (b,c) \in S \\ &\iff (a,c) \in S \circ R_2.\end{aligned}$$

Hence $S \circ R_1 \subset S \circ R_2$.

2. Suppose that $S_1 \subset S_2$. Then

$$\begin{aligned}(a,c) \in S_1 \circ R &\iff \text{there exists } b \in B \text{ such that } (a,b) \in R \text{ and } (b,c) \in S_1 \\ &\implies \text{there exists } b \in B \text{ such that } (a,b) \in R \text{ and } (b,c) \in S_2 \\ &\iff (a,c) \in S_2 \circ R.\end{aligned}$$

Hence $S_2 \circ R \subset S_1 \circ R$.

3. Suppose that $R_1 \subset R_2$ and $S_1 \subset S_2$. Then the previous parts imply that

$$\begin{aligned}S_1 \circ R_1 &\subset S_2 \circ R_1 \\ &\subset S_2 \circ R_2.\end{aligned}$$

□

Definition 1.1.1.9. Let A, B be sets and $R \in \mathcal{R}(A, B)$. We define the **converse of** R, denoted $R^* \in \mathcal{R}(B, A)$, by

$$R^* := \{(b, a) \in B \times A : (a, b) \in R\}.$$

Exercise 1.1.1.10. Let A, B be sets and R a relation between A and B. Then $(R^*)^* = R$.

Proof.

$$\begin{aligned} (a, b) \in (R^*)^* &\iff (b, a) \in R^* \\ &\iff (a, b) \in R. \end{aligned}$$

Hence $(R^*)^* = R$. □

Exercise 1.1.1.11. Let A, B, C be sets, $R \in \mathcal{R}(A, B)$ and $S \in \mathcal{R}(B, C)$. Then $(S \circ R)^* = R^* \circ S^*$.

Proof. We note that

$$\begin{aligned} (c, a) \in (S \circ R)^* &\iff (a, c) \in S \circ R \\ &\iff \text{there exists } b \in B \text{ such that } (a, b) \in R \text{ and } (b, c) \in S \\ &\iff \text{there exists } b \in B \text{ such that } (c, b) \in S^* \text{ and } (b, a) \in R^* \\ &\iff (c, a) \in R^* \circ S^*. \end{aligned}$$

Hence $(S \circ R)^* = R^* \circ S^*$. □

Exercise 1.1.1.12. Let A and B be sets and $(R_\gamma)_{\gamma \in \Gamma} \subset \mathcal{R}(A, B)$. Then

1. $\left(\bigcup_{\gamma \in \Gamma} R_\gamma \right)^* = \bigcup_{\gamma \in \Gamma} R_\gamma^*$
2. $\left(\bigcap_{\gamma \in \Gamma} R_\gamma \right)^* = \bigcap_{\gamma \in \Gamma} R_\gamma^*$

Proof.

1. We note that

$$\begin{aligned} (b, a) \in \left(\bigcup_{\gamma \in \Gamma} R_\gamma \right)^* &\iff (a, b) \in \bigcup_{\gamma \in \Gamma} R_\gamma \\ &\iff \text{there exists } \gamma_0 \in \Gamma \text{ such that } (a, b) \in R_{\gamma_0} \\ &\iff \text{there exists } \gamma_0 \in \Gamma \text{ such that } (b, a) \in R_{\gamma_0}^* \\ &\iff (b, a) \in \bigcup_{\gamma \in \Gamma} R_\gamma^*. \end{aligned}$$

   Hence $\left(\bigcup_{\gamma \in \Gamma} R_\gamma \right)^* = \bigcup_{\gamma \in \Gamma} R_\gamma^*$.

2. We note that

$$\begin{aligned} (b, a) \in \left(\bigcap_{\gamma \in \Gamma} R_\gamma \right)^* &\iff (a, b) \in \bigcap_{\gamma \in \Gamma} R_\gamma \\ &\iff \text{for each } \gamma \in \Gamma,\ (a, b) \in R_\gamma \\ &\iff \text{for each } \gamma \in \Gamma,\ (b, a) \in R_\gamma^* \\ &\iff (b, a) \in \bigcap_{\gamma \in \Gamma} R_\gamma^*. \end{aligned}$$

   Hence $\left(\bigcap_{\gamma \in \Gamma} R_\gamma \right)^* = \bigcap_{\gamma \in \Gamma} R_\gamma^*$.

□

1.1.2 Subrelations

Definition 1.1.2.1. Let $n \in \mathbb{N}$, A, B, E, F sets, $R \in \mathcal{R}(A, B)$ and $S \in \mathcal{R}(E, F)$. Then

- S is said to be a **restriction of** R if $E \subset A$ and $F \subset B$,
- R is said to be an **extension of** S if S is a restriction of R.

Definition 1.1.2.2. Let A, B be sets, $E \subset A$, $F \subset B$ and $R \in \mathcal{R}(A, B)$. We define the **restriction of** R **over** E **and** F, denoted $R|_{E\times F} \in \mathcal{R}(E, F)$, by $R|_{E\times F} := R \cap (E \times F)$.

Exercise 1.1.2.3. Let A, B be sets, $E \subset A$, $F \subset B$ and $R \in \mathcal{R}(A, B)$. Then $R|_B$ is a restriction of R.

Proof. Clear. fill out later □

maybe dont do this subsection, since relations are like morphisms, so unless you can find a categorical way to present restrictions on domain or codomain $f|_E$ or $f|^F$, just get rid of this subsection

Exercise 1.1.2.4. Let A, B be sets, $E \subset A$, $F \subset B$ and $R \in \mathcal{R}(A, B)$. Suppose that $E \subset A$ and $F \subset B$. Then $(R^*)|_{F\times E} = (R|_{E\times F})^*$.

Proof. We note that for each $b, a \in B \times A$,

$$\begin{aligned}(b, a) \in (R^*)|_{F\times E} &\iff (b, a) \in R^* \cap (F \times E)\\ &\iff (a, b) \in R \cap (E \times F)\\ &\iff (a, b) \in R|_{E\times F}.\end{aligned}$$

Hence $(R^*)|_{F\times E} = R|_{E\times F}$. □

1.1.3 Products of Relations

Definition 1.1.3.1. Let A, B, C, D be sets and $R \in \mathcal{R}(A, B), S \in \mathcal{R}(C, D)$. We define the **product relation of** R **and** S, denoted $R \times_{\text{Rel}} S \in \mathcal{R}(A \times C, B \times D)$, by

$$R \times_{\text{Rel}} S := \{((a, c), (b, d)) \in (A \times C) \times (B \times D) : (a, b) \in R \text{ and } (c, d) \in S\}.$$

Note 1.1.3.2. When the context is clear, we write $R \times S$ in place of $R \times_{\text{Rel}} S$.

Exercise 1.1.3.3. Let A, B, C, D be sets.

1. Let $R \in \mathcal{R}(A, B)$ and $(S_\gamma)_{\gamma\in\Gamma} \subset \mathcal{R}(C, D)$. Then
   (a) $R \times \left(\bigcup_{\gamma\in\Gamma} S_\gamma\right) = \bigcup_{\gamma\in\Gamma} (R \times S_\gamma)$
   (b) $R \times \left(\bigcap_{\gamma\in\Gamma} S_\gamma\right) = \bigcap_{\gamma\in\Gamma} (R \times S_\gamma)$
2. Let $(R_\gamma)_{\gamma\in\Gamma} \subset\in \mathcal{R}(A, B)$ and $S \in \mathcal{R}(C, D)$. Then
   (a) $\left(\bigcup_{\gamma\in\Gamma} R_\gamma\right) \times S = \bigcup_{\gamma\in\Gamma} (R_\gamma \times S)$
   (b) $\left(\bigcap_{\gamma\in\Gamma} R_\gamma\right) \times S = \bigcap_{\gamma\in\Gamma} (R_\gamma \times S)$

Proof.

1. (a) We note that

$$\begin{aligned}((a,c),(b,d)) \in R \times \left(\bigcup_{\gamma\in\Gamma} S_\gamma\right) &\iff (a,b) \in R \text{ and } (c,d) \in \bigcup_{\gamma\in\Gamma} S_\gamma \\ &\iff (a,b) \in R \text{ and there exists } \gamma_0 \in \Gamma \text{ such that } (c,d) \in S_{\gamma_0} \\ &\iff \text{there exists } \gamma_0 \in \Gamma \text{ such that } (a,b) \in R \text{ and } (c,d) \in S_{\gamma_0} \\ &\iff \text{there exists } \gamma_0 \in \Gamma \text{ such that } ((a,c),(b,d)) \in R \times S_{\gamma_0} \\ &\iff ((a,c),(b,d)) \in \bigcup_{\gamma\in\Gamma}(R \times S_\gamma).\end{aligned}$$

Hence $R \times \left(\bigcup_{\gamma\in\Gamma} S_\gamma\right) = \bigcup_{\gamma\in\Gamma}(R \times S_\gamma)$.

(b) We note that

$$\begin{aligned}((a,c),(b,d)) \in R \times \left(\bigcap_{\gamma\in\Gamma} S_\gamma\right) &\iff (a,b) \in R \text{ and } (c,d) \in \bigcap_{\gamma\in\Gamma} S_\gamma \\ &\iff (a,b) \in R \text{ for each } \gamma \in \Gamma,\ (c,d) \in S_\gamma \\ &\iff \text{for each } \gamma \in \Gamma,\ (a,b) \in R \text{ and } (c,d) \in S_\gamma \\ &\iff \text{for each } \gamma \in \Gamma,\ ((a,c),(b,d)) \in R \times S_\gamma \\ &\iff ((a,c),(b,d)) \in \bigcap_{\gamma\in\Gamma}(R \times S_\gamma).\end{aligned}$$

Hence $R \times \left(\bigcap_{\gamma\in\Gamma} S_\gamma\right) = \bigcap_{\gamma\in\Gamma}(R \times S_\gamma)$.

2. (a) We note that

$$\begin{aligned}((a,c),(b,d)) \in \left(\bigcup_{\gamma\in\Gamma} R_\gamma\right) \times S &\iff (a,b) \in \left(\bigcup_{\gamma\in\Gamma} R_\gamma\right) \text{ and } (c,d) \in S \\ &\iff \text{there exists } \gamma_0 \in \Gamma \text{ such that } (a,b) \in R_{\gamma_0} \text{ and } (c,d) \in S \\ &\iff \text{there exists } \gamma_0 \in \Gamma \text{ such that } ((a,c),(b,d)) \in R_{\gamma_0} \times S \\ &\iff ((a,c),(b,d)) \in \bigcup_{\gamma\in\Gamma}(R_\gamma \times S)\end{aligned}$$

Hence $\left(\bigcup_{\gamma\in\Gamma} R_\gamma\right) \times S = \bigcup_{\gamma\in\Gamma}(R_\gamma \times S)$.

(b) We note that

$$\begin{aligned}((a,c),(b,d)) \in \left(\bigcap_{\gamma\in\Gamma} R_\gamma\right) \times S &\iff (a,b) \in \left(\bigcap_{\gamma\in\Gamma} R_\gamma\right) \text{ and } (c,d) \in S \\ &\iff \text{for each } \gamma \in \Gamma,\ (a,b) \in R_\gamma \text{ and } (c,d) \in S \\ &\iff \text{for each } \gamma \in \Gamma,\ ((a,c),(b,d)) \in R_\gamma \times S \\ &\iff ((a,c),(b,d)) \in \bigcap_{\gamma\in\Gamma}(R_\gamma \times S)\end{aligned}$$

Hence $\left(\bigcap_{\gamma\in\Gamma} R_\gamma\right) \times S = \bigcap_{\gamma\in\Gamma}(R_\gamma \times S)$.

$\square$

1.2 n-ary Heterogeneous Relations

1.2.1 Introduction

can we define composition in an operadic way?

Definition 1.2.1.1. Let $n \in \mathbb{N}$ and $(A_j)_{j\in[n]}$ a collection of sets.

- Let $R \subset \prod_{j\in[n]} A_j$. Then R is said to be a **n-ary relation over** $(A_j)_{j\in[n]}$.
- We define $\mathcal{R}(A_j : j \in [n]) := \mathcal{P}(\prod_{j\in[n]} A_j)$

Definition 1.2.1.2. FIX rest of section!!!! (right now it is same as binary relations section) Let A, B, C be sets, $R \in \mathcal{R}(A, B)$, $S \in \mathcal{R}(B, C)$. We define the **composition relation of S with R**, denoted $S \circ_{\text{Rel}} R \in \mathcal{R}(A, C)$, by

$$S \circ_{\text{Rel}} R := \{(a, c) \in A \times C : \text{there exists } b \in B \text{ such that } (a, b) \in R \text{ and } (b, c) \in S\}.$$

Note 1.2.1.3. When the context is clear, we write $S \circ R$ in place of $S \circ_{\text{Rel}} R$.

Exercise 1.2.1.4. Let A, B, C, D be sets, R a relation between A and B, S relation between B and C and T relation between C and D. Then $(T \circ S) \circ R = T \circ (S \circ R)$.

Proof.

- Let $(a, d) \in (T \circ S) \circ R$. Then there exists $b \in B$ such that $(a, b) \in R$ and $(b, d) \in T \circ S$. Since $(b, d) \in T \circ S$, there exists $c \in C$ such that $(b, c) \in S$ and $(c, d) \in T$. Since $(a, b) \in R$ and $(b, c) \in S$, $(a, c) \in S \circ R$. Since $(a, c) \in S \circ R$ and $(c, d) \in T$, we have that $(a, d) \in T \circ (S \circ R)$. Since $(a, d) \in (T \circ S) \circ R$ is arbitrary, we have that for each $(a, d) \in (T \circ S) \circ R$, $(a, d) \in T \circ (S \circ R)$. Hence $(T \circ S) \circ R \subset T \circ (S \circ R)$.
- Similarly, $T \circ (S \circ R) \subset (T \circ S) \circ R$.

Since $(T \circ S) \circ R \subset T \circ (S \circ R)$ and $T \circ (S \circ R) \subset (T \circ S) \circ R$, we have that $(T \circ S) \circ R = T \circ (S \circ R)$. □

Definition 1.2.1.5. Let A be a set. We define the **diagonal relation on** A, denoted $\Delta_A \in \mathcal{R}(A, A)$, by

$$\Delta_A := \{(a, a) : a \in A\}.$$

Exercise 1.2.1.6. Let A, B be sets and $R \in \mathcal{R}(A, B)$. Then

1. $R \circ \Delta_A = R$
2. $\Delta_B \circ R = R$

Proof.

1.
   - Let $(a, b) \in R \circ \Delta_A$. Then there exists $a' \in A$ such that $(a, a') \in \Delta_A$ and $(a', b) \in R$. By definition of Δ_A, $a = a'$. Therefore $(a, b) \in R$. Since $(a, b) \in R \circ \Delta_A$ is arbitrary, we have that for each $(a, b) \in R \circ \Delta_A$, $(a, b) \in R$. Hence $R \circ \Delta_A \subset R$.
   - Let $(a, b) \in R$. Then $a \in A$. Therefore $(a, a) \in \Delta_A$. Since $(a, a) \in \Delta_A$ and $(a, b) \in R$, we have that $(a, b) \in R \circ \Delta_A$. Since $(a, b) \in R$ is arbitrary, we have that for each $(a, b) \in R$, $(a, b) \in R \circ \Delta_A$. Hence $R \circ \Delta_A \subset R$.

   Since $R \circ \Delta_A \subset R$ and $R \circ \Delta_A \subset R$, we have that $R \circ \Delta_A = R$.
2. Similar to (1).

□

Exercise 1.2.1.7. Let A, B, C be sets.

1. Let $R \in \mathcal{R}(A, B)$ and $(S_\gamma)_{\gamma\in\Gamma} \subset \mathcal{R}(B, C)$. Then
   (a) $\left(\bigcup_{\gamma\in\Gamma} S_\gamma\right) \circ R = \bigcup_{\gamma\in\Gamma} (S_\gamma \circ R)$,

(b) $\left(\bigcap_{\gamma\in\Gamma} S_\gamma\right)\circ R \subset \bigcap_{\gamma\in\Gamma}(S_\gamma \circ R)$

2. Let $(R_\gamma)_{\gamma\in\Gamma} \subset \mathcal{R}(A,B)$ and $S \in \mathcal{R}(B,C)$. Then

   (a) $S \circ \left(\bigcup_{\gamma\in\Gamma} R_\gamma\right) = \bigcup_{\gamma\in\Gamma}(S\circ R_\gamma)$.

   (b) $S \circ \left(\bigcap_{\gamma\in\Gamma} R_\gamma\right) \subset \bigcap_{\gamma\in\Gamma}(S\circ R_\gamma)$.

Proof.

1. (a) We note that

$$\begin{aligned}
(a,c) \in \left(\bigcup_{\gamma\in\Gamma} S_\gamma\right)\circ R &\iff \text{there exists } b\in B \text{ such that } (a,b)\in R \text{ and } (b,c)\in \bigcup_{\gamma\in\Gamma} S_\gamma \\
&\iff \text{there exists } b\in B \text{ and } \gamma_0\in\Gamma \text{ such that } (a,b)\in R \text{ and } (b,c)\in S_{\gamma_0} \\
&\iff \text{there exists } \gamma_0\in\Gamma \text{ such that } (a,c)\in S_{\gamma_0}\circ R \\
&\iff (a,c)\in \bigcup_{\gamma\in\Gamma}(S_\gamma\circ R).
\end{aligned}$$

   Hence $\left(\bigcup_{\gamma\in\Gamma} S_\gamma\right)\circ R = \bigcup_{\gamma\in\Gamma}(S_\gamma\circ R)$.

   (b) We note that

$$\begin{aligned}
(a,c) \in \left(\bigcap_{\gamma\in\Gamma} S_\gamma\right)\circ R &\iff \text{there exists } b\in B \text{ such that } (a,b)\in R \text{ and } (b,c)\in \bigcap_{\gamma\in\Gamma} S_\gamma \\
&\iff \text{there exists } b\in B \text{ such that } (a,b)\in R \text{ and for each } \gamma\in\Gamma \text{ and } (b,c)\in S_\gamma \\
&\implies \text{for each } \gamma\in\Gamma, \text{ there exists } b\in B \text{ such that } (a,b)\in R \text{ and } (b,c)\in S_\gamma \\
&\iff \text{for each } \gamma\in\Gamma,\ (a,c)\in S_\gamma\circ R \\
&\iff (a,c)\in \bigcap_{\gamma\in\Gamma}(S_\gamma\circ R).
\end{aligned}$$

   Hence $\left(\bigcap_{\gamma\in\Gamma} S_\gamma\right)\circ R = \bigcap_{\gamma\in\Gamma}(S_\gamma\circ R)$.

2. (a) We note that

$$\begin{aligned}
(a,c) \in S\circ\left(\bigcup_{\gamma\in\Gamma} R_\gamma\right) &\iff \text{there exists } b\in B \text{ such that } (a,b)\in \bigcup_{\gamma\in\Gamma} R_\gamma \text{ and } (b,c)\in S \\
&\iff \text{there exists } b\in B \text{ and } \gamma_0\in\Gamma \text{ such that } (a,b)\in R_{\gamma_0} \text{ and } (b,c)\in S \\
&\iff \text{there exists } \gamma_0\in\Gamma \text{ such that } (a,c)\in S\circ R_{\gamma_0} \\
&\iff (a,c)\in \bigcup_{\gamma\in\Gamma}(S\circ R_\gamma).
\end{aligned}$$

   Hence $S\circ\left(\bigcup_{\gamma\in\Gamma} R_\gamma\right) = \bigcup_{\gamma\in\Gamma}(S_\gamma\circ R)$.

(b) We note that

$$\begin{aligned}
(a,c) \in S \circ \left(\bigcap_{\gamma \in \Gamma} R_\gamma \right) &\iff \text{there exists } b \in B \text{ such that } (a,b) \in \bigcap_{\gamma \in \Gamma} R_\gamma \text{ and } (b,c) \in S \\
&\iff \text{there exists } b \in B \text{ such that for each } \gamma \in \Gamma,\ (a,b) \in R_\gamma \text{ and } (b,c) \in S \\
&\implies \text{for each } \gamma \in \Gamma, \text{ there exists } b \in B \text{ such that } (a,b) \in R_\gamma \text{ and } (b,c) \in S \\
&\iff \text{for each } \gamma \in \Gamma,\ (a,c) \in S \circ R_\gamma \\
&\iff (a,c) \in \bigcap_{\gamma \in \Gamma} (S \circ R_\gamma).
\end{aligned}$$

Hence $S \circ \left(\bigcap_{\gamma \in \Gamma} R_\gamma \right) \subset \bigcap_{\gamma \in \Gamma} (S_\gamma \circ R)$.

□

Exercise 1.2.1.8. Let A, B, C be sets.

1. Let $R_1, R_2 \in \mathcal{R}(A,B)$ and $S \in \mathcal{R}(B,C)$. If $R_1 \subset R_2$, then $S \circ R_1 \subset S \circ R_2$.
2. Let $R \in \mathcal{R}(A,B)$ and $S_1, S_2 \in \mathcal{R}(B,C)$. If $S_1 \subset S_2$, then $S_1 \circ R \subset S_2 \circ R$.
3. Let $R_1, R_2 \in \mathcal{R}(A,B)$ and $S_1, S_2 \in \mathcal{R}(B,C)$. If $R_1 \subset R_2$ and $S_1 \subset S_2$, then $S_1 \circ R_1 \subset S_2 \circ R_2$.

Proof.

1. Suppose that $R_1 \subset R_2$. Then

$$\begin{aligned}
(a,c) \in S \circ R_1 &\iff \text{there exists } b \in B \text{ such that } (a,b) \in R_1 \text{ and } (b,c) \in S \\
&\implies \text{there exists } b \in B \text{ such that } (a,b) \in R_2 \text{ and } (b,c) \in S \\
&\iff (a,c) \in S \circ R_2.
\end{aligned}$$

   Hence $S \circ R_1 \subset S \circ R_2$.

2. Suppose that $S_1 \subset S_2$. Then

$$\begin{aligned}
(a,c) \in S_1 \circ R &\iff \text{there exists } b \in B \text{ such that } (a,b) \in R \text{ and } (b,c) \in S_1 \\
&\implies \text{there exists } b \in B \text{ such that } (a,b) \in R \text{ and } (b,c) \in S_2 \\
&\iff (a,c) \in S_2 \circ R.
\end{aligned}$$

   Hence $S_2 \circ R \subset S_1 \circ R$.

3. Suppose that $R_1 \subset R_2$ and $S_1 \subset S_2$. Then the previous parts imply that

$$\begin{aligned}
S_1 \circ R_1 &\subset S_2 \circ R_1 \\
&\subset S_2 \circ R_2.
\end{aligned}$$

□

Definition 1.2.1.9. Let A, B be sets and $R \in \mathcal{R}(A,B)$. We define the **converse of** R, denoted $R^* \in \mathcal{R}(B,A)$, by

$$R^* := \{(b,a) \in B \times A : (a,b) \in R\}.$$

Exercise 1.2.1.10. Let A, B be sets and R a relation between A and B. Then $(R^*)^* = R$.

Proof.

$$\begin{aligned}
(a,b) \in (R^*)^* &\iff (b,a) \in R^* \\
&\iff (a,b) \in R.
\end{aligned}$$

Hence $(R^*)^* = R$. □

Exercise 1.2.1.11. Let A, B, C be sets, $R \in \mathcal{R}(A, B)$ and $S \in \mathcal{R}(B, C)$. Then $(S \circ R)^* = R^* \circ S^*$.

Proof. We note that

$$\begin{aligned}
(c, a) \in (S \circ R)^* &\iff (a, c) \in S \circ R \\
&\iff \text{there exists } b \in B \text{ such that } (a, b) \in R \text{ and } (b, c) \in S \\
&\iff \text{there exists } b \in B \text{ such that } (c, b) \in S^* \text{ and } (b, a) \in R^* \\
&\iff (c, a) \in R^* \circ S^*.
\end{aligned}$$

Hence $(S \circ R)^* = R^* \circ S^*$. □

Exercise 1.2.1.12. Let A and B be sets and $(R_\gamma)_{\gamma \in \Gamma} \subset \mathcal{R}(A, B)$. Then

1. $\left(\bigcup_{\gamma \in \Gamma} R_\gamma \right)^* = \bigcup_{\gamma \in \Gamma} R_\gamma^*$
2. $\left(\bigcap_{\gamma \in \Gamma} R_\gamma \right)^* = \bigcap_{\gamma \in \Gamma} R_\gamma^*$

Proof.

1. We note that

$$\begin{aligned}
(b, a) \in \left(\bigcup_{\gamma \in \Gamma} R_\gamma \right)^* &\iff (a, b) \in \bigcup_{\gamma \in \Gamma} R_\gamma \\
&\iff \text{there exists } \gamma_0 \in \Gamma \text{ such that } (a, b) \in R_{\gamma_0} \\
&\iff \text{there exists } \gamma_0 \in \Gamma \text{ such that } (b, a) \in R_{\gamma_0}^* \\
&\iff (b, a) \in \bigcup_{\gamma \in \Gamma} R_\gamma^*.
\end{aligned}$$

   Hence $\left(\bigcup_{\gamma \in \Gamma} R_\gamma \right)^* = \bigcup_{\gamma \in \Gamma} R_\gamma^*$.

2. We note that

$$\begin{aligned}
(b, a) \in \left(\bigcap_{\gamma \in \Gamma} R_\gamma \right)^* &\iff (a, b) \in \bigcap_{\gamma \in \Gamma} R_\gamma \\
&\iff \text{for each } \gamma \in \Gamma,\ (a, b) \in R_\gamma \\
&\iff \text{for each } \gamma \in \Gamma,\ (b, a) \in R_\gamma^* \\
&\iff (b, a) \in \bigcap_{\gamma \in \Gamma} R_\gamma^*.
\end{aligned}$$

   Hence $\left(\bigcap_{\gamma \in \Gamma} R_\gamma \right)^* = \bigcap_{\gamma \in \Gamma} R_\gamma^*$.

□

1.2.2 Subrelations

Definition 1.2.2.1. Let A, B be sets, $R \in \mathcal{R}(A, B)$, $E \subset A$ and $F \subset B$. We define the **subrelation of R between E and F**, denoted $R|_{E \times F} \in \mathcal{R}(E, F)$, by $R|_{E \times F}$

maybe dont do this subsection, since relations are like morphisms, so unless you can find a categorical way to present restrictions on domain or codomain $f|_E$ or $f|^F$, just get rid of this subsection

1.2.3 Products of Relations

Definition 1.2.3.1. Let A, B, C, D be sets and $R \in \mathcal{R}(A, B), S \in \mathcal{R}(C, D)$. We define the **product relation of R and S**, denoted $R \times_{\text{Rel}} S \in \mathcal{R}(A \times C, B \times D)$, by

$$R \times_{\text{Rel}} S := \{((a, c), (b, d)) \in (A \times C) \times (B \times D) : (a, b) \in R \text{ and } (c, d) \in S\}.$$

Note 1.2.3.2. When the context is clear, we write $R \times S$ in place of $R \times_{\text{Rel}} S$.

Exercise 1.2.3.3. Let A, B, C, D be sets.

1. Let $R \in \mathcal{R}(A, B)$ and $(S_\gamma)_{\gamma \in \Gamma} \subset \mathcal{R}(C, D)$. Then
   (a) $R \times \left(\bigcup_{\gamma \in \Gamma} S_\gamma \right) = \bigcup_{\gamma \in \Gamma} (R \times S_\gamma)$
   (b) $R \times \left(\bigcap_{\gamma \in \Gamma} S_\gamma \right) = \bigcap_{\gamma \in \Gamma} (R \times S_\gamma)$
2. Let $(R_\gamma)_{\gamma \in \Gamma} \subset\in \mathcal{R}(A, B)$ and $S \in \mathcal{R}(C, D)$. Then
   (a) $\left(\bigcup_{\gamma \in \Gamma} R_\gamma \right) \times S = \bigcup_{\gamma \in \Gamma} (R_\gamma \times S)$
   (b) $\left(\bigcap_{\gamma \in \Gamma} R_\gamma \right) \times S = \bigcap_{\gamma \in \Gamma} (R_\gamma \times S)$

Proof.

1. (a) We note that

$$\begin{aligned}
((a, c), (b, d)) \in R \times \left(\bigcup_{\gamma \in \Gamma} S_\gamma \right) &\iff (a, b) \in R \text{ and } (c, d) \in \bigcup_{\gamma \in \Gamma} S_\gamma \\
&\iff (a, b) \in R \text{ and there exists } \gamma_0 \in \Gamma \text{ such that } (c, d) \in S_{\gamma_0} \\
&\iff \text{there exists } \gamma_0 \in \Gamma \text{ such that } (a, b) \in R \text{ and } (c, d) \in S_{\gamma_0} \\
&\iff \text{there exists } \gamma_0 \in \Gamma \text{ such that } ((a, c), (b, d)) \in R \times S_{\gamma_0} \\
&\iff ((a, c), (b, d)) \in \bigcup_{\gamma \in \Gamma} (R \times S_\gamma).
\end{aligned}$$

   Hence $R \times \left(\bigcup_{\gamma \in \Gamma} S_\gamma \right) = \bigcup_{\gamma \in \Gamma} (R \times S_\gamma)$.

   (b) We note that

$$\begin{aligned}
((a, c), (b, d)) \in R \times \left(\bigcap_{\gamma \in \Gamma} S_\gamma \right) &\iff (a, b) \in R \text{ and } (c, d) \in \bigcap_{\gamma \in \Gamma} S_\gamma \\
&\iff (a, b) \in R \text{ for each } \gamma \in \Gamma,\ (c, d) \in S_\gamma \\
&\iff \text{for each } \gamma \in \Gamma,\ (a, b) \in R \text{ and } (c, d) \in S_\gamma \\
&\iff \text{for each } \gamma \in \Gamma,\ ((a, c), (b, d)) \in R \times S_\gamma \\
&\iff ((a, c), (b, d)) \in \bigcap_{\gamma \in \Gamma} (R \times S_\gamma).
\end{aligned}$$

   Hence $R \times \left(\bigcap_{\gamma \in \Gamma} S_\gamma \right) = \bigcap_{\gamma \in \Gamma} (R \times S_\gamma)$.

2. (a) We note that

$$\begin{aligned}((a,c),(b,d)) \in \left(\bigcup_{\gamma\in\Gamma} R_\gamma\right) \times S &\iff (a,b) \in \left(\bigcup_{\gamma\in\Gamma} R_\gamma\right) \text{ and } (c,d) \in S \\ &\iff \text{there exists } \gamma_0 \in \Gamma \text{ such that } (a,b) \in R_{\gamma_0} \text{ and } (c,d) \in S \\ &\iff \text{there exists } \gamma_0 \in \Gamma \text{ such that } ((a,c),(b,d)) \in R_{\gamma_0} \times S \\ &\iff ((a,c),(b,d)) \in \bigcup_{\gamma\in\Gamma} (R_\gamma \times S)\end{aligned}$$

Hence $\left(\bigcup_{\gamma\in\Gamma} R_\gamma\right) \times S = \bigcup_{\gamma\in\Gamma} (R_\gamma \times S)$.

(b) We note that

$$\begin{aligned}((a,c),(b,d)) \in \left(\bigcap_{\gamma\in\Gamma} R_\gamma\right) \times S &\iff (a,b) \in \left(\bigcap_{\gamma\in\Gamma} R_\gamma\right) \text{ and } (c,d) \in S \\ &\iff \text{for each } \gamma \in \Gamma,\ (a,b) \in R_\gamma \text{ and } (c,d) \in S \\ &\iff \text{for each } \gamma \in \Gamma,\ ((a,c),(b,d)) \in R_\gamma \times S \\ &\iff ((a,c),(b,d)) \in \bigcap_{\gamma\in\Gamma} (R_\gamma \times S)\end{aligned}$$

Hence $\left(\bigcap_{\gamma\in\Gamma} R_\gamma\right) \times S = \bigcap_{\gamma\in\Gamma} (R_\gamma \times S)$.

$\square$

1.3 n-ary Homogeneous Relations

1.3.1 Introduction

Definition 1.3.1.1. Let $n \in \mathbb{N}$ and A a set.

- Let $R \subset A^n$. Then R is said to be a **homogeneous n-ary relation on** A
- We define $\mathcal{R}^n(A) := \mathcal{P}(A^n)$.

1.3.2 Subrelations

Definition 1.3.2.1. Let $n \in \mathbb{N}$, A, B sets, $R \in \mathcal{R}^n(A)$ and $S \in \mathcal{R}^n(B)$. Then

- S is said to be a **restriction of** R if $B \subset A$,
- R is said to be an **extension of** S if S is a restriction of R.

Definition 1.3.2.2. Let $n \in \mathbb{N}$, A a set, $B \subset A$ and $R \in \mathcal{R}^n(A)$. We define the **restriction of** R **to** B, denoted $R|_B$, by $R|_B := R \cap B^n$.

Exercise 1.3.2.3. Let $n \in \mathbb{N}$, A a set, $B \subset A$ and $R \in \mathcal{R}^n(A)$. Then $R|_B$ is a restriction of R.

Proof. Clear. fill out later □

1.3.3 Product Relations

1.4 Binary Homogeneous Relations

1.4.1 Subrelations

Exercise 1.4.1.1. Let A be a set, $B \subset A$ and $R \in \mathcal{R}^2(A)$. Then $(R^*)|_B = (R|_B)^*$

Proof. Clear by prev ex in section on binary hetero relations. □

1.5 Equivalence Relations

Note 1.5.0.1. Let X be a set and $\nu \in \mathcal{R}^2(X)$. When the context is clear, we may write $a\nu b$ in place of $(a, b) \in \nu$.

Definition 1.5.0.2. Let X be a set. Suppose that $X \neq \varnothing$.

- Let $\theta \in \mathcal{R}^2(X)$. Then θ is said to be
  - **reflexive** if for each $a \in X$, $(a, a) \in \theta$
  - **symmetric** if for each $a, b \in X$, $(a, b) \in \theta$ implies that $(b, a) \in \theta$
  - **transitive** if for each $a, b, c \in X$, $(a, b) \in \theta$ and $(b, c) \in \theta$ implies that $(a, c) \in \theta$
  - an **equivalence relation on** X if θ is reflexive, symmetric and transitive
- We define $\mathrm{Equiv}(X) := \{\theta \in \mathcal{R}^2(X) : \theta \text{ is an equivalence relation on } X\}$

Exercise 1.5.0.3. Let X be a set and $\theta \in \mathcal{R}^2(X)$. Then

1. θ is reflexive iff $\Delta_X \subset \theta$
2. θ is symmetric iff $\theta^* \subset \theta$
3. θ is transitive iff $\theta \circ \theta \subset \theta$

Proof.

1.
   - $(\implies)$:
     Suppose that θ is reflexive. Let $(a, a) \in \Delta_X$. Then $a \in X$. Since θ is reflexive, $(a, a) \in \theta$. Since $(a, a) \in \Delta_X$ is arbitrary, we have that for each $(a, a) \in \Delta_X$, $(a, a) \in \theta$. Hence $\Delta_X \subset \theta$.
   - $(\impliedby)$:
     Suppose that $\Delta_X \subset \theta$. Let $a \in X$. By definition and assumption,
     $$\begin{aligned}(a, a) &\in \Delta_X \\ &\subset \theta.\end{aligned}$$
     Since $a \in X$ is arbitrary, we have that for each $a \in X$, $(a, a) \in \theta$. Hence θ is reflexive.
2.
   - $(\implies)$:
     Suppose that θ is symmetric. Let $(b, a) \in \theta^*$. By definition, $(a, b) \in \theta$. Since θ is symmetric, $(b, a) \in \theta$. Since $(b, a) \in \theta^*$ is arbitrary, we have that for each $(b, a) \in \theta^*$, $(b, a) \in \theta$. Hence $\theta^* \subset \theta$.
   - $(\impliedby)$:
     Suppose that $\theta^* \subset \theta$. Let $a, b \in X$. Suppose that $(a, b) \in \theta$. By definition and assumption,
     $$\begin{aligned}(b, a) &\in \theta^* \\ &\subset \theta.\end{aligned}$$
     Since $a, b \in X$ with $(a, b) \in \theta$ are arbitrary, we have that for each $a, b \in X$, $(a, b) \in \theta$ implies that $(b, a) \in \theta$. Hence θ is symmetric.
3.
   - $(\implies)$:
     Suppose that θ is transitive. Let $(a, c) \in \theta \circ \theta$. By definition, there exists $b \in X$ such that $(a, b) \in \theta$ and $(b, c) \in \theta$. Since θ is transitive, $(a, c) \in \theta$. Since $(a, c) \in \theta \circ \theta$ is arbitrary, we have that for each $(a, c) \in \theta \circ \theta$, $(a, c) \in \theta$. Hence $\theta \circ \theta \subset \theta$.
   - $(\impliedby)$:
     Suppose that $\theta \circ \theta \subset \theta$. Let $a, b, c \in X$. Suppose that $(a, b) \in \theta$ and $(b, c) \in \theta$. By definition and assumption,
     $$\begin{aligned}(a, c) &\in \theta \circ \theta \\ &\subset \theta.\end{aligned}$$
     Since $a, b, c \in X$ with $(a, b) \in \theta$ and $(b, c) \in \theta$ are arbitrary, we have that for each $a, b \in X$, $(a, b) \in \theta$ and $(b, c) \in \theta$ implies that $(a, c) \in \theta$. Hence θ is transitive.

□

Definition 1.5.0.4. Let X be a set. Suppose that $X \neq \varnothing$. We define the **universal relation on** X, denoted $\nabla_X \in \mathcal{R}^2(X)$, by $\nabla_X := X^2$.

Exercise 1.5.0.5. Let X be a set. Suppose that $X \neq \varnothing$. Then $\nabla_X \in \mathrm{Equiv}(X)$.

Proof. Let $a, b, c \in X$.

1. Since $a \in X$, $(a, a) \in \nabla_X$.
2. Suppose that $(a, b) \in \nabla_X$. Since $a, b \in X$, $(b, a) \in X^2$. Therefore $(a, b) \in \nabla_X$ implies that $(b, a) \in \nabla_X$.
3. Suppose that $(a, b), (b, c) \in \nabla_X$. Since $a, c \in X$, $(a, c) \in X^2$. Therefore $(a, b), (b, c) \in X^2$ implies that $(a, c) \in \nabla_X$.

□

Exercise 1.5.0.6. Let X be a set. Suppose that $X \neq \varnothing$. Then $\Delta_X \in \mathrm{Equiv}(X)$.

Proof. Let $a, b, c \in X$.

1. Since $a \in X$, $(a, a) \in \Delta_X$.
2. Suppose that $(a, b) \in \Delta_X$. Then $a = b$ and
$$\begin{aligned}(b, a) &= (a, a)\\ &\in \Delta_X.\end{aligned}$$
Therefore $(a, b) \in \Delta_X$ implies that $(b, a) \in \Delta_X$.
3. Suppose that $(a, b), (b, c) \in \Delta_X$. Then $a = b$ and $b = c$. Hence $a = c$ and
$$\begin{aligned}(a, c) &= (a, a)\\ &\in \Delta_X.\end{aligned}$$
Therefore $(a, b), (b, c) \in \Delta_X$ implies that $(a, c) \in \Delta_X$.

□

Exercise 1.5.0.7. Let X be a set. Suppose that $X \neq \varnothing$. Then for each $(\theta_\alpha)_{\alpha \in A} \subset \mathrm{Equiv}(X)$, $\bigcap_{\alpha \in A} \theta_\alpha \in \mathrm{Equiv}(X)$.

Proof. Let $(\theta_\alpha)_{\alpha \in A} \subset \mathrm{Equiv}(X)$. Set $\theta := \bigcap_{\alpha \in A} \theta_\alpha$. Let $a, b, c \in X$.

1. Let $\alpha \in A$. Since $\theta_\alpha \in \mathrm{Equiv}(X)$, $(a, a) \in \theta_\alpha$. Since $\alpha \in A$ is arbitrary, we have that for each $\alpha \in A$, $(a, a) \in \theta_\alpha$. Hence $(a, a) \in \theta$.
2. Suppose that $(a, b) \in \theta$. Let $\alpha \in A$. Then $(a, b) \in \theta_\alpha$. Since $\theta_\alpha \in \mathrm{Equiv}(X)$, we have that $(b, a) \in \theta_{a\backslash}$. Since $\alpha \in A$ is arbitrary, we have that for each $\alpha \in A$, $(b, a) \in \theta_\alpha$. Hence $(b, a) \in \theta$. Therefore $(a, b) \in \theta$ implies that $(b, a) \in \theta$.
3. Suppose that $(a, b) \in \theta$ and $(b, c) \in \theta$. Let $\alpha \in A$. Then $(a, b), (b, c) \in \theta_\alpha$. Since $\theta_\alpha \in \mathrm{Equiv}(X)$, $(a, c) \in \theta_\alpha$. Since $\alpha \in A$ is arbitrary, we have that for each $\alpha \in A$, $(a, c) \in \theta_\alpha$. Hence $(a, c) \in \theta$. Therefore $(a, b) \in \theta$ and $(b, c) \in \theta$ implies that $(a, c) \in \theta$.

Hence $\theta \in \mathrm{Equiv}(X)$. □

Definition 1.5.0.8. Let X be a set. Suppose that $X \neq \varnothing$. Let $\nu \in \mathcal{R}^2(X)$.

- We define the **equivalence relations containing** ν, denoted $\mathcal{E}(\nu)$, by $\mathcal{E}(\nu) := \{\theta \in \mathrm{Equiv}(X) : \nu \subset \theta\}$.
- We define the **equivalence relation on** X **generated by** ν, denoted $\epsilon(\nu) \in \mathcal{R}^2(X)$ by $\epsilon(\nu) := \bigcap_{\theta \in \mathcal{E}(\nu)} \theta$.

Note 1.5.0.9.

- A previous ex implies that $\nabla_X \in \mathcal{E}(\nu)$.
- A previous ex (about intersections of equiv relats) implies that $\epsilon(\nu) \in \mathcal{E}(\nu)$.

Exercise 1.5.0.10. Let X be a set. Suppose that $X \neq \varnothing$. Let $\nu \in \mathcal{R}^2(X)$. Define $(\nu_n)_{n\in\mathbb{N}_0} \subset \mathcal{R}^2(X)$ by

$$\nu_n := \begin{cases} \Delta_X \cup \nu \cup \nu^*, & n = 0 \\ \nu_{n-1} \circ \nu_{n-1}, & n \geq 1. \end{cases}$$

Then

1. (a) for each $n \in \mathbb{N}$, ν_n is reflexive
   (b) for each $n \in \mathbb{N}$, ν_n is symmetric
   (c) for each $n \in \mathbb{N}_0$, $\nu_n \subset \nu_{n+1}$
2. $\epsilon(\nu) = \bigcup_{n\in\mathbb{N}_0} \nu_n$.

Proof.

1. (a)
   - **Base Case:**
     Since
     $$\begin{aligned} \Delta_X &\subset \Delta_X \cup \nu \cup \nu^* \\ &= \nu_0, \end{aligned}$$
     a previous ex implies that ν_0 is reflexive.
   - **Induction Step:**
     Let $n \in \mathbb{N}_0$. Suppose that ν_n is reflexive. Then that same prev ex implies that $\Delta_X \subset \nu_n$. two prev ex in section on relations about composition with identity and subsets then imply that
     $$\begin{aligned} \Delta_X &= \Delta_X \circ \Delta_X \\ &\subset \nu_n \circ \nu_n \\ &= \nu_{n+1}. \end{aligned}$$
     Another application of that prev ex implies that ν_{n+1} is reflexive.

   By induction, for each $n \in \mathbb{N}_0$, ν_n is reflexive.

   (b)
   - **Base Case:**
     a previous ex about converse relations and unions, double converse and since $\Delta_X \in \mathrm{Equiv}(X)$ and an ex about symmetric relations implies that
     $$\begin{aligned} \nu_0^* &= (\Delta_X \cup \nu \cup \nu^*)^* \\ &= \Delta_X^* \cup \nu^* \cup (\nu^*)^* \\ &= \Delta_X^* \cup \nu^* \cup \nu \\ &\subset \Delta_X \cup \nu^* \cup \nu \\ &= \Delta_X \cup \nu \cup \nu^* \\ &= \nu_0. \end{aligned}$$
     that same previous ex about symmetric relations implies that ν_0 is symmetric.
   - **Induction Step:**
     Let $n \in \mathbb{N}_0$. Suppose that ν_n is symmetric. Then that same prev ex about symmetric and converse relations implies that $\nu_n^* \subset \nu_n$. Therefore prev exs imply
     $$\begin{aligned} \nu_{n+1}^* &= (\nu_n \circ \nu_n)^* \\ &= \nu_n^* \circ \nu_n^* \\ &\subset \nu_n \circ \nu_n. \end{aligned} \qquad = \nu_{n+1}.$$
     Another application of that prev ex implies that ν_{n+1} is symmetric.

   By induction, for each $n \in \mathbb{N}_0$, ν_n is symmetric.

(c) Let $n \in \mathbb{N}_0$. Since ν_n is reflexive, prev ex implies that $\Delta_X \subset \nu_n$. Therefore prev ex implies

$$\begin{aligned}\nu_n &= \Delta_X \circ \nu_n \\ &\subset \nu_n \circ \nu_n \\ &= \nu_{n+1}.\end{aligned}$$

Since $n \in \mathbb{N}_0$ is arbitrary, we have that for each $n \in \mathbb{N}_0$, $\nu_n \subset \nu_{n+1}$.

2. Set $\theta_0 := \bigcup_{n \in \mathbb{N}_0} \nu_n$.

- – **Base Case:**
  Since $\epsilon(\nu) \in \mathcal{E}(\nu)$, we have that $\epsilon(\nu) \in \mathrm{Equiv}(X)$ and $\nu \subset \epsilon(\nu)$. Since $\epsilon(\nu) \in \mathrm{Equiv}(X)$, we have that $\epsilon(\nu)$ is reflexive and symmetric. A ref ex then implies that $\Delta_X \subset \epsilon(\nu)$ and $\epsilon(\nu)^* \subset \epsilon(\nu)$. Therefore a prev ex about converse relations and subsets

$$\begin{aligned}\nu_0 &= \Delta_X \cup \nu \cup \nu^* \\ &\subset \epsilon(\nu) \cup \epsilon(\nu) \cup \epsilon(\nu)^* \\ &\subset \epsilon(\nu) \cup \epsilon(\nu) \cup \epsilon(\nu) \\ &= \epsilon(\nu).\end{aligned}$$

  – **Induction Step:**
  Let $n \in \mathbb{N}_0$. Suppose that $\nu_n \subset \epsilon(\nu)$. Then ref prev ex

$$\begin{aligned}\nu_{n+1} &= \nu_n \circ \nu_n \\ &\subset \epsilon(\nu) \circ \epsilon(\nu) \\ &= \epsilon(\nu).\end{aligned}$$

  By induction, for each $n \in \mathbb{N}_0$, $\nu_n \subset \epsilon(\nu)$. Therefore

$$\begin{aligned}\theta_0 &= \bigcup_{n \in \mathbb{N}_0} \nu_n \\ &\subset \epsilon(\nu).\end{aligned}$$

- – Since ν_0 is reflexive, we have that

$$\begin{aligned}\Delta_X &\subset \nu_0 \\ &\subset \bigcup_{n \in \mathbb{N}_0} \nu_n \\ &= \theta_0.\end{aligned}$$

  Hence θ_0 is reflexive.
  – Since for each $n \in \mathbb{N}_0$, ν_n is symmetric, we have that that ref exs for each $n \in \mathbb{N}_0$, $\nu_n^* \subset \nu_n$. Therefore

$$\begin{aligned}\theta_0^* &= (\bigcup_{n \in \mathbb{N}_0} \nu_n)^* \\ &= \bigcup_{n \in \mathbb{N}_0} \nu_n^* \\ &\subset \bigcup_{n \in \mathbb{N}_0} \nu_n \\ &= \theta_0.\end{aligned}$$

  Hence θ_0 is symmetric.

– Let $a, b, c \in X$. Suppose that $(a, b) \in \theta_0$ and $(b, c) \in \theta_0$. Then there exist $n_1, n_2 \in \mathbb{N}_0$ such that $(a, b) \in \nu_{n_1}$ and $(b, c) \in \nu_{n_2}$. Set $n_0 := \max\{n_1, n_2\}$. Since for each $n \in \mathbb{N}$, $\nu_n \subset \nu_{n+1}$, we have that

$$\begin{aligned}(a, b) &\in \nu_{n_1} \\ &\subset \nu_{n_0}.\end{aligned}$$

and

$$\begin{aligned}(b, c) &\in \nu_{n_2} \\ &\subset \nu_{n_0}.\end{aligned}$$

Since $(a, b), (b, c) \in \nu_{n_0}$, we have that

$$\begin{aligned}(a, c) &\in \nu_{n_0} \circ \nu_{n_0} \\ &= \nu_{n_0+1} \\ &\subset \bigcup_{n \in \mathbb{N}_0} \nu_n \\ &= \theta_0.\end{aligned}$$

Since $a, b, c \in X$ with $(a, b) \in \theta_0$ and $(b, c) \in \theta_0$ are arbitrary, we have that for each $a, b, c \in X$, $(a, b) \in \theta_0$ and $(b, c) \in \theta_0$ implies that $(a, c) \in \theta_0$. Hence θ_0 is transitive.

Since θ_0 is reflexive, symmetric and transitive, $\theta_0 \in \mathrm{Equiv}(X)$. Since

$$\begin{aligned}\nu &\subset \Delta_X \cup \nu \cup \nu^* \\ &= \nu_0 \\ &\subset \bigcup_{n \in \mathbb{N}_0} \nu_n \\ &= \theta_0,\end{aligned}$$

we have that $\theta_0 \in \mathcal{E}(\nu)$. Therefore

$$\begin{aligned}\epsilon(\nu) &= \bigcap_{\theta \in \mathcal{E}(\nu)} \theta \\ &\subset \theta_0.\end{aligned}$$

Since $\theta_0 \subset \epsilon(\nu)$ and $\epsilon(\nu) \subset \theta_0$, we have that $\theta_0 = \epsilon(\nu)$.

□

1.6 Products

Chapter 2

Ordered Sets

2.1 To Do

- Relabel exercises
- finish generated element subsection in galois connection section

at this point, there should be a simple structure capturing prosets and $\downarrow S$, measurable spaces and $\sigma(\mathcal{A})$, topological spaces and $\tau(\mathcal{E})$, groups/rings/$\cdots$/algebras and $\langle S\rangle/(E)/\cdots/(F)$. It seems like we need a category $\mathcal{C}$ with some structure, another category $\mathcal{C}_F$ which is like $\mathcal{C}$, but less structured and some forgetful-like functor $F : \mathcal{C} \to \mathcal{C}_F$. eg, $F : \mathbf{Pro} \to \mathbf{Set}$ or $F : \mathbf{Top} \to \mathbf{Set}$ and we need an associated poset $\mathcal{P}$ containing the structure forgotten by F, e.g. the lower sets of X or the set of topologies on X and we need a "generating" or "contained" object $\mathcal{E}$ in $\mathcal{C}_F$ in the sense that there is at least one object A in $\mathcal{C}$ and monomirphism $\iota_A : \mathcal{E} \to F(A)$ in $\mathcal{C}_F$. We then need a minimal element in the structure poset $\mathcal{P}$ in some universal sense relating to these monomorphisms. Ask people who know category theory

2.2 Prosets

2.2.1 Introduction

Definition 2.2.1.1. Proset:
Let X be a set and $\leq \subset X \times X$ a binary relation on X. Then

- $\leq$ is said to be a **preorder on** X if
  1. for each $a \in X$, $a \leq a$
  2. for each $a, b, c \in X$, $a \leq b$ and $b \leq c$ implies that $a \leq c$
- $(X, \leq)$ is said to be a **preordered set** or **proset** if $\leq$ is a preorder on X.

Definition 2.2.1.2. maybe just ref def about converse relation in set theory appendices section on heterogeneous binary relations
Let $(X, \leq)$ be a proset. We define the **dual order of** $\leq$ **on** X, denoted $\leq^*$, by $a \leq^* b$ iff $b \leq a$.

Exercise 2.2.1.3. Let $(X, \leq)$ be a proset. Then $\leq^*$ is a preorder on X.

Proof.

1. Let $a \in X$. Since $a \leq a$, we have that $a \leq^* a$.
2. Let $a, b, c \in X$. Suppose that $a \leq^* b$ and $b \leq^* c$. Then $b \leq a$ and $c \leq b$. Hence $c \leq a$. Thus $a \leq^* c$.

Therefore $\leq^*$ is a preorder on X. □

Definition 2.2.1.4. Let $(X, \leq)$ be a proset. We define $\sim_{\leq} \subset X \times X$ by $a \sim_{\leq} b$ iff $a \leq b$ and $b \leq a$.

Exercise 2.2.1.5. Let $(X, \leq)$ be a proset. Then $\sim_{\leq}$ is an equivalence relation on X.

Proof. Let $x, y, z \in X$.

1. Since $x \leq x$, $x \sim_{\leq} x$.
2. Suppose that $x \sim_{\leq} y$. Then $x \leq y$ and $y \leq x$. Thus $y \sim_{\leq} x$.
3. Suppose that $x \sim_{\leq} y$ and $y \sim_{\leq} z$. Then $x \leq y$, $y \leq x$, $y \leq z$ and $z \leq y$. Therefore $x \leq z$ and $z \leq x$. Hence $x \sim_{\leq} z$.

□

Exercise 2.2.1.6. Let $(X, \leq)$ be a proset. Then for each $x, y \in X$, $x \sim_{\leq} y$ iff $x \sim_{\leq^*} y$.

Proof. We have that for each $x, y \in X$,

$$\begin{aligned} x \sim_{\leq} y &\iff x \leq y \text{ and } y \leq x \\ &\iff y \leq^* x \text{ and } x \leq^* y \\ &\iff x \sim_{\leq^*} y. \end{aligned}$$

□

Exercise 2.2.1.7. Let X be a set. Then $(\mathcal{P}(X), \subset)$ is a proset.

Proof.

1. For each $A \in \mathcal{P}(X)$, $A \subset A$.
2. For each $A, B, C \in \mathcal{P}(X)$, if $A \subset B$ and $B \subset C$, then $A \subset C$.

Thus $\subset$ is a preorder on $\mathcal{P}(X)$. Hence $(\mathcal{P}(X), \subset)$ is a proset. □

Talk about presets/posests where the order is induced by a map and another preset/poset, i.e. Let $(Y, \leq)$ be a preset/poset and $f : X \to Y$. Define $f^* \leq$ on X by $x_1 f^* \leq x_2$ iff $f(x_1) \leq f(x_2)$. Similarly, If $(X, \leq)$ is a preset/poset and $f : X \to Y$. Define $f_* \leq$ on Y by $y_1 f_* \leq y_2$ iff there exists $x_1 \in f^{-1}(y_1)$ and $x_2 \in f^{-1}(y_2)$ such that $x_1 \leq x_2$. When are these well defined? I think the posetal reflection is an example of the second case.

2.2.2 Subprosets

Definition 2.2.2.1. ref defn in set theory appendices on restrictions of n-ary homogeneous relations

Note 2.2.2.2. Let $(X, \leq)$ be a proset and $A \subset X$. When the context is clear, we write $(A, \leq)$ in place of $(A, \leq|_A)$

2.2.3 Products

Definition 2.2.3.1. Let $(A, \leq_A)$ and $(B, \leq_B)$ be prosets. We define the

- **product preorder of $\leq_A$ and $\leq_B$ on $A \times B$**, denoted $\leq_A \otimes \leq_B$ by $(a_1, b_1) \leq_A \otimes \leq_B (a_2, b_2)$ iff $a_1 \leq_A a_2$ and $b_1 \leq_B b_2$.
- **product proset of $(A, \leq_A)$ and $(B, \leq_B)$**, denoted $(A, \leq_A) \otimes (B, \leq_B)$ by $(A, \leq_A) \otimes (B, \leq_B) := (A \times B, \leq_A \otimes \leq_B)$

Exercise 2.2.3.2. probably need to change notation since $\otimes$ might be reserved for something else. Let $(A, \leq_A)$ and $(B, \leq_B)$ be prosets. Then

1. $\leq_A \otimes \leq_B$ is a preorder on $A \times B$,
2. $(A, \leq_A) \otimes (B, \leq_B)$ is a proset.

Proof.

1. (a) Let $(a, b) \in A \times B$. Then $a \leq_A a$ and $b \leq_B b$. Therefore $(a, b) \leq_A \otimes \leq_B (a, b)$.
   (b) Let $(a_1, b_1), (a_2, b_2), (a_3, b_3) \in A \times B$. Suppose that $(a_1, b_1) \leq_A \otimes \leq_B (a_2, b_2)$ and $(a_2, b_2) \leq_A \otimes \leq_B (a_3, b_3)$. Then $a_1 \leq_A a_2$, $a_2 \leq_A a_3$, $b_1 \leq_B b_2$ and $b_2 \leq_B b_3$. Therefore $a_1 \leq_A a_3$ and $b_1 \leq_B b_3$. Hence $(a_1, b_1) \leq_A \otimes \leq_B (a_3, b_3)$.

   Hence $\leq_A \otimes \leq_B$ is a preorder on $A \times B$.
2. Since $\leq_A \otimes \leq_B$ is a preorder on $A \times B$, $(X, \leq)$ is a proset.

□

2.2.4 Quotients

2.3 Posets

2.3.1 Introduction

Definition 2.3.1.1. Poset:
Let $(X, \leq)$ be a proset. Then

- $\leq$ is said to be a **partial order on** X if for each $a, b \in X$, $a \leq b$ and $b \leq a$ implies that $a = b$
- $(X, \leq)$ is said to be a **partially ordered set** or **poset** if $\leq$ is a partial ordering on X.

Exercise 2.3.1.2. Let $(X, \leq)$ be a poset and $a, b \in X$. Then $a \sim_{\leq} b$ iff $a = b$.

Proof.

- $(\implies)$:
  Suppose that $a \sim_{\leq} b$. Then $a \leq b$ and $b \leq a$. Since $\leq$ is a partial order, $a = b$.
- $(\impliedby)$:
  Since $\sim_{\leq}$ is reflexive, $a = b$ implies that $a \sim_{\leq} b$.

□

Definition 2.3.1.3. Let $(X, \leq)$ be a proset. We define the

- **posetal reflection partial order on** $X/\sim_{\leq}$, denoted $\leq_{X/\sim_{\leq}} \subset X/\sim_{\leq} \times X/\sim_{\leq}$, by $y_1 \leq_{X/\sim_{\leq}} y_2$ iff there exists $x_1 \in y_1$ and $x_2 \in y_2$ such that $x_1 \leq x_2$.
- **the posetal reflection of** $(X, \leq)$, denoted $\mathrm{PosRefl}(X, \leq)$, by $\mathrm{PosRefl}(X, \leq) := (X/\sim_{\leq}, \leq_{X/\sim_{\leq}})$

Exercise 2.3.1.4. Let $(X, \leq)$ be a proset. Then $\mathrm{PosRefl}(X, \leq)$ is a poset.

Proof. Set $Y := X/\sim_{\leq}$ and $\leq_Y := X/\sim_{\leq}$.

1. (a) Let $y \in Y$. Then there exists $x \in y$. Since $x \leq x$, we have that $y \leq_Y y$.
   (b) Let $y_1, y_2, y_3 \in Y$. Suppose that $y_1 \leq_Y y_2$ and $y_2 \leq_Y y_3$. Then there exist $x_1 \in y_1$, $x_2 \in y_2$ and $x_3 \in y_3$ such that $x_1 \leq x_2$ and $x_2 \leq x_3$. Therefore $x_1 \leq x_3$. Hence $y_1 \leq_Y y_3$.

   Thus $\leq_Y$ is a preorder on Y.
2. Let $y_1, y_2 \in Y$. Suppose that $y_1 \leq_Y y_2$ and $y_2 \leq_Y y_1$. Then there exist $a_1, b_1 \in y_1$, $a_2, b_2 \in y_2$ such that $a_1 \leq a_2$ and $b_2 \leq b_1$. Since $a_1, b_1 \in y_1$ and $a_2, b_2 \in y_2$, $a_1 \sim_{\leq} b_1$ and $a_2 \sim_{\leq} b_2$. Thus

$$\begin{aligned} a_1 &\leq a_2 \\ &\leq b_2. \end{aligned}$$

   and

$$\begin{aligned} b_2 &\leq b_1 \\ &\leq a_1. \end{aligned}$$

   Hence $a_1 \sim_{\leq} b_2$ and

$$\begin{aligned} y_1 &= \pi(a_1) \\ &= \pi(b_2) \\ &= y_2. \end{aligned}$$

Therefore $\leq_Y$ is a partial order on Y. □

Exercise 2.3.1.5. Let X be a set. Then $(\mathcal{P}(X), \subset)$ is a poset.

Proof. Exercise 2.2.1.7 implies that $(\mathcal{P}(X), \subset)$ is a proset. We note that for each $A, B \in \mathcal{P}(X)$, $A \subset B$ and $B \subset A$ implies that $A = B$ ref def of equality in set theory notes?. □

2.4 Directed Sets

2.4.1 Introduction

Definition 2.4.1.1. Directed Set:
Let $(A, \leq)$ be a proset. Then

- $\leq$ is said to be a **direction on** A if for each $\alpha, \beta \in A$, $\mathrm{ub}(\{\alpha, \beta\}, \leq) \neq \varnothing$.
- $(A, \leq)$ is said to be a **directed set** if
  1. $A \neq \varnothing$
  2. $\leq$ is a direction on A

2.4.2 Subdirected Sets

Definition 2.4.2.1. Let $(A, \leq)$ be a proset and $B \subset A$. Then B is said to be

- **$\leq$-upward directed** if $(B, \leq|_B)$ is a directed set,
- **$\leq$-downard directed** if $(B, \leq^*|_B)$ is a directed set.

Note 2.4.2.2. When the context is clear, we may say "upward directed" and "downward directed" instead of "$\leq$-upward directed" and "$\leq$-downard directed" respectively.

Exercise 2.4.2.3. Let $(A, \leq)$ be a proset and $B \subset A$. Then

1. B is $\leq$-downward directed iff B is $\leq^*$-upward directed.
2. B is $\leq$-upward directed iff B is $\leq^*$-downward directed.

Proof.

1. We note that
$$\begin{aligned} B \text{ is } \leq\text{-downward directed} &\iff (B, \leq^*|_B) \text{ is a directed set} \\ &\iff B \text{ is } \leq^*\text{-upward directed.} \end{aligned}$$
2. We note that
$$\begin{aligned} B \text{ is } \leq\text{-upward directed} &\iff B \text{ is } (\leq^*)^*\text{-upward directed} && \text{ref ex in set theory section on binary hetero relations about converse r} \\ &\iff B \text{ is } \leq^*\text{-downward directed} && \text{ref previous part.} \end{aligned}$$

□

2.4.3 Products of Directed Sets

Exercise 2.4.3.1. Let $(A, \leq_A)$ and $(B, \leq_B)$ be directed sets. Then

1. $\leq_A \otimes \leq_B$ is a direction on $A \times B$,
2. $(A, \leq_A) \otimes (B, \leq_B)$ is a directed set.

Proof.

1. (a) Exercise **??** implies that $\leq_A \otimes \leq_B$ is a preorder of $A \times B$.
   (b) Let $(a_1, b_1), (a_2, b_2) \in A \times B$. Then there exist $a \in A$ and $b \in B$ such that $a_1, a_2 \leq_A a$ and $b_1, b_2 \leq_B b$. Hence $(a_1, b_1), (a_2, b_2) \leq_A \otimes \leq_B (a, b)$.

   Hence $\leq_A \otimes \leq_B$ is a direction on $A \times B$.
2. (a) Since $A \neq \varnothing$ and $B \neq \varnothing$, we have that $A \times B \neq \varnothing$.
   (b) From above, $\leq_A \otimes \leq_B$ is a direction on $A \times B$.

   Hence $(A \times B, \leq_A \otimes \leq_B)$ is a directed set.

□

2.5 Monotone Maps

2.5.1 Introduction

Definition 2.5.1.1. Let $(X, \leq_X)$ $(Y, \leq_Y)$ be prosets and $f : X \to Y$. Then f is said to be

- $(\leq_X, \leq_Y)$**-order preserving** or $(\leq_X, \leq_Y)$**-monotone** if for each $a, b \in X$, $a \leq_X b$ implies that $f(a) \leq_Y f(b)$.
- $(\leq_X, \leq_Y)$**-order reversing** or $(\leq_X, \leq_Y)$**-antitone** if for each $a, b \in X$, $a \leq_X b$ implies that $f(b) \leq_Y f(a)$.

Note 2.5.1.2. When the context is clear we say that f is "monotone" and "antitone" instead of "$(\leq_X, \leq_Y)$-monotone" and "$(\leq_X, \leq_Y)$-antitone" respectively.

Exercise 2.5.1.3. Let $(X, \leq_X)$ $(Y, \leq_Y)$ be prosets and $f : X \to Y$. Then

1. the folloing are equivalent:
   (a) f is $(\leq_X, \leq_Y)$-antitone
   (b) f is $(\leq_X, \leq_Y^*)$-monotone
   (c) f is $(\leq_X^*, \leq_Y)$-monotone
   (d) f is $(\leq_X^*, \leq_Y^*)$-antitone
2. the folloing are equivalent:
   (a) f is $(\leq_X, \leq_Y)$-monotone
   (b) f is $(\leq_X, \leq_Y^*)$-antitone
   (c) f is $(\leq_X^*, \leq_Y)$-antitone
   (d) f is $(\leq_X^*, \leq_Y^*)$-monotone

Proof.

1.
   - $(a) \implies (b)$:
     Suppose that f is $(\leq_X, \leq_Y)$-antitone. Let $a, b \in X$. Then
     $$\begin{aligned} a \leq_X b &\implies f(b) \leq_Y f(a) \\ &\iff f(a) \leq_Y^* f(b). \end{aligned}$$
     Since $a, b \in X$ are arbitrary, we have that for each $a, b \in X$, $a \leq_X b$ implies that $f(a) \leq_Y^* f(b)$. Hence f is $(\leq_X, \leq_Y^*)$-monotone.
   - $(b) \implies (c)$:
     Suppose that f is $(\leq_X, \leq_Y^*)$-monotone. Let $a, b \in X$. Then
     $$\begin{aligned} a \leq_X^* b &\iff b \leq_X^* a \\ &\implies f(b) \leq_Y^* f(a) \\ &\iff f(a) \leq_Y f(b). \end{aligned}$$
     Since $a, b \in X$ are arbitrary, we have that for each $a, b \in X$, $a \leq_X^* b$ implies that $f(a) \leq_Y f(b)$. Hence f is $(\leq_X^*, \leq_Y)$-monotone.
   - $(c) \implies (d)$:
     Suppose that f is $(\leq_X^*, \leq_Y)$-monotone. Let $a, b \in X$. Then
     $$\begin{aligned} a \leq_X^* b &\implies f(a) \leq_Y f(b) \\ &\iff f(b) \leq_Y^* f(a). \end{aligned}$$
     Since $a, b \in X$ are arbitrary, we have that for each $a, b \in X$, $a \leq_X^* b$ implies that $f(b) \leq_Y^* f(a)$. Hence f is $(\leq_X^*, \leq_Y^*)$-antitone.

- $(d) \implies (a)$:
  Suppose that f is $(\leq_X^*, \leq_Y^*)$-antitone. Let $a, b \in X$. Then

$$\begin{aligned} a \leq_X b &\iff b \leq_X^* a \\ &\implies f(a) \leq_Y^* f(b) \\ &\iff f(b) \leq_Y f(a). \end{aligned}$$

  Since $a, b \in X$ are arbitrary, we have that for each $a, b \in X$, $a \leq_X b$ implies that $f(b) \leq_Y f(a)$. Hence f is $(\leq_X, \leq_Y)$-antitone.

2. Part (1) implies that

$$\begin{aligned} f \text{ is } (\leq_X, (\leq_Y^*)^*)\text{-monotone} &\iff f \text{ is } (\leq_X, \leq_Y^*)\text{-antitone} \quad (\text{by } 1(b) \iff 1(a)) \\ &\iff f \text{ is } (\leq_X^*, (\leq_Y^*)^*)\text{-antitone} \quad (\text{by } 1(a) \iff 1(d)) \\ &\iff f \text{ is } (\leq_X^*, \leq_Y^*)\text{-monotone} \quad (\text{by } 1(d) \iff 1(c)). \end{aligned}$$

   Since $(\leq_Y^*)^* = \leq_Y$, we have that

$$\begin{aligned} f \text{ is } (\leq_X, \leq_Y)\text{-monotone} &\iff f \text{ is } (\leq_X, \leq_Y^*)\text{-antitone} \\ &\iff f \text{ is } (\leq_X^*, \leq_Y)\text{-antitone} \\ &\iff f \text{ is } (\leq_X^*, \leq_Y^*)\text{-monotone.} \end{aligned}$$

□

Exercise 2.5.1.4. Let $(X, \leq_X)$ $(Y, \leq_Y)$ be posets and $f : X \to Y$. Then f is $(\leq_X, \leq_Y)$-monotone iff f is $(\leq_X^*, \leq_Y^*)$-monotone.

Proof.

- $(\implies)$:
  Suppose that f is $(\leq_X, \leq_Y)$-monotone. Let $a, b \in X$. Suppose that $a \leq_X^* b$. Then $b \leq_X a$. Hence $f(b) \leq_Y f(a)$. Thus $f(a) \leq_Y^* f(b)$. Since $a, b \in X$ with $a \leq_X^* b$ are arbitrary, we have that for each $a, b \in X$, $a \leq_X^* b$ implies that $f(a) \leq_Y^* f(b)$. Therefore f is $(\leq_X^*, \leq_Y^*)$-monotone.

- $(\impliedby)$:
  Suppose that f is $(\leq_X^*, \leq_Y^*)$-monotone. Since $(\leq_X^*)^* = \leq_X$, the previous part implies that f is $(\leq_X, \leq_Y)$-monotone.

□

2.5.2 Galois Connections

ToDo: If this comes after groups rings, etc, give examples of defining the 'algebraic object generated by less structured objects' in terms of a galois connection. e.g. the subgroup/subring/upperset/sublattice/ideal/filter/sigma-alg/topology generated, subsets of set/powerset

Definition 2.5.2.1. Let $(X, \leq_X)$ and $(Y, \leq_Y)$ be posets, $f : X \to Y$ and $g : Y \to X$. Then

- (f, g) is said to be a $(\leq_X, \leq_Y)$**-monotone Galois connection** if for each $x \in X$ and $y \in Y$,

$$f(x) \leq_Y y \text{ iff } x \leq_X g(y),$$

- (f, g) is said to be a $(\leq_X, \leq_Y)$**-antitone Galois connection** if for each $x \in X$ and $y \in Y$,

$$f(x) \leq_Y y \text{ iff } g(y) \leq_X x,$$

- f is said to be a $(\leq_X, \leq_Y)$**-left adjoint of** g if (f, g) is a $(\leq_X, \leq_Y)$-monotone Galois connection,
- g is said to be a $(\leq_X, \leq_Y)$**-right adjoint of** f if (f, g) is a $(\leq_X, \leq_Y)$-monotone Galois connection.

- f is said to **have a $(\leq_X, \leq_Y)$-right adjoint** if there exists $f^* : Y \to X$ such that f_* is a $(\leq_X, \leq_Y)$-right adjoint of f.
- g is said to **have a $(\leq_X, \leq_Y)$-left adjoint** if there exists $g_* : X \to Y$ such that g^* is a $(\leq_X, \leq_Y)$-left adjoint of g.

Note 2.5.2.2. When the context is clear, we say

- "monotone (resp. antitone) Galois connection" instead of "$(\leq_X, \leq_Y)$-monotone (resp. $(\leq_X, \leq_Y)$-antitone) Galois connection"
- "left (resp. right) adjoint" instead of "$(\leq_X, \leq_Y)$-left (resp. $(\leq_X, \leq_Y)$-right) adjoint"

Question:
There is an interpretation of prosets, monotonic maps and galois connections in terms of information.

- Prosets are spaces that encode informational direction: $x_1 \leq_X x_2$ iff x_1 contains less information than x_2.
- Monotonic maps are those that preserve informational direction: if x_1 has less information than x_2, then $f(x_1)$ has less information than $f(x_2)$.
- Galois connections encode informational direction in different contexts as well as minimal and maximal changes in information: $f(x)$ has less information than y in the context of Y. More specifically,
  - * y contains at least as much information as x in the context of Y iff $f(x) \leq_Y y$.
    * $f(x)$ is the element y of Y such that
      1. y contains at least as much information as x in the context of Y, i.e. $f(x) \leq_Y y$.
      2. for each $y' \in Y$, if y contains at least as much information as x in the context of Y, then y' contains at least as much information as y in the context of Y, i.e. $f(x) \leq_Y y' \implies y \leq_X y'$.

      So that $f(x)$ is the object in Y such that the increase in information (information gain) from x to $f(x)$ in the context of Y is minimal. Equivalently, the change in information $\Delta_Y I(x, f(x))$ is minimal)
  - * x has less information than y in the context of X iff $x \leq_X g(y)$.
    * $g(y)$ is the element x of X such that
      1. y contains at least as much information as x in the context of X, i.e. $x \leq_X g(y)$.
      2. for each $x' \in X$, if y contains at least as much information as x' in the context of X, then x contains at least as much information as x' in the context of X, i.e. $x' \leq_X g(y) \implies x' \leq_X x$.

      So that $g(y)$ is the object in X such that the decrease in information (information loss) from y to $g(y)$ in the context of X is minimal. Equivalently, the change in information $\Delta_X I(y, g(y))$ is maximal (-1 is larger than -2)

This can be generalized to categories, functors and adjoint functors where

- $x_1 \leq_X x_2$ generalizes to $\mathrm{Hom}_X(x_2, x_1) \neq \varnothing$, i.e. x_2 has at least as much information as x_1 generalizes to x_1 is a processed version of x_1,
- a monotone map $f : X \to Y$ generalizes to a functor $F : X \to Y$
- a Galois connection (f, g) generalizes to an adjunction (F, G).

This categorical generalization also generalizes information theory and statistics. However, from most math sources, the typical language of adjunctions is in terms of efficiency instead of information change. For example $f(x)$ is the most 'efficient' way to obtain an object in Y from x which 'preserves x'. Similarly, $g(y)$ is the least wasteful way to obtain an object in X from y which ?????????? Could we rephrase this to describe dynamical systems, specifically, can we rephrase information or efficiency in terms of energy to capture describe physics? Something like the path in phase space from t_1 to t_2 is a right adjoint functor from the universe (a category) at a time t_1 to the universe at time t_2 so that the information loss is minimal. Maybe show that least action, or newtons equation is a manifestation of this.

Exercise 2.5.2.3. Let $(X, \leq_X)$ and $(Y, \leq_Y)$ be prosets, $f : X \to Y$ and $g : Y \to X$. Then

1. (f, g) is a $(\leq_X, \leq_Y)$-monotone Galois connection iff (g, f) is a $(\leq_Y^*, \leq_X^*)$-monotone Galois connection
2. (f, g) is a $(\leq_X, \leq_Y)$-antitone Galois connection iff (f, g) is a $(\leq_X^*, \leq_Y)$-monotone Galois connection

3. (f, g) is a $(\leq_X, \leq_Y)$-antitone Galois connection iff (g, f) is a $(\leq_Y, \leq_X)$-antitone Galois connection

Proof.

1. - $(\implies)$:
   Suppose that (f, g) is a $(\leq_X, \leq_Y)$-monotone Galois connection. Let $y \in Y$ and $x \in X$. Then by assumption,
   $$\begin{aligned} g(y) \leq_X^* x &\iff x \leq_X g(y) \\ &\iff f(x) \leq_Y y \\ &\iff y \leq_Y^* f(x). \end{aligned}$$
   Since $y \in Y$ and $x \in X$ are arbitrary, we have that for each $y \in Y$ and $x \in X$, $g(y) \leq_X^* x$ iff $y \leq_Y^* f(x)$. Therefore (g, f) is a $(\leq_Y^*, \leq_X^*)$-monotone Galois connection.
   - $(\impliedby)$:
   Suppose that (g, f) is a $(\leq_Y^*, \leq_X^*)$-monotone Galois connection. Let $x \in X$ and $y \in Y$. Then by assumption,
   $$\begin{aligned} f(x) \leq_Y y &\iff y \leq_Y^* f(x) \\ &\iff g(y) \leq_X^* x \\ &\iff x \leq_X g(y). \end{aligned}$$
   Since $x \in X$ and $y \in Y$ are arbitrary, we have that for each $x \in X$ and $y \in Y$, $f(x) \leq_Y y$ iff $x \leq_X g(y)$. Therefore (f, g) is a $(\leq_X, \leq_Y)$-monotone Galois connection.

2. - $(\implies)$:
   Suppose that (f, g) is a $(\leq_X, \leq_Y)$-antitone Galois connection. Let $x \in X$ and $y \in Y$. Then by assumption,
   $$\begin{aligned} f(x) \leq_Y y &\iff g(y) \leq_X x \\ &\iff x \leq_X^* g(y). \end{aligned}$$
   Since $x \in X$ and $y \in Y$ are arbitrary, we have that for each $x \in X$ and $y \in Y$, $f(x) \leq_Y y$ iff $x \leq_X^* g(y)$. Therefore (f, g) is a $(\leq_X^*, \leq_Y)$-monotone Galois connection.
   - $(\impliedby)$:
   Suppose that (f, g) is a $(\leq_X^*, \leq_Y)$-monotone Galois connection. Let $x \in X$ and $y \in Y$. Then by assumption,
   $$\begin{aligned} f(x) \leq_Y y &\iff x \leq_X^* g(y) \\ &\iff g(y) \leq_X x. \end{aligned}$$
   Since $x \in X$ and $y \in Y$ are arbitrary, we have that for each $x \in X$ and $y \in Y$, $f(x) \leq_Y y$ iff $g(y) \leq_X x$. Therefore (f, g) is a $(\leq_X, \leq_Y)$-antitone Galois connection.

3. By (1) and (2),
   $$\begin{aligned} (f, g) \text{ is a } (\leq_X, \leq_Y)\text{-antitone Galois connection} &\iff (f, g) \text{ is a } (\leq_X^*, \leq_Y)\text{-monotone Galois connection} \\ &\iff (g, f) \text{ is a } (\leq_Y^*, (\leq_X^*)^*)\text{-monotone Galois connection} \\ &\iff (g, f) \text{ is a } (\leq_Y^*, \leq_X)\text{-monotone Galois connection} \\ &\iff (g, f) \text{ is a } (\leq_Y, \leq_X)\text{-antitone Galois connection.} \end{aligned}$$

□

Exercise 2.5.2.4. Let $(X, \leq_X)$ and $(Y, \leq_Y)$ be prosets, $f : X \to Y$ and $g : Y \to X$. Then f is a $(\leq_X, \leq_Y)$-left adjoint of g iff g is a $(\leq_X, \leq_Y)$-right adjoint of f.

Proof. We have that
$$\begin{aligned} f \text{ is a } (\leq_X, \leq_Y)\text{-left adjoint of } g &\iff (f, g) \text{ is a } (\leq_X, \leq_Y)\text{-monotone Galois connection} \\ &\iff g \text{ is a } (\leq_X, \leq_Y)\text{-right adjoint of } f. \end{aligned}$$

□

Exercise 2.5.2.5. Let $(X, \leq_X)$ and $(Y, \leq_Y)$ be prosets, $f : X \to Y$ and $g : Y \to X$. Then

1. g is a $(\leq_X, \leq_Y)$-right adjoint of f iff g is a $(\leq_Y^*, \leq_X^*)$-left adjoint of f.
2. f is a $(\leq_X, \leq_Y)$-left adjoint of g iff f is a $(\leq_Y^*, \leq_X^*)$-right adjoint of g.

Proof.

1. Exercise 2.5.2.3 implies that that

$$\begin{aligned} g \text{ is a } (\leq_X, \leq_Y)\text{-right adjoint of } f &\iff (f, g) \text{ is a } (\leq_X, \leq_Y)\text{-monotone Galois connection} \\ &\iff (g, f) \text{ is a } (\leq_Y^*, \leq_X^*)\text{-monotone Galois connection} \\ &\iff g \text{ is a } (\leq_Y^*, \leq_X^*)\text{-left adjoint of } f. \end{aligned}$$

2. We have that

$$\begin{aligned} f \text{ is a } (\leq_X, \leq_Y)\text{-left adjoint of } g &\iff g \text{ is a } (\leq_X, \leq_Y)\text{-right adjoint of } f \quad \text{(by Exercise 2.5.2.4)} \\ &\iff g \text{ is a } (\leq_Y^*, \leq_X^*)\text{-left adjoint of } f \quad \text{(by prev part)} \\ &\iff f \text{ is a } (\leq_Y^*, \leq_X^*)\text{-right adjoint of } g \quad \text{(by Exercise 2.5.2.4).} \end{aligned}$$

□

Exercise 2.5.2.6. Let $(X, \leq_X)$ and $(Y, \leq_Y)$ be prosets, $f : X \to Y$ and $g : Y \to X$. Then

1. (f, g) is a $(\leq_X, \leq_Y)$-monotone Galois connection iff
   (a) i. $\mathrm{id}_X \leq_X g \circ f$
   ii. $f \circ g \leq_Y \mathrm{id}_Y$
   (b) f is $(\leq_X, \leq_Y)$-monotone and g is $(\leq_Y, \leq_X)$-monotone
2. (f, g) is a $(\leq_X, \leq_Y)$-antitone Galois connection iff
   (a) i. $g \circ f \leq_X \mathrm{id}_X$
   ii. $f \circ g \leq_Y \mathrm{id}_Y$
   (b) f is $(\leq_X, \leq_Y)$-antitone and g is $(\leq_Y, \leq_X)$-antitone

Proof.

1. • ($\Longrightarrow$):
   Suppose that (f, g) us a $(\leq_X, \leq_Y)$-monotone Galois connection.
   (a) i. Let $x \in X$. Define $y \in Y$ by $y := f(x)$. Since $f(x) \leq_Y f(x)$, we have that $f(x) \leq_Y y$. Therefore

$$\begin{aligned} x &\leq_X g(y) \\ &= g \circ f(x). \end{aligned}$$

   Since $x \in X$ is arbitrary, we have that for each $x \in X$, $x \leq_X g \circ f(x)$. Hence $\mathrm{id}_X \leq_X g \circ f$.
   ii. Let $y \in Y$. the previous ex implies that (g, f) is a $(\leq_Y^*, \leq_X^*)$-monotone Galois connection. Part 1(a) then implies that for each $y \in Y$, $y \leq_Y^* f \circ g(y)$. Therefore for each $y \in Y$, $f \circ g(y) \leq_Y y$. Hence $f \circ g \leq_Y \mathrm{id}_Y$.
   (b) – Let $a, b \in X$. Define $x \in X$ and $y \in Y$ by $x := a$ and $y := f(b)$. Suppose that $a \leq_X b$. Part 1(a) implies that

$$\begin{aligned} x &= a \\ &\leq_X b \\ &\leq_X g(f(x)) \\ &= g(y). \end{aligned}$$

Since (f, g) is a $(\leq_X, \leq_Y)$-monotone Galois connection, we have that

$$\begin{aligned} f(a) &= f(x) \\ &\leq_Y y \\ &= f(b). \end{aligned}$$

Since $a, b \in X$ with $a \leq_X b$ are arbitrary, we have that for each $a, b \in X$, $a \leq_X b$ implies that $f(a) \leq_Y f(b)$. Thus f is $(\leq_X, \leq_Y)$-monotone.

– Since (f, g) is a $(\leq_X, \leq_Y)$-monotone Galois connection, a previous ex implies that (g, f) is a $(\leq_Y^*, \leq_X^*)$-monotone Galois connection. The previous part implies that g is $(\leq_Y^*, \leq_X^*)$-monotone. Exercise 2.5.1.3 then implies that g is $(\leq_Y, \leq_X)$-monotone.

- $(\Longleftarrow)$:
  Suppose that
  (a) i. for each $x \in X$, $x \leq_X g \circ f(x)$
  ii. for each $y \in Y$, $f \circ g(y) \leq_Y y$
  (b) f is $(\leq_X, \leq_Y)$-monotone and g is $(\leq_Y, \leq_X)$-monotone.

  Let $x \in X$ and $y \in Y$.

  – $(\Longrightarrow)$:
  Suppose that $f(x) \leq_Y y$. Then by (a)i and (a)ii,

  $$\begin{aligned} x &\leq_X g \circ f(x) \\ &\leq_X g(y). \end{aligned}$$

  – $(\Longleftarrow)$:
  Suppose that $x \leq_X g(y)$. Then by (a)i and (a)ii,

  $$\begin{aligned} f(x) &\leq_Y f \circ g(y) \\ &\leq_Y y. \end{aligned}$$

  Since $x \in X$ and $y \in Y$ are arbitrary, we have that for each $x \in X$ and $y \in Y$,

  $$f(x) \leq_Y y \text{ iff } f(x) \leq_Y y.$$

  Therefore (f, g) is a $(\leq_X, \leq_Y)$-monotone Galois connection.

2. the previous ex implies that (f, g) is a $(\leq_X, \leq_Y)$-antitone Galois connection iff (f, g) is a $(\leq_X^*, \leq_Y)$-monotone Galois connection. Part (1) implies that (f, g) is a $(\leq_X^*, \leq_Y)$-monotone Galois connection iff

   (a) i. for each $x \in X$, $x \leq_X^* g \circ f(x)$
   ii. for each $y \in Y$, $f \circ g(y) \leq_Y y$
   (b) f is $(\leq_X^*, \leq_Y)$-monotone and g is $(\leq_Y, \leq_X^*)$-monotone.

   two previous exercise about monotonic functions and the dual order then imply that (f, g) is a $(\leq_X, \leq_Y)$-antitone Galois connection iff

   (a) i. for each $x \in X$, $g \circ f(x) \leq_X x$
   ii. for each $y \in Y$, $f \circ g(y) \leq_Y y$
   (b) f is $(\leq_X, \leq_Y)$-antitone and g is $(\leq_Y, \leq_X)$-antitone.

□

Exercise 2.5.2.7. Let $(X, \leq_X)$ and $(Y, \leq_Y)$ be prosets, $f : X \to Y$ and $g : Y \to X$. If (f, g) is a $(\leq_X, \leq_Y)$-monotone Galois connection or (f, g) is a $(\leq_X, \leq_Y)$-antitone Galois connection, then $f \sim_{\leq_Y} f \circ g \circ f$ and $g \sim_{\leq_X} g \circ f \circ g$.

Proof.

- Suppose that (f, g) a $(\leq_X, \leq_Y)$-monotone Galois connection.
  - Let $x \in X$.
    - Define $a \in X$ and $b \in Y$ by $a := g \circ f(x)$ and $b := f(x)$. Since

$$\begin{aligned} g \circ f(x) \leq g \circ f(x) &\iff a \leq_X g(b) \\ &\iff f(a) \leq_Y b \\ &\iff f \circ g \circ f(x) \leq_Y f(x), \end{aligned}$$

      we have that $f \circ g \circ f(x) \leq_Y f(x)$.
    - The prev ex implies that $x \leq_X g \circ f(x)$ and f is $(\leq_X, \leq_Y)$-monotone. Therefore $f(x) \leq_Y f \circ g \circ f(x)$.

    Since $f \circ g \circ f(x) \leq_Y f(x)$ and $f(x) \leq_Y f \circ g \circ f(x)$, we have that $f \circ g \circ f(x) \sim_{\leq_Y} f(x)$, Since $x \in X$ is arbitrary, we have that for each $x \in X$, $f \circ g \circ f(x) \sim_{\leq_Y} f(x)$. Therefore $f \sim_{\leq_Y} f \circ g \circ f$.
  - Let $y \in Y$.
    - Define $a \in X$ and $b \in Y$ by $a := g(y)$ and $b := f \circ g(y)$. Since

$$\begin{aligned} f \circ g(x) \leq_Y f \circ g(x) &\iff f(a) \leq_Y b \\ &\iff a \leq_X g(b) \\ &\iff g(y) \leq_X g \circ f \circ g(y), \end{aligned}$$

      we have that $g(y) \leq_X g \circ f \circ g(y)$.
    - The prev ex implies that $f \circ g(y) \leq_Y y$ and g is $(\leq_Y, \leq_X)$-monotone. Therefore $g \circ f \circ g(y) \leq_X g(y)$.

    Since $g(y) \leq_X g \circ f \circ g(y)$ and $g \circ f \circ g(y) \leq_X g(y)$, we have that $g(y) \sim_{\leq_X} g \circ f \circ g(y)$, Since $y \in Y$ is arbitrary, we have that for each $y \in Y$, $g(y) \sim_{\leq_X} g \circ f \circ g(y)$. Therefore $g \sim_{\leq_X} g \circ f \circ g$.
- Suppose that (f, g) a $(\leq_X, \leq_Y)$-antitone Galois connection. The prev ex then implies that (f, g) is a $(\leq_X^*, \leq_Y)$-monotone Galois connection. Part (1) then implies that $f \sim_{\leq_Y} f \circ g \circ f$ and $g \sim_{\leq_X^*} g \circ f \circ g$. Make previous ex that $x \sim_{\leq} y$ iff $x \sim_{\leq^*} y$ then implies that $g \sim_{\leq_X} g \circ f \circ g$.

FINSIH!! □

Exercise 2.5.2.8. Let $(X, \leq_X)$ and $(Y, \leq_Y)$ be posets, $f : X \to Y$ and $g : Y \to X$. If (f, g) is a $(\leq_X, \leq_Y)$-monotone Galois connection or (f, g) is a $(\leq_X, \leq_Y)$-antitone Galois connection, then $f = f \circ g \circ f$ and $g = g \circ f \circ g$.

Proof. Sppose that (f, g) is a $(\leq_X, \leq_Y)$-monotone Galois connection or (f, g) is a $(\leq_X, \leq_Y)$-antitone Galois connection. a prev ex implies that $f \sim_{\leq_Y} f \circ g \circ f$ and $g \sim_{\leq_X} g \circ f \circ g$. Since $(X, \leq_X)$ and $(Y, \leq_Y)$ are posets, we have that $f = f \circ g \circ f$ and $g = g \circ f \circ g$. □

Exercise 2.5.2.9. Let $(X, \leq_X)$ and $(Y, \leq_Y)$ be posets, $f : X \to Y$ and $g : Y \to X$.

1. For each $g' : Y \to X$, if g and g' are right adjoints of f, then $g = g'$.
2. For each $f' : X \to Y$, if f and f' are left adjoints of g, then $f = f'$.

Proof.

1. Let $g' : Y \to X$. Suppose that g and g' are right adjoints of f. hen (f, g) and (f, g') are Galois connections. Exercise 2.5.2.8 implies that $f \circ g' \circ f = f$ and for each $x \in X$, $g' \circ f(x) \leq_Y g \circ f(g' \circ f(x))$. Therefore for each $x \in X$,

$$\begin{aligned} g' \circ f(x) &\leq g \circ f(g' \circ f(x)) \\ &= g(f \circ g' \circ f(x)) \\ &= g(f(x)) \\ &= g \circ f(x). \end{aligned}$$

   Thus $g' \circ f \leq g \circ f$. Similarly, $g \circ f \leq g' \circ f$. Hence $g' \circ f = g \circ f$. Therefore

$$\begin{aligned} g &= g \circ f \circ g \\ &= g' \circ f \circ g. \end{aligned}$$

and

$$\begin{aligned} g' &= g' \circ f \circ g' \\ &= g \circ f \circ g'. \end{aligned}$$

Let $y \in Y$. Since $f \circ g'(y) \leq y$, we have that

$$\begin{aligned} g'(y) &= g \circ f \circ g'(y) \\ &= g(f \circ g'(y)) \\ &\leq g(y). \end{aligned}$$

Similarly, $g(y) \leq g'(y)$. Hence $g(y) = g'(y)$. Since $y \in Y$ is arbitrary, we have that for each $y \in Y$, $g(y) = g'(y)$. Thus $g = g'$.

2. Similar to (1).

TIDY UP!!!! □

Definition 2.5.2.10. Let $(X, \leq_X)$ and $(Y, \leq_Y)$ be posets, $f : X \to Y$ and $g : Y \to X$.

- If f has a $(\leq_X, \leq_Y)$-right adjoint, then we define the $(\leq_X, \leq_Y)$**-right adjoint of** f, denoted $\mathrm{RAdj}(f, (\leq_X, \leq_Y))$, to be the unique $g' : Y \to X$ such that g' is a $(\leq_X, \leq_Y)$-right adjoint of f.
- If g has a $(\leq_X, \leq_Y)$-left adjoint, then we define the $(\leq_X, \leq_Y)$**-left adjoint of** g, denoted $\mathrm{LAdj}(g, (\leq_X, \leq_Y))$, to be the unique $f' : X \to Y$ such that f' is a $(\leq_X, \leq_Y)$-left adjoint of g.

Note 2.5.2.11. When the context is clear, we write f^* and g_* in place of $\mathrm{RAdj}(f, (\leq_X, \leq_Y))$ and $\mathrm{LAdj}(g, (\leq_X, \leq_Y))$ respectively.

Exercise 2.5.2.12. Let $(X, \leq_X)$, $(Y, \leq_Y)$, $(Z, \leq_Z)$ be posets, $f : X \to Y$ and $g : Y \to Z$.

1. If f has a right adjoint and g has a right adjoint, then $g \circ f$ has a right adjoint and $(g \circ f)^* = f^* \circ g^*$.
2. If f has a left adjoint and g has a left adjoint, then $g \circ f$ has a left adjoint and $(g \circ f)_* = f_* \circ g_*$.

Proof.

1. Suppose that f has a $(\leq_X, \leq_Y)$-right adjoint and g has a $(\leq_X, \leq_Y)$-right adjoint. Since (f, f^*) is a $(\leq_X, \leq_Y)$-monotone Galois connection and (g, g^*) is a $(\leq_Y, \leq_Z)$-monotone Galois connection, we have that for each $x \in X$, $y \in Y$ and $z \in Z$, $x \leq_X f^*(y)$ iff $f(x) \leq_Y y$ and $y \leq_Y g^*(z)$ iff $g(y) \leq_Z z$. Hence for each $x \in X$ and $z \in Z$,

$$\begin{aligned} x \leq_X f^* \circ g^*(z) &\iff x \leq_X f^*(g^*(z)) \\ &\iff f(x) \leq_Y g^*(z) \\ &\iff g(f(x)) \leq_Z z \\ &\iff g \circ f(x) \leq_Z z. \end{aligned}$$

Thus $(g \circ f, f^* \circ g^*)$ is a $(\leq_X, \leq_Z)$-monotone Galois connection. Therefore $g \circ f$ has a $(\leq_X, \leq_Z)$-right adjoint and Exercise 2.5.2.9 implies that $(g \circ f)^* = f^* \circ g^*$.

2. Suppose that f has a $(\leq_Y, \leq_X)$-left adjoint and g has a $(\leq_Z, \leq_Y)$-left adjoint. Exercise 2.5.2.5 implies that $\mathrm{LAdj}(f, (\leq_Y, \leq_X)) = \mathrm{RAdj}(f, (\leq_X^*, \leq_Y^*))$ and $\mathrm{LAdj}(g, (\leq_Z, \leq_Y)) = \mathrm{RAdj}(g, (\leq_Y^*, \leq_Z^*))$. The previous part implies that $g \circ f$ has a $(\leq_X^*, \leq_Z^*)$-right adjoint and $\mathrm{RAdj}(g \circ f, (\leq_X^*, \leq_Z^*)) = \mathrm{LAdj}(f, (\leq_Y, \leq_X)) \circ \mathrm{LAdj}(g, (\leq_Z, \leq_Y))$. Exercise 2.5.2.5 then implies that $g \circ f$ has a $(\leq_Z, \leq_X)$-left adjoint and

$$\begin{aligned} \mathrm{LAdj}(g \circ f, (\leq_Z, \leq_X)) &= \mathrm{RAdj}(g \circ f, (\leq_X^*, \leq_Z^*)) \\ &= \mathrm{LAdj}(f, (\leq_Y, \leq_X)) \circ \mathrm{LAdj}(g, (\leq_Z, \leq_Y)). \end{aligned}$$

□

Exercise 2.5.2.13. Let X, Y be sets and $f : X \to Y$. We define $f_* : \mathcal{P}(X) \to \mathcal{P}(Y)$ and $f^* : \mathcal{P}(Y) \to \mathcal{P}(X)$ by $f_*(A) := f(A)$ and $f^*(B) := f^{-1}(B)$. Then (f_*, f^*) is a $(\subset_X, \subset_Y)$-monotone Galois connection.

Proof. We note that for each $A \in \mathcal{P}(X)$ and $B \in \mathcal{P}(Y)$,

$$\begin{aligned} f_*(A) \subset_Y B &\iff f(A) \subset_Y B \\ &\iff f^{-1}(f(A)) \subset_X f^{-1}(B) \quad \text{by (ref ex in set theory notes)} \\ &\iff A \subset_X f^{-1}(B) \quad \text{by (ref ex in set theory notes)} \\ &\iff A \subset_X f^*(B). \end{aligned}$$

Thus (f_*, f^*) is a $(\subset_X, \subset_Y)$-monotone Galois connection. □

Definition 2.5.2.14. maybe get rid of, ask if there is any point in rephrasing/renaming, possibly move right after definition of gallois connections, with discussion on 'information' Let $(X, \leq_X)$ and $(Y, \leq_Y)$ be posets, $f : X \to Y$ and $g : Y \to X$. Suppose that (f, g) is a $(\leq_X, \leq_Y)$-monotone Galois connection.

- Let $x \in X$. We define the **right** (f, g)**-connection of** x, denoted $\mathrm{RConn}_Y(x, (f, g)) \in Y$, by $\mathrm{RConn}_Y(x, (f, g)) := f(x)$.
- Let $y \in Y$. We define the **left** (f, g)**-connection of** y, denoted $\mathrm{LConn}_X(y, (f, g)) \in X$, by $\mathrm{LConn}_Y(x, (f, g)) := g(x)$.

2.6 Upper and Lower Sets

2.6.1 Introduction

Definition 2.6.1.1. Let $(X, \leq)$ be a proset and $A \subset X$. Then

- A is said to be a **$\leq$-upper set** if for each $a \in A$ and $x \in X$, $a \leq x$ implies that $x \in A$.
- A is said to be an **$\leq$-lower set** if for each $a \in A$ and $x \in X$, $x \leq a$ implies that $x \in A$.

Note 2.6.1.2. When the context is clear, we say A is a

- "upper set" instead of "$\leq$-upper set"
- "lower set" instead of "$\leq$-lower set"

Exercise 2.6.1.3. Let $(X, \leq)$ be a proset. Then

1. X is a $\leq$-upper set
2. X is a $\leq$-lower set

Proof.

1. Let $a, x \in X$. Suppose that $a \leq x$. By assumption, $x \in X$. Since $a, x \in X$ with $a \leq x$ are arbitrary, we have that for each $a, x \in X$, $a \leq x$ implies that $x \in A$. Hence X is a $\leq$-upper set.
2. Similar to (1).

□

Exercise 2.6.1.4. Let $(X, \leq)$ be a proset and $A \subset X$. Then

1. A is a $\leq$-upper set iff A is a $\leq^*$-lower set
2. A is a $\leq$-lower set iff A is a $\leq^*$-upper set

Proof.

1.
   - $(\implies)$:
     Suppose that A is a $\leq$-upper set. Let $a \in A$ and $x \in X$. Suppose that $x \leq^* a$. Then $a \leq x$. Since A is a $\leq$-upper set, $x \in A$. Since $a \in A$ and $x \in X$ with $x \leq^* a$ is arbitrary, we have that for each $a \in A$ and $x \in X$, $x \leq^* a$ implies that $x \in A$. Hence A is a $\leq^*$-lower set.
   - $(\impliedby)$:
     Suppose that A is a $\leq^*$-lower set. Let $a \in A$ and $x \in X$. Suppose that $a \leq x$. Then $x \leq^* a$. Since A is a $\leq^*$-lower set, $x \in A$. Since $a \in A$ and $x \in X$ with $a \leq x$ is arbitrary, we have that for each $a \in A$ and $x \in X$, $a \leq x$ implies that $x \in A$. Hence A is a $\leq$-upper set.
2. Similar to (1). USE DUALITY

□

Exercise 2.6.1.5. Let $(X, \leq)$ be a proset and $A \subset X$. Then

1. A is an upper set iff A^c is a lower set
2. A is a lower set iff A^c is an upper set

Proof.

1.
   - $(\implies)$:
     Suppose that A is an upper set. Let $b \in A^c$ and $x \in X$. Suppose that $x \leq b$. For the sake of contradiction, suppose that $x \in A$. Since $x \leq b$ and A is an upper set, $b \in A$. This is a contradiction since $b \in A^c$. Hence $x \in A^c$. Since $b \in A^c$ and $x \in X$ with $x \leq b$ are arbitrary, we have that for each $b \in A^c$ and $x \in X$, $x \leq b$ implies that $x \in A^c$. Thus A^c is a lower set.

- ($\Longleftarrow$):
  Suppose that A^c is a $\leq$-lower set. Exercise 2.6.1.4 implies that A^c is a $\leq^*$-upper set. The previous part implies that A is a $\leq^*$-lower set. Another application of Exercise 2.6.1.4 implies that A is a $\leq$-upper set.

2. Similar to (1).

□

Exercise 2.6.1.6. Let $(X, \leq)$ be a proset, $(E_\alpha)_{\alpha \in A} \subset \mathcal{P}(X)$.

1. If for each $\alpha \in A$, E_α is an upper set, then
   (a) $\bigcup\limits_{\alpha \in A} E_\alpha$ is an upper set
   (b) $\bigcap\limits_{\alpha \in A} E_\alpha$ is an upper set
2. If for each $\alpha \in A$, E_α is a lower set, then
   (a) $\bigcup\limits_{\alpha \in A} E_\alpha$ is a lower set
   (b) $\bigcap\limits_{\alpha \in A} E_\alpha$ is an lower set

Proof.

1. Suppose that for each $\alpha \in A$, E_α is an upper set. Set $E := \bigcup\limits_{\alpha \in A} E_\alpha$.
   (a) Let $e \in E$ and $x \in X$. Suppose that $e \leq x$. Since $e \in E$, there exists $\alpha \in A$ such that $e \in E_\alpha$. Since E_α is an upper set and $e \leq x$, we have that
   $$\begin{aligned} x &\in E_\alpha \\ &\subset \bigcup_{\alpha \in A} E_\alpha \\ &= E. \end{aligned}$$
   Since $e \in E$ and $x \in X$ with $e \leq x$ are arbitrary, we have that for each $e \in E$ and $x \in X$, $e \leq x$ implies that $x \in E$. Hence E is an upper set.
   (b) Let $\alpha \in A$. Since E_α is a $\leq$-upper set, Exercise 2.6.1.5 implies that E_α^c is a $\leq$-lower set. Exercise 2.6.1.4 then implies that E_α^c is a $\leq^*$-upper set. Since $\alpha \in A$ is arbitrary, we have that for each $\alpha \in A$, E_α^c is a $\leq^*$-upper set. The previous part implies that $\bigcup\limits_{\alpha \in A} E_\alpha^c$ is a $\leq^*$-upper set. Since $\bigcap\limits_{\alpha \in A} E_\alpha = \left(\bigcup\limits_{\alpha \in A} E_\alpha^c \right)^c$, Exercise 2.6.1.5 implies that $\bigcap\limits_{\alpha \in A} E_\alpha$ is a $\leq^*$-lower set. Exercise 2.6.1.4 then implies that $\bigcap\limits_{\alpha \in A} E_\alpha$ is a $\leq$-upper set.
2. Similar to (1).

□

Definition 2.6.1.7. Let $(X, \leq)$ be a proset and $S \subset X$.(define upper set generated by- as the generated element from a galois connection, then make exercise showing that $\uparrow(S, \leq) := \bigcap\limits_{U \in \mathcal{U}(S, \leq)} U$) We define the

- **$\leq$-upper sets containing** S, denoted $\mathcal{U}(S, \leq) \subset \mathcal{P}(X)$, by $\mathcal{U}(S, \leq) := \{U \subset X : U \text{ is a } \leq\text{-upper set and } S \subset U\}$
- **$\leq$-lower sets containing** S, denoted $\mathcal{L}(S, \leq) \subset \mathcal{P}(X)$, by $\mathcal{L}(S, \leq) := \{L \subset X : L \text{ is a } \leq\text{-lower set and } S \subset L\}$.
- **$\leq$-upper set generated by** S, denoted $\uparrow(S, \leq) \subset X$, by $\uparrow(S, \leq) := \bigcap\limits_{U \in \mathcal{U}(S, \leq)} U$
- **$\leq$-lower set generated by** S, denoted $\downarrow(S, \leq) \subset X$, by $\downarrow(S, \leq) := \bigcap\limits_{L \in \mathcal{L}(S, \leq)} L$

Note 2.6.1.8.

- When the context is clear, we write $\mathcal{U}(S), \mathcal{L}(S), \uparrow S$ and $\downarrow S$ in place of $\mathcal{U}(S, \leq), \mathcal{L}(S, \leq), \uparrow(S, \leq)$ and $\downarrow(S, \leq)$ respectively.
- If $S = \{s\}$, we write $\uparrow s$ and $\downarrow s$ in place of $\uparrow S$ and $\downarrow S$ respectively.
- Exercise 2.6.1.3 implies that $X \in \mathcal{U}(S)$ and $X \in \mathcal{L}(S)$.
- Exercise 2.6.1.6 implies that $\uparrow S \in \mathcal{U}(S)$ and $\downarrow S \in \mathcal{L}(S)$.

Exercise 2.6.1.9. Let $(X, \leq)$ be a proset and $S \subset X$. Then

1. $\mathcal{U}(S, \leq^*) = \mathcal{L}(S, \leq)$,
2. $\mathcal{L}(S, \leq^*) = \mathcal{U}(S, \leq)$,
3. $\uparrow(S, \leq^*) = \downarrow(S, \leq)$,
4. $\downarrow(S, \leq^*) = \uparrow(S, \leq)$.

Proof.

1.
   - Let $L \in \mathcal{U}(S, \leq^*)$. Then $S \subset L$ and L is a $\leq^*$-upper set. Exercise 2.6.1.4 then implies that L is a $\leq$-lower set. Hence $L \in \mathcal{L}(S, \leq)$. Since $L \in \mathcal{U}(S, \leq^*)$ is arbitrary, we have that for each $L \in \mathcal{U}(S, \leq^*)$, $L \in \mathcal{L}(S, \leq)$. Thus $\mathcal{U}(S, \leq^*) \subset \mathcal{L}(S, \leq)$.
   - Let $L \in \mathcal{L}(S, \leq)$. Then $S \subset L$ and L is a $\leq$-lower set. Another application of Exercise 2.6.1.4 implies that L is a $\leq^*$-upper set. Hence $L \in \mathcal{U}(S, \leq^*)$. Since $L \in \mathcal{L}(S, \leq)$ is arbitrary, we have that for each $L \in \mathcal{L}(S, \leq)$, $L \in \mathcal{U}(S, \leq^*)$. Thus $\mathcal{L}(S, \leq) \subset \mathcal{U}(S, \leq^*)$.

   Since $\mathcal{U}(S, \leq^*) \subset \mathcal{L}(S, \leq)$ and $\mathcal{L}(S, \leq) \subset \mathcal{U}(S, \leq^*)$, we have that $\mathcal{U}(S, \leq^*) = \mathcal{L}(S, \leq)$.
2. By (1), we have that
$$\begin{aligned}\mathcal{L}(S, \leq^*) &= \mathcal{U}(S, (\leq^*)^*) \\ &= \mathcal{U}(S, \leq).\end{aligned}$$
3. Part (1) implies that
$$\begin{aligned}\uparrow(S, \leq^*) &= \bigcap_{U \in \mathcal{U}(S, \leq^*)} U \\ &= \bigcap_{U \in \mathcal{L}(S, \leq)} U \\ &= \downarrow(S, \leq^*).\end{aligned}$$
4. Similar to (3).

□

Exercise 2.6.1.10. Let $(X, \leq)$ be a proset and $S \subset X$. Then

1. $S \in \mathcal{U}(S, \leq)$ iff $\uparrow S = S$
2. $S \in \mathcal{L}(S, \leq)$ iff $\downarrow S = S$

Proof.

1.
   - ($\Longrightarrow$):
     Suppose that $S \in \mathcal{U}(S, \leq)$. Then
$$\begin{aligned}\uparrow S &= \bigcap_{U \in \mathcal{U}(S, \leq)} U \\ &\subset S \\ &\subset \uparrow S.\end{aligned}$$
     Hence $\uparrow S = S$.

- ($\Longleftarrow$):
  Suppose that $\uparrow S = S$. Then

$$\begin{aligned} S &= \uparrow S \\ &\in \mathcal{U}(S, \leq). \end{aligned}$$

2. Similar to (1).

□

Exercise 2.6.1.11. Let $(X, \leq)$ be a proset and $a \in X$. Then

1. $\uparrow a = \{x \in X : a \leq x\}$
2. $\downarrow a = \{x \in X : x \leq a\}$

Proof.

1. Define $U \subset X$ by $U := \{x \in X : a \leq x\}$.

   - Let $u \in U$ and $x \in X$. Suppose that $u \leq x$. Then

$$\begin{aligned} a &\leq u \\ &\leq x. \end{aligned}$$

     Hence $x \in U$. Since $u \in U$ and $x \in X$ with $u \leq x$ are arbitrary, we have that for each $u \in U$ and $x \in X$, $u \leq x$ implies that $x \in U$. Hence U is an upper set. Since $a \leq a$, $a \in U$. By definition, $U \in \mathcal{U}(a)$. Therefore

$$\begin{aligned} \uparrow a &= \bigcap_{U' \in \mathcal{U}(a)} U' \\ &\subset U. \end{aligned}$$

   - Let $x \in U$ and $U' \in \mathcal{U}(a)$. Since $x \in U$, $a \leq x$. Since $U' \in \mathcal{U}(a)$, U' is an upper set and $a \in U'$. Thus $x \in U'$. Since $U' \in \mathcal{U}(a)$ is arbitrary, we have that for each $U' \in \mathcal{U}(a)$, $x \in U'$. Thus

$$\begin{aligned} x &\in \bigcap_{U' \in \mathcal{U}(a)} U' \\ &= \uparrow a. \end{aligned}$$

     Since $x \in U$ is arbitrary, we have that for each $x \in U$, $x \in \uparrow a$. Hence $U \subset \uparrow a$.

   Since $\uparrow a \subset U$ and $U \subset \uparrow a$, we have that $\uparrow a = U$.

2. Exercise 2.6.1.4 and (1) imply that

$$\begin{aligned} \downarrow(a, \leq) &= \uparrow(a, \leq^*) \\ &= \{x \in X : a \leq^* x\} \\ &= \{x \in X : x \leq a\}. \end{aligned}$$

□

Exercise 2.6.1.12. Let $(X, \leq)$ be a proset and $a, b \in X$.

1. If $\uparrow a = \uparrow b$, then $a \sim_{\leq} b$
2. If $\downarrow a = \downarrow b$, then $a \sim_{\leq} b$

follow up later with one about posets

Proof.

1. Suppose that $\uparrow a = \uparrow b$. Then

$$\begin{aligned} a &\in \uparrow a \\ &= \uparrow b. \end{aligned}$$

   Hence $a \leq b$. Similarly, $b \leq a$. Hence $a \sim_{\leq} b$.

2.

□

Exercise 2.6.1.13. Let $(X, \leq)$ be a poset and $a, b \in X$.

1. If $\uparrow a = \uparrow b$, then $a = b$
2. If $\downarrow a = \downarrow b$, then $a = b$

Proof.

1. Suppose that $\uparrow a = \uparrow b$. Exercise 2.6.1.12 then implies that $a \sim_{\leq} b$. Since $(X, \leq)$ is a poset, we have that $a = b$.
2. Similar to (1).

□

Exercise 2.6.1.14. Let $(X, \leq)$ be a proset and $S_1, S_2 \subset X$. If $S_1 \subset S_2$, then

1. $\uparrow S_1 \subset \uparrow S_2$
2. $\downarrow S_1 \subset \downarrow S_2$

Proof. Suppose that $S_1 \subset S_2$.

1. Since $\uparrow S_2 \in \mathcal{U}(S_2)$, $S_2 \subset \uparrow S_2$ and $\uparrow S_2$ is an upper set. Since

$$\begin{aligned} S_1 &\subset S_2 \\ &\subset \uparrow S_2, \end{aligned}$$

   we have that $\uparrow S_2 \in \mathcal{U}(S_1)$. Therefore

$$\begin{aligned} \uparrow S_1 &= \bigcap_{U \in \mathcal{U}(S_1)} U \\ &\subset \uparrow S_2. \end{aligned}$$

2. Part (1) and Exercise 2.6.1.9 implies that

$$\begin{aligned} \downarrow(S_1, \leq) &= \uparrow(S_1, \leq^*) \\ &\subset \uparrow(S_2, \leq^*) \\ &= \downarrow(S_2, \leq). \end{aligned}$$

□

Exercise 2.6.1.15. Let $(X, \leq)$ be a proset and $a, b \in X$. Then the following are equivalent:

1. $a \leq b$,
2. $\uparrow b \subset \uparrow a$,
3. $\downarrow a \subset \downarrow b$.

Proof.

1. (1) $\implies$ (2):
   Suppose that $a \leq b$. Since $\uparrow a \in \mathcal{U}(a)$, we have that $a \in \uparrow a$ and $\uparrow a$ is an upper set. Since $a \leq b$, we have that $b \in \uparrow a$. Hence $\uparrow a \in \mathcal{U}(b)$. Therefore
   $$\begin{aligned} \uparrow b &= \bigcap_{U \in \mathcal{U}(b)} U \\ &\subset \uparrow a. \end{aligned}$$

2. (2) $\implies$ (3):
   Suppose that $\uparrow(b, \leq) \subset \uparrow(a, \leq)$. Exercise 2.6.1.11 then implies that
   $$\begin{aligned} b &\in \uparrow(b, \leq) \\ &\subset \uparrow(a, \leq) \\ &= \{x \in X : a \leq x\}. \end{aligned}$$
   Hence $a \leq b$. Thus $b \leq^* a$. Exercise 2.6.1.9 and part (1) $\implies$ (2) then imply that
   $$\begin{aligned} \downarrow(a, \leq) &= \uparrow(a, \leq^*) \\ &\subset \uparrow(b, \leq^*) \\ &= \downarrow(b, \leq). \end{aligned}$$

3. (3) $\implies$ (1):
   Suppose that $\downarrow a \subset \downarrow b$. Exercise 2.6.1.11 then implies that
   $$\begin{aligned} a &\in \downarrow a \\ &\subset \downarrow b \\ &= \{x \in X : x \leq b\}. \end{aligned}$$
   Hence $a \leq b$.

□

Exercise 2.6.1.16. Let $(X, \leq)$ be a proset and $S \subset X$. Then

1. $\uparrow S = \bigcup_{s \in S} \uparrow s$
2. $\downarrow S = \bigcup_{s \in S} \downarrow s$

Proof.

1. Define $U \subset X$ by $U := \bigcup_{s \in S} \uparrow s$.
   - Since for each $s \in S$, $\uparrow s$ is an upper set, Exercise 2.6.1.6 implies that U is an upper set. Let $s \in S$. Then
     $$\begin{aligned} s &\in \uparrow s \\ &\subset \bigcup_{s \in S} \uparrow s \\ &= U. \end{aligned}$$
     Since $s \in S$ is arbitrary, we have that for each $s \in S$, $s \in U$. Hence $S \subset U$. Therefore $U \in \mathcal{U}(S)$ and
     $$\begin{aligned} \uparrow S &= \bigcap_{U' \in \mathcal{U}(S)} U' \\ &\subset U. \end{aligned}$$

- Let $x \in U$ and $U' \in \mathcal{U}(S)$. Since $x \in U$, there exists $s \in S$ such that $x \in \uparrow s$. Exercise 2.6.1.11 then implies that $s \leq x$. Since $U' \in \mathcal{U}(S)$, U' is an upper set and $S \subset U'$. Then

$$\begin{aligned} s &\in S \\ &\subset U'. \end{aligned}$$

Since U' is an upper set and $s \leq x$, $x \in U'$. Since $U' \in \mathcal{U}(S)$ is arbitrary, we have that for each $U' \in \mathcal{U}(S)$, $x \in U'$. Thus

$$\begin{aligned} x &\in \bigcap_{U' \in \mathcal{U}(S)} U' \\ &= \uparrow S. \end{aligned}$$

Since $x \in U$ is arbitrary, we have that for each $x \in U$, $x \in \uparrow S$. Hence $U \subset \uparrow S$.

Since $\uparrow S \subset U$ and $U \subset \uparrow S$, we have that $\uparrow S = U$.

2. Similar to (1).

□

Exercise 2.6.1.17. Let $(X, \leq)$ be a proset and $(E_\alpha)_{\alpha \in A} \subset \mathcal{P}(X)$. Then

1. $\uparrow \bigcup_{\alpha \in A} E_\alpha = \bigcup_{\alpha \in A} \uparrow E_\alpha$
2. $\downarrow \bigcup_{\alpha \in A} E_\alpha = \bigcup_{\alpha \in A} \downarrow E_\alpha$

Proof.

1. - Let $x \in \uparrow \bigcup_{\alpha \in A} E_\alpha$. Exercise 2.6.1.16 implies that there exists $y \in \bigcup_{\alpha \in A} E_\alpha$ such that $x \in \uparrow y$. Then there exists $\alpha_0 \in A$ such that $y \in E_{\alpha_0}$. Since $\{y\} \subset E_{\alpha_0}$, Exercise 2.6.1.14 implies that $\uparrow y \subset \uparrow E_{\alpha_0}$. Therefore

$$\begin{aligned} x &\in \uparrow y \\ &\subset \uparrow E_{\alpha_0} \\ &\subset \bigcup_{\alpha \in A} \uparrow E_\alpha. \end{aligned}$$

Since $x \in \uparrow \bigcup_{\alpha \in A} E_\alpha$ is arbitrary, we have that for each $x \in \uparrow \bigcup_{\alpha \in A} E_\alpha$, $x \in \bigcup_{\alpha \in A} \uparrow E_\alpha$. Hence $\uparrow \bigcup_{\alpha \in A} E_\alpha \subset \bigcup_{\alpha \in A} \uparrow E_\alpha$.

- Let $x \in \bigcup_{\alpha \in A} \uparrow E_\alpha$. Then there exists $\alpha_0 \in A$ such that $x \in \uparrow E_{\alpha_0}$. Exercise 2.6.1.16 implies that there exists $y \in E_{\alpha_0}$ such that $x \in \uparrow y$. Since

$$\begin{aligned} \{y\} &\subset E_{\alpha_0} \\ &\subset \bigcup_{\alpha \in A} E_\alpha, \end{aligned}$$

Exercise 2.6.1.14 implies that

$$\begin{aligned} x &\in \uparrow y \\ &\subset \uparrow \bigcup_{\alpha \in A} E_\alpha. \end{aligned}$$

Since $x \in \bigcup_{\alpha \in A} \uparrow E_\alpha$ is arbitrary, we have that for each $x \in \bigcup_{\alpha \in A} \uparrow E_\alpha$, $x \in \uparrow \bigcup_{\alpha \in A} E_\alpha$. Hence $\bigcup_{\alpha \in A} \uparrow E_\alpha \subset \uparrow \bigcup_{\alpha \in A} E_\alpha$.

Since $\uparrow \bigcup_{\alpha \in A} E_\alpha \subset \bigcup_{\alpha \in A} \uparrow E_\alpha$ and $\bigcup_{\alpha \in A} \uparrow E_\alpha \subset \uparrow \bigcup_{\alpha \in A} E_\alpha$, we have that $\uparrow \bigcup_{\alpha \in A} E_\alpha = \bigcup_{\alpha \in A} \uparrow E_\alpha$.

2. Similar to (1).

□

Exercise 2.6.1.18. Let $(A, \leq)$ be a proset and $B \subset A$.

1. If B is upward directed, then $\downarrow B$ is upward directed
2. If B is downward directed, then $\uparrow B$ is downward directed

Proof.

1. Suppose that B is upward directed.
   - Since B is upward directed, $(B, \leq |_B)$ is a directed set. Thus $B \neq \varnothing$. Thus there exists $x \in B$. Since $B \subset \downarrow B$, we have that $x \in \downarrow B$. Hence $\downarrow B \neq \varnothing$.
   - Let $a, b \in \downarrow B$. Exercise 2.6.1.16 implies that there exists $x_a, x_b \in B$ such that $a \in \downarrow x_a$ and $b \in \downarrow x_b$. Since B is upward directed, $(B, \leq |_B)$ is a directed set. Thus $\mathrm{ub}(\{x_a, x_b\}, \leq |_B) \neq \varnothing$. Therefore there exists $x_0 \in B$ such that $x_0 \geq x_a$ and $x_0 \geq x_b$. Hence
   $$\begin{aligned} x_0 &\geq x_a \\ &\geq a. \end{aligned}$$
   Similarly, $x_0 \geq b$. Since $B \subset \downarrow B$, $x_0 \in \downarrow B$. Thus $\mathrm{ub}(\{a, b\}, \leq |_{\downarrow B}) \neq \varnothing$. Since $a, b \in \downarrow B$ are arbitrary, we have that for each $a, b \in \downarrow B$, $\mathrm{ub}(\{a, b\}, \leq |_{\downarrow B}) \neq \varnothing$. Then $\leq |_{\downarrow B}$ is a direction on $\downarrow B$.

   Hence B is upward directed.
2. Suppose that B is $\leq$-downward directed. The prev ex implies that B is $\leq^*$-upward directed. The prev part implies that $\downarrow(B, \leq^*)$ is $\leq^*$-upward directed. A prev ex implies that $\downarrow(B, \leq^*) = \uparrow(B, \leq)$. Since $\uparrow(B, \leq)$ is $\leq^*$-upward directed, the prev ex implies that $\uparrow(B, \leq)$ is $\leq$-downward directed.

□

Exercise 2.6.1.19. Let $(X, \leq_X)$ $(Y, \leq_Y)$ be prosets and $f : X \to Y$. Then for each $U \subset Y$, U is a $\leq_Y$-upper set implies that $f^{-1}(U)$ is a $\leq_X$-upper set iff for each $L \subset Y$, L is a $\leq_Y$-lower set implies that $f^{-1}(L)$ is a $\leq_X$-lower set.

Proof.

- $(\implies)$:
  Suppose that for each $U \subset Y$, U is a $\leq_Y$-upper set implies that $f^{-1}(U)$ is a $\leq_X$-upper set. Let $L \subset Y$. Suppose that L is a $\leq_Y$-lower set. Exercise 2.6.1.5 then implies that L^c is a $\leq_Y$-upper set. By assumption, $f^{-1}(L^c)$ is a $\leq_X$-upper set. Another application of Exercise 2.6.1.5 implies that $f^{-1}(L^c)^c$ is a $\leq_X$-lower set. Since
  $$\begin{aligned} f^{-1}(L) &= [f^{-1}(L)^c]^c \\ &= f^{-1}(L^c)^c, \end{aligned}$$
  we have that $f^{-1}(L)$ is a $\leq_X$-lower set. Since $L \subset Y$ such that L is a $\leq_Y$-lower set is arbitrary, we have that for each $L \subset Y$, L is a $\leq_Y$-lower set implies that $f^{-1}(L)$ is a $\leq_X$-lower set.
- $(\impliedby)$:
  Similar to $(\implies)$.

□

Exercise 2.6.1.20. Let $(X, \leq_X)$ $(Y, \leq_Y)$ be prosets and $f : X \to Y$. If f is $(\leq_X, \leq_Y)$-monotone, then for each $U \subset Y$, U is a $\leq_Y$-upper set implies that $f^{-1}(U)$ is a $\leq_X$-upper set.

Proof. Suppose that f is $(\leq_X, \leq_Y)$-monotone. Let $U \subset Y$. Suppose that U is a $\leq_Y$-upper set. Let $a \in f^{-1}(U)$ and $x \in X$. Suppose that $a \leq x$. Since $a \in f^{-1}(U)$, $f(a) \in U$. Since f is monotone, $f(a) \leq f(x)$. Since U is a $\leq_Y$-upper set and $f(a) \in U$, we have that $f(x) \in U$. Hence $x \in f^{-1}(U)$. Since $a \in f^{-1}(U)$ and $x \in X$ with $a \leq x$ are arbitrary, we have that for each $a \in f^{-1}(U)$ and $x \in X$, $a \leq x$ implies that $x \in f^{-1}(U)$. Thus $f^{-1}(U)$ is a $\leq_X$-upper set. Since $U \subset Y$ such that U is a $\leq_Y$-upper set is arbitrary, we have that for each $U \subset Y$, U is a $\leq_Y$-upper set implies that $f^{-1}(U)$ is a $\leq_X$-upper set. □

Exercise 2.6.1.21. Let $(X, \leq_X)$ and $(Y, \leq_Y)$ be posets and $f : X \to Y$. Then f has a right adjoint iff for each $y \in Y$, there exists $x \in X$ such that $f^{-1}(\downarrow y) = \downarrow x$.

Proof.

- $(\implies)$:
  Suppose that f has a right adjoint. Then there exists $g : Y \to X$ such that (f, g) is a monotone Galois connection. a prev ex in galois connections on prosets section Exercise B.0.0.3 then implies that f and g are monotone, $\mathrm{id}_X \leq_X g \circ f$ and $f \circ g \leq_Y \mathrm{id}_Y$. Let $y \in Y$. Define $x \in X$ by $x := g(y)$.
  - We note that for each $a \in X$,
    $$\begin{aligned} a \in f^{-1}(\downarrow y) &\iff f(a) \in \downarrow y \\ &\iff f(a) \leq_Y y. \end{aligned}$$
    Let $a \in f^{-1}(\downarrow y)$. Then
    $$\begin{aligned} a &\leq_X g \circ f(a) \\ &= g(f(a)) \\ &\leq_X g(y) \\ &= x. \end{aligned}$$
    Therefore $a \in \downarrow x$. Since $a \in f^{-1}(\downarrow y)$ is arbitrary, we have that for each $a \in f^{-1}(\downarrow y)$, $a \in \downarrow x$. Hence $a \in f^{-1}(\downarrow y) \subset \downarrow x$. Then $f(a) \in \downarrow$.
  - Let $a \in \downarrow x$. Then $a \leq_X x$. Hence
    $$\begin{aligned} f(a) &\leq_Y f(x) \\ &= f(g(y)) \\ &= f \circ g(y) \\ &\leq_Y y. \end{aligned}$$
    Thus $f(a) \in \downarrow y$ and $a \in f^{-1}(\downarrow y)$. Since $a \in \downarrow x$ is arbitrary, we have that for each $a \in \downarrow x$, $a \in f^{-1}(\downarrow y)$. Hence $\downarrow x \subset f^{-1}(\downarrow y)$.

  Since $f^{-1}(\downarrow y) \subset \downarrow x$ and $\downarrow x \subset f^{-1}(\downarrow y)$, we have that $f^{-1}(\downarrow y) = \downarrow x$. Since $y \in Y$ is arbitrary, we have that for each $y \in Y$, there exists $x \in X$ such that $f^{-1}(\downarrow y) = \downarrow x$.

- $(\impliedby)$:
  Suppose that for each $y \in Y$, there exists $x \in X$ such that $f^{-1}(\downarrow y) = \downarrow x$. The axiom of choice implies that there exists $g : Y \to X$ such that for each $y \in Y$, $f^{-1}(\downarrow y) = \downarrow g(y)$. Then for each $x \in X$ and $y \in Y$,
  $$\begin{aligned} f(x) \leq_X y &\iff f(x) \in \downarrow y \\ &\iff x \in f^{-1}(\downarrow y) \\ &\iff x \in \downarrow g(y) \\ &\iff x \leq_Y g(y). \end{aligned}$$
  Thus (f, g) is a monotone Galois connection. Hence f has a right adjoint.

□

2.6.2 Maximal Elements

RELABEL, MAYBE RETITLE

Definition 2.6.2.1. Let $(X, \leq)$ be a proset and $a \in X$. Then a is said to be

- **$\leq$-maximal** if for each $x \in X$, $a \leq x$ implies that $x \sim_{\leq} a$.
- **$\leq$-minimal** if for each $x \in X$, $x \leq a$ implies that $x \sim_{\leq} a$.

Note 2.6.2.2. When the context is clear, we write "maximal" and "minimal" instead of "$\leq$-maximal" and "$\leq$-minimal" respectively.

Exercise 2.6.2.3. Let $(X, \leq)$ be a proset and $a \in X$. Then

1. a is maximal iff $\uparrow a = \pi_{X/\sim_{\leq}}(a)$
2. a is minimal iff $\downarrow a = \pi_{X/\sim_{\leq}}(a)$

Proof.

1. - ($\Longrightarrow$):
     Suppose that a is maximal.
     - Let $x \in \uparrow a$. Exercise 2.6.1.11 implies $a \leq x$. Since a is maximal, $x \sim_{\leq} a$. Thus $x \in \pi_{X/\sim_{\leq}}(a)$. Since $x \in \uparrow a$ is arbitrary, we have that for each $x \in \uparrow a$, $x \in \pi_{X/\sim_{\leq}}(a)$. Hence $\uparrow a \subset \pi_{X/\sim_{\leq}}(a)$.
     - Let $x \in \pi_{X/\sim_{\leq}}(a)$. Then $a \sim_{\leq} x$. Hence $a \leq x$ and $x \leq a$. Since $a \leq x$, we have that $x \in \uparrow a$. Since $x \in \pi_{X/\sim_{\leq}}(a)$ is arbitrary, we have that for each $x \in \pi_{X/\sim_{\leq}}(a)$, $x \in \uparrow a$. Hence $\pi_{X/\sim_{\leq}}(a) \subset \uparrow a$.

     Since $\uparrow a \subset \pi_{X/\sim_{\leq}}(a)$ and $\pi_{X/\sim_{\leq}}(a) \subset \uparrow a$, we have that $\uparrow a = \pi_{X/\sim_{\leq}}(a)$.
   - ($\Longleftarrow$):
     Suppose that $\uparrow a = \pi_{X/\sim_{\leq}}(a)$. Let $x \in X$. Suppose that $a \leq x$. Then
     $$\begin{aligned} x &\in \uparrow a \\ &= \pi_{X/\sim_{\leq}}(a). \end{aligned}$$
     Thus $x \sim_{\leq} a$. Since $x \in X$ with $a \leq x$ is arbitrary, we have that for each $x \in X$, $a \leq x$ implies that $x \sim_{\leq} a$. Hence a is maximal.
2. Similar to (1).

□

2.7 Suprema and Infima

2.7.1 Upper and Lower Bounds

Definition 2.7.1.1. maybe move to suprema infima section, or make a section on upper and lower bounds Let $(X, \leq)$ be a proset and $A \subset X$.

- Let $x \in X$. Then x is said to be a
  - **$\leq$-upper bound of** A if for each $a \in A$, $a \leq x$
  - **$\leq$-lower bound of** A if for each $a \in A$, $x \leq a$
- We define
  - $\mathrm{ub}(A, \leq) := \{x \in X : x \text{ is a } \leq\text{-upper bound of } A\}$
  - $\mathrm{lb}(A, \leq) := \{x \in X : x \text{ is a } \leq\text{-lower bound of } A\}$
- Then A is said to be

- **$\leq$-bounded above** if $\operatorname{ub}(A, \leq) \neq \varnothing$
- **$\leq$-bounded below** if $\operatorname{lb}(A, \leq) \neq \varnothing$

Note 2.7.1.2. When the context is clear, we write

- "upper bound" and "lower bound" instead of "$\leq$-upper bound" and "$\leq$-lower bound" respectively
- $\operatorname{ub} A$ and $\operatorname{lb} A$ in place of $\operatorname{ub}(A, \leq)$ and $\operatorname{lb}(A, \leq)$ respectively
- "bounded above" and "bounded below" instead of "$\leq$-bounded above" and "$\leq$-bounded below" respectively

Exercise 2.7.1.3. Let $(X, \leq)$ be a proset and $A \subset X$. Then

1. $\operatorname{ub}(A, \leq) = \operatorname{lb}(A, \leq^*)$
2. $\operatorname{ub}(A, \leq^*) = \operatorname{lb}(A, \leq)$

Proof.

1. We note that for each $x \in X$,

$$\begin{aligned} x \in \operatorname{ub}(A, \leq) &\iff \text{for each } a \in A,\ a \leq x \\ &\iff \text{for each } a \in A,\ x \leq^* a \\ &\iff x \in \operatorname{lb}(A, \leq^*). \end{aligned}$$

   Hence $\operatorname{ub}(A, \leq) = \operatorname{lb}(A, \leq^*)$.

2. Similar to (1) ref.

□

Exercise 2.7.1.4. Let $(X, \leq)$ be a proset and $A \subset X$. Then

1. $\operatorname{ub} A = \bigcap_{x \in A} \uparrow x$
2. $\operatorname{lb} A = \bigcap_{x \in A} \downarrow x$

Proof.

1.
   - Let $a \in \operatorname{ub} A$ and $x \in A$. Then $x \leq a$. Hence $a \in \uparrow x$. Since $x \in A$ is arbitrary, we have that for each $x \in A$, $a \in \uparrow x$. Thus $a \in \bigcap_{x \in A} \uparrow x$. Since $a \in \operatorname{ub} A$ is arbitrary, we have that for each $a \in \operatorname{ub} A$, $a \in \bigcap_{x \in A} \uparrow x$. Hence $\operatorname{ub} A \subset \bigcap_{x \in A} \uparrow x$.
   - Let $a \in \bigcap_{x \in A} \uparrow x$ and $x_0 \in A$. Then $a \in \uparrow x_0$. Hence $x_0 \leq a$. Since $x_0 \in A$ is arbitrary, we have that for each $x_0 \in A$, $x_0 \leq a$. Hence $a \in \operatorname{ub} A$. Since $a \in \bigcap_{x \in A} \uparrow x$ is arbitrary, we have that for each $a \in \bigcap_{x \in A} \uparrow x$, $a \in \operatorname{ub} A$. Thus $\bigcap_{x \in A} \uparrow x \subset \operatorname{ub} A$.

   Since $\operatorname{ub} A \subset \bigcap_{x \in A} \uparrow x$ and $\bigcap_{x \in A} \uparrow x \subset \operatorname{ub} A$, we have that $\operatorname{ub} A = \bigcap_{x \in A} \uparrow x$.

2. Similar to (1).

□

Exercise 2.7.1.5. Let $(X, \leq)$ be a proset and $A, B \subset X$. If $A \subset B$, then

1. $\operatorname{ub} B \subset \operatorname{ub} A$
2. $\operatorname{lb} B \subset \operatorname{lb} A$

Proof. Suppose that $A \subset B$.

1. Exercise 2.7.1.4 implies that

$$\begin{aligned}\operatorname{ub} B &= \bigcap_{x\in B} \uparrow x \\ &\subset \bigcap_{x\in A} \uparrow x \\ &= \operatorname{ub} A.\end{aligned}$$

2. Similar to (1). ref

□

Exercise 2.7.1.6. Let $(X, \leq)$ be a proset and $(E_\alpha)_{\alpha\in A} \subset \mathcal{P}(X)$. Then $\operatorname{ub} \bigcup_{\alpha\in A} E_\alpha = \bigcap_{\alpha\in A} \operatorname{ub} E_\alpha$.

Proof. Define $E_\vee \subset X$ by $E := \bigcup_{\alpha\in A} E_\alpha$. Exercise 2.7.1.4 then implies that

$$\begin{aligned}\operatorname{ub} \bigcup_{\alpha\in A} E_\alpha &= \bigcap_{x\in E} \uparrow x \\ &= \bigcap_{\alpha\in A} \left[\bigcap_{x\in E_\alpha} \uparrow x \right] \\ &= \bigcap_{\alpha\in A} \operatorname{ub} E_\alpha.\end{aligned}$$

□

Exercise 2.7.1.7. Let $(X, \leq)$ be a proset and $A, B \subset X$. Then

1. $\operatorname{ub}(B, \leq|_A) = \operatorname{ub}(B, \leq) \cap A$.
2. $\operatorname{lb}(B, \leq|_A) = \operatorname{lb}(B, \leq) \cap A$.

Proof.

1. We note that for each $x \in X$,

$$\begin{aligned}x \in \operatorname{ub}(B, \leq|_A) &\iff x \in A \text{ and for each } y \in B,\ y \leq|_A x \\ &\iff x \in A \text{ and for each } y \in B,\ y \leq x \\ &\iff x \in \operatorname{ub}(B, \leq) \cap A.\end{aligned}$$

2. We have that

$$\begin{aligned}\operatorname{lb}(B, \leq|_A) &= \operatorname{ub}(B, \leq|_A^*) \quad \text{(by Exercise 2.7.1.3)} \\ &= \operatorname{ub}(B, \leq^*|_A) \quad \text{(by ref ex in set theory notes in binary hetero relations)} \\ &= \operatorname{ub}(B, \leq^*) \cap A \quad \text{(by ref prev part)} \\ &= \operatorname{lb}(B, \leq) \cap A \quad \text{(by Exercise 2.7.1.3).}\end{aligned}$$

□

Exercise 2.7.1.8. Let $(X, \leq_X)$ and $(Y, \leq_Y)$ be prosets, $f : X \to Y$ and $g : Y \to X$.

1. If (f, g) is a $(\leq_X, \leq_Y)$-monotone Galois connection, then
   (a) for each $A \subset X$, $\operatorname{ub}(f(A), \leq_Y) = g^{-1}(\operatorname{ub}(A, \leq_X))$,
   (b) for each $B \subset Y$, $\operatorname{lb}(g(B), \leq_X) = f^{-1}(\operatorname{lb}(B, \leq_Y))$.

2. If (f, g) is a $(\leq_X, \leq_Y)$-antitone Galois connection, then
   (a) for each $A \subset X$, $\mathrm{ub}(f(A), \leq_Y) = g^{-1}(\mathrm{lb}(A, \leq_X))$,
   (b) for each $B \subset Y$, $\mathrm{ub}(g(B), \leq_X) = f^{-1}(\mathrm{lb}(B, \leq_Y))$.

Proof. Since (f, g) is a $(\leq_X, \leq_Y)$-monotone Galois connection, a prev ex implies that f is $(\leq_X, \leq_Y)$-monotonoe, g is $(\leq_Y, \leq_X)$-monotone, $\mathrm{id}_X \leq_X g \circ f$ and $f \circ g \leq_Y \mathrm{id}_Y$.

1. Suppose that (f, g) is a $(\leq_X, \leq_Y)$-monotone Galois connection.
   (a) Let $A \subset X$.
   - Let $y \in \mathrm{ub}(f(A), \leq_Y)$ and $a \in A$. Then $f(a) \leq_Y y$. Hence
   $$\begin{aligned} a &\leq_X g \circ f(a) \\ &= g(f(a)) \\ &\leq_X g(y). \end{aligned}$$
   Since $a \in A$ is arbitrary, we have that for each $a \in A$, $a \leq_X g(y)$. Thus $g(y) \in \mathrm{ub}(A, \leq_X)$. Hence $y \in g^{-1}(\mathrm{ub}(A, \leq_X))$. Since $y \in \mathrm{ub}(f(A), \leq_Y)$ is arbitrary, we have that for each $y \in \mathrm{ub}(f(A), \leq_Y)$, $y \in g^{-1}(\mathrm{ub}(A, \leq_X))$. Thus $\mathrm{ub}(f(A), \leq_Y) \subset g^{-1}(\mathrm{ub}(A, \leq_X))$.
   - Let $y \in g^{-1}(\mathrm{ub}(A, \leq_X))$. Then $g(y) \in \mathrm{ub}(A, \leq_X)$. Let $b \in f(A)$. Then there exists $a \in A$ such that $b = f(a)$. Then $a \leq_X g(y)$. Therefore
   $$\begin{aligned} b &= f(a) \\ &\leq_Y f(g(y)) \\ &= f \circ g(y) \\ &= \leq_Y y. \end{aligned}$$
   Since $b \in f(A)$ is arbitrary, we have that for each $b \in f(A)$, $b \leq_Y y$. Thus $y \in \mathrm{ub}(f(A), \leq_Y)$. Since $y \in g^{-1}(\mathrm{ub}(A, \leq_X))$ is arbitrary, we have that for each $y \in g^{-1}(\mathrm{ub}(A, \leq_X))$, $y \in \mathrm{ub}(f(A), \leq_Y)$. Hence $g^{-1}(\mathrm{ub}(A, \leq_X)) \subset \mathrm{ub}(f(A), \leq_Y)$.

   Since $\mathrm{ub}(f(A), \leq_Y) \subset g^{-1}(\mathrm{ub}(A, \leq_X))$ and $g^{-1}(\mathrm{ub}(A, \leq_X)) \subset \mathrm{ub}(f(A), \leq_Y)$, we have that
   $$\mathrm{ub}(f(A), \leq_Y) = g^{-1}(\mathrm{ub}(A, \leq_X)).$$
   (b) Since (f, g) is a $(\leq_X, \leq_Y)$-monotone Galois connection, a prev rex implies that (g, f) is a $(\leq_Y^*, \leq_X^*)$-monotone Galois connection. The previous part (ref) and Exercise 2.7.1.3 then imply that for each $B \subset Y$,
   $$\begin{aligned} \mathrm{lb}(g(B), \leq_X) &= \mathrm{ub}(g(B), \leq_X^*) \\ &= f^{-1}(\mathrm{ub}(B, \leq_Y^*)) \\ &= f^{-1}(\mathrm{lb}(B, \leq_Y)). \end{aligned}$$

2. Suppose that (f, g) is a $(\leq_X, \leq_Y)$-antitone Galois connection. A prev ex then implies that (f, g) is a $(\leq_X^*, \leq_Y)$-monotone Galois connection and (g, f) is a $(\leq_Y^*, \leq_X)$-monotone Galois connection. The prev part implies that
   (a) for each $A \subset X$,
   $$\begin{aligned} \mathrm{ub}(f(A), \leq_Y) &= g^{-1}(\mathrm{ub}(A, \leq_X^*)) \\ &= g^{-1}(\mathrm{lb}(A, \leq_X)). \end{aligned}$$
   (b) for each $B \subset Y$,
   $$\begin{aligned} \mathrm{ub}(g(B), \leq_X) &= f^{-1}(\mathrm{ub}(B, \leq_Y^*)) \\ &= f^{-1}(\mathrm{lb}(B, \leq_Y)). \end{aligned}$$

□

Exercise 2.7.1.9. Let $(X, \leq)$ be a proset. Define $f, g : \mathcal{P}(X) \to \mathcal{P}(X)$ by $f := \mathrm{ub}(\cdot, \leq)$ and $g := \mathrm{lb}(\cdot, \leq)$. Then (f, g) is a $(\subset^*, \subset^*)$-antitone Galois connection.

Proof.

1. make previous exercise implies that $A \subset g \circ f(A)$ and $A \subset f \circ g(A)$. Then for each $A \in \mathcal{P}(X)$, $g \circ f(A) \subset^* A$ and $f \circ g(A) \subset^* A$.
2. make previous exercises imply that f and g are $(\subset^*, \subset^*)$-antitone

the previous exercise implies that (f, g) FINISH!!! □

Exercise 2.7.1.10. Let $(X, \leq_X)$ $(Y, \leq_Y)$ be prosets, $f : X \to Y$ and $A \subset X$. If f is $(\leq_X, \leq_Y)$-monotone, then

1. $f(\mathrm{ub}(A, \leq_X)) \subset \mathrm{ub}(f(A), \leq_Y)$,
2. $f(\mathrm{lb}(A, \leq_X)) \subset \mathrm{lb}(f(A), \leq_Y)$

Proof. Suppose that f is $(\leq_X, \leq_Y)$-monotone.

1. Let $x \in \mathrm{ub}(A, \leq_X)$ and $y_0 \in f(A)$. Then there exists $x_0 \in A$ such that $f(x_0) = y$. Since $x \in \mathrm{ub}(A, \leq_X)$, $x_0 \leq x$. Since f is $(\leq_X, \leq_Y)$-monotone,

$$\begin{aligned} y_0 &= f(x_0) \\ &\leq f(x). \end{aligned}$$

   Since $y_0 \in f(A)$ is arbitrary, we have that for each $y_0 \in f(A)$, $y_0 \leq f(x)$. Therefore $f(x) \in \mathrm{ub}(F(A), \leq_Y)$. Since $x \in \mathrm{ub}(A, \leq_X)$ is arbitrary, ew have that for each $x \in \mathrm{ub}(A, \leq_X)$, $f(x) \in \mathrm{ub}(F(A), \leq_Y)$. Hence $f(\mathrm{ub}(A, \leq_X)) \subset \mathrm{ub}(f(A), \leq_Y)$.

2. Exercise 2.7.1.3 and the prev part imply that

$$\begin{aligned} f(\mathrm{lb}(A, \leq_X)) &= f(\mathrm{ub}(A, \leq_X^*)) \\ &\subset \mathrm{ub}(f(A), \leq_Y^*) \\ &= \mathrm{lb}(f(A), \leq_Y). \end{aligned}$$

□

2.7.2 SupSets and InfSets

Definition 2.7.2.1. Let $(X, \leq)$ be a proset and $A \subset X$.

- Let $x \in X$. Then x is said to be a
  - **$\leq$-supremum of** A or **$\leq$-least upper bound of** A, if
    1. $x \in \mathrm{ub}(A, \leq)$
    2. for each $y \in \mathrm{ub}(A, \leq)$, $x \leq y$
  - **$\leq$-infimum of** A or **$\leq$-greatest lower bound of** A, if
    1. $x \in \mathrm{lb}(A, \leq)$
    2. for each $y \in \mathrm{lb}(A, \leq)$, $y \leq x$.
-
  - We define $\mathrm{supSet}(A, \leq) \subset X$ by $\mathrm{supSet}(A, \leq) := \{x \in X : x \text{ is a } \leq\text{-supremum of } A\}$
  - We define $\mathrm{infSet}(A, \leq) \subset X$ by $\mathrm{infSet}(A, \leq) := \{x \in X : x \text{ is an } \leq\text{-infimum of } A\}$

Note 2.7.2.2. When the context is clear,

- we say "supremum" and "infimum" in place of "$\leq$-supremum" and "$\leq$-infimum" respectively,
- we write $\mathrm{supSet}\, A$ and $\mathrm{infSet}\, A$ in place of $\mathrm{supSet}(A, \leq)$ and $\mathrm{infSet}(A, \leq)$ respectively.

Exercise 2.7.2.3. NEED TO REORGANIZE AND REWORK EXERCISES ABOUT SUPREMA/INFIMA AND DUALITY Let $(X, \leq)$ be a poset and $A \subset X$. Then

1. (a) $\operatorname{supSet} A = \operatorname{ub} A \cap \operatorname{lb}(\operatorname{ub} A)$
   (b) $\operatorname{supSet} A = \operatorname{infSet}(\operatorname{ub} A)$
2. (a) $\operatorname{infSet} A = \operatorname{lb} A \cap \operatorname{ub}(\operatorname{lb} A)$
   (b) $\operatorname{infSet} A = \operatorname{supSet}(\operatorname{lb} A)$

Proof.

1. (a) We note that for each $x \in X$,

$$\begin{aligned} x \in \operatorname{supSet} A &\iff x \in \operatorname{ub} A \text{ and for each } y \in \operatorname{ub} A,\ x \leq y \\ &\iff x \in \operatorname{ub} A \text{ and } x \in \operatorname{lb}(\operatorname{ub} A) \\ &\iff x \in \operatorname{ub} A \cap \operatorname{lb}(\operatorname{ub} A). \end{aligned}$$

   Therefore $\operatorname{supSet} A = \operatorname{ub} A \cap \operatorname{lb}(\operatorname{ub} A)$.

   (b) Define $f, g : \mathcal{P}(X) \to \mathcal{P}(X)$ by $f := \operatorname{ub}(\cdot, \leq)$ and $g := \operatorname{lb}(\cdot, \leq)$. A previous exercis in poset galois connection subsection implies that (f, g) is a $(\subset^*, \subset^*)$-antitone Galois connection. Another prev ex in proset galois subsection implies that $\operatorname{ub}(\operatorname{lb}(\operatorname{ub} A)) \sim_{\subset} \operatorname{ub} A$. Therefore $\operatorname{ub}(\operatorname{lb}(\operatorname{ub} A)) \subset \operatorname{ub} A$ and $\operatorname{ub} A \subset \operatorname{ub}(\operatorname{lb}(\operatorname{ub} A))$. Hence $ubd(\operatorname{lb}(\operatorname{ub} A)) = \operatorname{ub} A$. Thus

$$\begin{aligned} \operatorname{supSet} A &= \operatorname{ub} A \cap \operatorname{lb}(\operatorname{ub} A) \quad (\text{by prev part (ref)}) \\ &= \operatorname{ub}(\operatorname{lb}(\operatorname{ub} A)) \cap \operatorname{lb}(\operatorname{ub} A) \quad (\text{by prev part (ref)}) \\ &= \operatorname{lb}(\operatorname{ub} A) \cap \operatorname{ub}(\operatorname{lb}(\operatorname{ub} A)) \\ &= \operatorname{infSet}(\operatorname{ub} A). \end{aligned}$$

2. Similar to (1) USE DUALITY???.

□

Exercise 2.7.2.4. Let $(X, \leq)$ be a poset and $A \subset X$. Then

1. $\operatorname{supSet}(A, \leq^*) = \operatorname{infSet}(A, \leq)$
2. $\operatorname{supSet}(A, \leq) = \operatorname{infSet}(A, \leq^*)$

Proof.

1. We have that

$$\begin{aligned} \operatorname{supSet}(A, \leq^*) &= \operatorname{ub}(A, \leq^*) \cap \operatorname{lb}(\operatorname{ub}(A, \leq^*), \leq^*) \quad (\text{by Exercise 2.7.2.3}) \\ &= \operatorname{lb}(A, \leq) \cap \operatorname{ub}(\operatorname{lb}(A, \leq), \leq) \quad (\text{by Exercise 2.7.1.3}) \\ &= \operatorname{infSet}(A, \leq)(\text{by Exercise 2.7.2.3}). \end{aligned}$$

2. We have that

$$\begin{aligned} \operatorname{supSet}(A, \leq) &= \operatorname{supSet}(A, (\leq^*)^*) \\ &= \operatorname{infSet}(A, \leq^*) \quad (\text{by prev part (ref prev part)}). \end{aligned}$$

Exercise 2.7.2.3 □

Exercise 2.7.2.5. Let $(X, \leq)$ be a poset and $A \subset X$. Then

1. for each $x, y \in \operatorname{supSet}(A, \leq)$, $x \sim_{\leq} y$.
2. for each $x, y \in \operatorname{infSet}(A, \leq)$, $x \sim_{\leq} y$.

Proof.

1. Let $x, y \in \operatorname{supSet}(A, \leq)$. Then $x, y \in \operatorname{ub}(A, \leq)$. Since $x \in \operatorname{supSet}(A, \leq)$ and $y \in \operatorname{ub}(A, \leq)$, $x \leq y$. Since $y \in \operatorname{supSet}(A, \leq)$ and $x \in \operatorname{ub}(A, \leq)$, $y \leq x$. Hence $x \sim_{\leq} y$. Since $x, y \in \operatorname{supSet}(A, \leq)$ are arbitrary, we have that for each $x, y \in \operatorname{supSet}(A, \leq)$, $x \sim_{\leq} y$.
2. We note that for each $x, y \in X$,

$$\begin{aligned} x, y \in \operatorname{infSet}(A, \leq) &\iff x, y \in \operatorname{supSet}(A, \leq^*) \quad \text{(by Exercise 2.7.2.4)} \\ &\implies x \sim_{\leq^*} y \quad \text{(by prev part (ref))} \\ &\iff x \sim_{\leq} y \quad \text{(by Exercise 2.2.1.6).} \end{aligned}$$

□

Exercise 2.7.2.6. Let $(X, \leq)$ be a poset and $A \subset X$. Then

1. for each $x, y \in \operatorname{supSet} A$, $x = y$.
2. for each $x, y \in \operatorname{infSet} A$, $x = y$.

Proof.

1. Let $x, y \in \operatorname{supSet} A$. Exercise 2.7.2.5 implies that $x \sim_{\leq} y$. Exercise 2.3.1.2 then implies that $x = y$.
2. Similar to (1).

□

Exercise 2.7.2.7. Let $(X, \leq)$ be a proset and $A, B \subset X$. Suppose that $\sup A$ and $\sup B$ exist. If $A \subset B$, then for each $s_A \in \operatorname{supSet} A$ and $s_B \in \operatorname{supSet} B$, $s_A \leq s_B$.

Proof. Suppose that $A \subset B$. Let $s_A \in \operatorname{supSet} A$ and $s_B \in \operatorname{supSet} B$. Then Exercise 2.7.1.5 implies that

$$\begin{aligned} s_B \operatorname{supSet} B \\ &\subset \operatorname{ub} B \\ &\subset \operatorname{ub} A. \end{aligned}$$

Since $s_A \in \operatorname{lb}(\operatorname{ub} A)$, we have that $s_A \leq s_B$. □

Exercise 2.7.2.8. Let $(X, \leq_X)$ and $(Y, \leq_Y)$ be prosets, $f : X \to Y$, $g : Y \to X$ and $A \subset X$. If (f, g) is a $(\leq_X, \leq_Y)$-monotone Galois connection, then $f(\operatorname{supSet}(A, \leq_X)) \subset \operatorname{supSet}(f(A), \leq_Y)$. can we say something about $\operatorname{infSet}(f(A), \leq_Y)$ ir $\operatorname{infSet}(g(B), \leq_X)$?

Proof. Suppose that (f, g) is a $(\leq_X, \leq_Y)$-monotone Galois connection. A prev ex in Galois connection subsection implies that f is $(\leq_X, \leq_Y)$-monotone, g is $(\leq_Y, \leq_X)$-monotone, $g \circ f \leq_X \operatorname{id}_X$ and $\operatorname{id}_Y \leq_Y f \circ g$. Let $s \in \operatorname{supSet}(A, \leq_X)$. Since $\operatorname{supSet}(A, \leq_X) \subset \operatorname{ub}(A, \leq_X)$, we have that $s \in \operatorname{ub}(A, \leq_X)$. Exercise 2.7.1.10 then implies that

$$\begin{aligned} f(s) &\in f(\operatorname{ub}(A, \leq_X)) \\ &\subset \operatorname{ub}(f(A), \leq_Y). \end{aligned}$$

Let $y \in \operatorname{ub}(f(A), \leq_Y)$. A prev ex in galois connection subsection imply that

$$\begin{aligned} y &\in \operatorname{ub}(f(A), \leq_Y) \\ &= g^{-1}(\operatorname{ub}(A, \leq_X)). \end{aligned}$$

Then $g(y) \in \operatorname{ub}(A, \leq_X)$. Since $s \in \operatorname{supSet}(A, \leq_X)$, we have that $s \leq_X g(y)$. Hence

$$\begin{aligned} f(s) &\leq_Y f(g(y)) \\ &= f \circ g(y) \\ &\leq_Y y. \end{aligned}$$

Since $y \in \operatorname{ub}(f(A), \leq_Y)$ is arbitrary, we have that for each $y \in \operatorname{ub}(f(A), \leq_Y)$, $f(s) \leq_Y y$. Thus $f(s) \in \operatorname{supSet}(f(A), \leq_Y)$. Since $s \in \operatorname{supSet}(A, \leq_X)$ is arbitrary, we have that for each $s \in \operatorname{supSet}(A, \leq_X)$, $f(s) \in \operatorname{supSet}(f(A), \leq_Y)$. Hence $f(\operatorname{supSet}(A, \leq_X)) \subset \operatorname{supSet}(f(A), \leq_Y)$. □

2.7.3 Suprema and Infima

relabel ex/def references

Definition 2.7.3.1. Let $(X, \leq)$ be a poset and $A \subset X$.

- If $\operatorname{supSet} A \neq \varnothing$, we define the **supremum of** A, denoted $\sup A$, to be the unique $x \in X$ such that $x \in \operatorname{supSet} A$.
- If $\operatorname{infSet} A \neq \varnothing$, we define the **infimum of** A, denoted $\inf A$, to be the unique $x \in X$ such that $x \in \operatorname{infSet} A$.
- We say that $\sup A$ **(resp.** $\inf A$**) exists** if $\operatorname{supSet} A \neq \varnothing$ (resp. $\operatorname{infSet} A \neq \varnothing$).

Exercise 2.7.3.2. Let $(X, \leq)$ be a poset and $A \subset X$. Then

1. (a) $\operatorname{supSet} A = \operatorname{infSet}(\operatorname{ub} A)$ move to supset section, just use galois connection fact that $f \sim_{\leq} f \circ g \circ f$ then make observation that the two subsets contain each other and are therefore equal, need to make ex that $(\mathcal{P}(X), \subset)$ is a proset beforehand
   (b) $\sup A$ exists iff $\inf(\operatorname{ub} A)$ exists
   (c) $\sup A$ exists implies that $\sup A = \inf(\operatorname{ub} A)$
2. (a) $\operatorname{infSet} A = \operatorname{supSet}(\operatorname{lb} A)$ move to proset section
   (b) $\inf A$ exists iff $\operatorname{supSet}(\operatorname{lb} A)$ exists
   (c) $\inf A$ exists implies that $\inf A = \sup(\operatorname{lb} A)$

Proof.

1. (a) A previous ex in subsection on suprema and infima implies that
   - $\operatorname{supSet} A = \operatorname{ub} A \cap \operatorname{lb}(\operatorname{ub} A)$
   - $\operatorname{infSet}(\operatorname{ub} A) = \operatorname{lb}(\operatorname{ub} A) \cap \operatorname{ub}(\operatorname{lb}(\operatorname{ub} A))$.

   Define $f, g : \mathcal{P}(X) \to \mathcal{P}(X)$ by $f := \operatorname{ub}(\cdot, \leq)$ and $g := \operatorname{lb}(\cdot, \leq)$. A previous exercis in poset galois connection subsection implies that (f, g) is a $(\subset^*, \subset^*)$-antitone Galois connection. Another prev ex in poset galois subsection implies that for each $A \in \mathcal{P}(X)$, $\operatorname{ub}(\operatorname{lb}(\operatorname{ub} A)) = \operatorname{ub} A$. Therefore

$$\begin{aligned}
\operatorname{supSet} A &= \operatorname{ub} A \cap \operatorname{lb}(\operatorname{ub} A) \\
&= \operatorname{ub}(\operatorname{lb}(\operatorname{ub} A)) \cap \operatorname{lb}(\operatorname{ub} A) \\
&= \operatorname{lb}(\operatorname{ub} A) \cap \operatorname{ub}(\operatorname{lb}(\operatorname{ub} A)) \\
&= \operatorname{infSet}(\operatorname{ub} A).
\end{aligned}$$

   (b) The previous part implies that

$$\begin{aligned}
\sup A \text{ exists} &\iff \operatorname{supSet} A \neq \varnothing \\
&\iff \operatorname{infSet}(\operatorname{ub} A) \neq \varnothing \\
&\iff \inf(\operatorname{ub} A) \text{ exists.}
\end{aligned}$$

   (c) Suppose that $\sup A$ exists. the previous part (a) then implies that $\inf(\operatorname{ub} A)$ exists. a previous ex implies that $\operatorname{supSet} A = \{\sup A\}$ and $\operatorname{infSet}(\operatorname{ub} A) = \{\inf(\operatorname{ub} A)\}$. The previous parts (a) then implies that

$$\begin{aligned}
\{\sup A\} &= \operatorname{supSet} A \\
&= \operatorname{infSet}(\operatorname{ub} A) \\
&= \{\inf(\operatorname{ub} A)\}.
\end{aligned}$$

   Hence $\sup A = \inf(\operatorname{ub} A)$.

2. Similar to (1).

□

Exercise 2.7.3.3. Let $(X, \leq)$ be a poset and $A \subset X$. Then

1. (a) $\sup(A, \leq)$ exists iff $\inf(A, \leq^*)$ exists
   (b) $\sup A$ exists implies that $\sup(A, \leq) = \inf(A, \leq^*)$
2. (a) $\operatorname{infSet} A = \operatorname{supSet}(\operatorname{lb}(A, \leq), \leq)$
   (b) $\inf(A, \leq)$ exists iff
      i. $\operatorname{supSet}(\operatorname{lb}(A, \leq), \leq)$ exists
      ii. $\sup(A, \leq^*)$ exists
   (c) $\inf(A <\leq)$ exists implies that
      i. $\inf(A, \leq) = \sup(\operatorname{lb}(A, \leq), \leq)$
      ii. $\inf(A, \leq) = \sup(A, \leq^*)$

Proof.

1. (a) A previous ex in subsection on suprema and infima implies that
   - $\operatorname{supSet} A = \operatorname{ub} A \cap \operatorname{lb}(\operatorname{ub} A)$
   - $\operatorname{infSet}(\operatorname{ub} A) = \operatorname{lb}(\operatorname{ub} A) \cap \operatorname{ub}(\operatorname{lb}(\operatorname{ub} A))$.

   Define $f, g : \mathcal{P}(X) \to \mathcal{P}(X)$ by $f := \operatorname{ub}(\cdot, \leq)$ and $g := \operatorname{lb}(\cdot, \leq)$. A previous exercis in poset galois connection subsection implies that (f, g) is a $(\subset^*, \subset^*)$-antitone Galois connection. Another prev ex in poset galois subsection implies that for each $A \in \mathcal{P}(X)$, $\operatorname{ub}(\operatorname{lb}(\operatorname{ub} A)) = \operatorname{ub} A$. Therefore

$$\begin{aligned}\operatorname{supSet} A &= \operatorname{ub} A \cap \operatorname{lb}(\operatorname{ub} A)\\ &= \operatorname{ub}(\operatorname{lb}(\operatorname{ub} A)) \cap \operatorname{lb}(\operatorname{ub} A)\\ &= \operatorname{lb}(\operatorname{ub} A) \cap \operatorname{ub}(\operatorname{lb}(\operatorname{ub} A))\\ &= \operatorname{infSet}(\operatorname{ub} A).\end{aligned}$$

   (b) i. The previous part implies that

$$\begin{aligned}\sup(A, \leq) \text{ exists} &\iff \operatorname{supSet}(A, \leq) \neq \varnothing\\ &\iff \operatorname{infSet}(\operatorname{ub}(A, \leq), \leq) \neq \varnothing\\ &\iff \inf(\operatorname{ub}(A, \leq), \leq) \text{ exists.}\end{aligned}$$

   ii.

$$\begin{aligned}\sup(A, \leq) \text{ exists} &\iff \operatorname{supSet}(A, \leq) \neq \varnothing\\ &\iff \operatorname{infSet}(A, \leq^*) \neq \varnothing \quad \text{(by Exercise 2.7.2.4)}\end{aligned}$$

   (c) Suppose that $\sup A$ exists. the previous part (a) then implies that $\inf(\operatorname{ub} A)$ exists. a previous ex implies that $\operatorname{supSet} A = \{\sup A\}$ and $\operatorname{infSet}(\operatorname{ub} A) = \{\inf(\operatorname{ub} A)\}$. The previous parts (a) then implies that

$$\begin{aligned}\{\sup A\} &= \operatorname{supSet} A\\ &= \operatorname{infSet}(\operatorname{ub} A)\\ &= \{\inf(\operatorname{ub} A)\}.\end{aligned}$$

   Hence $\sup A = \inf(\operatorname{ub} A)$.

2. Similar to (1).

□

Exercise 2.7.3.4. Let $(X, \leq)$ be a poset and $A, B \subset X$. Suppose that $\sup A$ and $\sup B$ exist. If $A \subset B$, then $\sup A \leq \sup B$.

Proof. Since $\sup A \in \operatorname{supSet} A$ and $\sup B \in \operatorname{supSet} B$, Exercise 2.7.2.7 implies that $\sup A \leq \sup B$. □

Exercise 2.7.3.5. Associativity of Supremum:
Let $(X, \leq)$ be a poset and $(E_\alpha)_{\alpha \in A} \subset \mathcal{P}(X)$.

1. Suppose that
   - for each $\alpha \in A$, $\sup E_\alpha$ exists
   - $\sup\limits_{\alpha \in A} \left[\sup E_\alpha \right]$ exists.

   Then $\sup \bigcup\limits_{\alpha \in A} E_\alpha$ exists and
   $$\sup \bigcup_{\alpha \in A} E_\alpha = \sup_{\alpha \in A} \left[\sup E_\alpha \right]$$
2. Suppose that
   - for each $\alpha \in A$, $\inf E_\alpha$ exists
   - $\inf\limits_{\alpha \in A} \left[\inf E_\alpha \right]$ exists.

   Then $\inf \bigcup\limits_{\alpha \in A} E_\alpha$ exists
   $$\inf \bigcup_{\alpha \in A} E_\alpha = \inf_{\alpha \in A} \left[\inf E_\alpha \right]$$

Proof.

1. Define $s \in X$ by $s := \sup\limits_{\alpha \in A} \left[\sup E_\alpha \right]$.
   - Let $x \in \bigcup\limits_{\alpha \in A} E_\alpha$. Then there exists $\alpha_0 \in A$ such that $x \in E_{\alpha_0}$. We note by definition, $s \in \text{ub}\{\sup E_\alpha : \alpha \in A\}$. Hence for each $\alpha \in A$, $s \geq \sup E_\alpha$. Since $\sup E_{\alpha_0} \in \text{ub}\, E_{\alpha_0}$, we have that
   $$\begin{aligned} x &\leq \sup E_{\alpha_0} \\ &\leq s. \end{aligned}$$
   Since $x \in \bigcup\limits_{\alpha \in A} E_\alpha$ is arbitrary, we have that for each $x \in \bigcup\limits_{\alpha \in A} E_\alpha$, $x \leq s$. Hence $s \in \text{ub} \bigcup\limits_{\alpha \in A} E_\alpha$.
   - Let $x \in \text{ub} \bigcup\limits_{\alpha \in A} E_\alpha$ and $\alpha_0 \in A$. Since $E_{\alpha_0} \subset \bigcup\limits_{\alpha \in A} E_\alpha$, Exercise 2.7.1.5 implies that
   $$\begin{aligned} x &\in \text{ub} \bigcup_{\alpha \in A} E_\alpha \\ &\subset \text{ub}\, E_{\alpha_0}. \end{aligned}$$
   Therefore $\sup E_{\alpha_0} \leq x$. Since $\alpha_0 \in A$ is arbitrary, we have that for each $\alpha_0 \in A$, $\sup E_{\alpha_0} \leq x$. Hence $x \in \text{ub}\{\sup E_\alpha : \alpha \in A\}$. Therefore $s \leq x$. Since $x \in \text{ub} \bigcup\limits_{\alpha \in A} E_\alpha$ is arbitrary, we have that for each $x \in \text{ub} \bigcup\limits_{\alpha \in A} E_\alpha$, $s \leq x$.

   Therefore $s \in \text{supSet} \bigcup\limits_{\alpha \in A} E_\alpha$ and $\sup \bigcup\limits_{\alpha \in A} E_\alpha = s$.
2. Similar to (1). Maybe fill out, use duality

□

Exercise 2.7.3.6. Let $(X, \leq)$ be a poset and $a, b \in X$. Then

1. $\sup\{a\}$ exists and $\sup\{a\} = a$

2. $a \leq b$ iff $\sup\{a, b\}$ exists and $b = \sup\{a, b\}$,
3. $\sup \downarrow a = a$,
4. $\inf \uparrow a = a$.

Proof.

1. Since $a \leq a$, $a \in \text{ub}\{a\}$. By definition, for each $x \in \text{ub}\{a\}$, $a \leq x$. Hence $a \in \text{supSet}\{a\}$. Thus $\sup\{a\}$ exists and $a = \sup\{a\}$.
2.
   - ($\Longrightarrow$):
     Suppose that $a \leq b$. Since $b \leq b$, we have that for each $c \in \{a, b\}$, $c \leq b$. Hence $b \in \text{lb}\{a, b\}$. Let $c \in \text{ub}\{a, b\}$. Then $b \leq c$. Since $c \in \text{ub}\{a, b\}$ is arbitrary, we have that for each $c \in \text{ub}\{a, b\}$, $b \leq c$. Hence $b \in \text{supSet}\{a, b\}$ and $b = \sup\{a, b\}$.
   - ($\Longleftarrow$):
     Suppose that $\sup\{a, b\}$ exists and $b = \sup\{a, b\}$. Then $b \in \text{ub}\{a, b\}$. Therefore $a \leq b$.
3. Since for each $x \in \downarrow a$, $x \leq a$, we have that $a \in \text{ub} \downarrow a$. Since $a \in \downarrow a$, we have that for each $x \in \text{ub} \downarrow a$, $a \leq x$. Hence $a = \sup \downarrow a$.
4. Similar to (3).

□

Exercise 2.7.3.7. Let $(X, \leq)$ be a poset. Then for each $a, b \in X$, $\sup\{a, b\}$ exists iff for each $n \in \mathbb{N}$ and $(a_k)_{k=1}^n \in X^n$, $\sup\{a_1, \ldots, a_n\}$ exists and

$$\sup\{a_1, \ldots, a_n\} = \begin{cases} a_1, & n = 1 \\ \sup\left\{\sup\{a_1, \ldots a_{n-1}\}, a_n\right\}, & n \geq 2. \end{cases}$$

Proof.

- ($\Longrightarrow$):
  Suppose that for each $a, b \in X$, $\sup\{a, b\}$ exists. Let $n \in \mathbb{N}$ and $(a_k)_{k=1}^n \in X^n$.
  - **Base Case:**
    Suppose that $n = 1$. Exercise 2.7.3.6 implies that $\sup\{a_1\}$ exists and $\sup\{a_1\} = a_1$.
  - **Induction Step:**
    Suppose that $n \geq 2$ and $\sup\{a_1, \ldots, a_{n-1}\}$ exists. Then by assumption, $\sup\left\{\sup\{a_1, \ldots a_{n-1}\}, a_n\right\}$ exists. Exercise 2.7.3.5 then implies that $\sup\{a_1, \ldots, a_n\}$ exists and

$$\begin{aligned} \sup\{a_1, \ldots, a_n\} &= \sup\left\{\{a_1, \ldots a_{n-1}\} \cup \{a_n\}\right\} \\ &= \sup\left\{\sup\{a_1, \ldots a_{n-1}\}, \sup\{a_n\}\right\} \\ &= \sup\left\{\sup\{a_1, \ldots a_{n-1}\}, a_n\right\}. \end{aligned}$$

  By induction, for each $n \in \mathbb{N}$ and $(a_k)_{k=1}^n \in X^n$, $\sup\{a_1, \ldots, a_n\}$ exists and

$$\sup\{a_1, \ldots, a_n\} = \begin{cases} a_1, & n = 1 \\ \sup\left\{\sup\{a_1, \ldots a_{n-1}\}, a_n\right\}, & n \geq 2. \end{cases}$$

- ($\Longleftarrow$):
  Clear.

□

Exercise 2.7.3.8. Let $(X, \leq)$ be a poset. Then

1. for each $a, b \in \operatorname{ub} X$, $a = b$,
2. for each $a \in \operatorname{ub} X$, $\sup X = a$,
3. for each $a, b \in \operatorname{lb} X$, $a = b$,
4. for each $a \in \operatorname{lb} X$, $\inf X = a$

Proof.

1. Let $a, b \in \operatorname{ub} X$. Since $a \in X$ and $b \in \operatorname{ub} X$, $a \leq b$. Similarly, $b \leq a$. Hence $a = b$.
2. Let $a, b \in \operatorname{ub} X$. the previous part implies that $a = b$. Since $b \in \operatorname{ub} X$ is arbitrary, we have that for each $b \in \operatorname{ub} X$, $a \leq b$. Thus $a = \sup X$.
3. Similar to (1) ref.
4. Similar to (2) ref.

□

Exercise 2.7.3.9. Let $(X, \leq_X)$ and $(Y, \leq_Y)$ be posets, $f : X \to Y$ and $g : Y \to X$. Suppose that (f, g) is a monotone Galois connection. Then

1. for each $A \subset X$, if $\sup(A, \leq_X)$ exists, then $\sup(f(A), \leq_Y)$ exists and $f(\sup(A, \leq_X)) = \sup(f(A), \leq_Y)$.
2. for each $B \subset Y$, if $\inf(B, \leq_Y)$ exists, then $g(\inf(B, \leq_Y)) = \inf(g(B), \leq_X)$.FIX!!!

Proof.

1. Suppose that $\sup(A, \leq_X)$ exists. Then $\operatorname{supSet}(A, \leq_X) \neq \varnothing$. Exercise 2.7.2.8 then implies that
$$\begin{aligned} \varnothing &\neq f(\operatorname{supSet}(A, \leq_X)) \\ &\subset \operatorname{supSet}(f(A), \leq_Y). \end{aligned}$$
Hence $\sup(f(A), \leq_Y)$ exists and $f(\sup(A, \leq_X)) = \sup(f(A), \leq_Y)$.
2. ???

□

Exercise 2.7.3.10. Let $(X, \leq_X)$ and $(Y, \leq_Y)$ be posets, $f : X \to Y$ and $g : Y \to X$. Suppose that (f, g) is a monotone Galois connection. Then

1. for each $x \subset X$, $f(x) = \inf(g^{-1}(\uparrow(x, \leq_X)), \leq_Y)$
2. for each $y \in Y$, $g(y) = \sup(f^{-1}(\downarrow(y, \leq_Y)), \leq_X)$

Proof.

1. Let $x \in X$. We note that for each $y \in Y$,
$$\begin{aligned} y \in g^{-1}(\uparrow(x, \leq_X)) &\iff g(y) \in \uparrow(x, \leq_X) \\ &\iff x \leq_X g(y) \\ &\iff f(x) \leq_Y y \quad \text{(since } (f, g) \text{ is a Galois connection)} \\ &\iff y \in \uparrow(f(x), \leq_Y). \end{aligned}$$
Thus $g^{-1}(\uparrow(x, \leq_X)) = \uparrow(f(x), \leq_Y)$. Exercise 2.7.3.6 then implies that
$$\begin{aligned} \inf(g^{-1}(\uparrow(x, \leq_X)), \leq_Y) &= \inf(\uparrow(f(x), \leq_Y), \leq_Y) \\ &= f(x). \end{aligned}$$
Since $x \in X$ is arbitrary, we have that for each $x \in X$, $f(x) = \inf(g^{-1}(\uparrow(x, \leq_X)), \leq_Y)$.

2. Let $y \in Y$. Since (f, g) is a $(\leq_X, \leq_Y)$-monotone Galois connection, Exercise B.0.0.3 ref prev ex in proset galois connection section implies that (g, f) is a $(\leq_Y^*, \leq_X^*)$-monotone Galois connection. The previous part and exercises in previous section on proset upper bounds implies that

$$\begin{aligned} g(y) &= \inf(f^{-1}(\uparrow(y, \leq_Y^*)), \leq_X^*) \\ &= \sup(f^{-1}(\downarrow(y, \leq_Y)), \leq_X). \end{aligned}$$

Since $y \in Y$ is arbitrary, we have that for each $y \in Y$, $g(y) = \sup(f^{-1}(\downarrow(y, \leq_Y)), \leq_X)$.

$\square$

Part II

Algebraic Structures

Chapter 3

Semigroups, Monoids and Groups

maybe move groups/rings etc after complete lattices? two choices: a) cover complete lattices after order and before other structures

(including lattices?) so that we can define the 'subgroup/ring/lattice generated by a subset' or a 'subquiver genereated by a sub set of vertices/edges' using the ideas of galois connection between two posets, one of which has extra structure b) cover complete lattices

after algebraic structures with lattices and note that we can capture all the subobjects with the notion of galois connection between complete lattices and give groups, rings etc as fundamental examples of complete lattices.
idk which is best option

3.1 Semigroups

3.2 Monoids

Definition 3.2.0.1. Let G be a set and $* : G \times G \to G$ (we write ab in place of $*(a, b)$). Then

1. $*$ is called a **binary operation** on G
2. $*$ is said to be **associative** if for each $x, y, z \in G$, $(xy)z = x(yz)$
3. $*$ is said to be **commutative** if for each $x, y \in G$, $xy = yx$

Definition 3.2.0.2. Let G be a set, $* : G \times G \to G$, $e, x, y \in G$. Then e is said to be an **identity element** if for each $x \in G$, $ex = xe = x$.

Definition 3.2.0.3. Let G be a set and $* : G \times G \to G$. Then G is said to be a **monoid** if

1. $*$ is associative
2. there exits $e \in G$ such that e is an identity element.

Exercise 3.2.0.4. Let G be a monoid. Then the identity element is unique.

Proof. Let $e, f \in G$. Suppose that e and f are identity elements. Then $e = ef = f$. □

Note 3.2.0.5. Unless otherwise specified, we will denote the identity element of a monoid by e.

Definition 3.2.0.6. Let G be a monoid, X a set and $* : G \times X \to X$ (we write gx in place of $*(g, x)$). Then $*$ is said to be a **monoid action** of G on X if for each $g, h \in G$ and $x \in X$,

1. $(gh)x = g(hx)$
2. $ex = x$

3.3 Groups

3.3.1 Direct Products

Definition 3.3.1.1. Let G, H be groups. Define a product $* : (G \times H) \times (G \times H) \to G \times H$ by

$$(x_1, y_1) * (x_2, y_2) = (x_1x_2, y_1y_2)$$

Then $(G \times H, *)$ is called the **direct product of G and H**.

Exercise 3.3.1.2. Let G, H be groups. Then the direct product $G \times H$ is a group.

Proof. Clear. □

Definition 3.3.1.3. Let G, H be groups. Define $\pi_G : G \times H \to G$ and $\pi_H : G \times H \to H$ by $\pi_G(x, y) = x$ and $\pi_H(x, y) = y$. Then π_G and π_H are respectively called the **projection maps onto G and H**.

Exercise 3.3.1.4. Let G, H be groups. Then

1. $\pi_G : G \times H \to G$ and $\pi_H : G \times H \to H$ are homomorphisms
2. $\ker \pi_G \cong H$ and $\ker \pi_H \cong G$

Proof.

1. Clear
2. Define $\iota_G : G \to \ker \pi_H$ by
$$\iota_G(x) = (x, e_H)$$
Then ι_G is an isomorphism. Similarly, we can define $\iota_H : H \to \ker \pi_G$ and show that it is an isomorphism.

□

Definition 3.3.1.5. Let G, H, K be groups, $\phi \in \mathrm{Hom}(G, K)$ and $\psi \in \mathrm{Hom}(H, K)$. We define $\phi \times \psi : G \times H \to K$ by $\phi \times \psi(x, y) = \phi(x)\psi(y)$

Exercise 3.3.1.6. Let G, H, K be groups, $\phi \in \mathrm{Hom}(G, K)$ and $\psi \in \mathrm{Hom}(H, K)$. If K is abelian, then $\phi \times \psi \in Hom(G \times H, K)$.

Proof. Let $x_1, x_2 \in G$ and $y_1, y_2 \in H$. Then

$$\begin{aligned}
\phi \times \psi[(x_1, y_1)(x_2, y_2)] &= \phi \times \psi(x_1x_2, y_1y_2) \\
&= \phi(x_1x_2)\psi(y_1y_2) \\
&= \phi(x_1)\phi(x_2)\psi(y_1)\psi(y_2) \\
&= \phi(x_1)\psi(y_1)\phi(x_2)\psi(y_2) \\
&= [\phi \times \psi(x_1, y_1)][\phi \times \psi(x_2, y_2)]
\end{aligned}$$

□

Exercise 3.3.1.7. Let G, H, K be groups and $\phi \in \mathrm{Hom}(G \times H, K)$. Then there exist $\phi_G \in \mathrm{Hom}(G, K)$, $\phi_H \in \mathrm{Hom}(H, K)$ such that $\phi_G \times \phi_H = \phi$.

Proof. Suppose that K is abelian. Define $\iota_G \in \mathrm{Hom}(G, \ker \pi_H)$ and $\iota_H \in \mathrm{Hom}(H, \ker \pi_G)$ as in part (2) of Exercise 3.3.1.4 Define $\phi_G \in \mathrm{Hom}(G, K)$ and $\phi_H \in \mathrm{Hom}(H, K)$ by $\phi_G = \phi \circ \iota_G$ and $\phi_H = \phi \circ \iota_H$. Let $(x, y) \in G \times H$. Then

$$\begin{aligned}
\phi_G \times \phi_H(x, y) &= \phi_G(x)\phi_H(y) \\
&= \phi \circ \iota_G(x)\phi \circ \iota_H(y) \\
&= \phi(x, e_H)\phi(e_G, y) \\
&= \phi(x, y)
\end{aligned}$$

So $\phi = \phi_G \times \phi_H$ □

Chapter 4

Rings

4.1 Introduction

Definition 4.1.0.1. Let R be a set and $+, * : R \times R \to R$ (we write $a + b$ and ab in place of $+(a, b)$ and $*(a, b)$ respectively). Then R is said to be a **ring** if for each $a, b, c \in R$,

1. R is an abelian group with respect to $+$. The identity element with respect to $+$ is denoted by 0.
2. R is a monoid with respect to $*$. The identity element of R with respect to $*$ is denoted 1.
3. R is commutative with respect to $*$.
4. $*$ distributes over $+$.

Definition 4.1.0.2. Let R be a ring and $I \subset R$. Then I is said to be an **ideal** of R if for each $a \in R$ and $x, y \in I$,

1. $x + y \in I$
2. $ax \in I$

Definition 4.1.0.3. Let R be a ring and $A, B \subset R$. We define the **product** of A and B, denoted AB, to be

$$AB = \left\{ \sum_{i=1}^{n} a_i b_i : a_i \in A, b_i \in B, n \in \mathbb{N} \right\}$$

Exercise 4.1.0.4. Let R be a ring and $I \subset R$. Then I is an ideal of R iff $RI \subset I$.

Proof. Suppose that $RI \subset I$. Let $a \in R$ and $x, y \in I$. Then by assumption $x + y = 1x + 1y \in I$ and $ax \in I$. So I is an ideal of R Conversely, suppose that I is an ideal of R. Let $a_1, \cdots, a_n \in R$ and $x_1, \cdots, x_n \in I$. Then by assumption, for each $i = 1, \cdots, n$, $a_i x_i \in I$ and therefore $\sum_{i=1}^{n} a_i b_i \in I$. Hence $RI \subset I$. $\square$

Chapter 5

Modules

5.1 Introduction

Definition 5.1.0.1. Let R be a ring, M a set, $+ : M \times M \to M$ and $* : R \times M \to M$ (we write rx in place of $*(r, x)$). Then M is said to be an R**-module** if

1. M is an abelian group with respect to $+$. The identity element of M with respect to $+$ is denoted by 0.
2. for each $r \in R$, $*(r, \cdot)$ is a group endomorphism of M
3. for each $x \in M$, $*(\cdot, x)$ is a group homomorphism from R to M
4. $*$ is a monoid action of R on M

Note 5.1.0.2. For the remainder of this section, we assume that R is a commutative ring.

Exercise 5.1.0.3. Let M be an R-module. Then for each $r \in R$ and $x \in M$,

1. $r0 = 0$
2. $0x = 0$
3. $(-1)x = -x$

Proof. Let $r \in R$ and $x \in M$. Then

1. $$\begin{aligned} r0 &= r(0+0) \\ &= r0 + r0 \end{aligned}$$

   which implies that $r0 = 0$.

2. $$\begin{aligned} 0x &= (0+0)x \\ &= 0x + 0x \end{aligned}$$

   which implies that $0x = 0$.

3. $$\begin{aligned} (-1)x + x &= (-1)x + 1x \\ &= (-1+1)x \\ &= 0x \\ &= 0 \end{aligned}$$

   which implies that $(-1)x = -x$.

□

Definition 5.1.0.4. Let M an R-module and $N \subset M$. Then N is said to be a **submodule** of M if for each $r \in R$ and $x, y \in N$, we have that $rx \in N$ and $x + y \in N$.

Definition 5.1.0.5. Let M be an R-module. We define $\mathcal{S}(M) = \{N \subset M : N \text{ is a submodule of } M\}$.

Exercise 5.1.0.6. Let M be an R-module and $N \in \mathcal{S}(M)$. Then N is a subgroup of M.

Proof. Let $x, y \in M$. Then $x - y = 1x + (-1)y \in N$. So N is a subgroup of M. □

Definition 5.1.0.7. Let M be an R-module and $N \in \mathcal{S}(M)$. We define

1. $M/N = \{x + N : x \in M\}$
2. $+ : M/N \times M/N \to M/N$ by
$$(x + N) + (y + N) = (x + y) + N$$
3. $* : R \times M/N \to M/N$ by
$$r(x + N) = (rx) + N$$

Under these operations (see next exercise), M/N is an R-module known as the **quotient module** of M by N.

Exercise 5.1.0.8. Let M be an R-module and $N \in \mathcal{S}(M)$. Then

1. the monoid action defined above is well defined
2. the quotient M/N is an R-module

Proof.

1. Let $r \in R$ and $x + N, y + N \in M/N$. Recall from group theory that $x + N = y + N$ iff $x - y \in N$. Suppose that $x + N = y + N$. Then $x - y \in N$ and there exists $n \in N$ such that $x - y = n$. Therefore
$$\begin{aligned} rx - ry &= r(x - y) \\ &= rn \\ &\in N \end{aligned}$$
So $rx + N = ry + N$.
2. Properties (1) - (4) in the definition of a module are easily shown to be satisfied for M/N since they are true for M.

□

Definition 5.1.0.9. Let M and N be R-modules and $\phi : M \to N$. Then ϕ is said to be a **module homomorphism** if for each $r \in R$ and $x, y \in M$

1. $\phi(rx) = r\phi(x)$
2. $\phi(x + y) = \phi(x) + \phi(y)$

Exercise 5.1.0.10. Let M and N be R-modules and $\phi : M \to N$. Then ϕ is a iff for each $r \in R$ and $x, y \in M$, $\phi(x+ry) = \phi(x)+r\phi(y)$.

Proof. Clear. □

Exercise 5.1.0.11. Let M and N be R-modules and $\phi : M \to N$ a homomorphism. Then

1. $\ker \phi$ is a submodule of M
2. $\operatorname{Im} \phi$ is a submodule of N

Proof. Let $r \in R$, $x, y \in \ker \phi$ and $w, z \in \operatorname{Im} \phi$. Then

1.
$$\begin{aligned}\phi(rx) &= r\phi(x)\\ &= r0\\ &= 0\end{aligned}$$

So $rx \in \ker\phi$. Group theory tells us that $\ker\phi$ is a subgroup of M, so $x+y \in \ker\phi$. Hence $\ker\phi$ is a submodule of M.

2. Similar.

□

Definition 5.1.0.12. Let M be an R-module and $A \subset M$. We define the **submodule of M generated by** A, denoted $\operatorname{span}(A)$, to be
$$\operatorname{span}(A) = \bigcap_{N \in \mathcal{S}(M)} N$$

Exercise 5.1.0.13. Let M be an R-module and $A \subset M$. Then $\operatorname{span}(A) \in \mathcal{S}(M)$

Proof. Let $r \in R$ and $x, y \in \operatorname{span}(A)$. Basic group theory tells us that $\operatorname{span}(A)$ is a subgroup of M. So $x+y \in \operatorname{span}(A)$. For $N \in \mathcal{S}(M)$, by definition we have $x \in N$ and therefore $rx \in N$. So $rx \in \operatorname{span}(A)$. Hence $\operatorname{span}(A)$ is a submodule of M. □

Exercise 5.1.0.14. Let M be an R-module and $A \subset M$. If $A \neq \varnothing$, then
$$\operatorname{span}(A) = \left\{ \sum_{i=1}^{n} r_i a_i : r_i \in R, a_i \in A, n \in \mathbb{N} \right\}$$

Proof. Clearly □

Definition 5.1.0.15. Let M

Chapter 6

Vector Spaces

Chapter 7

Fields

Chapter 8

Quivers

maybe move after lattice chapter, to be able to talk about subquivers generated by a subset of vertices and arrows

talk about quivers (i.e. categories without compostion) Defien subquivers, subquivers generated by a subset of vertices and arrows in complete lattice terms (i.e. using galois connections). Define prod and coprod quivers, quotient quivers etc,

8.1 Introduction

Definition 8.1.0.1. Let V, E, s, t be sets. Then (V, E, s, t) is said to be a **quiver** if $s, t : E \to V$.

Definition 8.1.0.2. Let $\mathcal{Q}$ be a quiver. Write $\mathcal{Q} = (V, E, s, t)$. We define the

- **vertices of** $\mathcal{Q}$, denoted $V_{\mathcal{Q}}$, by $V_{\mathcal{Q}} := V$.
- **edges of** $\mathcal{Q}$, denoted $E_{\mathcal{Q}}$, by $E_{\mathcal{Q}} := E$.
- **source function of** $\mathcal{Q}$, denoted $s_{\mathcal{Q}}$, by $s_{\mathcal{Q}} := s$.
- **target function of** $\mathcal{Q}$, denoted $t_{\mathcal{Q}}$, by $t_{\mathcal{Q}} := t$.

Note 8.1.0.3. Let $\mathcal{Q}$ be a quiver. When the context is clear, we write V, E, s, t in place of $V_{\mathcal{Q}}$, $E_{\mathcal{Q}}$, $s_{\mathcal{Q}}$, $t_{\mathcal{Q}}$ resepctively.

Definition 8.1.0.4. We define the **empty quiver**, denoted $\mathcal{Q}_0$ by $\mathcal{Q}_0 := (\varnothing, \varnothing, \varnothing, \varnothing)$.

Exercise 8.1.0.5. Define $\mathcal{Q}$ by $\mathcal{Q} := (\varnothing, \varnothing, \varnothing, \varnothing)$. Then $\mathcal{Q}$ is a quiver.

Proof. Since $\varnothing : \varnothing \to \varnothing$, $\mathcal{Q}$ is a quiver. □

Definition 8.1.0.6. Let $\mathcal{Q}$ and $\mathcal{L}$ be quivers, $\phi_v : V_{\mathcal{Q}} \to V_{\mathcal{L}}$ and $\phi_e : E_{\mathcal{Q}} \to E_{\mathcal{L}}$. Then (ϕ_e, ϕ_v) is said to be a $(\mathcal{Q}, \mathcal{L})$**-quiver morphism** if $s_{\mathcal{L}} \circ \phi_e = \phi_v \circ s_{\mathcal{Q}}$ and $t_{\mathcal{L}} \circ \phi_e = \phi_v \circ t_{\mathcal{Q}}$, i.e. the following diagrams commute:

$$\begin{tikzcd} E_{\mathcal{Q}} \arrow[r, "\phi_e"] \arrow[d, "s_{\mathcal{Q}}"'] & E_{\mathcal{L}} \arrow[d, "s_{\mathcal{L}}"] \\ V_{\mathcal{Q}} \arrow[r, "\phi_v"'] & V_{\mathcal{L}} \end{tikzcd} \qquad \begin{tikzcd} E_{\mathcal{Q}} \arrow[r, "\phi_e"] \arrow[d, "t_{\mathcal{Q}}"'] & E_{\mathcal{L}} \arrow[d, "t_{\mathcal{L}}"] \\ V_{\mathcal{Q}} \arrow[r, "\phi_v"'] & V_{\mathcal{L}} \end{tikzcd}$$

8.2 Subquivers

Definition 8.2.0.1. Let $\mathcal{Q}$ be a quiver.

- Let V, E, s, t be sets. Then (V, E, s, t) is said to be a **subquiver of** $\mathcal{Q}$ if
  1. (V, E, s, t) is a quiver,

2. $V \subset V_{\mathcal{Q}}$ and $E \subset E_{\mathcal{Q}}$,
3. (ι_V, ι_E) is a $((V, E, s, t), \mathcal{Q})$-quiver morphism.

- We define $\mathrm{Subquiv}(\mathcal{Q}) := \{\mathcal{L} : \mathcal{L} \text{ is a subquiver of } \mathcal{Q}\}$.
- We define the **subquiver relation on** $\mathrm{Subquiv}(\mathcal{Q})$, denoted $\leq_{\mathrm{Quiv}} \subset \mathrm{Subquiv}(\mathcal{Q}) \times \mathrm{Subquiv}(\mathcal{Q})$, by $\mathcal{L}_1 \leq_{\mathrm{Quiv}} \mathcal{L}_2$ iff $\mathcal{L}_1$ is a subquiver of $\mathcal{L}_2$.

Note 8.2.0.2. When the context is clear, we may write $\leq$ in place of $\leq_{\mathrm{Quiv}}$.

Exercise 8.2.0.3. $\mathcal{Q}$ be a quiver. Then $(\mathrm{Subquiv}(\mathcal{Q}), \leq_{\mathrm{Quiv}})$ is a poset.

Proof. FINISH!!! □

Exercise 8.2.0.4. Let $\mathcal{Q}$ and $\mathcal{L}$ be quivers. Then $\mathcal{L} \in \mathrm{Subquiv}(\mathcal{Q})$ iff

1. $V_{\mathcal{L}} \subset V_{\mathcal{Q}}$ and $E_{\mathcal{L}} \subset E_{\mathcal{Q}}$
2. $s_{\mathcal{Q}}|_{E_{\mathcal{L}}} = s_{\mathcal{L}}$ and $t_{\mathcal{Q}}|_{E_{\mathcal{L}}} = t_{\mathcal{L}}$

Proof.

- $(\implies)$:
  Suppose that $\mathcal{L} \in \mathrm{Subquiv}(\mathcal{Q})$. Then $V_{\mathcal{L}} \subset V_{\mathcal{Q}}$, $E_{\mathcal{L}} \subset E_{\mathcal{Q}}$ and $(\iota_{V_{\mathcal{L}}}, \iota_{E_{\mathcal{L}}})$ is a $(\mathcal{L}, \mathcal{Q})$-quiver morphism. Then $s_{\mathcal{Q}} \circ \iota_{E_{\mathcal{L}}} = \iota_{V_{\mathcal{L}}} \circ s_{\mathcal{L}}$ and $t_{\mathcal{Q}} \circ \iota_{E_{\mathcal{L}}} = \iota_{V_{\mathcal{L}}} \circ t_{\mathcal{L}}$, i.e. the following diagrams commute:

$$\begin{tikzcd} E_{\mathcal{L}} \arrow[r, "\iota_{E_{\mathcal{L}}}"] \arrow[d, "s_{\mathcal{L}}"'] & E_{\mathcal{Q}} \arrow[d, "s_{\mathcal{Q}}"] \\ V_{\mathcal{L}} \arrow[r, "\iota_{V_{\mathcal{L}}}"'] & V_{\mathcal{Q}} \end{tikzcd} \qquad \begin{tikzcd} E_{\mathcal{L}} \arrow[r, "\iota_{E_{\mathcal{L}}}"] \arrow[d, "t_{\mathcal{L}}"'] & E_{\mathcal{Q}} \arrow[d, "t_{\mathcal{Q}}"] \\ V_{\mathcal{L}} \arrow[r, "\iota_{V_{\mathcal{L}}}"'] & V_{\mathcal{Q}} \end{tikzcd}$$

  Therefore

$$\begin{aligned} s_{\mathcal{Q}}|_{E_{\mathcal{L}}} &= s_{\mathcal{Q}} \circ \iota_{E_{\mathcal{L}}} \\ &= \iota_{V_{\mathcal{L}}} \circ s_{\mathcal{L}} \\ &= s_{\mathcal{L}} \end{aligned}$$

  and

$$\begin{aligned} t_{\mathcal{Q}}|_{E_{\mathcal{L}}} &= t_{\mathcal{Q}} \circ \iota_{E_{\mathcal{L}}} \\ &= \iota_{V_{\mathcal{L}}} \circ t_{\mathcal{L}} \\ &= t_{\mathcal{L}}. \end{aligned}$$

- $(\impliedby)$:
  Suppose that

  1. $V_{\mathcal{L}} \subset V_{\mathcal{Q}}$ and $E_{\mathcal{L}} \subset E_{\mathcal{Q}}$,
  2. $s_{\mathcal{Q}}|_{E_{\mathcal{L}}} = s_{\mathcal{L}}$ and $t_{\mathcal{Q}}|_{E_{\mathcal{L}}} = t_{\mathcal{L}}$.

  Then

$$\begin{aligned} s_{\mathcal{Q}} \circ \iota_{E_{\mathcal{L}}} &= s_{\mathcal{Q}}|_{E_{\mathcal{L}}} \\ &= s_{\mathcal{L}} \\ &= \iota_{V_{\mathcal{L}}} \circ s_{\mathcal{L}} \end{aligned}$$

  and

$$\begin{aligned} t_{\mathcal{Q}} \circ \iota_{E_{\mathcal{L}}} &= t_{\mathcal{Q}}|_{E_{\mathcal{L}}} \\ &= t_{\mathcal{L}} \\ &= \iota_{V_{\mathcal{L}}} \circ t_{\mathcal{L}}. \end{aligned}$$

□

Exercise 8.2.0.5. Let $\mathcal{Q}$ be a quiver, $V \subset V_{\mathcal{Q}}$ and $E \subset E_{\mathcal{Q}}$. Then $(V, E, s_{\mathcal{Q}}|_E, t_{\mathcal{Q}}|_E) \in \mathrm{Subquiv}(\mathcal{Q})$ iff $s_{\mathcal{Q}}(E) \subset V$ and $t_{\mathcal{Q}}(E) \subset V$.

Proof. Set $\mathcal{L} := (V, E, s_{\mathcal{Q}}|_E, t_{\mathcal{Q}}|_E)$.

- $(\implies)$:
  Suppose that $\mathcal{L} \in \mathrm{Subquiv}(\mathcal{Q})$. Then $\mathcal{L}$ is a quiver and in particular, $s_{\mathcal{Q}}|_E : E \to V$ and $t_{\mathcal{Q}}|_E : E \to V$. Therefore $s_{\mathcal{Q}}(E) \subset V$ and $t_{\mathcal{Q}}(E) \subset V$.

- $(\impliedby)$
  Suppose that $s_{\mathcal{Q}}(E) \subset E$ and $t_{\mathcal{Q}}(E) \subset E$.

  1. Since $s_{\mathcal{Q}}(E) \subset V$ and $t_{\mathcal{Q}}(E) \subset V$, we have that $s_{\mathcal{Q}}|_E : E \to V$ and $t_{\mathcal{Q}}|_E : E \to V$. Therefore $\mathcal{L}$ is a quiver.
  2. (a) By assumption, $V_{\mathcal{L}} \subset V_{\mathcal{Q}}$ and $E_{\mathcal{L}} \subset E_{\mathcal{Q}}$.
     (b) By definition,

     $$\begin{aligned} s_{\mathcal{Q}}|_{E_{\mathcal{L}}} &= s_{\mathcal{Q}}|_E \\ &= s_{\mathcal{L}} \end{aligned}$$

     and

     $$\begin{aligned} t_{\mathcal{Q}}|_{E_{\mathcal{L}}} &= t_{\mathcal{Q}}|_E \\ &= t_{\mathcal{L}} \end{aligned}$$

     The prev ex then implies that $\mathcal{L} \in \mathrm{Subquiv}(\mathcal{Q})$.

□

Exercise 8.2.0.6. Let $\mathcal{Q}$ be a quiver and $(\mathcal{L}_\alpha)_{\alpha \in A} \subset \mathrm{Subquiv}(\mathcal{Q})$. Define V, E by $V := \bigcap_{\alpha \in A} V_{\mathcal{L}_\alpha}$ and $E := \bigcap_{\alpha \in A} E_{\mathcal{L}_\alpha}$. Then

1. $(V, E, s_{\mathcal{Q}}|_E, t_{\mathcal{Q}}|_E) \in \mathrm{Subquiv}(\mathcal{Q})$
2. $\inf((\mathcal{L}_\alpha)_{\alpha \in A}, \leq_{\mathrm{Quiv}}) = (V, E, s_{\mathcal{Q}}|_E, t_{\mathcal{Q}}|_E)$

Proof. Set $\mathcal{L} := (V, E, s_{\mathcal{Q}}|_E, t_{\mathcal{Q}}|_E)$. Since for each $\alpha \in A$, $\mathcal{L}_{\alpha_0} \in \mathrm{Subquiv}(\mathcal{Q})$, a prev ex implies that for each $\alpha \in A$, $V_{\mathcal{L}_\alpha} \subset V_{\mathcal{Q}}$, $E_{\mathcal{L}_\alpha} \subset E_{\mathcal{Q}}$, $s_{\mathcal{Q}}|_{E_{\mathcal{L}_\alpha}} = s_{\mathcal{L}_\alpha}$ and $t_{\mathcal{Q}}|_{E_{\mathcal{L}_\alpha}} = t_{\mathcal{L}_\alpha}$. Therefore

$$\begin{aligned} V &= \bigcap_{\alpha \in A} V_{\mathcal{L}_\alpha} \\ &\subset V_{\mathcal{Q}} \end{aligned}$$

and

$$\begin{aligned} E &= \bigcap_{\alpha \in A} E_{\mathcal{L}_\alpha} \\ &\subset E_{\mathcal{Q}}. \end{aligned}$$

1. Let $\alpha_0 \in A$. We note that

   $$\begin{aligned} V &= \bigcap_{\alpha \in A} V_{\mathcal{L}_\alpha} \\ &\subset V_{\mathcal{L}_{\alpha_0}} \end{aligned}$$

   and

   $$\begin{aligned} E &= \bigcap_{\alpha \in A} E_{\mathcal{L}_\alpha} \\ &\subset E_{\mathcal{L}_{\alpha_0}}. \end{aligned}$$

Since $V_{\mathcal{L}_{\alpha_0}} \subset V_{\mathcal{Q}}$, $E_{\mathcal{L}_{\alpha_0}} \subset E_{\mathcal{Q}}$, $s_{\mathcal{Q}}|_{E_{\mathcal{L}_{\alpha_0}}} = s_{\mathcal{L}_{\alpha_0}}$ and $t_{\mathcal{Q}}|_{E_{\mathcal{L}_{\alpha_0}}} = t_{\mathcal{L}_{\alpha_0}}$, we have that

$$\begin{aligned}(V_{\mathcal{L}_{\alpha_0}}, E_{\mathcal{L}_{\alpha_0}}, s_{\mathcal{Q}}|_{E_{\mathcal{L}_{\alpha_0}}}, t_{\mathcal{Q}}|_{E_{\mathcal{L}_{\alpha_0}}}) &= (V_{\mathcal{L}_{\alpha_0}}, E_{\mathcal{L}_{\alpha_0}}, s_{\mathcal{L}_{\alpha_0}}, t_{\mathcal{L}_{\alpha_0}}) \\ &= \mathcal{L}_{\alpha_0} \\ &\in \mathrm{Subquiv}(\mathcal{Q}).\end{aligned}$$

The prev ex then implies that $s_{\mathcal{Q}}(E_{\mathcal{L}_{\alpha_0}}) \subset V_{\mathcal{L}_{\alpha_0}}$ and $t_{\mathcal{Q}}(E_{\mathcal{L}_{\alpha_0}}) \subset V_{\mathcal{L}_{\alpha_0}}$. Therefore

$$\begin{aligned}s_{\mathcal{Q}}(E) &= s_{\mathcal{Q}}\left(\bigcap_{\alpha \in A} E_{\mathcal{L}_\alpha}\right) \\ &\subset s_{\mathcal{Q}}(E_{\mathcal{L}_{\alpha_0}}) \\ &\subset V_{\mathcal{L}_{\alpha_0}}\end{aligned}$$

and

$$\begin{aligned}t_{\mathcal{Q}}(E) &= t_{\mathcal{Q}}\left(\bigcap_{\alpha \in A} E_{\mathcal{L}_\alpha}\right) \\ &\subset t_{\mathcal{Q}}(E_{\mathcal{L}_{\alpha_0}}) \\ &\subset V_{\mathcal{L}_{\alpha_0}}.\end{aligned}$$

Since $\alpha_0 \in A$ is arbitrary, we have that for each $\alpha \in A$, $s_{\mathcal{Q}}(E) \subset V_{\mathcal{L}_\alpha}$ and $t_{\mathcal{Q}}(E) \subset V_{\mathcal{L}_\alpha}$. Hence

$$\begin{aligned}s_{\mathcal{Q}}(E) &\subset \bigcap_{\alpha \in A} V_{\mathcal{L}_\alpha} \\ &= V\end{aligned}$$

and

$$\begin{aligned}t_{\mathcal{Q}}(E) &\subset \bigcap_{\alpha \in A} V_{\mathcal{L}_\alpha} \\ &= V.\end{aligned}$$

The prev ex then implies that $\mathcal{L} \in \mathrm{Subquiv}(\mathcal{Q})$.

2. • Let $\alpha \in A$. Since

$$\begin{aligned}V_{\mathcal{L}} &= V \\ &\subset V_{\mathcal{L}_\alpha}\end{aligned}$$

and

$$\begin{aligned}E_{\mathcal{L}} &= E \\ &\subset E_{\mathcal{L}_\alpha},\end{aligned}$$

we have that

$$\begin{aligned}s_{\mathcal{L}} &= s_{\mathcal{Q}}|_{E_{\mathcal{L}}} \\ &= s_{\mathcal{Q}} \circ \iota_{E_{\mathcal{L}}}^{E_{\mathcal{Q}}} \\ &= s_{\mathcal{Q}} \circ (\iota_{E_{\mathcal{L}_\alpha}}^{E_{\mathcal{Q}}} \circ \iota_{E_{\mathcal{L}}}^{E_{\mathcal{L}_\alpha}}) \\ &= (s_{\mathcal{Q}} \circ \iota_{E_{\mathcal{L}_\alpha}}^{E_{\mathcal{Q}}}) \circ \iota_{E_{\mathcal{L}}}^{E_{\mathcal{L}_\alpha}} \\ &= s_{\mathcal{Q}}|_{E_{\mathcal{L}_\alpha}} \circ \iota_{E_{\mathcal{L}}}^{E_{\mathcal{L}_\alpha}} \\ &= s_{\mathcal{L}_\alpha} \circ \iota_{E_{\mathcal{L}}}^{E_{\mathcal{L}_\alpha}} \\ &= s_{\mathcal{L}_\alpha}|_{E_{\mathcal{L}}}.\end{aligned}$$

Similarly, $t_{\mathcal{L}} = t_{\mathcal{L}_\alpha}|_{E_{\mathcal{L}}}$. Therefore

(a) $V_{\mathcal{L}} \subset V_{\mathcal{L}_\alpha}$ and $E_{\mathcal{L}} \subset E_{\mathcal{L}_\alpha}$,
(b) $s_{\mathcal{L}_\alpha}|_{E_{\mathcal{L}}} = s_{\mathcal{L}}$ and $t_{\mathcal{L}_\alpha}|_{E_{\mathcal{L}}} = t_{\mathcal{L}}$.

A prev ex implies that $\mathcal{L} \in \mathrm{Subquiv}(\mathcal{L}_\alpha)$. Hence $\mathcal{L} \leq_{\mathrm{Quiv}} \mathcal{L}_\alpha$. Since $\alpha \in A$ is arbitrary, we have that for each $\alpha \in A$, $\mathcal{L} \leq_{\mathrm{Quiv}} \mathcal{L}_\alpha$. Thus $\mathcal{L} \in \mathrm{lb}((\mathcal{L}_\alpha)_{\alpha\in A}, \leq_{\mathrm{Quiv}})$.

- Let $\mathcal{L}' \in \mathrm{lb}((\mathcal{L}_\alpha)_{\alpha\in A}, \leq_{\mathrm{Quiv}})$ and $\alpha \in A$. Since $\mathcal{L}' \in \mathrm{lb}((\mathcal{L}_\alpha)_{\alpha\in A}, \leq_{\mathrm{Quiv}})$, we have that $\mathcal{L}' \leq_{\mathrm{Quiv}} \mathcal{L}_\alpha$. A prev ex then implies that $V_{\mathcal{L}'} \subset V_{\mathcal{L}_\alpha}$, $E_{\mathcal{L}'} \subset E_{\mathcal{L}_\alpha}$, $s_{\mathcal{L}_\alpha}|_{E_{\mathcal{L}'}} = s_{\mathcal{L}'}$ and $t_{\mathcal{L}_\alpha}|_{E_{\mathcal{L}'}} = t_{\mathcal{L}'}$. Since $\alpha \in A$ is arbitrary, we have that for each $\alpha \in A$, $V_{\mathcal{L}'} \subset V_{\mathcal{L}_\alpha}$, $E_{\mathcal{L}'} \subset E_{\mathcal{L}_\alpha}$, $s_{\mathcal{L}_\alpha}|_{E_{\mathcal{L}'}} = s_{\mathcal{L}'}$ and $t_{\mathcal{L}_\alpha}|_{E_{\mathcal{L}'}} = t_{\mathcal{L}'}$. Therefore $V_{\mathcal{L}'} \in \mathrm{lb}((V_{\mathcal{L}_\alpha})_{\alpha\in A}, \subset)$, $E_{\mathcal{L}'} \in \mathrm{lb}((E_{\mathcal{L}_\alpha})_{\alpha\in A}, \subset)$. Thus

$$\begin{aligned} V_{\mathcal{L}'} &\subset \inf((V_{\mathcal{L}_\alpha})_{\alpha\in A}, \subset) \\ &= \bigcap_{\alpha\in A} V_{\mathcal{L}_\alpha} \\ &= V. \end{aligned}$$

and

$$\begin{aligned} E_{\mathcal{L}'} &\subset \inf((E_{\mathcal{L}_\alpha})_{\alpha\in A}, \subset) \\ &= \bigcap_{\alpha\in A} E_{\mathcal{L}_\alpha} \\ &= E. \end{aligned}$$

Hence

$$\begin{aligned} s_{\mathcal{L}'} &= s_{\mathcal{Q}}|_{E_{\mathcal{L}'}} \\ &= s_{\mathcal{Q}} \circ \iota^{E_{\mathcal{Q}}}_{E_{\mathcal{L}'}} \\ &= s_{\mathcal{Q}} \circ (\iota^{E_{\mathcal{Q}}}_{E_{\mathcal{L}}} \circ \iota^{E_{\mathcal{L}}}_{E_{\mathcal{L}'}}) \\ &= (s_{\mathcal{Q}} \circ \iota^{E_{\mathcal{Q}}}_{E_{\mathcal{L}}}) \circ \iota^{E_{\mathcal{L}}}_{E_{\mathcal{L}'}} \\ &= s_{\mathcal{Q}}|_{E_{\mathcal{L}}} \circ \iota^{E_{\mathcal{L}}}_{E_{\mathcal{L}'}} \\ &= s_{\mathcal{L}} \circ \iota^{E_{\mathcal{L}}}_{E_{\mathcal{L}'}}. \end{aligned}$$

Similarly, $t_{\mathcal{L}'} = t_{\mathcal{L}} \circ \iota^{E_{\mathcal{L}}}_{E_{\mathcal{L}'}}$. a previous ex implies that $\mathcal{L}' \leq_{\mathrm{Quiv}} \mathcal{L}$. Since $\mathcal{L}' \in \mathrm{lb}((\mathcal{L}_\alpha)_{\alpha\in A}, \leq_{\mathrm{Quiv}})$ is arbitrary, we have that for each $\mathcal{L}' \in \mathrm{lb}((\mathcal{L}_\alpha)_{\alpha\in A}, \leq_{\mathrm{Quiv}})$, $\mathcal{L}' \leq_{\mathrm{Quiv}} \mathcal{L}$.

Thus $\inf((\mathcal{L}_\alpha)_{\alpha\in A}, \leq_{\mathrm{Quiv}}) = \mathcal{L}$.

□

Definition 8.2.0.7. Let $\mathcal{Q}$ be a quiver, $V \subset V_{\mathcal{Q}}$ and $E \subset E_{\mathcal{Q}}$. We define

- $\mathrm{Subquiv}(V, E, \mathcal{Q}) := \{\mathcal{L} \in \mathrm{Subquiv}(\mathcal{Q}) : V \subset V_{\mathcal{L}} \text{ and } E \subset E_{\mathcal{L}}\}$.
- the **subquiver of $\mathcal{Q}$ generated by V and E**, denoted $\mathrm{SubquivGen}(V, E, \mathcal{Q})$ by

$$\mathrm{SubquivGen}(V, E, \mathcal{Q}) := \inf(\mathrm{Subquiv}(V, E, \mathcal{Q}), \leq_{\mathrm{Quiv}}).$$

Definition 8.2.0.8. When the context is clear, we write $\mathrm{SubquivGen}(V, E)$ in place of $\mathrm{SubquivGen}(V, E, \mathcal{Q})$.

Chapter 9

Lattices

9.0.1 To Do

- maybe cover join/meet presemilattices, i.e. join/meet semilattices but with preorder instead of partial order, this would prbably need to move exercises around and rework stuff
- explain that we assume lattices are nonempty due to the convention in universal algebra (i guess for some slick way to state important results without having to make extra cases/exceptions?)

9.1 Introduction

9.1.1 Skew Semilattices

cite Leech, skew lattices in rings

Definition 9.1.1.1. Let X be a set and $\diamond : X \times X \to X$ a binary operator on X. Then

- $\diamond$ is said to be a **skew semilattice operator on** X if $\diamond$ is associative and idempotent.
- $(X, \diamond)$ is said to be a **skew semilattice** if
  1. $X \neq \varnothing$
  2. $\diamond$ is a skew semilattice operator on X

Definition 9.1.1.2. Let $(X, \diamond)$ be an skew semilattice. We define the

- **join preorder on** X **induced by** $\diamond$, denoted $\leq_\diamond^\vee \subset X \times X$, by $a \leq_\diamond^\vee b$ iff $(b \diamond a) \diamond b = b$
- **meet preorder on** X **induced by** $\diamond$, denoted $\leq_\diamond^\wedge \subset X \times X$, by $a \leq_\diamond^\wedge b$ iff $(a \diamond b) \diamond a = a$.

Exercise 9.1.1.3. Let $(X, \diamond)$ be an skew semilattice. Then

1. $\leq_\diamond^\vee$ is a preorder on X.
   Hint: If $cbc = c$ and $bab = b$, then
   (a) $c = (cb)(abc)(abc)$ and $c = cabc$
   (b) $cac = (ca)(ca)(bc)$
2. $\leq_\diamond^\wedge$ is a preorder on X.

Proof.

1. Let $a, b, c \in X$.

(a) Since $\diamond$ is idempotent, we have that

$$\begin{aligned}(a \diamond a) \diamond a &= a \diamond a \\ &= a\end{aligned}$$

and therefore $a \leq_\diamond^\vee a$.

(b) Suppose that $a \leq_\diamond^\vee b$ and $b \leq_\diamond^\vee c$. Then $(b \diamond a) \diamond b = b$ and $(c \diamond b) \diamond c = c$. Since $\diamond$ is associative and idempotent,

$$\begin{aligned}c &= cbc \\ &= cbabc \\ &= (cb)(abc) \\ &= (cb)(abc)(abc) \\ &= c(bab)cabc \\ &= cbcabc \\ &= (cbc)abc \\ &= cabc.\end{aligned}$$

Therefore

$$\begin{aligned}cac &= (ca)(c) \\ &= (ca)(cabc) \\ &= (ca)(ca)(bc) \\ &= (ca)(bc) \\ &= cabc \\ &= c.\end{aligned}$$

Hence $a \leq_\diamond^\vee c$.

So $\leq_\diamond^\vee$ is a preorder on X.

2. Similar to (1).

□

Exercise 9.1.1.4. Let $(X, \diamond)$ be an skew semilattice. Then

1. $(\leq_\diamond^\vee)^* = \leq_\diamond^\wedge$
2. $(\leq_\diamond^\wedge)^* = \leq_\diamond^\vee$

Proof.

1. Let $a, b \in X$. Then

$$\begin{aligned}a(\leq_\diamond^\vee)^* b &\iff b \leq_\diamond^\vee a \\ &\iff a \diamond b \diamond a = a \\ &\iff a \leq_\diamond^\wedge b.\end{aligned}$$

2. Similar to (1).

□

Exercise 9.1.1.5. Let $(X, \diamond)$ be an skew semilattice. Then for each $a, b \in X$,

1. $a \diamond b \in \mathrm{ub}(\{a, b\}, \leq_\diamond^\vee)$

2. $a \diamond b \in \mathrm{lb}(\{a, b\}, \leq_{\diamond}^{\wedge})$

Proof. Let $a, b \in X$.

1. We note that

$$\begin{aligned}(a \diamond b) \diamond [a \diamond (a \diamond b)] &= (a \diamond b) \diamond [(a \diamond a) \diamond b] \\ &= (a \diamond b) \diamond (a \diamond b) \\ &= a \diamond b.\end{aligned}$$

   Similarly, $[(a \diamond b) \diamond b] \diamond (a \diamond b) = a \diamond b$. Therefore $a \leq_{\diamond}^{\vee} a \diamond b$ and $b \leq_{\diamond}^{\vee} a \diamond b$. Hence $a \diamond b \in \mathrm{ub}(\{a, b\}, \leq_{\diamond}^{\vee})$.

2. FINISH!!!

□

9.1.2 Skew Semilattice Homomorphisms

Definition 9.1.2.1. Let $(X, \diamond_X)$ and $(Y, \diamond_Y)$ be skew semilattices and $f : X \to Y$. Then f is said to be a $(\diamond_X, \diamond_Y)$**-homormorphism** if for each $a, b \in X$, $f(a \diamond_X b) = f(a) \diamond_Y f(b)$.

Note 9.1.2.2. When the context is clear, we say homomorphism instead of $(\diamond_X, \diamond_Y)$-homomorphism.

Exercise 9.1.2.3. Let $(X, \diamond_X)$ and $(Y, \diamond_Y)$ be skew semilattices and $f : X \to Y$. If f is a $(\diamond_X, \diamond_Y)$-homomorphism, then

1. f is $(\leq_{\diamond_X}^{\vee}, \leq_{\diamond_Y}^{\vee})$-monotone
2. f is $(\leq_{\diamond_X}^{\wedge}, \leq_{\diamond_Y}^{\wedge})$-monotone

Proof. Suppose that f is a $(\diamond_X, \diamond_Y)$-homomorphism.

1. Let $a, b \in X$. Suppose that $a \leq_{\diamond_X}^{\vee} b$. Then $(b \diamond_X a) \diamond_X b = b$. Therefore

$$\begin{aligned}f(b) &= (b \diamond_X a) \diamond_X b \\ &= f(b \diamond_X a) \diamond_Y f(b) \\ &= (f(b) \diamond_Y f(a)) \diamond_Y f(b)\end{aligned}$$

   Thus $f(a) \leq_{\diamond_Y}^{\vee} f(b)$. Since $a, b \in X$ with $a \leq_{\diamond_X}^{\vee} b$ are arbitrary, we have that for each $a, b \in X$, $a \leq_{\diamond_X}^{\vee} b$ implies that $f(a) \leq_{\diamond_Y}^{\vee} f(b)$. Hence f is $(\leq_{\diamond_X}^{\vee}, \leq_{\diamond_Y}^{\vee})$-monotone.

2. use duality?, FINISH!!!

□

9.1.3 Skew Lattices

Definition 9.1.3.1. Let X be a set and $\vee, \wedge$ skew semilattice operators on X. Then $\vee$ and $\wedge$ are said to **satisfy the skew lattice absorption identities** if for each $a, b \in X$,

1. $a \wedge (a \vee b) = a$
2. $a \vee (a \wedge b) = a$
3. $(b \vee a) \wedge a = a$
4. $(b \wedge a) \vee a = a$

Exercise 9.1.3.2. Let X be a set and $\vee, \wedge$ skew semilattice operators on X. Then $\vee$ and $\wedge$ satisfy the skew lattice absorption identities iff for each $a, b \in X$,

1. $a \vee b = b$ iff $a \wedge b = a$
2. $a \vee b = a$ iff $a \wedge b = b$

Proof.

- ($\Longrightarrow$):
  Suppose that $\vee$ and $\wedge$ satisfy the skew lattice absorption identities. Let $a, b \in X$.

  1. – ($\Longrightarrow$):
     If $a \vee b = b$, then
     $$\begin{aligned} a \wedge b &= a \wedge (a \vee b) \\ &= a. \end{aligned}$$
     – ($\Longleftarrow$):
     If $a \wedge b = a$, then
     $$\begin{aligned} a \vee b &= (a \wedge b) \vee b \\ &= b. \end{aligned}$$
     Thus $a \vee b = b$ iff $a \wedge b = a$.
  2. Similar to (1)

- ($\Longleftarrow$):
  Suppose that for each $a, b \in X$,

  1. $a \vee b = b$ iff $a \wedge b = a$,
  2. $a \vee b = a$ iff $a \wedge b = b$.

  Let $a, b \in X$.

  1. Set $c := a \vee b$. By assumption,
     $$\begin{aligned} a \wedge (a \vee b) = a &\Longleftrightarrow a \wedge c = a \\ &\Longleftrightarrow a \vee c = c \\ &\Longleftrightarrow a \vee (a \vee b) = c \\ &\Longleftrightarrow (a \vee a) \vee b = c \\ &\Longleftrightarrow a \vee b = c \\ &\Longleftrightarrow c = c. \end{aligned}$$
     Since $c = c$, we have that $a \wedge (a \vee b) = a$.
  2. Similar to (1)
  3. Similar to (1)
  4. Similar to (1)

□

Exercise 9.1.3.3. Let X be a set and $\vee, \wedge$ skew semilattice operators on X. If $\vee$ and $\wedge$ satisfy the skew lattice absorption identities, then for each $a, b \in X$, $b \vee a \vee b = b$ iff $a \wedge b \wedge a = a$.
Hint: Exercise 9.1.3.2 implies that $b \vee (a \vee b) = b$ iff $b \wedge (a \vee b) = a \vee b$. Use, skew lattice absorption identities and associativity.

Proof. Suppose that $\vee$ and $\wedge$ satisfy the skew lattice absorption identities. Let $a, b \in X$.

- ($\Longrightarrow$):
  Suppose that $b \vee a \vee b = b$. Then Exercise 9.1.3.2 implies that $b \wedge (a \vee b) = (a \vee b)$. Therefore

$$\begin{aligned} a \wedge b &= (a \wedge b) \vee [(a \wedge b) \wedge (a \vee b)] \quad \text{(absorption)} \\ &= (a \wedge b) \vee [a \wedge (b \wedge [a \vee b])] \quad \text{(associativity)} \\ &= (a \wedge b) \vee [a \wedge (a \vee b)] \quad \text{(from earlier)} \\ &= (a \wedge b) \vee a \quad \text{(absorption).} \end{aligned}$$

  Another application of Exercise 9.1.3.2 implies that $(a \wedge b) \wedge a = a$.

- ($\Longleftarrow$):
  Similar to ($\Longrightarrow$).

□

Definition 9.1.3.4. Let X be a set and $\vee, \wedge$ skew semilattice operators on X. Then $(X, \vee, \wedge)$ is said to be a **skew lattice** if

1. $X \neq \varnothing$
2. $\vee$ and $\wedge$ satisfy the skew lattice absorption identities

Exercise 9.1.3.5. Let $(X, \vee, \wedge)$ be an skew lattice. Then

1. $\leq_{\vee}^{\vee} = \leq_{\wedge}^{\wedge}$.
2. $\leq_{\vee}^{\wedge} = \leq_{\wedge}^{\vee} = (\leq_{\vee}^{\vee})^*$.

Proof.

1. Let $a, b \in X$. Since $(X, \vee, \wedge)$ is an skew lattice, $\vee$ and $\wedge$ satisfy the skew lattice absorption identities. Exercise 9.1.3.3 then implies that

$$\begin{aligned} a \leq_{\vee}^{\vee} b &\iff b \vee a \vee b = b \\ &\iff a \wedge b \wedge a = a \\ &\iff a \leq_{\wedge}^{\wedge} b. \end{aligned}$$

   Since $a, b \in X$ are arbitrary, we have that for each $a, b \in X$, $a \leq_{\vee}^{\vee} b$ iff $a \leq_{\wedge}^{\wedge} b$. Hence $\leq_{\vee}^{\vee} = \leq_{\wedge}^{\wedge}$.

2. Exercise 9.1.1.4 and part (1) imply that

$$\begin{aligned} \leq_{\vee}^{\wedge} &= (\leq_{\vee}^{\vee})^* \\ &= (\leq_{\wedge}^{\wedge})^* \\ &= \leq_{\wedge}^{\vee}. \end{aligned}$$

□

Note 9.1.3.6. Let $(X, \vee, \wedge)$ be an skew lattice. When the context is clear, we write $\leq$ in place of $\leq_{\vee}^{\vee}$.

9.1.4 Semilattices

Definition 9.1.4.1. Let X be a set and $\diamond : X \times X \to X$. Then

- $\diamond$ is said to be a **semilattice operator on** X if $\diamond$ associative, idempotent and commutative.
- $(X, \diamond)$ is said to be a **semilattice** if
  1. $X \neq \varnothing$
  2. $\diamond$ is a semilattice operator on X.

Exercise 9.1.4.2. Let X be a set and $\diamond : X \times X \to X$. Then $\diamond$ is a semilattice operator on X iff

1. $\diamond$ is a skew semilattice operator on X,
2. $\diamond$ is commutative.

Proof. Clear by definition. □

Exercise 9.1.4.3. Let $(X, \diamond)$ be a semilattice. Then

1. for each $a, b \in X$,
   (a) $a \leq_\diamond^\vee b$ iff $a \diamond b = b$
   (b) $a \leq_\diamond^\wedge b$ iff $a \diamond b = a$
2. (a) $\leq_\diamond^\vee$ is a partial order on X
   (b) $\leq_\diamond^\wedge$ is a partial order on X

Proof.

1. Let $a, b \in X$.
   (a) Since $\diamond$ is commutative, associative and idempotent,
   $$\begin{aligned} a \leq_\diamond^\vee b &\iff (b \diamond a) \diamond b = b \\ &\iff (a \diamond b) \diamond b = b \\ &\iff a \diamond (b \diamond b) = b \\ &\iff a \diamond b = b. \end{aligned}$$
   (b) Similar to 1(a).
2. (a) Exercise 9.1.1.3 implies that $\leq_\diamond^\vee$ is a preorder on X. Let $a, b \in X$. Suppose that $a \leq_\diamond^\vee b$ and $b \leq_\diamond^\vee a$. Part (1) then implies that $a \diamond b = b$ and $b \diamond a = a$. Since $\diamond$ is commutative,
   $$\begin{aligned} a &= b \diamond a \\ &= a \diamond b \\ &= b. \end{aligned}$$
   Since $a, b \in X$ are arbitrary, we have that for each $a, b \in X$, if $a \leq_\diamond^\vee b$ and $b \leq_\diamond^\vee a$, then $a = b$. Thus $\leq_\diamond^\vee$ is a a partial order on X.
   (b) Similar to 2(a).

□

Exercise 9.1.4.4. Let $(X, \diamond)$ be a semilattice. Then for each $a, b \in X$,

1. $a \diamond b = \sup(\{a, b\}, \leq_\diamond^\vee)$
2. $a \diamond b = \inf(\{a, b\}, \leq_\diamond^\wedge)$

Proof. Let $a, b \in X$.

1. Exercise 9.1.1.5 implies that $a \diamond b \in \mathrm{ub}(\{a, b\}, \leq_\diamond^\vee)$. Let $c \in \mathrm{ub}(\{a, b\}, \leq_\diamond^\vee)$. Since $a \leq_\diamond^\vee c$ and $b \leq_\diamond^\vee c$, Exercise 9.1.4.3 implies that $a \diamond c = c$ and $b \diamond c = c$. Therefore

$$\begin{aligned}(a \diamond b) \diamond c &= a \diamond (b \diamond c) \\ &= a \diamond c \\ &= c.\end{aligned}$$

   Another application of Exercise 9.1.4.3 implies that $a \diamond b \leq_\diamond^\vee c$. Since $c \in \mathrm{ub}(\{a, b\}, \leq_\diamond^\vee)$ is arbitrary, we have that for each $c \in \mathrm{ub}(\{a, b\}, \leq_\diamond^\vee)$, $a \diamond b \leq_\diamond^\vee c$. Therefore $a \diamond b = \sup(\{a, b\}, \leq_\diamond^\vee)$.

2. FINISH!!!

□

Definition 9.1.4.5. Let $(X, \leq)$ be a poset. Then $(X, \leq)$ is said to be a

- **join semilattice** if
  1. $X \neq \varnothing$
  2. for each $a, b \in X$, $\sup\{a, b\}$ exists.
- **meet semilattice** if
  1. $X \neq \varnothing$
  2. for each $a, b \in X$, $\inf\{a, b\}$ exists.

Exercise 9.1.4.6. Let $(X, \leq)$ be a poset. Then

1. $(X, \leq)$ is a join semilattice iff $(X, \leq^*)$ is a meet semilattice
2. $(X, \leq^*)$ is a join semilattice iff $(X, \leq)$ is a meet semilattice

Proof.

1. UPDATE ONCE SUPREMA/INFIMA EXERCISES ARE REORGANIZED

$$\begin{aligned}(X, \leq) \text{ is a join semilattice} &\iff X \neq \varnothing \text{ and for each } a, b \in X, \sup(\{a, b\}, \leq) \text{ exists} \\ &\iff X \neq \varnothing \text{ and for each } a, b \in X, \inf(\{a, b\}, \leq^*) \text{ exists} \quad \text{(by Exercise B.0.0.3 ADD EXERCISE REF)} \\ &\iff (X, \leq) \text{ is a meet semilattice}\end{aligned}$$

2.

□

Exercise 9.1.4.7. Let $(X, \diamond)$ be a semilattice. Then

1. $(X, \leq_\diamond^\vee)$ is a join semilattice
2. $(X, \leq_\diamond^\wedge)$ is a meet semilattice

Proof. Exercise 9.1.4.4 implies that

1. For each $a, b \in X$, $\sup(\{a, b\}, \leq_\diamond^\vee) = a \diamond b$. Thus $(X, \leq_\diamond^\vee)$ is a join semilattice.
2. for each $a, b \in X$, $\inf(\{a, b\}, \leq_\diamond^\wedge) = a \diamond b$. Thus $(X, \leq_\diamond^\wedge)$ is a meet semilattice.

□

Definition 9.1.4.8. Let $(X, \leq)$ be a poset.

- If $(X, \leq)$ is a join semilattice, we define the **join operator on X induced by** $\leq$, denoted $\vee_{\leq} : X \times X \to X$, by $a \vee_{\leq} b := \sup\{a, b\}$.
- If $(X, \leq)$ is a meet semilattice, we define the **meet operator on X induced by** $\leq$, denoted $\wedge_{\leq} : X \times X \to X$, by $a \wedge_{\leq} b := \inf\{a, b\}$.

Exercise 9.1.4.9. Let $(X, \leq)$ be a poset.

1. If $(X, \leq)$ an join semilattice, then $\vee_{\leq}$ is a semilattice operator on X.
2. If $(X, \leq)$ an meet semilattice, then $\wedge_{\leq}$ is a semilattice operator on X.

Proof.

1. Suppose that $(X, \leq)$ an join semilattice.
   (a)
   - Let $a, b, c \in X$. Exercise 2.7.3.5 implies that
   $$\begin{aligned}(a \vee_{\leq} b) \vee_{\leq} c &= \sup\{\sup\{a, b\}, c\} \\ &= \sup\{\sup\{a, b\}, \sup\{c\}\} \\ &= \sup\{a, b, c\} \\ &= \sup\{\sup\{a\}, \sup\{b, c\}\} \\ &= \sup\{a, \sup\{b, c\}\} \\ &= a \vee_{\leq} (b \vee_{\leq} c).\end{aligned}$$
   Hence $\vee_{\leq}$ is associative.
   - Let $a \in X$. Then
   $$\begin{aligned}a \vee_{\leq} a &= \sup\{a, a\} \\ &= \sup\{a\} \\ &= a.\end{aligned}$$
   Hence $\vee_{\leq}$ is idempotent.
   - Let $a, b \in X$. Then
   $$\begin{aligned}a \vee_{\leq} b &= \sup\{a, b\} \\ &= \sup\{b, a\} \\ &= b \vee_{\leq} a.\end{aligned}$$
   Hence $\vee_{\leq}$ is commutative.

   Since $\vee_{\leq}$ is associative, idempotent and commutative, $\vee_{\leq}$ is a semilattice operator on X.
2. Similar to (1).

□

Exercise 9.1.4.10. Let $(X, \leq)$ be a poset.

1. If $(X, \leq)$ is a join semilattice, then
   (a) $\leq^{\vee}_{\vee_{\leq}} = \leq$,
   (b) $\leq^{\wedge}_{\vee_{\leq}} = \leq^{*}$.
2. If $(X, \leq)$ is a meet semilattice, then
   (a) $\leq^{\wedge}_{\wedge_{\leq}} = \leq$,
   (b) $\leq^{\vee}_{\wedge_{\leq}} = \leq^{*}$.

Proof.

1. Suppose that $(X, \leq)$ is a join semilattice.

   (a) Let $a, b \in X$. Then

$$\begin{aligned} a \leq^{\vee}_{\vee_{\leq}} b &\iff a \vee_{\leq} b = b \quad \text{(Exercise 9.1.4.3)} \\ &\iff \sup(\{a, b\}, \leq) = b \quad \text{(Definition 9.1.4.8)} \\ &\iff a \leq b \quad \text{(Exercise 2.7.3.6).} \end{aligned}$$

   Since $a, b \in X$ are arbitrary, we have that $\leq^{\vee}_{\vee_{\leq}} = \leq$.

   (b) Exercise 9.1.1.4 implies that

$$\begin{aligned} \leq^{\wedge}_{\vee_{\leq}} &= (\leq^{\vee}_{\vee_{\leq}})^* \\ &= \leq^* . \end{aligned}$$

2. Similar to (1).

□

Exercise 9.1.4.11. Let $(X, \diamond)$ be a semilattice. Then

1. $\leq^{\vee}_{\diamond}$ is the unique partial order $\leq$ on X such that $(X, \leq)$ is a join semilattice and for each $a, b \in X$, $a \diamond b = \sup(\{a, b\}, \leq)$.
2. $\leq^{\wedge}_{\diamond}$ is the unique partial order $\leq$ on X such that $(X, \leq)$ is an meet semilattice and for each $a, b \in X$, $a \diamond b = \inf(\{a, b\}, \leq)$.

Proof.

1. Let $\leq$ be partial order on X. Suppose that $(X, \leq)$ is a join semilattice and for each $a, b \in X$, $a \diamond b = \sup(\{a, b\}, \leq)$. Let $a, b \in X$. Then

$$\begin{aligned} a \leq b &\iff b = \sup(\{a, b\}, \leq) \quad \text{(Exercise 2.7.3.6)} \\ &\iff a \diamond b = b \quad \text{(by assumption)} \\ &\iff a \leq^{\vee}_{\diamond} b \quad \text{(Exercise 9.1.4.3).} \end{aligned}$$

   Since $a, b \in X$ are arbitrary, we have that for each $a, b \in X$, $a \leq b$ iff $a \leq^{\vee}_{\diamond} b$. Hence $\leq = \leq^{\vee}_{\diamond}$.

2. Similar to (1).

□

Exercise 9.1.4.12. Let $(X, \leq)$ be a poset.

1. If $(X, \leq)$ is a join semilattice, then $\vee_{\leq}$ is the unique semilattice operator $\diamond$ on X such that for each $a, b \in X$, $a \diamond b = \sup(\{a, b\}, \leq)$.
2. If $(X, \leq)$ is a meet semilattice, then $\wedge_{\leq}$ is the unique semilattice operator $\diamond$ on X such that for each $a, b \in X$, $a \diamond b = \inf(\{a, b\}, \leq)$.

Proof.

1. Let $\diamond : X \times X \to X$ be a semilattice operator on X. Suppose that $a, b \in X$, $a \diamond b = \sup(\{a, b\}, \leq)$. Then for each $a, b \in X$,

$$\begin{aligned} a \diamond b &= \sup(\{a, b\}, \leq) \quad \text{(by assumption)} \\ &= a \vee_{\leq} b \quad \text{(Definition 9.1.4.8).} \end{aligned}$$

   Hence $\diamond = \vee_{\leq}$.

2. Similar to (1).

□

9.1.5 Semilattice Homomorphisms

Definition 9.1.5.1. Let $(X, \leq_X), (Y, \leq_Y)$ be posets and $f : X \to Y$.

1. Suppose that $(X, \leq_X), (Y, \leq_Y)$ are join semilattices. Then f is said to be $(\leq_X, \leq_Y)$**-binary join preserving** if for each $a, b \in X$, $f(\sup(\{a, b\}, \leq_X)) = \sup(\{f(a), f(b)\}, \leq_Y)$.
2. Suppose that $(X, \leq_X), (Y, \leq_Y)$ are meet semilattices. Then f is said to be $(\leq_X, \leq_Y)$**-binary meet preserving** if for each $a, b \in X$, $f(\inf(\{a, b\}, \leq_X)) = \inf(\{f(a), f(b)\}, \leq_Y)$.

Exercise 9.1.5.2. Let $(X, \leq_X), (Y, \leq_Y)$ be posets and $f : X \to Y$.

1. Suppose that $(X, \leq_X), (Y, \leq_Y)$ are join semilattices. Then f is finite $(\leq_X, \leq_Y)$-join preserving iff f is a $(\vee_{\leq_X}, \vee_{\leq_Y})$-homomorphism.
2. Suppose that $(X, \leq_X), (Y, \leq_Y)$ are meet semilattices. Then f is finite $(\leq_X, \leq_Y)$-meet preserving iff f is a $(\wedge_{\leq_X}, \wedge_{\leq_Y})$-homomorphism.

Proof. Clear. □

Exercise 9.1.5.3. Let $(X, \diamond_X)$ and $(Y, \diamond_Y)$ be semilattices and $f : X \to Y$. Then the following are equivalent:

1. f is a $(\diamond_X, \diamond_Y)$-homomorphism
2. f is $(\leq^{\vee}_{\diamond_X}, \leq^{\vee}_{\diamond_Y})$-finite join preserving
3. f is $(\leq^{\wedge}_{\diamond_X}, \leq^{\wedge}_{\diamond_Y})$-finite meet preserving

Proof.

- (1) $\implies$ (2):
  Suppose that f is a $(\diamond_X, \diamond_Y)$-homomorphism. make/ref previous ex here implies that $\vee_{\leq^{\vee}_{\diamond_X}} = \diamond_X$ and $\vee_{\leq^{\vee}_{\diamond_Y}} = \diamond_Y$. Therefore f is a $(\vee_{\leq^{\vee}_{\diamond_X}}, \vee_{\leq^{\vee}_{\diamond_Y}})$-homomorphism. the previous ex implues that f is $(\leq^{\vee}_{\diamond_X}, \leq^{\vee}_{\diamond_Y})$-finite join preserving.
- (2) $\implies$ (3):
  Suppose that FINISH!!!
- (3) $\implies$ (1):

□

Exercise 9.1.5.4. Let $(X, \leq_X), (Y, \leq_Y)$ be posets and $f : X \to Y$.

1. Suppose that $(X, \leq_X)$ and $(Y, \leq_Y)$ are join semilattices. Then f is $(\leq_X, \leq_Y)$-monotone iff for each $a, b \in X$, $f(a) \vee_{\leq_Y} f(b) \leq_Y f(a \vee_{\leq_X} b)$.
2. Suppose that $(X, \leq_X)$ and $(Y, \leq_Y)$ are meet semilattices. Then f is $(\leq_X, \leq_Y)$-monotone iff for each $a, b \in X$, $f(a) \vee_{\leq_Y} f(b) \leq_Y f(a \vee_{\leq_X} b)$.

Proof.

1. - ($\implies$):
     Suppose that f is $(\leq_X, \leq_Y)$-monotone. Let $a, b \in X$. Since $a \vee_{\leq_X} b \in \mathrm{ub}(\{a, b\}, \leq_X)$, Exercise 2.7.1.10 implies that
     $$\begin{aligned} f(a \vee_{\leq_X} b) &\in f(\mathrm{ub}(\{a, b\}, \leq_X)) \\ &\subset \mathrm{ub}(f(\{a, b\}), \leq_Y) \\ &= \mathrm{ub}(\{f(a), f(b)\}, \leq_Y) \end{aligned}$$
     Therefore
     $$\begin{aligned} f(a) \vee_{\leq_Y} f(b) &= \sup(\{f(a), f(b)\}, \leq_Y) \\ &\leq f(a \vee_{\leq_X} b). \end{aligned}$$
     Since $a, b \in X$ are arbitrary, we have that for each $a, b \in X$, $f(a) \vee_{\leq_Y} f(b) \leq_Y f(a \vee_{\leq_X} b)$.

- ($\Longleftarrow$):
  Suppose that for each $a, b \in X$, $f(a) \vee_{\leq_Y} f(b) \leq_Y f(a \vee_{\leq_X} b)$. Let $a, b \in X$. Suppose that $a \leq_X b$. Exercise 2.7.3.6 then implies that $a \vee_{\leq_X} b = b$. By assumption,

$$\begin{aligned} f(a) \vee_{\leq_Y} f(b) &\leq_Y f(a \vee_{\leq_X} b) \\ &= f(b) \\ &\leq f(a) \vee_{\leq_Y} f(b). \end{aligned}$$

  Therefore $f(a) \vee_{\leq_Y} f(b) = f(b)$. Another application of Exercise 2.7.3.6 implies that $f(a) \leq_Y f(b)$. Since $a, b \in X$ with $a \leq b$ are arbitrary, we have that for each $a, b \in X$, $a \leq_X b$ implies that $f(a) \leq_Y f(b)$. Hence f is $(\leq_X, \leq_Y)$-monotone.

2. use duality FINISH!!!

□

Exercise 9.1.5.5. Let $(X, \leq_X), (Y, \leq_Y)$ be posets and $f : X \to Y$.

1. Suppose that $(X, \leq_X)$ and $(Y, \leq_Y)$ are join semilattices. If f binary join preserving, then f is $(\leq_X, \leq_Y)$-monotone.
2. Suppose that $(X, \leq_X)$ and $(Y, \leq_Y)$ are meet semilattices. If f binary meet preserving, then f is $(\leq_X, \leq_Y)$-monotone.

Proof.

1. Suppose that f is binary join preserving. Then the previous exercise implies that

$$\begin{aligned} \text{for each } a, b \in X,\ f(a) \vee_{\leq_Y} f(b) = f(a \vee_{\leq_X} b) &\implies \text{for each } a, b \in X,\ f(a) \vee_{\leq_Y} f(b) \leq f(a \vee_{\leq_X} b) \\ &\implies f \text{ is } (\leq_X, \leq_Y)\text{-monotone} \end{aligned}$$

2. Suppose that $(X, \leq_X)$ and $(Y, \leq_Y)$ are meet semilattices. If f preserves finite meets, then f is $(\leq_X, \leq_Y)$-monotone.

□

9.1.6 Lattices

Definition 9.1.6.1. Let X be a set and $\vee, \wedge$ semilattice operators on X. Then $\vee$ and $\wedge$ are said to **satisfy the lattice absorption identities** if for each $a, b \in X$,

1. $a \wedge (a \vee b) = a$
2. $a \vee (a \wedge b) = a$

Exercise 9.1.6.2. Let X be a set and $\vee, \wedge$ semilattice operators on X. Then $\vee$ and $\wedge$ satisfy the lattice absorption identities iff $\vee$ and $\wedge$ satisfy the skew lattice absorption identities.

Proof.

- ($\Longrightarrow$):
  Clear by commutativity.
- ($\Longleftarrow$):
  Immediate.

□

Definition 9.1.6.3. Let X be a set and $\vee, \wedge$ semilattice operators on X. Then $(X, \vee, \wedge)$ is said to be an **lattice** if

1. $X \neq \varnothing$
2. $\vee$ and $\wedge$ satisfy the skew lattice absorption identities.

Definition 9.1.6.4. Let $(X, \leq)$ be a poset. Then $(X, \leq)$ is said to be an **ordered lattice** if $(X, \leq)$ is a join semilattice and $(X, \leq)$ is a meet semilattice.

Exercise 9.1.6.5. Let $(X, \vee, \wedge)$ be a lattice. Then $(X, \leq_{\vee}^{\vee})$ is an ordered lattice.

Proof. Exercise 9.1.4.7 implies that $(X, \leq_{\vee}^{\vee})$ is a join semilattice and $(X, \leq_{\wedge}^{\wedge})$ is a meet semilattice. Exercise 9.1.3.5 implies that $\leq_{\vee}^{\vee} = \leq_{\wedge}^{\wedge}$. Therefore $(X, \leq_{\vee}^{\vee})$ is a join semilattice and $(X, \leq_{\vee}^{\vee})$ is a meet semilattice. Hence $(X, \leq_{\vee}^{\vee})$ is an ordered lattice. □

Exercise 9.1.6.6. Let $(X, \leq)$ be an ordered lattice. Then $(X, \vee_{\leq}, \wedge_{\leq})$ is a lattice.

Proof. Let $a, b \in X$.

1. - Since $\sup\{a, b\} \in \mathrm{ub}\{a, b\}$ is an upper bound of $\{a, b\}$, we have that $a \leq \sup\{a, b\}$. Since $a \leq a$, we have that $a \in \mathrm{lb}\{a, \sup\{a, b\}\}$. Hence $a \leq \inf\{a, \sup\{a, b\}\}$.
   - Since $\inf\{a, \sup\{a, b\}\} \in \mathrm{lb}\{a, \sup\{a, b\}\}$, we have that $\inf\{a, \sup\{a, b\}\} \leq a$.

   Since $a \leq \inf\{a, \sup\{a, b\}\}$ and $\inf\{a, \sup\{a, b\}\} \leq a$, we have that
$$\begin{aligned} a \wedge_{\leq} (a \vee_{\leq} b) &= \inf\{a, \sup\{a, b\}\} \\ &= a. \end{aligned}$$
2. Similarly, $a \vee_{\leq} (a \wedge_{\leq} b) = a$.

Since $a, b \in X$ are arbitrary, we have that for each $a, b \in X$,

1. $a \wedge_{\leq} (a \vee_{\leq} b) = a$
2. $a \vee_{\leq} (a \wedge_{\leq} b) = a$

Hence $\vee_{\leq}$ and $\wedge_{\leq}$ satisfy the lattice absorption identities. Therefore $(X, \vee_{\leq}, \wedge_{\leq})$ is an lattice. □

Exercise 9.1.6.7. Let $(X, \leq)$ be an ordered lattice. Then

1. $\leq_{\vee_{\leq}}^{\vee} = \leq_{\wedge_{\leq}}^{\wedge} = \leq$
2. $\leq_{\vee_{\leq}}^{\wedge} = \leq_{\wedge_{\leq}}^{\vee} = \leq^{*}$

Proof. Since $(X, \leq)$ is an ordered lattice, $(X, \leq)$ is a join semilattice $(X, \leq)$ is a meet semilattice. Exercise 9.1.4.10 then implies that

1. $$\begin{aligned} \leq_{\vee_{\leq}}^{\vee} &= \leq \\ &= \leq_{\wedge_{\leq}}^{\wedge} \end{aligned}$$
2. $$\begin{aligned} \leq_{\vee_{\leq}}^{\wedge} &= \leq^{*} \\ &= \leq_{\wedge_{\leq}}^{\vee} \end{aligned}$$

□

Exercise 9.1.6.8. Let $(X, \vee, \wedge)$ be an lattice. Then $\leq_{\vee}^{\vee}$ is the unique partial order $\leq$ on X such that $(X, \leq)$ is an ordered lattice and for each $a, b \in X$, $a \vee b = \sup(\{a, b\}, \leq)$.

Proof. Let $\leq$ be a partial order on X. Suppose that $(X, \leq)$ is an ordered lattice and for each $a, b \in X$, $a \vee b = \sup(\{a, b\}, \leq)$. Since $(X, \vee)$ is a semilattice, $(X, \leq_{\vee}^{\vee})$ and $(X, \leq)$ are join semilattices and for each $a, b \in X$, $a \vee b = \sup(\{a, b\}, \leq_{\vee}^{\vee})$ and $a \vee b = \sup(\{a, b\}, \leq)$, Exercise 9.1.4.11 implies that $\leq = \leq_{\vee}^{\vee}$. □

Exercise 9.1.6.9. Let $(X, \leq)$ be an ordered lattice. Then $\vee_{\leq}, \wedge_{\leq}$ are the unique semilattice operators $\vee, \wedge$ on X such that $(X, \vee, \wedge)$ is an lattice and for each $a, b \in X$, $a \vee b = \sup(\{a, b\}, \leq)$

Proof. Let $\vee, \wedge$ be semilattice operators on X. Suppose that $(X, \vee, \wedge)$ is a lattice and for each $a, b \in X$, $a \vee b = \sup(\{a, b\}, \leq)$. Then for each $a, b \in X$,

$$\begin{aligned} a \vee_{\leq} b &= \sup(\{a, b\}, \leq) \quad \text{(Definition 9.1.4.8)} \\ &= a \vee b \quad \text{(by assumption)}. \end{aligned}$$

Thus $\vee = \vee_{\leq}$. Since $(X, \vee, \wedge)$ is a lattice and for each $a, b \in X$, $a \vee b = \sup(\{a, b\}, \leq)$, Exercise 9.1.6.8 implies that $\leq_{\vee}^{\vee} = \leq$. Exercise 9.1.3.5 then implies that

$$\begin{aligned} \leq_{\wedge}^{\wedge} &= \leq_{\vee}^{\vee} \\ &= \leq . \end{aligned}$$

Therefore for each $a, b \in X$,

$$\begin{aligned} a \wedge b &= \inf(\{a, b\}, \leq_{\wedge}^{\wedge}) \quad \text{(Exercise 9.1.4.4)} \\ &= \inf(\{a, b\}, \leq) \\ &= a \wedge_{\leq} b \quad \text{(Definition 9.1.4.8)}. \end{aligned}$$

Hence $\wedge = \wedge_{\leq}$. □

Definition 9.1.6.10. Let $(X, \vee, \wedge)$ be a lattice. We define the

- $(\vee, \wedge)$**-dual join operator on** X, denoted $\vee^* : X \times X \to X$, by $\vee^* := \wedge$
- $(\vee, \wedge)$**-dual meet operator on** X, denoted $\vee^* : X \times X \to X$, by $\vee^* := \wedge$
- **dual lattice of** $(X, \vee, \wedge)$, denoted $(X, \vee, \wedge)^*$, by $(X, \vee, \wedge)^* := (X, \vee^*, \wedge^*)$.

Note 9.1.6.11. When the context is clear, we may say "dual join operator on X" and "dual meet operator on X" in place of "$(\vee, \wedge)$-dual join operator on X" and "$(\vee, \wedge)$-dual meet operator on X" respectively.

Exercise 9.1.6.12. Let $(X, \leq)$ be an ordered lattice. Then

1. $\vee_{\leq}^* = \vee_{\leq^*}$ and $\wedge_{\leq}^* = \wedge_{\leq^*}$
2. $(\vee^*)^* = \vee$ and $(\wedge^*)^* = \wedge)$

Proof.

1. For each $a, b \in X$,

$$\begin{aligned} a \vee_{\leq}^* b &= a \wedge_{\leq} b \\ &= \inf(\{a, b\}, \leq) \\ &= \sup(\{a, b\}, \leq^*) \quad \text{(by ref ex)} \\ &= a \vee_{\leq^*} b \end{aligned}$$

   Hence $\vee_{\leq}^* = \vee_{\leq^*}$. Similarly, $\wedge_{\leq}^* = \wedge_{\leq^*}$.

2. By Definition 9.1.6.10,

$$\begin{aligned} (\vee^*)^* &= \wedge^* \\ &= \vee. \end{aligned}$$

   Similarly, $(\wedge^*)^* = \wedge$.

□

Exercise 9.1.6.13. Let $(X, \vee, \wedge)$ be a lattice. Then $\leq_{\vee^*}^{\vee} = (\leq_{\vee}^{\vee})^*$.

Proof. We have that

$$\begin{aligned} \leq_{\vee^*}^{\vee} &= \leq_{\wedge}^{\vee} \quad \text{(by Definition B.0.0.4 ref prev definition)} \\ &= (\leq_{\vee}^{\vee})^* \quad \text{(by Exercise B.0.0.3 ref prev ex)} \end{aligned}$$

□

Note 9.1.6.14. We the context is clear, we write $\leq_X$ and $\leq_{X^*}$ in place of $\leq_{\vee}^{\vee}$ and $\leq_{\vee^*}^{\vee}$ respectively.

9.1.7 Lattice Homomorphisms

???

9.2 Fundamental Examples

9.2.1 Bounded Lattices

mayb define bounded prosets,

Definition 9.2.1.1. Let $(L, \vee, \wedge)$ be a lattice.

- Let $a \in L$. Then
  - a is said to be a $(\vee, \wedge)$**-zero** if for each $x \in L$, $x \vee a = x$.
  - a is said to be a $(\vee, \wedge)$**-one** if for each $x \in L$, $x \wedge a = x$
- Then
  - $(L, \vee, \wedge)$ is said to **have a zero** if there exists $a \in L$ such that a is a $(\vee, \wedge)$-zero
  - $(L, \vee, \wedge)$ is said to **have a one** if there exists $a \in L$ such that a is a $(\vee, \wedge)$-one

Note 9.2.1.2. We the context is clear, we say "zero" and "one" inplace of "$(\vee, \wedge)$-zero" and "$(\vee, \wedge)$-one" respectively.

Exercise 9.2.1.3. Let L be a lattice and $a, b \in L$.

1. If a, b are zeros, then $a = b$.
2. If a, b are ones, then $a = b$.

Proof.

1. Suppose that a, b are zeros. Then

$$\begin{aligned} a &= a \wedge b \\ &= b \wedge a \\ &= b. \end{aligned}$$

2. Similar to (1).

□

Definition 9.2.1.4. Let L be a lattice.

- If L has a zero, we define $0_L \in L$ to be the unique $a \in L$ such that a is a zero of L.
- If L has a one, we define $1_L \in L$ to be the unique $a \in L$ such that a is a one of L.

Note 9.2.1.5. When the context is clear, we may write 0 and 1 in place of 0_L and 1_L respectively.

Definition 9.2.1.6. Let $(L, \vee, \wedge)$ be a lattice. Then

- $(L, \vee, \wedge)$ is said to be a **lattice with** 0 if L has a $(\vee, \wedge)$-zero
- $(L, \vee, \wedge)$ is said to be a **lattice with** 1 if $(L, \vee, \wedge)$ has a $(\vee, \wedge)$-one
- $(L, \vee, \wedge, 0, 1)$ is said to be a **bounded lattice** if 0 is a $(\vee, \wedge)$-zero and 1 is a $(\vee, \wedge)$-one

Exercise 9.2.1.7. Let L be a lattice and $a \in L$. Then

1. a is a one of L iff $\sup L = a$.
2. a is a zero of L iff $\inf L = a$.

Proof.

1. - ($\Longrightarrow$):
   Suppose that a is a one of L. Let $x \in L$. Since $x \wedge a = x$, we have that $x \leq a$. Since $x \in L$ is arbitrary, we have that for each $x \in L$, $x \leq a$. Hence $a \in \operatorname{ub} L$. Exercise 2.7.3.8 then implies that $\sup L = a$.
   - ($\Longleftarrow$):
   Suppose that $\sup L = a$. Let $x \in L$. Then Since $a \in \operatorname{ub} L$, $x \leq a$. Therefore
   $$\begin{aligned} x \wedge a &= \inf\{x, a\} \\ &= x. \end{aligned}$$
   Since $x \in L$ is arbitrary, we have that for each $x \in L$, $x \wedge a = x$. Hence a is a one of L.
2. Similar to (1).

□

Exercise 9.2.1.8. Let $(L, \leq)$ be an ordered lattice. (maybe change to join/meet lattice) Then

1. $(L, \vee_{\leq}, \wedge_{\leq})$ is a lattice with 0 iff for each $A \subset L$, A is finite implies that $\sup A$ exists.
2. $(L, \vee_{\leq}, \wedge_{\leq})$ is a lattice with 1 iff for each $A \subset L$, A is finite implies that $\inf A$ exists.

Proof.

- ($\Longrightarrow$):
  Suppose that $(L, \vee_{\leq}, \wedge_{\leq})$ is a lattice with 0. Let $A \subset L$. Suppose that A is finite.
  - Suppose that $\#A = 0$. Then $A = \varnothing$. a previous ex implies that
  $$\begin{aligned} \sup A \text{ exists} &\iff \inf(\operatorname{ub} A) \text{ exists} \\ &\iff \inf(\operatorname{ub} \varnothing) \text{ exists} \\ &\iff \inf L \text{ exists.} \end{aligned}$$
  Since L has a zero, the previous ex implies that $\inf L$ exists. Therefore $\sup A$ exists.
  - Suppose that $\#A \geq 1$. Since $(L, \leq)$ is an join semilattice, we have that for each $a, b \in L$, $\sup\{a, b\}$ exists. Exercise 2.7.3.7 then implies that $\sup A$ exists.
- ($\Longleftarrow$):
  Suppose that for each $A \subset L$, A is finite implies that $\sup A$ exists. Set $z := \sup \varnothing$. a prev ex implies that $z = \inf L$. Then

FINISH!!!! □

9.2.2 Bounded Lattice Homomorphisms

Definition 9.2.2.1. move to section on semilattice homomorphisms or make section on semilattice homomorphisms with 0 or 1 and move it there Let $(X, \leq_X), (Y, \leq_Y)$ be posets and $f : X \to Y$.

1. Suppose that $(X, \leq_X), (Y, \leq_Y)$ are join semilattices with 0. Then f is said to be $(\leq_X, \leq_Y)$**-finite join preserving** if for each $A \subset X$, A is finite implies that $f(\sup(A, \leq_X)) = \sup(f(A), \leq_Y)$.
2. Suppose that $(X, \leq_X), (Y, \leq_Y)$ are meet semilattices with 1. Then f is said to be $(\leq_X, \leq_Y)$**-finite meet preserving** if for each $A \subset X$, A is finite implies that $f(\inf(A, \leq_X)) = \inf(f(A), \leq_Y)$.

Exercise 9.2.2.2. Let $(L, \leq_L)$ and $(M, \leq_M)$ be posets and $f : L \to M$.

1. Suppose that $(L, \leq_L)$ and $(M, \leq_M)$ are join semilattices with 0. Then f is $(\leq_L, \leq_M)$-finite join preserving iff f is $(\leq_L, \leq_M)$-binary join preserving and $f(0_L) = 0_M$.
2. Suppose that $(L, \leq_L)$ and $(M, \leq_M)$ are meet semilattices with 1. Then f is $(\leq_L, \leq_M)$-finite meet preserving iff f is $(\leq_L, \leq_M)$-binary meet preserving and $f(1_L) = 1_M$.

Proof.

1.
   - ($\Longrightarrow$):
     Suppose that f is $(\leq_L, \leq_M)$-finite join preserving. Then clearly f is $(\leq_L, \leq_M)$-binary join preserving. Since $\varnothing$ is finite, a prev ex about $\sup \varnothing = 0$ implies that

$$\begin{aligned} f(0_L) &= f(\sup(\varnothing, \leq_L)) \\ &= \sup(f(\varnothing), \leq_M) \\ &= \sup(\varnothing, \leq_M) \\ &= 0_M. \end{aligned}$$

   - ($\Longleftarrow$):
     Suppose that f is $(\leq_L, \leq_M)$-binary join preserving and $f(0_L) = 0_M$. Let $A \subset L$. Suppose that A is finite.
     - Suppose that $\#A = 0$. Then $A = \varnothing$ and

$$\begin{aligned} f(\sup(A, \leq_L)) &= f(\sup(\varnothing, \leq_L)) \\ &= f(0_L) \\ &= 0_M \\ &= \sup(\varnothing, \leq_M) \\ &= \sup(f(\varnothing), \leq_M) \\ &= \sup(f(A), \leq_M). \end{aligned}$$

     - Suppose that $\#A = 1$. Then there exists $a \in L$ such that $A = \{a\}$. Hence

$$\begin{aligned} f(\sup(A, \leq_L)) &= f(\sup(\{a\}, \leq_L)) \\ &= f(a) \\ &= \sup(\{f(a)\}, \leq_M) \\ &= \sup(f(A), \leq_M). \end{aligned}$$

     - Suppose that $\#A \geq 2$. Then there exists $n \in \mathbb{N}$ and $a_1, \ldots, a_n \in L$ such that $n \geq 2$ and $A = \{a_1, \ldots, a_n\}$. ref prev ex implies that $\sup(A, \leq_L) = (\cdots((a_1 \vee_{\leq_L} a_2) \vee_{\leq_L} a_3) \cdots a_n)$. Then

$$\begin{aligned} f(\sup(A, \leq_L)) &= f(\cdots((a_1 \vee_{\leq_L} a_2) \vee_{\leq_L} a_3) \vee_{\leq_L} \cdots \vee_{\leq_L} a_n) \\ &= \end{aligned}$$

       do by induction on $\#A$ so it is less messy.

2.

FINISH!!! □

9.2.3 Modular and Distributive Lattices

9.2.4 Introduction

Definition 9.2.4.1. Let L be a lattice. Then L is said to be a **distributive lattice** if for each $a, b, c \in L$, $a \vee (b \wedge c) = (a \vee b) \wedge (a \vee c)$.

Exercise 9.2.4.2. Let L be a lattice. Then for each $a, b, c \in L$,

1. $(a \wedge b) \vee (a \wedge c) \leq a \wedge (b \vee c)$.
2. $a \wedge (b \vee c) \leq (a \vee b) \wedge (a \vee c)$

Proof. Let $a, b, c \in L$.

1. - Since $a \wedge b \leq a$ and $a \wedge c \leq a$, we have that $a \in \mathrm{ub}\{a \wedge b, a \wedge c\}$. Therefore
$$\begin{aligned}(a \wedge b) \vee (a \wedge c) &= \sup\{a \wedge b, a \wedge c\} \\ &\leq a.\end{aligned}$$
   - Since
$$\begin{aligned}a \wedge b &\leq b \\ &\leq b \vee c\end{aligned}$$
   and
$$\begin{aligned}a \wedge c &\leq c \\ &\leq b \vee c,\end{aligned}$$
   we have that $b \vee c \in \mathrm{ub}\{a \wedge b, a \wedge c\}$. Therefore
$$\begin{aligned}(a \wedge b) \vee (a \wedge c) &= \sup\{a \wedge b, a \wedge c\} \\ &\leq b \vee c.\end{aligned}$$

   Then $(a \wedge b) \vee (a \wedge c) \in \mathrm{lb}\{a, b \vee c\}$. Hence
$$\begin{aligned}(a \wedge b) \vee (a \wedge c) &\leq \inf\{a, b \vee c\} \\ &= a \wedge (b \vee c).\end{aligned}$$

2. FINISH!!! (use duality)

□

Exercise 9.2.4.3. Let L be a lattice. Then for each $a, b, c \in L$,

1. $c \leq a$ implies that $(a \wedge b) \vee c \leq a \wedge (b \vee c)$
2. $a \leq c$ implies that $a \vee (b \wedge c) \leq (a \vee b) \wedge c$

Proof. Let $a, b, c \in L$.

1. Suppose that $c \leq a$. Then $a \wedge c = c$. The previous exercise then implies that
$$\begin{aligned}(a \wedge b) \vee c &= (a \wedge b) \vee (a \wedge c) \\ &\leq a \wedge (b \vee c).\end{aligned}$$

2. FINISH!!! use duality

□

Exercise 9.2.4.4. Let L be a lattice. Then for each $a, b, c \in L$,
$$(a \wedge b) \vee (b \wedge c) \vee (c \wedge a) \leq (a \vee b) \wedge (b \vee c) \wedge (c \vee a).$$

Proof. Let $a, b, c \in L$.

- We first note that
$$\begin{aligned}a \wedge b \leq a \leq a \vee b, \\ b \wedge c \leq b \leq a \vee b, \\ c \wedge a \leq a \leq a \vee b.\end{aligned}$$
  Therefore $a \vee b \in \mathrm{ub}\{a \wedge b, b \wedge c, c \wedge a\}$ and
$$\begin{aligned}(a \wedge b) \vee (b \wedge c) \vee (c \wedge a) &= \sup\{a \wedge b, b \wedge c, c \wedge a\} \\ &\leq a \vee b.\end{aligned}$$

- Similarly,

$$\begin{aligned} a \wedge b &\leq b \leq b \vee c, \\ b \wedge c &\leq c \leq b \vee c, \\ c \wedge a &\leq c \leq b \vee c. \end{aligned}$$

  Therefore $b \vee c \in \operatorname{ub}\{a \wedge b, b \wedge c, c \wedge a\}$ and

$$\begin{aligned} (a \wedge b) \vee (b \wedge c) \vee (c \wedge a) &= \sup\{a \wedge b, b \wedge c, c \wedge a\} \\ &\leq b \vee c. \end{aligned}$$

- Finally, we have that

$$\begin{aligned} a \wedge b &\leq a \leq c \vee a, \\ b \wedge c &\leq c \leq c \vee a, \\ c \wedge a &\leq c \leq c \vee a. \end{aligned}$$

  Therefore $c \vee a \in \operatorname{ub}\{a \wedge b, b \wedge c, c \wedge a\}$ and

$$\begin{aligned} (a \wedge b) \vee (b \wedge c) \vee (c \wedge a) &= \sup\{a \wedge b, b \wedge c, c \wedge a\} \\ &\leq c \vee a. \end{aligned}$$

Hence $(a \wedge b) \vee (b \wedge c) \vee (c \wedge a) \in \operatorname{lb}\{a \vee b, b \vee c, c \vee a\}$ and

$$\begin{aligned} (a \wedge b) \vee (b \wedge c) \vee (c \wedge a) &\leq \inf\{a \vee b, b \vee c, c \vee a\} \\ &= (a \vee b) \wedge (b \vee c) \wedge (c \vee a). \end{aligned}$$

□

Exercise 9.2.4.5. Let L be a lattice. Then the following are equivalent:

1. For each $a, b, c \in L$, $c \leq a$ implies that $a \wedge (b \vee c) = (a \wedge b) \vee c$.
2. For each $a, b, c \in L$, $c \leq a$ implies that $a \wedge (b \vee c) = (a \wedge b) \vee (a \wedge c)$.
3. For each $p, q, r \in L$, $p \wedge (q \vee (p \wedge r)) = (p \wedge q) \vee (p \wedge r)$.

Proof.

1. (1) $\implies$ (2):
   Suppose that for each $a, b, c \in L$, $c \leq a$ implies that $a \wedge (b \vee c) = (a \wedge b) \vee c$. Let $a, b, c \in L$. Suppose that $c \leq a$. Then $a \wedge c = c$. By assumption

$$\begin{aligned} a \wedge (b \vee c) &= (a \wedge b) \vee c \\ &= (a \wedge b) \vee (a \wedge c) \end{aligned}$$

   Since $a, b, c \in L$ with $c \leq a$ are arbitrary, we have that for each $a, b, c \in L$, $c \leq a$ implies that $a \wedge (b \vee c) = (a \wedge b) \vee (a \wedge c)$.

2. (2) $\implies$ (3):
   Suppose that for each $a, b, c \in L$, $c \leq a$ implies that $a \wedge (b \vee c) = (a \wedge b) \vee (a \wedge c)$. Let $p, q, r \in L$. Define $a, b, c \in L$ by $a := p$, $b := q$ and $c := p \wedge r$. Then

$$\begin{aligned} a \wedge c &= p \wedge (p \wedge r) \\ &= (p \wedge p) \wedge r \\ &= p \wedge r. \end{aligned}$$

By assumption,

$$\begin{aligned} p \wedge (q \vee (p \wedge r)) &= a \wedge (b \vee c) \\ &= (a \wedge b) \vee (a \wedge c) \\ &= (p \wedge q) \vee (p \wedge r). \end{aligned}$$

Since $p, q, r \in L$ are arbitrary, we have that for each $p, q, r \in L$, $p \wedge (q \vee (p \wedge r)) = (p \wedge q) \vee (p \wedge r)$.

3. (3) $\implies$ (1):
Suppose that for each $p, q, r \in L$, $p \wedge (q \vee (p \wedge r)) = (p \wedge q) \vee (p \wedge r)$. Let $a, b, c \in L$. Suppose that $c \leq a$. Define $p, q, r \in L$ by $p := a$, $q := b$ and $r := c$. Since $c \leq a$, we have that

$$\begin{aligned} p \wedge r &= a \wedge c \\ &= c. \end{aligned}$$

By assumption,

$$\begin{aligned} a \wedge (b \vee c) &= p \wedge (q \vee (p \wedge r)) \\ &= (p \wedge q) \vee (p \wedge r) \\ &= (a \wedge b) \vee c. \end{aligned}$$

Since $a, b, c \in L$ with $c \leq a$ are arbitrary, we have that for each $a, b, c \in L$, $c \leq a$ implies that $a \wedge (b \vee c) = (a \wedge b) \vee c$.

□

Definition 9.2.4.6. Let L be a lattice. Then L is said to be **modular** if for each $a, b, c \in L$,

$$c \leq a \text{ implies that } a \wedge (b \vee c) = (a \wedge b) \vee c.$$

Exercise 9.2.4.7. Let L be a lattice. Then the following are equivalent:

1. For each $a, b, c \in L$, $a \wedge (b \vee c) = (a \wedge b) \vee (a \wedge c)$.
2. For each $p, q, r \in L$, $p \vee (q \wedge r) = (p \vee q) \wedge (p \vee r)$.

Proof.

1. (1) $\implies$ (2):
Suppose that for each $a, b, c \in L$, $a \wedge (b \vee c) = (a \wedge b) \vee (a \wedge c)$. Let $p, q, r \in L$. Define $a, b, c \in L$ by $a := p \vee q$, $b := p$ and $c := r$. By absorption,

$$\begin{aligned} a \wedge b &= (p \vee q) \wedge p \\ &= p \wedge (p \vee q) \\ &= p. \end{aligned}$$

By assumption,

$$\begin{aligned} a \wedge c &= (p \vee q) \wedge r \\ &= r \wedge (p \vee q) \\ &= (r \wedge p) \vee (r \wedge q). \end{aligned}$$

Then by assumption and absorption,

$$\begin{aligned} (p \vee q) \wedge (p \vee r) &= a \wedge (b \vee c) \\ &= (a \wedge b) \vee (a \wedge c) \\ &= p \vee [(r \wedge p) \vee (r \wedge q)] \\ &= [p \vee (r \wedge p)] \vee (r \wedge q) \\ &= [p \vee (p \wedge r)] \vee (q \wedge r) \\ &= p \vee (q \wedge r). \end{aligned}$$

Since $p, q, r \in L$ are arbitrary, we have that for each $p, q, r \in L$, $p \vee (q \wedge r) = (p \vee q) \wedge (p \vee r)$.

2. (2) $\implies$ (1):
Suppose that for each $p, q, r \in L$, $p \vee (q \wedge r) = (p \vee q) \wedge (p \vee r)$. Let $a, b, c \in L$. Define $p, q, r \in L$ by $p := a \wedge b$, $q := a$ and $r := c$. By absorption,

$$\begin{aligned} p \vee q &= (a \wedge b) \vee a \\ &= a \vee (a \wedge b) \\ &= a. \end{aligned}$$

By assumption,

$$\begin{aligned} p \vee r &= (a \wedge b) \vee c \\ &= c \vee (a \wedge b) \\ &= (c \vee a) \wedge (c \vee b). \end{aligned}$$

Then by assumption and absorption,

$$\begin{aligned} (a \wedge b) \vee (a \wedge c) &= p \vee (q \wedge r) \\ &= (p \vee q) \wedge (p \vee r) \\ &= a \wedge [(c \vee a) \wedge (c \vee b)] \\ &= [a \wedge (c \vee a)] \wedge (c \vee b) \\ &= [a \wedge (a \vee c)] \wedge (b \vee c) \\ &= a \wedge (b \vee c). \end{aligned}$$

Since $a, b, c \in L$ are arbitrary, we have that for each $a, b, c \in L$, $a \wedge (b \vee c) = (a \wedge b) \vee (a \wedge c)$.

□

Definition 9.2.4.8. Let L be a lattice. Then L is said to be **distributive** if for each $a, b, c \in L$,

$$a \wedge (b \vee c) = (a \wedge b) \vee (a \wedge c).$$

Exercise 9.2.4.9. Let L be a lattice. If L is distributive, then L is modular.

Proof. Suppose that L is distributive. FINISH!!! □

Exercise 9.2.4.10. Let L be a lattice. Then for each $a, b, c \in L$, $a \vee (b \wedge c) = (a \vee b) \wedge (a \vee c)$ iff $a \wedge (b \vee c) = (a \wedge b) \vee (a \vee c)$.

Proof. Let $a, b, c \in L$.

- ($\implies$):
Suppose that $a \vee (b \wedge c) = (a \vee b) \wedge (a \vee c)$.
- ($\impliedby$):

□

9.2.5 Heyting Algebras

Definition 9.2.5.1. Let $(L, \vee, \wedge)$ be a lattice and $\to : L \times L \to L$. Then

- $\to$ is said to be a $(\vee, \wedge)$**-Heyting implication operator** if for each $a, x, y \in L$, $(x \wedge a) \leq_L y$ iff $x \leq_L (a \to y)$,
- $(L, \vee, \wedge, \to)$ is said to be a **Heyting algebra** if
  1. $(L, \vee, \wedge)$ is a lattice with 0,
  2. $\to : L \times L \to L$ is a $(L, \vee, \wedge)$-Heyting implication operator.

.

(ask if there is any way to show that L has a zero instead of showing it. Maybe there is a result saying that this is impossible in general.)

Note 9.2.5.2. When the context is clear, we say "Heyting implication operator" instead of "$(L, \vee, \wedge)$-Heyting implication operator" and we write L in place of $(L, \vee, \wedge, \to)$.

Exercise 9.2.5.3. Let L be a Heyting algebra. Then

1. L is a lattice with 1
2. for each $a \in L$, $a \to a = 1_L$.

(ask if there is any way to show that L has a zero instead of showing it. Maybe there is a result saying that this is impossible in general.)

Proof. Let $a \in L$. Define $y \in L$ by $y := a$. Then

$$\begin{aligned}(x \wedge a) &\leq a \\ &= y.\end{aligned}$$

Then by definition, we have that $x \leq a \to y$. Since $x \in L$ is arbitrary, we have that for each $x \in L$,

$$\begin{aligned}x &\leq (a \to y) \\ &= a \to a.\end{aligned}$$

Thus $a \to a$ is a one of L. Since $a \in L$ is arbitrary, we have that for each $a \in L$, $a \to a$ is a one of L.

1. Since L is a lattice, $L \neq \varnothing$. Thus there exists $a \in L$. Since $a \to a \in L$ and $a \to a$ is a one of L, L is a lattice with 1.
2. the part above implies that for each $a \in L$, $a \to a = 1_L$.

□

Exercise 9.2.5.4. Let $(L, \vee, \wedge)$ be a lattice with 0 and $\to: L \times L \to L$. Then $(L, \vee, \wedge, \to)$ is a Heyting algebra iff for each $a \in L$, $((\cdot \wedge a), (a \to \cdot))$ is a $(\leq_L, \leq_L)$-monotone Galois connection.
Can we get rid of assumption about having 0?

Proof. For each $a \in L$, define $f_a, g_a : L \to L$ by $f_a := \cdot \wedge a$ and $g_a := a \to \cdot$.

- $(\implies)$:
  Suppose that $(L, \vee, \wedge, \to)$ is a Heyting algebra. By definition, $\to: L \times L \to L$ is a $(L, \vee, \wedge)$-Heyting implication operator. Therefore, for each $a, x, y \in L$,

  $$\begin{aligned}f_a(x) \leq_L y &\iff (x \wedge a) \leq_L y \\ &\iff x \leq_L (a \to y) \\ &\iff x \leq_L g_a(y).\end{aligned}$$

  Thus for each $a \in L$, (f_a, g_a) is a $(\leq_L, \leq_L)$-monotone Galois connection.

- $(\impliedby)$:
  Suppose that for each $a \in L$, (f_a, g_a) is a $(\leq_L, \leq_L)$-monotone Galois connection.

  1. By assumption, $(L, \vee, \wedge)$ is a lattice with 0.
  2. We have that for each $a, x, y \in L$,

     $$\begin{aligned}(x \wedge a) \leq_L y &\iff f_a(x) \leq_L y \\ &\iff x \leq_L g_a(y) \quad \text{(by assumption)} \\ &\iff x \leq_L (a \to y).\end{aligned}$$

     Hence $\to: L \times L \to L$ is a $(\vee, \wedge)$-Heyting implication operator.

Thus $(L, \vee, \wedge, \to)$ is a Heyting algebra.

□

Exercise 9.2.5.5. Let L be a Heyting algebra. Then for each $a, b \in L$, $a \to b = \sup\{x \in L : x \wedge a \leq b\}$ (maybe write $\max A$? need to define $\max A$).

Proof. Let $a, b \in L$. Define $A \subset L$ by $A := \{x \in L : x \wedge a \leq b\}$. Define $y \in L$ by $y := b$. We note that

$$\begin{aligned} x \in A &\iff x \wedge a \leq b \\ &\iff x \wedge a \leq y \\ &\iff x \leq a \to y \quad \text{(by def)} \\ &\iff x \leq a \to b. \end{aligned}$$

1. Define $x_0 \in L$ by $x_0 := (a \to b)$. Since

$$\begin{aligned} x_0 &= a \to b \\ &\leq a \to b, \end{aligned}$$

   we have that $x_0 \in A$.

2. Let $x \in A$. Then

$$\begin{aligned} x &\leq a \to b \\ &= x_0. \end{aligned}$$

   Since $x \in A$ is arbitrary, we have that for each $x \in A$, $x \leq x_0$.

Exercise B.0.0.3 ref prev ex implies that $x_0 = \sup A$ (maybe write $\max A$? need to define $\max A$). □

Exercise 9.2.5.6. Let L be a Heyting algebra. Then for each $x, y \in L$, $x \leq y$ iff $x \to y = 1$.

Proof. Let $x, y \in L$. We note that $1 \wedge x = x$ and for each $a \in L$, $a = 1$ iff $1 \leq a$. Therefore

$$\begin{aligned} x \to y = 1 &\iff 1 \leq x \to y \\ &\iff 1 \wedge x \leq y \\ &\iff x \leq y. \end{aligned}$$

Since $x, y \in L$ are arbitrary, we have that for each $x, y \in L$, $x \leq y$ iff $x \to y = 1$. □

Definition 9.2.5.7. Let $(L, \vee, \wedge, \to)$ be a Heyting algebra. We define the **$\to$-negation operator**, denoted $\neg_\to : L \to L$, by $\neg_\to x := x \to 0$.

Note 9.2.5.8. When the context is clear, we write $\neg$ in place of $\neg_\to$.

Exercise 9.2.5.9. Let L be a Heyting algebra. Then $\neg : L \to L$ is a pseudocomplement.

Proof. FINISH!!!!, define pseudocomplement (and complement and orthocomplemented lattice, etc) □

9.2.6 Boolean Algebras

9.3 Lattice Structures

9.3.1 Irreducibility and Primality

Definition 9.3.1.1. Let $(L, \leq)$ be a poset.

- Suppose that $(L, \leq)$ is a join semilattice.
  - Let $x \in L$. Then x is said to be
    - $\leq$**-join irreducible** if
      1. x is not a zero of L,
      2. for each $a, b \in L$, $x = a \vee_{\leq} b$ implies that $x = a$ or $x = b$
    - $\leq$**-join prime**
      1. x is not a zero of L,
      2. if for each $a, b \in L$, $x \leq a \vee_{\leq} b$ implies that $x \leq a$ or $x \leq b$
  - We define
    - $\mathrm{irred}_{\vee}(L, \leq) := \{x \in L : x \text{ is } \leq\text{-join irreducible.}\}$
    - $\mathrm{prime}_{\vee}(L, \leq) := \{x \in L : x \text{ is } \leq\text{-join prime.}\}$
- Suppose that $(L, \leq)$ is a meet semilattice.
  - Let $x \in L$. Then x is said to be
    - $\leq$**-meet irreducible**
      1. x is not a one of L,
      2. if for each $a, b \in L$, $x = a \wedge_{\leq} b$ implies that $x = a$ or $x = b$
    - $\leq$**-meet prime** if
      1. x is not a one of L,
      2. for each $a, b \in L$, $a \wedge_{\leq} b \leq x$ implies that $a \leq x$ or $b \leq x$
  - We define
    - $\mathrm{irred}_{\wedge}(L, \leq) := \{x \in L : x \text{ is } \leq\text{-meet irreducible.}\}$
    - $\mathrm{prime}_{\wedge}(L, \leq) := \{x \in L : x \text{ is } \leq\text{-meet prime.}\}$

Note 9.3.1.2. When the context is clear,

- we say "join irreducible", "join prime", "meet irreducible" and "meet prime" in place of "$\leq$-join irreducible", "$\leq$-join prime", "$\leq$-meet irreducible" and "$\leq$-meet prime" respectively,
- we write $\mathrm{irred}_{\vee}(L)$, $\mathrm{prime}_{\vee}(L)$, $\mathrm{irred}_{\wedge}(L)$ and $\mathrm{prime}_{\wedge}(L)$ in place of $\mathrm{irred}_{\vee}(L, \leq)$, $\mathrm{prime}_{\vee}(L, \leq)$, $\mathrm{irred}_{\wedge}(L, \leq)$ and $\mathrm{prime}_{\wedge}(L, \leq)$ respectively

Exercise 9.3.1.3. Let $(L, \leq)$ be a poset.

1. Suppose that $(L, \leq)$ is a join semilattice. Then $\mathrm{prime}_{\vee}(L) \subset \mathrm{irred}_{\vee}(L)$.
2. Suppose that $(L, \leq)$ is a meet semilattice. Then $\mathrm{prime}_{\wedge}(L) \subset \mathrm{irred}_{\wedge}(L)$.

Proof.

1. Let $x \in \mathrm{prime}_{\vee}(L)$.
   (a) Since x is join prime, x is not at zero of L.
   (b) Let $a, b \in L$. Suppose that $x = a \vee b$. Then $x \leq a \vee b$. Since x is join prime, $x \leq a$ or $x \leq b$.

- Suppose that $x \leq a$. Then

$$\begin{aligned} a \vee b &= x \\ &\leq a \\ &\leq a \vee b \end{aligned}$$

  and therefore $x = a$. Thus $x \leq a$ implies that $x = a$.
- Similarly, $x \leq b$ implies that $x = b$.

Since $x \leq a$ or $x \leq b$, we have that $x = a$ or $x = b$. Since $a, b \in L$ with $x = a \vee b$ are arbitrary, we have that for each $a, b \in L$, $x = a \vee b$ implies that $x = a$ or $x = b$.

Thus $x \in \text{irred}_\vee(L)$. Since $x \in \text{prime}_\vee(L)$ is arbitrary, we have that for each $x \in \text{prime}_\vee(L)$, $x \in \text{irred}_\vee(L)$. Hence $\text{prime}_\vee(L) \subset \text{irred}_\vee(L)$.

2. Similar to (1) use duality.

□

9.3.2 Ideals and Filters

Definition 9.3.2.1. Let $(L, \vee, \wedge)$ be a lattice.

- Let $J \subset L$. Then
  - J is said to be an $(L, \vee, \wedge)$**-ideal** if
    1. $J \neq \varnothing$,
    2. for each $a, b \in J$, $a \vee b \in J$,
    3. for each $x \in L$ and $x \in J$, $x \wedge a \in J$,
  - J is said to be an $(L, \vee, \wedge)$**-filter** if
    1. $J \neq \varnothing$,
    2. for each $a, b \in J$, $a \wedge b \in J$,
    3. for each $x \in L$ and $a \in J$, $x \vee a \in J$.
- We define
  - $\mathcal{I}(L, \vee, \wedge) := \{I \subset L : I \text{ is an } (L, \vee, \wedge)\text{-ideal}\}$
  - $\mathcal{F}(L, \vee, \wedge) := \{F \subset L : F \text{ is a } (L, \vee, \wedge)\text{-filter}\}$

Note 9.3.2.2. When the context is clear,

- we may say "ideal" and "filter" instead of "$(L, \vee, \wedge)$-ideal" and "$(L, \vee, \wedge)$-filter" respectively,
- we may write $\mathcal{I}(L)$ and $\mathcal{F}(L)$ in place of $\mathcal{I}(L, \vee, \wedge)$ and $\mathcal{F}(L, \vee, \wedge)$ respectively.

Exercise 9.3.2.3. Let L be a lattice. If $L \neq \varnothing$, then

1. $L \in \mathcal{I}(L)$,
2. $L \in \mathcal{F}(L)$.

Proof.

1. (a) Since L is a lattice, $L \neq \varnothing$.
   (b) Clearly for each $a, b \in L$, $a \vee b \in L$
   (c) Clearly for each $x \in L$ and $b \in L$, $a \wedge b \in L$
2. Similar to (1) or use dualtity?.

□

Exercise 9.3.2.4. Let L be a lattice. Then for each $x \in L$,

1. $\downarrow x \in \mathcal{I}(L)$
2. $\uparrow x \in \mathcal{F}(L)$

Proof. Let $x \in L$.

1. (a) Since $x \in \downarrow x$, $\downarrow x \neq \varnothing$.
   (b) Let $a, b \in \downarrow x$. Then $a \leq x$ and $b \leq x$. Hence $x \in \mathrm{ub}(\{a, b\})$. Therefore
   $$\begin{aligned} a \vee b &= \sup\{a, b\} \\ &\leq x. \end{aligned}$$
   Thus $a \wedge b \in \downarrow x$. Since $a, b \in \downarrow x$ are arbitrary, we have that for each $a, b \in \downarrow x$, $a \vee b \in \downarrow x$.
   (c) Let $y \in L$ and $a \in \downarrow x$. Then $a \leq x$. Therefore
   $$\begin{aligned} y \wedge a &= \inf(\{y, a\}) \\ &\leq a \\ &\leq x. \end{aligned}$$
   Hence $y \wedge a \in \downarrow x$. Since $y \in L$ and $a \in \downarrow x$ are arbitrary, we have that for each $y \in L$ and $a \in \downarrow x$, $y \wedge a \in \downarrow x$.
2. use duality

□

Exercise 9.3.2.5. Let $(L, \vee, \wedge)$ be a lattice and $J \subset L$. Then

1. $J \in \mathcal{I}(L)$ iff $J \in \mathcal{F}(L^*)$,
2. $J \in \mathcal{F}(L)$ iff $J \in \mathcal{I}(L^*)$.

Proof.

1. • ($\Longrightarrow$):
   Suppose that $J \in \mathcal{I}(L)$.
   – Since $J \in \mathcal{I}(L, \vee, \wedge)$, $J \neq \varnothing$.
   – Since $J \in \mathcal{I}(L)$, for each $a, b \in J$,
   $$\begin{aligned} a \wedge^* b &= a \vee b \\ &\in J. \end{aligned}$$
   – Since $J \in \mathcal{I}(L)$, for each $x \in L$ and $a \in J$,
   $$\begin{aligned} &x \vee^* a \\ &\quad = x \wedge a \\ &\quad \in J. \end{aligned}$$
   Thus $J \in \mathcal{F}(L^*)$.
   • ($\Longleftarrow$) Suppose that $J \in \mathcal{F}(L^*)$. Then
2. Similar to (1) ref prev part

□

Exercise 9.3.2.6. Let L be a lattice and $J \subset L$. Then

1. (a) $J \in \mathcal{I}(L)$ iff J is a $\leq$-lower set and J is $\leq$-upward directed.
   (b) $J \in \mathcal{I}(L^*)$ iff J is a $\leq^*$-lower set and J is $\leq^*$-upward directed.
2. (a) $J \in \mathcal{F}(L)$ iff J is an $\leq$-upper set and J is $\leq$-downward directed.
   (b) $J \in \mathcal{F}(L^*)$ iff J is an $\leq^*$-upper set and J is $\leq^*$-downward directed.

FIX!!!
REWORK!!!!
THINK ABOUT WHAT STATEMENTS WE WANT AND SHORTEST WAY TO SHOW THEM

Proof.

1. (a)
   - ($\Longrightarrow$):
     Suppose that $J \in \mathcal{I}(L)$.
     - Let $x \in L$ and $a \in J$. Suppose that $x \leq a$. Then $x = \inf(\{a, b\}, \leq)$ and therefore
       $$\begin{aligned} x &= \inf(\{x, a\}, \leq) \\ &= x \wedge_{\leq} a \\ &\in J. \end{aligned}$$
       Since $a \in J$ and $x \in L$ with $x \leq a$ are arbitrary, we have that J is a down set.
     - * Since $J \in \mathcal{I}(L)$, we have that $J \neq \varnothing$.
       * Let $a, b \in J$. Since $J \in \mathcal{I}(L)$, we have that $a \vee b \in J$. Since $a \vee b = \sup(\{a, b\}, \leq)$ and $\{a, b\} \subset J$, we have that
       $$\begin{aligned} a \vee b &\in \mathrm{ub}(\{a, b\}, \leq) \cap J \\ &= \mathrm{ub}(\{a, b\}, \leq|_J) \quad \text{(by Exercise 2.7.1.7).} \end{aligned}$$
       Therefore $\mathrm{ub}(\{a, b\}, \leq|_J) \neq \varnothing$. Since $a, b \in J$ are arbitrary, we have that for each $a, b \in J$, $\mathrm{ub}(\{a, b\}, \leq|_J) \neq \varnothing$. Hence $\leq|_J$ is a direction on J.

     Thus $(J, \leq|_J)$ is a directed set. Therefore J is $\leq$-upward directed.
   - ($\Longleftarrow$):
     Suppose that J is a $\leq$-lower set and J is $\leq$-upward directed.
     i. Since J is $\leq$-upward directed, $(J, \leq|_J)$ is a directed set. Hence $J \neq \varnothing$.
     ii. Let $a, b \in J$. Since $(J, \leq|_J)$ is a directed set and $\{a, b\} \subset J$, we have that
     $$\begin{aligned} \mathrm{ub}(\{a, b\}, \leq) \cap J &= \mathrm{ub}(\{a, b\}, \leq|_J) \quad \text{(by Exercise 2.7.1.7)} \\ &\neq \varnothing. \end{aligned}$$
     Thus there exists
     $$c \in \mathrm{ub}(\{a, b\}, \leq) \cap J \tag{9.1}$$
     $$\subset \mathrm{ub}(\{a, b\}, \leq). \tag{9.2}$$
     Therefore $\sup(\{a, b\}, \leq) \leq c$. Since J is a $\leq$-lower set, $c \in J$ and $\sup(\{a, b\}, \leq) \leq c$, we have that
     $$\begin{aligned} a \vee b &= \sup(\{a, b\}, \leq) \\ &\in J. \end{aligned}$$
     Since $a, b \in J$ are arbitrary, we have that for each $a, b \in J$, $a \vee b \in J$.
     iii. Let $x \in L$ and $a \in J$. Then $x \wedge a \leq a$. Since J is a $\leq$-lower set, $x \wedge a \in J$. Since $x \in L$ and $a \in J$ are arbitrary, we have that for each $x \in L$ and $a \in J$, $x \wedge a \in J$.

   Therefore $J \in \mathcal{I}(L)$.

   (b) Exercise 9.1.6.13 and the previous part imply that $J \in \mathcal{I}(L^*)$ iff J is an $\leq^*$-lower set and J is $\leq^*$-upward directed.

2. (a) We have that

$$\begin{aligned} J \in \mathcal{F}(L) &\iff J \in \mathcal{I}(L^*) \quad \text{(by prev par} \\ &\iff J \text{ is an } \leq^*\text{-lower set and} \\ \iff J \text{ is an } \leq\text{-upper set and } J \text{ is } \leq\text{-downward directed} &\quad \text{(by prev ex in directed set section)} \end{aligned}$$

(b) Exercise 9.1.6.13 and the previous part imply that $J \in \mathcal{F}(L^*)$ iff J is an $\leq^*$-upper set and J is $\leq^*$-downward directed.

$$\begin{aligned} J \in \mathcal{F}(L^*) &\iff J \text{ is an } \leq^*\text{-upper set and } J \text{ is } \leq\text{-downward directed} \quad \text{(by prev part)} \\ &\iff J \text{ is a } \leq\text{-lower set and } J \text{ is } \leq\text{-upward directed} \\ &\iff J \text{ is a } \leq\text{-lower set and } J \text{ is } \leq\text{-upward directed} \end{aligned}$$

□

Exercise 9.3.2.7. Let L be a lattice.

1. If L has a zero, then for each $I \in \mathcal{I}(L)$, then $0 \in I$.
2. If L has a one, then for each $F \in \mathcal{F}(L)$, then $1 \in F$.

Proof.

1. Suppose that L has a zero. Let $J \in \mathcal{I}(L)$. Since $I \in \mathcal{I}(L)$, $I \neq \varnothing$ and a previous ex implies I is a lower set. Hence there exists $a \in I$. a prev ex implies that $0 \in \operatorname{lb} L$. Thus $0 \leq a$. Since I is a lower set, $0 \in I$.
2. Similar to (1).

□

Exercise 9.3.2.8. Let L be a lattice.

1. Let $(I_\alpha)_{\alpha \in A} \subset \mathcal{I}(L)$.
   (a) If $\bigcap_{\alpha \in A} I_\alpha \neq \varnothing$, then $\bigcap_{\alpha \in A} I_\alpha \in \mathcal{I}(L)$.
   (b) If $A = \{1, 2\}$, then $\bigcap_{\alpha \in A} I_\alpha \in \mathcal{I}(L)$.
   (c) If L has a zero, then $\bigcap_{\alpha \in A} I_\alpha \in \mathcal{I}(L)$.
2. Let $(F_\alpha)_{\alpha \in A} \subset \mathcal{F}(L)$.
   (a) If $\bigcap_{\alpha \in A} F_\alpha \neq \varnothing$, then $\bigcap_{\alpha \in A} F_\alpha \in \mathcal{F}(L)$.
   (b) If $A = \{1, 2\}$, then $\bigcap_{\alpha \in A} F_\alpha \in \mathcal{F}(L)$.
   (c) If L has a one, then $\bigcap_{\alpha \in A} F_\alpha \in \mathcal{F}(L)$.

Proof.

1. Set $I := \bigcap_{\alpha \in A} I_\alpha$.
   (a) Suppose that $I \neq \varnothing$.
      i. By assumption $I \neq \varnothing$.
      ii. Let $a, b \in I$ and $\alpha \in A$. Then $a, b \in I_\alpha$. Since $I_\alpha \in \mathcal{I}(L)$, $a \vee b \in I_\alpha$. Since $\alpha \in A$ is arbitrary, we have that for each $\alpha \in A$, $a \vee b \in I_\alpha$. Hence $a \vee b \in I$.

iii. Let $x \in L$, $a \in I$ and $\alpha \in A$. Then $a \in I_\alpha$. Since $I_\alpha \in \mathcal{I}(L)$, $x \in L$ and $a \in I_\alpha$, we have that $x \wedge a \in I_\alpha$. Since $\alpha \in A$ is arbitrary, we have that for each $\alpha \in A$, $x \wedge a \in I_\alpha$. Hence $x \wedge a \in I$. Since $x \in L$ and $a \in I$ are arbitrary, we have that for each $x \in L$ and $a \in I$, $x \wedge a \in I$.

Thus $I \in \mathcal{I}(L)$.

(b) Suppose that $A = \{1, 2\}$.

i. Since $I_1, I_2 \in \mathcal{I}(L)$, $I_1 \neq \varnothing$ and $I_2 \neq \varnothing$. Thus there exist $x_1 \in I_1$ and $x_2 \in I_2$. Since $I_1 \in \mathcal{I}(L)$, $x_2 \in L$ and $x_1 \in I_1$, we have that $x_1 \wedge x_2 \in I_1$. Similarly, $x_1 \wedge x_2 \in I_2$. Hence $x_1 \wedge x_2 \in I_1 \cap I_2$ and $I_1 \cap I_2 \neq \varnothing$. Part 1($a$) implies that $I_1 \cap I_2 \in \mathcal{I}(L)$.

(c) Suppose that L has a zero. Let $\alpha \in A$. Since $I_\alpha \in \mathcal{I}(L)$, a previous exercise implies that $0 \in I_\alpha$. Since $\alpha \in A$ is arbitrary, we have that for each $\alpha \in A$, $0 \in I_\alpha$. Hence $0 \in I$. Thus $I \neq \varnothing$. The previous part implies that $I \in \mathcal{I}(L)$.

2. Similar to (1).

□

Definition 9.3.2.9. (consider changing notation) Let L be a lattice and $A \subset L$.

- We define
  - $\text{Ideals}(A, L) := \{I \in \mathcal{I}(L) : S \subset I\}$
  - $\text{Filters}(A, L) := \{F \in \mathcal{F}(L) : S \subset F\}$
- We define
  - the L-ideal generated by A, denoted $\text{IdealGen}(A, L)$, by $\text{IdealGen}(A, L) := \bigcap_{I \in \text{Ideals}(A,L)} I$
  - the L-filter generated by A, denoted $\text{IdealGen}(A, L)$, by$\text{FilterGen}(A, L) := \bigcap_{F \in \text{Filters}(A,L)} F$

Note 9.3.2.10. the prev ex implies that $\text{IdealGen}(A, L) \in \text{Ideals}(A, L)$ and $\text{FilterGen}(A, L) \in \text{Filters}(A, L)$.

Exercise 9.3.2.11. Let L be a lattice and $B \subset L$.

1. If B is upward directed, then
   (a) $\downarrow B \in \mathcal{I}(L)$
   (b) $\text{IdealGen}(B) = \downarrow(B, \leq_L)$.
2. If B is downward directed, then
   (a) $\uparrow B \in \mathcal{F}(L)$
   (b) $\text{FilterGen}(B) = \uparrow(B, \leq_L)$.

Proof.

1. Suppose that B is upward directed.
   (a) Exercise B.0.0.3(ref ex) implies that $\downarrow B$ is upward directed and Exercise B.0.0.3 (ref ex) implies that $\downarrow B$ is an lower set. Exercise B.0.0.3 then implies that $\downarrow B \in \mathcal{I}(L)$.
   (b) The prev part implies that $\downarrow B \in \text{Ideals}(B, L)$. Therefore

$$\begin{aligned}\text{IdealGen}(B, L) &= \bigcap_{I \in \text{Ideals}(B,L)} I \\ &\subset \uparrow B.\end{aligned}$$

We note that for each $I \in \mathcal{I}(L)$, I is a lower set. Thus $\text{Ideals}(B, L) \subset \mathcal{L}(B, \leq_L)$. Hence

$$\begin{aligned}\downarrow B &= \bigcap_{\in \text{Ideals}(B,L)} I \\ &\subset \uparrow B.\end{aligned}$$

2. Suppose that B is $\leq$-downward directed. Exercise B.0.0.3 implies that B is $\leq^*$-upward directed. the prev part implies that $\uparrow(B, \leq^*) \in \mathcal{I}(L)$. Exercise B.0.0.3 then implies that $\uparrow(B, \leq^*) \in \mathcal{I}(L)$

   make new notation $\leq_L :=$ the unique partial order on L compatible with $\vee, \wedge$. Show $\leq_{L^*} = \leq_L^*$, make note that we write $\leq$ in place

□

Definition 9.3.2.12. define prefilter or filter basis, preideal or ideal basis as a downward directed/upward directed set, maybe don't need to, but look into what we need for topology.

Exercise 9.3.2.13. Let L be a lattice. Then

1. for each $I_1, I_2 \in \mathcal{I}(L)$, $\{x \in L : \text{there exist } a \in I_1 \text{ and } b \in I_2 \text{ such that } x \leq a \vee b\} \in \mathcal{I}(L)$.
2. for each $F_1, F_2 \in \mathcal{F}(L)$, $\{x \in L : \text{there exist } a \in F_1 \text{ and } b \in F_2 \text{ such that } a \vee b \leq x\} \in \mathcal{F}(L)$ (maybe FIX!!!).

Proof.

1. Let $I_1, I_2 \in \mathcal{I}(L)$. Set $I := \{x \in L : \text{there exist } a \in I_1 \text{ and } b \in I_2 \text{ such that } x \leq a \vee b\}$.
   (a) Since $I_1, I_2 \in \mathcal{I}(L)$, we have that $I_1 \neq \varnothing$ and $I_2 \neq \varnothing$. Hence there exist $x_1 \in I_1$ and $x_2 \in I_2$. Since $x_1 \leq x_1 \vee x_2$, we have that $x_1 \in I$. Thus $I \neq \varnothing$.
   (b) Let $a, b \in I$. Then there exist $x_a, x_b \in I_1$ and $y_a, y_b \in I_2$ such that $a \leq x_a \vee y_a$ and $b \leq x_b \vee y_b$. Define $z_1, z_2 \in L$ by $z_1 := x_a \vee x_b$ and $z_2 := y_a \vee y_b$. Since $I_1, I_2 \in \mathcal{I}(L)$, we have that $z_1 \in I_1$ and $z_2 \in I_2$. We note that
   $$\begin{aligned} a &\leq x_a \vee y_a \\ &\leq (x_a \vee y_a) \vee (x_b \vee y_b) \\ &= (x_a \vee x_b) \vee (y_a \vee y_b) \\ &= z_1 \vee z_2. \end{aligned}$$
   and similarly $b \leq z_1 \vee z_2$. Therefore $z_1 \vee z_2 \in \mathrm{ub}(\{a, b\}, \leq)$. Hence
   $$\begin{aligned} a \vee b &= \sup(\{a, b\}, \leq) \\ &\leq z_1 \vee z_2. \end{aligned}$$
   Thus $a \vee b \in I$. Since $a, b \in I$ are arbitrary, we have that for each $a, b \in I$, $a \vee b \in I$.
   (c) Let $x \in L$ and $y \in I$. Then there exists $a \in I_1$ and $b \in I_2$ such that $y \leq a \vee b$. Then
   $$\begin{aligned} x \wedge y &\leq y \\ &\leq a \vee b. \end{aligned}$$
   Therefore $x \wedge y \in I$. Since $x \in L$ and $y \in I$ are arbitrary, we have that for each $x \in L$ and $y \in I$, $x \wedge y \in I$.

   Therefore $I \in \mathcal{I}(L)$.
2. Similar to (1). FINISH!!!

□

Definition 9.3.2.14. Let L be a lattice. Define $\leq_{\mathcal{I}(L)} := \subset|_{\mathcal{I}(L)}$ (maybe standardize notation).

Exercise 9.3.2.15. Let L be a lattice.

1. Let $I_1, I_2 \in \mathcal{I}(L)$.
   (a) $\sup(\{I_1, I_2\}, \leq_{\mathcal{I}(L)}) = \{x \in L : \text{there exist } a \in I_1 \text{ and } b \in I_2 \text{ such that } x \leq a \vee b\}$.
   (b) $\inf(\{I_1, I_2\}, \leq_{\mathcal{I}(L)}) = I_1 \cap I_2$
2. Set $\leq_{\mathcal{F}(L)} := \subset|_{\mathcal{F}(L)}$ (maybe standardize notation). Let $F_1, F_2 \in \mathcal{F}(L)$. FIX!!!

(a) $\sup(\{F_1, F_2\}, \leq_{\mathcal{I}(L)}) = \{x \in L : \text{there exist } a \in F_1 \text{ and } b \in F_2 \text{ such that } x \leq a \vee b\}$.

(b) $\inf(\{F_1, F_2\}, \leq_{\mathcal{F}(L)}) = F_1 \cap F_2$

Proof.

1. (a) Set $I := \{x \in L : \text{there exist } a \in I_1 \text{ and } b \in I_2 \text{ such that } x \leq a \vee b\}$. (ref previous ex) implies that $I \in \mathcal{I}(L)$.

   i. • Let $a \in I_1$. Since $I_2 \in \mathcal{I}(L)$, $I_2 \neq \varnothing$. Hence there exists $b \in I_2$. Then $a \leq a \vee b$. Since $a \in I_1$, $b \in I_2$ and $a \leq a \vee b$, we have that $a \in I$. Since $a \in I_1$ is arbitrary, we have that for each $a \in I_1$, $a \in I$. Thus $I_1 \subset I$.
   • Similarly, $I_2 \subset I$.

   Since $I_1, I_2 \subset I$, we have that $I \in \mathrm{ub}(\{I_1, I_2\}, \leq_{\mathcal{I}(L)})$.

   ii. Let $K \in \mathrm{ub}(\{I_1, I_2\}, \leq_{\mathcal{I}(L)})$ and $x \in I$. Then $I_1, I_2 \subset K$ and there exist $a \in I_1$ and $b \in I_2$ such that $x \leq a \vee b$. Since $a \in I_1$, $b \in I_2$ and $I_1, I_2 \subset K$, we have that $a, b \in K$. Since $K \in \mathcal{I}(L)$, $a \vee b \in K$. Since $K \in \mathcal{I}(L)$ (ref a prev ex) implies that K is a lower set. Since K is a lower set, $a \vee b \in K$ and $x \leq a \vee b$, we have that $x \in K$. Since $x \in I$ is arbitrary, we have that for each $x \in I$, $x \in K$. Hence $I \subset K$. Since $K \in \mathrm{ub}(\{I_1, I_2\}, \leq_{\mathcal{I}(L)})$ is arbitrary, we have that for each $K \in \mathrm{ub}(\{I_1, I_2\}, \leq_{\mathcal{I}(L)})$, $I \leq_{\mathcal{I}(L)} K$.

   Since

   i. $I \in \mathrm{ub}(\{I_1, I_2\}, \leq_{\mathcal{I}(L)})$,

   ii. for each $K \in \mathrm{ub}(\{I_1, I_2\}, \leq_{\mathcal{I}(L)})$, $I \leq_{\mathcal{I}(L)} K$,

   we have that $I = \sup(\{I_1, I_2\}, \leq_{\mathcal{I}(L)})$.

   (b) (ref previous ex) implies that $I_1 \cap I_2 \in \mathcal{I}(L)$. Since $I_1 \cap I_2 \subset I_1$ and $I_1 \cap I_2 \subset I_2$, we have that $I_1 \cap I_2 \in \mathrm{lb}(\{I_1, I_2\}, \leq_{\mathcal{I}(L)})$. Since $(\mathcal{I}(L), \leq_{\mathcal{I}(L)})$ is a subposet of $(\mathcal{P}(L), \subset)$, (define subproset/subposet and make exercise for the following fact) implies that

$$\begin{aligned}\inf(\{I_1, I_2\}, \leq_{\mathcal{I}(L)}) &\leq \inf(\{I_1, I_2\}, \subset) \\ &= I_1 \cap I_2 \\ &\leq \inf(\{I_1, I_2\}, \leq_{\mathcal{I}(L)}).\end{aligned}$$

   Hence $I_1 \cap I_2 = \inf(\{I_1, I_2\}, \leq_{\mathcal{I}(L)})$.

2. Similar to (1). FINISH!!!

□

Definition 9.3.2.16. Let L be a lattice. Define $\vee_{\mathcal{I}(L)}, \wedge_{\mathcal{I}(L)} : \mathcal{I}(L) \times \mathcal{I}(L) \to \mathcal{I}(L)$ and $\vee_{\mathcal{F}(L)}, \wedge_{\mathcal{F}(L)} : \mathcal{F}(L) \times \mathcal{F}(L) \to \mathcal{F}(L)$ by

1. • $I_1 \vee_{\mathcal{I}(L)} I_2 := \{x \in L : \text{there exist } a \in I_1 \text{ and } b \in I_2 \text{ such that } x \leq a \vee b\}$
   • $I_1 \wedge_{\mathcal{I}(L)} I_2 := I_1 \cap I_2$.

2. FIX!!!

   • $F_1 \vee_{\mathcal{F}(L)} F_2 := \{x \in L : \text{there exist } a \in F_1 \text{ and } b \in F_2 \text{ such that } x \leq a \vee b\}$
   • $F_1 \wedge_{\mathcal{F}(L)} F_2 := F_1 \cap F_2$.

Exercise 9.3.2.17. Let L be a lattice. Then

1. $(\mathcal{I}(L), \vee_{\mathcal{I}(L)}, \wedge_{\mathcal{I}(L)})$ is a lattice

2. $(\mathcal{F}(L), \vee_{\mathcal{F}(L)}, \wedge_{\mathcal{F}(L)})$ is a lattice

Proof.

1. the previous exercise implies that $(\mathcal{I}(L), \leq_{\mathcal{I}(L)})$ is an ordered lattice. Exercise 9.1.6.6 then implies that $(\mathcal{I}(L), \vee_{\mathcal{I}(L)}, \wedge_{\mathcal{I}(L)})$ is a lattice.

2.

□

9.3.3 Properties of Ideals and Filters

Definition 9.3.3.1. Let L be a lattice.

- Let $I \in \mathcal{I}(L)$. Then I is said to be
  - **proper** if $I \neq L$.
  - **maximal** if I is maximal in $(\mathcal{I}(L) \setminus \{L\}, \subset)$.
  - **prime** if
    1. I is proper
    2. for each $a, b \in L$, $a \wedge b \in I$ implies that $a \in I$ or $b \in I$.
  - **principal** if there exists $a \in L$ such that $I = \downarrow a$.
- Let $F \in \mathcal{F}(L)$. Then F is said to be
  - **proper** if $F \neq L$.
  - **maximal** if F is maximal in $(\mathcal{F}(L) \setminus \{L\}, \subset)$.
  - **prime** if
    1. F is proper
    2. for each $a, b \in L$, $a \vee b \in F$ implies that $a \in F$ or $b \in F$.
  - **principal** if there exists $a \in L$ such that $F = \uparrow a$.

Exercise 9.3.3.2. Let L be a lattice.

1. For each $I \in \mathcal{I}(L)$.
   (a) if I is proper, then for each $x \in I$, x is not a one of L.
   (b) if I is prinicpal, then I is proper iff for each $x \in I$, x is not a one of L.
2. For each $F \in \mathcal{F}(L)$.
   (a) If F is proper, then for each $x \in F$, x is not a zero of L.
   (b) If F is prinicpal, then F is proper iff for each $x \in F$, x is not a zero of L.

Proof.

1. Let $I \in \mathcal{I}(L)$.
   (a) Suppose that I is proper. Let $x \in I$. For the sake of contradiction, suppose that x is a one of L. (previous ex) then implies that $(L, \leq) = \{x\}$. Let $y \in L$. Then $y \leq x$. Since I is a lower set, $x \in I$ and $y \leq x$, we have that $y \in I$. Since $y \in L$ is arbitrary, we have that for each $y \in L$, $y \in I$. Thus
   $$\begin{aligned} L &\subset I \\ &\subset L. \end{aligned}$$
   Thus $I = L$ and I is not proper. Therefore x is not a one of L. Since $x \in I$ is arbitrary, we have that for each $x \in I$, x is not a one of L.
   (b) Suppose that I is principal.
   - $(\implies)$:
     If I is proper, then $1(a)$ implies that for each $x \in I$, x is not a one of L.

- ($\Longleftarrow$):
  Suppose that I is not proper. Then $I = L$. Since I is principal, there exists $a \in I$ such that $I = \downarrow a$. Let $x \in L$. Since

$$\begin{aligned} x &\in L \\ &= I \\ &= \downarrow a, \end{aligned}$$

  we have that $x \leq a$. Hence

$$\begin{aligned} x \wedge a &= \inf\{x, a\} \\ &= x. \end{aligned}$$

  Since $x \in L$ is arbitrary, we have that for each $x \in L$, $x \wedge a = x$. Hence a is a one of L. Therefore I is not proper implies that there exists $a \in I$ such that a is a one of L. By contrapositive, we have that if for each $x \in I$, x is not a one of L, then I is proper.

2. FINISH!!

□

Exercise 9.3.3.3. Let L be a lattice and $x \in L$.

1. Then $\downarrow x \in \mathcal{I}_p(L)$ iff $x \in \text{prime}_\wedge(L)$.
2. Then $\uparrow x \in \mathcal{F}_p(L)$ iff $x \in \text{prime}_\vee(L)$.

Proof.

1. - ($\Longrightarrow$):
     Suppose that $\downarrow x \in \mathcal{I}_p(L)$. Then $\downarrow x$ is proper and for each $a, b \in L$, $a \wedge b \in \downarrow x$ implies that $a \in \downarrow x$ or $b \in \downarrow x$.
     - Since $\downarrow x$ is proper, previous exercise implies that x is not a one of L.
     - Let $a, b \in L$. Suppose that $a \wedge b \leq x$. Since $\downarrow x$ is a lower set, $x \in \downarrow x$ and $a \wedge b \leq x$, we have that $a \wedge b \in \downarrow x$. Therefore $a \in \downarrow x$ or $b \in \downarrow x$. Then $a \leq x$ or $b \leq x$. Since $a, b \in L$ with $a \wedge b \leq x$ are arbitrary, we have that for each $a, b \in L$, $a \wedge b \leq x$ implies that $a \leq x$ or $b \leq x$.

     Thus $x \in \text{prime}_\wedge(L)$.
   - ($\Longleftarrow$):
     Suppose that $x \in \text{prime}_\wedge(L)$.
     - For the sake of contradiction, suppose that $\downarrow x$ is not proper. Then $\downarrow x = L$. Thus $\text{ub}\, L = \{x\}$. A previous ex implies that x is a one of L. This is a contracition since $x \in \text{prime}_\wedge(L)$. Therefore $\downarrow x$ is proper.
     - Let $a, b \in L$. Suppose that $a \wedge b \in \downarrow x$. Then $a \wedge b \leq x$. Since $x \in \text{prime}_\wedge(L)$, $a \leq x$ or $b \leq x$. Hence $a \in \downarrow x$ or $b \in \downarrow x$. Since $a, b \in L$ with $a \wedge b \in \downarrow x$ are arbitrary, we have that for each $a, b \in L$, $a \wedge b \in \downarrow x$ implies that $a \in \downarrow x$ or $b \in \downarrow x$.

     Therefore $\downarrow x \in \mathcal{I}_p(L)$.
2. Similar to (1) FIX/CHECK/FINISH!!!

□

Exercise 9.3.3.4. Let L be a lattice.

1. For each $I \in \mathcal{I}(L)$, $I \in \mathcal{I}_p(L)$ iff $I^c \in \mathcal{F}(L)$.
2. For each $F \in \mathcal{F}(L)$, $F \in \mathcal{F}_p(L)$ iff $F^c \in \mathcal{I}(L)$.

Proof.

1. Let $I \in \mathcal{I}(L)$.

- ($\Longrightarrow$):
  Suppose that $I \in \mathcal{I}_p(L)$. Then I is proper and for each $a, b \in L$, $a \wedge b \in I$ implies that $a \in I$ or $b \in I$.
  (a) Since I is proper, $I \neq L$. Therefore $I^c \neq \varnothing$
  (b) Let $a, b \in I^c$. For the sake of contradiction, suppose that $a \wedge b \in I$. Since $I \in \mathcal{I}_p(L)$, $a \in I$ or $b \in I$. This is a contradiction since $a \in I^c$ and $b \in I^c$. Therefore $a \wedge b \notin I$. Hence $a \wedge b \in I^c$. Since $a, b \in I^c$ are arbitrary, we have that for each for each $a, b \in I^c$, $a \wedge b \in I^c$.
  (c) Since $I \in \mathcal{I}(L)$, a previous exercise implies that I is a lower set. an exercise in proset section implies that I^c is an upper set. Let $x \in L$ and $a \in I^c$. Since

$$\begin{aligned} x \vee a &\geq a \\ &\in I^c \end{aligned}$$

  and I^c is an upper set, we have that $x \vee a \in I^c$. Since $x \in L$ and $a \in I^c$ are arbitrary, we have that for each for each $x \in L$ and $a \in I^c$, $x \vee a \in I^c$.1

  Thus $I^c \in \mathcal{F}(L)$.
- ($\Longleftarrow$):
  Suppose that $I^c \in \mathcal{F}(L)$.
  (a) Since $I^c \in \mathcal{F}(L)$, $I^c \neq \varnothing$. Then $I \neq L$ and therefore I is proper.
  (b) Let $a, b \in L$. Suppose that $a \wedge b \in I$. For the sake of contradiction, suppose that $a \notin I$ and $b \notin I$. Then $a \in I^c$ and $b \in I^c$. Since $I^c \in \mathcal{F}(L)$, we have that $a \wedge b \in I^c$. This is a contradiction since $a \wedge b \in I$. Hence $a \in I$ or $b \in I$. Since $a, b \in L$ with $a \wedge b \in I$ are arbitrary, we have that for each $a, b \in L$, $a \wedge b \in I$ implies that $a \in I$ or $b \in I$.

  Thus $I \in \mathcal{I}_p(L)$.

2. use duality.

□

9.4 Contructions with Lattices

9.4.1 Sublattices

Definition 9.4.1.1. Let $(L, \vee_L, \wedge_L)$ and $(M, \vee_M, \wedge_M)$ be lattices. Then $(M, \vee_M, \wedge_M)$ is said to be a **sublattice of** $(L, \vee_L, \wedge_L)$ if

1. $M \subset L$
2. ι_M is a $((\vee_L, \wedge_L), (\vee_M, \wedge_M))$-homomorphism.

Exercise 9.4.1.2. Let $(L, \vee_L, \wedge_L)$ and $(M, \vee_M, \wedge_M)$ be lattices. Then $(M, \vee_M, \wedge_M)$ is a sublattice of L iff

1. $M \subset L$
2. $\vee_L|_{M^2} = \vee_M$ and $\wedge_L|_{M^2} = \wedge_M$.

Proof.

- ($\Longrightarrow$):
  Suppose that $(M, \vee_M, \wedge_M)$ is a sublattice of L.
  1. By definition, $M \subset L$.
  2. By definiton, ι_M is a $((\vee_L, \wedge_L), (\vee_M, \wedge_M))$-homomorphism. Then for each $a, b \in M$,

$$\begin{aligned} \vee_M(a, b) &= a \vee_M b \\ &= \iota_M(a \vee_M b) \\ &= \iota_M(a) \vee_L \iota_M(b) \\ &= a \vee_L b \\ &= \vee_L|_{M^2}(a, b). \end{aligned}$$

  Hence $\vee_L|_{M^2} = \vee_M$. Similarly, $\wedge_L|_{M^2} = \wedge_M$.

- ($\Longleftarrow$):
  Suppose that

  1. $M \subset L$
  2. $\vee_L|_{M^2} = \vee_M$ and $\wedge_L|_{M^2} = \wedge_M$.

  1. By assumption $M \subset L$.
  2. Since $\vee_L|_{M^2} = \vee_M$, we have that for each $a, b \in M$,

$$\begin{aligned}\iota_M(a \vee_M b) &= a \vee_M b \\ &= \vee_M(a, b) \\ &= \vee_L|_{M^2}(a, b) \\ &= \vee_L(a, b) \\ &= a \vee_L b.\end{aligned}$$

  Hence ι_M is a $(\vee_M, \vee_L)$-homomorphism. Similarly, ι_M is a $(\wedge_M, \wedge_L)$-homomorphism. Therefore ι_M is a $((\vee_L, \wedge_L), (\vee_M, \wedge_M))$-homomorphism.

Thus $(M, \vee_M, \wedge_M)$ is a sublattice of L.

FINISH!!!! □

need to make preliminary ex showing that if $M \subset L$, $M \neq \varnothing$ and $\vee(M^2), \wedge(M^2) \subset M$, then $(M, \vee|_{M^2}, \wedge|_{M^2})$ is a lattice

Exercise 9.4.1.3. Let $(L, \vee, \wedge)$ be a lattice and $M \subset L$. Then $(M, \vee|_{M^2}, \wedge|_{M^2})$ is a sublattice of $(L, \vee, \wedge)$ iff

1. $M \neq \varnothing$
2. $\vee(M^2), \wedge(M^2) \subset M$.

Proof. Set $\vee_M := \vee|_{M^2}$ and $\wedge_M := \wedge|_{M^2}$

- ($\Longrightarrow$):
  Suppose that $(M, \vee_M, \wedge_M)$ is a sublattice of $(L, \vee, \wedge)$. Since $(M, \vee_M, \wedge_M)$ is a sublattice of $(L, \vee, \wedge)$, we have that $(M, \vee_M, \wedge_M)$ is a lattice.

  1. Since $(M, \vee_M, \wedge_M)$ is a lattice, $M \neq \varnothing$.
  2. Since $(M, \vee_M, \wedge_M)$ is a lattice, $\vee_M(M^2), \wedge_M(M^2) \subset M$.

- ($\Longleftarrow$):
  Suppose that

  1. $M \neq \varnothing$
  2. $\vee(M^2), \wedge(M^2) \subset M$.

  1. By assumption $M \neq \varnothing$,
  2. Since $\vee, \wedge$ are semilattice operators, $\vee, \wedge$ satisfy the lattice absorption identities and $\vee(M^2), \wedge(M^2) \subset M$, we have that $\vee_M, \wedge_M$ are semilattice operators and $\vee|_M, \wedge|_M$ satisfy the lattice absorption identities.

  Thus $(M, \vee_M, \wedge_M)$ is a lattice. By assumption, $M \subset L$ and $\vee_M = \vee|_{M^2}$, $\wedge_M = \wedge|_{M^2}$, we have that $(M, \vee_M, \wedge_M)$ is a sublattice of $(L, \vee, \wedge)$.

□

9.4.2 Product Lattices

9.4.3 Coproduct Lattices

9.4.4 Quotient Lattices

Chapter 10

Complete Lattices

maybe move complete lattices to order chapter? two choices: a) cover complete lattices after order and before other structures (including lattices?) so that we can define the 'subgroup/ring/lattice generated by a subset' or a 'subquiver genereated by a sub set of vertices/edges' using the ideas of galois connection between two posets, one of which has extra structure b) cover complete lattices after algebraic structures with lattices and note that we can capture all the subobjects with the notion of galois connection between complete lattices and give groups, rings etc as fundamental examples of complete lattices.
idk which is best option, it might make more sense to keep complete lattices after lattices, since we can talk about completely prime ideals and other lattice-type objects, and we can give fundamental examples of complete lattices that were discussed prior like groups/rings/lattice ideals/etc

10.1 Introduction

Exercise 10.1.0.1. Let $(X, \leq)$ be a poset. Then for each $A \subset X$, $\sup A$ exists iff for each $A \subset X$, $\inf A$ exists.

Proof.

- $(\implies)$:
  Suppose that for each $A \subset X$, $\sup A$ exists. Let $A \subset X$. We note that

  $$\sup(\operatorname{lb} A) \text{ exists} \iff \inf A \text{ exists} \quad \text{(by Exercise 2.7.3.2).}$$

  Since $\operatorname{lb} A \subset X$, then by assumption, $\sup(\operatorname{lb} A)$ exists. Therefore $\inf A$ exists. Since $A \subset X$ is arbitrary, we have that for each $A \subset X$, $\inf A$ exists.

- $(\impliedby)$:
  Similar to $(\implies)$.

$\square$

Definition 10.1.0.2. Let $(X, \leq)$ be a poset.

- Then $(X, \leq)$ is said to be a **complete lattice** if for each $A \subset X$, $\sup A$ exists.
- If $(X, \leq)$ is a complete lattice, we define the **arbitrary join and meet operators on** X, denoted $\bigvee_{\leq} : \mathcal{P}(X) \to X$ and $\bigwedge_{\leq} : \mathcal{P}(X) \to X$ respectively, by $\bigvee_{\leq} A := \sup(A, \leq)$ and $\bigwedge_{\leq} A := \inf(A, \leq)$.

Note 10.1.0.3. When the context is clear, we may write $\bigvee$ and $\bigwedge$ in place of $\bigvee_{\leq}$ and $\bigwedge_{\leq}$ respectively.

Exercise 10.1.0.4. Let $(X, \leq)$ be a complete lattice. Then $(X, \vee_{\leq}, \wedge_{\leq}, \inf(L, \leq), \sup(L, \leq),)$ is a bounded lattice.

Proof. FIX!!!

1. make previous exercise implies that $\mathrm{lb}(\varnothing, \leq) = X$. Since $(X, \leq)$ is a complete lattice and $\varnothing \subset X$, there exists $x_1 \in X$ such that $x_1 := \inf \varnothing$. In particular, $X \neq \varnothing$.

2.
   - make previous ex saying $\inf \varnothing = \sup X$ and $\sup \varnothing = \inf X$ with the following logic

$$\begin{aligned} &\inf \varnothing \\ &\qquad \in \mathrm{lb}\, \varnothing \cap \mathrm{ub}(\mathrm{lb}\, \varnothing) \\ &\qquad = X \cap \mathrm{lb}(X) \\ &\qquad = \mathrm{lb}\, X. \end{aligned}$$

     ref an ex implying that $\mathrm{lb}\, X = \inf X$. Hence X is a one of X.
   - similar x_0

Clearly $\bigvee X$ is a one of X and $\bigwedge X$ is a zero of X. □

10.2 Complete Lattice Homomorphisms

Definition 10.2.0.1. Let $(X, \leq_X), (Y, \leq_Y)$ be complete lattices and $f : X \to Y$. Then f is said to

1. $(\leq_X, \leq_Y)$**-arbitrary join preserving** if for each $A \subset X$, $f(\sup(A, \leq_X)) = \sup(f(A), \leq_Y)$
2. $(\leq_X, \leq_Y)$**-arbitrary meet preserving** if for each $A \subset X$, $f(\inf(A, \leq_X)) = \inf(f(A), \leq_Y)$

Exercise 10.2.0.2. Let $(X, \leq_X), (Y, \leq_Y)$ be complete lattices and $f : X \to Y$. Then

1. f is $(\leq_X, \leq_Y)$-arbitrary join preserving iff f is $(\leq_X^*, \leq_Y^*)$-arbitrary meet preserving.
2. f is $(\leq_X, \leq_Y)$-arbitrary meet preserving iff f is $(\leq_X^*, \leq_Y^*)$-arbitrary join preserving.

Proof.

1.
   - $(\implies)$:
     Suppose that f is $(\leq_X, \leq_Y)$-arbitrary join preserving. Then a prev ex in poset section on suprema implies that for each $A \subset X$,

$$\begin{aligned} f(\inf(A, \leq_X^*)) &= f(\sup(A, \leq_X)) && (10.1) \\ &= \sup(f(A), \leq_Y) && (10.2) \\ &= \inf(f(A), \leq_Y^*). && (10.3) \end{aligned}$$

     Hence f is $(\leq_X^*, \leq_Y^*)$-arbitrary meet preserving.
   - $(\impliedby)$:
     Similar to $(\implies)$. ref.

2. Similar to (1). ref.

□

Exercise 10.2.0.3. Let $(X, \leq_X), (Y, \leq_Y)$ be complete lattices.

1. For each $f : X \to Y$, f is arbitrary join preserving iff f has a right adjoint.
2. For each $g : Y \to X$, g is arbitrary meet preserving iff g has a left adjoint.

Proof.

1. • ($\Longrightarrow$):
   Let $f : X \to Y$. Suppose that f is arbitrary join preserving. an ex in subsection on semilattice homomorphisms Exercise B.0.0.3 implies that f is monotone. Let $b \in Y$. Exercise 2.7.3.6 implies that $\sup \downarrow b = b$. Define $a \in X$ by $a := \sup f^{-1}(\downarrow b)$.
   – Let $x \in f^{-1}(\downarrow b)$. Then
   $$\begin{aligned} x &\leq \sup f^{-1}(\downarrow b) \\ &= a. \end{aligned}$$
   Thus $x \in \downarrow a$. Since $x \in f^{-1}(\downarrow b)$ is arbitrary, we have that for each $x \in f^{-1}(\downarrow b)$, $x \in \downarrow a$. Hence $f^{-1}(\downarrow b) \subset \downarrow a$.
   – We note that for each $x \in X$,
   $$\begin{aligned} x \in \downarrow a &\iff x \leq_X a \\ &\Longrightarrow f(x) \leq_Y f(a). \end{aligned}$$
   Let $x \in \downarrow a$. An exercise in basic set theory ex implies that $f(f^{-1}(\downarrow b)) \subset \downarrow b$. Exercise 2.7.3.4 then implies that
   $$\begin{aligned} f(x) &\leq_Y f(a) \\ &= f(\sup f^{-1}(\downarrow b)) \\ &= \sup f(f^{-1}(\downarrow b)) \\ &\leq \sup \downarrow b \\ &= b. \end{aligned}$$
   Then $f(x) \in \downarrow b$. Hence $x \in f^{-1}(\downarrow b)$. Since $x \in \downarrow a$ is arbitrary, we have that for each $x \in \downarrow a$, $x \in f^{-1}(\downarrow b)$. Hence $\downarrow a \subset f^{-1}(\downarrow b)$.

   Since $f^{-1}(\downarrow b) \subset \downarrow a$ and $\downarrow a \subset f^{-1}(\downarrow b)$, we have that $f^{-1}(\downarrow b) = \downarrow a$. Since $b \in Y$ is arbitrary, we have that for each $b \in Y$, there exists $a \in X$ such that $f^{-1}(\downarrow b) = \downarrow a$. A prev ex in proset section on galois connections Exercise B.0.0.3 then implies that f has a right adjoint.
   • ($\Longleftarrow$):
   Suppose that f has a right adjoint. Then there exists $f^* : Y \to X$ such that (f, f^*) is a monotone Galois connection. Exercise 2.7.3.9 implies that for each $A \subset X$, $f(\sup A) = \sup f(A)$. Hence f is arbitrary join preserving.

2. Let $g : Y \to X$. the prev part, the prev ex and an ex in galois connection poset subsection imply that
   $$\begin{aligned} g \text{ is } (\leq_Y, \leq_X)\text{-arbitrary meet preserving} &\iff g \text{ is } (\leq_Y^*, \leq_X^*)\text{-arbitrary join preserving} \\ &\iff g \text{ has a } (\leq_Y^*, \leq_X^*)\text{-right adjoint} \\ &\iff \text{there exists } g_* : X \to Y \text{ such that} \\ &\qquad (g, g_*) \text{ is a } (\leq_Y^*, \leq_X^*)\text{-monotone Galois connection} \\ &\iff \text{there exists } g_* : X \to Y \text{ such that} \\ &\qquad (g_*, g) \text{ is a } (\leq_X, \leq_Y)\text{-monotone Galois connection} \\ &\iff g \text{ has a } (\leq_X, \leq_Y)\text{-left adjoint.} \end{aligned}$$

   Since $g : Y \to X$ is arbitrary, we have that for each $g : Y \to X$, g is $(\leq_Y, \leq_X)$-arbitrary meet preserving iff there exists $g_* : Y \to X$ such that g_* is a $(\leq_X, \leq_Y)$-left adjoint of g.

□

Exercise 10.2.0.4. Let $(X, \leq_X), (Y, \leq_Y)$ be complete lattices, $f : X \to Y$ and $g : Y \to X$. Suppose that (f, g) is a monotone Galois connection.

1. (a)
   (b)

2. (a) $g(1_Y) = 1_X$ iff

(b)

Exercise 10.2.0.5. show that if g is a right adjoint, then $g(\inf \uparrow x) = \inf g(\uparrow x) = \inf \uparrow(g(x)) = g(x)$

Exercise 10.2.0.6.

try to capture idea of a forgetful functor in terms of posets. If we go from groups to sets, we 'lose information' a group with a relations (i.e. a quotient group) is somehow 'more informative' than the free group. So if I send a set to a group in the least informative way (i.e. in a way which preserves as much entropy as possible) then we should get the free group, anything else would be more informative, with a loss of entropy. So quotient objects of a free object can maybe be characterized by some loss of entropy.

For example in **Set**, $A \subset B$ (i.e A is a subobject of B) is equivalent to saying that A is a quotient of B in the opposite category, which is equivalent to the complete atomic boolean algebras, or the power set, so that if $\iota : A \to B$ is a monomorphism in **Set**, then $\iota^{-1} : \mathcal{P}(B) \to \mathcal{P}(A)$ is an epimorphism in complete atomic boolean lattices and we think of $\mathcal{P}(A)$ as a quotient of $\mathcal{P}(B)$. So in this sense $\mathcal{P}(A)$ has less information or is less informative that $\mathcal{P}(B)$ (i.e. A is less informative than B in $\mathbf{Set}^{\mathrm{op}}$.)

If (X, x) is a set X with some structure x, and (A, a) is in the same category $\mathcal{C}$ (which we order by information level) and we can think of (A, a) as a quotient of (X, x), so it is less informative as a structured object $(A, a) \leq_{\leq_{\mathrm{info}\mathcal{C}}} (X, x)$, then if we change the context (apply a covariant functor) to land in Set^{op} and think of A quotient of X, (i.e. A is subset of X in **Set**), so that $A \leq_{\mathrm{info}\mathbf{Set}^{\mathrm{op}}} B$. Then we would like an adjoint of F in the sense of a poset so that we obtain the other half a monotone galois connection $G : \mathbf{Set}^{\mathrm{op}} \to \mathcal{C}$. The idea here is that the order $\leq_{\mathrm{info}\mathbf{Set}^{\mathrm{op}}}$ on Set^{op} is a part of $\leq_{\mathrm{info}\mathcal{C}}$, that is $\leq_{\mathrm{info}\mathcal{C}}$ is $\leq_{\mathrm{info}\mathbf{Set}^{\mathrm{op}}}$ plus some extra order information

i.e. a restriction in some sense so $(A, a) \leq_{\mathcal{C}} (B, b)$ (i.e. there is a quotient from B to A, like free group to abelianization) implies $A \leq_{Set^{\mathrm{op}}} B$. So then that $\leq_{\mathcal{C}} \subset \leq_{Set^{\mathrm{op}}}$ (in some sense), so that $\mathrm{ub}((A, a), \leq_{\mathcal{C}}) \subset \mathrm{ub}(A, \leq_{Set^{\mathrm{op}}})$ in some sense as sets, so that $\inf(\mathrm{ub}(A, \leq_{Set^{\mathrm{op}}}) \leq_{Set^{\mathrm{op}}} \inf(\mathrm{ub}((A, a), \leq_{\mathcal{C}}))) = A$. Then if we have a right adjoint $G : (\mathbf{Set}^{\mathrm{op}}, \leq_{\mathbf{Set}^{\mathrm{op}}}) \to (\mathcal{C}, \leq_{\mathcal{C}})$, which preserves meets and is monotone, we would have that (considering $\leq_{\mathcal{C}}$ as a subset of $\leq_{Set^{\mathrm{op}}}$)

$$\begin{aligned} G(A) &= G(\inf \mathrm{ub}(A, \leq_{\mathcal{C}}), \leq_{\mathcal{C}}) \\ &= \inf(G(\mathrm{ub}(A, \leq_{\mathcal{C}}), \leq_{\mathcal{C}})) \\ &\geq \inf(G(\mathrm{ub}(A, \leq_{\mathcal{C}})), \leq_{\mathcal{C}}) \end{aligned}$$

idk, I think we can show this aligns with the (least informative object (B, b) in $\mathcal{C}$ which is "above" A, i.e. the structured object (B, b) generated by A), i.e. the structured object "above" A with the least entropy gained from the minimum or zero structured element in the poset.

Try to show that a disintegration of measure, i.e. a conditional probability is the kind of the opposite or dual of this, i.e. it is the highest structured object (markov kernel + data) sitting below an unstructured object (markov kernel of a measure), so that going from (kernel + data) to disintegration results in the least entropy lost, maybe this is related to kl divergence as a measure of entropy loss, see baez/leinster/fritz paper.

In the case of statistics and say maximum likeliood estimation, and kl divergence as loss of estimate $\hat{\theta}(x_j : j \in \mathbb{N}) : (X, P_\theta) \to (X, P_{\hat{\theta}})$, there shold be a categorical way of rephrasing so that the observed data $(x_j)_{j \in \mathbb{N}} \subset X$ could perhaps be viewed as maps $x_j : (\Lambda, \mu) \to (X, P_\theta)$, so that $\hat{\theta} : \left[\prod_{j \in \mathbb{N}} \mathrm{Hom}((\Lambda, \mu), (X, P_\theta))\right] \times (X, P_\theta) \to (X, P_{\hat{\theta}})$

We should be able to do the same in a different context, say the true group $(G, \cdot)$ is unknown, but we observe the elements of G and the fact that for each $a, b \in G$, $a \cdot b = b \cdot a$ (i.e. we are given the proposition that G is commutative), then what would be the best estimate for G under this information in the sense of minimizing info loss (i.e. maybe occams razor?)? Well it would probably be the free abelian group. Any further processed abelian group, (i.e. a quotient of the free abelian group) would lose more info, so even though the trivial group also satisfies the observation, it is suboptimal because we lose basically all info about G. Need a way to quantify info loss in this way. What would the analogouse 'data' be? here it seems we need a way to view relations about elements as a morphism into G from some other object, like in the case of statistics.

Vague idea: in two different contexts, say two different posets, we want to relate two objects in the ordering induced by $a \leq b$ iff b is a quotient of a, i.e. b is more informative, i.e. there is an arrow from a to b which exhibits entropy loss.

talk about the idea of the element in a complete lattice with more structure generated another element in a complete lattice with less structure. i.e. subgroup generated subset where subgroups orderd by inclusion, power set ordered by inclusion. Or topology generated by collection of subsets (i.e. subset of powerset) where topologies ordered by inclusion and double power set ordered by

inclusion. In these cases, both posets are complete lattices since arbitrary meets exist. If we consider the map g from the more structured space P2+ to the less structured space P1-, by sending an object to itself minus the structure, so a group to its underlying set, or a topology to itself considered as a subset of the power set, then meets are preserved. Thus g is a right adjoint to some left adjoint f from the less structured space P1- to the more structured space P2, so f preserves arbitrary joins. For an object1 in the less structured poset, we define the object in the more structured poset generated by obj1 to be the left adjoint of obj1, i.e. f(obj1). we can find this in many cases by noting that in P1-, obj1 is the sup of its upper bounds. Since sups are preserved by f, the generated structured object in P2+ is f(obj1) which (from a prev ex) is the sup of f(of the lower bounds of obj1) i.e. the sup of $f(\downarrow o1)$, which should be the sup of all upper bounds of f(obj1), which is the inf

ok in P_1 is less structured, P_2 is more structured, $f : P_1 \to P_2$ preserves joins, adjoint g preserves meets, then the object in P_2 generated by $a \in P_1$ is $f(a) = f(\sup \downarrow a) = \sup f(\downarrow a)$, or something like this

10.3 Properties of Ideals and Filters

10.3.1 Prime Ideals and Filters in Complete Lattices

Definition 10.3.1.1. Let L be a complete lattice.

- Let $I \in \mathcal{I}(L)$. Then I is said to be
  - **completely prime** if
    1. I is proper
    2. for each $A \subset L$, $\bigwedge A \in I$ implies that there exists $a \in A$ such that $a \in I$.
- Let $F \in \mathcal{F}(L)$. Then F is said to be
  - **completely prime** if
    1. F is proper
    2. for each $A \subset L$, $\bigvee A \in F$ implies that there exists $a \in A$ such that $a \in F$.

Exercise 10.3.1.2. Let L be a complete lattice.

1. For each $I \in \mathcal{I}(L)$, if I is completely prime, then $I \in \mathcal{I}_p(L)$.
2. For each $F \in \mathcal{F}(L)$, if F is completely prime, then $F \in \mathcal{F}_p(L)$.

Proof. Clear. fill out □

Exercise 10.3.1.3. Let L be a complete lattice.

1. For each $I \in \mathcal{I}_{cp}(L)$,
   (a) $\bigwedge I^c \in I^c$,
   (b) $\bigwedge I^c \in \text{prime}_\vee(L)$.
2. For each $F \in \mathcal{F}_{cp}(L)$,
   (a) $\bigvee F^c \in F^c$,
   (b) $\bigvee F^c \in \text{prime}_\wedge(L)$.

Proof.

1. Let $I \in \mathcal{I}_{cp}(L)$.
   (a) For the sake of contradiction, suppose that $\bigwedge I^c \in I$. Since I is completely prime, there exists $a \in I^c$ such that $a \in I$. This is a contradiction. Hence $\bigwedge I^c \notin I$. Thus $\bigwedge I^c \in I^c$.
   (b) i. Since $I \in \mathcal{I}(L)$, a previous ex implies that $0 \in I$. Since $\bigwedge I^c \in I^c$, we have that $\bigwedge I^c \neq 0$.

ii. Define $x \in L$ by $x := \bigwedge I^c$. Since $I \in \mathcal{I}(L)$, a prev ex implies that I is a lower set. Let $a, b \in L$. Suppose that $x \leq a \vee b$. For the sake of contradiction, suppose that $a \in I$ and $b \in I$. Since $I \in \mathcal{I}(L)$, we have that $a \vee b \in I$. Since I is a lower set and

$$\begin{aligned} x &\leq a \vee b \\ &\in I, \end{aligned}$$

we have that $x \in I$. This is a contradiction since $x \in I^c$. Hence $a \notin I$ or $b \notin I$. Thus $a \in I^c$ or $b \in I^c$. Since $x = \bigwedge I^c$, we have that $x \leq a$ or $x \leq b$. Since $a, b \in L$ with $x \leq a \vee b$ are arbitrary, we have that for each $a, b \in L$, $x \leq a \vee b$ implies that $x \leq a$ or $b \leq b$.

Hence $x \in \text{prime}_\vee(L)$.

2. use duality

□

Exercise 10.3.1.4. Let L be a complete lattice. Then

1. for each $x \in \text{prime}_\vee(L)$, $(\uparrow x)^c \in \mathcal{I}_{cp}(L)$.
2. for each $x \in \text{prime}_\wedge(L)$, $(\downarrow x)^c \in \mathcal{F}_{cp}(L)$.

Proof.

1. Let $x \in \text{prime}_\vee(L)$. a prev ex implies that $\uparrow x \in \mathcal{F}(L)$.
   (a) Since $x \in \text{prime}_\vee(L)$, $x \neq 0$. Hence $0 \in (\uparrow x)^c$. Thus $(\uparrow x)^c \neq \varnothing$.
   (b) Let $a, b \in (\uparrow x)^c$. For the sake of contradiction, suppose that $a \vee b \in \uparrow x$. Then $x \leq a \vee b$. Since $x \in \text{prime}_\vee(L)$, $x \leq a$ or $x \leq b$. Hence $a \in \uparrow x$ or $b \in \uparrow x$. This is a contraction since $a \in (\uparrow x)^c$ and $b \in (\uparrow x)^c$. Therefore $a \vee b \in (\uparrow x)^c$. Since $a, b \in (\uparrow x)^c$ are arbitrary, we have that for each $a, b \in (\uparrow x)^c$, $a \vee b \in (\uparrow x)^c$.
   (c) Let $y \in L$ and $a \in (\uparrow x)^c$. For the sake of contradiction, suppose that $y \wedge a \in \uparrow x$. Then

   $$\begin{aligned} x &\leq y \wedge a \\ &\leq a. \end{aligned}$$

   Thus $a \in \uparrow x$, which is a contradiction. Hence $y \wedge a \in (\uparrow x)^c$. Since $y \in L$ and $a \in (\uparrow x)^c$ are arbitrary, we have that for each $y \in L$ and $a \in (\uparrow x)^c$, $y \wedge a \in (\uparrow x)^c$.

   Thus $(\uparrow x)^c \in \mathcal{I}(L)$.
   (a) Since $\uparrow x \neq \varnothing$, we have that $(\uparrow x)^c \neq L$. Thus $(\uparrow x)^c$ is proper.
   (b) Let $A \subset L$. Suppose that $\bigwedge A \in (\uparrow x)^c$. For the sake of contradiction, suppose that for each $a \in A$, $a \notin (\uparrow x)^c$. Then for each $a \in A$, $a \in \uparrow x$. Hence for each $a \in A$, $x \leq a$. Therefore $x \in \text{lb}\, A$. Thus

   $$\begin{aligned} x &\leq \inf A \\ &= \bigwedge A. \end{aligned}$$

   Then $\bigwedge A \in \uparrow x$, which is a contradiction. Hence there exists $a \in A$ such that $a \in (\uparrow x)^c$. Since $A \subset L$ with $\bigwedge A \in (\uparrow x)^c$ is arbitrary, we have that for each $A \subset L$, $\bigwedge A \in (\uparrow x)^c$ implies that there exists $a \in A$ such that $a \in (\uparrow x)^c$.

   Thus $(\uparrow x)^c \in \mathcal{I}_{cp}(L)$.
2. similar or duality

□

maybe make ex in earlier section on ideals that $\downarrow x = \downarrow y$ iff $x = y$ and similar for $\uparrow x$. Then in this section, show that for complete lattice L, $\mathcal{F}_{cp}(L) \cong \text{prime}_\wedge(L)$ bijective.

10.4 Constructions with Complete Lattices

10.4.1 Complete Sublattices

Definition 10.4.1.1. define **complete sublattice** Let L, M be lattices

10.4.2 Products of Complete Lattices

10.4.3 Coproducts of Complete Lattices

10.4.4 Quotients of Complete Lattices

10.5 Fundamental Examples

10.5.1 Introduction

Exercise 10.5.1.1. Let L be a complete lattice.

10.5.2 Sets

10.5.3 Equivalence Relations

Proof. Let X be a set. Suppose that $X \neq \varnothing$. Then $(\text{Equiv}, \subset)$ is a complete lattice. □

10.5.4 Ideal and Filter Lattices

Exercise 10.5.4.1. Let L be a lattice. If L has a zero, then

1. $(\mathcal{I}(L), \leq_{\mathcal{I}(L)})$ is a complete lattice.
2. $(\mathcal{F}(L), \leq_{\mathcal{F}(L)})$ is a complete lattice.

10.5.5 Topological Spaces

10.6 Least Upper Bound Property

Definition 10.6.0.1. Let $(X, \leq)$ be a poset. Then $(X, \leq)$ is said to satisfy the

- **least upper bound (LUB) property** if for each $A \subset X$, if $A \neq \varnothing$ and A is bounded above, then there exists $x \in X$ such that $x = \sup A$
- **greatest lower bound (GLB) property** if for each $A \subset X$, if $A \neq \varnothing$ and A is bounded below, then there exists $x \in X$ such that $x = \inf A$.

Exercise 10.6.0.2. LUB iff GLB

Proof. FINISH!!!! □

Definition 10.6.0.3. Suplattice:
Let $(L, \leq)$ be a poset. Then $(L, \leq)$ is said to be a **suplattice** if for each $A \subset L$, there exists $x \in L$ such that $x = \sup A$.

10.6.1 Complete Lattice Homomorphisms

Chapter 11

Some Chapter

11.1 Closure Operators

Definition 11.1.0.1. Let A be a set and $C : \mathcal{P}(A) \to \mathcal{P}(A)$. Then C is said to be a **closure operator on** A if for each $X, Y \in \mathcal{P}(A)$,

1. $X \subset C(X)$,
2. $C^2(X) = C(X)$,
3. $X \subset Y$ implies that $C(X) \subset C(Y)$.

Exercise 11.1.0.2. Let A be a set and $C : \mathcal{P}(A) \to \mathcal{P}(A)$. Suppose that C is a closure operator on A. Then for each $(E_j)_{j \in J} \subset \mathcal{P}(A)$,

1. $C\left(\bigcap_{j \in J} E_j\right) \subset \bigcap_{k \in J} C(E_k)$,
2. $\bigcup_{k \in J} C(E_k) \subset C\left(\bigcup_{j \in J} E_j\right)$.

Proof. Let $(E_j)_{j \in J} \subset \mathcal{P}(A)$.

1. Let $k \in J$. Then $\bigcap_{j \in J} E_j \subset E_k$. So $C\left(\bigcap_{j \in J} E_j\right) \subset C(E_k)$. Since $k \in J$ is arbitrary, we have that
$$C\left(\bigcap_{j \in J} E_j\right) \subset \bigcap_{k \in J} C(E_k).$$
2. Let $k \in J$. Then $E_k \subset \bigcup_{j \in J} E_j$. Hence $C(E_k) \subset C\left(\bigcup_{j \in J} E_j\right)$. Since $k \in J$ is arbitrary, we have that
$$\bigcup_{k \in J} C(E_k) \subset C\left(\bigcup_{j \in J} E_j\right)$$

□

Definition 11.1.0.3. Let A be a set, $C : \mathcal{P}(A) \to \mathcal{P}(A)$ and $X \subset A$. Suppose that C is a closure operator on A. Then X is said to be C-closed if $C(X) = X$.

Definition 11.1.0.4. Let A be a set and $C : \mathcal{P}(A) \to \mathcal{P}(A)$. Suppose that C is a closure operator on A. We define the **lattice of C-closed subsets of** A, denoted $L_C(A) \subset \mathcal{P}(A)$, by
$$L_C(A) := \{X \subset A : X \text{ is } C\text{-closed}\}$$
.

Exercise 11.1.0.5. Let A be a set and $C : \mathcal{P}(A) \to \mathcal{P}(A)$. Suppose that C is a closure operator on A. Then

1. for each $(E_j)_{j\in J} \subset L_C(A)$, $\bigcap_{j\in J} E_j \in L_C(A)$ and $\bigcup_{j\in J} E_j \in L_C(A)$.
2. $(L_C(A), \subset)$ is a complete lattice define complete lattice

$$C\left(\bigcap_{j\in J} E_j\right) = \bigcap_{j\in J} E_j$$

and

$$C\left(\bigcup_{j\in J} E_j\right) = \bigcup_{j\in J} E_j.$$

Proof.

1. Let $(E_j)_{j\in J} \subset L_C(A)$.
   - A previous exercise Exercise B.0.0.3 implies that

$$\begin{aligned} C\left(\bigcap_{j\in J} E_j\right) &\subset \bigcap_{k\in J} C(E_k) \\ &= \bigcap_{k\in J} E_k \\ &\subset C\left(\bigcap_{k\in J} E_k\right). \end{aligned}$$

   Hence $C\left(\bigcap_{j\in J} E_j\right) = \bigcap_{k\in J} E_k$.
   - A previous exercise Exercise B.0.0.3 implies that

$$\begin{aligned} \bigcup_{k\in J} E_k &= \bigcup_{k\in J} C(E_k) \\ &\subset C\left(\bigcup_{j\in J} E_j\right) \\ &\subset \bigcap_{k\in J} C(E_k) \\ &= \bigcap_{k\in J} E_k \\ &\subset C\left(\bigcap_{k\in J} E_k\right). \end{aligned}$$

   Hence $C\left(\bigcap_{j\in J} E_j\right) = \bigcap_{k\in J} E_k$.
2.

FINISH!!!, don't need to show second part, □

Definition 11.1.0.6. then is said to be an **algebraic closure operator on** A if

Chapter 12

Universal Algebra

12.1 Introduction

Definition 12.1.0.1. Let $A, J \in \mathrm{Obj}(\mathbf{Set})$ be a set and $f \in \mathcal{F}^*(A)^J$. Then (A, f) is said to be an **algebra** if $A \neq \varnothing$ and $J \neq \varnothing$.

Definition 12.1.0.2. Let $\mathcal{A}$ be an algebra. Write $\mathcal{A} = (A, f)$. Set $J := \operatorname{dom} f$.

- We define the **universe of** $\mathcal{A}$, denoted $\operatorname{Uni}\mathcal{A}$, by $\operatorname{Uni}\mathcal{A} := A$.
- We define the **operations of** $\mathcal{A}$, denoted $\operatorname{Oper}\mathcal{A}$, by $\operatorname{Oper}\mathcal{A} := f$.
- We define the **type of** $\mathcal{A}$, denoted $\operatorname{Type}\mathcal{A} : J \to \mathbb{N}_0$, by $\operatorname{Type}\mathcal{A}(j) := \operatorname{arity} f_j$.

Definition 12.1.0.3. Let $\mathcal{A}$, $\mathcal{B}$ be algebras. Then $\mathcal{A}$ and $\mathcal{B}$ are said to be **type-similar**, denoted $\mathcal{A} \sim_{\mathrm{Type}} \mathcal{B}$, if $\operatorname{Type}\mathcal{A} = \operatorname{Type}\mathcal{B}$.

Definition 12.1.0.4. Let $(A, f), (B, g)$ be algebras and $\alpha : A \to B$. Suppose that $(A, f) \sim_{\mathrm{Type}} (B, g)$. Set $J := \operatorname{dom} f$ and $\rho := \operatorname{Type}(A, f)$. Write $f = (f_j)_{j \in J}$ and $g = (g_j)_{j \in J}$. Then α is said to be an $((A, f), (B, g))$**-homomorphism** if for each $j \in J$,

$$\alpha \circ f_j = g_j \circ \alpha^{\rho(j)}.$$

Exercise 12.1.0.5. Let $(A, f), (B, g), (C, h)$ be algebras and $\alpha : A \to B$, $\beta : B \to C$. Suppose that $(A, f) \sim_{\mathrm{Type}} (B, g), (C, h)$. Set $J := \operatorname{dom} f$ and $\rho := \operatorname{Type}(A, f)$. Write $f = (f_j)_{j \in J}$, $g = (g_j)_{j \in J}$ and $h = (h_j)_{j \in J}$. If α is a $((A, f), (B, g))$-homomorphism and β is a $((B, g), (C, h))$-homomorphism, then $\beta \circ \alpha$ is a $((A, f), (C, h))$-homomorphism.

Proof. Suppose that α is a $((A, f), (B, g))$-homomorphism and β is a $((B, g), (C, h))$-homomorphism. Let $j \in J$. Since α is a $((A, f), (B, g))$-homomorphism, $\alpha \circ f_j = g_j \circ \alpha^{\rho(j)}$. Since β is a $((B, g), (C, h))$-homomorphism, $\beta \circ g_j = h_j \circ \beta^{\rho(j)}$. Therefore

$$\begin{aligned}
(\beta \circ \alpha) \circ f_j &= \beta \circ (\alpha \circ f_j) \\
&= \beta \circ (g_j \circ \alpha^{\rho(j)}) \\
&= (\beta \circ g_j) \circ \alpha^{\rho(j)} \\
&= (h_j \circ \beta^{\rho(j)}) \circ \alpha^{\rho(j)} \\
&= h_j \circ (\beta^{\rho(j)} \circ \alpha^{\rho(j)}) \\
&= h_j \circ ([\beta \circ \alpha]^{\rho(j)}).
\end{aligned}$$

Since $j \in J$ is arbitrary, we have that for each $j \in J$ and $(\beta \circ \alpha) \circ f_j = h_j \circ ([\beta \circ \alpha]^{\rho(j)})$. Hence $\beta \circ \alpha$ is a $((A, f), (C, h))$-homomorphism. □

Exercise 12.1.0.6. Let (A, f) be and algebra. Set $J := \operatorname{dom} f$. Write $f = (f_j)_{j \in J}$. Then id_A is a $((A, f), (A, f))$-homomorphism.

Proof. Since for each $j \in J$,

$$\begin{aligned}
\mathrm{id}_A \circ f_j &= f_j \\
&= f_j \circ \mathrm{id}_{A^{\rho(j)}} \\
&= f_j \circ \mathrm{id}_A^{\rho(j)},
\end{aligned}$$

we have that id_A is a $((A, f), (A, f))$-homomorphism. □

Definition 12.1.0.7.

- Let J be a set and $\rho : J \to \mathbb{N}_0$. Then ρ is said to be a **type function**. We define Types $:= \{\rho : \rho \text{ is a type function}\}$.
- Let $\rho \in \text{Types}$. We define the **category of algebras with type** ρ, denoted $\mathbf{Alg}_\rho$, by
  - $\text{Obj}(\mathbf{Alg}_\rho) := \{\mathcal{A} : \mathcal{A} \text{ is an algebra and } \text{Type}\,\mathcal{A} = \rho\}$
  - For $(A, f), (B, g) \in \text{Obj}(\mathbf{Alg}_\rho)$, $\text{Hom}_{\mathbf{Alg}_\rho}(\mathcal{A}, \mathcal{B}) := \{\alpha \in B^A : \alpha \text{ is a } (\mathcal{A}, \mathcal{B})\text{-homomorphism.}\}$

Exercise 12.1.0.8. Let $\rho \in \text{Types}$. Then $\mathbf{Alg}_\rho$ is a category.

Proof. FINISH!!! □

12.2 Subalgebras

Definition 12.2.0.1. Let $\rho \in \text{Types}$ and $\mathcal{A}, \mathcal{B} \in \text{Obj}(\mathbf{Alg}_\rho)$. Write $\mathcal{A} = (A, f)$ and $\mathcal{B} = (B, g)$. Set $J := \operatorname{dom} \rho$. Write $f = (f_j)_{j \in J}$ and $g = (g_j)_{j \in J}$.

- Then $\mathcal{B}$ is said to be a **subalgebra of** $\mathcal{A}$ if
  1. $B \subset A$
  2. $\iota_B \in \text{Hom}_{\mathbf{Alg}_\rho}(\mathcal{B}, \mathcal{A})$.
- We define $\text{SubAlg}(\mathcal{A}) := \{\mathcal{B} \in \text{Obj}(\mathbf{Alg}_\rho) : \mathcal{A} \sim_{\text{Type}} \mathcal{B} \text{ and } \mathcal{B} \text{ is a subalgebra of } \mathcal{A}\}$.

Exercise 12.2.0.2. Let $\rho \in \text{Types}$ and $\mathcal{A}, \mathcal{B} \in \text{Obj}(\mathbf{Alg}_\rho)$. Write $\mathcal{A} = (A, f)$ and $\mathcal{B} = (B, g)$. Set $J := \operatorname{dom} \rho$. Write $f = (f_j)_{j \in J}$ and $g = (g_j)_{j \in J}$. Then $(B, g) \in \text{SubAlg}(A, f)$ is a subalgebra of (A, f) iff

1. $B \subset A$
2. for each $j \in J$, $f_j|_{B^{\rho(j)}} = g_j$.

Proof.

- $(\implies)$:
  Suppose that $\mathcal{B}$ is a subalgebra of $\mathcal{A}$.
  1. By definition, $B \subset A$.
  2. Let $j \in J$. Since $\mathcal{B}$ is a subalgebra of $\mathcal{A}$, $\iota_B \in \text{Hom}_{\mathbf{Alg}_\rho}(\mathcal{B}, \mathcal{A})$. Then
  $$\begin{aligned} f_j|_{B^{\rho(j)}} &= f_j \circ \iota_{B^{\rho(j)}} \\ &= f_j \circ \iota_B^{\rho(j)} \\ &= \iota_B \circ g_j \\ &= g_j. \end{aligned}$$
  Since $j \in J$ is arbitrary, we have that for each $j \in J$, $f_j|_{B^{\rho(j)}} = g_j$.
- $(\impliedby)$:
  Suppose that
  1. $B \subset A$
  2. for each $j \in J$, $f_j|_{B^{\rho(j)}} = g_j$.

  Let $j \in J$. Then
  $$\begin{aligned} \iota_B \circ g_j &= g_j \\ &= f_j|_{B^{\rho(j)}} \\ &= f_j \circ \iota_{B^{\rho(j)}} \\ &= f_j \circ \iota_B^{\rho(j)}. \end{aligned}$$
  Since $j \in J$ is arbitary, we have that for each $j \in J$, $\iota_B \circ g_j = f_j \circ \iota_B^{\rho(j)}$. Thus $\iota_B \in \text{Hom}_{\mathbf{Alg}_\rho}(\mathcal{B}, \mathcal{A})$.
  1. $B \subset A$
  2. Let $j \in J$ and $a \in B^{\rho(j)}$. Then
  $$\begin{aligned} f_j(\iota_B^{\rho(j)}(a)) &= f_j(\iota_{B^{\rho(j)}}(a)) \\ &= f_j \circ \iota_{B^{\rho(j)}}(a) \\ &= f_j|_{B^{\rho(j)}}(a) \\ &= g_j(a) \\ &= \iota_B(g_j(a)). \end{aligned}$$

□

Definition 12.2.0.3. Let $\rho \in \text{Types}$ and $(A, f), (B, g) \in \text{Obj}(\mathbf{Alg}_\rho)$. Set $J := \operatorname{dom} \rho$. Write $f = (f_j)_{j \in J}$ and $g = (g_j)_{j \in J}$.

- Then (B, g) is said to be a **subalgebra of** (A, f) if
  1. $B \subset A$
  2. for each $j \in J$, $f_j|_{B^{\rho(j)}} = g_j$.
- We define $\text{SubAlg}(\mathcal{A}) := \{\mathcal{B} \in \text{Obj}(\mathbf{Alg}_\rho) : \mathcal{A} \sim_{\text{Type}} \mathcal{B} \text{ and } \mathcal{B} \text{ is a subalgebra of } \mathcal{A}\}$.

Definition 12.2.0.4. Let $\rho \in \text{Types}$, $(A, f) \in \text{Obj}(\mathbf{Alg}_\rho)$ and $B \in \text{Obj}(\mathbf{Set})$.

- Then B is said to be a **subuniverse of** (A, f) if
  1. $B \subset A$,
  2. B is $\operatorname{Im} f$-closed.
- We define $\text{SubUni}(\mathcal{A}) := \{B \in \text{Obj}(\mathbf{Set}) : B \text{ is a subuniverse of } \mathcal{A}\}$.

Exercise 12.2.0.5. Let $\rho \in \text{Types}$ and $(A, f), (B, g) \in \text{Obj}(\mathbf{Alg}_\rho)$. If (B, g) is a subalgebra of (A, f), then B is a subuniverse of (A, f).

Proof. Set $J := \operatorname{dom} \rho$ and $\rho := \text{Type}(A, f)$. Suppose that (B, g) is a subalgebra of (A, f).

1. Since (B, g) is a subalgebra of (A, f), we have that $B \subset A$.
2. Let $j \in J$. Then for each $a \in B^\rho(j)$,
$$\begin{aligned} f_j(a_1, \ldots, a_{\rho(j)}) &= f_j|_{B^{\rho(j)}}(a_1, \ldots, a_{\rho(j)}) \\ &= g_j(a_1, \ldots, a_{\rho(j)}) \\ &\in B. \end{aligned}$$
Since $j \in J$ is arbitrary, we have that B is $\operatorname{Im} f$-closed. Thus B is a subuniverse of A.

□

Definition 12.2.0.6. Let $\mathcal{A}$ be an algebra and B a subuniverse of $\mathcal{A}$. Set $\mathcal{S}(B, \mathcal{A}) := \{S \subset A : S \text{ is a subuniverse of } \mathcal{A} \text{ and } B \subset S\}$. We define the **subuniverse of $\mathcal{A}$ generated by** B, denoted $\text{Sg}(B, \mathcal{A})$, by
$$\text{Sg}(B, \mathcal{A}) := \bigcap_{S \in \mathcal{S}(B, \mathcal{A})} S$$

show $\mathcal{S} \neq \varnothing$ and intersection of subiniverses is subuniverse

Exercise 12.2.0.7. Let (A, f) be an algebra and $B \subset A$. Then

1. $\text{Sg}(B, f)$ is a subuniverse of A
2. $B \subset \text{Sg}(B, f)$.

.

Proof.

1. Set $\mathcal{S} := \{S \subset A : S \text{ is an } f\text{-subuniverse of } A\}$. By construction, for each $S \in S$, S is f-closed. Since $\text{Sg}(B, f) = \bigcap_{S \in \mathcal{S}} S$, Exercise B.0.0.3 A previous exercise in the set theory section implies that $\text{Sg}(B, f)$ is f-closed. Hence $\text{Sg}(B, f)$ is an f-subuniverse of A.
2. By construction, for each $S \in S$, $B \subset S$. Thus
$$\begin{aligned} B &\subset \bigcap_{S \in \mathcal{S}} S \\ &= \text{Sg}(B, f). \end{aligned}$$

□

Exercise 12.2.0.8. Let (A, f) be an algebra. Then $\text{Sg}(\cdot, f)$ is an algebraic closure operator on A.

Proof. □

12.3 Congruences

Definition 12.3.0.1. Let $\rho \in \text{Types}$ and $(A, f) \in \text{Obj}(\mathbf{Alg}_\rho)$. Set $J := \text{dom}\,\rho$. Write $f = (f_j)_{j \in J}$.

- Let $\theta \in \text{Equiv}(A)$. Then θ is said to be a **congruence on** (A, f) if for each $j \in J$ and $a, b \in A^{\rho(j)}$, $a\theta^{\times\rho(j)}b$ implies that $f_j(a)\theta f_j(b)$.
- We define $\text{Cong}(A, f) := \{\theta \in \text{Equiv}(A) : \theta \text{ is a congruence on } (A, f)\}$.

Note 12.3.0.2. Question: θ is congruence iff $f_j^{\times 2}(\theta^{\times\rho(j)}) \subset \theta$, is there a generalization of equivalence relation to $\theta \subset A^n$ that is reflexive ($\Delta_{A,n} \subset \theta$) symmetric (invariant under S_n), and transitive (idk what the generalization of composition should be), then define congruence by

$$f_j^{\times n}(\theta^{\times\rho(j)}) \subset \theta$$

when $n = 3$, the question becomes what meaning could capture the statement that a set of three objects 'has sameness', but no two element subset of them 'has sameness' in the same way.

Exercise 12.3.0.3. Let $\rho \in \text{Types}$, $(A, f) \in \text{Obj}(\mathbf{Alg}_\rho)$ and $\theta \in \text{Cong}(A, f)$.

Proof. □

Definition 12.3.0.4. Let $\mathcal{A} \in \text{Obj}(\mathbf{Alg}_\rho)$ and $\theta \in \text{Cong}(\mathcal{A})$. Write $\mathcal{A} = (A, f)$, set $J := \text{dom}\,\rho$ and write $f = (f_j)_{j \in J}$. We define

- $f^\theta \in \mathcal{F}^*(A/\theta)^J$ by $f_j^\theta := \text{Factor}_{\theta^{\rho(j)},\theta}(f_j)$.
- **quotient algebra of $\mathcal{A}$ by** θ, denoted $\mathcal{A}/\theta$, by $\mathcal{A}/\theta := (A/\theta, f^\theta)$.

introduce factor notation here and apply it to analysis and other notes, aka if we fix $\pi_X : X \to X/\theta_X$, $\pi_Y : Y \to Y/\theta_Y$ and consider $f : X \to Y$ which is (θ_X, θ_Y)-invariant, then there exists a unique $\tilde{f} : X/\theta_X \to Y/\theta_Y$ such that $\pi_Y \circ f = \tilde{f} \circ \pi_{\theta_X}$, we define $\text{Factor}_{\theta_X,\theta_Y}(f) := \tilde{f}$.

I guess we can define a Factor map or something similar in other contexts, i.e. for other universal properties/limits/colimits since they all have some unique map involved as a factor of some expression related to the original map.

need to go over equiv relations in set theory notes

Exercise 12.3.0.5. Let $\rho \in \text{Type}$, $\mathcal{A} \in \text{Obj}(\mathbf{Alg}_\rho)$ and $\theta \in \text{Cong}(\mathcal{A})$. Then $\mathcal{A}/\theta \in \text{Obj}(\mathbf{Alg}_\rho)$.

Proof. Write $\mathcal{A} = (A, f)$. Set $J := \text{dom}\,\rho$. By assumption, $A, J \neq \varnothing$. Hence $(A/\theta, f^\theta)$ is an algebra. By definition, for each $j \in J$, $f_j \in \mathcal{F}^j(A)$ iff $f_j^\theta \in \mathcal{F}^j(A/\theta)$. Hence

$$\begin{aligned}\text{Type}(A/\theta, f^\theta) &= \text{Type}(A, f)\\ &= \rho.\end{aligned}$$

Thus $(A/\theta, f^\theta) \in \text{Obj}(\mathbf{Alg}_\rho)$. □

Appendix A

Summation

Definition A.0.0.1. Let $f : X \to [0, \infty)$, Then we define

$$\sum_{x \in X} f(x) := \sup_{\substack{F \subset X \\ F \text{ finite}}} \sum_{x \in F} f(x)$$

This definition coincides with the usual notion of summation when X is countable. For $f : X \to \mathbb{C}$, we can write $f = g + ih$ where $g, h : X \to \mathbb{R}$. If

$$\sum_{x \in X} |f(x)| < \infty,$$

then the same is true for g^+, g^-, h^+, h^-. In this case, we may define

$$\sum_{x \in X} f(x)$$

in the obvious way.

The following note justifies the notation $\sum_{x \in X} f(x)$ where $f : X \to \mathbb{C}$.

Note A.0.0.2. Let $f : X \to \mathbb{C}$ and $\alpha : X \to X$ a bijection. If $\sum_{x \in X} |f(x)| < \infty$, then $\sum_{x \in X} f(\alpha(x)) = \sum_{x \in X} f(x)$.

Appendix B

Asymptotic Notation

Definition B.0.0.1. Let X be a topological space, Y, Z be normed vector spaces, $f : X \to Y$, $g : X \to Z$ and $x_0 \in X \cup \{\infty\}$. Then we write

$$f = o(g) \qquad \text{as } x \to x_0$$

if for each $\epsilon > 0$, there exists $U \in \mathcal{N}(x_0)$ such that for each $x \in U$,

$$\|f(x)\| \leq \epsilon \|g(x)\|$$

Exercise B.0.0.2. Let X be a topological space, Y, Z be normed vector spaces, $f : X \to Y$, $g : X \to Z$ and $x_0 \in X \cup \{\infty\}$. If there exists $U \in \mathcal{N}(x_0)$ such that for each $x \in U \setminus \{x_0\}$, $g(x) > 0$, then

$$f = o(g) \text{ as } x \to x_0 \quad \text{iff} \quad \lim_{x \to x_0} \frac{\|f(x)\|}{\|g(x)\|} = 0$$

Exercise B.0.0.3. Let X and Y a be normed vector spaces, $A \subset X$ open and $f : A \to Y$. Suppose that $0 \in A$. If $f(h) = o(\|h\|)$ as $h \to 0$, then for each $h \in X$, $f(th) = o(|t|)$ as $t \to 0$.

Proof. Suppose that $f(h) = o(\|h\|)$ as $h \to 0$. Let $h \in X$ and $\epsilon > 0$. Choose $\delta' > 0$ such that for each $h' \in B(0, \delta')$, $h' \in A$ and

$$\|f(h')\| \leq \frac{\epsilon}{\|h\| + 1} \|h'\|$$

Choose $\delta > 0$ such that for each $t \in B(0, \delta)$, $th \in B(0, \delta')$. Let $t \in B(0, \delta)$. Then

$$\begin{aligned} \|f(th)\| &\leq \frac{\epsilon}{\|h\| + 1} |t| \|h\| \\ &< \epsilon |t| \end{aligned}$$

So $f(th) = o(|t|)$ as $t \to 0$. $\square$

Definition B.0.0.4. Let X be a topological space, Y, Z be normed vector spaces, $f : X \to Y$, $g : X \to Z$ and $x_0 \in X \cup \{\infty\}$. Then we write

$$f = O(g) \qquad \text{as } x \to x_0$$

if there exists $U \in \mathcal{N}(x_0)$ and $M \geq 0$ such that for each $x \in U$,

$$\|f(x)\| \leq M \|g(x)\|$$

Appendix C

Categories

move to notation?

Definition C.0.0.1. We define the category of topological measure spaces, denoted $\mathbf{TopMsr}_+$, by

- $\mathrm{Obj}(\mathbf{TopMsr}_+) := \{(X, \mu) : X \in \mathrm{Obj}(\mathbf{Top}) \text{ and } \mu \in M(X)\}$
- $\mathrm{Hom}_{\mathbf{TopMsr}_+}((X, \mu), (Y, \nu)) := \mathrm{Hom}_{\mathbf{Top}}(X, Y) \cap \mathrm{Hom}_{\mathbf{Msr}_+}((X, \mathcal{B}(X), \mu), (Y, \mathcal{B}(Y), \nu))$

Appendix D

Vector Spaces

it might be better to cover some category theory and write everything in terms of $\mathrm{Hom}_{\mathbf{Vect}_{\mathbb{K}}}$ and $\mathrm{Obj}(\mathbf{Vect}_{\mathbb{K}})$

D.1 Introduction

Definition D.1.0.1. Let X be a set, $\mathbb{K}$ a field, $+ : X \times X \to X$ and $\cdot : \mathbb{K} \times X \to X$. Then $(X, +, \cdot)$ is said to be a **$\mathbb{K}$-vector space** if

1. $(X, +)$ is an abelian group
2.

Definition D.1.0.2. Let $(X, +_X, \cdot_X)$ and $(E, +_E, \cdot_E)$ be vector spaces. Suppose that $E \subset X$. Then $(E, +_E, \cdot_E)$ is said to be a subspace of X if

1. $+_E = +_X|_{E \times E}$
2. $\cdot_E = \cdot_X|_{\mathbb{K} \times E}$

Exercise D.1.0.3. Let $(X, +_X, \cdot_X)$ and $(E, +_E, \cdot_E)$ be vector spaces. Suppose that $E \subset X$.

Exercise D.1.0.4. Let $(X, +, \cdot)$ be a vector space and $E \subset X$. Then E is a subspace of X

Definition D.1.0.5. Let X be a vector space and $(E_j)_{j \in J}$ a collection of subspaces of X. Then $\bigcap_{j \in J} E_j$ is a subspace of X.

Proof. Set $E := \bigcap_{j \in J} E_j$. Let $x, y \in E$ and $\lambda \in \mathbb{K}$. Then for each $j \in J$, $x, y \in E_j$. Since for each $j \in J$, E_j is a subspace of X, we have that for each $j \in J$, $x + \lambda y \in E_j$. Thus $x + \lambda y \in E$. Since $x, y \in E$ and $\lambda \in \mathbb{K}$ are arbitrary, (cite exercise here) we have that E is a subspace of X. □

Definition D.1.0.6. Let X, Y be vector spaces and $T : X \to Y$. Then T is said to be **linear** if for each $x_1, x_2 \in X$ and $\lambda \in \Lambda$,

1. $T(x_1 + x_2) = T(x_1) + T(x_2)$,
2. $T(\lambda x_1) = \lambda T(x_1)$.

We define $L(X; Y) := \{T : X \to Y : \ T \text{ is linear}\}$.

Exercise D.1.0.7. Let X, Y be vector spaces and $T : X \to Y$. Then T is linear iff for each $x_1, x_2 \in X$ and $\lambda \in \Lambda$,

$$T(x_1 + \lambda x_2) = T(x_1) + \lambda T(x_2)$$

Proof. Clear. (add details) □

Definition D.1.0.8. define addition/scalar multiplication of linear maps

Exercise D.1.0.9. Let X, Y be vector spaces. Then $L(X;Y)$ is a $\mathbb{K}$-vector space.

Proof. Clear □

Definition D.1.0.10. Let X be a vector space over $\mathbb{K}$ and $T : X \to \mathbb{K}$. Then T is said to be a **linear functional on** X if T is linear. We define the **dual space of** X, denoted X^*, by $X^* := \{T : X \to \mathbb{K} : T \text{ is linear}\}$.

Exercise D.1.0.11. Let X be a vector space. Then X^* is a vector space.

Proof. Clear. □

D.2 Bases

Definition D.2.0.1. Let X be a vector space and $(e_\alpha)_{\alpha \in A} \subset X$. Then $(e_\alpha)_{\alpha \in A}$ is said to be

- **linearly independent** if for each $(\alpha_j)_{j=1}^n \subset A$, $(\lambda_j)_{j=1}^n \subset \mathbb{K}$, $\sum_{j=1}^n \lambda_j e_{\alpha_j} = 0$ implies that for each $j \in [n]$, $\lambda_j = 0$.
- a **Hamel basis for** X if $(e_\alpha)_{\alpha \in A}$ is linearly independent and $\operatorname{span}(e_\alpha)_{\alpha \in A} = X$.

Exercise D.2.0.2. every vector space has a Hamel basis

Proof. □

Exercise D.2.0.3.

Exercise D.2.0.4. Let X be a $\mathbb{K}$-vector space and $x \in X$. Then $x = 0$ iff for each $\phi \in X^*$, $\phi(x) = 0$.

Proof.

- ($\Longrightarrow$):
  Suppose that $x = 0$. Linearity implies that for each $\phi \in X^*$ $\phi(x) = 0$.
- ($\Longleftarrow$):
  Conversely, suppose that $x \neq 0$. Define $\epsilon_x : \operatorname{span}(x) \to \mathbb{K}$ by $\epsilon_x(\lambda x) := \lambda$. Let $u, v \in \operatorname{span}(x)$. Then there exists $\lambda_u, \lambda_v \in \mathbb{K}$ such that $u = \lambda_u x$ and $v = \lambda_v x$. Suppose that $u = v$. Then

  $$\begin{aligned}(\lambda_u - \lambda_v)x &= \lambda_u x - \lambda_v x \\ &= u - v \\ &= 0\end{aligned}$$

  Since $x \neq 0$, we have that $\lambda_u - \lambda_v = 0$ and therefore $\lambda_u = \lambda_v$. Hence

  $$\begin{aligned}\lambda_u &= \epsilon_x(u) \\ &= \epsilon_x(v) \\ &= \lambda_v.\end{aligned}$$

  Thus ϵ_x is well defined.

□

D.3 Multilinear Maps

Definition D.3.0.1. Let $X_1, \cdots, X_n, Y$ be vector spaces and $T : \prod_{j=1}^{n} X_j \to \mathbb{K}$. Then T is said to be **multilinear** if for each $j_0 \in [n]$ and $(x_j)_{j=1}^n \in \prod_{j=1}^{n} X_j$, $T(x_1, \ldots, x_{j_0-1}, \cdot, x_{j_0+1})$ is linear.

$$L^n(X_1, \ldots, X_n; Y) = \left\{ T : \prod_{j=1}^{n} X_j \to Y : T \text{ is multilinear} \right\}$$

If $X_1 = \cdots = X_n = X$, we write $L^n(X; Y)$ in place of $L^n(X, \ldots, X; Y)$.

Definition D.3.0.2. define addition and scalar mult of multilinear maps

Exercise D.3.0.3. Let $X_1, \cdots, X_n, Y$ be vector spaces. Then $L^n(X_1, \ldots, X_n; Y)$ is a $\mathbb{K}$-vector space.

Proof. content... □

Exercise D.3.0.4. Let $X_1, \cdots, X_n, Y, Z$ be $\mathbb{K}$-vector spaces, $\alpha \in L^n(X_1, \ldots, X_n; Y)$ and $\phi \in L^1(Y; Z)$. Then $\phi \circ \alpha \in L^n(X_1, \ldots, X_n; Z)$.

Proof. Let $(x_j)_{j=1}^n \in \prod_{j=1}^{n} X_j$ and $j_0 \in [n]$. Define $f : X_{j_0} \to Y$ by

$$f(a) := \alpha(x_1, \ldots, x_{j_0-1}, a, x_{j_0+1}, \ldots, x_n)$$

Since $\alpha \in L^n(X_1, \ldots, X_n; Y)$, f is linear. Since ϕ is linear, and $\phi \circ f$ is linear. Since $(x_j)_{j=1}^n \in \prod_{j=1}^{n} X_j$ and $j_0 \in [n]$ are arbitrary, we have that $\phi \circ \alpha \in L^n(X_1, \ldots, X_n; Y)$. □

D.4 Tensor Products

Definition D.4.0.1. Let X, Y and T be vector spaces over $\mathbb{K}$ and $\alpha \in L^2(X, Y; T)$. Then (T, α) is said to be a **tensor product of X and Y** if for each vector space Z and $\beta \in L^2(X, Y; Z)$, there exists a unique $\phi \in L^1(T; Z)$ such that $\phi \circ \alpha = \beta$, i.e. the following diagram commutes:

$$\begin{tikzcd} X \times Y \arrow[r, "\alpha"] \arrow[dr, "\beta"'] & T \arrow[d, dashed, "\phi"] \\ & Z \end{tikzcd}$$

Exercise D.4.0.2. Let X, Y, S, T be vector spaces, $\alpha \in L^2(X, Y; S)$ and $\beta \in L^2(X, Y; T)$. Suppose that (S, α) and (T, β) are tensor products of X and Y. Then S and T are isomorphic.

Proof. Since (T, β) is a tensor product of X and Y, $\beta \in L^2(X, Y; T)$ there exists a unique $f \in L^1(T; T)$ such that $f\ circ\beta = \beta$, i.e. the following diagram commutes:

$$\begin{tikzcd} & T \arrow[dd, dashed, "f"] \\ X \times Y \arrow[ur, "\beta"] \arrow[dr, "\beta"'] & \\ & T \end{tikzcd}$$

Since $\mathrm{id}_T \in L^1(T; T)$ and $\mathrm{id}_T \circ \beta = \beta$, we have that $f = \mathrm{id}_T$. Since (S, α) is a tensor product of X and Y, there exists a unique $\phi : S \to T$ such that $\phi \circ \alpha = \beta$, i.e. the following diagram commutes:

$$\begin{tikzcd} X \times Y \arrow[r, "\alpha"] \arrow[dr, "\beta"'] & S \arrow[d, dashed, "\phi"] \\ & T \end{tikzcd}$$

Similarly, since (T, β) is a tensor product of X and Y, there exists a unique $\psi : T \to S$ such that $\psi \circ \beta = \alpha$, i.e. the following diagram commutes:

$$\begin{tikzcd} X \times Y \arrow[r, "\beta"] \arrow[dr, "\alpha"'] & T \arrow[d, dashed, "\psi"] \\ & S \end{tikzcd}$$

Therefore

$$\begin{aligned} (\phi \circ \psi) \circ \beta &= \phi \circ (\psi \circ \beta) \\ &= \phi \circ \alpha \\ &= \beta, \end{aligned}$$

i.e. the following diagram commutes:

$$\begin{tikzcd} & T \arrow[d, dashed, "\psi"] & \\ X \times Y \arrow[ur, "\beta"] \arrow[r, "\alpha"] \arrow[dr, "\beta"'] & S \arrow[d, dashed, "\phi"] & \Longrightarrow \\ & T & \end{tikzcd} \quad \begin{tikzcd} & T \arrow[dd, dashed, "\phi\circ\psi"] \\ X \times Y \arrow[ur, "\beta"] \arrow[dr, "\beta"'] & \\ & T \end{tikzcd}$$

By uniqueness of $f \in L^1(T; T)$, we have that

$$\begin{aligned} \mathrm{id}_T &= f \\ &= \phi \circ \psi \end{aligned}$$

A similar argument implies that $\psi \circ \phi = \mathrm{id}_S$. Hence ϕ and ψ are isomorphisms with $\phi^{-1} = \psi$. Hence S and T are isomorphic. $\square$

Definition D.4.0.3. Let X, Y be vector spaces, $x \in X$ and $y \in Y$. We define $x \otimes y : X^* \times Y^* \to \mathbb{K}$ by $x \otimes y(\phi, \psi) := \phi(x)\psi(y)$.

Exercise D.4.0.4. Let X, Y be vector spaces, $x \in X$ and $y \in Y$. Then $x \otimes y \in L^2(X^*, Y^*; \mathbb{K})$.

Proof. Let $\phi_1, \phi_2 \in X^*$, $\psi \in Y^*$ and $\lambda \in \mathbb{K}$. Then

$$\begin{aligned} x \otimes y(\phi_1 + \lambda\phi_2, \psi) &= [\phi_1 + \lambda\phi_2(x)]\psi(y) \\ &= \phi_1(x)\psi(y) + \lambda\phi_2(x)\psi(y) \\ &= x \otimes y(\phi_1, \psi) + \lambda x \otimes y(\phi_2, \psi) \end{aligned}$$

Since $\phi_1, \phi_2 \in X^*$, $\psi \in Y^*$ and $\lambda \in \mathbb{K}$ are arbitrary, we have that for each $\psi \in Y^*$, $x \otimes y(\cdot, \psi)$ is linear. Similarly for each $\phi \in X^*$, $x \otimes y(\phi, \cdot)$ is linear. Hence $x \otimes y$ is bilinear and $x \otimes y \in L^2(X^*, Y^*; \mathbb{K})$. □

Definition D.4.0.5. Let X, Y be vector spaces. We define

- the **tensor product of X and Y**, denoted $X \otimes Y \subset L^2(X^*, Y^*; \mathbb{K})$, by
$$X \otimes Y := \operatorname{span}(x \otimes y : x \in X \text{ and } y \in Y),$$
- the **tensor map**, denoted $\otimes : X \times Y \to X \otimes Y$, by $\otimes(x, y) := x \otimes y$.

Exercise D.4.0.6. Let X, Y be vector spaces, $(x_j)_{j=1}^n \subset X$ and $(y_j)_{j=1}^n \subset Y$. The following are equivalent:

1. $\sum_{j=1}^n x_j \otimes y_j = 0$
2. for each $\phi \in X^*$ and $\psi \in Y^*$, $\sum_{j=1}^n \phi(x_j)\psi(y_j) = 0$
3. for each $\phi \in X^*$, $\sum_{j=1}^n \phi(x_j)y_j = 0$
4. for each $\psi \in Y^*$, $\sum_{j=1}^n \psi(y_j)x_j = 0$

Proof.

1. $(1) \implies (2)$:
   Suppose that $\sum_{j=1}^n x_j \otimes y_j = 0$. Let $\phi \in X^*$ and $\psi \in Y^*$. Then
$$\begin{aligned} \sum_{j=1}^n \phi(x_j)\psi(y_j) &= \phi\left(\sum_{j=1}^n \psi(y_j)x_j\right) \\ &= \end{aligned}$$
2.
3.

□

Exercise D.4.0.7. Let X, Y be vector spaces. Then $(X \otimes Y, \otimes)$ is a tensor product of X and Y.

Proof. Let Z be a vector space and $\alpha \in L^2(X, Y; Z)$. Define $\phi : X \otimes Y \to Z$ by $\phi\left(\sum_{j=1}^n \lambda_j x_j \otimes y_j\right) := \sum_{j=1}^n \lambda_j \alpha(x_j, y_j)$.

- **(well defined):**
  Let $u \in X \otimes Y$. Then there exist $(\lambda_j)_{j=1}^n \subset \mathbb{K}$, $(x_j)_{j=1}^n \subset X$, $(y_j)_{j=1}^n \subset Y$ such that $u = \sum_{j=1}^n \lambda_j x_j \otimes y_j$. Suppose that $u = 0$. Let $\phi \in Z^*$. Then $\phi \circ \alpha \in L^2(X, Y; Z)$.

□

Bibliography

[1] Introduction to Algebra

[2] Introduction to Analysis

[3] Introduction to Fourier Analysis

[4] Introduction to Measure and Integration