

Similarity

Cosine Similarity

- If $X = Y$ (the two vectors have the same values)

$$\frac{X \cdot Y}{\|X\| \|Y\|} = \frac{X \cdot X}{\|X\| \|X\|} = \frac{x_1x_1 + \cdots + x_Nx_N}{\sqrt{x_1x_1 + \cdots + x_Nx_N} \cdot \sqrt{x_1x_1 + \cdots + x_Nx_N}} = 1$$

- If X is orthogonal to Y

$$- X = [-1, 3], Y = [3, 1]$$

$$\frac{X \cdot Y}{\|X\| \|Y\|} = \frac{(-1 \cdot 3) + (3 \cdot 1)}{\|X\| \|Y\|} = 0$$

- The length of vectors does not affect similarity

$$\frac{X \cdot Y}{\|X\| \|Y\|} = \frac{(5X) \cdot Y}{\|(5X)\| \|Y\|}$$

Euclidean Distance

- Euclidean distance is the length of the line connecting two points

$$\bullet d(X, Y) = \sqrt{\sum_i^N (x_i - y_i)^2}$$

- Example:

$$- X = [1, 2], Y = [3, 5]$$

$$- d(X, Y) = \sqrt{(1 - 3)^2 + (2 - 5)^2}$$

$$- d(X, Y) = \sqrt{13}$$

- If the vectors are the same, $d(X, X) = 0$

Manhattan Distance

$$\bullet d(X, Y) = \sum_i^N |x_i - y_i|$$

- Also called $L1$ or taxicab distance

- Measured in straight lines along each axis

- The number of blocks that you need to drive to get from one point to another

- Example:

$$- X = [1, 2], Y = [3, 5]$$

$$- d(X, Y) = |1 - 3| + |2 - 5| = 5$$

- If the vectors are the same, $d(X, X) = 0$

Minkowski Distance


- Euclidean and Manhattan distances are specific forms of the Minkowski distance

- $d(X, Y) = \left(\sum_i^N |x_i - y_i|^p \right)^{\frac{1}{p}}$

- If $p = 1$, we get the Manhattan
- If $p = 2$, we get Euclidean
- If $p = \infty$, we get Chebyshev

- $d(X, Y) = \max_i |x_i - y_i|$

Visualizing the Distances



Properties

- Euclidean distance and Manhattan distance both have the following properties
 - Symmetric: $d(X, Y) = d(Y, X)$
 - Positive
 - $d(X, Y) = 0$ if and only if the two vectors contain the same values
 - The triangle inequality holds for three points p, q, r . That is, $d(p, q) + d(q, r) \leq d(p, r)$

Binary Similarity – Jaccard Similarity Index

- If we have binary vectors (vectors only containing 1s and 0s), we can use specific similarities
- $J = \frac{M_{11}}{M_{01} + M_{10} + M_{11}}$
- M_{11} : The number of times that both vectors have a 1

- M_{01} : The number of times that vector A has a 0 when vector B has a 1
- M_{10} : The number of times that vector A has a 1 when vector B has a 0
- M_{00} : The number of times that both vectors have a 0
- Example:
 - $A = [0, 0, 0, 1, 1, 1]$
 - $B = [0, 1, 0, 1, 0, 0]$
 - $M_{11} = 1$
 - $M_{01} = 1$
 - $M_{10} = 2$
 - $J = \frac{1}{1+2+1} = 0.25$