

LETTER TO THE EDITOR

# A Presentation of Galaxies with Polar Structures

A.V. Mosenkov<sup>1</sup> and C.M. Garrett<sup>1</sup>

Department of Physics and Astronomy, N283 ESC, Brigham Young University, Provo, UT 84602, USA

Received November, 2023; accepted ???, 2023

## ABSTRACT

Certain galaxies exist that exhibit kinematically and photometrically decoupled systems that are inclined to an extreme angle as compared to the major planar axis of the host galaxy. Historically, these have been identified and studied, however, they are quite rare and difficult to catalogue. In this work, I examined a sample of over 10,000 galaxies from the Sloan Digital Sky Survey (SDSS) Stride 82 for the presence of galaxies with these polar structures described. With the deep SDSS Stride 82, the DESI Legacy Imaging Surveys, and the Hyper Suprime-Cam Subaru Strategic Program, 143 exceptional candidates with polar structures were found. The results indicate that the occurrence rate for which polar structures are found in galaxies is underestimated, and may total to 1.43%

**Key words.** galaxies: structure – galaxies: elliptical and lenticular – galaxies: formation – galaxies: evolution

## 1. Introduction

Polar-ring galaxies (PRGs) were identified first in a series of studies by [Schechter & Gunn \(1978\)](#), [Bertola & Galletta \(1978\)](#), and [Schweizer et al. \(1983\)](#), and are a rare type of galaxy that has a pronounced ring of stars and gas orbiting nearly orthogonal to the plane of the host galaxy. A more generalized class of polarized galactic structures were proposed which made space for polar/tilted discs, ([Bertola et al. 1999](#); [Mosenkov et al. 2020b](#)), polar bulges ([Corsini et al. 2012](#); [Reshetnikov et al. 2015](#)), halos that are perpendicular to the disc ([Crnojević et al. 2016](#); [Mosenkov et al. 2020a](#); [Martínez-Delgado et al. 2021](#)), and tidally disrupted material from nearby satellites ([Martínez-Delgado et al. 2010](#); [Müller et al. 2019](#); [Martínez-Delgado et al. 2023](#)). The class of polar structures has proven useful in exploring properties of dark matter ([Whitmore et al. 1987](#); [Sackett et al. 1994](#); [Combes & Arnaboldi 1996](#); [Iodice et al. 2003](#); [Moiseev et al. 2011](#); [Snaith et al. 2012](#); [Khoperskov et al. 2014](#); [Moiseev et al. 2015](#)) as well as galactic accretion dynamics ([Brook et al. 2008](#)) and merger events ([Bekki 1998](#); [Bournaud & Combes 2003](#)). To date, PRGs are generally understood to form from tidal accretion events, where matter from a donor galaxy is given to the host ([Schweizer et al. 1983](#); [Reshetnikov & Sotnikova 1997](#)) which also includes the disruption of a (gas-rich) satellite into the plane perpendicular of the host ([Rix & Katz 1991](#); [Katz & Rix 1992](#)), major mergers of galaxies ([Bekki 1997, 1998](#); [Bournaud & Combes 2003](#)), and lastly, cosmological filament accretion ([Macciò et al. 2006](#); [Brook et al. 2008](#)). Modern simulations conform the possibility of these structures forming in any of these manners ([Liao & Gao 2019](#); [Wright et al. 2021](#); [Smirnov et al. 2023](#)). While past catalogues of PRGs have experienced severe limitations in their photometric depth, more modern observations have allowed for much more careful analysis of structures previously unidentifiable. While it is unfortunate that many of these galaxies have very low average surface brightness, which has caused variance in predicted occurrence rates (See [Reshetnikov et al. \(2011\)](#), [Reshetnikov & Mosenkov \(2019\)](#)), polar structures are still discernible with deep imaging from the Sloan Digital Sky Survey. In this article, we aim to con-

tinue the study by [Mosenkov et al. \(2022\)](#), adding to ([Mosenkov et al. \(2024\)](#)) and the examination of Stride 82 data to identify additional, previously unrecognised galaxies with polar rings. Surprisingly, we have found a total of 143 galaxies with polar rings, polar bulges and polar halos, many more than previously spotted using regular imaging, demonstrating that PRGs are more common than expected.


## 2. Data and sample selection

A sample of 10,000 galaxies was obtained from the Siena Galaxy Atlas ([John Moustakas \(2020/2023\)](#)), which is a compilation of galaxies using the Hyperleda database. The galactic sample was randomly selected from the database. All of the images from the Siena Galaxy Atlas were downloaded from the DESI Legacy Survey and enhanced using the method described in [Mosenkov et al. \(2024\)](#). We note that the results of this classification are preliminary because they may be subject to our erroneous interpretation of the observed structures, so a quantitative analysis of the selected objects by means of host+ring decomposition (similar to that in [Mosenkov et al. 2022](#)) will be performed in a subsequent work. In this paper, this classification is provided only for reference, and we do not use this separation into sub-classes in the following sections. Here, we focus only on the statistics of polar rings or related objects, whereas in future paper, we will present an atlas of all selected candidates for galaxies with polar structures and explore their morphological features in detail. In Fig. 1, we present typical examples of candidates for galaxies with polar structures from the selected sample<sup>1</sup>.


## 3. Results and discussion

Out of 10,000 galaxies, 143 total were labeled as exhibiting polar structures. This is consistent with [Mosenkov et al. \(2024\)](#), where they gave an approximate value of 1-3%. The best polar structures are presented in Figure 1. Polar structures can be formed in

<sup>1</sup> Mosaic images for all candidates, along with their coordinates, are available in the supplementary material.



**Fig. 1.** Examples of candidates for galaxies with polar structures from our sample as seen in the DESI Legacy from left to right: a polar bulge structure, another galaxy with a polar bulge, a polar-halo galaxy, and a galaxy forming another polar bulge structure. The coordinates for each galaxy are given in the upper right corner of each plot, with the reference name in the upper left.



**Fig. 2.** Normalized histograms comparing *a)* apparent magnitude in the *r* band, *b)* redshift *z*, *c)* absolute magnitude in the *r* band, and *d)* galaxy colour *g-r*. The black filled histograms represent the entire Stripe 82 sample, while the SPRC+RM19 (see text) and our new candidates are shown as blue dashed and orange solid histograms, respectively. Plot *e)*: a colour-magnitude diagram (the underlying density plot depicts the distribution of the entire Stripe 82 sample of the 18,362 galaxies). All magnitudes have been corrected for K-correction (Chilingarian et al. 2010; Chilingarian & Zolotukhin 2012) and Galactic extinction using the conversions from Schlafly & Finkbeiner (2011) applied to  $E(B-V)$  from Schlegel et al. (1998). Plot *f)*: luminosity functions of our new PRG sample and the entire Stripe 82 sample calculated with Choloniewski’s method (Choloniewski 1986).

a variety of different methods. Firstly, polar structures can form via tidal accretion. As two galaxies interact gravitationally, polarities can form 3. Secondly, polar structures have been observed to form as repercussions of major merging. Extreme gravitational interactions yield such structures as can be viewed in 4. Thirdly, we can observe polar structures observe in starburst events. As gravitational perturbations or other phenomena trigger star burst in the host galaxy, either the gravitational perturbations themselves or the extreme increase in interstellar winds can generate polar structures. Such can be viewed in figure 5.


A final method we discuss in here (and this list is by no means exhaustive), is accretion from cosmic filaments. As a galaxy passes through filaments in large scale cosmological structures, it may develop polar structures as the intergalactic

gas can frictionally force polar structures out of the host galaxy, as seen in figure 6.


Out of the 143 candidates, we can identify 70 ‘strong’ candidates, which we consider to be those candidates that exhibit polar structures with a high confidence interval.

#### 4. Conclusions


In this letter, we present the findings from a systematic review over the course of 10 weeks to categorize 143 galaxies with polar structures, 70 of which are highly probable candidates. This entails a PRG occurrence rate of about 1.43%. This is highly consistent with the findings of Mosenkov et al. (2024), which means the occurrence rate of galaxies with polar structures are much




**Fig. 3.** A demonstration of tidal accretion– the target galaxy PGC 56239 is seen to gain material from a neighbor, which forms a structure perpendicular to the plane of the galaxy.



**Fig. 4.** A major merger is depicted, and polar structures are visible in all directions from the combining cores.



**Fig. 5.** The starburst event forms a ring of blue gas and star formation around the center, polar with respect to the plane of the galaxy.



**Fig. 6.** The bulge/core of the galaxy is pressure stripped as it moves through intergalactic gas in a cosmic filament.

higher than previously thought (Whitmore et al. 1990, Moiseev et al. 2011). These candidates will be used to train a neural network and conduct a rigorous analysis of polar structure galaxies, based on the largest sample to date. We stress, however, that these galaxies are only candidates for polar structures and are yet to be kinematically confirmed. In a future study, we will release a full catalogue of our polar structure candidates, alongside photometric decompositions of some candidate galaxies identified here.

**Acknowledgements.** Funding for the Sloan Digital Sky Survey IV has been provided by the Alfred P. Sloan Foundation, the U.S. Department of Energy Office of Science, and the Participating Institutions. SDSS-IV acknowledges support and resources from the Center for High-Performance Computing at the University of Utah. The SDSS website is [www.sdss.org](http://www.sdss.org). SDSS-IV is managed by the Astrophysical Research Consortium for the Participating Institutions of the SDSS Collaboration including the Brazilian Participation Group, the Carnegie Institution for Science, Carnegie Mellon University, the Chilean Participation Group, the French Participation Group, Harvard-Smithsonian Center for Astrophysics,

Instituto de Astrofísica de Canarias, The Johns Hopkins University, Kavli Institute for the Physics and Mathematics of the Universe (IPMU) / University of Tokyo, the Korean Participation Group, Lawrence Berkeley National Laboratory, Leibniz Institut für Astrophysik Potsdam (AIP), Max-Planck-Institut für Astronomie (MPIA Heidelberg), Max-Planck-Institut für Astrophysik (MPA Garching), Max-Planck-Institut für Extraterrestrische Physik (MPE), National Astronomical Observatories of China, New Mexico State University, New York University, University of Notre Dame, Observatório Nacional / MCTI, The Ohio State University, Pennsylvania State University, Shanghai Astronomical Observatory, United Kingdom Participation Group, Universidad Nacional Autónoma de México, University of Arizona, University of Colorado Boulder, University of Oxford, University of Portsmouth, University of Utah, University of Virginia, University of Washington, University of Wisconsin, Vanderbilt University, and Yale University. The Legacy Surveys consist of three individual and complementary projects: the Dark Energy Camera Legacy Survey (DECaLS; NOAO Proposal ID # 2014B-0404; PIs: David Schlegel and Arjun Dey), the Beijing-Arizona Sky Survey (BASS; NOAO Proposal ID # 2015A-0801; PIs: Zhou Xu and Xiaohui Fan), and the Mayall z-band Legacy Survey (MzLS; NOAO Proposal ID # 2016A-0453; PI: Arjun Dey). DECaLS, BASS and MzLS together include data obtained, respectively, at the Blanco telescope, Cerro Tololo Inter-American Observatory, National Optical Astronomy Observatory (NOAO); the Bok telescope, Steward Observatory, University of Arizona; and the Mayall tele-



scope, Kitt Peak National Observatory, NOAO. The Legacy Surveys project is honored to be permitted to conduct astronomical research on Iolkam Du'ag (Kitt Peak), a mountain with particular significance to the Tohono O'odham Nation. NOAO is operated by the Association of Universities for Research in Astronomy (AURA) under a cooperative agreement with the National Science Foundation. This project used data obtained with the Dark Energy Camera (DECam), which was constructed by the Dark Energy Survey (DES) collaboration. Funding for the DES Projects has been provided by the U.S. Department of Energy, the U.S. National Science Foundation, the Ministry of Science and Education of Spain, the Science and Technology Facilities Council of the United Kingdom, the Higher Education Funding Council for England, the National Center for Supercomputing Applications at the University of Illinois at Urbana-Champaign, the Kavli Institute of Cosmological Physics at the University of Chicago, Center for Cosmology and Astro-Particle Physics at the Ohio State University, the Mitchell Institute for Fundamental Physics and Astronomy at Texas A&M University, Financiadora de Estudos e Projetos, Fundacao Carlos Chagas Filho de Amparo, Financiadora de Estudos e Projetos, Fundacao Carlos Chagas Filho de Amparo, a Pesquisa do Estado do Rio de Janeiro, Conselho Nacional de Desenvolvimento Científico e Tecnológico and the Ministerio da Ciencia, Tecnologia e Inovacao, the Deutsche Forschungsgemeinschaft and the Collaborating Institutions in the Dark Energy Survey. The Collaborating Institutions are Argonne National Laboratory, the University of California at Santa Cruz, the University of Cambridge, Centro de Investigaciones Energeticas, Medioambientales y Tecnologicas-Madrid, the University of Chicago, University College London, the DES-Brazil Consortium, the University of Edinburgh, the Eidgenössische Technische Hochschule (ETH) Zurich, Fermi National Accelerator Laboratory, the University of Illinois at Urbana-Champaign, the Institut de Ciències de l'Espai (IEEC/CSIC), the Institut de Física d'Altes Energies, Lawrence Berkeley National Laboratory, the Ludwig-Maximilians Universität München and the associated Excellence Cluster Universe, the University of Michigan, the National Optical Astronomy Observatory, the University of Nottingham, the Ohio State University, the University of Pennsylvania, the University of Portsmouth, SLAC National Accelerator Laboratory, Stanford University, the University of Sussex, and Texas A&M University. BASS is a key project of the Telescope Access Program (TAP), which has been funded by the National Astronomical Observatories of China, the Chinese Academy of Sciences (the Strategic Priority Research Program "The Emergence of Cosmological Structures" Grant # XDB09000000), and the Special Fund for Astronomy from the Ministry of Finance. The BASS is also supported by the External Cooperation Program of Chinese Academy of Sciences (Grant # 114A11KYSB20160057), and Chinese National Natural Science Foundation (Grant # 11433005). The Legacy Survey team makes use of data products from the Near-Earth Object Wide-field Infrared Survey Explorer (NEOWISE), which is a project of the Jet Propulsion Laboratory/California Institute of Technology. NEOWISE is funded by the National Aeronautics and Space Administration. The Legacy Surveys imaging of the DESI footprint is supported by the Director, Office of Science, Office of High Energy Physics of the U.S. Department of Energy under Contract No. DE-AC02-05CH1123, by the National Energy Research Scientific Computing Center, a DOE Office of Science User Facility under the same contract; and by the U.S. National Science Foundation, Division of Astronomical Sciences under Contract No. AST-0950945 to NOAO. The Hyper Suprime-Cam (HSC) collaboration includes the astronomical communities of Japan and Taiwan, and Princeton University. The HSC instrumentation and software were developed by the National Astronomical Observatory of Japan (NAOJ), the Kavli Institute for the Physics and Mathematics of the Universe (Kavli IPMU), the University of Tokyo, the High Energy Accelerator Research Organization (KEK), the Academia Sinica Institute for Astronomy and Astrophysics in Taiwan (ASIAA), and Princeton University. Funding was contributed by the FIRST program from the Japanese Cabinet Office, the Ministry of Education, Culture, Sports, Science and Technology (MEXT), the Japan Society for the Promotion of Science (JSPS), Japan Science and Technology Agency (JST), the Toray Science Foundation, NAOJ, Kavli IPMU, KEK, ASIAA, and Princeton University. This paper makes use of software developed for Vera C. Rubin Observatory. We thank the Rubin Observatory for making their code available as free software at <http://pipelines.lsst.io/>. This paper is based on data collected at the Subaru Telescope and retrieved from the HSC data archive system, which is operated by the Subaru Telescope and Astronomy Data Center (ADC) at NAOJ. Data analysis was in part carried out with the cooperation of Center for Computational Astrophysics (CfCA), NAOJ. We are honored and grateful for the opportunity of observing the Universe from Maunakea, which has the cultural, historical, and natural significance in Hawaii.

## References

- Bekki, K. 1997, *The Astrophysical Journal*, 490, L37  
 Bekki, K. 1998, *ApJ*, 499, 635  
 Bekki, K. 1998, *The Astrophysical Journal*, 499, 635  
 Bertola, F., Corsini, E., Beltrán, J. V., et al. 1999, *AJ*, 519, L127

- Bertola, F. & Galletta, G. 1978, *ApJ*, 226, L115  
 Bournaud, F. & Combes, F. 2003, *A&A*, 401, 817  
 Brook, C. B., Governato, F., Quinn, T., et al. 2008, *ApJ*, 689, 678  
 Chilingarian, I. V., Melchior, A.-L., & Zolotukhin, I. Y. 2010, *MNRAS*, 405, 1409  
 Chilingarian, I. V. & Zolotukhin, I. Y. 2012, *MNRAS*, 419, 1727  
 Choloniewski, J. 1986, *MNRAS*, 223, 1  
 Combes, F. & Arnaboldi, M. 1996, *A&A*, 305, 763  
 Corsini, E. M., Méndez-Abreu, J., Pastorello, N., et al. 2012, *MNRAS*, 423, L79  
 Crnojević, D., Sand, D. J., Spekkens, K., et al. 2016, *ApJ*, 823, 19  
 Iodice, E., Arnaboldi, M., Bournaud, F., et al. 2003, *ApJ*, 585, 730  
 John Moustakas, D. L. 2020/2023, *Siena Galaxy Atlas 2020*  
 Katz, N. & Rix, H.-W. 1992, *ApJ*, 389, L55  
 Khoperskov, S. A., Moiseev, A. V., Khoperskov, A. V., & Saburova, A. S. 2014, *MNRAS*, 441, 2650  
 Liao, S. & Gao, L. 2019, *MNRAS*, 485, 464  
 Macciò, A. V., Moore, B., & Stadel, J. 2006, *ApJ*, 636, L25  
 Martínez-Delgado, D., Cooper, A. P., Román, J., et al. 2023, *A&A*, 671, A141  
 Martínez-Delgado, D., Gabany, R. J., Crawford, K., et al. 2010, *AJ*, 140, 962  
 Martínez-Delgado, D., Javier, R., Erkal, D., et al. 2021, *MNRAS*, 506, 5030  
 Moiseev, A., Khoperskov, S., Khoperskov, A., et al. 2015, *Baltic Astronomy*, 24, 76  
 Moiseev, A. V., Smirnova, K. I., Smirnova, A. A., & Reshetnikov, V. P. 2011, *MNRAS*, 418, 244  
 Mosenkov, A., Bahr, S., Reshetnikov, V., Shakespear, Z., & Smirnov, D. 2024, *Mosenkov, A., Rich, R. M., Koch, A., et al. 2020a, MNRAS*, 494, 1751  
 Mosenkov, A. V., Reshetnikov, V. P., Skryabina, M. N., & Shakespear, Z. 2022, *Research in Astronomy and Astrophysics*, 22, 115003  
 Mosenkov, A. V., Smirnov, A. A., Sil'chenko, O. K., et al. 2020b, *MNRAS*, 497, 2039  
 Müller, O., Vudragović, A., & Bílek, M. 2019, *A&A*, 632, L13  
 Reshetnikov, V., Faúndez-Abans, M., & de Oliveira-Abans, M. 2011, *Astronomy Letters*, 37, 171  
 Reshetnikov, V., Savchenko, S., Mosenkov, A., Sotnikova, N. Y., & Bizyaev, D. 2015, , 41, 748  
 Reshetnikov, V. & Sotnikova, N. 1997, *A&A*, 325, 933  
 Reshetnikov, V. P. & Mosenkov, A. V. 2019, *MNRAS*, 483, 1470  
 Rix, H. W. & Katz, N. 1991, in *Warped Disks and Inclined Rings around Galaxies*, 112  
 Sackett, P. D., Rix, H.-W., Jarvis, B. J., & Freeman, K. C. 1994, *ApJ*, 436, 629  
 Schechter, P. L. & Gunn, J. E. 1978, *AJ*, 83, 1360  
 Schlafly, E. F. & Finkbeiner, D. P. 2011, *ApJ*, 737, 103  
 Schlegel, D. J., Finkbeiner, D. P., & Davis, M. 1998, *ApJ*, 500, 525  
 Schweizer, F., Whitmore, B. C., & Rubin, V. 1983, *The Astronomical Journal*, 88, 909  
 Schweizer, F., Whitmore, B. C., & Rubin, V. C. 1983, *AJ*, 88, 909  
 Smirnov, D. V., Mosenkov, A. V., & Reshetnikov, V. P. 2023, *Monthly Notices of the Royal Astronomical Society*, 527, 4112  
 Snaith, O. N., Gibson, B. K., Brook, C. B., et al. 2012, *MNRAS*, 425, 1967  
 Whitmore, B. C., Lucas, R. A., McElroy, D. B., et al. 1990, *AJ*, 100, 1489  
 Whitmore, B. C., McElroy, D. B., & Schweizer, F. 1987, *ApJ*, 314, 439  
 Wright, R. J., Lagos, C. d. P., Power, C., & Correa, C. A. 2021, *MNRAS*, 504, 5702