

Multenions and Differential Invariants.—III.

BY ALEX. MCAULAY, PROFESSOR OF MATHEMATICS IN THE UNIVERSITY OF TASMANIA.

(Communicated by W. B. Hardy, Sec.R.S. Received July 20, 1921.)

§ 19. *SUGGESTED IMPROVEMENT OF SOME NOTATIONS AND TERMS.*—The reader has probably found that the terminology for multenion linities, adopted hitherto because of the writer's timid reluctance to depart from accepted meanings, is too inexpressive of the intended signification. A covariant linity fabricates a covariant multenion out of a contravariant. Let us call it a $c/\mathfrak{O}$ or a coexcontra (co-ex-contra); and similarly let us now denote a contravariant, comixt or contramixt linity by the name contraexco, coexco or contraexcontra, and by the symbol $\mathfrak{O}/c$, c/c or $\mathfrak{O}/\mathfrak{O}$ respectively. Let us also use parenthesis marks¹ before a letter such as ϕ , of the same general shapes as c and $\mathfrak{O}$ (omitting the solidus), to indicate the type of invariance, thus—let us say that

$$\phi = {}^{\circ}\phi, {}^{\circ}(\phi, ((\phi \text{ or }))\phi$$

indicates that ϕ is a

$$c/\mathfrak{O}, \quad \mathfrak{O}/c, \quad c/c \text{ or } \mathfrak{O}/\mathfrak{O}$$

respectively. Similarly, $p = {}^{\circ}p$ indicates that p is a covariant multenion and $q = {}^{\circ}q$ indicates that q is a contravariant multenion. If we actually insert all these indicating parenthesis marks, which we shall do but rarely, their mode of combination is simple. Thus

$$)p =))\phi {}^{\circ}(\psi (q$$

is an invariant equation. The features that render it so are (1) that the second mark of one letter, ϕ or ψ , is identical with the first mark of the next; these two marks virtually cancel one another; and (2) the proper mark of the resultant symbol p , of the whole operation, is the first written mark of the compound symbol $\phi\psi q$.

When with these notations we use the dash which indicates conjugacy, there is, unless we insert brackets, ambiguity in the meanings of $((F'$ and $))F'$ (the two mixed types as we have hitherto called them), because $((F')'$ is not of the same type c/c as $((F$, but of the type $\mathfrak{O}/\mathfrak{O}$. There is no such ambiguity with ${}^{\circ}\phi'$ and ${}^{\circ}(\phi'$ nor with $f^{\circ}(F$ and $f^{\circ})F$, where fx is a rational function of x . [Remember that $f^{\circ}\phi$ and $f^{\circ})\phi$ have no invariative meaning. I think it

¹Intended by author to be *commas*, upright and inverted. In future printing the author still thinks commas would be preferable to the bracket substitute.

best to adhere to the general rule, not rigidly prescribed, that $c/\mathfrak{O}$ and $\mathfrak{O}/c$ should be denoted by Greek letters and c/c and $\mathfrak{O}/\mathfrak{O}$ by Roman capitals.]

Our new notations and terms apply to other symbols than invariantive; for instance, to Γ_β . Thus

$$\Gamma_\beta = \mathfrak{)}\Gamma_\beta, \quad E'_\beta = \mathfrak{((}E'_\beta).$$

We shall retain the “cross” notation to indicate different symbols, and generally restrict it, without rigid prescription, to the same meanings as hitherto.

Our former $\bar{\Gamma}_\beta$ and $\bar{E}_\beta$ will henceforth not be used, because we have in §18 a new notation, according to which ${}^aE_\beta$ would have reference to a Weyl manifold, and would indicate that ${}^aE_\beta$ was the part of E_β not containing terms in λ ; and if λ is actually everywhere zero so that the manifold becomes a Riemann manifold, §18 provides the use of F_β in place of E_β . In the present paper we occupy ourselves exclusively with a Riemann manifold, and use the letters θ , l , F_β , Θ_β , Ω , C ($= \mathfrak{((}C = \Omega\theta^{-1}$) provided for that case.

A thick dot² will be used to denote that a symbol is a “density,” though occasionally, as with W (action density), and f a part of W below, we shall not consider it necessary to use the dot. Once for all, we define $\dot{\eta}'$ and $\dot{\theta}$ by the equations

$$\dot{\eta}' = k\eta^{-1}, \quad \dot{\theta} = l\theta^{-1}. \quad (1)$$

but we may regard it as a rule, unless the contrary is definitely stated, as here with $\dot{\eta}'$ and $\dot{\theta}$, that the dot attached to a symbol means that the undotted symbol has been multiplied by l (by k if we were dealing with a Weyl manifold). Thus $\dot{C}$, $\dot{G}$ stand for lC , lG , and again, if we introduce $\dot{\kappa}$ as a contravariant vector density, then κ will mean $l^{-1}\dot{\kappa}$.

With regard to $\dot{\theta}$ in (1), we shall use it as applied to a general multenion, say, to $w^\times = V_a q^\times$, but not exactly with the extensive meaning. Thus $\dot{\theta}u^\times$ means $l\theta^{-1}u^\times$ where θ^{-1} but not l is used in the extensive meaning. Thus, for instance, $\dot{\theta}V_3\alpha^\times\beta^\times\gamma^\times$ means $lV_3\theta^{-1}\alpha^\times\theta^{-1}\beta^\times\theta^{-1}\gamma^\times$, and not $l^3V_3\theta^{-1}\alpha^\times\theta^{-1}\beta^\times\theta^{-1}\gamma^\times$.

§ 20. MAXWELL'S EQUATIONS FOR MATTER IN BULK.—It is fully time to show how the special requirements of Relativity are met in detail in our method. In now introducing Maxwell's equations, let us throughout keep in mind the application to matter in bulk. If for no other reason this is desirable because thereby we are prepared to accept the beautiful

²It is necessary, because of papers despatched to England in the early months of 1922, to state that this dot replaced a *bar* placed over the symbol affected. The replacement was made in August, 1922, at the suggestion of the printers. It seems now (August) too late to replace the bar in the papers just mentioned.

explanation that Maxwell's theory affords of the known behaviour of light in uniaxal and biaxal crystals. When we compare results with Maxwell's we shall speak as if $n = 4$. Nevertheless, in the general argument, we shall continue to suppose n to have any positive integral value. When with $n = 4$ we put

$$\rho = xl_1 + yl_2 + zl_3 + tl_4$$

we shall speak of x, y, z as the spatial co-ordinates and t as the time, even when by transformation of all four co-ordinates t may mean something very different from the time as ordinarily understood.

Let Maxwell's vector potential be $\mathbf{A} = (A_1, A_2, A_3)$, electrostatic potential $= A_4$, magnetic induction $\mathbf{B} = (B_1, B_2, B_3)$, electric intensity $\mathbf{E} = (E_1, E_2, E_3)$, magnetic field strength $\mathbf{H} = (H_1, H_2, H_3)$, dielectric displacement $\mathbf{D} = (D_1, D_2, D_3)$, conduction current $\mathbf{K} = (K_1, K_2, K_3)$, electrostatic density $= K_4$, force per unit volume $\mathbf{F} = (F_1, F_2, F_3)$, power $(-\mathbf{SEK})$ per unit volume $= F_4$.

Also let λ be a covariant vector (better use $\lambda^\times$ in the future) given by

$$\lambda = \sum A_c l_c \quad (c = 1, 2, 3, 4),$$

$\omega^\times$ a covariant ${}_nV_2$ given by

$$\omega^\times = \sum B_1 l_2 l_3 + \sum E_1 l_4 l_1$$

(the meaning of each summation, of three terms, being sufficiently obvious), $\dot{\omega}$ a contravariant ${}_nV_2$ density given by

$$\dot{\omega} = \sum H_1 l_2 l_3 + \sum D_1 l_4 l_1,$$

$\dot{\kappa}$ a contravariant vector density given by

$$\dot{\kappa} = \sum K_c l_c \quad (c = 1, 2, 3, 4),$$

and $\dot{v}^\times$ a covariant vector density given by

$$\dot{v}^\times = \sum F_c l_c \quad (c = 1, 2, 3, 4).$$

Maxwell's quaternion equations

$$\mathbf{B} = V\nabla\mathbf{A}, \quad \mathbf{E} = -\nabla A_4 - \partial\mathbf{A}/c\partial t, \quad (1)$$

$$\mathbf{K} = V\nabla\mathbf{H} - \partial\mathbf{D}/\partial t, \quad \mathbf{K}_4 = -S\nabla\mathbf{D}, \quad (2)$$

$$\mathbf{F} = V\mathbf{KB} + K_4\mathbf{E}, \quad \mathbf{F}_4 = -S\mathbf{KE}, \quad (3)$$

are identical with the multenion equations³

$$\omega^\times = V_2 \nabla \lambda, \quad (1a)$$

$$\dot{\kappa} = V_1 \nabla \dot{\omega}, \quad (2a)$$

$$\dot{v}^\times = V_1 \dot{\kappa} \omega^\times. \quad (3a)$$

The equations (1a), (2a), (3a) are all invariantive.

³For a convenient, simple, systematic method of converting all ($n = 4$) multenion formulae to corresponding quaternion formulae, see a paper by the author (despatched a month ago, viz., November, 1922, to England), "The Mechanical Forces ... in an Electromagnetic Field."

[A digression. In Maxwell's presentation of the fundamental aspects of the electromagnetic field $V\nabla\mathbf{H}$ in equation (2) is looked upon as *the* electric current, consisting of two parts, conduction current $\mathbf{K}$ and dielectric current $\partial\mathbf{D}/\partial t$. In the new Relativity scheme, as given by (2a), we naturally interchange the *rôles* of $V\nabla\mathbf{H}$ and $\mathbf{K}$; that is to say, we look upon $\mathbf{K}$ as *the* current, consisting of two parts, a magnetic part, $V\nabla\mathbf{H}$, and an electric part, $-\partial\mathbf{D}/\partial t$. In the multenion presentation (see below for N and I) $V\nabla\mathbf{H}$ is $V_1\nabla(N\dot{\omega})$, $-\partial\mathbf{D}/\partial t$ is $NV_1\nabla(I\dot{\omega})$, and K_4 (multiplied into l_4) is $IV_1\nabla(I\dot{\omega})$. [In the last sentence N and I are used as if $N = {}^{\text{))}}N$, $I = {}^{\text{))}}I$; if, as usual, we take $N = {}^{\text{((}}N$, $I = {}^{\text{((}}I$, then in the last sentence each N should be N' and each I should be I' .] In the future, then, when we use the term current, Maxwell's conduction current will be meant. Maxwell does not put $\mathbf{F} = V\mathbf{KB}$ as we have asserted in (3), but instead $\mathbf{F} = V(V\nabla\mathbf{H})\mathbf{B}$.]

Before speaking of the relations between Maxwell's $(\mathbf{B}, \mathbf{E})$ and $(\mathbf{H}, \mathbf{D})$, that is, between $\omega^\times$ and $\dot{\omega}$, let us derive Lorentz's generalizations of Maxwell's equations to bodies in motion from the general principle of Relativity. The generalisations in nowise alter (1a), (2a), (3a). Put N_4 and I_4 for the normal and incident components with reference to l_4 , when the co-ordinates are Galilean. In reality we have

$$\left. \begin{array}{ll} \mathbf{A} \text{ is } N_4\lambda, & l_4A_4 \text{ is } l_4\lambda \\ \mathbf{B} \text{ is } N_4\omega^\times, & \mathbf{E} \text{ is } l_4\omega^\times \end{array} \right\} \quad (1b)$$

$$\left. \begin{array}{ll} \mathbf{H} \text{ is } N_4\dot{\omega}, & \mathbf{D} \text{ is } l_4\dot{\omega} \\ \mathbf{K} \text{ is } N_4\dot{\kappa}, & l_4K_4 \text{ is } l_4\dot{\kappa} \end{array} \right\} \quad (2b)$$

$$\mathbf{F} \text{ is } N_4\dot{v}^\times, \quad l_4F_4 \text{ is } l_4\dot{v}^\times. \quad (3b)$$

In the general θ (Riemann) manifold

$$\iota = dp/ds = dp/\sqrt{(V_0 dp \theta dp)}$$

degenerates when Galilean co-ordinates are used ($\theta = 1$) to

$$\begin{aligned} & (l_1 dx + l_2 dy + l_3 dz + l_4 dt) / \sqrt{(-dx^2 - dy^2 - dz^2 + dt^2)} \\ &= (1 - \sum \dot{x}^2)^{-\frac{1}{2}} \cdot (l_1 \dot{x} + l_2 \dot{y} + l_3 \dot{z} + l_4), \end{aligned}$$

of which we say nothing more here than that when the co-ordinates are so chosen that the particle is at rest ι (the contravariant world unit tangent vector to the world locus of a particle), degenerates to l_4 . Maxwell's equations, as given above, are to be supposed true only when the medium is at rest. The generalisations of N_4 and I_4 are thus N and I (or N' and I'), the normal and incident components with reference to ι ; normal *component* and normal *expression*, in this case, meaning the same, because

$V_0 \mathfrak{t} \theta \mathfrak{t} = 1$. Below N and I will always mean what has just been described with reference to $\mathfrak{t}$, unless explicit exception is made; (1b), (2b), (3b), become like (1a), (2a), (3a), all invariantive when we read $\mathfrak{t}$, N and I in place of l_4 , N_4 , and I_4 , in (1a) and (3a); and in (2a), $\mathfrak{t}$, N' and I' in place of l_4 , N_4 , and I_4 . [Remember that $N = (N, N' =)(\psi$, and similarly for I .]

The equations this process gives when a body is moving, and we are using Galilean co-ordinates, are Lorentz's well-known generalizations of Maxwell's quaternion equations (not of our multenion equations, which have been in no way altered by what has been said).

From (1a) and (2a) we have of course

$$V_3 \nabla \omega^\times = 0, \quad (4)$$

$$V_0 \nabla \dot{\mathbf{K}} = 0. \quad (5)$$

These must be implicitly contained in (1) and (2) and indeed are found to be

$$S \nabla \mathbf{B} = 0, \quad V \nabla \mathbf{E} + \partial \mathbf{B} / \partial t = 0, \quad (4a)$$

$$S \nabla \mathbf{K} = \partial K_4 / \partial t. \quad (5a)$$

Maxwell's relations between $(\mathbf{B}, \mathbf{E})$ and $(\mathbf{H}, \mathbf{D})$ are that $\mathbf{H}$ is a self-conjugate linity of $\mathbf{B}$, (permeability) $^{-1}$; and $\mathbf{D}$ a similar function of $\mathbf{E}$, capacity. In our case

$$\left. \begin{aligned} \dot{\omega} &= \psi \omega^\times, & \psi &= \psi' =)(\psi \\ I' \psi N &= 0 = N' \psi I \end{aligned} \right\} \quad (6)$$

Here ψ is a ${}_n V_2$ linity concerning which, $\psi = \psi'$ asserts that ψ is self-conjugate, $\psi =)(\psi$ asserts that it is contraexco, the dot implies that it is a density, and the two equations in the second line assert that ψ consists of two parts $N' \psi N + I' \psi I$, of which the first is confined to normal components with reference to $\mathfrak{t}$, and the other to incident components.

[Any c/c , say ϕ , may be expressed uniquely as the sum of four parts thus

$$\phi = (N + I) \phi (N + I) = N \phi N + I \phi I + N \phi I + I \phi N.$$

With a $\mathfrak{O}/c$ the four parts are obtained from $(N + I)'$ () $(N + I)$, and in the case of ψ two of these parts are asserted in (6) to be zero.

Digression on I and N .—I have forgotten to point out earlier that I' , I , N' and N all have subextensive significance, and that $(I - N)$ also has a close connection with extensive interpretation. In the neutral form we virtually deal with a Galilean manifold. If in such a manifold I , N are the incident and normal components with reference to a , we can, by a rotation, bring a to coincidence with l_1 , or some l_a , and then $(I - N)$ becomes the capital linity P_1 (which reverses the sign of l_1 in every term of q), introduced at the outset of our subject. The properties of I , N , $(I - N)$, may be based on this connection

with P_1 , through neutral to invariant forms. However, it is perfectly easy to prove directly that, say, I' , has subextensive significance, for we have defined thus

$$I'\beta = \alpha V_0 \eta \alpha \beta, \quad I'u = V_{a\alpha} V_{a-1} \eta \alpha u,$$

and to show that I' has subextensive significance we have to prove that

$$I'u = -V_a I' \xi V_{a-1} \xi u,$$

which is obvious as soon as stated.]

Maxwell's scheme also connects $\mathbf{K}$ with $\mathbf{E}$ by the self-conjugate linity conductivity. In our case express this first with Galilean co-ordinates. We have

$$\sum_{a=1}^3 E_a l_a = V_1 l_4 \omega^\times = R N_4 \dot{\mathbf{K}}$$

where R is put for resistivity. This generalizes to

$$N' \dot{\mathbf{K}} = \dot{\phi} V_1 I \omega^\times, \quad \dot{\phi} = N' \dot{\phi} N = \dot{\phi}' =)(\dot{\phi}. \quad (7)$$

where $\dot{\phi}$ is conductivity, a vector linity.

It appears to me an impressive fact that Maxwell's analysis of the electromagnetic field was so wonderfully true to the nature of things, that not one single feature of his scheme, so far considered, requires the slightest revision when examined with the powerful searchlight of Relativity. Only when the structure perforce comes into contact with Newtonian mechanics, as is the case in what we are about to consider, does it require, along with classical mechanics, to be amended.

For stress put

$$\dot{M} \alpha^\times = -\frac{1}{2} \alpha^\times V_0 \dot{\omega} \omega^\times + V_1 (V_1 \alpha^\times \dot{\omega}') \omega^\times. \quad (8)$$

We have

$$\dot{M}_9 \nabla_9 = -\frac{1}{2} \nabla V_0 \dot{\omega}' \omega^\times + V_1 (V_1 \nabla_9 \dot{\omega}') \omega_9^\times + v^\times$$

by (2a) and (3a). Now

$$V_1 (V_1 \nabla_9 \dot{\omega}') \omega_9^\times = \nabla_9 V_0 \dot{\omega}' \omega_9^\times - V_1 \dot{\omega}' V_3 \nabla \omega^\times$$

of which the last term is zero by (4). Hence

$$\left. \begin{aligned} v^\times &= \dot{M}_9 \nabla_9 + \frac{1}{2} (\nabla_8 - \nabla_9) V_0 \dot{\omega}'_8 \omega_9^\times \\ &= \dot{M}_9 \nabla_9 + \frac{1}{2} \nabla_9 V_0 \omega^\times \psi_9^{-1} \omega^\times \end{aligned} \right\} \quad (9)$$

by (5). By using $\dot{\omega}$ instead of $\omega^\times$ the last term on the right becomes $-\frac{1}{2} \nabla_9 V_0 \omega^\times \psi_9^{-1} \omega^\times$. Because of the factor l^{-1} in $\psi^{-1} = l^{-1} \psi^{-1}$ we may speak of ψ^{-1} as an *anti*-density.

§ 21. *STATIONARY ACTION IN A RIEMANN MANIFOLD.*—In our theorems of integration $d\dot{p}_c$ being equal to $V_c dp_1 dp_2 \dots dp_c$ is contravariant. But since

$$l db = l V_0 d\alpha dp_n$$

is an invariant, $d\alpha$, must be a covariant vector anti-density, and, similarly, $d\dot{p}_c$ in general (including db) is a covariant multienion anti-density.

(8) and (9) of §20 are unsatisfactory in that $\dot{M}\theta$ is not apparently self-conjugate, and from the second term on the right of (9) the force (cum power), per unit volume, cannot be expressed as a stress even of the non self-conjugate kind. But the pioneers of Relativity have developed a principle of stationary action which has for consequence that all of the equations of §20 may be retained, and that $\dot{M}$ of (8) *is* self-conjugate; and by what seems little more than a bold assumption as to the nature of things, they conclude that the intractable last term of (9) need not trouble us, because force generated by the field need not be begotten by stress; they give us something else instead, as will appear below. [I think I have here credited the pioneers with just a little more than they have set forth, but no more than such as, by small additions, can be included.]

Although in the main we copy their procedure, yet in two important respects we depart from it, and in doing so return much more nearly than they do to Maxwell's outlook. First, we keep in mind the application to matter in bulk; and, secondly, we abjure the use of λ *in toto*. Probably Heaviside, among those who studied these matters in the days before the advent of Relativity, as we know it to-day, emphasised as much as any writer the view that the potentials of λ have nothing to do with actualities, unless among such are to be reckoned the imperfections of our mathematics. In returning to the procedure I adopted in 1892,⁴ of taking Maxwell's $\mathbf{H}$ and $\mathbf{D}$ as the fundamental (three-dimensional) vectors of the field, instead of any of $\mathbf{A}$, $\mathbf{B}$, $\mathbf{C}$, $\mathbf{K}$, or $\mathbf{E}$, I find that the potentials can be completely ignored, that they are not wanted even as a mathematical crutch, and that the ignoring of them considerably simplifies the preliminary organising plans of our exploration. The electro-magnetic terms are all dependent on a single arbitrary scalar function of the ${}_nV_2$ called $\dot{\omega}$ in §20.

Let us recall that a particle of matter, with three dimensional extension, occupies a four-dimensional tube in the space-time continuum. The strength of this tube is the invariant constant (static) mass $d\dot{M} = l dM$ of the particle. The three dimensional anti-density covariant cross-section being $d\alpha$, an element of "length" of the tube being the proper time ds , corresponding to the contravariant vector dp , where $d\alpha$ and dp are both time-like, ι being the unit tangent vector dp/ds , db the four-dimensional volume of the element, we have

$$ds^2 = V_0 dp \theta dp, \quad \iota = dp/ds, \quad db = V_0 d\alpha dp. \quad (1)$$

and

$$d\dot{M} \cdot ds = m db, \quad (2)$$

⁴"On the Mathematical Theory of Electromagnetism," 'Phil. Trans.,' 1893.

where $\dot{m}$ or sometimes m is called the material density. The momentum is

$$\iota d\dot{M} = \iota \dot{m} (db/ds), \quad (3)$$

and

$$\iota \dot{m} = \dot{\mu}, \quad (4)$$

is called the momentum per unit volume. It is this vector whose magnitude multiplied by the cross-section yields the constant invariant strength, the quantity $d\dot{M}$. [In invariants we should say $m \times (ldb \div ds) = d\dot{M}$; so that perhaps $m\iota = \dot{\mu}$ rather than $\dot{\mu}$ should be called the momentum per unit volume.] We may note from the integration theorems, first for a finite portion of a tube, and next for a finite region of four dimensions, that

$$V_0 \nabla \dot{\mu} = 0. \quad (5)$$

This “equation of continuity” or “equation of conservation” of matter is given by Weyl, but seems nowhere to appear in Eddington’s Report.⁵

Our fundamental view of electricity is the same in principle, but we do not need to elaborate a notation such as $d\mathbf{E}^\bullet$, $e^\bullet$, etc., in connection with $\dot{\kappa}$ to serve the purposes of $d\dot{M}$, $\dot{m}$, etc., in connection with $\dot{\mu}$. The tubes in the case of electricity (strength $V_0 \dot{\kappa}' d\alpha$, just as strength of momentum tube $= V_0 \dot{\mu}' d\alpha$) are called current tubes. If we were dealing with the smallest portions of matter it would accord with the prevalent views of to-day to suppose the two sets of tubes to be identical, but for matter in bulk we must consider them to be independent. Fortunately this is without any influence on the deductions we have to make from our fundamental principles.

Our principle of stationary action is that

$$\delta \int^n W db = 0, \quad (6)$$

where W is an invariant scalar density (meaning that $l^{-1}W$ is invariant) function of position consisting of three parts

$$W = \dot{m} + f(\dot{\omega}, \theta) + l(c - \frac{1}{2}G), \quad (7)$$

the first, material, being $\dot{m}$ already described and equal to $\sqrt{(V_0 \dot{\mu} \theta \dot{\mu})}$; the second, electrical, a function $f(\dot{\omega}, \theta)$ of a ${}_n V_2$, $\dot{\omega}$, identified with Maxwell’s $\mathbf{H}$ and $\mathbf{D}$ as in §20, and of θ the fundamental linity of the manifold; and the third, gravitational, $l(c - \frac{1}{2}G)$ where G is the scalar curvature $-V_0 \zeta C \zeta$, the cosmological constant c possibly being zero, though Einstein teaches that it *must* differ from zero.

The variations implied by δ in (6) are of three independent kinds; (1) an arbitrary increment in θ

$$\delta \theta = \theta_0, \quad (8)$$

⁵For a more critical examination of equation (5) see the later paper by the author: “The Mechanical Forces ... Field” referred to above, footnote p. 164.

at every point of the manifold; (2) an arbitrary increment in $\dot{\omega}$

$$\delta\dot{\omega} = V_2\nabla v^\bullet \quad (v^\bullet = V_3w^\bullet) \quad (9)$$

at every point, of such a kind that the current tubes, that is to say $\dot{\kappa}$, which last is *defined* by

$$\dot{\kappa} = V_1\nabla\dot{\omega} \quad (10)$$

remain unvaried; and (3) an arbitrary displacement which shifts the tubes, momentum and current, together,

$$\delta p = \varepsilon \quad (11)$$

without alteration of their strengths, $V_0\dot{\mu}'d\alpha$, $V_0\dot{\kappa}'d\alpha$.

[*Added October, 1921.*—A consequence of remarks which occur below is so obvious that it is singular that only now have I happened to recognise it. If it is assumed that for the first two variations ($\delta\theta = \theta_0$ and $\delta\dot{\omega} = V_2\nabla v^\bullet$) the stationary condition (6) holds; then *it follows as a mathematical deduction* that the stationary condition also holds for the third variation ($\delta p = \varepsilon$) which shifts the tubes. Clearly then our postulated principle of stationary action should not include the *assumption* of the stationary condition for the third variation. The reader will easily justify the assertions that have just been made, if he will take as the proof of the equation $T_\xi^{(t)}\zeta = 0$, the indication of a proof, immediately following the identity (35) below. This equation $T_\xi^{(t)}\zeta = 0$ is virtually the necessary and sufficient condition for the existence of the stationary condition under the variation $\delta p = \varepsilon$.]

The integral of (6) extends over any arbitrary finite region at the boundary of which θ_0 , $\dot{\omega}$ and ε all vanish.

$\omega^\times$, eventually to be identified with Maxwell's $\mathbf{B}$ and $\mathbf{E}$ as in §20, is *defined* by

$$\omega^\times = -\dot{\omega}\nabla \quad (12)$$

where $\dot{\omega}\nabla$, as in Quaternions, means the differential operator for which $df = -V_0d\dot{\omega}\dot{\omega}\nabla f$, df being the increment in f due to an arbitrary increment $d\dot{\omega}$ in $\dot{\omega}$.

It is necessary to point out that if W consisted of $\dot{m}$ alone, *the condition* $\delta W = 0$ *would mean that a momentum tube is geodesic* because by (2) $\dot{m}db = d\dot{M}ds$, and by hypothesis $d\dot{M}$, the strength, is unaffected by the displacement ε of the tube. Indeed, following Weyl, we might use this identification to find the effects of ε and θ_0 on the variation of $\dot{m}$ in (6), but we pursue another course. It seems almost unnecessary to say that for a full treatment of matter in bulk W as above described is by no means adequate;

there are several scalar functions of position such as temperature and strain co-ordinates (adapted to the method of Relativity) that should be included in W for such a treatment.⁶ We may accept the inclusion of fluid and elastic solid stresses that Eddington so lucidly justifies in §37 of his report, despite the fact that his appeal to the kinetic theory of matter would by a precisionist be described as a departure from our elected task of dealing with matter in bulk.

We shall consider the effects of $w^\bullet$, ε and θ_0 in succession because that is the order of simplicity; θ_0 , $w^\bullet$ and ε all take part in the variation of f ; θ_0 and ε in that of $\dot{m}$; and θ_0 alone in that of G .

The variation due to $\dot{\omega}$ gives by (9) and (12)

$$0 = - \int \int V_0 \omega^\times \nabla w^\bullet db = - \int^{n-1} \int V_0 w^\bullet \omega^\times d\alpha + \int \int V_0 w^\bullet V_3 \nabla \omega^\times db,$$

whence, since $w^\bullet$ is zero at the boundary,

$$V_3 \nabla \omega^\times = 0. \quad (13)$$

The process, here exhibited, of making a ∇ , explicitly occurring in the integrand, apply to the whole integrand, and the consequent removal of the resulting expression (or the potentiality of such removal) to the boundary, is known as integration by parts.

The effect of ε we will first consider with reference to the current tubes. For the strength of every tube to remain unchanged by the displacement it suffices that $\dot{\omega}$ changes in precisely the manner that it changed in §15 when ε indicated a change of co-ordinates. Quoting then from §15, we first have

$$l^{-1} \delta' l = -\frac{1}{2} V_0 \xi \delta \theta \theta^{-1} \xi = V_0 \xi \chi_0 \xi = V_0 \nabla \varepsilon,$$

and next

$$\begin{aligned} \delta \dot{\omega} &= \dot{\omega} V_0 \nabla \varepsilon - V_2 \chi_0 \xi V_1 \xi \dot{\omega} \\ &= \dot{\omega} V_0 \nabla \varepsilon - V_2 \varepsilon_9 V_1 \nabla_9 \dot{\omega} = V_2 \nabla_9 V_3 \dot{\omega}' \varepsilon_9. \end{aligned}$$

For the future we shall write δ' for these changes to be quoted from §15 leaving δ for a second meaning thus (Weyl's notation)

$$\left. \begin{aligned} \delta'(\) &= (\text{value at } p + \varepsilon \text{ after displacement}) - (\text{value at } p \text{ before}) \\ \delta(\) &= (\text{value at } p \text{ after displacement}) - (\text{value at } p \text{ before,}) \end{aligned} \right\} \quad (14)$$

so that

$$\begin{aligned} \delta(\) &= \delta'(\) + V_0 \varepsilon \nabla \cdot (\), \\ \delta \dot{\omega} &= V_2 \nabla_9 V_3 \dot{\omega}' \varepsilon_9 + V_0 \varepsilon \nabla \cdot \dot{\omega}, \\ \delta f &= V_0 \omega_9^\times \delta \dot{\omega} = V_0 \omega_9^\times \nabla_9 V_3 \dot{\omega}' \varepsilon_9 - V_0 \cdot V_3 \dot{\omega}^\times \nabla_9 V_3 \dot{\omega}' \varepsilon + V_0 \varepsilon \nabla_9 V_0 \omega^\times \omega_9', \end{aligned}$$

because $V_3 \nabla \omega^\times = 0$. Thus $\delta f -$ (first term on right)

$$= V_0 \varepsilon V_1 \dot{\omega}' \omega^\times.$$

Hence

$$(\varepsilon \text{ part}) \quad \int \int \delta f db = \int \int V_0 \varepsilon V_1 \dot{\omega}' \omega^\times db. \quad (15)$$

⁶Treated in a series of three papers by the author; despatched within the past month (December, 1922) to England.

Next calculate the corresponding part from $\delta\dot{m}$, in which $\delta\dot{\mu}$ is obtained, as with $\delta\dot{\omega}$ above, but $\delta\theta = 0$. We shall for brevity use

$$\dot{m}^2 = V_0\dot{\mu}\theta\dot{\mu}, \quad m\iota = \dot{\mu},$$

but the reader would do well at the beginning to use only $\dot{m} = \sqrt{(V_0\dot{\mu}\theta\dot{\mu})}$ and not ι .

$$\delta\dot{m} = \frac{1}{2}\delta(\dot{m}^2)/\dot{m} = V_0\iota\theta(-\varepsilon_9V_0\dot{\mu}\nabla_9 + \dot{\mu}V_0\nabla\varepsilon + V_0\varepsilon\nabla\cdot\dot{\mu}).$$

Here prepare for integration by parts by altering the incidence of ∇ in the first two terms on the right, so as to apply to the whole term, and reject this part of each term. Thus $\delta\dot{m}$ — (the rejected parts)

$$\begin{aligned} &= V_0\varepsilon[(\theta\iota)_9V_0\dot{\mu}_9\nabla_9 - \nabla\dot{m} + \nabla_9V_0\dot{\mu}_9\theta\iota] \\ &= V_0\varepsilon(T_9\nabla_9 + \frac{1}{2}\nabla_9V_0\zeta\theta_9\theta^{-1}T\zeta) \end{aligned}$$

where

$$T^\bullet\alpha^\times = lT\alpha^\times = m\theta\iota V_0\alpha^\times\iota \quad [+ \text{ stress terms}] \quad (16)$$

(the stress terms meaning those inserted here from Eddington's argument in §37 of the Report). Thus

$$(\varepsilon \text{ part}) \quad \int \int \delta\dot{m}db = \int \int lV_0\varepsilon T_{(\zeta)}\zeta db \quad (17)$$

by equation (19) of §15.

T or $T^\bullet$ is called the material kinetic energy linity, or if the stress terms (say $T^{(s)}$) are included, it may be called the material stress-energy linity. The stress terms are self-conjugate and satisfy the equation $T^{(s)} = NT^{(s)}N$.

(15) and (17) are the only contributions of ε to (6), so that we have

$$lT_{(\zeta)}\zeta + V_1\dot{\kappa}'\omega^\times = 0. \quad (18)$$

We have now fully identified $\omega^\times$ with Maxwell's $\mathbf{B}$ and $\mathbf{E}$.

Of equations (1a), (2a), (3a), (4), (5), (6) of §20 we have decided that (1a) is quite unnecessary, is in fact better neglected than cared for (though, of course, it is a mere consequence of our present equation (13)). The other five are replaced in the present section by (10), (18), (13), (10), (12) respectively. [(10) does duty twice, since it has the obvious consequence $V_0\nabla\dot{\kappa} = 0$.]

We are now about to trace the consequences flowing from the fact that $l^{-1}f$ is an invariant. These are wholly satisfactory for empty space, but not for matter in bulk, for which, in accordance with Maxwell's principles, we must modify our view as to the nature of f , the electrical term of the action density. It is well to show, however, that it is this invariance of $l^{-1}f$ which causes the inadequacy of our present supposition that f is a function of $\dot{\omega}$ and θ only.

With our present (vacuum) assumption, we shall show that (18) can be added to, thus ($\kappa = l^{-1}\dot{\kappa}$)

$$-T_{(\zeta)}\zeta = V_1\kappa\omega^\times = 2(l^{-1}(\theta\mathbf{Q}f)_{(\zeta)}\zeta). \quad (19)$$

where, as in Quaternions, $\theta \mathbf{Q}$ means the self-conjugate linity differential operator given by

$$df = -V_0 d\theta \zeta \theta \mathbf{Q} f \zeta \quad (20)$$

where df is the increment in f due to the arbitrary increment $d\theta$ in θ . It is, therefore, not the linity $T^\bullet$, but, instead, the linity $T^{\bullet(t)}$ ($=$ total energy linity $= lT^{(t)}$) given by

$$T^{\bullet(t)} = T^\bullet + 2\theta \mathbf{Q} f \quad (21)$$

which may be supposed to be handed on through space by some kind of contact action, because from (19)

$$T_{(\zeta)}^{(t)} \zeta = 0. \quad (22)$$

As a matter of fact, we shall later, when considering gravitation, show that this equation can be transformed to the shape

$$\phi_9 \nabla_9 = 0.$$

We will now prove (19) on our present hypothesis [f as in (7)]. Let us, for brevity, use $v^\times$ as in (3a), §20; that is, we define $v^\times$ by

$$v^\times = V_1 \dot{\kappa}' \omega^\times = V_1 (V_1 \nabla \dot{\omega}') \omega^\times. \quad (23)$$

Since $l^{-1}f$ is invariant, and since, as we saw above, before equation (14), $l^{-1}\delta' l = V_0 \nabla \varepsilon$, where ε now, as in §15, expresses an infinitesimal change of co-ordinates, we have as with $\delta \dot{\omega}$ above in the equation following (14)

$$f V_0 \nabla \varepsilon_9 + V_0 \varepsilon \nabla f = \delta f = V_0 \omega^\times \delta \dot{\omega}' - V_0 \delta \theta \zeta \theta \mathbf{Q} f \zeta.$$

$\delta \dot{\omega}'$ is given in the equation following (14) and $\delta' \theta$ is given in §15 as $-\chi'_0 \theta - \theta \chi'_0$. Hence

$$\begin{aligned} f V_0 \nabla \varepsilon_9 + V_0 \varepsilon \nabla f &= V_0 \omega_9^\times (V_2 \nabla_9 V_3 \dot{\omega}' \varepsilon_9 + V_0 \varepsilon \nabla \cdot \dot{\omega}') \\ &+ V_0 (2\varepsilon_9 \theta \mathbf{Q} f \nabla_9 - \varepsilon \theta \mathbf{Q} f \zeta \theta_9 \zeta \cdot V_0 \varepsilon \nabla_9). \end{aligned} \quad (24)$$

Here ε and all the n^2 derivatives $(\nabla_9, \varepsilon_9)$ of ε are arbitrary, so that we may put arbitrary dummies $\gamma, \alpha^\times, \beta$ for $\varepsilon, \nabla_9, \varepsilon_9$ respectively. From the dummy γ we get

$$\nabla f = \nabla_9 V_0 \dot{\omega}'_9 \omega^\times - \nabla_9 V_0 \theta_9 \zeta \theta \mathbf{Q} f \zeta.$$

This identity, satisfied by f , does not depend on the invariance of f . It is easily seen to be a direct consequence of the assumption that f is a function of $\dot{\omega}$ and θ . A useful result is obtained from it by modifying the first term on the right, thus

$$\begin{aligned} \nabla_9 V_0 \dot{\omega}'_9 \omega^\times &= V_1 (V_3 \omega^\times \nabla_9) \dot{\omega}'_9 + v^\times & [\text{eq. (23)}] \\ &= V_1 (V_3 \omega_9^\times \nabla_9) \dot{\omega}'_9 + v^\times & [\text{eq. (13)}] \\ &= (-\dot{M}_9 \nabla_9 + \tfrac{1}{2} \nabla V_0 \dot{\omega}' \omega^\times) + v^\times & [\text{eq. (8) §20}], \end{aligned}$$

whence

$$\nabla f = -\dot{M}_9 \nabla_9 + \tfrac{1}{2} \nabla V_0 \dot{\omega}' \omega^\times + v^\times - \nabla_9 V_0 \theta_9 \zeta \theta \mathbf{Q} f \zeta. \quad (25)$$

From the dummy β above in (24)

$$\left. \begin{aligned} f\alpha^\times &= V_1(V_3\omega^\times\alpha^\times)\dot{\omega} + 2_\theta\mathbf{D}f\alpha^\times, \\ f &= -\dot{M} + \frac{1}{2}V_0\dot{\omega}\omega^\times + 2_\theta\mathbf{D}f \end{aligned} \right\} \quad (26)$$

This is an identity (remember $\omega^\times$ is defined as $-\omega \cdot \nabla f$) which we have proved f to satisfy by reason of the invariance of $l^{-1}f$. Putting $\alpha^\times$ in (26) = ∇_9 and subtracting (25), we get (19).

(26) shows that $\theta^{-1}\dot{M}$ is self-conjugate, which the reader will at first be inclined to claim as a highly satisfactory result. I am not inclined to dispute this so far as empty space is concerned, but for matter in bulk it is precisely what we do *not* desire. If $\theta^{-1}\dot{M}$ is self-conjugate for matter in bulk, then crystalline transmission of light cannot be explained on Maxwell's theory.

To prove this, choose co-ordinates such that $\iota = \iota_4$; this requires co-ordinates which at the point under consideration reduce the medium to rest. Examine the case when the co-ordinates are Galilean, and Maxwell's equations become (1a), (2a), (3a) of §20. To translate $\dot{M}\alpha^\times$ into quaternion form, identify the multenion $\iota_2\iota_3, \iota_3\iota_1, \iota_1\iota_2$ with the quaternion i, j, k and the multenion

$$\iota_1\iota_2\iota_3\iota_4 = \mathbf{v}$$

with Hamilton's $\sqrt{-1}$ in his bi-quaternion $p + q\sqrt{-1}$. Thus $\dot{\omega} = \mathbf{H} + \mathbf{vD}$ and $\omega^\times = \mathbf{B} + \mathbf{vE}$. Indicate a ${}_nV_1$ such as $\alpha^\times$ by (σ, s_4) where

$$\alpha^\times = \sum s_c \iota_c \quad (c = 1, 2, 3, 4), \quad \sigma = s_1i + s_2j + s_3k,$$

and indicate a ${}_nV_3$ such as $\mathbf{v}\alpha^\times$ by $\mathbf{v}(\sigma, s_4) = (\mathbf{v}\sigma, \mathbf{v}s_4)$. Thus [compare (3) and (3a) of §20]

$$\begin{aligned} V_1\alpha^\times\dot{\omega} &= V_1(\sigma, s_4)(\mathbf{H} + \mathbf{vD}) \\ &= (V\sigma\mathbf{H} + s_4\mathbf{D}, -S\sigma\mathbf{D}) = (\text{say})(\sigma', s'_4), \\ V_1(V_1\alpha^\times\dot{\omega})\omega^\times &= (V\sigma'\mathbf{B} + s'_4\mathbf{E}, -S\sigma'\mathbf{E}) \\ &= (V[V\sigma\mathbf{H} + s_4\mathbf{D}]\mathbf{B} - \mathbf{E}S\sigma\mathbf{D}, -S\mathbf{E}[\sigma\mathbf{H} + s_4\mathbf{D}]). \end{aligned}$$

adding $-\frac{1}{2}\alpha^\times V_0\dot{\omega}\omega^\times$ and making a slight quaternion reduction we obtain

$$\dot{M}\alpha^\times = \left(-\frac{1}{2}V[\mathbf{H}\sigma\mathbf{B} + \mathbf{E}\sigma\mathbf{D}] + \frac{1}{2}V\sigma V[\mathbf{HB} + \mathbf{ED}] + s_4V\mathbf{DB}, \quad S\sigma\mathbf{EH} - \frac{1}{2}s_4S[\mathbf{BH} + \mathbf{ED}]\right).$$

In the square array of sixteen scalars representing the linity $\dot{M}$ the fourth column contains the components of $V\mathbf{DB}$, the fourth row contains those of $-\mathbf{VEH}$, and the constituent in the fourth row and fourth column is $-\frac{1}{2}S(\mathbf{BH} + \mathbf{ED})$. The other nine scalars are those of the vector linity of σ in the expression for $\dot{M}\alpha^\times$ just written. Thus the conditions that $\theta^{-1}\dot{M}$ shall be self-conjugate are that

$$V\mathbf{DB} = \mathbf{VEH}, \quad V\mathbf{HB} = V\mathbf{DE}.$$

[These, of course, amount to saying that $\mathbf{D}$ is parallel to $\mathbf{E}$ and $\mathbf{B}$ to $\mathbf{H}$.]

Now in a plane light wave $\mathbf{VEH}$ is parallel to the ray and $\mathbf{VDB}$ is normal to the wave front, so that to say that $\mathbf{VDB} = \mathbf{VEH}$ is to deny the possibility of double and conical refraction.

Our assumption that f is a function of $\dot{\omega}$ and θ only is contrary to Maxwell's, which asserts that N and I enter into the relations between $\dot{\omega}$ and $\omega^\times$, that is ι , as well as $\dot{\omega}$ and θ is a constituent variable of f . We may take f to be an explicit function of $\dot{\omega}$, $\dot{\phi} = I\dot{\omega}$ and θ . Its general increment df , may be written

$$df = -V_0 d\dot{\omega} \cdot (\dot{\omega} \cdot \nabla) f - V_0 d\dot{\phi} \cdot (\dot{\phi} \cdot \nabla) f - V_0 d\theta \zeta (\theta \mathbf{Q}) f \zeta.$$

where $(\dot{\omega} \cdot \nabla)$, $(\dot{\phi} \cdot \nabla)$ and $(\theta \mathbf{Q})$ are all *partial* differential coefficient operators, whereas $\dot{\omega} \cdot \nabla$ and $\theta \mathbf{Q}$ by contrast are *total* differential operators. Thus $\dot{\omega} \cdot \nabla f$ comes partly from $(\dot{\omega} \cdot \nabla) f$ and partly from $(\dot{\phi} \cdot \nabla) f$, and again $\theta \mathbf{Q} f$ comes partly from $(\theta \mathbf{Q}) f$ and partly from $(\dot{\phi} \cdot \nabla) f$ (because θ is involved in $\dot{\phi}$). However, there is a part of $-V_0 d\dot{\omega} \cdot (\dot{\omega} \cdot \nabla) f$, involving $d\iota$, still standing over.

[In Maxwell's own case $(\dot{\phi} \cdot \nabla) f = (-I + N)\omega^\times$ and we might assume this equation to hold in general, though this is not done below.] Writing this outstanding part as $-V_0 d\iota \nabla f$ we shall have

$$df = -V_0 d\iota \nabla f - V_0 d\dot{\omega} \cdot \dot{\omega} \cdot \nabla f - V_0 d\theta \zeta \theta \mathbf{Q} f \zeta.$$

Thus using $d\dot{\omega}'$, for the moment, to stand for the increment of $\dot{\omega}$ depending on $d\iota$ we have

$$\dot{\omega}' = V_2 \iota V_1 \theta \iota \dot{\omega}',$$

$$d\dot{\omega}' = V_2 d\iota V_1 \theta \iota \dot{\omega}' + V_2 \iota V_1 \theta d\iota \dot{\omega}',$$

$$-V_0 d\iota \nabla f = df = -V_0 d\iota \zeta \{ V_1 [V_1 \theta \iota \dot{\omega}'] (\dot{\phi} \cdot \nabla) f + \theta V_1 \dot{\omega}' V_1 (\dot{\phi} \cdot \nabla) f \zeta \}.$$

Thus we may understand ∇f to mean as follows

$$\left. \begin{aligned} \nabla f &= \xi^\bullet \iota = N \{ V_1 [V_1 \theta \iota \dot{\omega}] (\psi^\bullet \cdot \nabla) f + \theta V_1 \dot{\omega} V_1 (\psi^\bullet \cdot \nabla) f \iota \}, \\ \text{where } \xi^\bullet \alpha &= N(\psi + \psi') \alpha, \quad \psi \alpha = \theta V_1 \dot{\omega} V_1 (\psi^\bullet \cdot \nabla) f \alpha, \\ \text{and } \xi^\bullet \iota &= ((\xi^\bullet \iota)). \end{aligned} \right\} \quad (23a)$$

[*Correction added September, 1921.*— N was absent from equation (23a) as given in the original paper. If N is omitted it is not necessarily true, but if N is included it is necessarily true, that $\xi^\bullet \iota = ((\xi^\bullet \iota))$. In the original presentation it was tacitly, and erroneously, assumed that $(\dot{\omega} \cdot \nabla) f$, $(\psi^\bullet \cdot \nabla) f$ and $(\theta \mathbf{Q}) f$ are always invariantive. They will doubtless be so when a proper invariantive form of f has been selected, and it is desirable to make such a selection when possible and convenient. We will now show that, whether such a selection has been made or not, ∇f , $\dot{\omega} \cdot \nabla f$ and $\theta \mathbf{Q} f$ are always invariantive, that is to say ∇f is a covariant vector density, $\dot{\omega} \cdot \nabla f$ is a covariant ${}_n V_2$, and $\theta \mathbf{Q} f$ is a

contraexco vector linity density. Let us put $\tau = x\iota$ where x is an arbitrary invariant scalar function of position which we shall *eventually* take to be $x = 1$. Thus

$$f = f(\dot{\omega}, \psi, \theta) = f(\dot{\omega}, \Gamma' \dot{\omega}, \theta) \quad (\text{A})$$

Γ' being that definite explicit function of τ and θ which is given by ($\Omega =$ a contravariant ${}_nV_2$ dummy)

$$\Gamma' \Omega = V_2 \tau V_1 \theta \tau \Omega / V_0 \tau \theta \tau. \quad (\text{B})$$

From this last $d\Gamma'$ can be expressed in terms of $d\theta$ and $d\tau$ (and it may be noticed in passing that since Γ' is homogeneous and of degree zero both in θ and in τ we have by Euler's theorem for homogeneous functions

$$V_0 \tau \nabla_9 \cdot \Gamma'_9 = 0 = V_0 \theta \zeta_{\theta \theta} \Omega_9 \cdot \Gamma'_9. \quad (\text{C})$$

the first of these explaining the presence of N in (23a), for it leads to $V_0 \tau \nabla f = 0$). We now have

$$\left. \begin{aligned} df &= -V_0 d\dot{\omega} \cdot (\dot{\omega} \cdot \nabla) f - V_0 d\psi \cdot (\psi^\bullet \cdot \nabla) f - V_0 d\theta \zeta (\theta \Omega f) \zeta, \\ &= -V_0 d\dot{\omega} \dot{\omega} \cdot \nabla f - V_0 d\tau \nabla f - V_0 d\theta \zeta \theta \Omega f \zeta \end{aligned} \right\} \quad (\text{D})$$

the second being derived from the first by finding $d\psi$ from

$$d\psi = \Gamma' d\dot{\omega} + d\Gamma' \cdot \dot{\omega},$$

$d\Gamma'$ coming from (B). In the second of (D) $d\dot{\omega}$, $d\tau$ and $d\theta$ are all *independently, arbitrary and invariantive* and therefore their coefficients,

$$\dot{\omega} \cdot \nabla f, \quad \tau \nabla f, \quad \theta \Omega f,$$

are *without ambiguity and are invariantive*. These last important deductions have been made possible by using τ instead of ι . τ (and therefore $d\tau$) is a general contravariant vector whereas ι is restricted by the equation $V_0 \iota \theta \iota = 1$.

From the above we obtain by straightforward filling in of detail that

$$x_\tau \nabla f = (1 - I)(\psi - \psi') \iota = \xi^\bullet \iota$$

as given in equation (23a) above. Since $x_\tau \nabla f$ is independent in meaning of x , it is natural as in (23a) to write it in the form ∇f .

I take this opportunity of suggesting what is the proper form of f *in vacuo* and inside matter. *In vacuo* we have

$$f = V_0 \dot{\omega} \theta^{-1} \dot{\omega} = l^{-1} V_0 \omega^\times \theta \omega^\times.$$

Now $\theta = N\theta N' + I\theta I'$ so that

$$lf = V_0 (N' \omega^\times) \theta (N' \omega^\times) + V_0 (I' \omega^\times) \theta (I' \omega^\times).$$

I would suggest that inside matter

$$lf = V_0 (N' \omega^\times) \xi (N' \omega^\times) + V_0 (I' \omega^\times) \xi (I' \omega^\times) \quad (\text{E})$$

where ξ is a function of θ and of position (p) only. Thus

$$lf = V_0 \omega^\times (N \xi N' + I \xi I') \omega^\times. \quad (\text{F})$$

Without loss of generality ξ may be replaced first by its self-conjugate part $\frac{1}{2}(\xi + \xi')$ since (E) is thereby unchanged, and next by the part

$$\frac{1}{2}[N(\xi + \xi')N' + I(\xi + \xi')I']$$

since (F) is thereby unchanged. With $n = 4$ the general ξ would involve 36 scalars, but the part we have finally reached involves only 12 scalars, namely 6 for $N(\xi + \xi')N'$ and 6 for $I(\xi + \xi')I'$. We may, as a special case, take $\xi = A\theta A'$ where A is a coexco linity function of position.]

We have to add to the right of (24), (25), and (26) the following marked (24a), (25a), (26a) respectively:

$$V_0 \varepsilon \xi^\bullet \iota \cdot V_0 \iota \nabla_9 - V_0 \varepsilon \nabla_8 \cdot V_0 \iota_8 \xi^\bullet \iota, \quad (24a)$$

$$-\nabla_8 \cdot V_0 \iota_8 \xi^\bullet \iota, \quad (25a)$$

$$\xi^\bullet \iota \cdot V_0 \iota \alpha^\times. \quad (26a)$$

(15) and (19) also are modified by $V_1 \kappa \omega^\times$ changing to

$$-lT_{(\zeta)}\zeta = V_1 \kappa \omega^\times - \nabla_9 V_0 \iota_9 \xi^\bullet \iota - (\xi^\bullet \iota)_9 V_0 \iota_9 \nabla_9. \quad (15a \text{ and } 19a)$$

I think there can be no doubt⁷ that this amendment is the true expression for the force, cum power, per unit volume. It will be observed that no change occurs in the total energy linity $T^{\bullet(t)} = T^\bullet + 2_\theta \mathbf{C}f$.

Pass now to the variation $\delta\theta = \theta_0$. There is no trouble in supplementing (15) and (17) with the θ_0 parts since no integration by parts is here involved. Thus since $\dot{m} = \sqrt{(V_0 \dot{\mu} \theta \dot{\mu})}$, $\delta\dot{m} = \frac{1}{2}V_0 \iota \theta_0 \dot{\mu}$, and

$$(\theta_0 \text{ part}) \quad \delta(\dot{m} + f) = -\frac{1}{2}V_0 \theta_0 \zeta \theta^{-1} T^{\bullet(t)} \zeta. \quad (27)$$

G and $G^\bullet (= lG)$ are functions of θ and its first and second derivatives (∇), and therefore of $\theta^\bullet$ and its first and second derivatives, where $\theta^\bullet = l\theta^{-1}$. Let $\theta^\bullet$ receive any infinitesimal increment $\delta\theta^\bullet$. At one part of our work we shall assume $\delta\theta^\bullet$ to have the meaning corresponding to $\delta\theta = \theta_0$ of equation (8); at another part $\delta\theta^\bullet$ will mean the actual increment $d\theta^\bullet = -V_0 \alpha \nabla \cdot \theta^\bullet$ of $\theta^\bullet$ on passing from the position p to the position $p + dp = p + \alpha$ in the manifold; on some other occasion it may mean some combination of these; in a word, $\delta\theta^\bullet$ is any conceivable defined infinitesimal increment of $\theta^\bullet$. The following two equations are true:—

$$G^\bullet = -V_0 \Omega \zeta \theta^\bullet \zeta = -V_0 \nabla (F_\zeta \theta^\bullet \zeta - \theta^\bullet F'_\zeta \zeta) - V_0 \zeta \theta^\bullet (F'_{\zeta_1} F'_\zeta - F'_\zeta F'_{\zeta_1}) \zeta_1, \quad (28)$$

$$-V_0 \Omega \zeta \delta\theta^\bullet \zeta = -V_0 \nabla (F_\zeta \delta\theta^\bullet \zeta - \delta\theta^\bullet F'_\zeta \zeta) - \delta \cdot V_0 \zeta \theta^\bullet (F'_{\zeta_1} F'_\zeta - F'_\zeta F'_{\zeta_1}) \zeta_1. \quad (29)$$

The great similarity between these equations is connected with the fact that the second term on the right of (28) is a function homogeneous and

⁷Especially in view of the paper twice referred to above: "The Mechanical Forces ... in an Electromagnetic Field."