

Multenions and Differential Invariants.—III.

By ALEX. MCAULAY, Professor of Mathematics in the University of Tasmania.

(Communicated by W. B. Hardy, Sec.R.S. Received July 20, 1921.)

§ 19. *Suggested Improvement of Some Notations and Terms.*—The reader has probably found that the terminology for multenion linities, adopted hitherto because of the writer's timid reluctance to depart from accepted meanings, is too inexpressive of the intended signification. A covariant linitity fabricates a covariant multenion out of a contravariant. Let us call it a $c/\varnothing$ or a coexcontra (co-ex-contra); and similarly let us now denote a contravariant, comixt or contramixt linitity by the name contraexco, coexco or contraexcontra, and by the symbol $\varnothing/c$, c/c or $\varnothing/\varnothing$ respectively. Let us also use parenthesis marks* before a letter such as ϕ , of the same general shapes as c and $\varnothing$ (omitting the solidus), to indicate the type of invariance, thus—let us say that

$$\phi = {}^{\circ}\phi, {}^{\circ}\phi, {}^{\circ}\phi \text{ or } {}^{\circ}\phi$$

indicates that ϕ is a

$$c/\varnothing, \varnothing/c, c/c \text{ or } \varnothing/\varnothing$$

respectively. Similarly, $p = {}^{\circ}p$ indicates that p is a covariant multenion and $q = {}^{\circ}q$ indicates that q is a contravariant multenion. If we actually insert all these indicating parenthesis marks, which we shall do but rarely, their mode of combination is simple. Thus

$${}^{\circ}p = {}^{\circ}\phi {}^{\circ}\psi {}^{\circ}q$$

is an invariant equation. The features that render it so are (1) that the second mark of one letter, ϕ or ψ , is identical with the first mark of the next; these two marks virtually cancel one another; and (2) the proper mark of the resultant symbol p , of the whole operation, is the first written mark of the compound symbol $\phi\psi q$.

When with these notations we use the dash which indicates conjugacy, there is, unless we insert brackets, ambiguity in the meanings of ${}^{\circ}F'$ and ${}^{\circ}F'$ (the two mixed types as we have hitherto called them), because $({}^{\circ}F)'$ is not of the same type c/c as ${}^{\circ}F$, but of the type $\varnothing/\varnothing$. There is no such ambiguity with ${}^{\circ}\phi'$ and ${}^{\circ}\phi'$ nor with $f({}^{\circ}F)$ and $f({}^{\circ}F)$, where fx is a rational function of x . [Remember that $f{}^{\circ}\phi$ and $f{}^{\circ}\phi$ have no invarientive meaning. I think it

* Intended by author to be commas, upright and inverted. In future printing the author still thinks commas would be preferable to the bracket substitute.

best to adhere to the general rule, not rigidly prescribed, that c/∂ and ∂/c should be denoted by Greek letters and c/c and ∂/∂ by Roman capitals.]

Our new notations and terms apply to other symbols than invariantive for instance, to Γ_β . Thus

$$\Gamma_\beta = {}^0\Gamma_\beta, \quad E_\beta' = {}^0(E_\beta').$$

We shall retain the "cross" notation to indicate different symbols, and generally restrict it, without rigid prescription, to the same meanings as hitherto.

Our former $\bar{\Gamma}_\beta$ and $\bar{E}_\beta$ will henceforth not be used, because we have in §18 a new notation, according to which ${}^aE_\beta$ would have reference to a Weyl manifold, and would indicate that ${}^aE_\beta$ was the part of E_β not containing terms in λ ; and if λ is actually everywhere zero so that the manifold becomes a Riemann manifold, §18 provides the use of F_β in place of E_β . In the present paper we occupy ourselves exclusively with a Riemann manifold, and use the letters $\theta, l, F_\beta, \Theta_\beta, \Omega, C (= {}^0C = \Omega\theta^{-1})$ provided for that case.

A thick dot* will be used to denote that a symbol is a "density," though occasionally, as with W (action density), and f a part of W below, we shall not consider it necessary to use the dot. Once for all, we define η^* and θ^* by the equations

$$\eta^* = k\eta^{-1}, \quad \theta^* = l\theta^{-1}, \quad (1)$$

but we may regard it as a rule, unless the contrary is definitely stated, as here with η^* and θ^* , that the dot attached to a symbol means that the undotted symbol has been multiplied by l (by k if we were dealing with a Weyl manifold). Thus C^*, G^* stand for lC, lG , and again, if we introduce κ^* as a contravariant vector density, then κ will mean $l^{-1}\kappa^*$.

With regard to θ^* in (1), we shall use it as applied to a general multenion, say, to $u^* = V_\alpha q^*$, but not exactly with the extensive meaning. Thus θ^*u^* means $l\theta^{-1}u^*$ where θ^{-1} but not l is used in the extensive meaning. Thus, for instance, $\theta^*V_3\alpha^*\beta^*\gamma^*$ means $lV_3\theta^{-1}\alpha^*\theta^{-1}\beta^*\theta^{-1}\gamma^*$, and not $l^3V_3\theta^{-1}\alpha^*\theta^{-1}\beta^*\theta^{-1}\gamma^*$.

§20. *Maxwell's Equations for Matter in Bulk.*—It is fully time to show how the special requirements of Relativity are met in detail in our method. In now introducing Maxwell's equations, let us throughout keep in mind the application to matter in bulk. If for no other reason this is desirable because thereby we are prepared to accept the beautiful

* It is necessary, because of papers despatched to England in the early months of 1922, to state that this dot replaced a bar placed over the symbol affected. The replacement was made in August, 1922, at the suggestion of the printers. It seems now (August) too late to replace the bar in the papers just mentioned.

explanation that Maxwell's theory affords of the known behaviour of light in uniaxal and biaxal crystals. When we compare results with Maxwell's we shall speak as if $n = 4$. Nevertheless, in the general argument, we shall continue to suppose n to have any positive integral value. When with $n = 4$ we put

$$\rho = x\iota_1 + y\iota_2 + z\iota_3 + t\iota_4$$

we shall speak of x, y, z as the spatial co-ordinates and t as the time, even when by transformation of all four co-ordinates t may mean something very different from the time as ordinarily understood.

Let Maxwell's vector potential be $\mathbf{A} = (A_1, A_2, A_3)$, electrostatic potential $= A_4$, magnetic induction $\mathbf{B} = (B_1, B_2, B_3)$, electric intensity $\mathbf{E} = (E_1, E_2, E_3)$, magnetic field strength $\mathbf{H} = (H_1, H_2, H_3)$, dielectric displacement $\mathbf{D} = (D_1, D_2, D_3)$, conduction current $\mathbf{K} = (K_1, K_2, K_3)$, electrostatic density $= K_4$, force per unit volume $\mathbf{F} = (F_1, F_2, F_3)$, power $(-\mathbf{SEK})$ per unit volume $= F_4$.

Also let λ be a covariant vector (better use λ^* in the future) given by

$$\lambda = \Sigma A_c \iota_c \quad (c = 1, 2, 3, 4),$$

ω^* a covariant ${}_nV_2$ given by

$$\omega^* = \Sigma B_1 \iota_2 \iota_3 + \Sigma E_1 \iota_4 \iota_1$$

(the meaning of each summation, of three terms, being sufficiently obvious)

ω^* a contravariant ${}_nV_2$ density given by

$$\omega^* = \Sigma H_1 \iota_2 \iota_3 + \Sigma D_1 \iota_4 \iota_1,$$

κ^* a contravariant vector density given by

$$\kappa^* = \Sigma K_c \iota_c \quad (c = 1, 2, 3, 4),$$

and ν^{**} a covariant vector density given by

$$\nu^{**} = \Sigma F_c \iota_c \quad (c = 1, 2, 3, 4).$$

Maxwell's quaternion equations

$$\mathbf{B} = \nabla \mathbf{A}, \quad \mathbf{E} = -\nabla A_4 - \partial \mathbf{A} / \partial t, \quad (1)$$

$$\mathbf{K} = \nabla \mathbf{H} - \partial \mathbf{D} / \partial t, \quad K_4 = -\mathbf{S} \nabla \mathbf{D}, \quad (2)$$

$$\mathbf{F} = \mathbf{VKB} + K_4 \mathbf{E}, \quad F_4 = -\mathbf{SKE}, \quad (3)$$

are identical with the multenion equations*

$$\omega^* = \mathbf{V}_2 \nabla \lambda, \quad (1a)$$

$$\kappa^* = \mathbf{V}_1 \nabla \omega^*, \quad (2a)$$

$$\nu^{**} = \mathbf{V}_1 \kappa^* \omega^*. \quad (3a)$$

The equations (1a), (2a), (3a) are all invariantive.

* For a convenient, simple, systematic method of converting all ($n = 4$) multenion formulæ to corresponding quaternion formulæ, see a paper by the author (despatched a month ago, viz., November, 1922, to England), "The Mechanical Forces . . . in an Electromagnetic Field."

[A digression. In Maxwell's presentation of the fundamental aspects of the electromagnetic field $V\nabla\mathbf{H}$ in equation (2) is looked upon as *the* electric current, consisting of two parts, conduction current $\mathbf{K}$ and dielectric current $\partial\mathbf{D}/\partial t$. In the new Relativity scheme, as given by (2a), we naturally interchange the rôles of $V\nabla\mathbf{H}$ and $\mathbf{K}$; that is to say, we look upon $\mathbf{K}$ as *the* current, consisting of two parts, a magnetic part, $V\nabla\mathbf{H}$, and an electric part, $-\partial\mathbf{D}/\partial t$. In the multenion presentation (see below for $\mathbf{N}$ and $\mathbf{I}$) $V\nabla\mathbf{H}$ is $V_1\nabla(N\omega^*)$, $-\partial\mathbf{D}/\partial t$ is $NV_1\nabla(I\omega^*)$, and $\mathbf{K}_4$ (multiplied into l_4) is $IV_1\nabla(I\omega^*)$. [In the last sentence $\mathbf{N}$ and $\mathbf{I}$ are used as if $\mathbf{N} = {}^N\mathbf{N}$, $\mathbf{I} = {}^N\mathbf{I}$; if, as usual, we take $\mathbf{N} = {}^N\mathbf{N}$, $\mathbf{I} = {}^N\mathbf{I}$, then in the last sentence each $\mathbf{N}$ should be $\mathbf{N}'$ and each $\mathbf{I}$ should be $\mathbf{I}'$.] In the future, then, when we use the term current, Maxwell's conduction current will be meant. Maxwell does not put $\mathbf{F} = V\mathbf{K}\mathbf{B}$ as we have asserted in (3), but instead $\mathbf{F} = V(V\nabla\mathbf{H})\mathbf{B}$.]

Before speaking of the relations between Maxwell's $(\mathbf{B}, \mathbf{E})$ and $(\mathbf{H}, \mathbf{D})$, that is, between ω^* and ω^* , let us derive Lorentz's generalizations of Maxwell's equations to bodies in motion from the general principle of Relativity. The generalisations in nowise alter (1a), (2a), (3a). Put $\mathbf{N}_4$ and $\mathbf{I}_4$ for the normal and incident components with reference to ι_4 , when the co-ordinates are Galilean. In reality we have

$$\left. \begin{array}{ll} \mathbf{A} \text{ is } \mathbf{N}_4\lambda, & \iota_4\mathbf{A}_4 \text{ is } \mathbf{I}_4\lambda \\ \mathbf{B} \text{ is } \mathbf{N}_4\omega^*, & \mathbf{E} \text{ is } \mathbf{I}_4\omega^* \end{array} \right\}, \quad (1b)$$

$$\left. \begin{array}{ll} \mathbf{H} \text{ is } \mathbf{N}_4\omega^*, & \mathbf{D} \text{ is } \mathbf{I}_4\omega^* \\ \mathbf{K} \text{ is } \mathbf{N}_4\kappa^*, & \iota_4\mathbf{K}_4 \text{ is } \mathbf{I}_4\kappa^* \end{array} \right\}, \quad (2b)$$

$$\mathbf{F} \text{ is } \mathbf{N}_4\nu^*, \quad \iota_4\mathbf{F}_4 \text{ is } \mathbf{I}_4\nu^*. \quad (3b)$$

In the general θ (Riemann) manifold

$$\iota = d\rho/ds = d\rho/\sqrt{(V_0d\rho\theta d\rho)}$$

degenerates when Galilean co-ordinates are used ($\theta = 1$) to

$$\begin{aligned} & (\iota_1dx + \iota_2dy + \iota_3dz + \iota_4dt)/\sqrt{(-dx^2 - dy^2 - dz^2 + dt^2)} \\ & = (1 - \Sigma \dot{x}^2)^{-\frac{1}{2}} \cdot (\iota_1\dot{x} + \iota_2\dot{y} + \iota_3\dot{z} + \iota_4), \end{aligned}$$

of which we say nothing more here than that when the co-ordinates are so chosen that the particle is at rest ι (the contravariant world unit tangent vector to the world locus of a particle), degenerates to ι_4 . Maxwell's equations, as given above, are to be supposed true only when the medium is at rest. The generalisations of $\mathbf{N}_4$ and $\mathbf{I}_4$ are thus $\mathbf{N}$ and $\mathbf{I}$ (or $\mathbf{N}'$ and $\mathbf{I}'$), the normal and incident components with reference to ι ; normal component and normal expression, in this case, meaning the same, because

$V_0 \iota \theta \iota = 1$. Below N and I will always mean what has just been described with reference to ι , unless explicit exception is made; (1b), (2b), (3b), become like (1a), (2a), (3a), all invariantive when we read ι , N and I in place of ι_4 , N_4 , and I_4 , in (1a) and (3a); and in (2a), ι , N' and I' in place of ι_4 , N_4 , and I_4 . [Remember that $N = \langle N, N' = \rangle (N')$, and similarly for I .]

The equations this process gives when a body is moving, and we are using Galilean co-ordinates, are Lorentz's well-known generalizations of Maxwell's quaternion equations (not of our multenion equations, which have been in no way altered by what has been said).

From (1a) and (2a) we have of course

$$V_3 \nabla \omega^* = 0, \quad (4)$$

$$V_0 \nabla \kappa^* = 0. \quad (5)$$

These must be implicitly contained in (1) and (2) and indeed are found to be

$$S \nabla \mathbf{B} = 0, \quad V \nabla \mathbf{E} + \partial \mathbf{B} / \partial t = 0, \quad (4a)$$

$$S \nabla \mathbf{K} = \partial K_4 / \partial t. \quad (5a)$$

Maxwell's relations between $(\mathbf{B}, \mathbf{E})$ and $(\mathbf{H}, \mathbf{D})$ are that $\mathbf{H}$ is a self-conjugate linity of $\mathbf{B}$, (permeability) $^{-1}$; and $\mathbf{D}$ a similar function of $\mathbf{E}$, capacity. In our case

$$\left. \begin{aligned} \omega^* &= \psi^* \omega^*, \quad \psi^* = \psi'^* = {}^N \psi^* \\ I' \psi^* N &= 0 = N' \psi^* I \end{aligned} \right\}. \quad (6)$$

Here ψ^* is a ${}_N V_2$ linity concerning which, $\psi^* = \psi'^*$ asserts that ψ^* is self-conjugate, $\psi^* = {}^N \psi^*$ asserts that it is contraexco, the dot implies that it is a density, and the two equations in the second line assert that ψ^* consists of two parts $N' \psi^* N + I' \psi^* I$, of which the first is confined to normal components with reference to ι , and the other to incident components.

[Any c/c , say ϕ , may be expressed uniquely as the sum of four parts thus

$$\phi = (N + I) \phi (N + I) = N \phi N + I \phi I + N \phi I + I \phi N.$$

With a c/c the four parts are obtained from $(N + I)' () (N + I)$,

and in the case of ψ^* two of these parts are asserted in (6) to be zero.

Digression on I and N.—I have forgotten to point out earlier that I' , I , N' and N all have subextensive significance, and that $(I - N)$ also has a close connection with extensive interpretation. In the neutral form we virtually deal with a Galilean manifold. If in such a manifold I , N are the incident and normal components with reference to α , we can, by a rotation, bring α to coincidence with ι_1 , or some ι_a , and then $(I - N)$ becomes the capital linity P_1 (which reverses the sign of ι_1 in every term of q), introduced at the outset of our subject. The properties of I , N , $(I - N)$, may be based on this connection

with P_1 , through neutral to invariant forms. However, it is perfectly easy to prove directly that, say, I' , has subextensive significance, for we have defined thus

$$I'\beta = \alpha V_0 \eta \alpha \beta, \quad I'u = V_a \alpha V_{a-1} \eta \alpha u,$$

and to show that I' has subextensive significance we have to prove that

$$I'u = -V_a I' \zeta V_{a-1} \zeta u,$$

which is obvious as soon as stated.]

Maxwell's scheme also connects $\mathbf{K}$ with $\mathbf{E}$ by the self-conjugate linity conductivity. In our case express this first with Galilean co-ordinates. We have

$$\sum_{a=1}^3 E_a t_a = V_1 t_4 \omega^* = R N_4 \kappa^*$$

where R is put for resistivity. This generalizes to

$$N' \kappa^* = \phi^* V_1 t \omega^*, \quad \phi^* = N' \phi^* N = \phi'^* = {}^{(1)}\phi^* \quad (7)$$

where ϕ^* is conductivity, a vector linity.

It appears to me an impressive fact that Maxwell's analysis of the electromagnetic field was so wonderfully true to the nature of things, that not one single feature of his scheme, so far considered, requires the slightest revision when examined with the powerful searchlight of Relativity. Only when the structure perforce comes into contact with Newtonian mechanics, as is the case in what we are about to consider, does it require, along with classical mechanics, to be amended.

For stress put

$$M^* \alpha^* = -\frac{1}{2} \alpha^* V_0 \omega^* \omega^* + V_1 (V_1 \alpha^* \omega^*) \omega^*. \quad (8)$$

We have

$$M^* \nabla_9 = -\frac{1}{2} \nabla V_0 \omega^* \omega^* + V_1 (V_1 \nabla_9 \omega^*) \omega_9^* + \nu^{*x}$$

by (2a) and (3a). Now

$$V_1 (V_1 \nabla_9 \omega^*) \omega_9^* = \nabla_9 V_0 \omega^* \omega_9^* - V_1 \omega^* V_3 \nabla \omega^*$$

of which the last term is zero by (4). Hence

$$\begin{aligned} \nu^{*x} &= M^* \nabla_9 + \frac{1}{2} (\nabla_8 - \nabla_9) V_0 \omega^* \omega_9^* \Big\}, \\ &= M^* \nabla_9 + \frac{1}{2} \nabla_9 V_0 \omega^* \psi^* \omega^* \end{aligned} \quad (9)$$

by (5). By using ω^* instead of ω^* the last term on the right becomes $-\frac{1}{2} \nabla_9 V_0 \omega^* \psi^{*-1} \omega^*$. Because of the factor l^{-1} in $\psi^{*-1} = l^{-1} \psi^{-1}$ we may speak of ψ^{*-1} as an *anti*-density.

§ 21. *Stationary Action in a Riemann Manifold.*—In our theorems of integration $d\rho_c$ being equal to $V_c d\rho_1 d\rho_2 \dots d\rho_c$ is contravariant. But since

$$ldb = lV_0 dx d\rho_n$$

is an invariant, dx must be a covariant vector anti-density, and, similarly, ds_z in general (including db) is a covariant multienion anti-density.

(8) and (9) of § 20 are unsatisfactory in that $M\theta$ is not apparently self-conjugate, and from the second term on the right of (9) the force (cum power), per unit volume, cannot be expressed as a stress even of the non self-conjugate kind. But the pioneers of Relativity have developed a principle of stationary action which has for consequence that all of the equations of § 20 may be retained, and that M^* of (8) is self-conjugate; and by what seems little more than a bold assumption as to the nature of things, they conclude that the intractable last term of (9) need not trouble us, because force generated by the field need not be begotten by stress; they give us something else instead, as will appear below. [I think I have here credited the pioneers with just a little more than they have set forth, but no more than such as, by small additions, can be included.]

Although in the main we copy their procedure, yet in two important respects we depart from it, and in doing so return much more nearly than they do to Maxwell's outlook. First, we keep in mind the application to matter in bulk; and, secondly, we abjure the use of λ *in toto*. Probably Heaviside, among those who studied these matters in the days before the advent of Relativity, as we know it to-day, emphasised as much as any writer the view that the potentials of λ have nothing to do with actualities, unless among such are to be reckoned the imperfections of our mathematics. In returning to the procedure I adopted in 1892,* of taking Maxwell's $\mathbf{H}$ and $\mathbf{D}$ as the fundamental (three-dimensional) vectors of the field, instead of any of $\mathbf{A}$, $\mathbf{B}$, $\mathbf{C}$, $\mathbf{K}$, or $\mathbf{E}$, I find that the potentials can be completely ignored, that they are not wanted even as a mathematical crutch, and that the ignoring of them considerably simplifies the preliminary organising plans of our exploration. The electro-magnetic terms are all dependent on a single arbitrary scalar function of the ${}_nV_2$ called ω^* in § 20.

Let us recall that a particle of matter, with three dimensional extension, occupies a four-dimensional tube in the space-time continuum. The strength of this tube is the invariant constant (static) mass $dM^* = ldM$ of the particle. The three dimensional anti-density covariant cross-section being $d\alpha$, an element of "length" of the tube being the proper time ds , corresponding to the contravariant vector $d\rho$, where $d\alpha$ and $d\rho$ are both time-like, ι being the unit tangent vector $d\rho/ds$, db the four-dimensional volume of the element, we have

$$ds^2 = V_0 d\rho \theta d\rho, \quad \iota = d\rho/ds, \quad db = V_0 d\alpha d\rho, \quad (1)$$

$$\text{and} \quad dM^* \cdot ds = m^* db, \quad (2)$$

* "On the Mathematical Theory of Electromagnetism," 'Phil. Trans.,' 1893.

where m^* or sometimes m is called the material density. The momentum is

$$dM^* = m^* (db/ds), \quad (3)$$

and $m^* = \mu^*$, (4)

is called the momentum per unit volume. It is this vector whose magnitude multiplied by the cross-section yields the constant invariant strength, the quantity dM^* . [In invariants we should say $m \times (ldb + ds) = dM^*$, so that perhaps $m = \mu$ rather than μ^* should be called the momentum per unit volume.] We may note from the integration theorems, first for a finite portion of a tube, and next for a finite region of four dimensions, that

$$V_0 \nabla \mu^* = 0. \quad (5)$$

This "equation of continuity" or "equation of conservation" of matter is given by Weyl, but seems nowhere to appear in Eddington's Report.*

Our fundamental view of electricity is the same in principle, but we do not need to elaborate a notation such as dE^* , e^* , etc., in connection with κ^* to serve the purposes of dM^* , m^* , etc., in connection with μ^* . The tubes in the case of electricity (strength $V_0 \kappa^* dx$, just as strength of momentum tube $= V_0 \mu^* dx$) are called current tubes. If we were dealing with the smallest portions of matter it would accord with the prevalent views of to-day to suppose the two sets of tubes to be identical, but for matter in bulk we must consider them to be independent. Fortunately this is without any influence on the deductions we have to make from our fundamental principles.

Our principle of stationary action is that

$$\delta \int^n W db = 0, \quad (6)$$

where W is an invariant scalar density (meaning that $l^{-1}W$ is invariant) function of position consisting of three parts

$$W = m^* + f(\omega^*, \theta) + l(c - \frac{1}{2}G), \quad (7)$$

the first, material, being m^* already described and equal to $\sqrt{(V_0 \mu^* \theta \mu^*)}$; the second, electrical, a function $f(\omega^*, \theta)$ of a ${}_n V_2$, ω^* , identified with Maxwell's $\mathbf{H}$ and $\mathbf{D}$ as in § 20, and of θ the fundamental linity of the manifold; and the third, gravitational, $l(c - \frac{1}{2}G)$ where G is the scalar curvature $-V_0 \zeta C \zeta$, the cosmological constant c possibly being zero, though Einstein teaches that it must differ from zero.

The variations implied by δ in (6) are of three independent kinds; (1) an arbitrary increment in θ

$$\delta \theta = \theta_0, \quad (8)$$

* For a more critical examination of equation (5) see the later paper by the author: "The Mechanical Forces . . . Field" referred to above, footnote p. 164.

at every point of the manifold; (2) an arbitrary increment in ω^*

$$\delta\omega^* = V_2\nabla w^*, \quad (w^* = V_3\omega^*) \quad (9)$$

at every point, of such a kind that the current tubes, that is to say κ^* , which last is *defined* by

$$\kappa^* = V_1\nabla\omega^* \quad (10)$$

remain unvaried; and (3) an arbitrary displacement which shifts the tubes, momentum and current, together,

$$\delta\rho = \epsilon \quad (11)$$

without alteration of their strengths, $V_0\mu^*d\alpha$, $V_0\kappa^*d\alpha$.

[*Added October, 1921.*—A consequence of remarks which occur below is so obvious that it is singular that only now have I happened to recognise it. If it is assumed that for the first two variations ($\delta\theta = \theta_0$ and $\delta\omega^* = V_2\nabla w^*$) the stationary condition (6) holds; then *it follows as a mathematical deduction* that the stationary condition also holds for the third variation ($\delta\rho = \epsilon$) which shifts the tubes. Clearly then our postulated principle of stationary action should not include the *assumption* of the stationary condition for the third variation. The reader will easily justify the assertions that have just been made, if he will take as the proof of the equation $T_\zeta^{(t)}\zeta = 0$, the indication of a proof, immediately following the identity (35) below. This equation $T_\zeta^{(t)}\zeta = 0$ is virtually the necessary and sufficient condition for the existence of the stationary condition under the variation $\delta\rho = \epsilon$.]

The integral of (6) extends over any arbitrary finite region at the boundary of which θ_0 , ω^* and ϵ all vanish.

ω^* , eventually to be identified with Maxwell's **B** and **E** as in § 20, is *defined* by

$$\omega^* = -\omega.\nabla \quad (12)$$

where $\omega.\nabla$, as in Quaternions, means the differential operator for which $df = -V_0d\omega^*.\omega.\nabla f$, df being the increment in f due to an arbitrary increment $d\omega^*$ in ω^* .

It is necessary to point out that if **W** consisted of m^* alone, the condition $\delta W = 0$ would mean that a momentum tube is *geodesic* because by (2) $m^*db = dM^*ds$, and by hypothesis dM^* the strength is unaffected by the displacement ϵ of the tube. Indeed, following Weyl, we might use this identification to find the effects of ϵ and θ_0 on the variation of m^* in (6), but we pursue another course. It seems almost unnecessary to say that for a full treatment of matter in bulk **W** as above described is by no means adequate;

there are several scalar functions of position such as temperature and strain co-ordinates (adapted to the method of Relativity) that should be included in W for such a treatment.* We may accept the inclusion of fluid and elastic solid stresses that Eddington so lucidly justifies in § 37 of his report, despite the fact that his appeal to the kinetic theory of matter would by a precisionist be described as a departure from our elected task of dealing with matter in bulk.

We shall consider the effects of w^* , ϵ and θ_0 in succession because that is the order of simplicity; θ_0 , w^* and ϵ all take part in the variation of f ; θ_0 and ϵ in that of m^* ; and θ_0 alone in that of G .

The variation due to ω^* gives by (9) and (12)

$$0 = -\int^n \int V_0 \omega^* \nabla w^* db = -\int^{n-1} \int V_0 w^* \omega^* d\alpha + \int^n \int V_0 w^* V_3 \nabla \omega^* db,$$

whence, since w^* is zero at the boundary,

$$V_3 \nabla \omega^* = 0. \quad (13)$$

The process, here exhibited, of making a ∇ , explicitly occurring in the integrand, apply to the whole integrand, and the consequent removal of the resulting expression (or the potentiality of such removal) to the boundary, is known as integration by parts.

The effect of ϵ we will first consider with reference to the current tubes. For the strength of every tube to remain unchanged by the displacement it suffices that ω^* changes in precisely the manner that it changed in § 15 when ϵ indicated a change of co-ordinates. Quoting then from § 15, we first have

$$l^{-1} \delta l = -\frac{1}{2} V_0 \xi \delta \theta \theta^{-1} \xi = V_0 \xi \chi_0 \xi = V_0 \nabla \epsilon,$$

and next

$$\begin{aligned} \delta \omega^* &= \omega^* V_0 \nabla \epsilon - V_2 \chi_0 \xi V_1 \xi \omega^* \\ &= \omega^* V_0 \nabla \epsilon - V_2 \epsilon_9 V_1 \nabla_9 \omega^* = V_2 \nabla_9 V_3 \omega^* \epsilon_9. \end{aligned}$$

For the future we shall write δ' for these changes to be quoted from § 15 leaving δ for a second meaning thus (Weyl's notation)

$$\left. \begin{aligned} \delta' () &= (\text{value at } \rho + \epsilon \text{ after displacement}) - (\text{value at } \rho \text{ before}) \\ \delta () &= (\text{value at } \rho \text{ after displacement}) - (\text{value at } \rho \text{ before}) \end{aligned} \right\} \quad (14)$$

so that

$$\delta () = \delta' () + V_0 \epsilon \nabla . ()$$

Thus

$$\delta \omega^* = V_2 \nabla_9 V_3 \omega^* \epsilon_9 + V_0 \epsilon \nabla . \omega^*,$$

$$\delta f = V_0 \omega^* \delta \omega^* = V_0 \omega_9 \times \nabla_9 V_3 \omega^* \epsilon_9 - V_0 . V_3 \omega^* \nabla_9 V_3 \omega^* \epsilon_9 + V_0 \epsilon \nabla_9 V_0 \omega^* \omega^*_9$$

because $V_3 \nabla \omega^* = 0$. Thus δf —(first term on right)

$$= V_0 \epsilon V_1 \kappa^* \omega^*.$$

Hence

$$(\epsilon \text{ part}) \quad \int^n \int \delta f db = \int^n \int V_0 \epsilon V_1 \kappa^* \omega^* db. \quad (15)$$

* Treated in a series of three papers by the author; despatched within the past month (December, 1922) to England.

Next calculate the corresponding part from δm^* , in which $\delta \mu^*$ is obtained, as with $\delta \omega^*$ above, but $\delta \theta = 0$. We shall for brevity use

$$m^{*2} = V_0 \mu^* \theta \mu^*, \quad m^* \iota = \mu^*,$$

but the reader would do well at the beginning to use only $m^* = \sqrt{(V_0 \mu^* \theta \mu^*)}$ and not ι .

$$\delta m^* = \frac{1}{2} \delta (m^{*2}) / m^* = V_0 \iota \theta (-\epsilon_9 V_0 \mu^* \nabla_9 + \mu^* V_0 \nabla_9 \epsilon + V_0 \epsilon \nabla \cdot \mu^*).$$

Here prepare for integration by parts by altering the incidence of ∇ in the first two terms on the right, so as to apply to the whole term, and reject this part of each term. Thus $\delta m^* -$ (the rejected parts)

$$= V_0 \epsilon [(\theta \iota)_9 V_0 \mu^* \nabla_9 - \nabla m^* + \nabla_9 V_0 \mu^* \theta \iota]$$

$$= V_0 \epsilon (T_9 \nabla_9 + \frac{1}{2} \nabla_9 V_0 \zeta \theta_9 \theta^{-1} T \cdot \zeta)$$

where

$$T \cdot \alpha^\times = l T \alpha^\times = m^* \theta \iota V_0 \alpha^\times \iota \quad [+ \text{stress terms}] \quad (16)$$

(the stress terms meaning those inserted here from Eddington's argument in § 37 of the Report). Thus

$$(\epsilon \text{ part}) \quad \int^n \delta m^* db = \int^n l V_0 \epsilon T_{(G)} \zeta db \quad (17)$$

by equation (19) of § 15.

T or T^* is called the material kinetic energy linity, or if the stress terms (say $T^{(s)}$) are included, it may be called the material stress-energy linity. The stress terms are self-conjugate and satisfy the equation $T^{(s)} = N T^{(s)} N$.

(15) and (17) are the only contributions of ϵ to (6), so that we have

$$l T_{(G)} \zeta + V_1 \kappa^* \omega^\times = 0. \quad (18)$$

We have now fully identified $\omega^\times$ with Maxwell's **B** and **E**.

Of equations (1a), (2a), (3a), (4), (5), (6) of § 20 we have decided that (1a) is quite unnecessary, is in fact better neglected than cared for (though, of course, it is a mere consequence of our present equation (13)). The other five are replaced in the present section by (10), (18), (13), (10), (12) respectively. [(10) does duty twice, since it has the obvious consequence $V_0 \nabla \kappa^* = 0$.]

We are now about to trace the consequences flowing from the fact that $l^{-1} f$ is an invariant. These are wholly satisfactory for empty space, but not for matter in bulk, for which, in accordance with Maxwell's principles, we must modify our view as to the nature of f , the electrical term of the action density. It is well to show, however, that it is this invariance of $l^{-1} f$ which causes the inadequacy of our present supposition that f is a function of ω^* and θ only.

With our present (vacuum) assumption, we shall show that (18) can be added to, thus ($\kappa = l^{-1} \kappa^*$)

$$-T_{(G)} \zeta = V_1 \kappa \omega^\times = 2 (l^{-1} \theta_{(G)} f)_{(G)} \zeta, \quad (19)$$

where, as in Quaternions, $\epsilon\mathbf{I}$ means the self-conjugate linity differential operator given by

$$df = -V_0 d\theta \zeta \epsilon \mathbf{I} f \zeta \quad (20)$$

where df is the increment in f due to the arbitrary increment $d\theta$ in θ . It is, therefore, not the linity T , but, instead, the linity $T^{(t)}$ ($=$ total energy linity $= lT^{(t)}$) given by

$$T^{(t)} = T + 2\theta \epsilon \mathbf{I} f \quad (21)$$

which may be supposed to be handed on through space by some kind of contact action, because from (19)

$$T_{(G)}^{(t)} \zeta = 0. \quad (22)$$

As a matter of fact, we shall later, when considering gravitation, show that this equation can be transformed to the shape

$$\phi_9 \nabla_9 = 0.$$

We will now prove (19) on our present hypothesis [f as in (7)]. Let us, for brevity, use ν^* as in (3a), § 20; that is, we define ν^* by

$$\nu^* = V_1 \kappa^* \omega^* = V_1 (V_1 \nabla \omega^*) \omega^*. \quad (23)$$

Since $l^{-1}f$ is invariant, and since, as we saw above, before equation (14), $l^{-1}\delta'l = V_0 \nabla \epsilon$, where ϵ now, as in § 15, expresses an infinitesimal change of co-ordinates, we have as with $\delta\omega^*$ above in the equation following (14)

$$f V_0 \nabla \epsilon + V_0 \epsilon \nabla f = \delta f = V_0 \omega^* \delta \omega^* - V_0 \delta \theta \zeta \epsilon \mathbf{I} f \zeta.$$

$\delta\omega^*$ is given in the equation following (14) and $\delta'\theta$ is given in § 15 as $-\chi_0' \theta - \theta \chi_0$. Hence

$$f V_0 \nabla_9 \epsilon_9 + V_0 \epsilon \nabla f = V_0 \omega^* (V_2 \nabla_9 V_3 \omega^* \epsilon_9 + V_0 \epsilon \nabla \cdot \omega^*) + V_0 (2\epsilon_9 \theta \epsilon \mathbf{I} f \nabla_9 - \theta \epsilon \mathbf{I} f \zeta \theta_9 \zeta \cdot V_0 \epsilon \nabla_9). \quad (24)$$

Here ϵ and all the n^2 derivatives (∇_9 , ϵ_9) of ϵ are arbitrary, so that we may put arbitrary dummies γ , α^* , β for ϵ , ∇_9 , ϵ_9 respectively. From the dummy γ we get

$$\nabla f = \nabla_9 V_0 \omega^* \omega^* - \nabla_9 V_0 \theta_9 \zeta \epsilon \mathbf{I} f \zeta.$$

This identity, satisfied by f , does not depend on the invariance of f . It is easily seen to be a direct consequence of the assumption that f is a function of ω^* and θ . A useful result is obtained from it by modifying the first term on the right, thus

$$\nabla_9 V_0 \omega^* \omega^* = V_1 (V_3 \omega^* \nabla_9) \omega^* + \nu^* \quad [\text{eq. (23)}]$$

$$= V_1 (V_3 \omega_9^* \nabla_9) \omega^* + \nu^* \quad [\text{eq. (13)}]$$

$$= (-M_9 \nabla_9 + \frac{1}{2} \nabla V_0 \omega^* \omega^*) + \nu^* \quad [\text{eq. (8) § 20}],$$

whence

$$\nabla f = -M_9 \nabla_9 + \frac{1}{2} \nabla V_0 \omega^* \omega^* + \nu^* - \nabla_9 V_0 \theta_9 \zeta \epsilon \mathbf{I} f \zeta. \quad (25)$$

From the dummy β above in (24)

$$\left. \begin{aligned} f\alpha^* &= V_1(V_3\omega^*\alpha^*)\omega^* + 2\theta_\theta \mathbf{C}f\alpha^* \\ \text{or } f &= -M^* + \frac{1}{2}V_0\omega^*\omega^* + 2\theta_\theta \mathbf{C}f \end{aligned} \right\}. \quad (26)$$

This is an identity (remember ω^* is defined as $-\omega \cdot \nabla f$) which we have proved f to satisfy by reason of the invariance of $l^{-1}f$. Putting α^* in (26) $= \nabla_9$ and subtracting (25), we get (19).

(26) shows that $\theta^{-1}M^*$ is self-conjugate, which the reader will at first be inclined to claim as a highly satisfactory result. I am not inclined to dispute this so far as empty space is concerned, but for matter in bulk it is precisely what we do *not* desire. If $\theta^{-1}M^*$ is self-conjugate for matter in bulk, then crystalline transmission of light cannot be explained on Maxwell's theory. To prove this, choose co-ordinates such that $\iota = \iota_4$; this requires co-ordinates which at the point under consideration reduce the medium to rest. Examine the case when the co-ordinates are Galilean, and Maxwell's equations become (1a), (2a), (3a) of § 20. To translate $M^*\alpha^*$ into quaternion form, identify the multenion $\iota_2\iota_3, \iota_3\iota_1, \iota_1\iota_2$ with the quaternion i, j, k and the multenion

$$\iota_1\iota_2\iota_3\iota_4 = \dot{v}$$

with Hamilton's $\sqrt{(-1)}$ in his bi-quaternion $p+q\sqrt{(-1)}$. Thus $\omega^* = \mathbf{H} + \dot{v}\mathbf{D}$ and $\omega^* = \mathbf{B} + \dot{v}\mathbf{E}$. Indicate a ${}_nV_1$ such as α^* by (σ, s_4) where

$$\alpha^* = \sum s_c \iota_c, \quad (c = 1, 2, 3, 4),$$

$$\sigma = s_1 i + s_2 j + s_3 k,$$

and indicate a ${}_nV_3$ such as $\dot{v}\alpha^*$ by $\dot{v}(\sigma, s_4) = (\dot{v}\sigma, \dot{v}s_4)$.

Thus [compare (3) and (3a) of § 20]

$$V_1\alpha^*\omega^* = V_1(\sigma, s_4)(\mathbf{H} + \dot{v}\mathbf{D})$$

$$= (V\sigma\mathbf{H} + s_4\mathbf{D}, -S\sigma\mathbf{D}) = (\text{say}) (\sigma', s_4'),$$

$$V_1(V_1\alpha^*\omega^*)\omega^* = (V\sigma'\mathbf{B} + s_4'\mathbf{E}, -S\sigma'\mathbf{E})$$

$$= (V[V\sigma\mathbf{H} + s_4\mathbf{D}]\mathbf{B} - \mathbf{E}S\sigma\mathbf{D}, -S\mathbf{E}[\sigma\mathbf{H} + s_4\mathbf{D}])$$

adding $-\frac{1}{2}\alpha^*V_0\omega^*\omega^*$ and making a slight quaternion reduction we obtain

$$M^*\alpha^* = (-\frac{1}{2}V[\mathbf{H}\sigma\mathbf{B} + \mathbf{E}\sigma\mathbf{D}] + \frac{1}{2}V\sigma V[\mathbf{H}\mathbf{B} + \mathbf{E}\mathbf{D}] + s_4V\mathbf{D}\mathbf{B}, \\ S\sigma\mathbf{E}\mathbf{H} - \frac{1}{2}s_4S[\mathbf{B}\mathbf{H} + \mathbf{E}\mathbf{D}]).$$

In the square array of sixteen scalars representing the linity M^* the fourth column contains the components of $V\mathbf{D}\mathbf{B}$, the fourth row contains those of $-\mathbf{V}\mathbf{E}\mathbf{H}$, and the constituent in the fourth row and fourth column is $-\frac{1}{2}S(\mathbf{B}\mathbf{H} + \mathbf{E}\mathbf{D})$. The other nine scalars are those of the vector linity of σ in the expression for $M^*\alpha^*$ just written. Thus the conditions that $\theta^{-1}M^*$ shall be self-conjugate are that

$$V\mathbf{D}\mathbf{B} = V\mathbf{E}\mathbf{H}, \quad V\mathbf{H}\mathbf{B} = V\mathbf{D}\mathbf{E}.$$

[These, of course, amount to saying that **D** is parallel to **E** and **B** to **H**.] Now in a plane light wave **VEH** is parallel to the ray and **VDB** is normal to the wave front, so that to say that **VDB = VEH** is to deny the possibility of double and conical refraction.

Our assumption that f is a function of ω^* and θ only is contrary to Maxwell's, which asserts that N and I enter into the relations between ω^* and $\omega^\times$, that is ι , as well as ω^* and θ is a constituent variable of f . We may take f to be an explicit function of ω^* , $\varphi^* = I\omega^*$ and θ . Its general increment df , may be written

$$df = -V_0 d\omega^* (\omega^* \nabla) f - V_0 d\varphi^* (\varphi^* \nabla) f - V_0 d\theta \zeta (\theta \nabla) f \zeta$$

where $(\omega^* \nabla)$, $(\varphi^* \nabla)$ and $(\theta \nabla)$ are all *partial* differential coefficient operators, whereas $\omega^* \nabla$ and $\theta \nabla$ by contrast are *total* differential operators. Thus $\omega^* \nabla f$ comes partly from $(\omega^* \nabla) f$ and partly from $(\varphi^* \nabla) f$, and again $\theta \nabla f$ comes partly from $(\theta \nabla) f$ and partly from $(\varphi^* \nabla) f$ (because θ is involved in φ^*). However, there is a part of $-V_0 d\varphi^* (\varphi^* \nabla) f$, involving $d\iota$, still standing over.

[In Maxwell's own case $(\varphi^* \nabla) f = (-I + N)\omega^\times$ and we might assume this equation to hold in general, though this is not done below.] Writing this outstanding part as $-V_0 d\iota \nabla f$ we shall have

$$df = -V_0 d\iota \nabla f - V_0 d\omega^* \omega^* \nabla f - V_0 d\theta \zeta \theta \nabla f \zeta.$$

Thus using $d\varphi^*$, for the moment, to stand for the increment of φ^* depending on $d\iota$ we have

$$\begin{aligned} \varphi^* &= V_2 \iota V_1 \theta \iota \omega^*, \\ d\varphi^* &= V_2 d\iota V_1 \theta \iota \omega^* + V_2 \iota V_1 \theta d\iota \omega^*, \\ -V_0 d\iota \nabla f &= df = -V_0 d\iota \{ V_1 [V_1 \theta \iota \omega^*] (\varphi^* \nabla) f + \theta V_1 \omega^* V_1 (\varphi^* \nabla) f \iota \}. \end{aligned}$$

Thus we may understand ∇f to mean as follows

$$\left. \begin{aligned} \nabla f &= \xi \cdot \iota = N \{ V_1 [V_1 \theta \iota \omega^*] (\varphi^* \nabla) f + \theta V_1 \omega^* V_1 (\varphi^* \nabla) f \iota \}, \\ \text{where } \xi \cdot \alpha &= N(\phi + \phi') \alpha, \text{ and } \phi \alpha = \theta V_1 \omega^* V_1 (\varphi^* \nabla) f \alpha, \\ \text{and } \xi \cdot \iota &= (\xi \cdot \iota), \end{aligned} \right\} \quad (23a)$$

[Correction added September, 1921.— N was absent from equation (23a) as given in the original paper. If N is omitted it is not necessarily true, but if N is included it is necessarily true, that $\xi \cdot \iota = (\xi \cdot \iota)$. In the original presentation it was tacitly, and erroneously, assumed that $(\omega^* \nabla) f$, $(\varphi^* \nabla) f$ and $(\theta \nabla) f$ are always invariantive. They will doubtless be so *when a proper invariantive form of f has been selected*, and it is desirable to make such a selection when possible and convenient. We will now show that, whether such a selection has been made or not, ∇f , $\omega^* \nabla f$ and $\theta \nabla f$ are always invariantive, that is to say ∇f is a covariant vector density, $\omega^* \nabla f$ is a covariant ${}_n V_2$, and $\theta \nabla f$ is a

contraexco vector linity density. Let us put $\tau = x$ where x is an arbitrary invariant scalar function of position which we shall *eventually* take to be $x = 1$. Thus

$$f = f(\omega^*, \varphi^*, \theta) = f(\omega^*, I'\omega^*, \theta) \quad (A)$$

I' being that definite explicit function of τ and θ which is given by ($\Omega =$ a contravariant ${}_nV_2$ dummy)

$$I'\Omega = V_2\tau V_1\theta\tau\Omega/V_0\tau\theta\tau. \quad (B)$$

From this last dI' can be expressed in terms of $d\theta$ and $d\tau$ (and it may be noticed in passing that since I' is homogeneous and of degree zero both in θ and in τ we have by Euler's theorem for homogeneous functions

$$V_0\tau_\tau\nabla_\theta \cdot I'_\theta = 0 = V_0\theta_\theta\xi_\theta(I'_\theta \cdot I'_\theta) \quad (C)$$

the first of these explaining the presence of N in (23a), for it leads to $V_0\tau_\tau\nabla f = 0$). We now have

$$\begin{aligned} df &= -V_0d\omega^*(\omega^*\nabla)f - V_0d\varphi^*(\varphi^*\nabla)f - V_0d\theta\xi(I'f)\xi \\ &= -V_0d\omega^*\omega^*\nabla f - V_0d\tau_\tau\nabla f - V_0d\theta\xi_\theta(I'f)\xi \end{aligned} \quad (D)$$

the second being derived from the first by finding $d\varphi^*$ from

$$d\varphi^* = I'd\omega^* + dI' \cdot \omega^*,$$

dI' coming from (B). In the second of (D) $d\omega^*$, $d\tau$ and $d\theta$ are all *independently, arbitrary and invariantive* and therefore their coefficients,

$$\omega^*\nabla f, \quad \tau_\tau\nabla f, \quad \theta_\theta\xi f,$$

are *without ambiguity and are invariantive*. These last important deductions have been made possible by using τ instead of ι . τ (and therefore $d\tau$) is a general contravariant vector whereas ι is restricted by the equation $V_0\iota\theta\iota = 1$. From the above we obtain by straightforward filling in of detail that

$$x_\tau\nabla f = (1 - I)(\phi - \phi')\iota = \xi\iota$$

as given in equation (23a) above. Since $x_\tau\nabla f$ is independent in meaning of x , it is natural as in (23a) to write it in the form ∇f .

I take this opportunity of suggesting what is the proper form of f *in vacuo* and inside matter. *In vacuo* we have

$$f = V_0\omega^*\theta^{-1}\omega^* = l^{-1}V_0\omega^*\theta\omega^*.$$

Now $\theta = N\theta N' + I\theta I'$ so that

$$lf = V_0(N'\omega^*)\theta(N'\omega^*) + V_0(I'\omega^*)\theta(I'\omega^*).$$

I would suggest that inside matter

$$lf = V_0(N'\omega^*)\xi(N'\omega^*) + V_0(I'\omega^*)\xi(I'\omega^*) \quad (E)$$

where ξ is a function of θ and of position (ρ) only. Thus

$$lf = V_0\omega^*(N\xi N' + I\xi I')\omega^*. \quad (F)$$

Without loss of generality ξ may be replaced first by its self-conjugate part $\frac{1}{2}(\xi + \xi')$ since (E) is thereby unchanged, and next by the part

$$\frac{1}{2}[N(\xi + \xi')N' + I(\xi + \xi')I']$$

since (F) is thereby unchanged. With $n = 4$ the general ξ would involve 36 scalars, but the part we have finally reached involves only 12 scalars, namely 6 for $N(\xi + \xi')N'$ and 6 for $I(\xi + \xi')I'$. We may, as a special case, take $\xi = A\theta A'$ where A is a coexco linity function of position.]

We have to add to the right of (24), (25), and (26) the following marked (24a), (25a), (26a) respectively.

$$V_0\epsilon\xi^{\bullet}\iota \cdot V_0\iota\nabla_9 - V_0\epsilon\nabla_8 \cdot V_0\iota_8\xi^{\bullet}\iota, \quad (24a)$$

$$-\nabla_8 \cdot V_0\iota_8\xi^{\bullet}\iota, \quad (25a)$$

$$\xi^{\bullet}\iota \cdot V_0\iota\alpha^{\times}, \quad (26a)$$

(15) and (19) also are modified by $V_1\kappa\omega^{\times}$ changing to

$$-lT_{(g)}\zeta = V_1\kappa^{\bullet}\omega^{\times} - \nabla_9 V_0\iota_9\xi^{\bullet}\iota - (\xi^{\bullet}\iota)_9 V_0\iota_9\nabla_9. \quad (15a) \text{ and } (19a)$$

I think there can be no doubt* that this amendment is the true expression for the force, cum power, per unit volume. It will be observed that no change occurs in the total energy linity $T^{(t)} = T + 2\theta_0\Omega f$.

Pass now to the variation $\delta\theta = \theta_0$. There is no trouble in supplementing (15) and (17) with the θ_0 parts since no integration by parts is here involved. Thus since $m^{\bullet} = \sqrt{(V_0\mu^{\bullet}\theta\mu^{\bullet})}$, $\delta m^{\bullet} = \frac{1}{2}V_0\iota\theta_0\mu^{\bullet}$, and

$$(\theta_0 \text{ part}) \quad \delta(m^{\bullet} + f) = -\frac{1}{2}V_0\theta_0\zeta\theta^{-1}T^{(t)}\zeta. \quad (27)$$

G and $G^{\bullet}$ ($= lG$) are functions of θ and its first and second derivatives (∇), and therefore of $\theta^{\bullet}$ and its first and second derivatives, where $\theta^{\bullet} = l\theta^{-1}$. Let $\theta^{\bullet}$ receive any infinitesimal increment $\delta\theta^{\bullet}$. At one part of our work we shall assume $\delta\theta^{\bullet}$ to have the meaning corresponding to $\delta\theta = \theta_0$ of equation (8); at another part $\delta\theta^{\bullet}$ will mean the actual increment $d\theta^{\bullet} = -V_0\alpha\nabla \cdot \theta^{\bullet}$ of $\theta^{\bullet}$ on passing from the position ρ to the position $\rho + d\rho = \rho + \alpha$ in the manifold; on some other occasion it may mean some combination of these; in a word, $\delta\theta^{\bullet}$ is any conceivable defined infinitesimal increment of $\theta^{\bullet}$. The following two equations are true:—

$$G^{\bullet} = -V_0\Omega\zeta\theta^{\bullet}\zeta = -V_0\nabla(F_{\zeta}\theta^{\bullet}\zeta - \theta^{\bullet}F_{\zeta}'\zeta) - V_0\zeta\theta^{\bullet}(F_{\zeta_1}'F_{\zeta}' - F_{\zeta}'F_{\zeta_1}')\zeta_1, \quad (28)$$

$$-V_0\Omega\zeta\delta\theta^{\bullet}\zeta = -V_0\nabla(F_{\zeta}\delta\theta^{\bullet}\zeta - \delta\theta^{\bullet}F_{\zeta}'\zeta) - \delta \cdot V_0\zeta\theta^{\bullet}(F_{\zeta_1}'F_{\zeta}' - F_{\zeta}'F_{\zeta_1}')\zeta_1. \quad (29)$$

The great similarity between these equations is connected with the fact that the second term on the right of (28) is a function homogeneous and

* Especially in view of the paper twice referred to above: "The Mechanical Forces . . . Field."

quadratic in the derivatives (∇_9, θ_9) , and also homogeneous and of degree -1 in θ ; (28) may, as a matter of fact, be derived from (29) by means of this property if attention is paid to symmetry in ∇_9, θ_9 , and ∇_8, θ_8 , in the quadratic terms of the form $(\nabla_9, \theta_9, \nabla_8, \theta_8)$. The proof of (29) is here outlined, and the proof of (28) is similar and simpler. From (24) of § 16 (remember $F_\zeta' \zeta = l^{-1} \nabla l$)

$$\begin{aligned} -V_0 \Omega \zeta \delta \theta \cdot \zeta &= -V_0 \nabla (F_\zeta \delta \theta \cdot \zeta - \delta \theta \cdot F_\zeta' \zeta) - V_0 \delta \theta \cdot \nabla_9 F_\zeta' \zeta + V_0 \delta \theta \cdot \nabla_9 \zeta F_\zeta' \nabla_9 \\ &\quad - V_0 \delta \theta \cdot \zeta (-F_{\zeta_1}' F_{\zeta'}' + F_{\zeta'}' F_{\zeta_1}') \zeta_1. \end{aligned}$$

Comparing this with (29), putting $F_\zeta' \zeta = \sigma$, and noting that

$$F_\zeta \theta \cdot \zeta = -\theta_9 \nabla_9, \quad \delta (F_\zeta \theta \cdot \zeta) = -(\delta \theta)_9 \nabla_9 \quad (30)$$

it will be seen that we still have to prove that

$$\begin{aligned} -\delta \cdot V_0 \zeta \theta \cdot (F_{\zeta_1}' F_{\zeta'}' \zeta_1 - F_{\zeta'}' \sigma) &= V_0 \sigma \delta (F_\zeta \theta \cdot \zeta) \\ &\quad + V_0 \delta \theta \cdot \nabla_9 \zeta F_\zeta' \nabla_9 + V_0 \delta \theta \cdot \zeta (F_{\zeta_1}' F_{\zeta'}' \zeta_1 - F_{\zeta'}' \sigma). \end{aligned}$$

The only term that gives any trouble is the first on the left. In it $F_{\zeta'}' \zeta_1$ and ζ may be interchanged because $F_{\zeta'}' \zeta_1$ is self-conjugate in ζ . Thus the term is

$$-\delta \cdot V_0 F_{\zeta'}' \zeta_1 \theta \cdot F_{\zeta_1}' \zeta = -2V_0 F_{\zeta'}' \zeta_1 \delta (\theta \cdot F_{\zeta_1}') \zeta + V_0 F_{\zeta'}' \zeta_1 (\delta \theta \cdot) F_{\zeta_1}' \zeta.$$

Here, in the first term on the right, we need only consider the self-conjugate part of $\delta (\theta \cdot F_{\zeta_1}')$ because $F_{\zeta'}' \zeta_1$ is self-conjugate in ζ . This self-conjugate part is simple for

$$\theta \cdot F_{\beta'}' + F_{\beta} \theta \cdot = l V_0 \beta \nabla \cdot \theta^{-1} = V_0 \beta \nabla \cdot \theta - \theta \cdot V_0 \beta \sigma.$$

When this is used the rest of the reduction is straightforward.

The first of (30) may be used to modify three terms in (28) and (29). Similarly, $F_\zeta' \zeta = l^{-1} \nabla l$ may be used to modify four terms.

Assuming now the truth of (28) and (29) the gravitational part of the action in (6) may be put

$$\int^n \int (cl - \frac{1}{2} G \cdot) db = -\frac{1}{2} \int^{n-1} \int V_0 (\theta_9 \nabla_9 + l^{-1} \nabla l) d\alpha + \int^n \int g \cdot db \quad (31)$$

where

$$g \cdot = cl + \frac{1}{2} V_0 \zeta \theta \cdot (F_{\zeta_1}' F_{\zeta'}' - F_{\zeta'}' F_{\zeta_1}') \zeta_1. \quad (32)$$

Using (27) and (29), the last for preparing $\delta g \cdot$ for integration by parts, we have since

$$l^{-1} \delta l = -\frac{1}{2} V_0 \zeta \theta_0 \theta^{-1} \zeta, \quad \delta \theta \cdot = -\theta \cdot (\theta_0 \theta^{-1} + \frac{1}{2} V_0 \zeta \theta_0 \theta^{-1} \zeta) \quad (33)$$

$$\text{that } 0 = \delta \int^n \int W db = -\frac{1}{2} \int^n \int db V_0 \theta \cdot \theta_0 \zeta \{T^{(t)} + (C + \frac{1}{2} V_0 \zeta_1 C \zeta_1) + c\} \zeta$$

where C stands for $\Omega \theta^{-1}$ and not for the conjugate $\theta^{-1} \Omega$ (we have in either case $V_0 \zeta C \zeta = -G$). Hence

$$T^{(t)} + (C + \frac{1}{2} V_0 \zeta C \zeta) + c = 0. \quad (34)$$

Since $T_{(g)}^{(t)} \zeta = 0$ by (22) it follows that

$$(C - \frac{1}{2} G)_{(g)} \zeta = C_{(g)} \zeta + \frac{1}{2} \nabla V_0 \zeta C \zeta = 0. \quad (35)$$

(35) has appeared incidentally in our application of stationary action, but it is really an identity satisfied by C that has nothing to do with stationary action. That this is the case is proved by help of (28) and (29), from the fact that G is an invariant in just the same manner as the identity (26), satisfied by f , was proved. Conversely, if we had chosen to establish this identity in the manner indicated (34) would furnish us with a second proof that $T_{(G)}^{(G)}\zeta = 0$.

From (29) we get another identity of importance in connection with the conservation of energy. Let the δ of (29) stand for $-V_0\alpha\nabla(\)$, and let (29) thus come to read $V_0\alpha\sigma^* = 0$. We have, of course, $\sigma^* = 0$. If the reader will write out in full the meaning of this identity he will find it can be expressed thus. [Contrary to the conventions of these papers, t^* is here used to denote a vector linity, because t is similarly used by writers on Relativity for the corresponding "pseudo-tensor."] If

$$2t^*\alpha^* = -\nabla_9 V_0\alpha^* (F_\zeta\theta_9\zeta - \theta_9 F_\zeta'\zeta) - \alpha^* V_0\zeta\theta^* (F_{\zeta_1}'F_{\zeta'} - F_{\zeta'}F_{\zeta_1}')\zeta_1 \quad (36)$$

we have the identity

$$2t^*\nabla_9 = \nabla_9 V_0\zeta\theta_9\theta^{-1} (C - \frac{1}{2}G)\zeta = -2(C - \frac{1}{2}G)_9\nabla_9 \quad (37)$$

by (35) just written, and (19) § 15.

The conservation of energy is regarded as the interpretation of the equation

$$0 = (T^{(G)} + t^*)_9\nabla_9 = (T^* + 2\theta \nabla f + t^*)_9\nabla_9 \quad (38)$$

where T^* comes from m^* (matter), $2\theta_9\nabla f$ comes from f (the electromagnetic field), and t^* comes from the gravitational field. t^* is the only one of our many symbols which is not invariantive. [The reader might suppose that the terms in (19a) above, which involve ∇_9 , are not invariantive, but it is not difficult to show directly that they are.]

A little reflection will show that the two identities (35) and (37) hold in Weyl's manifold as well as in Riemann's. For C , G and F_β we have, in the notation of § 18, to read aC , aG and ${}^aE_\beta$.

§ 22. *A Matter that has been Overlooked.*—I forgot to describe in the second paper a process I frequently use in my MS. work. It may prove useful to others, and is especially convenient when $n = 4$. I am reminded of it by the w^* ($= V_3w^*$) of (9) § 21. The use of w^* , a contravariant density may be replaced by a complementary $({}_nV_a$ replaced by a ${}_nV_{n-a}$) covariant multenion (not density); and, similarly, a covariant anti-density may be replaced by a complementary contravariant multenion; this latter is the replacement we are already familiar with of $d\rho_a$ by $d\varsigma_{n-a}$ and conversely.

$l\dot{u}$ (where $\dot{u} = u_1u_2 \dots u_n$) is a covariant multenion. More generally, $x\dot{u}$ where x is invariant is a covariant multenion; and its reciprocal (of form

$\gamma l^{-1} \dot{v}$ where γ is any invariant) is a contravariant multenion. It easily follows that the two linities

$$\mathbf{T} = l \dot{v} (), \quad \mathbf{T}^{-1} = l^{-1} \dot{v}^{-1} () \quad (1)$$

are the one a c/∂ and the other a ∂/c , that is, of the same types as θ and θ^{-1} . As an example, $\mathbf{T}\omega^*$ above ($n = 4$) is a covariant vector density; multiplying by l^{-1} , $\dot{v}\omega^*$ or $\omega^* \dot{v}^{-1}$ is a covariant vector, $= \sigma^*$ say. Thus, in (9),

$$\delta\omega^* = V_2 \nabla w^* = V_2 \nabla (\sigma^* \dot{v}) = V_2 \nabla \sigma^* \cdot \dot{v}.$$

[When $n = 4$ the complement of a ${}_4V_2$ is a second ${}_4V_2$.] This serves to indicate that our use of $V_2 \nabla w^*$ is not far different from the usual use of $\omega^* = V_2 \nabla \lambda$, $\delta\omega^* = V_2 \nabla \delta \lambda$. [As a matter of fact, the $f(\omega^*, \theta)$ used above, and the corresponding usual expression involving λ , can be shown to be "reciprocal" functions in Routh's sense.]

[*Added August, 1922.*—Let us illustrate both the methods of the present section and those of the "digression" in § 18 at the end of the second paper by considering the contractions of a certain identity,* namely,

$$V_3 \zeta \varpi'_\zeta \omega = 0, \quad (2)$$

or as we will here write it

$$V_3 \zeta \varpi'_\zeta \eta^{-1} V_2 \gamma^* \epsilon^* = 0 = V_3 \zeta (\varpi' \eta^{-1})_\zeta V_2 \gamma^* \epsilon^*, \quad (3)$$

where ϖ has the meaning given in § 16 above for a Weyl manifold. (The suffix ζ denotes absolute differentiation *proper*.) This may be contracted twice, the contractions being obtained from the modification of (3),

$$V_0 (V_3 \alpha \beta \delta) (V_3 \zeta \varpi'_\zeta \eta^{-1} V_2 \gamma^* \epsilon^*) = 0, \quad (3a)$$

by first setting $\delta, \epsilon^* = \zeta_1, \zeta_1$, and then setting $\beta, \gamma^* = \zeta_2, \zeta_2$; finally writing $V_2 \zeta_2 \zeta_1 = \zeta_2''$. The contractions are

$$V_2 \zeta_1 V_3 \zeta \varpi'_\zeta \eta^{-1} V_2 \gamma^* \zeta_1 = 0. \quad (4)$$

$$V_1 \zeta_2'' V_3 \zeta \varpi'_\zeta \eta^{-1} \zeta_2'' = 0. \quad (5)$$

(5) is the well-known identity (known, that is, in a Riemann manifold) numbered (35) in § 21 above. (3) asserts that a certain ${}_nV_3/{}_nV_2$ multenion linity (that is, a linity which acts solely on an arbitrary ${}_nV_2$ and converts it to a ${}_nV_3$) vanishes, (4) that a ${}_nV_2/{}_nV_1$ vanishes, and (5) that a vector vanishes.

Consider the number of scalar equations involved in the case of the Riemann manifold. The three identities are relations among the $\frac{1}{12} n^3 (n^2 - 1)$ first differential coefficients of ϖ . (3) is equivalent to ${}_nC_3 \cdot {}_nC_2$ scalar equations

* "A New Identity affecting Questions in Relativity; and some Cognate Matters." A paper despatched to the 'Phil. Mag.' in January, 1922, which as yet I have not seen in print.

(4) to ${}_nC_1 \cdot n$ such equations; and (5) to n such. The cases of $n = 3$ and $n = 4$ are particularly instructive. With $n = 3$ there are 18 differential coefficients and the other three numbers in order are 3, 9, 3; and when $n = 4$ there are 80 differential coefficients, and the three numbers in order are 24, 24, 4.

In the case of $n = 3$ there cannot, of course, really be nine independent equations resulting from the first contraction, since only three independent equations are given by (3). The details of this case are not difficult and may be left to the reader.

In the case of $n = 4$, the relativity case, the two numbers 24, 24 above suggest that perhaps the original equation (3) and its first contraction (4) are mathematically equivalent to one another, and this proves to be true. Put $F()$ for the $V_3 \zeta \omega \zeta' \eta^{-1}()$ occurring in (3), (4), (5), so that F is a $(c_n V_3)/(c_n V_2)$, from which we deduce that F' is a $(c_n V_2)/(c_n V_3)$, that $\Upsilon F' \Upsilon^{-1}$ is a $(c_n V_2)/(c_n V_3)$, and so on.

The contractions (4) and (5) of F are

$$H\gamma^* = V_2 \zeta F V_2 \gamma^* \zeta, \quad (4a)$$

where H is a $(c_n V_2)/(c_n V_1)$, and

$$\sigma^* = V_1 \zeta H \zeta = V_1 \zeta_2'' F \zeta_2'', \quad (5a)$$

H and σ^* , which are introduced for brevity, are defined by (4a) and (5a).

Now, presumptively, both F and H involve twenty-four independent scalars (F may be supposed by definition to involve the twenty-four, but H might possibly involve fewer, as when above we put $n = 3$), so that it appears probable that in the special case $n = 4$ (and also $n = 3$, but *certainly* not in general for $n > 4$) that not only can H be expressed in terms of F , but also F can be expressed in terms of H . Actually I find that

$$\Upsilon H' \Upsilon^{-1} \omega^* = \frac{1}{2} V_3 \omega^* \sigma^* + F \omega^*, \quad (6)$$

where ω^* is a dummy, and σ^* may be expressed either in terms of F or in terms of H . From (6), then, we easily express F in terms of H . I do not think I have found the simplest way to (6), namely by aid of a rectangular array of twenty-four scalars specifying F . [In the verification of (6) we need not cumber ourselves with the inconvenience of the signs in

$$\iota_1^2 = \iota_2^2 = \iota_3^2 = -\iota_4^2.$$

It is legitimate to work with i_1, i_2, i_3, i_4 , for which each $i_a^2 = -1$.]

The following can also be proved. If with $n = 4$, F is a $(c_4 V_2)/(c_4 V_2)$, that is a ${}_4V_2$ coexco linity, and if H is its first contraction and f its second according to the equations

$$\left. \begin{aligned} H\beta^* &= V_1 \zeta F V_2 \beta^* \zeta \\ f &= -V_0 \zeta H \zeta = -V_0 \zeta_2'' F \zeta_2'' \end{aligned} \right\} \quad (7)$$

then, putting C for the contractile part of F (so that $F - C$ is the non-contractile part) we have

$$\begin{aligned} C\omega^* &= \frac{1}{2} (F - TF'T^{-1} + \frac{1}{6}f)\omega^* \} \\ &= \frac{1}{2} V_2 \zeta H V_1 \omega^* \zeta - \frac{1}{6} f \cdot \omega^* \} \end{aligned} \quad (8)$$

where ω^* is a dummy. If in (7) and (8) F is put equal to $\varpi\eta^{-1}$, or $\varpi'\eta^{-1}$, or $\varpi\theta^{-1}$, the contractions are the familiar contractions of the curvature.

It should here be stated that throughout the above and also in general work with contractions the contractile parts can always be written down from the following easily verified statements. If (β, γ) , (β, γ^*) and (β^*, γ^*) stand for any bilinear invariant forms in two vectors then the contractile parts of the forms are

$$-n^{-1}(\zeta, \eta^{-1}\zeta) V_0 \beta \eta \gamma, \quad -n^{-1}(\zeta, \zeta) V_0 \beta \gamma^*, \quad -n^{-1}(\zeta, \eta\zeta) V_0 \beta^* \eta^{-1} \gamma^*$$

respectively.

With the design of trying to avoid future chaos in notation I venture to recommend a development of notation to users of the present methods.

The single letter symbol ω is not sufficient provision for the ${}_nV_2$ meaning. Also we have an insufficient supply of letter symbols for scalars. Let us release p, q, r, s, u, v, w for scalar use and provide several letters for the ${}_nV_2$ meaning. I recommend that most of the small letters in Clarendon type be ear-marked as follows. Use

p, q, r, s

for general multenions;

a, b, c

for the ${}_nV_a, {}_nV_b, {}_nV_c$ meanings;

f, g, h, i, j, k

for the ${}_nV_2$ meaning. [With $n = 4$ we may take

$$(i, j, k) = (\iota_2 \iota_3, \iota_3 \iota_1, \iota_1 \iota_2)].$$

Also use **l, m, n** for the ${}_nV_3$ meaning. **d, e, t, u, v, w, x, y, z** may be left for analogous uses as experience guides. Similarly occasional use may be made of capitals in Clarendon type, or perhaps such capitals had better be reserved for vectors in a quaternion system, as with **A, B, C, D, E, F, H, K** above (and, in other papers recently forwarded, **V** = velocity and **U** = unit radial vector).

Is it a vain hope that the publication of the present papers may encourage some young mathematician to "cultivate Hamilton's garden," his *linear associative vector algebra*, in spite of the genial sneers of Weyl and Eddington who virtually thank the Lord that *their* hands need not be soiled by this

"somewhat forbidding"*—hah—"orgy of"† agricultural labour? During 30 years and more I have often glanced over the hedge and wondered sadly why fellow workers should be content to turn over the, perhaps lighter, but incomparably less fertile, patches in the immediate neighbourhood of the garden.]

The Specific Heats of Air, Steam, and Carbon Dioxide.

By W. D. WOMERSLEY, M.A.

(Communicated by Sir Dugald Clark, F.R.S. Received January 6, 1923.)

In a recent number‡ is a criticism under the above title by Sir R. T. Glazebrook, of some figures given by me§ in a paper with the same title.

I am accused of not giving Holborn and Henning's own figures, because I state that part of my curve is filled in from the researches of Swann, and of Holborn and Henning. As Swann and Holborn and Henning are not in agreement, it seems evident that one set of figures cannot be used without some adjustment to the other set. These discrepancies were dealt with by the British Association Committee (of which Sir R. T. Glazebrook later became a member) in their 1908 Report, and the figures I give are in fairly close agreement with theirs for the low temperatures.

I state in my paper (p. 492) that I find Holborn and Henning about $7\frac{1}{2}$ per cent. too low at 800° C. for air and steam, and I assume that the same error applies to the carbon dioxide. (Prof. Callendar suggests their error may be as much as 10 per cent. at 1400° C.) I have, therefore, distributed this error over the range for which I have used Holborn and Henning's figures. On this account it may be anticipated "that the figures at the higher temperatures are higher than the corresponding figures due to Holborn." I believe it is usual, when quoting the results of other workers, to attach their names to the figures quoted; this I have not done.

Sir R. T. Glazebrook states that the slopes of the various lines in my figs. 4 and 5 give the volumetric heats. This is not the case, and as the meaning of

* Eddington's 'Report on Relativity,' preface to first edition, reproduced in second edition.

† English translation by Brose of Weyl's 'Space, Time, Matter,' p. 54.

‡ 'Roy. Soc. Proc.,' A, vol. 101, p. 112.

§ 'Roy. Soc. Proc.,' A, vol. 100, p. 483.