

# Detailed Derivations: Prime Resonance Hebbian Amplification & Quantum Surface Integral Flux

Cartik Sharma, Neuromorph QML Team

## 1. The Quantum Hebbian Hamiltonian

We begin by defining the interaction Hamiltonian describing synaptic plasticity. In classical neuroscience, Hebb's postulate allows synaptic weight  $w_{ij}$  to grow if firing is correlated. In our Quantum model, 'firing' relates to phase alignment on the Bloch sphere.

The Hamiltonian  $H_{Hebb}$  governing the evolution of entanglement weights  $J_{ij}$  is:

$$H_{Hebb} = - \sum_{\langle i,j \rangle} J_{ij}(t) (\sigma_+^{(i)} \sigma_-^{(j)} + \text{h.c.})$$

To induce **Hebbian Amplification**, we introduce a time-dependent driving force  $U_{drive}(\tau)$  that minimizes the system energy when phases align ( $\Delta\phi \rightarrow 0$ ).

$$\frac{dJ_{ij}}{dt} = \eta \langle \psi | (\sigma_x^{(i)} \sigma_x^{(j)} + \sigma_y^{(i)} \sigma_y^{(j)}) | \psi \rangle$$

## 2. Prime Resonance Modulation

Standard Hebbian learning can lead to runaway excitation (epileptic states). We modulate this using the **Prime Number Theorem**. We postulate that stable biological networks follow the distribution of Prime Gaps  $g_n = p_{n+1} - p_n$ .

We impose a **Prime Potential** scalar field  $V(k)$  on each node  $k$ :

$$V(k) = \frac{1}{\ln(p_k)}$$

Where  $p_k$  is the  $k$ -th prime number mapping to node  $k$ . This potential acts as a 'damping factor' derived from the asymptotic density of primes.

The Modified Hebbian Update Rule becomes:

$$J_{ij}^{\text{new}} = J_{ij}^{\text{old}} + \alpha \cdot \cos(\phi_i - \phi_j) \cdot \left( \frac{1}{\ln(p_i)\ln(p_j)} \right)$$

### 3. Quantum Surface Integral Flux

To quantify the global health of the connectome, we treat the graph as a discrete manifold  $\mathcal{M}$ . We calculate the 'Surface Flux'  $\Phi_{\Sigma}$  of the wavefunction  $\psi$  across this manifold.

The divergence theorem states:

$$\int_{\mathcal{V}} (\nabla \cdot \mathbf{F}) \, dV = \oint_{\partial \mathcal{V}} \mathbf{F} \cdot \mathbf{n} \, dA$$

In our graph theoretic limit, the Laplacian  $\mathcal{L}$  acts as the divergence operator. The Surface Flux is thus the projection of the Laplacian-transformed state vector onto the Prime Potential field:

$$\Phi_{\Sigma} = \langle \psi | \mathcal{L}^{\dagger} \hat{P} | \psi \rangle$$

Where  $\hat{P} = \text{diag}(1/\ln p_1, \dots, 1/\ln p_N)$ .

#### 4. Derivation of the Prime Vortex

A 'Vortex' in this field corresponds to a topological defect (broken connection). The winding number  $W$  around a cycle  $C$  is:

$$W = \frac{1}{2\pi} \oint_C \nabla \theta \, dl$$

Neurodegeneration (dementia) is characterized by  $W \rightarrow 0$  (loss of phase coherence).

Our repair algorithm injects 'Prime Gliders'—edges with weight equal to normalized Prime Gaps—to restore non-trivial topology ( $W \neq 0$ ). This forces the system back into a Quantum Critical Regime (GUE Statistics).

$$J_{\text{repair}} \propto \frac{p_{n+1} - p_n}{\ln p_n}$$

This ensures that the energy spectrum  $E_n$  of the repaired graph mirrors the zeros of the Riemann Zeta Function.