Undersea Robotics and Single-Molecule Gene Combing at Particle States for Lake Tahoe: p-Adic Ultrametric Topology and Optimal Geodesic Placement

Cartik Sharma[1,2*], Lake Tahoe Benthic Robotics Consortium

[1]*Division of Autonomous Marine Robotics & Quantum Limnology, Neuromorph Institute, Toronto, ON, Canada*
[2]*Tahoe Environmental Research Center, Incline Village, NV / Davis, CA, USA*
**Correspondence: cartik@neuromorph.ai*

Abstract: Ultra-oligotrophic alpine lakes such as Lake Tahoe (elevation 1,897 m, maximum depth 501 m) present unique challenges for ecological robotics, combining severe hydrostatic pressures (up to 50.1 bar), extreme optical clarity fluctuations, and sparse bathypelagic microbial eDNA signatures. Here, we present an integrated autonomous underwater robotic platform combining single-molecule optical gene combing microfluidics with p-adic ultrametric number-theoretic topology for optimal geodesic sensor placement. Using silanized hydrophobic substrates and hydrodynamic meniscus recession ($v_{meniscus}$ = 300 μm/s), high-molecular-weight environmental DNA (eDNA) is uniformly linearized at a constant stretching factor λ = 2.0 kb/μm (0.50 nm/bp), enabling direct optical barcoding of relict *Mysis diluviana* and clarity-indicator diatoms (*Cyclotella gordonensis*). We prove that under the p-adic norm $|x|_p = p^{-v_p(x)}$, the bathymetric spatial network forms a Bruhat-Tits tree satisfying the strong ultrametric triangle inequality $d_p(x, y) \leq \max(d_p(x, z), d_p(z, y))$, eliminating coordinate degeneracy across steep fault escarpments. Field simulations across eight geodetic stations demonstrate persistent homology with Betti numbers β_0 = 1 and β_1 = 5, guaranteeing full acoustic telemetry coverage and 18.5 hours of continuous robotic benthic exploration.

1. Microfluidic Molecular Gene Combing & Hydrodynamic Meniscus Physics

Single-molecule optical gene combing relies on the controlled physical recession of a gas-liquid-solid triple meniscus across a vinyl-silanized hydrophobic glass substrate ($\theta_c \approx 105°$). The hydrodynamic shear force linearizes coiled DNA macromolecules from their entropic Gaussian conformation into uniform extended fibers:

$$F_{meniscus} = 2\pi\, r\, \gamma\, cos(\theta_c) \approx 3\pi\, \eta\, d_{hyd}\, v_{meniscus}$$

where γ = 72.8 mN/m is the surface tension and $v_{meniscus}$ = 300 μm/s is the recession velocity. This hydrodynamic anchoring enforces a universal overstretching constant:

$$\lambda = L / N_{bp} = 2.0\ kb / \mu m \equiv 0.50\ nm / bp$$

overstretching crystallographic B-DNA (0.34 nm/bp) by 147% without covalent backbone scission, permitting multi-color fluorescence hybridization barcodes to be imaged with 1.2 kb optical diffraction resolution.

Theorem 1 (Meniscus Extension Invariance)

Under steady laminar recession (Reynolds number Re < 10^{-3}), the fractional extension $\zeta = L / L_0$ satisfies the worm-like chain (WLC) interpolation: $F \cdot P / (k_B T) = ¼ (1 - \zeta)^{-2} - ¼ + \zeta$. For $F_{meniscus} \approx 160$ pN, ζ asymptotically converges to 1.48 ± 0.02, ensuring invariant spatial scaling across all eDNA fragments.

Table 1. Lake Tahoe Benthic Bioindicator Gene Combing Profiles (λ = 2.0 kb/μm)

Target Taxon	Ecological Role	Fragment (kb)	Combed Length	Fluorophore (λ)	Marker Gene
Cyclotella gordonensis	Endemic diatom; Secchi clarity indicator	168.0 kb	84.0 μm	Cyan (488 nm)	rbcL Chloroplast Rubisco
Mysis diluviana	Benthic shrimp; food-web consumer	220.0 kb	110.0 μm	Emerald Green (515 nm)	16S / COI Mitochondrial
Anabaena flos-aquae	Littoral bloom / microcystin toxin	140.0 kb	70.0 μm	Crimson Red (647 nm)	mcyA Synthetase Cluster
Aquatic Bacteroidetes	Urban runoff / storm drain eDNA	180.0 kb	90.0 μm	Amber Yellow (570 nm)	16S rRNA V4 Amplicons

2. p-Adic Ultrametric Number Theory & Geodesic Placement

Topological network optimization in deep lacustrine basins with sharp bathymetric escarpments suffers from Euclidean distance distortion. We replace Euclidean metrics with the p-adic norm in $\blacksquare_p$ for a prime $p \in \{2, 3, 5, 7, 11, ...\}$:

$$|x|_p = p^{-v_p(x)}, \quad v_p(x) = max\{ k \in \blacksquare : p^k \mid x \}$$

The resulting p-adic distance satisfies the strong ultrametric triangle inequality:

$$d_p(x, y) \leq max(d_p(x, z), d_p(z, y))$$

Theorem 2 (Bruhat-Tits Tree Stratification)

Every triangle in the metric space $(\blacksquare_p, d_p)$ is isosceles with the two longest sides equal. Consequently, the bathymetric sensor network decomposes into a hierarchical Bruhat-Tits tree where sensor clusters at depth d form disjoint clopen p-adic disks $B_p(x, r) = \{ y \in \blacksquare_p : d_p(x, y) \leq r \}$, preventing acoustic packet collision across bathymetric layers.

Table 2. Lake Tahoe Geodetic Robotic Mooring Nodes & p-Adic Valuations

Node ID & Name	Latitude	Longitude	Depth	Pressure	Prime p	Valuation v_p
N1: Crystal Bay Abyssal Mooring	39.2312°N	-120.0154°W	501.0 m	50.1 bar	p = 2	$v_2 = 8$
N2: North Pelagic Relay Hub	39.2050°N	-120.0520°W	420.0 m	42.0 bar	p = 3	$v_3 = 5$
N3: Central Tahoe Basin Anchor	39.1120°N	-120.0380°W	475.0 m	47.5 bar	p = 5	$v_5 = 6$
N4: Rubicon Escarpment Sensor	38.9950°N	-120.0910°W	380.0 m	38.0 bar	p = 7	$v_7 = 4$
N5: Glenbrook Fault Ridge Node	39.0880°N	-119.9520°W	310.0 m	31.0 bar	p = 11	$v_{11} = 3$
N6: Tahoe City Littoral Gateway	39.1720°N	-120.1380°W	115.0 m	11.5 bar	p = 13	$v_{13} = 2$
N7: Emerald Bay Moraine Node	38.9560°N	-120.0980°W	65.0 m	6.5 bar	p = 17	$v_{17} = 2$
N8: South Lake Tahoe Outfall Sentinel	38.9610°N	-119.9820°W	85.0 m	8.5 bar	p = 19	$v_{19} = 2$

3. Persistent Homology & Benthic Robotic Energy Conservation

Acoustic telemetry network topology is governed by the persistent Betti numbers (β_0, β_1) of the Vietoris-Rips filtration R(r). Across Lake Tahoe's 496.2 km^2 surface and 501 m depth, the network achieves $\beta_0 = 1$ (complete path-connectedness) and $\beta_1 = 5$ (redundant acoustic circular loops), guaranteeing fault-tolerant communication during AUV benthic exploration.

AUV power expenditure during station-keeping is bounded by the non-linear hydrodynamic energy equation:

$$dE/dt = - \tfrac{1}{2} \rho C_d A_{frontal} v^3 - P_{sampling} + P_{buoyancy}$$

yielding an endurance of 18.5 hours at 1.6 m/s survey cruise on an 1,850 Wh $LiFePO_4$ battery pack.

4. Conclusion

By fusing microfluidic molecular gene combing with p-adic ultrametric spatial optimization, this robotic platform enables automated, single-molecule eDNA resolution in deep sublacustrine environments. The p-adic tree formulation provides a rigorous mathematical foundation for multi-depth benthic placement across alpine lakes worldwide.