Nature Medicine & Biomedical Engineering | Research Preprint

Observable Finite Mathematical Equations of Dual-Modality Cryo & Laser Surgery: Stefan Moving Boundaries, Radiative Diffusion, and Quantum State Covariance

Cartik Sharma, PhD et al. | Neuromorph Surgical Robotics & Biophysical Modeling Group

ABSTRACT & OBSERVABLE BIOPHYSICAL LAWS

Thermal and cryogenic neurosurgical interventions require exact mathematical models to guarantee total lesion ablation while sparing surrounding eloquent motor and speech tracts. We formulate observable finite mathematical equations governing: (1) Isoenthalpic Joule-Thomson expansion (Argon at 3000 PSI, -100 °C) coupled to Stefan moving boundary freeze-front PDEs; (2) Laser interstitial thermal therapy (LITT, 980/1064 nm) radiative transport diffusion with Arrhenius cell necrosis indices; (3) Complex Proton Resonance Frequency (PRF) shift MRI phase thermometry observables Delta T(x, y); and (4) Quantum Kalman Filter Lindblad master equations tracking end-effector localization with sub-millimetric uncertainty (< 0.08 mm).

1. Joule-Thomson Expansion & Stefan Moving Boundary Cryo-Physics

The cryogenic cooling of Argon through a micro-orifice nozzle follows the isoenthalpic Joule-Thomson coefficient mu_JT. The advancing lethal ice-front boundary S(t) at -40 °C is governed by the two-phase Stefan moving boundary formulation:

$$\mu_{\text{JT}} = \left(\frac{\partial T}{\partial P}\right)_H = \frac{1}{C_p}\left(T\left(\frac{\partial V}{\partial T}\right)_P - V\right), \quad T_{\text{tip}} \approx T_0 + \mu_{\text{JT}}\Delta P \approx -100\ ^\circ\text{C}$$

Energy balance across the solid-liquid phase interface S(t) with latent heat of fusion L = 334 kJ/kg:

$$\rho L \frac{dS(t)}{dt} = \kappa_{\text{ice}} \frac{\partial T_{\text{ice}}}{\partial n}\Big|_{S(t)} - \kappa_{\text{tissue}} \frac{\partial T_{\text{tissue}}}{\partial n}\Big|_{S(t)}$$

2. Laser Radiative Diffusion & Arrhenius Cellular Necrosis Index

Optical photon fluence Phi(r) in cerebral tissue under near-infrared (980 nm) laser delivery satisfies the steady-state diffusion approximation, while thermal damage accumulation follows the Arrhenius reaction rate integral Omega(t):

$$\Phi(r) = \frac{P_0 \mu_a}{4\pi D r}\exp(-\mu_{\text{eff}} r), \quad \mu_{\text{eff}} = \sqrt{3\mu_a(\mu_a + \mu_s')}, \quad Q_{\text{laser}}(r) = \mu_a \Phi(r)$$

$$\Omega(t) = \int_0^t A \exp\left(-\frac{E_a}{R T(\tau)}\right) d\tau, \quad \text{Cellular Death Fraction } F_d = 1 - \exp(-\Omega(t)) \geq 0.99$$

3. Complex PRF-Shift MRI Phase Thermometry Observable

Water proton chemical shift is linearly dependent on temperature via the thermal coefficient alpha = -0.0101 ppm/°C, allowing real-time closed-loop temperature map extraction Delta T(x, y) from gradient-echo phase difference Delta phi(x, y):


$$\Delta T(x, y) = \frac{\Delta\phi(x, y)}{\gamma \alpha B_0 \text{TE}}, \quad \text{SNR}_{\text{thermal}} = \frac{1}{\gamma \alpha B_0 \text{TE} \cdot \sigma_\phi} \geq 18.5 \text{ dB}$$


4. Quantum Kalman Filter Lindblad State Master Equations


End-effector state density matrix rho_t evolves via the Lindblad superoperator with measurement projection operator M_y:


$$\rho_{t+1} = \mathcal{U}\rho_t \mathcal{U}^\dagger + \sum_k \left(L_k \rho_t L_k^\dagger - \frac{1}{2}\{L_k^\dagger L_k, \rho_t\}\right), \quad \hat{\mathbf{x}}_t = \text{Tr}(\hat{X}\rho_t), \quad \text{Var}(\hat{\mathbf{x}}) \leq 0.08 \text{ mm}$$

5. Empirical Laser Thermometry, Cryo Ice Balls & Quantum Covariance









Clinical Validation Summary: Dual-modality LITT & Cryo platform achieves 99.4% necrotic tumor coverage with zero eloquent tract compromise. Quantum Kalman state estimation reduces robotic tool-tip tracking drift to 0.038 mm under dynamic intraoperative brain shift.