
Mathematical and numerical study of plasmonic structures with corners

Camille Carvalho¹

A.S. Bonnet-Ben Dhia¹, L. Chesnel², P. Ciarlet Jr.¹

¹POEMS, ENSTA ParisTech, Palaiseau, France

²CMAP École Polytechnique, INRIA Saclay Île-de-France, Palaiseau France




UC Merced, January 2015

What are surface plasmons ?


What are surface plasmons ?

Surface plasmons are **confined electromagnetic waves** at the interface between a metal and a dielectric.



What are surface plasmons ?


Surface plasmons are **confined electromagnetic waves** at the interface between a metal and a dielectric.



Such waves exist only if the permittivity or the permeability **changes sign**, which is possible **at optical frequencies**.

What are surface plasmons ?


Surface plasmons are **confined electromagnetic waves** at the interface between a metal and a dielectric.



Such waves exist only if the permittivity or the permeability **changes sign**, which is possible **at optical frequencies**.

What are surface plasmons ?


Surface plasmons are **confined electromagnetic waves** at the interface between a metal and a dielectric.



Such waves exist only if the permittivity or the permeability **changes sign**, which is possible **at optical frequencies**.


One simple model of permittivity: Drude's model (convention $e^{-i\omega t}$).

$$\varepsilon = \varepsilon^\gamma(\omega) := 1 - \frac{\omega_p^2}{\omega^2 + i\omega\gamma}$$



What are surface plasmons ?

Surface plasmons are **confined electromagnetic waves** at the interface between a metal and a dielectric.




Such waves exist only if the permittivity or the permeability **changes sign**, which is possible **at optical frequencies**.

One simple model of permittivity: Drude's model (convention $e^{-i\omega t}$).

$$\varepsilon = \varepsilon^\gamma(\omega) := 1 - \frac{\omega_p^2}{\omega^2 + i\omega\gamma}$$

At optical frequencies ($\gamma \ll \omega < \omega_p$),
 ε has a **negative** real part and a
neglectable imaginary part.



Applications


Applications

Guiding or confining surface plasmons in **nanophotonic devices** reveal a great interest in order to overcome the diffraction limit (optical antennas, high resolution imaging in near field, ...).



Applications


Guiding or confining surface plasmons in **nanophotonic devices** reveal a great interest in order to overcome the diffraction limit (optical antennas, high resolution imaging in near field, ...).



O'Connor et al., (2009)

Applications

Guiding or confining surface plasmons in **nanophotonic devices** reveal a great interest in order to overcome the diffraction limit (optical antennas, high resolution imaging in near field, ...).



O'Connor et al., (2009)

However these waves are very **sensitive to the geometry** of the interface between the two media.

Explicit computations of surface plasmons


Explicit computations of surface plasmons

For **simple geometries**, one can get the expression of surface plasmons.

Explicit computations of surface plasmons

For simple geometries, one can get the expression of surface plasmons.


For a 2D planar waveguide: one interface



Explicit computations of surface plasmons

For simple geometries, one can get the expression of surface plasmons.

For a 2D planar waveguide: one interface




Plasmons are solutions of Maxwell's equations in the time-harmonic regime, searched as $H_z(y)e^{i\beta x}$, $E_z(y)e^{i\beta x}$, $\beta \in \mathbb{R}$

Explicit computations of surface plasmons

For simple geometries, one can get the expression of surface plasmons.

For a 2D planar waveguide: one interface



Plasmons are solutions of Maxwell's equations in the time-harmonic regime, searched as $H_z(y)e^{i\beta x}$, $E_z(y)e^{i\beta x}$, $\beta \in \mathbb{R}$


$$\frac{d}{dy} \left(\varepsilon^{-1} \frac{dH_z}{dy} \right) + \left(\frac{\omega^2}{c^2} \mu - \beta^2 \varepsilon^{-1} \right) H_z = 0$$

$$\frac{d}{dy} \left(\mu^{-1} \frac{dE_z}{dy} \right) + \left(\frac{\omega^2}{c^2} \varepsilon - \beta^2 \mu^{-1} \right) E_z = 0$$

Explicit computations of surface plasmons

For simple geometries, one can get the expression of surface plasmons.

For a 2D planar waveguide: one interface



Plasmons are solutions of Maxwell's equations in the time-harmonic regime, searched as $H_z(y)e^{i\beta x}$, $E_z(y)e^{i\beta x}$, $\beta \in \mathbb{R}$


$$\frac{d}{dy} \left(\varepsilon^{-1} \frac{dH_z}{dy} \right) + \left(\frac{\omega^2}{c^2} \mu - \beta^2 \varepsilon^{-1} \right) H_z = 0$$

$$\frac{d}{dy} \left(\mu^{-1} \frac{dE_z}{dy} \right) + \left(\frac{\omega^2}{c^2} \varepsilon - \beta^2 \mu^{-1} \right) E_z = 0$$

Explicit computations of surface plasmons

For simple geometries, one can get the expression of surface plasmons.

For a 2D planar waveguide: one interface



Plasmons are solutions of Maxwell's equations in the time-harmonic regime, searched as $H_z(y)e^{i\beta x}$, $E_z(y)e^{i\beta x}$, $\beta \in \mathbb{R}$

$$\frac{d}{dy} \left(\varepsilon^{-1} \frac{dH_z}{dy} \right) + \left(\frac{\omega^2}{c^2} \mu - \beta^2 \varepsilon^{-1} \right) H_z = 0$$


$$\frac{d}{dy} \left(\mu^{-1} \frac{dE_z}{dy} \right) + \left(\frac{\omega^2}{c^2} \varepsilon - \beta^2 \mu^{-1} \right) E_z = 0$$

Surface plasmons exist because of the sign-changing coefficient in the principal part.

Explicit computations of surface plasmons

For simple geometries, one can get the expression of surface plasmons.

For a 2D planar waveguide: one interface




Plasmons are solutions of Maxwell's equations in the time-harmonic regime, searched as $H_z(y)e^{i\beta x}$, $E_z(y)e^{i\beta x}$, $\beta \in \mathbb{R}$

$$\frac{d}{dy} \left(\varepsilon^{-1} \frac{dH_z}{dy} \right) + \left(\frac{\omega^2}{c^2} \mu - \beta^2 \varepsilon^{-1} \right) H_z = 0$$

$$\frac{d}{dy} \left(\mu^{-1} \frac{dE_z}{dy} \right) + \left(\frac{\omega^2}{c^2} \varepsilon - \beta^2 \mu^{-1} \right) E_z = 0$$

Surface plasmons exist because of the sign-changing coefficient in the principal part.



Explicit computations of surface plasmons

For simple geometries, one can get the expression of surface plasmons.

For a 2D planar waveguide: two interfaces





Plasmons are solutions of Maxwell's equations in the time-harmonic regime, searched as $H_z(y)e^{i\beta x}$, $E_z(y)e^{i\beta x}$, $\beta \in \mathbb{R}$

$$\frac{d}{dy} \left(\varepsilon^{-1} \frac{dH_z}{dy} \right) + \left(\frac{\omega^2}{c^2} \mu - \beta^2 \varepsilon^{-1} \right) H_z = 0$$

$$\frac{d}{dy} \left(\mu^{-1} \frac{dE_z}{dy} \right) + \left(\frac{\omega^2}{c^2} \varepsilon - \beta^2 \mu^{-1} \right) E_z = 0$$

Surface plasmons exist because of the sign-changing coefficient in the principal part.



Explicit computations of surface plasmons

For simple geometries, one can get the expression of surface plasmons.

For a 2D planar waveguide: two interfaces





Plasmons are solutions of Maxwell's equations in the time-harmonic regime, searched as $H_z(y)e^{i\beta x}$, $E_z(y)e^{i\beta x}$, $\beta \in \mathbb{R}$

$$\frac{d}{dy} \left(\varepsilon^{-1} \frac{dH_z}{dy} \right) + \left(\frac{\omega^2}{c^2} \mu - \beta^2 \varepsilon^{-1} \right) H_z = 0$$

$$\frac{d}{dy} \left(\mu^{-1} \frac{dE_z}{dy} \right) + \left(\frac{\omega^2}{c^2} \varepsilon - \beta^2 \mu^{-1} \right) E_z = 0$$

Surface plasmons exist because of the sign-changing coefficient in the principal part.



How to compute surface plasmons in more complex geometries ?

Problems with sign-changing coefficients

Problems with sign-changing coefficients

Plasmonics belong the class of problems called **problems with sign-changing coefficients**.




Problems with sign-changing coefficients

Plasmonics belong the class of problems called **problems with sign-changing coefficients**.

Example: a 2D transmission problem in a bounded domain.

Find $u \in H_0^1(\Omega)$ such that:

$$\operatorname{div}(\varepsilon^{-1} \nabla u) + \omega^2 \mu u = -f \quad \text{in } \Omega$$




Problems with sign-changing coefficients

Plasmonics belong the class of problems called **problems with sign-changing coefficients**.

Example: a 2D transmission problem in a bounded domain.

$$\begin{aligned} & \text{Find } u \in H_0^1(\Omega) \text{ such that:} \\ & \operatorname{div}(\varepsilon^{-1} \nabla u) + \omega^2 \mu u = -f \quad \text{in } \Omega \end{aligned}$$



The variational formulation writes:

$$\begin{aligned} & \text{Find } u \in H_0^1(\Omega) \text{ such that:} \\ & a(u, v) + c(u, v) = \langle f, v \rangle \quad \forall v \in H_0^1(\Omega), \\ & a(u, v) = \int_{\Omega} \varepsilon^{-1} \nabla u \cdot \overline{\nabla v}, \quad c(u, v) = -\omega^2 \int_{\Omega} \mu u \bar{v}. \end{aligned}$$


Problems with sign-changing coefficients

Plasmonics belong the class of problems called **problems with sign-changing coefficients**.

Example: a 2D transmission problem in a bounded domain.

Find $u \in H_0^1(\Omega)$ such that:

$$\operatorname{div}(\varepsilon^{-1} \nabla u) + \omega^2 \mu u = -f \quad \text{in } \Omega$$



The variational formulation writes:

Find $u \in H_0^1(\Omega)$ such that:

$$a(u, v) + c(u, v) = \langle f, v \rangle \quad \forall v \in H_0^1(\Omega),$$

$$a(u, v) = \int_{\Omega} \varepsilon^{-1} \nabla u \cdot \overline{\nabla v}, \quad c(u, v) = -\omega^2 \int_{\Omega} \mu u \bar{v}.$$

$$a(u, u) \geq C \|u\|_{H_0^1(\Omega)}^2 \quad \forall u \in H_0^1(\Omega)$$

Classically, for $\varepsilon > 0$:

$a(\cdot, \cdot)$ is **coercive** and $c(\cdot, \cdot)$ is a **compact perturbation**: problem of **Fredholm type**.


Problems with sign-changing coefficients

Plasmonics belong the class of problems called **problems with sign-changing coefficients**.

Example: a 2D transmission problem in a bounded domain.

Find $u \in H_0^1(\Omega)$ such that:

$$\operatorname{div}(\varepsilon^{-1} \nabla u) + \omega^2 \mu u = -f \quad \text{in } \Omega$$



The variational formulation writes:

Find $u \in H_0^1(\Omega)$ such that:

$$a(u, v) + c(u, v) = \langle f, v \rangle \quad \forall v \in H_0^1(\Omega),$$

$$a(u, v) = \int_{\Omega} \varepsilon^{-1} \nabla u \cdot \overline{\nabla v}, \quad c(u, v) = -\omega^2 \int_{\Omega} \mu u \bar{v}.$$

$$a(u, u) \geq C \|u\|_{H_0^1(\Omega)}^2 \quad \forall u \in H_0^1(\Omega)$$

Classically, for $\varepsilon > 0$:

$a(\cdot, \cdot)$ is **coercive** and $c(\cdot, \cdot)$ is a **compact perturbation**: problem of **Fredholm type**.

When ε is sign-changing, the form a is no longer coercive !


Problems with sign-changing coefficients

Plasmonics belong the class of problems called **problems with sign-changing coefficients**.

Example: a 2D transmission problem in a bounded domain.

Find $u \in H_0^1(\Omega)$ such that:

$$\operatorname{div}(\varepsilon^{-1} \nabla u) + \omega^2 \mu u = -f \quad \text{in } \Omega$$



The variational formulation writes:

Find $u \in H_0^1(\Omega)$ such that:

$$a(u, v) + c(u, v) = \langle f, v \rangle \quad \forall v \in H_0^1(\Omega),$$

$$a(u, v) = \int_{\Omega} \varepsilon^{-1} \nabla u \cdot \overline{\nabla v}, \quad c(u, v) = -\omega^2 \int_{\Omega} \mu u \bar{v}.$$

$$a(u, u) \geq C \|u\|_{H_0^1(\Omega)}^2 \quad \forall u \in H_0^1(\Omega)$$

Classically, for $\varepsilon > 0$:

$a(\cdot, \cdot)$ is **coercive** and $c(\cdot, \cdot)$ is a **compact perturbation**: problem of **Fredholm type**.

When ε is sign-changing, the form a is no longer coercive !


The problem may be **ill-posed** for certain values of the **contrast** $\kappa_{\varepsilon} := \frac{\varepsilon_m}{\varepsilon_d} < 0$.

The T-coercivity approach

Find $u \in H_0^1(\Omega)$ such that:

$$a(u, v) + c(u, v) = \langle f, v \rangle \quad \forall v \in H_0^1(\Omega)$$

$$\kappa_\varepsilon := \frac{\varepsilon_m}{\varepsilon_d} < 0$$




The T-coercivity approach

Find $u \in H_0^1(\Omega)$ such that:

$$a(u, v) + c(u, v) = \langle f, v \rangle \quad \forall v \in H_0^1(\Omega)$$

$$\kappa_\varepsilon := \frac{\varepsilon_m}{\varepsilon_d} < 0$$

The **T-coercivity** theory allows to prove Fredholmness under some **conditions on the contrast κ_ε** and on the **geometry of Σ** .




The T-coercivity approach

Find $u \in H_0^1(\Omega)$ such that:

$$a(u, v) + c(u, v) = \langle f, v \rangle \quad \forall v \in H_0^1(\Omega)$$

$$\kappa_\varepsilon := \frac{\varepsilon_m}{\varepsilon_d} < 0$$

The **T-coercivity** theory allows to prove Fredholmness under some **conditions on the contrast κ_ε** and on the **geometry of Σ** .



Idea: build an **isomorphism \mathbf{T}** of $H_0^1(\Omega)$ such that the following problem is **coercive+compact**

Find $u \in H_0^1(\Omega)$ such that:

$$a(u, \mathbf{T}v) + c(u, \mathbf{T}v) = \langle f, \mathbf{T}v \rangle \quad \forall v \in H_0^1(\Omega)$$

Introduced in



Zwölf (2008).

Complete study in



Chesnel (2012) which gave the following result:


The T-coercivity approach

Find $u \in H_0^1(\Omega)$ such that:

$$a(u, v) + c(u, v) = \langle f, v \rangle \quad \forall v \in H_0^1(\Omega)$$

$$\kappa_\varepsilon := \frac{\varepsilon_m}{\varepsilon_d} < 0$$

The **T-coercivity** theory allows to prove Fredholmness under some **conditions on the contrast κ_ε** and on the **geometry of Σ** .



Idea: build an **isomorphism \mathbf{T}** of $H_0^1(\Omega)$ such that the following problem is **coercive+compact**

Find $u \in H_0^1(\Omega)$ such that:

$$a(u, \mathbf{T}v) + c(u, \mathbf{T}v) = \langle f, \mathbf{T}v \rangle \quad \forall v \in H_0^1(\Omega)$$

Introduced in



Zwölf (2008).

Complete study in



Chesnel (2012) which gave the following result:

The problem is of Fredholm type if and only if $\kappa_\varepsilon \notin I_c$, I_c is called **critical interval**.


The T-coercivity approach

Find $u \in H_0^1(\Omega)$ such that:

$$a(u, v) + c(u, v) = \langle f, v \rangle \quad \forall v \in H_0^1(\Omega)$$

$$\kappa_\varepsilon := \frac{\varepsilon_m}{\varepsilon_d} < 0$$

The **T-coercivity** theory allows to prove Fredholmness under some **conditions on the contrast κ_ε** and on the **geometry of Σ** .



Idea: build an **isomorphism \mathbf{T}** of $H_0^1(\Omega)$ such that the following problem is **coercive+compact**

Find $u \in H_0^1(\Omega)$ such that:

$$a(u, \mathbf{T}v) + c(u, \mathbf{T}v) = \langle f, \mathbf{T}v \rangle \quad \forall v \in H_0^1(\Omega)$$

Introduced in  Zwölf (2008).

Complete study in  Chesnel (2012) which gave the following result:

The problem is of Fredholm type if and only if $\kappa_\varepsilon \notin I_c$, I_c is called **critical interval**.

I_c always contain $\{-1\}$.


The T-coercivity approach

Find $u \in H_0^1(\Omega)$ such that:

$$a(u, v) + c(u, v) = \langle f, v \rangle \quad \forall v \in H_0^1(\Omega)$$

$$\kappa_\varepsilon := \frac{\varepsilon_m}{\varepsilon_d} < 0$$

The T-coercivity theory allows to prove Fredholmness under some conditions on the contrast κ_ε and on the geometry of Σ .



Idea: build an isomorphism T of $H_0^1(\Omega)$ such that the following problem is coercive+compact

Find $u \in H_0^1(\Omega)$ such that:

$$a(u, Tv) + c(u, Tv) = \langle f, Tv \rangle \quad \forall v \in H_0^1(\Omega)$$

Introduced in  Zwölf (2008).

Complete study in  Chesnel (2012) which gave the following result:

The problem is of Fredholm type if and only if $\kappa_\varepsilon \notin I_c$, I_c is called critical interval.

I_c always contain $\{-1\}$.


The case $\kappa_\varepsilon = -1$ is particularly problematic, we will not talk about it.



Ola (1995), Nguyen (2015).

Results

Find $u \in H_0^1(\Omega)$ such that:


$$\operatorname{div}(\varepsilon^{-1} \nabla u) + \omega^2 \mu u = -f \quad \text{in } \Omega$$


Results

Find $u \in H_0^1(\Omega)$ such that:

$$\operatorname{div}(\varepsilon^{-1} \nabla u) + \omega^2 \mu u = -f \quad \text{in } \Omega$$

«Outside the critical interval» $\kappa_\varepsilon \notin I_c$



Results

Find $u \in H_0^1(\Omega)$ such that:


$$\operatorname{div}(\varepsilon^{-1} \nabla u) + \omega^2 \mu u = -f \quad \text{in } \Omega$$

«Outside the critical interval» $\kappa_\varepsilon \notin I_c$

The problem is of Fredholm type. Characterization of I_c for any 2D configuration.



Bonnet-Ben Dhia, Chesnel, Ciarlet (2012).




Results

Find $u \in H_0^1(\Omega)$ such that:

$$\operatorname{div}(\varepsilon^{-1} \nabla u) + \omega^2 \mu u = -f \quad \text{in } \Omega$$

«Outside the critical interval» $\kappa_\varepsilon \notin I_c$



The problem is of Fredholm type. Characterization of I_c for any 2D configuration.



Bonnet-Ben Dhia, Chesnel, Ciarlet (2012).

Meshing conditions implying finite elements method convergence have been provided.




Bonnet-Ben Dhia, Ciarlet, Zwölf (2010), Nicaise, Venel (2011), Chesnel, Ciarlet (2013).

Results

Find $u \in H_0^1(\Omega)$ such that:

$$\operatorname{div}(\varepsilon^{-1} \nabla u) + \omega^2 \mu u = -f \quad \text{in } \Omega$$

«Outside the critical interval» $\kappa_\varepsilon \notin I_c$



The problem is of Fredholm type. Characterization of I_c for any 2D configuration.



Bonnet-Ben Dhia, Chesnel, Ciarlet (2012).

Meshing conditions implying finite elements method convergence have been provided.



Bonnet-Ben Dhia, Ciarlet, Zwölf (2010), Nicaise, Venel (2011), Chesnel, Ciarlet (2013).


Non optimal results for the discretization in the corners case.

Results

Find $u \in H_0^1(\Omega)$ such that:

$$\operatorname{div}(\varepsilon^{-1} \nabla u) + \omega^2 \mu u = -f \quad \text{in } \Omega$$

«Outside the critical interval» $\kappa_\varepsilon \notin I_c$



The problem is of Fredholm type. Characterization of I_c for any 2D configuration.



Bonnet-Ben Dhia, Chesnel, Ciarlet (2012).

Meshing conditions implying finite elements method convergence have been provided.



Bonnet-Ben Dhia, Ciarlet, Zwölf (2010), Nicaise, Venel (2011), Chesnel, Ciarlet (2013).

Non optimal results for the discretization in the corners case.


«Inside the critical interval» $\kappa_\varepsilon \in I_c$

Results

Find $u \in H_0^1(\Omega)$ such that:

$$\operatorname{div}(\varepsilon^{-1} \nabla u) + \omega^2 \mu u = -f \quad \text{in } \Omega$$

«Outside the critical interval» $\kappa_\varepsilon \notin I_c$



The problem is of Fredholm type. Characterization of I_c for any 2D configuration.



Bonnet-Ben Dhia, Chesnel, Ciarlet (2012).

Meshing conditions implying finite elements method convergence have been provided.



Bonnet-Ben Dhia, Ciarlet, Zwölf (2010), Nicaise, Venel (2011), Chesnel, Ciarlet (2013).

Non optimal results for the discretization in the corners case.

«Inside the critical interval» $\kappa_\varepsilon \in I_c$

The problem is no longer Fredholm due to hyper oscillating singularities at corners.

Fredholmness is recovered in a new functional framework taking into account the singularities.




Bonnet-Ben Dhia, Dauge, Ramdani (1999), Bonnet-Ben Dhia, Chesnel, Claeys (2013).

Results

Find $u \in H_0^1(\Omega)$ such that:

$$\operatorname{div}(\varepsilon^{-1} \nabla u) + \omega^2 \mu u = -f \quad \text{in } \Omega$$

«Outside the critical interval» $\kappa_\varepsilon \notin I_c$



The problem is of Fredholm type. Characterization of I_c for any 2D configuration.



Bonnet-Ben Dhia, Chesnel, Ciarlet (2012).

Meshing conditions implying finite elements method convergence have been provided.




Bonnet-Ben Dhia, Ciarlet, Zwölf (2010), Nicaise, Venel (2011), Chesnel, Ciarlet (2013).

Non optimal results for the discretization in the corners case.

«Inside the critical interval» $\kappa_\varepsilon \in I_c$

The problem is no longer Fredholm due to hyper oscillating singularities at corners.

Fredholmness is recovered in a new functional framework taking into account the singularities.




Bonnet-Ben Dhia, Dauge, Ramdani (1999), Bonnet-Ben Dhia, Chesnel, Claeys (2013).

No finite elements method convergence.

Results

Find $u \in H_0^1(\Omega)$ such that:

$$\operatorname{div}(\varepsilon^{-1} \nabla u) + \omega^2 \mu u = -f \quad \text{in } \Omega$$



«Outside the critical interval» $\kappa_\varepsilon \notin I_c$

The problem is of Fredholm type. Characterization of I_c for any 2D configuration.



Bonnet-Ben Dhia, Chesnel, Ciarlet (2012).

Meshing conditions implying finite elements method convergence have been provided.



Bonnet-Ben Dhia, Ciarlet, Dauge, Ramdani (2010), Melenk, Melenk, Melenk (2011), Gatica, Gatica, Gatica (2013).

Non optimal results for

Numerical method when the interface
presents corners ?

«Inside the critical int

The problem is no longer Fredholm due to singularities at corners.

Fredholmness is recovered in a new functional framework taking into account the singularities.



Bonnet-Ben Dhia, Dauge, Ramdani (1999), Bonnet-Ben Dhia, Chesnel, Claeys (2013).

No finite elements method convergence.

Goals of my work

Goals of my work

Develop a numerical method when the interface presents corners.

Goals of my work

Develop a numerical method when the interface presents corners.

«Outside the critical interval»



Goals of my work

Develop a numerical method when the interface presents corners.

«Outside the critical interval»

«Inside the critical interval»


Goals of my work

Develop a numerical method when the interface presents corners.

«Outside the critical interval»

«Inside the critical interval»

Application to two physical problems



Goals of my work

Develop a numerical method when the interface presents corners.

«Outside the critical interval»

To get FE convergence, one has to realize T-coercivity at the discrete level
(it can be ensured with a conformity of the mesh).

«Inside the critical interval»

Application to two physical problems



Goals of my work

Develop a numerical method when the interface presents corners.

«Outside the critical interval»

To get FE convergence, one has to realize T-coercivity at the discrete level
(it can be ensured with a conformity of the mesh).

«Inside the critical interval»

Analysis of the hyper oscillating singularity, and provide a FE-based numerical method.

Application to two physical problems



Goals of my work

Develop a numerical method when the interface presents corners.

«Outside the critical interval»

To get FE convergence, one has to realize T-coercivity at the discrete level
(it can be ensured with a conformity of the mesh).

«Inside the critical interval»

Analysis of the hyper oscillating singularity, and provide a FE-based numerical method.

Application to two physical problems

- 1) Scattering problem with a polygonal metallic inclusion.
- 2) The search of guided modes in a plasmonic waveguide.



Goals of my work

Develop a numerical method when the interface presents corners.

«Outside the critical interval»

To get FE convergence, one has to realize T-coercivity at the discrete level
(it can be ensured with a conformity of the mesh).

«Inside the critical interval»

Analysis of the hyper oscillating singularity, and provide a FE-based numerical method.

Application to two physical problems

- 1) Scattering problem with a polygonal metallic inclusion.
- 2) The search of guided modes in a plasmonic waveguide.



Goals of my work

Develop a numerical method when the interface presents corners.

«Outside the critical interval»

To get FE convergence, one has to realize T-coercivity at the discrete level
(it can be ensured with a conformity of the mesh).

«Inside the critical interval»

Analysis of the hyper oscillating singularity, and provide a FE-based numerical method.

Application to two physical problems

- 1) Scattering problem with a polygonal metallic inclusion.
- 2) The search of guided modes in a plasmonic waveguide.

Outline

- ✿ Part I: meshing rules for the corners case for $\kappa_\varepsilon \notin I_c$
- ✿ Part II: scattering problem with sign-changing coefficients
- ✿ Perspectives




Outline

- Part I: meshing rules for the corners case for $\kappa_\varepsilon \notin I_c$

Transmission problem and T-coercivity

Goal: build an **isomorphism \mathbf{T}** of $H_0^1(\Omega)$ such that the form a is **T-coercive**: $\forall u \in H_0^1(\Omega)$


$$a(u, \mathbf{T}u) \geq C\|u\|_{H_0^1(\Omega)}^2 \quad \text{with} \quad a(u, v) = \int_{\Omega} \varepsilon^{-1} \nabla u \cdot \overline{\nabla v}$$



Transmission problem and T-coercivity

Goal: build an **isomorphism \mathbf{T}** of $H_0^1(\Omega)$ such that the form a is **T-coercive**: $\forall u \in H_0^1(\Omega)$

$$a(u, \mathbf{T}u) \geq C\|u\|_{H_0^1(\Omega)}^2 \quad \text{with} \quad a(u, v) = \int_{\Omega} \varepsilon^{-1} \nabla u \cdot \overline{\nabla v}$$




1) **compensate** the change of sign of ε

$$\mathbf{T}u = \begin{cases} u_d & \text{in } \Omega_d \\ -u_m + \dots & \text{in } \Omega_m \end{cases}$$

Transmission problem and T-coercivity

Goal: build an **isomorphism \mathbf{T}** of $H_0^1(\Omega)$ such that the form a is **\mathbf{T} -coercive**: $\forall u \in H_0^1(\Omega)$

$$a(u, \mathbf{T}u) \geq C\|u\|_{H_0^1(\Omega)}^2 \quad \text{with} \quad a(u, v) = \int_{\Omega} \varepsilon^{-1} \nabla u \cdot \overline{\nabla v}$$




- 1) **compensate** the change of sign of ε
- 2) ensure the **matching** with a **transfert operator R**

$$\mathbf{T}u = \begin{cases} u_d & \text{in } \Omega_d \\ -u_m + 2Ru_d & \text{in } \Omega_m \end{cases}$$

Transmission problem and T-coercivity

Goal: build an **isomorphism \mathbf{T}** of $H_0^1(\Omega)$ such that the form a is **T-coercive**: $\forall u \in H_0^1(\Omega)$

$$a(u, \mathbf{T}u) \geq C\|u\|_{H_0^1(\Omega)}^2 \quad \text{with} \quad a(u, v) = \int_{\Omega} \varepsilon^{-1} \nabla u \cdot \overline{\nabla v}$$



- 1) **compensate** the change of sign of ε
- 2) ensure the **matching** with a **transfert operator R**


$$\mathbf{T}u = \begin{cases} u_d & \text{in } \Omega_d \\ -u_m + 2Ru_d & \text{in } \Omega_m \end{cases}$$

\mathbf{R} is a linear continuous operator from Ω_d into Ω_m such that $\mathbf{R}u|_{\Sigma} = u|_{\Sigma}$

Transmission problem and T-coercivity

Goal: build an **isomorphism \mathbf{T}** of $H_0^1(\Omega)$ such that the form a is **T-coercive**: $\forall u \in H_0^1(\Omega)$

$$a(u, \mathbf{T}u) \geq C\|u\|_{H_0^1(\Omega)}^2 \quad \text{with} \quad a(u, v) = \int_{\Omega} \varepsilon^{-1} \nabla u \cdot \overline{\nabla v}$$



- 1) **compensate** the change of sign of ε
- 2) ensure the **matching** with a **transfert operator R**

$$\mathbf{T}u = \begin{cases} u_d & \text{in } \Omega_d \\ -u_m + 2Ru_d & \text{in } \Omega_m \end{cases}$$


\mathbf{R} is a linear continuous operator from Ω_d into Ω_m such that $\mathbf{R}u|_{\Sigma} = u|_{\Sigma}$

$\mathbf{T} \circ \mathbf{T} = I_d$ then \mathbf{T} is an isomorphism of $H_0^1(\Omega)$.

Transmission problem and T-coercivity

Goal: build an **isomorphism \mathbf{T}** of $H_0^1(\Omega)$ such that the form a is **\mathbf{T} -coercive**: $\forall u \in H_0^1(\Omega)$

$$a(u, \mathbf{T}u) \geq C\|u\|_{H_0^1(\Omega)}^2 \quad \text{with} \quad a(u, v) = \int_{\Omega} \varepsilon^{-1} \nabla u \cdot \overline{\nabla v}$$



- 1) **compensate** the change of sign of ε
- 2) ensure the **matching** with a **transfert operator R**

$$\mathbf{T}u = \begin{cases} u_d & \text{in } \Omega_d \\ -u_m + 2\mathbf{R}u_d & \text{in } \Omega_m \end{cases} \quad \mathbf{T}'u = \begin{cases} u_d - 2\mathbf{R}'u_m & \text{in } \Omega_d \\ -u_m & \text{in } \Omega_m \end{cases}$$

\mathbf{R} is a linear continuous operator from Ω_d into Ω_m such that $\mathbf{R}u|_{\Sigma} = u|_{\Sigma}$


\mathbf{R}' is a linear continuous operator from Ω_m into Ω_d such that $\mathbf{R}'u|_{\Sigma} = u|_{\Sigma}$

$\mathbf{T} \circ \mathbf{T} = \mathbf{I}_d$ then \mathbf{T} is an isomorphism of $H_0^1(\Omega)$.

Transmission problem and T-coercivity

Goal: build an **isomorphism \mathbf{T}** of $H_0^1(\Omega)$ such that the form a is **\mathbf{T} -coercive**: $\forall u \in H_0^1(\Omega)$

$$a(u, \mathbf{T}u) \geq C\|u\|_{H_0^1(\Omega)}^2 \quad \text{with} \quad a(u, v) = \int_{\Omega} \varepsilon^{-1} \nabla u \cdot \overline{\nabla v}$$



- 1) compensate the change of sign of ε
- 2) ensure the matching with a **transfert operator R**

$$\mathbf{T}u = \begin{cases} u_d & \text{in } \Omega_d \\ -u_m + 2\mathbf{R}u_d & \text{in } \Omega_m \end{cases} \quad \mathbf{T}'u = \begin{cases} u_d - 2\mathbf{R}'u_m & \text{in } \Omega_d \\ -u_m & \text{in } \Omega_m \end{cases}$$

\mathbf{R} is a linear continuous operator from Ω_d into Ω_m such that $\mathbf{R}u|_{\Sigma} = u|_{\Sigma}$

\mathbf{R}' is a linear continuous operator from Ω_m into Ω_d such that $\mathbf{R}'u|_{\Sigma} = u|_{\Sigma}$


$\mathbf{T} \circ \mathbf{T} = \mathbf{I}_d$ then \mathbf{T} is an isomorphism of $H_0^1(\Omega)$.

Theorem. The form is **\mathbf{T} -coercive** if $\kappa_{\varepsilon} \notin [-\|\mathbf{R}'\|^2; -1/\|\mathbf{R}\|^2]$

Transmission problem and T-coercivity

Goal: build an **isomorphism \mathbf{T}** of $H_0^1(\Omega)$ such that the form a is **\mathbf{T} -coercive**: $\forall u \in H_0^1(\Omega)$

$$a(u, \mathbf{T}u) \geq C\|u\|_{H_0^1(\Omega)}^2 \quad \text{with} \quad a(u, v) = \int_{\Omega} \varepsilon^{-1} \nabla u \cdot \overline{\nabla v}$$



- 1) compensate the change of sign of ε
- 2) ensure the matching with a **transfert operator R**

$$\mathbf{T}u = \begin{cases} u_d & \text{in } \Omega_d \\ -u_m + 2\mathbf{R}u_d & \text{in } \Omega_m \end{cases} \quad \mathbf{T}'u = \begin{cases} u_d - 2\mathbf{R}'u_m & \text{in } \Omega_d \\ -u_m & \text{in } \Omega_m \end{cases}$$

\mathbf{R} is a linear continuous operator from Ω_d into Ω_m such that $\mathbf{R}u|_{\Sigma} = u|_{\Sigma}$

\mathbf{R}' is a linear continuous operator from Ω_m into Ω_d such that $\mathbf{R}'u|_{\Sigma} = u|_{\Sigma}$

$\mathbf{T} \circ \mathbf{T} = \mathbf{I}_d$ then \mathbf{T} is an isomorphism of $H_0^1(\Omega)$.


Theorem. The form is **\mathbf{T} -coercive** if $\kappa_{\varepsilon} \notin [-\|\mathbf{R}'\|^2; -1/\|\mathbf{R}\|^2]$

Next : build operators \mathbf{R} , \mathbf{R}' minimizing the interval. To do so we use elementary geometrical transforms (symmetries, angular dilations, ...).

$$\mathbf{R}u(x) = u(\mathcal{R}(x))$$

Some examples


For a **symmetric domain** (w.r.t. Σ) :



Some examples

For a **symmetric domain** (w.r.t. Σ) :

$$\mathbf{R}u(x) = \mathbf{R}'u(x) = u(\mathcal{S}_\Sigma(x)) \quad \|\mathbf{R}\|^2 = \|\mathbf{R}'\|^2 = 1$$




Some examples

For a **symmetric domain** (w.r.t. Σ) :

$$\mathbf{R}u(x) = \mathbf{R}'u(x) = u(\mathcal{S}_\Sigma(x)) \quad \|\mathbf{R}\|^2 = \|\mathbf{R}'\|^2 = 1$$

Theorem. The form is **T-coercive** $\iff \kappa_\varepsilon \neq -1$




Some examples


For a **symmetric domain** (w.r.t. Σ) :

$$\mathbf{R}u(x) = \mathbf{R}'u(x) = u(\mathcal{S}_\Sigma(x)) \quad \|\mathbf{R}\|^2 = \|\mathbf{R}'\|^2 = 1$$

Theorem. The form is **T-coercive** $\iff \kappa_\varepsilon \neq -1$



For an **angular sector** of aperture α :




Bonnet-Ben Dhia, Chesnel, Ciarlet (2012).

Some examples

For a **symmetric domain** (w.r.t. Σ) :


$$\mathbf{R}u(x) = \mathbf{R}'u(x) = u(\mathcal{S}_\Sigma(x)) \quad \|\mathbf{R}\|^2 = \|\mathbf{R}'\|^2 = 1$$

Theorem. The form is **T-coercive** $\iff \kappa_\varepsilon \neq -1$



For an **angular sector** of aperture α :

Operators \mathbf{R}, \mathbf{R}' can be built from the composition of a central symmetry and an angular dilation.




Bonnet-Ben Dhia, Chesnel, Ciarlet (2012).

Some examples

For a **symmetric domain** (w.r.t. Σ) :

$$\mathbf{R}u(x) = \mathbf{R}'u(x) = u(\mathcal{S}_\Sigma(x)) \quad \|\mathbf{R}\|^2 = \|\mathbf{R}'\|^2 = 1$$


Theorem. The form is **T-coercive** $\iff \kappa_\varepsilon \neq -1$



For an **angular sector** of aperture α :

Operators \mathbf{R}, \mathbf{R}' can be built from the composition of a central symmetry and an angular dilation.

$$\|\mathbf{R}\|^2 = \|\mathbf{R}'\|^2 = I_\alpha := \max\left(\frac{\alpha}{2\pi - \alpha}; \frac{2\pi - \alpha}{\alpha}\right) \quad I_\alpha \geqslant 1$$




Bonnet-Ben Dhia, Chesnel, Ciarlet (2012).

Some examples

For a **symmetric domain** (w.r.t. Σ) :

$$\mathbf{R}u(x) = \mathbf{R}'u(x) = u(\mathcal{S}_\Sigma(x)) \quad \|\mathbf{R}\|^2 = \|\mathbf{R}'\|^2 = 1$$


Theorem. The form is T-coercive $\iff \kappa_\varepsilon \neq -1$



For an **angular sector** of aperture α :

Operators \mathbf{R}, \mathbf{R}' can be built from the composition of a central symmetry and an angular dilation.

$$\|\mathbf{R}\|^2 = \|\mathbf{R}'\|^2 = I_\alpha := \max\left(\frac{\alpha}{2\pi - \alpha}; \frac{2\pi - \alpha}{\alpha}\right) \quad I_\alpha \geqslant 1$$



Theorem. The form is T-coercive $\iff \kappa_\varepsilon \notin I_c := [-I_\alpha; -1/I_\alpha]$




Bonnet-Ben Dhia, Chesnel, Ciarlet (2012).

Some examples

For a **symmetric domain** (w.r.t. Σ) :

$$\mathbf{R}u(x) = \mathbf{R}'u(x) = u(\mathcal{S}_\Sigma(x)) \quad \|\mathbf{R}\|^2 = \|\mathbf{R}'\|^2 = 1$$


Theorem. The form is T-coercive $\iff \kappa_\varepsilon \neq -1$



For an **angular sector** of aperture α :

Operators \mathbf{R}, \mathbf{R}' can be built from the composition of a central symmetry and an angular dilation.

$$\|\mathbf{R}\|^2 = \|\mathbf{R}'\|^2 = I_\alpha := \max\left(\frac{\alpha}{2\pi - \alpha}; \frac{2\pi - \alpha}{\alpha}\right) \quad I_\alpha \geq 1$$



Theorem. The form is T-coercive $\iff \kappa_\varepsilon \notin I_c := [-I_\alpha; -1/I_\alpha]$

If $\alpha \rightarrow 0$ or $\alpha \rightarrow 2\pi$ then $I_c \rightarrow \mathbb{R}^-$.




Bonnet-Ben Dhia, Chesnel, Ciarlet (2012).

Some examples

For a **symmetric domain** (w.r.t. Σ) :

$$\mathbf{R}u(x) = \mathbf{R}'u(x) = u(\mathcal{S}_\Sigma(x)) \quad \|\mathbf{R}\|^2 = \|\mathbf{R}'\|^2 = 1$$


Theorem. The form is T-coercive $\iff \kappa_\varepsilon \neq -1$



For an **angular sector** of aperture α :

Operators \mathbf{R}, \mathbf{R}' can be built from the composition of a central symmetry and an angular dilation.

$$\|\mathbf{R}\|^2 = \|\mathbf{R}'\|^2 = I_\alpha := \max\left(\frac{\alpha}{2\pi - \alpha}; \frac{2\pi - \alpha}{\alpha}\right) \quad I_\alpha \geqslant 1$$



Theorem. The form is T-coercive $\iff \kappa_\varepsilon \notin I_c := [-I_\alpha; -1/I_\alpha]$


If $\alpha \rightarrow 0$ or $\alpha \rightarrow 2\pi$ then $I_c \rightarrow \mathbb{R}^-$. If $\alpha \rightarrow \pi$ then $I_c \rightarrow \{-1\}$.



Bonnet-Ben Dhia, Chesnel, Ciarlet (2012).

Case of a polygonal interface

For a polygonal interface, one simply use **locally** the two previous cases.




Case of a polygonal interface

For a polygonal interface, one simply use **locally** the two previous cases.

$$\mathbf{T}u = \begin{cases} u_d & \text{in } \Omega_d \\ -u_m + 2 \sum_p \chi_p \mathbf{R}_p u_d & \text{in } \Omega_m \end{cases}$$


$(\chi_p)_p$ form a partition's unity in the neighborhood of the interface.



Case of a polygonal interface

For a polygonal interface, one simply use **locally** the two previous cases.

$$\mathbf{T}u = \begin{cases} u_d & \text{in } \Omega_d \\ -u_m + 2 \sum_p \chi_p \mathbf{R}_p u_d & \text{in } \Omega_m \end{cases}$$



$(\chi_p)_p$ form a partition's unity in the neighborhood of the interface.

Theoreme. The transmission problem

$$\begin{cases} \text{Find } u \in H_0^1(\Omega) \text{ such that:} \\ a(u, v) + c(u, v) = \langle f, v \rangle \quad \forall v \in H_0^1(\Omega), \\ a(u, v) = \int_{\Omega} \varepsilon^{-1} \nabla u \cdot \overline{\nabla v}, \quad c(u, v) = -\omega^2 \int_{\Omega} \mu u \overline{v}, \end{cases}$$


is Fredholm if and only if $\kappa_{\varepsilon} \notin I_c := [-I_{\alpha_{\max}}; -1/I_{\alpha_{\max}}]$.

$$I_{\alpha_{\max}} := \max_{\alpha} I_{\alpha}$$

Case of a polygonal interface

For a polygonal interface, one simply use **locally** the two previous cases.

$$\mathbf{T}u = \begin{cases} u_d & \text{in } \Omega_d \\ -u_m + 2 \sum_p \chi_p \mathbf{R}_p u_d & \text{in } \Omega_m \end{cases}$$



$(\chi_p)_p$ form a partition's unity in the neighborhood of the interface.

Theoreme. The transmission problem

$$\begin{cases} \text{Find } u \in H_0^1(\Omega) \text{ such that:} \\ a(u, v) + c(u, v) = \langle f, v \rangle \quad \forall v \in H_0^1(\Omega), \\ a(u, v) = \int_{\Omega} \varepsilon^{-1} \nabla u \cdot \overline{\nabla v}, \quad c(u, v) = -\omega^2 \int_{\Omega} \mu u \overline{v}, \end{cases}$$


is Fredholm if and only if $\kappa_{\varepsilon} \notin I_c := [-I_{\alpha_{\max}}; -1/I_{\alpha_{\max}}]$. Moreover if $\kappa_{\varepsilon} \notin I_c$ the problem is well-posed except for discrete sequence of frequencies ω .

$$I_{\alpha_{\max}} := \max_{\alpha} I_{\alpha}$$

Case of a polygonal interface

For a polygonal interface, one simply use **locally** the two previous cases.

$$\mathbf{T}u = \begin{cases} u_d & \text{in } \Omega_d \\ -u_m + 2 \sum_p \chi_p \mathbf{R}_p u_d & \text{in } \Omega_m \end{cases}$$



$(\chi_p)_p$ form a partition's unity in the neighborhood of the interface.

Theoreme. The transmission problem

$$\begin{cases} \text{Find } u \in H_0^1(\Omega) \text{ such that:} \\ a(u, v) + c(u, v) = \langle f, v \rangle \quad \forall v \in H_0^1(\Omega), \\ a(u, v) = \int_{\Omega} \varepsilon^{-1} \nabla u \cdot \overline{\nabla v}, \quad c(u, v) = -\omega^2 \int_{\Omega} \mu u \overline{v}, \end{cases}$$

is Fredholm if and only if $\kappa_{\varepsilon} \notin I_c := [-I_{\alpha_{\max}}; -1/I_{\alpha_{\max}}]$. Moreover if $\kappa_{\varepsilon} \notin I_c$ the problem is well-posed except for discrete sequence of frequencies ω .

$$I_{\alpha_{\max}} := \max_{\alpha} I_{\alpha}$$

What about the discretization ?

Discretization of the problem

If $\kappa_\varepsilon \notin I_c$ the the problem is of **Fredholm type**. Consider the case where there is a **unique** solution u .

Discretization of the problem

If $\kappa_\varepsilon \notin I_c$ the the problem is of **Fredholm type**. Consider the case where there is a **unique solution u** .

Find $u \in V := H_0^1(\Omega)$ such that:

$$a(u, v) + c(u, v) = \langle f, v \rangle \quad \forall v \in V$$

Discretization of the problem

If $\kappa_\varepsilon \notin I_c$ the the problem is of **Fredholm type**. Consider the case where there is a **unique solution u** .

Find $u \in V := H_0^1(\Omega)$ such that:

$$a(u, v) + c(u, v) = \langle f, v \rangle \quad \forall v \in V$$

Find $u_h \in V^h$ such that:

$$a(u_h, v_h) + c(u_h, v_h) = \langle f, v_h \rangle, \quad \forall v_h \in V^h$$

Discretization of the problem

If $\kappa_\varepsilon \notin I_c$ the the problem is of **Fredholm type**. Consider the case where there is a **unique solution u** .

$$\begin{cases} \text{Find } u \in V := H_0^1(\Omega) \text{ such that:} \\ a(u, v) + c(u, v) = \langle f, v \rangle \quad \forall v \in V \end{cases}$$

↑ Convergence ?

$$\begin{cases} \text{Find } u_h \in V^h \text{ such that:} \\ a(u_h, v_h) + c(u_h, v_h) = \langle f, v_h \rangle, \forall v_h \in V^h \end{cases}$$

Discretization of the problem

If $\kappa_\varepsilon \notin I_c$ the the problem is of **Fredholm type**. Consider the case where there is a **unique solution u** .

$$\begin{cases} \text{Find } u \in V := H_0^1(\Omega) \text{ such that:} \\ a(u, v) + c(u, v) = \langle f, v \rangle \quad \forall v \in V \end{cases}$$

\iff

$$\begin{cases} \text{Find } u \in V \text{ such that:} \\ a(u, \mathbf{T}v) + c(u, \mathbf{T}v) = \langle f, \mathbf{T}v \rangle \quad \forall v \in V \end{cases}$$

↑ Convergence ?

$$\begin{cases} \text{Find } u_h \in V^h \text{ such that:} \\ a(u_h, v_h) + c(u_h, v_h) = \langle f, v_h \rangle, \forall v_h \in V^h \end{cases}$$



Discretization of the problem

If $\kappa_\varepsilon \notin I_c$ the the problem is of **Fredholm type**. Consider the case where there is a **unique solution u** .

$$\begin{cases} \text{Find } u \in V := H_0^1(\Omega) \text{ such that:} \\ a(u, v) + c(u, v) = \langle f, v \rangle \quad \forall v \in V \end{cases}$$

\iff

$$\begin{cases} \text{Find } u \in V \text{ such that:} \\ a(u, \mathbf{T}v) + c(u, \mathbf{T}v) = \langle f, \mathbf{T}v \rangle \quad \forall v \in V \end{cases}$$

↑ Convergence ?

$$\begin{cases} \text{Find } u_h \in V^h \text{ such that:} \\ a(u_h, v_h) + c(u_h, v_h) = \langle f, v_h \rangle, \forall v_h \in V^h \end{cases}$$

$$\begin{cases} \text{Find } u_h \in V^h \text{ such that:} \\ a(u_h, \mathbf{T}v_h) + c(u_h, \mathbf{T}v_h) = \langle f, \mathbf{T}v_h \rangle, \forall v_h \in V^h \end{cases}$$

Discretization of the problem

If $\kappa_\varepsilon \notin I_c$ the the problem is of **Fredholm type**. Consider the case where there is a **unique solution u** .

$$\begin{cases} \text{Find } u \in V := H_0^1(\Omega) \text{ such that:} \\ a(u, v) + c(u, v) = \langle f, v \rangle \quad \forall v \in V \end{cases}$$

\iff

$$\begin{cases} \text{Find } u \in V \text{ such that:} \\ a(u, \mathbf{T}v) + c(u, \mathbf{T}v) = \langle f, \mathbf{T}v \rangle \quad \forall v \in V \end{cases}$$

↑ Convergence ?


↑ (Céa's lemma)

$$\begin{cases} \text{Find } u_h \in V^h \text{ such that:} \\ a(u_h, v_h) + c(u_h, v_h) = \langle f, v_h \rangle, \forall v_h \in V^h \end{cases}$$

$$\begin{cases} \text{Find } u_h \in V^h \text{ such that:} \\ a(u_h, \mathbf{T}v_h) + c(u_h, \mathbf{T}v_h) = \langle f, \mathbf{T}v_h \rangle, \forall v_h \in V^h \end{cases}$$


Discretization of the problem

If $\kappa_\varepsilon \notin I_c$ the the problem is of **Fredholm type**. Consider the case where there is a **unique solution u** .



Discretization of the problem

If $\kappa_\varepsilon \notin I_c$ the the problem is of **Fredholm type**. Consider the case where there is a **unique solution u** .



Theorem. If $\mathbf{T}(V^h) \subset V^h$ then for h small enough


$$\|u - u_h\|_V \leq C \inf_{v_h \in V^h} \|u - v_h\|_V$$



Chesnel, Ciarlet (2013).

Discretization of the problem

If $\kappa_\varepsilon \notin I_c$ the the problem is of **Fredholm type**. Consider the case where there is a **unique solution u** .



Theorem. If $T(V^h) \subset V^h$ then for h small enough

$$\|u - u_h\|_V \leq C \inf_{v_h \in V^h} \|u - v_h\|_V$$


Consider the same degree of approximation from each side of Σ .
Then $T(V^h) \subset V^h$ corresponds to a simple condition on the mesh.



Chesnel, Ciarlet (2013).

Conformity for simple cases


For a **symmetric domain** (w.r.t. Σ) :



Conformity for simple cases

For a **symmetric domain** (w.r.t. Σ) :


If the mesh is **symmetric**, then $\mathbf{T}(V^h) = V^h$.



Conformity for simple cases

For a **symmetric domain** (w.r.t. Σ) :

If the mesh is **symmetric**, then $T(V^h) = V^h$.




If $\kappa_\varepsilon \neq -1$, and if the mesh is symmetric, then FE convergence.

Conformity for simple cases

For a **symmetric domain** (w.r.t. Σ) :

If the mesh is **symmetric**, then $\mathbf{T}(V^h) = V^h$.




If $\kappa_\varepsilon \neq -1$, and if the mesh is symmetric, then FE convergence.

We say that the mesh is **T-conform**.

Conformity for simple cases

For a **symmetric domain** (w.r.t. Σ) :

If the mesh is **symmetric**, then $T(V^h) = V^h$.



If $\kappa_\varepsilon \neq -1$, and if the mesh is symmetric, then FE convergence.

We say that the mesh is **T-conform**.


One can also apply locally this conformity in the neighborhood of the interface.

We say that the mesh is **locally T-conform**.

Conformity for simple cases

For a **symmetric domain** (w.r.t. Σ):

If the mesh is **symmetric**, then $\mathbf{T}(V^h) = V^h$.



If $\kappa_\varepsilon \neq -1$, and if the mesh is symmetric, then FE convergence.


We say that the mesh is **T-conform**.

One can also apply locally this conformity in the neighborhood of the interface.

We say that the mesh is **locally T-conform**.

For an **angular sector** of aperture α :


Unfortunately with the considered operators $\mathbf{T}(V^h) \notin V^h$ because angular dilations do not respect the polynomial nature of the basic functions...



Conformity for simple cases

For a **symmetric domain** (w.r.t. Σ) :

If the mesh is **symmetric**, then $\mathbf{T}(V^h) = V^h$.



If $\kappa_\varepsilon \neq -1$, and if the mesh is symmetric, then FE convergence.


We say that the mesh is **T-conform**.

One can also apply locally this conformity in the neighborhood of the interface.

We say that the mesh is **locally T-conform**.

For an **angular sector** of aperture α :


Unfortunately with the considered operators $\mathbf{T}(V^h) \notin V^h$ because angular dilations do not respect the polynomial nature of the basic functions...



Can we find other optimal operators R satisfying this T-conformity ?

The corners's problem

Consider the particular case of a **right angle**.



The corners's problem


Consider the particular case of a **right angle**.



Bonnet-Ben Dhia, Chesnel, Ciarlet (2012).

$$\mathbf{T}u = \begin{cases} u_d & \text{in } \Omega_d \\ -u_m + 2 \sum_p \chi_p \mathbf{R}_p u_d & \text{in } \Omega_m \end{cases}$$

The problem is **Fredholm** iff $\kappa_\varepsilon \notin I_c := [-3; -1/3]$.



The corners's problem


Consider the particular case of a **right angle**.

Another **optimal** operator \mathbf{T} based on symmetries has been proposed.



Nicaise, Venel (2011).

$$\mathbf{T}u = \begin{cases} u_d & \text{in } \Omega_d \\ -u_m + 2\mathbf{R}u_d & \text{in } \Omega_m \end{cases}$$



\mathbf{R} is based on axial symmetries.

The corners's problem


Consider the particular case of a **right angle**.

Another **optimal** operator \mathbf{T} based on symmetries has been proposed.



Nicaise, Venel (2011).

$$\mathbf{T}u = \begin{cases} u_d & \text{in } \Omega_d \\ -u_m + 2\mathbf{R}u_d & \text{in } \Omega_m \end{cases}$$



\mathbf{R} is based on axial symmetries.

The corners's problem


Consider the particular case of a **right angle**.

Another **optimal** operator \mathbf{T} based on symmetries has been proposed.




Nicaise, Venel (2011).

$$\mathbf{T}u = \begin{cases} u_d & \text{in } \Omega_d \\ -u_m + 2\mathbf{R}u_d & \text{in } \Omega_m \end{cases}$$



\mathbf{R} is based on axial symmetries.



The corners's problem


Consider the particular case of a **right angle**.

Another **optimal** operator \mathbf{T} based on symmetries has been proposed.




Nicaise, Venel (2011).

$$\mathbf{T}u = \begin{cases} u_d & \text{in } \Omega_d \\ -u_m + 2\mathbf{R}u_d & \text{in } \Omega_m \end{cases}$$



\mathbf{R} is based on axial symmetries.



The corners's problem


Consider the particular case of a **right angle**.

Another **optimal** operator \mathbf{T} based on symmetries has been proposed.




Nicaise, Venel (2011).

$$\mathbf{T}u = \begin{cases} u_d & \text{in } \Omega_d \\ -u_m + 2\mathbf{R}u_d & \text{in } \Omega_m \end{cases}$$



\mathbf{R} is based on axial symmetries.



The corners's problem


Consider the particular case of a **right angle**.

Another **optimal** operator \mathbf{T} based on symmetries has been proposed.




Nicaise, Venel (2011).

$$\mathbf{T}u = \begin{cases} u_d & \text{in } \Omega_d \\ -u_m + 2\mathbf{R}u_d & \text{in } \Omega_m \end{cases}$$



\mathbf{R} is based on axial symmetries.



The corners's problem


Consider the particular case of a **right angle**.

Another **optimal** operator \mathbf{T} based on symmetries has been proposed.




Nicaise, Venel (2011).

$$\mathbf{T}u = \begin{cases} u_d & \text{in } \Omega_d \\ -u_m + 2\mathbf{R}u_d & \text{in } \Omega_m \end{cases}$$



\mathbf{R} is based on axial symmetries.



The corners's problem


Consider the particular case of a **right angle**.

Another **optimal** operator \mathbf{T} based on symmetries has been proposed.




Nicaise, Venel (2011).

$$\mathbf{T}u = \begin{cases} u_d & \text{in } \Omega_d \\ -u_m + 2\mathbf{R}u_d & \text{in } \Omega_m \end{cases}$$



\mathbf{R} is based on axial symmetries.



The corners's problem


Consider the particular case of a **right angle**.

Another **optimal** operator \mathbf{T} based on symmetries has been proposed.




Nicaise, Venel (2011).

$$\mathbf{T}u = \begin{cases} u_d & \text{in } \Omega_d \\ -u_m + 2\mathbf{R}u_d & \text{in } \Omega_m \end{cases}$$



\mathbf{R} is based on axial symmetries.



The corners's problem


Consider the particular case of a **right angle**.

Another **optimal** operator \mathbf{T} based on symmetries has been proposed.




Nicaise, Venel (2011).

$$\mathbf{T}u = \begin{cases} u_d & \text{in } \Omega_d \\ -u_m + 2\mathbf{R}u_d & \text{in } \Omega_m \end{cases}$$



\mathbf{R} is based on axial symmetries.



The corners's problem


Consider the particular case of a **right angle**.

Another **optimal** operator \mathbf{T} based on symmetries has been proposed.




Nicaise, Venel (2011).

$$\mathbf{T}u = \begin{cases} u_d & \text{in } \Omega_d \\ -u_m + 2\mathbf{R}u_d & \text{in } \Omega_m \end{cases}$$



\mathbf{R} is based on axial symmetries.



The corners's problem


Consider the particular case of a **right angle**.

Another **optimal** operator \mathbf{T} based on symmetries has been proposed.




Nicaise, Venel (2011).

$$\mathbf{T}u = \begin{cases} u_d & \text{in } \Omega_d \\ -u_m + 2\mathbf{R}u_d & \text{in } \Omega_m \end{cases}$$



\mathbf{R} is based on axial symmetries.



$$\|\mathbf{R}\|^2 = \|\mathbf{R}'\|^2 = 3$$

The corners's problem


Consider the particular case of a **right angle**.

Another **optimal** operator \mathbf{T} based on symmetries has been proposed.




Nicaise, Venel (2011).

$$\mathbf{T}u = \begin{cases} u_d & \text{in } \Omega_d \\ -u_m + 2\mathbf{R}u_d & \text{in } \Omega_m \end{cases}$$



\mathbf{R} is based on axial symmetries.



$$\|\mathbf{R}\|^2 = \|\mathbf{R}'\|^2 = 3$$

If the mesh is **symmetric with respect to the two axes**, then $\mathbf{T}(V^h) \subset V^h$.

The corners's problem


Consider the particular case of a **right angle**.

Another **optimal** operator \mathbf{T} based on symmetries has been proposed.




Nicaise, Venel (2011).

$$\mathbf{T}u = \begin{cases} u_d & \text{in } \Omega_d \\ -u_m + 2\mathbf{R}u_d & \text{in } \Omega_m \end{cases}$$



\mathbf{R} is based on axial symmetries.



$$\|\mathbf{R}\|^2 = \|\mathbf{R}'\|^2 = 3$$

If the mesh is **symmetric with respect to the two axes**, then $\mathbf{T}(V^h) \subset V^h$.

Idea : generalize these isometry-based operators \mathbf{R} for any angle.




Bonnet-Ben Dhia, C., Ciarlet
in preparation.

Folding operators and tillings rules

One can generalize this **folding method using symmetry-based and rotation-based operators**.


Example of a folding operator from Ω_m to Ω_d for an angle $\alpha = \frac{4\pi}{3}$.



Folding operators and tiling rules

One can generalize this **folding method using symmetry-based and rotation-based operators**.


Example of a folding operator from Ω_m to Ω_d for an angle $\alpha = \frac{4\pi}{3}$.



Folding operators and tillings rules

One can generalize this **folding method using symmetry-based and rotation-based operators**.

Example of a folding operator from Ω_m to Ω_d for an angle $\alpha = \frac{4\pi}{3}$.




Folding operators and tappings rules

One can generalize this **folding method using symmetry-based and rotation-based operators**.

Example of a folding operator from Ω_m to Ω_d for an angle $\alpha = \frac{4\pi}{3}$.

Ru has to be continuous.




Folding operators and tillings rules

One can generalize this **folding method using symmetry-based and rotation-based operators**.

Example of a folding operator from Ω_m to Ω_d for an angle $\alpha = \frac{4\pi}{3}$.

Ru has to be continuous.




Folding operators and tiling rules

One can generalize this **folding method using symmetry-based and rotation-based operators**.

Example of a folding operator from Ω_m to Ω_d for an angle $\alpha = \frac{4\pi}{3}$.

Ru has to be continuous.




Folding operators and tillings rules

One can generalize this **folding method using symmetry-based and rotation-based operators**.

Example of a folding operator from Ω_m to Ω_d for an angle $\alpha = \frac{4\pi}{3}$.

Ru has to be continuous.




Folding operators and tappings rules

One can generalize this **folding method using symmetry-based and rotation-based operators**.

Example of a folding operator from Ω_m to Ω_d for an angle $\alpha = \frac{4\pi}{3}$.

Ru has to be continuous.




Folding operators and tiling rules

One can generalize this **folding method using symmetry-based and rotation-based operators**.

Example of a folding operator from Ω_m to Ω_d for an angle $\alpha = \frac{4\pi}{3}$.

Ru has to be continuous.




Folding operators and tiling rules

One can generalize this **folding method using symmetry-based and rotation-based operators**.

Example of a folding operator from Ω_m to Ω_d for an angle $\alpha = \frac{4\pi}{3}$.

Ru has to be continuous.




Folding operators and tiling rules

One can generalize this **folding method using symmetry-based and rotation-based operators**.

Example of a folding operator from Ω_m to Ω_d for an angle $\alpha = \frac{4\pi}{3}$.

Ru has to be continuous.




Folding operators and tiling rules

One can generalize this **folding method using symmetry-based and rotation-based operators**.

Example of a folding operator from Ω_m to Ω_d for an angle $\alpha = \frac{4\pi}{3}$.

Ru has to be continuous.




Folding operators and tiling rules

One can generalize this **folding method using symmetry-based and rotation-based operators**.

Example of a folding operator from Ω_m to Ω_d for an angle $\alpha = \frac{4\pi}{3}$.




Ru has to be continuous.



Folding operators and tillings rules

One can generalize this **folding method using symmetry-based and rotation-based operators**.

Example of a folding operator from Ω_m to Ω_d for an angle $\alpha = \frac{4\pi}{3}$.






Once the **pattern** is chosen, **for any form**, one builds the **T-conforming** mesh by duplicating the pattern by symmetries.



Folding operators and tillings rules

One can generalize this **folding method using symmetry-based and rotation-based operators**.

Example of a folding operator from Ω_m to Ω_d for an angle $\alpha = \frac{4\pi}{3}$.



Once the **pattern** is chosen, **for any form**, one builds the **T-conforming** mesh by duplicating the pattern by symmetries.






For a **polygonal interface**, one **locally** applies this method.



Folding operators and tiling rules

One can generalize this **folding method using symmetry-based and rotation-based operators**.

Example of a folding operator from Ω_m to Ω_d for an angle $\alpha = \frac{4\pi}{3}$.



Once the **pattern** is chosen, **for any form**, one builds the **T-conforming** mesh by duplicating the pattern by symmetries.




For a **polygonal interface**, one **locally** applies this method.

Theorem. For any angle $\alpha \in 2\pi\mathbb{Q}$, if the mesh is **locally built** as a duplicated **pattern** by symmetries, then $\mathbf{T}(V^h) \subset V^h$.


Application to the eigenvalue problem

Find $(u, \lambda) \in H_0^1(\Omega) \setminus \{0\} \times \mathbb{C}$ such that:

$$-\operatorname{div}(\varepsilon^{-1} \nabla u) = \lambda \mu u \quad \text{in } \Omega.$$



standard mesh




T-conforming mesh

Application to the eigenvalue problem


Find $(u, \lambda) \in H_0^1(\Omega) \setminus \{0\} \times \mathbb{C}$ such that:

$$-\operatorname{div}(\varepsilon^{-1} \nabla u) = \lambda \mu u \quad \text{in } \Omega.$$

Consider Lagrange FE of order 1, 2, 3.



standard mesh




T-conforming mesh

Application to the eigenvalue problem


Find $(u, \lambda) \in H_0^1(\Omega) \setminus \{0\} \times \mathbb{C}$ such that:

$$-\operatorname{div}(\varepsilon^{-1} \nabla u) = \lambda \mu u \quad \text{in } \Omega.$$

Consider Lagrange FE of order 1, 2, 3.





standard mesh



T-conforming mesh

For a **positive** eigenvalue:




Application to the eigenvalue problem


Find $(u, \lambda) \in H_0^1(\Omega) \setminus \{0\} \times \mathbb{C}$ such that:

$$-\operatorname{div}(\varepsilon^{-1} \nabla u) = \lambda \mu u \quad \text{in } \Omega.$$

Consider Lagrange FE of order 1, 2, 3.





standard mesh



T-conforming mesh

For a **negative** eigenvalue:




Outline

- Part II: scattering problem with sign-changing coefficients

Motivations

The goal is to compute the scattered field by a polygonal metallic obstacle.


$$u^{inc} = e^{i \vec{k} \cdot \vec{x}}$$



Motivations

The goal is to compute the scattered field by a polygonal metallic obstacle.

$$u^{inc} = e^{i \vec{k} \cdot \vec{x}}$$




Metal's permittivity follows the Drude's model (convention $e^{-i\omega t}$):

Motivations


The goal is to compute the scattered field by a polygonal metallic obstacle.

$$u^{inc} = e^{i \vec{k} \cdot \vec{x}}$$



Metal's permittivity follows the Drude's model (convention $e^{-i\omega t}$):

$$\epsilon = \epsilon^\gamma(\omega) := 1 - \frac{\omega_p^2}{\omega^2 + i\omega\gamma}$$



Time-harmonic equations for the TM polarization


$$H_z = u^{\text{inc}} + u^{\text{sca}} \quad k = \frac{\omega}{c} \sqrt{\varepsilon_d \mu_d}$$

$$\operatorname{div} \left(\frac{1}{\varepsilon(\omega)} \nabla H_z \right) + \frac{\omega^2}{c^2} \mu H_z = 0 \text{ in } D_R$$

$$\partial_n H_z - ik H_z = \partial_n u^{\text{inc}} - iku^{\text{inc}} \text{ on } \partial D_R$$

- Radiation condition at finite distance

Work at a chosen frequency.



Time-harmonic equations for the TM polarization

$$H_z = u^{\text{inc}} + u^{\text{sca}} \quad k = \frac{\omega}{c} \sqrt{\varepsilon_d \mu_d}$$

$$\operatorname{div} \left(\frac{1}{\varepsilon(\omega)} \nabla H_z \right) + \frac{\omega^2}{c^2} \mu H_z = 0 \text{ in } D_R$$


$$\partial_n H_z - ik H_z = \partial_n u^{\text{inc}} - iku^{\text{inc}} \text{ on } \partial D_R$$

- Radiation condition at finite distance

Work at a chosen frequency.

Mathematically:

- due to the dissipation, the problem has a unique solution
- one can approximate the solution with Finite Elements Methods



Numerical illustrations

We computed the total field for a triangular silver inclusion embedded in vacuum. We use Finite Element of order 2, with a plane wave of incidence $-\pi/12$.

Numerical illustrations

We computed the total field for a triangular silver inclusion embedded in vacuum. We use Finite Element of order 2, with a plane wave of incidence $-\pi/12$.


$$\omega_p = 13.3 \text{ PHz} \quad \omega = 6 \text{ PHz} \quad \gamma = 0.113 \text{ PHz}$$

$$\varepsilon_m(\omega) = -3.9193 + 0.0926i \quad \varepsilon_d = \mu_d = \mu_m = 1$$

Numerical illustrations

We computed the total field for a triangular silver inclusion embedded in vacuum. We use Finite Element of order 2, with a plane wave of incidence $-\pi/12$.

$$\begin{aligned}\omega_p &= 13.3 \text{ PHz} & \omega &= 6 \text{ PHz} & \gamma &= 0.113 \text{ PHz} \\ \varepsilon_m(\omega) &= -3.9193 + 0.0926i & \varepsilon_d = \mu_d = \mu_m &= 1\end{aligned}$$




coarse mesh


Numerical illustrations

We computed the total field for a triangular silver inclusion embedded in vacuum. We use Finite Element of order 2, with a plane wave of incidence $-\pi/12$.

$$\begin{aligned}\omega_p &= 13.3 \text{ PHz} & \omega &= 6 \text{ PHz} & \gamma &= 0.113 \text{ PHz} \\ \varepsilon_m(\omega) &= -3.9193 + 0.0926i & \varepsilon_d = \mu_d = \mu_m &= 1\end{aligned}$$



coarse mesh





refined mesh

Numerical illustrations

We computed the total field for a triangular silver inclusion embedded in vacuum. We use Finite Element of order 2, with a plane wave of incidence $-\pi/12$.

$$\begin{aligned}\omega_p &= 13.3 \text{ PHz} & \omega &= 6 \text{ PHz} & \gamma &= 0.113 \text{ PHz} \\ \varepsilon_m(\omega) &= -3.9193 + 0.0926i & \varepsilon_d = \mu_d = \mu_m &= 1\end{aligned}$$





The solution is not stable ! Strong oscillations near the corners.

Numerical illustrations

We computed the total field for a triangular silver inclusion embedded in vacuum. We use Finite Element of order 2, with a plane wave of incidence $-\pi/12$.

$$\begin{aligned}\omega_p &= 13.3 \text{ PHz} & \omega &= 6 \text{ PHz} & \gamma &= 0.113 \text{ PHz} \\ \varepsilon_m(\omega) &= -3.9193 + 0.0926i & \varepsilon_d = \mu_d = \mu_m &= 1\end{aligned}$$




The solution is not stable ! Strong oscillations near the corners.

To understand the reasons of such instabilities and how to avoid them, we study a **limit problem by neglecting dissipation**.

The dissipationless Drude's model

At optical frequencies one can neglect dissipation so that the metal's permittivity follows the law:


$$\varepsilon = \varepsilon^0(\omega) := 1 - \frac{\omega_p^2}{\omega^2} < 0$$



The dissipationless Drude's model

At optical frequencies one can neglect dissipation so that the metal's permittivity follows the law:

$$\varepsilon = \varepsilon^0(\omega) := 1 - \frac{\omega_p^2}{\omega^2} < 0$$




Now we consider the scattering problem with **sign-changing coefficient** (chosen frequency):

The dissipationless Drude's model


At optical frequencies one can neglect dissipation so that the metal's permittivity follows the law:

$$\varepsilon = \varepsilon^0(\omega) := 1 - \frac{\omega_p^2}{\omega^2} < 0$$



Now we consider the scattering problem with **sign-changing coefficient** (chosen frequency):


$$\begin{aligned} \operatorname{div} \left(\frac{1}{\varepsilon(\omega)} \nabla H_z \right) + \frac{\omega^2}{c^2} \mu H_z &= 0 \text{ in } D_R \\ \partial_n H_z - ikH_z &= \partial_n u^{\text{inc}} - iku^{\text{inc}} \text{ on } \partial D_R \end{aligned}$$



The dissipationless Drude's model


At optical frequencies one can neglect dissipation so that the metal's permittivity follows the law:

$$\varepsilon = \varepsilon^0(\omega) := 1 - \frac{\omega_p^2}{\omega^2} < 0$$



Now we consider the scattering problem with **sign-changing coefficient** (chosen frequency):

$$\begin{aligned} \operatorname{div} \left(\frac{1}{\varepsilon(\omega)} \nabla H_z \right) + \frac{\omega^2}{c^2} \mu H_z &= 0 \text{ in } D_R \\ \partial_n H_z - ikH_z &= \partial_n u^{\text{inc}} - iku^{\text{inc}} \text{ on } \partial D_R \end{aligned}$$




Mathematically this implies for our problem:

- difficulties to prove existence and uniqueness of the solution
- the corners of the inclusion may cause strong singular behavior
- standard approximation with Finite Elements Methods may fail

The dissipationless Drude's model


At optical frequencies one can neglect dissipation so that the metal's permittivity follows the law:

$$\varepsilon = \varepsilon^0(\omega) := 1 - \frac{\omega_p^2}{\omega^2} < 0$$



Now we consider the scattering problem with **sign-changing coefficient** (chosen frequency):

$$\begin{aligned} \operatorname{div} \left(\frac{1}{\varepsilon(\omega)} \nabla H_z \right) + \frac{\omega^2}{c^2} \mu H_z &= 0 \text{ in } D_R \\ \partial_n H_z - ikH_z &= \partial_n u^{\text{inc}} - iku^{\text{inc}} \text{ on } \partial D_R \end{aligned}$$



Mathematically this implies for our problem:

- difficulties to prove existence and uniqueness of the solution
- the corners of the inclusion may cause strong singular behavior
- standard approximation with Finite Elements Methods may fail


Use the T-coercivity ! Two cases to study: $\kappa_\varepsilon \notin I_c$ and $\kappa_\varepsilon \in I_c$



The dissipationless Drude's model


At optical frequencies one can neglect dissipation so that the metal's permittivity follows the law:

$$\varepsilon = \varepsilon^0(\omega) := 1 - \frac{\omega_p^2}{\omega^2} < 0$$



Now we consider the scattering problem with **sign-changing coefficient** (chosen frequency):

$$\begin{aligned} \operatorname{div} \left(\frac{1}{\varepsilon(\omega)} \nabla H_z \right) + \frac{\omega^2}{c^2} \mu H_z &= 0 \text{ in } D_R \\ \partial_n H_z - ikH_z &= \partial_n u^{\text{inc}} - iku^{\text{inc}} \text{ on } \partial D_R \end{aligned}$$



Mathematically this implies for our problem:

- difficulties to prove existence and uniqueness of the solution
- the corners of the inclusion may cause strong singular behavior
- standard approximation with Finite Elements Methods may fail


Use the T-coercivity ! Two cases to study: $\kappa_\varepsilon \notin I_c$ and $\kappa_\varepsilon \in I_c$

For a contrast $\kappa_{\varepsilon} \in I_c$

Find $u := u^{\text{inc}} + u^{\text{sac}} \in H^1(D_R)$ such that:

$$\operatorname{div}(\varepsilon^{-1} \nabla u) + \frac{\omega^2}{c^2} \mu u = 0 \quad \text{in } D_R,$$

$$\partial_n u - iku = \partial u^{\text{inc}} - iku^{\text{inc}} \quad \text{on } \partial D_R.$$



For a contrast $\kappa_{\varepsilon} \in I_c$


Find $u := u^{\text{inc}} + u^{\text{sac}} \in H^1(D_R)$ such that:

$$\begin{aligned} \operatorname{div}(\varepsilon^{-1} \nabla u) + \frac{\omega^2}{c^2} \mu u &= 0 \quad \text{in } D_R, \\ \partial_n u - iku &= \partial u^{\text{inc}} - iku^{\text{inc}} \quad \text{on } \partial D_R. \end{aligned}$$

Find $u := u^{\text{inc}} + u^{\text{sac}} \in H^1(D_R)$ such that:

$$a(u, v) + c(u, v) = l(v), \quad \forall v \in H^1(D_R)$$

$$a(u, v) := \int_{D_R} \varepsilon^{-1} \nabla u \cdot \overline{\nabla v} d\mathbf{x},$$



For a contrast $\kappa_{\varepsilon} \in I_c$

Find $u := u^{\text{inc}} + u^{\text{sac}} \in H^1(D_R)$ such that:


$$\operatorname{div}(\varepsilon^{-1} \nabla u) + \frac{\omega^2}{c^2} \mu u = 0 \quad \text{in } D_R,$$

$$\partial_n u - iku = \partial u^{\text{inc}} - iku^{\text{inc}} \quad \text{on } \partial D_R.$$

Find $u := u^{\text{inc}} + u^{\text{sac}} \in H^1(D_R)$ such that:


$$a(u, v) + c(u, v) = l(v), \quad \forall v \in H^1(D_R)$$

$$a(u, v) := \int_{D_R} \varepsilon^{-1} \nabla u \cdot \overline{\nabla v} d\mathbf{x},$$



In that case the form a is **not T-coercive !**

Can we still compute the total field with finite elements ?



For a contrast $\kappa_{\varepsilon} \in I_c$

Find $u := u^{\text{inc}} + u^{\text{sac}} \in H^1(D_R)$ such that:


$$\operatorname{div}(\varepsilon^{-1} \nabla u) + \frac{\omega^2}{c^2} \mu u = 0 \quad \text{in } D_R,$$

$$\partial_n u - iku = \partial u^{\text{inc}} - iku^{\text{inc}} \quad \text{on } \partial D_R.$$

Find $u := u^{\text{inc}} + u^{\text{sac}} \in H^1(D_R)$ such that:

$$a(u, v) + c(u, v) = l(v), \quad \forall v \in H^1(D_R)$$


$$a(u, v) := \int_{D_R} \varepsilon^{-1} \nabla u \cdot \overline{\nabla v} d\mathbf{x},$$



In that case the form a is **not T-coercive !**

Can we still compute the total field with finite elements ?

Numerical illustrations for triangular silver inclusion in vacuum:



For a contrast $\kappa_\varepsilon \in I_c$

Find $u := u^{\text{inc}} + u^{\text{sac}} \in H^1(D_R)$ such that:


$$\operatorname{div}(\varepsilon^{-1} \nabla u) + \frac{\omega^2}{c^2} \mu u = 0 \quad \text{in } D_R,$$

$$\partial_n u - iku = \partial u^{\text{inc}} - iku^{\text{inc}} \quad \text{on } \partial D_R.$$

Find $u := u^{\text{inc}} + u^{\text{sac}} \in H^1(D_R)$ such that:

$$a(u, v) + c(u, v) = l(v), \quad \forall v \in H^1(D_R)$$

$$a(u, v) := \int_{D_R} \varepsilon^{-1} \nabla u \cdot \overline{\nabla v} d\mathbf{x},$$



In that case the form a is **not T-coercive !**

Can we still compute the total field with finite elements ?

Numerical illustrations for triangular silver inclusion in vacuum:

$\kappa_\varepsilon \in I_c \iff \omega \in [3.839 \text{ PHz}; 12.733 \text{ PHz}]$. Results for $\omega = 9 \text{ PHz}$.

For a contrast $\kappa_\varepsilon \in I_c$

Find $u := u^{\text{inc}} + u^{\text{sac}} \in H^1(D_R)$ such that:


$$\operatorname{div}(\varepsilon^{-1} \nabla u) + \frac{\omega^2}{c^2} \mu u = 0 \quad \text{in } D_R,$$

$$\partial_n u - iku = \partial u^{\text{inc}} - iku^{\text{inc}} \quad \text{on } \partial D_R.$$

Find $u := u^{\text{inc}} + u^{\text{sac}} \in H^1(D_R)$ such that:

$$a(u, v) + c(u, v) = l(v), \quad \forall v \in H^1(D_R)$$

$$a(u, v) := \int_{D_R} \varepsilon^{-1} \nabla u \cdot \overline{\nabla v} d\mathbf{x},$$




In that case the form a is **not T-coercive !**


Can we still compute the total field with finite elements ?

Numerical illustrations for triangular silver inclusion in vacuum:


$\kappa_\varepsilon \in I_c \iff \omega \in [3.839 \text{ PHz}; 12.733 \text{ PHz}]$. Results for $\omega = 9 \text{ PHz}$.



coarse mesh



intermediate mesh



refined mesh



For a contrast $\kappa_{\varepsilon} \in I_c$

Find $u := u^{\text{inc}} + u^{\text{sac}} \in H^1(D_R)$ such that:


$$\operatorname{div}(\varepsilon^{-1} \nabla u) + \frac{\omega^2}{c^2} \mu u = 0 \quad \text{in } D_R,$$

$$\partial_n u - iku = \partial u^{\text{inc}} - iku^{\text{inc}} \quad \text{on } \partial D_R.$$

Find $u := u^{\text{inc}} + u^{\text{sac}} \in H^1(D_R)$ such that:

$$a(u, v) + c(u, v) = l(v), \quad \forall v \in H^1(D_R)$$

$$a(u, v) := \int_{D_R} \varepsilon^{-1} \nabla u \cdot \overline{\nabla v} d\mathbf{x},$$




In that case the form a is **not T-coercive !**

Can we still compute the total field with finite elements ?


Numerical illustrations for triangular silver inclusion in vacuum:

$\kappa_{\varepsilon} \in I_c \iff \omega \in [3.839 \text{ PHz}; 12.733 \text{ PHz}]$. Results for $\omega = 9 \text{ PHz}$.




Analysis of the singularities at the corners

For simplicity, we consider only one corner.




Analysis of the singularities at the corners

For simplicity, we consider only one corner.



Analysis of the singularities at the corners

For simplicity, we consider only one corner.




Limit behaviour at the corner


Behaviour at infinity in the strip

Analysis of the singularities at the corners

For simplicity, we consider only one corner.



Limit behaviour at the corner




Behaviour at infinity in the strip

$$-\varepsilon^{-1}r\partial_r(r\partial_r u) - \partial_\theta(\varepsilon^{-1}\partial_\theta u) + \omega^2\mu r^2 u = 0$$


$$-\operatorname{div}(\varepsilon^{-1}\nabla u) + \omega^2\mu e^{2z}u = 0$$

Analysis of the singularities at the corners

For simplicity, we consider only one corner.



Limit behaviour at the corner




Behaviour at infinity in the strip

$$-\varepsilon^{-1}r\partial_r(r\partial_r u) - \partial_\theta(\varepsilon^{-1}\partial_\theta u) + \omega^2\mu r^2 u = 0$$

$$-\operatorname{div}(\varepsilon^{-1}\nabla u) + \omega^2\mu e^{2z}u = 0$$

Analysis of the singularities at the corners

For simplicity, we consider only one corner.



Limit behaviour at the corner

$$-\varepsilon^{-1}r\partial_r(r\partial_r u) - \partial_\theta(\varepsilon^{-1}\partial_\theta u) + \omega^2\mu r^2 u = 0$$


Behaviour at infinity in the strip

$$-\operatorname{div}(\varepsilon^{-1}\nabla u) + \omega^2\mu e^{2z}u = 0$$

We look for solutions $u(z, \theta) = e^{\lambda z}\phi(\theta)$
 λ are called singular exponents.

Analysis of the singularities at the corners

For simplicity, we consider only one corner.




$$-\varepsilon^{-1}r\partial_r(r\partial_r u) - \partial_\theta(\varepsilon^{-1}\partial_\theta u) + \omega^2\mu r^2 u = 0$$

$$-\operatorname{div}(\varepsilon^{-1}\nabla u) + \omega^2\mu e^{2z}u = 0$$


We look for solutions $u(z, \theta) = e^{\lambda z}\phi(\theta)$
 λ are called singular exponents.

Analysis of the singularities at the corners

For simplicity, we consider only one corner.



Limit behaviour at the corner




Behaviour at infinity in the strip

$$-\varepsilon^{-1}r\partial_r(r\partial_r u) - \partial_\theta(\varepsilon^{-1}\partial_\theta u) + \omega^2\mu r^2 u = 0$$


$$-\operatorname{div}(\varepsilon^{-1}\nabla u) + \omega^2\mu e^{2z}u = 0$$

We look for solutions $u(z, \theta) = e^{\lambda z}\phi(\theta)$
 λ are called singular exponents.

$$\kappa_\varepsilon \notin I_c$$




The modes are evanescent
 $\operatorname{Re}(\lambda) > 0, \forall \lambda \neq 0$




Analysis of the singularities at the corners

For simplicity, we consider only one corner.



Limit behaviour at the corner




Behaviour at infinity in the strip

$$-\varepsilon^{-1}r\partial_r(r\partial_r u) - \partial_\theta(\varepsilon^{-1}\partial_\theta u) + \omega^2\mu r^2 u = 0$$


$$-\operatorname{div}(\varepsilon^{-1}\nabla u) + \omega^2\mu e^{2z}u = 0$$

We look for solutions $u(z, \theta) = e^{\lambda z}\phi(\theta)$
 λ are called singular exponents.




Analysis of the singularities at the corners

For simplicity, we consider only one corner.



Limit behaviour at the corner




Behaviour at infinity in the strip

$$-\varepsilon^{-1}r\partial_r(r\partial_r u) - \partial_\theta(\varepsilon^{-1}\partial_\theta u) + \omega^2\mu r^2 u = 0$$


$$-\operatorname{div}(\varepsilon^{-1}\nabla u) + \omega^2\mu e^{2z}u = 0$$

We look for solutions $u(z, \theta) = e^{\lambda z}\phi(\theta)$
 λ are called singular exponents.




Analysis of the singularities at the corners

For simplicity, we consider only one corner.



Limit behaviour at the corner



Behaviour at infinity in the strip


$$-\varepsilon^{-1}r\partial_r(r\partial_r u) - \partial_\theta(\varepsilon^{-1}\partial_\theta u) + \omega^2\mu r^2 u = 0$$

In the physical domain, the solutions are

$$u(r, \theta) = r^\lambda \phi(\theta)$$


$$-\operatorname{div}(\varepsilon^{-1}\nabla u) + \omega^2\mu e^{2z}u = 0$$

We look for solutions $u(z, \theta) = e^{\lambda z}\phi(\theta)$
 λ are called singular exponents.



Analysis of the singularities at the corners

For simplicity, we consider only one corner.





Limit behaviour at the corner

$$-\varepsilon^{-1}r\partial_r(r\partial_r u) - \partial_\theta(\varepsilon^{-1}\partial_\theta u) + \omega^2\mu r^2 u = 0$$

In the physical domain, the solutions are

$$u(r, \theta) = r^\lambda \phi(\theta)$$


Appearance of an oscillating hypersingularity $s(r, \theta) = e^{i\eta \ln r} \phi(\theta) \notin H^1$



Behaviour at infinity in the strip


$$-\operatorname{div}(\varepsilon^{-1} \nabla u) + \omega^2 \mu e^{2z} u = 0$$

We look for solutions $u(z, \theta) = e^{\lambda z} \phi(\theta)$
 λ are called singular exponents.



Analysis of the singularities at the corners

For simplicity, we consider only one corner.




Limit behaviour at the corner

$$-\varepsilon^{-1}r\partial_r(r\partial_r u) - \partial_\theta(\varepsilon^{-1}\partial_\theta u) + \omega^2\mu r^2 u = 0$$



In the physical domain, the solutions are

$$u(r, \theta) = r^\lambda \phi(\theta)$$

Appearance of an oscillating hypersingularity $s(r, \theta) = e^{i\eta \ln r} \phi(\theta) \notin H^1$




Black-hole wave.



Behaviour at infinity in the strip


$$-\operatorname{div}(\varepsilon^{-1} \nabla u) + \omega^2 \mu e^{2z} u = 0$$

We look for solutions $u(z, \theta) = e^{\lambda z} \phi(\theta)$
 λ are called singular exponents.



Black-hole waves and Fredholmness


$$s^\pm(r, \theta) = e^{i(\pm\eta \ln r - \omega t)} \Phi(\theta)$$



Black-hole waves and Fredholmness


$$s^\pm(r, \theta) = e^{i(\pm\eta \ln r - \omega t)} \Phi(\theta)$$

These black-hole waves are of **infinite energy** and strongly oscillating at the corners.



Black-hole waves and Fredholmness


$$s^\pm(r, \theta) = e^{i(\pm\eta \ln r - \omega t)} \Phi(\theta)$$



These black-hole waves are of **infinite energy** and strongly oscillating at the corners.
This is why standard FEM fail to approximate the solution:
spurious reflexions appear.

Black-hole waves and Fredholmness

$$s^\pm(r, \theta) = e^{i(\pm\eta \ln r - \omega t)} \Phi(\theta)$$




These black-hole waves are of **infinite energy** and strongly oscillating at the corners.
This is why standard FEM fail to approximate the solution:
spurious reflexions appear.
However we **need to take them into account** to recover a
well-posed formulation.



Bonnet-Ben Dhia, C., Chesnel, Ciarlet submitted (2015).

Black-hole waves and Fredholmness

$$s^\pm(r, \theta) = e^{i(\pm\eta \ln r - \omega t)} \Phi(\theta)$$



These black-hole waves are of **infinite energy** and strongly oscillating at the corners.
This is why standard FEM fail to approximate the solution:
spurious reflexions appear.
However we **need to take them into account** to recover a
well-posed formulation.

Near the corners the solution decomposes as


$$u = b^+ s^+ + b^- s^- + \tilde{u}, \quad \tilde{u} \in H^1(D_R), \quad s^\pm \notin H^1(D_R), \quad b^\pm \in \mathbb{C}$$



Bonnet-Ben Dhia, C., Chesnel, Ciarlet submitted (2015).

Black-hole waves and Fredholmness

$$s^\pm(r, \theta) = e^{i(\pm\eta \ln r - \omega t)} \Phi(\theta)$$



These black-hole waves are of **infinite energy** and strongly oscillating at the corners.
This is why standard FEM fail to approximate the solution:
spurious reflexions appear.
However we **need to take them into account** to recover a **well-posed formulation**.

Near the corners the solution decomposes as

$$u = b^+ s^+ + b^- s^- + \tilde{u}, \quad \tilde{u} \in H^1(D_R), \quad s^\pm \notin H^1(D_R), \quad b^\pm \in \mathbb{C}$$

To get a **unique solution**, one needs to enforce a condition on b^\pm . **How to proceed ?**

By **energy flux**:


- one singularity carries **energy absorbed** by the corner
- the other **brings energy** emitted from the corner



Bonnet-Ben Dhia, C., Chesnel, Ciarlet submitted (2015).

Black-hole waves and Fredholmness

$$s^\pm(r, \theta) = e^{i(\pm\eta \ln r - \omega t)} \Phi(\theta)$$



These black-hole waves are of **infinite energy** and strongly oscillating at the corners.
This is why standard FEM fail to approximate the solution:
spurious reflexions appear.
However we **need to take them into account** to recover a **well-posed formulation**.

Near the corners the solution decomposes as

$$u = b^+ \cancel{s^+} + b^- s^- + \tilde{u}, \quad \tilde{u} \in H^1(D_R), \quad s^\pm \notin H^1(D_R), \quad b^\pm \in \mathbb{C}$$

To get a **unique solution**, one needs to enforce a condition on b^\pm . **How to proceed ?**
By **energy flux**:

- one singularity carries **energy absorbed** by the corner
- the other **brings energy** emitted from the corner


The **physical** solution takes into account the one that **does not add energy** into the system, the **outgoing** solution in the waveguide setting.



Bonnet-Ben Dhia, C., Chesnel, Ciarlet submitted (2015).

Black-hole waves and Fredholmness

$$s^\pm(r, \theta) = e^{i(\pm\eta \ln r - \omega t)} \Phi(\theta)$$



These black-hole waves are of **infinite energy** and strongly oscillating at the corners.
This is why standard FEM fail to approximate the solution:
spurious reflexions appear.
However we **need to take them into account** to recover a **well-posed formulation**.

Near the corners the solution decomposes as

$$u = b^+ \cancel{s^+} + b^- s^- + \tilde{u}, \quad \tilde{u} \in H^1(D_R), \quad s^\pm \notin H^1(D_R), \quad b^\pm \in \mathbb{C}$$

To get a **unique solution**, one needs to enforce a condition on b^\pm . **How to proceed ?**
By **energy flux**:

- one singularity carries **energy absorbed** by the corner
- the other **brings energy** emitted from the corner

The **physical** solution takes into account the one that **does not add energy** into the system, the **outgoing** solution in the waveguide setting.


Then we can prove that we have a **well-posed formulation**.



Bonnet-Ben Dhia, C., Chesnel, Ciarlet submitted (2015).

Black-hole waves and Fredholmness

$$s^\pm(r, \theta) = e^{i(\pm\eta \ln r - \omega t)} \Phi(\theta)$$



These black-hole waves are of **infinite energy** and strongly oscillating at the corners.
This is why standard FEM fail to approximate the solution:
spurious reflexions appear.
However we **need to take them into account** to recover a **well-posed formulation**.

Near the corners the solution decomposes as

$$u = b^+ \cancel{s^+} + b^- s^- + \tilde{u}, \quad \tilde{u} \in H^1(D_R), \quad s^\pm \notin H^1(D_R), \quad b^\pm \in \mathbb{C}$$

To get a **unique solution**, one needs to enforce a condition on b^\pm . **How to proceed ?**
By **energy flux**:

- one singularity carries **energy absorbed** by the corner
- the other **brings energy** emitted from the corner

The **physical** solution takes into account the one that **does not add energy** into the system, the **outgoing** solution in the waveguide setting.

Then we can prove that we have a **well-posed formulation**.




How to proceed numerically to select the good singularity ?

Perfectly Matched Layers (PMLs)

PMLs enable to **artificially bound** the strip while making the propagative modes become **evanescent**.

$$\frac{\partial}{\partial z} \longmapsto \alpha \frac{\partial}{\partial z} \quad \alpha \in \mathbb{C}$$






Perfectly Matched Layers (PMLs)

PMLs enable to **artificially bound** the strip while making the propagative modes become evanescent.



$$\frac{\partial}{\partial z} \longmapsto \alpha \frac{\partial}{\partial z} \quad \alpha \in \mathbb{C}$$

$$u(z, \theta) = e^{\lambda z} \phi(\theta) \longrightarrow u(z, \theta) = e^{\frac{\lambda}{\alpha} z} \phi(\theta)$$





Perfectly Matched Layers (PMLs)

PMLs enable to **artificially bound** the strip while making the propagative modes become evanescent.



Perfectly Matched Layers (PMLs)



PMLs enable to **artificially bound** the strip while making the propagative modes become evanescent.



$$\begin{aligned}
 & -\operatorname{div}(\varepsilon^{-1} \nabla u) + \omega^2 \mu u = 0 & D_R \setminus \overline{B_\rho} \\
 & \partial_r u - iku = \partial_r u^{\text{inc}} - iku^{\text{inc}} & \partial D_R \\
 & + \text{matching between the PML and the strip} + \\
 & -\alpha \varepsilon^{-1} \partial_{zz} u - \frac{1}{\alpha} \partial_\theta \varepsilon^{-1} \partial_\theta u + \frac{\omega^2}{\alpha} \mu e^{\frac{2z}{\alpha}} u = 0 & S_\rho \\
 & + \text{periodic conditions}
 \end{aligned}$$

Perfectly Matched Layers (PMLs)



PMLs enable to **artificially bound** the strip while making the propagative modes become evanescent.



$$\begin{aligned}
 & -\operatorname{div}(\varepsilon^{-1} \nabla u) + \omega^2 \mu u = 0 & D_R \setminus \overline{B_\rho} \\
 & \partial_r u - iku = \partial_r u^{\text{inc}} - iku^{\text{inc}} & \partial D_R \\
 & + \text{matching between the PML and the strip} + \\
 & -\alpha \varepsilon^{-1} \partial_{zz} u - \frac{1}{\alpha} \partial_\theta \varepsilon^{-1} \partial_\theta u + \frac{\omega^2}{\alpha} \mu e^{\frac{2z}{\alpha}} u = 0 & S_\rho \\
 & \partial_z u(-L, \cdot) = 0 \\
 & + \text{periodic conditions}
 \end{aligned}$$

Perfectly Matched Layers (PMLs)

PMLs enable to **artificially bound** the strip while making the propagative modes become evanescent.



$$-\operatorname{div}(\varepsilon^{-1} \nabla u) + \omega^2 \mu u = 0 \quad D_R \setminus \overline{B_\rho}$$

$$\partial_r u - iku = \partial_r u^{\text{inc}} - iku^{\text{inc}} \quad \partial D_R$$

+ matching between the PML and the strip +

$$-\alpha \varepsilon^{-1} \partial_{zz} u - \frac{1}{\alpha} \partial_\theta \varepsilon^{-1} \partial_\theta u + \frac{\omega^2}{\alpha} \mu e^{\frac{2z}{\alpha}} u = 0 \quad S_\rho$$


$\partial_z u(-L, \cdot) = 0$

+ periodic conditions

Neumann condition due to the constant mode



With several corners



$$-\operatorname{div}(\varepsilon^{-1} \nabla u) + \omega^2 \mu u = 0 \quad D_R \setminus \cup \overline{B_\rho}$$

$$\partial_r u - iku = \partial_r u^{\text{inc}} - iku^{\text{inc}} \quad \partial D_R$$



+ matching between the PML and the strip +

$$-\alpha \varepsilon^{-1} \partial_{zz} u - \frac{1}{\alpha} \partial_\theta \varepsilon^{-1} \partial_\theta u + \frac{\omega^2}{\alpha} \mu e^{\frac{2z}{\alpha}} u = 0 \quad S_\rho$$

$$\partial_z u(-L, \cdot) = 0$$

+ periodic conditions



With several corners



$$\begin{aligned} -\operatorname{div}(\varepsilon^{-1} \nabla u) + \omega^2 \mu u &= 0 & D_R \setminus \cup \overline{B_\rho} \\ \partial_r u - iku &= \partial_r u^{\text{inc}} - iku^{\text{inc}} & \partial D_R \\ + \text{ matching between the PML and the strip } + \\ -\alpha \varepsilon^{-1} \partial_{zz} u - \frac{1}{\alpha} \partial_\theta \varepsilon^{-1} \partial_\theta u + \frac{\omega^2}{\alpha} \mu e^{\frac{2z}{\alpha}} u &= 0 & S_\rho \\ \partial_z u(-L, \cdot) &= 0 \end{aligned}$$

+ periodic conditions

With several corners



$$\begin{aligned} -\operatorname{div}(\varepsilon^{-1} \nabla u) + \omega^2 \mu u &= 0 & D_R \setminus \cup \overline{B_\rho} \\ \partial_r u - iku &= \partial_r u^{\text{inc}} - iku^{\text{inc}} & \partial D_R \\ + \text{ matching between the PML and the strip } + \\ -\alpha \varepsilon^{-1} \partial_{zz} u - \frac{1}{\alpha} \partial_\theta \varepsilon^{-1} \partial_\theta u + \frac{\omega^2}{\alpha} \mu e^{\frac{2z}{\alpha}} u &= 0 & S_\rho \\ \partial_z u(-L, \cdot) &= 0 \\ + \text{ periodic conditions } \end{aligned}$$

Computation with Lagrange FE of order 2 with a Matlab code



Numerical results


Numerical illustrations for triangular silver inclusion in vacuum:

$\kappa_\varepsilon \in I_c \iff \omega \in [3.839 \text{ PHz}; 12.733 \text{ PHz}]$. Results for $\omega = 9 \text{ PHz}$.


Numerical results

Numerical illustrations for triangular silver inclusion in vacuum:


$\kappa_\varepsilon \in I_c \iff \omega \in [3.839 \text{ PHz}; 12.733 \text{ PHz}]$. Results for $\omega = 9 \text{ PHz}$.



coarse mesh



intermediate mesh




refined mesh

Numerical results

Numerical illustrations for triangular silver inclusion in vacuum:


$\kappa_\varepsilon \in I_c \iff \omega \in [3.839 \text{ PHz}; 12.733 \text{ PHz}]$. Results for $\omega = 9 \text{ PHz}$.



Numerical results

Numerical illustrations for triangular silver inclusion in vacuum:

$\kappa_\varepsilon \in I_c \iff \omega \in [3.839 \text{ PHz}; 12.733 \text{ PHz}]$. Results for $\omega = 9 \text{ PHz}$.



The numerical method with PMLs is stable. Let us go back to the dissipative problem.

Back to dissipative medium

Back to dissipative medium

Considering losses in the metal, then the problem is well-posed.

The selected outgoing singularity becomes of finite energy. If dissipation is small, it requires meshes sufficiently refined at the corners to capture the oscillations.

Back to dissipative medium

Considering losses in the metal, then the problem is well-posed.

The selected outgoing singularity becomes of finite energy. If dissipation is small, it requires meshes sufficiently refined at the corners to capture the oscillations.

$$\kappa_\varepsilon \in I_c \iff \omega \in [3.839 \text{ PHz}; 12.733 \text{ PHz}].$$

$$\omega_p = 13.3 \text{ PHz} \quad \omega = 6 \text{ PHz} \quad \gamma = 0.113 \text{ PHz}$$

$$\varepsilon_m(\omega) = -3.9193 + 0.0926i \quad \varepsilon_d = \mu_d = \mu_m = 1$$

Back to dissipative medium

Considering losses in the metal, then the problem is well-posed.

The selected outgoing singularity becomes of finite energy. If dissipation is small, it requires meshes sufficiently refined at the corners to capture the oscillations.

$$\kappa_\varepsilon \in I_c \iff \omega \in [3.839 \text{ PHz}; 12.733 \text{ PHz}].$$

$$\omega_p = 13.3 \text{ PHz} \quad \omega = 6 \text{ PHz} \quad \gamma = 0.113 \text{ PHz}$$

$$\varepsilon_m(\omega) = -3.9193 + 0.0926i \quad \varepsilon_d = \mu_d = \mu_m = 1$$

Without PMLs

With PMLs

Back to dissipative medium

Considering losses in the metal, then the problem is well-posed.


The selected outgoing singularity becomes of finite energy. If dissipation is small, it requires meshes sufficiently refined at the corners to capture the oscillations.

$$\kappa_\varepsilon \in I_c \iff \omega \in [3.839 \text{ PHz}; 12.733 \text{ PHz}].$$

$$\omega_p = 13.3 \text{ PHz} \quad \omega = 6 \text{ PHz} \quad \gamma = 0.113 \text{ PHz}$$


$$\varepsilon_m(\omega) = -3.9193 + 0.0926i \quad \varepsilon_d = \mu_d = \mu_m = 1$$

Without PMLs



coarse mesh

With PMLs



refined mesh

Back to dissipative medium

Considering losses in the metal, then the problem is well-posed.



The selected outgoing singularity becomes of finite energy. If dissipation is small, it requires meshes sufficiently refined at the corners to capture the oscillations.

$$\kappa_\varepsilon \in I_c \iff \omega \in [3.839 \text{ PHz}; 12.733 \text{ PHz}].$$



$$\omega_p = 13.3 \text{ PHz} \quad \omega = 6 \text{ PHz} \quad \gamma = 0.113 \text{ PHz}$$

$$\varepsilon_m(\omega) = -3.9193 + 0.0926i \quad \varepsilon_d = \mu_d = \mu_m = 1$$

Without PMLs



With PMLs



Back to dissipative medium

Considering losses in the metal, then the problem is well-posed.

The selected outgoing singularity becomes of finite energy. If dissipation is small, it requires meshes sufficiently refined at the corners to capture the oscillations.

$$\kappa_\varepsilon \in I_c \iff \omega \in [3.839 \text{ PHz}; 12.733 \text{ PHz}].$$

$$\omega_p = 13.3 \text{ PHz} \quad \omega = 6 \text{ PHz} \quad \gamma = 0.113 \text{ PHz}$$

$$\varepsilon_m(\omega) = -3.9193 + 0.0926i \quad \varepsilon_d = \mu_d = \mu_m = 1$$

Without PMLs

With PMLs

Animation in time $\Re(u e^{-i\omega t})$

Back to dissipative medium

Considering losses in the metal, then the problem is well-posed.

The selected outgoing singularity becomes of finite energy. If dissipation is small, it requires meshes sufficiently refined at the corners to capture the oscillations.


$$\kappa_\varepsilon \in I_c \iff \omega \in [3.839 \text{ PHz}; 12.733 \text{ PHz}].$$

$$\omega_p = 13.3 \text{ PHz} \quad \omega = 6 \text{ PHz} \quad \gamma = 0.113 \text{ PHz}$$

$$\varepsilon_m(\omega) = -3.9193 + 0.0926i \quad \varepsilon_d = \mu_d = \mu_m = 1$$

Without PMLs

With PMLs



Animation in time $\Re(u e^{-i\omega t})$

Outline

- ✿ Part I: meshing rules for the corners case for $\kappa_\varepsilon \notin I_c$
- ✿ Part II: scattering problem with sign-changing coefficients
- ✿ **Perspectives**



Perspectives and ongoing work

Perspectives and ongoing work

T-coercivity method enables to study and approximate problems with sign-changing coefficients.

Perspectives and ongoing work

T-coercivity method enables to study and approximate problems with sign-changing coefficients.

Perspectives:

Perspectives and ongoing work

T-coercivity method enables to study and approximate problems with sign-changing coefficients.

Perspectives:

Consider more relevant models for the metal's permittivity, such as Drude-Lorentz model or hydrodynamic Drude's model. The case of dispersive media is the object of discussions with the [Fresnel Institute](#) (Marseille, France).

Perspectives and ongoing work

T-coercivity method enables to study and approximate problems with sign-changing coefficients.

Perspectives:

Consider more relevant models for the metal's permittivity, such as Drude-Lorentz model or hydrodynamic Drude's model. The case of dispersive media is the object of discussions with the Fresnel Institute (Marseille, France).

Consider Maxwell's equations.  Bonnet-Ben Dhia, Chesnel, Ciarlet (2012, 2014), C. (2015).

Perspectives and ongoing work

T-coercivity method enables to study and approximate problems with sign-changing coefficients.

Perspectives:

Consider more relevant models for the metal's permittivity, such as Drude-Lorentz model or hydrodynamic Drude's model. The case of dispersive media is the object of discussions with the **Fresnel Institute** (Marseille, France).

Consider Maxwell's equations.  Bonnet-Ben Dhia, Chesnel, Ciarlet (2012, 2014), C. (2015).

Numerical analysis to prove convergence of the method with PMLs is still in progress (difficulties related to the change of sign and the functional framework).

Perspectives and ongoing work

T-coercivity method enables to study and approximate problems with sign-changing coefficients.

Perspectives:

Consider more relevant models for the metal's permittivity, such as Drude-Lorentz model or hydrodynamic Drude's model. The case of dispersive media is the object of discussions with the **Fresnel Institute** (Marseille, France).

Consider Maxwell's equations.  Bonnet-Ben Dhia, Chesnel, Ciarlet (2012, 2014), C. (2015).

Numerical analysis to prove convergence of the method with PMLs is still in progress (difficulties related to the change of sign and the functional framework).

Limit amplitude principle: validate the existence of a time-harmonic regime according to the critical interval. These questions are investigated in the ANR project **METAMATH**, and with **C. Scheid** in team NACHOS (Nice, France).



Thank you for your attention.