

---

---

# Mathematical and Numerical study of plasmonic structures with corners

Camille Carvalho

[camille.carvalho@ensta-paristech.fr](mailto:camille.carvalho@ensta-paristech.fr)

PhD Student in Applied Mathematics,  
under the direction of A.S. Bonnet-Ben Dhia and P. Ciarlet Jr.

<sup>1</sup>POEMS, UMR 7231 CNRS-INRIA-ENSTA, Palaiseau, France



UC Merced, October 2015

# About

---

I'm a french Ph.D student in team POEMS (Paris), specialized in the theoretical and numerical modeling of wave propagation.

# About

---

I'm a french Ph.D student in team POEMS (Paris), specialized in the theoretical and numerical modeling of wave propagation.

My work deals mainly with two problems in plasmonics:

- ❖ 2D Scattering problem by a metallic inclusion presenting corners embedded in a dielectric medium
- ❖ The modeling of plasmonic waveguides: characterization of the guided modes

# About

---

I'm a french Ph.D student in team POEMS (Paris), specialized in the theoretical and numerical modeling of wave propagation.


My work deals mainly with two problems in plasmonics:

- ❖ 2D Scattering problem by a metallic inclusion presenting corners embedded in a dielectric medium
- ❖ The modeling of plasmonic waveguides: characterization of the guided modes

Both theoretical and numerical aspects are investigated.

Both problems are tackled in the **time-harmonic regime**.


I'm more a mathematician than a physicist, I make some approximations in the model.



# Motivations of this work

The goal is to compute the scattered field by a polygonal metallic obstacle.


$$u^{inc} = e^{i \vec{k} \cdot \vec{x}}$$



# Motivations of this work

The goal is to compute the scattered field by a polygonal metallic obstacle.

$$u^{inc} = e^{i \vec{k} \cdot \vec{x}}$$




It is well-known that, for silver or gold, the permittivity follows the Drude's model (convention  $e^{-i\omega t}$ ):

# Motivations of this work

The goal is to compute the scattered field by a polygonal metallic obstacle.

$$u^{inc} = e^{i \vec{k} \cdot \vec{x}}$$




It is well-known that, for silver or gold, the permittivity follows the Drude's model (convention  $e^{-i\omega t}$ ):

$$\varepsilon = \varepsilon^\gamma(\omega) := 1 - \frac{\omega_p^2}{\omega^2 + i\omega\gamma}$$

# Motivations of this work

The goal is to compute the scattered field by a polygonal metallic obstacle.

$$u^{inc} = e^{i \vec{k} \cdot \vec{x}}$$




It is well-known that, for silver or gold, the permittivity follows the Drude's model (convention  $e^{-i\omega t}$ ):

$$\varepsilon = \varepsilon^\gamma(\omega) := 1 - \frac{\omega_p^2}{\omega^2 + i\omega\gamma}$$

# Motivations of this work

The goal is to compute the scattered field by a polygonal metallic obstacle.

$$u^{inc} = e^{i \vec{k} \cdot \vec{x}}$$



It is well-known that, for silver or gold, the permittivity follows the Drude's model (convention  $e^{-i\omega t}$ ):

$$\epsilon = \epsilon^\gamma(\omega) := 1 - \frac{\omega_p^2}{\omega^2 + i\omega\gamma}$$

# Time harmonic equations for the TM polarization

$$H_z = u^{\text{inc}} + u^{\text{sca}} \quad k = \frac{\omega}{c} \sqrt{\varepsilon_d \mu_d}$$

$$\begin{aligned} \operatorname{div} \left( \frac{1}{\varepsilon(\omega)} \nabla H_z \right) + \frac{\omega^2}{c^2} \mu H_z &= 0 \text{ in } B_R \\ \partial_n H_z - ik H_z &= \partial_n u^{\text{inc}} - iku^{\text{inc}} \text{ on } \Gamma_R \end{aligned}$$

- Radiation condition at finite distance


# Time harmonic equations for the TM polarization

$$H_z = u^{\text{inc}} + u^{\text{sca}} \quad k = \frac{\omega}{c} \sqrt{\varepsilon_d \mu_d}$$

$$\operatorname{div} \left( \frac{1}{\varepsilon(\omega)} \nabla H_z \right) + \frac{\omega^2}{c^2} \mu H_z = 0 \text{ in } B_R$$

$$\partial_n H_z - ik H_z = \partial_n u^{\text{inc}} - iku^{\text{inc}} \text{ on } \Gamma_R$$

- Radiation condition at finite distance




# Time harmonic equations for the TM polarization

$$H_z = u^{\text{inc}} + u^{\text{sca}} \quad k = \frac{\omega}{c} \sqrt{\varepsilon_d \mu_d}$$

$$\operatorname{div} \left( \frac{1}{\varepsilon(\omega)} \nabla H_z \right) + \frac{\omega^2}{c^2} \mu H_z = 0 \text{ in } B_R$$

$$\partial_n H_z - ik H_z = \partial_n u^{\text{inc}} - iku^{\text{inc}} \text{ on } \Gamma_R$$

- Radiation condition at finite distance



Mathematically:

- due to the dissipation, the problem has a unique solution
- one can approximate the solution with Finite Elements Methods

# Numerical illustrations

---

We computed the total field for a triangular silver inclusion embedded in vacuum. We use Finite Element of order 2, with a plane wave of incidence  $-\pi/12$ .

# Numerical illustrations

---

We computed the total field for a triangular silver inclusion embedded in vacuum. We use Finite Element of order 2, with a plane wave of incidence  $-\pi/12$ .


$$\omega_p = 13.3 \text{ PHz} \quad \omega = 6 \text{ PHz} \quad \gamma = 0.113 \text{ PHz}$$

$$\varepsilon_m(\omega) = -3.9193 + 0.0926i \quad \varepsilon_d = \mu_d = \mu_m = 1$$

# Numerical illustrations

We computed the total field for a triangular silver inclusion embedded in vacuum. We use Finite Element of order 2, with a plane wave of incidence  $-\pi/12$ .

$$\begin{aligned}\omega_p &= 13.3 \text{ PHz} & \omega &= 6 \text{ PHz} & \gamma &= 0.113 \text{ PHz} \\ \varepsilon_m(\omega) &= -3.9193 + 0.0926i & \varepsilon_d = \mu_d = \mu_m &= 1\end{aligned}$$




coarse mesh


# Numerical illustrations

We computed the total field for a triangular silver inclusion embedded in vacuum. We use Finite Element of order 2, with a plane wave of incidence  $-\pi/12$ .

$$\begin{aligned}\omega_p &= 13.3 \text{ PHz} & \omega &= 6 \text{ PHz} & \gamma &= 0.113 \text{ PHz} \\ \varepsilon_m(\omega) &= -3.9193 + 0.0926i & \varepsilon_d = \mu_d = \mu_m &= 1\end{aligned}$$



coarse mesh





refined mesh

# Numerical illustrations

We computed the total field for a triangular silver inclusion embedded in vacuum. We use Finite Element of order 2, with a plane wave of incidence  $-\pi/12$ .

$$\begin{aligned}\omega_p &= 13.3 \text{ PHz} & \omega &= 6 \text{ PHz} & \gamma &= 0.113 \text{ PHz} \\ \varepsilon_m(\omega) &= -3.9193 + 0.0926i & \varepsilon_d = \mu_d = \mu_m &= 1\end{aligned}$$





**The solution is not stable ! Strong oscillations near the corners.**

# Numerical illustrations

We computed the total field for a triangular silver inclusion embedded in vacuum. We use Finite Element of order 2, with a plane wave of incidence  $-\pi/12$ .

$$\begin{aligned}\omega_p &= 13.3 \text{ PHz} & \omega &= 6 \text{ PHz} & \gamma &= 0.113 \text{ PHz} \\ \varepsilon_m(\omega) &= -3.9193 + 0.0926i & \varepsilon_d = \mu_d = \mu_m &= 1\end{aligned}$$



**The solution is not stable ! Strong oscillations near the corners.**

To understand the reasons of such instabilities and how to avoid them, we study a **limit problem by neglecting dissipation**.

# The dissipationless Drude's model

---

At optical frequencies ( $\gamma \ll \omega < \omega_p$ ), one can neglect dissipation so that the metal's permittivity follows the law:

$$\varepsilon = \varepsilon^0(\omega) := 1 - \frac{\omega_p^2}{\omega^2} < 0$$

# The dissipationless Drude's model

---

At optical frequencies ( $\gamma \ll \omega < \omega_p$ ), one can neglect dissipation so that the metal's permittivity follows the law:

$$\varepsilon = \varepsilon^0(\omega) := 1 - \frac{\omega_p^2}{\omega^2} < 0$$

# The dissipationless Drude's model

---

At optical frequencies ( $\gamma \ll \omega < \omega_p$ ), one can neglect dissipation so that the metal's permittivity follows the law:

$$\varepsilon = \varepsilon^0(\omega) := 1 - \frac{\omega_p^2}{\omega^2} < 0$$

Mathematically this implies for our problem:

- difficulties to prove existence and uniqueness of the solution
- the corners of the inclusion may cause strong singular behavior
- standard approximation with Finite Elements Methods may fail

# The dissipationless Drude's model

At optical frequencies ( $\gamma \ll \omega < \omega_p$ ), one can neglect dissipation so that the metal's permittivity follows the law:

$$\varepsilon = \varepsilon^0(\omega) := 1 - \frac{\omega_p^2}{\omega^2} < 0$$

Mathematically this implies for our problem:

- difficulties to prove existence and uniqueness of the solution
- the corners of the inclusion may cause strong singular behavior
- standard approximation with Finite Elements Methods may fail

The goal of my Ph.D was to give an answer theoretically and numerically to this problem.

# The dissipationless Drude's model

At optical frequencies ( $\gamma \ll \omega < \omega_p$ ), one can neglect dissipation so that the metal's permittivity follows the law:

$$\varepsilon = \varepsilon^0(\omega) := 1 - \frac{\omega_p^2}{\omega^2} < 0$$

Mathematically this implies for our problem:

- difficulties to prove existence and uniqueness of the solution
- the corners of the inclusion may cause strong singular behavior
- standard approximation with Finite Elements Methods may fail

The goal of my Ph.D was to give an answer theoretically and numerically to this problem.

This work can be used to study scattering problems with negative index metamaterials.

# The dissipationless Drude's model

At optical frequencies ( $\gamma \ll \omega < \omega_p$ ), one can neglect dissipation so that the metal's permittivity follows the law:

$$\varepsilon = \varepsilon^0(\omega) := 1 - \frac{\omega_p^2}{\omega^2} < 0$$

Mathematically this implies for our problem:

- difficulties to prove existence and uniqueness of the solution
- the corners of the inclusion may cause strong singular behavior
- standard approximation with Finite Elements Methods may fail

The goal of my Ph.D was to give an answer theoretically and numerically to this problem.

This work can be used to study scattering problems with negative index metamaterials.

Once some answers have been given, we studied the guided modes of a dissipationless plasmonic waveguide.


# Outline

---

- \* Introduction
- \* **Scattering problem with sign-changing coefficients**
- \* Guided modes in a plasmonic waveguide

# Variational formulation

$$\left| \begin{array}{l} \text{Find } u := u^{\text{inc}} + u^{\text{sac}} \in H^1(D_R) \text{ such that: } H^1(D_R) := \{u | \int_{D_R} |u|^2 + |\nabla u|^2 d\mathbf{x} < +\infty\} \\ \text{div}(\varepsilon^{-1} \nabla u) + \frac{\omega^2}{c^2} \mu u = 0 \quad \text{in } D_R, \\ \partial_n u - iku = \partial u^{\text{inc}} - iku^{\text{inc}} \quad \text{on } \partial D_R. \end{array} \right.$$




# Variational formulation

$$\left| \begin{array}{l} \text{Find } u := u^{\text{inc}} + u^{\text{sac}} \in H^1(D_R) \text{ such that: } H^1(D_R) := \{u | \int_{D_R} |u|^2 + |\nabla u|^2 d\mathbf{x} < +\infty\} \\ \text{div}(\varepsilon^{-1} \nabla u) + \frac{\omega^2}{c^2} \mu u = 0 \quad \text{in } D_R, \\ \partial_n u - iku = \partial u^{\text{inc}} - iku^{\text{inc}} \quad \text{on } \partial D_R. \end{array} \right.$$


The variational formulation of the scattering problem writes:

$$\left| \begin{array}{l} \text{Find } u := u^{\text{inc}} + u^{\text{sac}} \in H^1(D_R) \text{ such that:} \\ a(u, v) + c(u, v) = l(v), \quad \forall v \in H^1(D_R) \end{array} \right.$$



# Variational formulation

$$\left| \begin{array}{l} \text{Find } u := u^{\text{inc}} + u^{\text{sac}} \in H^1(D_R) \text{ such that: } H^1(D_R) := \{u | \int_{D_R} |u|^2 + |\nabla u|^2 d\mathbf{x} < +\infty\} \\ \text{div}(\varepsilon^{-1} \nabla u) + \frac{\omega^2}{c^2} \mu u = 0 \quad \text{in } D_R, \\ \partial_n u - iku = \partial u^{\text{inc}} - iku^{\text{inc}} \quad \text{on } \partial D_R. \end{array} \right.$$




The variational formulation of the scattering problem writes:

$$\left| \begin{array}{l} \text{Find } u := u^{\text{inc}} + u^{\text{sac}} \in H^1(D_R) \text{ such that:} \\ a(u, v) + c(u, v) = l(v), \quad \forall v \in H^1(D_R) \\ a(u, v) := \int_{D_R} \varepsilon^{-1} \nabla u \cdot \overline{\nabla v} d\mathbf{x}, \quad c(u, v) := -\frac{\omega^2}{c^2} \int_{D_R} \mu u \bar{v} d\mathbf{x} - i \frac{k}{\varepsilon_d} \int_{\partial D_R} u \bar{v} d\sigma, \\ l(v) := \int_{\partial D_R} (\partial_n u^{\text{inc}} - iku^{\text{inc}}) \bar{v} d\sigma. \end{array} \right.$$

# Variational formulation

$$\left| \begin{array}{l} \text{Find } u := u^{\text{inc}} + u^{\text{sac}} \in H^1(D_R) \text{ such that: } H^1(D_R) := \{u | \int_{D_R} |u|^2 + |\nabla u|^2 d\mathbf{x} < +\infty\} \\ \text{div}(\varepsilon^{-1} \nabla u) + \frac{\omega^2}{c^2} \mu u = 0 \quad \text{in } D_R, \\ \partial_n u - iku = \partial u^{\text{inc}} - iku^{\text{inc}} \quad \text{on } \partial D_R. \end{array} \right.$$




The variational formulation of the scattering problem writes:

$$\left| \begin{array}{l} \text{Find } u := u^{\text{inc}} + u^{\text{sac}} \in H^1(D_R) \text{ such that:} \\ a(u, v) + c(u, v) = l(v), \quad \forall v \in H^1(D_R) \\ a(u, v) := \int_{D_R} \varepsilon^{-1} \nabla u \cdot \overline{\nabla v} d\mathbf{x}, \quad c(u, v) := -\frac{\omega^2}{c^2} \int_{D_R} \mu u \overline{v} d\mathbf{x} - i \frac{k}{\varepsilon_d} \int_{\partial D_R} u \overline{v} d\sigma, \\ l(v) := \int_{\partial D_R} (\partial_n u^{\text{inc}} - iku^{\text{inc}}) \overline{v} d\sigma. \end{array} \right.$$

Classically (when  $\varepsilon > 0$ ) one has:  $a$  is **coercive**  
 $c$  is a **compact** perturbation

# Variational formulation

$$\left| \begin{array}{l} \text{Find } u := u^{\text{inc}} + u^{\text{sac}} \in H^1(D_R) \text{ such that: } H^1(D_R) := \{u | \int_{D_R} |u|^2 + |\nabla u|^2 d\mathbf{x} < +\infty\} \\ \text{div}(\varepsilon^{-1} \nabla u) + \frac{\omega^2}{c^2} \mu u = 0 \quad \text{in } D_R, \\ \partial_n u - iku = \partial u^{\text{inc}} - iku^{\text{inc}} \quad \text{on } \partial D_R. \end{array} \right.$$



The variational formulation of the scattering problem writes:


$$\left| \begin{array}{l} \text{Find } u := u^{\text{inc}} + u^{\text{sac}} \in H^1(D_R) \text{ such that:} \\ a(u, v) + c(u, v) = l(v), \quad \forall v \in H^1(D_R) \\ a(u, v) := \int_{D_R} \varepsilon^{-1} \nabla u \cdot \overline{\nabla v} d\mathbf{x}, \quad c(u, v) := -\frac{\omega^2}{c^2} \int_{D_R} \mu u \bar{v} d\mathbf{x} - i \frac{k}{\varepsilon_d} \int_{\partial D_R} u \bar{v} d\sigma, \\ l(v) := \int_{\partial D_R} (\partial_n u^{\text{inc}} - iku^{\text{inc}}) \bar{v} d\sigma. \end{array} \right.$$

Classically (when  $\varepsilon > 0$ ) one has:  $a$  is **coercive**  
 $c$  is a **compact** perturbation

**Fredholm** problem (uniqueness deduced from classical scattering theory)

# Variational formulation

$$\left| \begin{array}{l} \text{Find } u := u^{\text{inc}} + u^{\text{sac}} \in H^1(D_R) \text{ such that: } H^1(D_R) := \{u | \int_{D_R} |u|^2 + |\nabla u|^2 d\mathbf{x} < +\infty\} \\ \text{div}(\varepsilon^{-1} \nabla u) + \frac{\omega^2}{c^2} \mu u = 0 \quad \text{in } D_R, \\ \partial_n u - iku = \partial u^{\text{inc}} - iku^{\text{inc}} \quad \text{on } \partial D_R. \end{array} \right.$$



The variational formulation of the scattering problem writes:

$$\left| \begin{array}{l} \text{Find } u := u^{\text{inc}} + u^{\text{sac}} \in H^1(D_R) \text{ such that:} \\ a(u, v) + c(u, v) = l(v), \quad \forall v \in H^1(D_R) \\ a(u, v) := \int_{D_R} \varepsilon^{-1} \nabla u \cdot \overline{\nabla v} d\mathbf{x}, \quad c(u, v) := -\frac{\omega^2}{c^2} \int_{D_R} \mu u \overline{v} d\mathbf{x} - i \frac{k}{\varepsilon_d} \int_{\partial D_R} u \overline{v} d\sigma, \\ l(v) := \int_{\partial D_R} (\partial_n u^{\text{inc}} - iku^{\text{inc}}) \overline{v} d\sigma. \end{array} \right.$$

Classically (when  $\varepsilon > 0$ ) one has:  **$a$  is coercive**  
 **$c$  is a compact perturbation**


Fredholm problem (uniqueness deduced from classical scattering theory)

Due to sign-changing coefficient, it is not guaranteed !

# Study of non coercive forms

Find  $u := u^{\text{inc}} + u^{\text{sac}} \in H^1(D_R)$  such that:

$$a(u, v) + c(u, v) = l(v), \quad \forall v \in H^1(D_R)$$




# Study of non coercive forms

Find  $u := u^{\text{inc}} + u^{\text{sac}} \in H^1(D_R)$  such that:

$$a(u, v) + c(u, v) = l(v), \quad \forall v \in H^1(D_R)$$

**Coercivity** can be satisfied **under some conditions** on  $\varepsilon$  and the geometry.



*T-coercivity for scalar interface problems between dielectrics and metamaterials,*  
Anne-Sophie Bonnet-Ben Dhia, Lucas Chesnel, Patrick Ciarlet, M2AN, 2012


# Study of non coercive forms

Find  $u := u^{\text{inc}} + u^{\text{sac}} \in H^1(D_R)$  such that:

$$a(u, v) + c(u, v) = l(v), \quad \forall v \in H^1(D_R)$$

Coercivity can be satisfied under some conditions on  $\varepsilon$  and the geometry.

In our case: YES if and only if  $\kappa_\varepsilon := \frac{\varepsilon_m}{\varepsilon_d} \notin I_c$




# Study of non coercive forms

Find  $u := u^{\text{inc}} + u^{\text{sac}} \in H^1(D_R)$  such that:

$$a(u, v) + c(u, v) = l(v), \quad \forall v \in H^1(D_R)$$

Coercivity can be satisfied under some conditions on  $\varepsilon$  and the geometry.

In our case: YES if and only if  $\kappa_\varepsilon := \frac{\varepsilon_m}{\varepsilon_d} \notin I_c$   $I_c$  is called critical interval.



T-coercivity for scalar interface problems between dielectrics and metamaterials,  
Anne-Sophie Bonnet-Ben Dhia, Lucas Chesnel, Patrick Ciarlet, M2AN, 2012


# Study of non coercive forms

Find  $u := u^{\text{inc}} + u^{\text{sac}} \in H^1(D_R)$  such that:


$$a(u, v) + c(u, v) = l(v), \quad \forall v \in H^1(D_R)$$

**Coercivity** can be satisfied **under some conditions** on  $\varepsilon$  and the geometry.

In our case: YES if and only if  $\kappa_\varepsilon := \frac{\varepsilon_m}{\varepsilon_d} \notin I_c$   $I_c$  is called **critical interval**.




If the interface is smooth:



$$I_c = \{-1\}$$

If the interface has corners:



$$I_c = \left[ \frac{\phi - 2\pi}{\phi}; \frac{\phi}{\phi - 2\pi} \right]$$

$$\phi \rightarrow 0, I_c \rightarrow \mathbb{R}^-$$

$$\phi \rightarrow \pi, I_c \rightarrow \{-1\}$$



*T-coercivity for scalar interface problems between dielectrics and metamaterials,*  
Anne-Sophie Bonnet-Ben Dhia, Lucas Chesnel, Patrick Ciarlet, M2AN, 2012


# Study of non coercive forms

Find  $u := u^{\text{inc}} + u^{\text{sac}} \in H^1(D_R)$  such that:


$$a(u, v) + c(u, v) = l(v), \quad \forall v \in H^1(D_R)$$

Coercivity can be satisfied under some conditions on  $\varepsilon$  and the geometry.

In our case: YES if and only if  $\kappa_\varepsilon := \frac{\varepsilon_m}{\varepsilon_d} \notin I_c$   $I_c$  is called critical interval.




If the interface is smooth:



$$I_c = \{-1\}$$

If the interface has corners:



$$I_c = \left[ \frac{\phi - 2\pi}{\phi}; \frac{\phi}{\phi - 2\pi} \right]$$

$$\phi \rightarrow 0, I_c \rightarrow \mathbb{R}^-$$

$$\phi \rightarrow \pi, I_c \rightarrow \{-1\}$$

Using Drude's model, the critical interval is related to a critical range of frequencies.



T-coercivity for scalar interface problems between dielectrics and metamaterials,  
Anne-Sophie Bonnet-Ben Dhia, Lucas Chesnel, Patrick Ciarlet, M2AN, 2012


# Study of non coercive forms

Find  $u := u^{\text{inc}} + u^{\text{sac}} \in H^1(D_R)$  such that:


$$a(u, v) + c(u, v) = l(v), \quad \forall v \in H^1(D_R)$$

**Coercivity** can be satisfied **under some conditions** on  $\varepsilon$  and the geometry.

In our case: YES if and only if  $\kappa_\varepsilon := \frac{\varepsilon_m}{\varepsilon_d} \notin I_c$   $I_c$  is called **critical interval**.




If the interface is smooth:



$$I_c = \{-1\}$$

If the interface has corners:



$$I_c = \left[ \frac{\phi - 2\pi}{\phi}; \frac{\phi}{\phi - 2\pi} \right]$$

$$\phi \rightarrow 0, I_c \rightarrow \mathbb{R}^-$$

$$\phi \rightarrow \pi, I_c \rightarrow \{-1\}$$

Using **Drude's model**, the critical interval is related to a **critical range of frequencies**.

$$\kappa_\varepsilon = -1 \iff \omega = \omega_{\text{sp}} := \frac{\omega_p}{\sqrt{\varepsilon_d + 1}}.$$



*T-coercivity for scalar interface problems between dielectrics and metamaterials,*  
Anne-Sophie Bonnet-Ben Dhia, Lucas Chesnel, Patrick Ciarlet, M2AN, 2012

# Two configurations

---

Outside  $I_c$   
(YES)

Inside  $I_c$   
(NO)

# Two configurations

Outside  $I_c$   
(YES)

The scattering problem has a **unique solution**  $u \in H^1(D_R)$   
**Finite Elements** converge (under some condition on the mesh)

Inside  $I_c$   
(NO)

# Two configurations

Outside  $I_c$   
(YES)

The scattering problem has a **unique solution**  $u \in H^1(D_R)$   
**Finite Elements** converge (under some condition on the mesh)

Inside  $I_c$   
(NO)

The scattering problem is **ill-posed** in  $H^1(D_R)$   
**No FEM** convergence

# Two configurations

Outside  $I_c$   
(YES)

The scattering problem has a **unique solution**  $u \in H^1(D_R)$   
**Finite Elements** converge (under some condition on the mesh)

Inside  $I_c$   
(NO)

The scattering problem is **ill-posed** in  $H^1(D_R)$   
**No FEM** convergence

Numerical illustrations for triangular silver inclusion in vacuum:

# Two configurations

Outside  $I_c$   
(YES)

The scattering problem has a **unique solution**  $u \in H^1(D_R)$   
**Finite Elements** converge (under some condition on the mesh)

Inside  $I_c$   
(NO)

The scattering problem is **ill-posed** in  $H^1(D_R)$   
**No FEM** convergence

Numerical illustrations for triangular silver inclusion in vacuum:

$$\kappa_\varepsilon \in I_c \iff \omega \in [3.839 \text{ PHz}; 12.733 \text{ PHz}]. \quad \omega_{\text{sp}} = 9.42 \text{ PHz.}$$

Results for  $\omega = 9$  PHz.

# Two configurations

Outside  $I_c$   
(YES)

The scattering problem has a **unique solution**  $u \in H^1(D_R)$   
**Finite Elements** converge (under some condition on the mesh)


Inside  $I_c$   
(NO)

The scattering problem is **ill-posed** in  $H^1(D_R)$   
**No FEM** convergence


Numerical illustrations for triangular silver inclusion in vacuum:

$$\kappa_\varepsilon \in I_c \iff \omega \in [3.839 \text{ PHz}; 12.733 \text{ PHz}]. \quad \omega_{\text{sp}} = 9.42 \text{ PHz.}$$


Results for  $\omega = 9 \text{ PHz.}$



coarse mesh



intermediate mesh



refined mesh

# Two configurations

Outside  $I_c$   
(YES)

The scattering problem has a **unique solution**  $u \in H^1(D_R)$   
**Finite Elements** converge (under some condition on the mesh)


Inside  $I_c$   
(NO)

The scattering problem is **ill-posed** in  $H^1(D_R)$   
**No FEM** convergence

Numerical illustrations for triangular silver inclusion in vacuum:

$$\kappa_\varepsilon \in I_c \iff \omega \in [3.839 \text{ PHz}; 12.733 \text{ PHz}]. \quad \omega_{\text{sp}} = 9.42 \text{ PHz.}$$


Results for  $\omega = 9 \text{ PHz.}$



# The appearance of black-hole waves

One proves that for a frequency **inside** the critical frequency range, **oscillating singularities appear at the corners** called **black-hole waves**. They can be interpreted as plasmonic waves propagating at the corners but never reach it:


$$s^\pm(r, \theta) = e^{i(\pm\eta \ln r - \omega t)} \Phi(\theta)$$



# The appearance of black-hole waves

One proves that for a frequency **inside** the critical frequency range, **oscillating singularities appear at the corners** called **black-hole waves**. They can be interpreted as plasmonic waves propagating at the corners but never reach it:

$$s^\pm(r, \theta) = e^{i(\pm\eta \ln r - \omega t)} \Phi(\theta)$$




These black-hole waves are of **infinite energy** and strongly oscillating at the corners. This is why standard FEM fail to approximate the solution: spurious reflexions appear. However we **need to take them into account** to recover a **well-posed formulation**.

# The appearance of black-hole waves

One proves that for a frequency **inside** the critical frequency range, **oscillating singularities appear at the corners** called **black-hole waves**. They can be interpreted as plasmonic waves propagating at the corners but never reach it:

$$s^\pm(r, \theta) = e^{i(\pm\eta \ln r - \omega t)} \Phi(\theta)$$



These black-hole waves are of **infinite energy** and strongly oscillating at the corners. This is why standard FEM fail to approximate the solution: spurious reflexions appear. However we **need to take them into account** to recover a **well-posed formulation**.

Near the corners the solution decomposes as

$$u = b^+ s^+ + b^- s^- + \tilde{u}, \quad \tilde{u} \in H^1(D_R), \quad s^\pm \notin H^1(D_R), \quad b^\pm \in \mathbb{C}$$

# Zoom at the corner


---

To simplify, let us consider a metallic inclusion with only one corner.

# Zoom at the corner


---

To simplify, let us consider a metallic inclusion with only one corner.




# Zoom at the corner

To simplify, let us consider a metallic inclusion with only one corner.



# Zoom at the corner

To simplify, let us consider a metallic inclusion with only one corner.



Asymptotic behaviour near the corner


$$\operatorname{div} \left( \frac{1}{\epsilon} \nabla u \right) + \omega^2 \mu u = 0$$

Behaviour at infinity in the waveguide

$$\operatorname{div} \left( \frac{1}{\epsilon} \nabla u \right) + \omega^2 \mu e^{2z} u = 0$$

# Zoom at the corner

To simplify, let us consider a metallic inclusion with only one corner.



Asymptotic behaviour near the corner


$$\operatorname{div} \left( \frac{1}{\epsilon} \nabla u \right) + \omega^2 \mu u = 0$$

Behaviour at infinity in the waveguide

$$\operatorname{div} \left( \frac{1}{\epsilon} \nabla u \right) + \omega^2 \mu e^{2z} u = 0$$

# Zoom at the corner

To simplify, let us consider a metallic inclusion with only one corner.



Asymptotic behaviour near the corner

$$\operatorname{div} \left( \frac{1}{\epsilon} \nabla u \right) + \omega^2 \mu u = 0$$


Behaviour at infinity in the waveguide

$$\operatorname{div} \left( \frac{1}{\epsilon} \nabla u \right) + \omega^2 \mu e^{2z} u = 0$$

Modal analysis: looking for the solutions

# Zoom at the corner

To simplify, let us consider a metallic inclusion with only one corner.



Asymptotic behaviour near the corner

$$\operatorname{div} \left( \frac{1}{\epsilon} \nabla u \right) + \omega^2 \mu u = 0$$

Behaviour at infinity in the waveguide


$$\operatorname{div} \left( \frac{1}{\epsilon} \nabla u \right) + \omega^2 \mu e^{2z} u = 0$$

Modal analysis: looking for the solutions

$$u(z, \theta) = e^{\lambda z} \phi(\theta)$$

# Zoom at the corner

To simplify, let us consider a metallic inclusion with only one corner.



Asymptotic behaviour near the corner

$$\operatorname{div} \left( \frac{1}{\epsilon} \nabla u \right) + \omega^2 \mu u = 0$$

Behaviour at infinity in the waveguide


$$\operatorname{div} \left( \frac{1}{\epsilon} \nabla u \right) + \omega^2 \mu e^{2z} u = 0$$

Modal analysis: looking for the solutions

$$u(z, \theta) = e^{\lambda z} \phi(\theta)$$


Appearance of two propagative modes

$$s^\pm(z, \theta) = e^{\pm i \kappa z} \phi(\theta)$$



# Zoom at the corner

To simplify, let us consider a metallic inclusion with only one corner.



Asymptotic behaviour near the corner

$$\operatorname{div} \left( \frac{1}{\epsilon} \nabla u \right) + \omega^2 \mu u = 0$$

In the physical domain, the solutions are

$$u(r, \theta) = r^\lambda \phi(\theta)$$

Behaviour at infinity in the waveguide


$$\operatorname{div} \left( \frac{1}{\epsilon} \nabla u \right) + \omega^2 \mu e^{2z} u = 0$$

Modal analysis: looking for the solutions

$$u(z, \theta) = e^{\lambda z} \phi(\theta)$$


Appearance of two propagative modes

$$s^\pm(z, \theta) = e^{\pm i \kappa z} \phi(\theta)$$



# Zoom at the corner

To simplify, let us consider a metallic inclusion with only one corner.



Asymptotic behaviour near the corner


$$\operatorname{div} \left( \frac{1}{\epsilon} \nabla u \right) + \omega^2 \mu u = 0$$

In the physical domain, the solutions are

$$u(r, \theta) = r^\lambda \phi(\theta)$$

Appearance of two **black-hole** waves

$$s^\pm(r, \theta) = r^{\pm i \kappa} \phi(\theta)$$



Behaviour at infinity in the waveguide


$$\operatorname{div} \left( \frac{1}{\epsilon} \nabla u \right) + \omega^2 \mu e^{2z} u = 0$$

Modal analysis: looking for the solutions

$$u(z, \theta) = e^{\lambda z} \phi(\theta)$$

Appearance of two **propagative** modes


$$s^\pm(z, \theta) = e^{\pm i \kappa z} \phi(\theta)$$



This is due to sign-changing coefficient !

# Zoom at the corner


To simplify, let us consider a metallic inclusion with only one corner.



One singularity carries energy absorbed by the corner, the other brings energy emitted from the corner.


Appearance of two **black-hole** waves

$$s^\pm(r, \theta) = r^{\pm i\kappa} \phi(\theta)$$



Appearance of two **propagative** modes


$$s^\pm(z, \theta) = e^{\pm i\kappa z} \phi(\theta)$$



This is due to sign-changing coefficient !

# Zoom at the corner

To simplify, let us consider a metallic inclusion with only one corner.




One singularity carries energy absorbed by the corner, the other brings energy emitted from the corner.

The **physical** solution takes into account the one that does not add energy into the system, the **outgoing** solution in the waveguide setting.


Appearance of two **black-hole** waves

$$s^\pm(r, \theta) = r^{\pm i\kappa} \phi(\theta)$$



Appearance of two **propagative** modes


$$s^\pm(z, \theta) = e^{\pm i\kappa z} \phi(\theta)$$



This is due to sign-changing coefficient !

# Zoom at the corner

To simplify, let us consider a metallic inclusion with only one corner.




One singularity carries energy absorbed by the corner, the other brings energy emitted from the corner.

The **physical** solution takes into account the one that does not add energy into the system, the **outgoing** solution in the waveguide setting.

How to proceed numerically to select the good singularity ?


Appearance of two **black-hole** waves

$$s^\pm(r, \theta) = r^{\pm i\kappa} \phi(\theta)$$



Appearance of two **propagative** modes

$$s^\pm(z, \theta) = e^{\pm i\kappa z} \phi(\theta)$$



This is due to sign-changing coefficient !

# An adapted numerical method

---


We put a PML in the Strip: then the propagatives modes become evanescent, and we can use FEM.

$$\frac{\partial}{\partial z} \longmapsto \alpha \frac{\partial}{\partial z} \quad \alpha \in \mathbb{C}, \quad 0 < \arg(\alpha) < \frac{\pi}{2}$$

# An adapted numerical method

We put a **PML** in the **Strip**: then the propagatives modes become evanescent, and we can use **FEM**.


$$\frac{\partial}{\partial z} \longmapsto \alpha \frac{\partial}{\partial z} \quad \alpha \in \mathbb{C}, \quad 0 < \arg(\alpha) < \frac{\pi}{2}$$






# An adapted numerical method

We put a **PML** in the **Strip**: then the propagatives modes become evanescent, and we can use **FEM**.



$$\frac{\partial}{\partial z} \longmapsto \alpha \frac{\partial}{\partial z} \quad \alpha \in \mathbb{C}, \quad 0 < \arg(\alpha) < \frac{\pi}{2}$$



$$u(z, \theta) = e^{\lambda z} \phi(\theta) \longrightarrow u(z, \theta) = e^{\frac{\lambda}{\alpha} z} \phi(\theta)$$




$$u(r, \theta) = r^\lambda \phi(\theta) \longrightarrow u(r, \theta) = r^{\frac{\lambda}{\alpha}} \phi(\theta)$$






# An adapted numerical method

We put a PML in the Strip: then the propagatives modes become evanescent, and we can use FEM.



$$\frac{\partial}{\partial z} \longmapsto \alpha \frac{\partial}{\partial z} \quad \alpha \in \mathbb{C}, \quad 0 < \arg(\alpha) < \frac{\pi}{2}$$



$$u(z, \theta) = e^{\lambda z} \phi(\theta) \longrightarrow u(z, \theta) = e^{\frac{\lambda}{\alpha} z} \phi(\theta)$$




$$u(r, \theta) = r^\lambda \phi(\theta) \longrightarrow u(r, \theta) = r^{\frac{\lambda}{\alpha}} \phi(\theta)$$



PML easier to implement in the Strip

# Implementation: splitting the domain



$$\operatorname{div} \left( \frac{1}{\epsilon} \nabla u \right) + \omega^2 c^{-2} \mu u = 0 \quad D_R \setminus \cup \text{PML}$$

$$\partial_n u - iku = \partial_n u^{\text{inc}} - iku^{\text{inc}} \quad \partial D_R$$

+ PML matching conditions +

$$\frac{\alpha}{\epsilon} \partial_{zz} u + \frac{1}{\alpha} \partial_\theta \frac{1}{\epsilon} \partial_\theta u + \frac{\omega^2}{\alpha c^2} \mu e^{\frac{2z}{\alpha}} u = 0 \quad \text{PML}$$

$$\partial_z u(-L, \cdot) = 0$$

+ periodic conditions

Resolution using Lagrange FE of order 2 with a Matlab code.


# Numerical results

---


$\kappa_\varepsilon \in I_c \iff \omega \in [3.839 \text{ PHz}; 12.733 \text{ PHz}]$ .  $\omega_{\text{sp}} = 9.42 \text{ PHz}$ . Results for  $\omega = 9 \text{ PHz}$ .

# Numerical results


$\kappa_\varepsilon \in I_c \iff \omega \in [3.839 \text{ PHz}; 12.733 \text{ PHz}]$ .  $\omega_{\text{sp}} = 9.42 \text{ PHz}$ . Results for  $\omega = 9 \text{ PHz}$ .



coarse mesh




intermediate mesh



refined mesh


# Numerical results

$\kappa_\varepsilon \in I_c \iff \omega \in [3.839 \text{ PHz}; 12.733 \text{ PHz}]$ .  $\omega_{\text{sp}} = 9.42 \text{ PHz}$ . Results for  $\omega = 9 \text{ PHz}$ .



# Numerical results

$\kappa_\varepsilon \in I_c \iff \omega \in [3.839 \text{ PHz}; 12.733 \text{ PHz}]$ .  $\omega_{\text{sp}} = 9.42 \text{ PHz}$ . Results for  $\omega = 9 \text{ PHz}$ .



The numerical method with PMLs is stable. Let us go back to the dissipative problem.

# Back to dissipative medium

---

# Back to dissipative medium

---

Considering losses in the metal, then the problem is well-posed.

The selected outgoing singularity becomes of finite energy. If dissipation is small, it requires meshes sufficiently refined at the corners to capture the oscillations.

# Back to dissipative medium

---

Considering losses in the metal, then the problem is well-posed.

The selected outgoing singularity becomes of finite energy. If dissipation is small, it requires meshes sufficiently refined at the corners to capture the oscillations.

$$\kappa_\varepsilon \in I_c \iff \omega \in [3.839 \text{ PHz}; 12.733 \text{ PHz}].$$

$$\omega_p = 13.3 \text{ PHz} \quad \omega = 6 \text{ PHz} \quad \gamma = 0.113 \text{ PHz}$$

$$\varepsilon_m(\omega) = -3.9193 + 0.0926i \quad \varepsilon_d = \mu_d = \mu_m = 1$$

# Back to dissipative medium

Considering losses in the metal, then the problem is well-posed.

The selected outgoing singularity becomes of finite energy. If dissipation is small, it requires meshes sufficiently refined at the corners to capture the oscillations.

$$\kappa_\varepsilon \in I_c \iff \omega \in [3.839 \text{ PHz}; 12.733 \text{ PHz}].$$

$$\omega_p = 13.3 \text{ PHz} \quad \omega = 6 \text{ PHz} \quad \gamma = 0.113 \text{ PHz}$$

$$\varepsilon_m(\omega) = -3.9193 + 0.0926i \quad \varepsilon_d = \mu_d = \mu_m = 1$$

Without PMLs

With PMLs

# Back to dissipative medium

Considering losses in the metal, then the problem is well-posed.


The selected outgoing singularity becomes of finite energy. If dissipation is small, it requires meshes sufficiently refined at the corners to capture the oscillations.

$$\kappa_\varepsilon \in I_c \iff \omega \in [3.839 \text{ PHz}; 12.733 \text{ PHz}].$$


$$\omega_p = 13.3 \text{ PHz} \quad \omega = 6 \text{ PHz} \quad \gamma = 0.113 \text{ PHz}$$

$$\varepsilon_m(\omega) = -3.9193 + 0.0926i \quad \varepsilon_d = \mu_d = \mu_m = 1$$

Without PMLs



coarse mesh



refined mesh

With PMLs



# Back to dissipative medium

Considering losses in the metal, then the problem is well-posed.


The selected outgoing singularity becomes of finite energy. If dissipation is small, it requires meshes sufficiently refined at the corners to capture the oscillations.

$$\kappa_\varepsilon \in I_c \iff \omega \in [3.839 \text{ PHz}; 12.733 \text{ PHz}].$$


$$\omega_p = 13.3 \text{ PHz} \quad \omega = 6 \text{ PHz} \quad \gamma = 0.113 \text{ PHz}$$

$$\varepsilon_m(\omega) = -3.9193 + 0.0926i \quad \varepsilon_d = \mu_d = \mu_m = 1$$

Without PMLs




coarse mesh




refined mesh

With PMLs



coarse mesh



refined mesh

# Perspectives and ongoing work

---

# Perspectives and ongoing work

---

Consider more relevant models for the metal permittivity, such as Drude-Lorentz model or hydrodynamic Drude's model.

# Perspectives and ongoing work

---

Consider more relevant models for the metal permittivity, such as Drude-Lorentz model or hydrodynamic Drude's model.

Numerical analysis to prove convergence of the method with PMLs is still in progress (difficulties related to the change of sign and the functional framework).

# Perspectives and ongoing work

---

Consider more relevant models for the metal permittivity, such as Drude-Lorentz model or hydrodynamic Drude's model.

Numerical analysis to prove convergence of the method with PMLs is still in progress (difficulties related to the change of sign and the functional framework).


This work is the object of an article (submitted very soon). The case of dispersive media is the object of discussions with the Fresnel Institute (Marseille, France).

# Outline


---

- \* Introduction
- \* Scattering problem with sign-changing coefficients
- \* **Guided modes in a plasmonic waveguide**


# Guided modes in a plasmonic waveguide




# Guided modes in a plasmonic waveguide



# Guided modes in a plasmonic waveguide




# Guided modes in a plasmonic waveguide



Metal permittivity modeled by the  
dissipationless Drude's model

$$\epsilon(\omega) = \epsilon_{\infty} \left( 1 - \frac{\omega_p^2}{\omega^2} \right)$$

# Guided modes in a plasmonic waveguide




Metal permittivity modeled by the  
**dissipationless Drude's model**

$$\epsilon(\omega) = \epsilon_\infty \left(1 - \frac{\omega_p^2}{\omega^2}\right)$$

We consider the time-harmonic scalar model of Maxwell's equations (simplified model).

# Guided modes in a plasmonic waveguide




Metal permittivity modeled by the  
dissipationless Drude's model

$$\epsilon(\omega) = \epsilon_\infty \left(1 - \frac{\omega_p^2}{\omega^2}\right)$$

We consider the time-harmonic scalar model of Maxwell's equations (simplified model).

Looking for propagative waves along  $z$ :  $u(x, y, z, t) = \tilde{u}(x, y)e^{i(\beta z - \omega t)}$   $\beta, \omega \in \mathbb{R}$

# Guided modes in a plasmonic waveguide




Metal permittivity modeled by the  
dissipationless Drude's model

$$\epsilon(\omega) = \epsilon_\infty \left(1 - \frac{\omega_p^2}{\omega^2}\right)$$

We consider the time-harmonic scalar model of Maxwell's equations (simplified model).

Looking for propagative waves along  $z$ :  $u(x, y, z, t) = \tilde{u}(x, y)e^{i(\beta z - \omega t)}$   $\beta, \omega \in \mathbb{R}$

# Guided modes in a plasmonic waveguide




Metal permittivity modeled by the  
dissipationless Drude's model

$$\epsilon(\omega) = \epsilon_\infty \left(1 - \frac{\omega_p^2}{\omega^2}\right)$$


We consider the time-harmonic scalar model of Maxwell's equations (simplified model).

Looking for propagative waves along  $z$ :  $u(x, y, z, t) = \tilde{u}(x, y)e^{i(\beta z - \omega t)}$   $\beta, \omega \in \mathbb{R}$

↓ Reduction to a 2D Eigenproblem  $A(\beta)\tilde{u} = \omega^2\tilde{u}$



# Guided modes in a plasmonic waveguide



Metal permittivity modeled by the  
dissipationless Drude's model

$$\epsilon(\omega) = \epsilon_\infty \left(1 - \frac{\omega_p^2}{\omega^2}\right)$$


We consider the time-harmonic scalar model of Maxwell's equations (simplified model).

Looking for propagative waves along  $z$ :  $u(x, y, z, t) = \tilde{u}(x, y)e^{i(\beta z - \omega t)}$   $\beta, \omega \in \mathbb{R}$

↓ Reduction to a 2D Eigenproblem


$$A(\beta)\tilde{u} = \omega^2\tilde{u}$$

$$\operatorname{div} \left( \frac{1}{\epsilon} \nabla \tilde{u} \right) - \frac{\beta^2}{\epsilon} \tilde{u} + \omega^2 \mu \tilde{u} = 0 \quad \Omega \quad \mu > 0$$



- linearized eigenproblem
- sign-changing coefficient
- corners

# Some numerical experiments (1)




For  $\beta \in \mathbb{R}$ , Find  $\tilde{u} \in V_0 \setminus \{0\}$ ,  $\omega \in \mathbb{C}$  s.t.  
 $A(\beta)\tilde{u} = \omega^2 \tilde{u}$

Numerical illustrations with FE

Parameters  $\epsilon_1 = 1$   $\epsilon_2 = -\frac{10}{7}$   $\beta = 1$

# Some numerical experiments (1)




For  $\beta \in \mathbb{R}$ , Find  $\tilde{u} \in V_0 \setminus \{0\}$ ,  $\omega \in \mathbb{C}$  s.t.  
 $A(\beta)\tilde{u} = \omega^2 \tilde{u}$


Numerical illustrations with FE

Parameters  $\epsilon_1 = 1$   $\epsilon_2 = -\frac{10}{7}$   $\beta = 1$

Coarse mesh




# Some numerical experiments (1)




For  $\beta \in \mathbb{R}$ , Find  $\tilde{u} \in V_0 \setminus \{0\}$ ,  $\omega \in \mathbb{C}$  s.t.  
 $A(\beta)\tilde{u} = \omega^2 \tilde{u}$

Numerical illustrations with FE

Parameters  $\epsilon_1 = 1$   $\epsilon_2 = -\frac{10}{7}$   $\beta = 1$




# Some numerical experiments (1)




For  $\beta \in \mathbb{R}$ , Find  $\tilde{u} \in V_0 \setminus \{0\}$ ,  $\omega \in \mathbb{C}$  s.t.  
 $A(\beta)\tilde{u} = \omega^2 \tilde{u}$

Numerical illustrations with FE

Parameters  $\epsilon_1 = 1$   $\epsilon_2 = -\frac{10}{7}$   $\beta = 1$




# Some numerical experiments (1)




For  $\beta \in \mathbb{R}$ , Find  $\tilde{u} \in V_0 \setminus \{0\}$ ,  $\omega \in \mathbb{C}$  s.t.  
 $A(\beta)\tilde{u} = \omega^2 \tilde{u}$

Numerical illustrations with FE


Parameters  $\epsilon_1 = 1$   $\epsilon_2 = -\frac{10}{7}$   $\beta = 1$



For a coarse mesh




# Some numerical experiments (1)





For  $\beta \in \mathbb{R}$ , Find  $\tilde{u} \in V_0 \setminus \{0\}$ ,  $\omega \in \mathbb{C}$  s.t.  
 $A(\beta)\tilde{u} = \omega^2 \tilde{u}$

Numerical illustrations with FE

Parameters  $\epsilon_1 = 1$   $\epsilon_2 = -\frac{10}{7}$   $\beta = 1$



For a coarse mesh




For a fine mesh

No convergence due to black-hole waves




# Some numerical experiments (1)




For  $\beta \in \mathbb{R}$ , Find  $\tilde{u} \in V_0 \setminus \{0\}$ ,  $\omega \in \mathbb{C}$  s.t.  
 $A(\beta)\tilde{u} = \omega^2 \tilde{u}$

Numerical illustrations with FE

Parameters  $\epsilon_1 = 1$   $\epsilon_2 = -\frac{10}{7}$   $\beta = 1$




For a coarse mesh




For a fine mesh

No convergence due to black-hole waves




# Some numerical experiments (1)




For  $\beta \in \mathbb{R}$ , Find  $\tilde{u} \in V_0 \setminus \{0\}$ ,  $\omega \in \mathbb{C}$  s.t.  
 $A(\beta)\tilde{u} = \omega^2 \tilde{u}$

Numerical illustrations with FE

Parameters  $\epsilon_1 = 1$   $\epsilon_2 = -\frac{10}{7}$   $\beta = 1$



For a coarse mesh




For a fine mesh

No convergence due to black-hole waves



Coarse mesh

Fine mesh

$\text{Im}(\omega^2)$




For a coarse mesh



For a fine mesh

# Some numerical experiments (1)




For  $\beta \in \mathbb{R}$ , Find  $\tilde{u} \in V_0 \setminus \{0\}$ ,  $\omega \in \mathbb{C}$  s.t.


$$A(\beta)\tilde{u} = \omega^2 \tilde{u}$$

Numerical illustrations with FE

Parameters  $\epsilon_1 = 1$   $\epsilon_2 = -\frac{10}{7}$   $\beta = 1$




For a coarse mesh




For a fine mesh


No convergence due to



No FEM convergence but some modes seem stable: guided modes are hidden in this spectrum !




For a coarse mesh




For a fine mesh

# Some numerical experiments (2)




For  $\beta \in \mathbb{R}$ , Find  $\tilde{u} \in V_0 \setminus \{0\}$ ,  $\omega \in \mathbb{C}$  s.t.  
 $A(\beta)\tilde{u} = \omega^2 \tilde{u}$

New numerical method which is stable, sorts the modes and reveals the guided modes.




# Some numerical experiments (2)




For  $\beta \in \mathbb{R}$ , Find  $\tilde{u} \in V_0 \setminus \{0\}$ ,  $\omega \in \mathbb{C}$  s.t.  
 $A(\beta)\tilde{u} = \omega^2 \tilde{u}$

New numerical method which is stable, sorts the modes and reveals the guided modes.




# Some numerical experiments (2)




For  $\beta \in \mathbb{R}$ , Find  $\tilde{u} \in V_0 \setminus \{0\}$ ,  $\omega \in \mathbb{C}$  s.t.  
 $A(\beta)\tilde{u} = \omega^2 \tilde{u}$

New numerical method which is stable, sorts the modes and reveals the guided modes.




# Some numerical experiments (2)



For  $\beta \in \mathbb{R}$ , Find  $\tilde{u} \in V_0 \setminus \{0\}$ ,  $\omega \in \mathbb{C}$  s.t.  
 $A(\beta)\tilde{u} = \omega^2 \tilde{u}$


New numerical method which is stable, sorts the modes and reveals the guided modes.



$\omega^2 \in \mathbb{R}$   
guided/evanescent  
modes


$\omega^2 \in \mathbb{C} \setminus \mathbb{R}$   
singular modes

# Some numerical experiments (2)



For  $\beta \in \mathbb{R}$ , Find  $\tilde{u} \in V_0 \setminus \{0\}$ ,  $\omega \in \mathbb{C}$  s.t.  
 $A(\beta)\tilde{u} = \omega^2 \tilde{u}$

New numerical method which is stable, sorts the modes and reveals the guided modes.



Explanation via energy technique

# Ongoing work

---

- ❖ Numerical analysis requires some conditions on the mesh
- ❖ Extensions to Maxwell's equations are currently in progress.
- ❖ Non linear eigenproblem (dissipationless Drude's model).
- ❖ What's happening when metamaterials ?
- ❖ Open plasmonic waveguides.

## Publications and communications

- ❖ 3 articles in preparation
- ❖ Several international communications: WAVES 13, PIERS 13, KOZVAWES 14, OWTNM 14, Leaky Days 15, WAVES 15.

---

---

Thank you for your attention.