






north creek through time
ei_obs
aridity_index
1.0
0.9
0.8
0.7
0.6
0.25
0.50
0.75
1.00
Year
2020
2015
2010
2005
2000
2010
2008
2007
2012
2013
2016
2018
2011
2005
2017
2006
2004
2009
2014
2023
2019
2022
2015
2020
2021



aridity_index
ei_obs
ratio
1.5
1.0
0.5
val
2000
2005
2010
2015
2020
2025
year



EB through time
ei_obs
aridity_index
1.00
0.75
0.50
0.25
0.00
0.00
0.25
0.50
0.75
1.00
2001
2002
2008
2007
2015
2000
2011
2012
2009
2013
Year
2020
2015
2010
2005
2000



aridity_index
ei_obs
ratio
1.00
0.75
0.50
0.25
0.00
val
2000
2005
2010
2015
2020
2025
year



RS02 through time
ei_obs
aridity_index
1.0
0.9
0.8
0.7
0.6
1
2
3
4
2015
2016
2014
2018
2013
2017
2004
2003
2009
2006
2011
2005
Year
2020
2015
2010
2005
2000



aridity_index
ei_obs
ratio
val
year
1
2
3
4
2000
2005
2010
2015
2020
2025



AB00 through time
evaporative_index
aridity_index
2.0
1.5
1.0
0.5
1
2
3
4
5
2007
2021
2002
2018
2014
2004
2015
2016
2013
2020
2012
2006
2003
2000
2001
2009
2008
2023
2017
2010
2005
2011
Year
2020
2015
2010
2005
2000



aridity_index
evaporative_index
ratio
val
year
1
2
3
4
5
2000
2005
2010
2015
2020
2025



Residuals vs Fitted
Residuals
0.2
0.1
0.0
−0.1
−0.2
24
6
10
0.42
0.44
0.46
0.48
Fitted values
lm(ratio ~ year)



Q−Q Residuals
Standardized residuals
Theoretical Quantiles
lm(ratio ~ year)
24
6
10
−2
−1
0
1
2



Scale−Location
24
10
6
1.5
1.0
0.5
0.0
√|Standardized residuals|
0.42
0.44
0.46
0.48
Fitted values
lm(ratio ~ year)



Residuals vs Leverage
Standardized residuals
Leverage
lm(ratio ~ year)
Cook's distance
0.5
0.5
24
6
10
2
1
0
−1
−2
0.00
0.05
0.10
0.15



Residuals vs Fitted
Residuals
Fitted values
lm(aridity_index ~ year)
8
15
3
2
1
0
−1
−2
2.65
2.70
2.75
2.80
2.85
2.90
2.95



Q−Q Residuals
Standardized residuals
Theoretical Quantiles
lm(aridity_index ~ year)
25
15
8
−2
−1
0
1
2
−1.5
−1.0
−0.5
0.0
0.5
1.0
1.5
2.0



Scale−Location
8
15
25
√|Standardized residuals|
0.0
0.2
0.4
0.6
0.8
1.0
1.2
2.65
2.70
2.75
2.80
2.85
2.90
2.95
Fitted values
lm(aridity_index ~ year)



Residuals vs Leverage
Standardized residuals
2
1
0
−1
3
24
25
Cook's distance
0.00
0.05
0.10
0.15
Leverage
lm(aridity_index ~ year)



Residuals vs Fitted
Residuals
1.0
0.5
0.0
−0.5
8
3
22
1.12
1.14
1.16
1.18
1.20
1.22
1.24
1.26
Fitted values
lm(evaporative_index ~ year)



Q−Q Residuals
Standardized residuals
Theoretical Quantiles
lm(evaporative_index ~ year)
8
3
22
−2
−1
0
1
2



Scale−Location
√|Standardized residuals|
0.0
0.5
1.0
1.5
8
3
22
1.12
1.14
1.16
1.18
1.20
1.22
1.24
1.26
Fitted values
lm(evaporative_index ~ year)



Residuals vs Leverage
Standardized residuals
Leverage
lm(evaporative_index ~ year)
Cook's distance
0.5
8
3
22
2
1
0
−1
0.00
0.05
0.10
0.15



w6 through time
1.0
0.8
0.6
0.4
ei_obs
0.25
0.50
0.75
1.00
aridity_index
Year
2020
2015
2010
2005
2000
2016
2022
2017
2020
2001
2015
2006
2004
2009
2002
2019


aridity_index
ei_obs
ratio
val
year
1.6
1.2
0.8
0.4
2000
2005
2010
2015
2020
2025


w5 through time
1.00
0.75
0.50
0.25
ei_obs
0.25
0.50
0.75
1.00
aridity_index
Year
2020
2015
2010
2005
2000
2016
2001
2019
2012
2009
2004
2002
2000


aridity_index
ei_obs
ratio
val
year
1.2
0.9
0.6
0.3
2000
2005
2010
2015
2020
2025


ER_RUS1
evaporative_index
aridity_index
1.0
0.8
0.6
0.4
0.50
0.75
1.00
1.25
1.50
Year
2020
2015
2010
2005
2000
2018
2002
2012
2021
2020
2016
2003
2022
2013
2001
2015
2000
2019
2004
2017
2009
2010
2011
2006
2008


aridity_index
evaporative_index
ratio
1.5
1.0
0.5
val
2000
2005
2010
2015
2020
2025
year