

Neural inspiratory time detection from surface diaphragm electromyography

L. Estrada^{1,2,3}, A. Torres^{1,2,3}, L. Sarlabous^{1,2,3}, R. Jané^{1,2,3}

¹ Institut de Bioenginyeria de Catalunya, The Barcelona Institute of Science and Technology, Barcelona, Spain

{lestrada, atorres, lsarlabous, rjane}@ibecbarcelona.eu

² Universitat Politècnica de Catalunya (UPC) – Barcelona Tech, Barcelona, Spain

³ Biomedical Research Networking Center in Bioengineering, Biomaterials and Nanomedicine (CIBER-BBN), Spain

1. Introduction

Electromyography is a technique for recording the electrical activity produced by respiratory muscles. The surface diaphragm electromyography (EMGdi) signal can be non-invasively recorded using electrodes in a bipolar configuration. The measurement of the EMGdi can be used to indirectly quantify the neural respiratory drive (NRD), which allows evaluating the level and pattern of the muscle activation [1]. Its use has contributed to improve patient-ventilation interaction in new modes of mechanical ventilation that assist patients in proportion to its respiratory efforts by evaluating the onset (n_{ton}) and offset (n_{toff}) of neural inspiratory time. Determining n_{ton} and n_{toff} can be challenging because muscle respiratory signals present a lower signal-to-noise ratio, when compared to conventional respiratory measures [2]. In this work, we present a novel method for determining the n_{ton} and n_{toff} of neural inspiratory time from EMGdi signals.

2. Materials and Methods

Ten healthy subjects carried out a respiratory protocol in which they breathe, progressively increasing their respiratory rate from 15 to 20, 24, 30 and 40 bpm every 16 respiratory cycles (Figure 1, left). To obtain the n_{ton} , a dynamic thresholding was established from the mode of the amplitude distribution of an EMGdi segment between two consecutive peaks by using the Gaussian kernel density estimation method. The determination of n_{toff} was considered as the time when the EMGdi signal has decreased to 70 % of its peak value (Figure 1, middle) [2]. To overcome the problem of the electrocardiographic interference, the fixed sample entropy (fSampEn), a robust technique against impulsive noise was used [1]. The n_{ton} and n_{toff} were compared to indirect estimates of neural onset and offset based on a zero-crossing criterion from respiratory airflow signal.






Figure 1. Left: Breathing protocol. Middle: Procedure for determining onset and offset time. Right: Comparison of onset and offset times detected in the airflow and EMGdi signals, in the respiratory protocol.

3. Results

The Bland–Altman analysis between the n_{ton} and n_{toff} detected in the airflow and EMGdi showed a global bias of 46 ms, with limits of agreement lower than 360 ms for onset and offset comparisons (Figure 1, right).

4. Conclusions

This study proposes a reliable method for estimating the neural inspiratory time (n_{ton} and n_{toff}), from surface EMGdi signals, which does not require filtering the cardiac interference preserving most of its spectral content.

Acknowledgments

This work was supported by the Spanish Ministry of Economy and Competitiveness through project DPI2015-68820R (MINECO/FEDER) and the CERCA Programme (Generalitat de Catalunya). L. Estrada was supported by the Instituto para la Formación y Aprovechamiento de Recursos Humanos and Secretaría Nacional de Ciencia, Tecnología e Innovación (IFARHU-SENACYT) from the Panama Government, under grant 270-2012-273.

References

- [1] Estrada L, Torres A, Sarlabous L, Jané R, Improvement in Neural Respiratory Drive Estimation from Diaphragm Electromyographic Signals using Fixed Sample Entropy. *IEEE Journal of Biomedical and Health Informatics*, vol 20, no 2, 2016, pp. 476–485 (ISSN: 2168-2194).
- [2] Estrada L, Torres A, Sarlabous L, Jané R. Onset and Offset Estimation of the Neural Inspiratory Time in Surface Diaphragm Electromyography: A Pilot Study in Healthy Subjects. *IEEE Journal of Biomedical and Health Informatics*, 2017. DOI: 10.1109/JBHI.2017.2672800 (ISSN: 2168-2194).