

# THE FRB 121102 CAMPAIGN: MULTI-OBSERVATORY RADIO BURST AND IMPLICATIONS FOR THE FRB POPULATION

C. J. LAW,<sup>1</sup> J. W. T. HESSELS,<sup>2,3</sup> R. S. WHARTON,<sup>4</sup> C. G. BASSA,<sup>2</sup> G. C. BOWER,<sup>5</sup> S. BURKE-SPOLAOR,<sup>6,7,8</sup>  
B. J. BUTLER,<sup>6</sup> T. CANTWELL,<sup>9</sup> S. H. CAREY,<sup>10</sup> S. CHATTERJEE,<sup>4</sup> J. M. CORDES,<sup>4</sup> P. DEMOREST,<sup>6</sup> R. FENDER,<sup>11</sup>  
K. GRAINGE,<sup>9</sup> J. HICKISH,<sup>12,10</sup> V. M. KASPI,<sup>13</sup> T. J. W. LAZIO,<sup>14</sup> M. A. MCCLAUGHLIN,<sup>7,8</sup> D. MICHILLI,<sup>2,3</sup> K. MOOLEY,<sup>11</sup>  
Y. C. PERROTT,<sup>10</sup> S. M. RANSOM,<sup>15</sup> N. RAZAVI-GHODS,<sup>10</sup> M. RUPEN,<sup>16</sup> A. SCAIFE,<sup>9</sup> P. SCOTT,<sup>10</sup> P. SCHOLZ,<sup>16</sup>  
A. SEYMOUR,<sup>17,18</sup> L. G. SPITLER,<sup>19</sup> S. P. TENDULKAR,<sup>13</sup> D. TITTERINGTON,<sup>10</sup> AND P. K. G. WILLIAMS<sup>20</sup>

<sup>1</sup>*Department of Astronomy and Radio Astronomy Lab, University of California, Berkeley, CA 94720, USA*

<sup>2</sup>*ASTRON, Netherlands Institute for Radio Astronomy, Postbus 2, 7990 AA, Dwingeloo, The Netherlands*

<sup>3</sup>*Anton Pannekoek Institute for Astronomy, University of Amsterdam, Science Park 904, 1098 XH Amsterdam, The Netherlands*

<sup>4</sup>*Cornell Center for Astrophysics and Planetary Science and Department of Astronomy, Cornell University, Ithaca, NY 14853, USA*

<sup>5</sup>*Academia Sinica Institute of Astronomy and Astrophysics, 645 N. A'ohoku Place, Hilo, HI 96720, USA*

<sup>6</sup>*National Radio Astronomy Observatory, Socorro, NM 87801, USA*

<sup>7</sup>*Department of Physics and Astronomy, West Virginia University, Morgantown, WV 26506, USA*

<sup>8</sup>*Center for Gravitational Waves and Cosmology, West Virginia University, Chestnut Ridge Research Building, Morgantown, WV 26505*

<sup>9</sup>*Jodrell Bank Centre for Astrophysics, Alan Turing Building, School of Physics & Astronomy, The University of Manchester, Oxford Road, Manchester M13 9PL, UK*

<sup>10</sup>*Astrophysics Group, Cavendish Laboratory, 19 J. J. Thomson Avenue, Cambridge CB3 0HE, UK*

<sup>11</sup>*Centre for Astrophysical Surveys, University of Oxford, Denys Wilkinson Building, Keble Road, Oxford OX1 3RH, UK*

<sup>12</sup>*Dept of Astronomy and Radio Astronomy Lab, University of California, Berkeley, CA 94720, USA*

<sup>13</sup>*Department of Physics and McGill Space Institute, McGill University, 3600 University St., Montreal, QC H3A 2T8, Canada*

<sup>14</sup>*Jet Propulsion Laboratory, California Institute of Technology, Pasadena, CA 91109, USA*

<sup>15</sup>*National Radio Astronomy Observatory, Charlottesville, VA 22903, USA*

<sup>16</sup>*National Research Council of Canada, Herzberg Astronomy and Astrophysics, Dominion Radio Astrophysical Observatory, P.O. Box 248, Penticton, BC V2A 6J9, Canada*

<sup>17</sup>*Arecibo Observatory, HC3 Box 53995, Arecibo, PR 00612, USA*

<sup>18</sup>*Max-Planck-Institut für Radioastronomie, Auf dem Hügel 69, Bonn, D-53121, Germany*

<sup>19</sup>*Max-Planck-Institut für Radioastronomie, Auf dem Hügel 69, D-53121 Bonn, Germany*

<sup>20</sup>*Harvard-Smithsonian Center for Astrophysics, Cambridge, MA, USA*

## ABSTRACT

The recent precision localization of FRB 121102 has helped identify a host galaxy, measure its distance, and establish it as a member of a truly new class of astrophysical source. We present results of the coordinated observing campaign that made this localization, including the first simultaneous detection of an FRB burst with multiple telescopes. Of the nine bursts detected by the Very Large Array at 3 GHz, four had simultaneous observing coverage at other observatories, and one was detected by the Arecibo Observatory at 1.4 GHz. We show that the burst spectra typically have a  $\sim 500$  MHz-scale envelope, plus fine spectral structure on MHz-frequency scales. We also calculate the energy distribution and temporal statistics for FRB 121102... \*population models\* \*slsn\* \*\*This source shows that optimal FRB survey strategies should...\*\*

## 1. INTRODUCTION

Fast Radio Bursts (FRBs) were discovered ten years ago as a millisecond-duration radio transient with an anomalously high dispersion measure (the “Lorimer burst”; Lorimer et al. 2007). Their large dispersion measure (DM) implied that they originated outside of our Galaxy, potentially at cosmological distances, and were orders of magnitude more luminous than any other millisecond radio transient (Thornton et al. 2013). Both their energetics and distance have inspired a wide variety of models and astrophysical applications (e.g., McQuinn 2014; Kulkarni et al. 2014; Connor et al. 2016b; Cordes & Wasserman 2016). However, that potential was limited by the lack of a definitive association of an FRB to an extragalactic host.

This paper is part of a series based on the first localization of an FRB and its unambiguous association to an extragalactic host (Chatterjee et al. 2017; Tendulkar et al. 2017; Marcote et al. 2017). FRB 121102, also known as the “repeating FRB”, was first detected in November 2012 by the Arecibo Observatory (Spitler et al. 2014). In mid 2015, new Arecibo observations revealed a series of bursts at the same DM ( $\sim 559$  pc  $\text{cm}^{-3}$ ) and sky position demonstrating that it repeats (Spitler et al. 2016). Beginning in August of 2015, we made the first of nine detections of FRB 121102 with the Very Large Array (Chatterjee et al. 2017) and localized it with a precision of  $0.1''$ . Subsequent radio and optical observing discovered more bursts from FRB 121102 and associated them with a persistent radio and optical source at a redshift of 0.193 with a precision of  $0.01''$  (Tendulkar et al. 2017; Marcote et al. 2017,  $\sim 40$  pc).

FRB 121102 has now been localized four orders of magnitude better than any other FRB and placed at a cosmological distance. Its lookback and luminosity distances are 746 and 972 Mpc, assuming a concordance cosmology with parameters given by Planck Collaboration et al. (2016). This association means that FRB 121102 is orders of magnitude more luminous than any other millisecond transient and that its DM traces both the host and intergalactic medium (Tendulkar et al. 2017). If FRB 121102 is representative of all FRBs, then we should expect them to be useful as probes of the intergalactic medium and their host galaxies. Thus, the confirmation of a cosmological distance for FRB 121102 has begun to fulfil the promise implied by the Lorimer burst.

Although new models of FRB origin have been already been developed (Kashiyama & Murase 2017; Metzger et al. 2017; Zhang 2017; Dokuchaev & Eroshenko 2017), we still do not know what causes them. *\*\*magnetar birth?* Also, it has not been demonstrated that FRB


121102 is representative of the overall FRB population. In fact, the repetition of its bursts is unique among all FRBs (Petroff et al. 2015), so it is natural to ask whether FRB 121102 is representative. An important first step is to demonstrate that the properties of FRB 121102 are consistent with the significant body of facts for the overall population (Macquart & Johnston 2015; Katz 2016). The repeating nature of FRB 121102 provides us with a sample of bursts covering a range of spectral properties, temporal properties, and luminosities that can be used to test its connection to the larger FRB population.

We can also assume that FRB 121102 is representative of all FRBs and use it to constrain the physical processes at play in the overall FRB population. In this case, we know that FRBs are apparently luminous, but if that is due to intrinsic processes, such as coherent, pulsar-like emission (Katz 2014; Luan & Goldreich 2014; Cordes & Wasserman 2016), or extrinsic effects like scintillation (Cordes et al. 2017). The paper presents an analysis of the spectral properties of VLA bursts implied by simultaneous observing at Effelsberg, GBT, AMI-LA, and Arecibo, including the first simultaneous detection of an FRB at two observatories. The repetition of FRB 121102 also has strong implications for calculations of their rate of occurrence (Connor et al. 2016a) and comparison to other classes of transient, such as superluminous supernovae (Tendulkar et al. 2017).

This paper opens by describing the multi-telescope observing campaign and a refined analysis of the nine VLA bursts. Next, we present the spectrum of a burst from FRB 121102 that was simultaneously detected by the VLA and Arecibo observatories. *\*\*Simulations of temporal statistics and luminosity distribution... \*\** *\*\*New estimate of FRB rate and comparison to other transient populations\*\** We close with a discussion of how the FRB 121102 burst spectra, host properties, burst rate estimates, and other properties inform new strategies for finding FRBs.

## 2. OBSERVATIONS

The data presented here were obtained from multiple programs and telescopes, but focuses on an observing campaign in August/September 2016. The central goal was to interferometrically localize FRB 121102 with the VLA in coordination with Arecibo Observatory, with others included on a best-effort basis. We coordinated observing between the VLA, Arecibo, Effelsberg, and AMI-LA telescopes, as shown in Figure 1. Below, we summarize these observations, with a focus on those conducted simultaneously with VLA burst detections from FRB 121102.



**Figure 1.** The top and bottom panels summarize the observing coverage and detections of FRB 121102 in the 2015 and 2016 campaigns. Symbols show days with observations with letters L, S, C, K referring to radio frequencies of 1.4, 3, 5, and 12 GHz. Circles highlight observations that detected bursts from FRB 121102. Multiple circles indicate multiple burst detections, except for Arecibo, which typically has multiple detections per observing session. The black dashed lines show the VLA burst detections with simultaneous coverage at other telescopes. The solid black line shows the simultaneous burst detection at VLA and Arecibo.

### 2.1. VLA

The FRB 121102 observing campaign started in late 2015 with a 10 hr campaign observed at 1.4 GHz in the compact D configuration. In April through May 2016, we conducted a 40 hr campaign at 3 GHz in the C and CnB configurations in coordination with Arecibo. We concluded with a 40 hr, coordinated campaign from August through September 2016 in the B configuration and during the move to the most extended A configuration. In the late-2016 campaign, the first 34 hours of VLA observations were made at 3 GHz, while the last 6 hours were observed at 6 GHz. This paper focuses on the data collected at 3 GHz, which includes all nine burst detections and is the widest bandwidth to have ever detected an FRB.

All VLA fast-sampled data were observed with 5 ms sampling, 256 channels, and dual-circular polarization (Law et al. 2015). The channel frequency width was set to maximize sensitivity to the known DM of FRB 121102 with the largest total bandwidth. The total bandwidth at L (1.4 GHz), S (3 GHz), and C (6 GHz) bands was 256 MHz, 1024 MHz, and 2048 MHz, respec-

tively. The 3 GHz data were recorded data in eight spectral windows with 32 channels each.

Observations in August and September were searched by a prototype version of *realfast*<sup>1</sup>. *realfast* is a real-time, fast imaging transient search system. The current prototype runs on CPU-based hardware that is normally dedicated to the VLA correlator; for this experiment, it runs the transient search pipeline software called *rtpipe* (<https://github.com/caseyjlaw/rtpipe>; Law et al. 2015). Images were formed for each integration with a DM grid of 0, 546, 556.9, 560, and 565  $\text{pc cm}^{-3}$  and a time resampling grid of 5, 10, 20, 40, and 80 ms. This DM grid was chosen to maintain 90% sensitivity to the nominal DM range of 540–570  $\text{pc cm}^{-3}$ , which includes the known FRB 121102 DM of 559  $\text{pc cm}^{-3}$  (Scholz et al. 2016). Gain calibration was made from observations of J0555+3948 by the “telcal” system, which uses phase-only calibration. A flux scale is calculated for each

<sup>1</sup> See <http://realfast.io>.

spectral window from an observation on **xx (Bryan)** and applied to all burst spectra.

Burst detections and localizations were made within hours of data being recorded. The transient search starts when data are recorded and proceeds slower than real-time, so we refer to it as “quasi real-time”. For each trial DM, integration, and time scale, we form an image and calculate the S/N ratio for the peak pixel in the dirty image. We empirically identified S/N thresholds of 6.4 and 7.4 as useful to capture data quality statistics and candidates for inspection, respectively. The higher threshold is relatively unlikely to be triggered by thermal noise in this configuration, so *realfast* generates a more detailed (and computationally intensive) candidate visualization that includes an image and spectrum. All visibilities are recorded so detailed analysis, including improved calibration and localization, can be conducted offline.

Computational (Jupyter) notebooks to reproduce the transient detection, localization, and analysis presented here can be found at <https://github.com/caseyjlaw/FRB121102>. Time cut-out visibility data and calibration products are available at <https://doi.org/10.7910/DVN/TLDKXG>. Original visibility data are available under VLA program codes 16A-459 and 16A-496 and can be downloaded at <http://archive.nrao.edu>.

## 2.2. Arecibo

During the late-2016 campaign, Arecibo observed with the L-wide receiver using the PUPPI pulsar backend. The observational frequency range was 1.15 to 1.73 GHz and frequency resolution was 1.5625 MHz. We recorded total full Stokes polarization intensity spectra with time at a resolution of 10.24  $\mu$ s. Each frequency channel was coherently dedispersed to 557 pc cm<sup>-3</sup>, thereby eliminating intra-channel dispersion smearing. The full width at half maximum (FWHM) beam size at band center is 3.3’.

In total, twelve Arecibo observations had some simultaneous coverage with the VLA. Four of those observations were simultaneous with bursts detected with the VLA and one of those observations detected the same VLA burst. During the first VLA burst with Arecibo coverage (MJD 57643), the PUPPI 1.4 GHz recording system failed so data were recorded at C band. No detection was made in those Arecibo data. Overall, there were many more bursts detected at Arecibo than with the VLA and a more detailed analysis of those bursts will be presented in a future paper.

## 2.3. Effelsberg

Effelsberg observations were conducted with the S60mm receiver at an observing frequency of 4.6 to

5.1 GHz. Total intensity spectra were recorded by the PFFTS backend in pulsar search mode with a time resolution of 65.5  $\mu$ s and 128 frequency channels. The system equivalent flux density is 18 Jy and a FWHM beam size of 2.4’ at 4.85 GHz.

Five Effelsberg observations had some simultaneous coverage with the VLA, of which two were simultaneous with VLA bursts. Unfortunately, due to a configuration error, a 100 MHz bandwidth filter centered at 4.85 GHz was in place for both of these sessions. The sensitivity was about two times worse than the nominal value. No burst was detected in either observation.

## 2.4. AMI-LA

We observed FRB 121102 with the Arcminute MicroKelvin Imager Large Array (AMI-LA; Zwart et al. 2008) for 3 hours each on four epochs starting at MJDs 57643, 57645, 57648, and 57649. Observations were made with the new digital correlator having 4096 channels across a 5 GHz bandwidth between 13–18 GHz with a 1 s integration time. The phase calibrator, J0518+3306, was observed every 12 minutes for about 1.5 minutes. The AMI-LA data were binned to eight 0.625 GHz channels and processed (RFI excision and calibration) with a fully-automated pipeline, AMI-REDUCE (e.g., Perrott et al. 2013). Daily measurements of 3C48 and 3C286 were used for the absolute flux calibration, which is good to about 10%.

We inspected the calibrated visibilities, and did not find any signal above 30 mJy in the 1s samples at and in the vicinity of the detected bursts. Concatenating and imaging the 12 hours of calibrated data with the CASA tasks *concat* and *clean* also does not yield any significant detection at the FRB location. Although the statistical  $3\sigma$  upper limit is 60  $\mu$ Jy, extended mJy-level sources in the field cause sidelobe confusion (the AMI-LA angular resolution is  $\sim 30''$ ), and the actual upper limit is larger. We introduced artificial point sources at the FRB location using the CASA *sm* tool, and found that these sources can be recovered as long as their peak flux densities are more than  $\sim 100\mu$ Jy. Hence, we place an upper limit of  $100 \pm 10\mu$ Jy on any quiescent or possible radio flaring on  $\sim$ days timescales from the FRB. This limit is similar to the flux density measured by the VLA (Chatterjee et al. 2017).

## 3. RESULTS

### 3.1. Multi-Observatory Burst Spectrum

Four of the VLA bursts were observed simultaneously with Arecibo and AMI-LA, while two were observed simultaneously with all four observatories (Figure 1). Of these bursts, only one VLA burst (on MJD

57648) was simultaneously detected by another observatory (Arecibo). The VLA and Arecibo burst detections had significances of  $25\sigma$  and  $39\sigma$  with integrated flux densities of 111 mJy and 14 mJy, respectively. Assuming a powerlaw flux density model ( $S_\nu \propto \nu^\alpha$ ), we find a spectral index  $\alpha \approx 2.7$ .

Figure 2 shows a dynamic spectrum for the 57648 burst with the VLA and AO data aligned in time. The temporal resolution of the VLA data (5 ms) is much coarser than that of Arecibo, so data were resampled to the same temporal grid after correcting for barycentric and dispersion measure offsets. The dispersion correction for both data sets assumed a DM of  $560.5 \text{ pc cm}^{-3}$ , which was typical of the source during the observing campaign (Hessels et al. 2017). The regridded dynamic spectrum has an apparent DM error that is evident both as a frequency-dependent time drift within the Arecibo observation and as an offset between the Arecibo and VLA bursts. Both of these drifts are consistent with an apparent DM of  $\sim 565 \text{ pc cm}^{-3}$ . This is consistent with modeling of the VLA spectrum described in §3.2.1.

Three of the four VLA bursts with simultaneous observing coverage by Arecibo were not detected at Arecibo. Numerous bursts were also detected by Arecibo during VLA observing, but were not detected by the VLA at 3 GHz. Considering this, and the order of magnitude difference in flux density for the one simultaneous burst detection, we conclude that the 3 GHz burst flux is not correlated with that at 1.4 GHz.

### 3.2. VLA Bursts

#### 3.2.1. Spectral Modeling

This paper refines the analysis of the nine VLA radio burst spectra described in Chatterjee et al. (2017) in a few ways. We use a better calibration scheme and have optimized the detection significance over a fine grid of DM ( $\Delta\text{DM} = 1 \text{ pc cm}^{-3}$ ). After calibration and flagging, the visibility phases were rotated to the best-fit location ((RA, Dec) = (05h31m58.70s, +33d08m52.5s); Chatterjee et al. 2017) to extract a Stokes I spectrum that maximizes the image S/N for each burst.

Table 1 summarizes the properties of all nine VLA bursts. Parameters such as integrated flux density, S/N, and Stokes V were measured from the burst properties integrated in frequency. Stokes V values estimated from the dual-circular feeds ( $(RR-LL)/(RR+LL) \approx 3\%$  and are comparable to the systematic error expected from the primary beam (Perley et al 2016, VLA memo). Given that systematic effects dominate the apparent circular polarization, we conclude that the true fractional circular polarization is less than 3%.


Figure 3 shows that the Stokes I spectra are generally characterized by a broad, Gaussian shape with inter-channel modulation as high as 100%. To better understand the spectra, we extracted dynamic radio spectra (time versus frequency) for 2d modeling with the MCMC sampler *emcee* (Foreman-Mackey et al. 2013). MCMC sampling requires a generative model for the data, so we wrote functions in Python to create 2d dynamic spectra. The dynamic spectra are generated in the spectral domain as a Gaussian envelope and in the temporal domain by a dispersive delay that scales as frequency squared. We further allow the pulses to have a finite temporal width. This brings the total number of parameters to 6 (Gaussian amplitude, center, width, start time, time width, and DM).

The 6-dimensional models were sampled with 100 walkers taking 500 steps. Typically, the first 200 steps were used to burn in the walkers and were ignored. A flat prior was used over all ranges with valid data. An additional requirement was that the integrated burst signal must have a detection significance higher than  $8\sigma$ . The inter-channel modulation (scintillation) was ignored in the model. The measured, off-burst noise value was typically 70 mJy per 5 ms integration and 4 MHz channel.

The parameters for the best model of the dynamic spectrum are given in Table 1 and the resulting Gaussian model overlaid on Figure 3. The typical burst has a spectral width of 500 MHz. All but two of the best-fit Gaussians are centered inside the 3 GHz band and some appear contained by the 1 GHz wide band. This is consistent with previous detections of FRB 121102 by Arecibo, which showed quasi-broadband structure (Scholz et al. 2016) and large variation in the implied spectral index (Spitler et al. 2014).

This analysis differs from traditional analysis in that it simultaneously models the spectral and temporal evolution of the burst. Since the burst crosses many time-frequency pixel boundaries, the temporal modeling is sensitive to temporal properties smaller than the 5 ms integration time. Table 1 shows that the typical burst width is measured as  $2 \pm 1 \text{ ms}$  (95% confidence interval); the brightest burst (57633.68) has a temporal width of  $2.05 \pm 0.02 \text{ ms}$ . The burst arrival times are typically measured with a precision of 1 ms; the brightest burst arrival time is measured with a precision of  $\sim 50 \mu\text{s}$ . Note that these errors are only accurate to the degree that the model represents the data. One potential bias in the model is that we do not model scintillation effects.

While most bursts are modeled with well-defined parameter probability distributions, one burst is an outlier. Figure 4 shows the scatterplot matrix for the MCMC run on burst 57646 (Foreman-Mackey 2016). Two clus-



**Figure 2.** Composite dynamic spectrum for the burst from FRB 121102 on MJD 57648 with data from the VLA and Arecibo observatories. Both data sets have been barycentered and corrected for an assumed dispersion measure of  $560.5 \text{ pc cm}^{-3}$ . The black line separates data from the VLA (2.5 to 3.5 GHz) and Arecibo (1.0 to 1.8 GHz). The amplitude scale is normalized by assuming the Arecibo to VLA gain ratio is 10.

**Table 1.** Properties of Bursts from FRB 121102

Date (MJD)	Image S/N	$S_{I,\text{int}}$ (mJy)	$S_V$ (mJy)	$DM_{\text{opt}}$ ( $\text{pc cm}^{-3}$ )	$dt$ (ms)	$S_{I,\text{peak}}$ (mJy)	Center (GHz)	FWHM (MHz)	$E_{\text{int}}$ ( $10^{38} \text{ erg}$ )
57623.74402686	38	194	+3	$567 \pm 2$	$2.0 \pm 0.2$	690	2.8	290	11
57633.67986367	179	1500	-35	$568.2 \pm 0.2$	$2.05 \pm 0.02$	3340	3.2	510	98
57633.69515938 <sup>a</sup>	15	69	+2	$562^{+4}_{-6}$	$2.5^{+0.9}_{-0.6}$	>430	<2.5	290	7
57638.49937435	12	55	+5	$567^{+7}_{-9}$	$1.3^{+1.4}_{-0.8}$	130	3.1	420	3
57643.45730263	100	508	-5	$565.6^{+0.6}_{-0.5}$	$1.9 \pm 0.1$	1170	2.8	510	34
57645.42958602	13	64	+3	$563^{+5}_{-4}$	$1.0 \pm 0.7$	170	2.8	380	4
57646.46600650 <sup>a</sup>	20	87	+1	$569 \pm 5$	$2.7^{+0.9}_{-1.4}$	>420	<2.5	430	10
57648.43691490 <sup>b</sup>	25	111	+9	$564 \pm 2$	$1.4^{+0.3}_{-0.4}$	260	2.8	470	7
57649.45175697	36	167	+1	$567 \pm 2$	$2.1 \pm 0.5$	290	3.0	690	12

<sup>a</sup>Best-fit Gaussian is not centered in 3 GHz band, so spectral parameters are limits.

<sup>b</sup>Detected simultaneously with Arecibo between 1.15 and 1.73 GHz.


ters of samples are identified for this burst: one narrow, low-DM and one wide, high-DM. The model seems to be too simple for this burst. This is consistent with high-resolution Arecibo burst spectra (Hessels et al. 2017), which show that bursts are composed of "sub-bursts" that drift in frequency.

### 3.2.2. Dispersion


The initial VLA detections were made with a matched-filter approach that allowed inter-DM sensitivity losses up to roughly 10% ( $\Delta DM = 10 \text{ pc cm}^{-3}$

Cordes & McLaughlin 2003). The MCMC modeling is more sophisticated in that it considers a wider range of parameters and models both data and observing effects (frequency-time pixel boundaries) simultaneously.

The top panel of Figure 5 shows the 95% confidence intervals on the DM for all nine VLA bursts as a function of burst time. The burst DMs are not consistent with any single value and tend to be larger than the long-term average of  $560.5 \text{ pc cm}^{-3}$ . The range and variance in DM observed by the VLA is similar to that reported for Arecibo observations of FRB 121102 in Scholz et al.



**Figure 3.** Spectra and best-fit model for nine bursts seen by the VLA from 2.5 to 3.5 GHz. Starting at the top left (moving right and then down), they correspond to bursts on MJDs 57623, 57633.68, 57633.70, 57638, 57643, 57645, 57646, 57648, and 57649. The solid line is a best-fit Gaussian model found through MCMC modeling. Note that bursts are detected in 5 ms images generated from dedispersed visibilities.



**Figure 4.** A scatterplot matrix that shows the correlation of every pair of parameters in the MCMC run for burst 57646. The equivalent plot for the other bursts show normally-distributed samples, but this burst shows significant structure in the samples.

(2016). As noted before, this is consistent with the idea that the DM measured at low resolution (e.g., by the VLA) is defined by true (extrinsic) dispersion and some spectral structure that changes between bursts. The VLA data demonstrate that the burst-dependent DM

effect is visible at frequencies near 3 GHz. Assuming a “true” DM of  $560.5 \text{ pc cm}^{-3}$ , the excess DM can be thought of as a frequency-dependent delay that varies between bursts with a delay rate up to  $2 \text{ ms GHz}^{-1}$ .


The bottom panel of Figure 5 compares the DM to the modeled temporal width of the bursts. There is a weak correlation between burst width and apparent DM. A change in width of  $\sim 1 \text{ ms}$  correlates with a change in apparent DM of approximately  $5 \text{ pc cm}^{-3}$ .

### 3.3. Temporal Statistics

The burst rate for FRB 121102 varied dramatically throughout the 2016 observing campaign (Figure ?). In the early-2016 campaign, we observed for 30 hours at S-band and no bursts were detected. In the late-2016 observing campaign, nine bursts were detected in the last 27 hours of S-band observing. Overall, the data quality is uniform and high, so the inhomogeneous burst distribution shows that the burst detection probability was not stationary. Assuming that the burst detection probability follows a Poisson distribution, the nondetection in the first half of S-band limits the FRB rate to  $R < 0.1 \text{ hour}^{-1}$  (95% confidence limit). The S-band burst rate was much higher during the late-2016 campaign,  $R = 0.33 \pm 0.1 \text{ hour}^{-1}$ .

### 3.4. Energy and Brightness Distribution

Knowing the source distance, we calculate the burst energy by integrating flux in time and frequency (Table



**Figure 5.** (Top) DM as a function of time for nine 3 GHz bursts detected by the VLA. The 95% confidence interval on DM is shown with a bar and the dashed line shows the best-fit  $DM = 560.5 \text{ pc cm}^{-3}$  inferred for Arecibo observations at 1.4 GHz. (Bottom) Burst DM versus temporal width for all nine VLA bursts.


1). Some VLA burst spectra seem to be contained by the 2.5 to 3.5 GHz band and most of them seem to have Gaussian envelopes that are well modeled by the emission within that band. Assuming that the Gaussian shape defines the emission window, the mean S-band flux density can be converted to a total energy with no further assumptions about their spectral properties.

Figure 6 shows the FRB 121102 burst energy cumulative distribution as seen by the VLA and calculated from prior observations by Arecibo and the GBT (Spitler et al. 2016; Scholz et al. 2016). The latter two energy distributions are scaled from the fluence by assuming that all burst energy is included by the observation. This likely underestimates the burst energy for some bursts by a factor of a few, but should not affect the slope of the distribution. The Arecibo and GBT energies were scaled by factors of 2 and 1 (help), respectively, to correct for the sensitivity loss due to the primary beam.

Broadly, the range of energies shown in Figure 6 is defined by the sensitivity of the telescope and the duration

of the observing campaign. The VLA distribution shown represents the 27-hour, late-2016 campaign, which is longer than the single-dish campaigns presented here. However, since the VLA observations have a relatively coarse time resolution, it is not as sensitive as Arecibo and the GBT, so no bursts were detected weaker than  $3 \times 10^{38} \text{ erg}$ . The rate upper limit (95% confidence) from the early-2016 VLA campaign shows that even identical observing campaigns have different detection rates.

We modeled the differential energy distribution,  $dN/d\log E$ , using a Poisson probability distribution with a rate function  $\lambda = AE^\alpha$ . Rather than trying to estimate a completeness limit for each energy distribution, we assume an effective detection limit of 0.9 times the weakest burst detected; the best-fit slope is weakly dependent on this, but general conclusions are robust. We directly sampled the likelihood distribution to estimate a best slope of  $\alpha_{\text{VLA}} = -0.7^{+0.3}_{-0.4}$ ,  $\alpha_{\text{AO}} = -0.8^{+0.3}_{-0.5}$ , and  $\alpha_{\text{GBT}} = -0.8^{+0.4}_{-0.5}$  (68% confidence interval). Expressed as a powerlaw function in  $dN/dE$ , the slope index is approximately  $-1.7$ .



**Figure 6.** The cumulative burst energy distributions for the VLA, Arecibo, and GBT are shown with dots connected by solid, dashed, and dotted lines, respectively. The Arecibo distribution is derived from the 11 1.4 GHz bursts reported in Spitler et al. (2016) and the GBT distribution is derived from the 5 1.4 GHz bursts reported in Scholz et al. (2016). An upper limit from the VLA nondetection in early 2016 is shown as a triangle.

## 4. DISCUSSION

### 4.1. Burst Spectra

We present the first simultaneous detection of an FRB with multiple telescopes and over frequencies from 1.2 to 3.5 GHz. The Arecibo burst flux density is an order of magnitude less than that seen by the VLA. At the same time, three other bursts from FRB 121102 had similar observing coverage but were not detected simultaneously. This is consistent with the spectral struc-



ture observed within the 3 GHz VLA band, which is typically limited to a Gaussian envelope with width of roughly 500 MHz. The center frequency of this envelope is a unique property of each burst and does not show an obvious pattern between bursts.

If FRB 121102 burst spectra are typical of the larger FRB population, then population models need to be generalized beyond the assumption of a spectral powerlaw. Most obviously, FRB 121102 implies that future multi-telescope searches for bursts are unlikely to simultaneously detect bursts in different bands. Second, since burst spectra have limited bands, the burst detection rate at one frequency may not represent that at another. That will require rate estimates for the FRB population to be made explicitly frequency-dependent. Finally, we note that the odds of detecting a burst will improve with bandwidth for bandwidths wider than their characteristic scale.

After correcting for barycentric and typical dispersion delays, the VLA/Arecibo burst spectrum shows a residual temporal drift. This drift can be interpreted as an excess DM of roughly  $5 \text{ pc cm}^{-3}$ . We also measure a burst-to-burst variation in DM within the whole sample of VLA 3 GHz bursts of a similar scale. This variation had been noted in earlier 1.4 GHz Arecibo observations of FRB 121102 (Scholz et al. 2016), but the VLA bursts demonstrate that this "excess", burst-specific DM is seen up to 3 GHz. The simultaneous VLA/Arecibo burst also shows that an individual burst's properties from 1.2 to 3.5 GHz are consistent with a single excess.

These features are generally consistent with intrinsic burst properties described in Hessels et al. (2017). That work uses bright, coherently-de-dispersed bursts to identify "sub-bursts" that preferentially drift to lower frequencies at later times. With high-resolution dynamic spectra, one can define a heuristic that distinguishes between a burst-specific dispersion-like effect and a more stable ("true") dispersion. Since the sub-bursts tend to drift to lower frequencies, they systematically bias DM measurements made with low-resolution data (like the VLA and earlier Arecibo and GBT observations). Interestingly, our modeling of the VLA burst dynamic spectra show a weak correlation between larger apparent DM and pulse width. This could be a signature of blending of many unresolved sub-bursts in a single VLA burst spectrum.

The burst-specific DM structure could be intrinsic to the emission mechanism or induced during propagation. Radio waves can be modified in a variety of ways (e.g., scintillation, scattering) and the duration of the emission in the source frame is not known (Cordes et al. 2016). Cordes et al. (2017) describe a model where plasma lens-

ing near the source of an FRB can magnify radio emission by orders of magnitude. That model argues that the simultaneous appearance of a burst from 1.2 to 3.5 GHz requires a lens with the focal frequency be  $\gtrsim 3.5 \text{ GHz}$ . This gives a constraint on the combination

$$\frac{DM_0}{a_{\text{AU}}^2} \frac{d_{sl} d_{lo}/d_{so}}{1 \text{ kpc}} \gtrsim (3.5/39.1)^2$$

where  $DM_0$ ,  $a$ , and  $d_*$  are the electron column density of the lens, size of the lens, and lensing distances, respectively. A lens placed at  $d_{sl} \approx 1 \text{ kpc}$  is consistent with typical properties of the interstellar medium. Lenses that are influenced by the source would be much nearer to the source  $d \ll 1 \text{ kpc}$ , requiring large values of  $\frac{DM_0}{a_{\text{AU}}^2}$ .

#### 4.2. Energy and Flux Distribution

The refined analysis shows that FRB 121102 has even higher isotropic energies than previously reported, reaching energies  $E_{\text{max}} \approx 10^{40} \text{ erg}$ . FRB 121102 has isotropic energies orders of magnitude higher than the strongest Crab pulsar pulses ( $\sim 10^{35} \text{ erg}$ ) and comparable brightness temperatures ( $T_b^{\text{Crab}} \sim 10^{41} \text{ K}$  versus  $T_b^{\text{FRB121102}} \sim 10^{38} \text{ K}$ ; Hankins et al. 2003; Katz 2014).

In §3.4 we compare the energy distribution of the 9 VLA bursts to those reported earlier by Arecibo and the GBT. All three energy distributions can be characterized as a powerlaw in  $dN/dE$  with slope  $\sim -1.7$ . This slope is seen even though the burst rate varies by almost an order of magnitude between observations and frequencies cover both 1.4 and 3 GHz bands. This suggests that the slope is related to the underlying physical process, rather than the state at any given time. The value of the energy distribution slope is similar to that observed in gamma-ray fluence of bursts from soft-gamma repeaters (Göğüş et al. 2000; Scholz & Kaspi 2011). This similarity has been used to argue that bursts are driven by self-regulated critical phenomena (slope =  $-5/3$ ; Aschwanden 2011).

While the surprising brightness of new FRB discoveries was once seen as a cause for concern (i.e., a sign of terrestrial interference), we may be able to use this to infer properties of the overall population (Lorimer et al. 2007; Ravi et al. 2016). When modeling the flux distribution as a powerlaw, an isotropic distribution of sources will have a powerlaw index,  $\alpha$ , of  $-1.5$  (the "Euclidean distribution"; Lyutikov et al. 2016). Prior to the measurement of the first GRB redshift, deviations from the Euclidean distribution were used to infer their cosmological distribution (e.g., the  $V/V_{\text{max}}$  test; Mao & Paczynski 1992; Fenimore & Bloom 1995). Multiple studies have suggested that FRBs have a "flat" flux distribution ( $-0.5 < \alpha < -0.9$ ; Vedantham et al. 2016;

Li et al. 2016; Lawrence et al. 2016). Others have noted that the flux distribution is not well modeled as a power-law, such that  $\alpha$  is effectively sensitivity (and telescope) dependent (Oppermann et al. 2016; CHIME Scientific Collaboration et al. 2017).

We now know that FRB 121102 is detectable with the VLA out to  $z = 0.7$ , which suggests that cosmological effects are even more important for understanding properties of the FRB population. CHIME Scientific Collaboration et al. (2017) demonstrate how redshift space distortions, time dilation, and spectral index ("k-correction") effects can flatten the flux distribution. To this complex scene, we add the fact that FRB 121102 burst spectra are not defined by a spectral index, which suggests that k-corrections are likely to be very difficult to calculate in practice. Rather than using wide band burst spectra to measure a single "typical" spectral index, we now must measure the burst rate as a function of frequency. These need not be measured simultaneously, but rather contemporaneously to insure they represent a similar state of activity of the FRB.

If we assume FRB 121102 is representative of the FRB population, we can estimate this frequency-dependent burst rate. The VLA and Arecibo observed intensively in a coordinated fashion during the late-2016 campaign. A complete search for bursts from the Arecibo is ongoing, but early results suggest that the burst rate at a fixed energy is much higher at 1.4 GHz than at 3 GHz. This suggests that high-redshift FRBs will have an apparently lower rate that flattens the  $dN/dF$  distribution in a manner similar to a k-correction. We can only infer the presence of this effect for observations near 1.4 GHz, since we have no estimate of the FRB 121102 burst rate outside the frequency range from 1.4 to 3 GHz. Propagation effects suggest that the rate will be lower at sub-GHz frequencies (Rajwade & Lorimer 2017). If so, the tension between constraints on  $\alpha$  near 800 MHz and 1.4 GHz may be resolved by a frequency-dependent FRB rate that peaks near 1.4 GHz.

#### 4.3. Repetition

The burst rate analysis presented in §3.3 shows that the detection probability of FRB 121102 is not stationary. As mentioned above, this could be caused by intrinsic events (e.g., magnetar outburst) or propagation effects. While a detailed analysis of temporal correlation will require more bursts and/or more sophistication, it is clear that FRB 121102 has significant correlation on short timescales (sometimes called a "red spectrum"; Connor et al. 2016a). Another way of stating this is that the detection of a burst improves odds of finding more bursts in near-future observations.

If this statistical property describes other FRBs, it weakens previous constraints on repetition (Petroff et al. 2015; Law et al. 2015). It is possible that other FRBs have repeating bursts, although calculating a limit requires knowing the temporal correlation function that defines the repetition. Repetition also implies that there are fewer physical sources of FRBs than implied by the burst detection rate.

FRBs like SGRs? 2016ApJ...826..226K

#### 4.4. FRB Rate

**Sarah:** can you update this as a per-SFR rate? Assuming that FRB 121102 is from the same population as the other FRBs, the measurement of a distance for the host of FRB 121102 allows us to re-evaluate the cosmic volume and event rate per galaxy for the FRB population. The estimated projected FRB rate  $R_p$  understates the true rate by a beaming fraction  $\Omega_b$  ( $\sim 10\%$ ; Tauris & Manchester 1998). For a comoving volume  $V(z)$  and galaxy number density  $\Phi(M)$ , the rate per galaxy is  $R_{FRB} = R_p / (\Omega_b \Phi(M) V(z))$ .

The first such rate estimate was made by Thornton et al. (2013), who used the measured DM to estimate a characteristic distance for their sample of 4 FRBs. They assumed that all of the extragalactic component of the DM was caused by the IGM and scaled as  $DM \approx z \times 1200 \text{ pc cm}^{-3}$  (Ioka 2003; Inoue 2004). They also calculated the number of galaxies by assuming a characteristic  $L_*$  galaxy (corresponding to stellar mass  $M_* \approx 10^{11} M_\odot$ ; Baldry et al. 2012).

**\*\*add error bar\*\*** Our calculation differs in that we have better estimates of all three parameters. First, the projected FRB rate is now believed to be closer to  $2 \times 10^3 \text{ sky}^{-1} \text{ day}^{-1}$  at high Galactic latitudes and flux densities brighter than 1 Jy ms (Lawrence et al. 2016; Champion et al. 2016; Rane et al. 2016). FRB 121102 is associated with a relatively small galaxy ( $5 \times 10^7 M_\odot$ ), which are roughly a factor of 100 more abundant ( $\Phi(M) \approx 10^{-2} \text{ Mpc}^{-3}$ ; Faber et al. 2007). Finally, the measured distance for FRB 121102 suggests that the extragalactic DM has comparable contributions from the IGM and host galaxy.

This reduces the characteristic distance by a factor of two and the volume by an order of magnitude. Considering these factors, we assume a characteristic redshift of the population that is twice that of FRB 121102 to estimate  $R_{FRB} \approx 10^{-4} (0.1/\Omega_b) \text{ galaxy}^{-1} \text{ year}^{-1}$ , which is two orders of magnitude lower than previously estimated (Thornton et al. 2013, assuming isotropic radiation;).

There are significant caveats to the comparison of this rate to the rates of other classes of transient. This isotropic FRB rate assumes a projected FRB rate at

a single (observationally defined) flux limit and that no bursts repeat. Using this calculation with data from more sensitive telescopes, we might infer a higher rate. However, if we assume that FRB 121102 is representative of the overall FRB population, then more sensitive observations would be more likely to find repeating FRBs that would effectively depress the estimate of the underlying isotropic rate per source. In this model, these two effects partially cancel, although without a physical model it is difficult to say how well.

#### 4.5. Observing Strategies

As discussed above, Connor et al. (2016a) note that the FRB repetition implies that the number of FRB-generating sources is smaller than implied by the number of bursts. In some scenarios, this means that there are areas on the sky that may contain no FRBs. Wide, shallow surveys are the preferred strategy for blind detection of FRBs. However, future efforts to localize FRBs are still most likely to succeed by targeting known FRBs to catch repetitions, assuming that all FRBs repeat.

### 5. CONCLUSIONS

Prior to being established as a cosmological source, the low Galactic latitude of FRB 121102 made it a somewhat compromised member of the FRB class. With its cosmological distance firmly established, FRB 121102 now serves as a new kind of standard by which FRBs are defined. That fact, combined with an abundance of data collected during its high activity state, allows us to think more generally about the physics of broader FRB population.

We presented the first multi-telescope detection (Arecibo and VLA) of an FRB. By detecting this burst from 1.2 to 4.5 GHz, we have demonstrated that some bursts have broad spectral structure. However, three other VLA bursts are undetected by Arecibo. That, combined with the measurement of spectra within the VLA band, suggest that the typical burst has a characteristic Gaussian spectral shape with width  $\sim 500$  MHz.

#### \*\*Flux/lum distribution\*\*

Analysis of the VLA burst times shows that the burst rate is highly variable. This shows that past constraints on FRB repetition are weaker than previously inferred. The combination of relatively narrow spectral structure, flat energy and flux distributions, and variable burst rate suggests that repeated observations, wide bandwidth, and large instantaneous field of view all improve the chance of detection.

New, arcsecond-scale localizations will continue to be highly informative, given that FRB hosts may be faint and can confuse the inference of a properties of the intergalactic medium (Deng & Zhang 2014). FRB 121102 was localized within hours by a prototype version of *realfast*, but an expanded *realfast* system is now under construction. This platform will search a TB/hour data stream in real time in parallel with ongoing VLA observations, potentially detecting and localizing multiple FRBs per year. **Sarah: any refinement to this statement?** VLASS+*realfast*?

### ACKNOWLEDGEMENTS

The National Radio Astronomy Observatory is a facility of the National Science Foundation operated under cooperative agreement by Associated Universities, Inc.. This research made use of Astropy, a community-developed core Python package for Astronomy (Astropy Collaboration, 2013). CJL is supported by the University of California Office of the President under Lab Fees Research Program Award 237863 and NSF award **\*\*number\*\***. MAM is supported by NSF award #1458952. KPM's research is supported by the Oxford Centre for Astrophysical Surveys which is funded through the Hintze Family Charitable Foundation. AS gratefully acknowledges support from the European Research Council under grant ERC-2012- StG-307215 LODESTONE. The AMI-LA telescope gratefully acknowledges support from the European Research Council under grant ERC-2012- StG-307215 LODESTONE, the UK Science and Technology Facilities Council (STFC) and the University of Cambridge. We acknowledge... Aspen Center for Physics Liam Connor

### REFERENCES

- Aschwendan, M. J. 2011, SoPh, 274, 99
- Baldry, I. K., et al. 2012, MNRAS, 421, 621
- Champion, D. J., et al. 2016, MNRAS, 460, L30
- Chatterjee, S., et al. 2017, Nature, 541, 58
- CHIME Scientific Collaboration et al. 2017, ArXiv e-prints
- Connor, L., Pen, U.-L., & Oppermann, N. 2016a, MNRAS, 458, L89
- Connor, L., Sievers, J., & Pen, U.-L. 2016b, MNRAS, 458, L19
- Cordes, J. M., & McLaughlin, M. A. 2003, ApJ, 596, 1142
- Cordes, J. M., & Wasserman, I. 2016, MNRAS, 457, 232
- Cordes, J. M., Wharton, R. S., Spitler, L. G., Chatterjee, S., & Wasserman, I. 2016, ArXiv e-prints
- Cordes et al. 2017, in prep

- Deng, W., & Zhang, B. 2014, *ApJL*, 783, L35
- Dokuchaev, V. I., & Eroshenko, Y. N. 2017, *ArXiv e-prints*, 1701.02492
- Faber, S. M., et al. 2007, *ApJ*, 665, 265
- Fenimore, E. E., & Bloom, J. S. 1995, *ApJ*, 453, 25
- Foreman-Mackey, D. 2016, *The Journal of Open Source Software*, 24
- Foreman-Mackey, D., Hogg, D. W., Lang, D., & Goodman, J. 2013, *PASP*, 125, 306
- Göğüş, E., Woods, P. M., Kouveliotou, C., van Paradijs, J., Briggs, M. S., Duncan, R. C., & Thompson, C. 2000, *ApJL*, 532, L121
- Hankins, T. H., Kern, J. S., Weatherall, J. C., & Eilek, J. A. 2003, *Nature*, 422, 141
- Hessels et al. 2017, in prep
- Inoue, S. 2004, *MNRAS*, 348, 999
- Ioka, K. 2003, *ApJL*, 598, L79
- Kashiyama, K., & Murase, K. 2017, *ArXiv e-prints*, 1701.04815
- Katz, J. I. 2014, *PhRvD*, 89, 103009
- . 2016, *Modern Physics Letters A*, 31, 1630013
- Kulkarni, S. R., Ofek, E. O., Neill, J. D., Zheng, Z., & Juric, M. 2014, *ApJ*, 797, 70
- Law, C. J., et al. 2015, *ApJ*, 807, 16
- Lawrence, E., Vander Wiel, S., Law, C. J., Burke Spolaor, S., & Bower, G. C. 2016, *ArXiv e-prints*
- Li, L., Huang, Y., Zhang, Z., Li, D., & Li, B. 2016, *ArXiv e-prints*
- Lorimer, D. R., Bailes, M., McLaughlin, M. A., Narkevic, D. J., & Crawford, F. 2007, *Science*, 318, 777
- Luan, J., & Goldreich, P. 2014, *ApJL*, 785, L26
- Lyutikov, M., Burzawa, L., & Popov, S. B. 2016, *MNRAS*, 462, 941
- Macquart, J.-P., & Johnston, S. 2015, *MNRAS*, 451, 3278
- Mao, S., & Paczynski, B. 1992, *ApJL*, 388, L45
- Marcote, B., et al. 2017, *ApJL*, 834, L8
- McQuinn, M. 2014, *ApJL*, 780, L33
- Metzger, B. D., Berger, E., & Margalit, B. 2017, *ArXiv e-prints*, 1701.02370
- Oppermann, N., Connor, L. D., & Pen, U.-L. 2016, *MNRAS*, 461, 984
- Perrott, Y. C., et al. 2013, *MNRAS*, 429, 3330
- Petroff, E., et al. 2015, *MNRAS*, 454, 457
- Planck Collaboration et al. 2016, *A&A*, 594, A13
- Rajwade, K. M., & Lorimer, D. R. 2017, *MNRAS*, 465, 2286
- Rane, A., Lorimer, D. R., Bates, S. D., McMann, N., McLaughlin, M. A., & Rajwade, K. 2016, *MNRAS*, 455, 2207
- Ravi, V., et al. 2016, *ArXiv e-prints*
- Scholz, P., & Kaspi, V. M. 2011, *ApJ*, 739, 94
- Scholz, P., et al. 2016, *ArXiv e-prints*
- Spitler, L. G., et al. 2014, *ApJ*, 790, 101
- . 2016, *Nature*, 531, 202
- Tauris, T. M., & Manchester, R. N. 1998, *MNRAS*, 298, 625
- Tendulkar, S. P., et al. 2017, *ApJL*, 834, L7
- Thornton, D., et al. 2013, *Science*, 341, 53
- Vedantham, H. K., Ravi, V., Hallinan, G., & Shannon, R. M. 2016, *ApJ*, 830, 75
- Zhang, B. 2017, *ArXiv e-prints*, 1701.04094