


# How can SaniPath build on the SFD's?

Suraja Raj, MPH

Center for Global Safe WASH at Emory University

## Case Study: Vellore, Tamil Nadu, India

- Two neighborhoods:
  - Cinna Allapuram (CAP) and Old Town (OT)
- Collaboration with Christian Medical College, Vellore, India and MAL-ED study



## **Study Objectives**

- Objective 1: To understand the dominant pathways of exposure to fecal contamination in two neighborhoods of Vellore, India.
  - Part 1: SaniPath Tool Deployment
- Objective 2: To quantify the associations between <u>household toilets</u> and <u>fecal sludge management</u> (FSM) with fecal contamination in different urban contexts
  - Part 2: Creating SFDs from SaniPath Data
  - Part 3: Extended SaniPath data collection and Spatial Analysis

## Part 1: Deployment of the SaniPath Tool

- Systematic, customizable method to collect relevant data on exposure to fecal contamination
- Help guide decision-making and advocacy surrounding urban sanitation
- Synthesize data using open-source software package

### SaniPath Field Methods

- Environmental Samples
  - 10 public area samples/ neighborhood, 25 HH samples
- Behavioral Surveys
  - Household surveys
     (100/neighborhood), School
     Surveys (4/neighborhood),
     community surveys
     (4/neighborhood)
- GPS data



### SaniPath Tool Exposure Assessment Analysis



Frequency of Municipal Drinking Water Contact in Shaibu (adults)

13.39%

12.95%

Other parameters:
intake volumes,
duration of


every day.

Piped Water (Adult)
Percent Exposed = 89%
Log10 Dose 3.1


exposure, etc.


Tool uses Bayesian analysis to estimate the distribution of environmental contamination and frequency of exposure.


#### **Environmental Contamination**




The mean dose and proportion of the population exposed are summarized from simulated distributions and displayed in risk profiles (left).












 Municipal piped drinking water and drain water posed the highest risks of exposure of adults and children living in CAP and Old Town

### Part 2: Building Shit Flows Diagrams







## Part 3: Spatial Analysis

- Kulldorff's Bernoulli Spatial Scan
- Microbial concentrations in environmental samples (outcome) and household survey characteristics (predictors) assessed for spatial clustering

## **Summary Points**

- The SaniPath Deployment showed Piped Drinking Water and Open Drains posed the highest risk of exposure in CAP and Old Town
- The SFDs showed that estimated household toilet coverage: 73% (CAP) vs. 32% (OT); estimated proportion of safely-managed excreta: 11% (CAP) vs. 2% (OT)
- Additional spatial analysis showed that areas with higher coverage of toilets and poor FSM had more pathogens in drains

## Acknowledgements

- Bill & Melinda Gates Foundation:
- Alyse Schrecongost, Erica Coppel, Radu Ban
- Center for Global Safe Water, Emory University:

Christine Moe, Clair Null, Peter Teunis, Monique Hennink, Kelly Baker, Amy Kirby, Julie Clennon, Juan Leon, Kate Robb, Habib Yakubu, Suraja Raj, Andrew Wang, Katherine Roguski

Christian Medical College, Vellore:

Prof. Gagandeep Kang, Dr. Venkata Mohan, Dr. Arun Kartikeyan, Sheela Roy; Study staff: Annai Gunasekaran, Priya Hemavathy, Daniel Beno, Ben Ghale, and all the field teams

- EpiTech Consulting: Nishant Kishore, Aaron White
- Research Triangle Institute: Matthew Scruggs, Megan Tulloch, Amir Mokhtari,
   Stephen Beaulieu
- Study Participants
- Paper in submission to AJTMH







## Thank You

For more information visit **SaniPath.org** 



Suraja Raj, MPH sraj@emory.edu

David M. Berendes, PhD, MSPH david.berendes@ce.gatech.edu