

Siphonophores from surface waters of the Colombian Pacific Ocean

JULIAN URIBE-PALOMINO¹, RAÚL LÓPEZ², MARK J. GIBBONS³, FELIPE GUSMÃO⁴
AND ANTHONY J. RICHARDSON^{1,5}

¹CSIRO Oceans and Atmosphere, Ecosciences Precinct, Dutton Park, Brisbane, QLD 4102, Australia, ²Programa de Biología Aplicada, Facultad de Ciencias Básicas y Aplicadas, Universidad Militar Nueva Granada, Campus Nueva Granada Km 2 vía Cajicá–Zipaquirá, Colombia, ³Biodiversity and Conservation Biology Department, University of the Western Cape, Private Bag X17, Bellville 7535, RSA, ⁴Department of Marine Science, Federal University of São Paulo, Santos, SP, Brazil, ⁵Centre for Applications in Natural Resource Mathematics (CARM), School of Mathematics and Physics, The University of Queensland, St Lucia, QLD 4072, Australia

Siphonophores are colonial hydrozoans that feed on zooplankton including fish larvae, and occur throughout the world's oceans from surface waters to ocean depths. Here we describe the composition of hyponeustonic siphonophores (0–3 m depth) from the tropical Colombian Pacific Ocean based on 131 plankton samples collected between June–October from 2001–2004. Samples were dominated by species of Calyptophorae, with only three species of Physonectae identified, consistent with their deeper depth distribution. Muggiaea atlantica, Chelophyes contorta, Diphyes dispar, and Eudoxoides mitra were the most common of the 21 species identified. We found moderate structuring of the siphonophore community by the salinity gradient from inshore to offshore, and greater richness during the night because of diel vertical migration. Temperature did not play a significant role in structuring siphonophore communities, perhaps because of the narrow temperature range observed (3.5 °C). We extend the known temperature and salinity range of several species, including M. atlantica up to temperatures of 28.6 °C and salinities down to 24.7. Interestingly, only polygastric stages of M. atlantica were found, suggesting the reproductive stage of M. atlantica in tropical waters might be found in deeper waters. Chelophyes appendiculata was rare in our study and C. contorta was common, providing evidence they have a potential allopatric relationship, with C. contorta replacing C. appendiculata in warm water. Finally, we found siphonophore abundance was positively related to the abundance of copepods and fish eggs, with the top 13 most abundant species all having positive correlations, suggesting siphonophore abundances are tightly controlled by their food.

Keywords: Gelatinous plankton, tropical zooplankton, cnidarian, biogeography, hyponeuston, plankton community, equatorial, diversity, temperature, salinity

Submitted 14 February 2017; accepted 18 December 2017

INTRODUCTION

Siphonophores are hydrozoans characterized by having complex colonies, comprising functional and specialized polymorphic units (zooids) that perform different tasks, including feeding and defence (gastrozooids), and reproduction (gonophores). The general body plan of siphonophores varies between taxonomic levels, and colonies range in size from a few centimetres to tens of metres (Mapstone, 2014). Some siphonophores display diel vertical migration, whilst others do not (Pugh, 1984; Mackie *et al.*, 1988; Mapstone, 2014). They play an important role in the ecology of marine systems: they reproduce quickly (Mackie *et al.*, 1988); they can be voracious predators of zooplankton and fish larvae (Alvariño, 1985; Purcell, 1985; Arai, 1988; Kršinić & Njire, 2001; Mills, 2001); they compete with fish for food (Arai, 1988; Buecher, 1999; Thibault-Botha *et al.*, 2013); and they are distributed throughout the water column, particularly in the deep sea (Silguero & Robison, 2000). When abundant,

they can impact economic activities such as tourism and marine aquaculture (Fosså *et al.*, 2003; Baxter *et al.*, 2011; Licandro *et al.*, 2012).

In the Pacific Ocean, there are many studies on the composition and distribution of siphonophores in temperate waters (Palma, 1973; 1999; Pagès *et al.*, 1990; Gasca & Suárez, 1992a, b; Palma & Rosales, 1995; Gasca, 2002; Palma & Apablaza, 2004; Palma & Silva, 2004; Apablaza & Palma, 2006; Palma *et al.*, 2007; Sanvicente-Añorve *et al.*, 2007, 2009; Lo *et al.*, 2012; 2013, 2014; Gamero-Mora *et al.*, 2015), but comparatively few from equatorial and tropical waters. Bigelow (1911) identified 40 siphonophores from samples collected during the Albatross expedition in October 1904 to March 1905, but few samples were collected from the Eastern Tropical Pacific. Alvariño (1971) also documented siphonophore species from the tropical-equatorial Eastern Pacific Ocean collected during two expeditions outside of the Colombian marine territory – Shellback (May–August 1952) and Capricorn (November 1952–February 1953). Andrade (2012) identified 15 species around Santa Clara Island (Ecuador) from hyponeustonic samples collected in September and November 2007, with *Muggiaea atlantica* and *Chelophyes appendiculata* being the dominant species at flood tide. The only work on the composition, distribution

Corresponding author:

J. Uribe-Palomino

Email: julian.uribepalomino@csiro.au

and abundance of siphonophores from neritic equatorial waters and from the Colombian Pacific Ocean is unpublished research from the thesis of Cely & Chiquillo (1993). They found 29 species, with *Chelophyes appendiculata*, *Diphyes dispar* and *Abylopsis eschscholtzii* dominant. This latter study was based on oblique samples collected in a single cruise conducted in waters adjacent to the coast in February–March 1991.

By comparison with the rest of the planktonic environment, the hyponeuston is poorly known and yet it provides an important habitat for early life history stages of fishes (Banse, 1964; Ahlstrom & Stevens, 1976; Jeong *et al.*, 2014), whose abundances may be impacted by siphonophores (Purcell, 1985). Here we provide new insights into the poorly known hyponeuston of the tropics, which are some of the most biodiverse areas of the world (Roy *et al.*, 1998; Boltovskoy, 1999; Harris *et al.*, 2000; Macpherson, 2002; Pierrot-Bults & Angel, 2013). We investigate siphonophore diversity within the hyponeuston for two major orders of siphonophores (Calycephorans and Physonects) and its relation to environmental variables. Samples were collected from coastal and oceanic waters of the tropical Colombian Pacific Ocean (CPO) over five cruises conducted over four consecutive years.

MATERIALS AND METHODS

Study region

The study area (339,000 km²) was the CPO and part of the Panamá Bight (CCCP, 2002) (Figure 1). In this region, trade winds are strongest from December to March

(Rodríguez-Rubio & Wolfgang, 2003), so all cruises were conducted outside peak upwelling season. The development of the Colombian Current (in neritic waters) (CCCP, 2002; Martínez *et al.*, 2007) and the presence of oceanic upwelling occur as a result of the presence of the NE trade winds in the CPO from November–March. These oceanographic features decline once the trade winds lose intensity toward the second half of the year (Wyrski, 1966; Rodríguez-Rubio & Wolfgang, 2003) when our study was conducted.

Sampling

Five cruises were conducted on board the Colombian Army research platforms *ARC-Malpelo* and *ARC-Gorgona* on 23 June–12 July 2001 (denoted Jun–Jul 2001), 27 August–15 September 2001 (Aug–Sep 2001), 3–22 September 2002 (Sep 2002), 2–21 September 2003 (Sep 2003), and 16 September–8 October 2004 (Sep–Oct 2004). Biological samples and environmental data were collected between 77–84°W and 2–6°N on a fixed grid of 35 stations, although not all stations were sampled during each cruise because of severe weather conditions and technical issues (Figure 2). Stations were 30 nautical miles apart in coastal waters and 60 nautical miles apart in oceanic areas. Samples were designated as being day (6 am–6 pm) or night according to the local time of arrival at the site. Two stations (49 and 111) were sampled more than once on the cruise conducted in Jun–Jul 2001.

At each station, sea surface temperature was measured with a reversing thermometer because it could be deployed close to the sea surface. Salinity was measured with a CTD (SeaBird electronics Series SEACAT) and the first reliable data were




Fig. 1. Map of the Colombian Pacific Basin showing the bathymetry and location of the sampling stations.




Fig. 2. Contour maps of temperature and salinity, and number of species of siphonophores found by site in the CPO during June–July 2001, August–September 2001, September 2002, September 2003 and September–October 2004.

at 4 m depth because of ship movement and waves. Samples of surface water were collected at each station with a 5 l Niskin bottle, and a 500 ml subsample of this was stored at -20°C for later analysis of nutrient concentration (ammonia, nitrite, nitrate, phosphate and silicate), following protocols of Garay *et al.* (1993).

To investigate the siphonophore community in the hyponeuston, we collected zooplankton samples in the first 0–3 m of water. Plankton samples were collected using a circular net of 0.5 m diameter and 363 μm mesh size, following a standardized circular trawling trajectory for 10 minutes. Samples were fixed and preserved in 5% buffered formalin.

In the laboratory, all siphonophores were identified from the samples using compound (Carl Zeiss S-1000 and Leica Zoom 2000) and dissecting microscopes (Carl Zeiss-Axiostar). Species were identified following Pagès *et al.* (1992) and Palma (1973) on the basis of whole organisms (Cystonectae) or portions of colonies, nectophores, eudoxids, gonophores and bracts (Physonectae and Calyphorae). Confirmation of the species was completed in 2005 following direct communication with the late Dr Francesc Pagès.

Statistical analyses

Most of our analyses are based on presence:absence of siphonophores as we could not estimate abundance robustly on all cruises because the flowmeter did not always work properly. For the two cruises with flowmeter data, estimates of siphonophore abundance for each species were obtained by summing the number of polygastric and reproductive stages. For Calyphorae, this was the greater of the number of anterior or posterior nectophores (or gonophores/bracts for eudoxids). For Physonectae, counts were based on pneumatophores; multiple bracts and nectophores without a pneumatophore were considered to originate from a single individual. We assessed how representative the presence:absence data were by correlating the frequency of occurrence of each species (based on presence:absence data) against abundance for cruises in Sep 2002 and Sep–Oct 2004.

To assess whether there might be bottom-up or top-down control between siphonophores and their prey, we obtained data on the abundance of copepods and fish eggs from unpublished work from the same net samples as the siphonophores. We assumed that bottom-up control would manifest as a positive relationship between siphonophore abundance (response) and copepods or fish eggs (predictors), and top-down control as a negative relationship between siphonophores and copepods or fish eggs (Richardson & Schoeman, 2004). Visual assessment of diagnostic plots showed that $\log_{10}(x + 1)$ transformation of both the response and predictors improved normality and homogeneity of variance assumptions. Data on fish egg and larval abundance were only available for the two cruises when flowmeter data were available: Sep 2002 and Sep–Oct 2004. Abundances were converted to densities (per 1000 m³), with a knowledge of the fraction of the sample examined (most commonly 1/8) and the volume of water filtered by the nets estimated using a flowmeter. Linear models were constructed in R and visualized using the effects package (Fox, 2003).

To map the data for each cruise, we constructed contour plots of sea surface temperature and salinity using kriging in Surfer® 8.04 (Golden software, LLC, 2003). To assess the variation of environmental variables, we built a linear model with Temperature and Salinity (untransformed) as response variables and included two factors, Region (Coastal (<600 m) or Oceanic) and Cruise (the five surveys), and their interaction.

Multivariate analyses

To examine the response of the siphonophore community to environmental conditions, we performed non-metric Multidimensional Scaling (nMDS) analysis (Clarke *et al.*, 2014). nMDS is an ordination technique that seeks to preserve the rank dissimilarities between objects (here samples) based

on a suite of variables (here species). On the 2-D ordinations, the closer two samples are, the more similar are their communities. We used the Bray–Curtis dissimilarity measure because it is robust to joint absences, so samples do not appear closer on the ordination because they have no species in common (Clarke *et al.*, 2014). We performed 1000 iterations using the metaMDS function in the R package vegan (Oksanen, 2011, 2017). We related the nMDS ordination to Region, Cruise, SST and Salinity; nutrient concentrations and chl-*a* were excluded because there were too many missing data. To aid interpretation, we included vector plots of environmental variables, species scores, and 95% confidence limits of centroids for Regions (Coastal vs Oceanic) and Time of Day (Day vs Night) on the ordination. To assess significant differences between groups and for continuous predictors, we performed a Permutational Multivariate Analysis of Variance (ADONIS) (Anderson, 2001). This gives the significance and the proportion of variance in the multivariate data (siphonophore communities) explained individually by each of the environmental predictors, added sequentially. To identify siphonophore species that were driving community differences, we used the Dufrene–Legendre Indicator Species Analysis (Dufrene & Legendre, 1997) in the R package labdsv. We used 0.05 as the level of significance in all statistical tests, unless otherwise specified.

RESULTS

Environmental conditions

There were relatively small differences in temperature in the tropical CPO during the study (Figure 2). Mean SST was 26.9 °C (± 0.57 °C, standard deviation) with a range of 3.5 °C, with coolest temperatures (25.5 °C) in Jun–Jul 2001 and Sep–Oct 2004 in offshore waters to the south-east of the CPO. Waters with temperature >27 °C were located between 80°W and the coast in Jun–Jul 2001, Aug–Sep 2001 and Sep 2002, and to the north-east of the CPO in Sep 2003 and Sep–Oct 2004.

The linear model for SST showed that there was no significant interaction between Region and Cruise, suggesting the temperature difference between Regions (Coastal and Oceanic) was similar among Cruises. Coastal waters were significantly warmer (27.2 ± 0.07 °C, mean \pm standard error) than Oceanic (26.7 ± 0.06 °C) waters, although the mean difference was only 0.5 °C (Figure 3A). Temperature also varied significantly among Cruises (Figure 3A): Jun–Jul 2001 (26.7 ± 0.1 °C), Aug–Sep 2001 (27.0 ± 0.1 °C), Sep 2002 (27.1 ± 0.1 °C), Sep 2003 (27.0 ± 0.1 °C) and Sep–Oct 2004 (26.7 ± 0.1 °C), but the effect size (i.e. maximum mean difference among cruises) was only 0.4 °C.

There was a relatively large range in salinity (10.07) among sites during the study (Figure 2). Low salinities were often found in the north-east section of the CPO and extended away from the coast. The lowest salinity (24.78) was recorded during Sep–Oct 2004 in the area off Cabo Corrientes, where a number of large rivers emerge. In general, concentrations of nutrients were often below detection levels, except for highest values near the San Juan River mouth in Sep 2002 (NH₄ 10.29 µg at. l⁻¹, NO₃ 27.8 µg at. l⁻¹, PO₄ 1.6 µg at. l⁻¹ and SiO₄ 40.9 µg at. l⁻¹; Table 1). Mean chlorophyll-*a*




Fig. 3. Final linear models of (A) Temperature and (B) Salinity. Means \pm confidence intervals are shown.

concentration was relatively high at times in coastal waters, up to 7.24 mg m^{-3} , but low offshore (Table 1).

The linear model for Salinity showed there was no significant interaction between Region and Cruise, suggesting the salinity difference between Regions (Coastal and Oceanic) was similar among Cruises. The main effect of Cruise was not significant, but Region was significant (Figure 3B). Coastal waters were significantly less saline (30.7 ± 0.3) than Oceanic waters (32.6 ± 0.1).

Siphonophore species present

A total of 21 species of siphonophores were recorded from surface waters of the CPO (Table 2). Nectophores and associated pneumatophores, and bracts were recorded for two agalmatids (*Agalma okenii* and *Agalma elegans*) and polygastric and reproductive stages for five calycophorans (*Abylopsis eschscholtzii*, *Diphyes dispar*, *Diphyes bojani*, *Eudoxoides mitra* and *Eudoxoides spiralis*). Of these, only the eudoxid stages of *D. bojani* were collected in Sep 2002 and Sep 2003, and only the polygastric stages of *E. spiralis* were recorded in oceanic waters in Aug–Sep 2001, Sep 2002 and Sep–Oct 2004. *Bassia bassensis* was identified only once from a single gonophore in Jun–Jul 2001, whilst the remaining 11 species of Calycophorae were identified from polygastric stages (mainly the anterior nectophore), and *Nanomia bijuga* from nectophores.

The eight most common species, in decreasing order of importance (from presence:absence data), were *D. dispar* (67.2%), *Chelophyes contorta* (51.1%), *Muggiaea atlantica* (39.7%), *E. mitra* (30.5%), *A. eschscholtzii* (19.8%), *D. bojani* (19.8%), *A. okenii* (16.0%) and *A. elegans* (13.0%) and were found in all five surveys. *Eudoxoides spiralis*, *Sulculeolaria monoica*, *Sulculeolaria quadrivalvis* and *Sulculeolaria turgida* were absent in Sep 2003, and *Physalia physalis* was absent in Aug–Sep 2001 (Table 2). *Bassia bassensis*, *Ceratocymba leuckartii*, *Chelophyes appendiculata*, *Hippopodus hippopus*, *Lensia campanella*, *N. bijuga*, *Sulculeolaria biloba* and *Sulculeolaria chuni* were recorded sporadically.

Siphonophore richness

The species richness of siphonophores varied spatially and temporally. In oceanic waters there were up to nine species in a sample and in coastal waters up to seven species in a sample (Figure 2). There was higher richness in coastal waters off Buenaventura in Jun–Jul 2001 and Sep 2003 (Figure 2). There appeared to be higher siphonophore species richness in Sep–Oct 2004 than in other cruises. In the final linear model of siphonophore Richness (response) and Region, Time of Day and Cruise (predictors), only the main effects were significant (pseudo $r^2 = 16.2\%$, Figure 4A). Richness was significantly higher in Oceanic (3.5 ± 0.08) than Coastal waters (2.2 ± 0.12). Richness was

Table 1. Summary of environmental variables by cruise.

Variable	23 Jun–12 Jul 2001	27 Aug–15 Sep 2001	03–22 Sep 2002	02–21 Sep 2003	16 Sep–08 Oct 2004	All 5 cruises 2001–2004
Temperature (°C)	25.4–27.7 min. – max. mean ± SE (n) 26.73 ± 0.1 (30)	26.1–27.8 min. – max. mean ± SE (n) 27.03 ± 0.09 (23)	26–27.8 min. – max. mean ± SE (n) 27.114 ± 0.09 (25)	25.9–28.9 min. – max. mean ± SE (n) 27.04 ± 0.13 (25)	25.64–28.09 min. – max. mean ± SE (n) 26.73 ± 0.1 (25)	25.4–28.9 min. – max. mean ± SE (n) 26.92 ± 0.05 (128)
Salinity	27.64–33.63 min. – max. mean ± SE (n) 32.51 ± 0.22 (30)	27.14–33.49 min. – max. mean ± SE (n) 31.75 ± 0.34 (23)	28.15–33.65 min. – max. mean ± SE (n) 31.96 ± 0.28 (25)	28.95–33.71 min. – max. mean ± SE (n) 31.82 ± 0.26 (25)	24.78–34.85 min. – max. mean ± SE (n) 31.22 ± 0.47 (25)	24.78–34.85 min. – max. mean ± SE (n) 31.89 ± 0.14 (128)
Ammonia (µg at. l ⁻¹)	BDL–2.83 min. – max. mean ± SE (n) 0.96 ± 0.15 (28)	BDL–1.63 min. – max. mean ± SE (n) 0.15 ± 0.08 (20)	BDL–10.29 min. – max. mean ± SE (n) 1.12 ± 0.47 (24)	BDL–1.05 min. – max. mean ± SE (n) 0.84 ± 0.08 (8)	BDL–2.05 min. – max. mean ± SE (n) 0.57 ± 0.09 (25)	BDL–10.29 min. – max. mean ± SE (n) 0.74 ± 0.12 (105)
Nitrites (µg at. l ⁻¹)	BDL–0.1 min. – max. mean ± SE (n) 0.01 ± 0.0 (28)	BDL–0.08 min. – max. mean ± SE (n) 0.02 ± 0.0 (20)	BDL min. – max. mean ± SE (n) 0.005 ± 0.00 (24)	BDL–0.1 min. – max. mean ± SE (n) 0.05 ± 0.01 (13)	BDL–0.05 min. – max. mean ± SE (n) 0.02 ± 0.0 (25)	BDL–0.10 min. – max. mean ± SE (n) 0.02 ± 0.0 (110)
Nitrates (µg at. l ⁻¹)	BDL–0.62 min. – max. mean ± SE (n) 0.10 ± 0.03 (28)	BDL–2.14 min. – max. mean ± SE (n) 0.47 ± 0.16 (20)	BDL–2.78 min. – max. mean ± SE (n) 1.21 ± 1.16 (24)	BDL–2.91 min. – max. mean ± SE (n) 0.55 ± 0.16 (20)	BDL–0.42 min. – max. mean ± SE (n) 0.07 ± 0.02 (25)	BDL–27.78 min. – max. mean ± SE (n) 0.46 ± 0.24 (117)
Phosphates (µg at. l ⁻¹)	BDL–0.24 min. – max. mean ± SE (n) 0.09 ± 0.01 (28)	BDL–0.49 min. – max. mean ± SE (n) 0.15 ± 0.03 (20)	BDL–1.57 min. – max. mean ± SE (n) 0.89 ± 0.04 (24)	BDL–0.89 min. – max. mean ± SE (n) 0.25 ± 0.05 (16)	BDL–0.38 min. – max. mean ± SE (n) 0.18 ± 0.02 (25)	BDL–1.57 min. – max. mean ± SE (n) 0.31 ± 0.03 (113)
Silicates (µg at. l ⁻¹)	BDL–26.81 min. – max. mean ± SE (n) 4.18 ± 1.09 (28)	BDL–42.43 min. – max. mean ± SE (n) 7.83 ± 2.9 (20)	BDL–40.94 min. – max. mean ± SE (n) 3.34 ± 1.66 (24)	BDL–7.63 min. – max. mean ± SE (n) 2.02 ± 0.61 (17)	BDL–8.47 min. – max. mean ± SE (n) 1.84 ± 0.41 (25)	BDL–42.43 min. – max. mean ± SE (n) 3.81 ± 0.7 (114)
Chlorophyll-a (mg m ⁻³)	0.02–1.11 min. – max. mean ± SE (n) 0.42 ± 0.05 (26)	0.08–4.56 min. – max. mean ± SE (n) 0.82 ± 0.27 (20)	0.08–7.24 min. – max. mean ± SE (n) 1.73 ± 0.57 (16)	0.01–1.38 min. – max. mean ± SE (n) 0.122 ± 0.07 (19)	0.17–0.98 min. – max. mean ± SE (n) 0.63 ± 0.04 (25)	0.01–7.24 min. – max. mean ± SE (n) 0.69 ± 0.11 (106)

SE, Standard Error; BDL, below detection level.

also significantly higher at Night (3.4 ± 0.08) than during the Day (2.4 ± 0.11). Richness varied significantly among cruises, with significantly higher Richness in Sep–Oct 2004 (4.6 ± 0.13) than Jun–Jul 2001 (2.8 ± 0.13), Aug–Sep 2001 (2.6 ± 0.16), Sep 2002 (2.5 ± 0.16) or Sep 2003 (2.6 ± 0.15).

Siphonophore communities

The nMDS ordination shows that there is a modest amount of structuring of the siphonophore community in the CPO (Figure 5). The Adonis model explained 10.8% of the variation in siphonophore communities, with Region (5.0%), Time of Day (3.7%) and Salinity (2.1%) all significant, and Cruise and Temperature not significant. Centroids for the Coastal vs Oceanic effect are close together, but their 95% confidence intervals do not overlap, suggesting that species toward the left of the ordination tend to be more Coastal and those toward the right tend to be more Oceanic. The Indicator Species Analysis showed that the species most different between Coastal and Oceanic samples were *C. contorta* (IV = 0.48, $P = 0.02$, Relative frequency of 41.0% Coastal vs 75.0% Oceanic), *E. mitra* (IV = 0.28, $P = 0.092$, 25.6% Coastal vs 44.1% Oceanic), *D. bojani* (IV = 0.28, $P = 0.01$, 7.7% Coastal vs 33.4% Oceanic), *A. eschscholtzii* (IV = 0.25, $P = 0.01$, 10.3% Coastal vs 32.4% Oceanic), *E. spiralis* (IV = 0.15, $P = 0.07$, 5.1% Coastal vs 19.1% Oceanic) and *S. monoica* (IV = 0.13, $P = 0.03$, 0% Coastal vs 13.2% Oceanic).

The centroids for Day vs Night are separated (although still relatively close together), implying some species are different during the Day and Night. Species that were most different between Day and Night were *M. atlantica* (IV = 0.36, $P = 0.07$, 35.9% Day vs 57.8% Night), *E. mitra* (IV = 0.34, $P = 0.01$, 20.9% Day vs 48.4% Night) and *S. turgida* (IV = 0.11, $P = 0.08$, 2.3% Day vs 12.5% Night) (see also Table 3), suggesting they vertically migrate.

The gradient of Salinity points toward the bottom right in the ordination, indicating *B. bassensis*, *E. spiralis*, *D. bojani*, *C. contorta* and *A. eschscholtzii* preferred more saline water, whereas *H. hippopus* and *D. dispar* preferred less saline waters. Siphonophore species were most common in waters with a salinity of 31–33.5 (Figure 6), and this could simply reflect that there was more water of this salinity in the region (Figure 2). Interestingly, *D. dispar* and *M. atlantica* were found in water with salinity as low as 24.7 (Figure 6). Two rare species (*S. chuni* and *H. hippopus*) were also found at lower salinities (29 and 24.8, respectively).

Temperature was not significant in the Adonis model. Siphonophore species were most common in waters of 26–27.5 °C (Figure 6), and this could also simply reflect that there was more water of this temperature in the region (Figure 2). Common species such as *D. dispar* and *M. atlantica* were ubiquitous throughout the study area (25.6–28.6 °C) waters (Figure 6).

Siphonophore abundances

For the two cruises where we had abundance data, there was a significant positive correlation between species mean abundance and relative frequency ($r = 0.90$, $N = 40$, $P = 0.00001$), providing support for the use of presence:absence data as a measure of relative abundance. Mean densities (per 1000 m³) indicate that the most abundant species were *D. dispar* (mean = 447.6 ± 201.2 , standard error), *E. mitra*

Table 2. Siphonophore species identified from the samples collected in the CPO 2001–2004, and in February–March 1991 (Cely & Chiquillo, 1993).

ABR. CODE	SPECIES	Authority	Feb–Mar 1991	23 Jun–12 Jul 2001		27 Aug–15 Sep 2001		03–22 Sep 2002		02–21 Sep 2003		16 Sep–08 Oct 2004	
			Coastal	Coastal	Oceanic	Coastal	Oceanic	Coastal	Oceanic	Coastal	Oceanic	Coastal	Oceanic
	SUBORDER CALYCOPHORAE												
Ab_esc	<i>Abylopsis eschscholtzii</i>	(Huxley, 1859)	P	P	P-R		P		P	P	P	P	P-R
Ba_bas	<i>Bassia bassensis</i>	(Quoy & Gaimard, 1833)	P-R		R								
Ce_leu	<i>Ceratocymba leuckartii</i>	(Huxley, 1859)								P			
Ch_ape	<i>Chelophyes appendiculata</i>	(Eschscholtz, 1829)	P	P							P		
Ch_con	<i>Chelophyes contorta</i>	(Lens & van Riemsdijk, 1908)	P	P	P	P	P	P	P	P	P	P	P
Di_dis	<i>Diphyes dispar</i>	(Chamisso & Eysenhardt, 1821)	P-R	P-R	P-R	P-R	P-R	P-R	P-R	P-R	P-R	P-R	P-R
Di_boj	<i>Diphyes bojani</i>	(Eschscholtz, 1825)			P-R		P-R		R	R	R	P	R
Eu_mit	<i>Eudoxoides mitra</i>	(Huxley, 1859)	R	P-R	P-R	P-R	P-R	R	P-R	R	P-R	P	P-R
Eu_spi	<i>Eudoxoides spiralis</i>	(Bigelow, 1911)	P-R	P	P-R	R	P		P				P
Hi_hip	<i>Hippopodius hippopus</i>	(Forsskål, 1776)	P		P							P	P
Le_cam	<i>Lensia campanella</i>	(Moser, 1917)	P								P		P
Mu_atl	<i>Muggiaea atlantica</i>	(Cunningham, 1892)	P-R	P	P	P	P	P	P	P	P	P	P
Su_bil	<i>Sulculeolaria biloba</i>	(Sars, 1846)	P										P
Su_chu	<i>Sulculeolaria chuni</i>	(Lens & van Riemsdijk, 1908)	P		P			P				P	
Su_mon	<i>Sulculeolaria monoica</i>	(Chun, 1888)			P		P		P				P
Su_qua	<i>Sulculeolaria quadrivalvis</i>	(de Blainville, 1830)	P	P	P	P		P				P	P
Su_tur	<i>Sulculeolaria turgida</i>	(Gegenbaur, 1854)	P		P	P			P			P	P
	<i>Sulculeolaria</i> spp.			P	P	P	P	P	P	P	P	P	P
	SUBORDER PHYSONECTAE												
Ag_ele	<i>Agalma elegans</i>	(Sars, 1846)	Pn	Pn	Pn-N	Pn		Pn	N	Pn		N	Pn-N
Ag_oke	<i>Agalma okenii</i>	(Eschscholtz, 1825)	Pn-N	Pn	Pn-Br		Pn	Br	Pn	Pn	Pn	Pn-N-Br	Pn-Br
Na_bij	<i>Nanomia bijuga</i>	(Delle Chiaje, 1844)	N						N	N	N		
	SUBORDER CYSTONECTAE												
Ph_phy	<i>Physalia physalis</i>	(Linnaeus, 1758)			Pn				Pn		Pn		Pn
Other species found in February–March 1991 (Cely & Chiquillo, 1993)						Authority							Stage
SUBORDER CALYCOPHORAE													
<i>Abylopsis tetragona</i>						(Otto, 1823)							P-R
<i>Enneagonum hyalinum</i>						(Quoy & Gaimard, 1827)							P
<i>Lensia subtilis</i>						(Chun, 1886)							P
<i>Lensia hardy</i>						(Totton, 1941)							P
<i>Lensia subtiloides</i>						(Lens & van Riemsdijk, 1908)							P
<i>Rosacea cymbiformis</i>						(Delle Chiaje, 1830)							R
SUBORDER PHYSONECTAE													
<i>Halistemma striata</i>						(Totton, 1965)							Pn
<i>Lychnagalma utricularia</i>						(Claus, 1879)							Pn
<i>Nanomia cara</i>						(Agassiz, 1865)							Pn
<i>Physophora hydrostatica</i>						(Forsskål, 1775)							Pn
SUBORDER CYSTONECTAE													
<i>Rhizophysa eysenhardtii</i>						(Gegenbaur, 1859)							Pn
<i>Rhizophysa filiformis</i> (as <i>Epibulia ritteriana</i>)						(Forsskål, 1775)							Pn

The presence of each species is shown based on the type of morphological stage found in the samples. For calycophorans: polygastric (P) and/or reproductive (R). For physonects: pneumatophore (Pn) and nectophore(s) (N). For cystonects: pneumatophore (Pn).




Fig. 4. Final linear models of (A) Siphonophore richness and (B) Siphonophore abundance for Sep 2002 and Sep–Oct 2004. Means \pm confidence intervals are shown.

(213.6 ± 85.5), *C. contorta* (189.7 ± 46) and *M. atlantica* (130.0 ± 39.7) (Table 4). Physonectae were less numerous, and of the two species of *Agalma* identified, *A. okenii* (53.3 ± 29.9) were slightly more numerous than *A. elegans* (40.1 ± 24.3). Densities of the reproductive stages of *D. dispar* were almost twice as high as those of the polygastric stages; 4.6 times in *E. mitra*, and almost 35 times more in *D. bojani*.

The linear model of Siphonophore abundance (response) showed that Cruise ($P = 0.0004$) and Time of Day ($P = 0.05$) were significant ($r^2 = 33.4\%$), but Region, Temperature and Salinity were not significant (Figure 4B). Siphonophore abundance was greater in Sep–Oct 2004 than in Sep 2002, and was greater at Night than during the Day.

For the two cruises with abundance data, there was a significant positive correlation between siphonophore




Fig. 5. A nMDS of the 21 siphonophore species, with species scores included. Environmental predictors are also plotted from the final Adonis model: Region (Coastal and Oceanic) and Time of Day (Day and Night) centroids and 95% confidence intervals superimposed, and the environmental vector Salinity.

Table 3. Observation of species (%) during the entire study, discriminated by day and night and by coastal and oceanic areas of the CPO.

Siphonophore species	Overall	Day	Night	Coastal	Oceanic
SUBORDER					
CYSTONECTAE					
<i>Physalia physalis</i>	5.34	0.07	0.04	0.02	0.07
SUBORDER					
PHYSONECTAE					
<i>Agalma elegans</i>	12.98	0.13	0.13	0.15	0.12
<i>Agalma okenii</i>	16.03	0.13	0.18	0.17	0.16
<i>Nanomia bijuga</i>	3.05	0	0.05	0.02	0.04
SUBORDER					
CALYCOPHORAE					
<i>Abylopsis eschscholtzii</i>	19.85	0.16	0.22	0.08	0.27
<i>Bassia bassensis</i>	0.76	0	0.01	0	0.01
<i>Ceratocymba leuckartii</i>	0.76	0.02	0	0.02	0
<i>Chelophyes appendiculata</i>	1.53	0	0.03	0.02	0.01
<i>Chelophyes contorta</i>	51.15	0.45	0.55	0.33	0.61
<i>Diphyes dispar</i>	67.18	0.67	0.67	0.65	0.69
<i>Diphyes bojani</i>	19.85	0.16	0.22	0.06	0.28
<i>Eudoxoides mitra</i>	30.53	0.16	0.41	0.21	0.36
<i>Eudoxoides spiralis</i>	11.45	0.11	0.12	0.04	0.16
<i>Hippopodius hippopus</i>	2.29	0.02	0.03	0.02	0.02
<i>Lensia campanella</i>	2.29	0	0.04	0	0.04
<i>Muggiaea atlantica</i>	39.69	0.27	0.49	0.38	0.41
<i>Sulculeolaria biloba</i>	0.76	0	0.01	0	0.01
<i>Sulculeolaria chuni</i>	3.05	0.02	0.04	0.06	0.01
<i>Sulculeolaria monoica</i>	6.87	0.05	0.08	0	0.11
<i>Sulculeolaria quadrivalvis</i>	6.11	0.04	0.08	0.10	0.04
<i>Sulculeolaria turgida</i>	6.87	0.02	0.11	0.04	0.08

DISCUSSION

Our study provides new insights into the ecology of siphonophores in the hyponeuston of the Equatorial Pacific Ocean. We show that there is a moderate structuring of the siphonophore community by environmental conditions in the CPO. This structuring is a result of the Coastal-Oceanic effect, reflected in the gradient of salinity from inshore to offshore, and also the greater siphonophore species richness during the night. The effect of temperature did not play a significant role in structuring siphonophore communities in the CPO, even though there is a temperature gradient present during each cruise (warmer inshore), perhaps because of the narrow range of temperature observed during the study (3.5 °C). Our analysis suggests potential bottom-up control of the total number of siphonophores, as their density is related positively to the density of fish eggs and copepods in Sep 2002 and Sep–Oct 2004.

This is the first study to describe the surface-dwelling siphonophores from both oceanic and coastal regions of the CPO. Of the 21 species of siphonophores recorded, 17 were also recorded in the unpublished study of Cely & Chiquillo (1993) from the coastal area of the CPO. Otherwise the




Fig. 6. Box and whisker plots of the occurrence of each siphonophore species identified during the five surveys in the Colombian Pacific Ocean in relation to temperature and salinity. The vertical bar is the median, the box is the 25th and 75th percentiles, and the whiskers represent the range. n is the number of stations where each species was present during the study.

Table 4. Abundance of siphonophores species found in the Colombian Pacific Ocean on September 2002 and Sep–Oct 2004 and associated environmental data.

Siphonophore species		Mean abundance \pm S.E. (Ind. 1000 m ⁻³) N = 40
SUBORDER		
CYSTONECTAE		
<i>Physalia physalis</i>		3.38 \pm 2.41
SUBORDER		
PHYSONECTAE		
<i>Agalma okenii</i>		53.28 \pm 29.93
<i>Agalma elegans</i>		40.08 \pm 24.34
SUBORDER		
CALYCOPHORAE		
<i>Abylopsis eschscholtzii</i>		30.82 \pm 10.07
<i>Chelophyes contorta</i>		189.75 \pm 45.91
<i>Diphyes dispar</i>	Polygastric	155.28 \pm 30.33
	Reproductive	292.34 \pm 189.83
	Total	447.62 \pm 201.24
<i>Diphyes bojani</i>	Polygastric	1.10 \pm 0.85
	Reproductive	38.43 \pm 15.28
	Total	39.53 \pm 15.23
<i>Eudoxoides mitra</i>	Polygastric	38.03 \pm 18.72
	Reproductive	175.59 \pm 74.04
	Total	213.63 \pm 85.52
<i>Eudoxoides spiralis</i>		12.65 \pm 9.48
<i>Hippopodius hippopus</i>		4.68 \pm 3.38
<i>Lensia campanella</i>		9.59 \pm 8.20
<i>Muggiaea atlantica</i>		130.04 \pm 39.69
<i>Sulculeolaria biloba</i>		1.48 \pm 1.48
<i>Sulculeolaria chuni</i>		4.31 \pm 3.01
<i>Sulculeolaria monoica</i>		4.74 \pm 3.54
<i>Sulculeolaria quadrivalvis</i>		5.84 \pm 3.57
<i>Sulculeolaria turgida</i>		32.76 \pm 23.19
Variables	Units	mean \pm S.E. N = 40
Salinity		31.30 \pm 0.32
Temperature	°C	26.93 \pm 0.08
Phosphates	$\mu\text{g-at. l}^{-1}$	0.45 \pm 0.06
Silicates	$\mu\text{g-at. l}^{-1}$	1.90 \pm 0.30
Chlorophyll- <i>a</i>	mg.m^{-3}	0.76 \pm 0.15
Abundance of copepods	ind. 1000 m ⁻³	572,990.53 \pm 324,417.15
Abundance of fish eggs	ind. 1000 m ⁻³	63,120.12 \pm 25,684.12
Total abundance of siphonophores	ind. 1000 m ⁻³	1224.17 \pm 274.16
Filtered volume by net	m ³	119.01 \pm 4.63

species observed here from the CPO have all been recorded previously in the Pacific Ocean (Alvariño, 1971; Pagès *et al.*, 1990). Species such as *A. eschscholtzii*, *Abylopsis tetragona*, *B. bassensis*, *C. appendiculata*, *C. contorta*, *E. spiralis*, *M. atlantica*, *S. chuni* and *S. quadrivalvis* are widely distributed in tropical waters of the Eastern Pacific (Gasca & Suárez, 1992a, b) and the Caribbean (Gasca, 1999), further west in the tropical Pacific (Pagès *et al.*, 1990; Lo *et al.*, 2012, 2014; Hsieh *et al.*, 2013) as well as cooler Chilean (Palma, 1999; Palma & Silva, 2004) and Peruvian (Ayón *et al.*, 2008) waters. Most species of siphonophore have a near global distribution (Mackie *et al.*, 1988), with all of the species found here reported from either the Atlantic (Pugh, 1999) or the Indian oceans (Daniel, 1974).

All species found in our study are epipelagic (Mackie *et al.*, 1988; Mapstone, 2014) and perform vertical migration (Pugh, 1984; Mackie *et al.*, 1988), which explains the higher richness

we found at night. Consistent with the idea that there is a peak in siphonophore richness in subtropical/tropical waters (Mackie *et al.*, 1988; Boltovskoy, 1999; Macpherson, 2002; Pierrot-Bults & Angel, 2013), we found more species (21 species) than in temperate waters such as off Chile (16 species, Palma, 1999) and in the Mediterranean Sea (12 species, Thibault-Botha *et al.*, 2013).

Our study provides several other interesting findings that we will describe in more detail.

Calycophorans dominate the hyponeuston

Our samples were dominated by species of Calycophorae, both in terms of abundance (the seven most abundant species were all calycophorans) and richness, with six times as many species present as Physonectae. Only two species of physonects were commonly encountered (*Agalma okenii* and *A. elegans*), with a third (*Nanomia bijuga*) present in only four samples. Although calycophorans are usually more common than the more fragile physonects in plankton samples (Pugh, 1974; Gasca & Suárez, 1992a, b; Mapstone, 2014), this effect could have been exaggerated because many physonect species are generally found deeper in the water column than calycophorans (Robison *et al.*, 1998; Silguero & Robison, 2000). Similarly, no physonectids were found in other eastern Pacific surface waters surrounding Easter Island (Palma, 1999) and Santa Clara Island in Ecuador (Andrade, 2012). Studies on siphonophores from other tropical, subtropical and temperate regions all indicate a predominance of calycophorans in the hyponeuston (Pagès *et al.*, 1990; Andrade, 2012; Thibault-Botha *et al.*, 2013; Jeong *et al.*, 2014).

Why are there fewer than expected siphonophore species in our study?

The number of species found in our study (21 species) is lower than other tropical studies in the eastern Pacific by Cely & Chiquillo (1993) or Gasca & Suárez (1992a) (29 species in both studies), both of which collected fewer samples than our study. Here are two potential explanations.

First, both Cely & Chiquillo (1993) and Gasca & Suárez (1992a) sampled throughout the water column and it is generally accepted that the number of siphonophore species in the hyponeuston is less than the number in the rest of the water column (Hempel & Weikert, 1972; Pagès *et al.*, 1990; Andrade, 2012). This is partly because physonects are relatively rare in the hyponeuston (Robison *et al.*, 1998; Silguero & Robison, 2000); we only recorded three species, although other factors might be at play.

The second potential explanation is that the Cely & Chiquillo (1993) and Gasca & Suárez (1992a) studies both included upwelling conditions, whereas there was no upwelling during our study. There is generally greater richness of siphonophore species in upwelled water, such as off the Peruvian coast (Ayón *et al.*, 2008), the Californian coast (Gasca & Suárez, 1992b), the Dome of Costa Rica (Gasca & Suárez, 1992a), the Benguela off South Africa (Thiriot, 1978; Gibbons & Thibault-Botha, 2002; Thibault-Botha & Gibbons, 2005). During the study by Cely & Chiquillo (1993) in the CPO, the combination of north-east trade winds (Rodríguez-Rubio & Wolfgang, 2003) and the development

of the Colombian current (CCCP, 2002) led to active upwelling. Further, we found only one species of *Lensia* (*L. campanella* was found only three times), whereas Cely & Chiquillo (1993) found three *Lensia* species (*L. subtilis*, *L. hardy*, *L. subtiloides*) during the upwelling season (Dec–Mar) in the CPO. Other studies in colder waters, such as Thibault-Botha *et al.* (2013) from the Bay of Marseilles, have found more (four) *Lensia* species. *Lensia* are often considered to be restricted to cooler waters (Mackie *et al.*, 1988; Mapstone, 2014). In tropical waters, *Lensia* might be more abundant in colder waters below the thermocline, and only reach the surface during active upwelling.

Potential allopatric relationship between *Chelophyes appendiculata* and *Chelophyes contorta*

It was surprising that *C. appendiculata* was rare in our study and that *C. contorta* was much more common, opposite to the findings of Andrade (2012) and of Cely & Chiquillo (1993). This difference can be explained by the suggested allopatric relationship in the distribution of *C. appendiculata* and *C. contorta*, with *C. appendiculata* replaced by *C. contorta* in warmer water (Alvariño, 1971). Mean sea temperatures were considerably cooler in the study by Andrade (2012) (22.8 °C) than the present study (26.9 °C), favouring *C. appendiculata* over *C. contorta*. Similarly, sea temperatures were also likely to be cooler in the study by Cely & Chiquillo (1993) because the CPO in Feb–Mar is influenced by trade winds (Wyrtki, 1966) that stimulate oceanic upwelling (Rodríguez-Rubio & Wolfgang, 2003) and lead to cooler waters, again favouring *C. appendiculata* over *C. contorta*.

Warmer upper thermal limit for *Muggiaea atlantica*

Although *M. atlantica* is a widely distributed species in warm and cold temperate waters (Alvariño, 1971; Mapstone, 2014) and is probably the best-studied siphonophore (Thibault-Botha *et al.*, 2004; Batistić *et al.*, 2013; Blackett *et al.*, 2014, 2015), our data indicate that *M. atlantica* can be found in temperatures up to 28.6 °C, far beyond the upper critical limit (24 °C) proposed by Batistić *et al.* (2013). This could suggest some form of local population adaptation, perhaps with regard to eudoxid production.

Whilst *M. atlantica* has a wide distribution in tropical shelf waters of Colombia (Cely & Chiquillo, 1993) and Panamá (Alvariño, 1971, 1974), it is also common in the Humboldt Current (Bigelow, 1911; Santander *et al.*, 1981) and in neritic and oceanic waters off Chile (Palma & Silva, 2004; Pavez *et al.*, 2010). Indeed, Palma & Silva (2004) extend the distribution of this species to sub-Antarctic latitudes of Chile, where they found it at a range of temperatures from 4–13 °C. Temperatures >10 °C seem to stimulate eudoxid production for this species, as has been found by Blackett *et al.* (2014) in an extensive study in the Western English Channel. However, in warmer waters *M. atlantica* typically is replaced by *Muggiaea kochii* (Will, 1844) (Russell, 1934; Alvariño, 1971; Carré & Carré, 1991), but this species was not observed in our study even though Alvariño (1974) recorded *M. kochii* in the Pacific waters of Panamá.

Lower salinity tolerances for some species

Siphonophores are an exclusively marine group and it is presumed that they have an aversion to low salinity water. Sanvicente-Añorve *et al.* (2007) investigated siphonophores in coastal waters of the Caribbean and extended the salinity tolerance of *D. bojani*, *E. mitra*, *D. dispar* and *M. atlantica* down to 30.7, but there was no water below this salinity in their study. Blackett *et al.* (2015) further extended the salinity tolerance of *M. atlantica* down to 28.5. We found *C. contorta*, *D. bojani* and *E. mitra* down to a salinity of ~27.5, and *D. dispar* and *M. atlantica* down to 24.7. These species thus have a lower salinity tolerance than previously recorded.

Reproductive and polygastric stages of calyphorans

We found that three dominant species of siphonophore (*D. dispar*, *D. bojani* and *E. mitra*) had more reproductive than polygastric stages, which suggests their populations are growing rapidly. For example, the number of reproductive stages of *D. dispar* were almost twice the number of polygastric stages in Sep 2002 and Sep–Oct 2004. Favourable environmental conditions for siphonophores stimulate the production of a high number of reproductive stages (Gasca, 1999). Interestingly, only polygastric stages of *M. atlantica* were found. The possibility that the reproductive stage of *M. atlantica* in tropical waters might be found in deeper waters where the temperature is more appropriate for its production warrants further study.

Siphonophore abundance is related to their prey abundance

We found siphonophore abundance was positively related to the abundance of copepods and fish eggs during Sep 2002 and Sep–Oct 2004. The top 13 most abundant species (*D. dispar*, *E. mitra*, *C. contorta*, *M. atlantica*, *A. okenii*, *A. elegans*, *D. bojani*, *S. turgida*, *A. eschscholtzii*, *E. spiralis*, *L. campanella*, *S. quadrivalvis* and *S. monoica*) all have positive correlations with both copepod abundance and fish eggs. These positive correlations imply bottom-up control of siphonophores by their common prey (Richardson & Schoeman, 2004). There is growing evidence that the abundance of siphonophores is related to their prey. In Mexican Caribbean waters, Sanvicente-Añorve *et al.* (2007) found a significant relationship between the community of siphonophores and the zooplankton wet biomass. In waters of the Mediterranean Sea, Fernández de Puelles *et al.* (2007) found a positive correlation between the abundance of *Muggiaea*, *Lensia*, *Eudoxoides* and *Abylopsis*, and abundance of copepods. In subtropical waters of Taiwan, Lo *et al.* (2014) suggested that the abundance and distribution of siphonophores are related to zooplankton biomass.

CONCLUSION

This study provides new insights into the poorly known hyponeuston of the tropics. The moderate structuring of the siphonophore community found in our study is likely to be related to the gradient of salinity from inshore to offshore in

waters from the CPO in the non-upwelling season. However, further research is required to determine the effect of environmental variables on the diversity, distribution and abundance of siphonophore species from the Equatorial Tropical Pacific during upwelling. A more comprehensive study of this group of cnidarians during upwelling season also might help to clarify the finding from our study related to the trophic relation between the siphonophores community and their most common prey (copepods, fish eggs) in the hyponeuston and in the water column from coastal and oceanic regions of the CPO.

SUPPLEMENTARY MATERIAL

The supplementary material for this article can be found at <https://doi.org/10.1017/S0025315417002065>.

ACKNOWLEDGEMENTS

We thank the Vicerrectoría de investigaciones of the Universidad Nueva Granada and the Centro Control Contaminación del Pacífico CCCP-DIMAR, for supporting the research on zooplankton from the Colombian Pacific Ocean. We also want to thank the crew of the Colombian Research Platforms (ARC-Malpelo and ARC-Providencia) for their invaluable help in collecting samples and environmental information in the field. Thanks to Eddie Farid (Gorgona drifter), Gloria Giraldo from the Library of the Universidad de Bogotá Jorge Tadeo Lozano for bibliographic support, Andrea Uribe-Palomino for her collaboration gathering data in Colombia and Frank Coman for helping to review the drafts of the manuscript. The authors would like to dedicate this article to the memory of Dr Francesc Pagès and his work on siphonophores around the globe. The authors thank the three anonymous reviewers for their time and constructive comments that contributed to improvement of the manuscript.

FINANCIAL SUPPORT

This study is part of the project CIAS 2004–016, entitled: ‘Variación espacio temporal del zooplankton durante los cruceros de julio y septiembre de 2005 de la serie ERFEN en el océano Pacífico colombiano’. This project was financially supported by the Vicerrectoría de Investigaciones from the Universidad Militar Nueva Granada – Colombia.

REFERENCES

- Ahlstrom E.H. and Stevens E. (1976) Report of neuston (surface) collections made on an extended CalCOFI cruise during May 1972. *California Cooperative Oceanic Fish. Investigations Report* 18, 167–180.
- Alvaríño A. (1971) Siphonophores of the Pacific with a review of the world distribution. *Bulletin of the Scripps Institution of Oceanography University of California* 16, 432.
- Alvaríño A. (1974) Distribution of siphonophores in the regions adjacent to the Suez and Panama Canals. *Fishery Bulletin* 72, 527–546.
- Alvaríño A. (1985) Predation in the plankton realm; mainly with reference to fish larvae. *Investigaciones Marinas CICIMAR* 2, 1–122.
- Anderson M.J. (2001) A new method for non-parametric multivariate analysis of variance. *Austral Ecology* 26, 32–46.
- Andrade C. (2012) Sifonóforos (Cnidaria, Hydrozoa) de aguas superficiales alrededor de la Isla Santa Clara, durante Septiembre y Noviembre del 2007. *Acta Oceanográfica del Pacífico* 17, 139–146.
- Apablaza P. and Palma S. (2006) Efecto de la zona de mínimo oxígeno sobre la migración vertical de zooplancton gelatinoso en la bahía de Mejillones. *Investigaciones Marinas* 34, 81–95.
- Arai M.N. (1988) Interactions of fish and pelagic coelenterates. *Canadian Journal of Zoology* 66, 1913–1927.
- Ayón P., Criales-Hernandez M.I., Schwaborn R. and Hirche H.J. (2008) Zooplankton research off Peru: a review. *Progress in Oceanography* 79, 238–255.
- Banase K. (1964) On the vertical distribution of zooplankton in the sea. *Progress in Oceanography* 2, 53–125.
- Batistić M., Lučić D., Carić M., Garić R., Licandro P. and Jasprica N. (2013) Did the alien calycophoran *Muggiaea atlantica* outcompete its native congeneric *M. kochii* in the marine lakes of Mljet Island (Croatia)? *Marine Ecology* 34(Suppl. 1), 3–13.
- Baxter E.J., Rodger H.D., McAllen R. and Doyle T.K. (2011) Gill disorders in marine-farmed salmon: investigating the role of hydrozoan jellyfish. *Aquaculture Environment Interactions* 1, 245–257.
- Bigelow H.B. (1911) The Siphonophorae. *Memoirs of the Museum of Comparative Zoology at Harvard College* 38, 173–401.
- Blackett M., Licandro P., Coombs S.H. and Lucas C.H. (2014) Long-term variability of the siphonophores *Muggiaea atlantica* and *M. kochii* in the Western English Channel. *Progress in Oceanography* 128, 1–14.
- Blackett M., Lucas C., Harmer R. and Licandro P. (2015) Population ecology of *Muggiaea atlantica* (Cnidaria, Siphonophora) in the Western English Channel. *Marine Ecology Progress Series* 535, 129–144.
- Boltovskoy D. (1999) *South Atlantic zooplankton*, vol. 1. Leiden: Backhuys.
- Buecher E. (1999) Appearance of *Chelophyes appendiculata* and *Abylopsis tetragona* (Cnidaria, Siphonophora) in the Bay of Villefranche, north-western Mediterranean. *Journal of Sea Research* 41, 295–307.
- Carré C. and Carré D. (1991) A complete life cycle of the calycophoran siphonophore *Muggiaea kochii* (Will) in the laboratory, under different temperature conditions: ecological implications. *Philosophical Transactions of the Royal Society B: Biological Sciences* 334, 27–32.
- CCCP (2002) *Compilación Oceanográfica de la Cuenca Pacífica colombiana*. Dirección General Marítima, Tumaco, Colombia.
- Cely H.A. and Chiquillo J.E. (1993) *Quetognatos, sifonóforos e hidromedusas de la región costera del Pacífico colombiano*. Thesis. Universidad de Bogotá Jorge Tadeo Lozano, Bogotá, Colombia.
- Clarke K.R., Gorley R.N., Somerfield P.J. and Warwick R.M. (2014) *Change in marine communities: an approach to statistical analysis and interpretation*, 3rd edition. Plymouth: PRIMER-E, 260 pp.
- Daniel R. (1974) *Siphonophora from the Indian Ocean*, Vol. 15, No. 4. Zoological Survey of India.
- Dufrene M. and Legendre P. (1997) Species assemblages and indicator species: the need for a flexible asymmetrical approach. *Ecological Monographs* 67, 345–366.
- Fernández de Puellas M.L., Alemany F. and Jansá J. (2007) Zooplankton time-series in the Balearic Sea (Western Mediterranean): variability during the decade 1994–2003. *Progress in Oceanography* 74, 329–354.

- Fosså J.H., Flood P.R., Olsen A.B. and Jensen F. (2003) Små og usynlige, men plagsomme maneter av arten *Muggiaea atlantica* (Small and invisible, but troublesome jellyfish of the species *Muggiaea atlantica*). *Fisken Og Havet (Fish and Sea)* 2, 99–103.
- Fox J. (2003) Effect displays in R for generalised linear models. *Journal of Statistical Software* 8, 1–27.
- Gamero-Mora E., Ceballos-Corona G., Gasca R. and Morales-Blake A. (2015) Analisis de la comunidad del zooplancton gelatinoso (Hydrozoa, Ctenophora, Thaliacea) en el Pacifico central mexicano, abril-mayo 2011. *Revista de Biología Marina y Oceanografía* 50, 111–124.
- Garay J., Panizzo L., Ramírez G. and Sánchez J. (1993) *Manual de técnicas analíticas de parámetros físico-químicos y contaminantes marinos*. Cartagena: Oceanográficas, Centro de Investigaciones Hidrográficas, CIOH.
- Gasca R. (1999) Siphonophores (Cnidaria) and summer mesoscale features in the Gulf of Mexico. *Bulletin of Marine Science* 65, 75–89.
- Gasca R. (2002) Lista faunística y bibliografía comentadas de los sifonóforos (Cnidaria?: Hydrozoa) de México. *Anales del Instituto de Biología, Universidad Nacional Autónoma de México* 73, 123–143.
- Gasca R. and Suárez E. (1992a) Sifonóforos (Cnidaria:Siphonophora) del Domo de Costa Rica. *Revista de Biología Tropical* 40, 125–130.
- Gasca R. and Suárez E. (1992b) Sifonóforos (Cnidaria: Hydrozoa) de la zona sudoccidental de la Península de Baja California, en invierno y verano durante El Niño 1983. *Revista de Investigación Científica* 3, 37–46.
- Gibbons M.J. and Thibault-Botha D. (2002) The match between ocean circulation and zoogeography of epipelagic siphonophores around southern Africa. *Journal of the Marine Biological Association of the United Kingdom* 82, 801–810.
- Golden software, LLC (2003) *Surfer (version 8.04)*. Windows platform. Golden, Colorado. www.goldensoftware.com
- Harris R., Wiebe P., Lenz J., Skjoldal H.R. and Huntley M. (2000) *ICES zooplankton methodology manual*. London: Academic Press.
- Hempel G. and Weikert H. (1972) The neuston of the sub-tropical and boreal North-east Atlantic: a review. *Marine Biology* 13, 70–88.
- Hsieh H.Y., Yu S.F. and Lo W.T. (2013) Influence of monsoon-driven hydrographic features on siphonophore assemblages in the Taiwan Strait, Western North Pacific Ocean. *Marine and Freshwater Research* 64, 348–358.
- Jeong H.G., Suh H.L., Lee W. and Soh H.Y. (2014) Seasonal variation of the neustonic zooplankton community in southern waters of Korea. *Ocean Science Journal* 49, 167–181.
- Kršinić F. and Njire J. (2001) An invasion by *Muggiaea atlantica* Cunningham 1892 in the northern Adriatic Sea in the summer of 1997 and the fate of small copepods. *Acta Adriatica* 42, 49–59.
- Licandro P., Souissi S., Ibanez F. and Carré C. (2012) Long-term variability and environmental preferences of calycophoran siphonophores in the Bay of Villefranche (north-western Mediterranean). *Progress in Oceanography* 97–100, 152–163.
- Lo W.T., Kang P.R. and Hsieh H.Y. (2012) Siphonophores from a transect off southern Taiwan between the Kuroshio current and South China sea. *Zoological Studies* 51, 1354–1366.
- Lo W.T., Yu S.F. and Hsieh H.Y. (2013) Effects of summer mesoscale hydrographic features on epipelagic siphonophore assemblages in the surrounding waters of Taiwan, Western North Pacific Ocean. *Journal of Oceanography* 69, 495–509.
- Lo W.T., Yu S.F. and Hsieh H.Y. (2014) Hydrographic processes driven by seasonal monsoon system affect siphonophore assemblages in tropical-subtropical waters (Western North Pacific Ocean). *PLoS ONE* 9, 1–19.
- Mackie G.O., Pugh P.R. and Purcell J.E. (1988) Siphonophore biology. *Advances in Marine Biology* 24, 97–262.
- Macpherson E. (2002) Large-scale species-richness gradients in the Atlantic Ocean. *Proceedings of the Royal Society of London B: Biological Sciences* 269, 1715–1720.
- Mapstone G.M. (2014) Global diversity and review of Siphonophorae (Cnidaria: Hydrozoa). *PLoS ONE* 9, 1–37.
- Martínez-Aguilar T.I., Giraldo López A. and Rodríguez-Rubio E. (2007) Zooplankton en la Corriente de Colombia, Pacífico Colombiano, durante Marzo de 2006. *Boletín Científico CCCP* 14, 69–82.
- Mills C.E. (2001) Jellyfish blooms: are populations increasing globally in response to changing ocean conditions? *Hydrobiologia* 451, 55–68.
- Oksanen J. (2011) Multivariate analysis of ecological communities in R: vegan tutorial. R package version 3.3.3.
- Oksanen J. (2017). Package ‘vegan’. <https://cran.r-project.org>. 292pp.
- Pagès F., Gili J.M. and Bouillon J. (1990) The siphonophores (Cnidaria, Hydrozoa) of Hansa Bay, Papua New Guinea. *Indo-Malayan Zoology* 6, 133–140.
- Pagès F., Gili J.M. and Bouillon J. (1992) Siphonophores (Cnidaria, Hydrozoa) of the Benguela Current (Southeastern Atlantic). *Scientia Marina* 56, 65–112.
- Palma S. (1973) Contribución al estudio de los sifonóforos encontrados frente a la costa de Valparaíso. I. taxonomía. *Investigaciones Marinas* 4, 17–88.
- Palma S. (1999) Sifonóforos (Cnidaria, Hydrozoa) de aguas superficiales de Isla de Pascua. *Investigaciones Marinas* 27, 19–23.
- Palma S. and Apablaza P. (2004) Abundancia estacional y distribución vertical del zooplancton gelatinoso carnívoro en una área de surgencia en el norte del sistema de la Corriente de Humboldt. *Investigaciones Marinas* 32, 49–70.
- Palma S., Apablaza P. and Soto D. (2007) Diversity and aggregation areas of planktonic cnidarians of the southern channels of Chile (Boca del Guafo to Pulluche Channel). *Investigaciones Marinas* 35, 71–82.
- Palma S. and Rosales S. (1995) Composición, distribución y abundancia estacional del macroplankton de la bahía de Valparaíso. *Investigaciones Marinas* 23, 49–66.
- Palma S. and Silva N. (2004) Distribution of siphonophores, chaetognaths, euphausiids and oceanographic conditions in the fjords and channels of southern Chile. *Deep-Sea Research Part II: Topical Studies in Oceanography* 51, 513–535.
- Pavez M.A., Landaeta M.F., Castro L.R. and Schneider W. (2010) Distribution of carnivorous gelatinous zooplankton in the upwelling zone off central Chile (austral spring 2001). *Journal of Plankton Research* 32, 1051–1065.
- Pierrot-Bults A.C. and Angel M.V. (2013) Pelagic biodiversity and biogeography of the Oceans. *Biology International* 51, 9–35.
- Pugh P.R. (1974) The vertical distribution of the siphonophores collected during the Sond Cruise, 1965. *Journal of the Marine Biological Association of the United Kingdom* 54, 25–90.
- Pugh P.R. (1984) The diel migrations and distributions within a mesopelagic community in the North East Atlantic. 7. Siphonophores. *Progress in Oceanography* 13, 461–489.
- Pugh P.R. (1999) Siphonophorae. In Boltovskoy D. (ed.) *South Atlantic zooplankton*. Leiden: Backhuys Publishers, pp. 467–513.

- Purcell J.E.** (1985) Predation of fish eggs and larvae by pelagic cnidarians and ctenophores. *Bulletin of Marine Science* 37, 739–755.
- Richardson A.J. and Schoeman D.S.** (2004) Climate impact on plankton ecosystems in the Northeast Atlantic. *Science* 305, 1609–1612.
- Robison B.H., Reisenbichler K.R., Sherlock R.E., Silguero J.M.B. and Chavez F.P.** (1998) Seasonal abundance of the siphonophore, *Nanomia bijuga*, in Monterey Bay. *Deep-Sea Research Part II: Topical Studies in Oceanography* 45, 1741–1751.
- Rodríguez-Rubio E. and Wolfgang S.** (2003) On the seasonal circulation within the Panama Bight derived from satellite observations of wind, altimetry and sea surface temperature. *Geophysical Research Letters* 30, 1–4.
- Roy K., Jablonski D., Valentine J.W. and Rosenberg G.** (1998) Marine latitudinal diversity gradients: tests of causal hypotheses. *Proceedings of the National Academy of Sciences USA* 95, 3699–3702.
- Russell F.S.** (1934) On the occurrence of the siphonophores *Muggiaea atlantica* Cunningham and *Muggiaea kochii* (Will) in the English Channel. *Journal of the Marine Biological Association of the United Kingdom* 19, 555–558.
- Santander H., Luyo R.G., Carrasco S., Véliz M. and de Castillo O.S.** (1981) Catálogo de zooplancton en el mar peruano. Primera parte: área Pisco-San Juan. *Boletín Instituto del Mar Del Perú* 6, 75.
- Sanvicente-Añorve L., Alba C., Alatorre M.A. and Flores-Coto C.** (2007) Cross-shelf and vertical distribution of siphonophore assemblages under the influence of freshwater outflows in the southern Gulf of Mexico. *Hydrobiologia* 586, 69–78.
- Sanvicente-Añorve L., Alba C., Flores-Coto C. and Castillo-Rivera M.** (2009) Siphonophores off a riverine system in the southern gulf of Mexico: factors affecting their distribution and spatial niche breadth and overlap. *Aquatic Ecology* 43, 423–435.
- Silguero J.M.B. and Robison B.H.** (2000) Seasonal abundance and vertical distribution of mesopelagic calycophoran siphonophores in Monterey Bay, CA. *Journal of Plankton Research* 22, 1139–1153.
- Thibault-Botha D. and Gibbons M.** (2005) Epipelagic siphonophores off the east coast of South Africa. *African Journal of Marine Science* 27, 129–139.
- Thibault-Botha D., Lutjeharms J.R.E. and Gibbons M.J.** (2004) Siphonophore assemblages along the east coast of South Africa: meso-scale distribution and temporal variations. *Journal of Plankton Research* 26, 1115–1128.
- Thibault-Botha D., Santo M. and Neuport M.** (2013) Gelatinous zooplankton in the Bay of Marseilles – potential control of the fish population over artificial reefs? *Rapport Commission Internationale pour l'Exploration Scientifique de La Méditerranée* 40.
- Thiriot A.** (1978) Zooplankton communities in the west African upwelling area. In Boje R. and Tomczak M. (eds) *Upwelling ecosystems*. Berlin: Springer-Verlag, pp. 32–61.
- and
- Wyrski K.** (1966) Oceanography of the Eastern Equatorial Pacific Ocean. *Oceanography and Marine Biology: An Annual Review* 4, 33–68.

Correspondence should be addressed to:

J. Uribe-Palomino
CSIRO Oceans and Atmosphere, Ecosciences Precinct, Dutton
Park, Brisbane, QLD 4102, Australia
email: julian.uribepalomino@csiro.au