

## Note

# Benthic scavengers and predators of jellyfish, material for a review

RON M. L. ATES\*

Govert Flinckstraat 19, 1506LL Zaandam, the Netherlands

Received 6 April 2016; Accepted 18 November 2016 Responsible Editor: Dhugal Lindsay

**Abstract:** Instances of predation and scavenging of jellyfish (pelagic Scyphozoa, Hydrozoa and Ctenophora) by benthic invertebrates are reviewed and presented. Sea anemones and decapod crabs are widespread as well as common predators or scavengers of jellyfish. Dead or dying jellyfish are also preyed upon by species of Echinodermata.

**Key words:** Actiniaria, Decapoda, Echinodermata, predation, scavenging, Scyphozoa, Hydrozoa, Ctenophora

Over the past decades the interaction between jellyfish and other animals received some attention (e.g. Arai 2005). Evidence had become available countering the previous prejudiced ideas about the negligible number of animal species preying on jellyfish. Nevertheless, jellyfish continue to be depicted occasionally (see Arai 2005 for examples) as dead ends in the food chains of the sea and/or as a rare source of food for predators (e.g. Condon et al. 2011), despite an increasing number of species of fishes (Redeke 1911, Luttenberger 1981, Gorelova & Grudtsev 1986, Kelley 1987, Ates 1988, Arai 1988, Zann 1988: 208, Cerqueira & Haimovici 1990, Hall 1992, Harbison 1993, Massuti et al. 1998, Purcell & Arai 2001, Bonaldo et al. 2004, Arai 2005, Orsi Relini et al. 2010a, Orsi Relini et al. 2010b, Chaves et al. 2010, Thaler 2012, Cardona et al. 2012, Milisenda et al. 2014, Battaglia et al. 2014, Sweetman et al. 2014, Dias & Almeida 2015), birds, reptiles and mammals (Dathe 1989, Ates 1991, Gronert 1992, Shiomi & Ogi 1992, Bell 1996, Zonfrillo 1997, Peglow 1998, Corsi 2000, Arai 2005, Suazo 2008, Cardona et al. 2012, Jarman et al. 2013, Melville 2013, Jones & Seminoff 2013), other cnidarians and ctenophores (Purcell 1991, Bamstedt et al. 1997, Bayha et al. 2012), polychaetes, helminths, molluscs and arthropods (Arai 2005, Stoner & Layman 2015) known to take them as prey.

Invertebrate species belonging to other groups have infrequently been mentioned as predators or scavengers of jellyfish. However, scuba-divers may encounter benthic animals consuming live or dead jellyfish on a regular basis. In Table 1 all records of sea anemones, corals, decapod crabs and echinoderms consuming live or dead jellyfish, known to me from

the literature or otherwise, are listed.

Jellyfish (pelagic Scyphozoa, Hydrozoa and Ctenophora) easily fall victim to benthic invertebrate predators such as sea anemones. For about thirty years I have been able to observe for example *Metridium senile* taking jellyfish, mainly *Aurelia aurita*, *Cyanea lamarckii* and *Pleurobrachia pileus*, in considerable numbers in the Oosterschelde estuary, southwestern Netherlands (unpublished observations). Although the predation and the scavenging of jellyfish appears not to be limited geographically, most records in Table 1 are from eastern Atlantic waters, doubtlessly because I am more familiar with the grey literature in Europe.

The predominant prey mentioned in Table 1 taken by sea anemones, decapod crabs and echinoderms alike appear to be species of Scyphozoa. Observations of *Pleurobrachia*, *Neoturris*, *Eutonina*, *Eucheilota* and even scyphozoan ephyrae (see Table 1) make clear that this predation is however not size-limited. As the consumption time of the larger jellyfish exceeds several times that of the mostly much smaller hydrozoans and ctenophores, the likelihood of encountering Scyphozoa being consumed is much higher.

Few records in Table 1 go beyond the mere noting of the actual consumption of the jellyfish. In fact many of them have a casual or even tentative character. For patterns to be recognized more detailed observations need to be published. Hopefully future observers will publish more data pertaining to their observations of jellyfish being eaten. Berryman (1984) suggested that the predation by the sea anemone *Sagartiogeton laceratus* on *A. aurita* may become noteworthy when swarms of jellyfish are locally concentrated due to currents and topography. However, when scuba-diving, predation by sea anemones *sensu lato* upon jellyfish may be observed in each and every location where the two occur together. When

\*Corresponding author: Ron M. L. Ates; E-mail, pseudocorynactis@zonnet.nl

**Table 1.** Benthic Cnidaria, decapod Crustacea and Echinodermata recorded to have consumed live or dead jellyfish.

PREDATOR OR SCAVENGER	PREY	LOCALITY	SOURCE
<b>Cnidaria Actiniaria</b>			
<i>Actinia equina</i> (Linnaeus, 1758)	<i>Pleurobrachia pileus</i> (Müller, 1776)	North Sea	Den Hartog 1963
<i>A. equina</i>	<i>Cyanea lamarckii</i> Péron & Lesueur, 1810	Channel Islands, England	S. Daly in: Wood 2013
<i>Actinia tenebrosa</i> Farquhar, 1898	Scyphozoa	Australian waters	Ayre 1984
<i>Actinoporus elegans</i> Duchassaing, 1850	<i>Aurelia</i> ?	Curaçao, Caribbean	Van der Vlugt 1981*
<i>Actinostola callosa</i> (Verrill, 1882)	<i>Periphylla periphylla</i> (Péron & Lesueur, 1810)	Norwegian waters	Jarms & Tiemann 2004
<i>Anthopleura elegantissima</i> (Brandt, 1835)	<i>Velella velella</i> (Linnaeus, 1758)	northeastern Pacific	Francis 1973; Wertheim 1984
<i>Diadumene cincta</i> Stephenson, 1925	<i>Eutonina indicans</i> (Romanes, 1876)	Oosterschelde, Netherlands	pers. comm. M. Vestjens 2002
<i>Diadumene leucolea</i> (Verrill, 1866)	<i>Chrysaora quinquecirrha</i> (DeSor, 1848)	western Atlantic	Cargo & Schultz 1967*; Cones & Haven 1969
<i>Metridium farcimen</i> (Brandt, 1835)	<i>Cyanea capillata</i> (Linnaeus, 1758)	northeastern Pacific	pers. comm. R. Harbo 2012
<i>Metridium senile</i> (Linnaeus, 1767)	<i>Aequorea victoria</i> (Murbach & Shearer, 1902)	northeastern Pacific	pers. comm. C.E. Mills 2000
<i>M. senile</i>	hydroid fragments	western Atlantic	Sebens & Koehl 1984
<i>M. senile</i>	<i>Aurelia aurita</i> (Linnaeus, 1758)	Grevelingen, Netherlands	this paper, see photo B
<i>M. senile</i>	<i>A. aurita</i> , <i>P. pileus</i> , <i>Chrysaora hysoscella</i> (Linnaeus, 1767)	Oosterschelde, Netherlands	Ates 2002
<i>M. senile</i>	<i>P. pileus</i>	North Sea	Greve 1972*
<i>Paraphelliactis pabista</i> Dunn, 1982	medusae?	northeastern Pacific	Fautin 1982
<i>Sagartia</i> sp.	<i>P. pileus</i>	North Sea	Greve 1972*
<i>Sagartia elegans</i> (Dalyell, 1848)	<i>C. hysoscella</i>	Oosterschelde, Netherlands	this paper, see photo C
<i>Sagartia troglodytes</i> (Price in Johnston, 1847)	<i>A. aurita</i> , <i>C. lamarckii</i> , <i>C. hysoscella</i>	Oosterschelde, Netherlands	Ates 2002
<i>S. troglodytes</i>	<i>E. indicans</i>	Oosterschelde, Netherlands	Ates 2014
<i>Sagartiogeton laceratus</i> (Dalyell, 1848)	<i>A. aurita</i>	North Sea	Berryman 1984
<i>Sagartiogeton undatus</i> (Müller, 1788)	<i>Aequorea vitrina</i> Gosse, 1853	Grevelingen, Netherlands	this paper, see photo D
<i>S. undatus</i>	cf. <i>Eucheilota</i>	Oosterschelde, Netherlands	pers. obs. July 2007
<i>Urticina lofotensis</i> (?)	<i>Aurelia labiata</i> Chamisso & Eysenhardt, 1821, <i>Phacellophora camtschatica</i> Brandt, 1835	?	pers. comm. P. Lovejoy Sept. 2000
<i>Urticina felina</i> (Linnaeus, 1761)	<i>Rhizostoma pulmo</i> (Macri, 1778), <i>P. pileus</i>	North Sea	Den Hartog 1963
<i>U. felina</i>	<i>P. pileus</i>	North Sea	Greve 1972*
<i>U. felina</i>	jellyfish ( <i>Aurelia</i> ?)	Scottish waters	Wood 2005
<i>U. felina</i>	<i>A. aurita</i> , <i>C. lamarckii</i> , <i>C. hysoscella</i>	Oosterschelde, Netherlands	Ates 2002
<i>U. felina</i>	<i>C. hysoscella</i>	North Sea	Schwanitz, in Jarms & Tiemann 2004
<i>U. felina</i>	<i>A. aurita</i>	southwestern Ireland	pers. comm. W. Northway, May 2011
<i>Urticinopsis antarcticus</i> (Verrill, 1922)	jellyfish	Antarctic	Dayton et al. 1974
<i>U. antarcticus</i>	<i>Desmonema</i> sp.	Antarctic	Conniff 2000
sea anemone	<i>Mastigias</i> ?	Lake Kakaban, Indonesia	this paper, see photo E
<i>Entacmaea</i> (?) <i>medusivora</i> Fautin & Fitt, 1991	<i>Mastigias papua</i> (Lesson, 1830), <i>A. aurita</i> , <i>Cassiopea ornata</i> Haeckel, 1880	Lake Kakaban, Indonesia	Hoeksema et al. 2014
sea anemones	<i>M. papua</i>	Palau, western Pacific	Hamner 1982
<i>Aiptasia pulchella</i> (?) = <i>E.</i> (?) <i>medusivora</i>	<i>M. papua</i>	Palau, western Pacific	Hamner & Hauri 1981; Fautin & Fitt 1991
sea anemones	<i>A. aurita</i> ephyrae	Waddenzee, Netherlands	Swennen 1956*
sea anemone	<i>Nemopilema nomurai</i> Kishinouye, 1922	Sea of Japan	Yamamoto et al. 2008
<b>Cnidaria other than Actiniaria</b>			
<i>Alcyonium digitatum</i> Linnaeus, 1758	<i>P. pileus</i>	North Sea	Greve 1972*
<i>Cerianthus lloydii</i> Gosse, 1859	cf. <i>Neoturris pileata</i>	Scottish waters	this paper, see photo A
<i>Fungia scruposa</i> Klunzinger, 1879	<i>A. aurita</i>	Red Sea	Alamaru et al. 2009
<i>Tubularia</i> sp.	<i>Lensia</i> sp.	?	Mackie 1966
<i>Tubularia larynx</i> Ellis & Solander, 1786	gelatinous plankton	Irish Sea	Gili et al. 1996

Table 1. Continued.

PREDATOR OR SCAVENGER	PREY	LOCALITY	SOURCE
<b>Crustacea Decapoda</b>			
<i>Aristeomorpha foliacea</i> (Risso, 1827)	Siphonophora	NW-Australia	Rainer 1992
<i>Aristeus virilis</i> (Bate, 1881)	Siphonophora	NW-Australia	Rainer 1992
<i>Callinectes sapidus</i> (Rathbun, 1896)	<i>C. capillata</i> , <i>Stomolophus meleagris</i> (Agassiz, 1862)	Gulf of Mexico	Farr 1978
<i>Cancer pagurus</i> Linnaeus, 1758	<i>A. aurita</i>	North Sea	Steinich 1973
<i>Cancer</i> sp.	<i>Linuche</i> sp.	Caribbean	Larson 1991
<i>Carcinus maenas</i> (Linnaeus, 1758)	<i>P. pileus</i>	North Sea	Greve 1972; Esser et al. 2004*
<i>C. maenas</i>	<i>A. aurita</i>	Kattegat	Rasmussen 1973
<i>C. maenas</i>	<i>A. aurita</i>	Baltic Sea	Lauckner 1980
<i>Crangon crangon</i> (Linnaeus, 1758)	<i>P. pileus</i>	North Sea	Greve 1972*, Esser et al. 2004*
<i>Chorilia longipes</i> Dana, 1852	<i>C. capillata</i>	northeastern Pacific	Harbo 2011
<i>Galathea squamifera</i> Leach, 1814	<i>P. pileus</i>	North Sea	Greve 1972*
<i>Haliporoides sibogae</i> (De Man, 1907)	Siphonophora	NW-Australia	Rainer 1992
<i>Heterocarpus sibogae</i> De Man, 1917	Siphonophora	NW-Australia	Rainer 1992
<i>Heterocarpus woodmasoni</i> Alcock, 1901	Siphonophora	NW-Australia	Rainer 1992
<i>Homarus gammarus</i> (Linnaeus, 1758)	<i>A. aurita</i>	Grevelingen, Netherlands	pers. comm. J. Bakker June 2013
<i>Hyas araneus</i> (Linnaeus, 1758)	<i>C. capillata</i>	Scottish waters	Ates & Hoyinck 2012
<i>H. araneus</i>	<i>A. aurita</i>	Scottish waters	pers. comm. W. Northway May 2011
<i>Jasus lalandii</i> (Milne-Edwards, 1837)	hydromedusae	?	Williamson, in Thomas 1963*
<i>Liocarcinus depurator</i> (Linnaeus, 1758)	<i>C. capillata</i>	Scottish waters	pers. comm. W. Northway Sept 2013
<i>Menippe mercenaria</i> (Say, 1818)	<i>Stomolophus</i>	Gulf of Mexico	Powell & Gunter 1968
<i>Munida rugosa</i> (Fabricius, 1775)	<i>A. aurita</i>	Scottish waters	Naylor 2011
<i>Necora puber</i> (Linnaeus, 1767)	<i>C. lamareckii</i>	Oosterschelde, Netherlands	this paper, see photo F
<i>N. puber</i>	<i>A. aurita</i>	Scottish waters	pers. comm. W. Northway Sept 2013
<i>Ocyropsis quadrata</i> (Fabricius, 1787)	<i>Physalia physalis</i> (Linnaeus, 1758)	Gulf of Mexico	Phillips et al. 1969
<i>Pagurus anachoretus</i> Risso, 1827	<i>Pelagia noctiluca</i> (Forskål, 1775)	Mgar-Ix-Ieni, Gozo, Malta	this paper, see photo G
<i>Pagurus bernhardus</i> (Linnaeus, 1758)	<i>P. pileus</i>	North Sea	Greve 1972*, Esser et al. 2004*
<i>P. bernhardus</i>	<i>A. aurita</i>	Scottish waters	Ates & Hoyinck 2012; this paper, see photo H
<i>P. bernhardus</i>	<i>C. capillata</i>	Scottish waters	pers. comm. W. Northway Sept 2013
<i>Pagurus floridanus</i> (Benedict, 1892), <i>Pagurus pollicaris</i> Say, 1817	<i>Cyanea</i> , <i>Stomolophus</i>	Gulf of Mexico	Phillips et al. 1969*
<i>Portunus holsatus</i> Fabricius, 1798	<i>P. pileus</i>	North Sea	Greve 1972*
<i>Plesiopenaeus edwardsianus</i> (Johnson, 1868)	Siphonophora	NW-Australia	Rainer 1992
<i>Pugettia producta</i> (Randall, 1840)	jellyfish	northeastern Pacific	pers. comm. A. Lamb 2012
<i>Scyphax ornatus</i> Dana, 1853	<i>Physalia</i> , <i>A. aurita</i> , <i>Veella</i>	New Zealand intertidal	Quilter 1987
<b>Crustacea other than Decapoda</b>			
<i>Balanus eburneus</i> Gould, 1841	<i>C. quinquecirrha</i> ephyrae	western Atlantic	Cones & Haven 1969
<i>Orchomenella obtusa</i> (Sars, 1891), <i>Munida tenuimana</i> Sars, 1872, decapod shrimps	<i>P. periphylla</i> , <i>C. capillata</i>	Norwegian waters	Sweetman et al. 2014
<b>Echinodermata</b>			
<i>Asterias rubens</i> Linnaeus, 1758	<i>C. capillata</i>	Norwegian waters	Moen & Svensen 2000
<i>A. rubens</i>	<i>A. aurita</i>	Scottish waters	Ates & Hoyinck 2012
<i>Ophiocomina nigra</i> (Abildgaard, 1789)	<i>A. aurita</i>	Scottish waters	Ates & Hoyinck 2012
<i>Ophiotrix fragilis</i> (Abildgaard, 1789)	<i>A. aurita</i>	Oosterschelde, Netherlands	Ates 2002
<i>O. fragilis</i>	<i>A. aurita</i>	Scottish waters	Ates & Hoyinck 2012
seastar ( <i>Anseropoda</i> ?)	jellyfish	California	Wu 1998
<i>Strongylocentrotus franciscanus</i> (Agassiz, 1863)	jellyfish	northeastern Pacific	pers. comm. A. Lamb 2012
ophiurids	<i>Aurelia limbata</i> (Brandt, 1835)	Sea of Japan	Miyake et al. 2002

\* aquarium observation






Plate 1.



offered a jellyfish, in an experimental context under water, a sea anemone will rarely fail to grab and devour it. In deeper water beyond the reach of scuba-divers, sea anemones are also known to prey heavily on jellyfish (Jarms & Tiemann 2004). As described by Jarms & Tiemann (2004), jellyfish well exceeding the size of the oral disc of the sea anemone may be ingested as a result of a considerable inflation of its mouth and its oral disc. In shallow water this is exactly the same (see e.g. Ates 2002 and S. Daly in Wood 2013).

Predation upon jellyfish by decapod crabs is equally widespread (Table 1). None of the records seem to indicate that decapod crabs first attack the gonads, the most nutritious part of the jellyfish (cf. Rasmussen 1973). Contrary to sea anemones, decapod crabs may be able to actively pursue jellyfish as has been reported by Esser et al. (2004) for *Pagurus bernhardus* jumping up to capture *P. pileus* from the water column. Decapod crabs also are able to protect and defend their jellyfish prey, e.g. by moving away from a disturbance with it. This has been documented for *Callinectes sapidus* by Farr (1978) and for *Hyas araneus* by Ates & Hoyinck (2012). Farr (1978) suggested that crabs may only prey on jellyfish in the absence of other prey. However, I can testify that in the majority of cases witnessed by me (see Table 1) other prey items were available, sometimes abundantly. It is not easy to have decapod crabs accept a jellyfish prey in situ as they are easily disturbed by the presence of a diver contrary to sea anemones. In cases where such an experiment succeeds, most crab species grab and eat the jellyfish.

Echinoderms have rarely been mentioned as scavengers of jellyfish. There can however be no doubt that *Ophiocoma nigra* plays an important role as such (Ates & Hoyinck 2012). During three dives in Loch Sunart (Scotland) Ates & Hoyinck (2012) observed at least twelve dead specimens of *A. aurita* in various stages of consumption by *O. nigra* and to a lesser extent by *Asterias rubens* and *Ophiothrix gracilis*. The jellyfish was always consumed from the outside in, leaving the highest part of the velum to be eaten last. There is no indication that echinoderm scavengers go for the gonads first, if these are

present.

As jellyfish often occur in swarms, their simultaneous death may lead to a high number of jellyfish carcasses raining down on the sea floor in a limited period of time at certain locations. In the last decade this has led to an increased interest in the impact of jellyfish carcasses on deep-sea ecology (e.g. Lebrato et al. 2012). Yamamoto et al. (2008) implicated one sea anemone, two decapod, one gastropod and one ophiurid species to be scavengers of *Nemopilema nomurai* carcasses in deep water of the Sea of Japan. In their experiments in a deep Norwegian fiord Sweetman et al. (2014) found evidence of scavenging by hagfish and some species of small crustaceans in high numbers consuming jellyfish carcasses in a matter of hours. If a high number of dead jellyfish fall down in a very short time scavengers may be too few to deal with their remains, as has been documented by Billett et al. (2006) and Lebrato et al. (2012). Billett et al. (2006) did not investigate the diet of the spider crab *Encephaloides armstrongi* Woodmason present in large numbers as can be seen on their picture 3A. Elsewhere spider crabs are known as predators or scavengers of jellyfish, see Table 1.

Incidentally some of the records of predation on jellyfish mentioned in the introduction relate to non-pelagic fishes. Luttenberger (1981), Chaves et al. (2010), Thaler (2012) and Dias & Almeida (2015) reported about coral reef fishes (e.g. species of *Chaetodon*, *Zebrasoma*, *Pomacanthus* and *Naso*) in different parts of the world benefitting from the influx of jellyfish into their realm. All these authors described the events as feeding frenzies, leading to the (near-)annihilation of the jellyfishes. This and my list of benthic invertebrate predators and scavengers (Table 1) contradicts the notion that “jellyfish are not readily consumed by other predators” (Condon et al. 2011). Quite the opposite may occur when jellyfish hit the coast or the sea floor. No barrier seems to exist for many benthic animals to consume dead or live jellyfish with relish. I have no doubt that proper fieldwork will reveal that jellyfish are being consumed by many more benthic scavengers and predators other than those contained in Table 1.

#### Plate 1.

- A. *Cerianthus lloydii* (Cnidaria: Anthozoa: Ceriantharia) engulfing *Neoturris* cf. *pileata* measuring less than 1.5 cm in diameter. Depth about 7 m. Picture taken on June 27, 2012 on the Lochaline Wall (Sound of Mull, Scotland) by R. M. L. Ates.
- B. A young *Aurelia aurita* measuring about 3 cm in diameter being taken by an immature *Metridium senile* in Dutch coastal waters. Depth unknown. Picture taken by R. Offermans on April 9, 2011.
- C. Two *Sagartia elegans* holding one *Chrysaora hysoscella*. Diameter of the sea anemones between 3 and 4 cm. Depth about 5 m. Picture taken by R. Offermans in the Oosterschelde estuary, southwestern Netherlands on July 17, 2015.
- D. *Sagartiogeton undatus* caught in the act of engulfing *Aequorea vitrina*, diameter about 5 cm. Depth about 10 m. Picture taken by R. M. L. Ates in July, 2005 in the Grevelingen, southwestern Netherlands.
- E. Unknown sea anemone consuming *Mastigias* sp. in Lake Kakaban, Kalimantan, Indonesia. Column height of the sea anemone about 5 cm. Depth unknown. Picture taken on October 2, 2010. Courtesy H. van Rijn.
- F. *Necora puber* eating *Cyanea lamarckii*, diameter less than 10 cm. Depth about 7 m. Picture taken in the Oosterschelde, Netherlands in August 2003 by R. M. L. Ates.
- G. Large numbers of *Pelagia noctiluca* may perish after entering Mgar-Ix-Ieni, Gozo, Malta due to southern winds. Their carcasses, less than 5 cm in length, are consumed by e.g. *Pagurus anachoretus*. Depth about 12 m. Picture taken in October 2002 by R. M. L. Ates.
- H. This mature *Pagurus bernhardus* is halfway through processing an *Aurelia aurita*. Depth about 3 m. Picture taken in June 2003 in Loch Fyne, Scotland by R. M. L. Ates.

In her discussion Arai (2005) expressed the hope that awareness of the predation on jellyfish would lead to more knowledge of their role in pelagic food webs. Likewise I express the hope that this awareness will be extended to the benthic domain where many potential predators and scavengers await live or dead jellyfish coming within their reaches.

### Acknowledgements

I thank Jos Bakker, Rick Harbo, Andy Lamb, Pat Lovejoy, Claudia Mills, Wendy Northway, Ron Offermans, Hans van Rijn and Mat Vestjens for sharing their observations. Additionally I thank Ron Offermans and Hans van Rijn for permission to use their photographs.

### References

- Alamaru A, Bronstein O, Loya Y, Dishon G (2009) Opportunistic feeding by the fungi coral *Fungia scruposa* on the moon jellyfish *Aurelia aurita*. *Coral Reefs* 28: 865.
- Arai MN (1988) Interactions of fish and pelagic coelenterates. *Can J Zool* 66: 1913–1927.
- Arai MN (2005) Predation on pelagic coelenterates: a review. *J Mar Biol Ass UK* 85: 523–536.
- Ates RML (1988) Medusivorous fishes, a review. *Zool Meded Leiden* 62: 29–42.
- Ates RML (1991) Predation on Cnidaria by vertebrates other than fishes. *Hydrobiol* 216/217: 305–307.
- Ates RML (2002) Kwallen als voedsel van zeeanemonen en van stormvogels. *Zeepaard* 62: 132–135.
- Ates RML (2014) Nieuwe waarnemingen van *Eutonina indicans* (Romanes, 1876) in Nederland. *Zeepaard* 74: 157–161.
- Ates RML, Hoyinck SEC (2012) Schijfkwallen in de voedselketen. *Zeepaard* 72: 126–131.
- Ayre DJ (1984) The sea anemone, *Actinia tenebrosa*: an opportunistic insectivore. *Ophelia* 23: 149–153.
- Bamstedt U, Ishii H, Martinussen B (1997) Is the scyphomedusa *Cyanea capillata* (L.) dependent on gelatinous prey for its early development? *Sarsia* 82: 269–273.
- Battaglia P, Musolino S, Esposito V, Ammendolia G, Consolia P, Andaloro F, Romeo T (2014) Feeding habits of juvenile fishes belonging to three medusivorous species (Centrolophidae and Nomeidae) from the Strait of Messina (central Mediterranean Sea). *Mar Biol Res* 10: 927–933.
- Bayha KM, Graham WM, Higgins JE, Fletcher HA (2012) Predation potential of the jellyfish *Drymonema larsoni* Bayha & Dawson (Scyphozoa: Drymonematidae) on the moon jellyfish *Aurelia* sp in the northern Gulf of Mexico. *Hydrobiol* 690: 189–197.
- Bell J (1996) Fulmars feeding on jellyfish. *Brit Birds* 89: 569.
- Berryman J (1984) Predation of *Sagartiogeton laceratus* upon *Aurelia aurita* in shallow water. *J Mar Biol Ass UK* 64: 725.
- Billett DSM, Bett BJ, Jacobs CL, Rouse IP, Wigham BD (2006) Mass deposition of jellyfish in the deep Arabian Sea. *Limnol Oceanogr* 51: 2077–2083.
- Bonaldo RM, Krajewski JP, Sazima I (2004) Does the association of young fishes with jellyfishes protect from predation? A report on a failure case due to damage to the jellyfish. *Neotrop Ichthyol* 2: 103–105.
- Cardona L, Álvarez de Quevedo I, Borrell A, Aguilar A (2012) Massive consumption of gelatinous plankton by Mediterranean apex predators. *PLoS ONE* 7(3): e31329.
- Cargo DG, Schultz LP (1967) Further observations on the biology of the sea nettle and jellyfishes in Chesapeake Bay. *Chesap Sci* 8(4): 209–220.
- Chaves LCT, Feitosa JLL, Pereira PHC (2010) First record of predation on jellyfish by butterflyfish on Brazilian rocky reefs. *Mar Biodiv Rec* 3: 3 pp.
- Cerqueira VR, Haimovici M (1990) Dinamica populacional do gordinho, *Peprilus parva*, no litoral sul do Brasil. *Rev Brasil Biol* 50: 599–613.
- Condon RH, Steinberg DK, Giorgio PA Del, Bouvier TC, Bronk DA, Graham WM, Ducklow HW (2011) Jellyfish blooms result in a major microbial respiratory sink of carbon in marine systems. *Proc Nat Acad Sci USA* 108: 10225–10230.
- Cones HN, Haven DS (1969) Distribution of *Chrysaora quinquecirrha* in the York River. *Chesap Sci* 10: 75–84.
- Conniff R (2000) Jelly bellies. *Nat Geogr Mag* 197(6): 82–101.
- Corsi F (2000) Inconsueta alimentazione del gabbiano reale mediterraneo *Larus cachinnans michahelis*. *Picus* 27: 99.
- Dathe H (1989) Zur Ernährung heimischer Möwen. *Ornith Mitt* 41: 15.
- Dayton PK, Robilliard GA, Paine RT, Dayton LB (1974) Biological accommodation in the benthic community at McMurdo Sound, Antarctica. *Ecol Monogr* 44: 105–128.
- Dias RM, Almeida DF (2015) First record of predation on jellyfish by an aggregation of French angelfish *Pomacanthus paru* on Brazilian rocky reefs. *Mar Biodiv Rec* 8: 3 pp.
- Esser M, Greve W, Boersma M (2004) Effects of temperature and the presence of benthic predators on the vertical distribution of the ctenophore *Pleurobrachia pileus*. *Mar Biol* 145: 595–601.
- Farr JA (1978) Blue crab predation on jellyfish. *Florida Sci* 41: 217–218.
- Fautin DG (1982) *Paraphelliactis pabista*, a new species of hormathiid sea anemone from abyssal northeastern Pacific waters. *Syesis* 15: 51–56.
- Fautin DG, Fitt WK (1991) A jellyfish-eating sea anemone from Palau: *Entacmaea medusivora*. *Hydrobiol* 216/217: 453–461.
- Francis L (1973) Intraspecific aggression and its effect on the distribution of *Anthopleura elegantissima* and some related sea anemones. *Biol Bull* 144: 73–92.
- Gili JM, Hughes RG, Alva V (1996) A quantitative study of feeding by the hydroid *Tubularia larynx*. *Sci Mar* 60: 43–54.
- Gorelova T, Grudtsev ME (1986). Feeding of flying fishes in the Atlantic ocean. *Okeanologiya* 27: 480–483.
- Greve W (1972) Ökologische Untersuchungen an *Pleurobrachia pileus*. *Helgol wiss Meeresunters* 23: 141–164.
- Gronert R (1992) Soms is een kwal om op te vreten. *Windbreker* 52: 9 & 53: 19.
- Hall H (1992) High seas' drifter. *Nat Hist* 192(12): 26–29.
- Hamner WM (1982) Strange world of Palau's salt lakes. *Nat Geogr Mag* 161: 264–282.
- Hamner WM, Hauri IR (1981) Long-distance horizontal migrations of zooplankton (Scyphomedusae: *Mastigias*). *Limnol Oceanogr* 26: 414–423.
- Harbison GR (1993) The potential of fishes for the control of gelatinous zooplankton. *ICES (Int Counc Explor Sea) CMS* (1993/L:74): 1–10.
- Harbo RM (2011) *Whelks to whales: coastal marine life of the Pacific Northwest*. Harbour Publishing Co Ltd, Madeira Park, BC, Canada, 328 pp.
- Hartog JC den (1963) De verspreiding van *Tealia felina*, de zeedahlia, in Nederland. *Zeepaard* 23: 75–82.
- Hoeksema BW, Tuti Y, Becking LE (2015) Mixed medusivory by the sea anemone *Entacmaea medusivora* (Anthozoa: Actiniaria) in Kakaban Lake, Indonesia. *Mar Biodivers* 45: 141–142.
- Jarman SN, McInnes J, Faux C, Polanowski AM, Marthick J, Deagle BE, Southwell C, Emmerson L (2013) Adélie penguin population diet monitoring by analysis of food DNA in scats. *PLoS ONE* 8(12): 11 pp.
- Jarms G, Tiemann H (2004) *Actinostola callosa*, a medusivorous sea anemone and its mass occurrence in the Lurefjord, Norway. *Helgol Mar Res* 58: 15–17.
- Jones TT, Seminoff JA (2013) Feeding biology, Advances from field-based observations, physiological studies, and molecular techniques. In: *The Biology of Sea Turtles III* (eds Wyneken J, Lohmann KJ,

- Musick JA). CRC Press, Boca Raton London New York, pp. 211–247.
- Kelley DF (1987) Food of bass in U.K. waters. *J Mar Biol Ass UK* 67: 275–286.
- Larson RJ (1991) Why jellyfish stick together. *Nat Hist* 100: 66–71.
- Lauckner G (1980) Diseases of Cnidaria. In: *Diseases of Marine Animals* (ed Kinne O). John Wiley & Sons, Chichester New York Brisbane Toronto, pp. 167–237.
- Lebrato M, Pitt KA, Sweetman KA, Jones DOB, Cartes JE, Oschlies A, Condon RH, Molinero JC, Adler L, Gaillard C, Loris D, Billett DSM (2012) Jelly-falls historic and recent observations: a review to drive future research directions. *Hydrobiol* 690: 227–245.
- Luttenberger F (1981) Nahrungsgewohnheiten einiger Fische in Korallenriffen des Roten Meeres. *Aquar Terr Zeitschr (DATZ)* 34(4): 131–134.
- Mackie GO (1966) Growth of the hydroid *Tubularia* in culture. In: *The Cnidaria and Their Evolution* (ed Rees WJ). Academic Press, New York, pp. 397–410.
- Massuti E, Deudero S, Sanchez P, Morales-Nin B (1998) Diet and feeding of dolphin (*Coryphaena hippurus*) in western Mediterranean waters. *Bull Mar Sci* 63: 329–341.
- Melville DS (2013) Variable (*Haematopus unicolor*) and pied oystercatchers (*H. finschi*) feeding on lion's mane jellyfish (*Cyanea* sp.). *Notorchis* 60: 309–310.
- Milisenda G, Rosa S, Fuentes VL, Boero F, Guglielmo L, Purcell JE, Piraino S (2014) Jellyfish as prey: frequency of predation and selective foraging of *Boops boops* (Vertebrata, Actinopterygii) on the mauve stinger *Pelagia noctiluca* (Cnidaria, Scyphozoa). *PLoS ONE* 9(4): 10 pp.
- Miyake H, Lindsay DJ, Hunt JC, Hamatsu T (2002) Scyphomedusa *Aurelia limbata* (Brandt, 1838) found in deep waters off Kushiro, Hokkaido, northern Japan. *Plankton Biol Ecol* 49: 44–46.
- Moen FE, Svensen E (2000) Dyreliv i havet, handbok i Norsk marin fauna. KOM Forlag, Kristiansand, 2nd ed, pp. 1–608.
- Naylor P (2011) Great British Marine Animals. Sound Diving Publ, Plymouth, UK, 3rd ed, 320 pp.
- Orsi Relini L, Garibaldi F, Lanteri L, Relini M (2010a) Medusivorous fishes of the Ligurian sea I. Chub mackerels and other pelagic fish species sometimes have the medusa *Pelagia noctiluca*. *Rapp Comm int Mer Médit* 39: 62.
- Orsi Relini L, Lanteri L, Garibaldi F (2010b) Medusivorous fishes of the Mediterranean. A coastal safety system against jellyfish blooms. *Biol Mar Mediterr* 17: 348–349.
- Peglow HG (1998) Quallengonaden als Silbermöwennahrung. *Ornith Mitt* 50: 124.
- Phillips PJ, Burke WD, Keener EJ (1969) Observations on the trophic significance of jellyfishes in Mississippi Sound with quantitative data on the associative behavior of small fish. *Trans Am Fish Soc* 1969(4): 703–712.
- Powell EH, Gunter G (1968) Observations on the stone crab *Menippe mercenaria* in the vicinity of Port Aransas, Texas. *Gulf Res Rep* 2: 285–299.
- Purcell JE (1991) A review of cnidarians and ctenophores feeding on competitors in the plankton. *Hydrobiol* 216/217: 335–342.
- Purcell JE, Arai MN (2001) Interactions of pelagic cnidarians and ctenophores with fish: a review. *Hydrobiol* 451: 27–44.
- Quilter CG (1987) Foraging activity of the sand beach isopod *Scyphax ornatus*. *New Zeal J Zool* 14: 433–439.
- Rainer SF (1992) Diet of prawns from the continental slope of north-western Australia. *Bull Mar Sci* 50: 258–274.
- Rasmussen E (1973) Systematics and ecology of the Isefjord marine fauna. *Ophelia* 11: 1–495.
- Redeke HC (1911) Natuurlijke historie onzer zeevisschen een handleiding ten gebruike bij het visscherij-onderwijs. C. de Boer, Den Helder, Nederland, 92 pp.
- Sebens KP, Koehl MAR (1984) Predation on zooplankton by the benthic anthozoans *Alcyonium siderium* and *Metridium senile* in the New England subtidal. *Mar Biol* 81: 255–271.
- Shiomi K, Ogi H (1992) Feeding ecology and body size dependence on diet of the sooty shearwater, *Puffinus griseus*, in the north Pacific. *Proc Nat Inst Polar Res Symp on Polar Biol* 5: 105–113.
- Steinich M (1973) Unterwasser-Blitztechnik. *Foto-Magazin* 25: 38–39.
- Stoner EW, Layman CA (2015) Bristle worms attack: benthic jellyfish are not trophic dead ends. *Front Ecol Env* 13: 226–227.
- Suazo CG (2008) Black-browed albatross foraging on jellyfish prey in the southeast Pacific coast, southern Chile. *Polar Biol* 31: 755–757.
- Sweetman AK, Smith CR, Dale T, Jones DOB (2014) Rapid scavenging of jellyfish carcasses reveals the importance of gelatinous material to deep-sea food webs. *Proc R Soc B* 281: 1–8.
- Swennen C (1956) Kwallen in gevangenschap. *Zeepaard* 16: 35–39.
- Thaler E (2012) Eine Qualle für alle! Ungewöhnliche Tischgesellschaften im Meer. *Koralle* 13(4): 78–81.
- Thomas LR (1963) Phyllosoma larvae associated with medusae. *Nature* 198: 208.
- Vlugt PJ van der (1981) *Actinoporus elegans*. *Zeeaquarium* 31(5): 5–9.
- Wertheim A (1984) The Intertidal Wilderness. Sierra Club Books, San Francisco, 156 pp.
- Wood C (2005) Seasearch Guide to Sea Anemones and Corals of Britain and Ireland. Marine Conservation Society, Ross-on-Wye, 128 pp.
- Wood C (2013) Sea Anemones and Corals of Britain and Ireland. Wild Nature Press, Plymouth, UK, 160 pp.
- Wu N (1998) Splendors of the Seas: The Photographs of Norbert Wu, Konemann UK Ltd, London, 252 pp.
- Yamamoto J, Hirose M, Ohtani T, Sugimoto K, Hirase K, Shimamoto N, Shimura T, Honda N, Fujimori Y, Mukai T (2008) Transportation of organic matter to the sea floor by carrion falls of the giant jellyfish *Nemopilema nomurai* in the Sea of Japan. *Mar Biol* 153: 311–317.
- Zann LP (1988) Marine Community Aquarium. TFH Publications, Neptune City, NJ 07783, 416 pp.
- Zonfrillo B (1997) Fulmars feeding on jellyfish and their parasites. *Brit Birds* 90: 293.