

## On a new species of deep-water Siphonophora, *Lensia multicristoides* sp. nov. from middle South China Sea

Zhang Jinbiao<sup>1</sup> and Lin Mao<sup>1</sup>

(Received December 6, 1986; accepted March 21, 1987)

**Abstract**—In this paper, a new species of deep-water Siphonophora, *Lensia multicristoides* sp. nov., collected from the middle South China Sea is described and compared with some similar species in the genus.

### INTRODUCTION

In the course of a detailed and systematic study of the deep-water Siphonophora collected from the water layers below 200 m in the middle South China Sea during the periods of April and December, 1984, a new deep-water Siphonophora, *Lensia* was found and has been named as *L. multicristoides* sp. nov.

*Lensia multicristoides* sp. nov. (Fig. 1)

*Type of specimens.* Holotype, one specimen. Paratype, four specimens (see Table 1).

Table 1. Type specimens

	Number	Date	Location	Depth (m)
Holotype	TIO-SC 021	Apr. 22, 1984	18°16.0'N, 113°57.6'E	4000-200
Paratype	TIO-SC 022	Apr. 22, 1984	18°16.0'N, 113°57.6'E	4000-200
	TIO-SC 023	Apr. 22, 1984	13°29.0'N, 115°2.8'E	1010-470
	TIO-SC 024	Apr. 28, 1984	18°0.6'N, 115°0.2'E	1017-526
	TIO-SC 025	Dec. 31, 1984	12°0.4'N, 115°0.1'E	1000-500

*Anterior nectophore.* 9.5-12.5 mm high, 3.5-5.2 mm wide. Long conical shaped. Seven longitudinal ridges. One dorsal and two ventral ridges, complete. Two ventro-laterals arrive at neither the apex nor the basal margin, but the terminate is at the level of the ostium. Two dorsal-laterals arrive at the apex but the terminates are above the ostium. The somatocyst, spindle-shaped, with a short stalk and usually with an oil drop, is a bit shorter than 1/3 of the length of the nectophore. Hydroecial cavity is at a moderate depth in comparison with others of *Lensia*, its top being nearly at the level of the ostium. Mouth-plate, in two halves, are larger.

1, Third Institute of Oceanography, State Oceanic Administration, Xiamen, China




Fig. 1. *Lensia multicristoides* sp. nov.




Fig. 2. Diagram of *Lensia multicristoides* sp. nov. with some similar species.

Posterior nectophore and eudoxid phase: unknown.

**Discussion.** There are 33 species (including subspecies) in the genus of *Lensia* (Alvarino and Wojtan, 1984; Bigelow, 1911; Daniel and Daniel, 1963; Leloup, 1934; Lens and Van Riemsdijk, 1908; Margulis, 1971; Rengarjan, 1973; Stepanyants, 1967; Stepanyants, 1970; Stepanyants, 1977; Totton, 1941; Totton, 1954; Totton, 1965; Totton and Bargmann, 1965), of which 15 have been described by Chinese researchers and 12 out of 15 were found in China Seas (Zhang, 1984; Zhang and Xu, 1980; Gao, 1982). This new species is similar in having seven ridges to *L. havock* Totton (Totton, 1941); *L. eugenioi* Alvarino et Wojtan (Alvarino et al., 1984); *L. landrumae* Alvarino et Wojtan (Alvarino et al., 1984); *L. hunter* Totton (Totton, 1941); *L. multicristata* (Moser) (Zhang, 1984), but differs from the latter in the characteristics as shown in Table 2 and Fig. 2.

**Distribution.** Found only in waters below 200 m in the middle South China Sea (12°–18°N, 114°–117°E).

Table 2. Comparison between *L. multicristoides* sp. nov. and some other similar species

Species	Dorsal-laterals	Ventro-laterals	Somatocyst
<i>L. havock</i> Totton 1941	complete, terminates arrive at ostium at 1/3 of dorsal ridge	complete, arrive at both apex and basal margin	spindle-shaped, shorter, about 1/3 of nectophore in length, with a stalk undistinguished
<i>L. landrumae</i> Alvarino et Wojtan, 1984	complete, arrive at ostium at 1/3 of dorsal ridge	incomplete, without reaching apex but bending to ventral side in ostial region onto basal margin	large, massive and globular-oval in shape, about 2/5 of nectophore in length
<i>L. eugenioi</i> Alvarino et Wojtan, 1984	complete, arrive at ostium at dorsal ridge	incomplete, arrive at apex but not ostium	short, pear-shaped, about 1/7 of nectophore in length
<i>L. hunter</i> Totton 1941	incomplete, not reach ostium	incomplete, without reaching apex but reaching basal margin	moderate, bilobed and asymmetrical, left side longer than the right, about 1/4 of nectophore in length
<i>L. multicristata</i> (Moser, 1925)	incomplete, not reach ostium	incomplete, reach neither apex nor basal margin	linear-shaped, about 1/2 of nectophore in length, with a stalk
<i>L. multicristoides</i> sp. nov.	incomplete, not reach ostium	incomplete, reach neither apex nor basal margin, but at the level of ostium	spindle-shaped, a little shorter than 1/3 of nectophore in length, with a distinct stalk

## REFERENCES

- Alvarino A. and J.M. Wojtan (1984) Three new species of *Lensia*, and description of the Eudoxia stages of *Lensia reticulata* and *Lensia lelouvetau* (Calycophorae: Siphonophorae). *Proc. Biol. Soc. Wash.*, **97**, 49-59.
- Bigelow H.B. (1911) The Siphonophorae, Rep. Sci. Res. Eastern Tropical Pacific Expedition Albatross. *Mem. Mus. Comp. Zool. Harvard College*, **38**, 173-401.
- Daniel A. and R. Daniel (1963) A new Siphonophore of the genus *Lensia* from the Bay of Bengal. *Ann. Mag. Nat. Hist.*, **13**, 621-623.
- Gao S.W. (1982) The medusae of the East China Sea. *Studia Marina Sinica* (in Chinese), No. 19, 100-108.
- Leloup E. (1934) Calycophorae (Siphonophora) of the tropical Atlantic Ocean and Australia. *Bull. Mus. R. Hist. Nat. Berlin* (in German), **10**, 1-87.
- Lens A. D. and T. van Riemsdijk (1908) The Siphonophora of the Siboga Expedition. *Siboga Exped., Monog.*, **9**, 1-130.
- Margulis R. J. (1971) Some data of distribution of the Siphonophora, genes *Lensia* (Sub-Order Calycophorae) in the Atlantic. *Oceanologia* (in Russian), **11**, 99-104.
- Rengarjan K. (1973) Siphonophores obtained during the cruises of "R.V. Varuna" from the west coast of India and the Laccadive Sea. *J. Mar. Biol. Ass. India*, **15**, 125-156.
- Stepanyants S. D. (1967) Siphonophora of the seas of the USSR and northwestern part of the Pacific Ocean. *Opredelitelipo faune USSR* (in Russian), **96**, 1-216.
- Stepanyants S. D. (1970) Siphonophores of the southern part of the Kurile-Kamchka Trench and the adjacent area. *Trudy Institute Oceanologia, Academiya Nauk, USSR* (in Russian), **96**, 222-235.
- Stepanyants S. D. (1977) Siphonophora of the central part of the Pacific Ocean. *Proceeding of Marine Fauna* (in Russian), **20**, 54-81.
- Totton A.K. (1941) New species of Siphonophora genus *Lensia* Totton 1932. *Ann. Mag. Nat. Hist.*, Ser 11, **8**, 145-168.
- Totton A. K. (1954) Siphonophora of the Indian Ocean together with systematic and biological notes on related specimens from other ocean. *Discovery Rep.*, **27**, 1-216.

- Totton A. K. (1965) A new species of *Lensia* (Siphonophora: Diphyidae) from the coastal waters of Vancouver, B.C. ; and its comparison with *Lensia achilles* Totton and another new species *Lensia cordata*. *Ann. Mag. Nat. Hist. Ser. 13*, **8**, 71-76.
- Totton A. K. and H.E. Bargmann (1965) A synopsis of the Siphonophora. *Transactions of the British Museum*, London, 1-230.
- Zhang J. B. (1984) The Calycophorae (Siphonophora) from tropical waters of the western Pacific Ocean. *Proceedings of the plankton from the tropical waters of the western Pacific Ocean*, Third Institute of Oceanography, NBO editor, China Ocean Press (in Chinese), Beijing, pp. 52-85.
- Zhang J. B. and Z.Z. Xu (1980) On the geographical distribution of the Siphonophores in the China Seas. *Universitatis Amoiensis Acta Scientiarum Naturalium* (in Chinese), **19**, 100-108.