




SHORT REPORT

Bibliography of Francesc Pagès (1986–2007)

COMPILED BY MARSH J. YOUNGBLUTH



Dr Francesc Pagès Hernández (Barcelona, 6 July 1962 – Palma de Mallorca, 5 May 2007).

References

- Riera T, Gili JM, Pagès F. 1986. Estudio cuantitativo y estacional de dos poblaciones de Cnidarios planctónicos frente de las costas de Barcelona (Mediterráneo Occidental): Ciclo entre 1966–67 y 1982–83. *Miscellanea Zoologica* 10:23–32.
- Gili JM, Pagès F. 1987. Pennatuláceos (Cnidaria, Anthozoa) recolectados en la plataforma continental Catalana (Mediterráneo occidental). *Miscellanea Zoológica* 11:25–39.
- Gili JM, Pagès F, Barangé M. 1987. Zoantarios (Cnidaria, Anthozoa) de la costa y de la plataforma continental catalanas (Mediterráneo occidental). *Miscellanea Zoológica* 11:13–24.
- Gili JM, Pagès F, Riera T. 1987. Distribución de las especies más frecuentes de sifonóforos calicóforos en la zona norte del Mediterráneo occidental. *Investigación Pesquera* 51:323–38.
- Gili JM, Pagès F, Vives F. 1987. Distribution and ecology of a population of planktonic cnidarians in the western Mediterranean. In: Bouillon J, Boero F, Cicogna F, Cornelius PFS, editors. *Modern Trends in the Systematics, Ecology and Evolution of Hydroids and Hydromedusae*. Oxford: Oxford University Press. p 157–70.
- Gili JM, Pagès F, Sabatès A, Ros JD. 1988. Small-scale distribution of a cnidarian population in the western Mediterranean. *Journal of Plankton Research* 10(3):385–401.
- Gili JM, Vervoort W, Pagès F. 1989. Hydroids from the West African coast: Guinea Bissau, Namibia and South Africa. *Scientia Marina* 53(1):67–112.
- Pagès F, Gili JM. 1989. Siphonophores (Cnidaria, Hydrozoa) collected during the “Magga Dan” Expedition (1966–67) from Africa to Antarctica. *Scientia Marina* 53(1):53–7.
- Pagès F, Gili JM, Bouillon J. 1989. The Siphonophores (Cnidaria, Hydrozoa) of Hansa Bay, Papua New Guinea. *Indo-Malayan Zoology* 6:133–40.
- Sabatès A, Gili JM, Pagès F. 1989. Relationship between zooplankton distribution, geographic characteristics and hydrographic patterns off the Catalan coast (Western Mediterranean). *Marine Biology* 103(2):153–9.
- Gili JM, Pagès F, Fusté X. 1991. Mesoscale coupling between spatial distribution of planktonic cnidarians and hydrographic features along the Galician coast (Northwestern Iberian Peninsula). *Scientia Marina* 55(2):419–26.
- Pagès F, Bouillon J, Gili JM. 1991. Four new species of Hydromedusae (Cnidaria, Hydrozoa) from the coast of south-western Africa. *Zoologica Scripta* 20(2):89–98.
- Pagès F, Verheye HM, Gili JM, Flos J. 1991. Short-term effects of coastal upwelling and wind reversals on epipelagic cnidarians in the southern Benguela ecosystem. *South African Journal of Marine Science* 10:203–11.
- Pagès F, Gili JM. 1991. Effects of large-scale advective processes on gelatinous zooplankton populations in the northern Benguela ecosystem. *Marine Ecology Progress Series* 75:205–15.
- Pagès F, Gili JM. 1991. Vertical distribution of epipelagic siphonophores at the confluence between Benguela waters and the Angola Current over 48 hours. *Hydrobiologia* 216/217(1):355–62.
- Pagès F, Gili JM. 1992. Influence of Agulhas waters on the population structure of planktonic cnidarians in the southern Benguela region. *Scientia Marina* 56(2):109–23.
- Pagès F, Gili JM. 1992. Siphonophores (Cnidaria, Hydrozoa) of the Benguela Current (southeastern Atlantic). *Scientia Marina* 56(1):65–112.
- Pagès F, Gili JM, Bouillon J. 1992. Medusae (Hydrozoa, Scyphozoa, Cubozoa) of the Benguela Current (southeastern Atlantic). *Scientia Marina* 56(1):1–64.
- Pagès F, Gili JM. 1992. Planktonic cnidarians of the Benguela Current: station data. *Scientia Marina* 56(1):113–44.
- Pagès F. 1992. Mesoscale coupling between planktonic cnidarian distribution and water masses during a temporal transition between active upwelling and abatement in the northern Benguela system. *South African Journal of Marine Science* 12(1):41–52.

- Pagès F, Gili JM. 1992. Influence of the thermocline on the vertical migration of medusae during a 48 h sampling period. *South African Journal of Zoology* 27(2):50–9.
- Pugh PR, Pagès F. 1993. A new species of *Clausophyes* (Siphonophora, Clausophyidae) with the redescription of *C. moserae* and *C. galeata*. *Journal of the Marine Biological Association of the UK* 73(3):595–608.
- Schiel S, Hagen W, Klauke U, Mizdalski E, Pagès F, Probyn T, et al. 1993. Zooplankton investigations. *Berichte zur Polarforschung* 121:75–86.
- Pagès F, Pugh PR, Gili JM. 1994. Macro- and megaplanktonic cnidarians collected in the eastern part of the Weddell Gyre during summer 1979. *Journal of the Marine Biological Association of the UK* 74(4):873–94.
- Pagès F, Kurbjeweit F. 1994. Vertical distribution and abundance of mesoplanktonic medusae and siphonophores from the Weddell Sea, Antarctica. *Polar Biology* 14(4):243–51.
- Carré D, Carré C, Gili JM, Pagès F. 1995. Asexual reproduction in the pelagic phase of *Clytia mccradyi* (Hydrozoa, Leptomedusae). *Scientia Marina* 59(2):193–202.
- Pugh PR, Pagès F. 1995. Is *Lensia reticulata* a diphyine species (Siphonophorae, Calyphorae, Diphyidae)? A re-description. *Scientia Marina* 59(2):181–92.
- Gili JM, Alvà V, Pagès F, Kloser H, Arntz WE. 1996. Benthic diatoms as the major food source in the sub-Antarctic marine hydroid *Silicularia rosea*. *Polar Biology* 16:507–12.
- Pagès F, González HE. 1996. Potential predatory impact of gelatinous zooplankton in Hardangerfjord (Norway). *Marine Ecology Progress Series* 139(1):69–77.
- Pagès F, González HE, González SR. 1996. Diet of the gelatinous zooplankton in Hardangerfjord (Norway) and potential predatory impact by *Aglantha digitale* (Trachymedusae). *Marine Ecology Progress Series* 139(1):69–77.
- Pagès F, Schnack-Schiel SB. 1996. Distribution patterns of the mesozooplankton, principally siphonophores and medusae, in the vicinity of the Antarctic Slope Front (Eastern Weddell Sea). *Journal of Marine Systems* 9(3–4):231–48.
- Pagès F, White MG, Rodhouse PG. 1996. Abundance of gelatinous carnivores in the nekton community of the Antarctic Polar Frontal Zone in summer 1994. *Marine Ecology Progress Series* 141(1):139–47.
- Gili JM, Bouillon J, Pagès F, Palanques A, Puig P, Heussner S. 1997. Submarine canyons as habitat of singular plankton populations in the Mediterranean: an ecological approach. *Mediterranean Marine Biodiversity Workshop*. Nicosia, (Cyprus). p 27–31.
- Jarre-Teichmann A, Brey T, Bathmann U, Dahm C, Dieckmann G, Gorny M, et al. 1997. Trophic flows in the benthic shelf community of the eastern Weddell Sea. In: Battaglia B, Valencia J, Walton D, editors. *Antarctic Communities: Species, Structure and Survival*. Cambridge: Cambridge University Press. p 118–34.
- Pagès F. 1997. The gelatinous zooplankton in the pelagic system of the Southern Ocean: a review. *Annales de l'Institut océanographique Paris* 73:139–58.
- Pagès F, Bouillon J. 1997. A redescription of *Paragotoea bathybia* Kramp 1942 (Hydroidomedusae: Corymorphidae) with a new diagnosis for the genus *Paragotoea*. *Scientia Marina* 61(4):487–93.
- Pugh PR, Pagès F, Boorman B. 1997. Vertical distribution and abundance of pelagic cnidarians in the eastern Weddell Sea, Antarctica. *Journal of the Marine Biological Association of the UK* 77(2):341–60.
- Pugh PR, Pagès F. 1997. A re-description of *Lensia asymmetrica* Stepanjants, 1970 (Siphonophorae, Diphyidae). *Scientia Marina* 61(2):153–61.
- Gili JM, Alvà V, Coma R, Orejas C, Pagès F, Ribes M, et al. 1998. The impact of the small benthic suspension feeders in shallow marine ecosystems: the hydroids as an example. *Zoologische Verhandlungen* 323:1–7.
- Gili JM, Bouillon J, Pagès F. 1998. A new species of *Krampella* (Hydrozoa, Hydroidomedusae, Tiarranidae) from the deep waters of Antikythira Strait (Cretan Sea, North East Mediterranean). *Scientia Marina* 62(1–2):135–9.
- Gili JM, Bouillon J, Pagès F, Palanques A, Puig P, Heussner S. 1998. Origin and biogeography of the deep-water Mediterranean Hydromedusae including the description of two new species collected in submarine canyons of Northwestern Mediterranean. *Scientia Marina* 62(1–2):113–34.
- Gili JM, Pagès F. 1998. Les meduses de les nostres costes. *Barcelona Verda* 59:22–3.
- Pagès F, Orejas C. 1999. Medusae, siphonophores and ctenophores of the Magellan region. *Scientia Marina* 63(1):51–7.
- Gili JM, Bouillon J, Pagès F, Palanques A, Puig P. 1999. Submarine canyons as habitats of prolific plankton populations: three new deep-sea Hydroidomedusae in the western Mediterranean. *Zoological Journal of the Linnean Society* 125:313–29.
- Pagès F, Pugh PR, Siegel V. 1999. The discovery of an Antarctic epipelagic medusan in the Mediterranean. *Journal of Plankton Research* 21(12):2431–5.
- Bouillon J, Gili JM, Pagès F, Isla E. 2000. *Amphinema modernisme*, a new Pandeid (Cnidaria: Anthomedusae) from the Southern Ocean. *Polar Biology* 23(1):34–7.
- Bouillon J, Pagès F, Gili JM, Palanques A, Puig P, Heussner S. 2000. Deep-water Hydromedusae from the Lacaze-Duthiers submarine canyon (Banyuls, Northwestern Mediterranean) and description of two new genera, *Guillea* and *Paratecladia*. *Scientia Marina* 64(1):87–95.
- Pagès F. 2000. Biological associations between barnacles and jellyfish with emphasis on the ectoparasitism of *Alepa pacifica* (Lepadomorpha) on *Diplumaris malayensis* (Scyphozoa). *Journal of Natural History* 34:2045–56.
- Gili JM, Pagès F, Bouillon J, Palanques A, Puig P, Heussner S, et al. 2000. A multidisciplinary approach to the understanding of hydromedusan populations inhabiting Mediterranean submarine canyons. *Deep Sea Research I* 47:1513–33.
- Bouillon J, Pagès F, Gili JM. 2001. New species of benthopelagic hydromedusae from the Weddell Sea. *Polar Biology* 24(11):839–45.
- Gili JM, Alfonso I, Gasol JM, Isla E, López-González P, Orejas C, et al. 2001. Pelagobenthic coupling and the role of benthic suspension feeders. *Berichte zur Polar- und Meeresforschung* 402:10–56.
- Gili JM, Pagès F, Font J. 2001. La superficie del mar labrada. *Investigación y Ciencia* 293:40–1.
- Graham WM, Pagès F, Hamner WM. 2001. A physical context for gelatinous zooplankton aggregations: a review. *Hydrobiologia* 451(1–3):199–212.
- Pagès F. 2001. Past and present anthropogenic factors promoting the invasion, colonization and dominance by jellyfish of a Spanish coastal lagoon. *CIESM Workshop Series* 14:69–71.
- Pagès F, González HE, Ramón M, Sobarzo M, Gili JM. 2001. Gelatinous zooplankton assemblages associated with water masses in the Humboldt Current System, and potential predatory impact by *Bassia bassensis* (Siphonophora: Calyphorae). *Marine Ecology Progress Series* 210:13–24.
- Pagès F. 2002. The status of three rare siphonophores (Cnidaria, Hydrozoa) described by Tamiji Kawamura: *Bathyphysa japonica*, *Athorybia longifolia* and *Forskalia misakiensis*. *Scientia Marina* 66(4):375–82.
- Pagès F, Pugh PR. 2002. Fuseudoxid: the elusive sexual stage of the calyphoran siphonophore *Crystalliphyes amygdalina*

- (Clausophyidae: Crystallophyinae). *Acta Zoologica* 83(4): 329–36.
- Gili JM, Rossi S, Pagès F, Teixidó N, Orejas C, López P. 2002. Una cadena trófica inesperada. *Investigación y Ciencia* 306:36–7.
- Maso M, Garces E, Pages F, Camp J. 2003. Drifting plastic debris as a potential vector for dispersing harmful algal bloom (HAB) species. *Scientia Marina* 67(1):107–11.
- Pagès F, Corbera J, Gili JM. 2003. Il·lustracions de meduses, cinc segles d'història. Mètode: Revista de difusió de la investigació de la Universitat de València 38:15–8.
- Bouillon J, Medel MD, Pagès F, Gili JM, Boero F, Gravili C. 2004. Fauna of the Mediterranean Hydrozoa. *Scientia Marina* 68(Suppl. 2):1–438.
- González HE, Giesecke R, Vargas CA, Pavez M, Iriarte J, Santibáñez P, et al. 2004. Carbon cycling through the pelagic food web in the northern Humboldt Current off Chile (23°S). *ICES Journal of Marine Science* 61(4):572–84.
- Medel MD, Pagès F, Bouillon J. 2004. Fauna of the Mediterranean Hydrozoa: introduction. *Scientia Marina* 68(2):5–15.
- Palanques A, García-Ladona E, Gomis D, Martín J, Marcos M, Pascual A, et al. 2005. General patterns of circulation, sediment fluxes and ecology of the Palamós (La Fonera) submarine canyon, northwestern Mediterranean. *Progress in Oceanography* 66(2–4):89–119.
- Uiblein F, Youngbluth M, Jacoby C, Pagès F, Picheral M, Gorsky G. 2005. In situ observations of deep-water fishes in four canyons off the Georges Bank, NW Atlantic. *Deep Sea* 2003. Conference on the governance and management of deep-sea fisheries. *FAO Fisheries Proceedings* 1/3:98–106.
- Pagès F, Flood P, Youngbluth M. 2006. Gelatinous zooplankton net-collected in the Gulf of Maine and adjacent submarine canyons: new species, new family (Jeanbouilloniidae), taxonomic remarks and some parasites. *Scientia Marina* 70(3):363–79.
- Fuentes V, Pagès F. 2006. Description of *Jubanyella plemmyris*, gen. nov. et sp. nov. (Hydrozoa: Narcomedusae) stranded off the Jubany Antarctic base, with a new diagnosis of the family Aeginidae. *Journal of Plankton Research* 28:959–63.
- Gili JM, Rossi S, Pagès F, Orejas C, González P, Arntz W. 2006. A new trophic link between the pelagic and benthic systems in the Southern Ocean. *Marine Ecology Progress Series* 322:43–9.
- Pagès F, Corbera J, Lindsay DJ. 2006. Piggybacking pycnogonids and parasitic narcomedusae on *Pandea rubra* (Anthomedusae, Pandeidae). *Plankton and Benthos Research* 1:132–40.
- Pagès F. 2006. L'oceà mesopelàgic vist des d'una bombolla d'aigua. *Omnis Cèl·lula* 11:20–5.
- Bouillon J, Gravili C, Pagès F, Gili JM, Boero F. 2006. An introduction to Hydrozoa. *Mémoires du Muséum National d'Histoire Naturelle Paris* 194:1–591.
- Hosia A, Pagès F. 2007. Unexpected new species of deep-water Hydroidomedusae from Korsfjorden, Norway. *Marine Biology* 151:177–84.

Editorial responsibility: Franz Uiblein