

Journal Entry - Week Five - Entry Four - Draft

Caspian Nicholls, u1027945

March 26, 2020

I give permission for portions of this work to be used as examples in future science communication course notes.

Article title: Search for octupole correlations in neutron-rich ^{148}Ce nucleus

Authors: Y. J. Chen, S. J. Zhu, J. H. Hamilton, A. V. Ramayya, J. K. Hwang, M. Sakhaee, Y. X. Luo, J. O. Rasmussen, K. Li, I. Y. Lee, X. L. Che, H. B. Ding, and M. L. Li

Reference: Phys. Rev. C 73, 054316, May 2006.

Link: [Article abstract](#).

Word count: 736.

Context



Theoretical calculations that make use of accepted models of the structure of nuclei predict that several nuclei with proton numbers Z around 56 and neutron numbers N around 88 should exhibit octupole correlations. These correlations occur most strongly when there is a strong interaction between nuclear orbitals that for principal, orbital angular momentum and total angular momentum quantum numbers n , l and j , have differences $\Delta n = 1$ and $\Delta l = \Delta j = 3$. The correlations themselves occur due to the octupole interaction component of the nuclear Hamiltonian having significant magnitude relative to the other terms in that Hamiltonian. The observation of octupole correlations in nuclei is evidence that the charge distribution of the nucleus may exhibit octupole deformation, so that its shape is no longer spherical. There remain various nuclei near $Z = 56$ and $N = 88$ for which experimental evidence of octupole correlations had not been recorded. Ultimately, observation of these correlations can provide useful information about the energy level structure and shape of these nuclei.

Purpose

In this work, the authors perform experiments to hunt for signs of octupole correlations in ^{148}Ce . This isotope has $Z = 58$ and $N = 90$, so it is one such nucleus that should exhibit this type of correlation. The authors also seek to extend the level scheme of this nucleus.

Approach



Gamma rays that were emitted from the excited primary fission fragments (which included excited forms of ^{148}Ce) of the spontaneous fission of ^{252}Cf were measured to study ^{148}Ce . The researchers analysed groups of three gamma rays that were emitted within a narrow enough time window to be confident that they originated from the same nucleus. They also considered the relative intensities of different gamma ray transitions and the total internal conversion coefficients, so that they could draw solid conclusions from their recorded data.

Contribution

20 new transitions and nine new levels were added to the energy level structure of ^{148}Ce as a result of their analysis. Spins and parities were able to be assigned (tentatively in some cases) for some of the newly discovered states and either confirmed or refined for some other existing levels. Internal conversion coefficients (where a high-energy electron is emitted instead of a gamma ray) for some transitions were also measured.

Analysis of the differences in energies between bands of opposite parity (positive and negative) led the authors to suggest that the octupole correlation they observed in ^{148}Ce has similarities to that observed within the nearby isotopes $^{144,146}\text{Ce}$. However, they found that different bands of energy levels within ^{148}Ce (which relate to energy states of the nucleus that arise when it is rotated, each with different rotational symmetries) may have octupole correlations with different strengths. This conclusion was also supported by further analysis of the relative transition energies and intensities within each rotational band.

Yet, analysis of the moment of inertia as a function of the rotational energy of the nucleus led the authors to suggest that the octupole deformation seen in this nucleus may be unstable, prompting further theoretical study.

Relevance

Following the disruptions caused by the COVID-19 pandemic, we have had to come up with an alternate path for my Honours project to pursue whilst we are not permitted to access ANU's experimental facilities. This back-up plan involves the analysis of data generated from an experiment that was performed with the aim of gauging octupole collectivity in the neutron-rich nuclei $^{150,152}\text{Ce}$, which are located near $Z = 56$ and $N = 88$, but remain unstudied. As such, having an understanding of what octupole correlations are and how they can be investigated is vital.

Quality

This work was clearly written and the analysis was thorough, as the authors describe in detail how they developed their final level scheme via gamma ray spectroscopic analysis of their collected data. I did have to refer to some other sources to clarify parts of the paper that I was not familiar with, but fortunately clear and precise explanations of these details were easy to come by. These authors also provided multiple pieces of evidence for their conclusion that they did observe octupole correlations in ^{148}Ce , making their conclusions trustworthy.

Questions/Directions

After reading this paper, it is clear to me that I need to brush up on my knowledge of the finer details of nuclear structure. The best way to do this would be referring back to the notes I have made when I took Physics of Matter last year. Before I start to analyse the experimental data I referred to in the *Relevance* section, reading more deeply about the theoretical models that predict the existence of these octupole correlations (starting from the references in this paper) would also be worthwhile.