

**DR.D.Y.PATIL INSTITUTE OF TECHNOLOGY,
PIMPRI, PUNE-18**

DEPARTMENT OF MATHEMATICS

Question bank for In semester exam

Semester II (2021-2022)

SUB: ENGINEERING MATHEMATICS III

Unit II

Fourier Transform and Z Transform

SR.NO	QUESTIONS	OPTION
1	<p>Given that $\int_0^\infty \frac{\sin t}{t} dt = \frac{\pi}{2}$ then Fourier sine transform $F_s(\lambda)$, $f(x) = \frac{1}{x}, x > 0$, is given by</p> <p>a) π b) π c) $\frac{\pi}{4}$ d) $\frac{\pi}{2}$</p>	D
2	<p>If $f(x) = \begin{cases} x, & 0 < x < 1 \\ 0, & x > 1 \end{cases}$ then Fourier sine transform $F_s(\lambda)$ of $f(x)$ is given by</p> <p>a) $\frac{\lambda \cos \lambda + \sin \lambda}{\lambda^2}$ b) $\frac{-\lambda \cos \lambda - \sin \lambda}{\lambda^2}$ c) $\frac{-\lambda \cos \lambda + \sin \lambda}{\lambda^2}$ d) $\frac{\cos \lambda}{\lambda^2}$</p>	C
3	<p>The Fourier cosine transform $F_c(\lambda)$ of $f(x) = e^{- x } - \infty < x < \infty$ is</p> <p>a) $\frac{\lambda}{1+\lambda^2}$ b) $\frac{1}{1+\lambda^2}$ c) $\frac{1}{1-\lambda^2}$ d) $\frac{-1}{1+\lambda^2}$</p>	B
4	<p>The Fourier cosine integral representation is</p> $\frac{4}{\pi} \int_0^\infty \frac{\sin \lambda - \lambda \cos \lambda}{\lambda} \cos \lambda x d\lambda = \begin{cases} 1 - x^2, & 0 \leq x \leq 1 \\ 0, & x > 1 \end{cases},$ <p>Then the value of the integral $\int_0^\infty \frac{\sin \lambda - \lambda \cos \lambda}{\lambda} \cos \frac{\lambda}{2} d\lambda$ is equal to,</p> <p>a) $-\frac{3\pi}{16}$ b) $\frac{3\pi}{16}$ c) $-\frac{3\pi}{8}$ d) $\frac{3\pi}{4}$</p>	B
5	<p>Find the Fourier Transform $F(\lambda)$ of $f(x) = \begin{cases} 1, & x < a \\ 0, & x > a \end{cases}$ is</p>	A

	$\text{a) } \frac{2 \sin \lambda a}{\lambda} \quad \text{b) } \frac{e^{-i\lambda a}}{\lambda} \quad \text{c) } \frac{e^{i\lambda a}}{\lambda} \quad \text{d) } \frac{2 \cos \lambda a}{\lambda}$	
6	<p>The Fourier sine transform $F_s(x)$ of $f(x) = \begin{cases} \frac{\pi}{2}, & 0 < x < \pi \\ 0, & x > \pi \end{cases}$ is</p> <p> $\text{a) } \frac{\pi}{2} \left(\frac{1 - \sin \lambda \pi}{\lambda} \right)$ $\text{b) } \frac{\pi}{2} \left(\frac{\cos \lambda \pi - 1}{\lambda} \right)$ $\text{c) } \frac{\pi}{2} \left(\frac{1 - \cos \lambda \pi}{\lambda} \right)$ $\text{d) } \left(\frac{\cos \lambda \pi}{\lambda} \right)$ </p>	C
7	<p>The Fourier sine transform $F_s(\lambda)$ of $f(x) = e^{-x} + e^{-2x}$</p> <p> $\text{a) } \frac{\lambda}{1+\lambda^2} + \frac{\lambda}{4+\lambda^2}$ $\text{b) } \frac{1}{1+\lambda^2} + \frac{1}{2+\lambda^2}$ $\text{c) } \frac{1}{1-\lambda^2} + \frac{1}{1+\lambda^2}$ $\text{d) } \frac{-1}{1+\lambda^2} + \frac{1}{4+\lambda^2}$ </p>	A
8	<p>The Fourier cosine transform $F_c(\lambda)$ of $f(x) = e^{- x } - \infty < x < \infty$ is</p> <p> $\text{a) } \frac{\lambda}{1+\lambda^2}$ $\text{b) } \frac{1}{1+\lambda^2}$ $\text{c) } \frac{1}{1-\lambda^2}$ $\text{d) } \frac{-1}{1+\lambda^2}$ </p>	B
9	<p>The Fourier cosine transform</p> $\int_0^\infty \frac{1 - \cos u}{u^2} \cos \lambda u du = \begin{cases} \frac{\pi}{2} (1 - \lambda), & 0 \leq \lambda \leq 1 \\ 0, & \lambda > 1 \end{cases}$ <p>is,</p> <p>then the value of the integral $\int_0^\infty \frac{\sin^2 z dz}{z^2}$ is</p> <p> $\text{a) } 1$ $\text{b) } \frac{\pi}{2}$ $\text{c) } 0$ $\text{d) } \frac{\pi}{4}$ </p>	B
10	<p>The Fourier Transform $F(\lambda)$ of $f(x) = \begin{cases} e^{-x}, & x > 0 \\ 0, & x < 0 \end{cases}$ is</p> <p> $\text{a) } \frac{1 - \lambda}{1 + \lambda^2}$ $\text{b) } \frac{1 - i\lambda}{1 + \lambda^2}$ $\text{c) } \frac{1 - i\lambda}{1 - \lambda^2}$ $\text{d) } \frac{1}{1 + \lambda^2}$ </p>	B
11	<p>The Fourier sine transform of $f(x) = \begin{cases} 1, & x < a \\ 0, & x > a \end{cases}$ is</p> <p> $\text{a) } \frac{1 - \cos \lambda a}{\lambda}$ $\text{b) } \frac{\sin \lambda a}{\lambda}$ $\text{c) } \frac{\cos \lambda a - 1}{\lambda}$ $\text{d) } \frac{\sin \lambda a}{a}$ </p>	A

12	<p>If the Fourier integral representation of $f(x)$ is</p> $\frac{2}{\pi} \int_0^{\infty} \frac{\sin \lambda \cos \lambda x}{\lambda} d\lambda = \begin{cases} 1, & x < 1 \\ 0, & x > 1 \end{cases}$ <p>then value of the integral $\int_0^{\infty} \frac{\sin \lambda}{\lambda} d\lambda$ is</p> <p>a) $\frac{\pi}{4}$ b) $\frac{\pi}{2}$ c) 0 d) 1</p>	B
13	<p>The Fourier sine representation,</p> $e^{-x} \cos x = \frac{2}{\pi} \int_0^{\infty} \frac{\lambda^3}{\lambda^4 + 4} \sin \lambda x d\lambda, \quad F_s(\lambda) \text{ is,}$ <p>a) $\frac{\lambda^2}{\lambda^4 + 4}$ b) $\frac{\lambda^3}{\lambda^4 + 4}$ c) $\frac{\lambda^4 + 4}{\lambda^3}$ d) $\frac{1}{\lambda^4 + 4}$</p>	B
14	<p>Given that $F_c(\lambda) = \int_0^{\infty} u^{m-1} \cos \lambda u du = \frac{\Gamma(m)}{\lambda^m} \cos \frac{\pi m}{2}$ is,</p> <p>Then the Fourier cosine transform $F_c(\lambda)$ of $f(x) = x^3, x > 0$ is given by,</p> <p>a) $\frac{6}{\lambda^4}$ b) $\frac{3}{\lambda^3}$ c) $\frac{4}{\lambda^2}$ d) $\frac{1}{\lambda^2}$</p>	A
15	<p>The solution of $f(x)$ of the integral equation,</p> $\int_0^{\infty} f(x) \cos \lambda x dx = e^{-\lambda}, \lambda > 0 \text{ is}$ <p>a) $\frac{2}{\pi} \left[\frac{e^{-x}}{(1+x^2)} \right]$ b) $\frac{2}{\pi} \left[\frac{x}{(1+x^2)} \right]$ c) $\frac{2}{\pi} \left[\frac{1}{(1-x^2)} \right]$ d) $\frac{2}{\pi} \left[\frac{1}{(1+x^2)} \right]$</p>	d
16	<p>The solution of $f(x)$ of the integral equation</p> $\int_0^{\infty} f(x) \sin \lambda x dx = \begin{cases} 1, & 0 \leq \lambda \leq 1 \\ 0, & \lambda \geq 1 \end{cases} \text{ is,}$ <p>a) $\frac{2}{\pi} \left[\frac{\cos x}{x} \right]$ b) $\frac{2}{\pi} \left[\frac{1 - \cos x}{x} \right]$ c) $\frac{2}{\pi} \left[\frac{1 + \sin x}{x} \right]$ d) $\frac{2}{\pi} \left[\frac{\sin x}{x} \right]$</p>	b
17	The solution of $f(x)$ of the integral equation	

	$\int_0^{\infty} f(x) \sin \lambda x dx = \begin{cases} 1 - \lambda, & 0 \leq \lambda \leq 1 \\ 0, & \lambda \geq 1 \end{cases} \text{ is,}$ $f(x) = \frac{2}{\pi} \int_0^1 (1 - \lambda) \sin \lambda x d\lambda$ then the value of f(x) is a) $\frac{2}{\pi} \left[\frac{1}{x} - \frac{\sin x}{x^2} \right]$ b) $\frac{2}{\pi} \left[\frac{1}{x} - \frac{\cos x}{x^2} \right]$ c) $\frac{2}{\pi} \left[\frac{1}{x} + \frac{\sin x}{x^2} \right]$ d) $\frac{2}{\pi} \left[-\frac{1}{x} + \frac{\sin x}{x^2} \right]$	
18	The solution of f(x) of the integral equation, $\int_0^{\infty} f(x) \sin \lambda x dx = e^{-\lambda}, \lambda > 0$ is a) $\frac{2}{\pi} \left[\frac{e^{-x}}{(1+x^2)} \right]$ b) $\frac{2}{\pi} \left[\frac{x}{(1+x^2)} \right]$ c) $\frac{2}{\pi} \left[\frac{1}{(1-x^2)} \right]$ d) $\frac{2}{\pi} \left[\frac{1}{(1+x^2)} \right]$	b
19	For the Fourier sine representation $\frac{12}{\pi} \int_0^{\infty} \frac{\lambda \sin \lambda x}{(4 + \lambda^2)(16 + \lambda^2)} d\lambda = e^{-3x} \sinh x,$ $x > 0, F_s(\lambda)$ is a) $\frac{6\lambda}{(4+\lambda^2)(16+\lambda^2)}$ b) $\frac{\lambda}{(4+\lambda^2)(16+\lambda^2)}$ c) $\frac{6\cos \lambda x}{(4+\lambda^2)(16+\lambda^2)}$ d) $\frac{1}{(4+\lambda^2)(16+\lambda^2)}$	A
20	In the Fourier integral representation of $\frac{1}{2\pi} \int_{-\infty}^{\infty} \left(\frac{e^{-i\lambda\pi} + 1}{1 - \lambda^2} \right) e^{i\lambda x} d\lambda = \begin{cases} \sin x, & 0 < x < \pi \\ 0, & x < 0 \text{ and } x > \pi \end{cases}$ $F(\lambda)$ is a) $\frac{1 + \lambda^2}{1 - i\lambda}$ b) $\frac{e^{-i\lambda}}{1 - \lambda^2}$ c) $\frac{e^{-i\lambda\pi} + 1}{1 - \lambda^2}$ d) $\frac{\sin \lambda}{1 - \lambda^2}$	c
Z- TRANSFORM		
1	If $f(k) = \sin \frac{\pi}{2} K, k \geq 0$, then Z transform of $\{\sin \frac{\pi}{2} k\}$ is given by a) $\frac{z}{Z^2 - 1}, z < 1$ b) $\frac{z^2}{Z^2 + 1}, z > 1$ C) $\frac{z}{Z^2 + 1}, z > 1$ D) $\frac{z}{Z^2 - 1}, z > 1$	c

2	<p>. If $f(k)=2^k \sin \frac{\pi}{2} K, k \geq 0$, then Z transform of $\{2^k \sin \frac{\pi}{2} k\}$ is given by</p> <p>a) $\frac{2z}{z^2-4}, z > 2$ b) $\frac{2z}{z^2-4}, z < 2$</p> <p>c) $\frac{2z}{z^2+4}, z > 2$ d) $\frac{2z}{z^2+4}, z < 2$</p>	D
3	<p>If $f(k)=k, k \geq 0$, then Z transform of $\{k\}$ is given by</p> <p>a) $\frac{z}{(z-1)^2}, z > 1$ b) $\frac{(z-1)^2}{z^2}, z > 1$</p> <p>c) $\frac{(z+1)^2}{z^2}, z > 1$ d) $\frac{z^2}{(z+1)^2}, z > 1$</p>	A
4	<p>If $z > 2$, $Z^{-1}\left[\frac{z}{(z-1)(z-2)}\right]$ is given by</p> <p>a) $2^k - 1, k \geq 0$ b) $2^{k+1} - 1, k > 1$</p> <p>c) $\frac{1^k}{2} - 1, k \geq 0$ d) $k - 1, k \geq 0$</p>	A
5	<p>For the difference equation $f(k+1) + \frac{1}{3}f(k) = \left(\frac{1}{3}\right)^k, k \geq 0, f(0) = 0$</p> <p>F(z) is given by</p> <p>a) $\frac{1}{\left(z - \frac{1}{3}\right)\left(z + \frac{1}{3}\right)}$ b) $\frac{z}{\left(z - \frac{1}{3}\right)\left(z + \frac{1}{3}\right)}$</p> <p>c) $\frac{z}{\left(z + \frac{1}{3}\right)\left(z + \frac{1}{2}\right)}$ d) $\frac{z}{\left(z - \frac{1}{3}\right)^2}$</p>	B
6	<p>If $f(k)=\frac{2^k}{k}, k \geq 1$, then Z transform of $\{\frac{2^k}{k}\}, k \geq 1$, is given by</p> <p>a) $-\log(1-2z^{-1})$ b) $\log(1-2z^{-1})$</p> <p>c) $-\log(1+2z^{-1})$ d) $\log(1+2z^{-1})$</p>	A

7	<p>If $f(k)=k5^k, k \geq 0$, then Z transform of $\{k5^k\}$ is given by</p> <p>a) $\frac{(z-5)^2}{5z}, z > 5$ b) $\frac{(z-5)^2}{z}, z > 5$</p> <p>c) $\frac{5z}{(z-5)^2}, z > 5$ d) $\frac{5z}{(z+5)^2}, z > 5$</p>	c
8	<p>Z-transform of $\{3^k e^{-2k}, k \geq 0\}$ is given by</p> <p>a) $\frac{z}{z-3e^2}$ b) $\frac{z}{z-3e^{-2}}$ c) $\frac{z}{z-2e^3}$ d) $\frac{z}{z+3e^2}$</p>	b
9	<p>For finding inverse Z-transform by inversion integral method of $F(z)=\frac{z}{(z-2)(z-3)}$ the residue of $z^{k-1}F(z)$ at the pole $z=2$ is</p> <p>a) $-(2)^k$ b) $(2)^k$ c) $-(3)^k$ d) $(3)^k$</p>	A
10	<p>For finding inverse Z-transform by inversion integral method of $F(z)=\frac{10z}{(z-1)(z-2)}$ the residue of $z^{k-1}F(z)$ at the pole $z=1$ is</p> <p>a) -10 b) 10^{k-1} c) 10 d) 10^k</p>	A
11	<p>If $\{X(k)\}=\{2^k\}*\{3^k\} \quad k \geq 0$ then $Z\{X(k)\}$ is given by</p> <p>a) $\left(\frac{z}{z-2}\right)\left(\frac{z}{z-3}\right), z > 3$ b) $\left(\frac{z}{z-2}\right)\left(\frac{z}{z-3}\right), z < 3$</p> <p>c) $\left(\frac{z}{z-2}\right)\left(\frac{z}{z-3}\right), z < 2$ d) $\left(\frac{z}{z-2}\right)\left(\frac{z}{z-3}\right), z > 2$</p>	A
12	<p>If $z < 1$, $Z^{-1}\left[\frac{z}{(z-1)(z-2)}\right]$ is given by</p> <p>a) $2^k - 1, k \geq 0$ b) $2^{k+1} - 1, k > 1$</p> <p>c) $1 - 2^k, k < 0$ d) $2 - 3^k, k < 0$</p>	c
13	<p>For the difference equation $y_k - 3y_{k-2} = 1, k \geq 0, Y(z)$ is given by</p>	D

	a) $\frac{z}{(z-1)(z^2-3)}$ b) $\frac{1}{(1-3z^2)}$ c) $\frac{z}{(z-1)(1-3z^2)}$ d) $\frac{z^3}{(z-1)(z^2-3)}$	
14	For finding inverse Z-transform by inversion integral method of $F(z) = \frac{z}{\left(z - \frac{1}{2}\right)\left(z - \frac{1}{5}\right)}$ the residue of $z^{k-1}F(z)$ at the pole $z = \frac{1}{2}$ is a) $\frac{10}{3} \left(\frac{1}{2}\right)^k$ b) $10 \left(\frac{1}{2}\right)^k$ c) $-10 \left(\frac{1}{2}\right)^k$ d) $\frac{1}{10} \left(\frac{1}{2}\right)^k$	A
15	If $f(k) = \frac{\sin ak}{k}, k \geq 0$, then Z transform of $\left\{ \frac{\sin ak}{k}, k \geq 0 \right\}$ is given by a) $\cot^{-1}(z + \cos a)$ b) $\cot^{-1}\left(\frac{z + \cos a}{\sin a}\right)$ c) $\cot^{-1}(z - \cos a)$ d) $\cot^{-1}\left(\frac{z - \cos a}{\sin a}\right)$	D
16	If $2 < z < 4$, $Z^{-1}\left[\frac{1}{(z-4)(z-2)}\right]$ is given by a) $-2^{k-1} - 4^{k-1}, k \leq 0, k \geq 1$ b) $2^{k-1} + 4^{k-1}, k \leq 0, k \geq 2$ c) $-2^{k-1} + 4^{k-1}, k \leq 0, k \leq 0$ d) $\frac{1}{2}[-(4)^{k-1} - (2)^{k-1}], k \leq 0, k \geq 1$	D
17	. For finding inverse Z-transform by inversion integral method of $F(z) = \frac{z}{\left(z - \frac{1}{4}\right)\left(z - \frac{1}{3}\right)}$ the residue of $z^{k-1}F(z)$ at the pole $z = \frac{1}{4}$ is a) $-\frac{1}{4} \left(\frac{1}{4}\right)^k$ b) $-12 \left(\frac{1}{4}\right)^k$ c) $-3 \left(\frac{1}{4}\right)^k$ d) $20 \left(\frac{1}{4}\right)^k$	B
18	If $ z > 2, k \geq 0$, inverse Z-transform of $\frac{z^2}{(z-2)^2}$ is given by a) $-(k+1)2^k$ b) $(k+1)2^k$ c) $(k+1)2^{-k}$ d) $(k-1)2^k$	B

19	<p>If $f(k) = \left(\frac{\pi}{2}\right)^k \cos \frac{\pi}{2} K, k \geq 0$, then Z transform of $\left\{\left(\frac{\pi}{2}\right)^k \cos \frac{\pi}{2} k\right\}$ is given by</p> <div style="display: flex; justify-content: space-around; align-items: flex-start;"> <div style="text-align: center;"> <p>a) $\frac{z^2}{Z^2 + \frac{\pi^2}{4}}, z > \frac{\pi}{2}$</p> <p>c) $\frac{z}{Z^2 + \frac{\pi^2}{4}}, z > \frac{\pi}{2}$</p> </div> <div style="text-align: center;"> <p>b) $\frac{z^2}{Z^2 - \frac{\pi^2}{4}}, z < \frac{\pi}{2}$</p> <p>d) $\frac{z}{Z^2 - \frac{\pi^2}{4}}, z > \frac{\pi}{2}$</p> </div> </div>	A
20	<p>. If $1 < z < 3$, $Z^{-1}\left[\frac{z}{(z-1)(z-3)}\right]$ is given by</p> <div style="display: flex; justify-content: space-around; align-items: flex-start;"> <div style="text-align: center;"> <p>a) $1 + 3^k, k > 0$</p> <p>c) $3^k - 1, k < 0$</p> </div> <div style="text-align: center;"> <p>b) $1 + 3^k, k < 0$</p> <p>d) $\frac{1}{2}(-3^k - 1), k < 0, k \geq 0$</p> </div> </div>	D