

Discrete and Algorithmic Geometry

Julian Pfeifle, UPC, 2014

Sheet 4

due on Monday, December 22, 2014

The **integral Gale complexity** of a polytope $P \subset \mathbb{R}^d$ with n vertices is

$$\text{igc}(P) = \min\{\|G\|_\infty : G \subset \mathbb{Z}^e \text{ is a Gale diagram of } P\},$$

where $e = n - d - 1$, $\|\mathcal{A}\|_\infty = \max\{\|v\|_\infty : v \in \mathcal{A}\}$ and $\|v\|_\infty = \max\{|v_i|\}$ for $v = (v_1, \dots, v_e)$.

While the existence of nonrational polytopes shows that $\text{igc}(P) = \infty$ is possible (since $\min \emptyset = \infty$), here we are concerned with the following problem:

Problem G. For $e \in \mathbb{Z}_{\geq 0}$ and $n, m \in \mathbb{Z}_{>0}$, determine

$$q(e, n, m) = \# \left\{ G \subset \mathbb{Z}^e : G \text{ is a Gale diagram of a polytope} \right. \\ \left. \text{with } n \text{ vertices and } \text{igc}(G) = m \right\} / \text{combinatorial equivalence}.$$

For example, $q(0, n, 0) = 1$ and $q(1, n, m) = q(1, n, 1)$ for all $m, n \geq 1$. In more down-to-earth terms, we want to solve the following problem:

Problem G*. Enumerate, up to combinatorial equivalence, all balanced configurations \mathcal{V} of n vectors in \mathbb{Z}^e whose coordinates are all at most m in absolute value, such that

- (1) the maximum m is achieved by some $v \in \mathcal{V}$,
- (2) and such that no hyperplane spanned by $e - 1$ of the vectors strictly separates exactly one vector from the others.

For this, recall that a vector configuration $\mathcal{V} = (v_1, \dots, v_n)$ is **balanced** if $\sum_i v_i = 0$; that no hyperplane defined by $e - 1$ elements of \mathcal{V} separates exactly one vector from the others iff the Gale dual of \mathcal{V} is in convex position; and that two vector configurations are **combinatorially equivalent** if they define the same oriented matroid.

Your job is to write a function in the **polymake** framework that calculates $q(e, n, m)$. Some considerations to keep in mind:

- Correctness is more important than efficiency, but efficiency is supremely important.
- Your code should be correct in all dimensions: no cutting corners by assuming $e = 2$!
- Your code should be able to calculate at least $q(2, 5, 2)$, $q(2, 6, 2)$ and $q(3, 6, 1)$.

You will need to think carefully about several independent aspects:

- (1) How do you iterate over all vector configurations?
- (2) Once you have generated a new configuration, how do you test whether you have already seen a combinatorially isomorphic copy?

My recommendation is to first implement quick-and-dirty solutions to these to test your code for correctness, then iteratively optimize. The choice of data structures will be very important, as well as the decision whether to call **polymake** methods on objects or not. Some methods make your work almost trivial (`P.callPolymakeMethod("LATTICE_POINTS")`, `P.give("VERTICES_IN_FACETS")`, `graph::isomorphic(VIF1, VIF2)`), but some of these are quite time-consuming if you call them often as they have to pass through the perl interface.

LOGISTICS

Please put your code into a directory `exercises/sheet4/code/your_name_here` in the repository; further instructions are in the README file there.

You will find a directory `reports/` with three subdirectories corresponding to Tuesday, December 16; Friday, December 19; and Monday, December 22. Please put a short text file detailing your progress into each of these directories at the appropriate date.

The more often you commit and pull your code, the more often I will be able to give you feedback!

At <https://github.com/julian-upc/discrete-geometry/wiki/2014-Sheet-4> there is a wiki where we can discuss any arising questions.

... AND BEYOND

This could turn into a research project, or even a paper!

- (1) Can you construct a family of rational polytopes with e fixed, but unbounded integral Gale complexity? Or is

$$\text{igc}_e(n) := \max \{ \text{igc}(P) : P \text{ rational, } n \text{ vertices, } \dim \text{Gale}(P) = e \}$$

always bounded?

- (2) If $\text{igc}_e(n)$ is bounded, how does the bound depend on e and n ?
- (3) More specifically, prove, disprove or improve: $\text{igc}_e(n) \in \Omega(\sqrt[e]{n})$.