

TITLE

Cassidy M. Wagner^{1*} and Brodie Pearson^{1*}

¹ Oregon State University ¶ Corresponding author * These authors contributed equally.

DOI: [10.xxxxxx/draft](https://doi.org/10.xxxxxx/draft)

Software

- [Review](#) ↗
- [Repository](#) ↗
- [Archive](#) ↗

Editor: [Open Journals](#) ↗

Reviewers:

- [@openjournals](#)

Submitted: 01 January 1970

Published: unpublished

License

Authors of papers retain copyright
and release the work under a
Creative Commons Attribution 4.0
International License ([CC BY 4.0](#)).

Summary

Fluid systems are everywhere, from small-scale engineering problems to planetary-and-larger-scale systems (atmosphere, ocean, galactic gas clouds). These systems are often turbulent, where motion is chaotic, unpredictable, and can only be characterized through statistical analyses. Spatial structure functions (SFs) are one such statistical analysis technique for turbulence, that require calculation of spatial differences in properties as a function of their separation distance. By combining and then averaging these spatial differences, various types of SF can be constructed to measure different critical properties of fluid flow. However, calculating SFs is often a cumbersome and computationally-intensive task tailored to the specific format of a given turbulence dataset.

non-specialist summary – accessible to oceanographers, turbulence people, fluids people

Earth's changing climate is driven by a small energy imbalance that makes up less than 1% of the total energy budget (CITE). Any deviations in the energy budget, no matter how small, may have a large impact on future climate prediction (CITE, maybe scenarios too). To minimize error in climate prediction it is necessary to investigate how ocean turbulence regulates the global energy budget through the transfer, or cascade, of energy across spatial scales and throughout the ocean.

Statement of need

FluidSFs is a flexible Python package for calculating spatial structure functions (SFs) in one, two, or three spatial dimensions from diverse fluid data sets. The package can construct user-defined SFs that utilize any fluid properties (e.g., velocity, vorticity, temperature, magnetic field etc.), including combinations of these properties and structure functions of arbitrary order. The flexibility of this package enables geophysical, astrophysical, and engineering applications... ADD EXAMPLES OF SF UTILITY BREADTH: e.g., quantifying the energy cycles within Earth's ocean ([Balwada et al., 2022](#); [pearson2019?](#)), Earth's atmosphere ([Lindborg, 1999](#)), and Jupiter's atmosphere ([Young & Read, 2017](#)), the intermittency of magnetohydrodynamic plasma turbulence ([Wan:2016?](#)), the anisotropy of flow over rough beds ([Coscarella et al., 2020](#)), the characteristics of ocean surface temperature ([Schloesser et al., 2016](#)), and the scaling laws of idealized 3D turbulence ([Iyer et al., 2020](#)).

Paragraph on package capabilities & limitations. Regularly-gridded data, Lat-lon gridded data, track/directional sampling, 1D-data, evenly-spaced, irregularly-spaced (what are limitations), binning, bootstrapping(?), local advection terms ([Pearson et al., 2021](#)), Bessel function examples(?) ([xie:2018?](#)) examples of time-averaging, SWOT application. What are limitations (can it take 2D data in a vector rather than array format? Can it calculate 2D or 3D maps of SF rather than just a function of $|r|$ magnitude?). Perhaps these don't need to be mentioned, or can be stated as future developments.

Contextualize package within other relevant software * flowsieve * fuchs 2022: matlab-based

41 GUI package that does third order structure functions among other things * # Acknowledge-
42 ments

43 References

- 44 Balwada, D., Xie, J.-H., Marino, R., & Feraco, F. (2022). Direct observational evidence
45 of an oceanic dual kinetic energy cascade and its seasonality. *Science Advances*, 8(41),
46 eabq2566.
- 47 Coscarella, F., Penna, N., Servidio, S., & Gaudio, R. (2020). Turbulence anisotropy and
48 intermittency in open-channel flows on rough beds. *Physics of Fluids*, 32(11).
- 49 Iyer, K. P., Sreenivasan, K. R., & Yeung, P. (2020). Scaling exponents saturate in three-
50 dimensional isotropic turbulence. *Physical Review Fluids*, 5(5), 054605.
- 51 Lindborg, E. (1999). Can the atmospheric kinetic energy spectrum be explained by two-
52 dimensional turbulence? *Journal of Fluid Mechanics*, 388, 259–288.
- 53 Pearson, B. C., Pearson, J. L., & Fox-Kemper, B. (2021). Advective structure functions
54 in anisotropic two-dimensional turbulence. *Journal of Fluid Mechanics*, 916, 49. <https://doi.org/10.1017/jfm.2021.247>
- 55
- 56 Schloesser, F., Cornillon, P., Donohue, K., Boussidi, B., & Iskin, E. (2016). Evaluation of
57 thermosalinograph and VIIRS data for the characterization of near-surface temperature
58 fields. *Journal of Atmospheric and Oceanic Technology*, 33(9), 1843–1858.
- 59 Young, R. M., & Read, P. L. (2017). Forward and inverse kinetic energy cascades in jupiter's
60 turbulent weather layer. *Nature Physics*, 13(11), 1135–1140.