

# Decision-Making for Land Conservation: A Derivative-Free Optimization Framework with Nonlinear Inputs

## Technical Appendix

Cassidy K. Buhler, Hande Y. Benson

Department of Decision Sciences and MIS, Drexel University  
cb3452@drexel.edu, hvb22@drexel.edu

### Data & Code

The data and code used in our paper can be found at the following repository <https://github.com/cassiebuhler/BBLand>. For an illustration of the file formats, see Figure 1.




Figure 1: Our software using PVA broken down by each step in the code.

### Parameters

The PVA component of our framework was implemented with *RAMAS GIS* (Akçakaya and Root 2013a) and *RAMAS Metapopulation* (Akçakaya and Root 2013b). In *RAMAS GIS*, we used *RAMAS GIS: Spatial Data* for generating our habitat and metapopulation, and *RAMAS GIS: Habitat Dy-*

*namics* to simulate the effects of parcels being protected. The parameters for *RAMAS GIS: Spatial Data* are in Table 1 and *RAMAS GIS: Habitat Dynamics* in Table 2. They are separated into their respective categories by where the input appears in the RAMAS interface.

Then, PVA was conducted in *RAMAS Metapopulation*, where each habitat configuration was evaluated for 100 years, and replicated 1000 times.

#### RAMAS GIS: Spatial Data

##### Habitat Relationships

HS function	[<ASC file>]
HS threshold	0.500
Neighborhood distance (cells)	1.00

##### Link to Metapopulation

Carrying capacity (K)	$ths * 4$
$R_{max}$	1.5
Initial abundance	$ths * 2$
Relative survival	$\max(1, ths * 1.2)$
Relative fecundity	$\max(1, ths * 1.2)$
Distances	Edge to edge

##### Default Population

Local threshold	0
Density dependency type	Scramble
Standard deviation of K	0

##### Dispersal

$a$	0.5
$b$	0.8
$c$	1
$D$	1

##### Correlation

$a$	0.8
$b$	2
$c$	1

Table 1: Parameters for RAMAS GIS: Spatial Data.  $ths$  is total habitat suitability. The formula for dispersal and correlation is (2)

## Density Dependency

Scramble is defined in the RAMAS User Manual (6.0) as “Logistic or Ricker type of density dependence, characteristic of the effect of scramble competition.”

The deterministic growth rate based on population size at time step  $t$  is the following

$$R(t) = R_{\max} * \exp\left(\frac{\ln(R_{\max}) * N_p}{K(t)}\right) \quad (1)$$

where  $N_p$  is the current total number of individuals in a population  $p$ , and  $K(t)$  is the carrying capacity at time  $t$ . Because our data is randomly generated, this type was also randomly selected.

## Dispersal and Correlation

The dispersal rate (migration) between population  $i$  and population  $j$  is calculated as

$$a * \exp\left(\frac{-D_{ij}^c}{b}\right). \quad (2)$$

This function is also used as the correlation of population fluctuation between population  $i$  and population  $j$ .

### RAMAS GIS: Habitat Dynamics

#### Habitat Changes

B - Time step	1
B - K change to next	same until next
B - F change to next	same until next
B - S change to next	same until next
Z - Time step	10

Table 2: Parameters for RAMAS GIS: Habitat Dynamics. We assume the habitat is B for the first 10 years, then use Z for the remaining 90 years.  $K$  is the carrying capacity,  $S$  is the relative survival rate, and  $F$  is relative fecundity rate. These values are dependent on *ths*, thus abundance will change from  $B$  to  $Z$ .

Finally, we provide the parameters for our *pygmo* optimization solver, ACO, in Table 3.

### Ant Colony Optimization

Generations	4
Kernel	$n$
Impstop	1000
Oracle	1e6

Table 3: *Generations* is the number of generations to run the algorithm, *Kernel* is the number of solutions stored in archive, *Impstop* stops the algorithm when this number of runs occurs without improvements, and *Oracle* is the penalty parameter.

Further details on implementation can be found in the README file in the code’s repository.

## Additional Figures

### Result Solutions


In the paper, we presented the figures for the solution to the  $10 \times 10$  constrained model. Here we include the  $Z$  configuration, except only where it is habitable. In Figure 2 we display the  $10 \times 10$  results and in Figure 3 has the  $20 \times 20$  results.

### PVA Metrics


In Figure 4, we visualize PVA metrics with an example of a habitat configuration that was predicted to go to extinction.

## References

- Akçakaya, H. R.; and Root, W. 2013a. RAMAS GIS: Linking spatial data with population viability analysis. *Applied Biomathematics, Setauket, NY*.
- Akçakaya, H. R.; and Root, W. 2013b. RAMAS Metapop: Viability analysis for stage-structured metapopulations (version 6). *Applied Biomathematics, Setauket, NY*.



Constrained Z Patches



Multi-Objective Z Patches






Figure 2:  $10 \times 10$  solutions from paper. The baseline habitat has 16 populations. The constrained model has 12 populations and the multi-objective model has 13 populations.



Multi-Objective Z Patches



Constrained Z Patches




Figure 3:  $20 \times 20$  solutions from paper. The baseline habitat has 35 populations. The constrained model has 43 populations and the multi-objective model has 48 populations.




Figure 4: Median time to quasi extinction is where the probability is 0.5. Expected minimum abundance is where the probability is 0.5. Risk of extinction is the probability when the threshold is 0.