

# Social and ecological determinants of Helicobacter pylori infection in the Philippines



CC Lee, AA Miller, TW McDade  
Department of Anthropology, Northwestern University



## Introduction

*Helicobacter pylori* (*H. pylori*) is a bacterium that infects the gastrointestinal tract, and without proper eradication, can lead to chronic infection.<sup>1</sup>

*H. pylori* infection is associated with gastrointestinal diseases such as peptic ulcer disease and gastric cancer.<sup>1</sup>

The route of transmission for *H. pylori* is unclear, however, evidence suggests the fecal-oral route is an important transmission pathway.<sup>2</sup>

Crowding, hygiene, water sources, socioeconomic status (SES), gender, and urbanicity have been identified as social and ecological determinants of *H. pylori* infection, however, risk factors vary across contexts.<sup>3-6</sup>

The objective of this poster is to discuss social and ecological determinants of *H. pylori* seropositivity in the Philippines to better understand the causes of infection within this context.

## Data

The Cebu Longitudinal Health and Nutrition Survey (CLHNS) is a longitudinal study of health outcomes for a cohort of individuals born between 1983 and 1984 in Metro Cebu, Philippines.<sup>7</sup> This study draws from data collected during the 2005 follow-up survey.

**Sociodemographic variables:** Assets, education, history of marriage, income, medical expenditure, sex, and urbanicity scale.

**Ecological variables:** Crowding, excrement near house level, garbage disposal location and method, hygiene index, neighborhood garbage level, toilet privacy, toilet type, and usual source of drinking water.

***H. pylori* seropositivity:** 128 randomly selected dried blood spots (DBS) were analyzed for *H. pylori* antibody levels using a protocol modified for DBS.<sup>8</sup>


Antibody levels < 12.2 EU/mL = seronegative. Levels > 15.3 EU/mL = seropositive. Four samples were equivocal and excluded.

## Sample Characteristics

**Table 1.** Descriptive statistics for characteristics of individuals included in the analysis.

Characteristic (N = 124)	n / N (%); Mean (SD)
<b>Seropositivity</b>	
Positive	40 / 124 (32%)
Negative	84 / 124 (68%)
<b>Sex</b>	
Female	66 / 124 (53%)
Male	58 / 124 (47%)
<b>Age</b>	
	21.0 (0.3)
<b>History of Marriage</b>	
Married	19 / 124 (15%)
Never married	105 / 124 (85%)


## Sociodemographic Variables



**Figure 1.** Distribution of *H. pylori* seropositivity across the social variables (A) assets, (B) history of marriage, (C) sex, and (D) urbanicity scale.

The odds of seropositivity were significantly greater for having a low asset score (OR = 5.70, 95% CI [1.94, 19.5]) compared to a high asset score, for having a history of marriage (OR = 2.78, 95% CI [1.02, 7.66]), for females (OR = 2.4, 95% CI [1.11, 5.40]) compared to males, and for medium urbanicity (OR = 3.56, 95% CI [1.37, 9.99]) compared to a high urbanicity.


## Ecological Variables



**Figure 2.** Distribution of *H. pylori* seropositivity across the ecological variables (A) excrement near house and (B) hygiene index.

The odds of seropositivity were significantly greater for having low (OR = 3.97, 95% CI [1.39, 13.3]) or high (OR = 4.73, 95% CI [1.61, 16.1]) excrement near house compared to no excrement near house and having medium (OR = 10.5, 95% CI [2.77, 68.6]) or low (OR = 10.7, 95% CI [2.66, 72.7]) hygiene index compared to a high hygiene index.

## Multivariable Regression



**Figure 3.** Results of multivariable logistic regression including assets (compared to high assets), hygiene index (compared to high hygiene), and sex (compared to males). Medium hygiene index (OR = 8.06, 95% CI [2.04, 54.1]) and low assets (OR = 4.25, 95% CI [1.23, 16.6]) and a remain associated with seropositivity.

## Conclusion and Discussion

Analysis of social and ecological determinants of *H. pylori* infection suggest that assets (SES) and hygiene were predictive of infection.

Education, income, medical expenditure, crowding, garbage disposal location and method, neighborhood garbage level, toilet type, and usual source of drinking water were not associated with seropositivity.

While there was a female sex bias for infection, sex itself was not predictive of seropositivity in multivariable analysis. This indicates that gender disparities in access to sanitation and hygiene infrastructure likely drive this relationship.

The increased odds of seropositivity for lower hygiene indices and higher levels of excrement near house supports the hypothesis that *H. pylori* is transmitted through the fecal-oral route.<sup>2</sup> The lack of association between seropositivity and usual source of drinking water suggests transmission via direct exposure to fecal matter rather than indirectly through drinking water.

Increased access to sanitation infrastructure, such as a centralized sewage management system,<sup>9</sup> could reduce the prevalence of *H. pylori* infections in Metro Cebu, Philippines.

## References

1. Mahachai et al. (2018). *Helicobacter pylori* management in ASEAN: The Bangkok consensus report: *Helicobacter pylori* management. *Journal of Gastroenterology and Hepatology*, 33(1), 37–56. <https://doi.org/10.1111/jgh.13911>
2. Bui et al. (2016). Serologic Evidence for Fecal–Oral Transmission of *Helicobacter pylori*. *The American Journal of Tropical Medicine and Hygiene*, 94(1), 82–88. <https://doi.org/10.4269/ajtmh.15-0297>
3. Khoder et al. (2019). Prevalence of *Helicobacter pylori* and Its Associated Factors among Healthy Asymptomatic Residents in the United Arab Emirates. *Pathogens*, 8(2), 44. <https://doi.org/10.3390/pathogens8020044>
4. Nurgalieva et al. (2002). *Helicobacter pylori* infection in Kazakhstan: Effect of water source and household hygiene. *The American Journal of Tropical Medicine and Hygiene*, 67(2), 201–206. <https://doi.org/10.4269/ajtmh.2002.67.201>
5. Emerenini et al. (2021). Prevalence and risk factors for *Helicobacter pylori* infection among children in Owerri, Nigeria. [https://doi.org/10.4103/njcp.njcp\\_687\\_20](https://doi.org/10.4103/njcp.njcp_687_20)
6. Cheng et al. (2009). Prevalence of *Helicobacter pylori* Infection and Identification of Risk Factors in Rural and Urban Beijing, China. *Helicobacter*, 14(2), 128–133. <https://doi.org/10.1111/j.1523-5378.2009.00668.x>
7. Adair et al. (2011). Cohort Profile: The Cebu Longitudinal Health and Nutrition Survey. *International Journal of Epidemiology*, 40(3), 619–625. <https://doi.org/10.1093/ije/dyq085>
8. Miller et al. (2022). A dried blood spot method for the quantification of *Helicobacter pylori* antibodies. *American Journal of Human Biology*, 34(S2). <https://doi.org/10.1002/ajhb.23740>
9. Ostojic et al. (2013). *Energizing Green Cities in Southeast Asia: Applying Sustainable Urban Energy and Emissions Planning*. The World Bank. <https://doi.org/10.1596/978-0-8213-9837-1>

## Acknowledgements

These findings are presented on behalf of the participants of the CLHNS who generously provided their time, samples, and information to support ongoing health research in the Philippines.

The study resulting in this presentation was assisted by grants from the Undergraduate Research Grant Program and Conference Travel Grant which are administered by Northwestern University's Office of Undergraduate Research, as well as a grant from the Baker Program in Undergraduate Research, which is administered by Northwestern University's Weinberg College of Arts and Sciences. However, the conclusions, opinions, and other statements in this presentation are the authors' and not necessarily those of the sponsoring institution.