

On Continuous Efron's Dice*

Cassio Neri

20 November 2011

Abstract

A die D_1 is said to be **stronger** than another die D_2 if the probability of D_1 beating D_2 is bigger than $1/2$. It is natural to expect that being stronger is a transitive relationship, that is, if D_1 is stronger than D_2 and D_2 is stronger than D_3 , then D_1 is stronger than D_3 . Efron's dice are a set of four dice invented by Bradley Efron which is a counter-example for the transitivity [1].

In this document, instead of a set of four dice, we consider a set of independent continuous random variables X_1, \dots, X_M such that $P(X_1 > X_2) = \dots = P(X_{M-1} > X_M) = P(X_M > X_1) = p$. Under certain technical assumptions, we shall see that $p \leq 3/4$.

1 Introduction

A more serious title for this document could be “On non-transitive continuous random variables”. However, I think that invoking the Efron's dice in the title is more appealing since this is a (literally) more concrete example of non-transitive random variables.

This document does not pretend to be a research paper. It considers the problem with a certain degree of mathematical rigorousness and generality and, at the same time, keeping things simple. Whenever generality is at the expense of clarity, the latter is favoured. A few remarks along the text indicate possible generalisations.

We are interested in a set of independent random variables X_1, \dots, X_M such that $P(X_1 > X_2) = \dots = P(X_{M-1} > X_M) = P(X_M > X_1) = p$. We shall call **non-transitive** to any set of random variables that satisfies this property. The probability p will be called the **winning** probability.

In the original Efron's dice problem, we have $M = 4$ and the random variables are the outcomes of four non-standard and specially built dice for which $p = 2/3$. The peculiarity of such dice is not that they are “unfair” and favour some faces in respect to others. All faces are equally probable but the numbers printed on them are not mutually distinct in $\{1, 2, 3, 4, 5, 6\}$. This is what makes some outcomes more likely than others. In other words, the probability space of each die is the set of six uniformly distributed faces which we can still number 1, ..., 6 but the random variables are not as those of a normal die where the i^{th} face is mapped to the number i . This fact will have a continuous counterparty as we shall see in Section 3.

Can the dice be built in order to get $p > 2/3$? A negative answer was given by Samson [2]. He also considered the generalization for M dice each of them having N equally probable faces. He showed that in the this case we have

$$p < \frac{3}{4} - \frac{1}{2N} - \frac{1}{4N^2},$$

and noted that $p \rightarrow 3/4$ as the number of sides N goes to infinity. Actually, we shall see that the upper bound $3/4$ holds for every non-transitive set of random variables provided some mild conditions on their probability distribution functions hold. Samson's work showed that we can get as close as we want to the bound $3/4$ but I don't know whether this bound can be reached.

*This work is licensed under a [Creative Commons Attribution-ShareAlike 4.0 International License](#)




Figure 2: If the square in 3rd row and 3rd column is white, then 11 others will be so.

contradiction remains the same in any case.

Taking a step back, let's try to understand what is special about the square in 3rd row and 3rd column.

First of all, in order to the proof to work, the special square under consideration must be in the i^{th} row and j^{th} column with $i = j$. Indeed, assuming $i \neq j$ and also that this square is black, following the previous argument, would make us conclude that $d_i^m > d_j^n$ with $i \neq j$. This inequality would not allow us to deduce the cyclical sequence of inequalities (1) where all terms have the same subscript. Hence we would not be able to deduce the absurd statement $d_i^m > d_i^m$.

Now, still following the proof, when the special square is white we conclude that there is at least a certain number of white squares. As a consequence, we get an upper bound for the number of black squares. To get the smallest upper bound, we need to choose the special square that maximises the white area.

Therefore, the proof needs the special square to be the one on the left-bottom to right-top diagonal of the grid that maximizes the rectangular white area spanned by this square. The closer the square is to the grid's centre, the bigger is this area. Hence, the square in the 3rd row and 3rd column satisfies our needs. The one on the 4th row and 4th column is equally adequate.

In preparation for the continuous case and overlooking the physical constraints on building dice, let's consider what happens if we increase the number of sides by splitting each of the 3rd and 4th faces (the *special ones*) into equally probable new faces. The new grid representation for such situation is shown on the left of Figure 3.

From the analysis of the proof we can immediately spot the the new special squares are the 4th and 5th on the left-bottom to right-top diagonal. Picking up the 4th one and assuming it's white, we obtain a white area of at least $3.5 \times 3 = 10.5$. This is illustrated on the right of Figure 3. The figure also emphasizes the extra grey area measuring 1.5 that was white in Figure 2. Actually, this area can be black, increasing the probability p by $3/2$ to $2/3 + 3/2 = 5/6$.

We stress that the previous arguments on the fictitious dice does not say that the upper bound for the winning probability is achieved. The problem is not the physical impossibility of building such dice because we could use another set of random variables. (Actually, we are already doing it by considering the grid with the Lebesgue measure.) The argument just gives just an upper bound for winning probability for a certain set of random variables.




Figure 3: Fictitious dice with 8 faces.

3 General non-transitive random variables

In this section we adapt the previous ideas to non-transitive random variables that are not necessarily discrete as in the dice case. We make two assumptions on the set of non-transitive random variables X_1, \dots, X_M , namely,

1. They are defined on the probability space $]0, 1[^M$ equipped with the Lebesgue σ -algebra and (probability) measure.
2. For all $(x_1, \dots, x_M) \in]0, 1[^M$, $X_m(x_1, \dots, x_M) = F_m(x_m)$, where $F_m :]0, 1[\rightarrow \mathbb{R}$ is non-decreasing.

These assumptions say that each X_m is a function of an uniformly distributed random variable x_m in $]0, 1[$. Notice the similarity with the original Efron's dice where the random variables are functions of uniformly random distributed faces of a die.

We shall prove that $3/4$ is an upper bound for the winning probability.

For each pair (m, n) , with $n \equiv m + 1 \pmod{M}$, the grid used in the previous argument becomes the square $]0, 1[\times]0, 1[$ where each point (x_m, x_n) is painted in black if $F_m(x_m) > F_n(x_n)$ and in white otherwise. In more mathematical terms, we consider the set $B_{m,n} = \{(x_m, x_n) \in]0, 1[\times]0, 1[; F_m(x_m) > F_n(x_n)\}$ whose Lebesgue measure is the winning probability $p = P(X_m > X_n)$ of X_m beating X_n .

The proof here also needs its special “square” which, for the continuous case, collapses into the single point $(1/2, 1/2)$.

Suppose that there exists a pair (m, n) , with $n \equiv m + 1 \pmod{M}$, such that $F_m(1/2) \leq F_n(1/2)$ (that is, $(1/2, 1/2)$ is white in this grid). Since F is non-decreasing we have

$$F_m(x_m) \leq F_m(1/2) \leq F_n(1/2) \leq F_n(x_n), \quad \forall x_m \in]0, 1/2[, \quad \forall x_n \in]1/2, 1[.$$

It follows that

$$A_{m,n} := \{x \in]0, 1[^M ; x_m \in]0, 1/2[, x_n \in]1/2, 1[\} \subset B_{m,n}^c.$$

Hence, $1/4 = P(A_{m,n}) \leq P(B_{m,n}^c) = P(X_m \leq X_n)$ and then, $p = P(X_m > X_n) \leq 3/4$.

We conclude that if $p > 3/4$, then $F_m(1/2) > F_n(1/2)$ for all (m, n) , such that $n \equiv m + 1 \pmod{M}$. Therefore,

$$F_1(1/2) > F_2(1/2) > \dots > F_M(1/2) > F_1(1/2)$$

which is absurd.

We end this document by noticing that we can replace the hypothesis on X_1, \dots, X_M stated at the beginning of this section by a simple assumption on their probability distribution functions. This means that the same bound on the winning probability holds for a larger collection of continuous random variables.

Indeed, let G_m be the probability distribution function of X_m and **assume that G_m is continuous and strictly increasing**. Then $G_m : \mathbb{R} \rightarrow]0, 1[$ is invertible with inverse $F_m := G_m^{-1} :]0, 1[\rightarrow \mathbb{R}$.

We denote by $x := (x_1, \dots, x_M)$ a generic point in the probability space $]0, 1[^M$ equipped with the Lebesgue σ -algebra and (probability) measure. Notice that the projections $\pi_m :]0, 1[^M \rightarrow]0, 1[$ given by $\pi_m(x) := x_m$, $m \in \{1, \dots, M\}$, are independent and uniformly distributed in $]0, 1[$.

Hence, the variables $\tilde{X}_1, \dots, \tilde{X}_M$ defined by $\tilde{X}_m(x) := F_m(x_m)$ for all $m \in \{1, \dots, M\}$ satisfy the conditions 1 and 2 above. Moreover \tilde{X}_m is equal in distribution to X_m since

$$P(\tilde{X}_m < y) = P(F_m(x_m) < y) = P(x_m < G_m(y)) = P(\pi_m(x) < G_m(y)) = G_m(y),$$

where the last equality is a consequence of π_m being uniformly distributed in $]0, 1[$.

Therefore, the result $p \leq 3/4$ still holds in this case.

I feel that these results can be extended further without the need for strict monotonicity of G_m but the proof must deal carefully with intervals where G_m is constant because they complicate the inversion of G_m .

References

- [1] Wikipedia, *Nontransitive dice*, http://en.wikipedia.org/wiki/Nontransitive_dice#Efron.27s_dice.
- [2] Ophir Samson, *Efrons Dice*, <http://ophir-research.com/Papers/Efron.pdf>.