

1. What three elimination matrices E_{21}, E_{31}, E_{32} put A into its upper triangular form $E_{32}E_{31}E_{21}A = U$? Multiply by $E_{32}^{-1}, E_{31}^{-1}, E_{21}^{-1}$ to factor A into L times U :

$$A = \begin{bmatrix} 1 & 0 & 1 \\ 2 & 2 & 2 \\ 3 & 4 & 5 \end{bmatrix}, L = E_{21}^{-1}, E_{31}^{-1}, E_{32}^{-1}$$

$$\begin{bmatrix} 1 & 0 & 1 \\ 2 & 2 & 2 \\ 3 & 4 & 5 \end{bmatrix} \xrightarrow{\textcircled{1} x-2+2} \begin{bmatrix} 1 & 0 & 1 \\ 0 & 2 & 0 \\ 3 & 4 & 5 \end{bmatrix} \xrightarrow{\textcircled{1} x-3+3} \begin{bmatrix} 1 & 0 & 1 \\ 0 & 2 & 0 \\ 0 & 4 & 2 \end{bmatrix} \xrightarrow{\textcircled{2} x-2+3} \begin{bmatrix} 1 & 0 & 1 \\ 0 & 2 & 0 \\ 0 & 0 & 2 \end{bmatrix}$$

$$E_{21} = \begin{bmatrix} 1 & 0 & 0 \\ -2 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \quad E_{31} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ -3 & 0 & 1 \end{bmatrix} \quad E_{32} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & -2 & 1 \end{bmatrix}$$

$$E_{32}E_{31}E_{21}A = U$$

$$A = E_{21}^{-1}E_{31}^{-1}E_{32}^{-1}U$$

$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & -2 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ -3 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 \\ -2 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ -2 & 1 & 0 \\ 1 & -2 & 1 \end{bmatrix}$$

$$\begin{bmatrix} 1 & 0 & 0 & | & 1 & 0 & 0 \\ -2 & 1 & 0 & | & 0 & 1 & 0 \\ 1 & -2 & 1 & | & 0 & 0 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 & | & 1 & 0 & 0 \\ 0 & 1 & 0 & | & 2 & 1 & 0 \\ 0 & 0 & 1 & | & 3 & 2 & 1 \end{bmatrix} \Rightarrow L = \begin{bmatrix} 1 & 0 & 0 \\ -2 & 1 & 0 \\ 1 & -2 & 1 \end{bmatrix} \#$$

2. Compute L and U for the symmetric matrix A :

$$A = \begin{bmatrix} a & a & a & a \\ a & b & b & b \\ a & b & c & c \\ a & b & c & d \end{bmatrix}$$

Find four conditions on a, b, c, d to get $A = LU$ with four pivots.

$$\begin{bmatrix} a & a & a & a \\ a & b & b & b \\ a & b & c & c \\ a & b & c & d \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 1 & 1 & 0 & 0 \\ 1 & 1 & 1 & 0 \\ 1 & 1 & 1 & 1 \end{bmatrix} \begin{bmatrix} a & a & a & a \\ 0 & b-a & b-a & b-a \\ 0 & 0 & c-b & c-b \\ 0 & 0 & 0 & d-c \end{bmatrix}$$

$$\textcircled{1} a \neq 0$$

$$\textcircled{2} a \neq b$$

$$\textcircled{3} b \neq c$$

$$\textcircled{4} c \neq d \#$$

3. Suppose A is rectangular (m by n) and S is symmetric (m by m).

(a) Transpose $A^T S A$ to show its symmetry. What shape is this matrix?

(b) Show why $A^T A$ has no negative numbers on its diagonal.

$$A^{m \times n} \cdot S^{m \times m} \cdot A^T_{n \times m}$$

(a) $A^T_{n \times m} \cdot S^{m \times m} \cdot A^{m \times n} = X_{n \times n} \Rightarrow \text{square}$

(b)
$$\begin{bmatrix} a_{11} & a_{12} & a_{13} & \dots & a_{1n} \\ a_{21} & & & & \\ \vdots & & & & \\ a_{m1} & & & & a_{mn} \end{bmatrix} \begin{bmatrix} a_{11} & a_{21} & \dots & a_{m1} \\ a_{12} & a_{22} & & \\ a_{13} & : & & \\ \vdots & : & & \\ a_{1n} & & & a_{mn} \end{bmatrix} = \begin{bmatrix} a_{11}^2 + a_{12}^2 + \dots + a_{1n}^2 & & & \\ & a_{21}^2 + a_{22}^2 + \dots + a_{2n}^2 & & \\ & & \ddots & \\ & & & a_{m1}^2 + a_{m2}^2 + \dots + a_{mn}^2 \end{bmatrix}$$

$$A \longrightarrow A^T$$

row 1	column 1
row 2	column 2
\vdots	
row x	column x

when $A \cdot A^T$, the number on its diagonal

will be (row x in A) \cdot (column x in A^T)

$$= (\text{row } x \text{ in } A) \cdot (\text{row } x \text{ in } A)$$

$$= (\text{row } x \text{ in } A)^2 \geq 0$$

$\therefore A A^T$ has no negative numbers on its diagonal.

4. Tridiagonal matrices have zero entries except on the main diagonal and the two adjacent diagonals. Factor these into $A = LU$ and $A = LDL^T$:

$$A = \begin{bmatrix} 1 & 1 & 0 \\ 1 & 2 & 1 \\ 0 & 1 & 2 \end{bmatrix} \text{ and } A = \begin{bmatrix} a & a & 0 \\ a & a+b & b \\ 0 & b & b+c \end{bmatrix}$$

$$A = \begin{bmatrix} 1 & 1 & 0 \\ 1 & 2 & 1 \\ 0 & 1 & 2 \end{bmatrix} \xrightarrow[\substack{E_{21} \\ \begin{bmatrix} 1 & 0 & 0 \\ -1 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}}]{\textcircled{1} \times (-1) + \textcircled{2}} \begin{bmatrix} 1 & 1 & 0 \\ 0 & 1 & 1 \\ 0 & 1 & 2 \end{bmatrix} \xrightarrow[\substack{E_{32} \\ \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & -1 & 1 \end{bmatrix}}]{\textcircled{2} \times (-1) + \textcircled{3}} \begin{bmatrix} 1 & 1 & 0 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{bmatrix}$$

$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & -1 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 \\ -1 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ -1 & 1 & 0 \\ 1 & -1 & 1 \end{bmatrix}$$

$$\begin{bmatrix} 1 & 0 & 0 & | & 1 & 0 & 0 \\ -1 & 1 & 0 & | & 0 & 1 & 0 \\ 1 & -1 & 1 & | & 0 & 0 & 1 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 0 & 0 & | & 1 & 0 & 0 \\ 0 & 1 & 0 & | & 1 & 1 & 0 \\ 0 & 0 & 1 & | & 0 & 1 & 1 \end{bmatrix}$$

$$A = \begin{bmatrix} 1 & 0 & 0 \\ 1 & 1 & 0 \\ 0 & 1 & 1 \end{bmatrix} \begin{bmatrix} 1 & 1 & 0 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 1 & 1 & 0 \\ 0 & 1 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 1 & 0 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{bmatrix} \#$$

$$A = \begin{bmatrix} a & a & 0 \\ a & a+b & b \\ 0 & b & b+c \end{bmatrix} \xrightarrow[\substack{\begin{bmatrix} 1 & 0 & 0 \\ -1 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}}]{\textcircled{1} \times (-1) + \textcircled{2}} \begin{bmatrix} a & a & 0 \\ 0 & b & b \\ 0 & b & b+c \end{bmatrix} \xrightarrow[\substack{\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & -1 & 1 \end{bmatrix}}]{\textcircled{2} \times (-1) + \textcircled{3}} \begin{bmatrix} a & a & 0 \\ 0 & b & b \\ 0 & 0 & c \end{bmatrix}$$

$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & -1 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 \\ -1 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ -1 & 1 & 0 \\ 1 & -1 & 1 \end{bmatrix}$$

$$\begin{bmatrix} 1 & 0 & 0 & | & 1 & 0 & 0 \\ -1 & 1 & 0 & | & 0 & 1 & 0 \\ 1 & -1 & 1 & | & 0 & 0 & 1 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 0 & 0 & | & 1 & 0 & 0 \\ 0 & 1 & 0 & | & 1 & 1 & 0 \\ 0 & 0 & 1 & | & 0 & 1 & 1 \end{bmatrix}$$

$$A = \begin{bmatrix} 1 & 0 & 0 \\ 1 & 1 & 0 \\ 0 & 1 & 1 \end{bmatrix} \begin{bmatrix} a & a & 0 \\ 0 & b & b \\ 0 & 0 & c \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 1 & 1 & 0 \\ 0 & 1 & 1 \end{bmatrix} \begin{bmatrix} a & 0 & 0 \\ 0 & b & 0 \\ 0 & 0 & c \end{bmatrix} \begin{bmatrix} 1 & 1 & 0 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{bmatrix} \#$$

5. Let $V = \{(a_1, a_2) : a_1, a_2 \in \mathbb{F}\}$ where \mathbb{F} is a field. Define addition of elements of V coordinatewise and for $c \in \mathbb{F}$ and $(a_1, a_2) \in V$, define,

$$c(a_1, a_2) = (a_1, 0)$$

Is V a vector space over \mathbb{F} with these operations? Justify your answer.

$$(a_1, a_2) \in V$$

$$c(a_1, a_2) = (a_1, 0) \notin \mathbb{F}$$

$$\text{If } c(a_1, a_2) \in \mathbb{F}$$

then $c(a_1, a_2)$ should equal (ca_1, ca_2) #

6. The matrix $A = \begin{bmatrix} 2 & -2 \\ 2 & -2 \end{bmatrix}$ is a "vector" in the space M of all 2 by 2 matrices. Write down the zero vector in this space, the vector $\frac{1}{2}A$, and the vector $-A$. What matrices are in the smallest subspace containing A ?

$$\frac{1}{2}A = \begin{bmatrix} 1 & -1 \\ 1 & -1 \end{bmatrix} \quad -A = \begin{bmatrix} -2 & 2 \\ -2 & 2 \end{bmatrix}$$

$$\text{zero vector} = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$$

$$\text{smallest subspace} = t \begin{bmatrix} 2 & -2 \\ 2 & -2 \end{bmatrix} \quad (t \in \mathbb{R})$$