

CHAPTER 6

Sampling of Continuous-Time Signals

Tutorial Problems

1. (a) Solution:

$$x_c(t) = \frac{5}{2}e^{j\frac{\pi}{6}}e^{j200\pi t} + \frac{5}{2}e^{-j\frac{\pi}{6}}e^{-j200\pi t} + \frac{2}{j}e^{j300\pi t} - \frac{2}{j}e^{-j300\pi t}$$

The spectra of $x_c(t)$ is:

$$X_c(j\Omega) = \begin{cases} \frac{5}{2}e^{j\frac{\pi}{6}}, & \Omega = 200\pi \\ \frac{5}{2}e^{-j\frac{\pi}{6}}, & \Omega = -200\pi \\ \frac{2}{j}, & \Omega = 300\pi \\ -\frac{2}{j}, & \Omega = -300\pi \\ 0, & \text{elsewhere} \end{cases}$$

The spectra $X(e^{j\omega})$ of $x[n]$ is:

$$X(e^{j\omega})|_{\omega=\Omega T} = F_s \sum_{k=-\infty}^{\infty} X_c(j\Omega - j2\pi k F_s)$$

$$X(e^{j\omega})|_{\omega=2\pi F T} = F_s \sum_{k=-\infty}^{\infty} X_c[j2\pi(F - k F_s)]$$

The signal can recovered for $x[n]$ if $F_s = 1$ KHz.

- (b) Solution:

The signal can recovered for $x[n]$ if $F_s = 500$ Hz.

- (c) Solution:

The signal can NOT recovered for $x[n]$ if $F_s = 100$ Hz.


(d) tba.

MATLAB script:


```
% P0601: Illustrates the alias distortion
close all; clc
Fs = 1e3; % Part (a)
% Fs = 500; % Part (b)
% Fs = 100; % Part (c)
T = 1/Fs;
FH = 150;
FL = FH+Fs;
F = -FL:50:FL;
X = zeros(1,length(F));
for k = -1:1;
    ind = F == -150+k*Fs; X(ind) = X(ind)-2/j;
    ind = F == -100+k*Fs; X(ind) = X(ind)+5/2*exp(-j*pi/6);
    ind = F == 100+k*Fs; X(ind) = X(ind)+5/2*exp(j*pi/6);
    ind = F == 150+k*Fs; X(ind) = X(ind)+2/j;
end
ind = X==0;
X(ind) = nan;
%% Plot:
hfa = figconfig('P0601a');
subplot(211)
stem(F*2*pi*T,abs(X),'filled')
ylim([0 max(abs(X))+0.5])
set(gca,'XTick',[-Fs*2*pi*T 0 Fs*2*pi*T])
set(gca,'XTickLabel',{'-\Omega_s*T','0','\Omega_s*T'})
xlabel('\omega','fontsize',LFS)
ylabel('|H(e^{j\omega})|','fontsize',LFS)
title('Magnitude Response','fontsize',TFS)
subplot(212)
stem(F*2*pi*T,angle(X),'filled')
set(gca,'XTick',[-Fs*2*pi*T 0 Fs*2*pi*T])
set(gca,'XTickLabel',{'-\Omega_s*T','0','\Omega_s*T'})
xlabel('\omega','fontsize',LFS)
ylabel('\angle H(e^{j\omega})|','fontsize',LFS)
title('Phase Response','fontsize',TFS)

hfb = figconfig('P0601b');
```

```
subplot(211)
stem(F,abs(X),'filled')
ylim([0 max(abs(X))+0.5])
set(gca,'XTick',[-Fs -Fs/2 0 Fs/2 Fs])
set(gca,'XTickLabel',{'-Fs','-Fs/2','0','Fs/2','Fs'})
xlabel('F','fontsize',LFS)
ylabel('|H(e^{j2\pi FT})|','fontsize',LFS)
title('Magnitude Response','fontsize',TFS)
subplot(212)
stem(F,angle(X),'filled')
set(gca,'XTick',[-Fs -Fs/2 0 Fs/2 Fs])
set(gca,'XTickLabel',{'-Fs','-Fs/2','0','Fs/2','Fs'})
xlabel('F','fontsize',LFS)
ylabel('\angle H(e^{j2\pi FT})','fontsize',LFS)
title('Phase Response','fontsize',TFS)
```




(a)




(b)

FIGURE 6.1: Spectra of $X(e^{j\omega})$ as a function of (a) ω in $\frac{\text{rad}}{\text{sam}}$ and (b) F in Hz when the sample rate is $F_s = 1 \text{ KHz}$.




(a)




(b)

FIGURE 6.2: Spectra of $X(e^{j\omega})$ as a function of (a) ω in $\frac{\text{rad}}{\text{sam}}$ and (b) F in Hz when the sample rate is $F_s = 500$ Hz.



(a)



(b)

FIGURE 6.3: Spectra of $X(e^{j\omega})$ as a function of (a) ω in $\frac{\text{rad}}{\text{sam}}$ and (b) F in Hz when the sample rate is $F_s = 100$ Hz.

2. (a) Solution:

The continuous signal $x_c(t)$ is:

$$x_c(t) = 5e^{-10|t|}$$

The sampled sequence $x[n]$ is:

$$x[n] = x_c(nT) = 5e^{-10|n|T} = 5a^{|n|}, \quad \text{define } a = e^{-10T}$$

The spectra $X(e^{j\omega})$ of $x[n]$ is:

$$X(e^{j\omega})|_{\omega=2\pi FT} = 5 \cdot \frac{1 - a^2}{1 - 2a \cos(2\pi FT) + a^2}$$




FIGURE 6.4: Magnitude response of $X(e^{j\omega})$ as a function of F in Hz when the sampling rate is $F_s = 100$.

(b) See plot below.

(c) See plot below.

(d) Solution:

For sampling rate $F_s = 100$ Hz, the signal $X_c(t)$ can be reasonably recovered from its samples $x[n]$.

MATLAB script:

```
% P0602: Plot the spectra of sampled sequence
close all; clc
% Fs = 100; % Part a
% Fs = 50; % Part b
Fs = 25; % Part c
T = 1/Fs;
```




FIGURE 6.5: Magnitude response of $X(e^{j\omega})$ as a function of F in Hz when the sampling rate is $F_s = 50$.




FIGURE 6.6: Magnitude response of $X(e^{j\omega})$ as a function of F in Hz when the sampling rate is $F_s = 25$.


```
a = exp(-10*T);
F = linspace(-150,150,1000);
X = 5*(1-a^2)./(1-2*a*cos(2*pi*F*T)+a^2);
%% Plot:
hfa = figconfg('P0602a','long');
plot(F,abs(X))
xlabel('F','fontsize',LFS)
ylabel('|H(e^{j2\pi F})|','fontsize',LFS)
title('Magnitude Response','fontsize',TFS)
legend(['Fs = ',num2str(Fs)],'location','best')
```


3. (a) Solution:


$$x[n] = x_c(0.05n) = 2 \cos(0.5\pi n - \frac{\pi}{3}) - 3 \sin(0.8\pi n)$$

(b) Solution:

$$y_r(t) = 2 \cos(10\pi t - 60^\circ) - 3 \sin(16\pi t)$$



(a)



(b)

FIGURE 6.7: (a) Plot of $x[n]$ and $x_c(t)$ and (b) plot of $y_r(t)$ when the continuous signal is sampled at $t = 0.05n$.

(c) Solution:

$$x[n] = x_c(0.1n) = 2 \cos(\pi n - \frac{\pi}{3}) - 3 \sin(1.6\pi n)$$

$$y_r(t) = 2 \cos(10\pi t - 60^\circ) + 3 \sin(4\pi t)$$

(d) Solution:

$$x[n] = x_c(0.5n) = 2 \cos(5\pi n - \frac{\pi}{3}) - 3 \sin(8\pi n)$$




FIGURE 6.8: (a) Plot of $x[n]$ and $x_c(t)$ and (b) plot of $y_r(t)$ when the continuous signal is sampled at $t = 0.1n$.

$$y_r(t) = 2 \cos(2\pi t - 60^\circ)$$

MATLAB script:

```
% P0603: Illustrate Ideal DAC
close all; clc
t1 = 0; t2 = 5;
dt = 1e-4;
t = t1:dt:t2;
xc = 2*cos(10*pi*t-pi/3)-3*sin(16*pi*t);
%% Part (a) and (b)
T = 0.05;
yr = 2*cos(10*pi*t-pi/3)-3*sin(16*pi*t);
%% Part (c)
% T = 0.1;
% % yr = 2*cos(10*pi*t)+3*sin(4*pi*t);
```




FIGURE 6.9: (a) Plot of $x[n]$ and $x_c(t)$ and (b) plot of $y_r(t)$ when the continuous signal is sampled at $t = 0.5n$.

```
% yr = 2*cos(10*pi*t-pi/3)+3*sin(4*pi*t);
%% Part (d)
% T = 0.5;
% yr = 2*cos(2*pi*t-pi/3);

% Sampling:
nT = t1:T:t2;
xn = 2*cos(10*pi*nT-pi/3)-3*sin(16*pi*nT);
%% Plot:
hfa = figconf('P0603a','long');
plot(t,xc); hold on
plot(nT,xn,'.','color','red')
xlabel('t','fontsize',LFS)
title(['Sampling Interval T = ',num2str(T)],'fontsize',TFS)
legend('x_c(t)', 'x[n]', 'location', 'northeast')
```

```

hfb = figconfig('P0603b','long');
plot(t,xc,t,yr,nT,xn,'.')
xlabel('t','fontsize',LFS)
title(['Sampling Interval T = ',num2str(T)],'fontsize',TFS)
legend('x_c(t)','y_r(t)','x[n]','location','northeast')

```

4. (a) Solution:

If $F_0 = 10, 20$, and 40 Hz, $2F_0 < F_s = 100$,

$$y_r(t) = x_c(t) = \cos(2\pi F_0 t + \theta_0)$$

- (b) tba.

- (c) Solution:

When $2F_0 = F_s$,

$$y_r(t) = 2 \cos \theta_0 \cdot \cos 2\pi F_0 t$$

5. The same to last problem P0604

6. MATLAB script:


```

% P0606: Sampling a linear FM signal
close all; clc
B = 10;
Fs = B;
tau = 10;
dt = 1e-4;
t = 0:dt:tau;
xc = sin(pi*B*t.^2/tau);
nT = 0:1/Fs:tau;
xn = sin(pi*B*nT.^2/tau);
ind = B*t/tau > Fs/2;
yr = xc;
yr(ind) = -sin(2*pi*(Fs-B*t(ind)/tau/2).*t(ind));
%% Plot:
hfa = figconfig('P0606a','long');
plot(t,xc)
xlabel('t (sec)','fontsize',LFS)
ylabel('x_c(t)','fontsize',LFS)


hfb = figconfig('P0606b','long');

```


```
plot(nT,xn,'.-');  
xlabel('t (sec)','fontsize',LFS)  
ylabel('x(nT)','fontsize',LFS)  
  
hfc = figconfig('P0606c','long');  
plot(t,yr); hold on  
plot(nT,xn,'.')  
ylim([-1 1])  
xlabel('t (sec)','fontsize',LFS)  
ylabel('x_r(t)','fontsize',LFS)
```



(a)



(b)



(c)

FIGURE 6.10: (a) Continuous signal $x_c(t)$, (b) sampled sequence $x[n]$, and (c) reconstructed signal $x_r(t)$.

7. (a) See plot below.




FIGURE 6.11: Plot of the signal $x_c(t)$.




FIGURE 6.12: Plot of the CTFT $X_c(j2\pi F)$ of signal $x_c(t)$.

(b) See plot below.



(c) See plot below.

(d) See plot below.

(e) See plot below.

MATLAB script:



```
% P0607: Sampling a exponential decaying signal
close all; clc
%% ii = 1, Fs = 1000; ii = 2, Fs = 5000
ii = 2;
t1 = -5e-3; t2 = 5e-3;
dt = 1e-5;
```

FIGURE 6.13: Plot of sampled signal $x_1[n]$.FIGURE 6.14: Plot of the CTFT $X_1(e^{j2\pi F})$ of sampled signal $x_1[n]$.

```

t = t1:dt:t2;
xc = exp(-1000*abs(t));
F = linspace(-2e3,2e3,1000);
Xc = 0.002./(1+(0.002*pi*F).^2);
Fs = [1e3 5e3];
nT = t1:1/Fs(ii):t2;
xn = exp(-1000*abs(nT));
a = exp(-1000/Fs(ii));
X = (1-a^2)./(1-2*a*cos(2*pi*F/Fs(ii))+a^2);
[G1,G2] = meshgrid(t,nT);
S = sinc(Fs(ii)*(G1-G2));
yr = xn*S;

%% Plot:
hfa = figconfig('P0607a1','long');
```




FIGURE 6.15: Plot of sampled signal $x_2[n]$.FIGURE 6.16: Plot of the CTFT $X_2(e^{j2\pi F})$ of sampled signal $x_2[n]$.

```

plot(t,xc)
xlabel('t (sec)', 'fontsize', LFS)
title('x_c(t)', 'fontsize', TFS)
hfb = figconfig('P0607a2', 'long');
plot(F/1000,abs(Xc))
xlabel('F (kHz)', 'fontsize', LFS)
title('|X_c(j2*\pi F)|', 'fontsize', TFS)

hfc = figconfig('P0607b1', 'long');
plot(t,xc, 'color', [1 1 1]*0.5);hold on
stem(nT,xn, 'filled')
xlabel('n', 'fontsize', LFS)
title(['x_', num2str(ii), '[n]'], 'fontsize', TFS)
hfd = figconfig('P0607b2', 'long');
plot(F/1000,abs(X))

```

FIGURE 6.17: Reconstructed signal $y_{r1}(t)$ from samples $x_1[n]$.FIGURE 6.18: Reconstructed signal $y_{r2}(t)$ from samples $x_2[n]$.

```

ylim([0 max(abs(X))])
xlabel('F (kHz)', 'fontsize', LFS)
title(['|X_', num2str(ii), '(e^{j2*\pi F})|'], 'fontsize', TFS)

hfe = figconfig('P0607e', 'long');
plot(t, yr, 'color', [1 1 1]*0.1); hold on
stem(nT, xn, 'filled')
ylim([min(yr) max(yr)])
xlabel('n', 'fontsize', LFS)
title(['y_{r', num2str(ii), '}[n]'], 'fontsize', TFS)

```

8. Proof:

$$\begin{aligned}
 x_c(t) &= \sum_{k=-\infty}^{\infty} c_k e^{j\frac{2\pi}{T_0}kt} \\
 x_c(nT) &= x_c(nT_0/N) = \sum_{k=-\infty}^{\infty} c_k e^{j\frac{2\pi}{T_0}k\frac{nT_0}{N}} = \sum_{k=-\infty}^{\infty} c_k e^{j\frac{2\pi}{N}kn} \\
 x[n] &= \sum_{k=0}^{N-1} \tilde{c}_k e^{j\frac{2\pi}{N}kn}
 \end{aligned}$$

Since we have $x[n] = x_c(nT)$, we require that

$$\begin{aligned}
 \sum_{k=-\infty}^{\infty} c_k e^{j\frac{2\pi}{N}kn} &= \sum_{k=0}^{N-1} \tilde{c}_k e^{j\frac{2\pi}{N}kn} \\
 \sum_{k=0}^{N-1} \tilde{c}_k e^{j\frac{2\pi}{N}kn} &= \sum_{k=0}^{N-1} \left(\sum_{\ell=-\infty}^{\infty} c_{k-\ell N} \right) e^{j\frac{2\pi}{N}kn} = \sum_{\ell=-\infty}^{\infty} \left(\sum_{k=0}^{N-1} c_{k-\ell N} e^{j\frac{2\pi}{N}kn} \right) \\
 &= \sum_{\ell=-\infty}^{\infty} \left(\sum_{k=0}^{N-1} c_{k-\ell N} e^{j\frac{2\pi}{N}kn} e^{j\frac{2\pi}{N}(-\ell N)} \right) \\
 &= \sum_{k=-\infty}^{\infty} c_k e^{j\frac{2\pi}{N}kn}
 \end{aligned}$$

Hence, we prove that

$$\tilde{c}_k = \sum_{\ell=-\infty}^{\infty} c_{k-\ell N}, \quad k = 0, \pm 1, \pm 2, \dots$$

9. Proof:

$$\begin{aligned}
 \sum_{n=-\infty}^{\infty} y[n] \delta[t - nT] * g_{BL}(t) &= \sum_{\tau=-\infty}^{\infty} \sum_{n=-\infty}^{\infty} y[n] \delta[\tau - nT] \cdot g_{BL}(t - \tau) \\
 &= \sum_{\tau=-\infty}^{\infty} \sum_{n=-\infty}^{\infty} y[n] \delta[t - \tau - nT] \cdot g_{BL}(\tau) \\
 &= \sum_{n=-\infty}^{\infty} y[n] \left(\sum_{\tau=-\infty}^{\infty} \delta[t - \tau - nT] \cdot g_{BL}(\tau) \right) \\
 &= \sum_{n=-\infty}^{\infty} y[n] \cdot g_{BL}(t - nT)
 \end{aligned}$$

10. (a) Solution:

The frequency response of $h_c(t)$ is:

$$H_c(j2\pi F) = \frac{\Omega_n^2}{\Omega_n^2 - (2\pi F)^2 + j2\zeta\Omega_n 2\pi F}$$

Its phase response is:

$$\angle H_c(j2\pi F) = -\tan^{-1} \frac{2\zeta\Omega_n 2\pi F}{\Omega_n^2 - (2\pi F)^2}$$




FIGURE 6.19: Plot of phase response $\angle H_c(j2\pi F)$.

(b) Solution:

The spectra of the sampled sequence is:

$$H(e^{j\omega}) = \frac{e^{-\zeta\Omega_n T} \sin(\Omega_n T \sqrt{1-\zeta^2}) e^{-j\omega}}{1 - 2e^{-\zeta\Omega_n T} \cos(\Omega_n T \sqrt{1-\zeta^2}) e^{-j\omega} + e^{-2\zeta\Omega_n T} e^{-2j\omega}}$$

Its phase response is:

$$\angle H(e^{j\omega}) = -\omega - \tan^{-1} \frac{2e^{-\zeta\Omega_n T} \cos(\Omega_n T \sqrt{1-\zeta^2}) \sin \omega - e^{-2\zeta\Omega_n T} \sin 2\omega}{1 - 2e^{-\zeta\Omega_n T} \cos(\Omega_n T \sqrt{1-\zeta^2}) \cos \omega + e^{-2\zeta\Omega_n T} \cos 2\omega}$$

The effective phase response is:

$$\angle H_{\text{eff}}(j2\pi F) = \begin{cases} \angle H(e^{j2\pi F T}), & |F| \leq \frac{F_s}{2} \\ 0, & |F| > \frac{F_s}{2} \end{cases}$$




FIGURE 6.20: Plot of the effective phase responses, $\angle H_{\text{eff}}(j2\pi F)$ for $F_s = 50$, 100, and 500 Hz.

(c) tba.

MATLAB script:

```
% P0610: Example 6.6: Second-order system
close all; clc
Fs = [50 100 500];
zeta = 0.3; Omega_n = 30*pi;
F = linspace(-50,50,1000);
Hc = Omega_n^2./((j*2*pi*F).^2+2*zeta*Omega_n*j*2*pi*F+Omega_n^2);
[FG FsG] = meshgrid(F,Fs);
H = Omega_n./sqrt(1-zeta^2)*exp(-zeta*Omega_n./FsG)...
    .*sin(Omega_n./FsG*sqrt(1-zeta^2)).*exp(-j*2*pi*FG./FsG)...
    ./ (1-2*exp(-zeta*Omega_n./FsG).*cos(Omega_n./FsG*sqrt(1-zeta^2))...
    .*exp(-j*2*pi.*FG./FsG)+exp(-2*zeta*Omega_n./FsG)...
    .*exp(-2*j*2*pi.*FG./FsG));
%% Plot:
hfb = figconfig('P0610a','small');
plot(F,angle(Hc));
xlabel('F (Hz)','fontsize',LFS)
title('Phase Response \angle H_c(j2\pi F)','fontsize',TFS)
hfc = figconfig('P0610b','small');
plot(F,angle(H));
xlabel('F (Hz)','fontsize',LFS)
title('Phase Response \angle H_{\text{eff}}(j2\pi F)','fontsize',TFS)
legend(['Fs = ',num2str(Fs(1))],['Fs = ',num2str(Fs(2))],...
```

```
[ 'Fs = ', num2str(Fs(3)) ], 'location', 'best' )
```

11. tba

12. (a) Solution:

The quantizer resolution is:

$$\frac{10\text{v}}{2^8} = 0.0390625\text{v}$$

(b) Solution:

$$\text{SQNR} = 10 \log_{10} \text{SQNR} = 6.02B + 1.76 = 6.02 \times 8 + 1.76 = 49.92\text{dB}$$

(c) Solution:

The sampling rate is:

$$F_s = \frac{2^{11}}{2^3} = 2^8 \text{sam/sec}$$

The folding frequency is $F_s/2 = 2^7$.

The Nyquist rate is 500.

(d) Solution:

The reconstructed signal $y_c(t)$ is:

$$y_c(t) = 2 \cos(200\pi t) - 3 \sin(12\pi t)$$

13. Proof:

(i) Linearity.

$$a_1 \cdot x_{\text{in}1}(nT) + a_2 \cdot x_{\text{in}2}(nT) = a_1 \cdot x_{\text{out}1}(t) + a_2 \cdot x_{\text{out}2}(t)$$

The S&H system follows the superposition property, and hence is a linear system.

(ii) Time-variance.

$$x_{\text{out}}(t - \tau) \neq x_{\text{out}}(t), \text{ if } t - \tau \notin [nT, (n+1)T]$$

Hence, the system is time-varying.

14. (a) Solution:

$$g_{\text{SH}}(t) = \begin{cases} 1, & 0 \leq t \leq T \\ 0, & \text{otherwise} \end{cases} \xrightarrow{\text{CTFT}} G_{\text{SH}}(j\Omega) = \frac{2 \sin(\Omega T/2)}{\Omega} e^{-j\Omega T/2}$$

$$H_r(j\Omega) = \begin{cases} \frac{\Omega T/2}{\sin(\Omega T/2)} \cdot e^{j\Omega T/2}, & |\Omega| < \pi/T \\ 0, & \text{otherwise} \end{cases}$$

The frequency response is:

$$H_r(e^{j\omega}) = \begin{cases} \frac{\omega/2}{\sin(\omega/2)} \cdot e^{j\omega/2}, & |\omega| < \pi \\ 0, & \text{otherwise} \end{cases}$$




FIGURE 6.21: Magnitude response of ideal digital filter $H_r(e^{j\omega})$.



(b) Solution: The magnitude response of $H_{\text{FIR}}(e^{j\omega})$ is:

$$\begin{aligned} |H_{\text{FIR}}(e^{j\omega})| &= \sqrt{\left(-\frac{1}{16} + \frac{9}{8} \cos \omega - \frac{1}{16} \cos 2\omega\right)^2 + \left(-\frac{9}{8} \sin \omega + \frac{1}{16} \sin 2\omega\right)^2} \\ &= \sqrt{\frac{1}{16^2} + \frac{9^2}{8^2} + \frac{1}{16^2} - 4 \times \frac{9}{8} \times 116 \cos \omega + \frac{2}{16^2} \cos 2\omega} \end{aligned}$$

(c) Solution: The magnitude response of $H_{\text{IIR}}(e^{j\omega})$ is:

$$|H_{\text{IIR}}(e^{j\omega})| = \frac{9}{\sqrt{(8 + \cos \omega)^2 + \sin^2 \omega}} = \frac{9}{\sqrt{1 + 8^2 + 2 \cos \omega}}$$

MATLAB script:

FIGURE 6.22: Magnitude response of low-order FIR filter $H_{\text{FIR}}(e^{j\omega})$.FIGURE 6.23: Magnitude response of low-order IIR filter $H_{\text{IIR}}(e^{j\omega})$.

```
% P0614: Investigate droop distortion compensation
close all; clc
w = linspace(-1,1,1000)*pi;
%% Part (a):
Hr = w/2./sin(w/2).*exp(j*w/2);
%% Part (b):
HFIR = -1/16+9/8*exp(-j*w)-1/16*exp(-2*j*w);
%% Part (c):
HIIR = 9./(8+exp(-j*w));
%% Plot:
hfa = figconfig('P0614a','small');
plot(w/pi,abs(Hr));
xlabel('\omega/\pi','fontsize',LFS)
```



```

title('Magnitude Response |H_r(e^{j\omega})|', 'fontsize', TFS)
hfb = figconfig('P0614b', 'small');
plot(w/pi, abs(HFIR));
xlabel('\omega/\pi', 'fontsize', LFS)
title('Magnitude Response |H_{FIR}(e^{j\omega})|', 'fontsize', TFS)
hfc = figconfig('P0614c', 'small');
plot(w/pi, abs(HIIR));
xlabel('\omega/\pi', 'fontsize', LFS)
title('Magnitude Response |H_{IIR}(e^{j\omega})|', 'fontsize', TFS)

```

15. Solution:

$$g_r(t) = \frac{\sin(\pi Bt)}{\pi Bt} \cos(2\pi F_c t) \quad (6.74)$$

The spectra is:

$$G_r(j2\pi F) = \begin{cases} 1, & |F| \in [F_L, F_H] \\ 0, & \text{otherwise} \end{cases}$$

Baseband spectra is:

$$G(j2\pi F) = \begin{cases} T, & |F| \leq B/2 \\ 0, & |F| > B/2 \end{cases}$$

The continuous time signal is

$$g(t) = \frac{\sin(\pi t B)}{\pi t B}$$

We can conclude that

$$G_r(j2\pi F) = G[j2\pi(F + F_c)] + G[j2\pi(F - F_c)]$$

Hence,

$$\begin{aligned} g_r(t) &= \frac{\sin(\pi Bt)}{\pi Bt} (e^{-j2\pi F_c t} + e^{j2\pi F_c t}) \\ &= \frac{\sin(\pi Bt)}{\pi Bt} \cos(2\pi F_c t) \end{aligned}$$

where

$$F_c = \frac{F_H + F_L}{2}, \quad B = F_H - F_L$$

16. (a) See plot below.




FIGURE 6.24: Spectrum of the sampled signal as a function of F Hz when the sampling rate is $F_s = 801$.

(b) See plot below.




FIGURE 6.25: Spectrum of the sampled signal as a function of F Hz when the sampling rate is $F_s = 201$.

(c) tba.

```
% P0616: Sampling Illustration
close all; clc
%% Part (a):
% Fs = 801;
%% Part (b):
Fs = 201;
dF = 1;
F = 0:dF:Fs;
```

```

X = zeros(size(F));
FcosF = [325 350 375 400];
Fcos = [1 2 3 4];
while any(FcosF > Fs/2)
    ind = FcosF > Fs/2;
    FcosF(ind) = abs(FcosF(ind) - Fs);
end
for jj = -1:1
for ii = 1:length(FcosF)
    ind = abs(F)==abs(FcosF(ii)+jj*Fs);
    X(ind) = X(ind) + Fcos(ii)*Fs;
end
end
ind = X==0;
X(ind) = nan;
%% Plot:
hfa = figconfg('P0616a','long');
stem(F,abs(X),'filled');
xlim([0 Fs])
set(gca,'Xtick',[0 Fs/2 Fs])
set(gca,'Xticklabel',{'0','Fs/2','Fs'})
set(gca,'XGrid','on')
xlabel('F','fontsize',LFS)
title(['|X(e^{j2\pi F/Fs})|', Fs = ',num2str(Fs)'],'fontsize',TFS)

```

17. Solution:

$$F'_L = 105 - 5 = 100\text{Hz}, \quad F'_H = 145 + 5 = 150\text{Hz}$$

The bandwidth is

$$B = F'_H - F'_L = 50\text{Hz}$$

The minimum sampling rate is computed by


$$\min F_s = 2F'_H / \lfloor F'_H / B \rfloor = 100\text{Hz}$$

MATLAB script:

```

% P0617: Sampling Illustration
close all; clc

```

FIGURE 6.26: Spectrum of the sampled signal as a function of F Hz.

```

FL = 105; FH = 145;
dF = 1;
F = -150:dF:150;
Fs = 100;
X = zeros(size(F));
XP = zeros(size(F));
XN = zeros(size(F));
for jj = -10:10
    ind = F > 105+jj*Fs & F < 145+jj*Fs;
    X(ind) = X(ind) + 1;
    XP(ind) = XP(ind) + 1;
    ind = F > -145+jj*Fs & F < -105+jj*Fs;
    X(ind) = X(ind) + 1;
    XN(ind) = XN(ind) + 1;
end
ind = X == 0;
X(ind) = nan;
%% Plot:
hfa = figconfig('P0617a','long');
plot(F,XP,F,XN);
ylim([0 1.5])
set(gca,'YTick',-1)
xlabel('F (Hz)','fontsize',LFS)
title(['|X(e^{j2\pi F/Fs})|', Fs = ',num2str(Fs)'],'fontsize',TFS)
legend('Right Replicas','Left Replicas','location','best')

```

18. Proof:

$$p_c(x, y) = \begin{cases} 1/A^2, & |x| < A/2, \quad |y| < A/2 \\ 0, & \text{otherwise} \end{cases} \quad (6.89)$$

$$P_c(F_x, F_y) = \frac{\sin(\pi F_x A)}{\pi F_x A} \times \frac{\sin(\pi F_y A)}{\pi F_y A} \quad (6.90)$$

$$\begin{aligned} P_c(F_x, F_y) &= \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} p_c(x, y) e^{-j2\pi(xF_x + yF_y)} dx dy \\ &= \int_{-\frac{A}{2}}^{\frac{A}{2}} \int_{-\frac{A}{2}}^{\frac{A}{2}} \frac{1}{A^2} e^{-j2\pi(xF_x + yF_y)} dx dy \\ &= \left(\int_{-\frac{A}{2}}^{\frac{A}{2}} \frac{1}{A} e^{-j2\pi x F_x} dx \right) \cdot \left(\int_{-\frac{A}{2}}^{\frac{A}{2}} \frac{1}{A} e^{-j2\pi y F_y} dy \right) \\ &= \frac{e^{-j2\pi x F_x} \Big|_{-\frac{A}{2}}^{\frac{A}{2}}}{A \cdot (-j2\pi F_x)} \cdot \frac{e^{-j2\pi y F_y} \Big|_{-\frac{A}{2}}^{\frac{A}{2}}}{A \cdot (-j2\pi F_y)} \\ &= \frac{\sin(\pi F_x A)}{\pi F_x A} \times \frac{\sin(\pi F_y A)}{\pi F_y A} \end{aligned}$$

19. (a) Solution:

$$s_c(x, y) = 3 \cos(2.4\pi x + 2.6\pi y) = 3 \cos(2.4\pi x) \cos(2.6\pi y) - 3 \sin(2.4\pi x) \sin(2.6\pi y)$$

$$s[m, n] = 3 \cos(0.8\pi m + 1.3\pi n)$$

$$s_r(x, y) = 3 \cos(1.6\pi x - 2.6\pi y)$$

(b) Solution:

$$s[m, n] = 3 \cos(1.2\pi m + 0.8667\pi n)$$

$$s_r(x, y) = 3 \cos(2.4\pi x - 1.4\pi y)$$

(c) Solution:

$$s[m, n] = 3 \cos(0.8\pi m + 0.8667\pi n)$$

$$s_r(x, y) = 3 \cos(2.4\pi x + 2.6\pi y)$$

20. (a) Solution:

$$s_{fa}[m, n] = \frac{1}{\Delta x \Delta y} \int_{m\Delta x - \frac{\Delta x}{2}}^{m\Delta x + \frac{\Delta x}{2}} \int_{n\Delta y - \frac{\Delta y}{2}}^{n\Delta y + \frac{\Delta y}{2}} s_c(x, y) dx dy$$

(b) tba

(c) tba

Basic Problems

21. Proof:

The sampler is:

$$x_{\text{out}}(t) = x_{\text{in}}(nT); \quad nT \leq t < (n+1)T, \quad \forall n$$

(i) Memoryless. The current system value is only related to the current time index and is not affected by previous system values. Hence, the sampler is memoryless.

(ii) Linearity.

$$a_1 \cdot x_{\text{in1}}(nT) + a_2 \cdot x_{\text{in2}}(nT) = a_1 \cdot x_{\text{out1}}(t) + a_2 \cdot x_{\text{out2}}(t)$$

The S&H system follows the superposition property, and hence is a linear system.

(iii) Time-variance.

$$x_{\text{out}}(t - \tau) \neq x_{\text{out}}(t), \text{ if } t - \tau \notin [nT, (n+1)T]$$

Hence, the system is time-varying.

22. Solution:

The spectra of the continuous signal $x_c(t)$ is

$$X_c(j2\pi F) = \begin{cases} 3, & F = 0 \\ j, & F = 8 \\ -j, & F = -8 \\ 5, & F = 12 \\ 5, & F = -12 \\ 0, & \text{otherwise} \end{cases}$$

The spectra of sampled sequence $x[n]$ is:

$$X(e^{j\omega}) \Big|_{\omega=2\pi F/F_s} = F_s \sum_{n=-\infty}^{\infty} X_c[j2\pi(F - nF_s)]$$

$x_c(t)$ can be recovered if (a) $F_s = 30$ Hz, and can NOT be recovered if (b) $F_s = 20$ Hz, (c) $F_s = 15$ Hz.

MATLAB script:


```

% P0622: Illustrates the alias distortion
close all; clc
Fs = 30; % Part (a)
% Fs = 20; % Part (b)
% Fs = 15; % Part (c)
T = 1/Fs;
FH = 24;
FL = FH+Fs;
F = -FL:1:FL;
X = zeros(1,length(F));
for k = -5:5;
ind = F == k*Fs; X(ind) = X(ind)+3*Fs;
ind = F == -12+k*Fs; X(ind) = X(ind)+5*Fs;
ind = F == -8+k*Fs; X(ind) = X(ind)-1/j*Fs;
ind = F == 8+k*Fs; X(ind) = X(ind)+1/j*Fs;
ind = F == 12+k*Fs; X(ind) = X(ind)+5*Fs;
end
ind = X==0;
X(ind) = nan;
%% Plot:
hfa = figconfig('P0622a');
subplot(211)
stem(F*2*pi*T,abs(X),'filled')
ylim([0 max(abs(X))+0.5])
set(gca,'XTick',[-Fs*2*pi*T 0 Fs*2*pi*T])
set(gca,'XTickLabel',{'-\Omega_s*T','0','\Omega_s*T'})
xlabel('\omega','fontsize',LFS)
ylabel('|H(e^{j\omega})|','fontsize',LFS)
title('Magnitude Response','fontsize',TFS)
subplot(212)
stem(F*2*pi*T,angle(X),'filled')
set(gca,'XTick',[-Fs*2*pi*T 0 Fs*2*pi*T])
set(gca,'XTickLabel',{'-\Omega_s*T','0','\Omega_s*T'})
xlabel('\omega','fontsize',LFS)
ylabel('\angle H(e^{j\omega})|','fontsize',LFS)
title('Phase Response','fontsize',TFS)


hfb = figconfig('P0622b');
subplot(211)
stem(F,abs(X),'filled')

```

```
ylim([0 max(abs(X))+0.5])
set(gca,'XTick',[-Fs -Fs/2 0 Fs/2 Fs])
set(gca,'XTickLabel',{'-Fs','Fs/2','0','Fs/2','Fs'})
xlabel('F','fontsize',LFS)
ylabel('|H(e^{j2\pi FT})|','fontsize',LFS)
title('Magnitude Response','fontsize',TFS)
subplot(212)
stem(F,angle(X),'filled')
set(gca,'XTick',[-Fs -Fs/2 0 Fs/2 Fs])
set(gca,'XTickLabel',{'-Fs','Fs/2','0','Fs/2','Fs'})
xlabel('F','fontsize',LFS)
ylabel('\angle H(e^{j2\pi FT})','fontsize',LFS)
title('Phase Response','fontsize',TFS)
```





(a)




(b)

FIGURE 6.27: Spectra of $X(e^{j\omega})$ as a function of (a) ω in $\frac{\text{rad}}{\text{sam}}$ and (b) F in Hz when the sample rate is $F_s = 30$ KHz.




(a)




(b)

FIGURE 6.28: Spectra of $X(e^{j\omega})$ as a function of (a) ω in $\frac{\text{rad}}{\text{sam}}$ and (b) F in Hz when the sample rate is $F_s = 20$ KHz.



(a)



(b)

FIGURE 6.29: Spectra of $X(e^{j\omega})$ as a function of (a) ω in $\frac{\text{rad}}{\text{sam}}$ and (b) F in Hz when the sample rate is $F_s = 15$ KHz.

23. Solution:

The spectra of the continuous signal $x_c(t)$ is:

$$X_c(j\Omega) = \begin{cases} 5, & \Omega = 40 \\ 3, & \Omega = -70 \end{cases}$$

The spectra of the sampled sequence $x[n]$ is:

$$X(e^{j\omega})|_{\omega=\Omega T} = \frac{1}{T} \sum_{k=-\infty}^{\infty} X_r[j(\Omega - k\Omega_s)]$$

The continuous signal $x_c(t)$ can be recovered if the sampling interval is (a) $T = 0.01$, (b) $T = 0.04$, and can NOT be recovered if the sampling interval is (c) $T = 0.1$.




FIGURE 6.30: Magnitude and phase responses of $X(e^{j\omega})$ as a function of ω in $\frac{\text{rad}}{\text{sam}}$ when the sampling interval is $T = 0.01$.

MATLAB script:

```
% P0623: Illustrates the alias distortion
close all; clc
% T = 0.01; % Part (a)
```




FIGURE 6.31: Magnitude and phase responses of $X(e^{j\omega})$ as a function of ω in $\frac{\text{rad}}{\text{sam}}$ when the sampling interval is $T = 0.04$.

```
% T = 0.04; % Part (b)
T = 0.1; % Part (c)
Fs = 1/T;
Omegas = 2*pi*Fs;
OmegaH = 70;
OmegaL = round(OmegaH+Omegas);
Omega = -OmegaL:1:OmegaL;
X = zeros(1,length(Omega));
for k = -10:10;
    ind = Omega == round(40+k*Omegas); X(ind) = X(ind)+5*Fs;
    ind = Omega == round(-70+k*Omegas); X(ind) = X(ind)+3*Fs;
end
ind = X==0;
X(ind) = nan;
%% Plot:
hfa = figconf('P0623a');
subplot(211)
stem(Omega*T,abs(X),'filled')
```




FIGURE 6.32: Magnitude and phase responses of $X(e^{j\omega})$ as a function of ω in $\frac{\text{rad}}{\text{sam}}$ when the sampling interval is $T = 0.1$.

```

ylim([0 max(abs(X))+0.5])
set(gca,'XTick',[-Omegas*T 0 Omegas*T])
set(gca,'XTickLabel',{'-\Omega_s*T','0','\Omega_s*T'})
xlabel('\omega','fontsize',LFS)
ylabel('|H(e^{j\omega})|','fontsize',LFS)
title('Magnitude Response','fontsize',TFS)
subplot(212)
stem(Omega*T,angle(X),'filled')
set(gca,'XTick',[-Omegas*T 0 Omegas*T])
set(gca,'XTickLabel',{'-\Omega_s*T','0','\Omega_s*T'})
xlabel('\omega','fontsize',LFS)
ylabel('\angle H(e^{j\omega})','fontsize',LFS)
title('Phase Response','fontsize',TFS)

```

24. Solution:

The spectra of the continuous signal $x_c(t)$ is:

$$X_c(j2\pi F) = \begin{cases} \frac{2}{5}|F| + 2, & |F| \leq 5 \\ -\frac{4}{5}|F| + 8, & 5 < |F| \leq 10 \\ 0, & \text{otherwise} \end{cases}$$

The spectra of the sampled sequence $x[n]$ is:

$$X(e^{j\omega})|_{\omega=2\pi F/F_s} = F_s \sum_{k=-\infty}^{\infty} X_c[j2\pi(F - kF_s)]$$

The signal $x_c(t)$ can NOT be recovered from $x[n]$ if the sampling rate is (a) $F_s = 10$ Hz and (b) $F_s = 15$ Hz and can be recovered if the sampling rate is (c) $F_s = 30$ Hz.




FIGURE 6.33: Magnitude response of $X(e^{j\omega})$ as a function of ω in $\frac{\text{rad}}{\text{sam}}$ when the sampling rate is $F_s = 10$.




FIGURE 6.34: Magnitude response of $X(e^{j\omega})$ as a function of ω in $\frac{\text{rad}}{\text{sam}}$ when the sampling rate is $F_s = 15$.

MATLAB script:




FIGURE 6.35: Magnitude response of $X(e^{j\omega})$ as a function of ω in $\frac{\text{rad}}{\text{sam}}$ when the sampling rate is $F_s = 30$.

```
% P0624: Illustrates the alias distortion
close all; clc
Fs = 10; % Part (a)
% Fs = 15; % Part (b)
% Fs = 30; % Part (c)
T = 1/Fs;
FH = 10;
FL = FH+Fs;
F = -FL:1:FL;
X = zeros(1,length(F));
for k = -5:5;
    ind = abs(F-k*Fs) <= 5;
    X(ind) = X(ind)+(2/5*abs(F(ind)-k*Fs)+2)*Fs;
    ind = abs(F-k*Fs) > 5 & abs(F-k*Fs) <= 10;
    X(ind) = X(ind)+(-4/5*abs(F(ind)-k*Fs)+8)*Fs;
end
%% Plot:
hfa = figconfig('P0624a','long');
plot(F*2*pi*T,abs(X))
ylim([0 max(abs(X))+1])
xlim([-FL*2*pi*T FL*2*pi*T])
set(gca,'XTick',[-Fs*2*pi*T 0 Fs*2*pi*T])
set(gca,'XTickLabel',{'-\Omega_s*T','0','\Omega_s*T'})
xlabel('\omega','fontsize',LFS)
ylabel('|H(e^{j\omega})|','fontsize',LFS)
title('Magnitude Response','fontsize',TFS)
```


25. Solution:

The spectra of the continuous signal $x_c(t)$ is:

$$X_c(j2\pi F) = \begin{cases} -5|F| + 10, & 0 \leq |F| \leq 1 \\ 5|F|, & 1 \leq |F| \leq 2 \\ 10, & 2 \leq |F| \leq 3 \end{cases}$$

The spectra of the sampled sequence $x[n]$ is:

$$X(e^{j2\pi F/F_s}) = F_s \sum_{k=-\infty}^{\infty} X_c[j2\pi(F - kF_s)]$$

The signal $x_c(t)$ can NOT be recovered from $x[n]$ when the sampling interval is (a) $T = 0.2$, (b) $T = 0.25$, (c) $T = 0.5$.




FIGURE 6.36: Magnitude response of $X(e^{j\omega})$ as a function of F in Hz when the sampling interval is $T = 0.2$.




FIGURE 6.37: Magnitude response of $X(e^{j\omega})$ as a function of F in Hz when the sampling interval is $T = 0.25$.

MATLAB script:




FIGURE 6.38: Magnitude response of $X(e^{j\omega})$ as a function of F in Hz when the sampling interval is $T = 0.5$.

```
% P0625: Illustrates the alias distortion
close all; clc
% T = 0.2; % Part (a)
% T = 0.25; % Part (b)
T = 0.5; % Part (c)
Fs = 1/T;
FH = 3;
FL = FH+Fs;
F = -FL:0.1:FL;
X = zeros(1,length(F));
for k = -5:5;
    ind = abs(F-k*Fs) <= 1;
    X(ind) = X(ind)+(-5*abs(F(ind)-k*Fs)+10)*Fs;
    ind = abs(F-k*Fs) > 1 & abs(F-k*Fs) <= 2;
    X(ind) = X(ind)+(5*abs(F(ind)-k*Fs))*Fs;
    ind = abs(F-k*Fs) > 2 & abs(F-k*Fs) <= 3;
    X(ind) = X(ind)+10*Fs;
end
%% Plot:
hfa = figconfg('P0625a','long');
plot(F,abs(X))
xlim([-FL FL])
ylim([0 max(abs(X))+1])
set(gca,'XTick',[-Fs -Fs/2 0 Fs/2 Fs])
set(gca,'XTickLabel',{'-Fs','-Fs/2','0','Fs/2','Fs'})
xlabel('F','fontsize',LFS)
ylabel('|H(e^{j2\pi FT})|','fontsize',LFS)
```

```
title('Magnitude Response','fontsize',TFS)
```

26. (a) Solution:


The sampled sequence $x[n]$ is:

$$x[n] = x_c(nT) = 3 \cos(0.3\pi n + \pi/4) + 3 \sin(0.8\pi n)$$


(b) Solution:

The constructed signal $y_r(t)$ is:

$$y_r(t) = 3 \cos(300\pi t + \pi/4) + 3 \sin(800\pi t)$$



(a)



(b)

FIGURE 6.39: (a) Plot of $x[n]$ and $x_c(t)$ and (b) plot of $y_r(t)$ when $F_1 = 150$ Hz and $F_2 = 400$ Hz.

(c) Solution:

The sampled sequence $x[n]$ is:

$$x[n] = x_c(nT) = 3 \cos(0.6\pi n + \pi/4) + 3 \sin(1.4\pi n)$$

The constructed signal $y_r(t)$ is:

$$y_r(t) = 3 \cos(600\pi t + \pi/4) - 3 \sin(600\pi t)$$

MATLAB script:




FIGURE 6.40: (a) Plot of $x[n]$ and $x_c(t)$ and (b) plot of $y_r(t)$ when $F_1 = 300$ Hz and $F_2 = 700$ Hz.

```
% P0626: Illustrate Ideal DAC
close all; clc
t1 = 0; t2 = 0.05;
dt = 1e-4;
t = t1:dt:t2;
T = 0.001;
Fs = 1/T;
% F1 = 150; F2 = 400; % Part (a) (b)
F1 = 300; F2 = 700; % Part (c)
xc = 3*cos(2*pi*F1*t+pi/4)+3*sin(2*pi*F2*t);
% Sampling:
nT = t1:T:t2;
xn = 3*cos(2*pi*F1*nT+pi/4)+3*sin(2*pi*F2*nT);
F1y = F1;
while F1y > Fs/2
    F1y = F1y-Fs;
end
F2y = F2;
```

```

while F2y > Fs/2
    F2y = F2y-Fs;
end
yr = 3*cos(2*pi*F1y*t+pi/4)+3*sin(2*pi*F2y*t);
%% Plot:
hfa = figconfg('P0626a','long');
plot(t,xc); hold on
plot(nT,xn, '.', 'color','red')
xlabel('t','fontsize',LFS)
title(['F_1 = ',num2str(F1),', F_2 = ',num2str(F2)],'fontsize',TFS)
legend('x_c(t)', 'x[n]', 'location','northeast')
hfb = figconfg('P0626b','long');
plot(t,xc,t,yr,nT,xn, '.')
xlabel('t','fontsize',LFS)
title(['F_1 = ',num2str(F1),', F_2 = ',num2str(F2)],'fontsize',TFS)
legend('x_c(t)', 'y_r(t)', 'x[n]', 'location','northeast')

```

27. (a) Solution:

$$B = \frac{64k}{8k} = 8\text{bits/sam}$$

The quantizer step is:

$$\frac{10v}{2^8} = 0.0390625v$$

(b) Solution:

The SQNR is:

$$\text{SQNR} = 10 \log_{10} \text{SQNR} = 6.02B + 1.76 = 6.02 * B + 1.76 = 49.92\text{dB}$$

(c) the folding frequency

Solution:

The folding frequency is $F_s/2 = 4k$.

(d) Solution:

The reconstructed signal $x_r(t)$ is:

$$x_r(t) = -5 \sin[6000\pi t - \pi/2]$$

28. Solution:

$$B = F_H - F_L = 20 - 18.1 = 1.9\text{KHz}$$

The minimum sampling frequency is:

$$\min F_s = 2F_H / \lfloor F_H / B \rfloor = 4\text{KHz}$$

29. Solution:

$$F'_L = F_L - 2\text{KHz} = 1000.5\text{KHz}$$

$$F'_H = F_H + 2\text{KHz} = 1048\text{KHz}$$

The bandwidth is:

$$B = F'_H - F'_L = 47.5\text{KHz}$$

The minimum sampling rate can be computed by

$$\min F_s = 2F'_H / \lfloor F'_H / B \rfloor = 95.2727273\text{KHz}$$




FIGURE 6.41: Baseband signal spectrum after sampling.

MATLAB script:

```
% P0629: Sampling Illustration
close all; clc
FL = 1002.5e3; FH = 1046e3;
dF = 10;
FG = 4e3;
FHH = FH + FG/2;
FLL = FL - FG/2;
Fs = 95.28e3;
F = -1.1*Fs:dF:1.1*Fs;
X = zeros(size(F));
XP = zeros(size(F));
```

```

XN = zeros(size(F));
for jj = -30:30
    ind = F > FL+jj*Fs & F < FH+jj*Fs;
    X(ind) = X(ind) + 1;
    XP(ind) = XP(ind) + 1;
    ind = F > -FH+jj*Fs & F < -FL+jj*Fs;
    X(ind) = X(ind) + 1;
    XN(ind) = XN(ind) + 1;
end
ind = X == 0;
X(ind) = nan;
%% Plot:
hfa = figconf('P0629a','long');
plot(F/1e3,XP,F/1e3,XN);
xlim([-Fs/1e3 Fs/1e3])
ylim([0 1.2])
set(gca,'YTick',-1)
xlabel('F (KHz)','fontsize',LFS)
set(gca,'XTick',[-Fs -Fs/2 0 Fs/2 Fs]/1e3)
set(gca,'XTickLabel',{'-Fs','Fs/2','0','Fs/2','Fs'})
title(['|X(e^{j2\pi F/Fs})|', Fs = ',num2str(Fs/1e3),' KHz']...
    , 'fontsize',TFS)
legend('Right Replicas','Left Replicas','location','best')

```

30. (a) Solution:

(i) when $\Delta x = \Delta y = 0.5$ meter, $F_{sx} = F_{sy} = 2$, the reconstructed signal is:

$$s_r(x, y) = 4 \cos(2\pi y)$$

(ii) when $\Delta x = \Delta y = 0.2$ meter, $F_{sx} = F_{sy} = 5$, the reconstructed signal is:

$$s_r(x, y) = 4 \cos(4\pi x) \cos(4\pi y)$$

(b) tba

31. tba.

Assessment Problems

32. Solution:

The spectra of the continuous signal $x_c(t)$ is:

$$X_c(j2\pi F) = \begin{cases} 2, & |F| = 2\text{KHz} \\ 3, & |F| = 3\text{KHz} \\ \frac{1}{2j}, & F = 7\text{KHz} \\ -\frac{1}{2j}, & F = -7\text{KHz} \\ 0, & \text{otherwise} \end{cases}$$

The spectra of the sampled sequence $x[n]$ is:

$$X(e^{j\omega})|_{\omega=2\pi F/F_s} = F_s \sum_{k=-\infty}^{\infty} X_c[j2\pi(F - kF_s)]$$

The signal $x_c(t)$ can be recovered from $x[n]$ if the sampling rate is (a) $F_s = 20$ and can NOT be recovered if the sampling rate is (b) $F_s = 10$, and (c) $F_s = 5$.

MATLAB script:

```
% P0632: Illustrates the alias distortion
close all; clc
% Fs = 20e3; % Part (a)
% Fs = 10e3; % Part (b)
Fs = 5e3; % Part (c)
T = 1/Fs;
FH = 7e3;
FL = FH+Fs;
F = -FL:10:FL;
X = zeros(1,length(F));
for k = -10:10;
    ind = F == -7e3+k*Fs; X(ind) = X(ind)-1/2/j*Fs;
    ind = F == -3e3+k*Fs; X(ind) = X(ind)+3*Fs;
    ind = F == -2e3+k*Fs; X(ind) = X(ind)+2*Fs;
    ind = F == 7e3+k*Fs; X(ind) = X(ind)+1/2/j*Fs;
    ind = F == 3e3+k*Fs; X(ind) = X(ind)+3*Fs;
    ind = F == 2e3+k*Fs; X(ind) = X(ind)+2*Fs;
end
ind = X==0;
```




```


X(ind) = nan;
%% Plot:
hfa = figconfig('P0632a');
subplot(211)
stem(F*2*pi*T,abs(X),'filled')
ylim([0 max(abs(X))+0.5])
xlim([-FL*2*pi*T FL*2*pi*T])
set(gca,'XTick',[-Fs*2*pi*T 0 Fs*2*pi*T])
set(gca,'XTickLabel',{'-\Omega_s*T','0','\Omega_s*T'})
xlabel('\omega','fontsize',LFS)
ylabel('|H(e^{j\omega})|','fontsize',LFS)
title('Magnitude Response','fontsize',TFS)
subplot(212)
stem(F*2*pi*T,angle(X),'filled')
% ylim([0 max(abs(X))+0.5])
xlim([-FL*2*pi*T FL*2*pi*T])
set(gca,'XTick',[-Fs*2*pi*T 0 Fs*2*pi*T])
set(gca,'XTickLabel',{'-\Omega_s*T','0','\Omega_s*T'})
xlabel('\omega','fontsize',LFS)
ylabel('\angle H(e^{j\omega})|','fontsize',LFS)
title('Phase Response','fontsize',TFS)

hfb = figconfig('P0632b');
subplot(211)
stem(F,abs(X),'filled')
ylim([0 max(abs(X))+0.5])
xlim([-FL FL])
set(gca,'XTick',[-Fs -Fs/2 0 Fs/2 Fs])
set(gca,'XTickLabel',{'-Fs','-Fs/2','0','Fs/2','Fs'})
xlabel('F','fontsize',LFS)
ylabel('|H(e^{j2\pi FT})|','fontsize',LFS)
title('Magnitude Response','fontsize',TFS)
subplot(212)
stem(F,angle(X),'filled')
xlim([-FL FL])
set(gca,'XTick',[-Fs -Fs/2 0 Fs/2 Fs])
set(gca,'XTickLabel',{'-Fs','-Fs/2','0','Fs/2','Fs'})
xlabel('F','fontsize',LFS)
ylabel('\angle H(e^{j2\pi FT})','fontsize',LFS)
title('Phase Response','fontsize',TFS)

```




(a)




(b)

FIGURE 6.42: Spectra of $X(e^{j\omega})$ as a function of (a) ω in $\frac{\text{rad}}{\text{sam}}$ and (b) F in Hz when the sample rate is $F_s = 20$ KHz.




(a)




(b)

FIGURE 6.43: Spectra of $X(e^{j\omega})$ as a function of (a) ω in $\frac{\text{rad}}{\text{sam}}$ and (b) F in Hz when the sample rate is $F_s = 10$ KHz.



(a)



(b)

FIGURE 6.44: Spectra of $X(e^{j\omega})$ as a function of (a) ω in $\frac{\text{rad}}{\text{sam}}$ and (b) F in Hz when the sample rate is $F_s = 5$ KHz.

33. Solution:

$$x_c(t) = 8 + 12e^{-j20\pi(t-1)} + 7e^{-j40\pi(t+1)} = x_c(t) = 8 + 12e^{-j20\pi t} + 7e^{-j40\pi t}$$

The spectra of the continuous signal $x_c(t)$ is:

$$X_c(j2\pi F) = \begin{cases} 8, & F = 0 \\ 12, & F = -10 \\ 7, & F = -20 \\ 0, & \text{otherwise} \end{cases}$$

The spectra of the sampled sequence $x[n]$ is:

$$X(e^{j\omega})|_{\omega=2\pi F/F_s} = F_s \sum_{k=-\infty}^{\infty} X_c[j2\pi(F - kF_s)]$$

The signal $x_c(t)$ can be recovered from $x[n]$ if the sampling rate is (a) $F_s = 50$ and can NOT be recovered if the sampling rate is (b) $F_s = 20$, and (c) $F_s = 10$.




FIGURE 6.45: Magnitude response of $X(e^{j\omega})$ as a function of F in Hz when the sampling rate is $F_s = 50$.

MATLAB script:

```
% P0633: Illustrates the alias distortion
close all; clc
% Fs = 50; % Part (a)
% Fs = 20; % Part (b)
Fs = 10; % Part (c)
```




FIGURE 6.46: Magnitude response of $X(e^{j\omega})$ as a function of F in Hz when the sampling rate is $F_s = 20$.




FIGURE 6.47: Magnitude response of $X(e^{j\omega})$ as a function of F in Hz when the sampling rate is $F_s = 10$.

```

T = 1/Fs;
FH = 20;
FL = FH+Fs;
F = -FL:1:FL;
X = zeros(1,length(F));
for k = -10:10;
    ind = F == k*Fs; X(ind) = X(ind)+8*Fs;
    ind = F == -20+k*Fs; X(ind) = X(ind)+7*Fs;
    ind = F == -10+k*Fs; X(ind) = X(ind)+12*Fs;
end
ind = X==0;
X(ind) = nan;

```

```

%% Plot:
hfa = figconfig('P0633a','long');
stem(F,abs(X),'filled')
ylim([0 max(abs(X))+0.5])
xlim([-FL FL])
set(gca,'XTick',[-Fs -Fs/2 0 Fs/2 Fs])
set(gca,'XTickLabel',{'-Fs','-Fs/2','0','Fs/2','Fs'})
xlabel('F','fontsize',LFS)
ylabel('|H(e^{j2\pi FT})|','fontsize',LFS)
title('Magnitude Response','fontsize',TFS)

```

34. Solution:

The spectra of the continuous signal $x_c(t)$ is:

$$X_c(j2\pi F) = \begin{cases} 3, & F = 0 \\ \frac{1}{j}, & F = 8 \\ -\frac{1}{j}, & F = -8 \\ 5, & |F| = 12 \\ 0, & \text{otherwise} \end{cases}$$

The spectra of the sampled sequence $x[n]$ is:

$$X(e^{j\omega})|_{\omega=2\pi F/F_s} = F_s \sum_{k=-\infty}^{\infty} X_c[j2\pi(F - kF_s)]$$

The signal $x_c(t)$ can be recovered from $x[n]$ if the sampling rate is (a) $F_s = 30$ Hz, and can NOT be recovered if the sampling rate is (b) $F_s = 20$ Hz, (c) $F_s = 15$ Hz.

MATLAB script:

```

% P0634: Illustrates the alias distortion
close all; clc
% Fs = 30; % Part (a)
% Fs = 20; % Part (b)
Fs = 15; % Part (c)
T = 1/Fs;
FH = 12;
FL = FH+Fs;
F = -FL:1:FL;

```

```


X = zeros(1,length(F));
for k = -10:10;
ind = F == k*Fs; X(ind) = X(ind)+3*Fs;
ind = F == -12+k*Fs; X(ind) = X(ind)+5*Fs;
ind = F == -8+k*Fs; X(ind) = X(ind)-1/j*Fs;
ind = F == 12+k*Fs; X(ind) = X(ind)+5*Fs;
ind = F == 8+k*Fs; X(ind) = X(ind)+1/j*Fs;
end
ind = X==0;
X(ind) = nan;
%% Plot:
hfa = figconfig('P0634a');
subplot(211)
stem(F*2*pi*T,abs(X),'filled')
ylim([0 max(abs(X))+0.5])
xlim([-FL*2*pi*T FL*2*pi*T])
set(gca,'XTick',[-Fs*2*pi*T 0 Fs*2*pi*T])
set(gca,'XTickLabel',{'-\Omega_s*T','0','\Omega_s*T'})
xlabel('\omega','fontsize',LFS)
ylabel('|H(e^{j\omega})|','fontsize',LFS)
title('Magnitude Response','fontsize',TFS)
subplot(212)
stem(F*2*pi*T,angle(X),'filled')
xlim([-FL*2*pi*T FL*2*pi*T])
set(gca,'XTick',[-Fs*2*pi*T 0 Fs*2*pi*T])
set(gca,'XTickLabel',{'-\Omega_s*T','0','\Omega_s*T'})
xlabel('\omega','fontsize',LFS)
ylabel('\angle H(e^{j\omega})|','fontsize',LFS)
title('Phase Response','fontsize',TFS)

hfb = figconfig('P0634b');
subplot(211)
stem(F,abs(X),'filled')
ylim([0 max(abs(X))+0.5])
xlim([-FL FL])
set(gca,'XTick',[-Fs -Fs/2 0 Fs/2 Fs])
set(gca,'XTickLabel',{'-Fs','-Fs/2','0','Fs/2','Fs'})
xlabel('F','fontsize',LFS)
ylabel('|H(e^{j2\pi FT})|','fontsize',LFS)
title('Magnitude Response','fontsize',TFS)


```



```
subplot(212)
stem(F,angle(X),'filled')
xlim([-FL FL])
set(gca,'XTick',[-Fs -Fs/2 0 Fs/2 Fs])
set(gca,'XTickLabel',{'-Fs','-Fs/2','0','Fs/2','Fs'})
xlabel('F','fontsize',LFS)
ylabel('\angle H(e^{j2\pi FT})','fontsize',LFS)
title('Phase Response','fontsize',TFS)
```




(a)




(b)

FIGURE 6.48: Spectra of $X(e^{j\omega})$ as a function of (a) ω in $\frac{\text{rad}}{\text{sam}}$ and (b) F in Hz when the sample rate is $F_s = 30$ KHz.




(a)




(b)

FIGURE 6.49: Spectra of $X(e^{j\omega})$ as a function of (a) ω in $\frac{\text{rad}}{\text{sam}}$ and (b) F in Hz when the sample rate is $F_s = 20$ KHz.



(a)



(b)

FIGURE 6.50: Spectra of $X(e^{j\omega})$ as a function of (a) ω in $\frac{\text{rad}}{\text{sam}}$ and (b) F in Hz when the sample rate is $F_s = 15$ KHz.

35. The same as P0623

36. Solution:

The spectra of the sampled sequence $x[n]$ is:

$$X(e^{j\omega})|_{\omega=\Omega/F_s} = F_s \sum_{k=-\infty}^{\infty} X_c[j(\Omega - 2\pi k F_s)]$$

The signal $x_c(t)$ can NOT be recovered when the sampling interval is (a) $T = \pi$, (b) $T = 0.5\pi$, (c) $T = 0.2\pi$.




FIGURE 6.51: Magnitude response of $X(e^{j\omega})$ as a function of ω in $\frac{\text{rad}}{\text{sam}}$ when the sampling interval is $T = \pi$.




FIGURE 6.52: Magnitude response of $X(e^{j\omega})$ as a function of ω in $\frac{\text{rad}}{\text{sam}}$ when the sampling interval is $T = 0.5\pi$.

MATLAB script:

```
% P0636: Plot the spectra of sampled sequence
close all; clc
T = pi; % Part a
% T = 0.5*pi; % Part b
```




FIGURE 6.53: Magnitude response of $X(e^{j\omega})$ as a function of ω in $\frac{\text{rad}}{\text{sam}}$ when the sampling interval is $T = 0.2\pi$.

```
% T = 0.2*pi; % Part c
Fs = 1/T;
Omeegas = 2*pi/T;
Omega = -2*Omeegas:0.01:2*Omeegas;
X = zeros(size(Omega));
for k = -10:1:10
    X = X + pi*exp(-abs(Omega-k*Omeegas))/T;
end
%% Plot:
hfa = figconf('P0636a','long');
plot(Omega*T,abs(X))
xlabel('\omega (rad/sam)','fontsize',LFS)
ylabel('|H(e^{j2\pi F})|','fontsize',LFS)
ylim([0 max(abs(X))*1.1])
xlim([-2*Omeegas*T 2*Omeegas*T])
title('Magnitude Response','fontsize',TFS)
```

37. Solution:

The spectra of the continuous signal $x_c(t)$ is:

$$X_c(j2\pi F) = \begin{cases} 1 - \frac{F^2}{25}, & |F| \leq 5 \\ 0, & \text{otherwise} \end{cases}$$

The spectra of the sampled sequence $x[n]$ is:

$$X(e^{j\omega})|_{\omega=2\pi F/F_s} = F_s \sum_{k=-\infty}^{\infty} X_c[j2\pi(F - kF_s)]$$

The continuous signal $x_c(t)$ can NOT be recovered from the sequence $x[n]$ when the sampling frequency is (a) $F_s = 6$ Hz, (b) $F_s = 4$ Hz, (c) $F_s = 2$ Hz.




FIGURE 6.54: Magnitude response of $X(e^{j\omega})$ as a function of ω in $\frac{\text{rad}}{\text{sam}}$ when the sampling rate is $F_s = 6$.




FIGURE 6.55: Magnitude response of $X(e^{j\omega})$ as a function of ω in $\frac{\text{rad}}{\text{sam}}$ when the sampling rate is $F_s = 4$.

MATLAB script:

```
% P0637: Illustrates the alias distortion
close all; clc
Fs = 6; % Part (a)
% Fs = 4; % Part (b)
% Fs = 2; % Part (c)
T = 1/Fs;
FH = 5;
FL = FH+Fs;
F = -FL:0.01:FL;
X = zeros(1,length(F));
for k = -5:5;
```




FIGURE 6.56: Magnitude response of $X(e^{j\omega})$ as a function of ω in $\frac{\text{rad}}{\text{sam}}$ when the sampling rate is $F_s = 2$.

```
ind = abs(F-k*Fs) <= 5;
X(ind) = X(ind)+(1-(F(ind)-k*Fs).^2/25)*Fs;
end
%% Plot:
hfa = figconfg('P0637a','long');
plot(F*2*pi*T,abs(X))
ylim([0 max(abs(X))+1])
xlim([-FL*2*pi*T FL*2*pi*T])
set(gca,'XTick',[-Fs*2*pi*T 0 Fs*2*pi*T])
set(gca,'XTickLabel',{'-\Omega_s*T','0','\Omega_s*T'})
xlabel('\omega (rad/sam)','fontsize',LFS)
ylabel('|H(e^{j\omega})|','fontsize',LFS)
title('Magnitude Response','fontsize',TFS)
```

38. Proof:

The ideal DAC is represented by

$$y_r(t) = \sum_{n=-\infty}^{\infty} y[n]g_{BL}(t - nT) \quad (6.43)$$

(i) Non-causality.

when $n < 0$, $y_r(t)$ is related to the future values, hence the system is non-causal.

(ii) Linearity.

when the input is $y[n] = a_1 y_1[n] + a_2 y_2[n]$, the output of the system is:

$$\begin{aligned} y_r(t) &= \sum_{n=-\infty}^{\infty} (a_1 y_1[n] + a_2 y_2[n]) \cdot g_{BL}(t - nT) \\ &= a_1 \sum_{n=-\infty}^{\infty} y_1[n] g_{BL}(t - nT) + a_2 \sum_{n=-\infty}^{\infty} y_2[n] g_{BL}(t - nT) \\ &= a_1 y_{r1}(t) + a_2 y_{r2}(t) \end{aligned}$$

which verifies that the system follows superposition property, hence the system is linear.

(iii) Time variance.

In general, we have

$$y_r(t - \tau) = \sum_{n=-\infty}^{\infty} y[n] g_{BL}(t - \tau - nT) \neq \sum_{n=-\infty}^{\infty} y[n] g_{BL}(t - nT) = y_r(t)$$

which proves the system is time varying.

39. (a) Solution:

The sampled sequence is:

$$x[n] = x_c(0.01n) = 10 + 3 \sin(0.2\pi n + \pi/3) + 5 \cos(0.4\pi n)$$

(b) Solution:

The reconstructed signal is:

$$y_r(t) = 10 + 3 \sin(20\pi t + \pi/3) + 5 \cos(40\pi t)$$

(c) Solution:

The sampled sequence is:

$$x[n] = x_c(0.05n) = 10 + 3 \sin(\pi n + \pi/3) + 5 \cos(2\pi n)$$

The reconstructed signal is:

$$y_r(t) = 10 + 3 \sin(20\pi t + \pi/3)$$

(d) Solution:

The sampled sequence is:

$$x[n] = x_c(0.1n) = 10 + 3 \sin(2\pi n + \pi/3) + 5 \cos(4\pi n)$$




FIGURE 6.57: (a) Plot of $x[n]$ and $x_c(t)$ and (b) plot of $y_r(t)$ when the signal is sampled at $t = 0.01n$.

The reconstructed signal is:

$$y_r(t) = 10 - \frac{3\sqrt{3}}{2} + 5 = 15 - \frac{3\sqrt{3}}{2}$$

MATLAB script:

```
% P0639: Illustrate Ideal DAC
close all; clc
t1 = 0; t2 = 1;
dt = 1e-4;
t = t1:dt:t2;
% T = 0.01; % Part (a) (b)
% T = 0.05; % Part (c)
T = 0.1; % Part (d)
Fs = 1/T;
F1 = 10; F2 = 20;
xc = 10 + 3*sin(2*pi*F1*t+pi/3)+5*cos(2*pi*F2*t);
% Sampling:
```




FIGURE 6.58: (a) Plot of $x[n]$ and $x_c(t)$ and (b) plot of $y_r(t)$ when the signal is sampled at $t = 0.05n$.

```

nT = t1:T:t2;
xn = 10 + 3*sin(2*pi*F1*nT+pi/3)+5*cos(2*pi*F2*nT);
F1y = F1;
while F1y > Fs/2
    F1y = F1y-Fs;
end
F2y = F2;
while F2y > Fs/2
    F2y = F2y-Fs;
end
yr = 10 + 3*sin(2*pi*F1y*t+pi/3)+5*cos(2*pi*F2y*t);
%% Plot:
hfa = figconf('P0639a','long');
plot(t,xc); hold on
plot(nT,xn,'.','color','red')
xlabel('t','fontsize',LFS)
title(['T = ',num2str(T)],'fontsize',TFS)
legend('x_c(t)', 'x[n]', 'location', 'northeast')

```




FIGURE 6.59: (a) Plot of $x[n]$ and $x_c(t)$ and (b) plot of $y_r(t)$ when the signal is sampled at $t = 0.1n$.

```
hfb = figconf('P0639b','long');
plot(t,xc,t,yr,nT,xn,'.')
xlabel('t','fontsize',LFS)
title(['T = ',num2str(T)],'fontsize',TFS)
legend('x_c(t)', 'y_r(t)', 'x[n]', 'location', 'northeast')
```

40. Solution:

The sampling frequency interval can be defined as

$$\frac{2F_H}{k} \leq F_s \leq \frac{2F}{k-1}$$

Hence, we compute as $11.2\text{Hz} \leq F_s \leq 12\text{Hz}$.

MATLAB script:

```
% P0640: Sampling Illustration
close all; clc
FL = 24; FH = 28;
dF = 0.01;
```




FIGURE 6.60: Spectra of the sampled sequence.

```

% Fs = 11.2;
% Fs = 12;
Fs = 11.5;
F = -FH:dF:FH;
X = zeros(size(F));
XP = zeros(size(F));
XN = zeros(size(F));
for jj = -10:10
    ind = F > FL+jj*Fs & F < FH+jj*Fs;
    X(ind) = X(ind) + 1;
    XP(ind) = XP(ind) + 1;
    ind = F > -FH+jj*Fs & F < -FL+jj*Fs;
    X(ind) = X(ind) + 1;
    XN(ind) = XN(ind) + 1;
end
ind = X == 0;
X(ind) = nan;
%% Plot:
hfa = figconfig('P0640a','long');
plot(F,XP,F,XN);
xlim([-FH FH])
ylim([0 1.2])
set(gca,'YTick',-1)
xlabel('F (Hz)','fontsize',LFS)
title(['|X(e^{j2\pi F/Fs})|', 'Fs = ',num2str(Fs),' Hz']...
    , 'fontsize',TFS)
legend('Right Replicas','Left Replicas','location','best')

```

41. Solution:

$$F'_L = 76 - 1 = 75\text{Hz}, \quad F'_H = 98 + 1 = 99\text{Hz}$$

The bandwidth is

$$B = F'_H - F'_L = 24\text{Hz}$$

The minimum sampling frequency is:

$$\min F_s = 2F'_H / \lceil F'_H / B \rceil = 49.5\text{Hz}$$




FIGURE 6.61: Plot of the resulting spectrum of the sampled signal over $[-100, 100]$ Hz range.

MATLAB script:

```
% P0641: Sampling Illustration
close all; clc
FL = 76; FH = 98;
dF = 0.01;
FG = 2;
FHH = FH + FG/2;
FLL = FL - FG/2;
Fs = 49.5;
F = -100:dF:100;
X = zeros(size(F));
XP = zeros(size(F));
XN = zeros(size(F));
for jj = -10:10
    ind = F > FL+jj*Fs & F < FH+jj*Fs;
    X(ind) = X(ind) + 1;
    XP(ind) = XP(ind) + 1;
```

```

ind = F > -FH+jj*Fs & F < -FL+jj*Fs;
X(ind) = X(ind) + 1;
XN(ind) = XN(ind) + 1;
end
ind = X == 0;
X(ind) = nan;
%% Plot:
hfa = figconfig('P0641a','long');
plot(F,XP,F,XN);
xlim([-100 100])
ylim([0 1.2])
set(gca,'YTick',-1)
xlabel('F (Hz)','fontsize',LFS)
title(['|X(e^{j2\pi F/Fs})|', Fs = ',num2str(Fs),' Hz']...
      , 'fontsize',TFS)
legend('Right Replicas','Left Replicas','location','best')

```

42. (a) Solution:

(i) when $F_{sx} = 25$ and $F_{sy} = 50$, the reconstructed signal is

$$s_r(x, y) = 2 \cos(2\pi x + 2\pi y)$$

(i) when $F_{sx} = 100$ and $F_{sy} = 200$, the reconstructed signal is

$$s_r(x, y) = 2 \cos(98\pi x + 198\pi y)$$

(b) tba

Review Problems

43. (a) MATLAB script:

```
function [xhat, t] = DAC1(x,N,T)
% Implementation of function (6.100)
% Input:
%       x: sample sequence x[n]
%       N: sample sequence index [0 N]
%       T: sampling time interval
% Output:
%       xhat: reconstructed time signal
%       t: time index
dt = T/50;
t = 0:dt:N*T;
n = 0:N;
[Gt Gn] = meshgrid(t,n);
xhat = x*sinc((Gt-Gn*T)/T);
```

(b) See plot below.

(c) See plot below.

(d) tba.

(e) tba.

(f) tba.

MATLAB script:

```
% P0643: Investigate approximate ideal reconstruction
close all; clc
%% Part (b):
ii = 4;
T = 1;
N = 5:5:20;
n = 0:N(ii);
x = ones(1,N(ii)+1);
[xhat t] = DAC1(x,N(ii),T);
xc = ones(size(t));
e1 = xc - xhat;
% Plot:
hfa = figconfig('P0643a','long');
plot(t,[xc;xhat;e1]); hold on
```



```

plot(n*T,x,'.')
xlabel('t','fontsize',LFS)
title(['N = ',num2str(N(ii)),' ', T = ',num2str(T)],'fontsize',TFS)
legend('x_c(t)', '\hat{x}_c(t)', 'e_1(t)', 'location', 'best')
%% Part (c):
NN = 3:20;
e1size = zeros(size(NN));
for jj = 1:length(NN)
    xtemp = ones(1,NN(jj)+1);
    [xhattemp ttemp] = DAC1(xtemp,NN(jj),T);
    Nthird = round(length(ttemp)/3);
    e1size(jj) = max(abs(1 - xhattemp(Nthird+1:2*Nthird)));
end
% Plot:
hfb = figconfig('P0643b','long');
stem(NN,e1size,'filled')
xlabel('N','fontsize',LFS)
title('Error Size','fontsize',TFS)

```

MATLAB script:

```

% P0643: Investigate approximate ideal reconstruction
close all; clc
%% Part (edf):
F = 1;
theta = 0;
T = 0.1;
ii = 4;
N = 5:5:20;
n = 0:N(ii);
x = cos(2*pi*F*n*T+theta);
[xhat t] = DAC1(x,N(ii),T);
xc = cos(2*pi*F*t+theta);
e1 = xc - xhat;
% Plot:
hfa = figconfig('P0643a','long');
plot(t,[xc;xhat;e1]); hold on
plot(n*T,x,'.')
xlabel('t','fontsize',LFS)
title(['N = ',num2str(N(ii)),' ', T = ',num2str(T)],'fontsize',TFS)

```

```

legend('x_c(t)', '\hat{x}_c(t)', 'e_1(t)', 'location', 'best')

NN = 3:20;
elsize = zeros(size(NN));
for jj = 1:length(NN)
    nn = 0:NN(jj);
    xtemp = cos(2*pi*F*nn*T+theta);
    [xhattemp ttemp] = DAC1(xtemp,NN(jj),T);
    Nthird = round(length(ttemp)/3);
    elsize(jj) = max(abs(cos(2*pi*F*ttemp(Nthird+1:2*Nthird))...
        - xhattemp(Nthird+1:2*Nthird))));
end
% Plot:
hfb = figconfig('P0643b','long');
stem(NN,elsize,'filled')
xlabel('N','fontsize',LFS)
title('Error Size','fontsize',TFS)

```








FIGURE 6.62: Plot of $x_c(t)$, $\hat{x}_c(t)$, and $e_1(t)$ for $T = 1$, and (a) $N = 5$, (b) $N = 10$, (c) $N = 15$, (d) $N = 20$.

FIGURE 6.63: Plot of the size of error as a function of N .

44. (a) MATLAB script:

```
function [xhat, t] = DAC2(x,N,K,T)
% Implementation of function (6.101) (6.102) and (6.103)
% Input:
%     x: sample sequence x[n]
%     N: sample sequence index [0 N]
%     K: argument employed in (6.101)
%     T: sampling time interval
% Output:
%     xhat: reconstructed time signal
%     t: time index
dt = T/50;
t = 0:dt:N*T;
n = 0:N;
[Gt Gn] = meshgrid(t,n);
xhat = x*gk(Gt-Gn*T,K,T);
%% Sub-functions:
function y = gk(t,K,T)
y = sinc(t/T);
ind = abs(t) >= K*T;
y(ind) = 0;
```

(b) See plot below.

(c) See plot below

(d) tba.

(e) tba.

(f) tba.

MATLAB script:

```
% P0644: Investigate approximated ideal reconstruction
close all; clc
%% Part (b):
ii = 1;
K = 20;
T = 1;
N = 5:5:20;
n = 0:N(ii);
x = ones(1,N(ii)+1);
[xhat t] = DAC2(x,N(ii),K,T);
```

```

xc = ones(size(t));
e1 = xc - xhat;
% Plot:
hfa = figconfig('P0644a','long');
plot(t,[xc;xhat;e1]); hold on
plot(n*T,x,'.')
xlabel('t','fontsize',LFS)
title(['N = ',num2str(N(ii)),',      K = ',num2str(K)],'fontsize',TFS)
legend('x_c(t)', '\hat{x}_c(t)', 'e_1(t)', 'location','best')
%% Part (c):
NN = 3:20;
e1size = zeros(size(NN));
for jj = 1:length(NN)
    xtemp = ones(1,NN(jj)+1);
    [xhattemp ttemp] = DAC2(xtemp,NN(jj),K,T);
    Nthird = round(length(ttemp)/3);
    e1size(jj) = max(abs(1 - xhattemp(Nthird+1:2*Nthird)));
end
% Plot:
hfb = figconfig('P0644b','long');
stem(NN,e1size,'filled')
xlabel('N','fontsize',LFS)
title(['Error Size,      K = ',num2str(K)],'fontsize',TFS)

MATLAB script:

% P0644: Investigate approximate ideal reconstruction
close all; clc
%% Part (def):
F = 1;
theta = 0;
T = 0.1;
K = 20;
ii = 4;
N = 5:5:20;
n = 0:N(ii);
x = cos(2*pi*F*n*T+theta);
[xhat t] = DAC2(x,N(ii),K,T);
xc = cos(2*pi*F*t+theta);
e1 = xc - xhat;
% Plot:

```

```

hfa = figconfig('P0644a','long');
plot(t,[xc;xhat;e1]); hold on
plot(n*T,x,'.')
xlabel('t','fontsize',LFS)
title(['N = ',num2str(N(ii)),',      T = ',num2str(T)],'fontsize',TFS)
legend('x_c(t)', '\hat{x}_c(t)', 'e_1(t)', 'location','best')

NN = 3:20;
e1size = zeros(size(NN));
for jj = 1:length(NN)
    nn = 0:NN(jj);
    xtemp = cos(2*pi*F*nn*T+theta);
    [xhattemp ttemp] = DAC2(xtemp,NN(jj),K,T);
    Nthird = round(length(ttemp)/3);
    e1size(jj) = max(abs(cos(2*pi*F*ttemp(Nthird+1:2*Nthird))...
        - xhattemp(Nthird+1:2*Nthird)));
end
% Plot:
hfb = figconfig('P0644b','long');
stem(NN,e1size,'filled')
xlabel('N','fontsize',LFS)
title('Error Size','fontsize',TFS)

```





FIGURE 6.64: Plot of $x_c(t)$, $\hat{x}_c(t)$, and $e_1(t)$ for $T = 1$, and (a) $N = 5$, (b) $N = 10$, (c) $N = 15$, (d) $N = 20$.

FIGURE 6.65: Plot of the size of error as a function of N .

45. (a) MATLAB script:

```
function [eH, e, eavg, evar] = QuantR(xn,B,N)
% Inputs:
%     xn:    sampled sequence
%     B:     # of bits representing one sample
%     N:     # of samples
% Outputs:
%     eH:    Normalized histogram of e[n]
%     e:     bins of histogram
%     eavg:  mean of e[n]
%     evar:  variance of e[n]
step = 1/(2^B);
xqn = round(xn/step)*step;
en = 2^B*(xn-xqn);
[eH e] = hist(en,100);
eH = eH/N;
eavg = mean(en);
evar = var(en);
```

(b) See plot below.

(c) See plot below.

(d) See plot below.

MATLAB script:

```
% P0645: Investigate quantization error
close all; clc
ii = 4;
B = [1 2 4 6];
N = 5e5;
n = 1:N;
xn = cos(n/11); % part (b)
% xn = 1/2*(cos(n/11)+cos(n/17)+sin(n/31)); % part (c)
% xn = rand(size(n)); % part (d)
[eH, e, eavg, evar] = QuantR(xn,B(ii),N);
% Plot:
hfa = figconfg('P0645a','long');
bar(e,eH)
xlabel('e[n]', 'fontsize', LFS)
title(['Normalized Histogram:   B = ', num2str(B(ii)), ...
```

```
    ', mean = ', num2str(eavg), ', var = ', ...  
    num2str(evar)], 'fontsize', TFS)
```







FIGURE 6.66: Plot the resulting distributions for quantization error $e[n]$ of $x[n]$ in part (a) for (a) 1 bit, (b) 2 bits, (c) 4 bits, (d) 6 bits.




FIGURE 6.67: Plot the resulting distributions for quantization error $e[n]$ of $x[n]$ in part (c) for (a) 1 bit, (b) 2 bits, (c) 4 bits, (d) 6 bits.




FIGURE 6.68: Plot the resulting distributions for quantization error $e[n]$ of $x[n]$ in part (d) for (a) 1 bit, (b) 2 bits, (c) 4 bits, (d) 6 bits.

46. (a) MATLAB script:

```
function [eH, e, eavg, evar] = QuantF(xn,B,N)
% Inputs:
%      xn:    sampled sequence
%      B:    # of bits representing one sample
%      N:    # of samples
% Outputs:
%      eH:    Normalized histogram of e[n]
%      e:    bins of histogram
%      eavg:  mean of e[n]
%      evar:  variance of e[n]
step = 1/(2^B);
xqn = fix(xn/step)*step;
en = 2^B*(xn-xqn);
[eH e] = hist(en,100);
eH = eH/N;
eavg = mean(en);
evar = var(en);
```

(b) See plot below.

(c) See plot below.

(d) See plot below.

MATLAB script:

```
% P0646: Investigate quantization error
close all; clc
ii = 4;
B = [1 2 4 6];
N = 5e5;
n = 1:N;
% xn = cos(n/11); % part (b)
% xn = 1/2*(cos(n/11)+cos(n/17)+sin(n/31)); % part (c)
xn = rand(size(n)); % part (d)
[eH, e, eavg, evar] = QuantF(xn,B(ii),N);
% Plot:
hfa = figconfg('P064a6','long');
bar(e,eH)
xlabel('e[n]', 'fontsize', LFS)
title(['Normalized Histogram:   B = ', num2str(B(ii)), ...
```

```
    ', mean = ', num2str(eavg), ', var = ', ...  
    num2str(evar)], 'fontsize', TFS)
```








FIGURE 6.69: Plot the resulting distributions for quantization error $e[n]$ of $x[n]$ in part (a) for (a) 1 bit, (b) 2 bits, (c) 4 bits, (d) 6 bits.







FIGURE 6.70: Plot the resulting distributions for quantization error $e[n]$ of $x[n]$ in part (c) for (a) 1 bit, (b) 2 bits, (c) 4 bits, (d) 6 bits.




(a)



(b)



(c)



(d)

FIGURE 6.71: Plot the resulting distributions for quantization error $e[n]$ of $x[n]$ in part (d) for (a) 1 bit, (b) 2 bits, (c) 4 bits, (d) 6 bits.

47. (a) See plot below.
 (b) See plot below.
 (c) tba.

MATLAB script:

```
% P0647: Investigate 2D sampling
close all; clc
xc = double(imread('DSP.png'));
%% Part (a):
xc_dim = size(xc);
sind = 5;
indx = mod(1:xc_dim(1),10) == sind;
indy = mod(1:xc_dim(2),10) == sind;
x = xc(indx,indy);
frescale = ones(10);
xs = kron(x,frescale);
% Plot:
hfa1 = figconfig('P0647a1');
imshow(xc)
hfa2 = figconfig('P0647a2');
imshow(x)
hfa3 = figconfig('P0647a3');
imshow(xs)
%% Part (b):
fave = fspecial('average',5); % change filter type and size
% fave = fspecial('gaussian',3);
yc = filter2(fave,xc);
y = yc(indx,indy);
ys = kron(y,frescale);
% Plot:
hfb1 = figconfig('P0647b1');
imshow(yc)
hfb2 = figconfig('P0647b2');
imshow(y)
hfb3 = figconfig('P0647b3');
imshow(ys)
```

A large, smooth, black serif font representing the letters 'DSP'.

(a)

A smaller, pixelated version of the 'DSP' text, where each letter is composed of a grid of black and white squares.

(b)

A pixelated version of the 'DSP' text, similar to (b), but with a slightly different arrangement of pixels, possibly representing a reconstructed signal.

(c)

FIGURE 6.72: Image show of (a) x_c , (b) x , (c) x_s .



(a)



(b)



(c)

FIGURE 6.73: Image show of (a) y_c , (b) y , (c) y_s .