

CHAPTER 4

Fourier Representation of Signals

Basic Problems

20. Solution:
 $x[n]$ is always periodic, and the fundamental period N is the least common multiple of N_1, N_2 .
21. (a) Solution:
 $x_1(t)$ is aperiodic.
- (b) Solution:
 $x_2[n]$ is periodic with fundamental period $N = 20$.
- (c) Solution:
 $x_3(t)$ is periodic with fundamental period $T = \frac{1}{500}$.
- (d) Solution:
 $x_4[n]$ is aperiodic.
- (e) Solution:
 $x_5[n]$ is periodic with fundamental period $N = 60$.
22. (a) Solution:
 $x[n]$ is periodic.
- (b) Solution:
The fundamental period of the sequence $x[n]$ is $N = 4$.

23. Solution:

$$c_k = \frac{A(1 - \cos \pi k)}{(\pi k)^2}$$

(a) See plots below.





FIGURE 4.1: (a) Magnitude spectra of $x(t)$ for $A = 1$ and $T_0 = 1$. (b) Phase spectra of $x(t)$ for $A = 1$ and $T_0 = 1$.

(b) See plots below.

FIGURE 4.2: (a) $x(t)$ for $A = 1$ and $T_0 = 1$. (b) $x_5(t)$. (c) $x_{10}(t)$. (d) $x_{30}(t)$.

24. (a) Solution:

$$X(F) = \frac{1}{j2\pi F} + \frac{2e^{-j2\pi F}}{(j2\pi F)^2} + \frac{2(e^{-j2\pi F} - 1)}{(j2\pi F)^3}$$

(b) Solution:

$$X(F) = \frac{-48j\pi^2 F}{(4\pi^2 F^2 + 12j\pi F - 9 - 4\pi^2)(4\pi^2 F^2 - 12j\pi F - 9 - 4\pi^2)}$$

(c) Solution:

$$X_3(F) = \begin{cases} 1, & -\frac{1}{2} \leq F \leq \frac{1}{2} \\ F + \frac{3}{2}, & -\frac{3}{2} \leq F \leq -\frac{1}{2} \\ -F + \frac{3}{2}, & \frac{1}{2} \leq F \leq \frac{3}{2} \\ 0, & \text{otherwise} \end{cases}$$




FIGURE 4.3: (a) Magnitude and phase spectra of signal $x_1(t) = (1 - t^2)[u(t) - u(t - 1)]$.




FIGURE 4.4: (a) Magnitude and phase spectra of signal $x_2(t) = e^{-3|t|} \sin 2\pi t$.




FIGURE 4.5: (a) Magnitude and phase spectra of signal $x_3(t) = \frac{\sin \pi t}{\pi t} \frac{\sin 2\pi t}{\pi t}$.

25. (a) Solution:




FIGURE 4.6: Sequences $\hat{x}_M[n]$. (a) $M = 1$. (b) $M = 2$. (c) $M = 3$. (d) $M = 4$.

26. (a) Solution:

$$c_k = \begin{cases} 0, & k = 5m \\ 6e^{-j\frac{\pi}{6}}, & k = 5m + 1 \\ 4e^{-j\frac{\pi}{3}}, & k = 5m + 2 \\ 4e^{j\frac{\pi}{3}}, & k = 5m + 3 \\ -6e^{j\frac{\pi}{6}}, & k = 5m + 4 \end{cases} \quad m = 0, \pm 1, \pm 2$$




FIGURE 4.7: Magnitude and phase spectra of periodic sequence $x_1[n] = 4 \cos(1.2\pi n + 60^\circ) + 6 \sin(0.4\pi n - 30^\circ)$.

(b) Solution:

$$c_k = \frac{1}{4} \left(1 + 2 \cos \frac{\pi}{4} \cos \frac{\pi}{2} k \right)$$





FIGURE 4.8: Magnitude and phase spectra of periodic sequence $x_2[n] = |\cos(0.25\pi n)|, 0 \leq n \leq 3$.


(c) Solution:

$$c_k = \frac{1}{8} \left[1 + 2 \cos\left(\frac{k\pi}{4}\right) + 2 \cos\left(\frac{3k\pi}{4}\right) + \cos(k\pi) \right]$$

FIGURE 4.9: Magnitude and phase spectra of periodic sequence $x_3[n]$.


(d) Solution:

$$c_k = \frac{1}{12} \left[1 + (1 - \sin(\frac{\pi}{4}))2 \cos(\frac{k\pi}{6}) + (1 - \sin(\frac{3\pi}{4}))2 \cos(\frac{k\pi}{2}) \right. \\ \left. + 2 \cos(\frac{2k\pi}{3}) + (1 - \sin(\frac{5\pi}{4}))2 \cos(\frac{5k\pi}{6}) + 2 \cos(k\pi) \right]$$

FIGURE 4.10: Magnitude and phase spectra of periodic sequence $x_4[n] = 1 - \sin(\pi n/4)$, $0 \leq n \leq 11$ (one period).

(e) Solution:

$$c_k = \frac{1}{7} \left(1 - 2e^{-j\frac{2\pi}{7}k} + e^{-j\frac{2\pi}{7}k \cdot 2} - e^{-j\frac{2\pi}{7}k \cdot 4} + 2e^{-j\frac{2\pi}{7}k \cdot 5} - e^{-j\frac{2\pi}{7}k \cdot 6} \right)$$

FIGURE 4.11: Magnitude and phase spectra of periodic sequence $x_5[n]$.

27. (a) Solution:

$$\frac{1}{N} \sum_{n=0}^{N-1} x[n - n_0] e^{-j\frac{2\pi}{N}kn} = e^{-j\frac{2\pi}{N}kn_0} a_k$$

(b) Solution:

$$\frac{1}{N} \sum_{n=0}^{N-1} (x[n] - x[n-1]) e^{-j\frac{2\pi}{N}kn} = a_k - e^{j\frac{2\pi}{N}k} a_k$$

(c) Solution:

$$\frac{1}{N} \sum_{n=0}^{N-1} (-1)^n x[n] e^{-j\frac{2\pi}{N}kn} = a_{k-\frac{N}{2}}$$

(d) tba

28. (a) Solution:

$$b_k = \sum_{m=0}^{N-1} a_m \cdot a_{m-k}^*$$

(b) Solution:

If a_k are real, we can claim that b_k are real as well.

29.

30. (a) Solution:

$$X_1(e^{j\omega}) = \frac{\frac{1}{3}e^{-j\omega}}{1 - \frac{1}{3}e^{-j\omega}}$$




FIGURE 4.12: Magnitude and phase response for sequence $x_1[n] = (1/3)^n u[n - 1]$.

(b) Solution:

$$X_2(e^{j\omega}) = \left(\frac{1}{2}\right)^5 \left[\frac{e^{-2j(\omega-\pi/4)}}{1 - \frac{1}{4}e^{-j(\omega-\pi/4)}} + \frac{e^{-2j(\omega+\pi/4)}}{1 - \frac{1}{4}e^{-j(\omega+\pi/4)}} \right]$$




FIGURE 4.13: Magnitude and phase response for sequence $x_2[n] = (1/4)^n \cos(\pi n/4) u[n - 2]$.

(c) Solution:

$$X_3(e^{j\omega}) = \begin{cases} \frac{16}{\pi^2} e^{-j4\omega}, & 0 \leq |\omega| \leq \frac{\pi^2}{4} \\ 0, & \frac{\pi^2}{4} < |\omega| < \pi \end{cases}$$

(d) Solution:

$$X_4(e^{j\omega}) = \frac{1}{2j} \left[\frac{1 - e^{-10j(\omega-0.1\pi)}}{1 - e^{-j(\omega-0.1\pi)}} - \frac{1 - e^{-10j(\omega+0.1\pi)}}{1 - e^{-j(\omega+0.1\pi)}} \right]$$




FIGURE 4.14: Magnitude and phase response for sequence $x_3[n] = \text{sinc}(2\pi n/8) * \text{sinc}\{2\pi(n - 4)/8\}$.





FIGURE 4.15: Magnitude and phase response for sequence $x_4[n] = \sin(0.1\pi n)(u[n] - u[n - 10])$.

(e) Solution:

$$X_5(e^{j\omega}) = \begin{cases} \frac{8}{\pi^3}(\frac{\pi^2}{2} - |\omega|), & 0 \leq |\omega| \leq |2\pi - \frac{\pi^2}{2}| \\ \frac{8(\pi-2)}{\pi^2}, & |2\pi - \frac{\pi^2}{2}| < |\omega| < \pi \end{cases}$$

FIGURE 4.16: Magnitude and phase response for sequence $x_5[n] = \text{sinc}^2(\pi n/4)$.

31. (a) Solution:

$$x_1[n] = \frac{1}{2\pi} \left(1 - 2 \cos \frac{\pi}{2} n \right)$$

(b) Solution:

$$x_2[n] = \frac{1}{5} \text{sinc} \left(\frac{n}{5} \right)$$

(c) Solution:

$$x_3[n] = \frac{-j}{n\pi} \cos\left(\frac{\pi}{2}n\right) + \frac{2j}{n^2\pi^2} \sin\left(\frac{\pi}{2}n\right)$$

(d) Solution:

$$x_4[n] = \frac{j}{\pi n} \left\{ 1 - \cos(\pi n) + \cos\left[\left(\omega_c + \frac{\Delta\omega}{2}\right)n\right] - \cos\left[\left(\omega_c - \frac{\Delta\omega}{2}\right)n\right] \right\}$$

32. (a) Solution:

$$X_1(e^{j\omega}) = 2e^{2j\omega} X(e^{j\omega}) + 3e^{-3j\omega} X(e^{j\omega})$$

(b) Solution:

$$X_2(e^{j\omega}) = \frac{1}{2} e^{j\frac{\pi}{6}} \delta\left(\omega - \frac{\pi}{5}\right) + \frac{1}{2} e^{-j\frac{\pi}{6}} \delta\left(\omega + \frac{\pi}{5}\right) + \frac{1}{2} e^{j\frac{\pi}{6}} X(e^{j(\omega - \frac{\pi}{5})}) + \frac{1}{2} e^{-j\frac{\pi}{6}} X(e^{j(\omega + \frac{\pi}{5})})$$

(c) Solution:

$$X_3(e^{j\omega}) = -2e^{j2(\omega - 0.5\pi)} X(e^{j(\omega - 0.5\pi)})$$

(d) Solution:

$$X_4(e^{j\omega}) = \frac{1}{2} X(e^{j\omega}) - \frac{1}{2} X(e^{j\omega})^*$$

(e) Solution:

$$X_5(e^{j\omega}) = X(e^{j(\omega-\frac{\pi}{2})})e^{j(\omega-\frac{\pi}{2})} + X(e^{j(\omega+\frac{\pi}{2})})e^{-j(\omega+\frac{\pi}{2})}$$

33. (a) Solution:

$$X_1(e^{j\omega}) = \frac{e^{2j(\omega-\frac{\pi}{2})}}{1 + 0.8e^{-j(\omega-\frac{\pi}{2})}}$$

(b) Solution:

$$X_2(e^{j\omega}) = \frac{\frac{1}{2}}{1 + 0.8e^{-j(\omega-0.4\pi)}} + \frac{\frac{1}{2}}{1 + 0.8e^{-j(\omega+0.4\pi)}}$$

(c) Solution:

$$X_3(e^{j\omega}) = \frac{1}{1 + 1.6 \cos(\omega) + 0.64}$$

(d) Solution:

$$X_4(e^{j\omega}) = \frac{1}{1 + 0.8e^{-j\frac{\omega}{2}}}$$

(e) Solution:

$$X_5(e^{j\omega}) = \frac{1}{1 - 0.8^2 e^{-j2\omega}}$$

34. (a) Solution:

$$X_R(e^{j\omega}), \quad \text{odd symmetric}$$

$$X_I(e^{j\omega}), \quad \text{even symmetric}$$

$$x_R[n] = 0,$$

$$x_I[n], \quad \text{nonsymmetric}$$

(b) Solution:

$$X_R(e^{j\omega}), \quad \text{odd symmetric}$$

$$X_I(e^{j\omega}), \quad \text{even symmetric}$$

$$x_R[n] = 0,$$

$$x_I[n], \quad \text{even symmetric}$$

(c) Solution:

$$X_R(e^{j\omega}), \quad \text{odd symmetric}$$

$$X_I(e^{j\omega}), \quad \text{even symmetric}$$

$$x_R[n] = 0,$$

$$x_I[n], \quad \text{odd symmetric}$$

35.

36. (a) See plot below.

(b)




FIGURE 4.17: Cross correlation $r_{xy}[\ell]$ plot of (a) $D = 10$. (b) $D = 20$. (c) $D = 50$.