

Predictors of END in Multivariable analysis

	Clinical (Multivariable Analysis)	Radiological (Multivariable Analysis)
Boulenoir et al. 2021	• None	• Suprabulbar iCAo pre-IVT: OR 4.0 (95% CI 1.3–12.2, P=0.015), the only one retained in the model after stepwise variable selection • Post-IVT: none
Boulenoir et al. 2022	END24h • IVT: aHR 2.01 (95% CI 1.07–3.92, P=0.03), adjusted for baseline NIHSS and onset-to-imaging time. END24h occurred earlier in IVT group (median imaging-to-END24h: 2.4h IQR 1.8–7.3 vs 8.3h IQR 4.0–15.3, P=0.04) • Non-atheromatous etiology: END24h less frequent in atheromatous cause (14%) vs cardioembolic (50%), dissection (31%), other/undetermined (35%), P<0.01 NB → the study assessed time-to-END (when will END occur), not END occurrence (if END will occur)	• None
Broccolini et al. 2024	• AFib: aOR 3.547 (95% CI 1.014–12.406, P=0.048), adjusted for age, antiplatelet, anticoagulant, dominant M2, Menon score, thrombolysis	• None
Gwak et al. 2021	IVT • aOR 7.68 (95% CI 2.52–23.43, P≤0.001) in Model 1, adjusted for Tmax >6 seconds ≥53.73 mL, onset-to-door time, IV tPA, and initial DWI volume • aOR 5.97 (95% CI 2.05–17.40, P≤0.001) in Model 2, adjusted for Tmax >6 seconds – DWI ≥32.77 mL, onset-to-door time, IV tPA, and initial DWI volume • aOR 14.94 (95% CI 4.18–53.33, P≤0.001) in Model 3, adjusted for Tmax 4–6 seconds ≥55.20 mL, onset-to-door time, IV tPA, and initial DWI volume	• Tmax >6s ≥53.73 mL (critical hypoperfusion): aOR 12.78 (95% CI 3.36–48.65, P<0.001); Model 1, adjusted for Tmax >6 seconds ≥53.73 mL, onset-to-door time, IV tPA, and initial DWI volume • Mismatch volume ≥32.77 mL (Tmax >6s - DWI): aOR 5.73 (95% CI 2.04–16.08, P=0.001); Model 2, adjusted for Tmax >6 seconds – DWI ≥32.77 mL, onset-to-door time, IV tPA, and initial DWI volume • Tmax 4–6s ≥55.20 mL (mild hypoperfusion): aOR 9.13 (95% CI 2.76–30.17, P<0.001); Model 3, adjusted for Tmax 4–6 seconds ≥55.20 mL, onset-to-door time, IV tPA, and initial DWI volume
Gwak et al. 2022	• IVT independently associated with END in both Model 1 and Model 2 (exact aOR values not reported in this paper; see Gwak et al. 2021 for statistics)	Poor perfusion profile • Defined as Wilcoxon-Mann-Whitney generalised OR >1.052 (Model 1): aOR 13.42 (95% CI: 4.38–41.15, P<0.001) • Defined as Wilcoxon-Mann-Whitney generalised OR >1.000 (Model 2): aOR 7.31 (95% CI 2.48-21.49, P<0.001) Perfusion profile = severity-weighted shift of Tmax strata distribution (Tmax 2–4s, 4–6s, 6–8s, 8–10s, >10s proportions vs Tmax>2s total), calculated via Wilcoxon-Mann-Whitney generalised OR comparing individual to cohort. Cutoffs based on Youden's index.
He et al. 2021	Higher baseline glucose: • aOR 1.02 (95% CI 1.01–1.03, P=0.009) in Model 1 (adjusted for SVS as continuous value, glucose, hypertension, smoking, cardiac embolism, undetermined stroke) • aOR 1.01 (95% CI 1.00–1.03, P=0.021) in Model 2 (adjusted for SVS ≥9.45 mm dichotomized, glucose, hypertension, smoking, cardiac embolism, undetermined stroke)	• Longer SVS: aOR 2.03 (95% CI 1.45–2.84, p<0.001); Model 1, adjusted for SVS value (continuous), baseline glucose, hypertension, current smoking, cardiac embolism, undetermined stroke) • SVS ≥9.45 mm: aOR 5.41 (95% CI 1.00–29.27, p=0.001); Model 2, adjusted for SVS ≥ 9.45 mm (dichotomized), baseline glucose, hypertension, current smoking, cardiac embolism, undetermined stroke)
Heitkamp et al. 2025	• Higher pre-stroke mRS: aOR 1.42 (95% CI 1.13–1.78, P=0.003), Adjusted for NIHSS at admission, pre stroke mRS, AFib, ICA occlusion, time from admission to groin puncture, tPA, general anesthesia, number of passes, successful recanalization TICl 2b-3, adverse events during EVT, ICH after 24h	• Longer admission-to-groin-puncture time: aOR 1.04 (95% CI 1.02–1.07, P=0.001) per minute • General anesthesia: aOR 1.68 (95% CI 1.08–2.63, P=0.022) • Higher number of passes: aOR 1.15 (95% CI 1.03–1.29, P=0.012) • Adverse events during EVT: aOR 1.89 (95% CI 1.19–3.01, P=0.007) • ICH within 24h: aOR 3.40 (95% CI 1.90–6.07, P<0.001) Adjusted for NIHSS, pre-stroke mRS, AFib, ICA occlusion, admission-to-groin time, tPA, general anesthesia, no. of passes, TICl 2b-3, adverse events during EVT, ICH within 24h • Of note, successful recanalization TICl 2b-3 was protective: aOR 0.29 (95% CI 0.17–0.50, P<0.001)
Hu et al. 2025	• Delayed antiplatelet therapy (>24h after IVT): aOR 1.46 (95% CI 1.17–1.84, P<0.001), adjusted for Tmax >4s volume, delayed antiplatelet therapy and occlusion site	• Isolated ICA occlusion: aOR 2.49 (95% CI 1.97–4.41, P<0.001), adjusted for Tmax >4s volume, delayed antiplatelet therapy and occlusion site
Itabashi et al. 2023	• No multivariable analysis done	• No multivariable analysis done
Kim et al. 2022	• History of Hypertension: aOR 1.49 (95% CI 1.03–2.14) • Higher baseline NIHSS: aOR 1.12 per 1-point increase (95% CI 1.02–1.23) Adjusted for age, male sex, occluded artery, NIHSS score, LKW to arrival, Hypertension, Diabetes, Advanced WMH, Baseline ASPECTS, collateral grade at baseline	• Extracranial ICA occlusion (without tandem): aOR 1.72 (95% CI 1.09–2.73). • Tandem occlusion: aOR 2.79 (95% CI 1.57–4.95) Adjusted for age, male sex, occluded artery, NIHSS score, LKW to arrival, Hypertension, Diabetes, Advanced WMH, Baseline ASPECTS, collateral grade at baseline
Lee et al. 2020	• No multivariable analysis done	• No multivariable analysis done
Mazya et al. 2018	• No multivariable analysis done	• No multivariable analysis done
Min et al. 2020	• None	• Insular lesions, compared with no insular lesions, were independently associated with END with infarct growth (OR 2.75, 95% CI 1.18–6.38, p = 0.02), and END-2 with infarct growth (OR 2.52, 95% CI 1.08–5.87, p = 0.03); included confounders: age, male sex, baseline NIHSS Notes • insular lesions associated with END with infarct growth, and not the other way around! • END was classified by presumed mechanism: END with infarct growth, END with new lesions, and END with swelling.
Saleem et al. 2020	• None	• None
Seners et al. 2021	• None	• More proximal occlusion (vs M2 reference): • distal M1 aOR 2.6 (1.4–5.0), proximal M1 aOR 4.2 (1.8–9.4), tandem aOR 5.1 (2.6–10.3), ICA-T/L aOR 21.0 (8.2–54.2) in Model 1, adjusted for sex, NIHSS, SBP, on-site EVT facility, occlusion site • distal M1 aOR 2.5 (1.2–5.1), proximal M1 aOR 5.2 (2.1–13.1), tandem aOR 4.5 (2.1–9.7), ICA-T/L aOR 16.0 (5.7–44.9) in Model 2, adjusted for sex, NIHSS, SBP, on-site EVT facility, occlusion site, thrombus length • Thrombus ≥9 mm: aOR 3.2 (95% CI 1.8–5.7, P=0.002) in Model 2, adjusted for sex, NIHSS, SBP, on-site EVT facility, occlusion site, thrombus length • Of note, perfusion parameters (Tmax volume, HIR) not associated with END.
Shi et al. 2025	• Highest quartile of admission SBP: aOR 2.47 (95% CI 1.47–4.29, P<0.001) • Highest quartile of 24h SBP variability: aOR 2.57 (95% CI 1.34–5.18, P<0.001) • Admission SBP >150 mmHg (ROC cutoff): AUC 0.822 (95% CI 0.749–0.895), sens 73.8%, spec 79.0%, PPV 32.6%, NPV 95.6% • SBP variability >24 mmHg (ROC cutoff): AUC 0.718 (95% CI 0.624–0.811), sens 73.8%, spec 62.9%, PPV 21.5%, NPV 94.6% • Combined SBP + SBP variability: AUC 0.848 (95% CI 0.775–0.922), sens 85.7%, spec 87.2%, PPV 47.2%, NPV 98.8%	• Tmax>6s volume highest quartile: aOR 2.09 (95% CI 1.28–5.89, P=0.001); Tmax>6s volumes categorized into quartiles with first quartile as reference.
Wang et al. 2022	NFL • aOR 1.178 (1.038–1.323, P=0.006) in Model 1, adjusted for age and sex • aOR 1.168 (1.034–1.320, P=0.012) in Model 2, adjusted for age, sex, NIHSS, TOAST classification, event to blood sampling (h), and infarct volume (Model 2) • AUC 0.850 (95% CI 0.731–0.970, P<0.001), sensitivity 73.7%, specificity 80% for NFL>55.03 pg/mL (cutoff)	• None