

Lowcountry Shrimp Dataset Deep Dive

August 4, 2025

Table of contents

Summary	3
About the Project	3
Collaborators	3
 I Definitions and Structure	 4
1 Shrimp Year Abundance Index Explanation	5
1.1 Shrimp Year	5
1.1.1 Table based on sample date and life stage	5
1.1.2 Environmental data	7
1.2 Graphic of abundances, temperatures, and assigned shrimp years	7
1.3 Abundance Index	8
1.4 About	9
 2 Data Processing and Directory Structure	 10
2.1 Directory organization	10
2.2 Data, data, data	10
2.3 About	11
 II Shrimp Data Exploration	 12
3 Postlarvae	14
3.1 Dataset	14
3.2 Graphics - Size Distributions	14
3.3 Graphics - Abundance and Size Boxplots	19
3.4 Tabular summary of abundance	19
3.5 Tabular summaries of size	21
3.5.1 Brown Shrimp	21
3.5.2 White Shrimp	22
3.6 About	23
 4 Juveniles	 25
4.1 Datasets	25
4.2 Graphics - Size Distributions	25

4.3	Graphics - Abundance/CPUE Boxplots	28
4.4	Tabular summaries of abundance	28
4.5	Tabular summaries of size	31
4.5.1	Brown Shrimp	31
4.5.2	White Shrimp	33
4.6	About	36
5	Subadults	37
5.1	Datasets	37
5.2	Graphics - Size Distributions	37
5.3	Graphics - Abundance/CPUE Boxplots	40
5.4	Tabular summaries of abundance	40
5.5	Tabular summaries of size	43
5.5.1	Brown Shrimp	43
5.5.2	White Shrimp	45
5.6	About	47
6	Adults - fisheries-independent	48
6.1	Datasets	49
6.2	Graphics - Abundance/Landings/CPUE Boxplots	49
6.3	Tabular summaries of abundance	51
6.4	About	53
7	Adults - landings and trip tickets	55
7.1	Datasets	55
7.2	Graphics - Abundance/Landings/CPUE Boxplots	56
7.3	Tabular summaries of abundance	60
7.4	About	65
III	Environmental Data Exploration	68
8	Water Temperature at Charleston Harbor	69
9	Salinity	70
IV	Summarizing to Abundance Index	71
10	Postlarval Stage	72
10.1	Tabular summaries of output datasets	72
10.1.1	Monthly	72
10.1.2	Yearly	73
10.2	About	74

11 Juvenile	76
11.1 Tabular summaries of output datasets	77
11.1.1 Monthly	77
11.1.2 Yearly	79
11.2 About	80
12 Subadults	82
12.1 Tabular summaries of output datasets	83
12.1.1 Monthly	83
12.1.2 Yearly	84
12.2 About	85
13 Adults - fishery-independent	86
13.1 test	86
13.2 another test	86
14 Adults - landings and trip tickets	87
14.1 test	87
 V Summarizing Environmental Data	 88
15 Water Temperature at Charleston Harbor	89
16 Salinity	90
 VI Relationships between datasets	 91
17 Temperature thresholds	92
18 Nursery Period Salinity	93
19 Regression Models	94

Summary

About the Project

Funded by the NERRS Science Collaborative, [link to page](#)

Collaborators

Name people here!

Part I

Definitions and Structure

1 Shrimp Year Abundance Index Explanation

The concept of **shrimp year** is somewhat complicated - a detailed explanation is below. If you prefer, jump straight to Table [1.1](#) and/or Figure [1.1](#).

1.1 Shrimp Year

When examining long-term datasets, it is convenient, if not necessary, to perform some aggregation of data to means or totals over some representative length of time (e.g. monthly mean temperature, annual precipitation total, monthly or annual totals of money spent on [anything related to budgets]).

In this project, we want to be able to link different life stages of shrimp to each other, and ask questions like “if postlarvae did well this year, did juveniles also do well? does that hold all the way through adults, and fisheries landings?”

Animal life cycles, inconveniently for us, do not necessarily align with calendar years. It may be (and for brown shrimp generally is the case) that young are spawned early in a calendar year and are adults late in that calendar year. However, white shrimp (and many other species) may be spawned in the middle or late part of a calendar year, and not be adults until the next calendar year. So, in order to relate those adults to the earlier life stages, we cannot use calendar year to represent the cohort of shrimp.

We use **shrimp year** to represent a single cohort of shrimp, and assign the calendar year in which that group of shrimp will live most of their lives. For brown shrimp, the shrimp year is the calendar year in which the individuals will show up in the fishery. For white shrimp, it is the year in which they go through most life stages; though adults will bleed into (and spawn in) the following calendar year.

Each survey in this project, based on its location of sampling, is expected to primarily catch a single life stage of each shrimp. Based on the month of sampling, the targeted life stage, and the shrimp species, we assigned shrimp year to each sampling event based on the following:

1.1.1 Table based on sample date and life stage

need to give it some formatting love, but this aligns with assignment

Table 1.1: Shrimp year assignments, based on month and calendar year of sampling, by species and life stage. ‘year’ represents the calendar year of sample date. ‘year - 1’ represents the calendar year prior to the sample date (e.g. 2004 when sample date is 2005). ‘year + 1’ represents the calendar year after the sample date (e.g. 2006 when sample date is 2005).

Month	Brown Shrimp				White Shrimp			
	Postlarvae	Juvenile	Subadult	Adult	Postlarvae	Juvenile	Subadult	Adult
1 - Jan	year	year	year	year	year	year	year - 1	year - 1
2 - Feb	year	year	year	year	year	year	year - 1	year - 1
3 - Mar	year	year	year	year	year	year	year - 1	year - 1
4 - Apr	year	year	year	year	year	year	year - 1	year - 1
5 - May	year	year	year	year	year	year	year - 1	year - 1
6 - Jun	removed	year	year	year	year	year	year - 1	year - 1
7 - Jul	removed	year	year	year	year	year	year	year
8 - Aug	removed	year	year	year	year	year	year	year
9 - Sep	removed	year + 1	year + 1	year	year	year	year	year
10 - Oct	removed	year + 1	year + 1	year	year	year	year	year
11 - Nov	removed	year + 1	year + 1	year	year	year	year	year
12 - Dec	removed	year + 1	year + 1	year	year	year	year	year

1.1.2 Environmental data

Winter temperatures and cold spells can impact shrimp growth and/or survival, and understanding these dynamics is important to shrimp fishery managers. Salinity during early life stages may also impact shrimp, and these effects extend to later stages of the life cycle.

Again, we need to aggregate data. **Winter** here is defined as December through February. For brown shrimp, the shrimp year impacted by winter temperatures is the calendar year containing that January and February. For white shrimp, we have assigned each winter metric separately to two life stages, as they occur in different years: postlarval winters and adult winters. Subadults and adults will be impacted by winter temperatures occurring a year after the winter temperatures that affect the postlarval and juvenile stages. See Figure 1.1. **Nursery period** is defined separately for each species, based on Batchelder et al. 2024*: April through July for brown shrimp, and June through October for white shrimp.

*Batchelder, L., Stone, J., Kimball, M., Pfirrmann, B., & Dunn, R. (2024). Phenology of penaeid shrimp nursery habitat use: Trends and environmental drivers over four decades. Marine Ecology Progress Series, 751, 79–95. <https://doi.org/10.3354/meps14741>

1.2 Graphic of abundances, temperatures, and assigned shrimp years

this will be what Kim showed the group at some point in the past, with 2 or 3 calendar years worth of data

need to actually process the data though and use quarto's embed functionality to show it here




Figure 1.1: Temperature and abundance data, colored by shrimp year, with irrelevant life stages grayed out.

group may want to insert conceptual diagram too when that is completed

1.3 Abundance Index

This project brought together a number of surveys by different organizations, with different methods, and different ways of quantifying shrimp abundance. In order to make these numbers comparable, while accounting for possible variation in the number of sampling stations and sample dates over time, we have adjusted abundance metrics to an index using the following steps. More details for individual datasets will be provided in that life stage's chapter in the "Summarizing to Abundance Index" portion of this book.

For each survey:

1. Assign shrimp year to each sample, based on month and species.
2. If a sampling event (site/station on a date) had replicates, average them.
3. Average data across all sites/stations on a sample day.
4. Average data across all sampling dates within each month = average abundance for the month.
5. Add the monthly averages, to get a total abundance for each shrimp year. This absorbs variability in timing for a year.
6. From total abundance per shrimp year, derive an index:
 - a. Take the square root of total abundance (deals some with extreme high numbers).
 - b. Mean-center the result from (a) across all shrimp years for the dataset.
 - c. Divide by the standard deviation across all shrimp years for the dataset.

talk about multiple datasets and how they were combined

1.4 About

Input .qmd file for this section was: **shrimp_year_explanations.qmd** and can be found in the main directory.

2 Data Processing and Directory Structure

This chapter primarily exists to guide other members of the working group in modifying, updating, or extracting information from this book.

2.1 Directory organization

- `_quarto.yml` is the main ‘table of contents’ of all `.qmd` files in order. Each chapter also has an ‘About’ section at the bottom identifying the exact file used to generate it, as well as details about the input and output datasets for the chapter.
- `_book/` is the subfolder with output files. Open any `.html` in here and you’ll have access to the full structure. The pdf output is also in this directory.

2.2 Data, data, data

- `data/raw` = time series, unmodified
- `data/intermediate` = time series with some simplifications for reporting and summarizing; generally outputs from the “Data Exploration” chapters of this book
- `data/processed` - as of this writing, I haven’t processed data yet. This folder will contain data that are aggregated to different levels:
 - month or season
 - shrimp year (based on month or season); these files will also contain a column for the abundance index

The older version of the processed data folder has the following subfolders:

- *01 Survey level* - averaged to month and then shrimp year, per survey. ALL files have the same columns: `survey`, `species`, `life_stage`, `shrimp_year`, `abundance_measure`, `abundance`, `sqrt_abundance`

- *02 Life Stage level* - averaged survey results to shrimp year x life stage, if relevant
Again all files have same columns. Survey, species, life_stage, shrimp_year, abundance, abundance_measure (if average of two surveys, will be told here), abundance, sqrt_abundance
- *03 Combined abundance* - combined all abundance files and calculated abundance index
- *04 Environmental Survey level*
- *05 Combined environmental*

2.3 About

Input .qmd file for this section was: **directory_structure.qmd** and can be found in the main directory.

Part II

Shrimp Data Exploration

Here, we go through each file to see the distribution of data and to make some graphics illustrating what is present in each dataset.

3 Postlarvae

3.1 Dataset

The only dataset representing postlarvae is from North Inlet-Winyah Bay.

3.2 Graphics - Size Distributions

In these plots, there is a panel for each shrimp species. Size is represented on the y-axis, and a cloud of points called a “beeswarm” has representation for every data point. Each point is colored by the numeric month in which the individual was caught, allowing us to see differences in size at different parts of the year and life-cycle.

There are three different measures of size in this dataset: Rostrum length, Telson length, and Uropod length. A graph is below for each.




Figure 3.1: Beeswarm plot of rostrum length in postlarval shrimp. Points represent measurements of individual shrimp, and are colored by the month in which they were captured. The y-axis represents length. More points spread out along the x-axis for a given length means that there were more individuals of that length captured than regions where points remain closer to the center.




Figure 3.2: Beeswarm plot of telson length in postlarval shrimp. Points represent measurements of individual shrimp, and are colored by the month in which they were captured. The y-axis represents length. More points spread out along the x-axis for a given length means that there were more individuals of that length captured than regions where points remain closer to the center.




Figure 3.3: Beeswarm plot of uropod length in postlarval shrimp. Points represent measurements of individual shrimp, and are colored by the month in which they were captured. The y-axis represents length. More points spread out along the x-axis for a given length means that there were more individuals of that length captured than regions where points remain closer to the center.

With brown shrimp, looks like we've got a couple of size classes in here - bigger shrimp earlier in the year, and smaller ones that are mostly later months.

The plot below shows the entire time series, with date (as year-month) along the x-axis and uropod length on the y-axis. Points are again colored by month.




Figure 3.4: Time series plot of uropod length in postlarval shrimp. Points represent measurements of individual shrimp, and are colored by the month in which they were captured.

3.3 Graphics - Abundance and Size Boxplots




Figure 3.5: Boxplots representing shrimp size and abundance Note the log10-scaled y-axes.

So even though we're seeing that pattern in brown shrimp of being so much bigger early in the year and smaller later, they're just really not all that common later in the year.

3.4 Tabular summary of abundance

Table 3.1: Abundance data frame summary

Table 3.1: Data summary

Name	postlarv_abund
Number of rows	1830
Number of columns	21
Column type frequency:	
character	6
Date	1

numeric	14
Group variables	None

Variable type: character

Table 3.2: Abundance data frame summary

skim_variable	n_missing	complete_rate	min	max	empty	n_unique	whitespace
replicate	0	1	1	10	0	13	0
date	0	1	0	10	6	915	0
season	0	1	4	6	0	4	0
ol_sal	0	1	0	7	2	181	0
ol_temp	0	1	0	7	2	241	0
notes	0	1	0	29	1746	22	0

Variable type: Date

Table 3.3: Abundance data frame summary

skim_variable	n_missing	complete_rate	min	max	median	n_unique
date2	6	1	1981-01-20	2017-12-30	1999-06-17	914

Variable type: numeric

Table 3.4: Abundance data frame summary

skim_variable	n_missing	complete_rate	mean	sd	p0	p25	p50	p75	p100
sample	0	1	458.00	264.21	1.0	229.25	458.00	686.75	915.00
year	0	1	1998.67	12.07	1900.0	1990.00	1999.00	2008.00	2017.00
month	0	1	6.48	3.45	1.0	3.00	6.00	9.00	12.00
day	0	1	15.58	8.83	0.0	8.00	16.00	23.00	31.00
dayofyear	0	1	181.63	105.57	0.0	90.25	181.00	271.75	365.00
week	0	1	26.96	15.08	1.0	14.00	27.00	40.00	53.00
bb_surface_salinity	0	1	31.59	5.08	5.9	30.83	33.20	34.50	38.60
bb_surface_temp	0	1	19.28	7.15	3.3	13.20	19.50	26.37	33.00
total_153_zoop_ind_0.3	0	1	9946.45	10692.28	0.0	3440.03	6531.50	12306.58	119401.90
vol.filt.0.13	7	1	32.36	4.76	0.0	29.36	32.47	35.74	45.68
total_ppl_density	9	1	0.20	0.63	0.0	0.00	0.00	0.13	8.40

Table 3.4: Abundance data frame summary

skim_variable	n_missing	complete_rate	mean	sd	p0	p25	p50	p75	p100
brown_density	9	1	0.05	0.19	0.0	0.00	0.00	0.03	3.77
white_density	9	1	0.05	0.29	0.0	0.00	0.00	0.00	8.19
pink_density	9	1	0.09	0.42	0.0	0.00	0.00	0.00	6.71

3.5 Tabular summaries of size

3.5.1 Brown Shrimp

Table 3.5: Brown shrimp size summary

Table 3.5: Data summary

Name	postlarv_size_brown
Number of rows	1952
Number of columns	16
Column type frequency:	
character	5
numeric	10
POSIXct	1
Group variables	None

Variable type: character

Table 3.6: Brown shrimp size summary

skim_variable	n_missing	complete_rate	min	max	empty	n_unique	whitespace
Shrimp_ID	0	1.00	6	8	0	1949	0
TowID	0	1.00	3	4	0	414	0
Replicate	0	1.00	1	1	0	2	0
Species	0	1.00	5	5	0	1	0
Notes	1830	0.06	1	34	0	33	0

Variable type: numeric

Table 3.7: Brown shrimp size summary

skim_variable	n_missing	complete_rate	mean	sd	p0	p25	p50	p75	p100
Cruise	0	1.00	510.58	192.75	32.00	388.00	521.00	606.00	909.00
Shrimp_ID_number	0	1.00	10.36	13.36	1.00	2.00	5.00	14.00	73.00
Surface_Salinity	2	1.00	26.58	7.84	5.90	23.10	29.60	32.38	38.60
Surface_Temperature	2	1.00	17.30	5.44	7.00	13.60	16.50	19.50	30.10
Rostrum_teeth	40	0.98	2.68	0.84	0.00	2.00	3.00	3.00	6.00
Rostrum_length	61	0.97	3141.90	415.89	1391.45	3018.81	3278.33	3415.37	4266.98
Telson_length	70	0.96	1516.31	179.36	749.62	1441.84	1567.86	1635.95	1872.39
Uropod_length	38	0.98	1768.37	214.82	799.73	1675.06	1822.97	1914.16	2324.15
Year	0	1.00	2001.33	7.81	1982.00	1996.00	2001.50	2005.00	2017.00
Month	0	1.00	4.26	2.31	1.00	3.00	4.00	4.00	12.00

Variable type: POSIXct

Table 3.8: Brown shrimp size summary

skim_variable	n_missing	complete_rate	min	max	median	n_unique
Date	0	1	1982-04-20	2017-10-03	2002-01-03	302

3.5.2 White Shrimp

Table 3.9: White shrimp size summary

Table 3.9: Data summary

Name	postlarv_size_white
Number of rows	1096
Number of columns	16
Column type frequency:	
character	5
numeric	10
POSIXct	1
Group variables	None

Variable type: character

Table 3.10: White shrimp size summary

skim_variable	n_missing	complete_rate	min	max	empty	n_unique	whitespace
Shrimp_ID	0	1.00	7	8	0	1087	0
TowID	0	1.00	3	4	0	249	0
Replicate	0	1.00	1	1	0	2	0
Species	0	1.00	5	5	0	1	0
Notes	1029	0.06	5	35	0	29	0

Variable type: numeric

Table 3.11: White shrimp size summary

skim_variable	n_missing	complete_rate	mean	sd	p0	p25	p50	p75	p100
Cruise	0	1.00	487.77	228.39	15.00	356.00	482.00	632.50	907.00
Shrimp_ID_number	0	1.00	7.28	8.19	1.00	2.00	4.00	10.00	47.00
Surface_Salinity	0	1.00	31.57	4.82	6.90	31.40	33.20	34.40	38.60
Surface_Temperature	0	1.00	26.03	2.43	13.90	25.20	26.70	27.50	30.10
Rostrum_teeth	16	0.99	1.65	0.79	0.00	1.00	2.00	2.00	6.00
Rostrum_length	27	0.98	1940.35	321.57	1107.64	1710.19	1831.93	2179.20	3514.92
Telson_length	28	0.97	1033.78	155.01	593.65	928.25	990.63	1144.55	1708.58
Uropod_length	18	0.98	1142.02	197.06	654.82	995.32	1081.15	1281.64	2075.20
Year	0	1.00	2000.24	9.24	1981.00	1995.00	2000.00	2006.00	2017.00
Month	0	1.00	6.65	1.16	3.00	6.00	6.00	7.00	10.00

Variable type: POSIXct

Table 3.12: White shrimp size summary

skim_variable	n_missing	complete_rate	min	max	median	n_unique
Date	0	1	1981-08-13	2017-09-01	2000-06-29	184

3.6 About

Input .qmd file for this section was: **raw__postlarv.qmd** and can be found in the main directory.

Input data file(s) - as-used; can be found in **data/raw/postlarv**:

- `Penaeus_PostLarvae_NInlet_1981_2017_wide_kac.csv`
- `Penaeus_Postlarval_Lengths_NInlet_1981 - 2017.xlsx`

Modifications to data:

- minor re-naming: inserted underscores rather than spaces; changed `Shrimp Species` to `Species`.

Generated data file(s): Data frames were modified as described above, and written to `data/intermediate/postlarval_dfs.RData` (the single `.RData` file contains multiple data frames):

- `postlarv_abund`
- `postlarv_size`

Other notes about data:

none

4 Juveniles

4.1 Datasets

There are two datasets that represent juveniles: SCDNR's Creek Trawls, and NIW NERR's Oyster Landing seines. We will explore each.

4.2 Graphics - Size Distributions

Beeswarm plots can be slow to render so I have subsetting both data frames here to 15,000 rows.




Figure 4.1: Beeswarm plot of juvenile shrimp length for the SCDNR data. Points represent measurements of individual shrimp, and are colored by the month in which they were captured. The y-axis represents length. More points spread out along the x-axis for a given length means that there were more individuals of that length captured than regions where points remain closer to the center.




Figure 4.2: Beeswarm plot of juvenile shrimp length for the Oyster Landing seine data. Points represent measurements of individual shrimp, and are colored by the month in which they were captured. The y-axis represents length. More points spread out along the x-axis for a given length means that there were more individuals of that length captured than regions where points remain closer to the center.

Looks like OL seines are generally catching smaller individuals of both species than the SCDNR creek trawls.

4.3 Graphics - Abundance/CPUE Boxplots




Figure 4.3: Boxplots representing shrimp abundance (as either CPUE or count, depending on the survey) by month. Note the log10-scaled y-axes.

Both surveys (unsurprisingly) show the same temporal pattern of abundance - aztecus in May/June, with setiferus later in the year.

4.4 Tabular summaries of abundance

SCDNR Creek Trawls

Table 4.1: Abundance (SC) data frame summary

Table 4.1: Data summary

Name	juv_sc_cpue
Number of rows	5619
Number of columns	13
Column type frequency:	

character	5
numeric	8
Group variables	None

Variable type: character

Table 4.2: Abundance (SC) data frame summary

skim_variable	n_missing	complete_rate	min	max	empty	n_unique	whitespace
StationCode	0	1	4	4	0	7	0
EstuaryCode	0	1	2	2	0	1	0
DTStart	0	1	8	10	0	621	0
SpCode	0	1	4	4	0	3	0
Species	0	1	4	5	0	3	0

Variable type: numeric

Table 4.3: Abundance (SC) data frame summary

skim_variable	n_missing	complete_rate	mean	sd	p0	p25	p50	p75	p100
X	0	1	2810.00	1622.21	1.00	1405.50	2810.00	4214.50	5619.00
Coll	0	1	20030686.47	33365.28	19801003.00	111032.00	31007.20	161081.20	231084.00
Year	0	1	2002.97	13.34	1980.00	1991.00	2003.00	2016.00	2023.00
Month	0	1	6.62	1.91	1.00	5.00	7.00	8.00	12.00
Day	0	1	15.29	8.38	1.00	9.00	15.00	22.00	31.00
CPUE	3	1	546.00	2690.49	0.00	0.00	0.00	82.00	62976.00
Lat	0	1	32.84	0.07	32.75	32.80	32.86	32.86	32.95
Long	0	1	-79.89	0.08	-79.99	-79.98	-79.88	-79.85	-79.76

Oyster Landing Seines

Table 4.4: Abundance (Oyster Landing) data frame summary

Table 4.4: Data summary

Name	juv_ol_count
Number of rows	1796
Number of columns	12

Column type frequency:	
character	4
Date	1
numeric	7
Group variables	
None	

Variable type: character

Table 4.5: Abundance (Oyster Landing) data frame summary

skim_variable	n_missing	complete_rate	min	max	empty	n_unique	whitespace
Protocol	0	1	7	8	0	3	0
WQ.Source	2	1	8	19	0	4	0
WQ.Time	2	1	0	5	36	74	0
Species	0	1	5	5	0	2	0

Variable type: Date

Table 4.6: Abundance (Oyster Landing) data frame summary

skim_variable	n_missing	complete_rate	min	max	median	n_unique
Date	0	1	1984-01-04	2023-12-21	2002-02-18	898

Variable type: numeric

Table 4.7: Abundance (Oyster Landing) data frame summary

skim_variable	n_missing	complete_rate	mean	sd	p0	p25	p50	p75	p100
Sample	0	1.00	449.76	259.63	1.0	225.0	449.5	675.00	899.0
Year	0	1.00	2002.63	11.71	1984.0	1993.0	2002.0	2012.00	2023.0
Month	0	1.00	6.60	3.26	1.0	4.0	7.0	9.00	12.0
Temp	94	0.95	20.47	6.91	3.8	14.5	21.6	26.90	31.4
Sal	106	0.94	31.82	4.78	2.3	30.0	33.3	35.00	38.7
Weight	489	0.73	625.27	2355.88	0.0	0.0	0.0	77.05	39196.1
Count	240	0.87	380.32	1326.23	0.0	0.0	0.0	62.00	22560.0

4.5 Tabular summaries of size

4.5.1 Brown Shrimp

SCDNR Creek Trawls

Table 4.8: Brown shrimp (SC) size summary

Table 4.8: Data summary

Name	sz_sc_brn
Number of rows	34818
Number of columns	14
Column type frequency:	
character	4
Date	1
numeric	9
Group variables	None

Variable type: character

Table 4.9: Brown shrimp (SC) size summary

skim_variable	n_missing	complete_rate	min	max	empty	n_unique	whitespace
DTStart	0	1	8	10	0	489	0
StationCode	0	1	4	4	0	7	0
SpCode	0	1	4	4	0	1	0
Species	0	1	5	5	0	1	0

Variable type: Date

Table 4.10: Brown shrimp (SC) size summary

skim_variable	n_missing	complete_rate	min	max	median	n_unique
Date	0	1	1980-05-08	2023-12-11	1996-06-12	489

Variable type: numeric

Table 4.11: Brown shrimp (SC) size summary

skim_variable	n_missing	complete_rate	mean	sd	p0	p25	p50	p75	p100
X	0	1.00	35073.55	22862.74	1.00	15867.25	32821.50	52650.75	81763.00
Coll	0	1.00	19981663.40	9955.91	19801003.00	19891005.00	19961014.20	20061026.20	20231080.00
Length	0	1.00	69.54	22.67	12.00	52.00	71.00	86.00	151.00
Lat	0	1.00	32.83	0.07	32.75	32.80	32.81	32.86	32.95
Long	0	1.00	-79.91	0.08	-79.99	-79.98	-79.96	-79.85	-79.76
TempS	372	0.99	27.61	2.92	8.40	25.90	28.00	29.70	36.70
SalinityS	913	0.97	17.58	5.21	0.00	14.00	18.00	21.00	28.00
Year	0	1.00	1998.07	12.00	1980.00	1989.00	1996.00	2006.00	2023.00
Month	0	1.00	5.81	0.86	1.00	5.00	6.00	6.00	12.00

Oyster Landing Seines

Table 4.12: Brown shrimp (Oyster Landing) size summary

Table 4.12: Data summary

Name	sz_ol_brn
Number of rows	15212
Number of columns	12
Column type frequency:	
character	5
Date	1
numeric	6
Group variables	None

Variable type: character

Table 4.13: Brown shrimp (Oyster Landing) size summary

skim_variable	n_missing	complete_rate	min	max	empty	n_unique	whitespace
Protocol	0	1	7	8	0	3	0
WQ.Source	0	1	8	19	0	4	0
WQ.Time	0	1	0	5	409	48	0
Species	0	1	5	5	0	1	0
Rep	0	1	4	6	0	100	0

Variable type: Date

Table 4.14: Brown shrimp (Oyster Landing) size summary

skim_variable	n_missing	complete_rate	min	max	median	n_unique
Date	0	1	1984-05-14	2023-08-29	1999-06-11	410

Variable type: numeric

Table 4.15: Brown shrimp (Oyster Landing) size summary

skim_variable	n_missing	complete_rate	mean	sd	p0	p25	p50	p75	p100
Sample	0	1.00	384.06	230.34	10.0	185	383.0	543	891.0
Year	0	1.00	1999.55	10.18	1984.0	1991	1999.0	2006	2023.0
Month	0	1.00	6.27	1.50	1.0	5	6.0	7	12.0
Temp	895	0.94	25.64	2.82	6.6	24	26.0	28	31.4
Sal	903	0.94	32.54	3.53	7.7	31	33.8	35	38.7
Length	0	1.00	21.20	6.70	0.0	16	21.0	26	71.0

4.5.2 White Shrimp

SCDNR Creek Trawls

Table 4.16: White shrimp (SC) size summary

Table 4.16: Data summary

Name	sz_sc_wht
Number of rows	46792
Number of columns	14
Column type frequency:	
character	4
Date	1
numeric	9
Group variables	None

Variable type: character

Table 4.17: White shrimp (SC) size summary

skim_variable	n_missing	complete_rate	min	max	empty	n_unique	whitespace
DTStart	0	1	8	10	0	467	0
StationCode	0	1	4	4	0	7	0
SpCode	0	1	4	4	0	1	0
Species	0	1	5	5	0	1	0

Variable type: Date

Table 4.18: White shrimp (SC) size summary

skim_variable	n_missing	complete_rate	min	max	median	n_unique
Date	0	1	1980-05-08	2023-12-11	2003-07-29	467

Variable type: numeric

Table 4.19: White shrimp (SC) size summary

skim_variable	n_missing	complete_rate	mean	sd	p0	p25	p50	p75	p100
X	0	1.00	45299.88	23170.58	7.00	25467.75	48030.50	65043.25	81764.00
Coll	0	1.00	20031152.32	4812.02	19801003.00	21067.20	31027.20	51040.20	231080.00
Length	1	1.00	53.26	21.32	8.00	37.00	51.00	66.00	152.00
Lat	0	1.00	32.84	0.07	32.75	32.80	32.81	32.86	32.95
Long	0	1.00	-79.90	0.08	-79.99	-79.98	-79.96	-79.85	-79.76
TempS	215	1.00	29.22	2.98	11.00	28.30	29.60	31.00	36.70
SalinityS	822	0.98	15.99	5.81	0.00	12.00	17.00	20.00	30.00
Year	0	1.00	2003.02	12.48	1980.00	1992.00	2003.00	2015.00	2023.00
Month	0	1.00	7.43	1.11	1.00	7.00	7.00	8.00	12.00

Oyster Landing Seines

Table 4.20: White shrimp (Oyster Landing) size summary

Table 4.20: Data summary

Name	sz_ol_wht
Number of rows	24387
Number of columns	12

Column type frequency:	
character	5
Date	1
numeric	6
Group variables	
None	

Variable type: character

Table 4.21: White shrimp (Oyster Landing) size summary

skim_variable	n_missing	complete_rate	min	max	empty	n_unique	whitespace
Protocol	0	1	7	8	0	3	0
WQ.Source	0	1	8	19	0	4	0
WQ.Time	0	1	0	5	874	51	0
Species	0	1	5	5	0	1	0
Rep	0	1	4	6	0	100	0

Variable type: Date

Table 4.22: White shrimp (Oyster Landing) size summary

skim_variable	n_missing	complete_rate	min	max	median	n_unique
Date	0	1	1984-07-12	2023-12-11	2000-09-11	436

Variable type: numeric

Table 4.23: White shrimp (Oyster Landing) size summary

skim_variable	n_missing	complete_rate	mean	sd	p0	p25	p50	p75	p100
Sample	0	1.00	421.82	222.99	14.0	236.0	414	592.0	898.0
Year	0	1.00	2001.01	9.94	1984.0	1993.0	2000	2008.0	2023.0
Month	0	1.00	8.45	1.39	1.0	7.0	8	10.0	12.0
Temp	2200	0.91	25.85	3.78	10.6	23.8	27	28.8	31.4
Sal	2305	0.91	32.81	3.96	5.4	31.8	34	35.1	38.7
Length	0	1.00	23.84	7.09	3.0	19.0	23	29.0	71.0

4.6 About

Input .qmd file for this section was: **raw__juv.qmd** and can be found in the main directory.

Input data file(s) - as-used; can be found in `data/raw/juv`:

- `SCDNR_CreekTrawlShrimpCPUE.csv`
- `SCDNR_CreekTrawlShrimpSize.csv`
- `Oyster Landing Seine Shrimp Dataset - 1984-2023 (BWP, 2025-04-24)(kac 2025-06-02).csv`

Modifications to data:

- All files: changed **Species** from provided scientific names to ‘brown’, ‘white’, and ‘pink’.
- Oyster Landing file: removed ‘DataGap’ from temp, salinity, and weight columns and turned these to numeric. Split data into counts data frame and sizes data frame. Pivoted sizes to long format.

Generated data file(s): Data frames were modified as described above, and written to `data/intermediate/juvenile_dfs.RData` (the single `.RData` file contains multiple data frames):

- `juv_sc_cpue`
- `juv_sc_size`
- `juv_ol_count`
- `juv_ol_size`

Other notes about data:

- Oyster Landing file used had a manual correction from Kim; original file had a typo - Sample 748, from 2017-07-20, LEN21, value of 221. Changed to 21.

5 Subadults

5.1 Datasets

There are two datasets that represent subadults: GADNR's EMTS sampling and SCDNR's Estuarine Trawls. We will explore each.

5.2 Graphics - Size Distributions

South Carolina's file has over 300,000 points, which makes for a beeswarm plot that is very large and slow to render. Georgia's file is "only" 65k and even it is too slow to render beeswarm plots. So rather than graphing the full datasets, I have randomly sampled 10,000 rows from GA and 15,000 rows from SC (because SC has three species).




Figure 5.1: Beeswarm plot of subadult shrimp length for the GADNR data. Points represent measurements of individual shrimp, and are colored by the month in which they were captured. The y-axis represents length. More points spread out along the x-axis for a given length means that there were more individuals of that length captured than regions where points remain closer to the center.

Size distribution by species in SCDNR Estuarine Trawls




Figure 5.2: Beeswarm plot of subadult shrimp length for the SCDNR data. Points represent measurements of individual shrimp, and are colored by the month in which they were captured. The y-axis represents length. More points spread out along the x-axis for a given length means that there were more individuals of that length captured than regions where points remain closer to the center.

5.3 Graphics - Abundance/CPUE Boxplots




Figure 5.3: Boxplots representing shrimp abundance (as either CPUE or count, depending on the survey) by month. Note the log10-scaled y-axes.

Both surveys (unsurprisingly) show the same temporal pattern of abundance - brown shrimp most abundant in June and July, also high in August; and May in SC. For white shrimp, we see a dip in June and (less so) July.

5.4 Tabular summaries of abundance

GADNR EMTS

Table 5.1: Abundance (GA) data frame summary

Table 5.1: Data summary

Name	subad_ga_count
Number of rows	39669
Number of columns	14

Column type frequency:	
character	2
Date	1
numeric	11
Group variables	None

Variable type: character

Table 5.2: Abundance (GA) data frame summary

skim_variable	n_missing	complete_rate	min	max	empty	n_unique	whitespace
RefNum	0	1	12	12	0	19828	0
Species	0	1	5	5	0	2	0

Variable type: Date

Table 5.3: Abundance (GA) data frame summary

skim_variable	n_missing	complete_rate	min	max	median	n_unique
GuessedDate	0	1	1975-12-19	2021-10-26	2000-01-10	3254

Variable type: numeric

Table 5.4: Abundance (GA) data frame summary

skim_variable	n_missing	complete_rate	mean	sd	p0	p25	p50	p75	p100
TotWt	3	1.00	1491.00	8839.01	0	0.00	27.22	680.39	1394789.25
TotNum	83	1.00	115.10	610.81	0	0.00	2.00	43.00	73800.00
SampleWt	13	1.00	382.37	716.10	0	0.00	27.22	680.39	65770.89
SampleNum	120	1.00	28.03	47.17	0	0.00	2.00	42.00	1220.00
NumMeas	21154	0.47	12.23	16.40	0	0.00	2.00	30.00	207.00
LbsperHr	12269	0.69	17.83	75.69	0	0.16	2.08	12.88	7060.50
NumperLb	12359	0.69	33.89	253.35	0	15.27	25.60	40.00	31751.50
NumFemales	2395	0.94	7.63	10.59	0	0.00	2.00	15.00	122.00
NumMales	2382	0.94	6.27	9.46	0	0.00	1.00	11.00	97.00
Year	0	1.00	1998.94	13.00	1975	1988.00	2000.00	2010.00	2021.00
Month	0	1.00	6.49	3.42	1	4.00	6.00	9.00	12.00

SCDNR Estuarine Trawls

Table 5.5: Abundance (SC) data frame summary

Table 5.5: Data summary

Name	subad_sc_cpue
Number of rows	21771
Number of columns	13
Column type frequency:	
character	5
numeric	8
Group variables	None

Variable type: character

Table 5.6: Abundance (SC) data frame summary

skim_variable	n_missing	complete_rate	min	max	empty	n_unique	whitespace
estuary	0	1	2	2	0	6	0
StationCode	0	1	4	4	0	24	0
DTStart	0	1	8	10	0	1647	0
SpCode	0	1	4	4	0	3	0
Species	0	1	4	5	0	3	0

Variable type: numeric

Table 5.7: Abundance (SC) data frame summary

skim_variable	n_missing	complete_rate	mean	sd	p0	p25	p50	p75	p100
X	0	1	10886.00	6284.89	1.000e+00	5443.50	10886.00	16328.50	21771.00
Coll	0	1	19986283.92	2808.61	1.979e+07	19890034.00	19970150.20	20090016.20	20230114.00
Year	0	1	1998.61	12.29	1.979e+03	1989.00	1997.00	2009.00	2023.00
Month	0	1	6.60	3.37	1.000e+00	4.00	6.00	9.00	12.00
Day	0	1	17.04	8.02	1.000e+00	11.00	18.00	23.00	31.00
CPUE	4	1	119.11	525.17	0.000e+00	0.00	0.50	24.00	20052.00
Latitude	0	1	32.57	0.23	3.215e+01	32.32	32.67	32.77	32.83
Longitude	0	1	-80.30	0.36	-	-80.65	-80.29	-79.92	-79.89
					8.085e+01				

5.5 Tabular summaries of size

5.5.1 Brown Shrimp

GADNR EMTS

Table 5.8: Brown shrimp (GA) size summary

Table 5.8: Data summary

Name	subad_ga_size_brown
Number of rows	11029
Number of columns	4
Column type frequency:	
character	3
numeric	1
Group variables	
None	

Variable type: character

Table 5.9: Brown shrimp (GA) size summary

skim_variable	n_missing	complete_rate	min	max	empty	n_unique	whitespace
RefNum	0	1	12	12	0	745	0
TowDate	0	1	8	10	0	308	0
Species	0	1	5	5	0	1	0

Variable type: numeric

Table 5.10: Brown shrimp (GA) size summary

skim_variable	n_missing	complete_rate	mean	sd	p0	p25	p50	p75	p100
Size	0	1	113.84	17.45	8	103	114	125	182

SCDNR Estuarine Trawls

Table 5.11: Brown shrimp (SC) size summary

Table 5.11: Data summary

Name	sz_sc_brn
Number of rows	66499
Number of columns	15
Column type frequency:	
character	5
numeric	10
Group variables	None

Variable type: character

Table 5.12: Brown shrimp (SC) size summary

skim_variable	n_missing	complete_rate	min	max	empty	n_unique	whitespace
SpCode	0	1	4	4	0	1	0
Sex	0	1	0	1	61324	4	0
estuary	0	1	2	2	0	6	0
StationCode	0	1	4	4	0	24	0
Species	0	1	5	5	0	1	0

Variable type: numeric

Table 5.13: Brown shrimp (SC) size summary

skim_variable	n_missing	complete_rate	mean	sd	p0	p25	p50	p75	p100
X	0	1.00	129622.3480691	211.000e+00	59387.50	126106.00	187401.50	304572.00	
Coll	0	1.00	19955732.852429	92.979e+07	19870163.00	19940120.20	1010143.20	20230106.00	
Length	0	1.00	102.12	19.99	2.100e+01	90.00	103.00	115.00	177.00
Year	0	1.00	1995.56	11.25	1.979e+03	1987.00	1994.00	2001.00	2023.00
Month	0	1.00	6.66	1.22	1.000e+00	6.00	6.00	7.00	12.00
Day	0	1.00	17.19	7.95	1.000e+00	10.00	19.00	24.00	31.00
Latitude	0	1.00	32.68	0.19	3.215e+01	32.67	32.77	32.80	32.83
Longitude	0	1.00	-80.12	0.30	-	-80.29	-79.97	-79.92	-79.89
TempB	1640	0.98	27.81	2.84	0.000e+00	27.00	28.40	29.40	32.40
SalinityB	2759	0.96	24.83	6.13	0.000e+00	21.00	26.00	30.00	39.00

5.5.2 White Shrimp

GADNR EMTS

Table 5.14: White shrimp (GA) size summary

Table 5.14: Data summary

Name	subad_ga_size_white
Number of rows	54505
Number of columns	4
Column type frequency:	
character	3
numeric	1
Group variables	None

Variable type: character

Table 5.15: White shrimp (GA) size summary

skim_variable	n_missing	complete_rate	min	max	empty	n_unique	whitespace
RefNum	0	1	12	12	0	2231	0
TowDate	0	1	8	10	0	726	0
Species	0	1	5	5	0	1	0

Variable type: numeric

Table 5.16: White shrimp (GA) size summary

skim_variable	n_missing	complete_rate	mean	sd	p0	p25	p50	p75	p100
Size	0	1	124.02	21.61	2	108	124	140	212

SCDNR Estuarine Trawls

Table 5.17: White shrimp (SC) size summary

Table 5.17: Data summary

Name	sz_sc_wht
Number of rows	224834
Number of columns	15
Column type frequency:	
character	5
numeric	10
Group variables	None

Variable type: character

Table 5.18: White shrimp (SC) size summary

skim_variable	n_missing	complete_rate	min	max	empty	n_unique	whitespace
SpCode	0	1	4	4	0	1	0
Sex	0	1	0	1	173246	4	0
estuary	0	1	2	2	0	6	0
StationCode	0	1	4	4	0	24	0
Species	0	1	5	5	0	1	0

Variable type: numeric

Table 5.19: White shrimp (SC) size summary

skim_variable	n_missing	complete_rate	mean	sd	p0	p25	p50	p75	p100
X	0	1.00	165616.5789820.947.000e+08	9692.25	172903.50244450.75307960.00				
Coll	0	1.00	19991381.293497.97.979e+07	9890293.09970176.20090085.20230114.00					
Length	0	1.00	113.38	22.51	1.100e+01	98.00	114.00	130.00	207.00
Year	0	1.00	1999.12	12.36	1.979e+03	1989.00	1997.00	2009.00	2023.00
Month	0	1.00	7.49	3.49	1.000e+00	4.00	8.00	10.00	12.00
Day	0	1.00	16.44	8.29	1.000e+00	10.00	17.00	23.00	31.00
Latitude	0	1.00	32.59	0.22	3.215e+01	32.46	32.67	32.77	32.83
Longitude	0	1.00	-80.27	0.35	-	-80.54	-80.24	-79.92	-79.89
TempB	4077	0.98	20.86	6.76	0.000e+00	15.00	20.60	27.90	32.40
SalinityB	4665	0.98	25.93	7.06	0.000e+00	22.00	27.00	31.00	209.00

5.6 About

Input .qmd file for this section was: **raw_subad.qmd** and can be found in the main directory.

Input data file(s) - as-used; can be found in `data/raw/subad`:

- `GADNR_EMST_PenaeidShrimp_Brown_Count_thru2021.xlsx`
- `GADNR_EMST_PenaeidShrimp_White_Count_thru2021.xlsx`
- `GADNR_EMST_PenaeidShrimp_SIZE_thru2021.xlsx`, sheet = “Brown shrimp”
- `GADNR_EMST_PenaeidShrimp_SIZE_thru2021.xlsx`, sheet = “White Shrimp”
- `SCDNR_EstuarineTrawlShrimpCPUE.csv`
- `SCDNR_EstuarineTrawlShrimpSize.csv`

Modifications to data:

- All files: changed **Species** from provided scientific names to ‘brown’, ‘white’, and ‘pink’.
- GA counts file: extracted date from the Reference Number, as date was not its own field.
- SC size file: filtered data to remove lengths >1000. There were 3 such rows.

Generated data file(s): Data frames were modified as described above, and written to `data/intermediate/subadult_dfs.RData` (the single `.RData` file contains multiple data frames):

- `subad_ga_count`
- `subad_ga_size`
- `subad_sc_cpue`
- `subad_sc_size`

Other notes about data:

- In GA count files, early exploration not shown here indicated that **NumperLb** was miscalculated when the sample weight wasn’t the typical 1360.78 - calculations in the spreadsheet used that value anyway. This issue seemed to happen when catch was low.

6 Adults - fisheries-independent

SEAMAP info here

About the Coastal Trawl Survey: <https://seamap.org/seamap-sa-coastal-trawl/>

Some excerpts:

- The standardized gear used since 1987 is a pair of 75 ft (22.9 m) mongoose-type Falcon trawl nets (manufactured by Beaufort Marine Supply, Beaufort, SC) without Turtle Excluder Devices. The body of the trawl net was constructed of #15 net twine with 47 mm stretch mesh.
- Historically, the catch from each net was processed separately and assigned a unique collection number, with data from both nets at each station pooled for analysis to form a standard unit of effort (tow). Beginning in 2021, only the catch from one net was processed with the tail bag of the other net fished untied.
- three sampling seasons: Spring (April - May), Summer (July - August), and Fall (September - November).

So it will probably be best to average to the event, because we've got events in 2022 with only one collection (prior to 2022, it was 2 collections per event).

Starting in 1989, which is when this file starts, sampling was pretty consistent - seasonally rather than monthly; daytime; 75ft nets.

There's also some inner vs. outer depth zone stuff to check. See below - 14,576 rows are from "inner" and only 534 from "outer", so will remove "outer" completely.

More on depth zones, from the Figure 1 caption on the Coastal Trawl website linked above: "The first map shows what's going on, and has this in the caption:"Inner (shallow) strata are sampled during all seasons (1990-present). Outer (deep) strata were sampled south in Spring (green) and north in Fall (yellow) from 1990-2000. In 1989, both depth zones were sampled over the entire latitudinal range in Spring and Fall."

NC, SC, GA, and FL are present in this dataset.

6.1 Datasets

There are two SEAMAP files: one of events, and one of catch data.

Looks like when no shrimp were caught in a net, the species is left blank and count and weight are both 0. So I'll widen the data frame, filling in 0s for NAs. Then lengthen back out so 0s are all accounted for.


Also filtering to "INNER" depth zone.

Weight and count were both recorded, even in subsamples, BUT "when large numbers of a species occurred in the processed catch", all individuals were weighed; a subsample was processed for abundance (typically 32-64 individuals); and count was estimated from the weight.


FOR THAT REASON I will use weights in my analyses, because it is the most likely piece to have been measured rather than estimated.

6.2 Graphics - Abundance/Landings/CPUE Boxplots


STILL IN PROGRESS




SEAMAP Shrimp Abundance per Collection



SEAMAP Shrimp Biomass per Collection





6.3 Tabular summaries of abundance

Brown Shrimp

Table 6.1: Brown shrimp SEAMAP summary

Table 6.1: Data summary

Name	filter(adsm_long, species...
Number of rows	11310
Number of columns	9
Column type frequency:	
character	2
Date	1
numeric	6
Group variables	None

Variable type: character

Table 6.2: Brown shrimp SEAMAP summary

skim_variable	n_missing	complete_rate	min	max	empty	n_unique	whitespace
Region	0	1	7	14	0	4	0
species	0	1	5	5	0	1	0

Variable type: Date

Table 6.3: Brown shrimp SEAMAP summary

skim_variable	n_missing	complete_rate	min	max	median	n_unique
Date	0	1	1989-04-12	2022-11-16	2005-07-19	994

Variable type: numeric

Table 6.4: Brown shrimp SEAMAP summary

skim_variable	n_missing	complete_rate	mean	sd	p0	p25	p50	p75	p100
Year	0	1	2004.80	9.08	1989	1997	2005	2012.00	2022.0
Month	0	1	7.30	2.41	4	5	7	10.00	11.0
Event	0	1	2005105.07	9090.43	1989001	1997302	2005309	2012542.50	2022559.0
Collection	0	1	20048332.60	798.12	1989000	1997030	2005031	20120542.70	20220559.0
Weight	0	1	0.18	0.92	0	0	0	0.03	30.4
Count	0	1	11.64	69.76	0	0	0	2.00	2983.0

White Shrimp

Table 6.5: White shrimp SEAMAP summary

Table 6.5: Data summary

Name	filter(adsm_long, species...
Number of rows	11310
Number of columns	9
Column type frequency:	
character	2
Date	1
numeric	6

Group variables	None
-----------------	------

Variable type: character

Table 6.6: White shrimp SEAMAP summary

skim_variable	n_missing	complete_rate	min	max	empty	n_unique	whitespace
Region	0	1	7	14	0	4	0
species	0	1	5	5	0	1	0

Variable type: Date

Table 6.7: White shrimp SEAMAP summary

skim_variable	n_missing	complete_rate	min	max	median	n_unique
Date	0	1	1989-04-12	2022-11-16	2005-07-19	994

Variable type: numeric

Table 6.8: White shrimp SEAMAP summary

skim_variable	n_missing	complete_rate	mean	sd	p0	p25	p50	p75	p100
Year	0	1	2004.80	9.08	1989	1997	2005	2012.00	2022.00
Month	0	1	7.30	2.41	4	5	7	10.00	11.00
Event	0	1	2005105.07	9090.43	1989001	1997302	2005309	2012542.50	2022559.00
Collection	0	1	20048332.60	798.12	1989000	1997030	2005031	20120542.75	20220559.00
Weight	1	1	1.00	2.86	0	0	0	0.46	44.07
Count	0	1	36.95	116.85	0	0	0	15.00	3641.00

6.4 About

Input .qmd file for this section was: **raw_ad_seamap.qmd** and can be found in the main directory.

Input data file(s) - as-used; can be found in `data/raw/ad_seamap`:

- `SEAMAP_ShrimpCatches_19892022.csv`

Modifications to data:

- Changed **Species** from provided names to ‘brown’, ‘white’, and ‘pink’.
- For **adsm_long** and **adsm_wide** data frames, filtered to only the “INNER” depth zone and filled in NAs with 0s, because when nothing was caught in a net, both columns in the original data were left blank.

Generated data file(s): Data frames were modified as described above, and written to **data/intermediate/adultSeamap_dfs.RData** (the single **.RData** file contains multiple data frames):

- **seamap** - the unfiltered data frame
- **adsm_wide**
- **adsm_long**

Other notes about data:

- Contains 4 states: GA, NC, SC, FL

7 Adults - landings and trip tickets

Commercial data here

7.1 Datasets

There are three files that represent commercial fisheries data for adult shrimp: Landings, from NMFS, at two different frequencies (annual for both species, and for white shrimp, fall vs. winter-spring-summer); and trip tickets, by 2-month blocks for both species.

For brown shrimp, calendar year is fine. White shrimp have different life cycles, and we are interested in spawning adults vs. non-spawning adults for each ‘shrimp year’; so we obtained the head-off white shrimp data file, split into seasons.

7.2 Graphics - Abundance/Landings/CPUE Boxplots




Figure 7.1: Time series plots of annual landings by species and state.




Figure 7.2: Time series plots of estimated CPUE by species and state, from trip ticket data. This plot uses free y-axes for overall context. Large differences probably signal changes in trip ticket collection and reporting, as states are in charge of their programs and may implement changes.




Figure 7.3: Time series plots of estimated CPUE by species and state, from trip ticket data. This plot is the same as that above, but it uses log-10 axes for easier comparison, especially at the low abundances. Large differences probably signal changes in trip ticket collection and reporting, as states are in charge of their programs and may implement changes.




Figure 7.4: Time series plots of white shrimp landings, faceted by season (1 Jan-Jul represents spring spawning adults; 2 Aug-Dec represents fall non-spawning adults).




Figure 7.5: Time series plots of white shrimp landings, colored by season (1 Jan-Jul represents spring spawning adults; 2 Aug-Dec represents fall non-spawning adults) but in the same panel and connected with a line. Jan-Jul points have been placed at March 1st each year, and Aug-Dec points placed at October 1st of each year.

7.3 Tabular summaries of abundance

Annual Landings (NMFS)

Brown Shrimp

Table 7.1: Brown shrimp annual landings summary

Table 7.1: Data summary

Name	filter(landings__annual, S...
Number of rows	120
Number of columns	10
Column type frequency:	
character	5
numeric	5

Group variables	None
-----------------	------

Variable type: character

Table 7.2: Brown shrimp annual landings summary

skim_variable	n_missing	complete_rate	min	max	empty	n_unique	whitespace
State	0	1	7	14	0	3	0
NMFS.Name	0	1	23	23	0	1	0
Collection	0	1	10	10	0	1	0
Confidentiality	0	1	6	6	0	1	0
Species	0	1	5	5	0	1	0

Variable type: numeric

Table 7.3: Brown shrimp annual landings summary

skim_variable	n_missing	complete_rate	mean	sd	p0	p25	p50	p75	p100
Year	0	1	1998.78	12.31	1978.00	1988.00	1999.50	2009.25	2019.00
Pounds	0	1	2059941.38	72035.51	1907.07	65917.75	1427067.57	20053.75	378157.00
Dollars	0	1	4039632.05	92856.65	9049.00	1461823.28	76552.09	83011.25	847514.00
Millions_of_dollars	0	1	4.04	3.69	0.06	1.46	2.88	4.98	18.85
Millions_of_Pounds	0	1	2.06	1.87	0.04	0.77	1.43	2.72	10.38

White Shrimp

Table 7.4: White shrimp annual landings summary

Table 7.4: Data summary

Name	filter(landings_annual, S...
Number of rows	120
Number of columns	10
Column type frequency:	
character	5
numeric	5

Group variables

None

Variable type: character

Table 7.5: White shrimp annual landings summary

skim_variable	n_missing	complete_rate	min	max	empty	n_unique	whitespace
State	0	1	7	14	0	3	0
NMFS.Name	0	1	23	23	0	1	0
Collection	0	1	10	10	0	1	0
Confidentiality	0	1	6	6	0	1	0
Species	0	1	5	5	0	1	0

Variable type: numeric

Table 7.6: White shrimp annual landings summary

skim_variable	n_missing	complete_rate	mean	sd	p0	p25	p50	p75	p100
Year	0	1	1998.78	12.31	1978.00	1988.00	1999.50	2009.25	2019.00
Pounds	0	1	3372708.21	71286.62	372.00	743573.73	144572.50	90372.75	172533.00
Dollars	0	1	8334883.55	49245.25	119.00	402430.73	160425.00	911377.23	688259.00
Millions_of_dollars	0	1	8.33	5.55	0.03	4.40	7.46	11.91	23.69
Millions_of_Pounds	0	1	3.37	2.17	0.01	1.74	3.14	4.59	9.47

Trip Ticket Landings**Brown Shrimp**

Table 7.7: Brown shrimp trip tickets summary

Table 7.7: Data summary

Name	filter(tripTick_abund, Sp...
Number of rows	983
Number of columns	15
Column type frequency:	
character	5
numeric	10

Group variables	None
-----------------	------

Variable type: character

Table 7.8: Brown shrimp trip tickets summary

skim_variable	n_missing	complete_rate	min	max	empty	n_unique	whitespace
Area.name	0	1	7	46	0	13	0
Common.Name	0	1	14	14	0	1	0
Species	0	1	5	5	0	1	0
Wave	0	1	9	9	0	6	0
State	0	1	0	2	237	5	0

Variable type: numeric

Table 7.9: Brown shrimp trip tickets summary

skim_variable	missing	complete_rate	mean	sd	p0	p25	p50	p75	p100
object	0	1	1883.32	1036.12	2.00	1032.00	1952.00	2785.00	3573.00
Year	0	1	2004.77	10.94	1978.00	1998.00	2007.00	2014.00	2020.00
WAVE	0	1	3.91	1.24	1.00	3.00	4.00	5.00	6.00
Lat	0	1	32.18	1.93	28.53	30.50	32.24	34.27	35.43
Long	0	1	-78.66	1.75	-	-80.35	-78.25	-77.36	-75.68
				81.21					
annland	0	1	287816.95	5768368.26	24.00	10378.01	41119.21	167878.98	8084396.00
numdeal	0	1	14777.14	67470.92	3.00	161.00	900.00	4312.00	919803.00
numfisher	0	1	14777.14	67470.92	3.00	161.00	900.00	4312.00	919803.00
CPUE_approx	0	1	140.18	294.14	0.52	13.27	42.87	149.40	4847.00
Wave_start	0	1	6.81	2.49	1.00	5.00	7.00	9.00	11.00

White Shrimp

Table 7.10: White shrimp trip tickets summary

Table 7.10: Data summary

Name	filter(tripTick_abund, Sp...
Number of rows	1636

Number of columns	15
Column type frequency:	
character	5
numeric	10
Group variables	None

Variable type: character

Table 7.11: White shrimp trip tickets summary

skim_variable	n_missing	complete_rate	min	max	empty	n_unique	whitespace
Area.name	0	1	7	46	0	14	0
Common.Name	0	1	14	14	0	1	0
Species	0	1	5	5	0	1	0
Wave	0	1	9	9	0	6	0
State	0	1	0	2	256	5	0

Variable type: numeric

Table 7.12: White shrimp trip tickets summary

skim_variable	n_missing	complete_rate	mean	sd	p0	p25	p50	p75	p100
object	0	1	1889.27	1018.10	6.00	1056.25	1938.50	2755.75	3575.00
Year	0	1	2004.90	10.65	1978.00	1998.00	2006.00	2014.00	2020.00
WAVE	0	1	3.87	1.68	1.00	3.00	4.00	5.00	6.00
Lat	0	1	31.77	1.92	25.25	30.50	31.33	33.19	35.43
Long	0	1	-79.26	1.76	-	-	-80.34	-78.25	-75.68
annland	0	1	349093.95	796531.34	81.21	80.50	29626.52	108025.30	314248.95
numdeal	0	1	15000.49	47907.43	16.00	309.75	1330.00	8266.50	648442.00
numfisher	0	1	15000.49	47907.43	16.00	309.75	1330.00	8266.50	648442.00
CPUE_approx	0	1	145.03	281.83	0.92	19.60	52.89	156.18	5202.00
Wave_start	0	1	6.75	3.35	1.00	5.00	7.00	9.00	11.00

Seasonal White Shrimp Landings

Table 7.13: White Shrimp (SC + GA) seasonal landings.

Table 7.13: Data summary

Name	landings_seas
Number of rows	90
Number of columns	6
Column type frequency:	
character	3
numeric	3
Group variables	None

Variable type: character

Table 7.14: White Shrimp (SC + GA) seasonal landings.

skim_variable	n_missing	complete_rate	min	max	empty	n_unique	whitespace
season	0	1	4	11	0	2	0
Species	0	1	5	5	0	1	0
Months	0	1	9	9	0	2	0

Variable type: numeric

Table 7.15: White Shrimp (SC + GA) seasonal landings.

skim_variable	n_missing	complete_rate	mean	sd	p0	p25	p50	p75	p100
Year	0	1	2001.0	13.06	1979	1990	2001.0	2012	2023
lbs_headOff	0	1	2593141.31799455	6474022	1169931	2164551	5354274	7863983	11699312
Period	0	1	1.5	0.50	1	1	1.5	2	2

7.4 About

Input .qmd file for this section was: **raw_ad_commercial.qmd** and can be found in the main directory.

Input data file(s) - as-used; can be found in **data/raw/ad_commercial**:

- **Nonconfidential Shrimp landings by species area year two month block.xlsx** - Trip Tickets. Commercial shrimp landings in NC, SC, GA, and FL, by 2-month block, as reported by commercial fishers and seafood processors. From state trip ticket data. per the data provider, this dataset probably best if we're interested in estimating CPUE. " Trip tickets, which have area fished, came online at different times (GA 1999, SC 2003, NC 1994). That is causing the differences. Also NC didn't always differentiate among the species (I think 96 to 98). "
- **Shrimp Landings NC SC GA from NMFS.csv** - both species, calendar year level (okay for brown shrimp, not white). Annual commercial shrimp landings in NC, SC, and GA, as reported by commercial fishers and seafood processors, and downloaded from [https://www.fisheries.noaa.gov/foss/f?p=215:200:::~::~](https://www.fisheries.noaa.gov/foss/f?p=215:200:::) per the data provider, this dataset is probably the better one for total landings.
- **scga_landings_headOff_whiteShrimp_byseason.csv** - white shrimp, by Fall vs. Winter-Spring-Summer. Combined white shrimp landings for SC and GA by two 'seasons'. The Fall is defined as Aug-Dec and the Win_Spr_Sum is defined as Jan-July. As such, these two seasons can be combined to come up with 'Annual' landings.

Modifications to data:

- All files: changed **Species** from provided names to 'brown', 'white', and 'pink'.
- Trip Tickets file: Calculated **CPUE_approx** as landings / number of dealers (number of dealers and number of fishers were the same). Made two new columns for two-month **WAVE**: **Wave** to identify the months, e.g. 1 **Jan-Feb**, 2 **Mar-Apr**, etc.; and **Wave_start** to identify the first month numerically, for better plotting of time series, e.g. Wave 1 (Jan-Feb) is assigned 1; Wave 2 (Mar-Apr) is assigned 3; etc. Added a column for **State**, based on mapping the lat/longs of each named location in the file during earlier exploration.
- Annual Landings from NMFS file: both **Pounds** and **Dollars** were read in as character due to commas in the number formatting; parsed these to numeric. Then calculated versions where the numbers will be smaller: **Millions_of_Dollars** and **Millions_of_Pounds**.
- Seasonal head-off landings of white shrimp: added two columns for better ordering, because 'Win_Spr_Sum' is Jan-July, and 'Fall' is Aug-Dec of each year. Added **Months** to show this, and **Period** where Fall is 1 and Win_Spr_Sum is 2. Otherwise Fall appears first in the data frame due to alphabetical ordering; but chronologically it should be 2nd.

Generated data file(s): Data frames were modified as described above, and written to `data/intermediate/adultCommercial_dfs.RData` (the single `.RData` file contains multiple data frames):

- `landings_annual`

- landings_seas
- tripTick_abund

Other notes about data:

- landings_seas only represents two states: GA and SC. The annual landings and trip ticket data frames also include NC.

Part III

Environmental Data Exploration

8 Water Temperature at Charleston Harbor

stuff

9 Salinity

from SCDNR trawls

Part IV

Summarizing to Abundance Index

10 Postlarval Stage

Setup

Compiling to monthly

Compiling to shrimp year

Saving files out

10.1 Tabular summaries of output datasets

10.1.1 Monthly

Table 10.1: Summary of monthly postlarval abundance data

Table 10.1: Data summary

Name	postlarv_abund_monthly
Number of rows	888
Number of columns	8
Column type frequency:	
character	4
numeric	4
Group variables	None

Variable type: character

Table 10.2: Summary of monthly postlarval abundance data

skim_variable	n_missing	complete_rate	min	max	empty	n_unique	whitespace
survey	0	1	6	6	0	1	0
species	0	1	5	5	0	2	0

Table 10.2: Summary of monthly postlarval abundance data

skim_variable	n_missing	complete_rate	min	max	empty	n_unique	whitespace
life_stage	0	1	10	10	0	1	0
abundance_measure	0	1	7	7	0	1	0

Variable type: numeric

Table 10.3: Summary of monthly postlarval abundance data

skim_variable	n_missing	complete_rate	mean	sd	p0	p25	p50	p75	p100
year	0	1	1999.00	10.68	1981	1990.00	1999.0	2008.00	2017.00
month	0	1	6.50	3.45	1	3.75	6.5	9.25	12.00
shrimp_year	0	1	4332.33	3638.30	1981	1994.00	2007.0	9999.00	9999.00
abundance	0	1	0.05	0.17	0	0.00	0.0	0.03	3.27

10.1.2 Yearly

Table 10.4: Summary of postlarval abundance data by shrimp year

Table 10.4: Data summary

Name	postlarv_yearly
Number of rows	74
Number of columns	9
Column type frequency:	
character	4
numeric	5
Group variables	None

Variable type: character

Table 10.5: Summary of postlarval abundance data by shrimp year

skim_variable	n_missing	complete_rate	min	max	empty	n_unique	whitespace
survey	0	1	6	6	0	1	0
species	0	1	5	5	0	2	0

Table 10.5: Summary of postlarval abundance data by shrimp year

skim_variable	n_missing	complete_rate	min	max	empty	n_unique	whitespace
life_stage	0	1	10	10	0	1	0
abundance_measure	0	1	7	7	0	1	0

Variable type: numeric

Table 10.6: Summary of postlarval abundance data by shrimp year

skim_variable	n_missing	complete_rate	mean	sd	p0	p25	p50	p75	p100
shrimp_year	0	1.00	1999.00	10.75	1981.00	1990.00	1999.00	2008.00	2017.00
abundance	0	1.00	0.54	0.64	0.01	0.13	0.33	0.82	4.01
sqrt_abundance	0	1.00	0.64	0.37	0.11	0.36	0.57	0.91	2.00
abundanceIndex_fullTimePeriod	0	1.00	0.00	0.99	-1.42	-0.73	-0.14	0.57	3.32
abundanceIndex_commonTimePeriod	0	1.00	0.00	0.99	-1.49	-0.78	-0.17	0.48	3.25

10.2 About

Input .qmd file for this section was: **process_postlarv.qmd** and can be found in the main directory.

Input data file(s) - in `data/intermediate/postlarval_dfs.RData` (the single .RData file contains multiple data frames):

- `postlarv_abund`
- `postlarv_size` (this data frame was ignored)

Modifications to data:

- there were some ‘year’ entries of 1900. These were removed. No data or dates were associated with these rows; presumably they were extra rows in the spreadsheet to aid in data entry.
- removed data for pink shrimp.

Generated data file(s): - fully processed files are in the `data/processed` directory:

- `postlarval_byMonth.csv` is summarized to species and month within year.
- `postlarval_byShrimpYear.csv` is summarized to species and shrimp year.

Other notes about data:

- there were some ‘year’ entries of 1900. These were removed. No data or dates were associated with these rows; presumably they were extra rows in the spreadsheet to aid in data entry.

11 Juvenile

There are two datasets that represent juveniles: SCDNR's Creek Trawls, and NIW NERR's Oyster Landing seines.

Setup

SC Creek Trawls - compile to monthly

Oyster Landing - compile to monthly

Combine monthly data frames

Make sure numbers are generally on the same scale before averaging.




Figure 11.1: Jittered graphic of the square-root of abundance by species and survey. The different surveys used different measures of abundance (SC Creek Trawls used CPUE; Oyster Landing used count) but they appear to be on the same scale, so averaging them should be fine.

Compile to Shrimp Year

First, get annual totals by survey.

Now average the annual values across surveys.

Write out.

11.1 Tabular summaries of output datasets

11.1.1 Monthly

Table 11.1: Summary of monthly juvenile abundance data, SC Creek Trawls

Table 11.1: Data summary

Name	juv_cpue_monthly
Number of rows	472
Number of columns	8
Column type frequency:	
character	4
numeric	4
Group variables	None

Variable type: character

Table 11.2: Summary of monthly juvenile abundance data, SC Creek Trawls

skim_variable	n_missing	complete_rate	min	max	empty	n_unique	whitespace
survey	0	1	15	15	0	1	0
species	0	1	5	5	0	2	0
life_stage	0	1	8	8	0	1	0
abundance_measure	0	1	4	4	0	1	0

Variable type: numeric

Table 11.3: Summary of monthly juvenile abundance data, SC Creek Trawls

skim_variable	n_missing	complete_rate	mean	sd	p0	p25	p50	p75	p100
year	0	1	2003.24	14.06	1980	1991.00	2003.0	2017.00	2023.00
month	0	1	6.85	2.11	1	5.00	7.0	8.00	12.00
shrimp_year	0	1	2003.34	14.08	1980	1991.00	2003.0	2017.00	2024.00
abundance	0	1	718.67	2074.63	0	0.43	56.2	481.33	23051.64

Table 11.4: Summary of monthly juvenile abundance data, Oyster Landing

Table 11.4: Data summary

Name	juv_ol_monthly
Number of rows	914

Number of columns	8
Column type frequency:	
character	4
numeric	4
Group variables	None

Variable type: character

Table 11.5: Summary of monthly juvenile abundance data, Oyster Landing

skim_variable	n_missing	complete_rate	min	max	empty	n_unique	whitespace
survey	0	1	13	13	0	1	0
species	0	1	5	5	0	2	0
life_stage	0	1	8	8	0	1	0
abundance_measure	0	1	5	5	0	1	0

Variable type: numeric

Table 11.6: Summary of monthly juvenile abundance data, Oyster Landing

skim_variable	n_missing	complete_rate	mean	sd	p0	p25	p50	p75	p100
year	0	1.00	2002.96	11.55	1984	1993	2003.0	2012.00	2023
month	0	1.00	6.59	3.35	1	4	7.0	9.00	12
shrimp_year	0	1.00	2003.12	11.56	1984	1993	2003.0	2013.00	2024
abundance	130	0.86	363.37	1086.03	0	0	0.5	101.12	11318

11.1.2 Yearly

Table 11.7: Summary of juvenile abundance data by shrimp year, after averaging SC and OL surveys.

Table 11.7: Data summary

Name	juv_combined_byShrimpYear...
Number of rows	89
Number of columns	9

Column type frequency:	
character	4
numeric	5
Group variables	
None	

Variable type: character

Table 11.8: Summary of juvenile abundance data by shrimp year, after averaging SC and OL surveys.

skim_variable	n_missing	complete_rate	min	max	empty	n_unique	whitespace
survey	0	1	29	29	0	1	0
species	0	1	5	5	0	2	0
life_stage	0	1	8	8	0	1	0
abundance_measure	0	1	52	52	0	1	0

Variable type: numeric

Table 11.9: Summary of juvenile abundance data by shrimp year, after averaging SC and OL surveys.

skim_variable	n_missing	complete_rate	mean	sd	p0	p25	p50	p75	p100
shrimp_year	0	1.00	2001.75	12.92	1980.00	1991.00	2002.00	2013.00	2024.00
abundance	0	1.00	3565.68	4478.64	0.43	594.59	1849.46	4813.75	24709.21
sqrt_abundance	0	1.00	50.12	32.65	0.65	24.38	43.01	69.38	157.19
abundanceIndex_fullTimePeriod	0	1.00	0.00	0.99	-1.94	-0.62	-0.15	0.55	2.62
abundanceIndex_commonTimePeriod	0	1.00	0.00	0.99	-1.91	-0.64	-0.17	0.61	2.52

11.2 About

Input .qmd file for this section was: **process_juv.qmd** and can be found in the main directory.

Input data file(s) - in `data/intermediate/juvenile_dfs.RData` (the single `.RData` file contains multiple data frames):

- `juv_sc_cpue`

- juv_sc_size
- juv_ol_count
- juv_ol_size

Modifications to data:

Generated data file(s): - fully processed files are in the **data/processed** directory:

- `juvenile_bySurveyAndMonth.csv` is summarized to species and month within year, by survey.
- `juvenile_bySurveyAndShrimpYear.csv` is summarized to species and shrimp year, by survey.
- `juvenile_byShrimpYear.csv` is summarized to species and shrimp year, as the average shrimp-year abundance across the two surveys. This is the file that will be used in further analyses.

Other notes about data:

- none

12 Subadults

There are two datasets that represent subadults: Georgia trawls and SC estuarine trawls.

Setup

GA Trawls - compile to monthly

SCDNR - compile to monthly

Combine monthly data frames

Make sure numbers are generally on the same scale before averaging.




Figure 12.1: Jittered graphic of the square-root of abundance by species and survey. The different surveys used different measures of abundance and appear to be on different enough scales that values will be normalized before averaging across surveys.

Compile to Shrimp Year

Write out.

12.1 Tabular summaries of output datasets

12.1.1 Monthly

Table 12.1: Summary of monthly subadult abundance data, GA trawls

Table 12.1: Data summary

Name	subad_ga_monthly
Number of rows	1054
Number of columns	8
Column type frequency:	
character	4
numeric	4
Group variables	None

Variable type: character

Table 12.2: Summary of monthly subadult abundance data, GA trawls

skim_variable	n_missing	complete_rate	min	max	empty	n_unique	whitespace
survey	0	1	7	7	0	1	0
species	0	1	5	5	0	2	0
life_stage	0	1	8	8	0	1	0
abundance_measure	0	1	12	12	0	1	0

Variable type: numeric

Table 12.3: Summary of monthly subadult abundance data, GA trawls

skim_variable	n_missing	complete_rate	mean	sd	p0	p25	p50	p75	p100
year	0	1	1998.82	13.00	1975	1988.00	1999.00	2010.00	2021.00
month	0	1	6.51	3.45	1	4.00	6.00	9.00	12.00
shrimp_year	0	1	1998.74	13.00	1975	1988.00	1999.00	2010.00	2022.00

Table 12.4: Summary of monthly juvenile abundance data, SC Estuarine Trawls

Error: object 'sub_sc_monthly' not found

Table 12.3: Summary of monthly subadult abundance data, GA trawls

skim_variable	n_missing	complete_rate	mean	sd	p0	p25	p50	p75	p100
abundance	0	1	1575.14	3021.84	0	2.09	426.06	2368.71	59513.63

12.1.2 Yearly

Table 12.5: Summary of subadult abundance data by shrimp year, after normalizing then averaging GA and SC surveys.

Table 12.5: Data summary

Name	subad_combined_byShrimpYe...
Number of rows	98
Number of columns	8
Column type frequency:	
character	4
numeric	4
Group variables	None

Variable type: character

Table 12.6: Summary of subadult abundance data by shrimp year, after normalizing then averaging GA and SC surveys.

skim_variable	n_missing	complete_rate	min	max	empty	n_unique	whitespace
survey	0	1	17	17	0	1	0
species	0	1	5	5	0	2	0
life_stage	0	1	8	8	0	1	0
abundance_measure	0	1	50	50	0	1	0

Variable type: numeric

Table 12.7: Summary of subadult abundance data by shrimp year, after normalizing then averaging GA and SC surveys.

skim_variable	n_missing	complete_rate	mean	sd	p0	p25	p50	p75	p100
shrimp_year	0	1.00	1999.50	14.22	1975.00	1987.25	1999.50	2011.75	2024.00
abundance	0	1.00	0.03	0.84	-2.86	-0.48	-0.01	0.47	3.87
abundanceIndex_fullTimePeriod	0	1.00	0.00	0.99	-3.47	-0.57	-0.04	0.54	4.39
abundanceIndex_commonTimePeriod	0	1.00	0.00	0.99	-1.90	-0.78	-0.01	0.61	2.47

12.2 About

Input .qmd file for this section was: **process_subad.qmd** and can be found in the main directory.

Input data file(s) - in `data/intermediate/subadult_dfs.RData` (the single `.RData` file contains multiple data frames):

- `subad_ga_count`
- `subad_ga_size`
- `subad_sc_cpue`
- `subad_sc_size`

Modifications to data:

Generated data file(s): - fully processed files are in the `data/processed` directory:

Other notes about data:

- none

13 Adults - fishery-independent

SEAMAP info here

13.1 test

rendering with freeze: auto

13.2 another test

[1] 4

14 Adults - landings and trip tickets

Commercial data here

14.1 test

code:

```
[1] 2
```

The end

Part V

Summarizing Environmental Data

15 Water Temperature at Charleston Harbor

stuff

16 Salinity

from SCDNR trawls

Part VI

Relationships between datasets

17 Temperature thresholds

some stuff

18 Nursery Period Salinity

stuff

19 Regression Models

Start to explore what happens when we do more than just correlations.