

# EDGAR v5.0 Global Air Pollutant Emissions for MOZART chemical mechanism

## Technical documentation

Caterina Mogno<sup>1,\*</sup> and Margaret R. Marvin<sup>1,2</sup>

<sup>1</sup>School of GeoSciences, The University of Edinburgh, Edinburgh, UK

<sup>2</sup>National Centre for Earth Observation, University of Edinburgh, Edinburgh, UK

\*c.mogno@ed.ac.uk

## Introduction

Emissions inventories need to be prepared to be ready to be used in chemical transport models (CTMs). They usually need ad-hoc preprocessing based on the chemical mechanism used in the CTM, including speciation of non-methane volatile organic compounds (VOCs). Here we report monthly EDGAR v5.0 global air pollutant emissions for the year 2015, speciated for the MOZART chemical mechanism. The dataset is also ready to be used in the WRF-Chem anthro\_emiss preprocessing tool with the MOZART-MOSAIC options. The folder contains:

1. edgar\_v5\_MOZART\_data.tar.gz: EDGAR v5.0 monthly emissions for the year 2015 (NetCDF format), speciated for MOZART chemical mechanism. Both total and individual sector emissions are included in each file.
2. edgarv5\_MOZART\_MOSAIC.inp: Input file for anthro\_emiss preprocessing tool for MOZART-MOSAIC options in WRF-Chem.
3. code folder: scripts used to prepare 1).
4. technical\_note\_EDGARv5\_MOZART.pdf: this technical documentation.

This technical documentation supports the dataset provided, and describes how the EDGAR v5.0 emission inventory was speciated for the MOZART chemical mechanism and prepared for use by the WRF-Chem model.

The dataset and the code provided are free to use under licence [CC-BY-SA 4.0](#)

## Inventory processing

### Pollutant species and sectors

EDGAR v.5.0 monthly global gridded air pollutant emissions at 0.1°x0.1° resolution for the year 2015 (Crippa et al., 2019b,a,c, 2020) have been downloaded from the repository at <https://data.jrc.ec.europa.eu/dataset/377801af-b094-4943-8fdc-f79a7c0c2d19#contact>. EDGAR v.5.0 considers the following air pollutant species: BC, CO, NH<sub>3</sub>, NO<sub>x</sub>, OC, PM<sub>2.5</sub>, PM<sub>10</sub>, SO<sub>2</sub> and total NMVOCs for 27 anthropogenic sectors. We consider here all sectors except for supersonic aviation (for which no monthly emissions are available). Table 1 list the sectors and their identification code. We also provide total emissions for each species, calculated from summing all the individual sector emissions for the given species. The original files downloaded have been aggregated by pollutant species in the form shown in Figure 1.

co									
xarray.Dataset									
► Dimensions: (lat: 1800, lon: 3600, time: 12)									
▼ Coordinates:									
time	(time)	int32	1	2	3	4	5	6	7
lat	(lat)	float32	-89.95	-89.85	...	89.85	89.95		
lon	(lon)	float32	0.05	0.15	0.25	...	359.85	359.95	
▼ Data variables:									
AWB	(time, lat, lon)	float32	...						
CHE	(time, lat, lon)	float32	...						
ENE	(time, lat, lon)	float32	...						
FFF	(time, lat, lon)	float32	...						
FOO-PAP	(time, lat, lon)	float32	...						
IND	(time, lat, lon)	float32	...						
IRO	(time, lat, lon)	float32	...						
NFE	(time, lat, lon)	float32	...						
NMM	(time, lat, lon)	float32	...						
PRO	(time, lat, lon)	float32	...						
RCO	(time, lat, lon)	float32	...						
REF-TRF	(time, lat, lon)	float32	...						
SWD-INC	(time, lat, lon)	float32	...						
TNR-Aviation-C...	(time, lat, lon)	float32	...						
TNR-Aviation-C...	(time, lat, lon)	float32	...						
TNR-Aviation-LTO	(time, lat, lon)	float32	...						
TNR-Other	(time, lat, lon)	float32	...						
TNR-Ship	(time, lat, lon)	float32	...						
TRO-noRES	(time, lat, lon)	float32	...						
TOTAL	(time, lat, lon)	float32	...						

Figure 1: Dataset structure for each emission file. Example for the CO species. Data variables are explained in Table 1.

## NMVOCs speciation to MOZART

Total NMVOCs in EDGAR v.5.0 are speciated to the MOZART chemical mechanism using the fractional mass contribution obtained from the EDGAR v4.3.2 NMVOCs as follows.

We first obtain sector specific monthly EDGAR v4.3.2 NMVOCs for year 2010 at  $0.1^\circ \times 0.1^\circ$  resolution Huang et al. (2017a,b) from the repository [https://data.jrc.ec.europa.eu/dataset/jrc-edgar-edgar\\_v432\\_voc\\_spec\\_gridmaps](https://data.jrc.ec.europa.eu/dataset/jrc-edgar-edgar_v432_voc_spec_gridmaps), and we reorganise them by pollutant species as in we did for original EDGAR v5.0 files in Figure 1. EDGAR v4.3.2 NMVOCs are speciated using the GEIA 25 NMVOCs groups. We relate these NMVOC groups to the MOZART chemistry using the mapping provided in Emmons et al. (2020) and reproduced in Table 2. Since EDGAR v4.3.2 NMVOCs emissions are provided in mass and not in molar units, we obtain a mass mapping of Table 2 by applying the molecular weights for the GEIA NMVOC groups. Molecular weights for the GEIA VOC species are taken from <http://www.globalchange.umd.edu/data/ceds/README-CEDS-VOC-speciation-2017-05-18.txt>. Figure 1 shows the resulting monthly MOZART VOC distributions described on a uniform  $0.1^\circ \times 0.1^\circ$  grid. We then calculate the gridded map of total NMVOCs mass by adding up the contribution of the individual MOZART species. For each MOZART VOCs species, we obtain a gridded map of its mass fractional contribution to total NMVOCs by dividing it by the gridded map of total NMVOC mass. We then apply these fractional gridded maps to the total EDGAR v5.0 NMVOCs file to obtain disaggregated NMVOCs emissions for the MOZART mechanism. The resulting EDGAR v5.0 MOZART VOCs are monthly gridded maps at  $0.1^\circ \times 0.1^\circ$  in the form of Figure 1. All the original EDGAR v5.0 NMVOC sectors are included in the MOZART NMVOCs dataset, except for manure management (MNM, ipcc1996:4B), which is not included in the EDGAR v4.3.2 NMVOCs dataset. In addition, since the definition of sectors is slightly different between EDGARv4.3.2 and EDGAR v5.0 sectors, we

sector name	sector id
Power industry	ENE
Oil refineries and Transformation industry	REF_TRF
Combustion for manufacturing	IND
Aviation landing&takeoff	TNR_Aviation_LTO
Aviation cruise	TNR_Aviation_CRS
Aviation climbing&descent	TNR_Aviation_CDS
Road transportation no resuspension	TRO_noRES
Road transportation resuspension	TRO_RES
Railways, pipelines, off-road transport	TNR_Other
Shipping	TNR_Ship
Energy for buildings	RCO
Fuel exploitation	PRO
Non-metallic minerals production	NMM
Chemical processes	CHE
Iron and steel production	IRO
Non-ferrous metals production	NFE
Non energy use of fuels	NEU
Solvents and products use	PRU_SOL
Food and Paper	FOO_PAP
Manure Management	MNM
Agricultural waste burning	AWB
Agricultural Soils	AGS
Solid waste landfills	SWD_LDF
Solid waste incineration	SWD_INC
Waste water handling	WWT
Fossil Fuel Fires	FFF

Table 1: Anthropogenic emissions sectors for EDGAR v5.0 air pollutant emissions. The sector\_id column is used as sector identifier in the dataset we provide. More information on sectors definition are available at EDGAR v5.0 air pollutant emissions official website ([Crippa et al., 2019b](#)).

needed to remap some of the NMVOCs sectors:

- Process emissions during production and application:  
PPA (EDGAR v.4.3.2) = CHE+FOO\_PAP+IRO+PRU-SOL+NMM (EDGAR v5.0)
- Waste solid and wastewater:  
SWD (EDGAR v.4.3.2) = SWD\_INC+SWD\_LDF+WWT (EDGAR v5.0)
- Oil refineries and Transformation industry:  
REF+TRF (EDGAR v.4.3.2) = REF\_TRF (EDGAR v.5.0)

The final speciated NMVOCs files that we report contain the mapped sectors (PPA,SWD,REF+TRF) and not the original EDGARv5.0 ones.

## Adapting the inventory emissions files for WRF-Chem use

We made the EDGAR v5.0 inventory speciated to MOZART ready to be used in WRF-Chem with the anthro-emiss utility (MOZART-MOSAIC options). We added to the data files monthly *date* and *datesec* variables, the attribute units:  $[kg\ m^{-2}\ s^{-1}]$  to each variable, and create the anthro-emiss input file (edgarv5-MOZART-MOSAIC.inp).

## Code information

All the data processing described above has been performed using scripts in Python 3.6, and it is summarised in the flowchart in Figure 2. For each step of the process the corresponding code is found

MOZART	GEIA/CEDS
C2H6	VOC02-ethane
C3H8	VOC03-propane
BIGALK	VOC04-butanes + VOC05-pentanes + VOC06-hexanes-pl + VOC18-esters + VOC19-ethers
C2H4	VOC07-ethene
C3H6	VOC08-propene
C2H2	VOC09-ethyne
BIGENE	VOC12-other-alke
BENZENE	VOC13-benzene
TOLUENE	VOC14-toluene
XYLENES	VOC15-xylene+VOC16-trimethylb+ VOC17-other-arom
CH2O	VOC21-methanal
CH3CHO	VOC22-other-alka
CH3OH	0.15*VOC01-alcohols
C2H5OH	0.85*VOC01-alcohols
CH3COCH3	0.2*VOC23-ketones
MEK	0.8*VOC23-ketones
HCOOH	0.5*VOC24-acids
CH3COOH	0.5*VOC24-acids

Table 2: Molar Mapping of NMVOCs species from GEIA/CEDS to MOZART chemistry. Adaptation of Table S3 from ([Emmons et al., 2020](#))

in the following scripts. For more details, see the individual Python scripts in the code folder.

- 1) download\_edgarv5.sh
- 2)edgarv5\_format.ipynb
- 3)download\_nmvoc\_edgarv432.sh
- 4) nmvoc4.3.2\_format.ipynb
- 5) CEDS\_MOZART\_VOCmap.xlsx
- 6) nmvoc4.3.2\_map\_mozart\_mass.ipynb
- 7), 8) nmvoc4.3.2\_map\_mozart\_fractions.ipynb
- 9): edgarv5\_NMVOC\_map\_sectors.xlsx; map\_v5\_to\_v4.3.2\_nmvoc\_sectors.ipynb
- 10) edgarv5\_nmvoc\_speciate\_to\_mozart.ipynb
- 11) add\_total\_emissions.ipynb
- 12) edgarv5\_to\_WRFChem\_anthroemiss.ipynb




Figure 2: Flowchart of the data processing for preparing EDGAR v5.0 air pollutant emissions for MOZART chemistry. Green rectangles represent EDGAR v5.0 data processing, orange rectangle EDGAR v4.3.2 NMVOCs data processing.

## References

- Monica Crippa, Diego Guizzardi, Marilena Muntean, Edwin Schaaf, and Gabriel Oreggioni. EDGAR v5.0 Global Air Pollutant Emissions. European Commission, Joint Research Centre (JRC) [Dataset], 2019a.
- Monica Crippa, Diego Guizzardi, Marilena Muntean, Edwin Schaaf, and Gabriel Oreggioni. EDGAR v5.0 Global Air Pollutant Emissions - Official Website. [https://edgar.jrc.ec.europa.eu/index.php/dataset\\_ap50](https://edgar.jrc.ec.europa.eu/index.php/dataset_ap50), 2019b.
- Monica Crippa, G Oreggioni, Diego Guizzardi, Marilena Muntean, Edwin Schaaf, Eleonora Lo Vullo, Efisio Solazzo, Fabio Monforti-Ferrario, Jos GJ Olivier, and Elisabetta Vignati. Fossil co2 and ghg emissions of all world countries. *Luxemburg: Publication Office of the European Union*, 2019c.
- Monica Crippa, Efisio Solazzo, Ganlin Huang, Diego Guizzardi, Ernest Koffi, Marilena Muntean, Christian Schieberle, Rainer Friedrich, and Greet Janssens-Maenhout. High resolution temporal profiles in the emissions database for global atmospheric research. *Scientific data*, 7(1):1–17, 2020.

- Louisa K Emmons, Rebecca H Schwantes, John J Orlando, Geoff Tyndall, Douglas Kinnison, Jean-François Lamarque, Daniel Marsh, Michael J Mills, Simone Tilmes, Charles Bardeen, et al. The chemistry mechanism in the community earth system model version 2 (cesm2). *Journal of Advances in Modeling Earth Systems*, 12(4), 2020.
- Ganlin Huang, Rosie Brook, Monica Crippa, Greet Janssens-Maenhout, Christian Schieberle, Chris Dore, Diego Guizzardi, Marilena Muntean, Edwin Schaaf, and Rainer Friedrich. Edgar v4.3.2 global speciated nmvocs emissions. european commission, joint research centre (jrc) [dataset], 2017a.
- Ganlin Huang, Rosie Brook, Monica Crippa, Greet Janssens-Maenhout, Christian Schieberle, Chris Dore, Diego Guizzardi, Marilena Muntean, Edwin Schaaf, and Rainer Friedrich. Speciation of anthropogenic emissions of non-methane volatile organic compounds: a global gridded data set for 1970–2012. *Atmospheric Chemistry and Physics*, 17(12):7683–7701, 2017b.