# SFX2_32

## Contents

# Sample information

ID: SFX2_32

Class: SF1_15yr

Disturbance year: 2011

Source: Visual interpretation of Landsat time series + auxiliary data (Tyukavina et al., 2022)

# Annual Landsat images (RGB SWIR1-NIR-GREEN)

RGB composites from 1984 to 2017 for the sample area (represented in white at the center) and its surroundings within a radius of 1 km.

1996 1997 1998 1999
2000 2001 2002 2003
2004 2005 2006 2007
2008 2009 2010 2011
2012 2013 2014 2015
2016 2017

# Auxiliary disturbance data

Auxiliary data for the identification of disturbance class for the sample area (represented by the red square) and for a 125 m radius around the sample.

# Hyperspectral RGB composite and LiDAR canopy height

RGB composite from the hyperspectral data (top) and LiDAR canopy height (bottom) for the sample area (represented by the red square) and for a 125 m radius around the sample.

## References

Turubanova, S., Potapov, P., Tyukavina, A., Hansen, M. (2018) Ongoing primary forest loss in Brazil, Democratic Republic of the Congo, and Indonesia. *Environmental Research Letters* https://doi.org/10.1088/1748-9326/aacd1c

Silva Junior, C.H.L., Heinrich, V.H.A., Freire, A.T.G., Broggio, I.S., Rosan, T.M., Doblas, J., Anderson, L.O., Rousseau, G.X., Shimabukuro, Y.E., Silva, C.A., House, J.I., Aragão, L.E.O.C. (2020) Benchmark maps of 33 years of secondary forest age for Brazil [Data set]. In Scientific Data (v2.0.0, Vol. 7, Number 269, https://doi.org/10.1038/s41597-020-00600-4). Zenodo. https://doi.org/10.5281/zenodo.3928660

Tyukavina, A., Potapov, P., Hansen, M.C., Pickens, A., Stehman, S., Turubanova, S., Parker, D., Zalles, V., Lima, A., Kommareddy, I., Song, X-P, Wang, L. and Harris, N. (2022) Global trends of forest loss due to fire, 2001-2019. *Frontiers in Remote Sensing* https://doi.org/10.3389/frsen.2022.825190

## Contact

For more information, contact catherine.almeida@unesp.br