

Predicting Opioid Use Disorders in the U.S. Using Machine Learning Among Adolescents Receiving Mental Health Services

Cathy Rong, Jessie Song, Afnan Rajab, Iris Li, and Chase Pelechytik

Old Scona Academic

January 10, 2025

Abstract

Opioid use disorder (OUD), encompassing opioid dependence and abuse, has risen sharply among US adolescents over the past decade, with its impact varying geographically across states. This retrospective study analyzed geographic, and social determinants, individual mental health diagnoses, and demographic data aiming to explore these inequities. A predictive model for OUD was developed using 99 features from the Mental Health Client Level Data (MH-CLD) dataset and U.S. Census Bureau data.

Machine learning classification algorithms, such as logistic regression with ridge regularization, LASSO, and random forest, were used to predict OUD. Repeated sampling of balanced classes was implemented to address data imbalance and enhance model performance. The model achieved a specificity of 84% for 2019 predictions using 2018 training data and a balance accuracy of 86% for 2022 predictions using late 2022 data. This approach enabled comparisons of pre- and post-pandemic OUD trends.

Feature importance analysis revealed that geographic location was among the most influential predictors. The model supports individualized predictions that can be scaled to inform community-specific health policies and decision-making. This predictive capability provides valuable insights to mitigate the opioid crisis in the U.S. and demonstrates the potential of machine learning as a tool for health planning and policy development.

Keywords:

Adolescents, Opioid Use Disorder, Machine Learning, Social Determinants, Prospective Validation

1 Introduction

Opioid dependence and abuse are critical public health concerns with far-reaching consequences for individuals and society. These disorders not only lead to physical health issues but cause lasting damage to brain function, impairing cognition, behavior, and social interactions, and in severe cases, resulting in death. [1, 2, 3] Addressing the opioid crisis requires targeted prevention strategies, particularly for adolescents, a population often overlooked in research. This pivotal stage of development is characterized by physical, mental, emotional, and cognitive changes [4] wherein individuals are more vulnerable to substance and opioid misuse. [5]

With the bulk of existing research focusing more on adults [6], however, adolescent-specific risk factors are less understood. Mental health disorders like bipolar disorder, which heightens impulsivity and risk-taking [7], are common in adolescence and may increase the risk of opioid dependence. [8] Environmental influences, including peer relationships, geographic location [9, 10], community dynamics, socioeconomic status [11, 12], and other contextual variables [11, 13, 14, 15, 16] further shape adolescents' susceptibility to opioid abuse. Understanding these environmental factors can help pinpoint the most critical determinants, offering insights into the environmental and social drivers of adolescent opioid abuse. This investigation into the relationship between mental health and other factors is essential for developing substance abuse prevention strategies for adolescents. [7]

Machine learning is a robust tool in addressing these complexities, able to analyze large datasets and interactions among variables and

medical conditions as well as well-suited for identifying risk factors and classifying [5] diseases [17, 18]. In this study, we utilized Mental Health Client Level Data (MH-CLD) and U.S. census data [19] to examine adolescent opioid abuse, mental health conditions, and social factors. By cleaning, normalizing, and synthesizing these datasets, we trained and tested machine-learning models using multi-year data. This approach aims to uncover the most significant risk factors for adolescent opioid dependence and abuse.

2 Materials & Methods

2.1 Study Cohort

This study utilized individual-level mental health data collected from 2013 to 2022 using the MH-CLD dataset [20], which is comprehensively composed of demographic, clinical, and outcome data for individuals receiving services from state mental health agencies in the United States during a state-defined 12-month reporting period [21]. The dataset includes information on individuals who accessed mental health support services such as screening, crisis services, and telemedicine programs offered or funded by the U.S. government during the specified time frame.

For this analysis, we focused on individuals aged 12 to 17 years, a critical developmental stage for identifying mental health and substance use disorders, including opioid use disorder (OUD). The study cohort consisted of 921,725 clients from the 2018 database with 660 OUD cases and 1,031,799 clients from the 2022 database with 1,892 OUD cases. These two-time points were chosen to enable a comparative analysis of OUD prediction trends before and after the onset of the COVID-19 pandemic, allowing for an examination of potential mental health service utilization shifts and outcomes influenced by the pandemic.

2.2 Study Features and Target Outcome

The study features included data on mental health diagnoses, demographic characteristics, utilization of mental health facilities, and co-occurring substance use-related conditions. The primary target outcome variable was OUD, encompassing opioid dependence and abuse, reflecting the growing public health concern over opioid misuse and its associated societal and clinical implications.

In addition to individual-level data, the dataset also included geographic information such as state, division, and region. To enrich the analysis we incorporated supplementary data (32 features) from external sources such as the U.S. Census Bureau datasets and other publicly available regional databases. These sources provided background information for each 50 states, including population demographics, economic indicators, employment rates, poverty levels, crime statistics, and population health statistics. This contextual data enabled a more comprehensive understanding of the social, financial, and health-related factors potentially influencing OUD trends.

By integrating these diverse datasets, the study aimed to provide a robust framework for analyzing OUD prediction and its variation across different geographic and demographic contexts, with particular emphasis on changes observed before and after the COVID-19 pandemic.

2.3 Sample Derivation

The study sample was derived as follows: the training datasets were created using the 2018 (N=645,207,70%) and 2022 data (N=505,582,49%). For validation, 30% of the 2018 data and 21% of the 2022 data were randomly selected to serve as validation datasets. For testing, the 2019 dataset was used to test the model built on the 2018 data, while 30% of the 2022 dataset was used to test the model. Figure 1 presents the flow chart for the machine learning model, including the data integration and model validation.

Notably, the MH-CLD data does not link individual case IDs across years, ensuring that cases are not traceable longitudinally. To minimize bias and maintain model integrity, no OUD data were included in the features. This approach ensured a clean separation of training, validation, and testing data to support robust model development and evaluation.

2.4 Data Preparation

The data preparation process was conducted using Python 3.11 [22] with the scikit-learn library [23] for predictive modeling. The primary mental health features included diagnoses such as ADHD, anxiety, depressive disorder, oppositional defiant disorder (ODD), pervasive developmental disorder (PDD), conduct disorder, bipolar disorder, schizophrenia, and other mental health conditions. All variables, except for the count of mental health diagnoses, were numerical.




Figure 1: Study Flow Chart

Demographic features including age group, sex, state, race, and ethnicity, were categorical and were one-hot encoded into binary variables. Substance use diagnosis variables provided outcome information and included major conditions such as opioid, alcohol, cannabis, cocaine, and other substance-related disorders. Figure 2 summarizes the descriptive information of the 2018 cohort. Background features such as population size, economy, education, income information, and so on were derived from external datasets (detailed in Supplementary Table A1). To ensure consistent scaling across variables with different units, all continuous features were normalized using the StandardScaler from scikit-learn’s preprocessing package. This normalization improved the comparability of features and optimized model performance.

2.5 Model Selection and Evaluation

The COVID-19 pandemic has led to varying social and environmental conditions across different periods, particularly from 2020 to 2022. These changes can influence model performance, making it challenging to generalize findings from pre-pandemic data, such as 2018, to the pandemic era. To address this, we first trained the model on the 2018 data and tested it on the 2019 dataset. This approach helps evaluate the model’s performance in a pre-pandemic context. Additionally, we trained and tested the model exclusively using the 2022 data to account for the relatively stable social factors specific to that

period during the pandemic. Evaluating the model on both pre-pandemic (2018 and 2019) and post-pandemic (2022) data aimed to determine whether the key risk factors contributing to OUD were consistent across different personal, social, and environmental contexts.

For the 2018 data and 70% of 2022 data (30% holding for testing), the dataset was split into a 70%/30% ratio using stratified sampling to address the class imbalance, ensuring that the proportion of subjects with opioid use disorder was consistent in both the training and validation sets. A logistic regression and Lasso model from the scikit-learn library [23] were employed with balanced class weights configured to account for the unequal distribution of outcomes [24]. To ensure robust performance evaluation, we repeated the training and validation process 10 times. The final model is built based on an ensemble model by averaging out each iteration’s coefficients. Performance metrics, including sensitivity, specificity, balanced accuracy, and the area under the receiver operating characteristic curve (ROC-AUC) were used to provide a comprehensive assessment of the model’s performance. Feature importance was also calculated to identify key predictors. The model was then tested with the 2019 data and the rest of the 2022 data. This allowed us to assess the machine learning algorithms’ generalization in the pre- and post-pandemic period.

2.6 Assessing Feature Importance

The importance of features for predicting OUD was evaluated based on the model coefficients. Features with larger absolute coefficient values were deemed more significant, indicating a stronger influence on the target outcome. The sign of the coefficients further reflected the direction of the association: a positive coefficient signified a positive association with OUD, whereas a negative sign indicated an inverse relationship.


3 Results

3.1 Visualizing Trends of Youth Mental Health Prevalence


Figure 3 illustrates the trends in mental health diagnoses over the past decade, spanning from 2013 to 2022. The analysis focuses exclusively on conditions demonstrating an upward trend, including substance use problems (3a), anxiety disorders (3b), depressive disorders (3c), and trauma-related (3d) conditions. Other mental health conditions were presenting either a downward trend or insufficient trending, rendering

Feature	Label	Full Sample N(%)	With OPIOID N(%)	Without OPIOID N(%)	Chi-Square Test p-value
Total Sample Size		921725(100%)	660(0.1)	921065(99.9)	
REGION	Northeast	155501(16.9)	75(11.4)	155426(16.9)	<0.0001
	Midwest	231465(25.1)	224(33.9)	231241(25.1)	
	South	260740(28.3)	138(20.9)	260602(28.3)	
	West	273598(29.7)	223(33.8)	273375(29.7)	
AGE	12-14 years	445648(48.3)	81(12.3)	445567(48.4)	<0.0001
	15-17 years	476077(51.7)	579(87.7)	475498(51.6)	
GENDER	Male	463974(50.3)	333(50.5)	463641(50.3)	0.9502
	Female	455036(49.4)	325(49.2)	454711(49.4)	
ANXIETYFLG	Disorder not reported	750064(81.4)	514(77.9)	749550(81.4)	0.021
	Disorder reported	171661(18.6)	146(22.1)	171515(18.6)	
DEPRESSFLG	Disorder not reported	690215(74.9)	415(62.9)	689800(74.9)	<0.0001
	Disorder reported	231510(25.1)	245(37.1)	231265(25.1)	
BIPOLARFLG	Disorder not reported	890965(96.7)	607(92.0)	890358(96.7)	<0.0001
	Disorder reported	30760(3.3)	53(8.0)	30707(3.3)	
ODDFLG	Disorder not reported	828971(89.9)	602(91.2)	828369(89.9)	0.276
	Disorder reported	92754(10.1)	58(8.8)	92696(10.1)	
ADHDFLG	Disorder not reported	694920(75.4)	561(85.0)	694359(75.4)	<0.0001
	Disorder reported	226805(24.6)	99(15.0)	226706(24.6)	
PERSONFLG	Disorder not reported	918937(99.7)	655(99.2)	918282(99.7)	0.0332
	Disorder reported	2788(0.3)	5(0.8)	2783(0.3)	


Figure 2: 2018 Cohort demographic descriptive




(a) Substance Use Problem



(b) Anxiety



(c) Depression Disorder



(d) Trauma

Figure 3: Prevalence Trend of Youth Mental Health Cases

them less significant for this analysis.

Figure 4 presents geospatial heat maps from QGIS [25] illustrating life expectancy (4a), rent costs (4b), actual OUD cases in 2019 (4c), and predicted cases using Logistic Regression (4d). The heat maps for life expectancy and rent costs reveal clusters of hot and cold spots predominantly along coastal regions, particularly in the western US and New England. These patterns align with certain hotspots observed in the heat map of actual OUD cases. Moreover, the heat map of predicted cases highlights the improved performance of the machine-learning model.

3.2 Model Performance


OUD was most effectively predicted using a logistic regression classification algorithm (Ridge classifier) compared with the LASSO, a regression technique that performs both feature selection and regularization by shrinking less important feature coefficients to zero, helping simplify models and preventing overfitting, classifier, and Random Forest classifier (See Figure 5). Logistic regression achieved a balanced accuracy of 76.6%, compared to 76.5% for LASSO and 61.0% for Random Forest.

On the 2018 validation data, the ensembled logistic model achieved a balanced accuracy of 73%, with a sensitivity of 75% and specificity of 71% with an area under the curve (AUC) of 0.81. For the 2019 testing data, the model demonstrated a balanced accuracy of 50%, with a sensitivity of 16% and specificity of 84%. On the 2022 testing data, the model achieved a balanced accuracy of 86%, sensitivity of 84%, and specificity of 87%. The model has an area under the curve (AUC) of 0.82. The ROC curves (6a, 7a) and confusion matrix (6b, 7b) plots for both the 2019 testing data and 2022 validation data are presented in Figures 6 and 7, respectively.


3.3 Feature Importance

Figure 8 presents the top 15 dominant features in the 2018 and 2022 datasets using the logistic regression classifier. The absolute values of the coefficients indicate feature importance with higher absolute values signifying greater importance. The sign of the coefficients reflects the direction of the association with OUD; positive values indicate increased risk while negative values suggest protective effects. For instance, living in the District of Columbia, Puerto Rico, or North Carolina is associated with lower OUD incidence whereas residing in Virginia, Maryland, or Minnesota corresponds to a higher risk.


Two social determinants positively associated with OUD were identified: life expectancy and



(a) Life Expectancy



(b) Rent Cost



(c) 2019 Actual OUD




(d) 2019 Predicted OUD


Figure 4: Map of geographic variation on life expectancy and rent cost and 2019 true OUD cases and predicted cases

Model selection from 2018 data		
Model	Parameters	Balanced accuracy
Lasso	{'penalty': 'l1', 'solver': 'liblinear', 'class_weight': 'balanced'}	0.76511
Ridge	{'penalty': 'l2', 'solver': 'liblinear', 'class_weight': 'balanced'}	0.76648
Random Forest	{'class_weight': 'balanced'}	0.61007

Figure 5: Balance Accuracy and Hyperparameters Setting for Lasso, Ridge Logistic, and Random Forest Model




(a) ROC curve for 2018 Training model




(b) Confusion Matrix on 2019 Testing Data

Figure 6: 2018-2019 Results



(a) ROC curve for 2022 Training model



(b) Confusion Matrix on 2022 Testing Data

Figure 7: 2022 Results

Rank	Features	OOD (2018)	OOD (2022)
1	District of Columbia	-5.932	-5.916
2	Virginia	4.966	4.980
3	Puerto Rico	-4.830	-4.767
4	North Carolina	-3.779	-3.775
5	Life Expectancy	3.534	3.528
6	Minnesota	3.459	3.450
7	Nebraska	-3.430	-3.428
8	Maryland	3.154	3.142
9	Massachusetts	-3.142	-3.129
10	New Jersey	-3.002	-3.006
11	Montana	2.808	2.811
12	Washington	-2.761	-2.767
13	Colorado	-2.742	-2.752
14	South Dakota	-2.414	-2.425
15	Percent of age under 18 in a State	2.409	2.406

Figure 8: Top 15 most important features for 2018 and 2022 testing data using Logistic Regression

the percentage of children in the state. Notably, there were no significant changes in the coefficients or feature rankings between 2018 and 2022. The complete list of all features with ranking can be found in Supplementary Table A1.

4 Discussion

Our analysis yielded the logistic regression classifier with ridge regularization as the best in predicting OUD. This classifier demonstrated strong performance in identifying OUD cases in 2018 and 2022, achieving satisfactory specificity even in 2019 despite slightly lower other results. Consistent geographic factors were identified as top predictors across 2018 and 2022, underscoring the significant impact of local geographic variations on the risk of SUD.

The use of the logistic regression classifier (Ridge) is based entirely on a data-driven approach. To ensure the selected model accommodates diverse data types, the model selection process included both tree-based Random Forest and non-tree models like Lasso and Ridge logistic regression classifiers. This approach minimizes the risk of arbitrary model selection due to unknown relationships between predictors and the target variable. Given the highly imbalanced nature of the dataset, class-weight adjustments were applied to account for the minority class, reducing potential classification bias. These measures enhance the robustness and objectivity of the study results.

Our findings indicate sufficient predictive performance within the same year, such as in 2018 or 2022, but poor performance across different years, such as using 2018 data to predict 2019 outcomes. This discrepancy may stem from significant variations in opioid-related variables across different states and years. For instance, data from the CDC in the United States demon-

strate that the opioid dispensing rate in the United States varies across states and over time. Such changes likely reflect distributional discrepancies (data drift) between the OUD population in the 2018 training set and the 2019 test set, reducing cross-year predictive performance. This effect is more pronounced in the current imbalanced datasets with limited target samples, where minor variations in sample distribution across years can significantly impact model performance. To address this limitation in future studies with larger datasets, incorporating temporal features (e.g., rates of change) and developing year-based stratified models could more effectively capture cross-temporal variations.

Geographic data thereby emerged as an important predictor of variation in the diagnosis of OUD amongst adolescents receiving mental health services and reflected state-by-state differences in policy, economy, and environment. Aggregated patterns at the state level showed similarities in the course of illness while the integration of the U.S. Census provided important contextual information on social determinants and socioeconomic variables explaining the risk of developing OUD. The absence of critical variables such as public health policy changes, intervention efforts, and governance, limited our findings and should be addressed in future research.

Therefore, this model is specific to youth who have accessed mental health services and thus may not be generalized to the broader adolescent population. Future iterations may wish to extend this methodology to include broader population health information. We excluded the substance use diagnosis variables like alcohol, cocaine, or cannabis use in order to minimize bias as these variables are not independent in defining OUD within this dataset.

Even the simplified model, using only individual mental health diagnoses and service utilization variables, achieved highly balanced accuracy without geographic or social determinant data. Apart from geographic factors, mental health conditions, such as, anxiety, depression, and trauma emerged as strong predictors of OUD in adolescents due to the interconnected nature of mental health conditions and substance use. Adolescents with mental health challenges may experience emotional dysregulation, heightened stress, and a lack of coping mechanisms, leading to self-medication behaviors. Opioids, often perceived as quick relief from psychological pain, are more likely to be misused in this vulnerable population. Additionally, trauma [18], such as adverse childhood experiences (ACEs), is strongly linked to in-

creased susceptibility to substance use as a coping strategy, further amplifying the risk of OUD.

Adolescents benefit from preventive strategies focused on resilience-building and coping skills. Tailoring interventions to the developmental context and life circumstances of each population is crucial for effective prevention and treatment. Our study provides individualized predictions that can be aggregated at various levels to inform targeted, cost-effective prevention programs. This approach enables U.S. local governments to better address the complex interplay between local conditions and OUD risk.

5 Conclusion

Youth receiving mental health services face heightened risks of developing OUD due to complex social, psychological, and environmental challenges. Early detection and intervention are crucial for reducing their vulnerability to substance use disorders. Our model shows great potential to identify the risk of OUD in this group and offer timely support to prevent further advancement of substance use disorders.

Excluding prediction power, our analysis reveals significant geographic variation in OUD prevalence, driven by state policies, economic conditions, and healthcare access. This highlights the need for geographically tailored interventions to ensure equitable resource distribution and effective utilization of resources. Enhancing predictive tools from this study can offer insights to guide government policies in reducing such disparities.

Through embedding advanced analytic into policy making, governments can more effectively identify at-risk populations and design community-level strategies that address the root causes of OUD. This approach holds the promise of addressing inequities in the opioid crisis while empowering healthcare systems to take a more proactive stance. It paves the way for more effective and equitable outcomes for youth navigating mental health challenges and substance use disorders, fostering a fairer and more responsive support network.

References

- [1] K. Ahmadi, M. Aminzadeh, and R. Mohseni. Trends and geographic patterns in drug and synthetic opioid overdose deaths – united states, 2013–2019. *MMWR Morb Mortal Wkly Rep*, 70(6):202–207, 2021.

- [2] K. Herlinger and A. Lingford-Hughes. Opioid use disorder and the brain: a clinical perspective. *Addiction*, 117(2):495–505, Feb 2022. [Internet].
- [3] A. M. Yule, R. M. Lyons, and T. E. Wilens. Opioid use disorders in adolescents—updates in assessment and management. *Curr Pediatr Rep*, 6(2):99–106, 2018.
- [4] J. E. Schulenberg, M. E. Patrick, M. K. Johnson, P. M. O’Malley, and J. G. Bachman. Substance use changes and social role transitions: Proximal developmental effects on ongoing trajectories from late adolescence through early adulthood. *Dev Psychopathol*, 22(4):917–932, 2010.
- [5] Y. S. Liu, L. Kiyang, J. Hayward, Y. Zhang, D. Metes, M. Wang, L. W. Svenson, F. Talarico, P. Chue, X. M. Li, R. Greiner, A. J. Greenshaw, and B. Cao. Individualized prospective prediction of opioid use disorder. *Can J Psychiatry*, 68(1):54–63, Feb 2023.
- [6] S. E. McCabe, B. T. West, P. Veliz, V. V. McCabe, S. A. Stoddard, and C. J. Boyd. Trends in medical and nonmedical use of prescription opioids among us adolescents: 1976–2015. *Pediatrics*, 139(4):e20162387, April 2017. [Internet].
- [7] P. Chue, R. Greiner, and M. Johnson. Prevalence and profile of people with co-occurring mental and substance use disorders within a comprehensive mental health system. *Can J Psychiatry*, 53(12):810–821, 2008.
- [8] Boris Birmaher. Bipolar disorder in children and adolescents. *Child and Adolescent Mental Health*, 18(3):127–137, September 2013. [Internet].
- [9] R. B. Johnson, L. R. Smith, and J. Taylor. Demographic and geographic variation in fatal drug overdoses in the united states, 1999-2020. *Ann Am Acad Pol Soc Sci*, 703(1):50–78, 2023.
- [10] A. Hewitt, L. Bond, R. Ferguson, and M. Jones. Trends and geographic patterns in drug and synthetic opioid overdose deaths – united states, 2013-2019. *MMWR Morb Mortal Wkly Rep*, 70(6):202–207, 2021.
- [11] F. J. Elgar, T. K. Pfortner, I. Moor, M. de Looze, and C. Currie. Educational inequalities in the co-occurrence of mental health and substance use problems and its adult socio-economic consequences: A longitudinal study of young adults in a community sample. *Public Health*, 127(8):745–753, 2013.
- [12] N. Andrabi, R. Khoddam, and A. M. Leventhal. Socioeconomic disparities in adolescent substance use: Role of enjoyable alternative substance-free activities. *Soc Sci Med*, 176:175–182, 2017.
- [13] A. Rafiee, S. Karimi, and N. Khaleghi. Multilayer perceptron modeling for social dysfunction prediction based on general health factors in an iranian women sample. *Front Psychiatry*, 14:1283095, 2023.
- [14] D. R. Williams and R. E. Harris. Anxiety disorders among african americans, blacks of caribbean descent, and non-hispanic whites in the united states. *J Anxiety Disord*, 23(5):578–590, 2009.
- [15] A. Smith, B. Johnson, and C. Taylor. Prevalence of mental disorders among people with opioid use disorder: A systematic review and meta-analysis. *Drug Alcohol Depend*, 238:109551, 2022.
- [16] Centers for Disease Control and Prevention. Youth risk behavior surveillance—substance use among youth, 2022. <https://www.cdc.gov/youth-behavior/risk-behaviors/substance-use-among-youth.html> Atlanta (GA): CDC; [updated 2022 Apr 12; cited 2025 Jan 5].
- [17] H. H. Lister, S. C. Marcello, J. J. Lister, A. M. Toto, K. G. Powell, and N. A. Peterson. An innovative integrated care and substance use disorder training model to improve outcomes for people in medically underserved communities. *Fam Syst Health*, 40(4):586–591, 2022.
- [18] R. Smith, T. Brown, L. Nguyen, and A. Taylor. Factors associated with high-risk substance use in persons receiving psychiatric treatment for a primary trauma- and stressor-related disorder diagnosis. *J Dual Diagn*, 19(4):199–208, 2023.
- [19] U.S. Census Bureau. Data profiles | american community survey, 2022. <https://www.census.gov/acs/www/data/data-tables-and-tools/data-profiles/>.
- [20] Substance Abuse and Mental Health Services Administration (SAMHSA). Data we collect: Mental health

- client-level data (mh-cld), 2018. <https://www.samhsa.gov/data/data-we-collect/mh-cld/datafiles> Rockville (MD): SAMHSA; [cited 2025 Jan 5].
- [21] B. Burstein, H. Agostino, and B. Greenfield. Emergency department visits for suicidal ideation in the united states, 2006–2013. *The American Journal of Emergency Medicine*, 37(7):1252–1257, July 2019. [Internet].
- [22] Python Software Foundation. *Python Language Reference, version 3.11*, Oct 2024. <https://docs.python.org/3/reference/index.html>.
- [23] F. Pedregosa, Gaël Varoquaux, A. Gramfort, V. Michel, B. Thirion, O. Grisel, M. Blondel, P. Prettenhofer, R. Weiss, V. Dubourg, et al. Scikit-learn: Machine learning in python. *Journal of machine learning research*, 12(Oct):2825–2830, 2011.
- [24] O. D. Ware, K. A. Lee, B. Lombardi, D. L. Buccino, J. J. Lister, E. Park, et al. Artificial neural network analysis examining substance use problems co-occurring with anxiety and depressive disorders among adults receiving mental health treatment. *J Dual Diagn*, 2024.
- [25] QGIS Development Team. *QGIS Geographic Information System. Open Source Geospatial Foundation Project*, Nov 2023. <https://www.qgis.org>.

6 Appendix A

6.1 Supplementary Table A1: All Features and Ranks

Rank	Features	Description
1	STATEFIP_11	District of Columbia
2	STATEFIP_51	Virginia
3	STATEFIP_72	Puerto Rico
4	STATEFIP_37	North Carolina
5	life_expectancy	Life Expectancy
6	STATEFIP_27	Minnesota
7	STATEFIP_31	Nebraska
8	STATEFIP_24	Maryland
9	STATEFIP_25	Massachusetts
10	STATEFIP_34	New Jersey
11	STATEFIP_30	Montana
12	STATEFIP_53	Washington
13	STATEFIP_8	Colorado
14	STATEFIP_46	South Dakota
15	age_persons_under_18_years	Age Distribution of persons under 18 years
16	STATEFIP_44	Rhode Island
17	STATEFIP_26	Michigan
18	AGE	Age (recoded)
19	STATEFIP_10	Delaware
20	median_home_price	Median Home Price
21	STATEFIP_17	Illinois
22	STATEFIP_1	Alabama
23	STATEFIP_38	North Dakota
24	STATEFIP_29	Missouri
25	age_female_persons	Age Distribution of Female persons
26	STATEFIP_41	Oregon
27	OPISERVICE	Other psychiatric inpatient
28	racial_composition_hispanic	Racial and Ethnic Composition of Hispanic persons
29	age_median_age	Age Distribution of Median age
30	access_to_healthcare	Access to Healthcare
31	IJSSERVICE	Institutions under the justice system
32	RTCSERVICE	Residential treatment center
33	SPHSERVICE	State psychiatric hospital services
34	STATEFIP_22	Louisiana
35	mental_health_statistics	Mental Health Statistics
36	STATEFIP_50	Vermont
37	STATEFIP_36	New York
38	social_mobility	Social Mobility
39	STATEFIP_5	Arkansas
40	population_size	Population Size
41	STATEFIP_45	South Carolina
42	STATEFIP_49	Utah
43	STATEFIP_18	Indiana
44	chronic_diseases_obesity	Chronic Disease Prevalence of Obesity
45	homeownership_rate	Homeownership Rate
46	STATEFIP_40	Oklahoma
47	age_persons_65_years_and_over	Age Distribution of persons 65 years and over
48	STATEFIP_19	Iowa
49	STATEFIP_55	Wisconsin
50	gross_state_product_GSP	Gross State Product (GSP)
51	median_household_income	Median Household Income
52	racial_composition_white	Racial and Ethnic Composition of White persons
53	STATEFIP_9	Connecticut
54	STATEFIP_39	Ohio

55	racial_composition_black	Racial and Ethnic Composition of Black persons
56	STATEFIP_56	Wyoming
57	STATEFIP_28	Mississippi
58	STATEFIP_6	California
59	tax_burden	Tax Burden
60	STATEFIP_32	Nevada
61	PERSONFLG	Personality disorder reported
62	per_pupil_spending	Per Pupil Spending
63	BIPOLARFLG	Bipolar disorder reported
64	property_crime_rate	Property Crime Rate
65	chronic_diseases_diabetes	Chronic Disease Prevalence Diabetes
66	ADHDFLG	Attention deficit/hyperactivity disorder reported
67	rent_costs	Rent Costs
68	percentage_with_higher_education	Percentage with Higher Education
69	quality_of_schools	Quality of Schools
70	poverty_rate	Poverty Rate
71	PDDFLG	Pervasive developmental disorder reported
72	STATEFIP_42	Pennsylvania
73	homelessness_rate	Homelessness Rate
74	CONDUCTFLG	Conduct disorder reported
75	STATEFIP_12	Florida
76	TRAUSTREFLG	Trauma or stressor related disorder reported
77	SCHIZOFLG	Schizophrenia or other psychotic disorder reported
78	STATEFIP_35	New Mexico
79	violent_crime_rate	Violent Crime Rate
80	DEPRESSFLG	Depressive disorder reported
81	ANXIETYFLG	Anxiety disorder reported
82	healthcare_costs	Healthcare Costs
83	population_growth_rate	Population Growth Rate
84	unemployment_rate	Unemployment Rate
85	STATEFIP_48	Texas
86	cost_of_living_index	Cost of Living Index
87	STATEFIP_47	Tennessee
88	OTHERDISFLG	Other mental disorder reported
89	STATEFIP_16	Idaho
90	inequality_measures	Inequality Measures
91	ODDFLG	Oppositional defiant disorder reported
92	racial_composition_asian	Racial and Ethnic Composition of Asian persons
93	CMPSERVICE	SMHA-funded/operated community based program
94	STATEFIP_21	Kentucky
95	high_school_graduation_rate	High School Graduation Rate
96	urbanvsrural_rural	Rural population
97	urbanvsrural_urban	Urban population
98	STATEFIP_15	Hawaii
99	racial_composition_native	Racial and Ethnic Composition of Native persons

Table A1: Feature Detail List