

Research Summary

Hongbo Cai

Supervisor: Arthur Kosowsky

Department of Physics and Astronomy, University of Pittsburgh

I. CMB LENSING RECONSTRUCTION ON THE CURVED SKY

Cosmic Microwave Background(CMB) photons are gravitationlly lensed by the large scale mass distribution. This effect is called CMB lensing. CMB temperature anisotropies(T) and polarization anisotropies(E,B) are distorted by CMB lensing. We can use deflection field or lensing potential to discribe CMB lensing[1]. It is expected to amongst the most powerful cosmological tools for ongoing and upcoming CMB experiments which will help find out premordial gravitationl waves, constrain cosmological parameters, constrain neutrino mass and understand the properties of dark energy and . So it is important to reconstruct the CMB lensing according to the oberved CMB temperature and polarization anisotropies. The process of reconstruct lensing potential is called CMB lensing reconstruction.

There are two aspects where we can see CMB lensing effects: 1.CMB lensing modulates CMB temperature and polarization power spectra(2 point power spectrum). Statistical anisotropy is induced. 2.CMB lensing produces higher-order correlations between the multiple moments. Off-diagonal mode-coupling between map harmonics can be seen. The off-diagonal mode coupling is proportional to lensing potential.[2]

$$\langle a_l^m b_{l'}^{m'} \rangle = C_l^{ab} \delta_{ll'} \delta_{m-m'} (-1)^m \quad (1)$$

$$\left. \langle a_l^m b_{l'}^{m'} \rangle \right|_{\text{lens}} = C_l^{ab} \delta_{ll'} \delta_{m-m'} (-1)^m + \sum_{LM} (-1)^M \begin{pmatrix} l & l' & L \\ m & m' & -M \end{pmatrix} f_{lLl'}^\alpha \phi_L^M \quad (2)$$

Eq(1) and Eq(2) are the covariance of unlensed and lensed CMB multipoles respectively. $\langle a_l^m$ and $\langle b_{l'}^{m'}$ are multipoles for T, E or B-mode multipoles. $\langle \phi_l^m$ is lensing potential multipole. As we can see, the multipole covariance gained an extra term which is proportional to lensing potential For primordial gravitationl waves, CMB lensing can be an important source of confusion[1]. Density perturbations in the linear regime generate only the so-called E-mode polarization[3]. Lensing converts E-mode polarization to B-mode polarization.[4]. Grav-

itationl waves can also produce B-mode polarization[5]. Removing the lensing-generated B-mode is called delensing.

To obtain lensing potential, we can take quadratic combinations of CMB fields and construct quadratic estimators with minimum variance[5]. On small angular scales $L > 10^2$, we are able to neglect the curvature of the sky and apply flat sky estimators[2]. Since lensing is most sensitive to the projected potential at $L < 10^2$ or several degrees on the sky, we need consider the curvature of the sky. In this study, we apply estimators for lensing potential on the curved sky provided by Ref[6] as shown is Eq(3):

$$d_L^{\alpha M} = \frac{A_L^\alpha}{\sqrt{L(L+1)}} \sum_{l_1 m_1} \sum_{l_2 m_2} (-1)^M \begin{pmatrix} l_1 & l_2 & L \\ m_1 & m_2 & -M \end{pmatrix} g_{l_1 l_2}^\alpha(L) a_{l_1}^{m_1} b_{l_2}^{m_2} \quad (3)$$

We reconstruct lensing potential and get noise properties for the full sky. This work now only works for the full sky using TT and EB estimators. And its performance is still limited by now.

This work is meant for improving the existing lensing reconstruction pipline in Advanced Atacama Cosmology Telescope(AdvACT) and Simons Obervatory(SO). I am a member of the both collaborations. I also learned lensing reconstruction technique from this work which is applied in the work related lensing reconstruction bias shown in the section 2 below.

I am now improving the performance of my code used in this work, trying to convert it to patchywork of CMB maps and implementing it into collaboration lensing reconstruction pipeline.

II. BIAS TO CMB LENSING RECONSTRUCTION FROM TEMPERATURE ANISOTROPIES DUE TO REIONIZATION KSZ

In this work, we are investigating a bias to CMB lensing reconstruction from temperature anisotropies due to the reionization kSZ effect(kinematic Sunyaev-Zel'dovich effect) based on simulation.

There are several ongoing and upcoming experiments, including Advanced Atacama Cosmology Telescope(AdvACT)[7], the South Pole Telescope-3G(SPT-3G)[8], the Simons Observatory[9], and CMB Stage-4(CMB-S4)[10]. For these experiments, the CMB lensing power spectrum will be measured with signal-to-noise(S/N) > 100 . At this precision level, we are required to consider more about biases in CMB reconstruction. Most of the biases

can be removed from primary CMB by a multifrequency component separation methods, but they don't work for kSZ effect, since kSZ effect preserves blackbody spectrum of the CMB.[11]

kSZ effect is the Doppler shift of CMB photons induced by Compton-scattering off moving electrons(bulk velocity).[12]. The kSZ signal has its own intrinsic non-Gaussianity and its correlation with CMB lensing field is non-zero.[13].

This kSZ effect produces a CMB temperature change, $\Theta^{\text{kSZ}}(\hat{\mathbf{n}}) = \Delta T^{\text{kSZ}}(\hat{\mathbf{n}})/T_{\text{CMB}}$,

$$\Theta^{\text{kSZ}}(\hat{\mathbf{n}}) = -\sigma_T \int \frac{d\eta}{1+z} e^{-\tau} n_e(\hat{\mathbf{n}}, \eta) \mathbf{v}_e \cdot \hat{\mathbf{n}} \quad (4)$$

/citeFerraro:2017fac

kSZ anisotropies can be produced when large fluctuation in electron density appears. There are two epochs when they can be produced: 1.a “late-time” contribution from redshifts $0 < z < 6$ in which inhomogeneities are large due to gravitational growth of structure 2.earlier during the epoch of reionization(from first stars, $6 < z < 20$) when hydrogen gets ionized again by the ultraviolet radiation of the first structures and the fluctuation electron density are caused by the fluctuations in the ionization fraction.[12] [14]. It is expected to be correlated with lensing field.

Simone Ferraro and J. Colin Hill have investigated the case of “late time” kSZ in [12]. According to their results, the bias induced by “late time” kSZ to CMB lensing auto-power spectrum measurements can be as large as approximately %1, %6, and %8 for Planck, CMB-S3, and CMB-S4, respectively, when using $l_{\text{max}} = 4000$, and about half of that for $l_{\text{max}} = 3000$. Thus, for CMB-S3 and CMB-S4 lensing measurements, the kSZ-induced bias cannot be neglected.

For the case of reionization kSZ, it could also contribute at some level. We are trying to estimate reionization-induced bias to CMB lensing reconstruction from temperature anisotropies based on lensed temperature anisotropies and reionization kSZ simulations from Websky[15] and reionization kSZ map from Ref[14]. In this study, we apply flat-sky reconstruction pipeline to cutout patchy lensed temperature maps with reionization kSZ and without reionization kSZ with different noise levels, beam sizes and l_{max} . We compare their reconstruction lensing auto-powerspectra and see how much bias the reionization kSZ induces. I am now simulating the bias and trying to understand the bias quantitatively. This

work is expected to be helpful for separating reionization kSZ signals from lensing signals.

-
- [1] Antony Lewis and Anthony Challinor. Weak gravitational lensing of the CMB. *Phys. Rept.*, 429:1–65, 2006. doi:10.1016/j.physrep.2006.03.002.
 - [2] Wayne Hu and Takemi Okamoto. Mass reconstruction with cmb polarization. *Astrophys. J.*, 574:566–574, 2002. doi:10.1086/341110.
 - [3] Marc Kamionkowski, Arthur Kosowsky, and Albert Stebbins. Statistics of cosmic microwave background polarization. *Phys. Rev.*, D55:7368–7388, 1997. doi:10.1103/PhysRevD.55.7368.
 - [4] Matias Zaldarriaga and Uros Seljak. Gravitational lensing effect on cosmic microwave background polarization. *Phys. Rev.*, D58:023003, 1998. doi:10.1103/PhysRevD.58.023003.
 - [5] Wayne Hu. Weak lensing of the CMB: A harmonic approach. *Phys. Rev.*, D62:043007, 2000. doi:10.1103/PhysRevD.62.043007.
 - [6] Takemi Okamoto and Wayne Hu. CMB lensing reconstruction on the full sky. *Phys. Rev.*, D67:083002, 2003. doi:10.1103/PhysRevD.67.083002.
 - [7] S. W. Henderson et al. Advanced ACTPol Cryogenic Detector Arrays and Readout. *J. Low. Temp. Phys.*, 184(3-4):772–779, 2016. doi:10.1007/s10909-016-1575-z.
 - [8] B. A. Benson et al. SPT-3G: A Next-Generation Cosmic Microwave Background Polarization Experiment on the South Pole Telescope. *Proc. SPIE Int. Soc. Opt. Eng.*, 9153:91531P, 2014. doi:10.1117/12.2057305.
 - [9] Peter Ade et al. The Simons Observatory: Science goals and forecasts. *JCAP*, 1902:056, 2019. doi:10.1088/1475-7516/2019/02/056.
 - [10] Kevork N. Abazajian et al. CMB-S4 Science Book, First Edition. 2016.
 - [11] Robert E. Smith, Carlos Hernandez-Monteagudo, and Uros Seljak. Impact of Scale Dependent Bias and Nonlinear Structure Growth on the ISW Effect: Angular Power Spectra. *Phys. Rev.*, D80:063528, 2009. doi:10.1103/PhysRevD.80.063528.
 - [12] Simone Ferraro and J. Colin Hill. Bias to CMB Lensing Reconstruction from Temperature Anisotropies due to Large-Scale Galaxy Motions. *Phys. Rev.*, D97(2):023512, 2018. doi:10.1103/PhysRevD.97.023512.
 - [13] Kendrick M. Smith and Simone Ferraro. Detecting Patchy Reionization in the Cosmic Microwave Background. *Phys. Rev. Lett.*, 119(2):021301, 2017. doi:

10.1103/PhysRevLett.119.021301.

- [14] Marcelo A. Alvarez. The Kinetic Sunyaev–Zel’dovich Effect From Reionization: Simulated Full-sky Maps at Arcminute Resolution. *Astrophys. J.*, 824(2):118, 2016. doi:10.3847/0004-637X/824/2/118.
- [15] George Stein, Marcelo A. Alvarez, J. Richard Bond, Alexander van Engelen, and Nicholas Battaglia. The Websky Extragalactic CMB Simulations. 2020.