

Blood Pressure Monitoring

CCG: C-1016 (CCG)

MCG Health
Chronic Care
28th Edition

- Assessment
- Quality Measures
- Plan of Care
- References
- Footnotes

Assessment

Q1. Patient: What was your most recent blood pressure reading? **QM**

- Reading was: _____
- Unknown [P1]
- No recent reading [P1]

Q2. Patient: What is your goal blood pressure range? **QM**

- My range is: _____
- Unknown [P1]

Q3. Patient: How often do you check your blood pressure? (Check all that apply.) **QM**

- 2 or more times a day [Q4]
- Once a day [Q4]
- 2 or more times a week [Q4] [P1]
- Once a week or less [Q4] [P1]
- After activity [Q4]
- When it seems like it may be high [Q4]
- Other, specify: _____ [Q4] [P1]
- Unknown [P1]

Q4. Patient: Why do you check your blood pressure? (Check all patient reports.) **QM**

- After a myocardial infarction
- After a stroke
- Coronary artery disease
- Diabetes
- Dizziness or vertigo
- High blood pressure
- Vital sign check
- Other, specify: _____ [P1]
- Unknown [P1]
- Not all appropriate reasons to check blood pressure reported [P1]

Q5. Patient: Do you measure your blood pressure at home?

- Yes
- No [P1]
- Unknown [P1]

Q6. Case Manager: Is patient's blood pressure within goal range? **QM**

- Yes
- No [P1]
- Unknown [P1]

Quality Measures

- Disease Management Quality Measures:
 - Information provided from this question may be requested by various accrediting organizations.

Plan of Care

Problems

P1. Blood pressure monitoring information needed

- **Goal:** Blood pressure self-monitoring improved

- **Interventions:**

- **ASSIST:** Identifying patient's goal priorities, willingness/ability to achieve goals, and preferences for communication and follow-up
- **ASSIST:** Goal setting that is specific, measurable, achievable, relevant, and with target completion time (eg, short/long-term, prioritized)
- **ASSIST:** Appointment scheduling with provider for evaluation of blood pressure problems or creation of self-care plan to include:
 - Obtaining home blood pressure monitor with correct cuff size for patient
 - Identifying blood pressure targets
- **EDUCATE:** Why blood pressure monitoring is important
- **EDUCATE:** When to check blood pressure
- **EDUCATE:** How to check blood pressure(1)(2)(3):
 - Take blood pressure medication at same time each day.
 - Avoid smoking, caffeine intake, and exercise for 30 minutes prior to taking blood pressure.
 - Fit and position cuff properly (cuff should be appropriate size for patient). Do not put cuff over clothing.
 - Support arm on flat surface at level of heart, with back supported and both feet on floor, for at least 5 minutes prior to taking blood pressure.
 - Measure blood pressure in same arm each reading.[A](1)
 - Blood pressure goal and the need to know own blood pressure from recent measurement
 - Keeping a diary of blood pressure readings and reviewing diary with provider at appointments
- **EDUCATE: When to call the provider:**
 - Blood pressure outside of provider's parameters
 - There is a 10 mm Hg or greater difference in pressure between right and left arm.[B](4)
- **COORDINATE:** Home care referral, if indicated
- **COORDINATE:** Follow-up contact to evaluate blood pressure monitoring, referral status, and need for further assistance(5)(6)
- **SEND:** How to check your blood pressure (Illustrated)  (Spanish)
- **SEND:** Blood pressure diary (Illustrated)  (Spanish)

References

1. Fetzter S. Vital signs. In: Perry AG, Potter PA, Ostendorf WR, editors. Nursing Interventions and Clinical Skills. 7th ed. Elsevier; 2020:101-132. [Context Link 1, 2, 3]
2. Wermers R. Hypertension. In: Harding MM, Kwong J, Hagler D, Reinisch C, editors. Lewis's Medical-Surgical Nursing: Assessment and Management of Clinical Problems. 12th ed. St. Louis, MO: Mosby; 2023:797-818. [Context Link 1]
3. Blood pressure measurement. In: Lippincott Nursing Procedures. 9th ed. Wolters Kluwer; 2023:80-83. [Context Link 1]
4. Fang JC, O'Gara PT. History and physical examination: an evidence-based approach. In: Libby P, Bonow RO, Mann DL, Tomaselli GF, Bhatt DL, Solomon SD, editors. Braunwald's Heart Disease: A Textbook of Cardiovascular Medicine. 12th ed. Elsevier; 2022:123-140. [Context Link 1, 2]
5. Care coordination and care transitions to support adherence. In: Perez R, Rogers S, Prince M, Fraser K, editors. Case Management Adherence Guidelines. 2020 ed. Case Management Society of America; 2022:43-56. [Context Link 1]
6. Monitoring and evaluation. In: Gillingham DC, editor. Foundations of Case Management. Blue Bayou Press; 2021:51-52. [Context Link 1]

Footnotes

[A] Patients may have up to a 10-point difference in readings between their two arms. For ongoing monitoring, blood pressure should be taken in the arm with the higher reading.(1) [A in Context Link 1]

[B] The difference in readings may be associated with subclavian artery disease, supraaortic stenosis, aortic coarctation, or aortic dissection.(4) [B in Context Link 1]

Last Update: 3/14/2024 3:32:10 AM

Build Number: 28.1.20240313235330.016020