

Depression

TC: TC-0032 (TC)

[Link to Codes](#)**MCG Health**Transitions of Care
28th Edition

- Patient Interview
 - Self-Care
 - Activity
 - Nutrition
 - Medicine
 - Complications
 - Follow-Up Care
- Case Management Evaluation
- Plan of Care
- Evidence Summary
- References
- Footnotes
- Codes

Patient Interview

Self-Care

Q1. Patient: What are you doing to keep your depression from getting worse? (Check all that apply.)

- Adhere to self-care plan
- Ask for help when feelings are overwhelming
- Avoid things that cause feeling worse
- Counseling or therapy attendance
- Exercise program
- Follow a daily care plan
- Follow up with doctors
- Get all follow-up laboratory tests when due
- Keep a diary about feelings, activities, eating times, and medicines taken
- Keep all doctor appointments
- Know when to contact the doctor or therapist
- Lose weight [Q2]
- Manage situation or event triggers
- Manage stress
- Meal planning (diet) changes
- Seek emergency care for signs of crisis plan not working
- Set realistic expectations for feeling better, managing symptoms, and recovery
- Stop using alcohol, tobacco, or drugs
- Support group attendance
- Take medicine
- Vaccinations
- Other, specify: _____ [P1]
- Nothing [P1]
- Unknown [P1] [P2]
- Not all care measures understood [P2]

Q2. Patient: What are you doing to lose weight? (Check all that apply.)

- Avoid sweets
- Be more active
- Eat fewer calories or less fat
- Meal planning (diet) changes
- Skip meals [P1] [P2] Nutrition Assessment, Adult [TC](#)
- Take diet supplements [P1] [P2] Nutrition Assessment, Adult [TC](#)

- Other, specify: _____ [P1] [P2] Nutrition Assessment, Adult TC
- Nothing [P1] [P2] Nutrition Assessment, Adult TC
- Unknown [P1] [P2] Nutrition Assessment, Adult TC
- Not applicable

Q3. Patient: Do you keep a diary of your mood and feelings?

- Yes
- No [P2]
- Unknown [P2]

Q4. Patient: Do you use tobacco, alcohol, or drugs? (Check all that apply.)

- Yes, tobacco use [P2] Tobacco Use TC
- Yes, alcohol use [P2] [P3] Alcohol Use TC
- Yes, drug use [P2] [P3] Substance Use TC
- No
- Unknown [P1] Tobacco Use TC Alcohol Use TC Substance Use TC

Q5. Patient: Are you having any problems with your self-care? (Check all that apply.)

- Affording medicine or supplies [P4] [P5] [P6] Financial Status and Benefits TC
- Behavior indicating cognitive impairment [P4] [P6] Cognitive Impairment TC Behavioral Health and Cognition Evaluation with Caregiver TC
- Blood pressure care [P4] [P6] Blood Pressure Monitoring TC
- Blood sugar care [P4] [P6] Blood Glucose Monitoring TC
- Blood thinner care [P4] [P6] Anticoagulation Management TC
- Bowel program [P4] [P6] Bowel Management TC
- Caregiver help needed [P4] [P6] [P7] Caregiver Resources Evaluation TC
- Community resource information needed [P4] [P6] Community Resources TC
- Exercising and staying active [P4] [P6] Exercise TC
- Fatigue or weakness [P4] [P6] Fatigue or Weakness TC
- Pain [P4] [P6] Pain Assessment TC
- Planning for long-term care needs [P4] [P6] Advance Care Planning TC Palliative or End-of-Life Evaluation TC
- Safety in the home [P4] [P6] Home Safety Evaluation TC
- Skin care management [P4] [P6] Pressure Injury Monitoring and Management TC
- Social support [P4] [P6] Social Support TC
- Taking medicine as prescribed [P4] [P6] [P8] Medication Adherence TC
- Transportation for care [P4] [P6] [P9] Transportation TC
- Trouble gaining weight [P4] [P6] Nutritional Status, Adult TC Nutritional Management to Gain Weight TC
- Trouble losing weight [P4] [P6] Nutritional Status, Adult TC Nutritional Management to Lose Weight TC
- Trouble making lifestyle changes [P4] [P6] Change Readiness TC
- Trouble understanding how to do self-care [P4] [P6] Health Literacy TC
- Other, specify: _____ [P4] [P6]
- No
- Unknown [P4] [P6]

Activity

Q1. Patient: What do you do to stay active? (Check all that apply.)

- Active range of motion exercises [Q2] [Q3]
- Aerobic exercise (eg, jogging, swimming, dancing) [Q2] [Q3]
- Low-impact exercises [Q2] [Q3]
- Strength training exercises [Q2] [Q3]
- Other, specify: _____ [Q2] [Q3]
- Nothing [P1] [P2]
- Unknown [Q2] [Q3] [P1] [P2]

Q2. Patient: How often do you exercise and for how long? (Check all that apply.)

- At least 3 times a week
- 1 or 2 times a week [P1] [P2]
- Inconsistently [P1] [P2]
- More than 30 minutes each session
- 15 to 30 minutes each session
- Less than 15 minutes each session [P1] [P2]
- Other, specify: _____ [P1] [P2]
- Unknown [P1] [P2]
- Not applicable

Q3. Patient: Do you have any problems when you exercise? (Check all that apply.)

- Chest pain [P1]
- Dizziness [P1]
- Trouble breathing [P1]
- Weakness [P1]
- Other, specify: _____ [P1]
- No
- Unknown [P2]
- Not applicable

Nutrition

Q1. Patient: What meal plan (diet) do you follow? (Check all that apply.)

- DASH Diet Knowledge Insufficient - DASH Diet [TC](#) Nutrition Assessment, Adult [TC](#)
- Heart healthy Diet Knowledge Insufficient - Heart Healthy [TC](#) Nutrition Assessment, Adult [TC](#)
- Kidney diet Diet Knowledge Insufficient - Renal Dysfunction [TC](#) Nutrition Assessment, Adult [TC](#)
- Low carb or low glycemic index Diet Knowledge Insufficient - Low Glycemic Index [TC](#) Nutrition Assessment, Adult [TC](#)
- Other, specify: _____ Nutrition Assessment, Adult [TC](#)
- None [P1] [P2] Nutrition Assessment, Adult [TC](#)
- Unknown [P1] [P2] Nutrition Assessment, Adult [TC](#)

Medicine

Q1. Patient: What medicine do you take for your depression? (Check all that apply.)

- Atypical antidepressants (eg, bupropion, mirtazapine, trazodone) [Q2] [P10] [P11]
- Monoamine oxidase inhibitors (MAOIs) (eg, isocarboxazid, phenelzine, tranylcypromine) [Q2] [P10] [P12]
- Serotonin-norepinephrine reuptake inhibitors (SNRIs) (eg, desvenlafaxine, duloxetine, venlafaxine) [Q2] [P10] [P13]
- Selective serotonin reuptake inhibitors (SSRIs) (eg, citalopram, fluoxetine, sertraline) [Q2] [P10] [P13]
- Tricyclic antidepressants (TCAs) (eg, amitriptyline, clomipramine, trimipramine) [Q2] [P10] [P14]
- Other, specify: _____ [Q2] [P10]
- None [P15]
- Unknown [P15]

Q2. Patient: Are you having side effects from your medicine? (Check all that apply.)

- Agitation [P8] [P10]
- Blurred vision [P8] [P10]
- Constipation [P8] [P10]
- Diarrhea [P8] [P10]
- Dizziness [P8] [P10]
- Dry mouth [P8] [P10]
- Fast heartbeat [P8] [P10]
- Headache [P8] [P10]
- Inability to sleep [P8] [P10]
- Inability to urinate [P8] [P10]
- Light sensitivity [P8] [P10]
- Nausea or vomiting [P8] [P10]
- Sexual dysfunction (abnormal ejaculation, impotence, or loss of orgasm) [P8] [P10]
- Sunburn [P8] [P10]
- Tiredness [P8] [P10]

- Weight loss or gain [P8] [P10]
- Other, specify: _____ [P8] [P10]
- None
- Unknown [P8] [P10]
- Not applicable

Q3. Patient: Are you taking any new medicines prescribed by the doctor?

- Yes, specify: _____ [P16]
- No, I do not have any new medicine.
- No, I haven't started taking the new medicine. Medication Adherence  TC
- Unknown Medication Adherence  TC

Q4. Patient: Are you taking any new medicines that you got without a prescription (eg, over-the-counter medicines, vitamins, minerals, or herbal or dietary supplements)?

- Yes, specify: _____ [P16]
- No
- Unknown

Q5. Patient: Are there any medicines that you are no longer taking?

- Yes, specify: _____ [P16]
- No
- Unknown

Q6. Patient: Are there any medicines that you are taking more or less of?

- Yes, specify: _____ [P16]
- No
- Unknown

Q7. Patient: Do you have a list of your current medicine information?

- Yes
- No [P16]
- Unknown

Q8. Patient: Do you have any medicines that have 1 refill or less?

- Yes [P8] [P17]
- No
- Unknown [P8] [P17]
- Not applicable

Q9. Patient: Do you ever miss taking your medicine?

- Yes [P8] Medication Adherence  TC
- No
- Unknown [P8] Medication Adherence  TC
- Not applicable

Complications

Q1. Patient: What are the signs of depression? (Check all that apply.)

- Aches and pains for no reason
- Changes in eating habits
- Crying spells for no reason
- Feeling tired, sluggish, or physically drained
- Feeling worthless or hopeless
- Little or no interest in activities or hobbies
- Little or no interest in being with friends and family
- Poor personal hygiene, not wanting to bathe, or not caring about what you wear
- Thoughts of death, suicide, or harming self or someone else
- Trouble concentrating or remembering things
- Trouble making decisions
- Trouble sleeping or sleeping too much
- Other, specify: _____ [P2]
- None [P2]
- Unknown [P2]

- Not all symptoms understood [P2]

Q2. Patient: Are you currently having symptoms of depression?

- Yes [P1]
- No
- Unknown [P1] [P2]

Q3. Patient: What do you do if you have signs of depression? (Check all that apply.)

- Avoid things that make me feel worse
- Call behavioral health counselor or therapist
- Call support system for help
- Exercise or do activity
- Get medical care
- Manage stress
- Take medicine, specify: _____
- Other, specify: _____ [P1] [P2]
- Nothing [P1] [P2]
- Unknown [P1] [P2]
- Not all care measures understood [P2]

Follow-Up Care

Q1. Patient: Are you supposed to have surgery or a procedure? If yes, when is it scheduled? (Check all that apply and include scheduled date.)

- Electroconvulsive therapy (ECT), date scheduled: _____ Preoperative Care
Coordination - Assessment and Planning [TC](#)
- Transcranial magnetic stimulation, date scheduled: _____ Outpatient Services -
Assessment and Planning [TC](#)
- Other, specify: _____ Preoperative Care Coordination - Assessment and Planning [TC](#)
- None
- Unknown [P2]

Q2. Patient: Has your doctor ordered any follow-up care? (Check all that apply.)

- Bright light therapy, specify: _____
- Dietitian Outpatient Services - Assessment and Planning [TC](#)
- Home care [P2] Outpatient Services - Assessment and Planning [TC](#)
- Laboratory testing, specify: _____ [Q3]
- Community services referral, specify: _____ Appointment Scheduling [TC](#)
- Behavioral health provider referral, specify: _____ Appointment Scheduling [TC](#)
- Psychiatrist referral, specify: _____ Appointment Scheduling [TC](#)
- Social worker referral, specify: _____ Appointment Scheduling [TC](#)
- Specialist referral, specify: _____ Appointment Scheduling [TC](#)
- Rehabilitation [P2] Outpatient Services - Assessment and Planning [TC](#)
- Other, specify: _____ [P2] Outpatient Services - Assessment and Planning [TC](#)
- None [P1]
- Unknown [P2]

Q3. Patient: When is your laboratory work due?

- This week
- 1 to 2 weeks
- 3 to 4 weeks
- 1 to 3 months [P2]
- 4 to 11 months [P18]
- 1 year or more [P18]
- Other, specify: _____ [P2] [P18]
- Unknown [P1] [P18]

Case Management Evaluation

Q1. Case Manager: Does the patient have a written self-care plan?

- Yes
- No [P1] [P6]
- Unknown [P1] [P6]

Q2. Case Manager: Is the patient at risk for further complications? (Check all that apply.)

- Adhering to self-care plan [P1] [P4] Change Readiness [TC](#)
- Affording medication or supplies [P1] [P4] [P5] Financial Status and Benefits [TC](#)
- Behavior indicating cognitive impairment [P1] [P4] Behavioral Health and Cognition Evaluation with Caregiver [TC](#)
- Caregiver help needed [P1] [P4] [P7] [P19] Caregiver Resources Evaluation [TC](#)
- Community resource information needed [P1] [P4] Community Resources [TC](#)
- Exercising and staying active [P1] [P4] Exercise [TC](#)
- Health literacy [P1] [P4] Health Literacy [TC](#) Home Care Coordination - Assessment and Planning [TC](#)
- Medication nonadherence [P1] [P4] Medication Adherence [TC](#) Home Care Coordination - Assessment and Planning [TC](#)
- Pain [P1] [P4] Pain Assessment [TC](#)
- Rehospitalization [P1] [P4] Hospitalization Risk Evaluation [TC](#)
- Safety in the home [P1] [P4] Home Safety Evaluation [TC](#) Home Care Coordination - Assessment and Planning [TC](#)
- Self-care education or training needed [P1] [P4] [P6] Home Care Coordination - Assessment and Planning [TC](#)
- Social support [P1] [P4] Social Support [TC](#)
- Transportation for care [P1] [P4] [P9] Transportation [TC](#)
- Trouble making lifestyle changes [P1] [P4] Change Readiness [TC](#)
- Other, specify: _____ [P1] [P4]
- None identified
- Unknown [P1] [P4]

Q3. Case Manager: Is continued care coordination needed?

- Yes
- No Transitions of Care Case Management Closure [TC](#)
- Unknown [P4]

Plan of Care

Problems

P1. Depression complication risk present

- **Goal:** Depression complication risk evaluated and managed **NNN**
 - **Interventions:**
 - **ASSIST:** Identifying patient's goal priorities, willingness/ability to achieve goals, and preferences for communication and follow-up
 - **ASSIST:** Goal setting that is specific, measurable, achievable, relevant, and with target completion time (eg, short/long-term, prioritized)
 - **ASSIST:** Appointment scheduling for evaluation of problems, comorbidities, and self-care plan[A]:
 - Recently hospitalized patient's evaluation within 1 week of discharge(1)**N**
 - Recently hospitalized patient's evaluation with outpatient healthcare providers (eg, nurse, social worker) within 1 to 3 days of discharge(2)(3)(4)
 - Appointments for patient receiving nonpharmacologic therapies should occur several times in first 3 months to determine if therapy is helping to improve symptoms.
 - Appointments for patient receiving pharmacologic therapy should occur within 2 weeks of initiating treatment, then every 2 to 3 months until patient is stable or in remission. More frequent follow-up may be appropriate for patients with high risk of suicidal ideation.
 - Appointment once a month or more for patient with questionable adherence to treatment plan
 - **EDUCATE:** When and how to get help for worsening depression problems
 - **EDUCATE:** Follow-up appointments for assessment of clinical status
 - **EDUCATE:** Treating depression can help improve outcomes of coexisting chronic medical conditions.
 - **EDUCATE:** Treatment of coexisting chronic medical conditions may alleviate or eliminate depression.(5)
 - **COORDINATE:** Specialist referral, if indicated
 - **COORDINATE:** Care team meeting, if needed(6)(7)
 - **COORDINATE:** Follow-up to review progress with self-care plan, referral status, and need for further assistance(2)(8)

- **RECONCILE:** Communicate care plan changes to entire care team.(9)(10)

P2. Depression self-care information needed

- **Goal:** Depression self-care information understood(11) **NNN**

o **Interventions:**

- **ASSIST:** Identifying patient's goal priorities, willingness/ability to achieve goals, and preferences for communication and follow-up
- **ASSIST:** Goal setting that is specific, measurable, achievable, relevant, and with target completion time (eg, short/long-term, prioritized)
- **ASSIST:** Support system (eg, family, friends) identification, enlistment, and education
- **ASSIST:** Obtaining a medical alert bracelet or necklace
- **EDUCATE:** Major depressive disorder, also known as clinical depression, is a debilitating mood disorder.(12)
 - It is characterized by a pervasive and persistent sadness, despair, and diminished ability to think or make decisions.
 - It is accompanied by low self-esteem and a loss of interest or the ability to experience pleasure.
 - It is a common condition among patients with chronic medical illnesses.
- **EDUCATE: Self-care plan information:**
 - Appointment scheduling and attendance
 - Adherence to plan necessary to achieve goals, and complications of nonadherence (eg, exacerbations, progression, comorbidities)(13)
 - Medication purpose, use, effects, and considerations (eg, timing before meals or exercise)(14)
 - Set small goals every day (eg, get out of bed at a certain time, finish one chore or project, or do something that brings pleasure like listen to music).(15)
 - Health checks, diary, and action plans (eg, write down thoughts, feelings, daily activities, eating times, and when medication taken)
 - Behavioral health treatment (eg, cognitive behavioral therapy (CBT), individual therapy, family therapy)(12)(16)
 - Symptom management
 - Relapse signs (eg, insomnia, decreased appetite, poor personal hygiene)(12)
 - Caregiver support plan, as needed
 - Crisis care plan: turn off distractions (eg, TV, radio), seek emergent care, call hotline, call police if escalates; caregivers advised to not argue(17)
 - Diet considerations (eg, limit caffeine and alcohol, avoid foods containing tyramine if taking an MAOI)(18)(19)
 - Exercise program within provider's recommendations(1)(20)(21) **N**
 - Sleep hygiene (eg, regular pattern, calm sleep environment)(22)
 - Stress management (eg, relaxation and coping techniques)
 - Substance use avoidance(5)(23)
 - Trigger avoidance (eg, high-stress situations, sleep deprivation)(1)(18)
 - Functional changes care (eg, adaptations and support for ADL and IADL)(24)
 - Comorbid condition management(5)
 - Wellness checkups (eg, vaccinations, screenings)(25)
 - Laboratory monitoring, as needed
 - Safety awareness and management (eg, remove suicide risks, de-escalation techniques)(26)(27)
 - Support group participation
 - Medical alert bracelet/necklace, obtain and wear
 - Tobacco cessation plan(25)
- **EDUCATE:** Complications with depression include:
 - Self-harm (eg, abuse, mutilation)
 - Substance misuse (eg, increased use, use interferes with ADL/IADL)
 - Social isolation
 - Sleep problems
- **EDUCATE: When to get emergency help(18):**
 - Heart attack (eg, persistent chest pain; pain radiating to the arms, shoulders, neck, back, and/or jaw; cool clammy skin)(28)
 - Stroke (eg, face drooping, arm weakness, speech problems, bad headache, vision changes, loss of balance or coordination)(29)
 - Suicidal or homicidal actions or plans(30)
 - Fatigue, extreme
 - Respiratory impairment (eg, gasping for air, choking, cyanosis, chest pain, severe dyspnea)(31)
 - Crisis plan not working
 - Delirium (eg, reduced awareness, agitation)
 - Hallucinations or delusions, increased or new
 - Psychotic behavior (eg, threatening)

- Seizures, new or complicated (eg, seizure longer than 5 minutes, multiple seizures with loss of consciousness, head injury during seizure)(32)
- **EDUCATE: When to call the provider for behavior changes(14):**
 - Depression (eg, feeling sad, hopeless, overwhelmed)(18)
 - Fatigue, unrelenting
 - Insomnia
 - Mood changes (eg, depression, anger, agitation)
 - Self-care changes (eg, ADL or IADL problems)(24)
 - Substance misuse (eg, increased use, use interferes with ADL/IADL)
 - Appointments missed (eg, support group, provider)
 - Cognitive changes (eg, thinking problems, forgetfulness worsens)
 - Life event, major (eg, job loss, death of someone close)
- **EDUCATE: When to call the provider(12)(14):**
 - Constipation
 - Gastrointestinal distress (eg, nausea, vomiting, diarrhea, cramping)
 - Heart rate outside of provider's parameters
 - Medication problems (eg, side effects, access, administration)
 - Medications newly ordered from other providers
 - Pain changes (eg, new or worsening, unresolved with treatment)
 - Pregnant or planning to get pregnant
 - Weight gain or loss without reason
- **COORDINATE:** Behavioral health services referral, if indicated
- **COORDINATE:** Psychiatrist referral, if indicated(1)
- **COORDINATE:** Home care referral, if indicated(9)(11)
- **COORDINATE:** Telehealth referral, if indicated(33)(34)
- **COORDINATE:** Crisis services referral, as needed
- **COORDINATE:** Outpatient imaging, tests, or procedures, if indicated[C]
- **COORDINATE:** Medical equipment and supplies, if indicated[D]
- **COORDINATE:** Tobacco use treatment, as needed
- **COORDINATE:** Chronic condition management referral, if indicated(35)
- **COORDINATE:** Rehabilitation therapy referral, if indicated
- **COORDINATE:** Dietitian referral, if indicated
- **COORDINATE:** Social worker referral, as needed(18)
- **COORDINATE:** Support group referral, as needed(36)
- **COORDINATE:** Community services referral, as needed(37)
- **COORDINATE:** Follow-up contact with patient to evaluate understanding and adherence to self-care plan, referral status, and need for further assistance(2)(8)
- **RECONCILE:** Self-care plans from involved care team members
- **RECONCILE:** Communicate care plan changes to entire care team.(9)(10)
- **SEND:** What you need to know - Depression  (Spanish)
- **SEND:** Your care plan - Depression  (Spanish)
- **SEND:** When to get help - Depression  (Spanish)
- **SEND:** Daily care for your condition - Depression  (Spanish)
- **SEND:** Be ready for your doctor visit - Depression  (Spanish)
- **SEND:** Be ready with questions for your doctor - Depression  (Spanish)
- **SEND:** Major Depressive Disorder, Adult: Discharge Information 
- **SEND:** Caregiver tips  (Spanish)
- **SEND:** Choosing a place to live  (Spanish)
- **SEND:** Help with sleeping or insomnia  (Spanish)
- **SEND:** Tips for good mental health  (Spanish)
- **SEND:** When you are pregnant - Depression  (Spanish)
- **SEND:** How to stop smoking  (Spanish)
- **SEND:** Symptoms diary (Illustrated)  (Spanish)
- **SEND:** Coronavirus disease 2019 (COVID-19) vaccines  (Spanish)
- **SEND:** Flu vaccine  (Spanish)

P3. Unhealthy behaviors potentially causing complications with treatment plan

- **Goal:** Information, support, and reinforcement of healthy behaviors to facilitate treatment plan understood(38)
 - **Interventions:**
 - **RECONCILE:** Treatment plan with patient's willingness and ability to facilitate in presence of unhealthy behaviors
 - **ASSIST:** Identifying types of unhealthy behaviors patients may exhibit, such as(39):
 - Aggressiveness, hostility, or abusiveness
 - Intentional nonadherence
 - Ineffective coping (eg, denial, poor insight, unmanageable anger, low self-esteem)
 - Self-care neglect (eg, not bathing, dressing, eating)
 - Social isolation or withdrawal
 - **COORDINATE:** Communication to care team members regarding potential problems with treatment plan related to patient's unhealthy behaviors
 - **ASSIST:** Appointment scheduling with care providers if unhealthy behaviors causing problems with patient's treatment plan
 - **ASSIST:** Creating behavior support plan to help facilitate treatment, including:
 - Identifying patient's goal priorities, willingness/ability to achieve goals, and preferences for communication and follow-up
 - Identifying factors that contribute to unhealthy behaviors and alternative behaviors that may help patient manage
 - Identifying support people who can help patient with behavior changes
 - Setting limits on patient's behavior and withdrawing attention from patient if behavior presents
 - Creation of emergency action plan if patient's unhealthy behavior escalates (eg, patient risk for harm to self or others)(40)
 - Providing positive reinforcement of healthy behaviors
 - **EDUCATE:** Unhealthy behaviors and potential consequences for treatment or outcomes
 - **EDUCATE:** Goals to help overcome unhealthy behaviors, including interventions, steps, and time frames
 - **EDUCATE:** Alternative coping mechanisms to help patient with behavior changes, such as(41)(42):
 - Social participation (eg, community groups, social organizations, family gatherings)
 - Recreation or hobbies participation
 - Behavioral health support (eg, counseling), role and benefit
 - Relaxation techniques (eg, deep breathing, guided imagery, music)
 - Stress management (eg, relaxation and coping techniques)
 - **EDUCATE:** When and how to get help for problems(43)
 - **EDUCATE:** Caregivers on managing unhealthy behaviors(41)(44):
 - Provide positive reinforcement for healthy behaviors.
 - Be aware of caregiver's personal feelings and their impact on patient's behavior.
 - Support the patient's self-responsibility (eg, independence with hygiene, taking medications, decision making).
 - Know signs of escalating behavior problems (eg, risk for harm to self or others) and when and how to get help.
 - Use agitation reduction techniques (eg, avoid overstimulation, provide privacy, encourage activities, learn to de-escalate).(45)
 - De-escalation techniques (eg, caregiver remains calm and provides emotional support, do not argue or threaten, stay serious, announce actions beforehand)(18)(39)
 - **COORDINATE:** Behavioral health services referral, if indicated
 - **COORDINATE:** Psychiatrist referral, if indicated
 - **COORDINATE:** Recreational therapy referral, if indicated
 - **COORDINATE:** Protective services or legal authorities, as needed
 - **COORDINATE:** Follow-up to evaluate status with behavior changes, treatment plan, referrals, and need for further assistance(2)(8)
 - **SEND:** Tips for good mental health 📄 (Spanish)
 - **SEND:** Caregiver tips 📄 (Spanish)

P4. Coordination of care needed for depression patient

- **Goal:** Self-care plan for depression communicated to all care team members(9)(11)
 - **Interventions:**
 - **RECONCILE:** Self-care plans from involved care team members
 - **ASSIST:** Identifying patient's goal priorities, willingness/ability to achieve goals, and preferences for communication and follow-up
 - **ASSIST:** Goal setting that is specific, measurable, achievable, relevant, and with target completion time (eg, short/long-term, prioritized)
 - **ASSIST:** Scheduling and preparing for provider appointments
 - **ASSIST:** Creation of a self-care plan between patient and providers to include(46)(47):
 - Identifying patient's self-care goals
 - Establishing daily management (medications, diet changes, health checks, and general wellness)

- Creating healthcare diary to track self-care information (eg, vital signs) and to keep written record of care plan changes (eg, changes in medication and when laboratory work is due)
- **ASSIST:** Communication between care team members
- **EDUCATE:** Care team members, contact information, and role in care planning
- **EDUCATE:** Self-care plan and when, where, and from whom to get help for problems with care(7)
- **COORDINATE:** Meeting with care team members, as needed
- **COORDINATE:** Behavioral health services referral, if indicated
- **COORDINATE:** Home care referral, if indicated
- **COORDINATE:** Psychiatrist referral, if indicated(1)
- **COORDINATE:** Rehabilitation therapy referral, if indicated
- **COORDINATE:** Community services referral, as needed
- **COORDINATE:** Social worker referral, as needed(18)
- **COORDINATE:** Follow-up contact with patient to evaluate understanding and adherence to self-care plan, referral status, and need for further assistance(2)(8)
- **COORDINATE:** Chronic condition management referral, if indicated(35)
- **COORDINATE:** Transmission of handoff communication
- **RECONCILE:** Communicate care plan changes to entire care team.(10)
- **SEND:** Your care plan - Depression 📄 (Spanish)
- **SEND:** What you need to know - Depression 📄 (Spanish)
- **SEND:** When to get help - Depression 📄 (Spanish)
- **SEND:** Daily care for your condition - Depression 📄 (Spanish)
- **SEND:** Be ready for your doctor visit - Depression 📄 (Spanish)
- **SEND:** Be ready with questions for your doctor - Depression 📄 (Spanish)

P5. Potential complication: financial issues present

- **Goal:** Financial resources identified and sufficient to continue care(48)(49)📄
 - **Interventions:**
 - **ASSIST:** Identifying healthcare benefits coverage from payers
 - **ASSIST:** Identifying availability of and eligibility for alternative financial resources
 - **ASSIST:** Communicating information on patient's healthcare benefits coverage with treatment team
 - **ASSIST:** Appointment scheduling and completing paperwork to support applications for financial support
 - **EDUCATE:** Healthcare benefits and out-of-pocket expenses
 - **EDUCATE:** Resources for financial assistance:
 - Administration for Children & Families: www.acf.hhs.gov
 - Centers for Medicare and Medicaid Services (CMS): www.medicare.gov/drug-coverage-part-d
 - Medicine Assistance Tool: <https://medicineassistancetool.org>
 - NeedyMeds. Find help with the cost of medicine: www.needymeds.org
 - Individual pharmaceutical company websites
 - **COORDINATE:** Financial assistance referral; examples include:
 - Area Agency on Aging
 - Disease advocacy
 - Financial counseling or assistance
 - Food banks or other sources for nutritional support and financial assistance
 - Government agencies
 - Legal aid
 - Long-term disability application
 - Medicaid application
 - Medicare application
 - Pharmacy assistance programs
 - Short-term disability application
 - Supplemental Security Income (SSI) or disability benefits application
 - Veterans Affairs (VA) benefits application
 - **COORDINATE:** Social worker referral, as needed(50)
 - **COORDINATE:** Follow-up for evaluation of referral status and need for further assistance(2)(10)
 - **Goal:** Financial issues addressed(48)
 - **Interventions:**
 - **RECONCILE:** Benefits from multiple sources, if needed
 - **ASSIST:** Obtaining preauthorization from health plan provider, if appropriate
 - **EDUCATE:** Source and type of financial support, how to access, and if follow-up needed
 - **COORDINATE:** Follow-up for evaluation of use of support programs or ongoing need for further assistance(2)(10)

P6. Potential complication: self-care needs exceed current ability.

- **Goal:** Self-care needs safely met during and after care transition(4)(51)[N](#)
 - **Interventions:**
 - **RECONCILE:** Patient's self-care needs and ability with transition plan; considerations include:
 - Complexity of self-care needs
 - Patient's preferences for next level of care
 - Care recommendations from care team members
 - Available and able caregiver at next level of care
 - Financial resources and out-of-pocket expenses
 - **ASSIST:** Identifying discharge options that will meet patient's self-care needs
 - **EDUCATE:** Care team's recommendation for next level of care, including:
 - Self-care needs and goals for care at next level of care
 - Role and purpose of resources at next level of care
 - Anticipated capability of support needed at next level of care (eg, skilled, trained)
 - Estimated time (duration) needed for support at next level of care
 - **EDUCATE:** Providers of care available to patient for transition (eg, list of facilities or providers)
 - **EDUCATE:** Options for alternative levels of care that may also support patient self-care
 - **EDUCATE:** Potential complications with proceeding with discharge plan that does not cover patient's self-care needs
 - **EDUCATE:** Financial resources and out-of-pocket expenses for various care levels
 - **COORDINATE:** Social worker referral, as needed(50)
 - **COORDINATE:** Home care referral, if indicated
 - **COORDINATE:** Community services referral, as needed
 - **COORDINATE:** Communication between care team members, patient's caregivers, alternative level-of-care providers, and financial resources (health plan)
 - **COORDINATE:** Follow-up contact to evaluate patient's discharge preference, referral status, and need for further assistance

P7. Potential complication: caregiver needed to support patient's self-care

- **Goal:** Caregiver support available and provided for safe transition to next level of care(44)(52)[N](#)
 - **Interventions:**
 - **ASSIST:** Support system (eg, family, friends) identification, enlistment, and education
 - **ASSIST:** Deploying strategies to reduce caregiver burden, as needed (eg, healthcare benefits, support groups, family leave)(53)
 - **EDUCATE:** Patient's self-care needs and goals for care at next level of care
 - **EDUCATE:** When and how to get help for problems with patient or patient care
 - **EDUCATE:** Caregiver's anticipated role (eg, assisting with ADL, performing treatments, managing appointments)
 - **EDUCATE:** Risk to long-term caregiver for depression, anxiety, sleep disturbance, and burnout(54)(55)
 - **EDUCATE:** Caregiver support(53):
 - Identification of other caregivers
 - Consideration of adult day care or respite program
 - Self-care needs (eg, medical, behavior health, social support)
 - **COORDINATE:** Care team meeting, if needed
 - **COORDINATE:** Educational program referral for caregiver
 - **COORDINATE:** Social worker referral, as needed
 - **COORDINATE:** Community services referral, as needed
 - **COORDINATE:** Follow-up contact to review caregiver resource needs, referral status, and need for additional support
 - **SEND:** Caregiver tips  (Spanish)

P8. Medication information needed

- **Goal:** Medication need, use, and management evaluated and problems resolved(51)(56)[N](#)
 - **Interventions:**
 - **RECONCILE:** Medication list. See Medication Reconciliation Tool  SR for more information.[N](#)
 - **ASSIST:** Creation of current and reconciled medication list
 - **ASSIST:** Providing reconciled medication list with all providers. See Medication Reconciliation Tool  SR for more information.
 - **ASSIST:** Identifying resources to pay for medications, if needed
 - **ASSIST:** Identifying activities that promote medication adherence
 - **ASSIST:** Communication with treating provider when another provider (eg, emergency or specialty provider) prescribes new medication
 - **ASSIST:** Appointment scheduling with provider to evaluate treatment outcomes
 - **EDUCATE: Reason for taking medication(57):**
 - Expected effects if the medication is working, and expected time for effects to occur

- How long medication will likely need to be taken
- Risks of nonadherence
- **EDUCATE:** Information on all medications(56)(57)(58)(59):
 - Medication purpose, dose, effects/side effects, and considerations (eg, managing missed doses, interactions with food/other drugs)
 - Name of provider who ordered medication
 - Administration techniques
 - Side effects, related management, and when to contact provider
 - **Get emergency help:** Adverse or allergic reactions (eg, wheezing, chest tightness, fever, itching, bad cough, blue skin color, hemorrhage, seizures, swelling of face, lips, tongue, or throat)
 - Common side effects include gastrointestinal changes, rash, dizziness, and headache.
 - Do not stop long-term medications abruptly due to risk of withdrawal symptoms.
 - Medication safety (eg, proper storage and disposal)(60)
 - Monitoring (eg, vital signs, laboratory checks) needed while taking medication
 - Talk to provider before taking any over-the-counter medications, vitamins, minerals, or herbal or dietary supplements.
 - Carry a list of medications at all times, particularly when visiting healthcare providers.(59)
- **EDUCATE:** Ways to improve following medication routine (eg, pillboxes, alarm reminders)
- **EDUCATE:** Pharmacy assistance programs, such as:
 - Centers for Medicare and Medicaid Services (CMS): www.medicare.gov/drug-coverage-part-d
 - Medicine Assistance Tool: <https://medicineassistancetool.org>
 - NeedyMeds: www.needymeds.org
 - Individual company websites
- **COORDINATE:** Follow-up on the medication plan:
 - Contact with prescribing providers to verify medication orders
 - Contact with pharmacy to confirm patient receipt of medication
 - Contact with patient to evaluate adherence to medication plan and need for more assistance
 - Laboratory testing, if indicated(61)
- **COORDINATE:** Home care referral, if indicated
- **COORDINATE:** Follow-up to evaluate adherence or problems with medication, provide additional education and support, and identify need for further assistance(2)(8)
- **RECONCILE:** Communicate reconciled medication list to all providers.
- **SEND:** Managing your medicines (Illustrated)  (Spanish)
- **Goal:** Medication safety understood and medication safety implemented(56)
 - **Interventions:**
 - **RECONCILE:** Medication list. See Medication Reconciliation Tool  SR for more information.
 - **ASSIST:** Communication with treating provider when another provider (eg, emergency or specialty provider) prescribes new medication
 - **ASSIST:** Communication with primary provider about prescription issues:
 - Polypharmacy
 - Food-drug and drug-drug issues
 - Adherence problems
 - **EDUCATE:** Medication safety instructions(62)(63):
 - Discard outdated medication.
 - Do not share prescription medication.
 - Store out of reach of children and in childproof containers.
 - Fill all prescriptions at the same pharmacy.
 - Store properly (in original containers and with other needs such as refrigeration or protection from sunlight).
 - Check with provider before taking over-the-counter medications, vitamins, minerals, or herbal or dietary supplements.
 - How to dispose of unused or expired medication
 - **EDUCATE:** Administration techniques
 - **EDUCATE:** De-prescribing or cessation of medications safely (eg, pharmacy-supported, provider monitoring planned) (64)
 - **COORDINATE:** Home care referral, if indicated
 - **COORDINATE:** Follow-up to evaluate adherence or problems with medication, provide additional education and support, evaluate referral status, and identify need for further assistance(2)(8)
 - **RECONCILE:** Communicate reconciled medication list to all providers.
 - **SEND:** Managing your medicines (Illustrated)  (Spanish)

P9. Potential complication: transportation insufficient for patient to access care

- **Goal:** Transportation issues addressed(4)(65) 

- **Interventions:**

- **ASSIST:** Identifying transportation resources and accessing as needed, such as:
 - Family, friends, neighbors, community group members
 - Public transportation (bus, train, access vans)
 - Private transportation (taxis, car service)
 - Pharmacy, equipment, or supply delivery services
- **ASSIST:** Completing transportation assistance forms, as needed
- **EDUCATE:** Local transportation services available and how to access them
- **EDUCATE:** Cost associated with available transportation
- **EDUCATE:** When and how to get help if transportation continues to be a problem
- **COORDINATE:** Communication between care team members on how patient is transported to and from service visits
- **COORDINATE:** Community services referral, as needed
- **COORDINATE:** Social worker referral, as needed(50)
- **COORDINATE:** Follow-up contact to evaluate referral status, use of transportation, and address any further transportation needs

P10. Depression medication management needed

- **Goal:** Medication information understood(59)

- **Interventions:**

- **ASSIST:** Identifying patient's goal priorities, willingness/ability to achieve goals, and preferences for communication and follow-up
- **ASSIST:** Goal setting that is specific, measurable, achievable, relevant, and with target completion time (eg, short/long-term, prioritized)
- **ASSIST:** Appointment scheduling with provider for discussion on appropriate medication
- **ASSIST:** Providing reconciled medication list to all providers. See Medication Reconciliation Tool  SR for more information.
- **EDUCATE: Purpose:** Importance of understanding what each medication is for, how it should be taken, and what to do if dose is missed
- **EDUCATE:** Medication indications, side effects, and considerations frequently change. See a drug reference for specific information not listed here.
- **EDUCATE:** Administration techniques:
 - Route (eg, oral, intramuscular (IM), intravenous (IV))
 - With or without foods or liquids (eg, milk)
 - Timing of administration
 - Appropriate measuring device (eg, liquid medications)
 - Proper storage
- **EDUCATE: Get emergency help:** Adverse or allergic reactions (eg, wheezing, chest tightness, fever, itching, bad cough, blue skin color, hemorrhage, seizures, swelling of face, lips, tongue, or throat)
- **EDUCATE: Side effects:** Management, and when to contact provider
 - Common side effects include gastrointestinal changes, dry mouth, fatigue, insomnia, rash, dizziness, and headache.
- **EDUCATE: Considerations:**
 - Discuss medication with provider if pregnant, planning a pregnancy, or breast-feeding.
 - Comorbidities may be contraindications to the use of a drug (eg, liver disease, diabetes).
 - Health checks, diary, and action plans (eg, blood pressure, weight, heart rate)
 - Safety measures (eg, sun protection, driving, or operating machinery)
 - Contact provider or pharmacist prior to taking any other over-the-counter or prescribed medication.
- **COORDINATE:** Clinical monitoring appropriate to medication use (eg, laboratory tests)
- **COORDINATE:** Communication between patient and pharmacy (for obtaining prescriptions, setting up home medication delivery, review for drug-drug interactions) 
- **COORDINATE:** Communication between care team members and medication list reconciliation
- **COORDINATE:** Follow-up to evaluate adherence or problems with medication, provide additional education and support, and identify need for further assistance(2)(8)
- **COORDINATE:** Laboratory testing, if indicated
- **COORDINATE:** Outpatient imaging, tests, or procedures, if indicated
- **COORDINATE:** Therapeutic drug level monitoring
- **SEND:** Managing your medicines (Illustrated)  (Spanish)

P11. Atypical antidepressant medication management needed

- **Goal:** Atypical antidepressant medication initiated and maintained

- **Interventions:**

- **ASSIST:** Appointment scheduling with prescribing provider for medication evaluation and refills

- **EDUCATE: Purpose:** Atypical antidepressant medications are used to decrease the frequency and intensity of depressive episodes.(18)
- **EDUCATE: Side effects:** Dry mouth, weight gain, insomnia(66)
 - Trazodone or mirtazapine may cause somnolence.
 - Trazodone may cause priapism.(14)
- **EDUCATE: Considerations:** Avoid bupropion in patients with seizure disorders.(67)
- **SEND:** Antidepressant medicines  (Spanish)

P12. Monoamine oxidase inhibitor (MAOI) medication management needed

- **Goal:** Monoamine oxidase inhibitor (MAOI) medication initiated and maintained(18)(68)
 - **Interventions:**
 - **ASSIST:** Appointment scheduling with prescribing provider for medication evaluation and refills
 - **EDUCATE: Purpose:** MAOIs are antidepressant medications that decrease the frequency and intensity of depressive episodes.(5)
 - **EDUCATE: Call provider immediately:** Serotonin syndrome and hypertensive crisis signs:
 - Serotonin syndrome (eg, tachycardia, diarrhea, hyperthermia, tremor, clonus, agitation, delirium)
 - Hypertensive crisis (eg, blood pressure of 180/120 mm Hg or higher; severe headache; motor or sensory deficits; somnolence; seizure; unexplained chest, abdominal, or back pain)
 - **EDUCATE: Side effects:** Jaundice, hallucinations, fainting, hyperexcitability, confusion, fever, disorientation, decreased sexual function
 - **EDUCATE: Considerations:**
 - Do not take any new medications (including over-the-counter medications) unless cleared with prescribing provider.
 - Life-threatening drug interactions may occur.[E]
 - Avoid foods with amino acid tyramine, as hypertensive crisis can occur if tyramine consumed with MAOIs. Tyramine is found in foods such as avocados, bananas, yogurt, aged cheese, yeast, and aged meat.(69)
 - Prevent sudden drop in blood pressure and related falls by rising slowly from sitting or lying down.
 - **SEND:** Antidepressant medicines  (Spanish)
 - **SEND:** How to read food labels (Illustrated)  (Spanish)

P13. Selective serotonin reuptake inhibitors (SSRIs) or serotonin-norepinephrine reuptake inhibitors (SNRIs) medication management needed

- **Goal:** Selective serotonin reuptake inhibitors (SSRIs) or serotonin-norepinephrine reuptake inhibitors (SNRIs) medication initiated and maintained
 - **Interventions:**
 - **ASSIST:** Appointment scheduling with prescribing provider for medication evaluation and refills
 - **EDUCATE: Purpose:** SSRIs and SNRIs are antidepressant medications.(5)
 - Prevent anxiety symptoms, reduce relapse risk, and have less safety risks compared to benzodiazepines(70)
 - Effects may take several weeks.(67)
 - **EDUCATE: Call provider immediately:** If taking SSRIs and MAOIs or TCAs and serotonin syndrome signs present(71)(72):
 - Serotonin syndrome (eg, tachycardia, diarrhea, hyperthermia, tremor, clonus, agitation, delirium)
 - **EDUCATE: Side effects:** Agitation, headache, postural hypotension, sexual dysfunction (eg, abnormal ejaculation, impotence, loss of orgasm)(66)
 - SNRIs have fewer sexual side effects than SSRIs.
 - **EDUCATE: Considerations(66)(73):**
 - Take dose early in the day to help avoid insomnia at night.
 - Use cautiously with cardiac conditions due to possible elevation in blood pressure with SNRIs.
 - **SEND:** Help with sleeping or insomnia  (Spanish)
 - **SEND:** Antidepressant medicines  (Spanish)

P14. Tricyclic antidepressants (TCAs) medication management needed

- **Goal:** Tricyclic antidepressants (TCAs) medication initiated and maintained(68)(73)
 - **Interventions:**
 - **ASSIST:** Appointment scheduling with prescribing provider for medication evaluation and refills
 - **EDUCATE: Purpose:** TCAs are antidepressant medications that decrease the frequency and intensity of depressive episodes.
 - TCAs decrease the frequency and intensity of panic attacks.
 - **EDUCATE: Call provider immediately:** If taking TCAs and MAOIs or SSRIs and serotonin syndrome signs present(71)(72):
 - Serotonin syndrome (eg, tachycardia, diarrhea, hyperthermia, tremor, clonus, agitation, delirium)

- **EDUCATE: Side effects:** Blurred vision, urinary retention, orthostatic hypotension, reduction in seizure threshold, tachycardia, light sensitivity, weight gain
- **EDUCATE: Considerations:** Avoid in patients with angle-closure glaucoma or urinary retention.

P15. Depression medication evaluation needed

- **Goal:** Depression medication evaluation completed and regimen established
 - **Interventions:**
 - **RECONCILE:** Medication list. See Medication Reconciliation Tool [SR](#) for more information.
 - **ASSIST:** Appointment scheduling with prescribing provider for evaluation of condition status
 - **EDUCATE:** When and how to get help for worsening depression problems
 - **EDUCATE:** Medication management needs common for depression
 - **COORDINATE:** Follow-up after provider contact for evaluation of medication changes
 - **RECONCILE:** Communicate reconciled medication list to all providers.(74)(75)

P16. Medication changes present

- **Goal:** Medication changes understood and medication taken as ordered
 - **Interventions:**
 - **ASSIST:** Identifying patient's goal priorities, willingness/ability to achieve goals, and preferences for communication and follow-up
 - **ASSIST:** Goal setting that is specific, measurable, achievable, relevant, and with target completion time (eg, short/long-term, prioritized)
 - **ASSIST:** Appointment scheduling with provider for evaluation of medication problems (clarification of dose or regimen)
 - **ASSIST:** Creation of current and reconciled medication list
 - **ASSIST:** Identifying activities that promote medication adherence
 - **ASSIST:** Communication with treating provider when another provider (eg, emergency or specialty provider) prescribes new medication
 - **EDUCATE: Reason for taking medication(57):**
 - Expected effects if the medication is working, and expected time for effects to occur
 - How long medication will likely need to be taken
 - Risks of nonadherence
 - **EDUCATE: Information on all medications(56)(57)(58)(59):**
 - Medication purpose, dose, effects/side effects, and considerations (eg, managing missed doses, interactions with food/other drugs)
 - Name of provider who ordered medication
 - Administration techniques
 - Side effects, related management, and when to contact provider
 - **Get emergency help:** Adverse or allergic reactions (eg, wheezing, chest tightness, fever, itching, bad cough, blue skin color, hemorrhage, seizures, swelling of face, lips, tongue, or throat)
 - Common side effects include gastrointestinal changes, rash, dizziness, and headache.
 - Do not stop long-term medications abruptly due to risk of withdrawal symptoms.
 - Medication safety (eg, proper storage and disposal)(60)
 - Monitoring (eg, vital signs, laboratory checks) needed while taking medication
 - Talk to provider before taking any over-the-counter medications, vitamins, minerals, or herbal or dietary supplements.
 - Carry a list of medications at all times, particularly when visiting healthcare providers.
 - **COORDINATE:** Home care referral, if indicated
 - **COORDINATE:** Communication between care team members and medication list reconciliation.(56)(74) See Medication Reconciliation Tool [SR](#) for more information.[N](#)
 - **COORDINATE:** Communication between patient and pharmacy (for obtaining prescriptions, setting up home medication delivery, review for drug-drug interactions)(76)[N](#)
 - **COORDINATE:** Follow-up contact for evaluation of medication understanding, referral status, and need for further assistance(2)(8)
 - **SEND:** Managing your medicines (Illustrated)  (Spanish)

P17. Medication refills needed

- **Goal:** Medications refilled
 - **Interventions:**
 - **RECONCILE:** Medication list. See Medication Reconciliation Tool [SR](#) for more information.
 - **ASSIST:** Appointment scheduling with prescribing provider for medication evaluation and refills
 - **ASSIST:** Obtaining refills in lieu of an appointment
 - **ASSIST:** Identifying resources to pay for medications, if needed

- **ASSIST:** Working with pharmacy for filling prescriptions (eg, changing pills to liquids, identifying financial support, delivery)
- **COORDINATE:** Prescription refill support (eg, pharmacy programs) referral
- **COORDINATE:** Social worker referral, as needed(18)
- **EDUCATE:** Refill medications prior to running out (1 to 2 weeks before).
- **EDUCATE:** Pharmacy assistance programs, such as:
 - Centers for Medicare and Medicaid Services (CMS): www.medicare.gov/drug-coverage-part-d
 - Medicine Assistance Tool: <https://medicineassistancetool.org>
 - NeedyMeds: www.needymeds.org
 - Individual company websites
- **COORDINATE:** Follow-up to evaluate continued need for refills, referral status, and need for further assistance(2)(8)
- **RECONCILE:** Communicate reconciled medication list to all providers.(74)(75)
- **SEND:** Managing your medicines (Illustrated)  (Spanish)

P18. Laboratory work coordination needed

- **Goal:** Laboratory work completed(51) 
- **Interventions:**
 - **RECONCILE:** Laboratory work needed from all providers
 - **COORDINATE:** Laboratory testing, if indicated
 - **ASSIST:** Identifying laboratory location and scheduling appointment, if needed
 - **ASSIST:** Identifying transportation resources and accessing as needed
 - **EDUCATE:** Purpose and importance of getting laboratory work
 - **EDUCATE:** Follow-up with provider for laboratory results
 - **COORDINATE:** Follow-up contact with provider after laboratory work done to evaluate patient's status
 - **COORDINATE:** Communicate laboratory results to all relevant providers.
 - **COORDINATE:** Follow-up to review laboratory results and follow-up care needs, and to evaluate referral status or need for further assistance(2)(8)
 - **SEND:** Reminder on laboratory work due date, location, and provider follow-up

P19. Potential complication with transition to post-acute care facility

- **Goal:** Potential complication with transition to post-acute care facility resolved(4)(51) 
- **Interventions:**
 - **RECONCILE:** Transition care plan (eg, preferences, abilities, resources)(77)
 - **COORDINATE:** Care team meeting, if needed(6)(7)
 - **ASSIST:** Identifying 3 facilities within reconciled level-of-care options:
 - Identifying facilities in or out of health plan network, if indicated
 - Site visit to facilities by family or healthcare proxy
 - **ASSIST:** Completion of admission paperwork by patient or designee
 - **COORDINATE:** Preauthorization with health plan(s), if indicated
 - **COORDINATE:** Post-acute care facility pre-transition evaluation, if indicated
 - **EDUCATE:** Level-of-care options meeting care team recommendations and within payment considerations
 - **EDUCATE:** Benefits for this level of care and any out-of-pocket expenses that may be patient's responsibility (eg, co-pays or deductibles)
 - **EDUCATE:** Post-acute care facility information (name, contact, and transfer date)
 - **EDUCATE:** Goals and expectations once transition complete
 - **EDUCATE:** Transition of care plan (including estimated timing of events)
 - **SEND:** Medical records to accepting facility:
 - History and physical
 - Transition of care summary
 - Last set of laboratory results
 - Discharge orders (including treatments, laboratory tests, medications)
 - Opioid prescription signed hard copy, if indicated
 - Results of tests or procedures
 - Information on consulting provider(s) and when follow-up is needed
 - Advance care planning information
 - **SEND:** Transition summary to primary care provider
 - **COORDINATE:** Confirmation of facility acceptance, confirming:
 - Bed availability
 - Admission date and time
 - Transport of patient to facility
 - Medical record and discharge orders received
 - **COORDINATE:** Disposition of patient's personal items
 - **COORDINATE:** Social worker referral, as needed(50)

- **COORDINATE:** Follow-up contact with facility or provider, if needed
- **COORDINATE:** Follow-up contact to evaluate for transition problems, referral status, and need for further assistance(2)(10)
- **RECONCILE:** Timing of care transition and self-care plan (eg, medication available, provider assessment)
- **RECONCILE:** Discharge and admission orders
- **SEND:** When you are going to a recovery facility for care 🇪🇸 (Spanish)
- **Goal:** Alternative care transition plan initiated if complication with post-acute facility transition cannot be resolved(51)(78)
 - **Interventions:**
 - **RECONCILE:** Patient (or designee) preferences, care team recommendations, and payment options for alternative care transition plan
 - **ASSIST:** Identifying alternative transition plans
 - **ASSIST:** Identifying resources to support alternative care transition plan
 - **EDUCATE:** Alternative levels of care besides post-acute care facility that will meet patient's needs
 - **EDUCATE:** Why post-acute care facility may be insufficient to meet patient's needs
 - **COORDINATE:** Care team meeting to discuss alternative care transition plan options(6)(7)
 - **COORDINATE:** Outpatient imaging, tests, or procedures, if indicated
 - **COORDINATE:** Home care referral, if indicated
 - **COORDINATE:** Community services referral, as needed
 - **COORDINATE:** Inpatient rehabilitation facility referral, if indicated
 - **COORDINATE:** End-of-life or palliative care referral, if indicated
 - **COORDINATE:** Assisted living referral, if indicated
 - **COORDINATE:** Long-term care services and support referral, as needed
 - **COORDINATE:** Follow-up contact to evaluate alternative discharge options, referral status, and need for further assistance(2)(10)

Evidence Summary

Quality standards: Standards indicate that programs should ensure that during care transitions the care plan is re-evaluated, modified, and communicated to appropriate care team members (eg, providers, caregivers) as needed to support the care transition.(9)(79)(80)(81)(82)

Quality standards: The American Case Management Association's Transitions of Care Standards indicate that patients' health risks should be assessed at each care episode or transition, a care management plan should be established and communicated across the care settings, and an evaluation should be done of the patient/family/caregiver's understanding of health status and self-care.(83)

Quality standards: The American Case Management Association's Transitions of Care Standards indicate that processes should be in place to identify patients at risk for care transition problems, patients' health risks should be assessed at each care episode or transition, and a care management plan should be established and communicated across care settings.(83)

Transition consideration: Several studies demonstrated a reduction in 30-day hospital readmissions when the patient follow-up was within 7 days of hospital discharge.(84)(85)(86) Earlier follow-up (eg, within 48 hours after hospital discharge) may also help reduce hospital readmissions for high-risk or vulnerable patients.(2) Follow-up appointment attendance rates were higher when appointments were scheduled before hospital discharge.(87)(88)

Quality standards: Standards indicate that programs should develop and communicate an individualized case management plan with the patient, facilitate and follow up with referrals, and assess the patient's progress with the care plan. Programs should monitor and communicate the patient's progress with the care plan, modifying the plan as needed, and continue to identify patient preferences, barriers, and adherence concerns. Program content should include an individualized self-care plan that includes patient preferences, prioritized goals, interventions with information on whom to contact in urgent situations, and scheduled follow-up. Programs should facilitate communication and collaboration with practitioners in a timely manner.(79)(81)(82)

Quality standards: The Case Management Society of America's Standards of Practice for Case Management principles include facilitation of informed decision making, counseling, and health education with patient self-care that is responsive to the patient's preferences (eg, culture, values, needs).(79)

Quality standards: The Case Management Society of America's Case Management Adherence Guide includes guidance for improving adherence via patient engagement.(89)

National guidelines: The 2018 Physical Activity Guidelines Advisory Committee advises that adults with chronic conditions or disabilities, who are able, should follow the key activity guidelines for adults. The guidelines recommend that adults should do at least 150 to 300 minutes a week of moderate-intensity or 75 to 150 minutes a week of vigorous-intensity aerobic physical activity, or an equivalent combination of moderate-intensity and vigorous-intensity aerobic activity. They should also do muscle-strengthening

activities on 2 or more days a week. Older adults should do multicomponent physical activity that includes balance training as well as aerobic and muscle-strengthening activities.(90)

Evidence: A retrospective cohort study of 17,629 patients found a significant reduction in healthcare utilization, hospital readmissions, and deaths when patients were discharged with home healthcare vs discharged to home without home healthcare.(91)

Evidence: A retrospective cohort study based on 1151 Medicare accountable care organization (ACO) patients suggests utilizing a transitional care planning approach to prioritize discharging patients to the least restrictive next level of care, as patients discharged to skilled nursing facilities were almost 5 times more likely to be readmitted to the hospital at 30 days compared to patients discharged to the home healthcare setting.(92)

Quality standards: Standards indicate that programs should have a process for identifying barriers to patients meeting goals or adhering to the case management plan and for following up with patients on the status of referrals.(79)

Quality standards: Standards indicate that programs should have a process for identifying barriers to patients meeting goals or adhering to the case management plan and for following up with patients on the status of referrals. Programs should ensure that during care transitions the care plan is re-evaluated, modified, and communicated to appropriate care team members (eg, providers, caregivers) as needed to support the care transition.(79)

Quality standards: Standards indicate that caregiver resources should be evaluated. Programs should ensure that during care transitions the care plan is re-evaluated, modified, and communicated to appropriate care team members (eg, providers, caregivers) as needed to support the care transition.(79)

Quality standards: Standards indicate that programs should have a process for identifying barriers to patients meeting goals or adhering to the case management plan and for following up with patients on the status of referrals. During transitions, medications should be documented with the patient and designee, and discrepancies should be communicated to the appropriate provider.(79)

Outcomes: Several studies support the role of the pharmacist in medication reconciliation to identify and resolve pharmacy-related problems and improve outcomes (eg, prevention of medication-related adverse events, discontinuation of inappropriate medication).(93)

Evidence: A prospective cohort study of 2655 hospitalized patients after discharge found that 24% of hospital medication changes were not followed, 30% of changed doses were filled with the incorrect dose, 27% of new medications were not filled, and 12% of discontinued medications were filled. Risk factors for medication change problems included increased out-of-pocket costs, not having medications filled before hospitalization, and discharge to a long-term care facility.(94)

Quality standards: Standards indicate that programs should collaborate on transitions with the discharge team, reassessing the appropriateness as needed; communicate transition plan progress with the patient and designee; and communicate care plan information with the receiving setting within a designated time frame. Programs should develop and communicate an individualized case management plan with the patient, facilitate and follow up with referrals, and assess the patient's progress with the care plan. Programs should ensure that during care transitions the care plan is re-evaluated, modified, and communicated to appropriate care team members (eg, providers, caregivers) as needed to support the care transition.(79)

References

1. Psychiatric guidelines. In: Cash JC, editor. Family Practice Guidelines. 6th ed. Springer Publishing; 2023:888-923. [Context Link 1, 2, 3, 4, 5]
2. Care coordination and care transitions to support adherence. In: Perez R, Rogers S, Prince M, Fraser K, editors. Case Management Adherence Guidelines. 2020 ed. Case Management Society of America; 2022:43-56. [Context Link 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16]
3. Morkisch N, et al. Components of the transitional care model (TCM) to reduce readmission in geriatric patients: a systematic review. BMC Geriatrics 2020;20(1):345. DOI: 10.1186/s12877-020-01747-w. [Context Link 1] View abstract...
4. Transitional planning: understanding levels and transitions of care. In: Powell SK, Tahan H, editors. Case Management a Practical Guide for Education and Practice. 4th ed. Philadelphia, PA: Wolters Kluwer, Lippincott, Williams & Wilkins; 2019:156-211. [Context Link 1, 2, 3, 4]
5. Reus VI. Psychiatric disorders. In: Loscalzo J, Fauci A, Kasper D, Hauser S, Longo D, Jameson JL, editors. Harrison's Principles of Internal Medicine. 21st ed. McGraw Hill Education; 2022:3540-3556. [Context Link 1, 2, 3, 4, 5, 6]
6. Case management standards and professional organizations. In: Powell SK, Tahan H, editors. Case Management a Practical Guide for Education and Practice. 4th ed. Philadelphia, PA: Wolters Kluwer, Lippincott, Williams & Wilkins; 2019:314-354. [Context Link 1, 2, 3]
7. Ralston JD, Wagner EH. Comprehensive chronic disease management. In: Goldman L, Schafer AI, editors. Goldman-Cecil Medicine. 26th ed. Elsevier; 2020:42-45.e2. [Context Link 1, 2, 3, 4]
8. Monitoring and evaluation. In: Gillingham DC, editor. Foundations of Case Management. Blue Bayou Press; 2021:51-52. [Context Link 1, 2, 3, 4, 5, 6, 7, 8, 9, 10]
9. Transitions of care. In: Gillingham DC, editor. Foundations of Case Management. Blue Bayou Press; 2021:137-144. [Context Link 1, 2, 3, 4, 5]
10. Joo JY, Liu MF. The experience of chronic illness transitional care: a qualitative systematic review. Clinical Nursing Research 2022;31(2):163-173. DOI: 10.1177/10547738211056166. [Context Link 1, 2, 3, 4, 5, 6, 7] View abstract...

11. Assessing the adult patient through application of the integrated case management complexity assessment grid. In: Fraser K, Perez R, Latour C, editors. *CMSA's Integrated Case Management*. New York, NY: Springer Publishing Company LLC; 2018:51-82. [Context Link 1, 2]
12. Chiappetta L. Mood disorders: depression. In: Fosbre CD, Varcarolis EM, Chiappetta L, editors. *Varcarolis' Essentials of Psychiatric-Mental Health Nursing*. 5th ed. Elsevier; 2023:207-234. [Context Link 1, 2, 3, 4]
13. Depression. NIH Publication No. 21-MH-8079 [Internet] National Institute of Mental Health. 2021 Accessed at: <https://www.nimh.nih.gov/health/publications/depression/>. [accessed 2023 Aug 10] [Context Link 1]
14. Depressive disorders. In: Morgan KI, Townsend MC, editors. *Davis Advantage for Psychiatric Mental Health Nursing*. 10th ed. Philadelphia, PA: F.A. Davis; 2021:479-518. [Context Link 1, 2, 3, 4]
15. Major depression. In: Swearingen PL, Wright JD, editors. *All-in-One Nursing Care Planning Resource*. 5th ed. St. Louis, MO: Elsevier; 2019:744-752. [Context Link 1]
16. Steele D. Depressive disorders. In: Steele D, editor. *Keltner's Psychiatric Nursing*. 9th ed. Elsevier; 2023:279-300. [Context Link 1]
17. Fosbre CD. Anger, aggression, and violence. In: Fosbre CD, Varcarolis EM, Chiappetta L, editors. *Varcarolis' Essentials of Psychiatric-Mental Health Nursing*. 5th ed. Elsevier; 2023:388-401. [Context Link 1]
18. Mood disorders and suicide. In: Videbeck SL, editor. *Psychiatric-Mental Health Nursing*. 9th ed. Wolters Kluwer; 2023:284-327. [Context Link 1, 2, 3, 4, 5, 6, 7, 8, 9, 10]
19. Neuropsychiatric disorders. In: Escott-Stump S, editor. *Nutrition and Diagnosis-Related Care*. 9th ed. Wolters Kluwer; 2022:276-378. [Context Link 1]
20. Greene RD, Cook A, Nowaskie D, Wang S. Neurological changes and depression 2019 update. *Clinics in Geriatric Medicine* 2020;36(2):297-313. DOI: 36. [Context Link 1, 2] View abstract...
21. Physical Activity and Sedentary Behaviour. WHO Guideline [Internet] World Health Organization. 2020 Nov Accessed at: <https://www.who.int/publications/i/item/9789240015128>. [accessed 2023 Aug 14] [Context Link 1]
22. Depression in Adults: Treatment and Management. NICE Guidance NG222 [Internet] National Institute for Health and Care Excellence. 2022 Jun Accessed at: <https://www.nice.org.uk/guidance>. [created 2022; accessed 2023 Aug 08] [Context Link 1]
23. Hides L, Quinn C, Stoyanov S, Kavanagh D, Baker A. Psychological interventions for co-occurring depression and substance use disorders. *Cochrane Database of Systematic Reviews* 2019, Issue 11. Art. No.: CD009501. DOI: 10.1002/14651858.CD009501.pub2. [Context Link 1] View abstract...
24. Precin P. Mood disorders. In: Atchison BJ, Dirette DP, editors. *Conditions in Occupational Therapy: Effect on Occupational Performance*. 6th ed. Wolters Kluwer; 2023:167-180. [Context Link 1, 2]
25. Mauk KL. Post-acute care. In: Larsen PD, editor. *Lubkin's Chronic Illness: Impact and Intervention*. 11th ed. Jones & Bartlett Learning; 2022:477-520. [Context Link 1, 2]
26. Miller TR, et al. Incidence and lethality of suicidal overdoses by drug class. *JAMA Network Open* 2020;3(3):e200607. DOI: 10.1001/jamanetworkopen.2020.0607. [Context Link 1] View abstract...
27. Okolie C, et al. Means restriction for the prevention of suicide by jumping. *Cochrane Database of Systematic Reviews* 2020, Issue 2. Art. No.: CD013543. DOI: 10.1002/14651858.CD013543. [Context Link 1] View abstract...
28. Shaffer RB. Coronary artery disease and acute coronary syndrome. In: Harding MM, Kwong J, Hagler D, Reinisch C, editors. *Lewis's Medical-Surgical Nursing: Assessment and Management of Clinical Problems*. 12th ed. St. Louis, MO: Mosby; 2023:819-858. [Context Link 1]
29. Bussard M. Stroke. In: Harding MM, Kwong J, Hagler D, Reinisch C, editors. *Lewis's Medical-Surgical Nursing: Assessment and Management of Clinical Problems*. 12th ed. St. Louis, MO: Mosby; 2023:1515-1538. [Context Link 1]
30. Fosbre CD. Suicidal thoughts and behaviors. In: Fosbre CD, Varcarolis EM, Chiappetta L, editors. *Varcarolis' Essentials of Psychiatric-Mental Health Nursing*. 5th ed. Elsevier; 2023:377-387. [Context Link 1]
31. Harding MM. Assessment and management: visual problems. In: Harding MM, Kwong J, Hagler D, Reinisch C, editors. *Lewis's Medical-Surgical Nursing: Assessment and Management of Clinical Problems*. 12th ed. St. Louis, MO: Mosby; 2023:407-435. [Context Link 1]
32. Neurologic care. In: Lynn P, editor. *Taylor's Clinical Nursing Skills*. 6th ed. Wolters Kluwer; 2023:1079-1113. [Context Link 1]
33. McQuaid JR, et al. The management of major depressive disorder: synopsis of the 2022 U.S. Department of Veterans Affairs and U.S. Department of Defense Clinical Practice Guideline. *Annals of Internal Medicine* 2022;175(10):1440-1451. DOI: 10.7326/M22-1603. [Context Link 1] View abstract...
34. Christensen LF, Moller AM, Hansen JP, Nielsen CT, Gildberg FA. Patients' and providers' experiences with video consultations used in the treatment of older patients with unipolar depression: A systematic review. *Journal of Psychiatric and Mental Health Nursing* 2020;27(3):258-271. DOI: 10.1111/jpm.12574. [Context Link 1] View abstract...
35. Hudon C, et al. Case management in primary care for frequent users of health care services: a realist synthesis. *Annals of Family Medicine* 2020;18(3):218-226. DOI: 10.1370/afm.2499. [Context Link 1, 2] View abstract...
36. Bartels SJ, DiMilia PR, Fortuna KL, Naslund JA. Integrated care for older adults with serious mental illness and medical comorbidity: evidence-based models and future research directions. *Clinics in Geriatric Medicine* 2020;36(2):341-352. DOI: 10.1016/j.cger.2019.11.012. [Context Link 1] View abstract...
37. Cohen DJ, Davis MM, Hall JD, Gilchrist EC, Miller BF. A Guidebook of Professional Practices for Behavioral Health and Primary Care Integration. The Academy Integrating Behavioral Health and Primary Care AHRQ Publication No. 14-0070-1-EF [Internet] Agency for Healthcare Research and Quality. 2015 Mar Accessed at: <https://www.ahrq.gov/>. [accessed 2023 Aug 15] [Context Link 1]
38. Motivational interviewing and shared decision making for the medically complex patient and family caregiver. In: Fraser K, Perez R, Latour C, editors. *CMSA's Integrated Case Management*. New York, NY: Springer Publishing Company LLC; 2018:149-176. [Context Link 1]

39. Anger, hostility, and aggression. In: Videbeck SL, editor. *Psychiatric-Mental Health Nursing*. 9th ed. Wolters Kluwer; 2023:175-187. [Context Link 1, 2]
40. Abuse and violence. In: Videbeck SL, editor. *Psychiatric-Mental Health Nursing*. 9th ed. Wolters Kluwer; 2023:188-204. [Context Link 1]
41. Assessment. In: Videbeck SL, editor. *Psychiatric-Mental Health Nursing*. 9th ed. Wolters Kluwer; 2023:130-144. [Context Link 1, 2]
42. Client's response to illness. In: Videbeck SL, editor. *Psychiatric-Mental Health Nursing*. 9th ed. Wolters Kluwer; 2023:118-129. [Context Link 1]
43. Greenberg SA. Chronic illness and older adults. In: Harding MM, Kwong J, Hagler D, Reinisch C, editors. *Lewis's Medical-Surgical Nursing: Assessment and Management of Clinical Problems*. 12th ed. St. Louis, MO: Mosby; 2023:63-80. [Context Link 1]
44. Martinez-Kratz M. Guide to planning care. In: Makic MB, Martinez-Kratz MR, editors. *Ackley and Ladwig's Nursing Diagnosis Handbook*. 13th ed. Elsevier; 2023:125-1051. [Context Link 1, 2]
45. Anxiety and anxiety disorders. In: Videbeck SL, editor. *Psychiatric-Mental Health Nursing*. 9th ed. Wolters Kluwer; 2023:222-240. [Context Link 1]
46. LeBlanc RG. Self-management. In: Larsen PD, editor. *Lubkin's Chronic Illness: Impact and Intervention*. 11th ed. Jones & Bartlett Learning; 2022:147-172. [Context Link 1]
47. Sochacki S. Adherence. In: Larsen PD, editor. *Lubkin's Chronic Illness: Impact and Intervention*. 11th ed. Jones & Bartlett Learning; 2022:269-306. [Context Link 1]
48. Addressing social determinants: clinical versus nonclinical issues. In: Fraser K, Perez R, Latour C, editors. *CMSA's Integrated Case Management*. New York, NY: Springer Publishing Company LLC; 2018:139-148. [Context Link 1, 2]
49. Rector C. Structure and economics of community/public health services. In: Rector C, Stanley MJ, editors. *Community and Public Health Nursing Promoting the Public's Health*. 10th ed. Wolters Kluwer; 2022:121-169. [Context Link 1]
50. Liebbeit D, Rutkowski R, Arbaje AI, Fields B, Werner NE. A scoping review of interventions for older adults transitioning from hospital to home. *Journal of the American Geriatrics Society* 2021;69(10):2950-2962. DOI: 10.1111/jgs.17323. [Context Link 1, 2, 3, 4] View abstract...
51. Interdisciplinary care teams fostering successful transitions of care. In: Fraser K, Perez R, Latour C, editors. *CMSA's Integrated Case Management*. New York, NY: Springer Publishing Company LLC; 2018:177-202. [Context Link 1, 2, 3, 4, 5]
52. Morris BC. Patient and caregiver teaching. In: Harding MM, Kwong J, Hagler D, Reinisch C, editors. *Lewis's Medical-Surgical Nursing: Assessment and Management of Clinical Problems*. 12th ed. St. Louis, MO: Mosby; 2023:49-62. [Context Link 1]
53. Messecar DC. Family caregiving. In: Boltz M, Capezuti EA, Zwicker D, Fulmer T, editors. *Evidence-Based Geriatric Nursing Protocols for Best Practice*. 6th ed. New York, NY: Springer; 2021:191-222. [Context Link 1, 2]
54. Lutz BJ, Camicia M. Family caregiving. In: Larsen PD, editor. *Lubkin's Chronic Illness: Impact and Intervention*. 11th ed. Jones & Bartlett Learning; 2022:329-378. [Context Link 1]
55. McCusker J, et al. Unmet needs of family caregivers of hospitalized older adults preparing for discharge home. *Chronic Illness* 2020;16(2):131-145. DOI: 10.1177/1742395318789467. [Context Link 1] View abstract...
56. Safe drug administration. In: Lippincott Williams & Wilkins, editor. *Nursing Drug Handbook*. 2023 ed. Lippincott Williams & Wilkins; 2023:12-22. [Context Link 1, 2, 3, 4, 5]
57. The case management process. In: Powell SK, Tahan H, editors. *Case Management a Practical Guide for Education and Practice*. 4th ed. Philadelphia, PA: Wolters Kluwer, Lippincott, Williams & Wilkins; 2019:92-118. [Context Link 1, 2, 3, 4, 5]
58. Managing Medicines for Adults Receiving Social Care in the Community. NICE Guidance NG67 [Internet] National Institute for Health and Care Excellence. 2017 Mar Accessed at: <https://www.nice.org.uk/guidance/>. [accessed 2023 Aug 08] [Context Link 1, 2]
59. The nursing process in drug therapy and patient safety. In: Tucker RG, editor. *Karch's Focus on Nursing Pharmacology*. 9th ed. Wolters Kluwer; 2023:47-57. [Context Link 1, 2, 3, 4]
60. Ostendorf WR. Preparation for safe medication administration. In: Perry AG, Potter PA, Ostendorf WR, editors. *Nursing Interventions and Clinical Skills*. 7th ed. Elsevier; 2020:551-567. [Context Link 1, 2]
61. Therapeutic drug monitoring guidelines. In: Lippincott Williams & Wilkins, editor. *Nursing Drug Handbook*. 2023 ed. Lippincott Williams & Wilkins; 2023:1660-1667. [Context Link 1]
62. Poison Prevention. [Internet] American Association of Poison Control Centers. Accessed at: <https://aapcc.org/Prevention/>. [accessed 2023 Sep 12] [Context Link 1]
63. A - W. In: Phelps LL, editor. *Nursing Diagnosis Reference Manual*. 12th ed. Wolters Kluwer; 2023:2-750. [Context Link 1]
64. Baruth JM, Gentry MT, Rummans TA, Miller DM, Burton MC. Polypharmacy in older adults: the role of the multidisciplinary team. *Hospital Practice* 2020;1-7. DOI: 10.1080/21548331.2019.1706995. [Context Link 1] View abstract...
65. Sutter RE, Hernandez L. Working with vulnerable people. In: Rector C, Stanley MJ, editors. *Community and Public Health Nursing Promoting the Public's Health*. 10th ed. Wolters Kluwer; 2022:667-688. [Context Link 1]
66. Antidepressants. In: Burchum JR, Rosenthal LD, editors. *Lehne's Pharmacology for Nursing Care*. 11th ed. Elsevier; 2022:351-371. [Context Link 1, 2, 3, 4]
67. Alphabetical listing of drugs by generic name. In: Lippincott Williams & Wilkins, editor. *Nursing Drug Handbook*. 2024 ed. Wolters Kluwer Health; 2024:63-1581. [Context Link 1, 2]
68. Antidepressant agents. In: Tucker RG, editor. *Karch's Focus on Nursing Pharmacology*. 9th ed. Wolters Kluwer; 2023:358-375. [Context Link 1, 2]
69. Psychopharmacology. In: Morgan KI, Townsend MC, editors. *Davis Advantage for Psychiatric Mental Health Nursing*. 10th ed. Philadelphia, PA: F.A. Davis; 2021:53-86. [Context Link 1]

70. Management of anxiety disorders. In: Burchum JR, Rosenthal LD, editors. *Lehne's Pharmacology for Nursing Care*. 11th ed. Elsevier; 2022:395-401. [Context Link 1]
71. Keltner NL, Vossos H. Antidepressant drugs. In: Steele D, editor. *Keltner's Psychiatric Nursing*. 9th ed. Elsevier; 2023:157-176. [Context Link 1, 2]
72. Serotonin syndrome: what you should know to protect your patient. In: Lippincott Williams & Wilkins, editor. *Nursing Drug Handbook*. 2023 ed. Lippincott Williams & Wilkins; 2023:1604. [Context Link 1, 2]
73. Selected therapeutic drug classifications. In: Lippincott Williams & Wilkins, editor. *Nursing Drug Handbook*. 2024 ed. Wolters Kluwer Health; 2024:23-62. [Context Link 1, 2]
74. Adverse drug reactions and medication errors. In: Burchum JR, Rosenthal LD, editors. *Lehne's Pharmacology for Nursing Care*. 11th ed. Elsevier; 2022:63-73. [Context Link 1, 2, 3]
75. Fischer SH, et al. Standardized assessment of medication reconciliation in post-acute care. *Journal of the American Geriatrics Society* 2022;70(4):1047-1056. DOI: 10.1111/jgs.17655. [Context Link 1, 2] View abstract...
76. Rhinehart HE, Snider MJ, James A, Blosser L. Improvement in the medication reconciliation postdischarge quality measure after implementation of a pharmacist-run service. *Journal of the American Pharmacists Association* 2020;60(2):391-396. DOI: 10.1016/j.japh.2019.10.006. [Context Link 1] View abstract...
77. Mayer RS, Noles A, Vinh D. Determination of postacute hospitalization level of care. *Medical Clinics of North America* 2020;104(2):345-357. DOI: 10.1016/j.mcna.2019.10.011. [Context Link 1] View abstract...
78. Levels of care and care settings. In: Gillingham DC, editor. *Foundations of Case Management*. Blue Bayou Press; 2021:131-136. [Context Link 1]
79. Standards of Practice for Case Management. [Internet] Case Management Society of America. 2022 Accessed at: <https://www.cmsa.org/>. [accessed 2023 Sep 21] [Context Link 1, 2, 3, 4, 5, 6, 7, 8]
80. Accreditation bodies, their standards, and requirements. In: Gillingham DC, editor. *Foundations of Case Management*. Blue Bayou Press; 2021:241-247. [Context Link 1]
81. Standards and Guidelines for the Accreditation and Certification for Case Management Accreditation. Effective for Surveys Beginning on or After July 1, 2020 [Internet] National Committee for Quality Assurance (NCQA). 2020 Accessed at: To purchase copies of this publication, visit <https://ncqa.org>. [accessed 2023 Oct 24] [Context Link 1, 2]
82. Standards and Guidelines for the Accreditation of Case Management for Long-Term Services and Supports Program. Effective for Surveys Beginning on or After July 1, 2024 [Internet] National Committee for Quality Assurance (NCQA). 2023 Accessed at: To purchase copies of this publication, visit <https://ncqa.org>. [accessed 2023 Sep 05] [Context Link 1, 2]
83. Transitions of Care Standards. A New Way Forward [Internet] American Case Management Association. Accessed at: <https://www.acmaweb.org/>. [accessed 2023 Sep 21] [Context Link 1, 2]
84. Nall RW, Herndon B, Mramba LK, Vogel-Anderson K, Hagen MG. An interprofessional primary care based transition of care clinic to reduce hospital readmission. *American Journal of Medicine* 2020;133(6):e260-e268. DOI: 10.1016/j.amjmed.2019.10.040. [Context Link 1] View abstract...
85. Nguyen HQ, et al. Association between post-hospital clinic and telephone follow-up provider visits with 30-day readmission risk in an integrated health system. *BMC Health Services Research* 2021;21(1):826. DOI: 10.1186/s12913-021-06848-9. [Context Link 1] View abstract...
86. Anderson A, et al. Follow-up post-discharge and readmission disparities among Medicare fee-for-service beneficiaries, 2018. *Journal of General Internal Medicine* 2022;37(12):3020-3028. DOI: 10.1007/s11606-022-07488-3. [Context Link 1] View abstract...
87. Marcondes FO, et al. Does scheduling a postdischarge visit with a primary care physician increase rates of follow-up and decrease readmissions? *Journal of Hospital Medicine* 2019;14:E37-E42. DOI: 10.12788/jhm.3309. [Context Link 1] View abstract...
88. Coppa K, Kim EJ, Oppenheim MI, Bock KR, Conigliaro J, Hirsch JS. Examination of post-discharge follow-up appointment status and 30-day readmission. *Journal of General Internal Medicine* 2021;36(5):1214-1221. DOI: 10.1007/s11606-020-06569-5. [Context Link 1] View abstract...
89. Improving adherence. In: Perez R, Rogers S, Prince M, Fraser K, editors. *Case Management Adherence Guidelines*. 2020 ed. Case Management Society of America; 2022:19-42. [Context Link 1]
90. Physical Activity Guidelines for Americans. [Internet] U.S. Department of Health and Human Services. 2nd ed; 2018 Accessed at: <https://health.gov/our-work>. [accessed 2023 Sep 18] [Context Link 1]
91. Xiao R, Miller JA, Zafirau WJ, Gorodeski EZ, Young JB. Impact of home health care on healthcare resource utilization following hospital discharge: a cohort study. *American Journal of Medicine* 2018;131(4):395-407. DOI: 10.1016/j.amjmed.2017.11.010. [Context Link 1] View abstract...
92. Chovanec KA, Arsene C, Beck A, Zachrich K, Liedel B, Wolff-Elliott J. Association of discharge disposition with outcomes. *Population Health Management* 2021;24(1):116-121. DOI: 10.1089/pop.2019.0176. [Context Link 1] View abstract...
93. Bethishou L, Herzik K, Fang N, Abdo C, Tomaszewski DM. The impact of the pharmacist on continuity of care during transitions of care: A systematic review. *Journal of the American Pharmacists Association* 2020;60(1):163-177.e2. DOI: 10.1016/j.japh.2019.06.020. [Context Link 1] View abstract...
94. Weir DL, et al. Challenges at care transitions: failure to follow medication changes made at hospital discharge. *American Journal of Medicine* 2019;132(10):1216-1224.e5. DOI: 10.1016/j.amjmed.2019.05.003. [Context Link 1] View abstract...
95. Gentry MT, Lapid MI, Rummans TA. Geriatric telepsychiatry: systematic review and policy considerations. *American Journal of Geriatric Psychiatry* 2019;27(2):109-127. DOI: 10.1016/j.jagp.2018.10.009. [Context Link 1] View abstract...

Footnotes

- [A] The timing of provider follow-up depends on the patient's disease severity, self-care complexity, patient/caregiver willingness and ability to provide patient care, and available social supports.(57) [A in Context Link 1]
- [B] Telehealth may improve access to and coordination of mental health care. Patients with behavioral health diagnoses may use telehealth or telepsychiatry to access mental health support or care (eg, to get assistance with problem-solving techniques or ask about presenting symptoms or medication use).(95) [B in Context Link 1]
- [C] Electroconvulsive therapy (ECT) may be useful for severe and psychotic depression, particularly if the patient is not responsive to pharmacologic treatment.(5)(20) [C in Context Link 1]
- [D] Bright light therapy is used as a first-line treatment for seasonal affective disorder (SAD).(96) [D in Context Link 1]
- [E] Medications to avoid include sympathomimetics (eg, amphetamines, combination cold tablets, nasal decongestants, and appetite suppressants), illicit drugs, narcotic medications (eg, meperidine, tramadol, and methadone), selective serotonin reuptake inhibitors (SSRIs), and tricyclic antidepressants (TCAs).(66) [E in Context Link 1]

Codes

ICD-10 Diagnosis: F06.32, F32.0, F32.1, F32.2, F32.3, F32.4, F32.9, F32.A, F33.0, F33.1, F33.2, F33.3, F33.40, F33.41, F33.9, F34.1, F43.21, F53.0, F53.1, O90.6, R45.88, Z91.51, Z91.52 [Hide]

DSM-5: F06.32, F32.0, F32.1, F32.2, F32.3, F32.4, F32.9, F32.A, F33.0, F33.1, F33.2, F33.3, F33.41, F33.9, F34.1, F43.21, R45.88 [Hide]