

Lung Transplant

ORG: S-1300 (ISC)

[Link to Codes](#)

MCG Health

Inpatient & Surgical

Care

28th Edition

- Care Planning - Inpatient Admission and Alternatives
 - Clinical Indications for Procedure
 - Alternatives to Procedure
 - Operative Status Criteria
 - Preoperative Care Planning
- Hospitalization
 - Optimal Recovery Course
 - Goal Length of Stay - **12 days postoperative**
 - Extended Stay
 - Hospital Care Planning
- Discharge
 - Discharge Planning
 - Discharge Destination
- Evidence Summary
 - Criteria
 - Length of Stay
 - Rationale
 - Related CMS Coverage Guidance
- References
- Footnotes
- Definitions
- Codes

Care Planning - Inpatient Admission and Alternatives

Clinical Indications for Procedure

- Procedure is indicated for **ALL** of the following(1)(2):[\[NNNNNN\]](#)
 - ☐ End-stage lung disease appropriate for transplant is present, as indicated by **1 or more** of the following:
 - ☐ COPD and **1 or more** of the following(1)(4):
 - BODE index greater than or equal to 7[A]
 - FEV1 less than 20% predicted
 - History of severe exacerbations (eg, acute hypercapnia (PCO₂ greater than 50 mm Hg (6.7 kPa)))
 - Chronic hypercapnia
 - Moderate to severe pulmonary hypertension (eg, mean pulmonary artery pressure greater than 35 mm Hg or mean right atrial pressure greater than 15 mm Hg)
 - ☐ Cystic fibrosis (CF) or non-CF bronchiectasis and **1 or more** of the following(1)(5)(6):
 - FEV1 less than 25% predicted
 - Rapidly declining lung function (eg, more than 30% decline over 12 months)
 - Frequent hospitalization (eg, greater than 28 days of hospitalization over last 12 months)
 - Exacerbation requiring mechanical ventilation
 - Hypoxemia
 - Hypercapnia (PCO₂ greater than 50 mm Hg (6.7 kPa))
 - Pulmonary hypertension (eg, echocardiogram evidence of pulmonary artery systolic pressure greater than 50 mm Hg or right ventricular dysfunction)
 - Recurrent hemoptysis not controlled by embolization
 - Refractory or recurrent pneumothorax
 - Ongoing weight loss or worsening nutritional status despite aggressive nutritional supplementation
 - World Health Organization functional class IV[B]
 - ☐ Interstitial lung disease (eg, idiopathic pulmonary fibrosis) and **1 or more** of the following[C](1)(7)(8):
 - Diffusion capacity less than 40% of predicted with decline of 10% or more over 6 months despite appropriate treatment
 - Forced vital capacity less than 80% of predicted with decrease in forced vital capacity greater than 10% over 6 months despite appropriate treatment

- Forced vital capacity less than 80% of predicted with decrease in forced vital capacity greater than 5% with radiographic progression
- Pulse oximetry less than 88% during 6-minute walk test
- Decrease in 6-minute walk test distance by more than 50 meters (165 feet) over 6 months
- Pulmonary hypertension
- Hospitalization due to acute exacerbation, related pneumothorax, or respiratory deterioration
- ☐ Pulmonary arterial hypertension and **1 or more** of the following^[D](1)(9)(10):
 - European Society of Cardiology/European Respiratory Society (ESC/ERS) high-risk category, or REVEAL score greater than 10 despite medical therapy (eg, combination therapy including IV or SC prostanoids)^[E]
 - Worsening Hypoxemia
 - Progressive renal or liver dysfunction due to pulmonary hypertension and right-sided heart failure
 - Life-threatening hemoptysis
- ☐ Lymphangioleiomyomatosis and **ALL** of the following(1)(11):
 - Severe findings, as indicated by **1 or more** of the following:
 - Severe lung function impairment (eg, FEV1 less than 30% predicted)
 - WHO functional class III^[F] or IV^[B]
 - Hypoxemia at rest
 - Pulmonary hypertension
 - Refractory pneumothorax
 - Treatment with mTOR inhibitor (eg, everolimus, sirolimus) has failed, is not tolerated, or is contraindicated.
- ☐ Pulmonary Langerhans cell histiocytosis and **1 or more** of the following^[G](12):
 - Severe lung function impairment
 - Severe limitation in exercise capacity
 - Hypoxemia at rest
- ☐ Sarcoidosis with WHO functional class III^[F] or IV^[B] and **1 or more** of the following(3)(13):
 - Pulmonary hypertension unresponsive to medical treatment⁽¹⁴⁾
 - Elevated right atrial pressure greater than 15 mm Hg
 - Hypoxemia at rest
- ☐ Graft vs host disease and **1 or more** of the following(15)(16):
 - Progressive lung damage resulting in severe compromise of activities of daily living
 - Life expectancy limited by lung disease
- ☐ Lung-limited lung cancer and **ALL** of the following(1):
 - Tumor is appropriate for transplant, as evidenced by **1 or more** of the following:
 - Adenocarcinoma in situ
 - Lepidic adenocarcinoma (previously called bronchoalveolar cell carcinoma)
 - Minimally invasive adenocarcinoma
 - Surgical therapy is not feasible (eg, multifocal disease or prohibitive pulmonary function).
 - Systemic therapies are contraindicated or have failed.
 - Lung transplant is anticipated to be curative.
 - Other end-stage lung disease (eg, prolonged acute respiratory distress syndrome, acute fulminant lung damage including post-COVID-19 infection) not expected to recover^[H](1)(17)(18)(20)(21)(22)
 - Retransplant after failure of primary transplant (eg, chronic lung allograft dysfunction, anastomotic stenosis, bronchiolitis obliterans, restrictive allograft syndrome)(23)
- Alternative to lung transplant is not available or not appropriate (eg, lung volume reduction surgery or bronchoscopic lung volume reduction for COPD, pulmonary thromboendarterectomy for patient with chronic pulmonary thromboembolic disease).(4)(9)(10)
- Medical therapy for condition does not exist or has failed.^[I]
- No contraindications to lung transplant present ^[J]

Alternatives to Procedure

- Alternatives include(1):
 - Medical management of COPD(4)
 - Medical management of pulmonary arterial hypertension(9)(10)(26)
 - Medical management of idiopathic pulmonary fibrosis(27)
 - Atrial septostomy for patient with pulmonary arterial hypertension unresponsive to medical therapy(28)
 - Lung volume reduction surgery for severe upper lobe predominant emphysema(4)(29)
 - Bronchoscopic lung volume reduction (eg, endobronchial valve, coil, or ablation; for severe emphysema with hyperinflation)(4)(30)(31)
 - Pulmonary thromboendarterectomy for chronic pulmonary thromboembolic pulmonary hypertension(32)
 - Soluble guanylate cyclase stimulator (eg, riociguat) for chronic pulmonary thromboembolic pulmonary hypertension(10)
 - Palliative care(4)(33)

Operative Status Criteria

Goal Length of Stay: 12 days postoperative

- Inpatient

Preoperative Care Planning

- Preoperative care planning needs may include(1)(2)(4)(8)(10)(34)(35)(36):
 - Preoperative evaluation, including:
 - Thorough evaluation by multidisciplinary lung transplant team for:
 - Severity of respiratory failure
 - Alternatives to lung transplant
 - Contraindications to lung transplant
 - Psychosocial evaluation including treatment adherence, mental health history, substance abuse, cognitive function, coping strategies, and social support(37)
 - Protocol as specified by transplant center
 - Routine preoperative evaluation. See Preoperative Education, Assessment, and Planning Tool [SR](#).
 - Diagnostic test scheduling, including:
 - Cardiac catheterization including pulmonary arterial pressure measurements(35)(38)
 - Carotid Doppler ultrasound(38)
 - Pulmonary function testing
 - 6-minute walk test
 - Cardiopulmonary exercise test with measurement of maximum oxygen uptake (VO₂ maximum)
 - Echocardiogram
 - Chest radiographic studies (may include chest x-ray, CT scan, CT angiogram, diaphragm fluoroscopy)(39)
 - Cardiac MRI (eg, for connective tissue disease or sarcoidosis with suspected myocardial involvement)(34)
 - Quantitative ventilation/perfusion lung scan(40)
 - Estimation of lung allograft size(39)
 - Laboratory testing
 - Hematologic evaluation for coagulopathy and thrombophilia (hypercoagulable state) (eg, for patient with lupus or other connective tissue disease, chronic thromboembolic pulmonary hypertension)(8)(34)
 - Nutrition evaluation(6)(41)
 - Dental evaluation
 - Evaluation for gastroesophageal reflux and aspiration risk (eg, esophageal manometry, pH monitoring, barium swallow)(8)(34)
 - Immunology screening, including blood and tissue typing
 - Detailed infectious disease screening, including(6)(42)(43):
 - Sputum cultures or bronchoscopy to rule out bacterial colonization with virulent multidrug-resistant organisms
 - Viral antibody titers, including HIV, cytomegalovirus, hepatitis B and C, varicella zoster
 - Viral antigen testing for hepatitis B, Epstein-Barr virus
 - HIV, hepatitis B, and hepatitis C nucleic acid testing
 - Nasopharyngeal PCR for severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2)
 - Purified protein derivative (PPD) or interferon gamma release assay (eg, QuantiFERON-TB Gold or T-SPOT.TB)
 - Screening for regional organisms as appropriate (eg, *Histoplasma* species in central United States, *Coccidioides* species in southeastern United States, *Blastomyces* species in midwestern, southern United States)(43)
 - Preoperative treatment, procedures, and stabilization, including addressing all treatable coexisting conditions before transplant
 - Vaccinations (eg, influenza; hepatitis; Streptococcus pneumoniae; SARS-CoV-2; measles, mumps, and rubella; Herpes varicella-zoster; Rotavirus)(43)
 - Preoperative pulmonary rehabilitation
 - Smoking cessation counseling
 - Extracorporeal membrane oxygenation support as bridge to transplant as indicated(1)(22)(35)(38)(44)(45)
 - Extracorporeal carbon dioxide removal support as a bridge to transplant as indicated (eg, for non-hypoxic hypercapnic respiratory failure)(22)
 - Mechanical ventilation as bridge to transplant in selected cases(22)(44)
 - Palliative care consultation(46)
 - Preoperative discharge planning as appropriate. See Discharge Planning in this guideline.

Hospitalization**Optimal Recovery Course**

Day	Level of Care	Clinical Status	Activity	Routes	Interventions	Medications
1	• OR to ICU • Social Determinants of	• Clinical Indications met[K]	• Bed rest	• IV fluids • IV medications	• Central lines • ECG • Oximetry	• Perioperative antibiotics

	- Health Assessment - Discharge planning	Intubated		NPO	- CBC, chemistries - CXR postoperatively - Possible extubation postoperatively - Oxygen - Chest tube	- Immunosuppressive medication and corticosteroids - Possible inhaled or parenteral vasoactive medication - Multimodal analgesia, possible PCA or epidural analgesics
2	- ICU - Social Determinants of Health Assessment	Hypotension absent	Bed rest	- Possible enteral feeds[L] - IV fluids - IV medications	- Central lines - Portable CXR - Chest tube - Oxygen - Oximetry	- Perioperative antibiotics - Immunosuppressive medication and corticosteroids - Possible inhaled or parenteral vasoactive medication - Multimodal analgesia
3	- ICU - Social Determinants of Health Assessment	No evidence of myocardial ischemia Dangerous arrhythmia absent	Bed rest	- Possible enteral feeds - IV fluids - IV medications	- Central lines - Possible chest tube - Oxygen - Oximetry	Epidural analgesics absent - Antibiotics absent - Immunosuppressive medication and corticosteroids - Possible inhaled or parenteral vasoactive medication - Possible PCA
4	- ICU - Social Determinants of Health Assessment	Hemodynamic stability	Bed rest	- Possible enteral feeds - IV fluids - IV medications	- Possible chest tube - Central lines - Oxygen - Oximetry	Wean inhaled or parenteral vasoactive medication PCA absent - Immunosuppressive medication and corticosteroids - Oral or IV analgesic agent
5	- ICU - Social Determinants of Health Assessment	Mechanical ventilation absent	Bed rest	- Possible enteral feeds - IV fluids - IV medications	Chest tube absent - Central lines - Oxygen - Oximetry	Inhaled or parenteral vasoactive medication absent - Immunosuppressive medication and corticosteroids - Oral or IV analgesic agent
6	- Possible intermediate care - Social Determinants of Health Assessment	Pain absent or managed	Dangle legs	Begin oral diet - IV fluids - IV medications	- Possible central lines - Oxygen - Oximetry	IV analgesic agent absent - Immunosuppressive medication and corticosteroids - Oral analgesic agent as needed

7	- Intermediate care or floor - Social Determinants of Health Assessment	Pulmonary edema absent or mild	Up to chair	- Advance diet - IV fluids - IV medications	Central lines absent - Oxygen - Oximetry	- Immunosuppressive medication and corticosteroids - Oral analgesic agent as needed
8	- Intermediate care or floor - Social Determinants of Health Assessment	Rejection absent or controlled Stable fluid balance	Limited ambulation	Oral medications - Advance diet - IV fluids or oral hydration	- Oxygen - Oximetry	- Immunosuppressive medication and corticosteroids conversion from intravenous to oral regimen - Oral analgesic agent as needed
9	- Intermediate care or floor - Social Determinants of Health Assessment	SaO₂ greater than 90% on room air	Limited ambulation	Oral hydration Oral medications - Advance diet	Taper oxygen - Oximetry	- Oral immunosuppressive medication and corticosteroids - Oral analgesic agent as needed
10	- Floor - Social Determinants of Health Assessment	Oxygenation stable with activity	Limited ambulation	Oral hydration Oral medications Oral diet	Oxygen absent - Oximetry - Immunosuppressant levels	- Oral immunosuppressive medication and corticosteroids - Oral analgesic agent as needed
11	- Floor - Social Determinants of Health Assessment	No evidence of postoperative or surgical site infection	Ambulatory or acceptable for next level of care	- Oral hydration - Oral medications - Oral diet	- Oximetry - Immunosuppressant levels	- Oral immunosuppressive medication and corticosteroids - Oral analgesic agent as needed
12	- Floor - Social Determinants of Health Assessment	Lung function studies acceptable	Ambulatory	- Oral hydration - Oral medications - Oral diet	- Oximetry - Immunosuppressant levels	Immunosuppressive medication regimen for next level of care initiated - Oral analgesic agent as needed
13	- Social Determinants of Health Assessment - Floor to discharge - Complete discharge planning	Hemodynamic stability Rejection absent No evidence of postoperative or surgical site infection SaO₂ greater than 90% on room air and stable with activity Pain absent or managed Discharge plans and education	Ambulatory or acceptable for next level of care	Oral hydration[M] Oral medications or regimen acceptable for next level of care Oral diet or acceptable for next level of care	- Oximetry - Immunosuppressant levels	Immunosuppressive medication regimen for next level of care established - Oral analgesic agent as needed

understood

(36)(38)(44)(48)(49)(50)(51)(52)(53)N

Recovery Milestones are indicated in **bold**.

Goal Length of Stay: 12 days postoperative

Note: Goal Length of Stay assumes optimal recovery, decision making, and care. Patients may be discharged to a lower level of care (either later than or sooner than the goal) when it is appropriate for their clinical status and care needs.

Extended Stay

Minimal (a few hours to 1 day), Brief (1 to 3 days), Moderate (4 to 7 days), and Prolonged (more than 7 days).

- Extended stay beyond goal length of stay may be needed for(36)(44)(48)(53):
 - Failure to meet discharge criteria (recovery milestones within final day in Optimal Recovery Course)
 - Expect brief stay extension.
 - Primary graft dysfunction or respiratory insufficiency(44)(45)(48)(54)
 - Anticipate continued mechanical ventilation.
 - Anticipate extracorporeal membrane oxygen support until lung function recovers.(22)
 - Respiratory insufficiency post extubation may require noninvasive ventilation or high-flow oxygenation to prevent reintubation.
 - Anticipate use of inhaled nitric oxide or epoprostenol for severe hypoxemia or pulmonary hypertension.
 - Expect moderate to prolonged stay extension.
 - Aspiration pneumonitis
 - Anticipate bedside swallow evaluation and possible barium swallow or functional endoscopic evaluation of swallowing (FEES).
 - Expect brief stay extension.
 - Infection (eg, blood stream infection, pneumonia, surgical wound, anastomosis)(43)(55)
 - Anticipate intravenous antifungal, antiviral, or antibiotic agents.
 - Anticipate diagnostic testing including CT chest, cultures, viral and fungal serology, galactomannan and beta-D glycan levels, and nucleic acid amplification test (eg, PCR).
 - Pneumonia may require bronchoscopy for bronchoalveolar lavage, transbronchial biopsy, and protected brush cultures.
 - Expect moderate stay extension.
 - Postoperative empyema(38)
 - Anticipate cultures, chest tube drainage, antibiotic treatment, intrapleural thrombolytic agent, and possible thoracoscopic decortication.
 - Expect moderate to prolonged stay extension.
 - Acute or hyperacute rejection(56)
 - Anticipate increased immunosuppression, plasma exchange, bronchoscopy, and lung biopsy.
 - Expect moderate stay extension.
 - Surgical complications(38)(57)(58)
 - Hemorrhage, pneumothorax, hemothorax, empyema, or persistent air leaks may require prolonged chest tube drainage.
 - Hemothorax may require correction of coagulopathy, transfusion, and possible surgical exploration.
 - Chylothorax may require dietary modification, thoracic duct embolization, or reoperation for thoracic duct ligation.
 - Expect brief to moderate stay extension.
 - Recurrent pleural effusion(58)
 - Recurrent effusions may require thoracentesis, catheter drainage, or thoracoscopic intervention.
 - Expect brief to moderate stay extension.
 - Heart-lung transplant(59)(60)
 - Complex surgery and complications such as phrenic or vagus nerve dysfunction, gastroparesis, and chylothorax may require longer inpatient care.
 - Anticipate longer dependence on mechanical ventilation.
 - Expect brief to moderate stay extension.
 - Lung-liver transplant(60)(61)
 - Complications from liver transplant (eg, hepatic artery or portal vein thrombosis, graft nonfunction, liver hematoma) may require longer inpatient care.
 - Post liver transplant graft reperfusion syndrome may cause hypotension and compromise lung function.
 - Vasopressor support and conservative fluid management required for lung allograft may compromise liver graft recovery.
 - Expect moderate stay extension.
 - Bronchial anastomotic defects, necrosis, or dehiscence(38)
 - Anticipate possible bronchoscopic intervention (eg, stent, cyanoacrylate glue) or surgical revision of anastomosis.

- Surgical intervention (eg, drainage, fistula closure, reanastomosis, or flap bronchoplasty) may be required.
- Expect brief to moderate stay extension.
- Pulmonary artery anastomotic complication (eg, thrombosis, kinking, torsion)(58)
 - Expect possible reoperation and revision of anastomosis.
 - Expect moderate to prolonged stay extension.
- Venous thromboembolism(62)
 - Anticipate CT or MR angiogram and anticoagulation.
 - Expect brief to moderate stay extension.
- Combined lung transplant with repair of underlying congenital heart lesion
 - Complex surgery and complications such as phrenic nerve dysfunction may require longer inpatient care.
 - Expect moderate to prolonged stay extension.
- Care for comorbidities
 - Patient with active comorbidities, such as chronic heart failure, liver disease, chronic renal disease, or diabetes mellitus may require continued inpatient care.
 - Expect brief to moderate stay extension.
- Postoperative atrial fibrillation(63)
 - Anticipate rate control (eg, calcium channel blocker, beta-blocker) or antiarrhythmic agent (eg, amiodarone, ibutilide), or cardioversion.
 - Expect brief to moderate stay extension.
- Medical complications of surgery (eg, acid base disturbance, electrolyte disorder, Acute kidney injury (stage 2) or Acute renal failure (stage 3 acute kidney injury), delirium)(57)(64)(65)
 - Anticipate evaluation and treatment of specific complication.
 - Anticipate possible renal replacement therapy for patient with acute renal failure.
 - Expect brief to moderate stay extension.
- High preoperative lung allocation score(66)(67)
 - Anticipate preoperative hospital and ICU days for stabilization of organ function.
 - Anticipate higher incidence of infection, acute renal failure, and need for inpatient dialysis.
 - Expect brief to prolonged stay extension.
- Transplant for COVID-19 infection(18)(19)
 - Anticipate prolonged need for postoperative mechanical ventilation and ICU care.
 - Expect prolonged stay extension.
- Severe malnutrition(68)
 - Expect brief to moderate stay extension.

See Common Complications and Conditions [ISC](#) for further information.

Hospital Care Planning

- Hospital evaluation and care needs may include(36)(38)(48):
 - Diagnostic test scheduling and completion, including:
 - CBC, chemistries monitoring
 - Chest CT scan and other chest imaging as indicated(69)
 - Transesophageal echocardiogram
 - Pulmonary function testing
 - Lower extremity Duplex ultrasound (for routine deep venous thrombosis screening)(70)
 - Treatment and procedure scheduling and completion, including:
 - Steroids
 - IV antibiotics(43)
 - Prophylactic anti-infective regimen, including antifungal, anti-CMV, and anti-*Pneumocystis jirovecii* agents(43)
 - Chest physiotherapy
 - Bronchoscopy
 - Thoracentesis
 - Establishment of immunosuppression protocols(71)
 - Right-sided heart catheterization(35)
 - Inhaled pulmonary vasodilator (eg, nitric oxide, epoprostenol) for pulmonary hypertension, right ventricular failure, or refractory hypoxia(72)
 - Multimodal analgesia (38)
 - Extracorporeal membrane oxygenation[N]
 - Consultation, assessment, and other services scheduling and completion, including:
 - Medical and surgical specialty consultations, as indicated
 - Dietitian(47)
 - Monitoring patient's status for deterioration and comorbid conditions (see Inpatient Monitoring and Assessment Tool [SR](#)); key items include(48):
 - Hemodynamic stability measurements, which may initially include:

- Blood pressure via arterial line
- Pulmonary artery pressure via right-sided heart catheter
- Central venous pressure
- Cardiac output and cardiac index
- Urine output
- Monitoring for arrhythmias
- Pleural drainage quantity and appearance and need for chest tubes
- Pulmonary status, including presence of:
 - Dyspnea or cough
 - Hypoxemia
 - Hypercarbia
 - Signs and symptoms of respiratory failure or diaphragm dysfunction
- Wound tenderness, swelling, or serous drainage from incision
- Pain management
- Renal function
- Gastrointestinal status
- Glycemic control (eg, due to high-dose corticosteroids)
- Signs and symptoms of infection or acute rejection, including:
 - Low-grade fever
 - Elevated WBC count
 - Decrease in FEV1 more than 10%
 - New infiltrates on chest x-ray
 - Pleuritic chest pain
 - New or increased sputum production
 - Purulent drainage from site of infection
 - Increasing oxygen requirements

Discharge

Discharge Planning

- Discharge planning includes^[O]:
 - Assessment of needs and planning for care, including⁽⁷⁴⁾⁽⁷⁵⁾:
 - Develop and modify treatment plan (involving multiple providers) as needed.
 - Evaluate and address preadmission functioning as needed.
 - Evaluate and address psychosocial status issues as indicated. See Psychosocial Assessment [SR](#) for further information.
 - Evaluate and address social determinants of health (eg, housing, food). See Social Determinants of Health Screening Tool [SR](#) for further information.⁽⁷³⁾
 - Evaluate and address patient or caregiver preferences as indicated.
 - Identify skilled services needed at next level of care, with specific attention to⁽⁷⁶⁾:
 - Medication management, adherence instruction, and side effects assessment⁽⁷⁷⁾
 - Ongoing education required
 - Psychosocial assessment, management, and referrals⁽⁷⁸⁾
 - Respiratory status assessment⁽⁷⁹⁾
 - Wound or dressing management
 - Early identification of anticipated discharge destination; options include⁽⁷⁵⁾⁽⁸⁰⁾:
 - Home, considerations include:
 - Home safety assessment. See Home Safety Assessment [SR](#) for further information.
 - ☐ Patient safe to go home; examples include⁽⁸¹⁾⁽⁸²⁾⁽⁸³⁾:
 - Medical status stable for patient's condition
 - Functional care can safely be provided with available resources.
 - Mental status stable for patient's condition
 - Medication availability confirmed and reconciliation complete
 - Patient/caregiver education completed with written discharge instructions provided
 - Community resources identified and referrals made, as needed
 - Home care arranged, if indicated
 - Necessary medical equipment delivery arranged or available in home, if indicated
 - Necessary medical supplies ordered, or patient/caregiver can obtain, if indicated
 - Access to follow-up care
 - Self-management ability if appropriate. See Activities of Daily Living (ADL) and Instrumental Activities of Daily Living (IADL) Assessment [SR](#) for further information.
 - Caregiver need, ability, and availability
 - Post-acute skilled care or custodial care as indicated. See Discharge Planning Tool [SR](#) for further information.

- Transitions of care plan complete, including(75)(80)(84):
 - Patient and caregiver education complete. See Lung Transplant: Patient Education for Clinicians [↗](#) SR for further information.
 - See Teach Back Tool [↗](#) SR for further information.
- ☐ Medication reconciliation completion includes(77)(85):
 - Compare patient's discharge list of medications (prescribed and over-the-counter) against provider's admission or transfer orders.
 - Assess each medication for correlation to disease state or medical condition.
 - Report medication discrepancies to prescribing provider, attending physician, and primary care provider, and ensure accurate medication order is identified.
 - Provide reconciled medication list to all treating providers.
 - Confirm that patient or caregiver can acquire medication.
 - Educate patient and caregiver.
 - Provide complete medication list to patient and caregiver.
 - Importance of presenting personal medication list to all providers at each care transition, including all provider appointments
 - Reason, dosage, and timing of medication (eg, use "teach-back" techniques)(86)
 - Encourage communication between patient, caregiver, and pharmacy for obtaining prescriptions, setting up home medication delivery, and reviewing for drug-drug interactions.
 - See Medication Reconciliation Tool [↗](#) SR for further information.
- Plan communicated to patient, caregiver, and all members of care team, including(87):
 - Inpatient care and service providers
 - Primary care provider
 - All post-discharge care and service providers
- Appointments planned or scheduled, which may include(76):
 - Primary care provider
 - Behavioral health provider
 - Pulmonologist
 - Rehabilitation therapy services(88)
 - Thoracic surgeon
 - Transplant center or team
 - Other
- Outpatient testing and procedure plans made, which may include(76):
 - Laboratory testing
 - Radiology
 - Spirometry testing
 - Other
- Referrals made for assistance or support, which may include:
 - Behavioral health services (eg, counseling)
 - Educational program (eg, chronic condition management, self-management)(89)
 - Financial, for follow-up care, medication, and transportation
 - Self-help or support groups
 - Tobacco use treatment(90)
 - Vocational rehabilitation(76)
 - Other
- Medical equipment and supplies coordinated (ie, delivered or delivery confirmed), which may include:
 - Antiembolic or compression stockings
 - Blood pressure cuff
 - Incentive spirometer(76)
 - Nutritional supplements
 - Wound care equipment and supplies(78)(91)
 - Other

Discharge Destination

- Post-hospital levels of admission may include:
 - Home.
 - Home healthcare. See Home Care Indications for Admission Section [↗](#) HC in Lung Transplant guideline in Home Care.

Evidence Summary

Criteria

The evidence for the clinical indications found in this guideline includes 19 published peer reviewed articles and 5 specialty society or other evidence-based guidelines.

For COPD, consensus guidelines state that lung transplant may be appropriate for patients with BODE score greater than 7 and additional factors, such as FEV1 less than 20% of predicted, recurrent hospitalizations, or moderate to severe pulmonary hypertension.(1)(2) **(EG 2)**

For cystic fibrosis, consensus guidelines state that lung transplant may be appropriate for patients with FEV1 less than 25% of predicted and signs of clinical deterioration, such as rapidly declining lung function, frequent hospitalizations, recurrent hemoptysis, recurrent pneumothorax, or World Health Organization functional class IV.(1)(2) **(EG 2)**

For interstitial lung disease, consensus guidelines state that lung transplant may be appropriate for patients with exercise-induced oxygen desaturation to SpO2 less than 88%, or signs of clinical deterioration such as decline in forced vital capacity, decline in diffusing capacity of the lungs for carbon monoxide (DLCO), or decline in 6-minute walking distance over a 6-month period despite treatment.(1)(2) **(EG 2)**

For pulmonary arterial hypertension, consensus guidelines state that lung transplant may be appropriate for patients with European Society of Cardiology/European Respiratory Society (ESC/ERS) high-risk category despite appropriate therapy, progressive hypoxemia, progressive liver or kidney injury, or life-threatening hemoptysis.(1)(2) **(EG 2)**

For lymphangioleiomyomatosis, consensus guidelines state that lung transplant may be appropriate for patients with FEV1 less than 30% of predicted, exertional dyspnea, hypoxemia at rest, pulmonary hypertension, refractory pneumothorax, or disease progression despite mTOR treatment.(1)(2) **(EG 2)**

For sarcoidosis, a narrative review states that lung transplant may be appropriate for patients with hypoxemia at rest, hypoxemia during exercise, pulmonary hypertension, forced vital capacity less than 80% of predicted, or diffusion capacity less than 40% of predicted.(3) **(EG 2)**

For other end-stage lung disease associated with conditions such as prolonged acute respiratory distress syndrome or pulmonary Langerhans cell histiocytosis, a consensus guideline states that lung transplant may be appropriate for patients with a high risk of death from lung disease within 2 years without transplant, and a high likelihood of post-transplant survival.(1) **(EG 2)**

Length of Stay

Analysis of 27,632 patients undergoing lung transplant (median age 60 years, 70% double lung) from the UNOS transplant registry found 25% discharged in 11 days or less.(49) **(EG 2)** Analysis of 4771 patients undergoing lung transplant (median age 61 years, 80% double lung) from the Scientific Registry of Transplant Recipients database reported 25% discharged in 12 days or less.(50) **(EG 2)** A retrospective study of 722 patients (mean age 53 years, 96% double lung) undergoing lung transplant reported that 25% were discharged in 11 days or less.(51) **(EG 2)** A retrospective study of 132 patients (median age 60 years, 70% double lung) undergoing lung transplant reported that 25% of patients were discharged within 10 days.(52) **(EG 2)** Analysis of national hospital discharge data shows 28% of hospitalized adult patients undergoing the principal procedure of lung transplant discharged in 12 days or less.(53) **(EG 3)**

Rationale

Use of this MCG care guideline helps the clinician identify, for a given procedure, which patient-specific factors and clinical conditions are appropriate for that procedure. The evidence-based clinical indication criteria assist the clinician in the decision to appropriately perform a procedure, evaluating whether the potential benefits of a procedure outweigh the potential risks. For Medicare enrollees, surgical MCG care guidelines also identify which procedures CMS has designated as inpatient only.

Use of these evidence-based clinical criteria to support decision making around the need for a given procedure is of benefit to the patient, as all procedures come with inherent risk that must be balanced by anticipated clinical benefit. Utilizing evidence-based clinical criteria enables a more accurate and patient-specific decision-making process. In addition, the use of evidence-based guidelines can help reduce unwarranted variation in care, such as divergent clinical thresholds to perform a procedure for clinically similar patients that vary across geographic regions, between facilities, and among individual clinicians.

Related CMS Coverage Guidance

This guideline supplements but does not replace, modify, or supersede existing Medicare regulations or applicable National Coverage Determinations (NCDs) or Local Coverage Determinations (LCDs).

Code of Federal Regulations (CFR): 42 CFR 412.3(92); 42 CFR 419.22(93); 42 CFR 422.101(94)

Internet-Only Manual (IOM) Citations: CMS IOM Publication 100-02, Medicare Benefit Policy Manual, Chapter 1 - Inpatient Hospital Services Covered Under Part A(95); CMS IOM Publication 100-02, Medicare Benefit Policy Manual, Chapter 6 - Hospital Services Covered Under Part B(96); CMS IOM Publication 100-02, Medicare Benefit Policy Manual, Chapter 15 - Covered Medical and Other Health Services(97); CMS IOM Publication 100-08, Medicare Program Integrity Manual, Chapter 6, Section 6.5 - Medical Review of Inpatient Hospital Claims for Part A Payment(98)

Medicare Coverage Determinations: Medicare Coverage Database(99)

References

1. Leard LE, et al. Consensus document for the selection of lung transplant candidates: An update from the International Society for Heart and Lung Transplantation. *Journal of Heart and Lung Transplantation* 2021;40(11):1349-1379. DOI: 10.1016/j.healun.2021.07.005. (Reaffirmed 2022 Jul) [Context Link 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20] View abstract...
2. Mannem H, Aversa M, Keller T, Kapnadak SG. The lung transplant candidate, indications, Timing, and selection criteria. *Clinics in Chest Medicine* 2023;44(1):15-33. DOI: 10.1016/j.ccm.2022.10.001. [Context Link 1, 2, 3, 4, 5, 6, 7, 8] View abstract...
3. Yserbyt J, et al. Solid organ transplantation in sarcoidosis. *Seminars in Respiratory and Critical Care Medicine* 2017;38(4):538-545. DOI: 10.1055/s-0037-1602383. [Context Link 1, 2] View abstract...
4. Global Strategy for the Diagnosis, Management, and Prevention of Chronic Obstructive Pulmonary Disease. 2023 report [Internet] Global Initiative for Chronic Obstructive Lung Disease (GOLD). 2023 Accessed at: <https://www.goldcopd.org>. [accessed 2023 Oct 19] [Context Link 1, 2, 3, 4, 5, 6, 7, 8]
5. Ramos KJ, et al. Lung transplant referral for individuals with cystic fibrosis: Cystic Fibrosis Foundation consensus guidelines. *Journal of Cystic Fibrosis* 2019;18(3):321-333. DOI: 10.1016/j.jcf.2019.03.002. [Context Link 1] View abstract...
6. Pilewski JM. Update on lung transplantation for cystic fibrosis. *Clinics in Chest Medicine* 2022;43(4):821-840. DOI: 10.1016/j.ccm.2022.07.002. [Context Link 1, 2, 3] View abstract...
7. Raghu G, et al. Diagnosis of idiopathic pulmonary fibrosis. an official ATS/ERS/JRS/ALAT clinical practice guideline. *American Journal of Respiratory and Critical Care Medicine* 2018;198(5):e44-e68. DOI: 10.1164/rccm.201807-1255ST. (Reaffirmed 2022 Jul) [Context Link 1, 2] View abstract...
8. Bermudez CA, Crespo MM. Managing connective tissue disease: how to select and facilitate successful transplantation. *Current Opinion in Organ Transplantation* 2022;27(3):191-197. DOI: 10.1097/MOT.0000000000000978. [Context Link 1, 2, 3, 4, 5] View abstract...
9. McLaughlin VV, et al. ACCF/AHA 2009 expert consensus document on pulmonary hypertension. *Journal of the American College of Cardiology* 2009;53(17):1573-1619. DOI: 10.1016/j.jacc.2009.01.004. (Reaffirmed 2022 May) [Context Link 1, 2, 3] View abstract...
10. Humbert M, et al. 2022 ESC/ERS Guidelines for the diagnosis and treatment of pulmonary hypertension. *European Heart Journal* 2022;43(38):3618-3731. DOI: 10.1093/eurheartj/ehac237. (Reaffirmed 2023 Mar) [Context Link 1, 2, 3, 4, 5, 6] View abstract...
11. Warrior K, Dilling DF. Lung transplantation for lymphangioleiomyomatosis. *Journal of Heart and Lung Transplantation* 2023;42(1):40-52. DOI: 10.1016/j.healun.2022.09.021. [Context Link 1] View abstract...
12. Wajda N, Zhu Z, Jandarov R, Dilling DF, Gupta N. Clinical outcomes and survival following lung transplantation in patients with pulmonary Langerhans cell histiocytosis. *Respirology* 2020;25(6):644-650. DOI: 10.1111/resp.13671. [Context Link 1] View abstract...
13. Le Pavec J, et al. Lung transplantation for sarcoidosis: outcome and prognostic factors. *European Respiratory Journal* 2021;58(2):Online. DOI: 10.1183/13993003.03358-2020. [Context Link 1] View abstract...
14. Savale L, et al. WASOG statement on the diagnosis and management of sarcoidosis-associated pulmonary hypertension. *European Respiratory Review* 2022;31(163):210165. DOI: 10.1183/16000617.0165-2021. [Context Link 1] View abstract...
15. Cheng GS, Edelman JD, Madtes DK, Martin PJ, Flowers ME. Outcomes of lung transplantation after allogeneic hematopoietic stem cell transplantation. *Biology of Blood and Marrow Transplantation* 2014;20(8):1169-1175. DOI: 10.1016/j.bbmt.2014.04.008. [Context Link 1] View abstract...
16. Kliman DS, Kotecha SR, Abelson DC, Snell GI, Glanville AR, Ma DDF. Favorable outcome of lung transplantation for severe pulmonary graft versus host disease: an Australian multicenter case series. *Transplantation* 2019;103(12):2602-2607. DOI: 10.1097/TP.0000000000002693. [Context Link 1] View abstract...
17. Aslam S, et al. Guidance from the International Society of Heart and Lung Transplantation Regarding the SARS CoV-2 Pandemic. [Internet] International Society of Heart and Lung Transplantation. 2021 Feb Accessed at: <https://www.isHLT.org/>. [accessed 2023 Oct 26] [Context Link 1, 2]
18. Avella D, Neumann H, Bharat A. Lung transplantation for coronavirus disease-2019 patients and coronavirus disease-2019 in lung transplant recipients. *Clinics in Chest Medicine* 2023;44(1):191-199. DOI: 10.1016/j.ccm.2022.11.002. [Context Link 1, 2, 3] View abstract...
19. Okumura K, Jyothula S, Kaleekal T, Dhand A. One-year outcomes of lung transplantation for COVID-19 associated end stage lung disease in the United States. *Clinical Infectious Diseases* 2023;76(12):2140-2147. DOI: 10.1093/cid/ciad072. [Context Link 1, 2] View abstract...
20. Holm AM, et al. Ethical considerations regarding heart and lung transplantation and mechanical circulatory support during the COVID-19 pandemic: an ISHLT COVID-19 Task Force statement. *Journal of Heart and Lung Transplantation* 2020;39(7):619-626. DOI: 10.1016/j.healun.2020.04.019. [Context Link 1, 2] View abstract...
21. Bharat A, Hoetzenecker K. Lung transplantation for acute respiratory distress syndrome. *Thoracic Surgery Clinics* 2022;32(2):135-142. DOI: 10.1016/j.thorsurg.2022.01.005. [Context Link 1, 2] View abstract...
22. Expert Consensus Panel, et al. The American Association for Thoracic Surgery (AATS) 2022 Expert Consensus Document: The use of mechanical circulatory support in lung transplantation. *Journal of Thoracic and Cardiovascular Surgery* 2023;165(1):301-326. DOI: 10.1016/j.jtcvs.2022.06.024. [Context Link 1, 2, 3, 4, 5, 6] View abstract...
23. Michel E, Galen Hartwig M, Sommer W. Lung retransplantation. *Thoracic Surgery Clinics* 2022;32(2):259-268. DOI: 10.1016/j.thorsurg.2021.12.001. [Context Link 1] View abstract...
24. Tang A, Thuita L, Siddiqui HU, et al. Urgently listed lung transplant patients have outcomes similar to those of electively listed patients. *Journal of Thoracic and Cardiovascular Surgery* 2020;S0022-5223(20):30997-1. DOI: <https://doi.org/10.1016/j.jtcvs.2020.02.140>. [Context Link 1]
25. Kanaparthi J, et al. Prior and perioperative revascularization does not affect survival in lung transplant patients. *Annals of Thoracic Surgery* 2020;109(6):1677-1683. DOI: 10.1016/j.athoracsur.2020.01.016. [Context Link 1] View abstract...
26. Taichman DB, et al. Pharmacologic therapy for pulmonary arterial hypertension in adults: CHEST guideline and expert panel report. *Chest* 2014;146(2):449-475. DOI: 10.1378/chest.14-0793. [Context Link 1] View abstract...
27. Kropski JA, Blackwell TS. Progress in understanding and treating idiopathic pulmonary fibrosis. *Annual Review of Medicine* 2019;70:211-224. DOI: 10.1146/annurev-med-041317-102715. [Context Link 1] View abstract...

28. Khan MS, et al. Use of balloon atrial septostomy in patients with advanced pulmonary arterial hypertension: a systematic review and meta-analysis. *Chest* 2019;156(1):53-63. DOI: 10.1016/j.chest.2019.03.003. [Context Link 1] View abstract...
29. van Agteren JE, Carson KV, Tiong LU, Smith BJ. Lung volume reduction surgery for diffuse emphysema. *Cochrane Database of Systematic Reviews* 2016, Issue 10. Art. No.: CD001001. DOI: 10.1002/14651858.CD001001.pub3. [Context Link 1] View abstract...
30. Garner JL, Shah PL. Lung volume reduction in pulmonary emphysema. *Seminars in Respiratory and Critical Care Medicine* 2020;41(6):874-885. DOI: 10.1055/s-0040-1702192. [Context Link 1] View abstract...
31. Criner GJ, et al. Improving lung function in severe heterogenous emphysema with the spiration valve system (EMPROVE). a multicenter, open-label randomized controlled clinical trial. *American Journal of Respiratory and Critical Care Medicine* 2019;200(11):1354-1362. DOI: 10.1164/rccm.201902-0383OC. [Context Link 1] View abstract...
32. Lang IM, Campean IA, Sadushi-Kolici R, Badr-Eslam R, Gerges C, Skoro-Sajer N. Chronic thromboembolic disease and chronic thromboembolic pulmonary hypertension. *Clinics in Chest Medicine* 2021;42(1):81-90. DOI: 10.1016/j.ccm.2020.11.014. [Context Link 1] View abstract...
33. Iyer AS, Sullivan DR, Lindell KO, Reinke LF. The role of palliative care in COPD. *Chest* 2021;Online. DOI: 10.1016/j.chest.2021.10.032. [Context Link 1] View abstract...
34. Crespo MM, et al. ISHLT consensus document on lung transplantation in patients with connective tissue disease: Part I: Epidemiology, assessment of extrapulmonary conditions, candidate evaluation, selection criteria, and pathology statements. *Journal of Heart and Lung Transplantation* 2021;40(11):1251-1266. DOI: 10.1016/j.healun.2021.07.014. (Reaffirmed 2022 Aug) [Context Link 1, 2, 3, 4] View abstract...
35. Bermudez CA, et al. ISHLT consensus document on lung transplantation in patients with connective tissue disease: Part II: Cardiac, surgical, perioperative, operative, and post-operative challenges and management statements. *Journal of Heart and Lung Transplantation* 2021;40(11):1267-1278. DOI: 10.1016/j.healun.2021.07.016. (Reaffirmed 2022 Aug) [Context Link 1, 2, 3, 4] View abstract...
36. Gray AL, Mulvihill MS, Hartwig MG. Lung transplantation at Duke. *Journal of Thoracic Disease* 2016;8(3):E185-E196. DOI: 10.21037/jtd.2016.02.08. [Context Link 1, 2, 3, 4] View abstract...
37. Dew MA, et al. The 2018 ISHLT/APM/AST/ICCAC/STSW recommendations for the psychosocial evaluation of adult cardiothoracic transplant candidates and candidates for long-term mechanical circulatory support. *Psychosomatics* 2018;59(5):415-440. DOI: 10.1016/j.psych.2018.04.003. [Context Link 1] View abstract...
38. Marczin N, et al. International consensus recommendations for anesthetic and intensive care management of lung transplantation. An EACTAIC, SCA, ISHLT, ESOT, ESTS, and AST approved document. *Journal of Heart and Lung Transplantation* 2021;40(11):1327-1348. DOI: 10.1016/j.healun.2021.07.012. [Context Link 1, 2, 3, 4, 5, 6, 7, 8, 9] View abstract...
39. Backhus LM, Mulligan MS, Ha R, Shriki JE, Mohammed TL. Imaging in lung transplantation: surgical considerations of donor and recipient. *Radiologic Clinics of North America* 2016;54(2):339-353. DOI: 10.1016/j.rcl.2015.09.013. [Context Link 1, 2] View abstract...
40. Pinho DF, Banga A, Torres F, Mathews D. Ventilation perfusion pulmonary scintigraphy in the evaluation of pre-and post-lung transplant patients. *Transplantation Reviews* 2019;33(2):107-114. DOI: 10.1016/j.tre.2018.10.003. [Context Link 1] View abstract...
41. Weber Gulling M, Schaefer M, Bishop-Simo L, Keller BC. Optimizing nutrition assessment to create better outcomes in lung transplant recipients: a review of current practices. *Nutrients* 2019;11(12):Online. DOI: 10.3390/nu11122884. [Context Link 1] View abstract...
42. Jones JM, et al. Assessing solid organ donors and monitoring transplant recipients for human immunodeficiency virus, hepatitis B virus, and hepatitis C virus infection - U.S. Public Health Service guideline, 2020. *MMWR - Recommendations and Reports* 2020;69(4):1-16. DOI: 10.15585/mmwr.rr6904a1. [Context Link 1] View abstract...
43. Magda G. Opportunistic infections post-lung transplantation: viral, fungal, and mycobacterial. *Clinics in Chest Medicine* 2023;44(1):159-177. DOI: 10.1016/j.ccm.2022.10.012. [Context Link 1, 2, 3, 4, 5, 6] View abstract...
44. Natalini JG, Clausen ES. Critical care management of the lung transplant recipient. *Clinics in Chest Medicine* 2023;44(1):105-119. DOI: 10.1016/j.ccm.2022.10.010. [Context Link 1, 2, 3, 4, 5] View abstract...
45. Bernhardt AM, Copeland H, Deswal A, Gluck J, Givertz MM. The International Society for Heart and Lung Transplantation/Heart Failure Society of America Guideline on acute mechanical circulatory support. *Journal of Heart and Lung Transplantation* 2023;42(4):e1-e64. DOI: 10.1016/j.healun.2022.10.028. [Context Link 1, 2] View abstract...
46. Scully BB, Nolley EP, Bush EL. Palliative care in lung transplantation. *Annals of Palliative Medicine* 2022;11(2):927-935. DOI: 10.21037/apm-21-170. [Context Link 1] View abstract...
47. Crespo MM, et al. ISHLT consensus document on lung transplantation in patients with connective tissue disease: Part III: Pharmacology, medical and surgical management of post-transplant extrapulmonary conditions statements. *Journal of Heart and Lung Transplantation* 2021;40(11):1279-1300. DOI: 10.1016/j.healun.2021.07.013. (Reaffirmed 2022 Mar) [Context Link 1, 2, 3] View abstract...
48. Di Nardo M, et al. Postoperative management of lung transplant recipients in the intensive care unit. *Anesthesiology* 2022;136(3):482-499. DOI: 10.1097/ALN.0000000000004054. [Context Link 1, 2, 3, 4, 5] View abstract...
49. Benissan-Messan DZ, et al. Lung transplantation in the septuagenarian can be successfully performed though long-term results impacted by diseases of aging. *Clinical Transplantation* 2022;Online. DOI: 10.1111/ctr.14593. [Context Link 1, 2] View abstract...
50. Drolen C, Cantu E, Goldberg HJ, Diamond JM, Courtwright A. Impact of the elimination of the donation service area on United States lung transplant practices and outcomes at high and low competition centers. *American Journal of Transplantation* 2020;20(12):3631-3638. DOI: 10.1111/ajt.16098. [Context Link 1, 2] View abstract...
51. Gerull WD, et al. Local versus distant lung donor procurement does not influence short-term clinical outcomes. *Journal of Thoracic and Cardiovascular Surgery* 2021;162(4):1284-1293 e. DOI: 10.1016/j.jtcvs.2020.07.115. [Context Link 1, 2] View abstract...
52. Halpern AL, et al. A comparison of frailty measures at listing to predict outcomes after lung transplantation. *Annals of Thoracic Surgery* 2020;109(1):233-240. DOI: 10.1016/j.athoracsur.2019.07.040. [Context Link 1, 2] View abstract...
53. Premier PINC AI™ Healthcare Database (PHD), 01/01/2019-12/31/2022. Premier, Inc. [Context Link 1, 2, 3]

54. Jin Z, Suen KC, Wang Z, Ma D. Review 2: Primary graft dysfunction after lung transplant-pathophysiology, clinical considerations and therapeutic targets. *Journal of Anesthesia* 2020;34(5):729-740. DOI: 10.1007/s00540-020-02823-6. [Context Link 1] View abstract...
55. Husain S, et al. The 2015 International Society for Heart and Lung Transplantation Guidelines for the management of fungal infections in mechanical circulatory support and cardiothoracic organ transplant recipients: Executive summary. *Journal of Heart and Lung Transplantation* 2016;35(3):261-282. DOI: 10.1016/j.healun.2016.01.007. [Context Link 1] View abstract...
56. Levine DJ, Hachem RR. Lung allograft rejection. *Thoracic Surgery Clinics* 2022;32(2):221-229. DOI: 10.1016/j.thorsurg.2021.12.003. [Context Link 1] View abstract...
57. Grewal HS, Thaniyavarn T, Arcasoy SM, Goldberg HJ. Common noninfectious complications following lung transplantation. *Clinics in Chest Medicine* 2023;44(1):179-190. DOI: 10.1016/j.ccm.2022.11.001. [Context Link 1, 2] View abstract...
58. Loor G, Mattar A, Schaheen L, Bremner RM. Surgical complications of lung transplantation. *Thoracic Surgery Clinics* 2022;32(2):197-209. DOI: 10.1016/j.thorsurg.2022.01.003. [Context Link 1, 2, 3] View abstract...
59. Pasupneti S, Dhillon G, Reitz B, Khush K. Combined heart lung transplantation: an updated review of the current literature. *Transplantation* 2017;101(10):2297-2302. DOI: 10.1097/TP.0000000000001820. [Context Link 1] View abstract...
60. Gadre S, Turowski J, Budev M. Overview of lung transplantation, heart-lung transplantation, liver-lung transplantation, and combined hematopoietic stem cell transplantation and lung transplantation. *Clinics in Chest Medicine* 2017;38(4):623-640. DOI: 10.1016/j.ccm.2017.07.004. [Context Link 1, 2] View abstract...
61. Han JL, Beal EW, Mumtaz K, Washburn K, Black SM. Combined liver-lung transplantation: Indications, outcomes, current experience and ethical issues. *Transplantation Reviews* 2019;33(2):99-106. DOI: 10.1016/j.trre.2018.11.002. [Context Link 1] View abstract...
62. Kristensen AW, Mortensen J, Berg RM. Pulmonary thromboembolism as a complication of lung transplantation. *Clinical Transplantation* 2017;31(4):e12922. DOI: 10.1111/ctr.12922. [Context Link 1] View abstract...
63. Barnes H, et al. Atrial flutter and fibrillation following lung transplantation: incidence, associations and a suggested therapeutic algorithm. *Heart, Lung & Circulation* 2020;29(10):1484-1492. DOI: 10.1016/j.hlc.2019.10.011. [Context Link 1] View abstract...
64. Nguyen AP, Gabriel RA, Golts E, Kistler EB, Schmidt U. Severity of acute kidney injury in the post-lung transplant patient is associated with higher healthcare resources and cost. *Journal of Cardiothoracic and Vascular Anesthesia* 2017;31(4):1361-1369. DOI: 10.1053/j.jvca.2017.02.004. [Context Link 1] View abstract...
65. Sher Y, Mooney J, Dhillon G, Lee R, Maldonado JR. Delirium after lung transplantation: Association with recipient characteristics, hospital resource utilization, and mortality. *Clinical Transplantation* 2017;31(5):e12966. DOI: 10.1111/ctr.12966. [Context Link 1] View abstract...
66. Banga A, et al. Hospital length of stay after lung transplantation: Independent predictors and association with early and late survival. *Journal of Heart and Lung Transplantation* 2017;36(3):289-296. DOI: 10.1016/j.healun.2016.07.020. [Context Link 1] View abstract...
67. Keller CA, et al. Utilization and cost analysis of lung transplantation and survival after 10 years of adapting the lung allocation score. *Transplantation* 2019;103(3):638-646. DOI: 10.1097/TP.0000000000002227. [Context Link 1] View abstract...
68. Bigelow B, Toci G, Etchill E, Krishnan A, Merlo C, Bush EL. Nutritional Risk Index: a predictive metric for mortality following lung transplant. *Annals of Thoracic Surgery* 2021;112(1):214-220. DOI: 10.1016/j.athoracsur.2020.08.014. [Context Link 1] View abstract...
69. Bin Saeedan M, Rizk A, Yadav R, Ghosh S. Imaging evaluation of airway complications after lung transplant. *Journal of Computer Assisted Tomography* 2020;44(3):314-327. DOI: 10.1097/RCT.0000000000000996. [Context Link 1] View abstract...
70. Jorge A, et al. Routine deep vein thrombosis screening after lung transplantation: Incidence and risk factors. *Journal of Thoracic and Cardiovascular Surgery* 2020;159(3):1142-1150. DOI: 10.1016/j.jtcvs.2019.08.030. [Context Link 1] View abstract...
71. Kotecha S, Ivulich S, Snell G. Review: immunosuppression for the lung transplant patient. *Journal of Thoracic Disease* 2021;13(11):6628-6644. DOI: 10.21037/jtd-2021-11. [Context Link 1] View abstract...
72. Ghadimi K, et al. Inhaled pulmonary vasodilator therapy in adult lung transplant: a randomized clinical trial. *JAMA Surgery* 2022;157(1):Online. DOI: 10.1001/jamasurg.2021.5856. [Context Link 1] View abstract...
73. Hudson T. The role of social determinates of health in discharge practices. *Nursing Clinics of North America* 2021;56(3):369-378. DOI: 10.1016/j.cnur.2021.04.004. [Context Link 1, 2] View abstract...
74. Mayer RS, Noles A, Vinh D. Determination of postacute hospitalization level of care. *Medical Clinics of North America* 2020;104(2):345-357. DOI: 10.1016/j.mcna.2019.10.011. [Context Link 1] View abstract...
75. Centers for Medicare and Medicaid Services. "Condition of participation: Discharge planning." 42 CFR Pt. 482.43. Washington, DC 2022 Oct [accessed 2023 Oct 30] Accessed at: <https://www.ecfr.gov/>. [Context Link 1, 2, 3]
76. Mondor EE. Lower respiratory problems. In: Harding MM, Kwong J, Hagler D, Reinisch C, editors. *Lewis's Medical-Surgical Nursing: Assessment and Management of Clinical Problems*. 12th ed. St. Louis, MO: Mosby; 2023:596-631. [Context Link 1, 2, 3, 4, 5]
77. Adverse drug reactions and medication errors. In: Burchum JR, Rosenthal LD, editors. *Lehne's Pharmacology for Nursing Care*. 11th ed. Elsevier; 2022:63-73. [Context Link 1, 2]
78. Hematologic and immunologic care plans. In: Gulanick M, Myers JL, editors. *Nursing Care Plans*. 10th ed. Elsevier; 2022:739-767. [Context Link 1, 2]
79. Bonaduce SJ. Assessment: respiratory system. In: Harding MM, Kwong J, Hagler D, Reinisch C, editors. *Lewis's Medical-Surgical Nursing: Assessment and Management of Clinical Problems*. 12th ed. St. Louis, MO: Mosby; 2023:515-538. [Context Link 1]
80. Saleeby J. Communication and collaboration. In: Perry AG, Potter PA, Ostendorf WR, editors. *Nursing Interventions and Clinical Skills*. 7th ed. Elsevier; 2020:9-21. [Context Link 1, 2]
81. Elements of Excellence in Transitions of Care (TOC). TOC Checklist [Internet] National Transitions of Care Coalition. Accessed at: <https://www.ntocc.org/> [accessed 2023 Aug 10] [Context Link 1]

82. Medical-surgical nursing. In: Hinkle JL, Cheever KH, Overbaugh KJ, editors. *Brunner & Suddarth's Textbook of Medical-Surgical Nursing*. 15th ed. Wolters Kluwer; 2022:33-55. [Context Link 1]
83. Transitions of care. In: Gillingham DC, editor. *Foundations of Case Management*. Blue Bayou Press; 2021:137-144. [Context Link 1]
84. Becker C, et al. Interventions to improve communication at hospital discharge and rates of readmission: a systematic review and meta-analysis. *JAMA Network Open* 2021;4(8):e2119346. DOI: 10.1001/jamanetworkopen.2021.19346. [Context Link 1] View abstract...
85. The nursing process in drug therapy and patient safety. In: Tucker RG, editor. *Karch's Focus on Nursing Pharmacology*. 9th ed. Wolters Kluwer; 2023:47-57. [Context Link 1]
86. Ostendorf WR. Preparation for safe medication administration. In: Perry AG, Potter PA, Ostendorf WR, editors. *Nursing Interventions and Clinical Skills*. 7th ed. Elsevier; 2020:551-567. [Context Link 1]
87. The case manager's roles, functions, and activities. In: Gillingham DC, editor. *Foundations of Case Management*. Blue Bayou Press; 2021:9-14. [Context Link 1]
88. Whiteson JH, Cohen JM, Prilik S. Chronic medical conditions: pulmonary disease, organ transplantation, and diabetes. In: Cifu DX, et al., editors. *Braddom's Physical Medicine and Rehabilitation*. 6th ed. Philadelphia, PA: Elsevier; 2021:535-567.e5. [Context Link 1]
89. Nursing diagnoses in alphabetical order. In: Doenges ME, Moorhouse MF, Murr AC, editors. *Nursing Diagnosis Manual: Planning, Individualizing, and Documenting Client Care*. 7th ed. F.A. Davis Company; 2022:40-1095. [Context Link 1]
90. Blitzer D, et al. Long term survival after lung transplantation: A single center experience. *Journal of Cardiac Surgery* 2020;35(2):273-278. DOI: 10.1111/jocs.14163. [Context Link 1] View abstract...
91. Management of patients with dermatologic disorders. In: Hinkle JL, Cheever KH, Overbaugh KJ, editors. *Brunner & Suddarth's Textbook of Medical-Surgical Nursing*. 15th ed. Wolters Kluwer; 2022:1815-1864. [Context Link 1]
92. Centers for Medicare and Medicaid Services. "Admissions." 42 CFR 412.3 Washington, DC 2023 Jul [accessed 2023 Oct 30] Accessed at: <https://www.ecfr.gov/current/title-42/chapter-IV/subchapter-B/part-412>. [Context Link 1]
93. Centers for Medicare and Medicaid Services. "Hospital services excluded from payment under the hospital outpatient prospective payment system." 42 CFR 419.22 Washington, DC 2023 Jul [accessed 2023 Oct 30] Accessed at: <https://www.ecfr.gov/>. [Context Link 1]
94. Centers for Medicare and Medicaid Services. "Requirements relating to basic benefits." 42 CFR 422.101 Washington, DC 2023 Jun 05 [accessed 2023 Oct 30] Accessed at: <https://www.ecfr.gov/>. [Context Link 1]
95. Centers for Medicare and Medicaid Services. Medicare Benefit Policy Manual. Chapter 1 - Inpatient Hospital Services Covered Under Part A Rev. 10892 [Internet] Centers for Medicare and Medicaid Services. 2021 Aug Accessed at: <https://www.cms.gov/manuals/>. [accessed 2023 Sep 08] [Context Link 1]
96. Centers for Medicare and Medicaid Services. Medicare Benefit Policy Manual. Chapter 6 - Hospital Services Covered Under Part B Rev. 10541 [Internet] Centers for Medicare and Medicaid Services. 2020 Dec Accessed at: <http://www.cms.gov/manuals/>. [accessed 2023 Nov 09] [Context Link 1]
97. Centers for Medicare and Medicaid Services. Medicare Benefit Policy Manual. Chapter 15 - Covered Medical and Other Health Services Rev. 12171 [Internet] Centers for Medicare and Medicaid Services. 2023 Aug 03 Accessed at: <https://www.cms.gov/Regulations-and-Guidance/Guidance/Manuals/>. [accessed 2023 Oct 25] [Context Link 1]
98. Centers for Medicare and Medicaid Services. Medicare Program Integrity Manual. Chapter 6, Section 6.5 - Medical Review of Inpatient Hospital Claims for Part A Payment Rev. 10365 [Internet] Centers for Medicare and Medicaid Services. 2020 Oct 02 Accessed at: <https://www.cms.gov/regulations-and-guidance/regulations-and-guidance>. [accessed 2023 Oct 20] [Context Link 1]
99. Medicare Coverage Database. [Internet] Centers for Medicare and Medicaid Services. Accessed at: <https://www.cms.gov/medicare-coverage-database/search.aspx>? Updated 2023 [accessed 2023 Oct 20] [Context Link 1]

Footnotes

[A] The BODE index includes the variables of BMI, airflow obstruction, dyspnea, and exercise capacity as assessed by 6-minute walking distance.(2)(4) An online calculator is available at <https://www.mdcalc.com/bode-index-copd-survival>. [A in Context Link 1]

[B] World Health Organization functional class IV describes patients with pulmonary hypertension who have signs of right-sided heart failure, symptoms of dyspnea at rest, and more severe symptoms with activity. [B in Context Link 1, 2, 3, 4]

[C] Interstitial lung disease refers to a category of nonmalignant diseases of the lung parenchyma characterized by inflammation, fibrosis, and destruction of the alveoli. These disorders can be secondary to inhalation injury, prior infection, connective tissue disease, or toxin exposure, but most cases are idiopathic (eg, idiopathic pulmonary fibrosis).(7)(8) [C in Context Link 1]

[D] Patients with pulmonary hypertension may be candidates for combined heart-lung transplant. Specific criteria include World Health Organization functional class IV symptoms, a cardiac index less than 2 L/min/m², and a mean right atrial pressure above 15 mm Hg.(1) [D in Context Link 1]

[E] The European Society of Cardiology/European Respiratory Society (ESC/ERS) high-risk category for pulmonary hypertension is predictive of a 1-year mortality greater than 10% and is characterized by clinical signs of right-sided heart failure with symptom progression, syncope, World Health Organization (WHO) functional class IV, echocardiogram findings of enlarged right atrium and pericardial effusion, and has severity criteria for 6-minute walk distance, cardiopulmonary exercise testing results, BNP, and hemodynamic data.(10) The REVEAL score includes variables for WHO pulmonary hypertension group, presence of renal insufficiency, WHO functional class, systemic and pulmonary artery pressures, BNP, and pulmonary function. A score greater than 10 is associated with a 1-year

mortality greater than 15%. A calculator for the REVEAL score is available at <https://www.mdcalc.com/reveal-registry-risk-score-pulmonary-arterial-hypertension-pah>. [E in Context Link 1]

[F] World Health Organization functional class III describes patients who have significant physical activity limitation due to pulmonary hypertension. Although asymptomatic at rest, they develop symptoms of dyspnea with less than ordinary activity. [F in Context Link 1, 2, 3]

[G] Langerhans cell histiocytosis is also known as histiocytosis X and eosinophilic granuloma. [G in Context Link 1]

[H] The latest International Society of Heart and Lung Transplantation (ISHLT) guideline has no formal recommendations regarding lung transplant criteria for patients with fulminant lung damage caused by COVID-19 infection; however, broad criteria that have been proposed by the ISHLT and applied at transplant centers include radiographic and clinical evidence of severe irreversible disease after allowing at least 28 days for possible recovery, absence of current infection (eg, 2 negative PCR tests by endotracheal aspirate or bronchoscopy 24 to 48 hours apart), single organ failure, and patient is otherwise appropriate for transplant.(1)(17)(18) Analysis of 400 patients undergoing lung transplant for COVID-19 respiratory failure (64% acute respiratory distress syndrome) from the UNOS registry showed a 1-year survival of 86.6%, which is comparable to the 86.3% survival for patients transplanted for other indications.(19) [H in Context Link 1]

[I] Patients with an irreversible acute lung injury, including those requiring mechanical ventilation or extracorporeal life support, may be candidates for transplant if they have no other acute organ dysfunction.(1)(20)(21)(24) [I in Context Link 1]

[J] Some relative contraindications, such as coronary artery disease, can be corrected prior to or during transplant.(25) [J in Context Link 1]

[K] See Clinical Indications for Procedure in this guideline. [K in Context Link 1]

[L] Enteral feeding and oral intake may be delayed in patients with connective tissue disease who have gastrointestinal involvement.(47) [L in Context Link 1]

[M] Some patients may have their hydration and nutritional needs met via alternative means (eg, percutaneous endoscopic gastrostomy or jejunostomy tube), especially those with connective tissue diseases that pose a high risk of postoperative aspiration.(47) [M in Context Link 1]

[N] Indications for postoperative extracorporeal membrane oxygenation include pulmonary hypertension with right ventricular dysfunction, primary graft dysfunction, or high risk for developing primary graft dysfunction (eg, lobar lung transplant, donor size discrepancy).(22) [N in Context Link 1]

[O] Discharge instructions should be given in the patient's and caregiver's native language using trained language interpreters whenever possible.(73) [O in Context Link 1]

Definitions

Acute kidney injury (stage 2)

- Acute kidney injury (stage 2) with significant clinical findings, as indicated by **ALL** of the following(1)(2)(3):
 - Acute[A] kidney injury (stage 2), as indicated by **1 or more** of the following[B]:
 - Within past 7 days, rise in serum creatinine between 2 and 2.9 times its baseline value (increase of 3 times baseline would qualify as stage 3 acute kidney injury)
 - Decreased urine output,[C] as indicated by **ALL** of the following:
 - Assessment of urine output appropriate (eg, adequate volume status, no urinary obstruction)
 - Urine output less than 0.5 mL/kg/hr for 12 hours
 - Significant clinical findings, as indicated by **1 or more** of the following[B]:
 - Hemodynamic instability
 - Worsening clinical status despite observation care (eg, rising creatinine or urine output remains less than 0.5 mL/kg/hr)
 - Altered mental status
 - Cardiac arrhythmias of immediate concern
 - Severe hypertension (SBP greater than 180 mm Hg or DBP greater than 120 mm Hg in adults or greater than the 95th percentile for age, gender, and height plus 30 mm Hg in pediatric patients)(4)(5)
 - Significant respiratory findings (eg, pulmonary edema) that are severe (eg, Hypoxemia) or persist despite observation care
 - Clinically significant electrolyte abnormality that is severe (eg, hyperkalemia with severe ECG findings)[D] or persists despite observation care
 - Clinically significant metabolic abnormality (eg, metabolic acidosis) that is severe or persists despite observation care

References

1. Kidney Disease: Improving Global Outcomes (KDIGO) Acute Kidney Injury Work Group. KDIGO clinical practice guideline for acute kidney injury. Kidney International. Supplement 2012;2(1):1-138. (Reaffirmed 2023 May)

2. Weisbord SD, Palevsky PM. Prevention and management of acute kidney injury. In: Yu AS, Chertow GM, Luyckx VA, Marsden PA, Skorecki K, Taal MW, editors. Brenner and Rector's The Kidney. 11th ed. Philadelphia, PA: Elsevier; 2020:940-977.e15.
3. Palevsky PM, et al. KDOQI US commentary on the 2012 KDIGO clinical practice guideline for acute kidney injury. American Journal of Kidney Diseases 2013;61(5):649-72. DOI: 10.1053/j.ajkd.2013.02.349.
4. Whelton PK, et al. 2017 ACC/AHA/AAPA/ABC/ACPM/AGS/APhA/ASH/ASPC/NMA/PCNA Guideline for the prevention, detection, evaluation, and management of high blood pressure in adults: a report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines. Circulation 2018;138(17):e484-e594. DOI: 10.1161/CIR.0000000000000596. (Reaffirmed 2023 Jan)
5. Flynn JT, et al. Clinical practice guideline for screening and management of high blood pressure in children and adolescents. Pediatrics 2017;140(3):e20171904. DOI: 10.1542/peds.2017-1904. (Reaffirmed 2022 Aug)
6. Mirvis DM, Goldberger AL. Electrocardiography. In: Libby P, Bonow RO, Mann DL, Tomaselli GF, Bhatt DL, Solomon SD, editors. Braunwald's Heart Disease: A Textbook of Cardiovascular Medicine. 12th ed. Elsevier; 2022:141-174.e3.
7. Mount DB. Fluid and electrolyte disturbances. In: Loscalzo J, Fauci A, Kasper D, Hauser S, Longo D, Jameson JL, editors. Harrison's Principles of Internal Medicine. 21st ed. McGraw Hill Education; 2022:338-356.

Footnotes

- A. A specialty society practice guideline defines acute kidney injury as a change in renal function that is known or presumed to have occurred over the previous 7 days.(1)
- B. In determining the stage of acute kidney injury (eg, stage 2 vs stage 3), patients should be staged according to the criteria that give them the highest stage.(1)
- C. The use of urine output as an indicator of acute kidney injury is important, as change in urine output can occur before a noticeable rise in creatinine.(1)(3) It is also suggested that urine output can identify patients with acute kidney injury that may otherwise be missed.(1)(3) However, urine output as a metric is only valid under the appropriate conditions. For example, the patient must not be hypovolemic, there must not be urinary obstruction, and the collection and measurement of urinary output must be complete and accurate.(1)(3) Use of urinary output to define acute kidney injury can be inaccurate in obese patients due to the nonlinear relationship between body weight and urine output. For example, in a 100-kg (220-lb) patient, urine output of 45 mL/hour over 12 hours (adequate output) would qualify as stage 2 acute kidney injury.(1)(3)
- D. ECG findings include peaked T-waves, PR interval greater than 0.20 seconds, QRS interval greater than 0.12 seconds, and arrhythmia.(6)(7)

Acute renal failure (stage 3 acute kidney injury)

- Acute[A] kidney injury (stage 3),[B] as indicated by **1 or more** of the following[C](1)(2)(3):
 - Within past 7 days, rise in serum creatinine to 3 times its baseline value or higher
 - Within past 7 days, rise in serum creatinine to at least 1.5 times its baseline value (increase of 50%) and serum creatinine is greater than 4 mg/dL (354 micromoles/L)
 - In child younger than 18 years, estimated glomerular filtration rate less than 35 mL/min/1.73m² (0.59 mL/sec/1.73m²)
-  eGFR - Pediatric Calculator and **1 or more** of the following:
 - Within past 7 days, rise in serum creatinine to at least 1.5 times its baseline value (increase of 50%)
 - Within past 48 hours, rise in serum creatinine greater than 0.3 mg/dL (26.5 micromoles/L)
- Decreased urine output,[D] as indicated by **ALL** of the following:
 - Assessment of urine output appropriate (eg, adequate volume status, no urinary obstruction)
 - Oligoanuria, as indicated by **1 or more** of the following:
 - Urine output less than 0.3 mL/kg/hour for 24 hours
 - Anuria (urine output less than 0.1 mL/kg/hour) for 12 hours
- Initiation of renal replacement therapy (RRT) required(4)

References

1. Kidney Disease: Improving Global Outcomes (KDIGO) Acute Kidney Injury Work Group. KDIGO clinical practice guideline for acute kidney injury. Kidney International. Supplement 2012;2(1):1-138. (Reaffirmed 2023 May)
2. Weisbord SD, Palevsky PM. Prevention and management of acute kidney injury. In: Yu AS, Chertow GM, Luyckx VA, Marsden PA, Skorecki K, Taal MW, editors. Brenner and Rector's The Kidney. 11th ed. Philadelphia, PA: Elsevier; 2020:940-977.e15.
3. Palevsky PM, et al. KDOQI US commentary on the 2012 KDIGO clinical practice guideline for acute kidney injury. American Journal of Kidney Diseases 2013;61(5):649-72. DOI: 10.1053/j.ajkd.2013.02.349.
4. Gaudry S, Palevsky PM, Dreyfuss D. Extracorporeal kidney-replacement therapy for acute kidney injury. New England Journal of Medicine 2022;386(10):964-975. DOI: 10.1056/NEJMra2104090.

Footnotes

- A. A specialty society practice guideline defines acute kidney injury as a change in renal function that is known or presumed to have occurred over the previous 7 days.(1)

- B. For purposes of these criteria, the term acute renal failure can be viewed as stage 3 acute kidney injury in the KDIGO classification system.(1) Of note, in the ICD-10 diagnostic coding system (main coding scheme used in labeling diagnoses), the relevant codes covering the clinical entity of significant acute kidney injury use the term acute kidney failure (eg, code N17.9).
- C. In determining the stage of acute kidney injury (eg, stage 2 vs stage 3), patients should be staged according to the criteria that give them the highest stage.(1)
- D. The use of urine output as an indicator of acute kidney injury is important, as change in urine output can occur before a noticeable rise in creatinine.(1)(3) It is also suggested that urine output can identify patients with acute kidney injury that may otherwise be missed.(1) (3) However, urine output as a metric is only valid under the appropriate conditions. For example, the patient must not be hypovolemic, there must not be urinary obstruction, and the collection and measurement of urinary output must be complete and accurate.(1)(3) Use of urinary output to define acute kidney injury can be inaccurate in obese patients due to the nonlinear relationship between body weight and urine output. For example, in a 100-kg (220-lb) patient, urine output of 45 mL/hour over 12 hours (adequate output) would qualify as stage 2 acute kidney injury.(1)(3)

Altered mental status

- Altered mental status (ie, different from baseline), as indicated by **1 or more** of the following(1)(2)(3)(4):
 - Confusional state (eg, disorientation, difficulty following commands, deficit in attention)
 - Lethargy (eg, awake or arousable, but with drowsiness; reduced awareness of self and environment)
 - Obtundation (ie, arousable only with strong stimuli, lessened interest in environment, slowed responses to stimulation)
 - Stupor (ie, may be arousable but patient does not return to normal baseline level of awareness)
 - Coma (ie, not arousable)

References

- Maher PJ. Confusion. In: Walls RM, editor. Rosen's Emergency Medicine: Concepts and Clinical Practice. 10th ed. Philadelphia, PA 19103-2899: Elsevier; 2023:126-131.e1.
- Lei C, Smith C. Depressed consciousness and coma. In: Walls RM, editor. Rosen's Emergency Medicine: Concepts and Clinical Practice. 10th ed. Philadelphia, PA 19103-2899: Elsevier; 2023:117-125.e1.
- Abraham G, Maher PJ. Delirium and dementia. In: Walls RM, editor. Rosen's Emergency Medicine: Concepts and Clinical Practice. 10th ed. Philadelphia, PA 19103-2899: Elsevier; 2023:1269-1280.e2.
- Schor NF. Neurologic evaluation. In: Kliegman RM, St. Geme JW, Blum NJ, Shah SS, Tasker RC, Wilson KM, editors. Nelson Textbook of Pediatrics. 21st ed. Philadelphia, PA: Elsevier; 2020:3053-3063.e1.

Cardiac arrhythmias of immediate concern

- Cardiac arrhythmias or findings of immediate concern, as indicated by **1 or more** of the following(1)(2)(3):
 - Heart rhythms that are inherently dangerous or unstable, as indicated by **1 or more** of the following(4)(5)(6):
 - Resuscitated ventricular fibrillation or cardiac arrest
 - Ventricular escape rhythm
 - Sustained ventricular tachycardia (30 seconds or more of ventricular rhythm at greater than 100 beats per minute)
 - Nonsustained ventricular tachycardia and **1 or more** of the following:
 - Suspected cardiac ischemia as cause or consequence of ventricular tachycardia
 - Acute myocarditis
 - Unstable cardiac conduction defects, as indicated by **1 or more** of the following(6)(7)(8):
 - Type II second-degree atrioventricular block
 - Third-degree atrioventricular block
 - New-onset left bundle branch block with suspected myocardial ischemia
 - Any heart rhythm and **1 or more** of the following(4)(5)(9)(10)(11):
 - Continuous long-term ECG monitoring needed (eg, initiation of drug requiring monitoring beyond observation care)
 - Patient has automatic implanted cardioverter-defibrillator that is repeatedly firing, malfunctioning, or in need of immediate adjustment of settings beyond scope of observation care.
 - Heart rhythms of concern due to **1 or more** of the following:
 - Hypotension
 - Respiratory distress
 - Association with other significant symptoms (eg, syncope)(9)(10)(12)

References

- Curtis AB, Tomaselli GF. Approach to the patient with cardiac arrhythmias. In: Libby P, Bonow RO, Mann DL, Tomaselli GF, Bhatt DL, Solomon SD, editors. Braunwald's Heart Disease: A Textbook of Cardiovascular Medicine. 12th ed. Elsevier; 2022:1145-1162.e7.
- Dalal AS, Van Hare GF. Disturbances of rate and rhythm of the heart. In: Kliegman RM, St. Geme JW, Blum NJ, Shah SS, Tasker RC, Wilson KM, editors. Nelson Textbook of Pediatrics. 21st ed. Philadelphia, PA: Elsevier; 2020:2436-2447.e1.
- Joglar JA, et al. 2023 HRS Expert Consensus Statement on the management of arrhythmias during pregnancy. Heart Rhythm 2023;20(10):Online. DOI: 10.1016/j.hrthm.2023.05.017.

4. El-Chami M, Fernando L. Ventricular arrhythmias. In: McKean SC, Ross JJ, Dressler DD, Scheurer DB, editors. Principles and Practice of Hospital Medicine. 2nd ed. New York, NY: McGraw-Hill Education; 2017:1003-1014.
5. Al-Khatib SM, et al. 2017 AHA/ACC/HRS guideline for management of patients with ventricular arrhythmias and the prevention of sudden cardiac death: a report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines and the Heart Rhythm Society. *Circulation* 2018;138(13):e272-e391. DOI: 10.1161/CIR.0000000000000549. (Reaffirmed 2023 Jul)
6. Epstein AE, et al. 2012 ACCF/AHA/HRS focused update incorporated into the ACCF/AHA/HRS 2008 guidelines for device-based therapy of cardiac rhythm abnormalities: a report of the American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines and the Heart Rhythm Society. *Circulation* 2013;127(3):e283-e352. DOI: 10.1161/CIR.0b013e318276ce9b. (Reaffirmed 2022 Jun)
7. Miller JM, Ellenbogen KA. Therapy for cardiac arrhythmias. In: Libby P, Bonow RO, Mann DL, Tomaselli GF, Bhatt DL, Solomon SD, editors. Braunwald's Heart Disease: A Textbook of Cardiovascular Medicine. 12th ed. Elsevier; 2022:1208-1244.
8. Park DS, Fishman GI. The cardiac conduction system. *Circulation* 2011;123(8):904-15. DOI: 10.1161/CIRCULATIONAHA.110.942284.
9. Joglar JA, et al. 2023 ACC/AHA/ACCP/HRS Guideline for the diagnosis and management of atrial fibrillation: a report of the American College of Cardiology/American Heart Association joint committee on clinical practice guidelines. *Journal of the American College of Cardiology* 2023 Nov;online. DOI: 10.1016/j.jacc.2023.08.017. (Reaffirmed 2023 Nov 30)
10. Chun EB, McGorisk GM. Supraventricular tachyarrhythmias. In: McKean SC, Ross JJ, Dressler DD, Scheurer DB, editors. Principles and Practice of Hospital Medicine. 2nd ed. New York, NY: McGraw-Hill Education; 2017:980-995.
11. Hoskins MH, De Lurgio DB. Pacemakers, defibrillators, and cardiac resynchronization devices in hospital medicine. In: McKean SC, Ross JJ, Dressler DD, Scheurer DB, editors. Principles and Practice of Hospital Medicine. 2nd ed. New York, NY: McGraw-Hill Education; 2017:1025-34.
12. Westerman S, Manogue M, Hoskins MH. Bradycardia. In: McKean SC, Ross JJ, Dressler DD, Scheurer DB, editors. Principles and Practice of Hospital Medicine. 2nd ed. New York, NY: McGraw-Hill Education; 2017:996-1002.

Dangerous arrhythmia absent

- Dangerous arrhythmia absent, as indicated by absence of **ALL** of the following(1)(2)(3)(4)(5)(6)(7)(8)(9)(10)(11):
 - Resuscitated ventricular fibrillation or cardiac arrest
 - Ventricular escape rhythm
 - Sustained (30 seconds or more of ventricular rhythm at greater than 100 beats per minute) ventricular tachycardia
 - Nonsustained ventricular tachycardia with myocarditis or ischemia
 - Type II second-degree atrioventricular block
 - Third-degree atrioventricular block
 - New-onset left bundle branch block with suspected myocardial ischemia
 - Heart rhythms of concern due to association with Hypotension, Respiratory distress, or other significant symptoms (eg, syncope)

References

1. Curtis AB, Tomaselli GF. Approach to the patient with cardiac arrhythmias. In: Libby P, Bonow RO, Mann DL, Tomaselli GF, Bhatt DL, Solomon SD, editors. Braunwald's Heart Disease: A Textbook of Cardiovascular Medicine. 12th ed. Elsevier; 2022:1145-1162.e7.
2. Dalal AS, Van Hare GF. Disturbances of rate and rhythm of the heart. In: Kliegman RM, St. Geme JW, Blum NJ, Shah SS, Tasker RC, Wilson KM, editors. Nelson Textbook of Pediatrics. 21st ed. Philadelphia, PA: Elsevier; 2020:2436-2447.e1.
3. El-Chami M, Fernando L. Ventricular arrhythmias. In: McKean SC, Ross JJ, Dressler DD, Scheurer DB, editors. Principles and Practice of Hospital Medicine. 2nd ed. New York, NY: McGraw-Hill Education; 2017:1003-1014.
4. Al-Khatib SM, et al. 2017 AHA/ACC/HRS guideline for management of patients with ventricular arrhythmias and the prevention of sudden cardiac death: a report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines and the Heart Rhythm Society. *Circulation* 2018;138(13):e272-e391. DOI: 10.1161/CIR.0000000000000549. (Reaffirmed 2023 Jul)
5. Epstein AE, et al. 2012 ACCF/AHA/HRS focused update incorporated into the ACCF/AHA/HRS 2008 guidelines for device-based therapy of cardiac rhythm abnormalities: a report of the American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines and the Heart Rhythm Society. *Circulation* 2013;127(3):e283-e352. DOI: 10.1161/CIR.0b013e318276ce9b. (Reaffirmed 2022 Jun)
6. Miller JM, Ellenbogen KA. Therapy for cardiac arrhythmias. In: Libby P, Bonow RO, Mann DL, Tomaselli GF, Bhatt DL, Solomon SD, editors. Braunwald's Heart Disease: A Textbook of Cardiovascular Medicine. 12th ed. Elsevier; 2022:1208-1244.
7. Park DS, Fishman GI. The cardiac conduction system. *Circulation* 2011;123(8):904-15. DOI: 10.1161/CIRCULATIONAHA.110.942284.
8. Joglar JA, et al. 2023 ACC/AHA/ACCP/HRS Guideline for the diagnosis and management of atrial fibrillation: a report of the American College of Cardiology/American Heart Association joint committee on clinical practice guidelines. *Journal of the American College of Cardiology* 2023 Nov;online. DOI: 10.1016/j.jacc.2023.08.017. (Reaffirmed 2023 Nov 30)
9. Chun EB, McGorisk GM. Supraventricular tachyarrhythmias. In: McKean SC, Ross JJ, Dressler DD, Scheurer DB, editors. Principles and Practice of Hospital Medicine. 2nd ed. New York, NY: McGraw-Hill Education; 2017:980-995.
10. Westerman S, Manogue M, Hoskins MH. Bradycardia. In: McKean SC, Ross JJ, Dressler DD, Scheurer DB, editors. Principles and Practice of Hospital Medicine. 2nd ed. New York, NY: McGraw-Hill Education; 2017:996-1002.
11. Joglar JA, et al. 2023 HRS Expert Consensus Statement on the management of arrhythmias during pregnancy. *Heart Rhythm* 2023;20(10):Online. DOI: 10.1016/j.hrthm.2023.05.017.

Hemodynamic instability

- Hemodynamic instability, as indicated by **1 or more** of the following⁽¹⁾⁽²⁾⁽³⁾⁽⁴⁾⁽⁵⁾⁽⁶⁾:
 - Vital sign abnormality not readily corrected by appropriate treatment, as indicated by **1 or more** of the following^[A]:
 - Tachycardia that persists despite appropriate treatment (eg, volume repletion, treatment of pain, treatment of underlying cause)
 - Hypotension that persists despite appropriate treatment (eg, volume repletion, treatment of underlying cause)
 - Orthostatic hypotension that persists despite appropriate treatment (eg, volume repletion)
 - Hypotension that is severe, as indicated by **ALL** of the following:
 - Hypotension
 - Severe hypotension, as indicated by **1 or more** of the following:
 - Lactate of 2.0 mmol/L (18 mg/dL) or more secondary to hypotension (ie, hypoperfusion)^[B](3)(6)
 - Metabolic acidosis (arterial or venous^[C] pH less than 7.35) not otherwise explained
 - Mean arterial pressure^{[A][D]} less than 65 mm Hg
 - IV inotropic or vasopressor medication required to maintain adequate blood pressure or perfusion

References

1. Puskas MA, Jones AE. Shock. In: Walls RM, editor. Rosen's Emergency Medicine: Concepts and Clinical Practice. 10th ed. Philadelphia, PA 19103-2899: Elsevier; 2023:34-41.e1.
2. Lewis J, Patel B. Shock. In: Gershel JC, Rauch DA, editors. Caring for the Hospitalized Child: A Handbook of Inpatient Pediatrics. 3rd ed. Elk Grove Village, IL: American Academy of Pediatrics; 2023:107-114.
3. Singer M, et al. The Third International Consensus definitions for sepsis and septic shock (Sepsis-3). Journal of the American Medical Association 2016;315(8):801-810. DOI: 10.1001/jama.2016.0287.
4. Turner DA, Cheifetz IM. Shock. In: Kliegman RM, St. Geme JW, Blum NJ, Shah SS, Tasker RC, Wilson KM, editors. Nelson Textbook of Pediatrics. 21st ed. Philadelphia, PA: Elsevier; 2020:572-583.e1.
5. Nyhan A. Cardiology. In: Kleinman K, McDaniel L, Molloy M, editors. The Harriet Lane Handbook: A Manual for Pediatric House Officers. 22nd ed. 202: Elsevier; 2021:145-188.e4.
6. Naidu SS, et al. SCAI SHOCK stage classification expert consensus update: a review and incorporation of validation studies: this statement was endorsed by the American College of Cardiology (ACC), American College of Emergency Physicians (ACEP), American Heart Association (AHA), European Society of Cardiology (ESC) Association for Acute Cardiovascular Care (ACVC), International Society for Heart and Lung Transplantation (ISHLT), Society of Critical Care Medicine (SCCM), and Society of Thoracic Surgeons (STS) in December 2021. Journal of the American College of Cardiology 2022;Online. DOI: 10.1016/j.jacc.2022.01.018.
7. Wardi G, Brice J, Correia M, Liu D, Self M, Tainter C. Demystifying lactate in the emergency department. Annals of Emergency Medicine 2020;75(2):287-298. DOI: 10.1016/j.annemergmed.2019.06.027.
8. Levitt DG, Levitt JE, Levitt MD. Quantitative assessment of blood lactate in shock: measure of hypoxia or beneficial energy source. BioMed Research International 2020;2020:2608318. DOI: 10.1155/2020/2608318.
9. Bakker J, Postelnicu R, Mukherjee V. Lactate: where are we now? Critical Care Clinics 2020;36(1):115-124. DOI: 10.1016/j.ccc.2019.08.009.
10. Kraut JA, Madias NE. Lactic acidosis. New England Journal of Medicine 2014;371(24):2309-2319. DOI: 10.1056/NEJMr1309483.
11. Olivieri L, Chasm R. Diabetic ketoacidosis in the pediatric emergency department. Emergency Medicine Clinics of North America 2013;31(3):755-773. DOI: 10.1016/j.emc.2013.05.004.
12. Maloney GE, Glauser JM. Diabetes mellitus and disorders of glucose homeostasis. In: Walls RM, editor. Rosen's Emergency Medicine: Concepts and Clinical Practice. 10th ed. Philadelphia, PA 19103-2899: Elsevier; 2023:1543-1558.e2.
13. Arnold TD, Miller M, van Wessem KP, Evans JA, Balogh ZJ. Base deficit from the first peripheral venous sample: a surrogate for arterial base deficit in the trauma bay. Journal of Trauma 2011;71(4):793-7; discussion 797. DOI: 10.1097/TA.0b013e31822ad694.
14. Malinoski DJ, Todd SR, Slone S, Mullins RJ, Schreiber MA. Correlation of central venous and arterial blood gas measurements in mechanically ventilated trauma patients. Archives of Surgery 2005;140(11):1122-5. DOI: 10.1001/archsurg.140.11.1122.
15. Zakrisson T, McFarlan A, Wu YY, Keshet I, Nathens A. Venous and arterial base deficits: do these agree in occult shock and in the elderly? A Bland-Altman analysis. Journal of Trauma and Acute Care Surgery 2013;74(3):936-9. DOI: 10.1097/TA.0b013e318282747a.

Footnotes

- A. Criteria based upon clinician-acquired numeric values (eg, vital signs, oxygen saturation) should be used if they are accurate reflections of the patient's condition. Transitory findings (eg, abnormal only upon initial emergency department intake or only one time out of multiple readings) that rapidly improve with no or minimal treatment usually do not reflect disease severity or risk for deterioration. This does not imply that an initial or one-time reading cannot ever be applicable. The goal is to separate erroneous or incidental findings from those that truly represent the patient's clinical picture.
- B. There are numerous causes of an elevated lactate level. The most common are cardiogenic or hypovolemic shock, severe heart failure, severe trauma, or sepsis. However, there are also other etiologies to consider such as vigorous exercise, seizures, liver disease, or medication use (eg, metformin, beta2-agonists); therefore, interpretation of elevated lactate levels requires consideration of the clinical context (eg, well-appearing post exercise vs hypotensive). The severity and persistence of the elevation can sometimes be helpful in this differentiation. In most instances of lactic acidosis, blood pH is less than 7.35, with a serum bicarbonate level 20 mEq/L (mmol/L) or lower. However, a coexisting respiratory or metabolic alkalosis can mask these findings.⁽⁶⁾⁽⁷⁾⁽⁸⁾⁽⁹⁾⁽¹⁰⁾

- C. Analyses have concluded that venous and arterial pH measurements are highly correlated, with small differences between them that are usually not clinically significant.(11)(12)(13)(14)(15) If a mixed acid-base disorder is suspected, an arterial blood gas is indicated. (12)
- D. The mean arterial pressure takes into account both systolic and diastolic blood pressure readings and is calculated as mean arterial pressure (MAP) equals $1/3 \text{ SBP} + 2/3 \text{ DBP}$.(1)(4) A calculator is available at <https://www.mdcalc.com/mean-arterial-pressure-map>.

Hemodynamic stability

- Hemodynamic stability, as indicated by **1 or more** of the following:
 - Hemodynamic abnormalities at baseline or acceptable for next level of care
 - Patient hemodynamically stable, as indicated by **ALL** of the following(1)(2)(3)(4)(5):
 - Tachycardia absent
 - Hypotension absent
 - No evidence of inadequate perfusion (eg, no myocardial ischemia)
 - No other hemodynamic abnormalities (eg, no Orthostatic hypotension)

References

- Puskarich MA, Jones AE. Shock. In: Walls RM, editor. Rosen's Emergency Medicine: Concepts and Clinical Practice. 10th ed. Philadelphia, PA 19103-2899: Elsevier; 2023:34-41.e1.
- Lewis J, Patel B. Shock. In: Gershel JC, Rauch DA, editors. Caring for the Hospitalized Child: A Handbook of Inpatient Pediatrics. 3rd ed. Elk Grove Village, IL: American Academy of Pediatrics; 2023:107-114.
- Ingbar DH, Thiele H. Cardiogenic shock and pulmonary edema. In: Loscalzo J, Fauci A, Kasper D, Hauser S, Longo D, Jameson JL, editors. Harrison's Principles of Internal Medicine. 21st ed. McGraw Hill Education; 2022:2250-2257.
- Brant EB, Seymour CW, Angus DC. Sepsis and septic shock. In: Loscalzo J, Fauci A, Kasper D, Hauser S, Longo D, Jameson JL, editors. Harrison's Principles of Internal Medicine. 21st ed. McGraw Hill Education; 2022:2241-2249.
- Singer M, et al. The Third International Consensus definitions for sepsis and septic shock (Sepsis-3). Journal of the American Medical Association 2016;315(8):801-810. DOI: 10.1001/jama.2016.0287.

Hypotension

- Hypotension, as indicated by **ALL** of the following(1)(2)(3)(4):
 - Not patient baseline (eg, healthy adult with low SBP) or intentional therapeutic goal (eg, low SBP as treatment goal in heart failure)
 - Low blood pressure, as indicated by **1 or more** of the following:
 - SBP less than 90 mm Hg in adult or child 10 years or older[A]
 - Mean arterial pressure[A][B] less than 70 mm Hg in adult or child 10 years or older
 - Decrease in baseline mean arterial pressure of 25% or more,[A][B] with significant signs or symptoms due to lower blood pressure (eg, near syncope, syncope, chest pain)
 - SBP less than sum of 70 mm Hg plus twice patient's age in years in child 1 to 9 years of age[A]
 - SBP less than 70 mm Hg in infant 1 to 11 months of age[A]

References

- Jones D, Di Francesco L. Hypotension. In: McKean SC, Ross JJ, Dressler DD, Scheurer DB, editors. Principles and Practice of Hospital Medicine. 2nd ed. New York, NY: McGraw-Hill Education; 2017:657-64.
- Massaro AF. Approach to the patient with shock. In: Loscalzo J, Fauci A, Kasper D, Hauser S, Longo D, Jameson JL, editors. Harrison's Principles of Internal Medicine. 21st ed. McGraw Hill Education; 2022:2235-2241.
- Horeczko T. Pediatric cardiac disorders. In: Walls RM, editor. Rosen's Emergency Medicine: Concepts and Clinical Practice. 10th ed. Philadelphia, PA 19103-2899: Elsevier; 2023:2109-2131.e1.
- Singh S, Holmes JF. Pediatric trauma. In: Walls RM, editor. Rosen's Emergency Medicine: Concepts and Clinical Practice. 10th ed. Philadelphia, PA 19103-2899: Elsevier; 2023:2052-2066.e3.

Footnotes

- A. Criteria based upon clinician acquired numeric values (eg, vital signs, oxygen saturation) should be used if they are accurate reflections of the patient's condition. Transitory findings (eg, abnormal only upon initial emergency department intake or only one time out of multiple readings) that rapidly improve with no or minimal treatment usually do not reflect disease severity or risk for deterioration. This does not imply that an initial or one-time reading cannot ever be applicable. The goal is to separate erroneous or incidental findings from those that truly represent the patient's clinical picture.
- B. The mean arterial pressure takes into account both systolic and diastolic blood pressure readings and is calculated as Mean Arterial Pressure (MAP) = $1/3 \text{ SBP} + 2/3 \text{ DBP}$. A calculator is available at <https://www.mdcalc.com/mean-arterial-pressure-map>.

Hypotension absent

- Hypotension absent^[A], as indicated by **1 or more** of the following⁽¹⁾⁽²⁾⁽³⁾⁽⁴⁾:
 - SBP greater than or equal to 90 mm Hg in adult or child 10 years or older
 - Mean arterial pressure^[B] greater than or equal to 70 mm Hg in adult or child 10 years or older
 - Mean arterial pressure^[B] at patient's baseline (eg, healthy adult with low SBP), at intentional therapeutic goal (eg, patient with heart failure), or acceptable for next level of care (eg, blood pressure stable and no significant signs or symptoms due to low blood pressure)
 - SBP greater than or equal to sum of 70 mm Hg plus twice patient's age in years in child 1 to 9 years of age
 - SBP greater than or equal to 70 mm Hg in infant 1 to 11 months of age

References

- Jones D, Di Francesco L. Hypotension. In: McKean SC, Ross JJ, Dressler DD, Scheurer DB, editors. Principles and Practice of Hospital Medicine. 2nd ed. New York, NY: McGraw-Hill Education; 2017:657-64.
- Massaro AF. Approach to the patient with shock. In: Loscalzo J, Fauci A, Kasper D, Hauser S, Longo D, Jameson JL, editors. Harrison's Principles of Internal Medicine. 21st ed. McGraw Hill Education; 2022:2235-2241.
- Horeczko T. Pediatric cardiac disorders. In: Walls RM, editor. Rosen's Emergency Medicine: Concepts and Clinical Practice. 10th ed. Philadelphia, PA 19103-2899: Elsevier; 2023:2109-2131.e1.
- Singh S, Holmes JF. Pediatric trauma. In: Walls RM, editor. Rosen's Emergency Medicine: Concepts and Clinical Practice. 10th ed. Philadelphia, PA 19103-2899: Elsevier; 2023:2052-2066.e3.

Footnotes

- Criteria based upon clinician-acquired numeric values (eg, vital signs, oxygen saturation) should be used if they are accurate reflections of the patient's condition. Transitory findings (eg, abnormal only upon initial emergency department intake or only one time out of multiple readings) that rapidly improve with no or minimal treatment usually do not reflect disease severity or risk for deterioration. This does not imply that an initial or one-time reading cannot ever be applicable. The goal is to separate erroneous or incidental findings from those that truly represent the patient's clinical picture.
- The mean arterial pressure takes into account both systolic and diastolic blood pressure readings and is calculated as Mean Arterial Pressure (MAP) = $1/3 \text{ SBP} + 2/3 \text{ DBP}$.

Hypoxemia

- Hypoxemia, as indicated by **1 or more** of the following⁽¹⁾:
 - Patient without baseline need for supplemental oxygen with oxygen saturation (SaO₂) of less than 90% or arterial blood gas partial pressure of oxygen (PO₂) of less than 60 mm Hg (8.0 kPa) on room air^{[A][B]}
 - Patient without baseline need for supplemental oxygen who now requires supplemental oxygen to keep SaO₂ greater than 89% or PO₂ greater than 59 mm Hg (7.9 kPa)^[A]
 - Patient with baseline need for supplemental oxygen who now requires increased supplemental oxygen to maintain oxygenation at baseline or acceptable level

References

- Dezube R, Lechtzin N. Hypoxia. In: McKean SC, Ross JJ, Dressler DD, Scheurer DB, editors. Principles and Practice of Hospital Medicine. 2nd ed. New York, NY: McGraw-Hill Education; 2017:669-675.
- Jamali H, et al. Racial disparity in oxygen saturation measurements by pulse oximetry: evidence and implications. Annals of the American Thoracic Society 2022;19(12):1951-1964. DOI: 10.1513/AnnalsATS.202203-270CME.
- Rojas-Camayo J, et al. Reference values for oxygen saturation from sea level to the highest human habitation in the Andes in acclimatized persons. Thorax 2018;73(8):776-778. DOI: 10.1136/thoraxjnl-2017-210598.

Footnotes

- Pulse oximetry (SpO₂) may underestimate arterial saturation (SaO₂), especially in people with dark skin pigmentation, thick skin, cold body temperature, or who are wearing colored nail polish, and thus may not accurately detect hypoxemia.⁽²⁾ Where appropriate, pulse oximetry should be measured after warming fingers or removing nail polish. Pulse oximetry results should be assessed within the context of the patient's clinical presentation, and performance of an arterial blood gas considered for a more accurate assessment of oxygenation if indicated.⁽²⁾ Specific target values may require adjustment when care is delivered at high altitude.⁽¹⁾⁽³⁾
- Criteria based upon clinician-acquired numeric values (eg, vital signs, oxygen saturation) should be used if they are accurate reflections of the patient's condition. Transitory findings (eg, abnormal only upon initial emergency department intake or only one time out of multiple readings) that rapidly improve with no or minimal treatment usually do not reflect disease severity or risk for deterioration. This does not imply that an initial or one-time reading cannot ever be applicable. The goal is to separate erroneous or incidental findings from those that truly represent the patient's clinical picture.

Multimodal analgesia

- Multimodal analgesia involves the utilization of 2 or more analgesic agents with different mechanisms of action in order to provide additive or synergistic pain control, while minimizing side effects and reliance on opioids.(1)(2)(3)

References

1. D'Souza RS, Johnson RL. Regional and multimodal treatments of perioperative pain. In: Benzon H, editor. Practical Management of Pain. 6th ed. Philadelphia, PA 19103-2899: Elsevier; 2023:355-373.
2. George S, Johns M. Review of nonopioid multimodal analgesia for surgical and trauma patients. American Journal of Health-System Pharmacy 2020;77(24):2052-2063. DOI: 10.1093/ajhp/zxaa301.
3. Finnerup NB. Nonnarcotic methods of pain management. New England Journal of Medicine 2019;380(25):2440-2448. DOI: 10.1056/NEJMr1807061.

No contraindications to lung transplant present

- No contraindications to lung transplant present, as indicated by **ALL** of the following(1)(2)(3)(4)(5):
 - No malignancy that has high risk of recurrence or cancer-related mortality[A]
 - No untreatable failure or dysfunction of another major organ system unless combined organ transplants can be performed[B]
 - No stroke, acute coronary syndrome, or myocardial infarction within 30 days[C](1)
 - No incurable extrapulmonary infection
 - No active *Mycobacterium tuberculosis* infection
 - No HIV infection with detectable viral load[D](1)
 - No uncontrolled systemic (extrapulmonary) disease[E](7)
 - No bilateral diaphragmatic paralysis or severe diaphragmatic weakness(7)
 - No severe tracheobronchomalacia
 - No severe transpleural systemic to bronchial artery collaterals or aorto-pulmonary collaterals
 - BMI not over 35  BMI Calculator
 - No uncorrectable bleeding diathesis
 - No active alcohol, tobacco, or illicit substance misuse or dependence[F](1)
 - No unaddressed barriers to ability to comply with immunosuppression protocol and follow-up requirements (eg, adequate social support, no severe uncontrolled psychiatric illness)
 - No history of persistent nonadherence
 - No progressive cognitive impairment(1)
 - No severe swallowing dysfunction with recurrent aspiration that is not amenable to therapy (eg, in patient with systemic sclerosis or Sjogren syndrome)(7)

References

1. Leard LE, et al. Consensus document for the selection of lung transplant candidates: An update from the International Society for Heart and Lung Transplantation. Journal of Heart and Lung Transplantation 2021;40(11):1349-1379. DOI: 10.1016/j.healun.2021.07.005. (Reaffirmed 2022 Jul)
2. Martin AK, Renew JR, Jayaraman AL, Murray AW, Fritz AV, Ramakrishna H. Analysis of outcomes in lung transplantation. Journal of Cardiothoracic and Vascular Anesthesia 2019;33(5):1455-1466. DOI: 10.1053/j.jvca.2018.06.014.
3. Rossano JW, Goldfarb SB. Pediatric heart and heart-lung transplantation. In: Kliegman RM, St. Geme JW, Blum NJ, Shah SS, Tasker RC, Wilson KM, editors. Nelson Textbook of Pediatrics. 21st ed. Philadelphia, PA: Elsevier; 2020:2483-2487.e1.
4. Faro A, Weymann A. Transplantation. Pediatric Clinics of North America 2016;63(4):709-734. DOI: 10.1016/j.pcl.2016.04.010.
5. Bryant R, Morales D, Schechter M. Pediatric lung transplantation. Seminars in Pediatric Surgery 2017;26(4):213-216. DOI: 10.1053/j.sempedsurg.2017.07.005.
6. Kanaparthi J, et al. Prior and perioperative revascularization does not affect survival in lung transplant patients. Annals of Thoracic Surgery 2020;109(6):1677-1683. DOI: 10.1016/j.athoracsurg.2020.01.016.
7. Crespo MM, et al. ISHLT consensus document on lung transplantation in patients with connective tissue disease: Part I: Epidemiology, assessment of extrapulmonary conditions, candidate evaluation, selection criteria, and pathology statements. Journal of Heart and Lung Transplantation 2021;40(11):1251-1266. DOI: 10.1016/j.healun.2021.07.014. (Reaffirmed 2022 Aug)

Footnotes

- A. Transplant may be considered for selected cases of lung-limited cancer (eg, adenocarcinoma in situ, lepidic adenocarcinoma (previously called bronchoalveolar cell carcinoma), or minimally invasive adenocarcinoma) if surgical resection is not otherwise feasible, systemic therapies are contraindicated or have failed, and transplant is thought to be curative.
- B. Patients who required cardiac interventions before transplant are not precluded from lung transplant.(6)
- C. Patients with recent type 2 myocardial infarction (demand ischemia) are not precluded from lung transplant.(1)
- D. Patients with HIV infection and undetectable viral load are not precluded from lung transplant because case series show these patients have similar 5-year survival rates as those without HIV. Coordination of the antiretroviral and immunosuppressive regimens is needed due to risk of drug interactions.(1)

- E. Examples include progressive neuromuscular disease (eg, amyotrophic lateral sclerosis), refractory collagen vascular disease (eg, systemic lupus erythematosus or systemic scleroderma not confined to the lung, rheumatoid arthritis with unstable atlantoaxial instability at risk for spinal chord compression, or severe arthropathy with inability to perform perioperative pulmonary rehabilitation), amyloidosis, or refractory sarcoidosis not confined to the lung.(1)(2)(7) Some patients with cardiac involvement may be considered for combined heart-lung transplant.(7)
- F. The duration of abstinence is discretionary and dependent on the psychosocial evaluation by the transplant center.(1)

Orthostatic hypotension

- Orthostatic hypotension,[A][B] as indicated by **1 or more** of the following(1)(2)(3):
 - Fall in SBP of 20 mm Hg or more 1 to 3 minutes after patient sits or stands from recumbent position
 - Fall in DBP of 10 mm Hg or more 1 to 3 minutes after patient sits or stands from recumbent position

References

1. Shibao C, Lipsitz LA, Biaggioni I, American Society of Hypertension Writing Group. Evaluation and treatment of orthostatic hypotension. Journal of the American Society of Hypertension 2013 Jul-Aug;7(4):317-324. DOI: 10.1016/j.jash.2013.04.006.
2. Dalal AS, Van Hare GF. Syncope. In: Kliegman RM, St. Geme JW, Blum NJ, Shah SS, Tasker RC, Wilson KM, editors. Nelson Textbook of Pediatrics. 21st ed. Philadelphia, PA: Elsevier; 2020:566-571.e1.
3. Fang JC, O'Gara PT. History and physical examination: an evidence-based approach. In: Libby P, Bonow RO, Mann DL, Tomaselli GF, Bhatt DL, Solomon SD, editors. Braunwald's Heart Disease: A Textbook of Cardiovascular Medicine. 12th ed. Elsevier; 2022:123-140.

Footnotes

- A. Concomitant measurements of the heart rate are important to measure to help diagnose subtypes of orthostatic hypotension (eg, the lack of a compensatory increase in heart rate is typical of autonomic failure and an exaggerated tachycardia may be reflective of volume depletion). However, the heart rate is not a component of the definition of orthostatic hypotension which relies upon blood pressure alone.(1)(2)(3)
- B. Criteria based upon clinician acquired numeric values (eg, vital signs, oxygen saturation) should be used if they are accurate reflections of the patient's condition. Transitory findings (eg, abnormal only upon initial emergency department intake or only one time out of multiple readings) that rapidly improve with no or minimal treatment usually do not reflect disease severity or risk for deterioration. This does not imply that an initial or one-time reading cannot ever be applicable. The goal is to separate erroneous or incidental findings from those that truly represent the patient's clinical picture.

Respiratory distress

- Respiratory distress, as indicated by **ALL** of the following(1)(2):
 - Patient with **1 or more** of the following:
 - Dyspnea (difficulty breathing)
 - Tachypnea
 - Abnormal breathing pattern (eg, chest retractions)
 - Other evidence of difficulty breathing
 - Evidence of respiratory compromise, as indicated by **1 or more** of the following:
 - Hypoxemia
 - Altered mental status
 - Other evidence of respiratory compromise (eg, respiratory acidosis)

References

1. Braithwaite SA, Wessel AL. Dyspnea. In: Walls RM, editor. Rosen's Emergency Medicine: Concepts and Clinical Practice. 10th ed. Philadelphia, PA 19103-2899: Elsevier; 2023:194-201.e2.
2. Naureckas ET, Solway J. Disturbances of respiratory function. In: Loscalzo J, Fauci A, Kasper D, Hauser S, Longo D, Jameson JL, editors. Harrison's Principles of Internal Medicine. 21st ed. McGraw Hill Education; 2022:2133-2139.

Social Determinants of Health Assessment

- Risk of poor health outcomes may be increased by the presence of **1 or more** of the following social determinants of health(1)(2)(3)(4):
 - Housing insecurity, as indicated by **1 or more** of the following:
 - Individual or caregiver's current living situation is **1 or more** of the following(5):
 - Does not have own housing (eg, staying in a hotel, shelter, or with others)
 - Has own housing (eg, house, apartment), but at risk of losing it in the future (ie, behind on rent or mortgage)
 - Has own housing (eg, house, apartment), but has lived in 3 or more places in past year
 - Current housing has **1 or more** of the following:
 - Electrical appliances (eg, stove, refrigerator) not working or unavailable

- Insufficient heating or cooling
- Insufficient ventilation
- Lead paint or pipes
- Mold
- Pests (eg, bugs) or rodents
- Smoke detectors not working or unavailable
- Food insecurity, as indicated by **1 or more** of the following(6):
 - In the past year, individual or caregiver ran out of food and did not have money to buy more food.
 - In the past year, individual or caregiver worried that they would run out of food before they received money to buy more food.
- Insufficient transportation, as indicated by **1 or more** of the following(7):
 - In the past year, individual or caregiver missed medical appointments or could not get medications due to lack of transportation.
 - In the past year, individual or caregiver missed nonmedical activities, work, or could not get things needed for daily living due to lack of transportation.
- Insufficient utilities, as indicated by **1 or more** of the following(8):
 - Utilities (eg, electricity, water, gas, or oil) are currently shut off or unavailable.
 - In the past year, electric, water, gas, or oil company threatened to shut off services.
- Personal safety risk, as indicated by **2 or more** of the following(6):
 - Individual is sometimes or frequently physically hurt by another person (including family member).
 - Individual is sometimes or frequently insulted or talked down to by another person (including family member).
 - Individual is sometimes or frequently threatened with physical harm by another person (including family member).
 - Individual is sometimes or frequently screamed or cursed at by another person (including family member).
- Insufficient dependent care, as indicated by **1 or more** of the following:
 - In the past year, individual or caregiver was unable to work due to lack of dependent care.
 - In the past year, individual or caregiver was unable to work more (additional) hours due to lack of dependent care.
 - In the past year, individual or caregiver missed medical appointments or could not get medications due to lack of dependent care.
 - In the past year, individual or caregiver missed nonmedical activities (eg, school, church, social activity) due to lack of dependent care.
- Depression risk, as indicated by **ALL** of the following:
 - In the past 2 weeks, individual had little interest or pleasure in normal activities on at least several days.
 - In the past 2 weeks, individual felt down, depressed, or hopeless on at least several days.

References

1. Scanlon A, Reinisch C. Social determinants of health. In: Harding MM, Kwong J, Hagler D, Reinisch C, editors. *Lewis's Medical-Surgical Nursing: Assessment and Management of Clinical Problems*. 12th ed. St. Louis, MO: Mosby; 2023:18-20.
2. Moen M, Storr C, German D, Friedmann E, Johantgen M. A review of tools to screen for social determinants of health in the United States: a practice brief. *Population Health Management* 2020;23(6):422-429. DOI: 10.1089/pop.2019.0158.
3. Daniel-Robinson L, Moore JE. Innovation and Opportunities to Address Social Determinants of Health in Medicaid Managed Care. [Internet] Institute for Medicaid Innovation. 2019 Jan Accessed at: <https://medicaidinnovation.org/emerging-topics/>. [accessed 2023 Oct 20]
4. Billioux A, Verlander K, Anthony S, Alley D. Standardized Screening for Health-Related Social Needs in Clinical Settings: the Accountable Health Communities Screening Tool. [Internet] National Academy of Sciences. 2017 May Accessed at: <https://nam.edu/>. [accessed 2023 Sep 18]
5. Sandel M, et al. Unstable housing and caregiver and child health in renter families. *Pediatrics* 2018;14(2):e20172199. DOI: 10.1542/peds.2017-2199.
6. Children's HealthWatch Survey. Screening Instrument [Internet] Children's HealthWatch. 2020 Sep Accessed at: <https://childrenshealthwatch.org/>. [accessed 2023 Sep 11]
7. PRAPARE®: Protocol for Responding to and Assessing Patient Assets, Risks, and Experiences Screening Tool. [Internet] Association of Asian Pacific Community Health Organizations (AAPCHO) and National Association of Community Health Centers (NACHC). 2016 Sep Accessed at: <https://prapare.org/the-prapare-screening-tool/>. [accessed 2023 Sep 21]
8. Cook JT, et al. A brief indicator of household energy security: associations with food security, child health, and child development in US infants and toddlers. *Pediatrics* 2008;122(4):e867-75. DOI: 10.1542/peds.2008-0286.

Tachycardia

- Tachycardia,^{[A][B]} as indicated by **1 or more** of the following(1)(2):
 - Heart rate greater than 100 beats per minute in adult or child age 6 years or older
 - Heart rate greater than 115 beats per minute in child 3 to 5 years of age
 - Heart rate greater than 125 beats per minute in child 1 or 2 years of age
 - Heart rate greater than 130 beats per minute in infant 6 to 11 months of age
 - Heart rate greater than 150 beats per minute in infant 3 to 5 months of age
 - Heart rate greater than 160 beats per minute in infant 1 or 2 months of age

References

1. Southmayd GL. Tachycardia. In: McKean SC, Ross JJ, Dressler DD, Scheurer DB, editors. Principles and Practice of Hospital Medicine. 2nd ed. New York, NY: McGraw-Hill Education; 2017:729-739.
2. Pediatric parameters and equipment. In: Kleinman K, McDaniel L, Molloy M, editors. The Harriet Lane Handbook: A Manual for Pediatric House Officers. 22nd ed. 202: Elsevier; 2021:frontpiece tables.

Footnotes

- A. Criteria based upon clinician acquired numeric values (eg, vital signs, oxygen saturation) should be used if they are accurate reflections of the patient's condition. Transitory findings (eg, abnormal only upon initial emergency department intake or only one time out of multiple readings) that rapidly improve with no or minimal treatment usually do not reflect disease severity or risk for deterioration. This does not imply that an initial or one-time reading cannot ever be applicable. The goal is to separate erroneous or incidental findings from those that truly represent the patient's clinical picture.
- B. Interpretation of heart rate requires clinical judgment and consideration of several patient-specific factors, such as the patient's baseline heart rate, medications, and clinical impact. For example, an elderly patient on a beta-blocker medication with a baseline resting heart rate of 60 beats per minute may be clinically tachycardic at a heart rate of 94 beats per minute. Likewise, a patient who is upset, in pain, or nervous in the emergency department with a heart rate of 106 beats per minute may meet the technical definition of tachycardia, but this tachycardia (absent associated findings such as chest pain or hypotension) may not be clinically important. The numeric values included in this definition are provided to allow for consistency in terms of a technical definition of the term tachycardia. Whether a heart rate above or below the technical threshold is clinically meaningful is a matter of persistence, context, and clinical judgment.

Tachycardia absent

- Tachycardia[A][B] absent, as indicated by **1 or more** of the following(1)(2):
 - Heart rate less than or equal to 100 beats per minute in adult or child 6 years or older
 - Heart rate less than or equal to 115 beats per minute in child 3 to 5 years of age
 - Heart rate less than or equal to 125 beats per minute in child 1 or 2 years of age
 - Heart rate less than or equal to 130 beats per minute in infant 6 to 11 months of age
 - Heart rate less than or equal to 150 beats per minute in infant 3 to 5 months of age
 - Heart rate less than or equal to 160 beats per minute in infant 1 or 2 months of age

References

1. Southmayd GL. Tachycardia. In: McKean SC, Ross JJ, Dressler DD, Scheurer DB, editors. Principles and Practice of Hospital Medicine. 2nd ed. New York, NY: McGraw-Hill Education; 2017:729-739.
2. Pediatric parameters and equipment. In: Kleinman K, McDaniel L, Molloy M, editors. The Harriet Lane Handbook: A Manual for Pediatric House Officers. 22nd ed. 202: Elsevier; 2021:frontpiece tables.

Footnotes

- A. Criteria based upon clinician acquired numeric values (eg, vital signs, oxygen saturation) should be used if they are accurate reflections of the patient's condition. Transitory findings (eg, abnormal only upon initial emergency department intake or only one time out of multiple readings) that rapidly improve with no or minimal treatment usually do not reflect disease severity or risk for deterioration. This does not imply that an initial or one-time reading cannot ever be applicable. The goal is to separate erroneous or incidental findings from those that truly represent the patient's clinical picture.
- B. Interpretation of heart rate requires clinical judgment and consideration of several patient-specific factors, such as the patient's baseline heart rate, medications, and clinical impact. For example, an elderly patient on a beta-blocker medication with a baseline resting heart rate of 60 beats per minute may be clinically tachycardic at a heart rate of 94 beats per minute. Likewise, a patient who is upset, in pain, or nervous in the emergency department with a heart rate of 106 beats per minute may meet the technical definition of tachycardia, but this tachycardia (absent associated findings such as chest pain or hypotension) may not be clinically important. The numeric values included in this definition are provided to allow for consistency in terms of a technical definition of the term tachycardia. Whether a heart rate above or below the technical threshold is clinically meaningful is a matter of persistence, context, and clinical judgment.

Tachypnea

- Tachypnea,[A][B] as indicated by respiratory rate of **1 or more** of the following(2)(3):
 - Greater than 18 breaths per minute in adult or child 13 years of age or older[A]
 - Greater than 22 breaths per minute in child 6 to 12 years of age[A]
 - Greater than 25 breaths per minute in child 3 to 5 years of age[A]
 - Greater than 30 breaths per minute in child 1 or 2 years of age[A]
 - Greater than 40 breaths per minute in infant 6 to 11 months of age[A]

- Greater than 45 breaths per minute in infant 3 to 5 months of age[A]
- Greater than 60 breaths per minute in infant 1 or 2 months of age[A]

References

1. Schriger DL. Approach to the patient with abnormal vital signs. In: Goldman L, Schafer AI, editors. Goldman-Cecil Medicine. 26th ed. Elsevier; 2020:28-31.e2.
2. Braithwaite SA, Wessel AL. Dyspnea. In: Walls RM, editor. Rosen's Emergency Medicine: Concepts and Clinical Practice. 10th ed. Philadelphia, PA 19103-2899: Elsevier; 2023:194-201.e2.
3. Pediatric parameters and equipment. In: Kleinman K, McDaniel L, Molloy M, editors. The Harriet Lane Handbook: A Manual for Pediatric House Officers. 22nd ed. 202: Elsevier; 2021:frontpiece tables.

Footnotes

- A. Criteria based upon clinician acquired numeric values (eg, vital signs, oxygen saturation) should be used if they are accurate reflections of the patient's condition. Transitory findings (eg, abnormal only upon initial emergency department intake or only one time out of multiple readings) that rapidly improve with no or minimal treatment usually do not reflect disease severity or risk for deterioration. This does not imply that an initial or one-time reading cannot ever be applicable. The goal is to separate erroneous or incidental findings from those that truly represent the patient's clinical picture.
- B. Interpretation of respiratory rate requires clinical judgment and consideration of several patient-specific factors, such as the patient's baseline respiratory rate and whether the respiratory rate is indicative of significant underlying respiratory or other pathology. A mildly increased respiratory rate (eg, 20 to 22 in an adult), absent other indications of serious illness (eg, hypoxemia, metabolic acidosis, decreased tidal volume, pain), may be transitory and not clinically important. The numeric values included in this definition are provided to allow for consistency in terms of a technical definition of the term tachypnea. Whether a respiratory rate above the technical threshold is clinically meaningful is a matter of persistence, context, and clinical judgment. In addition, the measurement of the respiratory rate is subject to error. A medical textbook states that because there can be significant breath to breath variation in respiratory rate, the rate should be assessed for at least 30 seconds, preferably for a full minute.⁽¹⁾ This same textbook also states that new technologies designed to measure the respiratory rate have not proved to be clinically useful.⁽¹⁾

Codes

ICD-10 Diagnosis: D86.0, D86.2, E84.0, I27.0, I27.20, I27.21, I27.22, I27.23, I27.24, I27.29, I27.81, I27.83, I27.89, I27.9, J43.0, J43.1, J43.2, J43.8, J43.9, J44.1, J44.9, J47.0, J47.1, J47.9, J84.10, J84.112, J84.81, J84.82, J84.83, J84.89 [Hide]

ICD-10 Procedure: 0BYC0Z0, 0BYC0Z1, 0BYC0Z2, 0BYD0Z0, 0BYD0Z1, 0BYD0Z2, 0BYF0Z0, 0BYF0Z1, 0BYF0Z2, 0BYG0Z0, 0BYG0Z1, 0BYG0Z2, 0BYH0Z0, 0BYH0Z1, 0BYH0Z2, 0BYJ0Z0, 0BYJ0Z1, 0BYJ0Z2, 0BYK0Z0, 0BYK0Z1, 0BYK0Z2, 0BYL0Z0, 0BYL0Z1, 0BYL0Z2, 0BYM0Z0, 0BYM0Z1, 0BYM0Z2 [Hide]

CPT®: 32851, 32852, 32853, 32854, 33935

CPT copyright 2023 American Medical Association. All rights reserved.

MCG Health
Inpatient & Surgical Care 28th Edition
Copyright © 2024 MCG Health, LLC
All Rights Reserved

Last Update: 3/14/2024 9:41:34 AM
Build Number: 28.1.20240313235330.016020