

eGFR Calculator

Glomerular filtration rate (GFR) is the best overall index of kidney function. Normal GFR varies according to age, sex, and body size, and declines with age.

The National Kidney Foundation (NKF) and the American Society of Nephrology (ASN) convened a Task Force in 2021 to focus on the use of race when estimating GFR. The joint NKF/ASN Task Force recommends using the CKD-EPI Creatinine Equation (2021) to estimate GFR. More information regarding this recommendation may be found in this [article highlighting the Task Force's recommendations](#).

This eGFR Calculator is intended for use only by health care professionals. For more information about measuring your kidney health, please refer to the following resources:

- [Kidney Tests](#)
- [Estimated Glomerular Filtration Rate \(eGFR\)](#)
- [Serum Creatinine](#)
- [Cystatin C](#)

For specific questions about your results or kidney health, be sure to ask your health care professional.

Serum Creatinine:		☒ mg/dL ☐ µmol/L
Serum Cystatin C:		mg/L
Age:		Years
Gender:	☒ Male ☐ Female	
Standardized Assays:	☒ Yes ☐ No ☐ Not Sure	
Adjust for body surface area:	☐ Yes ☒ No ☐ Not Sure	

Calculate

Results

Feedback

CKD-EPI creatinine equation (2021)		mL/min/1.73m ²
CKD-EPI creatinine-cystatin equation (2021)		mL/min/1.73m ²
CKD-EPI cystatin C equation (2012)		mL/min/1.73m ²

For persons under 18 years of age, use the pediatric GFR calculator.

Is it CKD?

Either of the following must be present for ≥3 months to be CKD:

- GFR less than 60 ≥3 months
- ACR ≥30 mg/g or other markers of kidney damage

Click to learn [more](#).

Equation used to estimate GFR?

- ☒ CKD-EPI Creatinine (2021)
☐ CKD-EPI Creatinine-Cystatin C (2021)
☐ CKD-EPI Cystatin C (2012)

What is the patient's ACR?†

- ☒ <30 mg/g ☐ <3 mg/mmol
☐ 30-300 mg/g ☐ 3-30 mg/mmol
☐ >300 mg/g ☐ >30 mg/mmol

Based on the information supplied:

GFR category is:†

ACR category is:**

CKD classification is:

Risk of progression is:



Frequency of monitoring
should be:

Referral to a nephrologist
is:

† Neither the category of GFR nor the category of albuminuria alone can fully capture prognosis of CKD. Persistent and increased albuminuria has been shown to be an independent risk factor for CKD progression.

*In the absence of evidence of kidney damage, neither GFR category G1 nor G2 fulfill the criteria for CKD.

**ACR 30-300 mg/g for > 3 months indicates CKD.

Additional Information

Formulas

- [CKD-EPI Creatinine Equation \(2021\)](#)
- [CKD-EPI Creatinine-Cystatin Equation \(2021\)](#)
- [CKD-EPI Cystatin C Equation \(2012\)](#)
- [MDRD Study Equation](#)
- Learn more about CKD-EPI Collaboration and estimating GFR at www.ckdepi.org.



About CKD

- [Criteria for CKD?](#)
- [Classification of CKD?](#)
- [Why classify CKD?](#)
- [Explore Case Studies](#)

[CKD Risk Map](#)

[Clinician Tools](#)

[Kidney Failure Risk Equation](#)

[Guidelines](#)

Calculators

Use our GFR calculators to estimate GFR for adults or children.

- [eGFR Calculator](#)

- [Pediatric GFR Calculator](#)
- [Cockcroft-Gault formula](#) (*use for drug research only*)
- [Pediatric Chronic Kidney Disease Risk Calculator](#) (*used by nephrologists and other healthcare providers only*)
- [Kidney Failure Risk Equation](#)
- [eGFR Calculator App for iPhone/iPad](#)
- [FAQs About GFR Estimates](#)

