

forall*x*

BRISTOL

Version from December 7, 2020
Download [here](#).
Please redownload every week.

forall x : Bristol

Textbook for

PHIL10032: Logic and Critical Thinking 2020

Also used in

PHIL10014: Introduction to Formal Logic 2020

By **P. D. Magnus**

Tim Button

with additions by

J. Robert Loftis

Robert Trueman

remixed and revised by

Aaron Thomas-Bolduc

Richard Zach

and by

Catrin Campbell-Moore

Last updated December 7, 2020

This book is based on **forallx: Calgary** by

Aaron Thomas-Bolduc & Richard Zach
University of Calgary

used under a **CC BY-SA 4.0** license, which is based on **forallx: Cambridge**, by

Tim Button
University of Cambridge

used under a **CC BY-SA 3.0** license, which is based in turn on **forallx**, by

P.D. Magnus
University at Albany, State University of New York

used under a **CC BY-SA 3.0** license.
and was remixed, revised, & expanded by

Aaron Thomas-Bolduc & Richard Zach
University of Calgary
then altered by

Catrin Campbell-Moore
University of Bristol Further contributors: Stuart Presnell, Richard Pettigrew.

It includes additional material from **forallx** by P.D. Magnus and *Metatheory* by Tim Button, both used under a **CC BY-SA 3.0** license, and from **forallx: Lorain County Remix**, by Cathal Woods and J. Robert Loftis, used under a **CC BY-SA 4.0** license.

This work is licensed under a **Creative Commons Attribution-ShareAlike 4.0** license. You are free to copy and redistribute the material in any medium or format, and remix, transform, and build upon the material for any purpose, even commercially, under the following terms:

- You must give appropriate credit, provide a link to the license, and indicate if changes were made. You may do so in any reasonable manner, but not in any way that suggests the licensor endorses you or your use.
- If you remix, transform, or build upon the material, you must distribute your contributions under the same license as the original.

The \LaTeX source for this book is available on **GitHub**.

Preface

The book is divided into nine parts. Part **I** introduces the topic and notions of logic in an informal way, without introducing a formal language yet. Parts **II–IV** concern truth-functional languages. In it, sentences are formed from basic sentences using a number of connectives (‘or’, ‘and’, ‘not’, ‘if . . . then’) which just combine sentences into more complicated ones. We discuss logical notions such as entailment in two ways: semantically, using the method of truth tables (in Part **III**) and proof-theoretically, using a system of formal derivations (in Part **IV**). Parts **VI–VIII** deal with a more complicated language, that of first-order logic. It includes, in addition to the connectives of truth-functional logic, also names, predicates, identity, and the so-called quantifiers. These additional elements of the language make it much more expressive than the truth-functional language, and we’ll spend a fair amount of time investigating just how much one can express in it. Again, logical notions for the language of first-order logic are defined semantically, using interpretations, and proof-theoretically, using a more complex version of the formal derivation system introduced in Part **IV**. Part **X** covers two advanced topics: that of conjunctive and disjunctive normal forms and the expressive completeness of the truth-functional connectives, and the soundness of natural deduction for TFL. Part **XI** discusses the extension of TFL by non-truth-functional operators for possibility and necessity: modal logic.

In the appendices you’ll find a discussion of alternative notations for the languages we discuss in this text, of alternative derivation systems, and a quick reference listing most of the important rules and definitions. The central terms are listed in a glossary at the very end.

This book is based on a text originally written by P. D. Magnus in the version revised and expanded by Tim Button. It also includes

some material (mainly exercises) by J. Robert Loftis. The material in Part **XI** is based on notes by Robert Trueman, and the material in Part ?? on two chapters from Tim Button's open text *Metatheory*. Aaron Thomas-Bolduc and Richard Zach have combined elements of these texts into the present version, changed some of the terminology and examples, rewritten some sections, and added material of their own. Catrin Campbell-Moore has then made very minor alterations for the Bristol course. The resulting text is licensed under a Creative Commons Attribution-ShareAlike 4.0 license.

Contents

Preface	iii
---------	-----

I	Arguments	1
----------	------------------	----------

1	Arguments	2
2	The scope of logic	11
3	Other logical notions	24

II	Truth-functional logic	31
-----------	-------------------------------	-----------

4	First steps to symbolization	32
5	Connectives	36
6	Sentences of TFL	47
7	Symbolising complex sentences	55
8	Ambiguity	60
9	Use and mention	65

III	Truth tables	68
------------	---------------------	-----------

10	Truth rules for the connectives of TFL	69
11	Truth-functional connectives	75
12	Complete truth tables	79
13	Validity in TFL	89
14	Other logical notions	98
15	Truth table shortcuts	103

16	Partial truth tables	108
17	TFL vs English connectives	114

IV Natural deduction for TFL **116**

18	The very idea of natural deduction	117
19	The First Basic Rules for TFL	120
20	More Basic Rules for TFL	138
21	Proofs and Validity	153
22	Additional rules for TFL	157
23	Derived rules	162
24	Soundness and completeness	167
25	Proof-theoretic concepts	175

V Fallacies of Reasoning **178**

26	Logical fallacies	179
27	Probabilistic fallacies	185

VI First-order logic **195**

28	Building blocks of FOL	196
29	Sentences with one quantifier	206
30	Multiple quantifiers	218
31	Sentences of FOL	228
32	Ambiguity	235

VII Interpretations **240**

33	Extensionality	241
34	Truth in FOL	246
35	Semantic concepts	256
36	Using interpretations	258
37	Reasoning about all interpretations	264

VIII Natural deduction for FOL **268**

38	Basic rules for FOL	269
39	Conversion of quantifiers	285
40	Derived rules	287
41	Proof-theoretic and semantic concepts	289

IX Identity **293**

42	Identity	294
43	Definite descriptions	299
44	Semantics for FOL with identity	306
45	Rules for identity	309

X Metatheory (Non-examinable) **314**

46	Normal forms	315
47	Expressive Adequacy	320
48	Soundness	325

XI Modal logic (non-examinable) **331**

49	Introducing modal logic	332
50	Natural deduction for ML	335
51	Semantics for ML	347

Appendices **359**

A	Symbolic notation	359
B	Alternative proof systems	362
C	Quick reference	368

	Glossary	378
--	----------	-----

PART I

Arguments

CHAPTER 1

Arguments

1.1 Introduction

Much of philosophical practice is about argument and analysis. Arguing in support of or against some position, or understanding someone else's argument. Logic is the study of the practice of argument and analysis, abstracted from the specific details of a particular case.

In everyday language, we sometimes use the word 'argument' to talk about belligerent shouting matches. If you and a friend have an argument in this sense, things are not going well between the two of you. Logic is not concerned with such teeth-gnashing and hair-pulling. They are not arguments, in our sense; they are disagreements.

An argument, as we will understand it, is something more like this:

- If I acted of my own free will, then I could have acted otherwise.
- I could not have acted otherwise.
- Therefore: I did not act of my own free will.

We here have a series of sentences which may either be true or false. The final sentence, "I did not act of my own free will." expresses the *conclusion* of the argument. The two sentences before that are the *premises* of the argument. In a good argument, the conclusion follows from the premises. If you believe the premises then the argument should lead you to believing the conclusion.

Logic provides the ideal model of good argument: rational argument without rhetoric. The logical study of an argument can show whether it supports its conclusion or is flawed. Logic focuses only on the statements presented and the relationships between them. Extraneous

factors are set aside: unspoken assumptions, additional connotations of words, appeal to emotions.

Logical thinking can help us to work out the intended interpretation of a text, and to find alternative unintended interpretations. This can be helpful when reading someone else's writing, and essential when we are trying to write unambiguously. Logical analysis can help us to find ambiguity and alternative interpretations, and to write in a precise and unambiguous way that can only be interpreted as we intend. These are vital skills used in all philosophy as well as in life more generally.

This Part discusses arguments in natural languages like English. Throughout this textbook we will also consider arguments in formal languages and say what it is for those to be valid or invalid. We want formal validity, as defined in the formal language, to have at least some of the important features of natural-language validity.

1.2 Finding the components of an argument

Arguments consist of a list of *premises* along with a *conclusion*. In a good argument, the conclusion will follow from the premises. Our standard way to present them is:

- Premise 1
- Premise 2
- ...
- Premise n
- Therefore: Conclusion

For example

- If I acted of my own free will, then I could have acted otherwise.
- I could not have acted otherwise.
- Therefore: Therefore: I did not act of my own free will.

Often arguments are presented simply in a paragraph of text, or in a speech or article, and we first have to work out what the premises and conclusions are. Sometimes it's easy, for example:

If I acted of my own free will, then I could have acted otherwise. But, I could not have acted otherwise. So, I did not act of my own free will.

But often it is a significant piece of work to work out the premises and conclusion of an argument.

Many arguments start with premises, and end with a conclusion, but not all of them. It might start with the conclusion:

We should not have a second Brexit referendum. A second Brexit referendum would erode the very basis of democracy by suggesting that rule by the majority is an insufficient condition for democratic legitimacy.

Or it might have been presented with the conclusion in the middle:

Since the first Brexit referendum was made under false pretences, **the voters deserve a further say on any final deal agreed with Brussels.** After all, decisions as big as this need to have the public support, which has to come from a referendum.

Sometimes premises or the conclusion may be clauses in a sentence. A complete argument may even be contained in a single sentence:

The butler has an alibi; so they cannot have done it.

This argument has one premise, followed immediately by its conclusion. One particular kind of sentence can be confusing. Consider:

- If the murder weapon was a gun, then Prof. Plum did it.

These conditional, or “if-then”, statements might look like it expresses the argument, but in itself it does not. It’s just stating a fact, albeit a conditional fact. It might also be used in an argument, even as the conclusion of the argument:

- If I have free will, then there is some event that I could have caused to go differently.
- If determinism is true, then there is no event that I could have caused to go differently.
- Therefore: If determinism is true, I do not have free will.

As a guideline, there are some words you can look for which are often used to indicate whether something is a premise or conclusion:

Words often used to indicate an argument's conclusion:

so, therefore, hence, thus, accordingly, consequently

Words often used to indicate a premise:

since, because, as, given that, recalling that, after all


In analysing an argument, there is no substitute for a good nose. Whenever you come across an argument in a piece of philosophy you read, be it lecture notes, primary text, or secondary text, or in a newspaper article or on the internet, practice identifying the premises and conclusion.

Sometimes, though, people aren't giving arguments but are simply presenting facts or stating their opinion. For example, the following do not contain arguments, they're not trying to convince us of anything.

- I don't like cats. I think they're evil.
- Hundreds of vulnerable children as young as 10, who have spent most of their lives in the UK, are having their applications for British citizenship denied for failing to pass the government's controversial 'good character' test.

1.3 Intermediate Conclusions

We said an argument is given by a collection of *premises* along with a single *conclusion*. We might represent this as something like:



The premises are working together to lead to the conclusion.

But sometimes in the process of someone making an argument someone will make use of *intermediate conclusions*. Such arguments might have a structure more like:



However, we say that an argument is only something of the first kind. So what do we say about the second kind of thing? We can consider it two ways. We might consider it as an argument from premise 1, 2 and 3 to the conclusion. Or alternatively we can think of it as two arguments of the first kind chained together, one from premise 1 and 2 to the intermediate conclusion, and the second from the intermediate conclusion and premise 3 to the final conclusion.

1.4 Sentences

What kinds of things are the premises and conclusions of arguments? They are sentences which can either be true or false. Such sentences are called **DECLARATIVE SENTENCES**.

There are many other kinds of sentences, for example:

Questions ‘Are you sleepy yet?’ is an interrogative sentence. Although you might be sleepy or you might be alert, the question itself is neither true nor false. For this reason, questions will not count as declarative sentences. Suppose you answer the question: ‘I am not sleepy.’ This is either true or false, and so it is a declarative sentence. Generally, *questions* will not count as declarative sentences, but *answers* will.

‘What is this course about?’ is not a declarative sentence (in our sense). ‘No one knows what this course is about’ is a declarative sentence.

Imperatives Commands are often phrased as imperatives like ‘Wake up!’, ‘Sit up straight’, and so on. These are imperative sentences. Although it might be good for you to sit up straight or it might

not, the command is neither true nor false and it is thus not a declarative sentence. Note, however, that commands are not always phrased as imperatives. ‘You will respect my authority’ *is* either true or false— either you will or you will not— and so it counts as a declarative sentences.

Exclamations ‘Ouch!’ is sometimes called an exclamatory sentence, but it is not the sort of thing which is true or false. ‘That hurt!’, however, is a declarative sentence.

Our focus is only on *declarative sentences* — those sentences which can be true or false — for example ‘spiders have eight legs’. We typically drop the term ‘declarative’ and simply call them sentences, but bear in mind that it is only these sorts of sentences that are relevant in this textbook.

You should not confuse the idea of a sentence that can be true or false with the difference between fact and opinion. Often, sentences in logic will express things that would count as facts— such as ‘spiders have eight legs’ or ‘Kierkegaard liked almonds.’ They can also express things that you might think of as matters of opinion—such as, ‘Almonds are tasty.’ In other words, a sentence is not disqualified from being part of an argument because we don’t know if it is true or false, or because its truth or falsity is a matter of opinion. All that matters is whether it is the sort of thing that could be true or false. If it is, it can play the role of premise or conclusion.

When you are reading a text and putting it in our standard form you should make sure that your premises and conclusions are declarative sentences. You should also make them as clear as possible. Each premise and the conclusion should be able to be read and understood independently. Any context from the original paragraph should be copied over to each of the premises and conclusions. For example:

Donating to charity no strings attached is the most effective way to do so. So if you are going to donate to charity, you should do it this way.

When presenting this we should fill out “this way” with the relevant way. So I’d write:

- Donating to charity no strings attached is the most effective way to do so.
- Therefore: If you are going to donate to charity, you should do so no strings attached.

Practice exercises

At the end of some chapters, there are exercises that review and explore the material covered in the chapter. The problem sheet you need to complete is constructed from these exercises, but the book offers some additional practice if you want more. There is no substitute for actually working through some problems. This course isn't about memorizing facts but about developing a way of thinking.

So here's the first exercise.

A.

1. Are arguments always presented in our standard form?
2. Do conclusions always come after the premises in arguments in texts?
3. Might premises and conclusions be clauses within sentences?
4. Can questions be premises?

B. Write down the conclusion of each of these arguments:

1. It is sunny. So I should take my sunglasses.
2. It must have been sunny. I did wear my sunglasses, after all.
3. No one but you has had their hands in the cookie-jar. And the scene of the crime is littered with cookie-crumbs. You're the culprit!
4. Miss Scarlett and Professor Plum were in the study at the time of the murder. Reverend Green had the candlestick in the ballroom, and we know that there is no blood on his hands. Hence Colonel Mustard did it in the kitchen with the lead-piping. Recall, after all, that the gun had not been fired.
5. Since I do not know that I am not under the spell of a malicious demon, I do not know that this table exists. After all, if I know that this table exists, then I know that I am not under the spell of a malicious demon.
6. Cutting the interest rate will have no effect on the stock market this time round as people have been expecting a rate cut all along. This factor has already been reflected in the market.
7. Virgin would then dominate the rail system. Is that something the government should worry about? Not necessarily. The industry is regulated, and one powerful company might at least offer a more coherent schedule of services than the present arrangement has produced. The reason the industry was broken up into more than 100 companies at privatisation was not operational, but political:

the Conservative government thought it would thus be harder to renationalise. *The Economist* 16.12.2000; used on *critical thinking web*

8. The idea that being vegetarian is better for the environment has, over the last forty years, become a piece of conventional wisdom. But it is simply wrong. A paper from Carnegie Mellon University researchers published this week finds that the diets recommended by the Dietary Guidelines for Americans, which include more fruits and vegetables and less meat, exacts a greater environmental toll than the typical American diet. Shifting to the diets recommended by Dietary Guidelines for American would increase energy use by 38 percent, water use by ten percent and greenhouse gas emissions by six percent, according to the paper.
9. There are no hard numbers, but the evidence from Asia's expatriate community is unequivocal. Three years after its handover from Britain to China, Hong Kong is unlearning English. The city's gweilos (Cantonese for "ghost men") must go to ever greater lengths to catch the oldest taxi driver available to maximize their chances of comprehension. Hotel managers are complaining that they can no longer find enough English- speakers to act as receptionists. Departing tourists, polled at the airport, voice growing frustration at not being understood.

The Economist 20.1.2001, used in *Critical Thinking Web*

C. Write each of the following arguments in the standard form.

- x. It might surprise you, but denoting to charity no strings attached is the most effective way to do so. So if you are going to donate to charity, you should do it this way.

- Answer:

- Denoting to charity no strings attached is the most effective way to do so.
- Therefore: If you are going to donate to charity, you should do so no strings attached.

1. It is sunny. So I should take my sunglasses.
2. It must have been sunny. I did wear my sunglasses, after all.
3. No one but you has had their hands in the cookie-jar. And the scene of the crime is littered with cookie-crumbs. You're the culprit!

4. Kate didn't write it. If Kate or David wrote it, it will be reliable; and it isn't.
5. Since I do not know that I am not under the spell of a malicious demon, I do not know that this table exists. After all, if I know that this table exists, then I know that I am not under the spell of a malicious demon.
6. Miss Scarlett and Professor Plum were in the study at the time of the murder. And Reverend Green had the candlestick in the ballroom, and we know that there is no blood on his hands. Hence Colonel Mustard did it in the kitchen with the lead-piping. Recall, after all, that the gun had not been fired.

CHAPTER 2

The scope of logic

2.1 Consequence and validity

In §1, we talked about arguments, i.e., a collection of sentences (the premises), followed by a single sentence (the conclusion). We said that some words, such as “therefore,” indicate which sentence is supposed to be the conclusion. “Therefore,” of course, suggests that there is a connection between the premises and the conclusion, namely that the conclusion *follows from*, or *is a consequence of*, the premises.

This notion of consequence is one of the primary things logic is concerned with. One might even say that logic is the science of what follows from what. Logic develops theories and tools that tell us when a sentence follows from some others.

What about the following argument:

- Either the butler or the gardener did it.
- The butler didn’t do it.
- Therefore: The gardener did it.

We don’t have any context for what the sentences in this argument refer to. Perhaps you suspect that “did it” here means “was the perpetrator” of some unspecified crime. You might imagine that the argument occurs in a mystery novel or TV show, perhaps spoken by a detective working through the evidence. But even without having any of this information, you probably agree that the argument is a good one in the sense that

whatever the premises refer to, if they are both true, the conclusion cannot but be true as well. If the first premise is true, i.e., it's true that "the butler did it or the gardener did it," then at least one of them "did it," whatever that means. And if the second premise is true, then the butler did not "do it." That leaves only one option: "the gardener did it" must be true. Here, the conclusion follows from the premises. We call arguments that have this property **VALID**.

By way of contrast, consider the following argument:

- If the driver did it, the maid didn't do it.
- The maid didn't do it.
- Therefore: The driver did it.

We still have no idea what is being talked about here. But, again, you probably agree that this argument is different from the previous one in an important respect. If the premises are true, it is not guaranteed that the conclusion is also true. The premises of this argument do not rule out, by themselves, that someone other than the maid or the driver "did it." So there is a case where both premises are true, and yet the driver didn't do it, i.e., the conclusion is not true. In this second argument, the conclusion does not follow from the premises. If, like in this argument, the conclusion does not follow from the premises, we say it is **INVALID**.

2.2 Validity

How did we determine that the second argument is invalid? We pointed to a case in which the premises are true and in which the conclusion is not. This was the scenario where neither the driver nor the maid did it, but some third person did. Let's call such a case a **COUNTEREXAMPLE** to the argument. If there is a counterexample to an argument, the conclusion cannot be a consequence of the premises. For the conclusion to be a consequence of the premises, the truth of the premises must guarantee the truth of the conclusion. If there is a counterexample, the truth of the premises does not guarantee the truth of the conclusion.

As logicians, we want to be able to determine when the conclusion of an argument follows from the premises. And the conclusion is a consequence of the premises if there is no counterexample—no case where the premises are all true but the conclusion is not. This motivates a definition:

A sentence Y is a **CONSEQUENCE** of sentences X_1, \dots, X_n if and only if there is no case where X_1, \dots, X_n are all true and Y is not true. (We then also say that Y **FOLLOWS FROM** X_1, \dots, X_n or that X_1, \dots, X_n **ENTAIL** Y .)

We said that arguments where the conclusion is a consequence of the premises are called valid, and those where the conclusion isn't a consequence of the premises are invalid. That is:

An argument is **VALID** if and only if there is no case where all the premises are true and the conclusion false.

An argument is **INVALID** if and only if it is not valid. That is, there is some case where all the premises are true and the conclusion is false.

2.3 Cases and types of validity

The “definitions” from the previous section are incomplete: it does not tell us what a “case” is or what it means to be “true in a case.” So far we’ve only seen an example: a hypothetical scenario involving three people. Of the three people in the scenario—a driver, a maid, and some third person—the driver and maid didn’t do it, but the third person did. In this scenario, as described, the driver didn’t do it, and so it is a case in which the sentence “the driver did it” is not true. The premises of our second argument are true, but the conclusion is not true: the scenario is a counterexample.

One thing that logicians do is to make the notion of “case” more precise, and investigate which arguments are valid when “case” is made precise in one way or another. If we take “case” to mean “hypothetical scenario” like the counterexample to the second argument, it’s clear that the first argument counts as valid. If we imagine a scenario in which either the butler or the gardener did it, and also the butler didn’t do it, we are automatically imagining a scenario in which the gardener did it. So any hypothetical scenario in which the premises of our first argument are true automatically makes the conclusion of our first argument true. This makes the first argument valid.

Making “case” more specific by interpreting it as “hypothetical sce-

nario” is an advance. But it is not the end of the story. The first problem is that we don’t know what to count as a hypothetical scenario. Are they limited by the laws of physics? By what is conceivable, in a very general sense? What answers we give to these questions determine which arguments we count as valid.

Suppose the answer to the first question is “yes.” Consider the following argument:

- The spaceship *Rocinante* took six hours to reach Jupiter from Tycho space station.
- Therefore: The distance between Tycho space station and Jupiter is less than 14 billion kilometers.

A counterexample to this argument would be a scenario in which the *Rocinante* makes a trip of over 14 billion kilometers in 6 hours, exceeding the speed of light. Since such a scenario is incompatible with the laws of physics, there is no such scenario if hypothetical scenarios have to conform to the laws of physics. If hypothetical scenarios are not limited by the laws of physics, however, there is a counterexample: a scenario where the *Rocinante* travels faster than the speed of light.

Suppose the answer to the second question is “yes,” and consider another argument:

- Priya is an ophthalmologist.
- Therefore: Priya is an eye doctor.

If we’re allowing only conceivable scenarios, this is also a valid argument. If you imagine Priya being an ophthalmologist, you thereby imagine Priya being an eye doctor. That’s just what “ophthalmologist” and “eye doctor” mean. A scenario where Priya is an ophthalmologist but not an eye doctor is ruled out by the conceptual connection between these words.

When we consider cases of various kinds in order to evaluate the validity of an argument, we will make a few assumptions. The first assumption is that every case makes every sentence true or false—at least, every sentence in the argument under consideration. So imagined scenarios have to specify all relevant facts. Any imagined scenario which leaves it undetermined if a sentence in our argument is true will not be considered as a potential counterexample.

Depending on what kinds of cases we consider as potential counterexamples, then, we arrive at different notions of consequence and validity. We might call an argument **NOMOLOGICALLY VALID** if there are

no counterexamples that don't violate the laws of nature, and an argument **CONCEPTUALLY VALID** if there are no counterexamples that don't violate conceptual connections between words. For both of these notions of validity, aspects of the world (e.g., what the laws of nature are) and aspects of the meaning of the sentences in the argument (e.g., that "ophthalmologist" just means a kind of eye doctor) figure into whether an argument is valid.

One distinguishing feature of *logical* consequence, however, is that it should not depend on the content of the premises and conclusion, but only on their logical form. In other words, as logicians we want to develop a theory that can make finer-grained distinctions still. For instance, both

- Either Priya is an ophthalmologist or a dentist.
- Priya isn't a dentist.
- Therefore: Priya is an eye doctor.

and

- Either Priya is an ophthalmologist or a dentist.
- Priya isn't a dentist.
- Therefore: Priya is an ophthalmologist.

are valid arguments. But while the validity of the first depends on the content (i.e., the meaning of "ophthalmologist" and "eye doctor"), the second does not. The second argument is **FORMALLY VALID**. We can describe the "form" of this argument as a pattern, something like this:

- Either a is an F or a G .
- a isn't an F .
- Therefore: a is a G .

Here, a , F , and G are placeholders for appropriate expressions that, when substituted for a , F , and G , turn the pattern into an argument consisting of sentences. For instance,

- Either Mei is a mathematician or a botanist.
- Mei isn't a botanist.
- Therefore: Mei is a mathematician.

is an argument of the same form, but the first argument above is not: we would have to replace F by different expressions (once by "ophthalmologist" and once by "eye doctor") to obtain it from the pattern.

Moreover, the first argument is not formally valid. *Its* form is this:

- Either a is an F or a G .
- a isn't an F .
- Therefore: a is a H .

In this pattern we can replace F by “ophthalmologist” and H by “eye doctor” to obtain the original argument. But here is another argument of the same form:

- Either Mei is a mathematician or a botanist.
- Mei isn't a botanist.
- Therefore: Mei is an acrobat.

This argument is clearly not valid, since we can imagine a mathematician named Mei who is not an acrobat.

Our strategy as logicians will be to come up with a notion of “case” on which an argument turns out to be valid if it is formally valid. Clearly such a notion of “case” will have to violate not just some laws of nature but some laws of English. Since the first argument is invalid in this sense, we must allow as counterexample a case where Priya is an ophthalmologist but not an eye doctor. That case is not a conceivable situation: it is ruled out by the meanings of “ophthalmologist” and “eye doctor.”

2.4 Sound arguments

Before we go on and execute this strategy, a few clarifications. Arguments in our sense, as conclusions which (supposedly) follow from premises, are of course used all the time in everyday, philosophical and scientific discourse. When they are, arguments are given to support or even prove their conclusions. Now, if an argument is valid, it will support its conclusion, but *only if* its premises are all true. Validity rules out the possibility that the premises are true and the conclusion is not true at the same time. It does not, by itself, rule out the possibility that the conclusion is not true, period. In other words, it is perfectly possible for a valid argument to have a conclusion that isn't true!

Consider this example:

- Oranges are either fruit or musical instruments.
- Oranges are not fruit.
- Therefore: Oranges are musical instruments.

The conclusion of this argument is ridiculous. Nevertheless, it follows from the premises. *If* both premises are true, *then* the conclusion just has to be true. So the argument is valid.

Conversely, having true premises and a true conclusion is not enough to make an argument valid. Consider this example:

- London is in England.
- Beijing is in China.
- Therefore: Paris is in France.

The premises and conclusion of this argument are, as a matter of fact, all true, but the argument is invalid. If Paris were to declare independence from the rest of France, then the conclusion would no longer be true, even though both of the premises would remain true. Thus, there is a case where the premises of this argument are true without the conclusion being true. So the argument is invalid.

The important thing to remember is that validity is not about the actual truth or falsity of the sentences in the argument. It is about whether it is *possible* for all the premises to be true and the conclusion to be not true at the same time (in some hypothetical case). What is in fact the case has no special role to play; and what the facts are does not determine whether an argument is valid or not.¹ Nothing about the way things are can by itself determine if an argument is valid. It is often said that logic doesn't care about feelings. Actually, it doesn't care about facts, either.

When we use an argument to prove that its conclusion *is true*, then, we need two things. First, we need the argument to be valid, i.e., we need the conclusion to follow from the premises. But we also need the premises to be true. We will say that an argument is **SOUND** if and only if it is both valid and all of its premises are true.

The flip side of this is that when you want to rebut an argument, you have two options: you can show that (one or more of) the premises are not true, or you can show that the argument is not valid. Logic, however, will only help you with the latter!

2.5 Missing premises

If someone you disagree with makes an invalid argument, you might be tempted just to dismiss it as obviously incorrect. But it's more useful

¹Well, there is one case where it does: if the premises are in fact true and the conclusion is in fact not true, then we live in a counterexample; so the argument is invalid.

(and more charitable) to consider whether there are missing premises that could be filled in that make the argument better. Perhaps the author was assuming that this premise was so obvious that it didn't need to be stated.

For example an author might make the following argument:

- I could not have acted otherwise.
- Therefore: I did not act of my own free will.

This argument is invalid. But, it can be made valid by addition of the premise:

- If I could not have acted otherwise, I did not act of my own free will.

But be careful when you're filling in 'missing' premises. The aim is to help improve the argument, to make it more convincing, so you can assess it fairly. Only add extra premises that seem reasonable, or that you think the original author would agree with. There's no point in adding absurd or unreasonable premises, or premises that the author wouldn't endorse. Then you just create a *strawman* argument – a caricature of the original argument.

“Just how charitable are you supposed to be when criticizing the views of an opponent? If there are obvious contradictions in the opponent's case, then of course you should point them out, forcefully. If there are somewhat hidden contradictions, you should carefully expose them to view—and then dump on them. But the search for hidden contradictions often crosses the line into nitpicking, sea-lawyering, and—as we have seen—outright parody. The thrill of the chase and the conviction that your opponent has to be harboring a confusion somewhere encourages uncharitable interpretation, which gives you an easy target to attack. But such easy targets are typically irrelevant to the real issues at stake and simply waste everybody's time and patience, even if they give amusement to your supporters.”

Daniel C. Dennett (2013). “Intuition Pumps And Other Tools for Thinking”.

Dennett formulates the following four rules (named after Anatol Rapoport) for “how to compose a successful critical commentary”:

1. You should attempt to re-express your target's position so clearly, vividly, and fairly that your target says, "Thanks, I wish I'd thought of putting it that way."
2. You should list any points of agreement (especially if they are not matters of general or widespread agreement).
3. You should mention anything you have learned from your target.
4. Only then are you permitted to say so much as a word of rebuttal or criticism

2.6 Ampliative Arguments

Sometimes there are no plausible missing premises you could add to someone's argument to make it valid. However, this doesn't necessarily mean that the author was wrong or mistaken. Deductively valid arguments with plausible premises are good arguments, but they aren't the only good arguments there are. This is just as well, since many arguments we give in our everyday lives are not deductively valid, even after filling in plausible missing premises. Here's an example:

- In January 1997, it rained in London.
- In January 1998, it rained in London.
- In January 1999, it rained in London.
- In January 2000, it rained in London.
- Therefore: It rains every January in London.

This argument generalises from observations about several cases to a conclusion about all cases—in each year listed, it rained in January, so it does in every year. Such arguments are called **INDUCTIVE** arguments. The argument could be made stronger by adding additional premises before drawing the conclusion: In January 2001, it rained in London; In January 2002... But, however many premises of this form we add, the argument will remain invalid. Even if it has rained in London in every January thus far, it remains possible that London will stay dry next January. The point of all this is that inductive arguments—even good inductive arguments—are not (deductively) valid. They are not watertight. The premises might make the conclusion very likely, but they don't absolutely guarantee its truth. Unlikely though it might be, it is possible for their conclusion to be false, even when all of their premises are true.

Inductive arguments of the sort just given belong to a species of argument called **AMPLIATIVE ARGUMENTS**. This means that the conclusion goes beyond what you find in the premises. That is, the premises don't guarantee, or entail, the conclusion. They do, however, provide some support for it. These arguments are deductively invalid. They may be good and useful, however it is important to know the difference.

In this book, we will set aside the question of what makes for a good ampliative argument and focus instead on sorting the deductively valid arguments from the deductively invalid ones. But we pause here to mention some further forms of ampliative argument.

Inductive arguments, like the one we saw above, allow one to infer from a series of observed cases to a generalization that covers them: from all observed *F*s have been *G*s, we infer all *F*s are *G*s. We use these all the time. Every time I've drunk water from my tap, it's quenched my thirst; therefore, every time I ever drink water from my tap, it will quench my thirst. Every time I've stroked my neighbour's cat, it hasn't bitten me; therefore, every time I ever stroke my neighbour's cat, it won't bite me. And it's a form of arguments much beloved by scientists. Every time we've measured the acceleration of a body falling, it's matched Newton's theory, therefore, all bodies are governed by Newton's theory. The premises of these argument seem to make their conclusions likely without guaranteeing them. The areas of philosophy called inductive logic or confirmation theory try to make precise what that means and why it's true. And of course inductive arguments can go wrong. Before I visited Australia, every swan I'd ever seen was white, and so I concluded that all swans were white; but when I visited Australia, I realised my conclusion was wrong, because some swans there are black.

A closely related, but different form of argument, is **STATISTICAL**. Here, we start with an observation about the proportion of *F*s that are *G*s in a sample that we've observed, and we infer that the same proportion of *F*s are *G*s in general. So, for instance, if I poll 1,000 people in Scotland eligible to vote in a second independence referendum, and 600 say that they'll vote yes, I might infer that 60% of all eligible voters will vote yes. Or if I test 1,000,000 people in England for an active infection, and 20,000 test positive, I might infer that 2% of the whole population has an active infection. How good these argument are depends on a number of things, and these are studied by statisticians. For instance, suppose you picked the 1,000 Scottish voters entirely at random from an anonymised version of the electoral register. But suppose that, when

you deanonymised, you learned that, by chance, all of the people you'd picked were over 65, or they all lived on the Isle of Skye. Then you might worry that your sample, though random, was unrepresentative of the population as a whole. This question is a genuine concern for randomised controlled trials in medicine.

Abductive arguments provide an inference from a phenomenon you've observed to the *best explanation* of that phenomenon: from *E*, and the best explanation of *E* is *H*, you might conclude *H*. Again, this is extremely widespread. A classic sort of example would be the inferences that detectives draw during their investigations. They look at the evidence and the possible explanations of it, and they tend to conclude in favour of the best one. And similarly for doctors looking at a patient's suite of symptoms and trying to discover what ails them. Another important example comes from science. Here is Charles Darwin explaining what convinces him of his theory of natural selection:

"It can hardly be supposed that a false theory would explain, in so satisfactory a manner as does the theory of natural selection, the several large classes of facts above specified. It has recently been objected that this is an unsafe method of arguing; but it is a method used in judging of the common events of life, and has often been used by the greatest natural philosophers."

(Charles Darwin, *On the origin of species by means of natural selection* (6th ed.). London: John Murray)

Practice exercises

A.

1. What kind of things are valid or invalid?
2. When is an argument said to be valid?
3. When is an argument said to be sound?

B. Are the following valid? If it is invalid, describe a counterexample.

- x.
 - Every good zoo has a giraffe.
 - It is a zoo.
 - Therefore: It has a giraffe.
- Invalid. It is a zoo, but not a good one. (And has a giraffe.)

1.
 - Everyone in group 1 handed in their homework.
 - Jenny is in group 1.
 - Therefore: Jenny handed in her homework.
2.
 - If she won the lottery then she is rich.
 - She is rich.
 - Therefore: She won the lottery.
3.
 - Most people are scared of spiders.
 - Therefore: Oscar is scared of spiders.
4.
 - She is a donkey.
 - Therefore: She does not talk.

C. Which of the following arguments are conceptually valid? Which are conceptually invalid?

- Socrates is a man.
- All men are carrots.
- Therefore: Socrates is a carrot.

- Abe Lincoln was either born in Illinois or he was once president.
- Abe Lincoln was never president.
- Therefore: Abe Lincoln was born in Illinois.

- If I pull the trigger, Abe Lincoln will die.
- I do not pull the trigger.
- Therefore: Abe Lincoln will not die.

- Abe Lincoln was either from France or from Luxemborg.
- Abe Lincoln was not from Luxemborg.
- Therefore: Abe Lincoln was from France.

- If the world were to end today, then I would not need to get up tomorrow morning.
- I will need to get up tomorrow morning.
- Therefore: The world will not end today.

- Joe is now 19 years old.
- Joe is now 87 years old.
- Therefore: Bob is now 20 years old.

D. Could there be:

1. A valid argument that has one false premise and one true premise?
2. A valid argument that has only false premises?
3. A valid argument with only false premises and a false conclusion?
4. An invalid argument that can be made valid by the addition of a new premise?
5. A valid argument that can be made invalid by the addition of a new premise?

In each case: if so, give an example; if not, explain why not.

CHAPTER 3

Other logical notions

In §2, we introduced the ideas of consequence and of valid argument. This is one of the most important ideas in logic. In this section, we will introduce are some similarly important ideas. They all rely, as did validity, on the idea that sentences are true (or not) in cases. For the rest of this section, we'll take cases in the sense of conceivable scenario, i.e., in the sense in which we used them to define conceptual validity. The points we made about different kinds of validity can be made about our new notions along similar lines: if we use a different idea of what counts as a “case” we will get different notions. And as logicians we will, eventually, consider a more permissive definition of case than we do here.

3.1 Joint possibility

Consider these two sentences:

- B1. Jane's only brother is shorter than her.
- B2. Jane's only brother is taller than her.

Logic alone cannot tell us which, if either, of these sentences is true. Yet we can say that *if* the first sentence (B1) is true, *then* the second sentence (B2) must be false. Similarly, if B2 is true, then B1 must be false. There is no possible scenario where both sentences are true to-

gether. These sentences are incompatible with each other, they cannot all be true at the same time. This motivates the following definition:

Sentences are **JOINTLY POSSIBLE** if and only if there is a case where they are all true together.

B1 and B2 are *jointly impossible*, while, say, the following two sentences are jointly possible:

- B1. Jane's only brother is shorter than her.
- B2. Jane's only brother is younger than her.

We can ask about the joint possibility of any number of sentences. For example, consider the following four sentences:

- G1. There are at least four giraffes at the wild animal park.
- G2. There are exactly seven gorillas at the wild animal park.
- G3. There are not more than two martians at the wild animal park.
- G4. Every giraffe at the wild animal park is a martian.

G1 and G4 together entail that there are at least four martian giraffes at the park. This conflicts with G3, which implies that there are no more than two martian giraffes there. So the sentences G1–G4 are jointly impossible. They cannot all be true together. (Note that the sentences G1, G3 and G4 are jointly impossible. But if sentences are already jointly impossible, adding an extra sentence to the mix cannot make them jointly possible!)

3.2 Necessary truths, necessary falsehoods, and contingency

In assessing arguments for validity, we care about what would be true *if* the premises were true, but some sentences just *must* be true. Consider these sentences:

- 1. It is raining.
- 2. Either it is raining here, or it is not.
- 3. It is both raining here and not raining here.

In order to know if sentence 1 is true, you would need to look outside or check the weather channel. It might be true; it might be false. A

sentence which is capable of being true and capable of being false (in different circumstances, of course) is called **CONTINGENT**.

Sentence 2 is different. You do not need to look outside to know that it is true. Regardless of what the weather is like, it is either raining or it is not. That is a **NECESSARY TRUTH**.

Equally, you do not need to check the weather to determine whether or not sentence 3 is true. It must be false, simply as a matter of logic. It might be raining here and not raining across town; it might be raining now but stop raining even as you finish this sentence; but it is impossible for it to be both raining and not raining in the same place and at the same time. So, whatever the world is like, it is not both raining here and not raining here. It is a **NECESSARY FALSEHOOD**.

Something might *always* be true and still be contingent. For instance, if there never were a time when the universe contained fewer than seven things, then the sentence 'At least seven things exist' would always be true. Yet the sentence is contingent: the world could have been much, much smaller than it is, and then the sentence would have been false.

3.3 Necessary equivalence

We can also ask about the logical relations *between* two sentences. For example:

- John went to the store after he washed the dishes.
- John washed the dishes before he went to the store.

These two sentences are both contingent, since John might not have gone to the store or washed dishes at all. Yet they must have the same truth-value. If either of the sentences is true, then they both are; if either of the sentences is false, then they both are. When two sentences have the same truth value in every case, we say that they are **NECESSARILY EQUIVALENT**.

Summary of logical notions

- ▷ An argument is **VALID** if there is no case where the premises are all true and the conclusion is not; it is **INVALID** otherwise.
- ▷ A **NECESSARY TRUTH** is a sentence that is true in every case.
- ▷ A **NECESSARY FALSEHOOD** is a sentence that is false in every case.
- ▷ A **CONTINGENT SENTENCE** is a sentence that is neither a necessary truth nor a necessary falsehood; a sentence that is true in some case and false in some other case.
- ▷ Two sentences are **NECESSARILY EQUIVALENT** if, in every case, they are both true or both false.
- ▷ A collection of sentences is **JOINTLY POSSIBLE** if there is a case where they are all true together; it is **JOINTLY IMPOSSIBLE** otherwise.

Practice exercises

A. For each of the following: Is it a necessary truth, a necessary falsehood, or contingent?

1. Caesar crossed the Rubicon.
2. Someone once crossed the Rubicon.
3. No one has ever crossed the Rubicon.
4. If Caesar crossed the Rubicon, then someone has.
5. Even though Caesar crossed the Rubicon, no one has ever crossed the Rubicon.
6. If anyone has ever crossed the Rubicon, it was Caesar.

B. For each of the following: Is it a necessary truth, a necessary falsehood, or contingent?

1. Elephants dissolve in water.
2. Wood is a light, durable substance useful for building things.
3. If wood were a good building material, it would be useful for building things.

4. I live in a three story building that is two stories tall.
5. If gerbils were mammals they would nurse their young.

C. Which of the following pairs of sentences are necessarily equivalent?

1. Elephants dissolve in water.
If you put an elephant in water, it will disintegrate.
2. All mammals dissolve in water.
If you put an elephant in water, it will disintegrate.
3. George Bush was the 43rd president.
Barack Obama is the 44th president.
4. Barack Obama is the 44th president.
Barack Obama was president immediately after the 43rd president.
5. Elephants dissolve in water.
All mammals dissolve in water.

D. Which of the following pairs of sentences are necessarily equivalent?

1. Thelonious Monk played piano.
John Coltrane played tenor sax.
2. Thelonious Monk played gigs with John Coltrane.
John Coltrane played gigs with Thelonious Monk.
3. All professional piano players have big hands.
Piano player Bud Powell had big hands.
4. Bud Powell suffered from severe mental illness.
All piano players suffer from severe mental illness.
5. John Coltrane was deeply religious.
John Coltrane viewed music as an expression of spirituality.

E. Consider the following sentences:

- G₁ There are at least four giraffes at the wild animal park.
- G₂ There are exactly seven gorillas at the wild animal park.
- G₃ There are not more than two Martians at the wild animal park.
- G₄ Every giraffe at the wild animal park is a Martian.

Now consider each of the following collections of sentences. Which are jointly possible? Which are jointly impossible?

1. Sentences G₂, G₃, and G₄

2. Sentences G_1 , G_3 , and G_4
3. Sentences G_1 , G_2 , and G_4
4. Sentences G_1 , G_2 , and G_3

F. Consider the following sentences.

M_1 All people are mortal.

M_2 Socrates is a person.

M_3 Socrates will never die.

M_4 Socrates is mortal.

Which combinations of sentences are jointly possible? Mark each “possible” or “impossible.”

1. Sentences M_1 , M_2 , and M_3
2. Sentences M_2 , M_3 , and M_4
3. Sentences M_2 and M_3
4. Sentences M_1 and M_4
5. Sentences M_1 , M_2 , M_3 , and M_4

G. Which of the following is possible? If it is possible, give an example. If it is not possible, explain why.

1. A valid argument that has one false premise and one true premise
2. A valid argument that has a false conclusion
3. A valid argument, the conclusion of which is a necessary falsehood
4. An invalid argument, the conclusion of which is a necessary truth
5. A necessary truth that is contingent
6. Two necessarily equivalent sentences, both of which are necessary truths
7. Two necessarily equivalent sentences, one of which is a necessary truth and one of which is contingent
8. Two necessarily equivalent sentences that together are jointly impossible
9. A jointly possible collection of sentences that contains a necessary falsehood
10. A jointly impossible set of sentences that contains a necessary truth

H. Which of the following is possible? If it is possible, give an example. If it is not possible, explain why.

1. A valid argument, whose premises are all necessary truths, and whose conclusion is contingent
2. A valid argument with true premises and a false conclusion
3. A jointly possible collection of sentences that contains two sentences that are not necessarily equivalent
4. A jointly possible collection of sentences, all of which are contingent
5. A false necessary truth
6. A valid argument with false premises
7. A necessarily equivalent pair of sentences that are not jointly possible
8. A necessary truth that is also a necessary falsehood
9. A jointly possible collection of sentences that are all necessary falsehoods

PART II

Truth- functional logic

CHAPTER 4

First steps to symbolization

4.1 Validity in virtue of form

Consider this argument:

- It is raining outside.
- If it is raining outside, then Jenny is miserable.
- Therefore: Jenny is miserable.

and another argument:

- Jenny is an anarcho-syndicalist.
- If Jenny is an anarcho-syndicalist, then Dipan is an avid reader of Tolstoy.
- Therefore: Dipan is an avid reader of Tolstoy.

Both arguments are valid, and there is a straightforward sense in which we can say that they share a common structure. We might express the structure thus:

- A
- If A, then B
- Therefore: B

This looks like an excellent argument *structure*. Indeed, surely any argument with this *structure* will be valid.

What about:

- Jenny is miserable.
- If it is raining outside, then Jenny is miserable.
- Therefore: It is raining outside.

The form of this argument is:

- B
- If A then B
- Therefore: A

Arguments of this form are generally invalid.

Be careful, though, not every argument of this form is sure to be invalid. It's possible to have an argument of this form that's valid – see if you can work out how! But most arguments of this form are invalid.

Consider an argument like:

- Jenny is happy or sad.
- Jenny is not happy.
- Therefore: Jenny is sad.

Again, this is a valid argument. The structure here is something like:

- A or B
- not- A
- Therefore: B

A superb structure! Here is another example:

- It's not the case that Jim both studied hard and acted in lots of plays.
- Jim studied hard
- Therefore: Jim did not act in lots of plays.

This valid argument has a structure which we might represent thus:

- not- $(A$ and $B)$
- A
- Therefore: not- B

These examples illustrate an important idea, which we might describe as *validity in virtue of form*. The validity of the arguments just considered has nothing very much to do with the meanings of English expressions like 'Jenny is miserable', 'Dipan is an avid reader of Tolstoy', or 'Jim acted in lots of plays'. If it has to do with meanings at all, it is with the meanings of phrases like 'and', 'or', 'not,' and 'if... , then... '.

In Parts II–IV, we are going to develop a formal language which allows us to symbolize many arguments in such a way as to show that they are valid in virtue of their form. That language will be *truth-functional logic*, or TFL.

It will have sentences like

$$(A \ \& \ (B \rightarrow \neg C))$$

4.2 Validity for special reasons

There are plenty of arguments that are valid, but not for reasons relating to their form. Take an example:

- Juanita is a vixen
- Therefore: Juanita is a fox

It is impossible for the premise to be true and the conclusion false. So the argument is valid. However, the validity is not related to the form of the argument. Here is an invalid argument with the same form:

- Juanita is a vixen
- Therefore: Juanita is a cathedral

This might suggest that the validity of the first argument *is* keyed to the meaning of the words ‘vixen’ and ‘fox’. But, whether or not that is right, it is not simply the *shape* of the argument that makes it valid. Equally, consider the argument:

- The sculpture is green all over.
- Therefore: The sculpture is not red all over.

Again, it seems impossible for the premise to be true and the conclusion false, for nothing can be both green all over and red all over. So the argument is valid, but here is an invalid argument with the same form:

- The sculpture is green all over.
- Therefore: The sculpture is not shiny all over.

The argument is invalid, since it is possible to be green all over and shiny all over. (One might paint their nails with an elegant shiny green varnish.) Plausibly, the validity of the first argument is keyed to the way that colours (or colour-words) interact, but, whether or not that is right, it is not simply the *shape* of the argument that makes it valid.

The important moral can be stated as follows. *At best, TFL will help us to understand arguments that are valid due to their form.*

4.3 Atomic sentences

We started isolating the form of an argument, in §4.1, by replacing *sub-sentences* of sentences with individual letters. Thus in the first example of this section, ‘it is raining outside’ is a subsentence of ‘If it is raining outside, then Jenny is miserable’, and we replaced this subsentence with ‘A’.

Our artificial language, TFL, pursues this idea absolutely ruthlessly. We start with some *atomic sentences*. These will be the basic building blocks out of which more complex sentences are built. We will use uppercase Roman letters for atomic sentences of TFL (except for X , Y , and Z which we reserve for metavariables). There are only twenty-three letters A – W , but there is no limit to the number of atomic sentences that we might want to consider. By adding subscripts to letters, we obtain new atomic sentences. So, here are five different atomic sentences of TFL:

$$A, P, P_1, P_2, A_{234}$$

We will use atomic sentences to represent, or *symbolize*, certain English sentences. To do this, we provide a **SYMBOLIZATION KEY**, such as the following:

A : It is raining outside
 C : Jenny is miserable

In doing this, we are not fixing this symbolization *once and for all*. We are just saying that, for the time being, we will think of the atomic sentence of TFL, A , as symbolizing the English sentence ‘It is raining outside’, and the atomic sentence of TFL, C , as symbolizing the English sentence ‘Jenny is miserable’. Later, when we are dealing with different sentences or different arguments, we can provide a new symbolization key; as it might be:

A : Jenny is an anarcho-syndicalist
 C : Dipan is an avid reader of Tolstoy

It is important to understand that whatever structure an English sentence might have is lost when it is symbolized by an atomic sentence of TFL. From the point of view of TFL, an atomic sentence is just a letter. It can be used to build more complex sentences, but it cannot be taken apart.

CHAPTER 5

Connectives

In the previous chapter, we considered symbolizing fairly basic English sentences with atomic sentences of TFL. This leaves us wanting to deal with the English expressions ‘and’, ‘or’, ‘not’, and so forth. These are *connectives*—they can be used to form new sentences out of old ones. In TFL, we will make use of logical connectives to build complex sentences from atomic components. There are five logical connectives in TFL. This table summarises them, and they are explained throughout this section.

symbol	what it is called	rough meaning
\neg	negation	‘It is not the case that...’
$\&$	conjunction	‘Both... and ...’
\vee	disjunction	‘... or ...’
\rightarrow	conditional	‘If ... then ...’
\leftrightarrow	biconditional	‘... if and only if ...’

These are not the only connectives of English of interest. Others are, e.g., ‘unless’, ‘neither ... nor ...’, and ‘because’. We will see that the first two can be expressed by the connectives we will discuss, while the last cannot. ‘Because’, in contrast to the others, is not *truth functional*.

5.1 Negation

Consider how we might symbolize these sentences:

1. Mary is in Barcelona.
2. It is not the case that Mary is in Barcelona.

3. Mary is not in Barcelona.

In order to symbolize sentence **1**, we will need an atomic sentence. We might offer this symbolization key:

B: Mary is in Barcelona.

Since sentence **2** is obviously related to sentence **1**, we will not want to symbolize it with a completely different sentence. Roughly, sentence **2** means something like ‘It is not the case that *B*’. In order to symbolize this, we need a symbol for negation. We will use ‘ \neg ’. Now we can symbolize sentence **2** with $\neg B$.

Sentence **3** also contains the word ‘not’, and it is obviously equivalent to sentence **2**. As such, we can also symbolize it with $\neg B$.

If a sentence can be paraphrased as ‘it is not the case that *X*’ it can be symbolised as $\neg X$.

It will help to offer a few more examples:

4. The widget can be replaced.
5. The widget is irreplaceable.
6. The widget is not irreplaceable.

Let us use the following representation key:

R: The widget is replaceable

Sentence **4** can now be symbolized by *R*. Moving on to sentence **5**: saying the widget is irreplaceable means that it is not the case that the widget is replaceable. So even though sentence **5** does not contain the word ‘not’, we will symbolize it as follows: $\neg R$.

Sentence **6** can be paraphrased as ‘It is not the case that the widget is irreplaceable.’ Which can again be paraphrased as ‘It is not the case that it is not the case that the widget is replaceable’. So we might symbolize this English sentence with the TFL sentence $\neg\neg R$.

But some care is needed when handling negations. Consider:

7. Jane is happy.
8. Jane is unhappy.

If we let the TFL-sentence *H* symbolize ‘Jane is happy’, then we can symbolize sentence **7** as *H*. However, it would be a mistake to symbolize

sentence 8 with $\neg H$. If Jane is unhappy, then she is not happy; but sentence 8 does not mean the same thing as ‘It is not the case that Jane is happy’. Jane might be neither happy nor unhappy; she might be in a state of blank indifference. In order to symbolize sentence 8, then, we would need a new atomic sentence of TFL.

5.2 Conjunction

Consider these sentences:

9. Adam is athletic.
10. Barbara is athletic.
11. Adam is athletic, and Barbara is also athletic.

We will need separate atomic sentences of TFL to symbolize sentences 9 and 10; perhaps

A: Adam is athletic.

B: Barbara is athletic.

Sentence 9 can now be symbolized as *A*, and sentence 10 can be symbolized as *B*. Sentence 11 roughly says ‘A and B’. We need another symbol, to deal with ‘and’. We will use ‘&’. Thus we will symbolize it as (*A* & *B*). This connective is called **CONJUNCTION**. We also say that *A* and *B* are the two **CONJUNCTS** of the conjunction (*A* & *B*).

If a sentence can be paraphrased as ‘*X* and *Y*’
it can be symbolised as *X* & *Y*.

Notice that we make no attempt to symbolize the word ‘also’ in sentence 11. Words like ‘both’ and ‘also’ function to draw our attention to the fact that two things are being conjoined. Maybe they affect the emphasis of a sentence, but we will not (and cannot) symbolize such things in TFL.

Some more examples will bring out this point:

12. Barbara is athletic and energetic.
13. Barbara and Adam are both athletic.
14. Although Barbara is energetic, she is not athletic.
15. Adam is athletic, but Barbara is more athletic than him.

Sentence 12 is obviously a conjunction. The sentence says two things (about Barbara). In English, it is permissible to refer to Barbara only once. It *might* be tempting to think that we need to symbolize sentence 12 with something along the lines of '*B* and energetic'. This would be a mistake. Once we symbolize part of a sentence as *B*, any further structure is lost, as *B* is an atomic sentence of TFL. Conversely, 'energetic' is not an English sentence at all. What we are aiming for is something like '*B* and Barbara is energetic'. So we need to add another sentence letter to the symbolization key. Let *E* symbolize 'Barbara is energetic'. Now the entire sentence can be symbolized as (*B* & *E*).

Sentence 13 says one thing about two different subjects. It says of both Barbara and Adam that they are athletic, even though in English we use the word 'athletic' only once. The sentence can be paraphrased as 'Barbara is athletic, and Adam is athletic'. We can symbolize this in TFL as (*B* & *A*), using the same symbolization key that we have been using.

Sentence 14 is slightly more complicated. The word 'although' sets up a contrast between the first part of the sentence and the second part. Nevertheless, the sentence tells us both that Barbara is energetic and that she is not athletic. In order to make each of the conjuncts an atomic sentence, we need to replace 'she' with 'Barbara'. So we can paraphrase sentence 14 as, '*Both* Barbara is energetic, *and* Barbara is not athletic'. The second conjunct contains a negation, so we paraphrase further: '*Both* Barbara is energetic *and it is not the case that* Barbara is athletic'. Now we can symbolize this with the TFL sentence (*E* & $\neg B$). Note that we have lost all sorts of nuance in this symbolization. There is a distinct difference in tone between sentence 14 and 'Both Barbara is energetic and it is not the case that Barbara is athletic'. TFL does not (and cannot) preserve these nuances.

Sentence 15 raises similar issues. There is a contrastive structure, but this is not something that TFL can deal with. So we can paraphrase the sentence as '*Both* Adam is athletic, *and* Barbara is more athletic than Adam'. (Notice that we once again replace the pronoun 'him' with 'Adam'.) How should we deal with the second conjunct? We already have the sentence letter *A*, which is being used to symbolize 'Adam is athletic', and the sentence *B* which is being used to symbolize 'Barbara is athletic'; but neither of these concerns their relative athleticism. So, to symbolize the entire sentence, we need a new sentence letter. Let the TFL sentence *R* symbolize the English sentence 'Barbara is more athletic than Adam'. Now we can symbolize sentence 15 by (*A* & *R*).

We can add these to our toolbox for symbolisation:

If a sentence can be paraphrased as

- ▷ ‘ X and Y ’
- ▷ ‘ X but Y ’
- ▷ ‘Both X and Y ’
- ▷ ‘Although X , Y ’

it can be symbolised as $X \& Y$.

5.3 Disjunction

Consider these sentences:

16. Fatima will play videogames, or she will watch movies.
17. Fatima or Omar will play videogames.

For these sentences we can use this symbolization key:

F : Fatima will play videogames.

O : Omar will play videogames.

M : Fatima will watch movies.

However, we will again need to introduce a new symbol. Sentence **16** is symbolized by $(F \vee M)$. The connective is called **DISJUNCTION**. We also say that F and M are the **DISJUNCTS** of the disjunction $(F \vee M)$.

Sentence **17** is only slightly more complicated. There are two subjects, but the English sentence only gives the verb once. However, we can paraphrase sentence **17** as ‘Fatima will play videogames, or Omar will play videogames’. Now we can obviously symbolize it by $(F \vee O)$ again.

If a sentence can be paraphrased as ‘ X or Y ’
it can be symbolised as $X \vee Y$.

X and Y need to be the sorts of things that can be whole sentences. We cannot simply take X to be ‘Fatima’, instead we first need to paraphrase it as in **16**

Sometimes in English, the word ‘or’ is used in a way that excludes the possibility that both disjuncts are true. This is called an **EXCLUSIVE**

OR. An *exclusive or* is clearly intended when it says, on a restaurant menu, ‘Entrees come with either soup or salad’: you may have soup; you may have salad; but, if you want *both* soup *and* salad, then you have to pay extra.

At other times, the word ‘or’ allows for the possibility that both disjuncts might be true. This is probably the case with sentence 17, above. Fatima might play videogames alone, Omar might play videogames alone, or they might both play. Sentence 17 merely says that *at least* one of them plays videogames. This is called an **INCLUSIVE OR**. The TFL symbol ‘ \vee ’ always symbolizes an *inclusive or*.

TFL can symbolise the exclusive or, it just doesn’t do it with ‘ \vee ’. We will discuss this in §??

5.4 Conditional

Consider the sentence

18. If Jean is in Paris, then she is in France.

Let’s use the following symbolization key:

P : Jean is in Paris.
 F : Jean is in France

Sentence 18 is roughly of this form: ‘if P , then F ’. We will use the symbol ‘ \rightarrow ’ to symbolize this ‘if... then...’ structure. So we symbolize sentence 18 by $(P \rightarrow F)$. The connective ‘ \rightarrow ’ is called **THE CONDITIONAL**. Here, P is called the **ANTECEDENT** of the conditional $(P \rightarrow F)$, and F is called the **CONSEQUENT**.

If a sentence can be paraphrased as
‘If X , then Y ’ it can be symbolised as $X \rightarrow Y$.

Now consider the sentences

19. If Jean is in Paris, she’s in France.
20. Jean is in France if she is in Paris.

Sentence 19 and 20 are just rephrasings of 18. So we will again symbolise them as $(P \rightarrow F)$.

Now consider

21. Jean is in France only if she is in Paris.

21 is also a conditional. In 18–20 we took as an antecedent the part of sentence that has the word ‘if’ in it. It might then be tempting to do the same here and symbolize this as $(P \rightarrow F)$. That would be a mistake. Your knowledge of geography tells you that sentence 18 is unproblematically true: there is no way for Jean to be in Paris that doesn’t involve Jean being in France. But sentence 21 is not so straightforward: were Jean in Dieppe, Lyons, or Toulouse, Jean would be in France without being in Paris, thereby rendering sentence 21 false. Since geography alone dictates the truth of sentence 18, whereas travel plans (say) are needed to know the truth of sentence 21, they must mean different things. In fact, sentence 21 can be paraphrased as ‘If Jean is in France, then Jean is in Paris’. So we can symbolize it by $(F \rightarrow P)$: the other way around to 18.

If a sentence can be paraphrased as

- ▷ ‘If X , then Y ’
- ▷ ‘If X , Y ’
- ▷ ‘ Y if X ’
- ▷ ‘ X only if Y ’

it can be symbolised as $X \rightarrow Y$.

It is important to bear in mind that the connective ‘ \rightarrow ’ tells us only that, if the antecedent is true, then the consequent is true. It says nothing about a *causal* connection between two events (for example). In fact, we lose a huge amount when we use \rightarrow to symbolize English conditionals. We will return to this in §§11.1 and 17.

5.5 Biconditional

Consider these sentences:

22. Laika is a dog only if she is a mammal
23. Laika is a dog if she is a mammal
24. Laika is a dog if and only if she is a mammal

We will use the following symbolization key:

D : Laika is a dog

M : Laika is a mammal

Sentence 22, for reasons discussed above, can be symbolized by $D \rightarrow M$.

Sentence 23 is importantly different. It can be paraphrased as, ‘If Laika is a mammal then Laika is a dog’. So it can be symbolized by $M \rightarrow D$.

Sentence 24 says something stronger than either 22 or 23. It can be paraphrased as ‘Laika is a dog if Laika is a mammal, and Laika is a dog only if Laika is a mammal’. This is just the conjunction of sentences 22 and 23. So we can symbolize it as $((D \rightarrow M) \& (M \rightarrow D))$. We call this a **BICONDITIONAL**, because it entails the conditional in both directions.

We could treat every biconditional this way. So, just as we do not need a new TFL symbol to deal with *exclusive or*, we do not really need a new TFL symbol to deal with biconditionals. Because the biconditional occurs so often, however, we will use the symbol ‘ \leftrightarrow ’ for it. We can then symbolize sentence 24 with the TFL sentence $(D \leftrightarrow M)$.

The expression ‘if and only if’ occurs a lot especially in philosophy, mathematics, and logic. For brevity, we can abbreviate it with the snappier word ‘iff’. We will follow this practice. So ‘if’ with only *one* ‘f’ is the English conditional. But ‘iff’ with *two* ‘f’s is the English biconditional. Armed with this we can say:

If a sentence can be paraphrased as ‘ X if and only if Y ’ it can be symbolised as $X \leftrightarrow Y$.

A word of caution. Ordinary speakers of English often use ‘if ..., then...’ when they really mean to use something more like ‘...if and only if ...’. Perhaps your parents told you, when you were a child: ‘if you don’t eat your greens, you won’t get any dessert’. Suppose you ate your greens, but that your parents refused to give you any dessert, on the grounds that they were only committed to the *conditional* (roughly ‘if you get dessert, then you will have eaten your greens’), rather than the biconditional (roughly, ‘you get dessert iff you eat your greens’). Well, a tantrum would rightly ensue. So, be aware of this when interpreting people; but in your own writing, make sure you use the biconditional iff you mean to.

Practice exercises

A. Using the symbolization key given, symbolize each English sentence in TFL.

- A*: Mister Ace was murdered.
- B*: The butler did it.
- C*: The cook did it.
- D*: The Duchess is lying.
- E*: Mister Edge was murdered.
- F*: The murder weapon was a frying pan.

1. Mister Ace and Mister Edge were murdered.
2. The butler or the cook did it.
3. The butler didn't do it.
4. If Mister Ace was murdered, then the cook did it.
5. The cook did it if Mister Edge was murdered.
6. Either the butler did it, or the Duchess is lying.
7. The cook did it only if the Duchess is lying.
8. If the murder weapon was a frying pan, then the culprit was the cook.
9. If the murder weapon was not a frying pan, then the culprit was either the cook or the butler.
10. Of course the Duchess is lying!

B. Using the symbolization key given, symbolize each English sentence in TFL.

- E*₁: Ava is an electrician.
- E*₂: Harrison is an electrician.
- F*₁: Ava is a firefighter.
- F*₂: Harrison is a firefighter.
- S*₁: Ava is satisfied with her career.
- S*₂: Harrison is satisfied with his career.

1. Ava and Harrison are both electricians.
2. If Ava is a firefighter, then she is satisfied with her career.
3. Harrison is an unsatisfied electrician.
4. Harrison is satisfied only if he is a firefighter.

C. Give a symbolization key and symbolize the following English sentences in TFL.

1. Alice and Bob are both spies.
2. If either Alice or Bob is a spy, then the code has been broken.
3. If neither Alice nor Bob is a spy, then the code remains unbroken.
4. The German embassy will be in an uproar, unless someone has broken the code.
5. Either the code has been broken or it has not, but the German embassy will be in an uproar regardless.
6. Either Alice or Bob is a spy, but not both.

D. Give a symbolization key and symbolize the following English sentences in TFL.

1. If there is food to be found in the pridelands, then Rafiki will talk about squashed bananas.
2. Rafiki will talk about squashed bananas unless Simba is alive.
3. Rafiki will either talk about squashed bananas or he won't, but there is food to be found in the pridelands regardless.
4. Scar will remain as king if and only if there is food to be found in the pridelands.
5. If Simba is alive, then Scar will not remain as king.

E. For each argument, write a symbolization key and symbolize all of the sentences of the argument in TFL.

1. If Dorothy plays the piano in the morning, then Roger wakes up cranky. Dorothy plays piano in the morning unless she is distracted. So if Roger does not wake up cranky, then Dorothy must be distracted.
2. It will either rain or snow on Tuesday. If it rains, Neville will be sad. If it snows, Neville will be cold. Therefore, Neville will either be sad or cold on Tuesday.
3. If Zoog remembered to do his chores, then things are clean but not neat. If he forgot, then things are neat but not clean. Therefore, things are either neat or clean; but not both.

F. For each argument, write a symbolization key and symbolize the argument as well as possible in TFL. The part of the passage in *italics* is there to provide context for the argument, and doesn't need to be symbolized.

1. It is going to rain soon. I know because my leg is hurting, and my leg hurts if it's going to rain.

2. *Spider-man tries to figure out the bad guy's plan.* If Doctor Octopus gets the uranium, he will blackmail the city. I am certain of this because if Doctor Octopus gets the uranium, he can make a dirty bomb, and if he can make a dirty bomb, he will blackmail the city.
3. *A westerner tries to predict the policies of the Chinese government.* If the Chinese government cannot solve the water shortages in Beijing, they will have to move the capital. They don't want to move the capital. Therefore they must solve the water shortage. But the only way to solve the water shortage is to divert almost all the water from the Yangzi river northward. Therefore the Chinese government will go with the project to divert water from the south to the north.

CHAPTER 6

Sentences of TFL

We have seen connectives in TFL, but there is another part of the language of TFL: brackets. These provide TFL with its grammar.

Sentences of TFL will be things like:

$$A \vee (B \& C)$$

$$\neg(A \& B)$$

There are two purposes of grammar.

The first is to avoid nonsense. Just as in English we want to avoid

The and dog brown or is.

in TFL we want to avoid

$$A \& \vee B \rightarrow$$

which would be nonsense.

The second purpose is to avoid ambiguity. Just as we wish to avoid

John's tired and Sue's tall or Rob's short

in English, in TFL we wish to avoid ambiguity. Brackets are sort of like punctuation in English, they help us know what goes with what.

1. John's tired and Sue's tall or Rob's short

By adding commas this can either be read as:

2. John's tired, and Sue's tall or Rob's short.
3. John's tired and Sue's tall, or Rob's short

In TFL we follow mathematics in using brackets to do this.

4. John's tired and Sue's tall or Rob's short

By adding commas this can either be read as:

5. John's tired, and Sue's tall or Rob's short.
6. John's tired and Sue's tall, or Rob's short

In mathematics,

7. $9 + 3 \times 4$

can either be read as:

8. $9 + (3 \times 4)$ $(= 9 + 12 = 21)$
9. $(9 + 3) \times 4$ $(= 12 \times 4 = 48)$

In TFL,

10. $A \& B \vee C$

can either be read as:

11. $A \& (B \vee C)$
12. $(A \& B) \vee C$

In fact, we will say that $A \& B \vee C$ is not a sentence of TFL at all. It is only expressions which have proper grammar which count as sentences.

To make this precise, this chapter offers a formal definition of what it is to be a sentence in TFL.

6.1 Syntactic rules of TFL

We give rules for what counts as a sentence which will together give us a definition.

We start with the rule:

1. Any uppercase Roman letters A – W , or with subscripts, e.g., $A_1, B_3, A_{100}, J_{375}$, are sentences of TFL.
These are called atomic sentences.

We only permit use of A – W because X , Y , and Z are reserved for metavariables.

Our second rule says:

2. If X is a sentence of TFL, then so is $\neg X$.

By rule 1, we know that A is a sentence. Rule 2 then allows us to conclude that $\neg A$ is also a sentence. We could then apply it again and conclude that $\neg\neg A$ is also a sentence.

FORMATION TREES help us keep track of this process. For the case of $\neg\neg A$ this would be:



Our third rule says:

3. If X and Y are sentences, then so is $(X \& Y)$.

By rule 1, B_1 and D are both sentences. So rule 3 allows us to conclude that $(B_1 \& D)$ is a sentence. We might then apply rule 2 to conclude that $\neg(B_1 \& D)$ is also a sentence.




We then give similar rules for each of our other connectives: \vee , \rightarrow and \leftrightarrow .


We summarise this in a definition:

1. Every atomic sentence is a sentence.
2. If X is a sentence, then $\neg X$ is a sentence.
3. If X and Y are sentences, then $(X \& Y)$ is a sentence.
4. If X and Y are sentences, then $(X \vee Y)$ is a sentence.
5. If X and Y are sentences, then $(X \rightarrow Y)$ is a sentence.
6. If X and Y are sentences, then $(X \leftrightarrow Y)$ is a sentence.
7. Nothing else is a sentence.

For example, consider $(A \& (B \vee C))$ we can check this is a sentence by drawing the following formation tree:



Each of the steps here tracks one of the rules of what it is to be a sentence. So we can conclude that this is a sentence of TFL. This also helps us see how to read it. It has a different formation tree from $((A \& B) \vee C)$:



$A \& B \vee C$ is not a sentence of TFL. It cannot be broken up into smaller sentences by any of our rules. It needs brackets to be a well-formed sentence. The different formations will be important when we describe truth-tables for these sentences. $((A \& B) \vee C)$ and $(A \& (B \vee C))$ will differ in when they are true.

Definitions like this are called *recursive*. Recursive definitions begin with some specifiable base elements, and then present ways to generate

indefinitely many more elements by compounding together previously established ones. To give you a better idea of what a recursive definition is, we can give a recursive definition of the idea of *an ancestor of mine*. We specify a base clause.

- My parents are ancestors of mine.

and then offer further clauses like:

- If x is an ancestor of mine, then x 's parents are ancestors of mine.
- Nothing else is an ancestor of mine.

Using this definition, we can easily check to see whether someone is my ancestor: just check whether she is the parent of the parent of... one of my parents. And the same is true for our recursive definition of sentences of TFL. Just as the recursive definition allows complex sentences to be built up from simpler parts, the definition allows us to decompose sentences into their simpler parts. Once we get down to atomic sentences, then we know we are ok.

One more example: consider $\neg(P \ \& \ \neg(\neg Q \vee P))$ we can check this is a sentence by drawing the following formation tree:



each of the steps here tracks one of the rules of what it is to be a sentence. So we can conclude that this is a sentence of TFL. This also helps us see how to read it. The will be important when we consider the circumstances under which a particular sentence would be true or false. The sentence $\neg Q$ is true if and only if the sentence Q is false, and so on through the structure of the sentence, until we arrive at the atomic components. We will return to this point in Part III.

This displays clearly the recursive structure of the tree. The sentences further up the tree are formed by one of the formation rules

from the sentences further down the tree. The main connective of each sentence is in red.

When drawing these trees we have highlighted a particular connective on each of our nodes. We call that connective the **MAIN CONNECTIVE** of the sentence.

The **MAIN CONNECTIVE** of sentence is the last connective that was introduced in the construction of the sentence.

In the case of $((\neg E \vee F) \rightarrow \neg \neg G)$, the main connective is \rightarrow . Here we can see that the whole sentence can be described in the form $(X \rightarrow Y)$ with both X and Y being complete sentences (put $X = (\neg E \vee F)$ and $Y = \neg \neg G$). That's enough to see that \rightarrow is the main operator. In the case of $\neg \neg \neg D$, the main logical operator is the very first \neg sign. This is because we can see the sentence as having the form $\neg X$ with X being the complete sentence $\neg \neg D$. In the case of $(P \& \neg(\neg Q \vee R))$, the main logical operator is $\&$: it's an $(X \& Y)$ with X as P and Y as $\neg(\neg Q \vee R)$.

We also say:

The **SCOPE** of an instance of a connective is the subsentence for which it is the main connective.

For example, in:

$$(P \& (\neg(R \& Q) \leftrightarrow P))$$

The scope of the \neg is $\neg(R \& Q)$.

To see this, we can draw the formation tree:



$\neg(R \ \& \ Q)$ is the subsentence in which \neg is the main logical operator. We have worked that out by drawing the formation tree and finding the lowest sentence in which that connective instance appears.

We might informally describe the scope as the part of the whole sentence that that connective cares about. For the purposes of the connective $\&$ in $(A \vee (B \ \& \ \neg C))$, the truth value of A is irrelevant. It is only what is going on with B and $\neg C$ that matters for it. The notion of scope will be very important when we look at First Order Logic.

6.2 Bracketing conventions

Strictly speaking, $A \ \& \ B$ is not a sentence of TFL. When we introduce a connective $\&, \vee, \rightarrow$ or \leftrightarrow , strictly speaking, we must include brackets. It is only $(A \ \& \ B)$. The reason for this rule is that we might use $(A \ \& \ B)$ as a subsentence in a more complicated sentence. For example, we might want to negate $(A \ \& \ B)$, obtaining $\neg(A \ \& \ B)$. If we just had $A \ \& \ B$ without the brackets and put a negation in front of it, we would have $\neg A \ \& \ B$. It is most natural to read this as meaning the same thing as $(\neg A \ \& \ B)$, but this may be very different from $\neg(A \ \& \ B)$.

When working with TFL, however, it will make our lives easier if we are sometimes a little less than strict. So, here are some convenient conventions.

First, we allow ourselves to omit the *outermost* brackets of a sentence. Thus we allow ourselves to write $A \ \& \ B$ instead of the sentence $(A \ \& \ B)$. However, we must remember to put the brackets back in, when we want to embed the sentence into a more complicated sentence!

Second, it can be a bit painful to stare at long sentences with many nested pairs of brackets. To make things a bit easier on the eyes, we will allow ourselves to use square brackets, '[' and ']', instead of rounded ones. So there is no logical difference between $(P \ \vee \ Q)$ and $[P \ \vee \ Q]$, for example.

Combining these two conventions, we can rewrite the unwieldy sentence

$$(((H \rightarrow I) \vee (I \rightarrow H)) \ \& \ (J \vee K))$$

rather more clearly as follows:

$$[(H \rightarrow I) \vee (I \rightarrow H)] \ \& \ (J \vee K)$$

The scope of each connective is now much easier to pick out.

Practice exercises

A. For each of the following: (a) Is it a sentence of TFL, strictly speaking? (b) Is it a sentence of TFL, allowing for our relaxed bracketing conventions?

1. (A)
2. $J_{374} \vee \neg J_{374}$
3. $\neg\neg\neg\neg F$
4. $\neg \& S$
5. $(G \& \neg G)$
6. $(A \rightarrow (A \& \neg F)) \vee (D \leftrightarrow E)$
7. $[(Z \leftrightarrow S) \rightarrow W] \& [J \vee C]$
8. $(F \leftrightarrow \neg D \rightarrow J) \vee (C \& D)$

B. Are there any sentences of TFL that contain no atomic sentences? Explain your answer.

C. What is the scope of each connective in the sentence

$$[(H \rightarrow I) \vee (I \rightarrow H)] \& (J \vee K)$$

CHAPTER 7

Symbolising complex sentences

In §5 we discussed how to symbolise sentences. But we mostly focused on simple sentences. Sentences of English, though, might end up having much more complex structure.

Our general strategy will be:

1. See if the sentence can be reformulated naturally with ‘if’, ‘and’ etc between sentences. If not, use an atomic sentence.
2. Replace the ‘and’ with &, or as appropriate (with brackets).
3. Repeat the procedure with the sentences connected by &.

We can now see some cases where the brackets are very important.

Consider, for example, how negation might interact with conjunction. Consider:

1. It’s not the case that you will get both soup and salad.
2. You will not get soup but you will get salad.

Sentence 1 can be paraphrased as ‘It is not the case that: both you will get soup and you will get salad’. Using this symbolization key:

- S_1 : You will get soup.
 S_2 : You will get salad.

We would symbolize ‘both you will get soup and you will get salad’ as $(S_1 \& S_2)$. To symbolize sentence 1, then, we simply negate the whole sentence, thus: $\neg(S_1 \& S_2)$.

Sentence 2 is a conjunction: you *will not* get soup, and you *will* get salad. ‘You will not get soup’ is symbolized by $\neg S_1$. So to symbolize sentence 2 itself, we offer $(\neg S_1 \& S_2)$.

These English sentences are very different, and their symbolizations differ accordingly. In one of them, the entire conjunction is negated. In the other, just one conjunct is negated. Brackets help us to keep track of things like the *scope* of the negation.

3. You will not have soup, or you will not have salad.
4. You will have neither soup nor salad.
5. You get either soup or salad (but not both).

Using the same symbolization key as before, sentence 3 can be paraphrased in this way: ‘Either *it is not the case that* you get soup, or *it is not the case that* you get salad’. To symbolize this in TFL, we need both disjunction and negation. ‘It is not the case that you get soup’ is symbolized by $\neg S_1$. ‘It is not the case that you get salad’ is symbolized by $\neg S_2$. So sentence 3 itself is symbolized by $(\neg S_1 \vee \neg S_2)$.

Sentence 4 also requires negation. It can be paraphrased as, ‘*It is not the case that* either you get soup or you get salad’. Since this negates the entire disjunction, we symbolize sentence 4 with $\neg(S_1 \vee S_2)$.

Sentence 5 is an *exclusive or*. We can break the sentence into two parts. The first part says that you get one or the other. We symbolize this as $(S_1 \vee S_2)$. The second part says that you do not get both. We can paraphrase this as: ‘It is not the case both that you get soup and that you get salad’. Using both negation and conjunction, we symbolize this with $\neg(S_1 \& S_2)$. Now we just need to put the two parts together. As we saw above, ‘but’ can usually be symbolized with $\&$. Sentence 5 can thus be symbolized as $((S_1 \vee S_2) \& \neg(S_1 \& S_2))$.

Sometimes in English it is ambiguous whether the ‘or’ is to be understood in English inclusively or exclusively. When symbolising into TFL, one needs to decide if you want to add ‘but not both’ or not.

It is also important to note that even though the TFL symbol ‘ \vee ’ always symbolizes *inclusive or*, we can symbolize an *exclusive or* in TFL. We just have to use a few of our other symbols as well.

- ▷ If a sentence can be paraphrased as ‘neither X or Y ’ it can be symbolised as $\neg(X \vee Y)$.
- ▷ If a sentence paraphrased as ‘either X or Y , but not both’ (an *exclusive or*) it can be symbolised as $((X \vee Y) \& \neg(X \& Y))$.

7.1 Unless

An especially difficult case is when we use the English-language connective ‘unless’:

6. Unless you wear a jacket, you will catch a cold.
7. You will catch a cold unless you wear a jacket.

These two sentences are equivalent. They are also equivalent to the following:

8. If you do not wear a jacket, then you will catch a cold.
9. If you do not catch a cold, then you wore a jacket.
10. Either you will wear a jacket or you will catch a cold.

And we know how to symbolise these sentences. We will use the symbolization key:

J : You will wear a jacket.

D : You will catch a cold.

and can then give the symbolizations $\neg J \rightarrow D$, $\neg D \rightarrow J$ and $J \vee D$.

All three are correct symbolizations. Indeed, in chapter ?? we will see that all three symbolizations are equivalent in TFL.

If a sentence can be paraphrased as ‘Unless X , Y ’, then it can be symbolized as $(X \vee Y)$.

Again, though, there is a little complication. ‘Unless’ can be symbolized as a conditional; but as we said above, people often use the conditional (on its own) when they mean to use the biconditional. Equally, ‘unless’ can be symbolized as a disjunction; but there are two kinds of disjunction (exclusive and inclusive). So it will not surprise you to discover that ordinary speakers of English often use ‘unless’ to mean

something more like the biconditional, or like exclusive disjunction. Suppose someone says: 'I will go running unless it rains'. They probably mean something like 'I will go running iff it does not rain' (i.e. the biconditional), or 'either I will go running or it will rain, but not both' (i.e. exclusive disjunction). Again: be aware of this when interpreting what other people have said, but be precise in your writing.

Practice exercises

A. Using the symbolization key given, symbolize each English sentence in TFL.

M: Those creatures are men in suits.

C: Those creatures are chimpanzees.

G: Those creatures are gorillas.

1. Those creatures are not men in suits.
2. Those creatures are men in suits, or they are not.
3. Those creatures are either gorillas or chimpanzees.
4. Those creatures are neither gorillas nor chimpanzees.
5. If those creatures are chimpanzees, then they are neither gorillas nor men in suits.
6. Unless those creatures are men in suits, they are either chimpanzees or they are gorillas.

B. Using the symbolization key given, symbolize each English sentence in TFL.

A: Mister Ace was murdered.

B: The butler did it.

C: The cook did it.

D: The Duchess is lying.

E: Mister Edge was murdered.

F: The murder weapon was a frying pan.

1. Either Mister Ace or Mister Edge was murdered.
2. If Mister Ace was murdered, then the cook did it.
3. If Mister Edge was murdered, then the cook did not do it.
4. Either the butler did it, or the Duchess is lying.
5. The cook did it only if the Duchess is lying.
6. If the murder weapon was a frying pan, then the culprit must have been the cook.

7. If the murder weapon was not a frying pan, then the culprit was either the cook or the butler.
8. Mister Ace was murdered if and only if Mister Edge was not murdered.
9. The Duchess is lying, unless it was Mister Edge who was murdered.
10. If Mister Ace was murdered, he was done in with a frying pan.
11. Of course the Duchess is lying!

C. For each argument, write a symbolization key and symbolize the argument as well as possible in TFL. The part of the passage in italics is there to provide context for the argument, and doesn't need to be symbolized.

1. It is going to rain soon. I know because my leg is hurting, and my leg hurts if it's going to rain.
2. *Spider-man tries to figure out the bad guy's plan.* If Doctor Octopus gets the uranium, he will blackmail the city. I am certain of this because if Doctor Octopus gets the uranium, he can make a dirty bomb, and if he can make a dirty bomb, he will blackmail the city.
3. *A westerner tries to predict the policies of the Chinese government.* If the Chinese government cannot solve the water shortages in Beijing, they will have to move the capital. They don't want to move the capital. Therefore they must solve the water shortage. But the only way to solve the water shortage is to divert almost all the water from the Yangzi river northward. Therefore the Chinese government will go with the project to divert water from the south to the north.

CHAPTER 8

Ambiguity

In English, sentences can be **AMBIGUOUS**, i.e., they can have more than one meaning. There are many sources of ambiguity. One is *lexical ambiguity*: a sentence can contain words which have more than one meaning. For instance, ‘bank’ can mean the bank of a river, or a financial institution. So I might say that ‘I went to the bank’ when I took a stroll along the river, or when I went to deposit a check. Depending on the situation, a different meaning of ‘bank’ is intended, and so the sentence, when uttered in these different contexts, expresses different meanings.

A different kind of ambiguity is *structural ambiguity*. This arises when a sentence can be interpreted in different ways, and depending on the interpretation, a different meaning is selected. A famous example due to Noam Chomsky is the following:

- Flying planes can be dangerous.

There is one reading in which ‘flying’ is used as an adjective which modifies ‘planes’. In this sense, what’s claimed to be dangerous are airplanes which are in the process of flying. In another reading, ‘flying’ is a gerund: what’s claimed to be dangerous is the act of flying a plane. In the first case, you might use the sentence to warn someone who’s about to launch a hot air balloon. In the second case, you might use it to counsel someone against becoming a pilot.

When the sentence is uttered, usually only one meaning is intended. Which of the possible meanings an utterance of a sentence intends is determined by context, or sometimes by how it is uttered (which parts of the sentence are stressed, for instance). Often one interpretation is much more likely to be intended, and in that case it will even be difficult

to “see” the unintended reading. This is often the reason why a joke works, as in this example from Groucho Marx:

- One morning I shot an elephant in my pajamas.
- How he got in my pajamas, I don’t know.

Ambiguity is related to, but not the same as, vagueness. An adjective, for instance ‘rich’ or ‘tall,’ is **VAGUE** when it is not always possible to determine if it applies or not. For instance, a person who’s 6 ft 4 in (1.9 m) tall is pretty clearly tall, but a building that size is tiny. Here, context has a role to play in determining what the clear cases and clear non-cases are (‘tall for a person,’ ‘tall for a basketball player,’ ‘tall for a building’). Even when the context is clear, however, there will still be cases that fall in a middle range.

In TFL, we generally aim to avoid ambiguity. We will try to give our symbolization keys in such a way that they do not use ambiguous words or disambiguate them if a word has different meanings. So, e.g., your symbolization key will need two different sentence letters for ‘Rebecca went to the (money) bank’ and ‘Rebecca went to the (river) bank.’ Vagueness is harder to avoid. Since we have stipulated that every case (and later, every valuation) must make every basic sentence (or sentence letter) either true or false and nothing in between, we cannot accommodate borderline cases in TFL.

It is an important feature of sentences of TFL that they *cannot* be structurally ambiguous. Every sentence of TFL can be read in one, and only one, way. This feature of TFL is also a strength. If an English sentence is ambiguous, TFL can help us make clear what the different meanings are. Although we are pretty good at dealing with ambiguity in everyday conversation, avoiding it can sometimes be terribly important. Logic can then be usefully applied: it helps philosopher express their thoughts clearly, mathematicians to state their theorems rigorously, and software engineers to specify loop conditions, database queries, or verification criteria unambiguously.

Stating things without ambiguity is of crucial importance in the law as well. Here, ambiguity can, without exaggeration, be a matter of life and death. Here is a famous example of where a death sentence hinged on the interpretation of an ambiguity in the law. Roger Casement (1864–1916) was a British diplomat who was famous in his time for publicizing human-rights violations in the Congo and Peru (for which he was knighted in 1911). He was also an Irish nationalist. In 1914–16, Casement secretly travelled to Germany, with which Britain

was at war at the time, and tried to recruit Irish prisoners of war to fight against Britain and for Irish independence. Upon his return to Ireland, he was captured by the British and tried for high treason.

The law under which Casement was tried is the *Treason Act of 1351*. That act specifies what counts as treason, and so the prosecution had to establish at trial that Casement's actions met the criteria set forth in the Treason Act. The relevant passage stipulated that someone is guilty of treason

if a man is adherent to the King's enemies in his realm,
giving to them aid and comfort in the realm, or elsewhere.

Casement's defense hinged on the last comma in this sentence, which is not present in the original French text of the law from 1351. It was not under dispute that Casement had been 'adherent to the King's enemies', but the question was whether being adherent to the King's enemies constituted treason only when it was done in the realm, or also when it was done abroad. The defense argued that the law was ambiguous. The claimed ambiguity hinged on whether 'or elsewhere' attaches only to 'giving aid and comfort to the King's enemies' (the natural reading without the comma), or to both 'being adherent to the King's enemies' and 'giving aid and comfort to the King's enemies' (the natural reading with the comma). Although the former interpretation might seem far fetched, the argument in its favor was actually not unpersuasive. Nevertheless, the court decided that the passage should be read with the comma, so Casement's antics in Germany were treasonous, and he was sentenced to death. Casement himself wrote that he was 'hanged by a comma'.

We can use TFL to symbolize both readings of the passage, and thus to provide a disambiguation. First, we need a symbolization key:

- A*: Casement was adherent to the King's enemies in the realm.
- G*: Casement gave aid and comfort to the King's enemies in the realm.
- B*: Casement was adherent to the King's enemies abroad.
- H*: Casement gave aid and comfort to the King's enemies abroad.

The interpretation according to which Casement's behavior was not treasonous is this:

- $A \vee (G \vee H)$

The interpretation which got him executed, on the other hand, can be symbolized by:

$$\bullet (A \vee B) \vee (G \vee H)$$

Remember that in the case we're dealing with Casement, was adherent to the King's enemies abroad (B is true), but not in the realm, and he did not give the King's enemies aid or comfort in or outside the realm (A , G , and H are false).

One common source of structural ambiguity in English arises from its lack of parentheses. For instance, if I say 'I like movies that are not long and boring', you will most likely think that what I dislike are movies that are long and boring. A less likely, but possible, interpretation is that I like movies that are both (a) not long and (b) boring. The first reading is more likely because who likes boring movies? But what about 'I like dishes that are not sweet and flavorful'? Here, the more likely interpretation is that I like savory, flavorful dishes. (Of course, I could have said that better, e.g., 'I like dishes that are not sweet, yet flavorful'.) Similar ambiguities result from the interaction of 'and' with 'or'. For instance, suppose I ask you to send me a picture of a small and dangerous or stealthy animal. Would a leopard count? It's stealthy, but not small. So it depends whether I'm looking for small animals that are dangerous or stealthy (leopard doesn't count), or whether I'm after either a small, dangerous animal or a stealthy animal (of any size).

These kinds of ambiguities are called *scope ambiguities*, since they depend on whether or not a connective is in the scope of another. For instance, the sentence, '*Avengers: Endgame* is not long and boring' is ambiguous between:

1. *Avengers: Endgame* is not: both long and boring.
2. *Avengers: Endgame* is both: not long and boring.

Sentence 2 is certainly false, since *Avengers: Endgame* is over three hours long. Whether you think 1 is true depends on if you think it is boring or not. We can use the symbolization key:

B : *Avengers: Endgame* is boring.

L : *Avengers: Endgame* is long.

Sentence 1 can now be symbolized as ' $\neg(L \& B)$ ', whereas sentence 2 would be ' $\neg L \& B$ '. In the first case, the '&' is in the scope of ' \neg ', in the second case ' \neg ' is in the scope of '&'.

The sentence ‘Tai Lung is small and dangerous or stealthy’ is ambiguous between:

3. Tai Lung is either both small and dangerous or stealthy.
4. Tai Lung is both small and either dangerous or stealthy.

We can use the following symbolization key:

D: Tai Lung is dangerous.

S: Tai Lung is small.

T: Tai Lung is stealthy.

The symbolization of sentence 3 is ‘ $(S \& D) \vee T$ ’ and that of sentence 4 is ‘ $S \& (D \vee T)$ ’. In the first, & is in the scope of \vee , and in the second \vee is in the scope of &.

Practice exercises

A. The following sentences are ambiguous. Give symbolization keys for each and symbolize the different readings.

1. Haskell is a birder and enjoys watching cranes.
2. The zoo has lions or tigers and bears.
3. The flower is not red or fragrant.

CHAPTER 9

Use and mention

In this Part, we have talked a lot *about* sentences. So we should pause to explain an important, and very general, point.

9.1 Quotation conventions

Consider these two sentences:

- Justin Trudeau is the Prime Minister.
- The expression ‘Justin Trudeau’ is composed of two uppercase letters and eleven lowercase letters

When we want to talk about the Prime Minister, we *use* his name. When we want to talk about the Prime Minister’s name, we *mention* that name, which we do by putting it in quotation marks.

There is a general point here. When we want to talk about things in the world, we just *use* words. When we want to talk about words, we typically have to *mention* those words. We need to indicate that we are mentioning them, rather than using them. To do this, some convention is needed. We can put them in quotation marks, or display them centrally in the page (say). So this sentence:

- ‘Justin Trudeau’ is the Prime Minister.

says that some *expression* is the Prime Minister. That’s false. The *man* is the Prime Minister; his *name* isn’t. Conversely, this sentence:

- Justin Trudeau is composed of two uppercase letters and eleven lowercase letters.

also says something false: Justin Trudeau is a man, made of flesh rather than letters. One final example:

- “‘Justin Trudeau’” is the name of ‘Justin Trudeau’.

On the left-hand-side, here, we have the name of a name. On the right hand side, we have a name. Perhaps this kind of sentence only occurs in logic textbooks, but it is true nonetheless.

Those are just general rules for quotation, and you should observe them carefully in all your work! To be clear, the quotation-marks here do not indicate indirect speech. They indicate that you are moving from talking about an object, to talking about the name of that object.

9.2 Quotation in Logic

Since we are introducing a formal language, we never really *use* the language in the way we would use English or German; we are always mentioning it. And it’s clear from the way it’s written that it’s not part of English, so logicians often drop the quotation marks around logic. Really we should say

- ‘ D ’ is an atomic sentence of TFL.
- ‘ $(D \ \& \ (B \vee C))$ ’ is a sentence of TFL.

But we don’t generally bother, and will typically just write

- D is an atomic sentence of TFL.
- $(D \ \& \ (B \vee C))$ is a sentence of TFL.

This textbook follows this convention.

9.3 Metavariables

Often we do not just want to talk about *specific* expressions of TFL. We also want to be able to talk about *any arbitrary* sentence of TFL. Indeed, we had to do this in §6, when we presented the recursive definition of a sentence of TFL. We used letters from the end of the alphabet to do this, namely:

$$X, Y, Z, X_1, Y_{100}, \dots$$

These symbols do not belong to TFL. Rather, they are part of our (augmented) metalanguage that we use to talk about *any* expression of our formal language. To repeat the second clause of the recursive definition of a sentence of TFL, we said:

2. If X is a sentence, then $\neg X$ is a sentence.

This talks about *arbitrary* sentences. If we had instead offered:

- If A is a sentence, then $\neg A$ is a sentence.

this would not have allowed us to determine whether $\neg B$ is a sentence, it would only tell us about the specific case for A . We don't want to say something about A specifically, but about all sentences. For example this should also give us that

- If $(A \vee (B \& C))$ is a sentence, then $\neg (A \vee (B \& C))$ is a sentence.

We get this by using a metavariable X . It stands in place of an arbitrary expression.

To emphasize, then:

X is a symbol (called a **METAVARIABLE**) in augmented English, which we use to talk about any expression of our formal language. A is a particular atomic sentence of TFL.

PART III

Truth tables

CHAPTER 10

Truth rules for the connectives of TFL

We now move to looking at when sentences of TFL are true or false. We will give precise rules for determining this. The important feature of truth functional logic is that the truth value of a complex sentence, such as $A \vee (B \& C)$ is determined just by the truths of its component parts, that is A , B and C . If we're told whether A , B and C are true or false, then we will be able to say whether $A \vee (B \& C)$ is true or false.

To be able to do this, we need to describe how the truth values are to be combined. We work through each of our connectives describing the rules governing it.

10.1 Negation

Consider:

1. Bristol is not in France.
2. Bristol is not in England.

'Bristol is in France' is false, so 'Bristol is not in France' is true. 'Bristol is in England' is true, so 'Bristol is not in England' is false.

In general, to know whether a sentence of the form $\neg X$ is true. This depends on whether X is true or not in the way:

- If X is true, then $\neg X$ is false.
- If X is false, then $\neg X$ is true.

We record this in shorthand:

If X is:		then $\neg X$ is:
T	\leadsto	F
F	\leadsto	T

We have abbreviated ‘True’ with ‘T’ and ‘False’ with ‘F’. (But just to be clear, the two truth values are True and False; the truth values are not *letters*!)

10.2 Conjunction

Recall that $X \& Y$ was used to symbolise ‘ X and Y ’.

Consider:

3. She can speak German and French.

If she can speak German and she can speak French, then this is true, but otherwise it is false.

More generally, the rule governing $\&$ is:

- If X and Y are both true, then $X \& Y$ is true.
- Otherwise, $X \& Y$ is false.

Which we summarise

If X is:	and Y is:		then $X \& Y$ is:
T	T	\leadsto	T
T	F	\leadsto	F
F	T	\leadsto	F
F	F	\leadsto	F

Note that conjunction is *symmetrical*. The truth value for $X \& Y$ is always the same as the truth value for $Y \& X$.

10.3 Disjunction

Disjunction is a bit more subtle. Consider:

4. She can speak German or French.

If she cannot speak either German or French, then this is false. If she can speak German but not French, then it is true, and if she can speak French but not German it is also true. We have the general rules:

- ▷ If X and Y are both false, then $X \vee Y$ is false.
- ▷ If X is true and Y are false, then $X \vee Y$ is true.
- ▷ If X is false and Y are true, then $X \vee Y$ is true.

But what if she can speak both? Is it true or false? We have already pointed out that in English there are two kinds of disjunctions: an *inclusive* and an *exclusive* one.

For the inclusive or, we might whisper a “or both” after it; whereas for the exclusive or, we’d want to whisper a “but not both”:

5. She speaks German or French (or both).
6. She speaks German or French (but not both).

In logic there can be no ambiguity. We choose that \vee stands for the *inclusive or*. That is, we give the final rule:

- ▷ If X and Y are both true, then $X \vee Y$ is true.

So, when doing symbolisations, one should only symbolise a sentence as $X \vee Y$ if it is to be read as the *inclusive or*. To symbolise the exclusive or, you need to use the more complex sentence: $(X \vee Y) \ \& \ \neg(X \ \& \ Y)$, which essentially makes explicit the whispered “but not both”. Sometimes the English is ambiguous, in which case one should point that out when symbolising and give the two alternative symbolisations.

Consider:

7. She either ate pizza or pasta.

Maybe this is an exclusive or, though an alternative treatment is to take this to be an inclusive or, but note that there’s a implicit, or missing, premise: that she didn’t eat both.

To summarise the rules for \vee :

If X is:	and Y is:		then $X \vee Y$ is:
T	T	\leadsto	T
T	F	\leadsto	T
F	T	\leadsto	T
F	F	\leadsto	F

Like conjunction, disjunction is symmetrical.

10.4 Conditional

Suppose you are a bartender considering the conditional

8. If she is drinking a beer, then she is over eighteen.

Here is what we'll say about this conditional:

- If she's drinking beer and is sixteen, then it is false. (You should kick her out of the bar.)
- If she is drinking beer and is nineteen, then it's true.
- If she is drinking coke, then it's true. (No need to check her age)

We summarise these rules:

If X is:	and Y is:		then $X \rightarrow Y$ is:
T	T	\leadsto	T
T	F	\leadsto	F
F	T	\leadsto	T
F	F	\leadsto	T

In this case, it's very important to remember which way around it goes. The TF-line is different to the FT-line.

This is why the terms 'antecedent' and 'consequent' are so useful.

In $X \rightarrow Y$, X is called the **ANTECEDENT**, and Y the **CONSEQUENT**.

We can redescribe this rule: If the antecedent is true and the consequent false, then the conditional sentence is false, otherwise it is true.

The TFL connective \rightarrow is *stipulated* to be governed by these rules. It is sometimes called the **MATERIAL CONDITIONAL** to highlight that it is governed by these rules.

Subjunctive conditionals

Some English sentences that have the form ‘if...then...’ do not fit with these rules. Consider

9. If the UK had voted to remain in the EU, then the moon would be made of cheese.

This has a false antecedent (we voted to exit), and a false consequent (I certainly can’t fly). But the whole sentence would be judged to be true. So this doesn’t fit with the rule we’ve given for \rightarrow .

There are two kinds of sentences of the form ‘if..., then...’ in English: indicative and subjunctive ones. Consider the two sentences:

10. If Oswald didn’t kill Kennedy, someone else did.
11. If Oswald hadn’t killed Kenney, someone else would have.

The former uses the **INDICATIVE CONDITIONAL** and is plausibly true. After all, we know that Kennedy was assassinated. The latter uses the **SUBJUNCTIVE CONDITIONAL** and is plausibly false. Unless there was a conspiracy plot, if Oswald hadn’t have killed Kenney then Kennedy wouldn’t have been assassinated. The term **COUNTERFACTUAL CONDITIONAL** is also often used.¹

Whilst we use the English ‘if...then...’ for both sentences **10** and **11**, they are very different kinds of conditionals. The conditional of TFL does a terrible job at symbolising the subjunctive conditional which aren’t governed by these rules. In fact TFL just doesn’t have the resources to consider subjunctive conditionals, they are not “truth functional”. We will discuss this in the next chapter, but for now we simply note that if you have to symbolise a subjunctive conditional in TFL it is then a matter of judgement whether to do it as $X \rightarrow Y$, or to simply use an atomic sentence.

The material conditional does a much better job at symbolising the indicative conditionals, though it is a matter of substantial philosophical debate about exactly how well they do. We briefly mention some of the worries in §17.

¹Strictly speaking, we can use subjunctive conditionals with true antecedents, whereas a counterfactual conditional has to have a false antecedent (to be “counter to the fact”).

10.5 Biconditional

Consider

12. Sue is coming to the party if and only if Maya is coming.

If Sue and Maya are both coming, then this is true. If they're both not coming, then it is also true. If only one of them is coming, then it is false.

If X is:	and Y is:		then $X \leftrightarrow Y$ is:
T	T	\sim	T
T	F	\sim	F
F	T	\sim	F
F	F	\sim	T

You can see a summary of the rules for all the connectives, for ease of reference, in the 'Quick Reference Appendix' §3.2

CHAPTER 11

Truth- functional connectives

In this chapter, we reflect on truth-functional logic and the connectives we've used.

11.1 Non truth-functional connectives

Let's introduce an important idea.

A connective is **TRUTH-FUNCTIONAL** iff the truth value of a sentence with that connective as its main connective is uniquely determined by the truth value(s) of the constituent sentence(s).

Every connective in TFL is truth-functional. We were able to give rules to determine what the truth value of a sentence $\neg X$ is depending only on the truth value of X . The truth value of X uniquely determines the truth value of $\neg X$. The same is true for all the other connectives of TFL ($\&$, \vee , \rightarrow , \leftrightarrow). This is what gives TFL its name: it is *truth-functional logic*.

This then means that to determine the truth value of any TFL sen-

tence, we only need to know the truth value of the atomic sentences it includes. We will see exactly how to do this in §12.

In plenty of languages there are connectives that are not truth-functional. We here describe just a few:

Necessarily

In English, for example, we can form a new sentence from any simpler sentence by prefixing it with ‘It is necessarily the case that...’. The truth value of this new sentence is not fixed solely by the truth value of the original sentence. For consider two true sentences:

1. $2 + 2 = 4$
2. Shostakovich wrote fifteen string quartets

Whereas it is necessarily the case that $2 + 2 = 4$, it is not *necessarily* the case that Shostakovich wrote fifteen string quartets. If Shostakovich had died earlier, he would have failed to finish Quartet no. 15; if he had lived longer, he might have written a few more. So ‘It is necessarily the case that...’ is a connective of English, but it is not *truth-functional*.

Subjunctive conditionals

We said that \rightarrow was pretty bad at capturing *subjunctive conditionals* of English. The problem is that a subjunctive conditional is not truth functional. Consider the two sentences:

3. If Mitt Romney had won the 2012 election, then he would have been the 45th President of the USA.
4. If Mitt Romney had won the 2012 election, then he would have turned into a helium-filled balloon and floated away into the night sky.

Sentence 3 is true; sentence 4 is false, but both have false antecedents and false consequents. So the truth value of the whole sentence is not uniquely determined by the truth value of the parts.

\rightarrow is the best that can be done at symbolising subjunctive conditionals of English in TFL. TFL just doesn’t have the required resources as the subjunctive conditional is not truth functional.

11.2 Symbolizing versus translating

All of the connectives of TFL are truth-functional, but more than that: they really do nothing *but* map us between truth values.

When we symbolize a sentence or an argument in TFL, we ignore everything *besides* the contribution that the truth values of a component might make to the truth value of the whole. There are subtleties to our ordinary claims that far outstrip their mere truth values. Sarcasm; poetry; snide implicature; emphasis; these are important parts of everyday discourse, but none of this is retained in TFL. As remarked in §5, TFL cannot capture the subtle differences between the following English sentences:

1. Dana is a logician and Dana is a nice person
2. Although Dana is a logician, Dana is a nice person
3. Dana is a logician despite being a nice person
4. Dana is a nice person, but also a logician
5. Dana's being a logician notwithstanding, he is a nice person

All of the above sentences will be symbolized with the same TFL sentence, perhaps ' $L \ \& \ N$ '.

We keep saying that we use TFL sentences to *symbolize* English sentences. Many other textbooks talk about *translating* English sentences into TFL. However, a good translation should preserve certain facets of meaning, and—as we have just pointed out—TFL just cannot do that. This is why we will speak of *symbolizing* English sentences, rather than of *translating* them.

This affects how we should understand our symbolization keys. Consider a key like:

L: Dana is a logician.

N: Dana is a nice person.

Other textbooks will understand this as a stipulation that the TFL sentence '*L*' should *mean* that Dana is a logician, and that the TFL sentence '*N*' should *mean* that Dana is a nice person, but TFL just is totally unequipped to deal with *meaning*. The preceding symbolization key is doing no more and no less than stipulating that the TFL sentence '*L*' should take the same truth value as the English sentence 'Dana is a logician' (whatever that might be), and that the TFL sentence '*N*' should take the same truth value as the English sentence 'Dana is a nice person' (whatever that might be).

When we treat a TFL sentence as *symbolizing* an English sentence, we are stipulating that the TFL sentence is to take the same truth value as that English sentence.

CHAPTER 12

Complete truth tables

In §10 we described how the truth values of two sentences, X and Y , should combine to determine the truth of a sentence such as $X \vee Y$.

We will now describe how to extend this reasoning to determine the truths of more complex sentences such as $(\neg I \rightarrow H) \& H$.

So far, we have considered assigning truth values to TFL sentences indirectly. We have said, for example, that a TFL sentence such as ' B ' is to take the same truth value as the English sentence 'Ben is happy' (whatever that truth value may be), but we can also assign truth values *directly*. We can simply stipulate that ' B ' is to be true, or stipulate that it is to be false.

A **VALUATION** is any assignment of truth values to particular atomic sentences of TFL.

We describe how to fill out **TRUTH TABLES**. Each row of a truth table is a valuation. The entire truth table represents all possible valuations; thus the truth table provides us with a means to calculate the truth values of complex sentences, on each possible valuation. This is easiest to explain by example.

12.1 A worked example

Consider the sentence $(\neg I \ \& \ H) \rightarrow H$. We will give a *truth table* which lists all the valuations and says whether this sentence is true or false on each of them. The valuations assign either True or False to each atomic sentence. In this case we have two atomic sentences, I and H , so we have four possible valuations, each of which is a line in the truth table:

I	H	$(\neg I \ \& \ H) \rightarrow H$
T	T	
T	F	
F	T	
F	F	

Our job is to fill out the truth values of $(\neg I \ \& \ H) \rightarrow H$.

Here the formation tree will help us know what to do (see §6.1):



The truth rule for \neg tells us how the truth value of $\neg I$ depends on the truth of I . Then the rule for $\&$ tells us how the truth value of $\neg I \ \& \ H$ depends on the truths of $\neg I$ and H ; and finally, the rule for \rightarrow tells us how the truth value of $(\neg I \ \& \ H) \rightarrow H$ depends on those of $\neg I \ \& \ H$ and H .

So to work out the truth values of $(\neg I \ \& \ H) \rightarrow H$ we first need to work out the truth values of $\neg I$ and $\neg I \ \& \ H$.

So, we expand our truth table with columns for each of these.

I	H	$\neg I$	$(\neg I \ \& \ H)$	$(\neg I \ \& \ H) \rightarrow H$
T	T			
T	F			
F	T			
F	F			

The first step is $\neg I$. To be able to do this, we will use the truth rule we specified for \neg :

If X is		then $\neg X$ is
T	\leadsto	F
F	\leadsto	T

Now, we can fill out:

I	H	$\neg I$	$(\neg I \ \& \ H)$	$(\neg I \ \& \ H) \rightarrow H$
T	T	a=F		
T	F	b=F		
F	T	c=T		
F	F	d=T		
★				

We worked these out by:

- ▷ For ‘a’: Look at the column for I , and see it’s T, so by our rule (T-line), we fill out a=F.
- ▷ For ‘b’: Look at the column for I , and see it’s T, so by our rule (T-line), we fill out b=F.
- ▷ For ‘c’: Look at the column for I , and see it’s F, so by our rule (F-line), we fill out c=T.
- ▷ For ‘d’: Look at the column for I , and see it’s F, so by our rule (F-line), we fill out d=T.

The next step is to consider $\neg I \ \& \ H$. For this we will use the truth rule for $\&$:

If X is	and Y is		then $X \ \& \ Y$ is
T	T	\leadsto	T
T	F	\leadsto	F
F	T	\leadsto	F
F	F	\leadsto	F

Now, we can fill out:

I	H	$\neg I$	$(\neg I \ \& \ H)$	$(\neg I \ \& \ H) \rightarrow H$
T	T	F	a=F	
T	F	F	b=F	
F	T	T	c=T	
F	F	T	d=F	
★	★			

We worked these out by:

- For 'a': Look at the column for $\neg I$ and the column for H , we have F and T; so by our rule (FT-line), we fill out $a=F$.
- For 'b': Look at the column for $\neg I$ and the column for H , we have F and F; so by our rule (FF-line), we fill out $a=F$.
- For 'c': Look at the column for $\neg I$ and the column for H , we have T and T; so by our rule (TT-line), we fill out $a=T$.
- For 'd': Look at the column for $\neg I$ and the column for H , we have T and F; so by our rule (TF-line), we fill out $a=F$.

Now, finally, we need to look at $(\neg I \ \& \ H) \rightarrow H$, and will use the truth rule for \rightarrow :

If X is	and Y is		then $X \rightarrow Y$ is
T	T	\leadsto	T
T	F	\leadsto	F
F	T	\leadsto	T
F	F	\leadsto	T

Now, we can fill out:

I	H	$\neg I$	$(\neg I \ \& \ H)$	$(\neg I \ \& \ H) \rightarrow H$
T	T	F	F	a=T
T	F	F	F	b=T
F	T	T	T	c=T
F	F	T	F	d=T
★consequent			★antecedent	

We worked these out by:

- For 'a': In the column for $(\neg I \ \& \ H)$ (which is our antecedent) we have F; and in the column for H (our consequent) we have T. For the conditional, it's very important to bear in mind which order they come in. We are looking at antecedent then consequent, so the rule line we are looking at is FT (the antecedent is False and the consequent is True rather than visa versa). And we get $a=T$.
- For 'b': In the column for $(\neg I \ \& \ H)$ we have F; and in the column for H we have F. So by our rule for \rightarrow (FF-line) we have $b=T$.
- For 'c': In the column for $(\neg I \ \& \ H)$ we have T; and in the column for H we have T. So by our rule for \rightarrow (TT-line) we have $b=T$.

- For ‘d’: In the column for $(\neg I \ \& \ H)$ we have F; and in the column for H we have F. So by our rule for \rightarrow (FF-line) we have $d=T$.

When we do these in practice, our sentences can have many sub-sentences, consider, e.g.

$$\neg(B \ \& \ (\neg B \leftrightarrow \neg A)))$$

A tool for fitting our truth tables on a page is not to put all the sub-sentences out as separate entire columns to calculate, but to simply list the values underneath the main connective of the sentence. This is simply to keep the truth table more concise.

So instead of writing

A	B	$\neg B$	$\neg A$	$\neg B \leftrightarrow \neg A$	$(B \ \& \ (\neg B \leftrightarrow \neg A)))$	$\neg(B \ \& \ (\neg B \leftrightarrow \neg A)))$
T	T	F	F	T	T	F
T	F	T	F	F	F	T
F	T	F	T	F	F	T
F	F	T	T	T	F	T

I can write:

A	B	$\neg(B \ \& \ (\neg B \leftrightarrow \neg A)))$			
T	T	F	T	F	T
T	F	T	F	T	F
F	T	T	F	F	T
F	F	T	F	T	T

and highlight the column under the main connective, which provides our final answer. You should feel free to use whichever method you find easier.

There are tricks that mean you can miss out some of the gaps by using strategies such as: knowing that X is false is already enough to see that $X \ \& \ Y$ is false, so we don’t need to continue to work out the truth value of Y . These “shortcuts” are discussed in §16, but we won’t read it in this course.

12.2 The possible valuations

The size of the complete truth table depends on the number of different atomic sentences in the table. A sentence that contains only one atomic sentence requires only two rows, as in the characteristic truth table for

negation. This is true even if the same letter is repeated many times, as in the sentence ‘ $[(C \leftrightarrow C) \rightarrow C] \& \neg(C \rightarrow C)$ ’. The complete truth table requires only two lines because there are only two possibilities: ‘ C ’ can be true or it can be false. The truth table for this sentence looks like this:

C	$[(C \leftrightarrow C) \rightarrow C] \& \neg (C \rightarrow C)$								
T	T	T	T	T	T	FF	T	T	T
F	F	T	F	F	F	FF	F	T	F

Looking at the column underneath the main logical operator, we see that the sentence is false on both rows of the table; i.e., the sentence is false regardless of whether ‘ C ’ is true or false. It is false on every valuation.

A sentence that contains two atomic sentences requires four lines for a complete truth table, as in the characteristic truth tables for our binary connectives, and as in the complete truth table for ‘ $(H \& I) \rightarrow H$ ’.

A sentence that contains three atomic sentences requires eight lines:

M	N	P	$M \& (N \vee P)$							
T	T	T	T	T	T	T	T	T	T	T
T	T	F	T	T	T	T	F	T	T	F
T	F	T	T	T	F	T	T	T	T	T
T	F	F	T	F	F	F	F	F	F	F
F	T	T	F	F	T	T	T	T	T	T
F	T	F	F	F	T	T	F	T	T	F
F	F	T	F	F	F	T	T	T	T	T
F	F	F	F	F	F	F	F	F	F	F

From this table, we know that the sentence ‘ $M \& (N \vee P)$ ’ can be true or false, depending on the truth values of ‘ M ’, ‘ N ’, and ‘ P ’.

A complete truth table for a sentence that contains four different atomic sentences requires 16 lines. Five letters, 32 lines. Six letters, 64 lines. And so on. To be perfectly general: If a complete truth table has n different atomic sentences, then it must have 2^n lines.

In order to fill in the columns of a complete truth table, begin with the right-most atomic sentence and alternate between ‘T’ and ‘F’. In the next column to the left, write two ‘T’s, write two ‘F’s, and repeat. For the third atomic sentence, write four ‘T’s followed by four ‘F’s. This yields an eight line truth table like the one above. For a 16 line truth table, the next column of atomic sentences should have eight ‘T’s followed

by eight ‘F’s. For a 32 line table, the next column would have 16 ‘T’s followed by 16 ‘F’s, and so on.

12.3 More about brackets

Consider these two sentences:

$$((A \& B) \& C)$$

$$(A \& (B \& C))$$

These are truth functionally equivalent. Consequently, it will never make any difference from the perspective of truth value – which is all that TFL cares about (see §11) – which of the two sentences we assert (or deny). Even though the order of the brackets does not matter as to their truth, we should not just drop them. The expression

$$A \& B \& C$$

is ambiguous between the two sentences above. The same observation holds for disjunctions. The following sentences are logically equivalent:

$$((A \vee B) \vee C)$$

$$(A \vee (B \vee C))$$

But we should not simply write:

$$A \vee B \vee C$$

In fact, it is a specific fact about the characteristic truth table of \vee and $\&$ that guarantees that any two conjunctions (or disjunctions) of the same sentences are truth functionally equivalent, however you place the brackets. *But be careful.* These two sentences have *different* truth tables:

$$((A \rightarrow B) \rightarrow C)$$

$$(A \rightarrow (B \rightarrow C))$$

So if we were to write:

$$A \rightarrow B \rightarrow C$$

it would be dangerously ambiguous. So we must not do the same with conditionals. Equally, these sentences have different truth tables:

$$((A \vee B) \& C)$$

$$(A \vee (B \& C))$$

So if we were to write:

$$A \vee B \& C$$

it would be dangerously ambiguous. *Never write this.* The moral is: never drop brackets.

Practice exercises

A. Draw formation trees and offer complete truth tables for each of the following:

1. $A \rightarrow A$
2. $C \rightarrow \neg C$
3. $(A \leftrightarrow B) \leftrightarrow \neg(A \leftrightarrow \neg B)$
4. $(A \rightarrow B) \vee (B \rightarrow A)$
5. $(A \& B) \rightarrow (B \vee A)$
6. $\neg(A \vee B) \leftrightarrow (\neg A \& \neg B)$
7. $[(A \& B) \& \neg(A \& B)] \& C$
8. $[(A \& B) \& C] \rightarrow B$
9. $\neg[(C \vee A) \vee B]$

B. Check all the claims made in introducing the new notational conventions in §12.3, i.e. show that:

1. ‘ $((A \& B) \& C)$ ’ and ‘ $(A \& (B \& C))$ ’ have the same truth table
2. ‘ $((A \vee B) \vee C)$ ’ and ‘ $(A \vee (B \vee C))$ ’ have the same truth table
3. ‘ $((A \vee B) \& C)$ ’ and ‘ $(A \vee (B \& C))$ ’ do not have the same truth table
4. ‘ $((A \rightarrow B) \rightarrow C)$ ’ and ‘ $(A \rightarrow (B \rightarrow C))$ ’ do not have the same truth table

Also, check whether:

5. ‘ $((A \leftrightarrow B) \leftrightarrow C)$ ’ and ‘ $(A \leftrightarrow (B \leftrightarrow C))$ ’ have the same truth table

C. Write complete truth tables for the following sentences and mark the column that represents the possible truth values for the whole sentence.

1. $\neg(S \leftrightarrow (P \rightarrow S))$
2. $\neg[(X \& Y) \vee (X \vee Y)]$
3. $(A \rightarrow B) \leftrightarrow (\neg B \leftrightarrow \neg A)$
4. $[C \leftrightarrow (D \vee E)] \& \neg C$
5. $\neg(G \& (B \& H)) \leftrightarrow (G \vee (B \vee H))$

D. Write complete truth tables for the following sentences and mark the column that represents the possible truth values for the whole sentence.

1. $(D \& \neg D) \rightarrow G$
2. $(\neg P \vee \neg M) \leftrightarrow M$
3. $\neg\neg(\neg A \& \neg B)$
4. $[(D \& R) \rightarrow I] \rightarrow \neg(D \vee R)$
5. $\neg[(D \leftrightarrow O) \leftrightarrow A] \rightarrow (\neg D \& O)$

E. Can you think of sentences with the following truth table:

	<i>A</i>	<i>B</i>		?
	T	T		T
1.	T	F		T
	F	T		T
	F	F		T

	<i>A</i>	<i>B</i>		?
	T	T		F
2.	T	F		F
	F	T		T
	F	F		F

	<i>A</i>	<i>B</i>		?
	T	T		T
3.	T	F		F
	F	T		T
	F	F		T

F. There are in fact sixteen different possible columns if there are two atomic sentences.

1. Can you explain why?

2. Can you show for each of these combinations that there is a sentence of TFL with that column describing its truth.
3. Can you show there's always a formula just using \neg and $\&$ with that column describing its truth.

If you want additional practice, you can construct truth tables for any of the sentences and arguments in the exercises for the previous chapter.

CHAPTER 13

Validity in TFL

In the previous section, we introduced the idea of a valuation and showed how to determine the truth value of any TFL sentence, on any valuation, using a truth table. In this section, we will introduce some related ideas, and show how to use truth tables to test whether or not they apply.

13.1 Validity

Logic is particularly useful for evaluating arguments, in particular to help us see if an argument is *valid*: logic can help us see if an argument is valid in virtue of its form.

Now, since we are going to be talking about arguments a lot, we will introduce some abbreviation. An argument had the form:

- X_1
- X_2
- \dots
- X_n
- Therefore: Y

We can write this more concisely as:

$$X_1, X_2, \dots, X_n \therefore Y$$

The \therefore symbol can be read out loud as ‘therefore’. (When typing any answers out in word, you can replace it with ‘Therefore’.)

This is an argument, which may be valid or not.

In §2 we introduce the notion of validity: an argument is valid if it’s impossible for all the premises to be true and the conclusion false. We investigated validity as it applies to English language arguments, but we also have TFL arguments. The logical substitute notion of validity replaces ‘impossible’ by ‘*logically* impossible’, where logical impossibility is characterised by there being no valuations with the property of interest.

The TFL argument ‘ $X_1, X_2, \dots, X_n \therefore Y$ ’ is **VALID** iff there are no valuations where the premises are true and the conclusion false.

Validity is a property of an *argument*. If we want to talk about the conclusion being a consequence of the premises, or *following from*, we use the term ‘*entails*’.

The TFL sentences X_1, X_2, \dots, X_n **ENTAIL** the sentence Y iff there are no valuations where all of X_1, X_2, \dots, X_n true and Y false.

An argument is valid if and only if its premises entail its conclusion. It is easy to check for this with a truth table. Consider the argument:

$$\neg L \rightarrow (J \vee L), \neg L \therefore J$$

We need to check whether there is any valuation which makes both $\neg L \rightarrow (J \vee L)$ and $\neg L$ true whilst making J false. So we use a truth table:

J	L	$\neg L \rightarrow (J \vee L)$			$\neg L$	J
T	T	F	T	T	F	T
T	F	T	T	T	T	T
F	T	F	T	T	F	F
F	F	T	F	F	T	F

The only row on which both ‘ $\neg L \rightarrow (J \vee L)$ ’ and ‘ $\neg L$ ’ are true is the second row, and that is a row on which ‘ J ’ is also true. So ‘ $\neg L \rightarrow (J \vee L)$ ’ and ‘ $\neg L$ ’ entail ‘ J ’.

In short, we have a way to test for the validity of English arguments. First, we symbolize them in TFL, as having premises X_1, X_2, \dots, X_n , and conclusion Y . Then we check whether they are valid using truth tables.

‘Entails’ versus ‘ \rightarrow ’

We now want to compare and contrast ‘entails’ and ‘ \rightarrow ’.

Observe: X entails Y iff there is no valuation of the atomic sentences that makes X true and Y false.

Observe: $X \rightarrow Y$ is a tautology iff there is no valuation of the atomic sentences that makes $X \rightarrow Y$ false. Since a conditional is true except when its antecedent is true and its consequent false, $X \rightarrow Y$ is a tautology iff there is no valuation that makes X true and Y false.

Combining these two observations, we see that $X \rightarrow Y$ is a tautology iff X entails Y . But there is a really, really important difference between entailment and ‘ \rightarrow ’: ‘ \rightarrow ’ is a sentential connective of TFL. ‘Entails’ is a word in English. When ‘ \rightarrow ’ is flanked with two TFL sentences, the result is a longer TFL sentence. By contrast, when we use ‘entails’, we are expressing that there is a relationship between the surrounding TFL sentences.

13.2 Formal validity and English arguments

We can now use the tool of TFL to investigate whether an argument of English is formally valid.

Consider an argument like:

If Jones signed the contract under duress, then the contract is void. But since it was not signed under duress, the contract is not void.¹

We are interested in determining if this is valid or not. Here’s a general strategy to help us:

¹This sort of example is discussed in <https://lawpublications.barry.edu/cgi/viewcontent.cgi?article=1026&context=barrylrev>.

1. Find the structure of the argument.
Identify the premises and conclusion.
2. Symbolise the argument in TFL.
3. Check if the TFL argument is valid.
 - Using truth tables to look for a valuation providing a counter example. If there is no such valuation, then it is valid.
 - Or, use natural deduction to show that it is valid.

Following this procedure, then, we need to first find the structure of the argument. It has two premises and a conclusion:

- If Jones signed the contract under duress, then the contract is void.
- Jones did not sign the contract under duress.
- Therefore: The contract is not void.

We are interested in whether it is valid. I.e. is there a situation where the premises are true and the conclusion is false. Perhaps in this case you can work it out, but if not TFL provides us a tool for thinking through all the possibilities. We therefore move to stage 2 and symbolise the sentences of the argument in TFL as well as we can.

- $D \rightarrow V$
- $\neg D$
- Therefore: $\neg V$

And moving to stage 3, we check whether the TFL argument is valid. We do this with truth tables. Later in the course we'll see Natural Deduction which will provide an alternative method for checking validity.

So we need to fill in truth table with all the sentences involved in the argument. In this case each sentence has a simple structure, if we draw out the formation trees they just have one step, e.g.

$$\begin{array}{c}
 D \rightarrow V \\
 \swarrow \quad \searrow \\
 D \quad V
 \end{array}$$

So our truth table doesn't need any additional "calculation" lines. We can directly fill it in, following our truth rules

D	V	Premise	Premise	Concln	
		$D \rightarrow V$	$\neg D$	$\neg V$	
T	T	T	F	F	
T	F	F	F	T	↯ counterexample
F	T	T	T	F	
F	F	T	T	T	

And we can see that this argument is invalid. The valuation providing the counter example is line 3: $\frac{D \quad V}{F \quad T}$

This allows us to read of a possibility where the premises are true but the conclusion false: It might be that the contract is void without having been made under duress. E.g. mental illness, misrepresentation,...

This has then allowed us to see that the original English argument is invalid. In fact, the argument followed an invalid pattern that is commonly found. It is a common fallacy called "denying the consequent".

This argument is invalid. So we can accept the premises and still reject the conclusion. If we have a valid argument, then the conclusion can only be rejected if the premises are rejected. In a valid argument our discussion needs to be focused on the premises.

In fact in this case we can give a fixed-up, related, valid argument

- $V \rightarrow D$
- $\neg D$
- Therefore: $\neg V$

This uses the \rightarrow the other way around and is valid. It is an argument form called Modus Tollens. However, presenting this argument and showing it's valid isn't enough to convince me of the conclusion that the contract is void: the argument uses a false premise: $V \rightarrow D$. Or at least, this is a premise that I won't accept without further justification: look, if there was misrepresentation in the trial then $V \rightarrow D$ is false. Give me a way of ruling out misrepresentation if you want to get to the conclusion that the contract is not void.

Another example: disjunctive reasoning

The murder weapon was the dagger or the rope. So Mrs. Peacock is guilty. After all, if the murder weapon was

the dagger then she's guilty. And if it was the rope, she's guilty.

First we find the argument structure:

- The murder weapon was the dagger or the rope.
- If the murder weapon was the dagger then Mrs. Peacock is guilty.
- If the murder weapon was the rope then Mrs. Peacock is guilty.
- Therefore: Mrs. Peacock is guilty.

We then symbolise each constituent sentence into TFL:

- $D \vee R$
- $D \rightarrow G$
- $R \rightarrow G$
- Therefore: G

Now, we check if the TFL argument is valid using truth tables:

			Premise	Premise	Premise	Concln
D	G	R	$D \vee G$	$D \rightarrow G$	$R \rightarrow G$	G
T	T	T	T	T	T	T
T	T	F	T	T	T	T
T	F	T	T	F	F	F
T	F	F	T	F	T	F
F	T	T	T	T	T	T
F	T	F	T	T	T	T
F	F	T	F	T	F	F
F	F	F	F	T	T	F

And we see that this argument is valid: there is no valuation (=line) where the premises are all true and the conclusion false.

13.3 Other kinds of validity

We have reached an important milestone: a test for the validity of arguments! However, we should not get carried away just yet. It is important to understand the limits of our achievement.

This allows us to investigate the formal validity of an argument. More precisely, whether its TFL-form is a valid form.

If the TFL-form of an argument is valid, then the argument is also conceptually valid, nomologically valid, and valid in any other (plausible) sense. We cannot conceive of possibilities that do not follow the laws of logic.

If the TFL symbolisation of an argument is valid (in TFL), then the original argument is formally valid, and valid in all the other senses.

However, some arguments may have their TFL forms being invalid but are nonetheless conceptually valid. For example:

- That is a triangle.
- Therefore: That has three sides.

This argument is conceptually valid. But it would be symbolised in TFL simply as:

- A
- Therefore: B

and looking at the truth tables (which are very easy to do!) we see:

		Premise	Concln	
A	B	A	B	
T	T	T	T	↔ counterexample valuation, not conceptually possible
T	F	T	F	
F	T	F	T	
F	F	F	F	

The valuation in line 2 gives us a counterexample: it makes the premise, A , true, and the conclusion, B , false. This valuation, however, violates the conceptual connections between words in English. Triangles simply have to have three sides. The “case” being described by this valuation makes it true that it is a triangle but false that it has three sides. That is not a conceptual possibility. The argument is conceptually valid, but its TFL symbolisation is invalid.

If the TFL symbolisation is valid, the argument must be conceptually valid too. But some conceptually valid arguments have invalid TFL symbolisations.

Consider:

- If Jones signed the contract under duress, then the contract is void.
- Jones did not sign the contract under duress.
- Therefore: The contract is not void.

which we symbolised in TFL as:

$$D \rightarrow V, \neg D \therefore \neg V$$

which is invalid. The valuation providing the counterexample is:

$\frac{D \quad V}{F \quad T}$ To see if the argument is *conceptually* valid, one should then investigate whether the valuations providing counterexamples to the formal validity are conceivable possibilities. Here, we're probably interested in something like legal validity, which governs conditions under which a contract is void. We now know we should investigate whether the possibility of it being void but not signed under duress is compatible with the legal laws. This will depend on what the legal laws are. Actually, a contract can be void without being signed under duress, for example due to misrepresentation or mental incapacity. But were the laws different, and the only way to be void was to be signed under duress, then the argument would have been legally valid despite being formally invalid.

There is one loose end to tie up. In fact, TFL is in't all there is to formal validity. If the TFL symbolisation of an argument is valid, then the original argument is formally valid. But consider the following argument: Consider

- Socrates is a man.
- All men are mortal.
- Therefore: Socrates is mortal.

When we symbolise this in TFL, each of the sentences is simply an atomic sentence, so we get:

$$A, B \therefore C$$

which is invalid. The problem isn't the symbolisation we gave. It is the best we could do *in TFL*. The problem is that TFL is limited in the "form" it can see, after all. We will improve on this when we move to First Order Logic in §28. This argument is formally valid in virtue of its FOL form rather than TFL form.

Practice exercises

A. Use truth tables to determine whether each argument is valid or invalid.

1. $A \rightarrow A \therefore A$
2. $A \rightarrow (A \& \neg A) \therefore \neg A$
3. $A \vee (B \rightarrow A) \therefore \neg A \rightarrow \neg B$
4. $A \vee B, B \vee C, \neg A \therefore B \& C$
5. $(B \& A) \rightarrow C, (C \& A) \rightarrow B \therefore (C \& B) \rightarrow A$

B. Determine whether each argument is valid or invalid, using a complete truth table.

1. $A \rightarrow B, B \therefore A$
2. $A \leftrightarrow B, B \leftrightarrow C \therefore A \leftrightarrow C$
3. $A \rightarrow B, A \rightarrow C \therefore B \rightarrow C$
4. $A \rightarrow B, B \rightarrow A \therefore A \leftrightarrow B$

C. Determine whether each argument is valid or invalid, using a complete truth table.

1. $A \vee [A \rightarrow (A \leftrightarrow A)] \therefore A$
2. $A \vee B, B \vee C, \neg B \therefore A \& C$
3. $A \rightarrow B, \neg A \therefore \neg B$
4. $A, B \therefore \neg(A \rightarrow \neg B)$
5. $\neg(A \& B), A \vee B, A \leftrightarrow B \therefore C$

CHAPTER 14

Other logical notions

14.1 Tautologies and contradictions

We have similar substitutes for the notions we introduced in §3. There we explained *necessary truth* and *necessary falsity*. Both notions have surrogates in TFL. We will start with a surrogate for necessary truth.

X is a **TAUTOLOGY** iff it is true on every valuation.

We can determine whether a sentence is a tautology just by using truth tables. If the sentence is true on every line of a complete truth table, then it is true on every valuation, so it is a tautology. In the example of §12, ‘ $(H \ \& \ I) \rightarrow H$ ’ is a tautology.

Our main example of a tautology is $A \vee \neg A$. Whatever the weather is like you know it’s either raining or not. Similarly here, whether or not A is true, we already know $A \vee \neg A$. We can check this with a truth table.

A	$A \vee \neg A$
T	T T F T
F	F T T F

Since there is a T under the main connective on every line of the truth table, $A \vee \neg A$ is true, whatever the valuation is, i.e. whether A is true or false.

This is only, though, a *surrogate* for necessary truth. There are some necessary truths that we cannot adequately symbolize in TFL. An example is ‘ $2 + 2 = 4$ ’. This *must* be true, but if we try to symbolize it in TFL, the best we can offer is an atomic sentence, and no atomic sentence is a tautology. Still, if we can adequately symbolize some English sentence using a TFL sentence which is a tautology, then that English sentence expresses a necessary truth.

We have a similar surrogate for necessary falsity:

X is a **CONTRADICTION** iff it is false on every valuation.

We can determine whether a sentence is a contradiction just by using truth tables. If the sentence is false on every line of a complete truth table, then it is false on every valuation, so it is a contradiction.

Our core example of a contradiction is $A \ \& \ \neg A$. Whether A is true or false, $A \ \& \ \neg A$ is false. This can again be checked using truth tables, observing that there is an F under the main connective in each line of the truth table.

14.2 Logical equivalence

Here is a similar useful notion:

X and Y are **LOGICALLY EQUIVALENT** iff, for every valuation, their truth values agree, i.e. if there is no valuation in which they have opposite truth values.

We have already made use of this notion, in effect, in §12.3; the point was that ‘ $(A \ \& \ B) \ \& \ C$ ’ and ‘ $A \ \& \ (B \ \& \ C)$ ’ are logically equivalent. Again, it is easy to test for logical equivalence using truth tables. Consider the sentences ‘ $\neg(P \vee Q)$ ’ and ‘ $\neg P \ \& \ \neg Q$ ’. Are they logically equivalent? To find out, we construct a truth table.

P	Q	$\neg(P \vee Q)$	$\neg P \ \& \ \neg Q$
T	T	F	T T T
T	F	F	T T F
F	T	F	F F T
F	F	T	F F F

Look at the columns for the main logical operators; negation for the first sentence, conjunction for the second. On the first three rows, both

are false. On the final row, both are true. Since they match on every row, the two sentences are logically equivalent.

14.3 Consistency

In §3, we said that sentences are jointly possible iff it is possible for all of them to be true at once. We can offer a surrogate for this notion too:

X_1, X_2, \dots, X_n are **CONSISTENT** iff there is some valuation which makes them all true.

Derivatively, sentences are jointly logically inconsistent if there is no valuation that makes them all true. Again, it is easy to test for joint logical consistency using truth tables.

14.4 These notions and the English variants

Just as in §13.2 we said we can use the logical notion of validity in TFL to help with various notions of validity of English, the same can be said for these other logical notions.

For example, if you show that the TFL symbolisation of an English sentence is a tautology, you can conclude that it is a necessary truth. However, there are different kinds of ‘necessary truths’. Some sentences like ‘A triangle has three sides’ are necessary truths, but are not tautologies. But the tool of TFL can help.

Practice exercises

A. Revisit your answers to §12A. Determine which sentences were tautologies, which were contradictions, and which were neither tautologies nor contradictions.

B. Use truth tables to determine whether these sentences are jointly consistent, or jointly inconsistent:

1. $A \rightarrow A, \neg A \rightarrow \neg A, A \& A, A \vee A$
2. $A \vee B, A \rightarrow C, B \rightarrow C$
3. $B \& (C \vee A), A \rightarrow B, \neg(B \vee C)$
4. $A \leftrightarrow (B \vee C), C \rightarrow \neg A, A \rightarrow \neg B$

C. Determine whether each sentence is a tautology, a contradiction, or a contingent sentence, using a complete truth table.

1. $\neg B \& B$
2. $\neg D \vee D$
3. $(A \& B) \vee (B \& A)$
4. $\neg[A \rightarrow (B \rightarrow A)]$
5. $A \leftrightarrow [A \rightarrow (B \& \neg B)]$
6. $[(A \& B) \leftrightarrow B] \rightarrow (A \rightarrow B)$

D. Determine whether each the following sentences are logically equivalent using complete truth tables. If the two sentences really are logically equivalent, write “equivalent.” Otherwise write, “Not equivalent.”

1. A and $\neg A$
2. $A \& \neg A$ and $\neg B \leftrightarrow B$
3. $[(A \vee B) \vee C]$ and $[A \vee (B \vee C)]$
4. $A \vee (B \& C)$ and $(A \vee B) \& (A \vee C)$
5. $[A \& (A \vee B)] \rightarrow B$ and $A \rightarrow B$

E. Determine whether each the following sentences are logically equivalent using complete truth tables. If the two sentences really are equivalent, write “equivalent.” Otherwise write, “not equivalent.”

1. $A \rightarrow A$ and $A \leftrightarrow A$
2. $\neg(A \rightarrow B)$ and $\neg A \rightarrow \neg B$
3. $A \vee B$ and $\neg A \rightarrow B$
4. $(A \rightarrow B) \rightarrow C$ and $A \rightarrow (B \rightarrow C)$
5. $A \leftrightarrow (B \leftrightarrow C)$ and $A \& (B \& C)$

F. Determine whether each collection of sentences is jointly consistent or jointly inconsistent using a complete truth table.

1. $A \& \neg B, \neg(A \rightarrow B), B \rightarrow A$
2. $A \vee B, A \rightarrow \neg A, B \rightarrow \neg B$
3. $\neg(\neg A \vee B), A \rightarrow \neg C, A \rightarrow (B \rightarrow C)$
4. $A \rightarrow B, A \& \neg B$
5. $A \rightarrow (B \rightarrow C), (A \rightarrow B) \rightarrow C, A \rightarrow C$

G. Determine whether each collection of sentences is jointly consistent or jointly inconsistent, using a complete truth table.

1. $\neg B, A \rightarrow B, A$
2. $\neg(A \vee B), A \leftrightarrow B, B \rightarrow A$
3. $A \vee B, \neg B, \neg B \rightarrow \neg A$
4. $A \leftrightarrow B, \neg B \vee \neg A, A \rightarrow B$
5. $(A \vee B) \vee C, \neg A \vee \neg B, \neg C \vee \neg B$

H. Answer each of the questions below and justify your answer.

1. Suppose X is true and Y is true
Is $X \vee Y$ true.
2. Suppose X is false
Is $X \vee Y$ true?
3. Suppose X is false
Is $X \rightarrow Y$ true.
4. Suppose that X, Y and Z are jointly inconsistent.
What can you say about $X \& Y \& Y$?
5. Suppose X is a tautology and $X \therefore Y$ is a valid argument.
What can you say about Y ?
6. Suppose that $(X \& Y) \rightarrow Z$ is not a tautology.
What can you say about whether $X, Y \therefore Z$ is valid?
7. Suppose that X is a contradiction.
What can you say about whether $X, Y \models Z$?
8. Suppose that X and Y are logically equivalent.
What can you say about $(X \vee Y)$?
9. Suppose that X and Y are *not* logically equivalent.
What can you say about $X \vee Y$?

I. Consider the following principle:

- Suppose X and Y are logically equivalent. Suppose an argument contains X (either as a premise, or as the conclusion). The validity of the argument would be unaffected, if we replaced X with Y .

Is this principle correct? Explain your answer.

CHAPTER 15

Truth table shortcuts

Not taught in course. This knowledge will not be needed for the exam. (Although you can use these techniques in the exam if you wish.) With practice, you will quickly become adept at filling out truth tables. In this section, we want to give you some permissible shortcuts to help you along the way.

15.1 Working through truth tables

You will quickly find that you do not need to copy the truth value of each atomic sentence, but can simply refer back to them. So you can speed things up by writing:

P	Q	$(P \vee Q) \leftrightarrow \neg P$	
T	T	T	F F
T	F	T	F F
F	T	T	T T
F	F	F	F T

You also know for sure that a disjunction is true whenever one of the disjuncts is true. So if you find a true disjunct, there is no need to work out the truth values of the other disjuncts. Thus you might offer:

P	Q	$(\neg P \vee \neg Q) \vee \neg P$		
T	T	F	F	F
T	F	F	T	T
F	T			T
F	F			T

Equally, you know for sure that a conjunction is false whenever one of the conjuncts is false. So if you find a false conjunct, there is no need to work out the truth value of the other conjunct. Thus you might offer:

P	Q	$\neg(P \& \neg Q) \& \neg P$		
T	T			F
T	F			F
F	T	T	F	T
F	F	T	F	T

A similar short cut is available for conditionals. You immediately know that a conditional is true if either its consequent is true, or its antecedent is false. Thus you might present:

P	Q	$((P \rightarrow Q) \rightarrow P) \rightarrow P$		
T	T			T
T	F			T
F	T	T	F	T
F	F	T	F	T

So ‘ $((P \rightarrow Q) \rightarrow P) \rightarrow P$ ’ is a tautology. In fact, it is an instance of *Peirce’s Law*, named after Charles Sanders Peirce.

15.2 Testing for validity and entailment

When we use truth tables to test for validity or entailment, we are checking for *bad* lines: lines where the premises are all true and the conclusion is false. Note:

- Any line where the conclusion is true is not a bad line.
- Any line where some premise is false is not a bad line.

Since *all* we are doing is looking for bad lines, we should bear this in mind. So: if we find a line where the conclusion is true, we do not need to evaluate anything else on that line: that line definitely isn’t bad. Likewise, if we find a line where some premise is false, we do not need to evaluate anything else on that line.

With this in mind, consider how we might test the following for validity:

$$\neg L \rightarrow (J \vee L), \neg L \therefore J$$

The *first* thing we should do is evaluate the conclusion. If we find that the conclusion is *true* on some line, then that is not a bad line. So we can simply ignore the rest of the line. So at our first stage, we are left with something like:

<i>J</i>	<i>L</i>	$\neg L \rightarrow (J \vee L)$	$\neg L$	<i>J</i>
T	T			T
T	F			T
F	T	?	?	F
F	F	?	?	F

where the blanks indicate that we are not going to bother doing any more investigation (since the line is not bad) and the question-marks indicate that we need to keep investigating.

The easiest premise to evaluate is the second, so we next do that:

<i>J</i>	<i>L</i>	$\neg L \rightarrow (J \vee L)$	$\neg L$	<i>J</i>
T	T			T
T	F			T
F	T		F	F
F	F	?	T	F

Note that we no longer need to consider the third line on the table: it will not be a bad line, because (at least) one of premises is false on that line. Finally, we complete the truth table:

<i>J</i>	<i>L</i>	$\neg L \rightarrow (J \vee L)$	$\neg L$	<i>J</i>
T	T			T
T	F			T
F	T		F	F
F	F	T F F	T	F

The truth table has no bad lines, so the argument is valid. (Any valuation on which all the premises are true is a valuation on which the conclusion is true.)

It might be worth illustrating the tactic again. Let us check whether the following argument is valid

$$A \vee B, \neg(A \& C), \neg(B \& \neg D) \therefore (\neg C \vee D)$$

At the first stage, we determine the truth value of the conclusion. Since this is a disjunction, it is true whenever either disjunct is true, so we can speed things along a bit. We can then ignore every line apart from the few lines where the conclusion is false.

A	B	C	D	$A \vee B$	$\neg(A \& C)$	$\neg(B \& \neg D)$	$(\neg C \vee D)$
T	T	T	T				T
T	T	T	F	?	?	?	F F
T	T	F	T				T
T	T	F	F				T T
T	F	T	T				T
T	F	T	F	?	?	?	F F
T	F	F	T				T
T	F	F	F				T T
F	T	T	T				T
F	T	T	F	?	?	?	F F
F	T	F	T				T
F	T	F	F				T T
F	F	T	T				T
F	F	T	F	?	?	?	F F
F	F	F	T				T
F	F	F	F				T T

We must now evaluate the premises. We use shortcuts where we can:

A	B	C	D	$A \vee B$	$\neg(A \& C)$	$\neg(B \& \neg D)$	$(\neg C \vee D)$
T	T	T	T				T
T	T	T	F	T	F T		F F
T	T	F	T				T
T	T	F	F				T T
T	F	T	T				T
T	F	T	F	T	F T		F F
T	F	F	T				T
T	F	F	F				T T
F	T	T	T				T
F	T	T	F	T	T F	F TT	F F
F	T	F	T				T
F	T	F	F				T T
F	F	T	T				T
F	F	T	F	F			F F
F	F	F	T				T
F	F	F	F				T T

If we had used no shortcuts, we would have had to write 256 'T's or 'F's on this table. Using shortcuts, we only had to write 37. We have saved ourselves a *lot* of work.

We have been discussing shortcuts in testing for logical validity, but exactly the same shortcuts can be used in testing for entailment. By employing a similar notion of *bad* lines, you can save yourself a huge amount of work.

Practice exercises

A. Using shortcuts, determine whether each sentence is a tautology, a contradiction, or neither.

1. $\neg B \ \& \ B$
2. $\neg D \vee D$
3. $(A \ \& \ B) \vee (B \ \& \ A)$
4. $\neg[A \rightarrow (B \rightarrow A)]$
5. $A \leftrightarrow [A \rightarrow (B \ \& \ \neg B)]$
6. $\neg(A \ \& \ B) \leftrightarrow A$
7. $A \rightarrow (B \vee C)$
8. $(A \ \& \ \neg A) \rightarrow (B \vee C)$
9. $(B \ \& \ D) \leftrightarrow [A \leftrightarrow (A \vee C)]$

CHAPTER 16

Partial truth tables

Not taught in course. This knowledge will not be needed for the exam. (Although you can use these techniques in the exam if you wish.)

Sometimes, we do not need to know what happens on every line of a truth table. Sometimes, just a line or two will do.

Tautology. In order to show that a sentence is a tautology, we need to show that it is true on every valuation. That is to say, we need to know that it comes out true on every line of the truth table. So we need a complete truth table.

To show that a sentence is *not* a tautology, however, we only need one line: a line on which the sentence is false. Therefore, in order to show that some sentence is not a tautology, it is enough to provide a single valuation—a single line of the truth table—which makes the sentence false.

Suppose that we want to show that the sentence ' $(U \& T) \rightarrow (S \& W)$ ' is *not* a tautology. We set up a **PARTIAL TRUTH TABLE**:

S	T	U	W	$(U \& T) \rightarrow (S \& W)$
				F

We have only left space for one line, rather than 16, since we are only looking for one line on which the sentence is false. For just that reason, we have filled in 'F' for the entire sentence.

The main logical operator of the sentence is a conditional. In order for the conditional to be false, the antecedent must be true and the consequent must be false. So we fill these in on the table:

S	T	U	W	$(U \& T) \rightarrow (S \& W)$
				T F F

In order for the ' $(U \& T)$ ' to be true, both ' U ' and ' T ' must be true.

S	T	U	W	$(U \& T) \rightarrow (S \& W)$
	T	T		T T T F F

Now we just need to make ' $(S \& W)$ ' false. To do this, we need to make at least one of ' S ' and ' W ' false. We can make both ' S ' and ' W ' false if we want. All that matters is that the whole sentence turns out false on this line. Making an arbitrary decision, we finish the table in this way:

S	T	U	W	$(U \& T) \rightarrow (S \& W)$
F	T	T	F	T T T F F F F

We now have a partial truth table, which shows that ' $(U \& T) \rightarrow (S \& W)$ ' is not a tautology. Put otherwise, we have shown that there is a valuation which makes ' $(U \& T) \rightarrow (S \& W)$ ' false, namely, the valuation which makes ' S ' false, ' T ' true, ' U ' true and ' W ' false.

Contradiction. Showing that something is a contradiction requires a complete truth table: we need to show that there is no valuation which makes the sentence true; that is, we need to show that the sentence is false on every line of the truth table.

However, to show that something is *not* a contradiction, all we need to do is find a valuation which makes the sentence true, and a single line of a truth table will suffice. We can illustrate this with the same example.

S	T	U	W	$(U \& T) \rightarrow (S \& W)$
				T

To make the sentence true, it will suffice to ensure that the antecedent is false. Since the antecedent is a conjunction, we can just make one of them false. For no particular reason, we choose to make ' U ' false; and then we can assign whatever truth value we like to the other atomic sentences.

S	T	U	W	$(U \& T) \rightarrow (S \& W)$
F	T	F	F	F F T T F F F

Truth functional equivalence. To show that two sentences are logically equivalent, we must show that the sentences have the same truth value on every valuation. So this requires a complete truth table.

To show that two sentences are *not* logically equivalent, we only need to show that there is a valuation on which they have different truth values. So this requires only a one-line partial truth table: make the table so that one sentence is true and the other false.

Consistency. To show that some sentences are jointly consistent, we must show that there is a valuation which makes all of the sentences true, so this requires only a partial truth table with a single line.

To show that some sentences are jointly inconsistent, we must show that there is no valuation which makes all of the sentence true. So this requires a complete truth table: You must show that on every row of the table at least one of the sentences is false.

Validity. To show that an argument is valid, we must show that there is no valuation which makes all of the premises true and the conclusion false. So this requires a complete truth table. (Likewise for entailment.)

To show that argument is *invalid*, we must show that there is a valuation which makes all of the premises true and the conclusion false. So this requires only a one-line partial truth table on which all of the premises are true and the conclusion is false. (Likewise for a failure of entailment.)

This table summarises what is required:

	Yes	No
tautology?	complete truth table	one-line partial truth table
contradiction?	complete truth table	one-line partial truth table
equivalent?	complete truth table	one-line partial truth table
consistent?	one-line partial truth table	complete truth table
valid?	complete truth table	one-line partial truth table
entailment?	complete truth table	one-line partial truth table

Practice exercises

A. Use complete or partial truth tables (as appropriate) to determine whether these pairs of sentences are logically equivalent:

1. $A, \neg A$
2. $A, A \vee A$
3. $A \rightarrow A, A \leftrightarrow A$
4. $A \vee \neg B, A \rightarrow B$
5. $A \& \neg A, \neg B \leftrightarrow B$
6. $\neg(A \& B), \neg A \vee \neg B$
7. $\neg(A \rightarrow B), \neg A \rightarrow \neg B$
8. $(A \rightarrow B), (\neg B \rightarrow \neg A)$

B. Use complete or partial truth tables (as appropriate) to determine whether these sentences are jointly consistent, or jointly inconsistent:

1. $A \& B, C \rightarrow \neg B, C$
2. $A \rightarrow B, B \rightarrow C, A, \neg C$
3. $A \vee B, B \vee C, C \rightarrow \neg A$
4. $A, B, C, \neg D, \neg E, F$
5. $A \& (B \vee C), \neg(A \& C), \neg(B \& C)$
6. $A \rightarrow B, B \rightarrow C, \neg(A \rightarrow C)$

C. Use complete or partial truth tables (as appropriate) to determine whether each argument is valid or invalid:

1. $A \vee [A \rightarrow (A \leftrightarrow A)] \therefore A$
2. $A \leftrightarrow \neg(B \leftrightarrow A) \therefore A$
3. $A \rightarrow B, B \therefore A$
4. $A \vee B, B \vee C, \neg B \therefore A \& C$
5. $A \leftrightarrow B, B \leftrightarrow C \therefore A \leftrightarrow C$

D. Determine whether each sentence is a tautology, a contradiction, or a contingent sentence. Justify your answer with a complete or partial truth table where appropriate.

1. $A \rightarrow \neg A$
2. $A \rightarrow (A \& (A \vee B))$
3. $(A \rightarrow B) \leftrightarrow (B \rightarrow A)$
4. $A \rightarrow \neg(A \& (A \vee B))$
5. $\neg B \rightarrow [(\neg A \& A) \vee B]$
6. $\neg(A \vee B) \leftrightarrow (\neg A \& \neg B)$
7. $[(A \& B) \& C] \rightarrow B$
8. $\neg[(C \vee A) \vee B]$
9. $[(A \& B) \& \neg(A \& B)] \& C$

$$10. (A \& B) \rightarrow [(A \& C) \vee (B \& D)]$$

E. Determine whether each sentence is a tautology, a contradiction, or a contingent sentence. Justify your answer with a complete or partial truth table where appropriate.

1. $\neg(A \vee A)$
2. $(A \rightarrow B) \vee (B \rightarrow A)$
3. $[(A \rightarrow B) \rightarrow A] \rightarrow A$
4. $\neg[(A \rightarrow B) \vee (B \rightarrow A)]$
5. $(A \& B) \vee (A \vee B)$
6. $\neg(A \& B) \leftrightarrow A$
7. $A \rightarrow (B \vee C)$
8. $(A \& \neg A) \rightarrow (B \vee C)$
9. $(B \& D) \leftrightarrow [A \leftrightarrow (A \vee C)]$
10. $\neg[(A \rightarrow B) \vee (C \rightarrow D)]$

F. Determine whether each the following pairs of sentences are logically equivalent using complete truth tables. If the two sentences really are logically equivalent, write “equivalent.” Otherwise write, “not equivalent.”

1. A and $A \vee A$
2. A and $A \& A$
3. $A \vee \neg B$ and $A \rightarrow B$
4. $(A \rightarrow B)$ and $(\neg B \rightarrow \neg A)$
5. $\neg(A \& B)$ and $\neg A \vee \neg B$
6. $((U \rightarrow (X \vee X)) \vee U)$ and $\neg(X \& (X \& U))$
7. $((C \& (N \leftrightarrow C)) \leftrightarrow C)$ and $(\neg\neg\neg N \rightarrow C)$
8. $[(A \vee B) \& C]$ and $[A \vee (B \& C)]$
9. $((L \& C) \& I)$ and $L \vee C$

G. Determine whether each collection of sentences is jointly consistent or jointly inconsistent. Justify your answer with a complete or partial truth table where appropriate.

1. $A \rightarrow A, \neg A \rightarrow \neg A, A \& A, A \vee A$
2. $A \rightarrow \neg A, \neg A \rightarrow A$
3. $A \vee B, A \rightarrow C, B \rightarrow C$
4. $A \vee B, A \rightarrow C, B \rightarrow C, \neg C$
5. $B \& (C \vee A), A \rightarrow B, \neg(B \vee C)$

6. $(A \leftrightarrow B) \rightarrow B, B \rightarrow \neg(A \leftrightarrow B), A \vee B$
7. $A \leftrightarrow (B \vee C), C \rightarrow \neg A, A \rightarrow \neg B$
8. $A \leftrightarrow B, \neg B \vee \neg A, A \rightarrow B$
9. $A \leftrightarrow B, A \rightarrow C, B \rightarrow D, \neg(C \vee D)$
10. $\neg(A \& \neg B), B \rightarrow \neg A, \neg B$

H. Determine whether each argument is valid or invalid. Justify your answer with a complete or partial truth table where appropriate.

1. $A \rightarrow (A \& \neg A) \therefore \neg A$
2. $A \vee B, A \rightarrow B, B \rightarrow A \therefore A \leftrightarrow B$
3. $A \vee (B \rightarrow A) \therefore \neg A \rightarrow \neg B$
4. $A \vee B, A \rightarrow B, B \rightarrow A \therefore A \& B$
5. $(B \& A) \rightarrow C, (C \& A) \rightarrow B \therefore (C \& B) \rightarrow A$
6. $\neg(\neg A \vee \neg B), A \rightarrow \neg C \therefore A \rightarrow (B \rightarrow C)$
7. $A \& (B \rightarrow C), \neg C \& (\neg B \rightarrow \neg A) \therefore C \& \neg C$
8. $A \& B, \neg A \rightarrow \neg C, B \rightarrow \neg D \therefore A \vee B$
9. $A \rightarrow B \therefore (A \& B) \vee (\neg A \& \neg B)$
10. $\neg A \rightarrow B, \neg B \rightarrow C, \neg C \rightarrow A \therefore \neg A \rightarrow (\neg B \vee \neg C)$

I. Determine whether each argument is valid or invalid. Justify your answer with a complete or partial truth table where appropriate.

1. $A \leftrightarrow \neg(B \leftrightarrow A) \therefore A$
2. $A \vee B, B \vee C, \neg A \therefore B \& C$
3. $A \rightarrow C, E \rightarrow (D \vee B), B \rightarrow \neg D \therefore (A \vee C) \vee (B \rightarrow (E \& D))$
4. $A \vee B, C \rightarrow A, C \rightarrow B \therefore A \rightarrow (B \rightarrow C)$
5. $A \rightarrow B, \neg B \vee A \therefore A \leftrightarrow B$

CHAPTER 17

TFL vs English connectives

Not covered in this course. This is just a taster.

Consider the sentence:

1. Jan is neither bald nor not-bald.

To symbolize this sentence in TFL, we would offer something like ' $\neg J \& \neg \neg J$ '. This is a contradiction (check this with a truth-table), but sentence **1** does not itself seem like a contradiction; for we might have happily go on to add 'Jan is on the borderline of baldness'!

Third, consider the following sentence:

2. It's not the case that, if God exists, She answers malevolent prayers.

Symbolizing this in TFL, we would offer something like ' $\neg(G \rightarrow M)$ '. Now, ' $\neg(G \rightarrow M)$ ' entails ' G ' (again, check this with a truth table). So if we symbolize sentence **2** in TFL, it seems to entail that God exists. But that's strange: surely even an atheist can accept sentence **2**, without contradicting herself!

One lesson of this is that the symbolization of 2 as ' $\neg(G \rightarrow M)$ ' shows that 2 does not express what we intend. Perhaps we should rephrase it as

3. If God exists, She does not answer malevolent prayers.

and symbolize 3 as ' $G \rightarrow \neg M$ '. Now, if atheists are right, and there is no God, then ' G ' is false and so ' $G \rightarrow \neg M$ ' is true, and the puzzle disappears. However, if ' G ' is false, ' $G \rightarrow M$ ', i.e. 'If God exists, She answers malevolent prayers', is *also* true!

In different ways, these examples highlight some of the limits of working with a language (like TFL) that can *only* handle truth-functional connectives. Moreover, these limits give rise to some interesting questions in philosophical logic. The case of Jan's baldness (or otherwise) raises the general question of what logic we should use when dealing with *vague* discourse. The case of the atheist raises the question of how to deal with the (so-called) *paradoxes of the material conditional*. Part of the purpose of this course is to equip you with the tools to explore these questions of *philosophical logic*. But we have to walk before we can run; we have to become proficient in using TFL, before we can adequately discuss its limits, and consider alternatives.

PART IV

Natural deduction for TFL

CHAPTER 18

The very idea of natural deduction

Way back in §2, we said that an argument is valid iff it is impossible to make all of the premises true and the conclusion false.

In the case of TFL, this led us to develop truth tables. Each line of a complete truth table corresponds to a valuation. So, when faced with a TFL argument, we have a very direct way to assess whether it is possible to make all of the premises true and the conclusion false: just thrash through the truth table.

However, truth tables do not necessarily give us much *insight*. Consider two arguments in TFL:

$$P \vee Q, \neg P \therefore Q$$

$$P \rightarrow Q, P \therefore Q$$

Clearly, these are valid arguments. You can confirm that they are valid by constructing four-line truth tables, but we might say that they make use of different *forms* of reasoning. It might be nice to keep track of these different forms of inference.

One aim of a *natural deduction system* is to show that particular arguments are valid, in a way that allows us to understand the reasoning that the arguments might involve. We begin with very basic rules of

inference. These rules can be combined to offer more complicated arguments. Indeed, with just a small starter pack of rules of inference, we hope to capture all valid arguments.

This is a very different way of thinking about arguments.

With truth tables, we directly consider different ways to make sentences true or false. With natural deduction systems, we manipulate sentences in accordance with rules that we have set down as good rules. The latter promises to give us a better insight—or at least, a different insight—into how arguments work.

The move to natural deduction might be motivated by more than the search for insight. It might also be motivated by *necessity*. Once our arguments involve 5 atomic sentences, a truth table test for validity will require 32 lines of truth table. That’s quite a lot to check. But sometimes we might want to check such arguments.

- Alice, or Betty, or Carys, or Dan, or Ella stole the teacher’s pen.
- It wasn’t Alice.
- It wasn’t Betty,
- It wasn’t Carys,
- It wasn’t Dan
- ∴ It was Ella.

$$A \vee (B \vee (C \vee (D \vee E))), \neg A, \neg B, \neg C, \neg D \therefore E$$

And that will increase exponentially as more atomic sentences get added. Once an argument involves 20 atomic sentences,

- Alice, or Betty, or Carys, . . . , or Uli or Volker stole the teacher’s pen.
- It wasn’t Alice.
- It wasn’t Betty,
- ⋮
- It wasn’t Uli,
- Therefore: It was Volker.

This argument is also valid—as you might be able to tell—but to test it requires a truth table with $2^{20} = 1048576$ lines. In principle, we can set a machine to grind through truth tables and report back when it is finished. In practice, complicated arguments in TFL can become *intractable* if we use truth tables.

When we get to first-order logic (FOL) (beginning in chapter 28) the problem gets dramatically worse. There is nothing like the truth

table test for FOL. To assess whether or not an argument is valid, we have to reason about *all* interpretations, but, as we will see, there are infinitely many possible interpretations. We cannot even in principle set a machine to grind through infinitely many possible interpretations and report back when it is finished: it will *never* finish. We either need to come up with some more efficient way of reasoning about all interpretations, or we need to look for something different. We will be looking for something different; and we will develop natural deduction.¹

The modern development of natural deduction dates from simultaneous and unrelated papers by Gerhard Gentzen and Stanisław Jaśkowski (both in 1934). However, the natural deduction system that we will consider is based largely around work by Frederic Fitch (first published in 1952).

Natural deduction selects a few basic rules of inference and natural forms of reasoning and encodes these into a proof system. We will now see natural deduction for TFL. This system will form the basis also for natural deduction for FOL, which will also add rules for the quantifiers.

¹There are, in fact, systems that codify ways to reason about all possible interpretations which can be used for FOL in a similar way to the way we use truth tables for TFL. They were developed in the 1950s by Evert Beth and Jaakko Hintikka, but we will not follow this path.

CHAPTER 19

The First Basic Rules for TFL

The idea of natural deduction is how you might go about trying to convince an interlocutor that your argument is valid. You would generally do this by breaking your argument into smaller steps that are obviously valid and piecing these together.

Suppose you're trying to convince someone that $A \rightarrow (B \& C), A \therefore B$ is valid, and they don't see it. You can help them by breaking it up into two steps: first use Modus Ponens to get $B \& C$ and then note that B follows from $B \& C$. In more detail:

Grant me that $A \rightarrow (B \& C)$ and A are true. Then what else do we know to be true? Here's something: $B \& C$. Why? Because in general Modus Ponens is an excellent argument pattern: any argument of the form $X, X \rightarrow Y \therefore Y$ is valid: there are no valuations where X and $X \rightarrow Y$ are true but Y is false. So if our premises $A \rightarrow (B \& C)$ and A are true, then $B \& C$ must also be true: that's just Modus Ponens. So now from our supposition of $A \rightarrow (B \& C)$ and A , we now also know $B \& C$. What else follows from these three statements? Here's something: B . Why? Well, $B \& C$ is true; so certainly B must be true. So we know that from $A \rightarrow (B \& C)$ and A we can conclude B by walking someone through these two steps. This will be enough to show

that $A \rightarrow (B \& C)$, $A \therefore B$ is valid.

To keep track of what assumptions have been made and steps of the argument we will give precise forms that this argument should be written:

1		$A \rightarrow (B \& C)$	
2		A	
3		$B \& C$	Modus Ponens (which will be called $\rightarrow E$) 1, 2
4		B	&E 3

Each new line follows from the previous lines by an application of one of our simple steps of inference, e.g. Modus Ponens. We will always justify what allowed us to conclude the next line by adding a citation next to the line, e.g. “&E 3”. We are allowed to write B on line 4 because it follows from line 3 by an application of the rule &E.

You might think of it as a bag you’re collecting things to be accepted in. You have to grant the premises, they go in the bag for free, then we give certain rules that allow us to add additional statements which must be true so long as the other things already in the bag are true.

We will develop this **NATURAL DEDUCTION** system by specifying some basic rules of the system. For each connective, there will be **INTRODUCTION** rules, that allow us to prove a sentence that has that connective as the main logical operator, and **ELIMINATION** rules, that allow us to prove something given a sentence that has that connective as the main logical operator.

19.1 The idea of a formal proof

A *formal proof* is a sequence of sentences, some of which are marked as being initial assumptions (or premises). The last line of the formal proof is the conclusion. (Henceforth, we will simply call these ‘proofs’, but you should be aware that there are *informal proofs* too.)

As an illustration, consider:

$$\neg(A \vee B) \therefore \neg A \& \neg B$$

We will start a proof by writing the premise:

1		$\neg(A \vee B)$
---	--	------------------

Note that we have numbered the premise, since we will want to refer back to it. Indeed, every line on a proof is numbered, so that we can refer back to it.

Note also that we have drawn a line underneath the premise. Everything written above the line is an *assumption*. Everything written below the line will either be something which follows from the assumptions, or it will be some new assumption. We are hoping to conclude that ' $\neg A \ \& \ \neg B$ '; so we are hoping ultimately to conclude our proof with

$$n \quad | \quad \neg A \ \& \ \neg B$$

for some number n . It doesn't matter which line we end on, but we would obviously prefer a short proof to a long one.

Similarly, suppose we wanted to consider:

$$A \vee B, \neg(A \ \& \ C), \neg(B \ \& \ \neg D) \therefore \neg C \vee D$$

The argument has three premises, so we start by writing them all down, numbered, and drawing a line under them:

$$\begin{array}{l|l} 1 & A \vee B \\ 2 & \neg(A \ \& \ C) \\ 3 & \neg(B \ \& \ \neg D) \\ \hline \end{array}$$

and we are hoping to conclude with some line:

$$n \quad | \quad \neg C \vee D$$

If we can find a sequence of lines, each of which follows from the earlier lines by one of our rules, then this is a proof that the argument is valid, or we more simply say it is a proof of the argument.

All that remains to do is to explain each of the rules that we can use along the way from premises to conclusion.

It would be nice to present our rules by going through the various connectives and giving introduction and elimination rules. However, some rules will involve the tool of *subproofs*, which introduces some complexity. So we begin just by giving the rules that don't do so.

19.2 Reiteration

The very first rule is so breathtakingly obvious that it is surprising we bother with it at all.

If you already have shown something in the course of a proof, the *reiteration rule* allows you to repeat it on a new line. For example:

4		$A \& B$	
\vdots		\vdots	
10		$A \& B$	R 4

This indicates that we have written ' $A \& B$ ' on line 4. Now, at some later line—line 10, for example—we have decided that we want to repeat this. So we write it down again. We also add a citation which justifies what we have written. In this case, we write 'R', to indicate that we are using the reiteration rule, and we write '4', to indicate that we have applied it to line 4.

Here is a general expression of the rule:

		\vdots	
m		X	
		\vdots	
		X	R m

The point is that, if any sentence X occurs on some line, then we can repeat X on later lines. Each line of our proof must be justified by some rule, and here we have 'R m '. This means: Reiteration, applied to line m .

Two things need emphasising. First ' X ' is not a sentence of TFL. Rather, it is a symbol in the metalanguage, which we use when we want to talk about any sentence of TFL (see §9). Second, and similarly, ' m ' is not a numeral that will appear on a proof. Rather, it is a symbol in the metalanguage, which we use when we want to talk about any line number of a proof. In an actual proof, the lines are numbered '1', '2', '3', and so forth. But when we define the rule, we use variables to underscore the point that the rule may be applied at any point.

Why might this be useful? For example, we can now show $A \therefore A$ is valid using the proof:

1		A	
2		A	R 2

The rule really becomes useful, though, once we are dealing with sub-proofs, which we will see in the next chapter.

19.3 Modus Ponens

Consider the following argument:

If Jane is smart then she is fast. Jane is smart. \therefore Jane is fast.

This argument is certainly valid. In fact any argument of the form

$$X \rightarrow Y, X \therefore Y$$

is valid. We introduce a rule of natural deduction that encodes this idea. This is called *Modus Ponens*.

We introduce a rule of Natural Deduction which allows us to make this reasoning step. We will call it the “Conditional Elimination” rule (\rightarrow E). This choice of name is because we start with something including the connective \rightarrow and we derive something without the connective, that is we have *eliminated* the \rightarrow connective. For each connective we will have introduction and elimination rules, however we will wait until the next chapter to see Conditional Introduction.

In a simple use of this rule, we might just use it to derive from the premises $S \rightarrow F$ and S the conclusion F :

1		$S \rightarrow F$	
2		S	
3		F	\rightarrow E 1, 2

This would then be a natural deduction proof that $S \rightarrow F, S \therefore F$ is valid.

Each line, except for the premises which are taken as assumptions, has to be labelled with the rule it used. So here, we write “ \rightarrow E 1,2” to

say that we obtained line 3 by use of this rule \rightarrow Elimination applied to lines 1 and 2.

We can also apply the rule when our $X \rightarrow Y$ and X are not themselves premises but have themselves been derived in the course of the proof.

1	Premise 1	
2	Premise 2	
	\vdots	
8	$S \rightarrow F$	some rule
	\vdots	
15	S	another rule
	\vdots	
23	F	\rightarrow E 8, 15

It also can be that they appear in a different order, or that one appears in the premises, for example:

1	S	
	\vdots	
8	$S \rightarrow F$	
	\vdots	
23	F	\rightarrow E 8, 1

We write our general rule as:

		⋮	
m		$X \rightarrow Y$	
		⋮	
n		X	
		⋮	
		Y	$\rightarrow E\ m, n$

We can apply it to any X and Y . For example,

1		$(A \vee B) \rightarrow \neg F$	
2		$(A \vee B)$	
3		$\neg F$	$\rightarrow E\ 1, 2$

In this, X is $(A \vee B)$, and Y is $\neg F$.

We would typically now move to introducing Conditional Introduction. However, we will first do all the other rules of the system, because Conditional Introduction involves additional complexity.

19.4 Conjunction Introduction

Suppose we want to show that Ludwig is both reactionary and libertarian. One obvious way to do this would be as follows: first we show that Ludwig is reactionary; then we show that Ludwig is libertarian; then we put these two demonstrations together, to obtain the conjunction.

Our natural deduction system will capture this thought straightforwardly. In the example given, we might adopt the following symbolization key:

- R : Ludwig is reactionary
- L : Ludwig is libertarian

Perhaps we are working through a proof, and we have obtained ' R ' on line 8 and ' L ' on line 15. Then on any subsequent line we can obtain ' $R \ \& \ L$ '. For example our proof might contain the following lines:

8		R	
15		L	
23		$R \& L$	$\&I\ 8, 15$

Note that every line of our proof must either be an assumption, or must be justified by some rule. We cite ‘ $\&I\ 8, 15$ ’ here to indicate that the line is obtained by the rule of conjunction introduction ($\&I$) applied to lines 8 and 15. More generally, here is our conjunction introduction rule:

		\vdots	
m		X	
		\vdots	
n		Y	
		\vdots	
		$X \& Y$	$\&I\ m, n$

Two things need emphasising.

First ‘ X ’ and ‘ Y ’ are metavariables. They are not particular sentences of TFL but are there to play the role of any particular sentence (see §9).

Similarly, ‘ m ’ is not a numeral that will appear on a proof. Rather, it is a symbol in the metalanguage, which we use when we want to talk about any line number of a proof. In an actual proof, the lines are numbered ‘1’, ‘2’, ‘3’, and so forth. But when we define the rule, we use variables to underscore the point that the rule may be applied at any point.

To be clear, the statement of the rule is *schematic*. It is not itself a proof. ‘ X ’ and ‘ Y ’ are not sentences of TFL. Rather, they are symbols in the metalanguage, which we use when we want to talk about any sentence of TFL (see §9). Similarly, ‘ m ’ and ‘ n ’ are not a numerals that will appear on any actual proof. Rather, they are symbols in the metalanguage, which we use when we want to talk about any line number of any proof. In an actual proof, the lines are numbered ‘1’, ‘2’, ‘3’, and so forth, but when we define the rule, we use variables to emphasize that the rule may be applied at any point. The rule requires only that

we have both conjuncts available to us somewhere in the proof. They can be separated from one another, and they can appear in any order.

The rule is called ‘conjunction *introduction*’ because it introduces the symbol ‘&’ into our proof where it may have been absent.

19.5 Conjunction Elimination

Correspondingly, we have a rule that *eliminates* that symbol. Suppose you have shown that Ludwig is both reactionary and libertarian. You are entitled to conclude that Ludwig is reactionary. Equally, you are entitled to conclude that Ludwig is libertarian. Putting this together, we obtain our conjunction elimination rule(s):

		⋮	
m		$X \& Y$	
		⋮	
		X	$\&E\ m$

and equally:

		⋮	
m		$X \& Y$	
		⋮	
		Y	$\&E\ m$

The point is simply that, when you have a conjunction on some line of a proof, you can obtain either of the conjuncts by &E. (One point, might be worth emphasising: you can only apply this rule when conjunction is the main logical operator. So you cannot infer ‘ D ’ just from ‘ $C \vee (D \& E)$ ’!)

Even with just these two rules, we can start to see some of the power of our formal proof system. Consider:

- $[(A \vee B) \rightarrow (C \vee D)] \& \neg(E \vee F)$

- Therefore: $\neg(E \vee F) \ \& \ [(A \vee B) \rightarrow (C \vee D)]$

The main logical operator in both the premise and conclusion of this argument is ‘&’. In order to provide a proof, we begin by writing down the premise, which is our assumption. We draw a line below this: everything after this line must follow from our assumptions by (repeated applications of) our rules of inference. So the beginning of the proof looks like this:

$$1 \quad \left| \begin{array}{l} [(A \vee B) \rightarrow (C \vee D)] \ \& \ \neg(E \vee F) \end{array} \right.$$

From the premise, we can get each of the conjuncts by &E. The proof now looks like this:

$$\begin{array}{l|l} 1 & [(A \vee B) \rightarrow (C \vee D)] \ \& \ \neg(E \vee F) \\ 2 & [(A \vee B) \rightarrow (C \vee D)] & \text{\&E 1} \\ 3 & \neg(E \vee F) & \text{\&E 1} \end{array}$$

So by applying the &I rule to lines 3 and 2 (in that order), we arrive at the desired conclusion. The finished proof looks like this:

$$\begin{array}{l|l} 1 & [(A \vee B) \rightarrow (C \vee D)] \ \& \ \neg(E \vee F) \\ 2 & [(A \vee B) \rightarrow (C \vee D)] & \text{\&E 1} \\ 3 & \neg(E \vee F) & \text{\&E 1} \\ 4 & \neg(E \vee F) \ \& \ [(A \vee B) \rightarrow (C \vee D)] & \text{\&I 3, 2} \end{array}$$

This is a very simple proof, but it shows how we can chain rules of proof together into longer proofs. In passing, note that investigating this argument with a truth table would have required a staggering 256 lines; our formal proof required only four lines.

It is worth giving another example. Way back in §12.3, we noted that this argument is valid:

$$A \ \& \ (B \ \& \ C) \ \therefore \ (A \ \& \ B) \ \& \ C$$

To provide a proof corresponding with this argument, we start by writing:

$$1 \quad \left| \begin{array}{l} A \ \& \ (B \ \& \ C) \end{array} \right.$$

From the premise, we can get each of the conjuncts by applying &E twice. We can then apply &E twice more, so our proof looks like:

1	$A \& (B \& C)$	
2	A	&E 1
3	$B \& C$	&E 1
4	B	&E 3
5	C	&E 3

But now we can merrily reintroduce conjunctions in the order we wanted them, so that our final proof is:

1	$A \& (B \& C)$	
2	A	&E 1
3	$B \& C$	&E 1
4	B	&E 3
5	C	&E 3
6	$A \& B$	&I 2, 4
7	$(A \& B) \& C$	&I 6, 5

Recall that our official definition of sentences in TFL only allowed conjunctions with two conjuncts. The proof just given suggests that we could drop inner brackets in all of our proofs. However, this is not standard, and we will not do this. Instead, we will maintain our more austere bracketing conventions. (Though we will still allow ourselves to drop outermost brackets, for legibility.)

Let me offer one final illustration. When using the &I rule, there is no requirement that it is applied to two different sentences. So we can formally prove ' A ' from ' A ' as follows:

1	A	
2	$A \& A$	&I 1, 1
3	A	&E 2

Simple, but effective. In fact this shows that we didn't need to have the rule of Reiteration as we could always argue by &I then &E. But for ease we will allow Reiteration as a basic rule so you don't have to argue this way.

19.6 Biconditional

For the biconditional we encode the idea that $X \leftrightarrow Y$ is equivalent to $(X \rightarrow Y) \& (Y \rightarrow X)$.

The rules we give, then, are very closely related to the rules for conjunction: Two elimination rules

$$\begin{array}{c|l}
 & \vdots \\
 m & X \leftrightarrow Y \\
 & \vdots \\
 & X \rightarrow Y \quad \leftrightarrow E \ m
 \end{array}$$

and

$$\begin{array}{c|l}
 & \vdots \\
 m & X \leftrightarrow Y \\
 & \vdots \\
 & Y \rightarrow X \quad \leftrightarrow E \ m
 \end{array}$$

and an introduction rule:

$$\begin{array}{c|l}
 & \vdots \\
 m & X \rightarrow Y \\
 & \vdots \\
 n & Y \rightarrow X \\
 & \vdots \\
 & X \leftrightarrow Y \quad \leftrightarrow I \ m, \ n
 \end{array}$$

19.7 Disjunction Introduction

Suppose Ludwig is reactionary. Then Ludwig is reactionary or libertarian. After all, to say that Ludwig is reactionary or libertarian is to say something weaker than to say that Ludwig is reactionary.

Let me emphasize this point. Suppose Ludwig is reactionary. It follows that Ludwig is reactionary *or* he is a kumquat. Equally, it follows that Ludwig is reactionary *or* that kumquats are the only fruit. Equally, it follows that Ludwig is reactionary *or* that God is dead. Many of these are strange inferences to draw, but there is nothing *logically* wrong with them (even if they maybe violate all sorts of implicit conversational norms).

Armed with all this, we present the disjunction introduction rule(s):

$$\begin{array}{c|l}
 & \vdots \\
 m & X \\
 & \vdots \\
 & X \vee Y \qquad \vee I \ m
 \end{array}$$

and

$$\begin{array}{c|l}
 & \vdots \\
 m & X \\
 & \vdots \\
 & Y \vee X \qquad \vee I \ m
 \end{array}$$

Notice that Y can be *any* sentence whatsoever, so the following is a perfectly acceptable proof:

$$\begin{array}{c|l}
 1 & M \\
 \hline
 2 & M \vee ([(A \leftrightarrow B) \rightarrow (C \& D)] \leftrightarrow [E \& F]) \qquad \vee I \ 1
 \end{array}$$

Using a truth table to show this would have taken 128 lines.

19.8 Law of Excluded Middle

We will actually add another rule for how to introduce a disjunction. There are special kinds of disjunctions that don't need further justification: sentences of the form $X \vee \neg X$. In §14.1 we saw that $A \vee \neg A$ is a tautology: it is true on all valuations. More generally, any sentence of the form $X \vee \neg X$ is a tautology. The rule *Law of Excluded Middle* encodes this fact: it simply says that you are always allowed to write $X \vee \neg X$:

$$\begin{array}{c|l}
 & \vdots \\
 & X \vee \neg X \qquad \text{LEM}
 \end{array}$$

As always, X can be whatever you want, e.g. $(A \& (B \rightarrow C))$. Then the rule tells us, e.g. that you can write $(A \& (B \rightarrow C)) \vee \neg(A \& (B \rightarrow C))$ on any line of the proof.

To see the rule in action, consider:

$$A \rightarrow B, \neg A \rightarrow B \therefore B$$

We can give the following proof to show this is valid:

1	$A \rightarrow B$	
2	$\neg A \rightarrow B$	
3	$A \vee \neg A$	LEM
4	B	$\vee E$ 1, 2, 3

The law of Excluded Middle is often used in combination with Disjunction Elimination.

However, we will not yet introduce Disjunction Elimination. It will be introduced in §20.2

19.9 Contradiction

Instead of tackling negation directly, we will first think about *contradiction*.

Contradiction Introduction

An effective form of argument is to argue your opponent into contradicting themselves. At that point, you have them on the ropes. They have to give up at least one of their assumptions. We are going to make use of this idea in our proof system, by adding a new symbol, ' \perp ', to our proofs. This should be read as something like 'contradiction!' or 'reductio!' or 'but that's absurd!' The rule for introducing this symbol is that we can use it whenever we explicitly contradict ourselves, i.e. whenever we find both a sentence and its negation appearing in our proof:

		⋮	
m		X	
		⋮	
n		$\neg X$	
		⋮	
		\perp	$\perp I\ m,\ n$

It does not matter what order the sentence and its negation appear in, and they do not need to appear on adjacent lines.

Contradiction Elimination

Our elimination rule for ‘ \perp ’ is known as *ex falso quod libet*, or *explosion*. This means ‘anything follows from a contradiction’, and the idea is precisely that: if we obtained a contradiction, symbolized by ‘ \perp ’, then we can infer whatever we like. How can this be motivated, as a rule of argumentation? Well, consider the English rhetorical device ‘...and if *that’s* true, I’ll eat my hat’. Since contradictions simply cannot be true, if one *is* true then not only will I eat my hat, I’ll have it too.¹ Here is the formal rule:

		⋮	
m		\perp	
		X	$\perp E\ m$

Note that X can be *any* sentence whatsoever.

A final remark. We have said that ‘ \perp ’ should be read as something like ‘contradiction!’ but this does not tell us much about the symbol. There are, roughly, three ways to approach the symbol.

- We might regard ‘ \perp ’ as a new atomic sentence of TFL, but one which can only ever have the truth value False.

¹Thanks to Adam Caulton for this.

- We might regard ' \perp ' as an abbreviation for some canonical contradiction, such as ' $A \ \& \ \neg A$ '. This will have the same effect as the above—obviously, ' $A \ \& \ \neg A$ ' only ever has the truth value False—but it means that, officially, we do not need to add a new symbol to TFL.
- We might regard ' \perp ', not as a symbol of TFL, but as something more like a *punctuation mark* that appears in our proofs. (It is on a par with the line numbers and the vertical lines, say.)

There is something very philosophically attractive about the third option, but here we will *officially* adopt the first. ' \perp ' is to be read as a sentence letter that is always false. This means that we can manipulate it, in our proofs, just like any other sentence.

19.10 Strategies

We have a few more rules to introduce, but at this point we pause to mention a strategies to come up with proofs yourself:

Work backwards from what you want. The ultimate goal is to obtain the conclusion. Look at the conclusion and ask what the introduction rule is for its main logical operator. This gives you an idea of what should happen *just before* the last line of the proof. Then you can treat this line as if it were your goal. Ask what you could do to get to this new goal.

For example: If your conclusion is a conditional $X \ \& \ Y$, plan to use the $\&I$ rule. This requires finding both X and Y .

Work forwards from what you have. When you are starting a proof, look at the premises; later, look at the sentences that you have obtained so far. Think about the elimination rules for the main operators of these sentences. These will often tell you what your options are.

For a short proof, you might be able to eliminate the premises and introduce the conclusion. A long proof is formally just a number of short proofs linked together, so you can fill the gap by alternately working back from the conclusion and forward from the premises.

Sometimes, though, this won't yet be possible. For that, we need to finish off our list of the basic rules of natural deduction.

Practice exercises

A. The following ‘proofs’ are *incorrect*. Explain the mistakes they make.

1	$A \& (B \& C)$	
2	$(B \vee C) \rightarrow D$	
3	B	&E 1
4	$B \vee C$	\vee I 3
5	D	\rightarrow E 4, 2

B. The following proofs are missing their citations (rule and line numbers). Add them, to turn them into *bona fide* proofs. Additionally, write down the argument that corresponds to each proof.

1	$P \& S$	
2	$S \rightarrow R$	
3	S	
4	R	
5	$R \vee E$	

C. Give a proof corresponding to each of the following arguments:

1. $A \& (B \vee C) \therefore (B \vee C) \& A$
2. $(A \vee \neg A) \rightarrow B \therefore B$

CHAPTER 20

More Basic Rules for TFL

To introduce a conditional we need to introduce another kind of thing: a sub-proof.

20.1 Conditional Introduction

The following argument is valid:

Ludwig is reactionary. Therefore if Ludwig is libertarian,
then Ludwig is both reactionary *and* libertarian.

If someone doubted that this was valid, we might try to convince them otherwise by explaining ourselves as follows:

Assume that Ludwig is reactionary. Now, *additionally* assume that Ludwig is libertarian. Then by conjunction introduction—which we just discussed—Ludwig is both reactionary and libertarian. Of course, that’s conditional on the assumption that Ludwig is libertarian. But this just means that, if Ludwig is libertarian, then he is both reactionary and libertarian.

We might write this in a form that is closer to our natural deduction format:

1	Ludwig is reactionary	
2	Suppose that he is libertarian	
3	Then he is both libertarian and reactionary	&I 1, 2
4	Thus, <i>if</i> he is libertarian, then he is both libertarian and reactionary.	→I 2–3

The natural deduction format of lines and indentations is there to replace the words and context like “suppose that”. The “suppose that” used on line 2 is represented in the formal system by the indentation and additional line. And like the premises, this line does not need to be justified, it is taken as an assumption, and a line underneath it is drawn. What comes underneath this, on line 3, is still indented, and is within the context of the supposition that he is a libertarian. But once we move to line 4, the additional assumption is no longer in place. It has been *discharged*.

Now let’s present this again a little more formally: We started with one premise, ‘Ludwig is reactionary’, thus:

1	<u>R</u>
---	----------

The next thing we did is to make an *additional* assumption (‘Ludwig is libertarian’), for the sake of argument. To indicate that we are no longer dealing *merely* with our original assumption (‘R’), but with some additional assumption, we continue our proof as follows:

1	<u>R</u>	
2	<u>L</u>	

Note that we are *not* claiming, on line 2, to have proved ‘L’ from line 1, so we do not need to write in any justification for the additional assumption on line 2. We do, however, need to mark that it is an additional assumption. We do this by drawing a line under it (to indicate that it is an assumption) and by indenting it with a further vertical line (to indicate that it is additional).

With this extra assumption in place, we are in a position to use &I. So we can continue our proof:

1		R	
2			L
3			$R \& L$ &I 1, 2

So we have now shown that, on the additional assumption, ‘ L ’, we can obtain ‘ $R \& L$ ’. We can therefore conclude that, if ‘ L ’ obtains, then so does ‘ $R \& L$ ’. Or, to put it more briefly, we can conclude ‘ $L \rightarrow (R \& L)$ ’:

1		R	
2			L
3			$R \& L$ &I 1, 2
4		$L \rightarrow (R \& L)$	\rightarrow I 2–3

Observe that we have dropped back to using one vertical line. We have *discharged* the additional assumption, ‘ L ’, since the conditional itself follows just from our original assumption, ‘ R ’.

The general pattern at work here is the following. We first make an additional assumption, X ; and from that additional assumption, we prove Y . In that case, we know the following: If X , then Y . This is wrapped up in the rule for conditional introduction:

m			X	
			\vdots	
n			Y	
		$X \rightarrow Y$		\rightarrow I m – n

There can be as many or as few lines as you like between lines m and n .

It will help to offer a second illustration of \rightarrow I in action. Suppose we want to consider the following:

$$P \rightarrow Q, Q \rightarrow R \therefore P \rightarrow R$$

We start by listing *both* of our premises. Then, since we want to arrive at a conditional (namely, ‘ $P \rightarrow R$ ’), we additionally assume the antecedent to that conditional. Thus our main proof starts:

1		$P \rightarrow Q$	
2		$Q \rightarrow R$	
<hr/>			
3			P
<hr/>			

Note that we have made ‘ P ’ available, by treating it as an additional assumption, but now, we can use \rightarrow E on the first premise. This will yield ‘ Q ’. We can then use \rightarrow E on the second premise. So, by assuming ‘ P ’ we were able to prove ‘ R ’, so we apply the \rightarrow I rule—discharging ‘ P ’—and finish the proof. Putting all this together, we have:

1		$P \rightarrow Q$	
2		$Q \rightarrow R$	
<hr/>			
3			P
<hr/>			
4			Q \rightarrow E 1, 3
5			R \rightarrow E 2, 4
6		$P \rightarrow R$	\rightarrow I 3–5

The subproof also doesn’t need to start immediately. For example:

1		$(P \rightarrow Q) \& O$	
2		$Q \rightarrow R$	
<hr/>			
3		$P \rightarrow Q$	$\&$ E 1
<hr/>			
4			P
<hr/>			
5			Q \rightarrow E 3, 4
6			R \rightarrow E 2, 5
7		$P \rightarrow R$	\rightarrow I 4–6

20.2 Disjunction Elimination

The disjunction elimination rule also makes use of subproofs.

Suppose that Ludwig is reactionary or he is libertarian. What can you conclude? Not that Ludwig is reactionary; it might be that he is libertarian instead. Equally, not that Ludwig is libertarian; for he might merely be reactionary. Disjunctions, just by themselves, are hard to work with.

But suppose that we could somehow show both of the following: first, that Ludwig's being reactionary entails that he is an Austrian economist: second, that Ludwig's being libertarian entails that he is an Austrian economist. Then if we know that Ludwig is reactionary or libertarian, then we know that, whichever he is, Ludwig is an Austrian economist. This insight can be expressed in the following rule, which is our disjunction elimination (\vee E) rule:

m	$X \vee Y$	
i	X	
	\vdots	
j	Z	
k	Y	
	\vdots	
l	Z	
	Z	\vee E $m, i-j, k-l$

This is obviously a bit clunkier to write down than our previous rules, but the point is fairly simple. Suppose we have some disjunction, $X \vee Y$. Suppose we have two subproofs, showing us that Z follows from the assumption that X , and that Z follows from the assumption that Y . Then we can infer Z itself. As usual, there can be as many lines as you like between i and j , and as many lines as you like between k and l . Moreover, the subproofs and the disjunction can come in any order, and do not have to be adjacent.

Some examples might help illustrate this. Consider this argument:

$$(P \ \& \ Q) \vee (P \ \& \ R) \therefore P$$

A proof corresponding to this argument is:

1		$(P \ \& \ Q) \vee (P \ \& \ R)$	
2		$P \ \& \ Q$	
3		P	&E 2
4		$P \ \& \ R$	
5		P	&E 4
6		P	\vee E 1, 2–3, 4–5

Consider the following brain teaser:¹

Three people are standing in a row looking at each other.



Alice



Bob



Charlie

Alice is happy. Charlie is not happy. Is there someone who is happy who is looking at someone who is not happy?

... Think about it!

... Answer: Yes. Our Disjunction Elimination rule along with the Law of Excluded Middle allow us to show this. We can demonstrate this in the following argument, which we present in a pseudo-formal style.²

¹Originally by Hector Levesque.

²Though, actually, this is most naturally formulated as a validity claim of First Order Logic. We'll walk through the formal proof as formulated in First Order Logic in §38.2.

1	Bob is either happy or he's not happy
2	<u>Suppose Bob is happy</u>
3	Then happy Bob is looking at not-happy Charlie
4	So someone who is happy is looking at someone who is not
5	<u>Suppose Bob is not happy</u>
6	Then happy Alice is looking at not-happy Bob
7	So someone who is happy is looking at someone who is not
8	Therefore, someone who is happy is looking at someone who is not.

Are you convinced now that someone who is happy is looking at someone who is not happy? If not, find a friend and work through it together. Sometimes it can really help to try walking through the argument together.

Coming up with this sort of argument does just take that moment of inspiration to see how this argument will go (that's why it's a brain teaser). This is often the case with arguments that involve the law of excluded middle. We pick it out of nowhere and have to use our inspiration to see how it might be useful. But hopefully, with more examples you'll become familiar with cases where it might be of use. A strategy that might help is: it's a backup option if everything else fails. If it doesn't look like there's any elimination rules to use on your premises or any introduction rules that can get you to your conclusion, then perhaps LEM is the way forwards.

One more example:

$$P \therefore (P \ \& \ D) \vee (P \ \& \ \neg D)$$

Here is a proof corresponding with the argument:

1	P	
2	$D \vee \neg D$	LEM
3	D	
4	$P \& D$	&I 1, 3
5	$(P \& D) \vee (P \& \neg D)$	\vee I 4
6	$\neg D$	
7	$P \& \neg D$	&I 1, 6
8	$(P \& D) \vee (P \& \neg D)$	\vee I 7
9	$(P \& D) \vee (P \& \neg D)$	\vee E 2, 3–5, 6–8

20.3 Negation

Negation Introduction

If assuming something leads you to a contradiction, then the assumption must be wrong. This thought motivates the following rule:

m		X	
		\vdots	
n		\perp	
	$\neg X$		\neg I m – n

To see this in practice, and interacting with negation, consider this proof:

1	D	
2	$\neg D$	
3	\perp	\perp I 1, 2
4	$\neg\neg D$	\neg I 2–3

Proof by Contradiction

The next rule is quite similar. If assuming that something is false leads you to a contradiction, then that assumption must be wrong — and so that ‘something’ must in fact be true.

m			$\neg X$	
			\vdots	
n			\perp	
				X
				PbC $m-n$

This is called the method of *proof by contradiction*, or *indirect proof*. It allows us to prove something by assuming its negation and showing that it leads to contradiction. This technique is very common in mathematics.³

For example:

1		$\neg\neg D$	
2			$\neg D$
3			\perp
			\perp I 1, 2
4		D	PbC 2-3

There is no explicit rule for Negation Elimination. Though both Proof by Contradiction and Contradiction Introduction involve eliminating negations.

20.4 Additional assumptions and subproofs

The rules we have just seen involve the idea of making additional assumptions. These need to be handled with some care.

Consider this proof:

³For example it is used to show that $\sqrt{2}$ is irrational. We begin by assuming that $\sqrt{2}$ is rational (i.e. that there is a fraction $\frac{p}{q}$ whose square is 2) and then derive contradiction from this assumption.

1		A	
2			B
3			B R 2
4		$B \rightarrow B$	\rightarrow I 2–3

This is perfectly in keeping with the rules we have laid down already, and it should not seem particularly strange. Since ' $B \rightarrow B$ ' is a tautology, no particular premises should be required to prove it.

But suppose we now tried to continue the proof as follows:

1		A	
2			B
3			B R 2
4		$B \rightarrow B$	\rightarrow I 2–3
5		B	naughty attempt to invoke \rightarrow E 4, 3

If we were allowed to do this, it would be a disaster. It would allow us to prove any atomic sentence letter from any other atomic sentence letter. However, if you tell me that Anne is fast (symbolized by ' A '), we shouldn't be able to conclude that Queen Boudica stood twenty-feet tall (symbolized by ' B ')! We must be prohibited from doing this, but how are we to implement the prohibition?

We can describe the process of making an additional assumption as one of performing a *subproof*: a subsidiary proof within the main proof. When we start a subproof, we draw another vertical line to indicate that we are no longer in the main proof. Then we write in the assumption upon which the subproof will be based. A subproof can be thought of as essentially posing this question: *what could we show, if we also make this additional assumption?*

When we are working within the subproof, we can refer to the additional assumption that we made in introducing the subproof, and to anything that we obtained from our original assumptions. (After all, those original assumptions are still in effect.) At some point though, we will want to stop working with the additional assumption: we will want to return from the subproof to the main proof. To indicate that

we have returned to the main proof, the vertical line for the subproof comes to an end. At this point, we say that the subproof is closed:

A subproof is **CLOSED** when the vertical line for the subproof comes to an end. At that point we say the assumption has been **DISCHARGED**

We typically do this when we use one of our rules that involve subproofs, such as \rightarrow I. We introduced the assumption X to allow us to conclude Y ; and this reasoning allows us to close the subproof and conclude $X \rightarrow Y$, which no longer relies on the assumption X . Having closed a subproof, we have set aside the additional assumption, so it will be illegitimate to draw upon anything that depends upon that additional assumption. Thus we stipulate:

Any rule whose citation requires mentioning individual lines can mention any earlier lines, *except* for those lines which occur within a closed subproof.

Put another way: you cannot refer back to anything that was obtained using discharged assumptions

This stipulation rules out the disastrous attempted proof above. The rule of \rightarrow E requires that we cite two individual lines from earlier in the proof. In the purported proof, above, one of these lines (namely, line 4) occurs within a subproof that has (by line 6) been closed. This is illegitimate.

Subproofs, then, allow us to think about what we could show, if we made additional assumptions. The point to take away from this is not surprising—in the course of a proof, we have to keep very careful track of what assumptions we are making, at any given moment. Our proof system does this very graphically. (Indeed, that's precisely why we have chosen to use *this* proof system.)

Once we have started thinking about what we can show by making additional assumptions, nothing stops us from posing the question of what we could show if we were to make *even more* assumptions. This might motivate us to introduce a subproof within a subproof. Here is an example which only uses the rules of proof that we have considered so far:

1	A	
2	B	
3	C	
4	$A \& B$	$\&I$ 1, 2
5	$C \rightarrow (A \& B)$	$\rightarrow I$ 3–4
6	$B \rightarrow (C \rightarrow (A \& B))$	$\rightarrow I$ 2–5

Notice that the citation on line 4 refers back to the initial assumption (on line 1) and an assumption of a subproof (on line 2). This is perfectly in order, since neither assumption has been discharged at the time (i.e. by line 4).

Again, though, we need to keep careful track of what we are assuming at any given moment. Suppose we tried to continue the proof as follows:

1	A	
2	B	
3	C	
4	$A \& B$	$\&I$ 1, 2
5	$C \rightarrow (A \& B)$	$\rightarrow I$ 3–4
6	$B \rightarrow (C \rightarrow (A \& B))$	$\rightarrow I$ 2–5
7	$C \rightarrow (A \& B)$	naughty attempt to invoke $\rightarrow I$ 3–4

This would be awful. If we tell you that Anne is smart, you should not be able to infer that, if Cath is smart (symbolized by ‘ C ’) then *both* Anne is smart and Queen Boudica stood 20-feet tall! But this is just what such a proof would suggest, if it were permissible.

The essential problem is that the subproof that began with the assumption ‘ C ’ depended crucially on the fact that we had assumed ‘ B ’ on line 2. By line 6, we have *discharged* the assumption ‘ B ’: we have stopped asking ourselves what we could show, if we also assumed ‘ B ’. So it is simply cheating, to try to help ourselves (on line 7) to the subproof that began with the assumption ‘ C ’. Thus we stipulate, much as before:

Any rule whose citation requires mentioning an entire subproof can mention any earlier subproof, *except* for those subproofs which occur within some *other* closed subproof.

The attempted disastrous proof violates this stipulation. The subproof of lines 3–4 occurs within a subproof that ends on line 5. So it cannot be invoked in line 7.

It is always permissible to open a subproof with any assumption. However, there is some strategy involved in picking a useful assumption. Starting a subproof with an arbitrary, wacky assumption would just waste lines of the proof. In order to obtain a conditional by \rightarrow I, for instance, you must assume the antecedent of the conditional in a subproof.

Equally, it is always permissible to close a subproof and discharge its assumptions. However, it will not be helpful to do so until you have reached something useful.

20.5 Proof Strategies

These are all of the basic rules for the proof system for TFL. For ease of reference, they're listed again in appendix C.

There is no simple recipe for proofs, and there is no substitute for practice. Here, though, are some rules of thumb and strategies to keep in mind.

Work backwards from what you want. The ultimate goal is to obtain the conclusion. Look at the conclusion and ask what the introduction rule is for its main logical operator. This gives you an idea of what should happen *just before* the last line of the proof. Then you can treat this line as if it were your goal. Ask what you could do to get to this new goal.

For example: If your conclusion is a conditional $X \rightarrow Y$, plan to use the \rightarrow I rule. This requires starting a subproof in which you assume X . The subproof ought to end with Y . So, what can you do to get Y ?

Work forwards from what you have. When you are starting a proof, look at the premises; later, look at the sentences that you have obtained

so far. Think about the elimination rules for the main operators of these sentences. These will often tell you what your options are.

For a short proof, you might be able to eliminate the premises and introduce the conclusion. A long proof is formally just a number of short proofs linked together, so you can fill the gap by alternately working back from the conclusion and forward from the premises.

Try proceeding indirectly. If you cannot find a way to show X directly, try starting by assuming $\neg X$. If a contradiction follows, then you will be able to obtain X by PbC.

Law of Excluded Middle. If you're hitting a blank, try seeing if there's some instance of LEM that might help you. These arguments do just need some inspiration to see the instance of LEM that'll be helpful.

Persist. Try different things. If one approach fails, then try something else.

If the argument is actually valid (which is defined using truth-tables) there will be a proof of it somehow...

Practice exercises

A. The following two 'proofs' are *incorrect*. Explain the mistakes they make.

1	$\neg L \rightarrow (A \ \& \ L)$	
2	$L \vee \neg L$	LEM
3	$\neg L$	
4	A	\rightarrow E 1, 3
5	L	
6	\perp	\perp I 5, 3
7	A	\perp E 6
8	A	\vee E 2, 3–4, 5–7

1		$A \& (B \& C)$	
2		$(B \vee C) \rightarrow D$	
3		B	&E 1
4		$B \vee C$	\vee I 3
5		D	\rightarrow E 4, 2

B. The following three proofs are missing their citations (rule and line numbers). Add them, to turn them into *bona fide* proofs. Additionally, write down the argument that corresponds to each proof.

1		$P \& S$
2		$S \rightarrow R$
3		S
4		R
5		$R \vee E$

1		$A \rightarrow D$
2		$A \& B$
3		A
4		D
5		$D \vee E$
6		$(A \& B) \rightarrow (D \vee E)$

1		$\neg L \rightarrow (J \vee L)$
2		$\neg L$
3		$J \vee L$
4		J
5		J
6		$J \rightarrow J$
7		L
8		\perp
9		J
10		$L \rightarrow J$
11		J

CHAPTER 21

Proofs and Validity

The system of rules we have set up is not just a game. It helps us understand the validity of arguments.

An argument $X_1, X_2, \dots, X_n \therefore Y$ may have a proof in the system of natural deduction. Such a proof may look something like:

1		X_1
2		X_2
\vdots		
n		X_n
<hr/>		
		\vdots
		Y

That is, it will start with the premises as assumptions, and proceed following the rules we have given and finishing with the conclusion. It might also have subproofs along the way, something like:

1		X_1	
2		X_2	
n		X_n	
<hr/>			
		\vdots	
m		Some additional assumption	
<hr/>			
		\vdots	
m		Maybe some more assumptions	
<hr/>			
		\vdots	
		\vdots	
		\vdots	
		Y	

But any subproofs need to have been closed by the time we get to Y .
 So, for example if we gave

1		A	
<hr/>			
2		B	
<hr/>			
3		B	R 2

this does not count as a proof corresponding to the argument $A \therefore B$.
 But

1		A	
<hr/>			
2		B	
<hr/>			
3		B	R 2
4		$B \rightarrow B$	\rightarrow I 2–3

counts as a proof corresponding to $A \therefore B \rightarrow B$ as the subproof has
 been closed on line 4.

So we have now said when we have a proof in our natural deduction system that corresponds to a particular argument. If we can find a proof then we know that the argument is valid.

If there is a proof in natural deduction corresponding to the argument $X_1 \dots X_n \therefore Y$, then this argument $X_1 \dots X_n \therefore Y$ is valid.

This property of our proof system is called **SOUNDNESS**. It holds because we only chose rules that matched valid reasoning steps. Recall that $A \rightarrow B, B \therefore A$ is invalid. Had we added a rule such as

	\vdots	
n	$X \rightarrow Y$	
	\vdots	
m	Y	
	\vdots	
	X	Do not do this. Does not follow from n, m

we would then have been able to construct a proof corresponding to the invalid argument $A \rightarrow B, B \therefore A$. We do not have such a rule in our system. All the rules we gave in our system will result in proofs of valid arguments.

We can actually strengthen the link between proofs corresponding to arguments and those argument's validity:

If an argument is valid, then there is a proof of it in natural deduction.

This property of our proof system is called **COMPLETENESS**.

So for every valid argument there will be some proof. This doesn't mean it is always easy to come up with such a proof, but there will be one. Persist!

Practice exercises

A. Show that each of the following arguments are valid using natural deduction:

1. $J \rightarrow \neg J \therefore \neg J$
2. $Q \rightarrow (Q \ \& \ \neg Q) \therefore \neg Q$
3. $A \rightarrow (B \rightarrow C) \therefore (A \ \& \ B) \rightarrow C$
4. $K \ \& \ L \therefore K \leftrightarrow L$
5. $(C \ \& \ D) \vee E \therefore E \vee D$
6. $A \leftrightarrow B, B \leftrightarrow C \therefore A \leftrightarrow C$
7. $\neg F \rightarrow G, F \rightarrow H \therefore G \vee H$
8. $(Z \ \& \ K) \vee (K \ \& \ M), K \rightarrow D \therefore D$
9. $P \ \& \ (Q \vee R), P \rightarrow \neg R \therefore Q \vee E$
10. $S \leftrightarrow T \therefore S \leftrightarrow (T \vee S)$
11. $\neg(P \rightarrow Q) \therefore \neg Q$
12. $\neg(P \rightarrow Q) \therefore P$

CHAPTER 22

Additional rules for TFL

In §19, we introduced the basic rules of our proof system for TFL. In this section, we will add some additional rules to our system. These will make our system much easier to work with. (However, in §23 we will see that they are not strictly speaking *necessary*.)

22.1 Disjunctive syllogism

Here is a very natural argument form.

Elizabeth is in Massachusetts or in DC. She is not in DC.
So, she is in Massachusetts.

This inference pattern is called *disjunctive syllogism*. We add it to our proof system as follows:

m	$X \vee Y$	
n	$\neg X$	
	Y	DS m, n

and

m	$X \vee Y$	
n	$\neg Y$	
	X	DS m, n

As usual, the disjunction and the negation of one disjunct may occur in either order and need not be adjacent.

22.2 Modus tollens

Another useful pattern of inference is embodied in the following argument:

If Mitt has won the election, then he is in the White House.
He is not in the White House. So he has not won the election.

This inference pattern is called *modus tollens*. The corresponding rule is:

m	$X \rightarrow Y$	
n	$\neg Y$	
	$\neg X$	MT m, n

As usual, the premises may occur in either order.

22.3 Double-negation

A sentence $\neg\neg X$ is always logically equivalent to X . We can add rules to our system that encode this idea: allowing us to immediately eliminate or introduce double negations:

m	$\neg\neg X$	
	X	DNE m

m	X	
	$\neg\neg X$	DNI m

That said, you should be aware that in ordinary language we can sometimes speak in a way that is similar to, but not quite, a double negation. Consider: ‘Jane is not *unhappy*’. Arguably, one cannot infer ‘Jane is happy’. Perhaps the speaker is using this unusual indirect phrasing to draw attention to the possible difference between ‘unhappy’ and ‘not unhappy’. Perhaps what they mean to suggest is that ‘Jane is in a state of profound indifference’. Here, then, ‘Jane is unhappy’ should not be thought of as equivalent to ‘It is not the case that Jane is happy’, and it should not be symbolised as $\neg H$ but should rather be a separate atomic sentence. So ‘Jane is not unhappy’ is not then seen as a double negation.

22.4 De Morgan Rules

Our final additional rules are called De Morgan’s Laws. (These are named after Augustus De Morgan.) The shape of the rules should be familiar from truth tables.

The first De Morgan rule is:

m	$\neg(X \& Y)$	
	$\neg X \vee \neg Y$	DeM m

The second De Morgan is the reverse of the first:

m	$\neg X \vee \neg Y$	
	$\neg(X \& Y)$	DeM m

The third De Morgan rule is the *dual* of the first:

$$\begin{array}{l|l}
 m & \neg(X \vee Y) \\
 & \neg X \ \& \ \neg Y \qquad \text{DeM } m
 \end{array}$$

And the fourth is the reverse of the third:

$$\begin{array}{l|l}
 m & \neg X \ \& \ \neg Y \\
 & \neg(X \vee Y) \qquad \text{DeM } m
 \end{array}$$

There are many more rules one could add to the system as derived rules. But these are all the ones we'll introduce.

Practice exercises

A. The following proofs are missing their citations (rule and line numbers). Add them wherever they are required:

$$\begin{array}{l|l}
 1 & W \rightarrow \neg B \\
 2 & A \ \& \ W \\
 3 & B \vee (J \ \& \ K) \\
 \hline
 4 & W \\
 5 & \neg B \\
 6 & J \ \& \ K \\
 7 & K
 \end{array}$$

$$\begin{array}{l|l}
 1 & \neg O \rightarrow L \\
 2 & L \vee \neg O \\
 \hline
 3 & \neg L \\
 \hline
 4 & \neg O \\
 5 & L \\
 6 & \perp \\
 7 & \neg L \rightarrow \perp \\
 8 & \neg \neg L \\
 9 & L
 \end{array}$$

1	$Z \rightarrow (C \ \& \ \neg N)$
2	$\neg Z \rightarrow (N \ \& \ \neg C)$
3	$Z \vee \neg Z$
4	Z
5	$C \ \& \ \neg N$
6	C
7	$N \vee C$
8	$Z \rightarrow (N \vee C)$
9	$\neg Z$
10	$N \ \& \ \neg C$
11	N
12	$N \vee C$
13	$\neg Z \rightarrow (N \vee C)$
14	$N \vee C$

B. Give a proof for each of these arguments:

1. $E \vee F, F \vee G, \neg F \therefore E \ \& \ G$
2. $M \vee (N \rightarrow M) \therefore \neg M \rightarrow \neg N$
3. $(M \vee N) \ \& \ (O \vee P), N \rightarrow P, \neg P \therefore M \ \& \ O$
4. $(X \ \& \ Y) \vee (X \ \& \ Z), \neg(X \ \& \ D), D \vee M \therefore M$

CHAPTER 23

Derived rules

In this section, we will see why we introduced the rules of our proof system in two separate batches. In particular, we want to show that the additional rules of §22 are not strictly speaking necessary, but can be derived from the basic rules of §19.

23.1 Derivation of Disjunctive syllogism

Suppose that you are in a proof, and you have something of this form:

$$\begin{array}{l|l} m & X \vee Y \\ n & \neg X \end{array}$$

You now want, on line k , to prove Y . You can do this with the rule of DS, introduced in §22, but equally well, you can do this with the *basic* rules of §19:

m	$X \vee Y$	
n	$\neg X$	
k	X	
$k+1$	\perp	$\perp I\ k, n$
$k+2$	Y	$\perp E\ k+1$
$k+3$	Y	
$k+4$	Y	$R\ k+3$
$k+5$	Y	$\vee E\ m, k-k+2, k+3-k+4$

To be clear: this is not a proof. Rather, it is a proof *scheme*. (This is why we use letters like m and k to label the lines of the proof rather than numbers.) Whatever sentences of TFL we plugged in for ‘ X ’ or ‘ Y ’, and whatever lines we were working on, we could produce a bona fide proof. So you can think of this as a recipe for producing proofs.

Indeed, it is a recipe which shows us that, anything we can prove using the rule DS, we can prove (with a few more lines) using just the other rules of §19.

23.2 Derivation of Modus Tollens

Suppose in the course of your proof you already have $X \rightarrow Y$, say on line m , and $\neg Y$ on line n . At some later line, k , you want to get $\neg X$. You can do this with the rule of Modus Tollens (MT), introduced in §22. But you could also do this with the *basic* rules of §19:

m	$X \rightarrow Y$	
n	$\neg Y$	
k	X	
$k+1$	Y	$\rightarrow E\ m, k$
$k+2$	\perp	$\perp I\ k+1, n$
$k+3$	$\neg X$	$\neg I\ k-k+2$

Again, the rule of MT can be derived from the *basic* rules of §19.

23.3 Derivation of Double-negation rules

Consider the following deduction schema:

m	X	
j	$\neg X$	
$j+1$	\perp	$\perp I \ j, m$
$j+2$	$\neg\neg X$	$\neg I \ j-(j+1)$

and

m	$\neg\neg X$	
j	$\neg X$	
$j+1$	\perp	$\perp I \ j, m$
$j+2$	X	$PbC \ j-(j+1)$

So again, we can derive the double negations rules from the *basic* rules of §19.

23.4 Derivation of De Morgan rules

Here is a demonstration of how we could derive the first De Morgan rule:

m	$\neg(X \& Y)$	
j	$X \vee \neg X$	LEM
k	X	
$k+1$	Y	
$k+2$	$X \& Y$	$\&I\ k, k+1$
$k+3$	\perp	$\perp I\ k+2, m$
$k+4$	$\neg Y$	$\neg I\ k+1-k+3$
$k+5$	$\neg X \vee \neg Y$	$\vee I\ k+4$
$k+6$	$\neg X$	
$k+7$	$\neg X \vee \neg Y$	$\vee I\ k+6$
$k+8$	$\neg X \vee \neg Y$	$\vee E\ j, k-k+5, k+6-k+7$

Here is a demonstration of how we could derive the second De Morgan rule:

m	$\neg X \vee \neg Y$	
k	$X \& Y$	
$k+1$	X	$\&E\ k$
$k+2$	Y	$\&E\ k$
$k+3$	$\neg X$	
$k+4$	\perp	$\perp I\ k+1, k+3$
$k+5$	$\neg Y$	
$k+6$	\perp	$\perp I\ k+2, k+5$
$k+7$	\perp	$\vee E\ m, k+3-k+4, k+5-k+6$
$k+8$	$\neg(X \& Y)$	$\neg I\ k-k+7$

Similar demonstrations can be offered explaining how we could derive the third and fourth De Morgan rules. These are left as exercises.

Practice exercises

A. Provide proof schemes that justify the addition of the third and fourth De Morgan rules as derived rules.

B. The proofs you offered in response to the practice exercises of §§22–25 used derived rules. Replace the use of derived rules, in such proofs, with only basic rules. You will find some ‘repetition’ in the resulting proofs; in such cases, offer a streamlined proof using only basic rules. (This will give you a sense, both of the power of derived rules, and of how all the rules interact.)

CHAPTER 24

Soundness and completeness

(Non-examinable) A very important result:

A TFL argument is valid if and only if it can be given a proof in this natural deduction system.

In this chapter, we explain a bit more about how such an argument would go. Soundness is proved in more detail in §48.

24.1 Entails

For this chapter we will make use of a further symbols:

We use the symbol \models as shorthand for ‘entails’. Rather than saying that the TFL sentences X_1, X_2, \dots and X_n together entail Y , we will abbreviate this by:

$$X_1, X_2, \dots, X_n \models Y$$

The symbol ‘ \models ’ is known as *the double-turnstile*, since it looks like a turnstile with two horizontal beams.

Let me be clear. ‘ \models ’ is not a symbol of TFL. Rather, it is a symbol of our metalanguage, augmented English (recall the difference between object language and metalanguage from §9). So the metalanguage sentence:

$$\bullet P, P \rightarrow Q \models Q$$

is just an abbreviation for the English sentence:

- The TFL sentences ‘ P ’ and ‘ $P \rightarrow Q$ ’ entail ‘ Q ’

Note that there is no limit on the number of TFL sentences that can be mentioned before the symbol ‘ \models ’. Indeed, we can even consider the limiting case:

$$\models Y$$

24.2 Proves

The following expression:

$$X_1, X_2, \dots, X_n \vdash Y$$

means that there is some proof which starts with assumptions among X_1, X_2, \dots, X_n and ends with Y (and contains no undischarged assumptions other than those we started with). Derivatively, we will write:

$$\vdash X$$

to mean that there is a proof of X with no assumptions.

The symbol ‘ \vdash ’ is called the *single turnstile*. We want to emphasize that this is not the double turnstile symbol (‘ \models ’) that we introduced to symbolize entailment. The single turnstile, ‘ \vdash ’, concerns the existence of proofs; the double turnstile, ‘ \models ’, concerns the existence of valuations (or interpretations, when used for FOL). *They are very different notions.*

24.3 Their equivalence

However, it turns out that they are equivalent. That is:

$$\begin{aligned} X_1, X_2, \dots, X_n \vdash Y \\ \text{if and only if} \\ X_1, X_2, \dots, X_n \models Y \end{aligned}$$

A full proof here goes well beyond the scope of this book. However, we can sketch what it would be like.

Soundness

This argument from \vdash to \models is the problem of **SOUNDNESS**. A proof system is **SOUND** if there are no derivations of arguments that can be shown invalid by truth tables. Demonstrating that the proof system is sound would require showing that *any* possible proof is the proof of a valid argument. It would not be enough simply to succeed when trying to prove many valid arguments and to fail when trying to prove invalid ones.

The proof that we will sketch depends on the fact that we initially defined a sentence of TFL using a recursive definition (see p. 50). We could have also used recursive definitions to define a proper proof in TFL and a proper truth table. (Although we didn't.) If we had these definitions, we could then use a *recursive proof* to show the soundness of TFL. A recursive proof works the same way as a recursive definition. With the recursive definition, we identified a group of base elements that were stipulated to be examples of the thing we were trying to define. In the case of a TFL sentence, the base class was the set of sentence letters A, B, C, \dots . We just announced that these were sentences. The second step of a recursive definition is to say that anything that is built up from your base class using certain rules also counts as an example of the thing you are defining. In the case of a definition of a sentence, the rules corresponded to the five sentential connectives (see p. 50). Once you have established a recursive definition, you can use that definition to show that all the members of the class you have defined have a certain property. You simply prove that the property is true of the members of the base class, and then you prove that the rules for extending the base class don't change the property. This is what it means to give a recursive proof.

Even though we don't have a recursive definition of a proof in TFL, we can sketch how a recursive proof of the soundness of TFL would go. Imagine a base class of one-line proofs, one for each of our basic rules of inference. The members of this class would look like this $X, Y \vdash X \& Y$; $X \& Y \vdash X$; $X \vee Y, \neg X \vdash Y \dots$ etc. Since some rules have a couple different forms, we would have to have add some members to this base class, for instance $X \& Y \vdash Y$. Notice that these are all statements in the metalanguage. The proof that TFL is sound is not a part of TFL, because TFL does not have the power to talk about itself.

You can use truth tables to prove to yourself that each of these one-line proofs in this base class is valid $_{\models}$. For instance the proof $X, Y \vdash X \& Y$ corresponds to a truth table that shows $X, Y \models X \& Y$. This

establishes the first part of our recursive proof.

The next step is to show that adding lines to any proof will never change a valid_F proof into an invalid_F one. We would need to do this for each of our basic rules of inference. So, for instance, for &I we need to show that for any proof $X_1, \dots, X_n \vdash Y$ adding a line where we use &I to infer $Z \& V$, where $Z \& V$ can be legitimately inferred from X_1, \dots, X_n, Y , would not change a valid proof into an invalid proof. But wait, if we can legitimately derive $Z \& V$ from these premises, then Z and V must be already available in the proof. They are either already among X_1, \dots, X_n, B , or can be legitimately derived from them. As such, any truth table line in which the premises are true must be a truth table line in which Z and V are true. According to the characteristic truth table for $\&$, this means that $Z \& V$ is also true on that line. Therefore, $Z \& V$ validly follows from the premises. This means that using the &E rule to extend a valid proof produces another valid proof.

In order to show that the proof system is sound, we would need to show this for the other inference rules. Since the derived rules are consequences of the basic rules, it would suffice to provide similar arguments for the other basic rules. This tedious exercise falls beyond the scope of this book.

So we have shown that $X \vdash Y$ implies $X \models Y$. What about the other direction, that is why think that *every* argument that can be shown valid using truth tables can also be proven using a derivation.

Completeness

This is the problem of completeness. A proof system has the property of **COMPLETENESS** if and only if there is a derivation of every semantically valid argument. Proving that a system is complete is generally harder than proving that it is sound. Proving that a system is sound amounts to showing that all of the rules of your proof system work the way they are supposed to. Showing that a system is complete means showing that you have included *all* the rules you need, that you haven't left any out. Showing this is beyond the scope of this book. The important point is that, happily, the proof system for TFL is both sound and complete. This is not the case for all proof systems or all formal languages. Because it is true of TFL, we can choose to give proofs or give truth tables—whichever is easier for the task at hand.

24.4 Other Semantic and Proof Theoretic Notions

Now we know that the proof theoretic and truth-table methods are equivalent, we can use them interchangeably depending on which is more useful.

We can also use proof theoretic methods for determining other logical notions, such as being consistent. We summarise how one would define them in 24.1

In fact, we can give general guidelines about when it's best to give proofs and when it is best to give truth tables. We do this in 24.2:

Practice exercises

A. Use either a derivation or a truth table for each of the following.

1. Show that $A \rightarrow [(B \& C) \vee D] \rightarrow A$ is a tautology.
2. Show that $A \rightarrow (A \rightarrow B)$ is not a tautology
3. Show that the sentence $A \rightarrow \neg A$ is not a contradiction.
4. Show that the sentence $A \leftrightarrow \neg A$ is a contradiction.
5. Show that the sentence $\neg(W \rightarrow (J \vee J))$ is contingent
6. Show that the sentence $\neg(X \vee (Y \vee Z)) \vee (X \vee (Y \vee Z))$ is not contingent
7. Show that the sentence $B \rightarrow \neg S$ is equivalent to the sentence $\neg\neg B \rightarrow \neg S$
8. Show that the sentence $\neg(X \vee O)$ is not equivalent to the sentence $X \& O$
9. Show that the sentences $\neg(A \vee B)$, C , $C \rightarrow A$ are jointly inconsistent.
10. Show that the sentences $\neg(A \vee B)$, $\neg B$, $B \rightarrow A$ are jointly consistent
11. Show that $\neg(A \vee (B \vee C)) \therefore \neg C$ is valid.
12. Show that $\neg(A \& (B \vee C)) \therefore \neg C$ is invalid.

B. Use either a derivation or a truth table for each of the following.

Concept	Truth table (semantic) definition	Proof-theoretic (syntactic) definition
Tautology	A sentence whose truth table only has Ts under the main connective	A sentence that can be derived without any premises.
Contradiction	A sentence whose truth table only has Fs under the main connective	A sentence whose negation can be derived without any premises
Contingent sentence	A sentence whose truth table contains both Ts and Fs under the main connective	A sentence that is not a theorem or contradiction
Equivalent sentences	The columns under the main connectives are identical.	The sentences can be derived from each other
Inconsistent sentences	Sentences which do not have a single line in their truth table where they are all true.	Sentences from which one can derive a contradiction
Consistent sentences	Sentences which have at least one line in their truth table where they are all true.	Sentences which are not inconsistent
Valid argument	An argument whose truth table has no lines where there are all Ts under main connectives for the premises and an F under the main connective for the conclusion.	An argument where one can derive the conclusion from the premises

Table 24.1: Two ways to define logical concepts.

Logical property	To prove it present	To prove it absent
Being a tautology	Derive the sentence	Find the false line in the truth table for the sentence
Being a contradiction	Derive the negation of the sentence	Find the true line in the truth table for the sentence
Contingency	Find a false line and a true line in the truth table for the sentence	Prove the sentence or its negation
Equivalence	Derive each sentence from the other	Find a line in the truth tables for the sentence where they have different values
Consistency	Find a line in truth table for the sentence where they all are true	Derive a contradiction from the sentences
Validity	Derive the conclusion from the premises	Find no line in the truth table where the premises are true and the conclusion false.

Table 24.2: When to provide a truth table and when to provide a proof.

1. Show that $A \rightarrow (B \rightarrow A)$ is a tautology
2. Show that $\neg(((N \leftrightarrow Q) \vee Q) \vee N)$ is not a tautology
3. Show that $Z \vee (\neg Z \leftrightarrow Z)$ is contingent
4. show that $(L \leftrightarrow ((N \rightarrow N) \rightarrow L)) \vee H$ is not contingent
5. Show that $(A \leftrightarrow A) \& (B \& \neg B)$ is a contradiction
6. Show that $(B \leftrightarrow (C \vee B))$ is not a contradiction.
7. Show that $((\neg X \leftrightarrow X) \vee X)$ is equivalent to X
8. Show that $F \& (K \& R)$ is not equivalent to $(F \leftrightarrow (K \leftrightarrow R))$

9. Show that the sentences $\neg(W \rightarrow W), (W \leftrightarrow W) \& W, E \vee (W \rightarrow \neg(E \& W))$ are inconsistent.
10. Show that the sentences $\neg R \vee C, (C \& R) \rightarrow \neg R, (\neg(R \vee R) \rightarrow R)$ are consistent.
11. Show that $\neg\neg(C \leftrightarrow \neg C), ((G \vee C) \vee G) \therefore ((G \rightarrow C) \& G)$ is valid.
12. Show that $\neg\neg L, (C \rightarrow \neg L) \rightarrow C \therefore \neg C$ is invalid.

CHAPTER 25

Proof-theoretic concepts

(Non-examinable)

Armed with our ‘ \vdash ’ symbol, we can introduce some new terminology.

X is a **THEOREM** iff $\vdash X$

To illustrate this, suppose we want to prove that ‘ $\neg(A \ \& \ \neg A)$ ’ is a theorem. So we must start our proof without *any* assumptions. However, since we want to prove a sentence whose main logical operator is a negation, we will want to immediately begin a subproof, with the additional assumption ‘ $A \ \& \ \neg A$ ’, and show that this leads to contradiction. All told, then, the proof looks like this:

1		$A \ \& \ \neg A$	
2		A	&E 1
3		$\neg A$	&E 1
4		\perp	\perp I 2, 3
5		$(A \ \& \ \neg A) \rightarrow \perp$	\rightarrow I 1–4
6		$\neg(A \ \& \ \neg A)$	\neg I 5

We have therefore proved ‘ $\neg(A \ \& \ \neg A)$ ’ on no (undischarged) assumptions. This particular theorem is an instance of what is sometimes called *the Law of Non-Contradiction*.

To show that something is a theorem, you just have to find a suitable proof. It is typically much harder to show that something is *not* a theorem. To do this, you would have to demonstrate, not just that certain proof strategies fail, but that *no* proof is possible. Even if you fail in trying to prove a sentence in a thousand different ways, perhaps the proof is just too long and complex for you to make out. Perhaps you just didn’t try hard enough.

Here is another new bit of terminology:

Two sentences X and Y are **PROVABLY EQUIVALENT** iff each can be proved from the other; i.e., both $X \vdash Y$ and $Y \vdash X$.

As in the case of showing that a sentence is a theorem, it is relatively easy to show that two sentences are provably equivalent: it just requires a pair of proofs. Showing that sentences are *not* provably equivalent would be much harder: it is just as hard as showing that a sentence is not a theorem.

Here is a third, related, bit of terminology:

The sentences X_1, X_2, \dots, X_n are **PROVABLY INCONSISTENT** iff a contradiction can be proved from them, i.e. $X_1, X_2, \dots, X_n \vdash \perp$. If they are not **INCONSISTENT**, we call them **PROVABLY CONSISTENT**.

It is easy to show that some sentences are provably inconsistent: you just need to prove a contradiction from assuming all the sentences. Showing that some sentences are not provably inconsistent is much harder. It would require more than just providing a proof or two; it would require showing that no proof of a certain kind is *possible*.

This table summarises whether one or two proofs suffice, or whether we must reason about all possible proofs.

	Yes	No
theorem?	one proof	all possible proofs
inconsistent?	one proof	all possible proofs
equivalent?	two proofs	all possible proofs
consistent?	all possible proofs	one proof

Practice exercises

A. Show that each of the following sentences is a theorem:

1. $O \rightarrow O$
2. $N \vee \neg N$
3. $J \leftrightarrow [J \vee (L \& \neg L)]$
4. $((A \rightarrow B) \rightarrow A) \rightarrow A$

B. Provide proofs to show each of the following:

1. $C \rightarrow (E \& G), \neg C \rightarrow G \vdash G$
2. $M \& (\neg N \rightarrow \neg M) \vdash (N \& M) \vee \neg M$
3. $(Z \& K) \leftrightarrow (Y \& M), D \& (D \rightarrow M) \vdash Y \rightarrow Z$
4. $(W \vee X) \vee (Y \vee Z), X \rightarrow Y, \neg Z \vdash W \vee Y$

C. Show that each of the following pairs of sentences are provably equivalent:

1. $R \leftrightarrow E, E \leftrightarrow R$
2. $G, \neg\neg\neg\neg G$
3. $T \rightarrow S, \neg S \rightarrow \neg T$
4. $U \rightarrow I, \neg(U \& \neg I)$
5. $\neg(C \rightarrow D), C \& \neg D$
6. $\neg G \leftrightarrow H, \neg(G \leftrightarrow H)$

D. If you know that $X \vdash Y$, what can you say about $(X \& Z) \vdash Y$? What about $(X \vee Z) \vdash Y$? Explain your answers.

E. In this chapter, we claimed that it is just as hard to show that two sentences are not provably equivalent, as it is to show that a sentence is not a theorem. Why did we claim this? (*Hint:* think of a sentence that would be a theorem iff X and Y were provably equivalent.)

PART V

Fallacies of Reasoning

CHAPTER 26

Logical fallacies

So far, we've been looking at arguments given in English, and we've been symbolising them and applying truth table tests to discover whether or not they are valid. But among the invalid ones, we haven't been distinguishing much. We said in Chapter 2 that there are various invalid arguments that are nonetheless good arguments. For instance, there are inductive arguments: All Fs we've seen so far are Gs; therefore, all Fs are Gs. There are statistical arguments: $x\%$ of Fs we've seen so far are Gs; therefore, $x\%$ of all Fs are Gs. And there are abductive arguments: of all available hypotheses, H best explains our evidence; therefore, H is true. In this chapter, we pay a little more attention to the bad invalid arguments. Why, you might ask? Well, it's good to know your enemy. And bad invalid arguments come in many shapes and sizes, and it's helpful to be able to characterise them. This makes it easier for you to spot them in the wild. When you feel uneasy about an argument, and you show that it is not valid, you might wish to ask whether it belongs to one of the standard sorts of bad invalid argument. If it does, that will help you to combat it.

26.1 Affirming the consequent

We know that the following form of argument is valid:

- If X , then Y

- X
- Therefore: Y .

This form is called *modus ponens*, or as we used it in our natural deduction system, $\rightarrow E$. But here's a form of argument that you sometimes see, and it's not valid. It's called *affirming the consequent*.

- If X , then Y
- Y
- Therefore: X .

Here's an example:

- If the universe were created by a benevolent God, people would behave altruistically, at least some of the time.
- People do behave altruistically, at least some of the time.
- Therefore: The universe were created by a benevolent God.

This is invalid because it's possible for the premises to be true and the conclusion false. The reason is that the first premise tells us one possible reason why people would behave altruistically, namely, that they are part of a universe created by a benevolent God. But it doesn't rule out that there are other possible reasons. It might be that altruistic behaviour has evolved by natural selection because altruistic groups are better able to cooperate and thus increase their survival chances and their evolutionary fitness.

26.2 Quantifier shift fallacy

Here is an argument that is sometimes ascribed to Aristotle (though it isn't at all clear this is what Aristotle actually meant):

- Every agent acts for an end.
- Therefore: Therefore, there is some (one) end for which every agent acts.

And here's an argument from St Thomas Aquinas's Third Way (the third among his five ways to argue for the existence of God):

- Everything at some time fails to exist.
- Therefore: There must be some (one) time at which everything fails to exist.

In both cases, the same thing goes wrong. The first premise says that, for every thing, there is some other thing related to the first thing in a particular way. The conclusion says that there is some one thing that is related to everything in that way. So Aristotle is said to argue that, since for every agent there is some end for which that agent acts, there must be a single end for which all agents act. And Aquinas is said to argue that, since for every thing, there is some time at which it does not exist, there must be some time at which everything does not exist. In both cases, they are committing a quantifier shift fallacy. It's perfectly possible that each agent has a different end for which they act. That would make Aristotle's premise true and his conclusion false. It's perfectly possible that each thing exists for four years, say, and at any time, some thing exists. That would make Aquinas' premise true and his conclusion false.

We'll return to this topic in §??.

26.3 Ad hominem fallacy

The name comes from Latin, where 'ad hominem' means 'to the person'. This form of argument concludes against a claim by attacking the person or group of people who put forward the claim, even when the attack is quite irrelevant to the truth of the claim in question. It is very obviously fallacious, but that doesn't stop it from being rather widespread in philosophy! Here's a rather unsubtle example:

- The original proponent of the theory of natural selection was Charles Darwin.
- Charles Darwin ate meat.
- Eating meat is wrong.
- Therefore: The theory of natural selection is wrong.

This is unsubtle because Darwin's morality is surely irrelevant to the correctness of his theory of evolution. But there are other cases where things aren't quite so clear cut. For instance, suppose you criticise the actions of your government. You then learn that a virulently racist hate group has also criticised the government for those same actions. Is it a good objection to your criticism to point out that evil people hold the same view? The American philosopher Joshua Blanchard calls this *the problem of unwelcome epistemic company*.

It is surely not a good argument just to point out that someone evil agrees with you. After all, presumably all evil people in history

agree with me that the sky is blue! But perhaps the argument has more force if the conclusion on which they agree with me is a moral one? Still, however, it doesn't seem to be a good argument. Here is Noam Chomsky explaining why:

“[I]t doesn't bother me if I happen to agree with the mainstream media [in criticizing the Soviet Union]. Trotsky [...] was charged in the 1930s with agreeing with the fascists in his condemnation of the Soviet Union. And he pointed out that his critique was true; he wasn't going to abandon it if somebody else happened to say it for different reasons.”

(Chomsky, Q&A after ‘Manufacturing Consent: The Political Economy of the Mass Media’, The University of Wisconsin—Madison, March 15th, 1989 (<https://chomsky.info/19890315/>))

And Chomsky is right. The point is that you might think something for one reason, while the evil person thinks it for another reason. For instance, you and an evil person might agree that there should be no death penalty for some particularly horrible crime. You might think it because you think there should be no death penalty for any crime at all; they might think it because they think that the horrible crime should not in fact be a crime at all.

Nonetheless, unwelcome epistemic company should give us pause. It should force us to think about what our reasons are for our conclusions, and what the evil person's reasons are, so that we can ensure our reasons are good.

26.4 Begging the question

Sometimes this is called ‘petitio principii’. Both are terrible names for it, but sadly we're stuck with them. The original name in the Greek means ‘assuming the conclusion’. This was then mistranslated into Latin as ‘petitio principii’, and this was mistranslated again into English as ‘begging the question’. Nowadays, people who haven't studied philosophy often use ‘beg the question’ to mean ‘raise the question’. Philosophers sometime get very upset about this, which is a bit silly, given the history of the term in philosophy. In any case, in the philosophers' sense, you beg the question when you argue for a conclusion, but you sneak that very conclusion itself into one of your premises. As a re-

sult, your argument actually ends up being valid, but it won't convince anyone who didn't already believe the conclusion.

This is a particular problem in philosophy, since often we are not giving arguments for a conclusion in order to convince people of that conclusion — often we all agree with the conclusion, but we're trying to figure out how you might justify it rather than simply take it for granted. For instance, suppose we try to answer the sceptic who says that the external world doesn't really exist (maybe it's Descartes talking about his evil demon in the *Meditations*). So we give an argument for the existence of the external world. The conclusion of our argument is that the external world exists. We don't expect that there's any real person who needs this argument to convince them that it exists. Rather, we give it to show that our core beliefs are justified and not just accepted on faith or because we're told they're true when we're young.

A good example of begging the question is G. E. Moore's famous argument against the sceptic. Moore held out his hands and said:

- Here is one hand.
- And here is another.
- Therefore: There are at least two external objects in the world.

But of course the sceptic won't accept the first or second premise here. Moore has assumed his conclusion in his premises.

Practice exercises

A. Give your own example of each of the following fallacies:

1. Affirming the consequent
2. Quantifier shift fallacy
3. Ad hominem fallacy
4. Begging the question

B. Identify which fallacy the following arguments commit:

1. John Locke thought that the criterion for personal identity is psychological continuity. But he wrote in support of slavery in *The Fundamental Constitutions of Carolina*. Therefore, personal identity cannot consist in psychological continuity.
2. Cigarettes are deadly, since cigarettes will kill you.

3. When it's about to rain, cows go under nearby trees. And indeed they are going under the trees. So it must be about to rain.
4. Everything that happens must have a cause, and so there must be some first thing that causes everything else.¹

¹This is sometimes (mis)attributed to the early modern British empiricist philosopher, John Locke.

CHAPTER 27

Probabilistic fallacies

Let's now move on to a couple of probabilistic fallacies. These are arguments that appeal to probabilities in some way, but they go badly wrong by misunderstanding how probabilities work. While you encounter this sort of reasoning less often in philosophy texts, they come up all the time in everyday life. Misuse of probabilities is very widespread and can lead to disastrous decision-making.

27.1 Base rate fallacy

This is perhaps the most famous probabilistic fallacy. It's famous partly because two social psychologists called Amos Tversky and Daniel Kahneman surveyed a group of students at Harvard Medical School in the 1970s, and a large proportion of them failed to identify it as a fallacy! Given that the argument, at least in the form presented to them and presented here, concerns the chance of having a disease, that is definitely cause for concern!

Our example of the base rate fallacy needs a bit of a backstory. Here goes:

Disease X A new test is developed for disease X. It has a sensitivity of 99% and a specificity of 99%. That means that, if you have the disease, your chance of testing positive is 99%, and if you don't have the disease your chance of

testing negative is 99%. The incidence of the disease in the population is 0.01%. That is, 1 in 10,000 people have it. You happen to be passing a testing station and you take a test. It comes back positive. What is the chance that you have disease X?

Many people (including lots of those Harvard medical students) answer 99%. A good deal more give an answer above 10%. The correct answer is around 1%! So anyone who argued for one of the higher answers has committed a fallacy of probabilistic reasoning.

Why is the answer around 1%? The easiest way to see why is to ask how these probabilities would play out in a population of 1,000,000 people. The incidence of the disease is 0.01% (i.e. 1 in 10,000), so 100 people in this population have the disease, while 999,900 do not. Now suppose that they are all tested. Of the 100 who have the disease, 99% will test positive and 1% will test negative. So 99 will test positive and 1 will test negative. Of the 999,900 who do not have the disease, 99% will test negative and 1% will test positive. So 989,901 will test negative, and 9,999 will test positive. So, $99 + 9,999 = 10,098$ will test positive in total. Of those, 9,999 will not have the disease, and 99 will have it. 99 is around 1% of 10,098. So, if you test positive, you've a 1% chance of being someone with the disease who got a true positive test, and 99% chance of being someone without the disease who got a false positive test.

It is called the base rate fallacy because people who commit it ignore how uncommon the disease is in the population—that's the base rate of the disease in the population. Because the disease is so uncommon, you're very unlikely to have it. And so if you get a positive test it's much more likely that it's a false positive and you don't have the disease than it is that it's a true positive and you do have the disease. This is one reason why doctors are so worried about the specificity and sensitivity of their tests.

27.2 Simpson's Paradox

This fallacy arises when people see a trend in two separate samples and infer that the same trend will occur when the samples are merged. Let's see what this means in a particular example. Suppose I have kidney stones. I go to my doctor and I ask what sort of treatment she'd recommend. She tells me that there are two available: Treatment 1 and Treatment 2. She also tells me that doctors divide kidney stones into two

types: small ones and large ones. She tells me that, when treating small kidney stones, Treatment 1 has been successful a greater proportion of the time it's been used than Treatment 2. And she also tells me that, when treating large kidney stones, Treatment 1 has been successful a greater proportion of the time it's been used than Treatment 2. 'OK,' I say, 'sounds like a no-brainer—I should go for Treatment 1, because it has been more successful a greater proportion of the time it's been used than Treatment 2.' Unfortunately, though, this reasoning is bad.

It's easy to see why it's tempting, but it's nonetheless bad probabilistic reasoning. It's tempting because it looks a lot like a valid form of reasoning in truth-functional logic:

- X or Y .
- If X , then Z .
- If Y , then Z .
- Therefore: Z .

This is basically what we used in the natural deduction rule $\vee E$. It is a valid argument form (check with truth-tables!). And my reasoning about the two treatments seems very close to this. I reason:

- For small kidney stones, Treatment 1 has been successful a greater proportion of the time it's been used than Treatment 2.
- For large kidney stones, Treatment 1 has been successful a greater proportion of the time it's been used than Treatment 2.
- Kidney stones are either small or large.
- Therefore: For kidney stones in general, Treatment 1 has been successful a greater proportion of the time it's been used than Treatment 2.

But this reasoning is bad. How do we show that? Well, remember that, to show that an argument is invalid in truth-functional logic, we describe a case in which the premises are true and the conclusion false. Here, we give a case in which the probabilities are as first and second premise say they are, and the third premise is true, but the probabilities are not as the conclusion says they are. Here's the case: suppose the following table gives the success rates of each treatment for each size of kidney stone.

	Treatment 1	Treatment 2
Small	9 out of 10 (90%)	80 out of 100 (80%)
Large	70 out of 100 (70%)	6 out of 10 (60%)
Total	79 out of 110 (72%)	86 out of 110 (78%)

So, Treatment 1 has a higher success rate than Treatment 2 in treating small kidney stones, and a higher success rate in treating large kidney stones; but it has a lower success rate in treating kidney stones! Such cases are known as instances of Simpson’s paradox. It’s paradoxical, because intuitively it doesn’t seem like these cases should be possible. But they are!

27.3 How do probabilities work?

We said above that the base rate fallacy and Simpson’s Paradox arise because people misunderstand how probabilities work. So how do they work? This is a huge question and the subject matter of all of statistics. But we can introduce the fundamental ideas here. This can end up looking a bit mathsy, but if you’re not terribly comfortable with maths, it’s still very worthwhile to work through. It’s very useful to be able to understand how probabilities work and to be able to identify when reasoning with them goes wrong. People sometimes say ‘You can prove anything with statistics’. That is really very false. But people do often make bad probabilistic or statistical arguments for their conclusions, and knowing how probabilities work really helps you to spot when they’re doing that.

Let’s start a little abstractly, and then we’ll move on to real examples. So suppose you have two atomic sentences, A and B . As we know, there are four possible valuations. That is, there are four ways to assign truth values to these atomic sentences. (Recall, in a truth table, we have a line for each valuation.)

A	B
T	T
T	F
F	T
F	F

Each valuation represents a possible “case”. The first row represents the worlds where A and B are both true, for instance. When we assign probabilities, we assign numbers to each valuation. And we make sure of two things:

- first, each number is either 0 or 1 or something in between them—e.g. 0.1, or 0.7, or 0.1987362;

- ▷ second, when we add up the numbers we assign to all the valuations, they add up to 1.

A **PROBABILITY ASSIGNMENT** assigns a number to every valuation ensuring:

- ▷ each number is either 0 or 1 or something in between them—e.g. 0.1, or 0.7, or 0.1987362;
- ▷ when we add up the numbers we assign to all the valuations, they add up to 1.

So, for instance, here are four probability assignments (p_1 , p_2 , p_3 , p_4). Notice that, in each case, the assignments add up to 1; that is, the numbers in each column add up to 1.

A	B	p_1	p_2	p_3	p_4
T	T	0.1	0.5	0.25	0.9
T	F	0.3	0.2	0.25	0
F	T	0.2	0.1	0.25	0
F	F	0.4	0.2	0.25	0.1

OK, so that gives our assignments of probabilities to the valuations. But now we want to know the probabilities of A , B , or any sentence built up from A and B using truth-functional connectives. How do we get that?

The probability of a sentence, X , is the sum of the probabilities assigned to the valuations where X is true.

To get the probability of a sentence X , we look at the probabilities assigned to each of the valuations that make it true, and we add them up. So for example, when we are looking at the probability assignment p_1 above, and are asking what is $p_1(A)$, we need to sum up the probability assignments of the valuations where A is true:

A	B	p_1	Is A true?
T	T	0.1	✓
T	F	0.3	✓
F	T	0.2	✗
F	F	0.4	✗

So

$$p_1(A) = 0.1 + 0.3 = 0.4$$

since A is true at the first two valuations, and p_1 gives them probability 0.1 and 0.3 respectively.

Similarly,

$$p_1(B) = 0.1 + 0.2 = 0.3.$$

What about the probability of $A \vee B$ and the probability of $A \& B$? Well, let's look at the truth tables:

A	B	$A \vee B$	p_1	p_2
T	T	T	0.1	0.6
T	F	T	0.3	0.2
F	T	T	0.2	0
F	F	F	0.4	0.3

$A \vee B$ is true on the first three valuations, and false on the final valuation. So

$$\begin{aligned} p_1(A \vee B) &= 0.1 + 0.3 + 0.2 = 0.6 \\ p_2(A \vee B) &= 0.6 + 0.2 + 0 = 0.8 \end{aligned}$$

Whereas $A \& B$ is true only on the first valuation, so:

$$\begin{aligned} p_1(A \& B) &= 0.1 \\ p_2(A \& B) &= 0.6 \end{aligned}$$

General properties

We can also see some general things about probabilities of sentences which are logically related:

Tautology If X is a tautology, then $p(X) = 1$.

A tautology is true in all valuations. So when calculating its probability, we add up the probabilities assigned to *all* valuations. And since we required that a probability assignment will always sum to 1, this will be 1.

Contradiction If X is a contradiction, then $p(X) = 0$.

The probability of X will be the sum of the probabilities assigned to each valuation where it is true. But there are no such valuations, so when we calculate the probability of X , there is nothing to sum up, and we just get 0.

Negation For any sentence X , $p(\neg X) = 1 - p(X)$.

That is, you get the probability of the negation of a sentence by subtracting the probability of the sentence from 1. Why? Well, the probability of X is the sum of the probabilities of the valuations at which X is true, while the probability of $\neg X$ is the sum of the probabilities of the valuations at which X is false. Since the probabilities assigned to all the valuations must add up to 1, we get $p(X) + p(\neg X) = 1$, and so $p(\neg X) = 1 - p(X)$.

Entailment For any sentences X and Y , if X entails Y , i.e., $X \therefore Y$ is valid, then $p(X) \leq p(Y)$.

If X entails Y , then any valuation where X is true, so is Y . So when we sum the probabilities assigned to the valuations where X is true, and compare it to the same for Y , we see that anything that contributed to the probability of X also contributed to the probability of Y . So the total value for Y is greater than or equal to that of X .

General Additivity For any sentences X and Y , $p(X \vee Y) = p(X) + p(Y) - p(X \& Y)$.

It's worth thinking this through. The probability of $X \vee Y$ is the sum of the probabilities of the valuations at which $X \vee Y$ is true; that is, the valuations at which X or Y or both are true. Now, first, add up the probabilities of the valuations at which X is true. That gives $p(X)$. Second, add up the probabilities of the valuations at which Y is true. That gives $p(Y)$. But at this point we've double counted those valuations at which X and Y are both true. We counted them once when we calculated $p(X)$ and we counted them again when we calculated $p(Y)$. So we need to subtract one lot of them. But of course the valuations at which X and Y are both true are exactly the valuations at which $X \& Y$ is true. So we subtract $p(X \& Y)$ from $p(X) + p(Y)$. So $p(X \vee Y) = p(X) + p(Y) - p(X \& Y)$.

Conditional Probabilities

So that gives us the probabilities of different sentences. There is one further probabilistic notion that is very important. It's the notion of a conditional probability. Because often we don't just ask how likely X is; we ask how likely X is *given that Y is true*. For instance, I ask how likely I am to have the disease *given I tested positive*. Or I ask how likely it is that we'll play rounders on the Downs *given that it is raining*. So, if we

have two sentences, X and Y , and a probability assignment p , we write $p(X|Y)$ for the probability of X given Y . Now, how do we calculate $p(X|Y)$? Well, here's the definition:

$$p(X|Y) = \frac{p(X \& Y)}{p(Y)}$$

provided $p(Y)$ is not 0.

Let's try to unpack that. Here's one way to see what's happening: the probability of X given Y is the proportion of the probability of Y that is assigned to valuations that make X true as well. The probability assigned to Y is obviously $p(Y)$. And the probability assigned to valuations that make X true as well as Y is just the probability of $X \& Y$, which is $p(X \& Y)$. So the proportion of the probability of Y that is assigned to valuations that make X true as well is just $\frac{p(X \& Y)}{p(Y)}$, which is how the definition defines $p(X|Y)$.

So let's see it in action. Sticking with probability assignment p_1 , what's the probability of A given B ?

$$p_1(A|B) = \frac{p_1(A \& B)}{p_1(B)} = \frac{0.1}{0.3} = \frac{1}{3} = 0.333 \dots$$

The proportion of the probability of B that is assigned to valuations at which A is also true is one-third.

Finally, we can state what is perhaps the most important fact in the theory of probability. It's called Bayes' Theorem, and it was first stated by the Reverend Thomas Bayes, an eighteenth century Presbyterian clergyman who lived in Kent. It's beautifully simple, but incredibly powerful:¹

¹The proof is quite short, but don't worry if you don't follow it. Here it goes. By definition

$$p(A)p(B|A) = p(A \& B) \quad \text{and} \quad p(\neg A)(B|\neg A) = p(\neg A \& B)$$

So

$$\frac{p(A) \times p(B|A)}{(p(A) \times p(B|A)) + (p(\neg A) \times p(B|\neg A))} = \frac{p(A \& B)}{p(A \& B) + p(\neg A \& B)}$$

But

$$p(A \& B) + p(\neg A \& B) = p(B)$$

So

$$\frac{p(A \& B)}{p(A \& B) + p(\neg A \& B)} = \frac{p(A \& B)}{p(B)} = p(A|B)$$

Bayes' Theorem:

$$p(A|B) = \frac{p(A) \times p(B|A)}{(p(A) \times p(B|A)) + (p(\neg A) \times p(B|\neg A))}$$

So this allows you to calculate $p(A|B)$ if you know $p(A)$ (and therefore you also know $p(\neg A)$), $p(B|A)$, and $p(B|\neg A)$. And it turns out that often we do know these latter three values. In fact, let's see all of this in action in the base rate fallacy, where Bayes' Theorem plays a big role.

Let A stand for *I have the disease*. Let B stand for *I test positive*. Then, the way the example is set out, we're told some information about the probability assignment:

- ▷ We're told that the incidence of the disease in the population is 1 in 10,000. So $p(A) = 0.0001$.
- ▷ We're told that the sensitivity and specificity of the disease is 99%. So $p(B|A) = 0.99$ and $p(\neg B|\neg A) = 0.99$, so $p(B|\neg A) = 0.01$.

This gives us enough to use Bayes' Theorem to calculate the probability I have the disease given that I test positive, i.e., $p(A|B)$. According to Bayes' Theorem:

$$p(A|B) = \frac{p(A) \times p(B|A)}{(p(A) \times p(B|A)) + (p(\neg A) \times p(B|\neg A))} = \frac{0.0001 \times 0.99}{0.0001 \times 0.99 + 0.9999 \times 0.01} \approx 0.0098$$

And this is what we calculated above as well.

Practice exercises

A. Look at probability assignment p_2 above. Calculate the following probabilities.

1. $p_2(A)$
2. $p_2(B)$
3. $p_2(A \& B)$
4. $p_2(A \vee B)$

5. $p_2(B \rightarrow A)$

6. $p_2(A|B)$

7. $p_2(B|A)$

If you feel you still need more practice, try doing the same for p_3 and p_4 .

B. Suppose I give you the following probabilities:

$$p(A) = 0.3 \quad p(B|A) = 0.6 \quad p(B|\neg A) = 0.4$$

Use them and Bayes' Theorem to calculate $p(A|B)$.

C. Go back to the Base Rate Fallacy and show using Bayes' Theorem that the answer is 1%.

D. Explain why the following facts hold for any sentences X, Y .

1. $p(X) \leq p(X \vee Y)$

2. $p(X \& Y) \leq p(X)$

3. If there is no valuation where X and Y are both true, $p(X \vee Y) = p(X) + p(Y)$.

PART VI

First-order logic

CHAPTER 28

Building blocks of FOL

28.1 The need to decompose sentences

Consider the following argument, which is obviously valid in English:

1. Willard is a logician.
2. All logicians wear funny hats.
 - Therefore: Willard wears a funny hat.

To symbolize it in TFL, we might offer a symbolization key:

- L*: Willard is a logician.
A: All logicians wear funny hats.
F: Willard wears a funny hat.

And the argument itself becomes:

$$L, A \therefore F$$

This is *invalid* in TFL, but the original English argument is clearly valid.

The problem is not that we have made a mistake while symbolizing the argument. This is the best symbolization we can give *in TFL*. The problem lies with TFL itself. ‘All logicians wear funny hats’ is about both logicians and hat-wearing. By not retaining this structure in our symbolization, we lose the connection between Willard’s being a logician and Willard’s wearing a hat.

The basic units of TFL are atomic sentences, and TFL cannot decompose these. To symbolize arguments like the preceding one, we will have to develop a new logical language which will allow us to *split the atom*. We will call this language *first-order logic*, or *FOL*.

The details of FOL will be explained throughout this chapter, but here is the basic idea for splitting the atom.

First, we have *names*. In FOL, we indicate these with lowercase italic letters. For instance, we might let *b* stand for Bertie, or let *i* stand for Willard.

Second, we have predicates. English predicates are expressions like ‘_____ is a dog’ or ‘_____ is a logician’. These are not complete sentences by themselves. In order to make a complete sentence, we need to fill in the gap. We need to say something like ‘Bertie is a dog’ or ‘Willard is a logician’. In FOL, we indicate predicates with uppercase italic letters. For instance, we might let the FOL predicate *D* symbolize the English predicate ‘_____ is a dog’. Then the expression *Db* will be a sentence in FOL, which symbolizes the English sentence ‘Bertie is a dog’. Equally, we might let the FOL predicate *L* symbolize the English predicate ‘_____ is a logician’. Then the expression *Li* will symbolize the English sentence ‘Willard is a logician’.

Third, we have quantifiers. For instance, \exists will roughly convey ‘There is at least one ...’. So we might symbolize the English sentence ‘there is a dog’ with the FOL sentence $\exists xDx$, which we would naturally read out-loud as ‘there is at least one thing, *x*, such that *x* is a dog’.

That is the general idea, but FOL is significantly more subtle than TFL, so we will come at it slowly.

28.2 Names

In English, a *singular term* is a word or phrase that refers to a *specific* person, place, or thing. The word ‘dog’ is not a singular term, because there are a great many dogs. The phrase ‘Bertie’ is a singular term, because it refers to a specific terrier. Likewise, the phrase ‘Philip’s dog Bertie’ is a singular term, because it refers to a specific little terrier.

Proper names are a particularly important kind of singular term. These are expressions that pick out individuals without describing them. The name ‘Emerson’ is a proper name, and the name alone does not tell you anything about Emerson. Of course, some names are traditionally given to boys and other are traditionally given to girls. If

‘Hilary’ is used as a singular term, you might guess that it refers to a woman. You might, though, be guessing wrongly. Indeed, the name does not necessarily mean that the person referred to is even a person: Hilary might be a giraffe, for all you could tell just from the name.

In FOL, our **NAMES** are lower-case letters a through to r . We can add subscripts if we want to use some letter more than once. So here are some singular terms in FOL:

$$a, b, c, \dots, r, a_1, f_{32}, j_{390}, m_{12}$$

These should be thought of along the lines of proper names in English, but with one difference. ‘Tim Button’ is a proper name, but there are several people with this name. (Equally, there are at least two people with the name ‘P.D. Magnus’.) We live with this kind of ambiguity in English, allowing context to individuate the fact that ‘Tim Button’ refers to an author of this book, and not some other Tim. In FOL, we do not tolerate any such ambiguity. Each name must pick out *exactly* one thing. (However, two different names may pick out the same thing.)

As with TFL, we can provide symbolization keys. These indicate, temporarily, what a name will pick out. So we might offer:

e : Elsa

g : Gregor

m : Marybeth

28.3 Predicates

The simplest predicates are properties of individuals. They are things you can say about an object. Here are some examples of English predicates:

_____ is a dog

_____ is a member of Monty Python

A piano fell on _____

In general, you can think about predicates as things which combine with singular terms to make sentences. Conversely, you can start with sentences and make predicates out of them by removing terms. Consider the sentence, ‘Vinnie borrowed the family car from Nunzio.’ By removing a singular term, we can obtain any of three different predicates:

_____ borrowed the family car from Nunzio
Vinnie borrowed _____ from Nunzio
Vinnie borrowed the family car from _____

In FOL, **PREDICATES** are capital letters A through Z , with or without subscripts. We might write a symbolization key for predicates thus:

Ax : _____ _{x} is angry
 Hx : _____ _{x} is happy

(Why the subscripts on the gaps? We will return to this in the next section.)

If we combine our two symbolization keys, we can start to symbolize some English sentences that use these names and predicates in combination. For example, consider the English sentences:

1. Elsa is angry.
2. Gregor and Marybeth are angry.
3. If Elsa is angry, then so are Gregor and Marybeth.

Sentence **1** is straightforward: we symbolize it by Ae .

Sentence **2**: this is a conjunction of two simpler sentences. The simple sentences can be symbolized just by Ag and Am . Then we help ourselves to our resources from TFL, and symbolize the entire sentence by $Ag \& Am$. This illustrates an important point: FOL has all of the truth-functional connectives of TFL.

Sentence **3**: this is a conditional, whose antecedent is sentence **1** and whose consequent is sentence **2**, so we can symbolize this with $Ae \rightarrow (Ag \& Am)$.

Sometimes the truth functional nature of a sentence isn't explicit. For example suppose we wish to symbolise:

4. Betty is a red car.

This sentence can be paraphrased as 'Betty is red and Betty is a car'. And we can symbolise this as $Rb \& Cb$ using the symbolisation key

b : Betty
 Rx : _____ _{x} is red
 Cx : _____ _{x} is a car

28.4 Many-placed predicates

All of the predicates that we have considered so far concern properties that objects might have. Those predicates have one gap in them, and to make a sentence, we simply need to slot in one term. They are **ONE-PLACE** predicates.

However, other predicates concern the *relation* between two things. Here are some examples of relational predicates in English:

____ loves ____
____ is to the left of ____
____ is in debt to ____

These are **TWO-PLACE** predicates. They need to be filled in with two terms in order to make a sentence. Conversely, if we start with an English sentence containing many singular terms, we can remove two singular terms, to obtain different two-place predicates. Consider the sentence ‘Vinnie borrowed the family car from Nunzio’. By deleting two singular terms, we can obtain any of three different two-place predicates

Vinnie borrowed ____ from ____
____ borrowed the family car from ____
____ borrowed ____ from Nunzio

and by removing all three singular terms, we obtain a **THREE-PLACE** predicate:

____ borrowed ____ from ____

Indeed, there is no in principle upper limit on the number of places that our predicates may contain.

Now there is a little foible with the above. We have used the same symbol, ‘____’, to indicate a gap formed by deleting a term from a sentence. However (as Frege emphasized), these are *different* gaps. To obtain a sentence, we can fill them in with the same term, but we can equally fill them in with different terms, and in various different orders. The following are all perfectly good sentences, and they all mean very different things:

Karl loves Karl
Karl loves Imre
Imre loves Karl
Imre loves Imre

The point is that we need to keep track of the gaps in predicates, so that we can keep track of how we are filling them in.

To keep track of the gaps, we will label them. The labelling conventions we will adopt are best explained by example. Suppose we want to symbolize the following sentences:

5. Karl loves Imre.
6. Imre loves himself.
7. Karl loves Imre, but not vice versa.
8. Karl is loved by Imre.

We will start with the following symbolisation key:

domain: people

i: Imre

k: Karl

Lxy: ______x loves ______y

Sentence 5 will now be symbolized by *Lki*.

Sentence 6 can be paraphrased as ‘Imre loves Imre’. It can now be symbolized by *Lii*.

Sentence 7 is a conjunction. We might paraphrase it as ‘Karl loves Imre, and Imre does not love Karl’. It can now be symbolized by *Lki* & $\neg Lik$.

Sentence 8 might be paraphrased by ‘Imre loves Karl’. It can then be symbolized by *Lik*. Of course, this slurs over the difference in tone between the active and passive voice; such nuances are lost in FOL.

This last example, though, highlights something important. Suppose we add to our symbolization key the following:

Mxy: ______y loves ______x

Here, we have used the same English word (‘loves’) as we used in our symbolization key for *Lxy*. However, we have swapped the order of the *gaps* around (just look closely at those little subscripts!) So *Mki* and *Lik* now *both* symbolize ‘Imre loves Karl’. *Mik* and *Lki* now *both* symbolize ‘Karl loves Imre’. Since love can be unrequited, these are very different claims.

The moral is simple. When we are dealing with predicates with more than one place, we need to pay careful attention to the order of the places.

28.5 Quantifiers

We are now ready to introduce quantifiers. Consider these sentences:

9. Everyone is happy.
10. Someone is angry.

It might be tempting to symbolize sentence 9 as $He \& Hg \& Hm$. Yet this would only say that Elsa, Gregor, and Marybeth are happy. We want to say that *everyone* is happy, even those with no names. In order to do this, we introduce the \forall symbol. This is called the **UNIVERSAL QUANTIFIER**.

A quantifier must always be followed by a **VARIABLE**. In FOL, variables are italic lowercase letters s through z , with or without subscripts. So we might symbolize sentence 9 as $\forall x Hx$. The variable x is serving as a kind of placeholder. The expression $\forall x$ intuitively means that you can pick anyone and put them in as x . The subsequent Hx indicates, of that thing you picked out, that it is happy.

It should be pointed out that there is no special reason to use x rather than some other variable. The sentences $\forall x Hx$, $\forall y Hy$, $\forall z Hz$, and $\forall x_5 Hx_5$ use different variables, but they will all be logically equivalent.

To symbolize sentence 10, we introduce another new symbol: the **EXISTENTIAL QUANTIFIER**, \exists . Like the universal quantifier, the existential quantifier requires a variable. Sentence 10 can be symbolized by $\exists x Ax$. Whereas $\forall x Ax$ is read naturally as ‘for all x , x is angry’, $\exists x Ax$ is read naturally as ‘there is something, x , such that x is angry’. Once again, the variable is a kind of placeholder; we could just as easily have symbolized sentence 10 with $\exists z Az$, $\exists w_{256} Aw_{256}$, or whatever.

Some more examples will help. Consider these further sentences:

11. No one is angry.
12. There is someone who is not happy.
13. Not everyone is happy.

Sentence 11 can be paraphrased as, ‘It is not the case that someone is angry’. We can then symbolize it using negation and an existential quantifier: $\neg \exists x Ax$. Yet sentence 11 could also be paraphrased as, ‘Everyone is not angry’. With this in mind, it can be symbolized using negation and a universal quantifier: $\forall x \neg Ax$. Both of these are acceptable symbolizations. Indeed, it will transpire that, in general, $\forall x \neg X$ is logically equivalent to $\neg \exists x X$. (Notice that we have here returned

to the practice of using X as a metavariable, from §9.) Symbolizing a sentence one way, rather than the other, might seem more ‘natural’ in some contexts, but it is not much more than a matter of taste.

Sentence 12 is most naturally paraphrased as, ‘There is some x , such that x is not happy’. This then becomes $\exists x \neg Hx$. Of course, we could equally have written $\neg \forall x Hx$, which we would naturally read as ‘it is not the case that everyone is happy’. That too would be a perfectly adequate symbolization of sentence 13.

28.6 Domains

Given the symbolization key we have been using, $\forall x Hx$ symbolizes ‘Everyone is happy’. Who is included in this *everyone*? When we use sentences like this in English, we usually do not mean everyone now alive on the Earth. We certainly do not mean everyone who was ever alive or who will ever live. We usually mean something more modest: everyone now in the building, everyone enrolled in the ballet class, or whatever.

In order to eliminate this ambiguity, we will need to specify a **DOMAIN**. The domain is the collection of things that we are talking about. So if we want to talk about people in Chicago, we define the domain to be people in Chicago. We write this at the beginning of the symbolization key, like this:

domain: people in Chicago

The quantifiers *range over* the domain. Given this domain, $\forall x$ is to be read roughly as ‘Every person in Chicago is such that...’ and $\exists x$ is to be read roughly as ‘Some person in Chicago is such that...’.

In FOL, the domain must always include at least one thing. Moreover, in English we can infer ‘something is angry’ from ‘Gregor is angry’. In FOL, then, we will want to be able to infer $\exists x Ax$ from A_g . So we will insist that each name must pick out exactly one thing in the domain. If we want to name people in places beside Chicago, then we need to include those people in the domain.

A domain must have *at least* one member. A name must pick out *exactly* one member of the domain, but a member of the domain may be picked out by one name, many names, or none at all.

Even allowing for a domain with just one member can produce some strange results. Suppose we have this as a symbolization key:

domain: the Eiffel Tower

Px : ______x is in Paris.

The sentence $\forall xPx$ might be paraphrased in English as ‘Everything is in Paris.’ Yet that would be misleading. It means that everything *in the domain* is in Paris. This domain contains only the Eiffel Tower, so with this symbolization key $\forall xPx$ just means that the Eiffel Tower is in Paris.

Non-referring terms

In FOL, each name must pick out exactly one member of the domain. A name cannot refer to more than one thing—it is a *singular* term. Each name must still pick out *something*. This is connected to a classic philosophical problem: the so-called problem of non-referring terms.

Medieval philosophers typically used sentences about the *chimera* to exemplify this problem. Chimera is a mythological creature; it does not really exist. Consider these two sentences:

14. Chimera is angry.
15. Chimera is not angry.

It is tempting just to define a name to mean ‘chimera.’ The symbolization key would look like this:

domain: creatures on Earth

Ax : ______x is angry.

c : chimera

We could then symbolize sentence 14 as Ac and sentence 15 as $\neg Ac$.

Problems will arise when we ask whether these sentences are true or false.

One option is to say that sentence 14 is not true, because there is no chimera. If sentence 14 is false because it talks about a non-existent thing, then sentence 15 is false for the same reason. Yet this would mean that Ac and $\neg Ac$ would both be false. Given the truth conditions for negation, this cannot be the case.

Since we cannot say that they are both false, what should we do? Another option is to say that sentence 14 is *meaningless* because it talks about a non-existent thing. So Ac would be a meaningful expression in FOL for some interpretations but not for others. Yet this would make

our formal language hostage to particular interpretations. Since we are interested in logical form, we want to consider the logical force of a sentence like *Ac* apart from any particular interpretation. If *Ac* were sometimes meaningful and sometimes meaningless, we could not do that.

This is the *problem of non-referring terms*, and we will return to it later (see p. 299.) The important point for now is that each name of FOL *must* refer to something in the domain, although the domain can contain any things we like. If we want to symbolize arguments about mythological creatures, then we must define a domain that includes them. This option is important if we want to consider the logic of stories. We can symbolize a sentence like ‘Sherlock Holmes lived at 221B Baker Street’ by including fictional characters like Sherlock Holmes in our domain.

CHAPTER 29

Sentences with one quantifier

We now have all of the pieces of FOL. Symbolizing more complicated sentences will only be a matter of knowing the right way to combine predicates, names, quantifiers, and connectives. There is a knack to this, and there is no substitute for practice.

29.1 Common quantifier phrases

Consider these sentences:

1. Every coin in my pocket is a quarter.
2. Some coin on the table is a dime.
3. Not all the coins on the table are dimes.
4. None of the coins in my pocket are dimes.

In providing a symbolization key, we need to specify a domain. Since we are talking about coins in my pocket and on the table, the domain must at least contain all of those coins. Since we are not talking about anything besides coins, we let the domain be all coins. Since we are not talking about any specific coins, we do not need to deal with any names. So here is our key:

domain: all coins

Px : ______x is in my pocket

Tx : ______x is on the table

Qx : _____ _{x} is a quarter

Dx : _____ _{x} is a dime

Sentence 1 is most naturally symbolized using a universal quantifier. The universal quantifier says something about everything in the domain, not just about the coins in my pocket. Sentence 1 can be paraphrased as ‘for any coin, *if* that coin is in my pocket *then* it is a quarter’. So we can symbolize it as $\forall x(Px \rightarrow Qx)$.

Since sentence 1 is about coins that are both in my pocket *and* that are quarters, it might be tempting to symbolize it using a conjunction. However, the sentence $\forall x(Px \& Qx)$ would symbolize the sentence ‘every coin is both a quarter and in my pocket’. This obviously means something very different than sentence 1. And so we see:

If a sentence can be paraphrased in English as

‘every F is G’,
‘all Fs are Gs’, or
‘any F is a G’,

it can be symbolised as can be symbolized as

$\forall x(Fx \rightarrow Gx)$.

Sentence 2 is most naturally symbolized using an existential quantifier. It can be paraphrased as ‘there is some coin which is both on the table and which is a dime’. So we can symbolize it as $\exists x(Tx \& Dx)$.

Notice that we needed to use a conditional with the universal quantifier, but we used a conjunction with the existential quantifier. Suppose we had instead written $\exists x(Tx \rightarrow Dx)$. That would mean that there is some object in the domain of which $(Tx \rightarrow Dx)$ is true. Recall that, in TFL, $X \rightarrow Y$ is logically equivalent (in TFL) to $\neg X \vee Y$. This equivalence will also hold in FOL. So $\exists x(Tx \rightarrow Dx)$ is true if there is some object in the domain, such that $(\neg Tx \vee Dx)$ is true of that object. That is, $\exists x(Tx \rightarrow Dx)$ is true if some coin is *either* not on the table *or* is a dime. Of course there is a coin that is not on the table: there are coins lots of other places. So it is *very easy* for $\exists x(Tx \rightarrow Dx)$ to be true. A conditional will usually be the natural connective to use with a universal quantifier, but a conditional within the scope of an existential quantifier tends to say something very weak indeed. As a general rule of thumb, do not put conditionals in the scope of existential quantifiers unless you are sure that you need one.

If a sentence can be paraphrased in English as

‘some F is G’,
‘there is some F that is G’,
‘some F is G’, or
‘there is at least one F that is a G’

it can be symbolised as can be symbolized as

$$\exists x(Fx \& Gx).$$

Sentence 3 can be paraphrased as, ‘It is not the case that every coin on the table is a dime’. So we can symbolize it by $\neg\forall x(Tx \rightarrow Dx)$. You might look at sentence 3 and paraphrase it instead as, ‘Some coin on the table is not a dime’. You would then symbolize it by $\exists x(Tx \& \neg Dx)$. Although it is probably not immediately obvious yet, these two sentences are logically equivalent. (This is due to the logical equivalence between $\neg\forall xX$ and $\exists x\neg X$, mentioned in §28, along with the equivalence between $\neg(X \rightarrow Y)$ and $X \& \neg Y$.)

If a sentence can be paraphrased in English as

‘not all Fs are Gs’,

it can be symbolised as can be symbolized as

$$\neg\forall x(Fx \rightarrow Gx), \text{ or } \\ \exists x(Fx \& \neg Gx).$$

Sentence 4 can be paraphrased as, ‘It is not the case that there is some dime in my pocket’. This can be symbolized by $\neg\exists x(Px \& Dx)$. It might also be paraphrased as, ‘Everything in my pocket is a non-dime’, and then could be symbolized by $\forall x(Px \rightarrow \neg Dx)$. Again the two symbolizations are logically equivalent; both are correct symbolizations of sentence 4.

If a sentence can be paraphrased in English as

‘no Fs are Gs’,

it can be symbolised as can be symbolized as

$$\neg \exists x(Fx \ \& \ Gx), \text{ or} \\ \forall x(Fx \rightarrow \neg Gx).$$

29.2 Empty predicates

In §28, we emphasized that a name must pick out exactly one object in the domain. However, a predicate need not apply to anything in the domain. A predicate that applies to nothing in the domain is called an **EMPTY PREDICATE**. This is worth exploring.

Suppose we want to symbolize these two sentences:

5. Every monkey knows sign language
6. Some monkey knows sign language

It is possible to write the symbolization key for these sentences in this way:

domain: animals

Mx : _____ _{x} is a monkey.

Sx : _____ _{x} knows sign language.

Sentence 5 can now be symbolized by $\forall x(Mx \rightarrow Sx)$. Sentence 6 can be symbolized as $\exists x(Mx \ \& \ Sx)$.

It is tempting to say that sentence 5 *entails* sentence 6. That is, we might think that it is impossible for it to be the case that every monkey knows sign language, without its also being the case that some monkey knows sign language, but this would be a mistake. It is possible for the sentence $\forall x(Mx \rightarrow Sx)$ to be true even though the sentence $\exists x(Mx \ \& \ Sx)$ is false.

How can this be? The answer comes from considering whether these sentences would be true or false *if there were no monkeys*. If there were no monkeys at all (in the domain), then $\forall x(Mx \rightarrow Sx)$ would be *vacuously* true: take any monkey you like—it knows sign language! But if there were no monkeys at all (in the domain), then $\exists x(Mx \ \& \ Sx)$ would be false.

Another example will help to bring this home. Suppose we extend the above symbolization key, by adding:

Rx : _____ _{x} is a refrigerator

Now consider the sentence $\forall x(Rx \rightarrow Mx)$. This symbolizes ‘every refrigerator is a monkey’. This sentence is true, given our symbolization key, which is counterintuitive, since we (presumably) do not want to say that there are a whole bunch of refrigerator monkeys. It is important to remember, though, that $\forall x(Rx \rightarrow Mx)$ is true iff any member of the domain that is a refrigerator is a monkey. Since the domain is *animals*, there are no refrigerators in the domain. Again, then, the sentence is *vacuously* true.

If you were actually dealing with the sentence ‘All refrigerators are monkeys’, then you would most likely want to include kitchen appliances in the domain. Then the predicate R would not be empty and the sentence $\forall x(Rx \rightarrow Mx)$ would be false.

When F is an empty predicate, a sentence $\forall x(Fx \rightarrow \dots)$ will be vacuously true.

29.3 Picking a domain

The appropriate symbolization of an English language sentence in FOL will depend on the symbolization key. Choosing a key can be difficult. Suppose we want to symbolize the English sentence:

7. Every rose has a thorn.

We might offer this symbolization key:

Rx : _____ _{x} is a rose

Tx : _____ _{x} has a thorn

It is tempting to say that sentence 7 should be symbolized as $\forall x(Rx \rightarrow Tx)$, but we have not yet chosen a domain. If the domain contains all roses, this would be a good symbolization. Yet if the domain is merely *things on my kitchen table*, then $\forall x(Rx \rightarrow Tx)$ would only come close to covering the fact that every rose *on my kitchen table* has a thorn. If there are no roses on my kitchen table, the sentence would be trivially true. This is not what we want. To symbolize sentence 7 adequately, we need to include all the roses in the domain, but now we have two options.

First, we can restrict the domain to include all roses but *only* roses. Then sentence 7 can, if we like, be symbolized with $\forall xTx$. This is true iff everything in the domain has a thorn; since the domain is just the roses, this is true iff every rose has a thorn. By restricting the domain, we have been able to symbolize our English sentence with a very short sentence of FOL. So this approach can save us trouble, if every sentence that we want to deal with is about roses.

Second, we can let the domain contain things besides roses: rhododendrons; rats; rifles; whatever., and we will certainly need to include a more expansive domain if we simultaneously want to symbolize sentences like:

8. Every cowboy sings a sad, sad song.

Our domain must now include both all the roses (so that we can symbolize sentence 7) and all the cowboys (so that we can symbolize sentence 8). So we might offer the following symbolization key:

domain: people and plants

Cx : ______x is a cowboy

Sx : ______x sings a sad, sad song

Rx : ______x is a rose

Tx : ______x has a thorn

Now we will have to symbolize sentence 7 with $\forall x(Rx \rightarrow Tx)$, since $\forall xTx$ would symbolize the sentence ‘every person or plant has a thorn’. Similarly, we will have to symbolize sentence 8 with $\forall x(Cx \rightarrow Sx)$.

In general, the universal quantifier can be used to symbolize the English expression ‘everyone’ if the domain only contains people. If there are people and other things in the domain, then ‘everyone’ must be treated as ‘every person’.

29.4 The utility of paraphrase

When symbolizing English sentences in FOL, it is important to understand the structure of the sentences you want to symbolize. What matters is the final symbolization in FOL, and sometimes you will be able to move from an English language sentence directly to a sentence of FOL. Other times, it helps to paraphrase the sentence one or more times. Each successive paraphrase should move from the original sentence closer to something that you can easily symbolize directly in FOL.

For the next several examples, we will use this symbolization key:

domain: people

Bx : ______x is a bassist.

Rx : ______x is a rock star.

k : Kim Deal

Now consider these sentences:

9. If Kim Deal is a bassist, then she is a rock star.
10. If a person is a bassist, then she is a rock star.

The same words appear as the consequent in sentences **9** and **10** ('... she is a rock star'), but they mean very different things. To make this clear, it often helps to paraphrase the original sentences, removing pronouns.

Sentence **9** can be paraphrased as, 'If Kim Deal is a bassist, then *Kim Deal* is a rockstar'. This can obviously be symbolized as $Bk \rightarrow Rk$.

Sentence **10** must be paraphrased differently: 'If a person is a bassist, then *that person* is a rock star'. This sentence is not about any particular person, so we need a variable. As an intermediate step, we can paraphrase this as, 'For any person x , if x is a bassist, then x is a rockstar'. Now this can be symbolized as $\forall x(Bx \rightarrow Rx)$. This is the same sentence we would have used to symbolize 'Everyone who is a bassist is a rock star'. On reflection, that is surely true iff sentence **10** is true, as we would hope.

Consider these further sentences:

11. If anyone is a bassist, then Kim Deal is a rock star.
12. If anyone is a bassist, then she is a rock star.

The same words appear as the antecedent in sentences **11** and **12** ('If anyone is a bassist...'), but it can be tricky to work out how to symbolize these two uses. Again, paraphrase will come to our aid.

Sentence **11** can be paraphrased, 'If there is at least one bassist, then Kim Deal is a rock star'. It is now clear that this is a conditional whose antecedent is a quantified expression; so we can symbolize the entire sentence with a conditional as the main logical operator: $\exists x Bx \rightarrow Rk$.

Sentence **12** can be paraphrased, 'For all people x , if x is a bassist, then x is a rock star'. Or, in more natural English, it can be paraphrased by 'All bassists are rock stars'. It is best symbolized as $\forall x(Bx \rightarrow Rx)$, just like sentence **10**.

The moral is that the English words 'any' and 'anyone' should typically be symbolized using quantifiers, and if you are having a hard time

determining whether to use an existential or a universal quantifier, try paraphrasing the sentence with an English sentence that uses words *besides* ‘any’ or ‘anyone’.

29.5 Quantifiers and scope

Continuing the example, suppose we want to symbolize these sentences:

13. If everyone is a bassist, then Lars is a bassist
14. Everyone is such that, if they are a bassist, then Lars is a bassist.

To symbolize these sentences, we will have to add a new name to the symbolization key, namely:

l : Lars

Sentence 13 is a conditional, whose antecedent is ‘everyone is a bassist’, so we will symbolize it with $\forall x Bx \rightarrow Bl$. This sentence is *necessarily* true: if *everyone* is indeed a bassist, then take any one you like—for example Lars—and he will be a bassist.

Sentence 14, by contrast, might best be paraphrased by ‘every person x is such that, if x is a bassist, then Lars is a bassist’. This is symbolized by $\forall x (Bx \rightarrow Bl)$. This sentence is false; Kim Deal is a bassist. So Bk is true, but Lars is not a bassist, so Bl is false. Accordingly, $Bk \rightarrow Bl$ will be false, so $\forall x (Bx \rightarrow Bl)$ will be false as well.

In short, $\forall x Bx \rightarrow Bl$ and $\forall x (Bx \rightarrow Bl)$ are very different sentences. We can explain the difference in terms of the *scope* of the quantifier. The scope of quantification is very much like the scope of negation, which we considered when discussing TFL, and it will help to explain it in this way.

In the sentence $\neg Bk \rightarrow Bl$, the scope of \neg is just the antecedent of the conditional. We are saying something like: if Bk is false, then Bl is true. Similarly, in the sentence $\forall x Bx \rightarrow Bl$, the scope of $\forall x$ is just the antecedent of the conditional. We are saying something like: if Bx is true of *everything*, then Bl is also true.

In the sentence $\neg (Bk \rightarrow Bl)$, the scope of \neg is the entire sentence. We are saying something like: $(Bk \rightarrow Bl)$ is false. Similarly, in the sentence $\forall x (Bx \rightarrow Bl)$, the scope of $\forall x$ is the entire sentence. We are saying something like: $(Bx \rightarrow Bl)$ is true of *everything*.

The moral of the story is simple. When you are using conditionals, be very careful to make sure that you have sorted out the scope correctly.

29.6 Ambiguous predicates

Suppose we just want to symbolize this sentence:

15. Adina is a skilled surgeon.

Let the domain be people, let Kx mean ‘ x is a skilled surgeon’, and let a mean Adina. Sentence 15 is simply Ka .

Suppose instead that we want to symbolize this argument:

The hospital will only hire a skilled surgeon. All surgeons are greedy. Billy is a surgeon, but is not skilled. Therefore, Billy is greedy, but the hospital will not hire him.

We need to distinguish being a *skilled surgeon* from merely being a *surgeon*. So we define this symbolization key:

domain: people

Gx : _____ x is greedy.

Hx : The hospital will hire _____ x .

Rx : _____ x is a surgeon.

Kx : _____ x is skilled.

b : Billy

Now the argument can be symbolized in this way:

- $\forall x[\neg(Rx \ \& \ Kx) \rightarrow \neg Hx]$
- $\forall x(Rx \rightarrow Gx)$
- $Rb \ \& \ \neg Kb$
- Therefore: $Gb \ \& \ \neg Hb$

Next suppose that we want to symbolize this argument:

Carol is a skilled surgeon and a tennis player. Therefore, Carol is a skilled tennis player.

If we start with the symbolization key we used for the previous argument, we could add a predicate (let Tx mean ‘ x is a tennis player’) and a name (let c mean Carol). Then the argument becomes:

- $(Rc \ \& \ Kc) \ \& \ Tc$
- Therefore: $Tc \ \& \ Kc$

This symbolization is a disaster! It takes what in English is a terrible argument and symbolizes it as a valid argument in FOL. The problem is that there is a difference between being *skilled as a surgeon* and *skilled as a tennis player*. Symbolizing this argument correctly requires two separate predicates, one for each type of skill. If we let K_1x mean ‘ x is skilled as a surgeon’ and K_2x mean ‘ x is skilled as a tennis player,’ then we can symbolize the argument in this way:

- $(Rc \ \& \ K_1c) \ \& \ Tc$
- Therefore: $Tc \ \& \ K_2c$

Like the English language argument it symbolizes, this is invalid.

The moral of these examples is that you need to be careful of symbolizing predicates in an ambiguous way. Similar problems can arise with predicates like *good*, *bad*, *big*, and *small*. Just as skilled surgeons and skilled tennis players have different skills, big dogs, big mice, and big problems are big in different ways.

Is it enough to have a predicate that means ‘ x is a skilled surgeon’, rather than two predicates ‘ x is skilled’ and ‘ x is a surgeon’? Sometimes. As sentence 15 shows, sometimes we do not need to distinguish between skilled surgeons and other surgeons.

Must we always distinguish between different ways of being skilled, good, bad, or big? No. As the argument about Billy shows, sometimes we only need to talk about one kind of skill. If you are symbolizing an argument that is just about dogs, it is fine to define a predicate that means ‘ x is big.’ If the domain includes dogs and mice, however, it is probably best to make the predicate mean ‘ x is big for a dog.’

Practice exercises

A. Here are the syllogistic figures identified by Aristotle and his successors, along with their medieval names:

- **Barbara.** All G are F. All H are G. So: All H are F
- **Celarent.** No G are F. All H are G. So: No H are F
- **Ferio.** No G are F. Some H is G. So: Some H is not F
- **Darii.** All G are F. Some H is G. So: Some H is F.
- **Camestres.** All F are G. No H are G. So: No H are F.
- **Cesare.** No F are G. All H are G. So: No H are F.
- **Baroko.** All F are G. Some H is not G. So: Some H is not F.
- **Festino.** No F are G. Some H are G. So: Some H is not F.

- **Datisi.** All G are F. Some G is H. So: Some H is F.
- **Disamis.** Some G is F. All G are H. So: Some H is F.
- **Ferison.** No G are F. Some G is H. So: Some H is not F.
- **Bokardo.** Some G is not F. All G are H. So: Some H is not F.
- **Camenes.** All F are G. No G are H So: No H is F.
- **Dimaris.** Some F is G. All G are H. So: Some H is F.
- **Fresison.** No F are G. Some G is H. So: Some H is not F.

Symbolize each argument in FOL.

B. Using the following symbolization key:

domain: people

Kx : ______x knows the combination to the safe

Sx : ______x is a spy

Vx : ______x is a vegetarian

h : Hofthor

i : Ingmar

symbolize the following sentences in FOL:

1. Neither Hofthor nor Ingmar is a vegetarian.
2. No spy knows the combination to the safe.
3. No one knows the combination to the safe unless Ingmar does.
4. Hofthor is a spy, but no vegetarian is a spy.

C. Using this symbolization key:

domain: all animals

Ax : ______x is an alligator.

Mx : ______x is a monkey.

Rx : ______x is a reptile.

Zx : ______x lives at the zoo.

a : Amos

b : Bouncer

c : Cleo

symbolize each of the following sentences in FOL:

1. Amos, Bouncer, and Cleo all live at the zoo.
2. Bouncer is a reptile, but not an alligator.
3. Some reptile lives at the zoo.
4. Every alligator is a reptile.
5. Any animal that lives at the zoo is either a monkey or an alligator.

6. There are reptiles which are not alligators.
7. If any animal is a reptile, then Amos is.
8. If any animal is an alligator, then it is a reptile.

D. For each argument, write a symbolization key and symbolize the argument in FOL.

1. Willard is a logician. All logicians wear funny hats. So Willard wears a funny hat
2. Nothing on my desk escapes my attention. There is a computer on my desk. As such, there is a computer that does not escape my attention.
3. All my dreams are black and white. Old TV shows are in black and white. Therefore, some of my dreams are old TV shows.
4. Neither Holmes nor Watson has been to Australia. A person could see a kangaroo only if they had been to Australia or to a zoo. Although Watson has not seen a kangaroo, Holmes has. Therefore, Holmes has been to a zoo.
5. No one expects the Spanish Inquisition. No one knows the troubles I've seen. Therefore, anyone who expects the Spanish Inquisition knows the troubles I've seen.
6. All babies are illogical. Nobody who is illogical can manage a crocodile. Berthold is a baby. Therefore, Berthold is unable to manage a crocodile.

CHAPTER 30

Multiple quantifiers

We see more power of FOL when quantifiers start stacking on top of one another.

Consider

1. Someone loves everyone.

Before considering how to symbolise that, we start off by symbolising the related sentence:

2. John loves everyone.

This can be symbolised as $\forall x Ljx$, using the symbolisation key:

domain: all people

j : John

Lxy : _____ x loves _____ y

This gives us an insight into how to symbolise **1**: it's like **2** except it might not be John who loves everyone, **1** just says that there is *someone*, y , such that y loves everyone. We will thus symbolise it $\exists y \forall x Lyx$.

In the earlier examples we always used x as our variables; but we here had to use y because x is already taken. We don't want to say $\exists x \forall x Lxx$ as it's not clear how one should read this sentence as we need to identify which variables come with which quantifiers.

30.1 The order of quantifiers

Consider the sentence ‘everyone loves someone’. This is potentially ambiguous. It might mean either of the following:

3. For every person x , there is some person that x loves
4. There is some particular person whom every person loves

Sentence 3 can be symbolized by $\forall x\exists yLxy$, and would be true of a love-triangle. For example, suppose that our domain of discourse is restricted to Imre, Juan and Karl. Suppose also that Karl loves Imre but not Juan, that Imre loves Juan but not Karl, and that Juan loves Karl but not Imre. Then sentence 3 is true.

Sentence 4 is symbolized by $\exists y\forall xLxy$. Sentence 4 is *not* true in the situation just described. Again, suppose that our domain of discourse is restricted to Imre, Juan and Karl. This requires that all of Juan, Imre and Karl converge on (at least) one object of love.

The point of the example is to illustrate that the order of the quantifiers matters a great deal. Indeed, to switch them around is called a *quantifier shift fallacy*. Here is an example, which comes up in various forms throughout the philosophical literature:

- For every person, there is some truth they cannot know. $(\forall\exists)$
- Therefore: There is some truth that no person can know. $(\exists\forall)$

This argument form is obviously invalid. It’s just as bad as:¹

- Every dog has its day. $(\forall\exists)$
- Therefore: There is a day for all the dogs. $(\exists\forall)$

The moral is: take great care with the order of quantification.

30.2 Stepping-stones to symbolization

Once we have the possibility of multiple quantifiers, representation in FOL can quickly start to become a bit tricky. When you are trying to symbolize a complex sentence, we recommend laying down several stepping stones. As usual, this idea is best illustrated by example. Consider this representation key:

domain: people and dogs

¹Thanks to Rob Trueman for the example.

Dx : ______x is a dog
 Fxy : ______x is a friend of ______y
 Oxy : ______x owns ______y
 g : Geraldo

Now let's try to symbolize these sentences:

5. Geraldo is a dog owner.
6. Someone is a dog owner.
7. All of Geraldo's friends are dog owners.
8. Every dog owner is a friend of a dog owner.
9. Every dog owner's friend owns a dog of a friend.

Sentence 5 can be paraphrased as, 'There is a dog that Geraldo owns'. This can be symbolized by $\exists x(Dx \ \& \ Ogx)$.

Sentence 6 can be paraphrased as, 'There is some y such that y is a dog owner'. Dealing with part of this, we might write $\exists y(y \text{ is a dog owner})$. Now the fragment we have left as ' y is a dog owner' is much like sentence 5, except that it is not specifically about Geraldo. So we can symbolize sentence 6 by:

$$\exists y \exists x (Dx \ \& \ Oyx)$$

We should pause to clarify something here. In working out how to symbolize the last sentence, we wrote down $\exists y(y \text{ is a dog owner})$. To be very clear: this is *neither* an FOL sentence *nor* an English sentence: it uses bits of FOL (\exists , y) and bits of English ('dog owner'). It is really is *just a stepping-stone* on the way to symbolizing the entire English sentence with a FOL sentence. You should regard it as a bit of rough-working-out, on a par with the doodles that you might absent-mindedly draw in the margin of this book, whilst you are concentrating fiercely on some problem.

Sentence 7 can be paraphrased as, 'Everyone who is a friend of Geraldo is a dog owner'. Using our stepping-stone tactic, we might write

$$\forall x [F x g \rightarrow x \text{ is a dog owner}]$$

Now the fragment that we have left to deal with, ' x is a dog owner', is structurally just like sentence 5. However, it would be a mistake for us simply to write

$$\forall x [F x g \rightarrow \exists x (Dx \ \& \ Oxx)]$$

for we would here have a *clash of variables*. The scope of the universal quantifier, $\forall x$, is the entire conditional, so the x in Dx should be

governed by that, but Dx also falls under the scope of the existential quantifier $\exists x$, so the x in Dx should be governed by that. Now confusion reigns: which x are we talking about? Suddenly the sentence becomes ambiguous (if it is even meaningful at all), and logicians hate ambiguity. The broad moral is that a single variable cannot serve two quantifier-masters simultaneously.

To continue our symbolization, then, we must choose some different variable for our existential quantifier. What we want is something like:

$$\forall x[Fxg \rightarrow \exists z(Dz \& Oxz)]$$

This adequately symbolizes sentence 7.

Sentence 8 can be paraphrased as ‘For any x that is a dog owner, there is a dog owner who x is a friend of’. Using our stepping-stone tactic, this becomes

$$\forall x[x \text{ is a dog owner} \rightarrow \exists y(y \text{ is a dog owner} \& Fxy)]$$

Completing the symbolization, we end up with

$$\forall x[\exists z(Dz \& Oxz) \rightarrow \exists y(\exists z(Dz \& Oyz) \& Fxy)]$$

Note that we have used the same letter, z , in both the antecedent and the consequent of the conditional, but that these are governed by two different quantifiers. This is ok: there is no clash here, because it is clear which quantifier that variable falls under. We might graphically represent the scope of the quantifiers thus:

$$\begin{array}{c} \text{scope of ‘}\forall x\text{’} \\ \overbrace{\hspace{15em}} \\ \text{scope of ‘}\exists y\text{’} \\ \overbrace{\hspace{10em}} \\ \text{scope of 1st ‘}\exists z\text{’} \quad \text{scope of 2nd ‘}\exists z\text{’} \\ \overbrace{\hspace{5em}} \quad \overbrace{\hspace{5em}} \\ \forall x[\exists z(Dz \& Oxz) \rightarrow \exists y(\exists z(Dz \& Oyz) \& Fxy)] \end{array}$$

This shows that no variable is being forced to serve two masters simultaneously.

Sentence 9 is the trickiest yet. First we paraphrase it as ‘For any x that is a friend of a dog owner, x owns a dog which is also owned by a friend of x ’. Using our stepping-stone tactic, this becomes:

$$\forall x[x \text{ is a friend of a dog owner} \rightarrow \\ x \text{ owns a dog which is owned by a friend of } x]$$

Breaking this down a bit more:

$$\forall x [\exists y (Fxy \ \& \ y \text{ is a dog owner}) \rightarrow \exists y (Dy \ \& \ Oxy \ \& \ y \text{ is owned by a friend of } x)]$$

And a bit more:

$$\forall x [\exists y (Fxy \ \& \ \exists z (Dz \ \& \ Oyz)) \rightarrow \exists y (Dy \ \& \ Oxy \ \& \ \exists z (Fzx \ \& \ Ozy))]$$

And we are done!

30.3 Supressed quantifiers

Logic can often help to get clear on the meanings of English claims, especially where the quantifiers are left implicit or their order is ambiguous or unclear. The clarity of expression and thinking afforded by FOL can give you a significant advantage in argument, as can be seen in the following takedown by British political philosopher Mary Astell (1666–1731) of her contemporary, the theologian William Nicholls. In Discourse IV: The Duty of Wives to their Husbands of his *The Duty of Inferiors towards their Superiors, in Five Practical Discourses* (London 1701), Nicholls argued that women are naturally inferior to men. In the preface to the 3rd edition of her treatise *Some Reflections upon Marriage, Occasion'd by the Duke and Duchess of Mazarine's Case; which is also considered*, Astell responded as follows:

'Tis true, thro' Want of Learning, and of that Superior Genius which Men as Men lay claim to, she [Astell] was ignorant of the *Natural Inferiority* of our Sex, which our Masters lay down as a Self-Evident and Fundamental Truth. She saw nothing in the Reason of Things, to make this either a Principle or a Conclusion, but much to the contrary; it being Sedition at least, if not Treason to assert it in this Reign.

For if by the Natural Superiority of their Sex, they mean that *every* Man is by Nature superior to *every* Woman, which is the obvious meaning, and that which must be stuck to if they would speak Sense, it wou'd be a Sin in *any* Woman to have Dominion over *any* Man, and the greatest Queen ought not to command but to obey her Footman, because

no Municipal Laws can supersede or change the Law of Nature; so that if the Dominion of the Men be such, the *Salique Law*,² as unjust as *English Men* have ever thought it, ought to take place over all the Earth, and the most glorious Reigns in the *English, Danish, Castilian*, and other Annals, were wicked Violations of the Law of Nature!

If they mean that *some* Men are superior to *some* Women this is no great Discovery; had they turn'd the Tables they might have seen that *some* Women are Superior to *some* Men. Or had they been pleased to remember their Oaths of Allegiance and Supremacy, they might have known that *One* Woman is superior to *All* the Men in these Nations, or else they have sworn to very little purpose.³ And it must not be suppos'd, that their Reason and Religion wou'd suffer them to take Oaths, contrary to the Laws of Nature and Reason of things.⁴

We can symbolize the different interpretations Astell offers of Nicholls' claim that men are superior to women: He either meant that every man is superior to every woman, i.e.,

$$\forall x(Mx \rightarrow \forall y(Wy \rightarrow Sxy))$$

or that some men are superior to some women,

$$\exists x(Mx \ \& \ \exists y(Wy \ \& \ Sxy)).$$

The latter is true, but so is

$$\exists y(Wy \ \& \ \exists x(Mx \ \& \ Syx)).$$

(some women are superior to some men), so that would be “no great discovery.” In fact, since the Queen is superior to all her subjects, it's even true that some woman is superior to every man, i.e.,

$$\exists y(Wy \wedge \forall x(Mx \rightarrow Syx)).$$

But this is incompatible with the “obvious meaning” of Nicholls' claim, i.e., the first reading. So what Nicholls claims amounts to treason against the Queen!

²The Salique law was the common law of France which prohibited the crown be passed on to female heirs.

³In 1706, England was ruled by Queen Anne.

⁴Mary Astell, *Reflections upon Marriage*, 1706 Preface, iii–iv, and Mary Astell, *Political Writings*, ed. Patricia Springborg, Cambridge University Press, 1996, 9–10.

Practice exercises

Warning: some of these problems require you to use identity, which hasn't yet been introduced. See Chapter IX.

A. Using this symbolization key:

domain: all animals

Ax : ______x is an alligator

Mx : ______x is a monkey

Rx : ______x is a reptile

Zx : ______x lives at the zoo

Lxy : ______x loves ______y

a : Amos

b : Bouncer

c : Cleo

symbolize each of the following sentences in FOL:

1. If Cleo loves Bouncer, then Bouncer is a monkey.
2. If both Bouncer and Cleo are alligators, then Amos loves them both.
3. Cleo loves a reptile.
4. Bouncer loves all the monkeys that live at the zoo.
5. All the monkeys that Amos loves love him back.
6. Every monkey that Cleo loves is also loved by Amos.
7. There is a monkey that loves Bouncer, but sadly Bouncer does not reciprocate this love.

B. Using the following symbolization key:

domain: all animals

Dx : ______x is a dog

Sx : ______x likes samurai movies

Lxy : ______x is larger than ______y

r : Rave

h : Shane

d : Daisy

symbolize the following sentences in FOL:

1. Rave is a dog who likes samurai movies.
2. Rave, Shane, and Daisy are all dogs.
3. Shane is larger than Rave, and Daisy is larger than Shane.

4. All dogs like samurai movies.
5. Only dogs like samurai movies.
6. There is a dog that is larger than Shane.
7. If there is a dog larger than Daisy, then there is a dog larger than Shane.
8. No animal that likes samurai movies is larger than Shane.
9. No dog is larger than Daisy.
10. Any animal that dislikes samurai movies is larger than Rave.
11. There is an animal that is between Rave and Shane in size.
12. There is no dog that is between Rave and Shane in size.
13. No dog is larger than itself.
14. Every dog is larger than some dog.
15. There is an animal that is smaller than every dog.
16. If there is an animal that is larger than any dog, then that animal does not like samurai movies.

C. Using the symbolization key given, symbolize each English-language sentence into FOL.

domain: candies

Cx : ______x has chocolate in it.

Mx : ______x has marzipan in it.

Sx : ______x has sugar in it.

Tx : Boris has tried ______x .

Bxy : ______x is better than ______y .

1. Boris has never tried any candy.
2. Marzipan is always made with sugar.
3. Some candy is sugar-free.
4. The very best candy is chocolate.
5. No candy is better than itself.
6. Boris has never tried sugar-free chocolate.
7. Boris has tried marzipan and chocolate, but never together.
8. Any candy with chocolate is better than any candy without it.
9. Any candy with chocolate and marzipan is better than any candy that lacks both.

D. Using the following symbolization key:

domain: people and dishes at a potluck

Rx : ______x has run out.

Tx : ______x is on the table.

Fx : ______x is food.
 Px : ______x is a person.
 Lxy : ______x likes ______y .
 e : Eli
 f : Francesca
 g : the guacamole

symbolize the following English sentences in FOL:

1. All the food is on the table.
2. If the guacamole has not run out, then it is on the table.
3. Everyone likes the guacamole.
4. If anyone likes the guacamole, then Eli does.
5. Francesca only likes the dishes that have run out.
6. Francesca likes no one, and no one likes Francesca.
7. Eli likes anyone who likes the guacamole.
8. Eli likes anyone who likes the people that he likes.
9. If there is a person on the table already, then all of the food must have run out.

E. Using the following symbolization key:

domain: people

Dx : ______x dances ballet.
 Fx : ______x is female.
 Mx : ______x is male.
 Cxy : ______x is a child of ______y .
 Sxy : ______x is a sibling of ______y .
 e : Elmer
 j : Jane
 p : Patrick

symbolize the following sentences in FOL:

1. All of Patrick's children are ballet dancers.
2. Jane is Patrick's daughter.
3. Patrick has a daughter.
4. Jane is an only child.
5. All of Patrick's sons dance ballet.
6. Patrick has no sons.
7. Jane is Elmer's niece.
8. Patrick is Elmer's brother.
9. Patrick's brothers have no children.

10. Jane is an aunt.
11. Everyone who dances ballet has a brother who also dances ballet.
12. Every woman who dances ballet is the child of someone who dances ballet.

CHAPTER 31

Sentences of FOL

We know how to represent English sentences in FOL. The time has finally come to define the notion of a *sentence* of FOL.

31.1 Expressions

There are six kinds of symbols in FOL:

Predicates $A, B, C, \dots, Z,$ with subscripts, as needed:
 $A_1, Z_2, A_{25}, J_{375}, \dots$ ¹

Names $a, b, c, \dots, t,$ or with subscripts, as needed $a_1, b_{224}, h_7, m_{32}, \dots$

Variables $x, y, z,$ or with subscripts, as needed $x_1, y_1, z_1, x_2, \dots$

¹Each predicate will have a number of places associated with it. We should thus really introduce:

Zero-Place Predicates = Atomic sentences of TFL $A^0, B^0, \dots, Z^0,$ with subscripts, as needed: $A_1^0, Z_2^0, A_{25}^0, J_{375}^0, \dots$

One-Place Predicates $A^1, B^1, \dots, Z^1,$ with subscripts, as needed: $A_1^1, Z_2^1, A_{25}^1, J_{375}^1, \dots$

Two-Place Predicates $A^2, B^2, \dots, Z^2,$ with subscripts, as needed: $A_1^2, Z_2^2, A_{25}^2, J_{375}^2, \dots$

Three-Place Predicates $A^3, B^3, \dots, Z^3,$ with subscripts, as needed: $A_1^3, Z_2^3, A_{25}^3, J_{375}^3, \dots$
etc. We drop the superscripts for ease.

Connectives $\neg, \&, \vee, \rightarrow, \leftrightarrow$

Brackets $(,)$

Quantifiers \forall, \exists

We define an **EXPRESSION OF FOL** as any string of symbols of FOL. Take any of the symbols of FOL and write them down, in any order, and you have an expression.

31.2 Formulas

In §6, we went straight from the statement of the vocabulary of TFL to the definition of a sentence of TFL. In FOL, we will have to go via an intermediary stage: via the notion of a **FORMULA**. The intuitive idea is that a formula is any sentence, or anything which can be turned into a sentence by adding quantifiers out front. But this will take some unpacking.

As we did for TFL, we will present a recursive definition of a formula of FOL.

The starting point of this is the notion of an *atomic formula*. In TFL we started our definition with the notion of an atomic sentence, which were just given to us in our vocabulary. In FOL, the starting point of our definition is the notion of an atomic formula. Atomic formulas will be given by the following definition:

If Z is an n -place predicate and t_1, \dots, t_n are either variables or names, then $Zt_1 \dots t_n$ is an **ATOMIC FORMULA**.

For example, if D is a one-place predicate (we might have introduced it to symbolise ‘_____x is a dog’), and L is a two-place predicate (we might have introduced it to symbolise ‘_____x loves _____y’), then the following are atomic formulas:

$$Db, Dx, Dy, Lki, Lkx, Lyz$$

Formulas are constructed by starting with these and using either our TFL connectives or our quantifiers.

We can now give the recursive definition of what it is to be a formula of FOL.

1. If Z is an n -place predicate and t_1, \dots, t_n are either variables or names, then $Zt_1 \dots t_n$ is a formula.
These are called **ATOMIC FORMULAS**.
2. If X is a formula, then $\neg X$ is a formula.
3. If X and Y are formulas, then
 - a) $(X \ \& \ Y)$ is a formula,
 - b) $(X \vee Y)$ is a formula,
 - c) $(X \rightarrow Y)$ is a formula, and
 - d) $(X \leftrightarrow Y)$ is a formula.
4. If X is a formula, v is a variable and $\forall v$ and $\exists v$ do not already appear in X , then
 - a) $\forall v X$ is a formula
 - b) $\exists v X$ is a formula.
5. Nothing else is a formula.

As for TFL, we start out with some formulas, such as Dx or Db , and we can construct more complicated formulas with our connectives, e.g.

$$\begin{aligned} &(Dx \ \& \ Db), \\ &\neg(Dx \ \& \ Db) \\ &(\neg(Dx \ \& \ Db) \rightarrow Lxy) \end{aligned}$$

And we can display their construction using our formation trees, as in 6.1.

$$\begin{array}{c} (\neg(Dx \ \& \ Db) \rightarrow Lxy) \\ \swarrow \quad \searrow \\ \neg(Dx \ \& \ Db) \quad Lxy \\ | \\ (Dx \ \& \ Db) \\ \swarrow \quad \searrow \\ Dx \quad Db \end{array}$$

This is exactly as in the case of TFL, the only difference is that the “leaves” of the tree have more structure to them: they’re predicates applied to names or variables rather than simply the single atomic sentences that we had in TFL.

The new clauses here are in 4. This lets us put $\forall x$ in front of a formula, e.g. Bx to construct a formula $\forall x Bx$. We can also add quantifiers when the formula was already more complicated, e.g., we can construct a formula

$$\forall x(\neg(Dx \ \& \ Db) \rightarrow Lxy).$$

(v is a metavariable for variables, it could be x , y , z or x_{3000} . Recall §9.3.)

We could also have added an existential quantifier, \exists , to construct

$$\exists x(\neg(Dx \ \& \ Db) \rightarrow Lxy).$$

We can also do it with other variables, e.g.

$$\forall y(\neg(Dx \ \& \ Db) \rightarrow Lxy).$$


We can then add further quantifiers to *these* new formula, to construct, e.g.

$$\exists y \forall x(\neg(Dx \ \& \ Db) \rightarrow Lxy).$$

We can again display the structure and construction of the sentence perspicuously by presenting a formation tree:

$$\begin{array}{c} \exists y \forall x(\neg(Dx \ \& \ Db) \rightarrow Lxy) \\ | \\ \forall x(\neg(Dx \ \& \ Db) \rightarrow Lxy) \\ | \\ (\neg(Dx \ \& \ Db) \rightarrow Lxy) \\ / \quad \backslash \\ \neg(Dx \ \& \ Db) \quad Lxy \\ | \\ (Dx \ \& \ Db) \\ / \quad \backslash \\ Dx \quad Db \end{array}$$

Let’s just see one more example:



Moving up the formation tree is following one of the rules of the recursive specification of what it is to be a sentence.


Why in 4 did we have the restriction that $\exists v$ or $\forall v$ is not already in X ? This is to ensure that variables only serve one master at any one time (see §30). Otherwise we could see that $\forall x Rxx$ is a sentence and then conclude that $\exists x \forall x Rxx$, which is not a good sentence. However, note that $\exists x Cx \vee \forall x Bx$ is a sentence. Here the variable x is used in both quantifiers, but there's no ambiguity because the sentence was constructed by combining the two sentences $\exists x Cx$ and $\forall x Bx$ with a connective, \vee . The *scope* of $\exists x$ is just $\exists x Cx$; it doesn't look "over" the connective to where $\forall x$ is used. So the two uses of x are kept separate and no problems arise. However, to avoid any potential worries it is generally a good idea to use different variables when symbolising sentences; in this case one could equally well give $\exists x Cx \vee \forall y By$ as a symbolisation.

The notions of scope and main logical operators that were given in ?? equally applies to FOL but now the main logical operator might be a quantifier. These were:

The **MAIN LOGICAL OPERATOR** in a sentence is the operator that was introduced last when that sentence was constructed using the recursion rules.

The **SCOPE** of a logical operator in a sentence is the formula for which that operator is the main logical operator.

We can graphically illustrate scopes as follows:



31.3 Sentences

Recall that we are largely concerned in logic with assertoric sentences: sentences that can be either true or false. Many formulas are not sentences. Consider the following symbolization key:

domain: people

Lxy : ---_x loves ---_y

b : Boris

Consider the atomic formula Lzz . Can it be true or false? You might think that it will be true just in case the person named by z loves themself, in the same way that Lbb is true just in case Boris (the person named by b) loves himself. *However, z is a variable, and does not name anyone or any thing.*

Of course, if we put an existential quantifier out front, obtaining $\exists zLzz$, then this would be true iff someone loves herself. Equally, if we wrote $\forall zLzz$, this would be true iff everyone loves themself. The point is that we need a quantifier to tell us how to deal with a variable.

Let's make this idea precise.

A **BOUND VARIABLE** is an occurrence of a variable v that is within the scope of either $\forall v$ or $\exists v$.

A **FREE VARIABLE** is any variable that is not bound.

For example, consider the formula

$$\forall x(Ex \vee Dy) \rightarrow \exists z(Ex \rightarrow Lzx)$$

The scope of the universal quantifier $\forall x$ is $\forall x(Ex \vee Dy)$, so the first x is bound by the universal quantifier. However, the second and third occurrence of x are free. Equally, the y is free. The scope of the existential quantifier $\exists z$ is $(Ex \rightarrow Lzx)$, so z is bound.

Finally we can say the following.

A **SENTENCE** of FOL is any formula of FOL that contains no free variables.

31.4 Bracketing conventions

We will adopt the same notational conventions governing brackets that we did for TFL (see §6 and §12.3.): we may omit the outermost brackets of a formula.

Practice exercises

A. Identify which variables are bound and which are free.

1. $\exists x Lxy \ \& \ \forall y Lyx$
2. $\forall x Ax \ \& \ Bx$
3. $\forall x (Ax \ \& \ Bx) \ \& \ \forall y (Cx \ \& \ Dy)$
4. $\forall x \exists y [Rxy \rightarrow (Jz \ \& \ Kx)] \vee Ryx$
5. $\forall x_1 (Mx_2 \leftrightarrow Lx_2x_1) \ \& \ \exists x_2 Lx_3x_2$

CHAPTER 32

Ambiguity

In chapter 8 we discussed the fact that sentences of English can be ambiguous, and pointed out that sentences of TFL are not. One important application of this fact is that the structural ambiguity of English sentences can often, and usefully, be straightened out using different symbolizations. One common source of ambiguity is *scope ambiguity*, where the English sentence does not make it clear which logical word is supposed to be in the scope of which other. Multiple interpretations are possible. In FOL, every connective and quantifier has a well-determined scope, and so whether or not one of them occurs in the scope of another in a given sentence of FOL is always determined.

Negation

For instance, consider the English idiom,

1. Everything that glitters is not gold.

If we think of this sentence as of the form ‘every F is not G ’ where Fx symbolizes ‘_____ x glitters’ and Gx is ‘_____ x is not gold’, we would symbolize it as:

$$\bullet \forall x(Fx \rightarrow \neg Gx),$$

in other words, we symbolize it the same way as we would ‘Nothing that glitters is gold’. But the idiom does not mean that! It means that one should not assume that just because something glitters, it is gold; not everything that appears valuable is in fact valuable. To capture the actual meaning of the idiom, we would have to symbolize it instead as we would ‘Not everything that glitters is gold’, i.e., in the following way:

- $\neg\forall x(Fx \rightarrow Gx)$

Compare the first of these with the previous symbolization: again we see that the difference in the two meanings of the ambiguous sentence lies in whether the ‘ \neg ’ is in the scope of the ‘ \forall ’ (in the first symbolization) or ‘ \forall ’ is in the scope of ‘ \neg ’ (in the second).

Of course we can alternatively symbolize the two readings using existential quantifiers as well:

- $\neg\exists x(Fx \& Gx)$
- $\exists x(Fx \& \neg Gx)$

Only

Let’s first say a bit explicitly about symbolising ‘only’ before discussing an ambiguity that arises with it. Consider ‘only’, as in:

2. Only dimes are on the table.

How should we symbolize this? A good strategy is to consider when the sentence would be false. If we are saying that only dimes are on the table, we are excluding all the cases where something on the table is a non-dime. So we can symbolize the sentence the same way we would symbolize ‘No non-dimes are on the table.’ Remembering the lesson we just learned, and symbolizing ‘ x is a non-dime’ as ‘ $\neg D(x)$ ’, the possible symbolizations are: ‘ $\neg\exists x(T(x) \& \neg D(x))$ ’, or alternatively: ‘ $\forall x(T(x) \rightarrow \neg\neg D(x))$ ’. Since double negations cancel out, the second is just as good as ‘ $\forall x(T(x) \rightarrow D(x))$ ’. In other words, ‘Only dimes are on the table’ and ‘Everything on the table is a dime’ are symbolized the same way.

If a sentence can be paraphrased in English as

‘only F s are G s’,

it can be symbolised as can be symbolized as

$$\neg\exists x(G(x) \& \neg F(x)), \text{ or} \\ \forall x(G(x) \rightarrow F(x))$$

Now let’s look at an ambiguity that arises:

Consider the sentence:

3. Only young cats are playful.

According to our schema, we would symbolize it this way:

- $\forall x(Px \rightarrow (Yx \ \& \ Cx))$

The meaning of this sentence of FOL is something like, ‘If an animal is playful, it is a young cat’. (Assuming that the domain is animals, of course.) This is probably not what’s intended in uttering sentence 3, however. It’s more likely that we want to say that old cats are not playful. In other words, what we mean to say is that if something is a cat and playful, it must be young. This would be symbolized as:

- $\forall x((Cx \ \& \ Px) \rightarrow Yx)$

There is even a third reading! Suppose we’re talking about young animals and their characteristics. And suppose you wanted to say that of all the young animals, only the cats are playful. You could symbolize this reading as:

- $\forall x((Yx \ \& \ Px) \rightarrow Cx)$

Each of the last two readings can be made salient in English by placing the stress appropriately. For instance, to suggest the last reading, you would say ‘Only young *cats* are playful’, and to get the other reading you would say ‘Only *young* cats are playful’. The very first reading can be indicated by stressing both ‘young’ and ‘cats’: ‘Only *young cats* are playful’ (but not old cats, or dogs of any age).

Quantifier Order

In sections 30.1 and 30.3 we discussed the importance of the order of quantifiers. This is relevant here because, in English, the order of quantifiers is sometimes not completely determined. When both universal (‘all’) and existential (‘some’, ‘a’) quantifiers are involved, this can result in scope ambiguities. Consider:

4. Everyone went to see a movie.

This sentence is ambiguous. In one interpretation, it means that there is a single movie that everyone went to see. In the other, it means that everyone went to see some movie or other, but not necessarily the same one. The two readings can be symbolized, respectively, by

- $\exists x(Mx \ \& \ \forall y(Py \rightarrow Sy, x))$

- $\forall y(Py \rightarrow \exists x(Mx \& Sy, x))$

We assume here that the domain contains (at least) people and movies, and the symbolization key,

Py : ______y is a person,
 Mx : ______x is a movie
 Sy, x : ______y went to see ______x.

In the first reading, we say that the existential quantifier has *wide scope* (and its scope contains the universal quantifier, which has *narrow scope*), and the other way round in the second.

Negation and Definite Descriptions

In chapter 43, we will encounter another scope ambiguity, arising from definite descriptions interacting with negation. Consider Russell's own example:

5. The King of France is not bald.

If the definite description has wide scope, and we are interpreting the 'not' as an 'inner' negation (as we said before), sentence 5 is interpreted to assert the existence of a single King of France, to whom we are ascribing non-baldness. In this reading, it is symbolized as ' $\exists x[Kx \& \forall y(Ky \rightarrow x = y)) \& \neg Bx]$ '. In the other reading, the 'not' denies the sentence 'The King of France is bald', and we would symbolize it as: ' $\neg \exists x[Kx \& \forall y(Ky \rightarrow x = y)) \& Bx]$ '. In the first case, we say that the definite description has wide scope and in the second that it has narrow scope.

Practice exercises

A. Each of the following sentences is ambiguous. Provide a symbolization key for each, and symbolize all readings.

1. Noone likes a quitter.
2. CSI found only red hair at the scene.
3. Smith's murderer hasn't been arrested.

B. Russell gave the following example in his paper 'On Denoting':

I have heard of a touchy owner of a yacht to whom a guest, on first seeing it, remarked, 'I thought your yacht was larger than it is'; and the owner replied, 'No, my yacht is not larger than it is'.

Explain what's going on.

PART VII

Interpretations

CHAPTER 33

Extensionality

Recall that TFL is a truth-functional language. Its connectives are all truth-functional, and *all* that we can do with TFL is key sentences to particular truth values. We can do this *directly*. For example, we might stipulate that the TFL sentence P is to be true. Alternatively, we can do this *indirectly*, offering a symbolization key, e.g.:

P : Big Ben is in London

Now recall from §11 that this should be taken to mean:

- The TFL sentence P is to take the same truth value as the English sentence ‘Big Ben is in London’ (whatever that truth value may be)

The point that we emphasized is that TFL cannot handle differences in meaning that go beyond mere differences in truth value.

33.1 Symbolizing versus translating

FOL has some similar limitations, but it goes beyond mere truth values, since it enables us to split up sentences into terms, predicates and quantifier expressions. This enables us to consider what is *true of* some particular object, or of some or all objects. But we can do no more than that.

When we provide a symbolization key for some FOL predicates, such as:

Cx : _____ $_x$ teaches Logic III in Calgary

we do not carry the *meaning* of the English predicate across into our FOL predicate. We are simply stipulating something like the following:

- Cx and ‘_____x teaches Logic III in Calgary’ are to be *true of* exactly the same things.

So, in particular:

- Cx is to be true of all and only those things which teach Logic III in Calgary (whatever those things might be).

This is an indirect stipulation. Alternatively, we can directly stipulate which objects a predicate should be true of. For example, we can stipulate that Cx is to be true of Richard Zach, and Richard Zach alone. As it happens, this direct stipulation would have the same effect as the indirect stipulation. Note, however, that the English predicates ‘_____ is Richard Zach’ and ‘_____ teaches Logic III in Calgary’ have very different meanings!

The point is that FOL does not give us any resources for dealing with nuances of meaning. When we interpret FOL, all we are considering is what the predicates are true of, regardless of whether we specify these things directly or indirectly. The things a predicate is true of are known as the **EXTENSION** of that predicate. We say that FOL is an **EXTENSIONAL LANGUAGE** because FOL does not represent differences of meaning between predicates that have the same extension.

For this reason, we say only that FOL sentences *symbolize* English sentences. It is doubtful that we are *translating* English into FOL, as translations should preserve meanings, and not just extensions.

33.2 A word on extensions

We can stipulate directly what predicates are to be true of, so it is worth noting that our stipulations can be as arbitrary as we like. For example, we could stipulate that Hx should be true of, and only of, the following objects:

Justin Trudeau
the number π
every top-F key on every piano ever made

Now, the objects that we have listed have nothing particularly in common. But this doesn’t matter. Logic doesn’t care about what strikes us

mere humans as ‘natural’ or ‘similar’. Armed with this interpretation of Hx , suppose we now add to our symbolization key:

j : Justin Trudeau

r : Rachel Notley

p : the number π

Then Hj and Hp will both be true, on this interpretation, but Hr will be false, since Rachel Notley was not among the stipulated objects.

33.3 Many-place predicates

All of this is quite easy to understand when it comes to one-place predicates, but it gets messier when we consider two-place predicates. Consider a symbolization key like:

Lxy : _____ x loves _____ y

Given what we said above, this symbolization key should be read as saying:

- Lxy and ‘_____ x loves _____ y ’ are to be true of exactly the same things

So, in particular:

- Lxy is to be true of x and y (in that order) iff x loves y .

It is important that we insist upon the order here, since love—famously—is not always reciprocated. (Note that ‘ x ’ and ‘ y ’ here are symbols of augmented English, and that they are being *used*. By contrast, x and y are symbols of FOL, and they are being *mentioned*.)

That is an indirect stipulation. What about a direct stipulation? This is slightly harder. If we *simply* list objects that fall under Lxy , we will not know whether they are the lover or the beloved (or both). We have to find a way to include the order in our explicit stipulation.

To do this, we can specify that two-place predicates are true of *pairs* of objects, where the order of the pair is important. Thus we might stipulate that Bxy is to be true of, and only of, the following pairs of objects:

⟨Lenin, Marx⟩

⟨Heidegger, Sartre⟩

⟨Sartre, Heidegger⟩

Here the angle-brackets keep us informed concerning order. Suppose we now add the following stipulations:

l: Lenin
m: Marx
h: Heidegger
r: Sartre

Then *Blm* will be true, since $\langle \text{Lenin}, \text{Marx} \rangle$ was in our explicit list, but *Bml* will be false, since $\langle \text{Marx}, \text{Lenin} \rangle$ was not in our list. However, both *Bhr* and *Brh* will be true, since both $\langle \text{Heidegger}, \text{Sartre} \rangle$ and $\langle \text{Sartre}, \text{Heidegger} \rangle$ are in our explicit list.

To make these ideas more precise, we would need to develop some *set theory*. That would give us some precise tools for dealing with extensions and with ordered pairs (and ordered triples, etc.). However, set theory is not covered in this book, so we will leave these ideas at an imprecise level. Nevertheless, the general idea should be clear.

33.4 Interpretation

We defined a **VALUATION** in TFL as any assignment of truth and falsity to atomic sentences. In FOL, we are going to define an **INTERPRETATION** as consisting of three things:

- the specification of a domain
- for each name that we care to consider, an assignment of exactly one object within the domain
- for each predicate that we care to consider, a specification of what things (in what order) the predicate is to be true of

The symbolization keys that we considered in Part **VI** consequently give us one very convenient way to present an interpretation. We will continue to use them throughout this chapter. However, it is sometimes also convenient to present an interpretation *diagrammatically*.


Suppose we want to consider just a single two-place predicate, *Rxy*. Then we can represent it just by drawing an arrow between two objects, and stipulate that *Rxy* is to hold of *x* and *y* just in case there is an arrow running from *x* to *y* in our diagram. As an example, we might offer:



This would be suitable to characterize an interpretation whose domain is the first four positive whole numbers, and which interprets Rxy as being true of and only of:

$$\langle 1, 2 \rangle, \langle 2, 3 \rangle, \langle 3, 4 \rangle, \langle 4, 1 \rangle, \langle 1, 3 \rangle$$

Equally we might offer:



for an interpretation with the same domain, which interprets Rxy as being true of and only of:

$$\langle 1, 3 \rangle, \langle 3, 1 \rangle, \langle 3, 4 \rangle, \langle 1, 1 \rangle, \langle 3, 3 \rangle$$

If we wanted, we could make our diagrams more complex. For example, we could add names as labels for particular objects. Equally, to symbolize the extension of a one-place predicate, we might simply draw a ring around some particular objects and stipulate that the thus encircled objects (and only them) are to fall under the predicate Hx , say.

CHAPTER 34

Truth in FOL

We know what interpretations are. Since, among other things, they tell us which predicates are true of which objects, they will provide us with an account of the truth of atomic sentences. However, we must also present a detailed account of what it is for an arbitrary FOL sentence to be true or false in an interpretation.

But we defined what a sentence was by first specifying what a formula is. Formulas like Hx aren't the sorts of things that are true or false in interpretations. Only sentences are true or false. But if we provide extra information we can determine the truth of Hx : we need to specify what x refers to. This is done by using a variable assignment:

A variable assignment specifies an object for each variable.

We define whether a formula is true or false *under a variable assignment*. We know from §31 that there are three kinds of formulas in FOL:

- atomic formulas
- formulas whose main logical operator is a sentential connective
- formulas whose main logical operator is a quantifier

We need to explain truth for all three kinds of formula.

We will provide a completely general explanation in this section. However, to try to keep the explanation comprehensible, we will, at several points, use the following interpretation:

domain: all people born before 2000CE

a : Aristotle

b : Beyoncé

Px : _____ _{x} is a philosopher
 Rxy : _____ _{x} was born before _____ _{y}

This will be our *go-to example* in what follows.

34.1 Atomic formulas

Atomic formulas are things like Px , Pb or Rax .

An atomic sentence like Pb is checked for truth just by consulting our interpretation: Px is ‘_____ _{x} is a philosopher’, so if we’re looking at Pb we fill out the gap with Beyoncé, and Beyoncé is not a philosopher, so Pb is false.

What about Px ? This reads something like ‘they are a philosopher’. The question is who ‘they’ refers to, or in the logic terms: who x is. This depends on a variable assignment. Our variable assignment needs to give an object in our domain for the variable x . For example, it might give Beyoncé, then since Beyoncé is not a philosopher, Px would be false on this variable assignment. Our variable assignment doesn’t need to specify one of the objects that are named, it can give us anyone in our domain, e.g. Queen Elizabeth II. Under the variable assignment which assigns x Queen Elizabeth II, Px is false: Queen Elizabeth II is not a philosopher.

Likewise, on this interpretation, Rab is true iff the object named by a was born before the object named by b . Well, Aristotle was born before Beyoncé. So Rab is true. Equally, Raa is false: Aristotle was not born before Aristotle. How about Rax ? Well what does our variable assignment specify for x ? If we have a variable assignment where x is Queen Elizabeth II, then Rax is true: Aristotle was born before Queen Elizabeth II.

Dealing with atomic sentences, then, is very intuitive. When R is an n -place predicate and t_1, t_2, \dots, t_n are names or variables, then

$Rt_1t_2 \dots t_n$ is true in an interpretation under a variable assignment **iff**

R is true of the objects referred to by t_1, t_2, \dots, t_n in that interpretation under that variable assignment (considered in that order)

34.2 Sentential connectives

We saw in §31 that FOL formulas can be built up from simpler ones using the truth-functional connectives that were familiar from TFL. The rules governing these truth-functional connectives are *exactly* the same as they were when we considered TFL. Here they are:

$X \& Y$ is true in an interpretation under a variable assignment
iff

both X and Y is true in that interpretation under that variable assignment

$X \vee Y$ is true in an interpretation under a variable assignment
iff

either X is true or Y is true in that interpretation under that variable assignment

$\neg X$ is true in an interpretation under a variable assignment **iff**
 X is false in that interpretation under that variable assignment

$X \rightarrow Y$ is true in an interpretation under a variable assignment
iff

either X is false or Y is true in that interpretation under that variable assignment

$X \leftrightarrow Y$ is true in an interpretation under a variable assignment
iff

X has the same truth value as Y in that interpretation under that variable assignment

This is just another presentation of the truth rules we gave for the connectives in TFL; it just does so in a slightly different way. Some examples will probably help to illustrate the idea. On our go-to interpretation:

- Pa is true
- $Rab \& Pb$ is false because, although Rab is true, Pb is false
- $\neg Pa$ is false
- $Pa \& \neg(Pb \& Rab)$ is true, because Pa is true and Pb is false, so $Pb \& Rab$ is false, thus $\neg(Pb \& Rab)$ is also true.

Make sure you understand these examples.


We can also carry variable assignments around with us. Consider a variable assignment which assigns David Hume to x . Then

- Px is true under this variable assignment: David Hume was a philosopher
- Bxa is false under this variable assignment: David Hume was born after Aristotle
- $Px \rightarrow Bxa$ is false under this variable assignment: Px is true and Bxa is false, so by our rule for \rightarrow , $Px \rightarrow Bxa$ is false.

34.3 When the main logical operator is a quantifier

The exciting innovation in FOL, though, is the use of *quantifiers*.

Consider the following interpretation:



domain: People in above picture (Alice, Bob, Cathy and Denny)

Hx : _____ $_x$ has horns (Bob)

Sx : _____ $_x$ carrying a sword (Alice and Bob)

Cx : _____ $_x$ is looking at a computer (Alice and Cathy)

Is $\exists x Sx$ true? To check this we see if there is a choice of an object for x which gives us a variable assignment under which Sx is true. Consider assigning Alice to x , which we can as shorthand write by $x \mapsto \text{Alice}$. Under this variable assignment, Sx is true: Alice does have a sword. So $\exists x Sx$ is true. There is a choice of an object in our domain for x under which Sx is true.

What about $\forall x Sx$? This is true iff Sx is true under any choice of a person for x . Let's go through them.

	Sx
$x \mapsto \text{Alice}$	T
$x \mapsto \text{Bob}$	T
$x \mapsto \text{Cathy}$	F
$x \mapsto \text{Denny}$	F

So $\forall x Sx$ is false: it is not the case that Sx is true under any choice of an object for x : when we have an assignment of Cathy to x , Sx is false.


What about $\forall x(Sx \rightarrow (Hx \vee Cx))$

	Sx	Hx	Cx	$Hx \vee Cx$	$Sx \rightarrow (Hx \vee Cx)$
$x \mapsto \text{Alice}$	T	F	T	T	T
$x \mapsto \text{Bob}$	T	T	F	T	T
$x \mapsto \text{Cathy}$	F	F	T	T	T
$x \mapsto \text{Denny}$	F	F	F	F	T

So $Sx \rightarrow (Hx \vee Cx)$ is true under every assignment of an object to the variable x . And so $\forall x(Sx \rightarrow (Hx \vee Cx))$ is true.

We have to tread more carefully once we start having multiple quantifiers. Let's walk through some cases.

Consider a new interpretation:



domain: Numbers 1, 2, 3 and 4.

Rxy : There is an arrow from $______x$ to $______y$ in the diagram.

Sx : There is a square around $______x$ in the diagram.

Is $\exists x \forall y Rxy$ true? We need to find some choice of an object for x where $\forall y Rxy$ is true under that choice ("variable assignment"). Let's (with foresight) chose the number 3 for x ($x \mapsto 3$). Is $\forall y Rxy$ true under this variable assignment $x \mapsto 3$? To check this we need to do something more with our variable assignment: we need to extend it with a choice of an object for y . Moreover, we need to think about all ways of picking an object for y , while we've fixed x as the number 3. Consider, e.g. $y \mapsto 1$. Then we have to evaluate whether Rxy is true with $x \mapsto 3, y \mapsto 1$. This is true iff there is an arrow from 3 (x) to 1 (y),

and there is such an arrow. We can work through all the cases and see that all of them have an arrow from 3, so Rxy is true for any choice of object for y :

	Rxy
$x \mapsto 3 \quad y \mapsto 1$	T
$x \mapsto 3 \quad y \mapsto 2$	T
$x \mapsto 3 \quad y \mapsto 3$	T
$x \mapsto 3 \quad y \mapsto 4$	T

So under any way of extending our variable assignment of $x \mapsto 3$ by choosing an object for y results in a variable assignment on which Rxy true. This tells us that $\forall y Rxy$ is true under the variable assignment $x \mapsto 3$. And *that* tells us that $\exists x \forall y Rxy$ is true: there's an assignment of the variable x under which the constituent formula $\forall y Rxy$ is true.

What about $\exists x \exists y (Rxy \ \& \ Ryx)$? To show that it is true we will want to choose an object that we can assign to x under which $\exists y (Rxy \ \& \ Ryx)$ is true. Let's consider $x \mapsto 3$ (again I'm using my foresight of what will come to choose carefully). Now is $\exists y (Rxy \ \& \ Ryx)$ true under the variable assignment $x \mapsto 3$? We need to find an extension of this which chooses an object for y under which $Rxy \ \& \ Ryx$ is true. Consider $y \mapsto 3$. We now have a variable assignment $x \mapsto 3, y \mapsto 3$. They are different variables but there's nothing stopping them denoting the same object. And we can then consider whether $Rxy \ \& \ Ryx$ is true under this interpretation. Well, Rxy is true: 3 does have an arrow to 3. And Ryx is also true: 3 does have an arrow to 3. So by our clause for $\&$, $Rxy \ \& \ Ryx$ is true under this variable assignment $x \mapsto 3, y \mapsto 3$. And so $\exists y (Rxy \ \& \ Ryx)$ is true under the variable assignment $x \mapsto 3$. And so $\exists x \exists y (Rxy \ \& \ Ryx)$ is true in this interpretation.

One more example: $\forall x (Sx \rightarrow \exists y Rxy)$? To check this is true we will need to go through each of our objects for x and see that $Sx \rightarrow \exists y Rxy$ is true under that interpretation. We can already see if Sx is true under each variable assignment, and if we find that Sx is false that's enough information to determine that $Sx \rightarrow \exists y Rxy$ is true (check the definition of truth for \rightarrow to see this):

	Sx	$\exists y Rxy$	$Sx \rightarrow \exists y Rxy$
$x \mapsto 1$	T	?	?
$x \mapsto 2$	F	?	T
$x \mapsto 3$	T	?	?
$x \mapsto 4$	F	?	T

So we need to check whether $\exists y Rxy$ is true under the variable assignments $x \mapsto 1$ and $x \mapsto 3$.

Consider $x \mapsto 1$. We can find an object for y where Rxy is true under that variable assignment: consider $y \mapsto 2$. Since there is an arrow from 1 to 2, Rxy is true in the variable assignment $x \mapsto 1$ and $y \mapsto 2$. Thus $\exists y Rxy$ is true on the variable assignment $x \mapsto 1$. We're also able to do something similar for $x \mapsto 3$:

	Rxy
$x \mapsto 1 \quad y \mapsto 2$	T
$x \mapsto 3 \quad y \mapsto 3$	T

So we now have

	Sx	$\exists y Rxy$	$Sx \rightarrow \exists y Rxy$
$x \mapsto 1$	T	T	T
$x \mapsto 2$	F	?	T
$x \mapsto 3$	T	T	T
$x \mapsto 4$	F	?	T

So $\forall x (Sx \rightarrow \exists y Rxy)$ is true. Informally we might say this as: for every number that has a square around it has an arrow going out of it.

One final example: $\forall x \forall y Rxy$. To check this we will need to consider all choices for x and all choices for y and check Rxy is true on all of them. There are 16 such choices. But we won't have to go through them all: it'll be false. Consider $x \mapsto 1$ and $y \mapsto 4$. Rxy is false under this variable assignment: there is no arrow from 1 to 4. Thus $\forall y Rxy$ is false on the variable assignment $x \mapsto 1$. And so $\forall x \forall y Rxy$ is false in the interpretation.

Let's now give a formal definition of the idea we've been using here. Quantified formulas like $\exists y Rxy$ still need to be given truth conditions *relative to a variable assignment*, because we'll need to specify an assignment of an object for the free variable x . So we define when $\exists v X$ is true under a variable assignment α , which might be, e.g. $x \mapsto 1$. To give a general definition, though we might also consider whether $\forall y Rxy$ is true under a variable assignment $x \mapsto 1, y \mapsto 4$. This is slightly odd: we're considering $\exists y Rxy$ but have been told who y refers to already. However when we evaluate it we simply ignore whatever our given variable assignment tells us about y : we consider variable assignments that modify the assignment by changing what is assigned to y . And by modifying it to $x \mapsto 1, y \mapsto 2$ we have a variable assignment under which

Rxy is true, so $\exists y Rxy$ is true under the original variable assignment $x \mapsto 1, y \mapsto 4$.

Similarly consider $\exists x Sx$ under the variable assignment $x \mapsto 1$. To evaluate this we consider modification of this variable assignment which assign other objects to x . Under the variable assignment $x \mapsto 2$, Sx is true. So under our original variable assignment $\exists x Sx$ is true: it didn't matter what our original variable assignment was, we ignored this and considered variants to evaluate its truth.

This is a general feature: Sentences, which have all variables bound, have truth values independent of any variable assignment they're evaluated with: when we have a quantifier like $\exists x$ or $\forall x$ we ignore whatever our original variable assignment told us about x . So when all our variables are bound by quantifiers, all the original components of our variable assignment are ignored. To summarise: Sentences are simply true or false in interpretations, variable assignments don't matter.

Now for our formal definition:

$\forall v X$ is true under a variable assignment α
iff X is true under *every* variable assignment that is the result of modifying/extending α with a choice of an object in our domain for v .

$\exists v X$ is true under a variable assignment α
iff X is true under *some* variable assignment that is the result of modifying/extending α with a choice of an object in our domain for v .

To be clear: all this is doing is formalizing the intuitive idea expressed in our examples. The result is a bit ugly, and the final definition might look a bit opaque. Hopefully, though, the *spirit* of the idea is clear.

Practice exercises

A. Consider the following interpretation:

- The domain comprises only Corwin and Benedict
- Ax is to be true of both Corwin and Benedict
- Bx is to be true of Benedict only
- Nx is to be true of no one
- c is to refer to Corwin

Determine whether each of the following sentences is true or false in that interpretation:

1. Bc
2. $Ac \leftrightarrow \neg Nc$
3. $Nc \rightarrow (Ac \vee Bc)$
4. $\forall xAx$
5. $\forall x\neg Bx$
6. $\exists x(Ax \& Bx)$
7. $\exists x(Ax \rightarrow Nx)$
8. $\forall x(Nx \vee \neg Nx)$
9. $\exists xBx \rightarrow \forall xAx$


B. Consider the following interpretation:

- The domain comprises only Lemmy, Courtney and Eddy
- Gx is to be true of Lemmy, Courtney and Eddy.
- Hx is to be true of and only of Courtney
- Mx is to be true of and only of Lemmy and Eddy
- c is to refer to Courtney
- e is to refer to Eddy

Determine whether each of the following sentences is true or false in that interpretation:

1. Hc
2. He
3. $Mc \vee Me$
4. $Gc \vee \neg Gc$
5. $Mc \rightarrow Gc$
6. $\exists xHx$
7. $\forall xHx$
8. $\exists x\neg Mx$
9. $\exists x(Hx \& Gx)$
10. $\exists x(Mx \& Gx)$
11. $\forall x(Hx \vee Mx)$
12. $\exists xHx \& \exists xMx$
13. $\forall x(Hx \leftrightarrow \neg Mx)$
14. $\exists xGx \& \exists x\neg Gx$
15. $\forall x\exists y(Gx \& Hy)$

C. Following the diagram conventions introduced at the end of §33, consider the following interpretation:



Determine whether each of the following sentences is true or false in that interpretation:

1. $\exists x Rxx$
2. $\forall x Rxx$
3. $\exists x \forall y Rxy$
4. $\exists x \forall y Ryx$
5. $\forall x \forall y \forall z ((Rxy \ \& \ Ryz) \rightarrow Rxz)$
6. $\forall x \forall y \forall z ((Rxy \ \& \ Rxz) \rightarrow Ryz)$
7. $\exists x \forall y \neg Rxy$
8. $\forall x (\exists y Rxy \rightarrow \exists y Ryx)$
9. $\exists x \exists y (\neg x = y \ \& \ Rxy \ \& \ Ryx)$
10. $\exists x \forall y (Rxy \leftrightarrow x = y)$
11. $\exists x \forall y (Ryx \leftrightarrow x = y)$
12. $\exists x \exists y (\neg x = y \ \& \ Rxy \ \& \ \forall z (Rzx \leftrightarrow y = z))$

CHAPTER 35

Semantic concepts

Offering a precise definition of truth in FOL was more than a little fiddly, but now that we are done, we can define various central logical notions. These will look very similar to the definitions we offered for TFL. However, remember that they concern *interpretations*, rather than valuations.

$$X_1, X_2, \dots, X_n \therefore Z$$

is **VALID** iff there is no interpretation in which all of X_1, X_2, \dots, X_n are true and in which Z is false.

The other logical notions also have corresponding definitions in FOL:

- An FOL sentence X is a **LOGICAL TRUTH** iff X is true in every interpretation.
- X is a **CONTRADICTION** iff X is false in every interpretation.
- Two FOL sentences X and Y are **LOGICALLY EQUIVALENT** iff they are true in exactly the same interpretations as each other.
- The FOL sentences X_1, X_2, \dots, X_n are **JOINTLY LOGICALLY CONSISTENT** iff there is some interpretation in which all of the sentences are true. They are **JOINTLY LOGICALLY INCONSISTENT** iff there is no such interpretation.

CHAPTER 36

Using interpretations

36.1 Logical truths and contradictions

Suppose we want to show that $\exists x Axx \rightarrow Bd$ is *not* a logical truth. This requires showing that the sentence is not true in every interpretation; i.e., that it is false in some interpretation. If we can provide just one interpretation in which the sentence is false, then we will have shown that the sentence is not a logical truth.

In order for $\exists x Axx \rightarrow Bd$ to be false, the antecedent ($\exists x Axx$) must be true, and the consequent (Bd) must be false. To construct such an interpretation, we start by specifying a domain. Keeping the domain small makes it easier to specify what the predicates will be true of, so we will start with a domain that has just one member. For concreteness, let's say it is the city of Paris.

domain: Paris

The name d must refer to something in the domain, so we have no option but:

d : Paris

Recall that we want $\exists x Axx$ to be true, so we want all members of the domain to be paired with themselves in the extension of A . We can just offer:

Axy : _____ _{x} is identical with _____ _{y}

Now Add is true, so it is surely true that $\exists x Axx$. Next, we want Bd to be false, so the referent of d must not be in the extension of B . We might simply offer:

Bx : _____ _{x} is in Germany

Now we have an interpretation where $\exists x Axx$ is true, but where Bd is false. So there is an interpretation where $\exists x Axx \rightarrow Bd$ is false. So $\exists x Axx \rightarrow Bd$ is not a logical truth.

We can just as easily show that $\exists x Axx \rightarrow Bd$ is not a contradiction. We need only specify an interpretation in which $\exists x Axx \rightarrow Bd$ is true; i.e., an interpretation in which either $\exists x Axx$ is false or Bd is true. Here is one:

domain: Paris

d : Paris

Axy : _____ _{x} is identical with _____ _{y}

Bx : _____ _{x} is in France

This shows that there is an interpretation where $\exists x Axx \rightarrow Bd$ is true. So $\exists x Axx \rightarrow Bd$ is not a contradiction.

36.2 Logical equivalence

Suppose we want to show that $\forall x Sx$ and $\exists x Sx$ are not logically equivalent. We need to construct an interpretation in which the two sentences have different truth values; we want one of them to be true and the other to be false. We start by specifying a domain. Again, we make the domain small so that we can specify extensions easily. In this case, we will need at least two objects. (If we chose a domain with only one member, the two sentences would end up with the same truth value. In order to see why, try constructing some partial interpretations with one-member domains.) For concreteness, let's take:

domain: Ornette Coleman, Miles Davis

We can make $\exists x Sx$ true by including something in the extension of S , and we can make $\forall x Sx$ false by leaving something out of the extension of S . For concreteness we will offer:

Sx : _____ _{x} plays saxophone

Now $\exists xSx$ is true, because Sx is true of Ornette Coleman. Slightly more precisely, extend our interpretation by allowing c to name Ornette Coleman. Sc is true in this extended interpretation, so $\exists xSx$ was true in the original interpretation. Similarly, $\forall xSx$ is false, because Sx is false of Miles Davis. Slightly more precisely, extend our interpretation by allowing d to name Miles Davis, and Sd is false in this extended interpretation, so $\forall xSx$ was false in the original interpretation. We have provided a counter-interpretation to the claim that $\forall xSx$ and $\exists xSx$ are logically equivalent.

To show that X is not a logical truth, it suffices to find an interpretation where X is false.

To show that X is not a contradiction, it suffices to find an interpretation where X is true.

To show that X and Y are not logically equivalent, it suffices to find an interpretation where one is true and the other is false.

36.3 Validity, entailment and consistency

To test for validity, entailment, or consistency, we typically need to produce interpretations that determine the truth value of several sentences simultaneously.

Consider the following argument in FOL:

$$\exists x(Gx \rightarrow Ga) \therefore \exists xGx \rightarrow Ga$$

To show that this is invalid, we must make the premise true and the conclusion false. The conclusion is a conditional, so to make it false, the antecedent must be true and the consequent must be false. Clearly, our domain must contain two objects. Let's try:

domain: Karl Marx, Ludwig von Mises

Gx : _____ _{x} hated communism

a : Karl Marx

Given that Marx wrote *The Communist Manifesto*, Ga is plainly false in this interpretation. But von Mises famously hated communism, so $\exists xGx$ is true in this interpretation. Hence $\exists xGx \rightarrow Ga$ is false, as required.

Does this interpretation make the premise true? Yes it does! Note that $Ga \rightarrow Ga$ is true. (Indeed, it is a logical truth.) But then certainly

$\exists x(Gx \rightarrow Ga)$ is true, so the premise is true, and the conclusion is false, in this interpretation. The argument is therefore invalid.

In passing, note that we have also shown that $\exists x(Gx \rightarrow Ga)$ does *not* entail $\exists xGx \rightarrow Ga$. Equally, we have shown that the sentences $\exists x(Gx \rightarrow Ga)$ and $\neg(\exists xGx \rightarrow Ga)$ are jointly consistent.

Let's consider a second example. Consider:

$$\forall x\exists yLxy \therefore \exists y\forall xLxy$$


Again, we want to show that this is invalid. To do this, we must make the premises true and the conclusion false. Here is a suggestion:

domain: UK citizens currently in a civil partnership with another UK citizen

Lxy : ______x is in a civil partnership with ______y

The premise is clearly true on this interpretation. Anyone in the domain is a UK citizen in a civil partnership with some other UK citizen. That other citizen will also, then, be in the domain. So for everyone in the domain, there will be someone (else) in the domain with whom they are in a civil partnership. Hence $\forall x\exists yLxy$ is true. However, the conclusion is clearly false, for that would require that there is some single person who is in a civil partnership with everyone in the domain, and there is no such person, so the argument is invalid. We observe immediately that the sentences $\forall x\exists yLxy$ and $\neg\exists y\forall xLxy$ are jointly consistent and that $\forall x\exists yLxy$ does not entail $\exists y\forall xLxy$.

For our third example, we'll mix things up a bit. In §33, we described how we can present some interpretations using diagrams. For example:



Using the conventions employed in §33, the domain of this interpretation is the first three positive whole numbers, and Rxy is true of x and y just in case there is an arrow from x to y in our diagram. Here are some sentences that the interpretation makes true:

- $\forall x\exists yRyx$

- $\exists x \forall y Rxy$ witness 1
- $\exists x \forall y (Ryx \leftrightarrow x = y)$ witness 1
- $\exists x \exists y \exists z ((\neg y = z \ \& \ Rxy) \ \& \ Rzx)$ witness 2
- $\exists x \forall y \neg Rxy$ witness 3
- $\exists x (\exists y Ryx \ \& \ \neg \exists y Rxy)$ witness 3

This immediately shows that all of the preceding six sentences are jointly consistent. We can use this observation to generate *invalid* arguments, e.g.:

$$\begin{aligned} & \forall x \exists y Ryx, \exists x \forall y Rxy \therefore \forall x \exists y Rxy \\ & \exists x \forall y Rxy, \exists x \forall y \neg Rxy \therefore \neg \exists x \exists y \exists z (\neg y = z \ \& \ (Rxy \ \& \ Rzx)) \end{aligned}$$

and many more besides.

To show that $X_1, X_2, \dots, X_n \therefore Z$ is invalid, it suffices to find an interpretation where all of X_1, X_2, \dots, X_n are true and where Z is false.

That same interpretation will show that X_1, X_2, \dots, X_n do not entail Z .

It will also show that $X_1, X_2, \dots, X_n, \neg Z$ are jointly consistent.

When you provide an interpretation to refute a claim—to logical truth, say, or to entailment—this is sometimes called providing a *counter-interpretation* (or providing a *counter-model*).

Practice exercises

A. Show that each of the following is neither a logical truth nor a contradiction:

1. $Da \ \& \ Db$
2. $\exists x T x h$
3. $Pm \ \& \ \neg \forall x P x$
4. $\forall z J z \leftrightarrow \exists y J y$
5. $\forall x (W x m n \vee \exists y L x y)$
6. $\exists x (G x \rightarrow \forall y M y)$
7. $\exists x (x = h \ \& \ x = i)$

B. Show that the following pairs of sentences are not logically equivalent.

1. Ja, Ka
2. $\exists x Jx, Jm$
3. $\forall x Rxx, \exists x Rxx$
4. $\exists x Px \rightarrow Qc, \exists x (Px \rightarrow Qc)$
5. $\forall x (Px \rightarrow \neg Qx), \exists x (Px \& \neg Qx)$
6. $\exists x (Px \& Qx), \exists x (Px \rightarrow Qx)$
7. $\forall x (Px \rightarrow Qx), \forall x (Px \& Qx)$
8. $\forall x \exists y Rxy, \exists x \forall y Rxy$
9. $\forall x \exists y Rxy, \forall x \exists y Ryx$

C. Show that the following sentences are jointly consistent:

1. $Ma, \neg Na, Pa, \neg Qa$
2. $Lee, Leg, \neg Lge, \neg Lgg$
3. $\neg (Ma \& \exists x Ax), Ma \vee Fa, \forall x (Fx \rightarrow Ax)$
4. $Ma \vee Mb, Ma \rightarrow \forall x \neg Mx$
5. $\forall y Gy, \forall x (Gx \rightarrow Hx), \exists y \neg Iy$
6. $\exists x (Bx \vee Ax), \forall x \neg Cx, \forall x [(Ax \& Bx) \rightarrow Cx]$
7. $\exists x Xx, \exists x Yx, \forall x (Xx \leftrightarrow \neg Yx)$
8. $\forall x (Px \vee Qx), \exists x \neg (Qx \& Px)$
9. $\exists z (Nz \& Oz), \forall x \forall y (Oxy \rightarrow Oyx)$
10. $\neg \exists x \forall y Rxy, \forall x \exists y Rxy$
11. $\neg Raa, \forall x (x = a \vee Rxa)$
12. $\forall x \forall y \forall z [(x = y \vee y = z) \vee x = z], \exists x \exists y \neg x = y$
13. $\exists x \exists y ((Zx \& Zy) \& x = y), \neg Zd, d = e$

D. Show that the following arguments are invalid:

1. $\forall x (Ax \rightarrow Bx) \therefore \exists x Bx$
2. $\forall x (Rx \rightarrow Dx), \forall x (Rx \rightarrow Fx) \therefore \exists x (Dx \& Fx)$
3. $\exists x (Px \rightarrow Qx) \therefore \exists x Px$
4. $Na \& Nb \& Nc \therefore \forall x Nx$
5. $Rde, \exists x Rxd \therefore Red$
6. $\exists x (Ex \& Fx), \exists x Fx \rightarrow \exists x Gx \therefore \exists x (Ex \& Gx)$
7. $\forall x Oxc, \forall x Ocx \therefore \forall x Oxx$
8. $\exists x (Jx \& Kx), \exists x \neg Kx, \exists x \neg Jx \therefore \exists x (\neg Jx \& \neg Kx)$
9. $Lab \rightarrow \forall x Lxb, \exists x Lxb \therefore Lbb$
10. $\forall x (Dx \rightarrow \exists y Tyx) \therefore \exists y \exists z \neg y = z$

CHAPTER 37

Reasoning about all interpretations

37.1 Logical truths and contradictions

We can show that a sentence is *not* a logical truth just by providing one carefully specified interpretation: an interpretation in which the sentence is false. To show that something is a logical truth, on the other hand, it would not be enough to construct ten, one hundred, or even a thousand interpretations in which the sentence is true. A sentence is only a logical truth if it is true in *every* interpretation, and there are infinitely many interpretations. We need to reason about all of them, and we cannot do this by dealing with them one by one!

Sometimes, we can reason about all interpretations fairly easily. For example, we can offer a relatively simple argument that $Raa \leftrightarrow Raa$ is a logical truth:

Any relevant interpretation will give Raa a truth value. If Raa is true in an interpretation, then $Raa \leftrightarrow Raa$ is true in that interpretation. If Raa is false in an interpretation, then $Raa \leftrightarrow Raa$ is true in that interpretation. These are the only alternatives. So $Raa \leftrightarrow Raa$ is true in every interpretation. Therefore, it is a logical truth.

This argument is valid, of course, and its conclusion is true. However, it is not an argument in FOL. Rather, it is an argument in English *about* FOL: it is an argument in the metalanguage.

Note another feature of the argument. Since the sentence in question contained no quantifiers, we did not need to think about how to interpret a and R ; the point was just that, however we interpreted them, Raa would have some truth value or other. (We could ultimately have given the same argument concerning TFL sentences.)

Here is another bit of reasoning. Consider the sentence $\forall x(Rxx \leftrightarrow Rxx)$. Again, it should obviously be a logical truth, but to say precisely why is quite a challenge. We cannot say that $Rxx \leftrightarrow Rxx$ is true in every interpretation, since $Rxx \leftrightarrow Rxx$ is not even a *sentence* of FOL (remember that x is a variable, not a name). So we have to be a bit cleverer.

Consider some arbitrary interpretation. Consider some arbitrary member of the domain, which, for convenience, we will call *obbie*, and suppose we extend our original interpretation by adding a new name, c , to name *obbie*. Then either Rcc will be true or it will be false. If Rcc is true, then $Rcc \leftrightarrow Rcc$ is true. If Rcc is false, then $Rcc \leftrightarrow Rcc$ will be true. So either way, $Rcc \leftrightarrow Rcc$ is true. Since there was nothing special about *obbie*—we might have chosen any object—we see that no matter how we extend our original interpretation by allowing c to name some new object, $Rcc \leftrightarrow Rcc$ will be true in the new interpretation. So $\forall x(Rxx \leftrightarrow Rxx)$ was true in the original interpretation. But we chose our interpretation arbitrarily, so $\forall x(Rxx \leftrightarrow Rxx)$ is true in every interpretation. It is therefore a logical truth.

This is quite longwinded, but, as things stand, there is no alternative. In order to show that a sentence is a logical truth, we must reason about *all* interpretations.

37.2 Other cases

Similar points hold of other cases too. Thus, we must reason about all interpretations if we want to show:

- that a sentence is a contradiction; for this requires that it is false in *every* interpretation.

- that two sentences are logically equivalent; for this requires that they have the same truth value in *every* interpretation.
- that some sentences are jointly inconsistent; for this requires that there is no interpretation in which all of those sentences are true together; i.e. that, in *every* interpretation, at least one of those sentences is false.
- that an argument is valid; for this requires that the conclusion is true in *every* interpretation where the premises are true.
- that some sentences entail another sentence.

The problem is that, with the tools available to you so far, reasoning about all interpretations is a serious challenge! Let's take just one more example. Here is an argument which is obviously valid:

$$\forall x(Hx \& Jx) \therefore \forall xHx$$

After all, if everything is both H and J, then everything is H. But we can only show that the argument is valid by considering what must be true in every interpretation in which the premise is true. To show this, we would have to reason as follows:

Consider an arbitrary interpretation in which the premise $\forall x(Hx \& Jx)$ is true. It follows that, however we expand the interpretation with a new name, for example c , $Hc \& Jc$ will be true in this new interpretation. Hc will, then, also be true in this new interpretation. But since this held for *any* way of expanding the interpretation, it must be that $\forall xHx$ is true in the old interpretation. We've assumed nothing about the interpretation except that it was one in which $\forall x(Hx \& Jx)$ is true, so any interpretation in which $\forall x(Hx \& Jx)$ is true is one in which $\forall xHx$ is true. The argument is valid!

Even for a simple argument like this one, the reasoning is somewhat complicated. For longer arguments, the reasoning can be extremely torturous.

The following table summarises whether a single (counter-)interpretation suffices, or whether we must reason about all interpretations.

	Yes	No
logical truth?	all interpretations	one counter-interpretation
contradiction?	all interpretations	one counter-interpretation
equivalent?	all interpretations	one counter-interpretation
consistent?	one interpretation	all interpretations
valid?	all interpretations	one counter-interpretation
entailment?	all interpretations	one counter-interpretation

This might usefully be compared with the table at the end of §16. The key difference resides in the fact that TFL concerns truth tables, whereas FOL concerns interpretations. This difference is deeply important, since each truth-table only ever has finitely many lines, so that a complete truth table is a relatively tractable object. By contrast, there are infinitely many interpretations for any given sentence(s), so that reasoning about all interpretations can be a deeply tricky business.

PART VIII

Natural deduction for FOL

CHAPTER 38

Basic rules for FOL

FOL makes use of all of the connectives of TFL. So proofs in FOL will use all of the basic and derived rules from Part IV. We will also use the proof-theoretic notions (particularly, the symbol ‘ \vdash ’) introduced there. However, we will also need some new basic rules to govern the quantifiers.

38.1 Universal elimination

Consider:

- Everyone is happy
- Therefore: Catrin is happy.

This is a valid argument. Generally, then, from the claim that everything is F, you can infer that any particular thing is F. You name it; it’s F. So the following should be fine:

1	$\forall xFx$	
2	Fa	$\forall E$ 1

We obtained line 2 by dropping the universal quantifier and replacing ‘ x ’ with ‘ a ’.

This isn’t restricted to simple properties. Consider the following argument:

- Every cat is sleeping.
- Therefore: If Fluffy is a cat, then she is sleeping.

This is a valid argument. And it will be allowed by our rule $\forall E$:

1	$\forall x(Cx \rightarrow Sx)$	
2	$Cf \rightarrow Sf$	$\forall E$ 1

Note here that we have to replace two instances of ‘ x ’ with our name: ‘ f ’ or Fluffy. Indeed it would have been fine to do with any name. We can even do it with names we already have. Consider the following:

- Pavel owes everyone money.
- Therefore: Pavel owes himself money.

We symbolise this as:

- $\forall xOpx$
- Therefore: Opp

This is valid: the premise says *everything* in the domain owes money to Pavel; and Pavel is something in the domain. So it implies that Pavel owes money to himself. A closely related sentence is:

1. Pavel owes money to everyone *else*.

and that will not be symbolised as $\forall xOxx$; however, we do not yet have the resources to symbolise at. In §IX we will introduce identity which will allow us to formalise it properly.

This argument is also directly allowed by our rule $\forall E$:

1	$\forall xOpx$	
2	Opp	$\forall E$ 1

We can now give our general rule using the notation from §34: Whenever you have a sentence $\forall xX(\dots x \dots x \dots)$, for example $\forall xFx$, $\forall x(Cx \rightarrow Sx)$, $\forall xOpx$; one can conclude that we have the sentence which is obtained by stripping of the quantifier and replacing the variable by a name, be it $a, b, c \dots$. So we could derive Fa , $Cf \rightarrow Sf$ or Opp .

Here is the formal specification of the universal elimination rule ($\forall E$):

m	$\forall x X(\dots x \dots x \dots)$	
	$X(\dots c \dots c \dots)$	$\forall E\ m$

The point is that you can obtain any *substitution instance* of a universally quantified formula: replace every instance of the quantified variable with any name you like.

I should emphasize that (as with every elimination rule) you can only apply the $\forall E$ rule when the universal quantifier is the main logical operator. Thus the following is outright banned:

1	$\forall x Bx \rightarrow Bk$	
2	$Bb \rightarrow Bk$	naughtily attempting to invoke $\forall E\ 1$

This is illegitimate, since ‘ $\forall x$ ’ is not the main logical operator in line 1. (If you need a reminder as to why this sort of inference should be banned, reread §29.)

38.2 Existential introduction

The following argument is valid:

- Catrin is happy
- Therefore: Someone is happy.

This is the idea of our existential introduction rule: from the claim that some particular thing is F, you can infer that something is F:

1	Fa	
2	$\exists x Fx$	$\exists I\ 1$

We obtained line 2 by replacing the name ‘ a ’ with the variable ‘ x ’ and adding $\exists x$ in front of the sentence. This will be permissible by our rule of $\exists I$.

This isn’t restricted to simple properties.

- Bob is a money and knows sign language.
- Therefore: There is a monkey who knows sign language.

1		$Mb \ \& \ Sb$	
2		$\exists x(Mx \ \& \ Sx)$	$\exists I \ 1$

Or even

- Catrin is friends with someone who is friends with everyone.
- Therefore: Someone is friends with someone who is friends with everyone.

1		$\exists x(Fcx \ \& \ \forall yFxy)$	
2		$\exists z\exists x(Fzx \ \& \ \forall yFxy)$	$\exists I \ 1$

We replaced the name, ‘*c*’ with the variable ‘*z*’ and added $\exists z$ at the beginning of the sentence.

This rule will now allow us to carefully work through our brain teaser from §20.2

Three people are standing in a row looking at eachother.



Alice



Bob



Charlie

Alice is happy. Charlie is not happy. Is there someone who is happy who is looking at someone who is not happy?

Answer: Yes.

We can formalise this argument as:

$$Lab, Lbc, Ha, \neg Hc \ \therefore \ \exists x\exists y(Hx \ \& \ (Lxy \ \& \ \neg Hx)).$$

And we can show it is valid. In §20.2 we wrote this in a pseudo-formal style.

1	Bob is either happy or he's not happy	
2	<table><tr><td>Suppose Bob is happy</td></tr></table>	Suppose Bob is happy
Suppose Bob is happy		
3	Then happy Bob is looking at unhappy Charlie	
4	So someone who is happy is looking at someone who is not happy.	
5	<table><tr><td>Suppose Bob is unhappy</td></tr></table>	Suppose Bob is unhappy
Suppose Bob is unhappy		
6	Then happy Alice is looking at unhappy Bob	
7	So someone who is happy is looking at someone who is not happy.	
8	Therefore, someone who is happy is looking at someone who is not happy.	

We can now fill out the details of this to see that it's a formal proof:

1		Lab	
2		Lbc	
3		Ha	
4		$\neg Hc$	
5		$Hb \vee \neg Hb$	LEM
6			Hb
7			$Lbc \& \neg Hc$ &I 2, 6
8			$Hb \& (Lbc \& \neg Hc)$ &I 7, 6
9			$\exists y(Hb \& (Lby \& \neg Hy))$ \exists I 8
10			$\exists x \exists y(Hx \& (Lxy \& \neg Hx))$ \exists I 9
11			$\neg Hb$
12			$Lab \& \neg Hb$ &I 11, 1
13			$Ha \& (Lab \& \neg Hb)$ &I 3, 12
14			$\exists y(Ha \& (Lay \& \neg Hy))$ \exists I 13
15			$\exists x \exists y(Hx \& (Lxy \& \neg Hx))$ \exists I 14
16		$\exists x \exists y(Hx \& Lxy \& \neg Hx)$	\vee E 5, 6–10, 11–15

Consider the following example:

- Narcissus loves himself.
- Therefore: There is someone who loves Narcissus.

This is a valid argument. The formalised version, which will be allowed by our rule is:

1		Lnn	
2		$\exists x Lxn$	\exists I 1

This shows us that we do not have to replace *all* instances of the name with the variable. Though of course we can if we wish: we could also deduce there is someone who loves himself.

To give our rule in general we need to introduce some new notation for this ability to replace just some of our instances of the name: If X is a sentence containing the name c , we can emphasize this by writing ' $X(\dots c \dots c \dots)$ '. We will write ' $X(\dots x \dots c \dots)$ ' to indicate any formula obtained by replacing *some or all* of the instances of the name c with the variable x . Armed with this, our introduction rule is:

m	$X(\dots c \dots c \dots)$ $\exists x X(\dots x \dots c \dots)$	$\exists I\ m$
x must not occur in $X(\dots c \dots c \dots)$		

All the cases we've seen in this section follow this rule.

You might have noticed the additional constraint that's added to the rule. It is part of the rule; so if you are asked to write the rule $\exists I$ you must include this constraint. However, you will not need to worry about it in practice. It's simply there to guarantee that applications of the rule yield sentences of FOL. If the rule were not there we would be allowed to argue as follows:

1	$\exists x Lnx$	
2	$\exists x \exists x Lxx$	naughtily attempting to invoke $\exists I\ 1$

But this expression on line 2 contains clashing variables. It will not count as a sentence of FOL.

38.3 Empty domains

The following proof combines our two new rules for quantifiers:

1	$\forall x Fx$	
2	Fa	$\forall E\ 1$
3	$\exists x Fx$	$\exists I\ 2$

Could this be a bad proof? If anything exists at all, then certainly we can infer that something is F, from the fact that everything is F. But what if *nothing* exists at all? Then it is surely vacuously true that everything

is F; however, it does not follow that something is F, for there is nothing to *be* F. So if we claim that, as a matter of logic alone, ‘ $\exists xFx$ ’ follows from ‘ $\forall xFx$ ’, then we are claiming that, as a matter of *logic alone*, there is something rather than nothing. This might strike us as a bit odd.

Actually, we are already committed to this oddity. In §28, we stipulated that domains in FOL must have at least one member. We then defined a logical truth (of FOL) as a sentence which is true in every interpretation. Since ‘ $\exists x(Ax \vee \neg Ax)$ ’ will be true in every interpretation, this *also* had the effect of stipulating that it is a matter of logic that there is something rather than nothing.

Since it is far from clear that logic should tell us that there must be something rather than nothing, we might well be cheating a bit here.

If we refuse to cheat, though, then we pay a high cost. Here are three things that we want to hold on to:

- $\forall xFx \vdash Fa$: after all, that was $\forall E$.
- $Fa \vdash \exists xFx$: after all, that was $\exists I$.
- the ability to copy-and-paste proofs together: after all, reasoning works by putting lots of little steps together into rather big chains.

If we get what we want on all three counts, then we have to countenance that $\forall xFx \vdash \exists xFx$. So, if we get what we want on all three counts, the proof system alone tells us that there is something rather than nothing. And if we refuse to accept that, then we have to surrender one of the three things that we want to hold on to!

Before we start thinking about which to surrender, we might want to ask how *much* of a cheat this is. Granted, it may make it harder to engage in theological debates about why there is something rather than nothing. But the rest of the time, we will get along just fine. So maybe we should just regard our proof system (and FOL, more generally) as having a very slightly limited purview. If we ever want to allow for the possibility of *nothing*, then we will have to cast around for a more complicated proof system. But for as long as we are content to ignore that possibility, our proof system is perfectly in order. (As, similarly, is the stipulation that every domain must contain at least one object.)

38.4 Universal introduction

Suppose you had shown of each particular thing that it is F (and that there are no other things to consider). Then you would be justified in

claiming that everything is F. This would motivate the following proof rule. If you had established each and every single substitution instance of ‘ $\forall xFx$ ’, then you can infer ‘ $\forall xFx$ ’.

Unfortunately, that rule would be utterly unusable. To establish each and every single substitution instance would require proving ‘ Fa ’, ‘ Fb ’, ..., ‘ Fj_2 ’, ..., ‘ Fr_{79002} ’, ..., and so on. Indeed, since there are infinitely many names in FOL, this process would never come to an end. So we could never apply that rule. We need to be a bit more cunning in coming up with our rule for introducing universal quantification.

Our cunning thought will be inspired by considering:

$$\forall xFx \therefore \forall yFy$$

This argument should *obviously* be valid. After all, alphabetical variation ought to be a matter of taste, and of no logical consequence. But how might our proof system reflect this? Suppose we begin a proof thus:

$$\begin{array}{l|l} 1 & \forall xFx \\ \hline 2 & Fa \qquad \forall E\ 1 \end{array}$$

We have proved ‘ Fa ’. And, of course, nothing stops us from using the same justification to prove ‘ Fb ’, ‘ Fc ’, ..., ‘ Fj_2 ’, ..., ‘ Fr_{79002} ’, ..., and so on until we run out of space, time, or patience. But reflecting on this, we see that there is a way to prove Fc , for any name c . And if we can do it for *any* thing, we should surely be able to say that ‘ F ’ is true of *everything*. This therefore justifies us in inferring ‘ $\forall yFy$ ’, thus:

$$\begin{array}{l|l} 1 & \forall xFx \\ \hline 2 & Fa \qquad \forall E\ 1 \\ 3 & \forall yFy \qquad \forall I\ 2 \end{array}$$

The crucial thought here is that ‘ a ’ was just some *arbitrary* name. There was nothing special about it—we might have chosen any other name—and still the proof would be fine. And this crucial thought motivates the universal introduction rule ($\forall I$):

$$\begin{array}{c|l}
 m & X(\dots c \dots c \dots) \\
 & \forall x X(\dots x \dots x \dots) \quad \forall I \ m
 \end{array}$$

c must not occur in any undischarged assumption
 x must not occur in $X(\dots c \dots c \dots)$

A crucial aspect of this rule, though, is bound up in the first constraint. This constraint ensures that we are always reasoning at a sufficiently general level. To see the constraint in action, consider this terrible argument:

Everyone loves Kylie Minogue; therefore everyone loves themselves.

We might symbolize this obviously invalid inference pattern as:

$$\forall x Lxk \therefore \forall x Lxx$$

Now, suppose we tried to offer a proof that vindicates this argument:

$$\begin{array}{c|l}
 1 & \forall x Lxk \\
 \hline
 2 & Lkk \quad \forall E \ 1 \\
 3 & \forall x Lxx \quad \text{naughtily attempting to invoke } \forall I \ 2
 \end{array}$$

This is not allowed, because ‘ k ’ occurred already in an undischarged assumption, namely, on line 1. The crucial point is that, if we have made any assumptions about the object we are working with, then we are not reasoning generally enough to license $\forall I$.

Although the name may not occur in any *undischarged* assumption, it may occur as a discharged assumption. That is, it may occur in a subproof that we have already closed. For example:

$$\begin{array}{c|l}
 1 & \begin{array}{c|l} & Gd \\ \hline & Gd \end{array} \\
 2 & \quad \quad \quad R \ 1 \\
 3 & Gd \rightarrow Gd \quad \rightarrow I \ 1-2 \\
 4 & \forall z (Gz \rightarrow Gz) \quad \forall I \ 3
 \end{array}$$

This tells us that ' $\forall z(Gz \rightarrow Gz)$ ' is a *theorem*. And that is as it should be.

38.5 Existential elimination

Suppose we know that *something* is F. The problem is that simply knowing this does not tell us which thing is F. So it would seem that from ' $\exists xFx$ ' we cannot immediately conclude ' Fa ', ' $F_{e_{23}}$ ', or any other substitution instance of the sentence. What can we do?

Suppose we know that something is F, and that everything which is F is G. In (almost) natural English, we might reason thus:

Since something is F, there is some particular thing which is an F. We do not know anything about it, other than that it's an F, but for convenience, let's call it 'obbie'. So: obbie is F. Since everything which is F is G, it follows that obbie is G. But since obbie is G, it follows that something is G. And nothing depended on which object, exactly, obbie was. So, something is G.

We might try to capture this reasoning pattern in a proof as follows:

1	$\exists xFx$	
2	$\forall x(Fx \rightarrow Gx)$	
3	<div style="border-left: 1px solid black; padding-left: 10px;">Fo</div>	
4	<div style="border-left: 1px solid black; padding-left: 10px;">$Fo \rightarrow Go$</div>	$\forall E$ 2
5	<div style="border-left: 1px solid black; padding-left: 10px;">Go</div>	$\rightarrow E$ 4, 3
6	<div style="border-left: 1px solid black; padding-left: 10px;">$\exists xGx$</div>	$\exists I$ 5
7	$\exists xGx$	$\exists E$ 1, 3–6

Breaking this down: we started by writing down our assumptions. At line 3, we made an additional assumption: ' Fo '. This was just a substitution instance of ' $\exists xFx$ '. On this assumption, we established ' $\exists xGx$ '. Note that we had made no *special* assumptions about the object named by ' o '; we had *only* assumed that it satisfies ' Fx '. So nothing depends upon which object it is. And line 1 told us that *something* satisfies ' Fx ', so our reasoning pattern was perfectly general. We can discharge the specific assumption ' Fo ', and simply infer ' $\exists xGx$ ' on its own.

Putting this together, we obtain the existential elimination rule ($\exists E$):

m	$\exists x X(\dots x \dots x \dots)$	
i	$X(\dots c \dots c \dots)$	
	\vdots	
j	Y	
	Y	$\exists E\ m, i-j$

c must not occur in any assumption undischarged before line i
 c must not occur in $\exists x X(\dots x \dots x \dots)$
 c must not occur in Y

As with universal introduction, the constraints are extremely important. To see why, consider the following terrible argument:

Tim Button is a lecturer. There is someone who is not a lecturer. So Tim Button is both a lecturer and not a lecturer.

We might symbolize this obviously invalid inference pattern as follows:

$$Lb, \exists x \neg Lx \therefore Lb \ \& \ \neg Lb$$

Now, suppose we tried to offer a proof that vindicates this argument:

1	Lb	
2	$\exists x \neg Lx$	
3	$\neg Lb$	
4	$Lb \ \& \ \neg Lb$	$\&I\ 1, 3$
5	$Lb \ \& \ \neg Lb$	naughtily attempting to invoke $\exists E\ 2, 3-4$

The last line of the proof is not allowed. The name that we used in our substitution instance for ' $\exists x \neg Lx$ ' on line 3, namely ' b ', occurs in line 4. The following proof would be no better:

1		Lb	
2		$\exists x \neg Lx$	
3			$\neg Lb$
4			$Lb \ \& \ \neg Lb$ &I 1, 3
5			$\exists x (Lx \ \& \ \neg Lx)$ \exists I 4
6		$\exists x (Lx \ \& \ \neg Lx)$	naughtily attempting to invoke \exists E 2, 3–5

The last line of the proof would still not be allowed. For the name that we used in our substitution instance for ‘ $\exists x \neg Lx$ ’, namely ‘ b ’, occurs in an undischarged assumption, namely line 1.

The moral of the story is this. *If you want to squeeze information out of an existential quantifier, choose a new name for your substitution instance.* That way, you can guarantee that you meet all the constraints on the rule for \exists E.

Practice exercises

A. The following two ‘proofs’ are *incorrect*. Explain why both are incorrect. Also, provide interpretations which would invalidate the fallacious argument forms the ‘proofs’ enshrine:

1		$\forall x Rxx$		1		$\forall x \exists y Rxy$	
2		Raa	\forall E 1	2		$\exists y Ray$	\forall E 1
3		$\forall y Ray$	\forall I 2	3			Raa
4		$\forall x \forall y Rxy$	\forall I 3	4			$\exists x Rxx$ \exists I 3
				5		$\exists x Rxx$	\exists E 2, 3–4

B. The following three proofs are missing their citations (rule and line numbers). Add them, to turn them into bona fide proofs.

1	$\forall x \exists y (Rxy \vee Ryx)$
2	$\forall x \neg Rmx$
3	$\exists y (Rmy \vee Rym)$
4	$Rma \vee Ram$
5	$\neg Rma$
6	Ram
7	$\exists x Rxm$
8	$\exists x Rxm$

1	$\forall x (\exists y Lxy \rightarrow \forall z Lzx)$
2	Lab
3	$\exists y Lay \rightarrow \forall z Lza$
4	$\exists y Lay$
5	$\forall z Lza$
6	Lca
7	$\exists y Lcy \rightarrow \forall z Lzc$
8	$\exists y Lcy$
9	$\forall z Lzc$
10	Lcc
11	$\forall x Lxx$

1	$\forall x (Jx \rightarrow Kx)$
2	$\exists x \forall y Lxy$
3	$\forall x Jx$
4	$\forall y Lay$
5	Laa
6	Ja
7	$Ja \rightarrow Ka$
8	Ka
9	$Ka \& Laa$
10	$\exists x (Kx \& Lxx)$
11	$\exists x (Kx \& Lxx)$

C. In §29 problem A, we considered fifteen syllogistic figures of Aristotelian logic. Provide proofs for each of the argument forms. NB: You will find it *much* easier if you symbolize (for example) ‘No F is G’ as ‘ $\forall x (Fx \rightarrow \neg Gx)$ ’.

D. Aristotle and his successors identified other syllogistic forms which depended upon ‘existential import’. Symbolize each of the following argument forms in FOL and offer proofs.

- **Barbari.** Something is H. All G are F. All H are G. So: Some H is F
- **Celarent.** Something is H. No G are F. All H are G. So: Some H is not F
- **Cesaro.** Something is H. No F are G. All H are G. So: Some H is not F.
- **Camestros.** Something is H. All F are G. No H are G. So: Some H is not F.
- **Felapton.** Something is G. No G are F. All G are H. So: Some H is not F.
- **Darapti.** Something is G. All G are F. All G are H. So: Some H is F.
- **Calemos.** Something is H. All F are G. No G are H. So: Some H is not F.
- **Fesapo.** Something is G. No F is G. All G are H. So: Some H is not F.
- **Bamalip.** Something is F. All F are G. All G are H. So: Some H are F.

E. Provide a proof of each claim.

1. $\vdash \forall x Fx \vee \neg \forall x Fx$
2. $\vdash \forall z (Pz \vee \neg Pz)$
3. $\forall x (Ax \rightarrow Bx), \exists x Ax \vdash \exists x Bx$
4. $\forall x (Mx \leftrightarrow Nx), Ma \ \& \ \exists x Rxa \vdash \exists x Nx$
5. $\forall x \forall y Gxy \vdash \exists x Gxx$
6. $\vdash \forall x Rxx \rightarrow \exists x \exists y Rxy$
7. $\vdash \forall y \exists x (Qy \rightarrow Qx)$
8. $Na \rightarrow \forall x (Mx \leftrightarrow Ma), Ma, \neg Mb \vdash \neg Na$
9. $\forall x \forall y (Gxy \rightarrow Gyx) \vdash \forall x \forall y (Gxy \leftrightarrow Gyx)$
10. $\forall x (\neg Mx \vee Ljx), \forall x (Bx \rightarrow Ljx), \forall x (Mx \vee Bx) \vdash \forall x Ljx$

F. Write a symbolization key for the following argument, symbolize it, and prove it:

There is someone who likes everyone who likes everyone that she likes. Therefore, there is someone who likes herself.

G. Show that each pair of sentences is provably equivalent.

1. $\forall x (Ax \rightarrow \neg Bx), \neg \exists x (Ax \ \& \ Bx)$
2. $\forall x (\neg Ax \rightarrow Bd), \forall x Ax \vee Bd$
3. $\exists x Px \rightarrow Qc, \forall x (Px \rightarrow Qc)$

H. For each of the following pairs of sentences: If they are provably equivalent, give proofs to show this. If they are not, construct an interpretation to show that they are not logically equivalent.

1. $\forall xPx \rightarrow Qc, \forall x(Px \rightarrow Qc)$
2. $\forall x\forall y\forall zBxyz, \forall xBxxx$
3. $\forall x\forall yDxy, \forall y\forall xDxy$
4. $\exists x\forall yDxy, \forall y\exists xDxy$
5. $\forall x(Rca \leftrightarrow Rxa), Rca \leftrightarrow \forall xRxa$

I. For each of the following arguments: If it is valid in FOL, give a proof. If it is invalid, construct an interpretation to show that it is invalid.

1. $\exists y\forall xRxy \therefore \forall x\exists yRxy$
2. $\forall x\exists yRxy \therefore \exists y\forall xRxy$
3. $\exists x(Px \& \neg Qx) \therefore \forall x(Px \rightarrow \neg Qx)$
4. $\forall x(Sx \rightarrow Ta), Sd \therefore Ta$
5. $\forall x(Ax \rightarrow Bx), \forall x(Bx \rightarrow Cx) \therefore \forall x(Ax \rightarrow Cx)$
6. $\exists x(Dx \vee Ex), \forall x(Dx \rightarrow Fx) \therefore \exists x(Dx \& Fx)$
7. $\forall x\forall y(Rxy \vee Ryx) \therefore Rjj$
8. $\exists x\exists y(Rxy \vee Ryx) \therefore Rjj$
9. $\forall xPx \rightarrow \forall xQx, \exists x\neg Px \therefore \exists x\neg Qx$
10. $\exists xMx \rightarrow \exists xNx, \neg\exists xNx \therefore \forall x\neg Mx$

CHAPTER 39

Conversion of quantifiers

In this section, we will add some additional rules to the basic rules of the previous section. These govern the interaction of quantifiers and negation.

In §28, we noted that $\neg\exists xX$ is logically equivalent to $\forall x\neg X$. We will add some rules to our proof system that govern this. In particular, we add:

$$\begin{array}{c|c} m & \begin{array}{l} \forall x\neg X \\ \neg\exists xX \end{array} \\ \hline & \text{CQ } m \end{array}$$

and

$$\begin{array}{c|c} m & \begin{array}{l} \neg\exists xX \\ \forall x\neg X \end{array} \\ \hline & \text{CQ } m \end{array}$$

Equally, we add:

$$\begin{array}{c|l}
 m & \begin{array}{l} \exists x \neg X \\ \neg \forall x X \end{array} \\
 & \text{CQ } m
 \end{array}$$

and

$$\begin{array}{c|l}
 m & \begin{array}{l} \neg \forall x X \\ \exists x \neg X \end{array} \\
 & \text{CQ } m
 \end{array}$$

Practice exercises

A. Show in each case that the sentences are provably inconsistent:

1. $Sa \rightarrow Tm, Tm \rightarrow Sa, Tm \& \neg Sa$
2. $\neg \exists x Rxa, \forall x \forall y Ryx$
3. $\neg \exists x \exists y Lxy, Laa$
4. $\forall x (Px \rightarrow Qx), \forall z (Pz \rightarrow Rz), \forall y Py, \neg Qa \& \neg Rb$

B. Show that each pair of sentences is provably equivalent:

1. $\forall x (Ax \rightarrow \neg Bx), \neg \exists x (Ax \& Bx)$
2. $\forall x (\neg Ax \rightarrow Bx), \forall x Ax \vee Bx$

C. In §29, we considered what happens when we move quantifiers ‘across’ various logical operators. Show that each pair of sentences is provably equivalent:

1. $\forall x (Fx \& Ga), \forall x Fx \& Ga$
2. $\exists x (Fx \vee Ga), \exists x Fx \vee Ga$
3. $\forall x (Ga \rightarrow Fx), Ga \rightarrow \forall x Fx$
4. $\forall x (Fx \rightarrow Ga), \exists x Fx \rightarrow Ga$
5. $\exists x (Ga \rightarrow Fx), Ga \rightarrow \exists x Fx$
6. $\exists x (Fx \rightarrow Ga), \forall x Fx \rightarrow Ga$

NB: the variable ‘x’ does not occur in ‘Ga’.

When all the quantifiers occur at the beginning of a sentence, that sentence is said to be in *prenex normal form*. These equivalences are sometimes called *prenexing rules*, since they give us a means for putting any sentence into prenex normal form.

CHAPTER 40

Derived rules

As in the case of TFL, we first introduced some rules for FOL as basic (in §38), and then added some further rules for conversion of quantifiers (in §39). In fact, the CQ rules should be regarded as *derived* rules, for they can be derived from the *basic* rules of §38. (The point here is as in §23.) Here is a justification for the first CQ rule:

1	$\forall x \neg Ax$	
2	$\exists x Ax$	
3	Ac	
4	$\neg Ac$	$\forall E$ 1
5	\perp	$\perp I$ 3, 4
6	\perp	$\exists E$ 2, 3–5
7	$\neg \exists x Ax$	$\neg I$ 2–6

Here is a justification of the third CQ rule:

CHAPTER 41

Proof-theoretic and semantic concepts

We have used two different turnstiles in this book. This:

$$X_1, X_2, \dots, X_n \vdash C$$

means that there is some proof which starts with assumptions X_1, X_2, \dots, X_n and ends with C (and no undischarged assumptions other than X_1, X_2, \dots, X_n). This is a *proof-theoretic notion*.

By contrast, this:

$$X_1, X_2, \dots, X_n \models C$$

means that there is no valuation (or interpretation) which makes all of X_1, X_2, \dots, X_n true and makes C false. This concerns assignments of truth and falsity to sentences. It is a *semantic notion*.

It cannot be emphasized enough that these are different notions. But we can emphasize it a bit more: *They are different notions*.

Once you have internalised this point, continue reading.

Although our semantic and proof-theoretic notions are different, there is a deep connection between them. To explain this connection, we will start by considering the relationship between logical truths and theorems.

To show that a sentence is a theorem, you need only produce a proof. Granted, it may be hard to produce a twenty line proof, but it is not so hard to check each line of the proof and confirm that it is legitimate; and if each line of the proof individually is legitimate, then the whole proof is legitimate. Showing that a sentence is a logical truth, though, requires reasoning about all possible interpretations. Given a choice between showing that a sentence is a theorem and showing that it is a logical truth, it would be easier to show that it is a theorem.

Contrawise, to show that a sentence is *not* a theorem is hard. We would need to reason about all (possible) proofs. That is very difficult. However, to show that a sentence is not a logical truth, you need only construct an interpretation in which the sentence is false. Granted, it may be hard to come up with the interpretation; but once you have done so, it is relatively straightforward to check what truth value it assigns to a sentence. Given a choice between showing that a sentence is not a theorem and showing that it is not a logical truth, it would be easier to show that it is not a logical truth.

Fortunately, *a sentence is a theorem if and only if it is a logical truth*. As a result, if we provide a proof of X on no assumptions, and thus show that X is a theorem, we can legitimately infer that X is a logical truth; i.e., $\models X$. Similarly, if we construct an interpretation in which X is false and thus show that it is not a logical truth, it follows that X is not a theorem.

More generally, we have the following powerful result:

$$X_1, X_2, \dots, X_n \vdash Y \text{ iff } X_1, X_2, \dots, X_n \models Y$$

This shows that, whilst provability and entailment are *different* notions, they are extensionally equivalent. As such:

- An argument is *valid* iff *the conclusion can be proved from the premises*.
- Two sentences are *logically equivalent* iff they are *provably equivalent*.
- Sentences are *provably consistent* iff they are *not provably inconsistent*.

For this reason, you can pick and choose when to think in terms of proofs and when to think in terms of valuations/interpretations, doing whichever is easier for a given task. The table on the next page summarises which is (usually) easier.

It is intuitive that provability and semantic entailment should agree. But—let us repeat this—do not be fooled by the similarity of the symbols ‘ \models ’ and ‘ \vdash ’. These two symbols have very different meanings. The

fact that provability and semantic entailment agree is not an easy result to come by.

In fact, demonstrating that provability and semantic entailment agree is, very decisively, the point at which introductory logic becomes intermediate logic.

	Yes	No
Is X a logical truth?	give a proof which shows $\vdash X$	give an interpretation in which X is false
Is X a contradiction?	give a proof which shows $\vdash \neg X$	give an interpretation in which X is true
Are X and Y equivalent?	give two proofs, one for $X \vdash Y$ and one for $Y \vdash X$	give an interpretation in which X and Y have different truth values
Are X_1, X_2, \dots, X_n jointly consistent?	give an interpretation in which all of X_1, X_2, \dots, X_n are true	prove a contradiction from assumptions X_1, X_2, \dots, X_n
Is $X_1, X_2, \dots, X_n \therefore C$ valid?	give a proof with assumptions X_1, X_2, \dots, X_n and concluding with C	give an interpretation in which each of X_1, X_2, \dots, X_n is true and C is false

PART IX

Identity

CHAPTER 42

Identity

Consider this sentence:

1. Pavel owes money to everyone

Let the domain be people; this will allow us to symbolize ‘everyone’ as a universal quantifier. Offering the symbolization key:

Oxy : ______x owes money to ______y
 p : Pavel

we can symbolize sentence 1 by ‘ $\forall xOp_x$ ’. But this has a (perhaps) odd consequence. It requires that Pavel owes money to every member of the domain (whatever the domain may be). The domain certainly includes Pavel. So this entails that Pavel owes money to himself.

Perhaps we meant to say:

2. Pavel owes money to everyone *else*
3. Pavel owes money to everyone *other than* Pavel
4. Pavel owes money to everyone *except* Pavel himself

but we do not know how to deal with the italicised words yet. The solution is to add another symbol to FOL.

42.1 Adding identity

The new symbol we add is ‘=’. This is a symbol that we can use for *identity*.

We will then be able symbolise

5. Clark Kent is Superman.

as $k = s$, using the symbolisation key

k : Clark Kent

s : Superman

This will also be a symbolisations of paraphrases of 5:

6. Clark Kent and Superman are the same person.

7. Clark Kent is identical to Superman.

Using \doteq we will now be able to symbolise sentences 2–4. All of these sentences can be paraphrased as ‘Everyone who is not Pavel is owed money by Pavel’. Paraphrasing some more, we get: ‘For all x , if x is not Pavel, then x is owed money by Pavel’. Now that we are armed with our new identity symbol, we can symbolize this as ‘ $\forall x(\neg x = p \rightarrow Opx)$ ’.

In addition to sentences that use the word ‘else’, ‘other than’ and ‘except’, identity will be helpful when symbolizing some sentences that contain the words ‘besides’ and ‘only.’ Consider these examples:

8. No one besides Pavel owes money to Hikaru.

9. Only Pavel owes Hikaru money.

Let ‘ h ’ name Hikaru. Sentence 8 can be paraphrased as, ‘No one who is not Pavel owes money to Hikaru’. This can be symbolized by ‘ $\neg \exists x(\neg x = p \ \& \ O x h)$ ’. Equally, sentence 9 can be paraphrased as ‘for all x , if x owes money to Hikaru, then x is Pavel’. It can then be symbolized as ‘ $\forall x(O x h \rightarrow x = p)$ ’.

Sentence 9 can be treated similarly, but there is one subtlety here. Do either sentence 8 or 9 entail that Pavel himself owes money to Hikaru?

42.2 There are at least...

We will now look at more that we can do armed with our new identity symbol. We can also use identity to say how many things there are of a particular kind. For example, consider these sentences:

10. There is at least one apple

11. There are at least two apples

12. There are at least three apples

We will use the symbolization key:

Ax : _____ _{x} is an apple

Sentence 10 does not require identity. It can be adequately symbolized by ' $\exists xAx$ ': There is an apple; perhaps many, but at least one.

It might be tempting to also symbolize sentence 11 without identity. Yet consider the sentence ' $\exists x\exists y(Ax \ \& \ Ay)$ '. Roughly, this says that there is some apple x in the domain and some apple y in the domain. Since nothing precludes these from being one and the same apple, this would be true even if there were only one apple. In order to make sure that we are dealing with *different* apples, we need an identity predicate. Sentence 11 needs to say that the two apples that exist are not identical, so it can be symbolized by ' $\exists x\exists y((Ax \ \& \ Ay) \ \& \ \neg x=y)$ '.

Sentence 12 requires talking about three different apples. Now we need three existential quantifiers, and we need to make sure that each will pick out something different: ' $\exists x\exists y\exists z[((Ax \ \& \ Ay) \ \& \ Az) \ \& \ ((\neg x=y \ \& \ \neg y=z) \ \& \ \neg x=z)]$ '.

42.3 There are at most. . .

Now consider these sentences:

13. There is at most one apple
14. There are at most two apples

Sentence 13 can be paraphrased as, 'It is not the case that there are at least *two* apples'. This is just the negation of sentence 11:

$$\neg\exists x\exists y[(Ax \ \& \ Ay) \ \& \ \neg x=y]$$

But sentence 13 can also be approached in another way. It means that if you pick out an object and it's an apple, and then you pick out an object and it's also an apple, you must have picked out the same object both times. With this in mind, it can be symbolized by

$$\forall x\forall y[(Ax \ \& \ Ay) \rightarrow x=y]$$

The two sentences will turn out to be logically equivalent.

In a similar way, sentence 14 can be approached in two equivalent ways. It can be paraphrased as, 'It is not the case that there are *three* or more distinct apples', so we can offer:

$$\neg\exists x\exists y\exists z(Ax \ \& \ Ay \ \& \ Az \ \& \ \neg x=y \ \& \ \neg y=z \ \& \ \neg x=z)$$

Alternatively we can read it as saying that if you pick out an apple, and an apple, and an apple, then you will have picked out (at least) one of these objects more than once. Thus:

$$\forall x \forall y \forall z [(Ax \ \& \ Ay \ \& \ Az) \rightarrow (x=y \vee x=z \vee y=z)]$$

42.4 There are exactly...

We can now consider precise statements, like:

15. There is exactly one apple.
16. There are exactly two apples.
17. There are exactly three apples.

Sentence 15 can be paraphrased as, ‘There is *at least* one apple and there is *at most* one apple’. This is just the conjunction of sentence 10 and sentence 13. So we can offer:

$$\exists x Ax \ \& \ \forall x \forall y [(Ax \ \& \ Ay) \rightarrow x=y]$$

But it is perhaps more straightforward to paraphrase sentence 15 as, ‘There is a thing x which is an apple, and everything which is an apple is just x itself’. Thought of in this way, we offer:

$$\exists x [Ax \ \& \ \forall y (Ay \rightarrow x=y)]$$

Similarly, sentence 16 may be paraphrased as, ‘There are *at least* two apples, and there are *at most* two apples’. Thus we could offer

$$\begin{aligned} &\exists x \exists y ((Ax \ \& \ Ay) \ \& \ \neg x=y) \ \& \\ &\forall x \forall y \forall z [((Ax \ \& \ Ay) \ \& \ Az) \rightarrow ((x=y \vee x=z) \vee y=z)] \end{aligned}$$

More efficiently, though, we can paraphrase it as ‘There are at least two different apples, and every apple is one of those two apples’. Then we offer:

$$\exists x \exists y [((Ax \ \& \ Ay) \ \& \ \neg x=y) \ \& \ \forall z (Az \rightarrow (x=z \vee y=z))]$$

Finally, consider these sentence:

18. There are exactly two things
19. There are exactly two objects

It might be tempting to add a predicate to our symbolization key, to symbolize the English predicate ‘_____ is a thing’ or ‘_____ is an object’, but this is unnecessary. Words like ‘thing’ and ‘object’ do not sort wheat from chaff: they apply trivially to everything, which is to say, they apply trivially to every thing. So we can symbolize either sentence with either of the following:

$$\begin{aligned} \exists x \exists y \neg x=y \ \& \ \neg \exists x \exists y \exists z ((\neg x=y \ \& \ \neg y=z) \ \& \ \neg x=z) \\ \exists x \exists y [\neg x=y \ \& \ \forall z (x=z \vee y=z)] \end{aligned}$$

Practice exercises

A. Explain why:

- ‘ $\exists x \forall y (Ay \leftrightarrow x=y)$ ’ is a good symbolization of ‘there is exactly one apple’.
- ‘ $\exists x \exists y [\neg x=y \ \& \ \forall z (Az \leftrightarrow (x=z \vee y=z))]$ ’ is a good symbolization of ‘there are exactly two apples’.

CHAPTER 43

Definite descriptions

Consider sentences like:

1. Nick is the traitor.
2. The traitor went to Cambridge.
3. The traitor is the deputy

These are definite descriptions: they are meant to pick out a *unique* object. They should be contrasted with *indefinite* descriptions, such as ‘Nick is *a* traitor’. They should equally be contrasted with *generics*, such as ‘*The* whale is a mammal’ (it’s inappropriate to ask *which* whale). The question we face is: how should we deal with definite descriptions in FOL?

43.1 Treating definite descriptions as terms

One option would be to introduce new names whenever we come across a definite description. This is probably not a great idea. We know that *the* traitor—whoever it is—is indeed *a* traitor. We want to preserve that information in our symbolization.

A second option would be to use a *new* definite description operator, such as ‘*t*’. The idea would be to symbolize ‘the F’ as ‘ ιxFx ’; or to symbolize ‘the G’ as ‘ ιxGx ’, etc. Expression of the form ιxXx would then behave like names. If we followed this path, then using the following symbolization key:

domain: people

Tx : ______x is a traitor

Dx : ______x is a deputy

Cx : ______x went to Cambridge

n : Nick

We could symbolize sentence 1 with ' $\neg Tx = n$ ', sentence 2 with ' $C \neg Tx$ ', and sentence 3 with ' $\neg Tx = \neg Dx$ '.

However, it would be nice if we didn't have to add a new symbol to FOL. And indeed, we might be able to make do without one.

43.2 Russell's analysis

Bertrand Russell offered an analysis of definite descriptions. Very briefly put, he observed that, when we say 'the F' in the context of a definite description, our aim is to pick out the *one and only* thing that is F (in the appropriate context). Thus Russell analysed the notion of a definite description as follows:¹

the F is G **iff** there is at least one F, *and*
there is at most one F, *and*
every F is G

Note a very important feature of this analysis: '*the*' *does not appear on the right-side of the equivalence*. Russell is aiming to provide an understanding of definite descriptions in terms that do not presuppose them.

Now, one might worry that we can say 'the table is brown' without implying that there is one and only one table in the universe. But this is not (yet) a fantastic counterexample to Russell's analysis. The domain of discourse is likely to be restricted by context (e.g. to objects in my line of sight).

If we accept Russell's analysis of definite descriptions, then we can symbolize sentences of the form 'the F is G' using our strategy for numerical quantification in FOL. After all, we can deal with the three conjuncts on the right-hand side of Russell's analysis as follows:

$$\exists xFx \ \& \ \forall x\forall y((Fx \ \& \ Fy) \rightarrow x=y) \ \& \ \forall x(Fx \rightarrow Gx)$$

In fact, we could express the same point rather more crisply, by recognizing that the first two conjuncts just amount to the claim that there

¹ Bertrand Russell, 'On Denoting', 1905, *Mind* 14, pp. 479–93; also Russell, *Introduction to Mathematical Philosophy*, 1919, London: Allen and Unwin, ch. 16.

is *exactly* one F, and that the last conjunct tells us that that object is F. So, equivalently, we could offer:

$$\exists x[(Fx \ \& \ \forall y(Fy \rightarrow x=y)) \ \& \ Gx]$$

Using these sorts of techniques, we can now symbolize sentences 1–3 without using any new-fangled fancy operator, such as ‘ ι ’.

Sentence 1 is exactly like the examples we have just considered. So we would symbolize it by ‘ $\exists x(Tx \ \& \ \forall y(Ty \rightarrow x=y) \ \& \ x=n)$ ’.

Sentence 2 poses no problems either: ‘ $\exists x(Tx \ \& \ \forall y(Ty \rightarrow x=y) \ \& \ Cx)$ ’.

Sentence 3 is a little trickier, because it links two definite descriptions. But, deploying Russell’s analysis, it can be paraphrased by ‘there is exactly one traitor, x, and there is exactly one deputy, y, and $x=y$ ’. So we can symbolize it by:

$$\exists x\exists y([Tx \ \& \ \forall z(Tz \rightarrow x=z)] \ \& \ [Dy \ \& \ \forall z(Dz \rightarrow y=z)] \ \& \ x=y)$$

Note that we have made sure that the formula ‘ $x=y$ ’ falls within the scope of both quantifiers!

43.3 Empty definite descriptions

One of the nice features of Russell’s analysis is that it allows us to handle *empty* definite descriptions neatly.

France has no king at present. Now, if we were to introduce a name, ‘ k ’, to name the present King of France, then everything would go wrong: remember from §28 that a name must always pick out some object in the domain, and whatever we choose as our domain, it will contain no present kings of France.

Russell’s analysis neatly avoids this problem. Russell tells us to treat definite descriptions using predicates and quantifiers, instead of names. Since predicates can be empty (see §29), this means that no difficulty now arises when the definite description is empty.

Indeed, Russell’s analysis helpfully highlights two ways to go wrong in a claim involving a definite description. To adapt an example from Stephen Neale (1990),² suppose Alex claims:

4. I am dating the present king of France.

Using the following symbolization key:

²Neale, *Descriptions*, 1990, Cambridge: MIT Press.

a: Alex

Kx : ______{*x*} is a present king of France

Dxy : ______{*x*} is dating ______{*y*}

Sentence 4 would be symbolized by ' $\exists x(\forall y(Ky \leftrightarrow x=y) \& Dax)$ '. Now, this can be false in (at least) two ways, corresponding to these two different sentences:

5. There is no one who is both the present King of France and such that he and Alex are dating.
6. There is a unique present King of France, but Alex is not dating him.

Sentence 5 might be paraphrased by 'It is not the case that: the present King of France and Alex are dating'. It will then be symbolized by ' $\neg \exists x[(Kx \& \forall y(Ky \rightarrow x=y)) \& Dax]$ '. We might call this *outer* negation, since the negation governs the entire sentence. Note that it will be true if there is no present King of France.

Sentence 6 can be symbolized by ' $\exists x((Kx \& \forall y(Ky \rightarrow x=y)) \& \neg Dax)$ '. We might call this *inner* negation, since the negation occurs within the scope of the definite description. Note that its truth requires that there is a present King of France, albeit one who is not dating Alex.

43.4 The adequacy of Russell's analysis

How good is Russell's analysis of definite descriptions? This question has generated a substantial philosophical literature, but we will restrict ourselves to two observations.

One worry focusses on Russell's treatment of empty definite descriptions. If there are no Fs, then on Russell's analysis, both 'the F is G' is and 'the F is non-G' are false. P.F. Strawson suggested that such sentences should not be regarded as false, exactly.³ Rather, they involve presupposition failure, and need to be regarded as *neither* true *nor* false.

If we agree with Strawson here, we will need to revise our logic. For, in our logic, there are only two truth values (True and False), and every sentence is assigned exactly one of these truth values.

But there is room to disagree with Strawson. Strawson is appealing to some linguistic intuitions, but it is not clear that they are very robust. For example: isn't it just *false*, not 'gappy', that Tim is dating the present King of France?

³P.F. Strawson, 'On Referring', 1950, *Mind* 59, pp. 320–34.

Keith Donnellan raised a second sort of worry, which (very roughly) can be brought out by thinking about a case of mistaken identity.⁴ Two men stand in the corner: a very tall man drinking what looks like a gin martini; and a very short man drinking what looks like a pint of water. Seeing them, Malika says:

7. The gin-drinker is very tall!

Russell's analysis will have us render Malika's sentence as:

7'. There is exactly one gin-drinker [in the corner], and whoever is a gin-drinker [in the corner] is very tall.

Now suppose that the very tall man is actually drinking *water* from a martini glass; whereas the very short man is drinking a pint of (neat) gin. By Russell's analysis, Malika has said something false, but don't we want to say that Malika has said something *true*?

Again, one might wonder how clear our intuitions are on this case. We can all agree that Malika intended to pick out a particular man, and say something true of him (that he was tall). On Russell's analysis, she actually picked out a different man (the short one), and consequently said something false of him. But maybe advocates of Russell's analysis only need to explain *why* Malika's intentions were frustrated, and so why she said something false. This is easy enough to do: Malika said something false because she had false beliefs about the men's drinks; if Malika's beliefs about the drinks had been true, then she would have said something true.⁵

To say much more here would lead us into deep philosophical waters. That would be no bad thing, but for now it would distract us from the immediate purpose of learning formal logic. So, for now, we will stick with Russell's analysis of definite descriptions, when it comes to putting things into FOL. It is certainly the best that we can offer, without significantly revising our logic, and it is quite defensible as an analysis.

Practice exercises

A. Using the following symbolization key:

⁴Keith Donnellan, 'Reference and Definite Descriptions', 1966, *Philosophical Review* 77, pp. 281–304.

⁵Interested parties should read Saul Kripke, 'Speaker Reference and Semantic Reference', 1977, in French et al (eds.), *Contemporary Perspectives in the Philosophy of Language*, Minneapolis: University of Minnesota Press, pp. 6–27.

domain: people

Kx : ______x knows the combination to the safe.

Sx : ______x is a spy.

Vx : ______x is a vegetarian.

Txy : ______x trusts ______y .

h : Hofthor

i : Ingmar

symbolize the following sentences in FOL:

1. Hofthor trusts a vegetarian.
2. Everyone who trusts Ingmar trusts a vegetarian.
3. Everyone who trusts Ingmar trusts someone who trusts a vegetarian.
4. Only Ingmar knows the combination to the safe.
5. Ingmar trusts Hofthor, but no one else.
6. The person who knows the combination to the safe is a vegetarian.
7. The person who knows the combination to the safe is not a spy.

B. Using the following symbolization key:

domain: cards in a standard deck

Bx : ______x is black.

Cx : ______x is a club.

Dx : ______x is a deuce.

Jx : ______x is a jack.

Mx : ______x is a man with an axe.

Ox : ______x is one-eyed.

Wx : ______x is wild.

symbolize each sentence in FOL:

1. All clubs are black cards.
2. There are no wild cards.
3. There are at least two clubs.
4. There is more than one one-eyed jack.
5. There are at most two one-eyed jacks.
6. There are two black jacks.
7. There are four deuces.
8. The deuce of clubs is a black card.
9. One-eyed jacks and the man with the axe are wild.
10. If the deuce of clubs is wild, then there is exactly one wild card.
11. The man with the axe is not a jack.

12. The deuce of clubs is not the man with the axe.

C. Using the following symbolization key:

domain: animals in the world

Bx : ______x is in Farmer Brown's field.

Hx : ______x is a horse.

Px : ______x is a Pegasus.

Wx : ______x has wings.

symbolize the following sentences in FOL:

1. There are at least three horses in the world.
2. There are at least three animals in the world.
3. There is more than one horse in Farmer Brown's field.
4. There are three horses in Farmer Brown's field.
5. There is a single winged creature in Farmer Brown's field; any other creatures in the field must be wingless.
6. The Pegasus is a winged horse.
7. The animal in Farmer Brown's field is not a horse.
8. The horse in Farmer Brown's field does not have wings.

D. In this chapter, we symbolized 'Nick is the traitor' by ' $\exists x(Tx \ \& \ \forall y(Ty \rightarrow x=y) \ \& \ x=n)$ '. Two equally good symbolizations would be:

- $Tn \ \& \ \forall y(Ty \rightarrow n=y)$
- $\forall y(Ty \leftrightarrow y=n)$

Explain why these would be equally good symbolizations.

CHAPTER 44

Semantics for FOL with identity

FOL with identity extends FOL as we presented it earlier.

44.1 Sentences of FOL with identity

When we add the identity symbol to FOL, we add a new kind of atomic sentence, for example $a = b$.

We simply do this by adding a new kind of atomic sentence:

If a and b are names, then $a = b$ is an atomic sentence.

This is added to the other clauses of what it is to be a sentence as they were given in §31

We can now see that $\forall x(\neg x = p \rightarrow Op x)$ is a sentence, as it could be constructed as follows:

$$\begin{array}{c}
 \forall x(\neg x=p \rightarrow Op x) \\
 | \\
 (\neg c=p \rightarrow Op c) \\
 \swarrow \quad \searrow \\
 \neg c=p \quad Op c \\
 | \\
 c=p
 \end{array}$$

Be aware that when you see $\neg a=b$ the negation is attached to the whole sentence $a=b$, not to a . So, you should not write $a=\neg b$. This is not a sentence. Sometimes you might see $a \neq b$, and this is short hand for $\neg a=b$.

Sometimes you might come across cases where you might feel tempted to write something like $a = \neg b$. Such temptation should be avoided. For example, in our semi-formalised English we might say ‘for all x , if x is not p , then Ax ’. But note that this should be $\forall x(\neg x=p \rightarrow Ax)$, and should not be written with $x=\neg p$.

44.2 Semantics for identity

Now that we have added the identity symbol to FOL, we simply need to expand our notion of truth from Part VII to also account for atomic sentences like $a=b$. Our clause that we add is:

$a=b$ is true **iff** a and b name the same object in that interpretation.

So on our go-to interpretation from Part VII,

domain: all people born before 2000CE

a : Aristotle

b : Beyoncé

Px : ---_x is a philosopher

Rxy : ---_x was born before ---_y

we have that $a=b$ is false: Aristotle and Beyoncé are different people, so ‘ a ’ and ‘ b ’ name different objects.

Consider the following interpretation

domain: All celestial bodies

e : The evening star
 m : The morning star

It turns out that The Morning Star is *the same object as* The Evening Star: they are names for Venus. So here we have two names for the same object. We thus have $e = m$ is true on this interpretation.

Identity becomes particularly useful when we have quantifiers. Suppose we have

domain: Alfred, Billy, Carys
 a : Alfred
 b : Billy
 c : Carys

Remember, to check if $\exists x X(\dots x \dots x \dots)$ is true we first add a new name, let's use d , and we see if there is some way of extending the domain so that $X(\dots d \dots d \dots)$ is true. So, let's see if $\exists x x = a$ is true: we add a new name d and consider the extended interpretation with

d : Alfred

Then $d = a$ is true on this interpretation. So there is some way of interpreting ' d ' where $d = a$ is true; and thus $\exists x x = a$ is true.

On this interpretation we can also see that $\forall x (x = a \vee x = b \vee x = c)$ is true. Why? We add a new name ' d '. There are three ways we can extend our original interpretation:

1. d : Alfred
2. d : Billy
3. d : Carys

On the first of these, $d = a$ is true, on the second $d = b$ is true, and on the third $d = c$ is true. So on each of these interpretations, $d = a \vee d = b \vee d = c$ is true, and thus $\forall x (x = a \vee x = b \vee x = c)$.

CHAPTER 45

Rules for identity

If two names refer to the same object, then swapping one name for another will not change the truth value of any sentence. So, in particular, if ‘*a*’ and ‘*b*’ name the same object, then all of the following will be valid:

$$Aa \therefore Ab$$

$$Ab \therefore Aa$$

$$Raa \therefore Rbb$$

$$Raa \therefore Rab$$

$$Rca \therefore Rcb$$

$$\forall x Rxa \therefore \forall x Rxb$$

We capture this idea in our elimination rule. If you have established ‘*a* = *b*’, then anything that is true of the object named by ‘*a*’ must also be true of the object named by ‘*b*’. For any sentence with ‘*a*’ in it, you can replace some or all of the occurrences of ‘*a*’ with ‘*b*’ and produce an equivalent sentence. For example, from ‘*Raa*’ and ‘*a* = *b*’, you are justified in inferring ‘*Rab*’, ‘*Rba*’ or ‘*Rbb*’. More generally:

m	$a=b$	
n	$X(\dots a \dots a \dots)$	
	$X(\dots b \dots a \dots)$	$\Rightarrow \exists m, n$

The notation here is as for \exists I. So $X(\dots a \dots a \dots)$ is a formula containing the name a , and $X(\dots b \dots a \dots)$ is a formula obtained by replacing one or more instances of the name a with the name b . Lines m and n can occur in either order, and do not need to be adjacent, but we always cite the statement of identity first. Symmetrically, we allow:

m	$a=b$	
n	$X(\dots b \dots b \dots)$	
	$X(\dots a \dots b \dots)$	$\Rightarrow \exists m, n$

This rule is sometimes called *Leibniz's Law*, after Gottfried Leibniz.

Some philosophers have believed the reverse of this claim. That is, they have believed that when exactly the same sentences (not containing \equiv) are true of two objects, then they are really just one and the same object after all. This is a highly controversial philosophical claim (sometimes called the *identity of indiscernibles*) and our logic will not subscribe to it; we allow that exactly the same things might be true of two *distinct* objects.

To bring this out, consider the following interpretation:

domain: P.D. Magnus, Tim Button

a : P.D. Magnus

b : Tim Button

- For every primitive predicate we care to consider, that predicate is true of *nothing*.

Suppose ' A ' is a one-place predicate; then ' Aa ' is false and ' Ab ' is false, so ' $Aa \leftrightarrow Ab$ ' is true. Similarly, if ' R ' is a two-place predicate, then ' Raa ' is false and ' Rab ' is false, so that ' $Raa \leftrightarrow Rab$ ' is true. And so it goes: every atomic sentence not involving \equiv is false, so every biconditional linking such sentences is true. For all that, Tim Button and P.D. Magnus are two distinct people, not one and the same!

Since we are not subscribing to the thesis of identity of indiscernibles, no matter how much you learn about two objects, we cannot prove that they are identical. That is unless, of course, you learn that the two objects are, in fact, identical, but then the proof will hardly be very illuminating.

The consequence of this, for our proof system, is that there are no sentences that do not already contain the identity predicate that could justify the conclusion ' $a=b$ '. This means that the identity introduction rule will not justify ' $a=b$ ', or any other identity claim containing two different names.

However, every object is identical to itself. No premises, then, are required in order to conclude that something is identical to itself. So this will be the identity introduction rule:

$$\begin{array}{|l} c=c \\ \hline \end{array} \quad \equiv\text{I}$$

Notice that like the Law of Excluded Middle, this rule does not require referring to any prior lines of the proof. For any name c , you can write $c=c$ on any point, with only the $\equiv\text{I}$ rule as justification.

To see the rules in action, we will prove some quick results. First, we will prove that identity is *symmetric*:

1	$a=b$	
2	$a=a$	$\equiv\text{I}$
3	$b=a$	$\equiv\text{E } 1, 2$
4	$a=b \rightarrow b=a$	$\rightarrow\text{I } 1-3$
5	$\forall y(a=y \rightarrow y=a)$	$\forall\text{I } 4$
6	$\forall x\forall y(x=y \rightarrow y=x)$	$\forall\text{I } 5$

We obtain line 3 by replacing one instance of ' a ' in line 2 with an instance of ' b '; this is justified given ' $a=b$ '.

Second, we will prove that identity is *transitive*:

1		$a = b \ \& \ b = c$	
2		$a = b$	&E 1
3		$b = c$	&E 1
4		$a = c$	\equiv E 2, 3
5		$(a = b \ \& \ b = c) \rightarrow a = c$	\rightarrow I 1–4
6		$\forall z((a = b \ \& \ b = z) \rightarrow a = z)$	\forall I 5
7		$\forall y \forall z((a = y \ \& \ y = z) \rightarrow a = z)$	\forall I 6
8		$\forall x \forall y \forall z((x = y \ \& \ y = z) \rightarrow x = z)$	\forall I 7

We obtain line 4 by replacing ‘ b ’ in line 3 with ‘ a ’; this is justified given ‘ $a = b$ ’.

Practice exercises

A. Provide a proof of each claim.

1. $Pa \vee Qb, Qb \rightarrow b = c, \neg Pa \vdash Qc$
2. $m = n \vee n = o, An \vdash Am \vee Ao$
3. $\forall x \ x = m, Rma \vdash \exists x Rxx$
4. $\forall x \forall y (Rxy \rightarrow x = y) \vdash Rab \rightarrow Rba$
5. $\neg \exists x \neg x = m \vdash \forall x \forall y (Px \rightarrow Py)$
6. $\exists x Jx, \exists x \neg Jx \vdash \exists x \exists y \neg x = y$
7. $\forall x (x = n \leftrightarrow Mx), \forall x (Ox \vee \neg Mx) \vdash On$
8. $\exists x Dx, \forall x (x = p \leftrightarrow Dx) \vdash Dp$
9. $\exists x [(Kx \ \& \ \forall y (Ky \rightarrow x = y)) \ \& \ Bx], Kd \vdash Bd$
10. $\vdash Pa \rightarrow \forall x (Px \vee \neg x = a)$

B. Show that the following are provably equivalent:

- $\exists x ([Fx \ \& \ \forall y (Fy \rightarrow x = y)] \ \& \ x = n)$
- $Fn \ \& \ \forall y (Fy \rightarrow n = y)$

And hence that both have a decent claim to symbolize the English sentence ‘Nick is the F’.

C. In §IX, we claimed that the following are logically equivalent symbolizations of the English sentence ‘there is exactly one F’:

- $\exists x Fx \ \& \ \forall x \forall y [(Fx \ \& \ Fy) \rightarrow x=y]$
- $\exists x [Fx \ \& \ \forall y (Fy \rightarrow x=y)]$
- $\exists x \forall y (Fy \leftrightarrow x=y)$

Show that they are all provably equivalent. (*Hint:* to show that three claims are provably equivalent, it suffices to show that the first proves the second, the second proves the third and the third proves the first; think about why.)

D. Symbolize the following argument

There is exactly one F. There is exactly one G. Nothing is both F and G. So: there are exactly two things that are either F or G.

And offer a proof of it.

PART X

Metatheory (Non- examinable)

CHAPTER 46

Normal forms

46.1 Disjunctive Normal Form

Sometimes it is useful to consider sentences of a particularly simple form. For instance, we might consider sentences in which \neg only attaches to atomic sentences, or those which are combinations of atomic sentences and negated atomic sentences using only $\&$. A relatively general but still simple form is that where a sentence is a disjunction of conjunctions of atomic or negated atomic sentences. When such a sentence is constructed, we start with atomic sentences, then (perhaps) attach negations, then (perhaps) combine using $\&$, and finally (perhaps) combine using \vee .

Let's say that a sentence is in **DISJUNCTIVE NORMAL FORM** *iff* it meets all of the following conditions:

- (DNF1) No connectives occur in the sentence other than negations, conjunctions and disjunctions;
- (DNF2) Every occurrence of negation has minimal scope (i.e. any ' \neg ' is immediately followed by an atomic sentence);
- (DNF3) No disjunction occurs within the scope of any conjunction.

So, here are some sentences in disjunctive normal form:

$$\begin{aligned} &A \\ &(A \& \neg B \& C) \\ &(A \& B) \vee (A \& \neg B) \\ &(A \& B) \vee (A \& B \& C \& \neg D \& \neg E) \\ &A \vee (C \& \neg P_{234} \& P_{233} \& Q) \vee \neg B \end{aligned}$$

Note that we have here broken one of the maxims of this book and *temporarily* allowed ourselves to employ the relaxed bracketing-conventions that allow conjunctions and disjunctions to be of arbitrary length. These conventions make it easier to see when a sentence is in disjunctive normal form. We will continue to help ourselves to these relaxed conventions, without further comment.

To further illustrate the idea of disjunctive normal form, we will introduce some more notation. We write ‘ $\pm A$ ’ to indicate that A is an atomic sentence which may or may not be prefaced with an occurrence of negation. Then a sentence in disjunctive normal form has the following shape:

$$(\pm A_1 \wedge \dots \wedge \pm A_i) \vee (\pm A_{i+1} \wedge \dots \wedge \pm A_j) \vee \dots \vee (\pm A_{m+1} \wedge \dots \wedge \pm A_n)$$

We now know what it is for a sentence to be in disjunctive normal form. The result that we are aiming at is:

Disjunctive Normal Form Theorem. For any sentence, there is a logically equivalent sentence in disjunctive normal form.

Henceforth, we will abbreviate ‘Disjunctive Normal Form’ by ‘DNF’.

46.2 Proof of DNF Theorem via truth tables

Our first proof of the DNF Theorem employs truth tables. We will first illustrate the technique for finding an equivalent sentence in DNF, and then turn this illustration into a rigorous proof.

Let’s suppose we have some sentence, S , which contains three atomic sentences, ‘ A ’, ‘ B ’ and ‘ C ’. The very first thing to do is fill out a complete truth table for S . Maybe we end up with this:

A	B	C	S
T	T	T	T
T	T	F	F
T	F	T	T
T	F	F	F
F	T	T	F
F	T	F	F
F	F	T	T
F	F	F	T

As it happens, S is true on four lines of its truth table, namely lines 1, 3, 7 and 8. Corresponding to each of those lines, we will write down four sentences, whose only connectives are negations and conjunctions, where every negation has minimal scope:

- ' $A \& B \& C$ ' which is true on line 1 (and only then)
- ' $A \& \neg B \& C$ ' which is true on line 3 (and only then)
- ' $\neg A \& \neg B \& C$ ' which is true on line 7 (and only then)
- ' $\neg A \& \neg B \& \neg C$ ' which is true on line 8 (and only then)

We now combine all of these conjunctions using \vee , like so:

$$(A \& B \& C) \vee (A \& \neg B \& C) \vee (\neg A \& \neg B \& C) \vee (\neg A \& \neg B \& \neg C)$$

This gives us a sentence in DNF which is true on exactly those lines where one of the disjuncts is true, i.e. it is true on (and only on) lines 1, 3, 7, and 8. So this sentence has exactly the same truth table as S . So we have a sentence in DNF that is logically equivalent to S , which is exactly what we wanted!

Now, the strategy that we just adopted did not depend on the specifics of S ; it is perfectly general. Consequently, we can use it to obtain a simple proof of the DNF Theorem.

Pick any arbitrary sentence, S , and let A_1, \dots, A_n be the atomic sentences that occur in S . To obtain a sentence in DNF that is logically equivalent S , we consider S 's truth table. There are two cases to consider:

1. S is false on every line of its truth table. Then, S is a contradiction. In that case, the contradiction $(A_1 \& \neg A_1)$ is in DNF and logically equivalent to S .
2. S is true on at least one line of its truth table. For each line i of the truth table, let B_i be a conjunction of the form

$$(\pm A_1 \wedge \dots \wedge \pm A_n)$$

where the following rules determine whether or not to include a negation in front of each atomic sentence:

$$\begin{aligned} A_m &\text{ is a conjunct of } B_i \text{ iff } A_m \text{ is true on line } i \\ \neg A_m &\text{ is a conjunct of } B_i \text{ iff } A_m \text{ is false on line } i \end{aligned}$$

Given these rules, B_i is true on (and only on) line i of the truth table which considers all possible valuations of A_1, \dots, A_n (i.e. S 's truth table).

Next, let i_1, i_2, \dots, i_m be the numbers of the lines of the truth table where S is *true*. Now let D be the sentence:

$$B_{i_1} \vee B_{i_2} \vee \dots \vee B_{i_m}$$

Since S is true on at least one line of its truth table, D is indeed well-defined; and in the limiting case where S is true on exactly one line of its truth table, D is just B_{i_1} , for some i_1 .

By construction, D is in DNF. Moreover, by construction, for each line i of the truth table: S is true on line i of the truth table *iff* one of D 's disjuncts (namely, B_i) is true on, and only on, line i . Hence S and D have the same truth table, and so are logically equivalent.

These two cases are exhaustive and, either way, we have a sentence in DNF that is logically equivalent to S .

So we have proved the DNF Theorem. Before we say any more, though, we should immediately flag that we are hereby returning to the austere definition of a (TFL) sentence, according to which we can assume that any conjunction has exactly two conjuncts, and any disjunction has exactly two disjuncts.

46.3 Conjunctive Normal Form

So far in this chapter, we have discussed *disjunctive* normal form. It may not come as a surprise to hear that there is also such a thing as *conjunctive normal form* (CNF).

The definition of CNF is exactly analogous to the definition of DNF. So, a sentence is in CNF *iff* it meets all of the following conditions:

- (CNF1) No connectives occur in the sentence other than negations, conjunctions and disjunctions;
- (CNF2) Every occurrence of negation has minimal scope;
- (CNF3) No conjunction occurs within the scope of any disjunction.

Generally, then, a sentence in CNF looks like this

$$(\pm A_1 \vee \dots \vee \pm A_i) \wedge (\pm A_{i+1} \vee \dots \vee \pm A_j) \wedge \dots \wedge (\pm A_{m+1} \vee \dots \vee \pm A_n)$$

where each A_k is an atomic sentence.

We can now prove another normal form theorem:

Conjunctive Normal Form Theorem. For any sentence, there is a logically equivalent sentence in conjunctive normal form.

Given a TFL sentence, S , we begin by writing down the complete truth table for S .

If S is *true* on every line of the truth table, then S and $(A_1 \vee \neg A_1)$ are logically equivalent.

If S is *false* on at least one line of the truth table then, for every line on the truth table where S is false, write down a disjunction $(\pm A_1 \vee \dots \vee \pm A_n)$ which is *false* on (and only on) that line. Let C be the conjunction of all of these disjuncts; by construction, C is in CNF and S and C are logically equivalent.

Practice exercises

A. Consider the following sentences:

1. $(A \rightarrow \neg B)$
2. $\neg(A \leftrightarrow B)$
3. $(\neg A \vee \neg(A \& B))$
4. $(\neg(A \rightarrow B) \& (A \rightarrow C))$
5. $(\neg(A \vee B) \leftrightarrow ((\neg C \& \neg A) \rightarrow \neg B))$
6. $((\neg(A \& \neg B) \rightarrow C) \& \neg(A \& D))$

For each sentence, find a logically equivalent sentence in DNF and one in CNF.

CHAPTER 47

Expressive Adequacy

Of our connectives, \neg attaches to a single sentences, and the others all combine exactly two sentences. We may also introduce the idea of an n -place connective. For example, we could consider a three-place connective, ' \heartsuit ', and stipulate that it is to have the following characteristic truth table:

A	B	C	$\heartsuit(A, B, C)$
T	T	T	F
T	T	F	T
T	F	T	T
T	F	F	F
F	T	T	F
F	T	F	T
F	F	T	F
F	F	F	F

Probably this new connective would not correspond with any natural English expression (at least not in the way that '&' corresponds with 'and'). But a question arises: if we wanted to employ a connective with this characteristic truth table, must we add a *new* connective to TFL? Or can we get by with the connectives we *already have*?

Let us make this question more precise. Say that some connectives are **JOINTLY EXPRESSIVELY ADEQUATE** *iff*, for any possible truth table,

there is a sentence containing only those connectives with that truth table.

The general point is, when we are armed with some jointly expressively adequate connectives, no characteristic truth table lies beyond our grasp. And in fact, we are in luck.

Expressive Adequacy Theorem. The connectives of TFL are jointly expressively adequate. Indeed, the following pairs of connectives are jointly expressively adequate:

1. ‘ \neg ’ and ‘ \vee ’
2. ‘ \neg ’ and ‘ $\&$ ’
3. ‘ \neg ’ and ‘ \rightarrow ’

Given any truth table, we can use the method of proving the DNF Theorem (or the CNF Theorem) via truth tables, to write down a scheme which has the same truth table. For example, employing the truth table method for proving the DNF Theorem, we find that the following scheme has the same characteristic truth table as $\heartsuit(A, B, C)$, above:

$$(A \& B \& \neg C) \vee (A \& \neg B \& C) \vee (\neg A \& B \& \neg C)$$

It follows that the connectives of TFL are jointly expressively adequate. We now prove each of the subsidiary results.

Subsidiary Result 1: expressive adequacy of ‘ \neg ’ and ‘ \vee ’. Observe that the scheme that we generate, using the truth table method of proving the DNF Theorem, will only contain the connectives ‘ \neg ’, ‘ $\&$ ’ and ‘ \vee ’. So it suffices to show that there is an equivalent scheme which contains only ‘ \neg ’ and ‘ \vee ’. To show do this, we simply consider that

$$(A \& B) \quad \text{and} \quad \neg(\neg A \vee \neg B)$$

are logically equivalent.

Subsidiary Result 2: expressive adequacy of ‘ \neg ’ and ‘ $\&$ ’. Exactly as in Subsidiary Result 1, making use of the fact that

$$(A \vee B) \quad \text{and} \quad \neg(\neg A \& \neg B)$$

are logically equivalent.

Subsidiary Result 3: expressive adequacy of ‘ \neg ’ and ‘ \rightarrow ’. Exactly as in Subsidiary Result 1, making use of these equivalences instead:

$$\begin{aligned} (A \vee B) \quad \text{and} \quad (\neg A \rightarrow B) \\ (A \& B) \quad \text{and} \quad \neg(A \rightarrow \neg B) \end{aligned}$$

Alternatively, we could simply rely upon one of the other two subsidiary results, and (repeatedly) invoke only one of these two equivalences.

In short, there is never any *need* to add new connectives to TFL. Indeed, there is already some redundancy among the connectives we have: we could have made do with just two connectives, if we had been feeling really austere.

47.1 Individually expressively adequate connectives

In fact, some two-place connectives are *individually* expressively adequate. These connectives are not standardly included in TFL, since they are rather cumbersome to use. But their existence shows that, if we had wanted to, we could have defined a truth-functional language that was expressively adequate, which contained only a single primitive connective.

The first such connective we will consider is ‘ \uparrow ’, which has the following characteristic truth table.

A	B	$A \uparrow B$
T	T	F
T	F	T
F	T	T
F	F	T

This is often called ‘the Sheffer stroke’, after Henry Sheffer, who used it to show how to reduce the number of logical connectives in Russell and Whitehead’s *Principia Mathematica*.¹ (In fact, Charles Sanders Peirce had anticipated Sheffer by about 30 years, but never published his results.)² It is quite common, as well, to call it ‘nand’, since its characteristic truth table is the negation of the truth table for ‘&’.

‘ \uparrow ’ is expressively adequate all by itself.

The Expressive Adequacy Theorem tells us that ‘ \neg ’ and ‘ \vee ’ are jointly expressively adequate. So it suffices to show that, given any scheme which contains only those two connectives, we can rewrite it as a logically equivalent scheme which contains only ‘ \uparrow ’. As in the proof

¹Sheffer, ‘A Set of Five Independent Postulates for Boolean Algebras, with application to logical constants,’ (1913, *Transactions of the American Mathematical Society* 14.4)

²See Peirce, ‘A Boolean Algebra with One Constant’, which dates to c.1880; and Peirce’s *Collected Papers*, 4.264–5.

of the subsidiary cases of the Expressive Adequacy Theorem, then, we simply apply the following equivalences:

$$\neg A \quad \text{and} \quad (A \uparrow A) \\ (A \vee B) \quad \text{and} \quad ((A \uparrow A) \uparrow (B \uparrow B))$$

to the Subsidiary Result 1.

Similarly, we can consider the connective ‘ \downarrow ’:

A	B	$A \downarrow B$
T	T	F
T	F	F
F	T	F
F	F	T

This is sometimes called the ‘Peirce arrow’ (Peirce himself called it ‘ampheck’). More often, though, it is called ‘nor’, since its characteristic truth table is the negation of ‘ \vee ’, that is, of ‘neither ... nor ...’.

‘ \downarrow ’ is expressively adequate all by itself.

As in the previous result for \uparrow , although invoking the equivalences:

$$\neg A \quad \text{and} \quad (A \downarrow A) \\ (A \& B) \quad \text{and} \quad ((A \downarrow A) \downarrow (B \downarrow B))$$

and Subsidiary Result 2.

47.2 Failures of expressive adequacy

In fact, the *only* two-place connectives which are individually expressively adequate are ‘ \uparrow ’ and ‘ \downarrow ’. But how would we show this? More generally, how can we show that some connectives are *not* jointly expressively adequate?

The obvious thing to do is to try to find some truth table which we *cannot* express, using just the given connectives. But there is a bit of an art to this.

To make this concrete, let’s consider the question of whether ‘ \vee ’ is expressively adequate all by itself. After a little reflection, it should be clear that it is not. In particular, it should be clear that any scheme which only contains disjunctions cannot have the same truth table as negation, i.e.:

A	$\neg A$
T	F
F	T

The intuitive reason, why this should be so, is simple: the top line of the desired truth table needs to have the value False; but the top line of any truth table for a scheme which *only* contains \vee will always be True. The same is true for $\&$, \rightarrow , and \leftrightarrow .

‘ \vee ’, ‘ $\&$ ’, ‘ \rightarrow ’, and ‘ \leftrightarrow ’ are not expressively adequate by themselves.

In fact, the following is true:

The *only* two-place connectives that are expressively adequate by themselves are ‘ \uparrow ’ and ‘ \downarrow ’.

This is of course harder to prove than for the primitive connectives. For instance, the “exclusive or” connective does not have a T in the first line of its characteristic truth table, and so the method used above no longer suffices to show that it cannot express all truth tables. It is also harder to show that, e.g., ‘ \leftrightarrow ’ and ‘ \neg ’ *together* are not expressively adequate.

CHAPTER 48

Soundness

In this chapter we relate TFL's semantics to its natural deduction *proof system* (as defined in Part IV). We will prove that the formal proof system is safe: you can only prove sentences from premises from which they actually follow. Intuitively, a formal proof system is sound iff it does not allow you to prove any invalid arguments. This is obviously a highly desirable property. It tells us that our proof system will never lead us astray. Indeed, if our proof system were not sound, then we would not be able to trust our proofs. The aim of this chapter is to prove that our proof system is sound.

Let's make the idea more precise. We'll abbreviate a list of sentences using the greek letter Γ ('gamma'). A formal proof system is **SOUND** (relative to a given semantics) *iff*, whenever there is a formal proof of Z from assumptions among Γ , then Γ genuinely entails Z (given that semantics). Otherwise put, to prove that TFL's proof system is sound, we need to prove the following

Soundness Theorem. For any sentences Γ and Z : if $\Gamma \vdash Z$, then $\Gamma \models Z$

To prove this, we will check each of the rules of TFL's proof system individually. We want to show that no application of those rules ever leads us astray. Since a proof just involves repeated application of those rules, this will show that no proof ever leads us astray. Or at least, that is the general idea.

To begin with, we must make the idea of 'leading us astray' more precise. Say that a line of a proof is **SHINY** iff the assumptions on which that line depends tautologically entail the sentence on that line. The

word ‘shiny’ is not standard among logicians, but it will help us with our discussions. To illustrate the idea, consider the following:

1		$F \rightarrow (G \& H)$	
2			F
3			$G \& H$ $\rightarrow E$ 1, 2
4			G $\& E$ 3
5		$F \rightarrow G$	$\rightarrow I$ 2–4

Line 1 is shiny iff $F \rightarrow (G \& H) \models F \rightarrow (G \& H)$. You should be easily convinced that line 1 is, indeed, shiny! Similarly, line 4 is shiny iff $F \rightarrow (G \& H), F \models G$. Again, it is easy to check that line 4 is shiny. As is every line in this TFL-proof. We want to show that this is no coincidence. That is, we want to prove:

Shininess Lemma. Every line of every TFL-proof is shiny.

Then we will know that we have never gone astray, on any line of a proof. Indeed, given the Shininess Lemma, it will be easy to prove the Soundness Theorem:

Proof. Suppose $\Gamma \vdash Z$. Then there is a TFL-proof, with Z appearing on its last line, whose only undischarged assumptions are among Γ . The Shininess Lemma tells us that every line on every TFL-proof is shiny. So this last line is shiny, i.e. $\Gamma \models Z$. QED

It remains to prove the Shininess Lemma.

To do this, we observe that every line of any TFL-proof is obtained by applying some rule. So what we want to show is that no application of a rule of TFL’s proof system will lead us astray. More precisely, say that a rule of inference is **rule-sound** iff for all TFL-proofs, if we obtain a line on a TFL-proof by applying that rule, and every earlier line in the TFL-proof is shiny, then our new line is also shiny. What we need to show is that *every* rule in TFL’s proof system is rule-sound.

We will do this in the next section. But having demonstrated the rule-soundness of every rule, the Shininess Lemma will follow immediately:

Proof. Fix any line, line n , on any TFL-proof. The sentence written on line n must be obtained using a formal inference rule which is rule-sound. This is to say that, if every earlier line is shiny, then line n itself

is shiny. Hence, by strong induction on the length of TFL-proofs, every line of every TFL-proof is shiny. QED

Note that this proof appeals to a principle of strong induction on the length of TFL-proofs. This is the first time we have seen that principle, and you should pause to confirm that it is, indeed, justified.

It remains to show that every rule is rule-sound. This is not difficult, but it is time-consuming, since we need to check each rule individually, and TFL's proof system has plenty of rules! To speed up the process marginally, we will introduce a convenient abbreviation: ' Δ_i ' ('delta') will abbreviate the assumptions (if any) on which line i depends in our TFL-proof (context will indicate which TFL-proof we have in mind).

Introducing an assumption is rule-sound.

If X is introduced as an assumption on line n , then X is among Δ_n , and so $\Delta_n \models X$.

&I is rule-sound.

Proof. Consider any application of &I in any TFL-proof, i.e., something like:

i	X	
j	Y	
n	$X \& Y$	&I i, j

To show that &I is rule-sound, we assume that every line before line n is shiny; and we aim to show that line n is shiny, i.e. that $\Delta_n \models X \& Y$.

So, let v be any valuation that makes all of Δ_n true.

We first show that v makes X true. To prove this, note that all of Δ_i are among Δ_n . By hypothesis, line i is shiny. So any valuation that makes all of Δ_i true makes X true. Since v makes all of Δ_i true, it makes X true too.

We can similarly see that v makes Y true.

So v makes X true and v makes Y true. Consequently, v makes $X \& Y$ true. So any valuation that makes all of the sentences among Δ_n true also makes $X \& Y$ true. That is: line n is shiny. QED

All of the remaining lemmas establishing rule-soundness will have, essentially, the same structure as this one did.

$\&E$ is rule-sound.

Proof. Assume that every line before line n on some TFL-proof is shiny, and that $\&E$ is used on line n . So the situation is:

i	$X \& Y$	
n	X	$\&E\ i$

(or perhaps with Y on line n instead; but similar reasoning will apply in that case). Let v be any valuation that makes all of Δ_n true. Note that all of Δ_i are among Δ_n . By hypothesis, line i is shiny. So any valuation that makes all of Δ_i true makes $X \& Y$ true. So v makes $X \& Y$ true, and hence makes X true. So $\Delta_n \models X$. QED

$\vee I$ is rule-sound.

We leave this as an exercise.

$\vee E$ is rule-sound.

Proof. Assume that every line before line n on some TFL-proof is shiny, and that $\vee E$ is used on line n . So the situation is:

m	$X \vee Y$	
i	X	
j	Z	
k	Y	
l	Z	
n	Z	$\vee E\ m, i-j, k-l$

Let v be any valuation that makes all of Δ_n true. Note that all of Δ_m are among Δ_n . By hypothesis, line m is shiny. So any valuation that makes Δ_n true makes $X \vee Y$ true. So in particular, v makes $X \vee Y$ true, and hence either v makes X true, or v makes Y true. We now reason through these two cases:

Case 1: v makes X true. All of Δ_i are among Δ_n , with the possible exception of X . Since v makes all of Δ_n true, and also makes X true, v makes all of Δ_i true. Now, by assumption, line j is shiny; so $\Delta_j \models Z$. But the sentences Δ_i are just the sentences Δ_j , so $\Delta_i \models Z$. So, any valuation that makes all of Δ_i true makes Z true. But v is just such a valuation. So v makes Z true.

Case 2: v makes Y true. Reasoning in exactly the same way, considering lines k and l , v makes Z true.

Either way, v makes Z true. So $\Delta_n \models Z$. QED

$\neg E$ is rule-sound.

Proof. Assume that every line before line n on some TFL-proof is shiny, and that $\neg E$ is used on line n . So the situation is:

i	X	
j	$\neg X$	
n	\perp	$\neg I\ i, j$

Note that all of Δ_i and all of Δ_j are among Δ_n . By hypothesis, lines i and j are shiny. So any valuation which makes all of Δ_n true would have to make both X and $\neg X$ true. But no valuation can do that. So no valuation makes all of Δ_n true. So $\Delta_n \models \perp$, vacuously. QED

$\perp I$ is rule-sound.

We leave this as an exercise.

$\neg I$ is rule-sound.

Proof. Assume that every line before line n on some TFL-proof is shiny, and that $\neg I$ is used on line n . So the situation is:

i	<div style="border-right: 1px solid black; padding-right: 10px;">X</div>	
j	<div style="border-right: 1px solid black; padding-right: 10px;">\perp</div>	
n	$\neg X$	$\neg I\ i-j$

Let v be any valuation that makes all of Δ_n true. Note that all of Δ_n are among Δ_i , with the possible exception of X itself. By hypothesis, line j is shiny. But no valuation can make ' \perp ' true, so no valuation can make all of Δ_j true. Since the sentences Δ_i are just the sentences Δ_j , no valuation can make all of Δ_i true. Since v makes all of Δ_n true, it must therefore make X false, and so make $\neg X$ true. So $\Delta_n \models \neg X$. QED

$\neg I$, $\perp E$, $\rightarrow I$, $\rightarrow E$, $\leftrightarrow I$, and $\leftrightarrow E$ are all rule-sound.

We leave these as exercises.

This establishes that all the basic rules of our proof system are rule-sound. Finally, we show:

All of the derived rules of our proof system are rule-sound.

Proof. Suppose that we used a derived rule to obtain some sentence, X , on line n of some TFL-proof, and that every earlier line is shiny. Every use of a derived rule can be replaced (at the cost of long-windedness) with multiple uses of basic rules. That is to say, we could have used basic rules to write X on some line $n + k$, without introducing any further assumptions. So, applying our individual results that all basic rules are rule-sound several times ($k + 1$ times, in fact), we can see that line $n + k$ is shiny. Hence the derived rule is rule-sound. QED

And that's that! We have shown that every rule—basic or otherwise—is rule-sound, which is all that we required to establish the Shininess Lemma, and hence the Soundness Theorem.

But it might help to round off this chapter if we repeat my informal explanation of what we have done. A formal proof is just a sequence—of arbitrary length—of applications of rules. We have shown that any application of any rule will not lead you astray. It follows (by induction) that no formal proof will lead you astray. That is: our proof system is sound.

Practice exercises

A. Complete the Lemmas left as exercises in this chapter. That is, show that the remaining rules are rule-sound.

PART XI

Modal logic (non- examenable)

CHAPTER 49

Introducing modal logic

Modal logic (ML) is the logic of *modalities*, ways in which a statement can be true. *Necessity* and *possibility* are two such modalities: a statement can be true, but it can also be necessarily true (true no matter how the world might have been). For instance, logical truths are not just true because of some accidental feature of the world, but true come what may. A possible statement may not actually be true, but it might have been true. We use \Box to express necessity, and \Diamond to express possibility. So you can read $\Box X$ as *It is necessarily the case that X*, and $\Diamond X$ as *It is possibly the case that X*.

There are lots of different kinds of necessity. It is *humanly impossible* for me to run at 100mph. Given the sorts of creatures that we are, no human can do that. But still, it isn't *physically impossible* for me to run that fast. We haven't got the technology to do it yet, but it is surely physically possible to swap my biological legs for robotic ones which could run at 100mph. By contrast, it is physically impossible for me to run faster than the speed of light. The laws of physics forbid any object from accelerating up to that speed. But even that isn't *logically impossible*. It isn't a contradiction to imagine that the laws of physics might have been different, and that they might have allowed objects to move faster than light.

Which kind of modality does ML deal with? *All of them!* ML is a very flexible tool. We start with a basic set of rules that govern \Box and \Diamond , and then add more rules to fit whatever kind of modality we

are interested in. In fact, ML is so flexible that we do not even have to think of \Box and \Diamond as expressing *necessity* and *possibility*. We might instead read \Box as expressing *provability*, so that $\Box X$ means *It is provable that X*, and $\Diamond X$ means *It is not refutable that X*. Similarly, we can interpret \Box to mean *S knows that X* or *S believes that X*. Or we might read \Box as expressing *moral obligation*, so that $\Box X$ means *It is morally obligatory that X*, and $\Diamond X$ means *It is morally permissible that X*. All we would need to do is cook up the right rules for these different readings of \Box and \Diamond .

A modal formula is one that includes modal operators such as \Box and \Diamond . Depending on the interpretation we assign to \Box and \Diamond , different modal formulas will be provable or valid. For instance, $\Box X \rightarrow X$ might say that “if X is necessary, it is true,” if \Box is interpreted as necessity. It might express “if X is known, then it is true,” if \Box expresses known truth. Under both these interpretations, $\Box X \rightarrow X$ is valid: All necessary propositions are true come what may, so are true in the actual world. And if a proposition is known to be true, it must be true (one can’t know something that’s false). However, when \Box is interpreted as “it is believed that” or “it ought to be the case that,” $\Box X \rightarrow X$ is not valid: We can believe false propositions. Not every proposition that ought to be true is in fact true, e.g., “Every murderer will be brought to justice.” This *ought* to be true, but it isn’t.

We will consider different kinds of systems of ML. They differ in the rules of proof allowed, and in the semantics we use to define our logical notions. The different systems we’ll consider are called **K**, **T**, **S4**, and **S5**. **K** is the basic system; everything that is valid or provable in **K** is also provable in the others. But there are some things that **K** does not prove, such as the formula $\Box A \rightarrow A$ for sentence letter A . So **K** is not an appropriate modal logic for necessity and possibility (where $\Box X \rightarrow X$ should be provable). This is provable in the system **T**, so **T** is more appropriate when dealing with necessity and possibility, but less appropriate when dealing with belief or obligation, since then $\Box X \rightarrow X$ should *not* (always) be provable. The perhaps best system of ML for necessity and possibility, and in any case the most widely accepted, is the strongest of the systems we consider, **S5**.

49.1 The Language of ML

In order to do modal logic, we have to do two things. First, we want to learn how to prove things in ML. Second, we want to see how to construct interpretations for ML. But before we can do either of these

things, we need to explain how to construct sentences in ML.

The language of ML is an extension of TFL. We could have started with FOL, which would have given us Quantified Modal Logic (QML). QML is much more powerful than ML, but it is also much, much more complicated. So we are going to keep things simple, and start with TFL.

Just like TFL, ML starts with an infinite stock of *atoms*. These are written as capital letters, with or without numerical subscripts: A , B , $\dots A_1$, B_1 , \dots . We then take all of the rules about how to make sentences from TFL, and add two more for \Box and \Diamond :

- (1) Every atom of ML is a sentence of ML.
- (2) If X is a sentence of ML, then $\neg X$ is a sentence of ML.
- (3) If X and Y are sentences of ML, then $(X \& Y)$ is a sentence of ML.
- (4) If X and Y are sentences of ML, then $(X \vee Y)$ is a sentence of ML.
- (5) If X and Y are sentences of ML, then $(X \rightarrow Y)$ is a sentence of ML.
- (6) If X and Y are sentences of ML, then $(X \leftrightarrow Y)$ is a sentence of ML.
- (7) If X is a sentence of ML, then $\Box X$ is a sentence of ML.
- (8) If X is a sentence of ML, then $\Diamond X$ is a sentence of ML.
- (9) Nothing else is a sentence of ML.

Here are some examples of ML sentences:

$$A, P \vee Q, \Box A, C \vee \Box D, \Box \Box (A \rightarrow R), \Box \Diamond (S \& (Z \leftrightarrow (\Box W \vee \Diamond Q)))$$

CHAPTER 50

Natural deduction for ML

Now that we know how to make sentences in ML, we can look at how to *prove* things in ML. We will use \vdash to express provability. So $X_1, X_2, \dots X_n \vdash Z$ means that Z can be proven from $X_1, X_2, \dots X_n$. However, we will be looking at a number of different systems of ML, and so it will be useful to add a subscript to indicate which system we are working with. So for example, if we want to say that we can prove Z from $X_1, X_2, \dots X_n$ *in system \mathbf{K}* , we will write: $X_1, X_2, \dots X_n \vdash_{\mathbf{K}} Z$.

50.1 System \mathbf{K}

We start with a particularly simple system called \mathbf{K} , in honour of the philosopher and logician Saul Kripke. \mathbf{K} includes all of the natural deduction rules from TFL, including the derived rules as well as the basic ones. \mathbf{K} then adds a special kind of subproof, plus two new basic rules for \Box .

The special kind of subproof looks like an ordinary subproof, except it has a \Box in its assumption line instead of a formula. We call them *strict subproofs*—they allow us to reason and prove things about alternate possibilities. What we can prove inside a strict subproof holds in any

alternate possibility, in particular, in alternate possibilities where the assumptions in force in our proof may not hold. In a strict subproofs, all assumptions are disregarded, and we are not allowed to appeal to any lines outside the strict subproof (except as allowed by the modal rules given below).

The $\Box I$ rule allows us to derive a formula $\Box X$ if we can derive X inside a strict subproof. It is our fundamental method of introducing \Box into proofs. The basic idea is simple enough: if X is a theorem, then $\Box X$ should be a theorem too. (Remember that to call X a theorem is to say that we can prove X without relying on any undischarged assumptions.)

Suppose we wanted to prove $\Box(A \rightarrow A)$. The first thing we need to do is prove that $A \rightarrow A$ is a theorem. You already know how to do that using TFL. You simply present a proof of $A \rightarrow A$ which doesn't start with any premises, like this:

1			A	
2			A	R 1
3		$A \rightarrow A$		$\rightarrow I$ 1-2

But to apply $\Box I$, we need to have proven the formula inside a strict subproof. Since our proof of $A \rightarrow A$ makes use of no assumptions at all, this is possible.

1			\Box	
2				A
3				A
4			$A \rightarrow A$	$\rightarrow I$ 2-3
5		$\Box(A \rightarrow A)$		$\Box I$ 1-4

m	<div style="border-bottom: 1px solid black; width: 100%;"></div> <div style="text-align: center; width: 100%;">□</div>	
n	<div style="border-left: 1px solid black; padding-left: 10px; vertical-align: middle;"> <div style="border-bottom: 1px solid black; width: 100%;"></div> <div style="text-align: center; width: 100%;">X</div> </div>	
	□X	□I $m-n$

No line above line m may be cited by any rule within the strict subproof begun at line m unless the rule explicitly allows it.

It is essential to emphasise that in strict subproof you cannot use any rule which appeals to anything you proved outside of the strict subproof. There are exceptions, e.g., the $\Box E$ rule below. These rules will explicitly state that they can be used inside strict subproofs and cite lines outside the strict subproof. This restriction is essential, otherwise we would get terrible results. For example, we could provide the following proof to vindicate $A \therefore \Box A$:

$$\begin{array}{r|l}
1 & A \\
\hline
2 & \square \\
3 & A \quad \text{incorrect use of R 1} \\
\hline
4 & \square A \quad \square I \text{ 2-3}
\end{array}$$

This is not a legitimate proof, because at line 3 we appealed to line 1, even though line 1 comes before the beginning of the strict subproof at line 2.

We said above that a strict subproof allows us to reason about arbitrary alternate possible situations. What can be proved in a strict subproof holds in all alternate possible situations, and so is necessary. This is the idea behind the $\Box I$ rule. On the other hand, if we've assumed that something is necessary, we have therewith assumed that it is true in all alternate possible situations. Hence, we have the rule $\Box E$:

m		$\Box X$	
			\Box
			X
n			$\Box E\ m$

$\Box E$ can only be applied if line m (containing $\Box A$) lies *outside* of the strict subproof in which line n falls, and this strict subproof is not itself part of a strict subproof not containing m .

$\Box E$ allows you to assert X inside a strict subproof if you have $\Box X$ outside the strict subproof. The restriction means that you can only do this in the first strict subproof, you cannot apply the $\Box E$ rule inside a nested strict subproof. So the following is not allowed:

1		$\Box X$	
2			\Box
3			\Box
4			X
			incorrect use of $\Box E\ 1$

The incorrect use of $\Box E$ on line 4 violates the condition, because although line 1 lies outside the strict subproof in which line 4 falls, the strict subproof containing line 4 lies inside the strict subproof beginning on line 2 which does not contain line 1.

Let's begin with an example.

1		$\Box A$	
2		$\Box B$	
3			\Box
4			A
5			B
6			$A \ \& \ B$
7		$\Box(A \wedge B)$	$\Box I\ 3-7$

We can also mix regular subproofs and strict subproofs:

1	$\Box(A \rightarrow B)$			
2		$\Box A$		
3			\Box	
4			A	$\Box E\ m$
5			$A \rightarrow B$	$\Box E\ 1$
6			B	$\rightarrow E\ 4, 5$
7		$\Box B$		
8	$\Box A \rightarrow \Box B$			$\rightarrow I\ 2-7$

This is called the *Distribution Rule*, because it tells us that \Box ‘distributes’ over \rightarrow .

The rules $\Box I$ and $\Box E$ look simple enough, and indeed **K** is a very simple system! But **K** is more powerful than you might have thought. You can prove a fair few things in it.

50.2 Possibility

In the last subsection, we looked at all of the basic rules for **K**. But you might have noticed that all of these rules were about necessity, \Box , and none of them were about possibility, \Diamond . That’s because we can *define* possibility in terms of necessity:

$$\Diamond X =_{df} \neg \Box \neg X$$

In other words, to say that X is *possibly true*, is to say that X is *not necessarily false*. As a result, it isn’t really essential to add a \Diamond , a special symbol for possibility, into system **K**. Still, the system will be *much* easier to use if we do, and so we will add the following definitional rules:

m	$\neg\Box\neg X$	
	$\Diamond X$	Def \Diamond m
m	$\Diamond X$	
	$\neg\Box\neg X$	Def \Diamond m

Importantly, you should not think of these rules as any real addition to **K**: they just record the way that \Diamond is defined in terms of \Box .

If we wanted, we could leave our rules for **K** here. But it will be helpful to add some *Modal Conversion* rules, which give us some more ways of flipping between \Box and \Diamond :

m	$\neg\Box X$	
	$\Diamond\neg X$	MC m
m	$\Diamond\neg X$	
	$\neg\Box X$	MC m
m	$\neg\Diamond X$	
	$\Box\neg X$	MC m
m	$\Box\neg X$	
	$\neg\Diamond X$	MC m

These Modal Conversion Rules are also no addition to the power of **K**, because they can be derived from the basic rules, along with the definition of \Diamond .

In system **K**, using Def \Diamond (or the modal conversion rules), one can prove $\Diamond A \leftrightarrow \neg\Box\neg A$. When laying out system **K**, we started with \Box as our primitive modal symbol, and then defined \Diamond in terms of it. But if we had preferred, we could have started with \Diamond as our primitive, and then defined \Box as follows: $\Box X =_{df} \neg\Diamond\neg X$. There is, then, no sense in which necessity is somehow more *fundamental* than possibility. Necessity and

possibility are exactly as fundamental as each other.

50.3 System T

So far we have focussed on **K**, which is a very simple modal system. **K** is so weak that it will not even let you prove X from $\Box X$. But if we are thinking of \Box as expressing *necessity*, then we will want to be able to make this inference: if X is *necessarily true*, then it must surely be *true*!

This leads us to a new system, **T**, which we get by adding the following rule to **K**:

$$\begin{array}{c|c} m & \Box X \\ n & X \end{array} \quad \text{RT } m$$

The line n on which rule RT is applied must *not* lie in a strict subproof that begins after line m .

The restriction on rule **T** is in a way the opposite of the restriction on $\Box E$: you can *only* use $\Box E$ in a nested strict subproof, but you cannot use **T** in a nested strict subproof.

We can prove things in **T** which we could not prove in **K**, e.g., $\Box A \rightarrow A$.

50.4 System S4

T allows you to strip away the necessity boxes: from $\Box X$, you may infer X . But what if we wanted to add extra boxes? That is, what if we wanted to go from $\Box X$ to $\Box\Box X$? Well, that would be no problem, if we had proved $\Box X$ by applying $\Box I$ to a strict subproof of X which itself does not use $\Box E$. In that case, X is a tautology, and by nesting the strict subproof inside another strict subproof and applying $\Box I$ again, we can

prove $\Box\Box X$. For example, we could prove $\Box\Box(P \rightarrow P)$ like this:

1			\Box			
2				\Box		
3					P	
4					P	R 3
5				$P \rightarrow P$	\rightarrow I 3–4	
6			$\Box(P \rightarrow P)$	\Box I 2–5		
7		$\Box\Box(P \rightarrow P)$	\Box I 1–6			

But what if we didn't prove $\Box X$ in this restricted way, but used $\Box E$ inside the strict subproof of X . If we put that strict subproof inside another strict subproof, the requirement of rule $\Box E$ to not cite a line containing $\Box X$ which lies in another strict subproof that has not yet concluded, is violated. Or what if $\Box X$ were just an assumption we started our proof with? Could we infer $\Box\Box X$ then? Not in **T**, we couldn't. And this might well strike you as a limitation of **T**, at least if we are reading \Box as expressing *necessity*. It seems intuitive that if X is necessarily true, then it couldn't have *failed* to be necessarily true.

This leads us to another new system, **S4**, which we get by adding the following rule to **T**:

m		$\Box X$	
			\Box
n		$\Box X$	R4 m

Note that **R4** can only be applied if line m (containing $\Box A$) lies outside of the strict subproof in which line n falls, and this strict subproof is not itself part of a strict subproof not containing n .

Rule **R4** looks just like $\Box E$, except that instead of yielding X from $\Box X$ it yields $\Box X$ inside a strict subproof. The restriction is the same, however: **R4** allows us to “import” $\Box X$ into a strict subproof, but not into a strict subproof itself nested inside a strict subproof. However, if

that is necessary, an additional application of **R4** would have the same result.

Now we can prove even more results. For instance:

1			$\Box A$	
2				\Box
3				$\Box A$ R4 1
4			$\Box \Box A$	$\Box I$ 2–3
5		$\Box A \rightarrow \Box \Box A$		$\rightarrow I$ 1–6

Similarly, we can prove $\Diamond \Diamond A \rightarrow \Diamond A$. This shows us that as well as letting us *add* extra *boxes*, **S4** lets us *delete* extra *diamonds*: from $\Diamond \Diamond X$, you can always infer $\Diamond X$.

50.5 System S5

In **S4**, we can always add a box in front of another box. But **S4** does not automatically let us add a box in front of a *diamond*. That is, **S4** does not generally permit the inference from $\Diamond X$ to $\Box \Diamond X$. But again, that might strike you as a shortcoming, at least if you are reading \Box and \Diamond as expressing *necessity* and *possibility*. It seems intuitive that if X is possibly true, then it couldn't have *failed* to be possibly true.

This leads us to our final modal system, **S5**, which we get by adding the following rule to **S4**:

m		$\neg \Box X$	
			\Box
n			$\neg \Box X$ R5 m

Rule **R5** can only be applied if line m (containing $\neg \Box X$) lies outside of the strict subproof in which line n falls, and this strict subproof is not itself part of a strict subproof not containing line m .

This rule allows us to show, for instance, that $\Diamond \Box A \vdash_{S5} \Box A$:

1	$\Diamond \Box A$	
	<hr/>	
2	$\neg \Box \neg \Box A$	Def \Diamond 1
3	$\neg \Box A$	
	<hr/>	
4	\Box	
	<hr/>	
5	$\neg \Box A$	R5 3
6	$\Box \neg \Box A$	\Box I 4–5
7	\perp	\neg E 2, 6
8	$\Box A$	IP 3–7

So, as well as adding boxes in front of diamonds, we can also delete diamonds in front of boxes.

We got **S5** just by adding the rule R5 rule to **S4**. In fact, we could have added rule R5 to **T** alone, and leave out rule R4). Everything we can prove by rule R4 can also be proved using RT together with R5.

For instance, here is a proof that shows $\Box A \vdash_{S5} \Box\Box A$ without using R4:

1	$\Box A$	
2	$\Box \neg \Box A$	
3	$\neg \Box A$	RT 2
4	\perp	$\neg E$ 1, 3
5	$\neg \Box \neg \Box A$	$\neg I$ 2–4
6	\Box	
7	$\neg \Box A$	
8	\Box	
9	$\neg \Box A$	R5 7
10	$\Box \neg \Box A$	$\Box I$ 8–9
11	$\neg \Box \neg \Box A$	R5 5
12	\perp	$\neg E$ 10, 11
13	$\Box A$	PbC 7–12
14	$\Box\Box A$	$\Box I$ 6–13

S5 is *strictly stronger* than **S4**: there are things which can be proved in **S5**, but not in **S4** (e.g., $\Diamond\Box A \rightarrow \Box A$).

The important point about **S5** can be put like this: if you have a long string of boxes and diamonds, in any combination whatsoever, you can delete all but the last of them. So for example, $\Diamond\Box\Diamond\Box\Box\Diamond\Box A$ can be simplified down to just $\Box A$.

Practice exercises

A. Provide proofs for the following:

- $\Box(A \& B) \vdash_K \Box A \& \Box B$
- $\Box A \& \Box B \vdash_K \Box(A \& B)$
- $\Box A \vee \Box B \vdash_K \Box(A \vee B)$
- $\Box(A \leftrightarrow B) \vdash_K \Box A \leftrightarrow \Box B$

B. Provide proofs for the following (without using Modal Conversion!):

1. $\neg \Box A \vdash_{\mathbf{K}} \Diamond \neg A$
2. $\Diamond \neg A \vdash_{\mathbf{K}} \neg \Box A$
3. $\neg \Diamond A \vdash_{\mathbf{K}} \Box \neg A$
4. $\Box \neg A \vdash_{\mathbf{K}} \neg \Diamond A$

C. Provide proofs of the following (and now feel free to use Modal Conversion!):

1. $\Box(A \rightarrow B), \Diamond A \vdash_{\mathbf{K}} \Diamond B$
2. $\Box A \vdash_{\mathbf{K}} \neg \Diamond \neg A$
3. $\neg \Diamond \neg A \vdash_{\mathbf{K}} \Box A$

D. Provide proofs for the following:

1. $P \vdash_{\mathbf{T}} \Diamond P$
2. $\vdash_{\mathbf{T}} (A \& B) \vee (\neg \Box A \vee \neg \Box B)$

E. Provide proofs for the following:

1. $\Box(\Box A \rightarrow B), \Box(\Box B \rightarrow C), \Box A \vdash_{\mathbf{S4}} \Box \Box C$
2. $\Box A \vdash_{\mathbf{S4}} \Box(\Box A \vee B)$
3. $\Diamond \Diamond A \vdash_{\mathbf{S4}} \Diamond A$

F. Provide proofs in **S5** for the following:

1. $\neg \Box \neg A, \Diamond B \vdash_{\mathbf{S5}} \Box(\Diamond A \& \Diamond B)$
2. $A \vdash_{\mathbf{S5}} \Box \Diamond A$
3. $\Diamond \Diamond A \vdash_{\mathbf{S5}} \Diamond A$

CHAPTER 51

Semantics for ML

So far, we have focussed on laying out various systems of Natural Deduction for ML. Now we will look at the *semantics* for ML. A semantics for a language is a method for assigning truth-values to the sentences in that language. So a semantics for ML is a method for assigning truth-values to the sentences of ML.

51.1 Interpretations of ML

The big idea behind the semantics for ML is this. In ML, sentences are not just true or false, full stop. A sentence is true or false *at a given possible world*, and a single sentence may well be true at some worlds and false at others. We then say that $\Box X$ is true iff X is true at *every* world, and $\Diamond X$ is true iff X is true at *some* world.

That's the big idea, but we need to refine it and make it more precise. To do this, we need to introduce the idea of an *interpretation* of ML. The first thing you need to include in an interpretation is a collection of *possible worlds*. Now, at this point you might well want to ask: What exactly is a possible world? The intuitive idea is that a possible world is another way that this world could have been. But what exactly does that mean? This is an excellent philosophical question, and we will look at it in a lot of detail later. But we do not need to worry too much about it right now. As far as the formal logic goes, possible worlds can be

anything you like. All that matters is that you supply each interpretation with a non-empty collection of things labelled **POSSIBLE WORLDS**.

Once you have chosen your collection of possible worlds, you need to find some way of determining which sentences of ML are true at which possible worlds. To do that, we need to introduce the notion of a *valuation function*. Those of you who have studied some maths will already be familiar with the general idea of a function. But for those of you who haven't, a function is a mathematical entity which maps arguments to values. That might sound a little bit abstract, but some familiar examples will help. Take the function $x + 1$. This is a function which takes in a number as argument, and then spits out the next number as value. So if you feed in the number 1 as an argument, the function $x + 1$ will spit out the number 2 as a value; if you feed in 2, it will spit out 3; if you feed in 3, it will spit out 4 ... Or here is another example: the function $x + y$. This time, you have to feed two arguments into this function if you want it to return a value: if you feed in 2 and 3 as your arguments, it spits out 5; if you feed in 1003 and 2005, it spits out 3008; and so on.

A valuation function for ML takes in a sentence and a world as its arguments, and then returns a truth-value as its value. So if v is a valuation function and w is a possible world, $v_w(X)$ is whatever truth-value v maps X and w to: if $v_w(X) = F$, then X is false at world w on valuation v ; if $v_w(X) = T$, then X is true at world w on valuation v .

These valuation functions are allowed to map any *atomic* sentence to any truth-value at any world. But there are rules about which truth-values more complex sentences get assigned at a world. Here are the rules for the connectives from TFL:

- (1) $v_w(\neg X) = T$ iff: $v_w(X) = F$
- (2) $v_w(X \& Y) = T$ iff: $v_w(X) = T$ and $v_w(Y) = T$
- (3) $v_w(X \vee Y) = T$ iff: $v_w(X) = T$ or $v_w(Y) = T$, or both
- (4) $v_w(X \rightarrow Y) = T$ iff: $v_w(X) = F$ or $v_w(Y) = T$, or both
- (5) $v_w(X \leftrightarrow Y) = T$ iff: $v_w(X) = T$ and $v_w(Y) = T$, or $v_w(X) = F$ and $v_w(Y) = F$

So far, these rules should all look very familiar. Essentially, they all work exactly like the truth-tables for TFL. The only difference is that these truth-table rules have to be applied over and over again, to one world at a time.

But what are the rules for the new modal operators, \Box and \Diamond ? The most obvious idea would be to give rules like these:

$$\nu_w(\Box X) = T \text{ iff } \forall w' (\nu_{w'}(X) = T)$$

$$\nu_w(\Diamond X) = T \text{ iff } \exists w' (\nu_{w'}(X) = T)$$

This is just the fancy formal way of writing out the idea that $\Box X$ is true at w just in case X is true at *every* world, and $\Diamond X$ is true at w just in case X is true at *some* world.

However, while these rules are nice and simple, they turn out not to be quite as useful as we would like. As we mentioned, ML is meant to be a very flexible tool. It is meant to be a general framework for dealing with lots of different kinds of necessity. As a result, we want our semantic rules for \Box and \Diamond to be a bit less rigid. We can do this by introducing another new idea: *accessibility relations*.

An accessibility relation, R , is a relation between possible worlds. Roughly, to say that Rw_1w_2 (in English: world w_1 *accesses* world w_2) is to say that w_2 is possible *relative to* w_1 . In other words, by introducing accessibility relations, we open up the idea that a given world might be possible relative to some worlds but not others. This turns out to be a *very* fruitful idea when studying modal systems. We can now give the following semantic rules for \Box and \Diamond :


$$(6) \quad \nu_{w_1}(\Box X) = T \text{ iff } \forall w_2 (Rw_1w_2 \rightarrow \nu_{w_2}(X) = T)$$

$$(7) \quad \nu_{w_1}(\Diamond X) = T \text{ iff } \exists w_2 (Rw_1w_2 \ \& \ \nu_{w_2}(X) = T)$$

Or in plain English: $\Box X$ is true in world w_1 iff X is true in every world that is possible relative to w_1 ; and $\Diamond X$ is true in world w_1 iff X is true in some world that is possible relative to w_1 .

So, there we have it. An interpretation for ML consists of three things: a collection of possible worlds, W ; an accessibility relation, R ; and a valuation function, ν . The collection of ‘possible worlds’ can really be a collection of anything you like. It really doesn’t matter, so long as W isn’t empty. (For many purposes, it is helpful just to take a collection of numbers to be your collection of worlds.) And for now, at least, R can be any relation between the worlds in W that you like. It could be a relation which every world in W bears to every world in W , or one which no world bears to any world, or anything in between. And lastly, ν can map any atomic sentence of ML to any truth-value at any world. All that matters is that it follows the rules (1)–(7) when it comes to the more complex sentences.

Let's look at an example. It is often helpful to present interpretations of ML as diagrams, like this:



Here is how to read the interpretation off from this diagram. It contains just two worlds, 1 and 2. The arrows between the worlds indicate the accessibility relation. So 1 and 2 both access 1, but neither 1 nor 2 accesses 2. The boxes at each world let us know which atomic sentences are true at each world: A is true at 1 but false at 2; B is false at 1 but true at 2. You may only write an atomic sentence or the negation of an atomic sentence into one of these boxes. We can figure out what truth-values the more complex sentences get at each world from that. For example, on this interpretation all of the following sentences are true at w_1 :

$$A \ \& \ \neg B, \ B \rightarrow A, \ \Diamond A, \ \Box \neg B$$

If you don't like thinking diagrammatically, then you can also present an interpretation like this:

$$W: 1, 2$$

$$R: \langle 1, 1 \rangle, \langle 2, 1 \rangle$$

$$v_1(A) = T, v_2(B) = F, v_2(A) = F, v_2(B) = T$$

You will get the chance to cook up some interpretations of your own shortly, when we start looking at *counter-interpretations*.

51.2 A Semantics for System K

We can now extend all of the semantic concepts of TFL to cover ML:

- ▷ $X_1, X_2, \dots, X_n \therefore Z$ is **MODALLY VALID** iff there is no world in any interpretation at which X_1, X_2, \dots, X_n are all true and Z is false.
- ▷ X is a **MODAL TRUTH** iff X is true at every world in every interpretation.
- ▷ X is a **MODAL CONTRADICTION** iff X is false at every world in every interpretation.
- ▷ X is **MODALLY SATISFIABLE** iff X is true at some world in some interpretation.

(From now on we will drop the explicit ‘modal’ qualifications, since they can be taken as read.)

We can also extend our use of \models . However, we need to add subscripts again, just as we did with \vdash . So, when we want to say that $X_1, X_2, \dots, X_n \therefore Z$ is valid, we will write: $X_1, X_2, \dots, X_n \models_{\mathbf{K}} Z$.

Let’s get more of a feel for this semantics by presenting some counter-interpretations. Consider the following (false) claim:

$$\neg A \models_{\mathbf{K}} \neg \Diamond A$$

In order to present a counter-interpretation to this claim, we need to cook up an interpretation which makes $\neg A$ true at some world w , and $\neg \Diamond A$ false at w . Here is one such interpretation, presented diagrammatically:



It is easy to see that this will work as a counter-interpretation for our claim. First, $\neg A$ is true at world 1. And second, $\neg \Diamond A$ is false at 1: A is true at 2, and 2 is accessible from 1. So there is some world in this interpretation where $\neg A$ is true and $\neg \Diamond A$ is false, so it is not the case that $\neg A \models_{\mathbf{K}} \neg \Diamond A$.

Why did we choose the subscript **K**? Well, it turns out that there is an important relationship between system **K** and the definition of validity we have just given. In particular, we have the following two results:

- ▷ If $X_1, X_2, \dots, X_n \vdash_{\mathbf{K}} Z$, then $X_1, X_2, \dots, X_n \models_{\mathbf{K}} Z$
- ▷ If $X_1, X_2, \dots, X_n \models_{\mathbf{K}} Z$, then $X_1, X_2, \dots, X_n \vdash_{\mathbf{K}} Z$

The first result is known as a *soundness* result, since it tells us that the rules of \mathbf{K} are good, sound rules: if you can vindicate an argument by giving a proof for it using system \mathbf{K} , then that argument really is valid. The second result is known as a *completeness* result, since it tells us that the rules of \mathbf{K} are broad enough to capture all of the valid arguments: if an argument is valid, then it will be possible to offer a proof in \mathbf{K} which vindicates it.

Now, it is one thing to state these results, quite another to prove them. However, we will not try to prove them here. But the idea behind the proof of soundness will perhaps make clearer how strict subproofs work.

In a strict subproof, we are not allowed to make use of any information from outside the strict subproof, except what we import into the strict subproof using $\Box E$. If we've assumed or proved $\Box X$, by $\Box E$, we can use X inside a strict subproof. And in \mathbf{K} , that is the only way to import a formula into a strict subproof. So everything that can be proved inside a strict subproof must follow from formulas X where outside the strict subproof we have $\Box X$. Let's imagine that we are reasoning about what's true in a possible world in some interpretation. If we know that $\Box X$ is true in that possible world, we know that X is true in all accessible worlds. So, everything proved inside a strict subproof is true in all accessible possible worlds. That is why $\Box I$ is a sound rule.

51.3 A Semantics for System T

A few moments ago, we said that system \mathbf{K} is sound and complete. Where does that leave the other modal systems we looked at, namely \mathbf{T} , $\mathbf{S4}$ and $\mathbf{S5}$? Well, they are all *unsound*, relative to the definition of validity we gave above. For example, all of these systems allow us to infer A from $\Box A$, even though $\Box A \not\models_{\mathbf{K}} A$.

Does that mean that these systems are a waste of time? Not at all! These systems are only unsound *relative to the definition of validity we gave above*. (Or to use symbols, they are unsound relative to $\models_{\mathbf{K}}$.) So when we are dealing with these stronger modal systems, we just need to modify our definition of validity to fit. This is where accessibility relations come in really handy.

When we introduced the idea of an accessibility relation, we said that it could be any relation between worlds that you like: you could have it relating every world to every world, no world to any world, or anything in between. That is how we were thinking of accessibility relations in our definition of \models_K . But if we wanted, we could start putting some restrictions on the accessibility relation. In particular, we might insist that it has to be *reflexive*:

$$\triangleright \forall w Rww$$

In English: every world accesses itself. Or in terms of relative possibility: every world is possible relative to itself. If we imposed this restriction, we could introduce a new consequence relation, \models_T , as follows:

$X_1, X_2, \dots, X_n \models_T Z$ iff there is no world in any interpretation *which has a reflexive accessibility relation*, at which X_1, X_2, \dots, X_n are all true and Z is false

We have attached the **T** subscript to \models because it turns out that system **T** is sound and complete relative to this new definition of validity:

- \triangleright If $X_1, X_2, \dots, X_n \vdash_T Z$, then $X_1, X_2, \dots, X_n \models_T Z$
- \triangleright If $X_1, X_2, \dots, X_n \models_T Z$, then $X_1, X_2, \dots, X_n \vdash_T Z$

As before, we will not try to prove these soundness and completeness results. However, it is relatively easy to see how insisting that the accessibility relation must be reflexive will vindicate the **RT** rule:

m	$\Box X$	
X		RT m

To see this, just imagine trying to cook up a counter-interpretation to this claim:

$$\Box X \models_T X$$

We would need to construct a world, w , at which $\Box X$ was true, but X was false. Now, if $\Box X$ is true at w , then X must be true at every world w accesses. But since the accessibility relation is reflexive, w accesses w . So X must be true at w . But now X must be true *and* false at w . Contradiction!

51.4 A Semantics for S4

How else might we tweak our definition of validity? Well, we might also stipulate that the accessibility relation has to be *transitive*:

$$\triangleright \forall w_1 \forall w_2 \forall w_3 ((Rw_1 w_2 \ \& \ Rw_2 w_3) \rightarrow Rw_1 w_3)$$

In English: if w_1 accesses w_2 , and w_2 accesses w_3 , then w_1 accesses w_3 . Or in terms of relative possibility: if w_3 is possible relative to w_2 , and w_2 is possible relative to w_1 , then w_3 is possible relative to w_1 . If we added this restriction on our accessibility relation, we could introduce a new consequence relation, \models_{S4} , as follows:

$X_1, X_2, \dots X_n \models_{S4} Z$ iff there is no world in any interpretation *which has a reflexive and transitive accessibility relation*, at which $X_1, X_2, \dots X_n$ are all true and Z is false

We have attached the **S4** subscript to \models because it turns out that system **S4** is sound and complete relative to this new definition of validity:

- \triangleright If $X_1, X_2, \dots X_n \vdash_{S4} Z$, then $X_1, X_2, \dots X_n \models_{S4} Z$
- \triangleright If $X_1, X_2, \dots X_n \models_{S4} Z$, then $X_1, X_2, \dots X_n \vdash_{S4} Z$

As before, we will not try to prove these soundness and completeness results. However, it is relatively easy to see how insisting that the accessibility relation must be transitive will vindicate the **S4** rule:

$$\begin{array}{c|c|c} m & \Box X & \\ \hline & \Box & \\ & \hline & \Box X & R4 \ m \end{array}$$

The idea behind strict subproofs, remember, is that they are ways to prove things that must be true in all accessible worlds. So the **R4** rule means that whenever $\Box X$ is true, $\Box X$ must also be true in every accessible world. In other words, we must have $\Box X \models_{S4} \Box \Box X$.

To see this, just imagine trying to cook up a counter-interpretation to this claim:

$$\Box X \models_{S4} \Box \Box X$$

We would need to construct a world, w_1 , at which $\Box X$ was true, but $\Box\Box X$ was false. Now, if $\Box\Box X$ is false at w_1 , then w_1 must access some world, w_2 , at which $\Box X$ is false. Equally, if $\Box X$ is false at w_2 , then w_2 must access some world, w_3 , at which X is false. We just said that w_1 accesses w_2 , and w_2 accesses w_3 . So since we are now insisting that the accessibility relation be transitive, w_1 must access w_3 . And as $\Box X$ is true at w_1 , and w_3 is accessible from w_1 , it follows that X must be true at w_3 . So X is true *and* false at w_3 . Contradiction!

51.5 A Semantics for S5

Let's put one more restriction on the accessibility relation. This time, let's insist that it must also be *symmetric*:

$$\triangleright \forall w_1 \forall w_2 (Rw_1 w_2 \rightarrow R w_2 w_1)$$

In English: if w_1 accesses w_2 , then w_2 accesses w_1 . Or in terms of relative possibility: if w_2 is possible relative to w_1 , then w_1 is possible relative to w_2 . Logicians call a relation that is reflexive, symmetric, and transitive an *equivalence* relation. We can now define a new consequence relation, \models_{S5} , as follows:

$X_1, X_2, \dots, X_n \models_{S5} Z$ iff there is no world in any interpretation *whose accessibility relation is an equivalence relation*, at which X_1, X_2, \dots, X_n are all true and Z is false

We have attached the **S5** subscript to \models because it turns out that system **S5** is sound and complete relative to this new definition of validity:

- \triangleright If $X_1, X_2, \dots, X_n \vdash_{S5} Z$, then $X_1, X_2, \dots, X_n \models_{S5} Z$
- \triangleright If $X_1, X_2, \dots, X_n \models_{S5} Z$, then $X_1, X_2, \dots, X_n \vdash_{S5} Z$

As before, we will not try to prove these soundness and completeness results here. However, it is relatively easy to see how insisting that the accessibility relation must be an equivalence relation will vindicate the **R5** rule:

m		$\neg\Box X$	
			\Box
			$\neg\Box X$
			R5 m

The rule says that if X is not necessary, i.e., false in some accessible world, it is also not necessary in any accessible possible world, i.e., we have $\neg\Box X \vdash_{S5} \Box\neg\Box X$.

To see this, just imagine trying to cook up a counter-interpretation to this claim:

$$\neg\Box X \vdash_{S5} \Box\neg\Box X$$

We would need to construct a world, w_1 , at which $\neg\Box X$ was true, but $\Box\neg\Box X$ was false. Now, if $\neg\Box X$ is true at w_1 , then w_1 must access some world, w_2 , at which X is false. Equally, if $\Box\neg\Box X$ is false at w_1 , then w_1 must access some world, w_3 , at which $\neg\Box X$ is false. Since we are now insisting that the accessibility relation is an equivalence relation, and hence symmetric, we can infer that w_3 accesses w_1 . Thus, w_3 accesses w_1 , and w_1 accesses w_2 . Again, since we are now insisting that the accessibility relation is an equivalence relation, and hence transitive, we can infer that w_3 accesses w_2 . But earlier we said that $\neg\Box X$ is false at w_3 , which implies that X is true at every world which w_3 accesses. So X is true *and* false at w_2 . Contradiction!

In the definition of \vdash_{S5} , we stipulated that the accessibility relation must be an equivalence relation. But it turns out that there is another way of getting a notion of validity fit for **S5**. Rather than stipulating that the accessibility relation be an equivalence relation, we can instead stipulate that it be a *universal* relation:

$$\triangleright \forall w_1 \forall w_2 R w_1 w_2$$

In English: every world accesses every world. Or in terms of relative possibility: every world is possible relative to every world. Using this restriction on the accessibility relation, we could have defined \vdash_{S5} like this:

$X_1, X_2, \dots X_n \vdash_{S5} Z$ iff there is no world in any interpretation which has a *universal accessibility relation*, at which $X_1, X_2, \dots X_n$ are all true and Z is false.

If we defined \models_{S5} like this, we would still get the same soundness and completeness results for **S5**. What does this tell us? Well, it means that if we are dealing with a notion of necessity according to which *every* world is possible relative to *every* world, then we should use **S5**. What is more, most philosophers assume that the notions of necessity that they are most concerned with, like *logical necessity* and *metaphysical necessity*, are of exactly this kind. So **S5** is the modal system that most philosophers use most of the time.

Practice exercises

A. Present counter-interpretations to the following false claims:

1. $\neg P \models_K \neg \Diamond P$
2. $\Box(P \vee Q) \models_K \Box P \vee \Box Q$
3. $\models_K \neg \Box(A \ \& \ \neg A)$
4. $\Box A \models_K A$

B. Present counter-interpretations to the following false claims:

1. $\Diamond A \models_{S4} \Box \Diamond A$
2. $\Diamond A, \Box(\Diamond A \rightarrow B) \models_{S4} \Box B$

C. Present counter-interpretations to the following false claims:

1. $\Box(M \rightarrow O), \Diamond M \models_T O$
2. $\Box A \models_T \Box \Box A$

Further reading

Modal logic is a large subfield of logic. We have only scratched the surface. If you want to learn more about modal logic, here are some textbooks you might consult.

- Hughes, G. E., & Cresswell, M. J. (1996). *A New Introduction to Modal Logic*, Oxford: Routledge.
- Priest, G. (2008). *An Introduction to Non-Classical Logic*, 2nd ed., Cambridge: Cambridge University Press.
- Garson, J. W. (2013). *Modal Logic for Philosophers*, 2nd ed., Cambridge: Cambridge University Press.

None of these authors formulate their modal proof systems in quite the way we did, but the closest formulation is given by Garson.

Appendices

APPENDIX A

Symbolic notation

1.1 Alternative nomenclature

Truth-functional logic. TFL goes by other names. Sometimes it is called *sentential logic*, because it deals fundamentally with sentences. Sometimes it is called *propositional logic*, on the idea that it deals fundamentally with propositions. We have stuck with *truth-functional logic*, to emphasize the fact that it deals only with assignments of truth and falsity to sentences, and that its connectives are all truth-functional.

First-order logic. FOL goes by other names. Sometimes it is called *predicate logic*, because it allows us to apply predicates to objects. Sometimes it is called *quantified logic*, because it makes use of quantifiers.

Formulas. Some texts call formulas *well-formed formulas*. Since ‘well-formed formula’ is such a long and cumbersome phrase, they then abbreviate this as *wff*. This is both barbarous and unnecessary (such texts do not countenance ‘ill-formed formulas’). We have stuck with ‘formula’.

In §6, we defined *sentences* of TFL. These are also sometimes called ‘formulas’ (or ‘well-formed formulas’) since in TFL, unlike FOL, there is no distinction between a formula and a sentence.

Valuations. Some texts call valuations *truth-assignments*, or *truth-value assignments*.

Expressive adequacy. Some texts describe TFL as *truth-functionally complete*, rather than expressively adequate.

n-place predicates. We have chosen to call predicates ‘one-place’, ‘two-place’, ‘three-place’, etc. Other texts respectively call them ‘monadic’, ‘dyadic’, ‘triadic’, etc. Still other texts call them ‘unary’, ‘binary’, ‘ternary’, etc.

Names. In FOL, we have used ‘*a*’, ‘*b*’, ‘*c*’, for names. Some texts call these ‘constants’. Other texts do not mark any difference between names and variables in the syntax. Those texts focus simply on whether the symbol occurs *bound* or *unbound*.

Domains. Some texts describe a domain as a ‘domain of discourse’, or a ‘universe of discourse’.

1.2 Alternative symbols

In the history of formal logic, different symbols have been used at different times and by different authors. Often, authors were forced to use notation that their printers could typeset.

This appendix presents some common symbols, so that you can recognize them if you encounter them in an article or in another book.

Negation. Two commonly used symbols are the *hoe*, ‘ \neg ’, and the *swung dash* or *tilda*, ‘ \sim ’. There are some issues typing ‘ \neg ’ on a keyboard, and ‘ \sim ’ is perfectly acceptable for you to use. In some more advanced formal systems it is necessary to distinguish between two kinds of negation; the distinction is sometimes represented by using both ‘ \neg ’ and ‘ \sim ’. Older texts sometimes indicate negation by a line over the formula being negated, e.g., $\overline{A \& B}$. Some texts use ‘ $x \neq y$ ’ to abbreviate ‘ $\neg x = y$ ’.

Disjunction. The symbol ‘ \vee ’ is typically used to symbolize inclusive disjunction. One etymology is from the Latin word ‘vel’, meaning ‘or’.

Conjunction. The two symbols commonly used for conjunction are *wedge*, ‘ \wedge ’, and *ampersand*, ‘ $\&$ ’. The ampersand is a decorative form of the Latin word ‘et’, which means ‘and’. (Its etymology still lingers in certain fonts, particularly in italic fonts; thus an italic ampersand might appear as ‘ $\&$ ’.) We have chosen to use it to allow for easier typing on a keyboard during these online-heavy times. However there are some substantial reservations about this choice. This symbol is commonly used in natural English writing (e.g. ‘Smith & Sons’), and so even though it is a natural choice, many logicians use a different symbol to avoid confusion between the object and metalanguage: as a symbol in a formal system, the ampersand is not the English word ‘ $\&$ ’. The most common choice now is ‘ \wedge ’, which is a counterpart to the symbol used for disjunction. Sometimes a single dot, ‘ \cdot ’, is used. In some older texts, there is no symbol for conjunction at all; ‘ A and B ’ is simply written ‘ AB ’.

Material Conditional. There are two common symbols for the material conditional: the *arrow*, ‘ \rightarrow ’, and the *hook*, ‘ \supset ’.

Material Biconditional. The *double-headed arrow*, ‘ \leftrightarrow ’, is used in systems that use the arrow to represent the material conditional. Systems that use the hook for the conditional typically use the *triple bar*, ‘ \equiv ’, for the biconditional.

Quantifiers. The universal quantifier is typically symbolized as a rotated ‘ A ’, and the existential quantifier as a rotated, ‘ E ’. In some texts, there is no separate symbol for the universal quantifier. Instead, the variable is just written in parentheses in front of the formula that it binds. For example, they might write ‘ $(x)Px$ ’ where we would write ‘ $\forall x Px$ ’.

These alternative typographies are summarised below:

negation	\neg, \sim
conjunction	$\wedge, \&, \cdot$
disjunction	\vee
conditional	\rightarrow, \supset
biconditional	\leftrightarrow, \equiv
universal quantifier	$\forall x, (x)$

APPENDIX B

Alternative proof systems

In formulating our natural deduction system, we treated certain rules of natural deduction as *basic*, and others as *derived*. However, we could equally well have taken various different rules as basic or derived. We will illustrate this point by considering some alternative treatments of disjunction, negation, and the quantifiers. We will also explain why we have made the choices that we have.

2.1 Alternative disjunction elimination

Some systems take DS as their basic rule for disjunction elimination. Such systems can then treat the $\vee E$ rule as a derived rule. For they might offer the following proof scheme:

m	$X \vee Y$	
i	X	
	\vdots	
j	C	
k	Y	
	\vdots	
l	C	
n	$X \rightarrow C$	$\rightarrow I\ i-j$
$n+1$	$Y \rightarrow C$	$\rightarrow I\ k-l$
$n+2$	$C \vee \neg C$	LEM
$n+3$	C	
$n+4$	C	R $n+3$
$n+5$	$\neg C$	
$n+6$	X	
$n+7$	C	$\rightarrow E\ n, n+6$
$n+8$	\perp	$\perp I\ n+7, n+5$
$n+9$	$\neg X$	$\neg I\ n+6-n+8$
$n+10$	Y	DS $m, n+9$
$n+11$	C	$\rightarrow E\ n+1, n+10$
$n+12$	C	$\vee E\ n+2, n+3-n+4, n+5-n+11$

So why did we choose to take $\vee E$ as basic, rather than DS?¹ Our reasoning is that DS involves the use of ‘ \neg ’ in the statement of the rule. It is in some sense ‘cleaner’ for our disjunction elimination rule to avoid mentioning *other* connectives. The rule $\vee E$ we use is also closely connected to the rule $\exists E$. Whereas there is no such analogy with DS.

¹P.D. Magnus’s original version of this book went the other way.

2.2 Alternative negation rules

Some systems take the following rule as their basic negation introduction rule:

$$\begin{array}{c|c|c} m & & X \\ n-1 & & Y \\ n & & \neg Y \\ & \neg X & \neg I^* \ m-n \end{array}$$

and the following as their basic negation elimination rule:

$$\begin{array}{c|c|c} m & & -X \\ n-1 & & Y \\ n & & -Y \\ & X & -E^*_{m-n} \end{array}$$

Using these two rules, we could have derived all of the rules governing negation and contradiction that we have taken as basic (i.e. $\perp I$, $\perp E$, $\neg I$ and $\neg E$). Indeed, we could have avoided all use of the symbol ‘ \perp ’ altogether. Negation would have had a single introduction and elimination rule, and would have behaved much more like the other connectives.

The resulting system would have had fewer rules than ours. So why did we chose to separate out contradiction, and to use an explicit rule LEM?²

Our first reason is that adding the symbol ‘ \perp ’ to our natural deduction system makes proofs considerably easier to work with.

Our second reason is that a lot of fascinating philosophical discussion has focussed on the acceptability or otherwise of *law of excluded middle* (i.e. LEM) and *explosion* (i.e. $\perp E$). By treating these as separate rules in the proof system, we will be in a better position to engage with that philosophical discussion. In particular: having invoked these rules explicitly, it will be much easier for us to know what a system which lacked these rules would look like.

²Again, P.D. Magnus's original version of this book went the other way.

2.3 Alternative quantification rules

An alternative approach to the quantifiers is to take as basic the rules for $\forall I$ and $\forall E$ from §38, and also two CQ rule which allow us to move from $\forall x\neg X$ to $\neg\exists xX$ and vice versa.³

Taking only these rules as basic, we could have derived the $\exists I$ and $\exists E$ rules provided in §38. To derive the $\exists I$ rule is fairly simple. Suppose X contains the name c , and contains no instances of the variable x , and that we want to do the following:

m	$X(\dots c \dots c \dots)$
k	$\exists xX(\dots x \dots c \dots)$

This is not yet permitted, since in this new system, we do not have the $\exists I$ rule. We can, however, offer the following:

m	$X(\dots c \dots c \dots)$	
$m + 1$	$\neg\exists xX(\dots x \dots c \dots)$	
$m + 2$	$\forall x\neg X(\dots x \dots c \dots)$	CQ $m + 1$
$m + 3$	$\neg X(\dots c \dots c \dots)$	$\forall E\ m + 2$
$m + 4$	\perp	$\perp I\ m, m + 3$
$m + 5$	$\neg\neg\exists xX(\dots x \dots c \dots)$	$\neg I\ m + 1 - m + 4$
$m + 6$	$\exists xX(\dots x \dots c \dots)$	DNE $m + 5$

To derive the $\exists E$ rule is rather more subtle. This is because the $\exists E$ rule has an important constraint (as, indeed, does the $\forall I$ rule), and we need to make sure that we are respecting it. So, suppose we are in a situation where we *want* to do the following:

m	$\exists xX(\dots x \dots x \dots)$
i	$X(\dots c \dots c \dots)$
j	Y
k	Y

³Warren Goldfarb follows this line in *Deductive Logic*, 2003, Hackett Publishing Co.

where c does not occur in any undischarged assumptions, or in Y , or in $\exists xX(\dots x \dots x \dots)$. Ordinarily, we would be allowed to use the $\exists E$ rule; but we are not here assuming that we have access to this rule as a basic rule. Nevertheless, we could offer the following, more complicated derivation:

m	$\exists xX(\dots x \dots x \dots)$	
i	$X(\dots c \dots c \dots)$	
j	Y	
k	$X(\dots c \dots c \dots) \rightarrow Y$	$\rightarrow I\ i-j$
$k+1$	$\neg Y$	
$k+2$	$\neg X(\dots c \dots c \dots)$	MT $k, k+1$
$k+3$	$\forall x\neg X(\dots x \dots x \dots)$	$\forall I\ k+2$
$k+4$	$\neg\exists xX(\dots x \dots x \dots)$	CQ $k+3$
$k+5$	\perp	$\perp I\ m, k+4$
$k+6$	$\neg\neg Y$	$\neg I\ k+1-k+5$
$k+7$	Y	DNE $k+6$

We are permitted to use $\forall I$ on line $k+3$ because c does not occur in any undischarged assumptions or in Y . The entries on lines $k+4$ and $k+1$ contradict each other, because c does not occur in $\exists xX(\dots x \dots x \dots)$.

Armed with these derived rules, we could now go on to derive the two remaining CQ rules, exactly as in §40.

So, why did we start with all of the quantifier rules as basic, and then derive the CQ rules?

Our first reason is that it seems more intuitive to treat the quantifiers as on a par with one another, giving them their own basic rules for introduction and elimination.

Our second reason relates to the discussion of alternative negation rules. In the derivations of the rules of $\exists I$ and $\exists E$ that we have offered in this section, we invoked DNE. This is a derived rule, whose derivation essentially depends upon the use of LEM. But, as we mentioned earlier, LEM is a contentious rule. So, if we want to move to a system

which abandons LEM, but which still allows us to use existential quantifiers, we will want to take the introduction and elimination rules for the quantifiers as basic, and take the CQ rules as derived. (Indeed, in a system without LEM, we will be *unable* to derive the CQ rule which moves from $\neg\forall xX$ to $\exists x\neg X$.)

APPENDIX C

Quick reference

3.1 Sentences of TFL

Definition of being a sentence of TFL:

1. Every atomic sentence is a sentence.
 $\triangleright A, B, C \dots, W$, or with subscripts $A_1, B_3, A_{100}, J_{375}$
2. If X is a sentence, then $\neg X$ is a sentence.
3. If X and Y are sentences, then $(X \& Y)$ is a sentence.
4. If X and Y are sentences, then $(X \vee Y)$ is a sentence.
5. If X and Y are sentences, then $(X \rightarrow Y)$ is a sentence.
6. If X and Y are sentences, then $(X \leftrightarrow Y)$ is a sentence.
7. Nothing else is a sentence.

3.2 Truth Rules for Connectives in TFL

\neg :	$\begin{array}{l} T \rightsquigarrow F \\ F \rightsquigarrow T \end{array}$	$\&$:	$\begin{array}{l} T, T \rightsquigarrow T \\ T, F \rightsquigarrow F \\ F, T \rightsquigarrow F \\ F, F \rightsquigarrow F \end{array}$	\vee :	$\begin{array}{l} T, T \rightsquigarrow T \\ T, F \rightsquigarrow T \\ F, T \rightsquigarrow T \\ F, F \rightsquigarrow F \end{array}$

\rightarrow :	T, T \leadsto T	\leftrightarrow :	T, T \leadsto T
	T, F \leadsto F		T, F \leadsto F
	F, T \leadsto T		F, T \leadsto F
	F, F \leadsto T		F, F \leadsto T

3.3 Symbolization

Rough Meaning of the TFL Connectives

symbol	name	rough meaning
\neg	negation	'It is not the case that...'
$\&$	conjunction	'... and ...'
\vee	disjunction	'... or ...'
\rightarrow	conditional	'If ... then ...'
\leftrightarrow	biconditional	'... if and only if ...'

Sentential Connectives

It is not the case that P	$\neg P$
P or Q	$(P \vee Q)$
P and Q	$(P \& Q)$
If P, then Q	$(P \rightarrow Q)$
P if and only if Q	$(P \leftrightarrow Q)$

Further symbolisation help:

Neither P nor Q	$\neg(P \vee Q)$ or $(\neg P \& \neg Q)$
P but Q	$(P \& Q)$
P unless Q	$(P \vee Q)$
P only if Q	$(P \rightarrow Q)$

Predicates

All Fs are Gs	$\forall x(Fx \rightarrow Gx)$
Some Fs are Gs	$\exists x(Fx \& Gx)$
Not all Fs are Gs	$\neg\forall x(Fx \rightarrow Gx)$ or $\exists x(Fx \& \neg Gx)$
No Fs are Gs	$\forall x(Fx \rightarrow \neg Gx)$ or $\neg\exists x(Fx \& Gx)$

Identity

Only c is G	$\forall x(Gx \rightarrow x=c)$ or perhaps \leftrightarrow .
Everything besides c is G	$\forall x(\neg x=c \rightarrow Gx)$
The F is G	$\exists x(Fx \& \forall y(Fy \rightarrow x=y) \& Gx)$
It is not the case that the F is G	$\neg \exists x(Fx \& \forall y(Fy \rightarrow x=y) \& Gx)$
The F is non- G	$\exists x(Fx \& \forall y(Fy \rightarrow x=y) \& \neg Gx)$

3.4 Using identity to symbolize quantities

There are at least _____ Fs.

- one: $\exists xFx$
 two: $\exists x_1\exists x_2(Fx_1 \& Fx_2 \& \neg x_1=x_2)$
 three: $\exists x_1\exists x_2\exists x_3(Fx_1 \& Fx_2 \& Fx_3 \& \neg x_1=x_2 \& \neg x_1=x_3 \& \neg x_2=x_3)$
 four: $\exists x_1\exists x_2\exists x_3\exists x_4(Fx_1 \& Fx_2 \& Fx_3 \& Fx_4 \& \neg x_1=x_2 \& \neg x_1=x_3 \& \neg x_1=x_4 \& \neg x_2=x_3 \& \neg x_2=x_4 \& \neg x_3=x_4)$
 n : $\exists x_1 \dots \exists x_n(Fx_1 \& \dots \& Fx_n \& \neg x_1=x_2 \& \dots \& \neg x_{n-1}=x_n)$

There are at most _____ Fs.

One way to say ‘there are at most n Fs’ is to put a negation sign in front of the symbolization for ‘there are at least $n + 1$ Fs’. Equivalently, we can offer:

- one: $\forall x_1\forall x_2[(Fx_1 \& Fx_2) \rightarrow x_1=x_2]$
 two: $\forall x_1\forall x_2\forall x_3[(Fx_1 \& Fx_2 \& Fx_3) \rightarrow (x_1=x_2 \vee x_1=x_3 \vee x_2=x_3)]$
 three: $\forall x_1\forall x_2\forall x_3\forall x_4[(Fx_1 \& Fx_2 \& Fx_3 \& Fx_4) \rightarrow (x_1=x_2 \vee x_1=x_3 \vee x_1=x_4 \vee x_2=x_3 \vee x_2=x_4 \vee x_3=x_4)]$
 n : $\forall x_1 \dots \forall x_{n+1}[(Fx_1 \& \dots \& Fx_{n+1}) \rightarrow (x_1=x_2 \vee \dots \vee x_n=x_{n+1})]$

There are exactly _____ Fs.

One way to say ‘there are exactly n Fs’ is to conjoin two of the symbolizations above and say ‘there are at least n Fs and there are at most n Fs.’ The following equivalent formulae are shorter:

- zero: $\forall x\neg Fx$
 one: $\exists x[Fx \& \forall y(Fy \rightarrow x=y)]$
 two: $\exists x_1\exists x_2[Fx_1 \& Fx_2 \& \neg x_1=x_2 \& \forall y(Fy \rightarrow (y=x_1 \vee y=x_2))]$
 three: $\exists x_1\exists x_2\exists x_3[Fx_1 \& Fx_2 \& Fx_3 \& \neg x_1=x_2 \& \neg x_1=x_3 \& \neg x_2=x_3 \& \forall y(Fy \rightarrow (y=x_1 \vee y=x_2 \vee y=x_3))]$

$$n: \exists x_1 \dots \exists x_n [Fx_1 \& \dots \& Fx_n \& \neg x_1 = x_2 \& \dots \& \neg x_{n-1} = x_n \& \\ \forall y (Fy \rightarrow (y = x_1 \vee \dots \vee y = x_n))]$$

3.5 Basic deduction rules for TFL

Conjunction

m	X		m	$X \& Y$	
n	Y			X	$\&E\ m$
	$X \& Y$	$\&I\ m, n$	m	$X \& Y$	
				Y	$\&E\ m$

Disjunction

m	X		m	$X \vee Y$	
	$X \vee Y$	$\vee I\ m$	i	X	
				\vdots	
m	X		j	Z	
	$Y \vee X$	$\vee I\ m$	k	Y	
				\vdots	
			l	Z	
				Z	$\vee E\ m, i-j, k-l$

Conditional

m	X		m	$X \rightarrow Y$	
	\vdots		n	X	
n	Y			Y	$\rightarrow E\ m, n$
	$X \rightarrow Y$	$\rightarrow I\ m-n$			

Contradiction

m	X		m	\perp	
n	$\neg X$			X	$\perp E\ m$
	\perp	$\perp I\ m, n$			

Negation

m			X	
			\vdots	
n			\perp	
			$\neg X$	$\neg\text{I } m-n$

m			$\neg X$	
			\vdots	
n			\perp	
			X	PbC $m-n$

Reiteration

m		X	
		X	R m

Law of Excluded Middle

	$X \vee \neg X$	LEM
--	-----------------	-----

3.6 Derived rules for TFL

Disjunctive syllogism

m	$X \vee Y$	
n	$\neg X$	
	Y	DS m, n

m	$X \vee Y$	
n	$\neg Y$	
	X	DS m, n

Modus Tollens

m	$X \rightarrow Y$	
n	$\neg Y$	
	$\neg X$	MT m, n

Double-negation elimination

m	$\neg\neg X$	
	X	DNE m

De Morgan Rules

m	$\neg(X \vee Y)$	
	$\neg X \ \& \ \neg Y$	DeM m

m	$\neg X \ \& \ \neg Y$	
	$\neg(X \vee Y)$	DeM m

m	$\neg(X \ \& \ Y)$	
	$\neg X \vee \neg Y$	DeM m

m	$\neg X \vee \neg Y$	
	$\neg(X \ \& \ Y)$	DeM m

3.7 Basic deduction rules for FOL

Universal elimination

m	$\forall x X(\dots x \dots x \dots)$	
	$X(\dots c \dots c \dots)$	$\forall E\ m$

x must not occur in $X(\dots c \dots c \dots)$

Universal introduction

m	$X(\dots c \dots c \dots)$	
	$\forall x X(\dots x \dots x \dots)$	$\forall I\ m$

c must not occur in any undischarged assumption

x must not occur in $X(\dots c \dots c \dots)$

Existential elimination

m	$\exists x X(\dots x \dots x \dots)$	
i	$X(\dots c \dots c \dots)$	
	\vdots	
j	Y	
	Y	$\exists E\ m, i-j$

Existential introduction

m	$X(\dots c \dots c \dots)$	
	$\exists x X(\dots x \dots c \dots)$	$\exists I\ m$

c must not occur in any undischarged assumption, in $\exists x X(\dots x \dots x \dots)$, or in Y

Identity introduction

$c = c$	$= I$
---------	-------

Identity elimination

m	$a = b$	
n	$X(\dots a \dots a \dots)$	
	$X(\dots b \dots a \dots)$	$= E\ m, n$

m	$a = b$	
n	$X(\dots b \dots b \dots)$	
	$X(\dots a \dots b \dots)$	$= E\ m, n$

3.8 Derived rules for FOL

$$m \left| \begin{array}{l} \forall x \neg X \\ \neg \exists x X \end{array} \right. \quad \text{CQ } m$$

$$m \left| \begin{array}{l} \neg \exists x X \\ \forall x \neg X \end{array} \right. \quad \text{CQ } m$$

$$m \left| \begin{array}{l} \exists x \neg X \\ \neg \forall x X \end{array} \right. \quad \text{CQ } m$$

$$m \left| \begin{array}{l} \neg \forall x X \\ \exists x \neg X \end{array} \right. \quad \text{CQ } m$$

Glossary

antecedent The sentence on the left side of a **conditional**.

argument a connected series of sentences, divided into **premises** and **conclusion**.

atomic sentence A sentence used to represent a basic sentence; a single letter in TFL, or a predicate symbol followed by names in FOL.

biconditional The symbol \leftrightarrow , used to represent words and phrases that function like the English phrase “if and only if”; or a sentence formed using this connective..

bound variable an occurrence of a variable in a **formula of FOL** which is in the scope of a quantifier followed by the same variable.

completeness A property held by logical systems if and only if \models implies \vdash . 170

conclusion the last sentence in an **argument**.

conclusion indicator a word or phrase such as “therefore” used to indicate that what follows is the conclusion of an argument.

conditional The symbol \rightarrow , used to represent words and phrases that function like the English phrase “if ... then ...”; a sentence formed by using this symbol.

conjunct A sentence joined to another by a **conjunction**.

conjunction The symbol $\&$, used to represent words and phrases that function like the English word “and”; or a sentence formed using that symbol.

conjunctive normal form (DNF) a sentence which is a conjunction of disjunctions of atomic sentences or negated atomic sentences.

connective A logical operator in TFL used to combine **atomic sentences** into larger sentences.

consequent The sentence on the right side of a **conditional**.

consistency (in TFL) A property held by sentences if and only if there is some **valuation** that makes all the sentences true.

contingent sentence A sentence that is neither a **necessary truth** nor a **necessary falsehood**; a sentence that in some case is true and in some other case, false.

contradiction (of FOL) A **sentence of FOL** that is false in every **interpretation**.

contradiction (of TFL) A sentence that is false on every **valuation**.

disjunct A sentence joined to another by a **disjunction**.

disjunction The connective \vee , used to represent words and phrases that function like the English word “or” in its inclusive sense; or a sentence formed by using this connective.

disjunctive normal form (DNF) a sentence which is a disjunction of conjunctions of atomic sentences or negated atomic sentences.

domain the collection of objects assumed for a symbolization in FOL, or that gives the range of the quantifiers in an **interpretation**.

empty predicate a **predicate** that applies to no object in the **domain**.

entailment (in TFL) Some sentences entail another if and only if no **valuation** makes all the former sentences true and the latter false, i.e. if the corresponding argument is valid..

existential quantifier the symbol \exists of FOL used to symbolize existence; $\exists x Fx$ is true iff at least one member of the domain is F .

expressive adequacy property of a collection of connectives which holds iff every possible truth table is the truth table of a sentence involving only those connectives.

formation tree A tree showing the structure of a sentence and its sub-sentences..

formula of FOL A string of symbols in FOL that can be built up according to the recursive rules given on p. 230.

free variable an occurrence of a variable in a **formula of FOL** which is not a **bound variable**.

interpretation a specification of a **domain** together with the objects the **names** pick out and which objects the **predicates** are true of.

invalid A property of arguments that holds when the conclusion is not a consequence of the premises; the opposite of **valid**.

joint possibility A property possessed by some sentences when they are all true in a single case.

logical consistency (in FOL) A property held by **sentence of FOLs** if and only if some **interpretation** makes all the sentences true.

logical equivalence (in FOL) A property held by pairs of **sentence of FOLs** if and only if the sentences have the same truth value in every **interpretation**..

logical equivalence (in TFL) A property held by pairs of sentences if and only if the sentences have the same truth value on every valuation.

logical truth A **sentence of FOL** that is true in every **interpretation**.
Note that the equivalent notion for TFL is ‘tautology’.

logical validity (in FOL) A property held by arguments if and only if no **interpretation** makes all premises true and the conclusion false.

logical validity (in TFL) A property held by arguments if and only if no **valuation** makes all premises true and the conclusion false.

main connective The last connective that you add when you assemble a sentence using the recursive definition..

metavariables A variable in the metalanguage that can represent any sentence in the object language..

name a symbol of FOL used to pick out an object of the **domain**.

necessary equivalence A property held by a pair of sentences that, in every case, are either both true or both false.

necessary falsehood A sentence that is false in every case.

necessary truth A sentence that is true in every case.

negation The symbol \neg , used to represent words and phrases that function like the English word “not”.

predicate a symbol of FOL used to symbolize a property or relation.

premise a sentence in an **argument** other than the **conclusion**.

premise indicator a word or phrase such as “because” used to indicate that what follows is the premise of an argument.

provable equivalence A property held by pairs of statements if and only if there is a derivation which takes you from each one to the other one.

provable inconsistency Sentences are provably inconsistent iff a contradiction can be derived from them.

scope The subsentence of which that instance of the connective is the main connective..

sentence of FOL a expression of FOL constructed according to the recursive rules..

sentence of TFL A string of symbols in TFL that can be built up according to the recursive rules given on p. 50.

sound A property of arguments that holds if the argument is valid and has all true premises.

soundness A property held by logical systems if and only if \vdash implies \models . 169

symbolization key A list that shows which English sentences are represented by which **atomic sentences** in TFL.

tautology A sentence that is true on every **valuation**.

theorem A sentence that can be proved without any premises.

truth table A table that gives all the possible **truth values** for a **sentence of TFL** or sentences in TFL, with a line for every possible **valuation** of all atomic sentences.

truth value One of the two logical values sentences can have: True and False.

truth-functional connective an operator that builds larger sentences out of smaller ones and fixes the **truth value** of the resulting sentence based only on the truth value of the component sentences.

universal quantifier the symbol \forall of FOL used to symbolize generality; $\forall x Fx$ is true iff every member of the domain is F .

valid A property of arguments where there conclusion is a consequence of the premises.

valuation An assignment of **truth values** to particular atomic **sentence of TFLs**.

variable a symbol of FOL used following quantifiers and as placeholders in atomic formulas; lowercase letters between s and z .

In the Introduction to his volume *Symbolic Logic*, Charles Lutwidge Dodson advised: “When you come to any passage you don’t understand, *read it again*: if you *still* don’t understand it, *read it again*: if you fail, even after *three* readings, very likely your brain is getting a little tired. In that case, put the book away, and take to other occupations, and next day, when you come to it fresh, you will very likely find that it is *quite* easy.”

The same might be said for this volume, although readers are forgiven if they take a break for snacks after *two* readings.