NATIONAL ALLOTMENTS JOURNAL.

ON

YOUR ALLOTMENT USE



ALLOTMENT FERTILIZER

ITS GRANULAR FORM MAKES IT EASY TO DISTRIBUTE AND PLEASANT TO HANDLE

I.C.I. ALLOTMENT FERTILIZER IS DISTRIBUTED BY PLANT PROTECTION YALDING

THE POTATO EELWORM

By R. P. THOMAS, DIP. HORT.

County Instructor, Monmountains is the most serious pest of the potato, and infestation is lapidly on the increase.

The sprouts from the seed tubers are late coming through the and the barrent spindly and stunted. the sprouts from the seed tubers are late coming through the foliage is doubt growth is very weak, spindly and stunted. The foliage is dull and pale and at an early date they curl and wilt.

The root svetaniand pale and at an early date they curl and wilt.

The root svetaniand pale and at an early date they curl and wilt. The root system is much reduced and the tubers produced are is much reduced and the tubers containing lew in number and small in size. Cysts (hard cases containing are easily, and small in size. Cysta easily seen on the rootlets.

Cysts are present on infected soil and the potato excretes a befance which sent on infected soil and the eggs which burst Systance which stimulates the hatching of the eggs which burst the cysts at the cysts and the potato excited a specific pen, the cysts at the cysts he larvae (grub stage) bore just below the surface layers of the balance of the plant. larvae (grub stage) bore just below the surface layers of appear, and in a few weeks the females become swollen and flaskhaped. Later the body portion breaks through the outside of the body portion breaks through the body portion breaks through the body portion breaks through the outside of the body portion breaks through the breaks through the breaks through the breaks through the breaks t comes. After fertilisation the female degenerates until it becomes a cyst filled with eggs. The cysts at first are creamywhite in colour, but later they become brown, and drop off into he soil, where they remain until the following spring.

Control Measures

Control Measures application of Calcium Cyanamide, applied at the rate of soil per 10 per per 10 perch plot. This should be thoroughly mixed with about three plot. Soil about three to four weeks prior to planting, and when the Mond is moderately dry. Nore. Eelworms attack the rootlets and not the tubers.

THE TENURE PROBLEM—MINISTER'S VIEW

TENURE PROBLEM—MINISTER'S VIA TONING ALEXANDER, M.P. for one of the Sheffield Divisions, a Movement, one-time First Movement, one-time Movement, o onlinent leader in the Co-operative Movement, one-time First the Admin the Co-operative Movement and of the Allotment Movements that of the Admiralty, and a good friend of the Allotments that the Admiralty, and a good friend of the Allotments that tote specially the about the insecurity of allotments that the specially the special spe Mote specially to Mr. Morrison, Minister of Agriculture, about

The following reply is, on the whole, a very reasonable statewish it to be. One can detect an undertone of sympathy decause and the feeling that if it rested with of the position as it is, although it is not so good as we wish it to be as it is, although it is not so good with Morrison along the cannot help feeling that if it rested with Morrison alone he would go a long way towards meeting our

have already had under consideration proposals submitted during 1937, designed to National Allotments Society, Ltd., during 1937, designed to