

Formalization of a BS in Measurement Theory

Lars Warren Ericson

Catskills Research Company

September 2026

CMU-CS-26-XXX

School of Computer Science

Carnegie Mellon University

Pittsburgh, PA 15213

Source repository: <https://github.com/catskillsresearch/fabbro>

Abstract

This report presents a reconstructed Carnegie Mellon undergraduate path through the Mathematical Sciences 21-xxx catalog, sequenced so that Dana Scott’s 1964 measurement theorems (1.1–4.1) and their extension to infinite Boolean algebras can be proved end to end. Each course contributes one problem that requires the full toolkit of that course rather than a single technique in isolation. Solutions are written in English and then formalized in Lean 4 / Mathlib. The accompanying library `BSinMeasurementTheory` is sorry-free under the pinned toolchain `leanprover/lean4:v4.34.0-rc1` and introduces no project axioms beyond Mathlib’s classical footprint. Large-language-model assistance was used in drafting solutions, but every accepted declaration is checked by Lean’s kernel. The complete source and a reproducible build pipeline are publicly available with this report.

Copyright © 2026 Lars Warren Ericson

Keywords: Lean 4; formal verification; measurement theory; undergraduate curriculum; Scott 1964; Boolean algebras; Stone spaces

Contents

1	Introduction	2
2	Syllabus	2
3	Phase 1: Language and linear structure	4
3.1	21-120/21-122 — Differential and Integral Calculus / Integration and Approximation	4
3.2	21-127 — Concepts of Mathematics	7
3.3	21-241 — Matrices and Linear Transformations	8
3.4	21-228 — Discrete Mathematics	11
4	Phase 2: Analysis, algebras, and formal proof	14
4.1	21-355 — Principles of Real Analysis I	15
4.2	21-373 — Algebraic Structures	18
4.3	21-321 — Interactive Theorem Proving	21
5	Phase 3: Duality and set theory	23
5.1	21-292 — Operations Research I	24
5.2	21-329 — Set Theory	27
6	Phase 4: Stone spaces and measures	30
6.1	21-651 — General Topology	30
6.2	21-720 — Measure and Integration	33
7	Phase 5: The missing extension	36
7.1	21-640 — Introduction to Functional Analysis	36
8	Phase 6: Assembly and consistency check	39
8.1	21-410/21-599 — Independent Study	39
8.2	21-322 — Topics in Formal Mathematics (opportunistic)	42
9	References	44
A	Lean module index	44

1 Introduction

Measurement theory asks when qualitative comparisons can be represented by real numbers. Scott’s 1964 paper organizes several such representation questions around a finite theory of homogeneous linear inequalities, with applications to intransitive indifference, ordered utility differences, and subjective probability. The infinite direction, which Scott announced but did not carry out, runs through Stone spaces, positive functionals, Hahn–Banach extension, and Riesz representation.

The pedagogical claim of this report is that a complete undergraduate mathematics major at Carnegie Mellon already contains the pieces of that argument, if the courses are taken in the right order and each is asked to do real work. The syllabus below is that ordering. It is framed as a Bachelor of Science in Measurement Theory: not a new degree program, but a reading of the existing catalog as a single connected proof.

The student of record for the problem set is Giovanni Fabbro. Each problem is built so that quoting a theorem from a later course is not available: the calculus problem uses the fundamental theorem, comparison, and Taylor remainders; the linear-algebra problem speaks the dual-space language of Scott’s Theorem 1.1; the discrete-mathematics problem is the combinatorial counting behind Theorems 1.2 and 3.2; and the independent-study problem assembles set theory, topology, measure, and functional analysis into the infinite extension of Theorem 4.1.

The formalizations live in the Lean library `BSinMeasurementTheory`. Each course snippet is a self-contained module. The root file `BSinMeasurementTheory.lean` imports all of them. The English arguments occupy the body of this report, in syllabus order; the Lean sources are collected in the appendix.

2 Syllabus

Phase	Course	Title	Role toward Scott (1964)
1	21-120 / 21-122	Differential and Integral Calculus / Integration and Approximation	FTC, comparison, integration by parts, Taylor remainder
1	21-127	Concepts of Mathematics	Equivalence relations, explicit bijections, induction
1	21-241	Matrices and Linear Transformations	Bases, dual bases, and the vector-space language of Theorem 1.1
1	21-228	Discrete Mathematics	Permutations, multisets, and the counting behind Theorems 1.2 and 3.2

Phase	Course	Title	Role toward Scott (1964)
2	21-355	Principles of Real Analysis I	Completeness, limsup/liminf, and compactness from axioms
2	21-373	Algebraic Structures	Atoms and characteristic functions of finite Boolean algebras
2	21-321	Interactive Theorem Proving	Formal statements of Theorems 1.1 and 1.2, and their equivalence
3	21-292	Operations Research I	Simplex and LP duality, deriving Kuhn–Tucker / Theorem 1.1 from the dual
3	21-329	Set Theory	Zorn’s lemma, ultrafilters, and cardinal arithmetic
4	21-651	General Topology	Stone spaces, compactness via Tychonoff, Stone representation
4	21-720	Measure and Integration	Riesz representation and the induced finitely additive measure on B
5	21-640	Introduction to Functional Analysis	Hahn–Banach extension, the step Scott cites and leaves unproved
6	21-410 / 21-599	Independent Study	Full reconstruction of the infinite-algebra extension of Theorem 4.1
6	21-322	Topics in Formal Mathematics	Proof that the finite atomic picture is what the infinite machinery reduces to

Every course problem is solved in two layers: an English mathematical argument, and a Lean 4 formalization using Mathlib, with numerical examples re-checked in Lean when they appear. Course 21-322

is opportunistic: it should be taken in a term that overlaps real analysis or order theory in Lean, but the standing problem is independent of that term's advertised topic.

The remainder of this report is the problem set itself, in syllabus order.

3 Phase 1: Language and linear structure

Calculus, proof language, dual bases, and combinatorial cancellation.

3.1 21-120/21-122 — Differential and Integral Calculus / Integration and Approximation

Let $f(x) = \int_0^x \frac{\sin(t^2)}{1+t^2} dt$.

- Show f is differentiable everywhere and find $f'(x)$.
- Determine whether $\lim_{x \rightarrow \infty} f(x)$ exists, and if so, bound it using comparison and integration by parts on the tail.
- Find the Taylor series of f about $x = 0$ through the x^9 term, and use it to estimate $f(0.5)$ to four decimal places with an explicit error bound.

Requires: FTC, limits/comparison tests, integration by parts, power series, Taylor remainder estimation.

Mathematical Solution

(a) Differentiability and Derivative

Let $g(t) = \frac{\sin(t^2)}{1+t^2}$. Since $t \mapsto \sin(t^2)$ and $t \mapsto 1+t^2$ are continuous on $\mathbb{R}$ and $1+t^2 \geq 1 > 0$ for all $t \in \mathbb{R}$, $g(t)$ is everywhere continuous. By the Fundamental Theorem of Calculus:

$$f(x) = \int_0^x g(t) dt$$

is differentiable for all $x \in \mathbb{R}$, and:

$$f'(x) = g(x) = \frac{\sin(x^2)}{1+x^2}.$$

(b) Existence of $\lim_{x \rightarrow \infty} f(x)$ and Tail Bound

- Existence:** For all $t \geq 0$, $|\sin(t^2)| \leq 1$, so

$$\left| \frac{\sin(t^2)}{1+t^2} \right| \leq \frac{1}{1+t^2}.$$

Since $\int_0^\infty \frac{dt}{1+t^2} = [\arctan(t)]_0^\infty = \frac{\pi}{2} < \infty$, the improper integral $\int_0^\infty \frac{\sin(t^2)}{1+t^2} dt$ is absolutely convergent. Hence, $\lim_{x \rightarrow \infty} f(x)$ exists.

2. **Tail Bound via Integration by Parts:** For any cutoff $M > 0$, decompose the limit into:

$$\lim_{x \rightarrow \infty} f(x) = \int_0^M \frac{\sin(t^2)}{1+t^2} dt + \int_M^\infty \frac{\sin(t^2)}{1+t^2} dt.$$

On the tail, rewrite $\sin(t^2) dt = \frac{1}{2t} d(-\cos(t^2))$. Integrating by parts with $u = \frac{1}{2t(1+t^2)}$ and $dv = 2t \sin(t^2) dt$:

$$\begin{aligned} \int_M^\infty \frac{\sin(t^2)}{1+t^2} dt &= \left[-\frac{\cos(t^2)}{2t(1+t^2)} \right]_M^\infty - \int_M^\infty (-\cos(t^2)) \left(-\frac{1+3t^2}{2t^2(1+t^2)^2} \right) dt \\ &= \frac{\cos(M^2)}{2M(1+M^2)} - \int_M^\infty \cos(t^2) \frac{1+3t^2}{2t^2(1+t^2)^2} dt. \end{aligned}$$

Since $|\cos(t^2)| \leq 1$ and $\frac{1+3t^2}{2t^2(1+t^2)^2} = -\frac{d}{dt} \left(\frac{1}{2t(1+t^2)} \right) > 0$:

$$\left| \int_M^\infty \frac{\sin(t^2)}{1+t^2} dt \right| \leq \frac{1}{2M(1+M^2)} + \int_M^\infty \left(-\frac{d}{dt} \frac{1}{2t(1+t^2)} \right) dt = \frac{1}{M(1+M^2)}.$$

Combining with the comparison bound on $[0, M]$:

$$\left| \lim_{x \rightarrow \infty} f(x) \right| \leq \arctan(M) + \frac{1}{M(1+M^2)}.$$

For example, setting $M = 1$ yields $|f(\infty)| \leq \frac{\pi}{4} + \frac{1}{2} \approx 1.2854$.

(c) Taylor Series and Numerical Estimation of $f(0.5)$

1. **Taylor Expansion:** Expanding the integrand around $t = 0$:

$$\sin(t^2) = t^2 - \frac{t^6}{6} + O(t^{10}), \quad \frac{1}{1+t^2} = 1 - t^2 + t^4 - t^6 + O(t^8)$$

Multiplying through terms up to degree 8:

$$g(t) = \left(t^2 - \frac{t^6}{6} \right) (1 - t^2 + t^4 - t^6) + O(t^{10}) = t^2 - t^4 + \frac{5}{6}t^6 - \frac{5}{6}t^8 + O(t^{10}).$$

Integrating term-by-term from 0 to x :

$$f(x) = \frac{x^3}{3} - \frac{x^5}{5} + \frac{5}{42}x^7 - \frac{5}{54}x^9 + O(x^{11}).$$

2. **Estimation at $x = 0.5$:**

$$\begin{aligned} \frac{(0.5)^3}{3} &= \frac{1}{24} \approx 0.041667 \\ -\frac{(0.5)^5}{5} &= -\frac{1}{160} = -0.006250 \end{aligned}$$

$$\frac{5(0.5)^7}{42} = \frac{5}{5376} \approx 0.000930$$

$$-\frac{5(0.5)^9}{54} = -\frac{5}{27648} \approx -0.000181$$

Summing these gives:

$$T_9(0.5) \approx 0.041667 - 0.006250 + 0.000930 - 0.000181 = 0.036166 \approx 0.0362.$$

3. **Explicit Error Bound:** On $[0, 0.5]$, the series for the integrand is alternating with terms decreasing monotonically in magnitude. By the alternating series estimation theorem, the truncation error in the integrand is bounded by the magnitude of the next term:

$$R_{\text{integrand}}(t) \leq \left(1 - \frac{1}{6} + \frac{1}{120}\right) t^{10} = \frac{101}{120} t^{10}.$$

Integrating yields:

$$|R_f(0.5)| \leq \int_0^{0.5} \frac{101}{120} t^{10} dt = \frac{101}{1320} (0.5)^{11} = \frac{101}{1320 \times 2048} \approx 3.73 \times 10^{-5} < 0.0001.$$

Thus, to four decimal places:

$$f(0.5) \approx 0.0362 \quad (\text{with error} < 4 \times 10^{-5}).$$

Lean 4 Formal Proof (Part a)

Below is the complete Lean 4 proof using Mathlib. It defines f , proves the continuity of the integrand, establishes differentiability everywhere, and computes the derivative using the Fundamental Theorem of Calculus with **no sorry**.

Lean 4 formalization: `appendix module BSinMeasurementTheory/Course21120FTC.lean`.

We can implement and verify the exact Taylor series calculations in Lean 4.

Lean's `norm_num` tactic can fully verify exact arithmetic over $\mathbb{Q}$ without any `sorry`, while `#eval` allows you to inspect the decimal (`Float`) representations directly in the Lean InfoView.

Here is the complete, self-contained Lean 4 code:

Lean 4 formalization: `appendix module BSinMeasurementTheory/Course21120Taylor.lean`.

Summary of what Lean verifies here:

1. Term-by-term calculation:

- $x^3/3 = 1/24 \approx 0.041667$
- $x^5/5 = 1/160 = 0.006250$
- $5x^7/42 = 5/5376 \approx 0.000930$
- $5x^9/54 = 5/27648 \approx 0.000181$

2. **Exact value:** $T_9(1/2) = \frac{34997}{967680} \approx 0.0361659$.

3. **Rounding:** Lean proves $0.03615 < T_9(1/2) < 0.03625$, confirming it rounds to 0.0362.

4. **Error bound:** Lean proves the remainder bound $\frac{101}{2703360} \approx 3.74 \times 10^{-5} < 10^{-4}$.

3.2 21-127 — Concepts of Mathematics

Define a relation $\sim$ on $\mathbb{Z} \times (\mathbb{Z} \setminus \{0\})$ by $(a, b) \sim (c, d) \iff ad = bc$.

- (a) Prove $\sim$ is an equivalence relation.
- (b) Prove the induced quotient set is in bijection with $\mathbb{Q}$, exhibiting the bijection explicitly and proving it well-defined, injective, and surjective.
- (c) Prove by strong induction that every nonempty subset of $\mathbb{Q}_{\geq 0}$ that is well-ordered under the usual order is either empty or has infinitely many elements — or find the flaw in that claim and prove the correct statement instead, using contradiction.

Requires: relations/equivalence classes, functions/bijections, induction, proof by contradiction, and — the real test — recognizing a false claim rather than forcing a proof of it.

Analysis and Mathematical Discussion

(a) Equivalence Relation

We define $S = \mathbb{Z} \times (\mathbb{Z} \setminus \{0\})$ and $(a, b) \sim (c, d) \iff ad = bc$: - **Reflexivity:** $ab = ba$ holds by commutativity of multiplication in $\mathbb{Z}$. - **Symmetry:** $ad = bc \implies cb = da$, so $(c, d) \sim (a, b)$. - **Transitivity:** If $ad = bc$ and $cf = de$, multiplying gives $(af)d = (ad)f = (bc)f = b(cf) = b(de) = (be)d$. Since $d \neq 0$ and $\mathbb{Z}$ is an integral domain (has no zero divisors), $(af - be)d = 0 \implies af = be$.

(b) Bijection with $\mathbb{Q}$

The canonical map is $F(a, b) = \frac{a}{b} \in \mathbb{Q}$: - **Well-defined:** If $ad = bc$, then $\frac{a}{b} = \frac{c}{d}$ in $\mathbb{Q}$ because clearing denominators yields $ad - bc = 0$. - **Injective:** If $\frac{a}{b} = \frac{c}{d}$ in $\mathbb{Q}$, then $ad = bc$ in $\mathbb{Z}$, meaning $[(a, b)] = [(c, d)]$. - **Surjective:** Every $q \in \mathbb{Q}$ has canonical numerator $q.\text{num} \in \mathbb{Z}$ and denominator $q.\text{den} \in \mathbb{N}^+$, with $q = \frac{q.\text{num}}{q.\text{den}}$. Since $q.\text{den} \neq 0$, $F([(q.\text{num}, q.\text{den})]) = q$.

(c) The Flaw in the Claim and the Correct Statement

The prompt poses the challenge: > “Prove by strong induction that every nonempty subset of $\mathbb{Q}_{\geq 0}$ that is well-ordered under the usual order is either empty or has infinitely many elements — or find the flaw in that claim and prove the correct statement instead, using contradiction.”

The Flaws in the Claim: 1. **Syntactic contradiction:** A subset assumed to be *nonempty* cannot be *empty*. 2. **Mathematical falsehood:** The claim asserts that any nonempty well-ordered subset of $\mathbb{Q}_{\geq 0}$ must be infinite. This is completely false. Any finite nonempty subset of $\mathbb{Q}_{\geq 0}$ (for instance, the singleton $\{0\}$, or $\{0, 1\}$, or $\{1/2\}$) is: - A subset of $\mathbb{Q}_{\geq 0}$, - Nonempty, - Well-ordered under the standard $\leq$ order (every nonempty subset has a least element), - Yet strictly **finite**, not infinite. 3. **Why strong induction fails here (the trap):** An attempted proof by induction would try to construct an infinite sequence $x_0 < x_1 < x_2 < \dots$ by repeatedly taking successors. This fails because a finite well-ordered subset has a **maximal element** with no strictly greater element in the set.

The Correct Theorems: 1. **Refutation by contradiction:** The claim is false. Assuming the claim holds for $S = \{0\}$ implies $\{0\} = \emptyset$ or $\{0\}$ is infinite, both of which immediately contradict the properties of singletons. 2. **Positive statement via induction:** Every finite nonempty subset of $\mathbb{Q}$ (and hence of $\mathbb{Q}_{\geq 0}$) has a least element, and is therefore well-ordered. We formalize this by induction on lists.

Complete Lean 4 Code

Lean 4 formalization: `appendix module BSinMeasurementTheory/Course21127.lean.`

3.3 21-241 — Matrices and Linear Transformations

Let V be a finite-dimensional real vector space, $C \subset V$ a convex cone with nonempty interior, and $v \notin C$.

- Construct a basis of V and its dual basis of V^* explicitly for the case $V = \mathbb{R}^3$, $C = \{x : x_1, x_2, x_3 \geq 0\}$.
- Prove that a linear functional $\varphi \in V^*$ separating v from C (i.e., $\varphi(v) < 0 \leq \varphi(c)$ for all $c \in C$) corresponds to a specific vector under the standard inner product, and compute it for a concrete $v \notin C$.
- Prove that the set of linear functionals nonnegative on all of C is itself a convex cone in V^* , and identify its dual basis representation.

Requires: basis/dual basis construction, linear transformations, the vector-space language Scott's Theorem 1.1 is stated in — not just matrix computation.

Prove it in English and then in Lean 4, using as much Mathlib as possible. No uses of sorry, admit or axiom are allowed. If you do any numerical calculations for examples to illustrate the proof, reproduce them after the proof in Lean.

Mathematical Exposition

Geometric Separation and Scott's Theorem 1.1

Let V be a finite-dimensional real vector space. A non-empty subset $C \subseteq V$ is a **convex cone** if: 1. For all $x, y \in C$, $x + y \in C$ (closed under addition). 2. For all $x \in C$ and all $\lambda \geq 0$, $\lambda x \in C$ (closed under non-negative scaling).

Scott's Theorem 1.1 (the geometric Hahn–Banach separation theorem for convex cones) establishes that if $C \subset V$ is a convex cone with non-empty interior $\text{int}(C) \neq \emptyset$, and $v \in V \setminus C$, then there exists a non-zero linear functional $\varphi \in V^*$ strictly separating v from $\text{int}(C)$ and non-strictly separating v from C :

$$\varphi(v) < 0 \leq \varphi(c), \quad \forall c \in C.$$

The separation level must be 0: if $\varphi(c) \geq \alpha$ for all $c \in C$, then for any $\lambda > 0$, $\varphi(\lambda c) = \lambda \varphi(c) \geq \alpha \implies \varphi(c) \geq \alpha/\lambda$. Sending $\lambda \rightarrow \infty$ forces $\varphi(c) \geq 0$. Furthermore, since $0 \in \overline{C}$, $\alpha \leq \varphi(0) = 0$.

Part (a): Basis and Dual Basis Construction

Let $V = \mathbb{R}^3$ and $C = \{x = (x_0, x_1, x_2) \in \mathbb{R}^3 \mid x_0 \geq 0, x_1 \geq 0, x_2 \geq 0\}$ be the non-negative orthant.

- 1. Primal Basis of V :** Define the standard basis vectors $e_0, e_1, e_2 \in V$ by:

$$e_i(j) = \delta_{ij} = \begin{cases} 1 & \text{if } i = j, \\ 0 & \text{if } i \neq j. \end{cases}$$

Explicitly:

$$e_0 = (1, 0, 0), \quad e_1 = (0, 1, 0), \quad e_2 = (0, 0, 1).$$

- **Spanning:** Any vector $x \in V$ can be uniquely expressed as:

$$x = \sum_{i=0}^2 x_i e_i.$$

- **Linear Independence:** If $\sum_{i=0}^2 c_i e_i = 0$, evaluating the j -th coordinate gives $c_j = 0$ for each $j \in \{0, 1, 2\}$.
- 2. **Dual Basis of V^* :** The algebraic dual space is $V^* = \text{Hom}_{\mathbb{R}}(V, \mathbb{R})$. The dual basis $\{e_0^*, e_1^*, e_2^*\} \subset V^*$ is defined by the coordinate projections:

$$e_i^*(x) := x_i = \text{proj}_i(x).$$

- **Duality relation:**

$$e_i^*(e_j) = e_j(i) = \delta_{ij}.$$

- **Expansion in the dual basis:** For any linear map $\varphi \in V^*$, by linearity:

$$\varphi(x) = \varphi\left(\sum_{i=0}^2 x_i e_i\right) = \sum_{i=0}^2 x_i \varphi(e_i) = \sum_{i=0}^2 \varphi(e_i) e_i^*(x),$$

$$\text{meaning } \varphi = \sum_{i=0}^2 \varphi(e_i) e_i^*.$$

Part (b): Inner Product Representation and Concrete Separation

1. **Riesz Isomorphism under the Standard Inner Product:** Equip V with the standard Euclidean inner product:

$$\langle u, x \rangle = \sum_{i=0}^2 u_i x_i.$$

Every vector $u \in V$ induces a linear functional $\varphi_u(x) = \langle u, x \rangle$. Conversely, every linear functional $\varphi \in V^*$ uniquely corresponds to the vector:

$$u_\varphi := \sum_{i=0}^2 \varphi(e_i) e_i = (\varphi(e_0), \varphi(e_1), \varphi(e_2)).$$

Indeed:

$$\langle u_\varphi, x \rangle = \sum_{i=0}^2 \varphi(e_i) x_i = \varphi\left(\sum_{i=0}^2 x_i e_i\right) = \varphi(x).$$

2. **Non-negativity on C :**

$$\forall c \in C, \langle u, c \rangle \geq 0 \iff \forall i \in \{0, 1, 2\}, u_i \geq 0.$$

- ($\implies$) Since $e_i \in C$, we have $u_i = \langle u, e_i \rangle \geq 0$.
- ($\impliedby$) If $u_i \geq 0$ for all i and $c \in C$ (so $c_i \geq 0$), then $\langle u, c \rangle = \sum_{i=0}^2 u_i c_i \geq 0$.

3. **Separating Vector for $v \notin C$:** If $v \notin C$, then by definition $\exists k \in \{0, 1, 2\}$ such that $v_k < 0$. Choosing $u = e_k$, we verify:

- For all $c \in C$: $\langle u, c \rangle = \langle e_k, c \rangle = c_k \geq 0$.

- At v : $\langle u, v \rangle = \langle e_k, v \rangle = v_k < 0$. Thus $\varphi(x) = \langle e_k, x \rangle = e_k^*(x)$ is the desired separating functional.

4. **Concrete Numerical Example:** Let $v = (-1, 1, 1)$.

- $v \notin C$ because $v_0 = -1 < 0$.
- Choose $u = e_0 = (1, 0, 0)$.
- Separation check:

$$\langle u, v \rangle = 1 \cdot (-1) + 0 \cdot 1 + 0 \cdot 1 = -1 < 0.$$

- Cone non-negativity: for any $c = (c_0, c_1, c_2) \in C$,

$$\langle u, c \rangle = c_0 \geq 0.$$

Part (c): The Dual Cone and its Dual Basis Representation

1. **The Dual Cone is a Convex Cone:** The dual cone of C is defined by:

$$C^* := \{\varphi \in V^* \mid \forall c \in C, \varphi(c) \geq 0\}.$$

- **Closed under non-negative scaling:** Let $\varphi \in C^*$ and $\lambda \geq 0$. For any $c \in C$:

$$(\lambda\varphi)(c) = \lambda \cdot \varphi(c) \geq 0 \quad (\text{product of non-negative reals}).$$

- **Closed under addition:** Let $\varphi, \psi \in C^*$. For any $c \in C$:

$$(\varphi + \psi)(c) = \varphi(c) + \psi(c) \geq 0 + 0 = 0.$$

- **Conic combinations:** If $\varphi, \psi \in C^*$ and $a, b \geq 0$, then $a\varphi + b\psi \in C^*$. Thus C^* is a convex cone in V^* .

2. **Dual Basis Representation:** Using the dual basis expansion from Part (a), any $\varphi \in V^*$ is $\varphi = \sum_{i=0}^2 a_i e_i^*$, where $a_i = \varphi(e_i)$.

- If $\varphi \in C^*$, then since $e_i \in C$, $a_i = \varphi(e_i) \geq 0$ for all i .
- Conversely, if $a_i \geq 0$ for all i , then for any $c \in C$:

$$\varphi(c) = \sum_{i=0}^2 a_i e_i^*(c) = \sum_{i=0}^2 a_i c_i \geq 0.$$

Therefore:

$$C^* = \left\{ \sum_{i=0}^2 a_i e_i^* \mid a_0, a_1, a_2 \geq 0 \right\}.$$

Under the dual basis $\{e_0^*, e_1^*, e_2^*\}$, the dual cone C^* is canonically identified with the non-negative orthant $\mathbb{R}_{\geq 0}^3$.

Fully Verified Lean 4 Formalization

Lean 4 formalization: appendix module `BSinMeasurementTheory/Course21241.lean`.

Verification of Numerical Calculations

Reproducing the concrete example computations performed in the proofs:

1. **Choice of point $v \notin C$:**

$$v = (-1, 1, 1) \implies v_0 = -1 < 0 \implies v \notin C.$$

2. **Separating vector $u \in V$:**

$$u = e_0 = (1, 0, 0).$$

3. **Inner product value at v :**

$$\langle u, v \rangle = \sum_{i=0}^2 u_i v_i = (1)(-1) + (0)(1) + (0)(1) = -1 < 0.$$

4. **Inner product on arbitrary $c \in C$:**

$$\langle u, c \rangle = \sum_{i=0}^2 u_i c_i = (1)c_0 + (0)c_1 + (0)c_2 = c_0 \geq 0 \quad (\text{since } c \in C \implies c_i \geq 0 \forall i).$$

5. **Linear functional separation:**

$$\varphi_u(v) = -1 < 0 \leq c_0 = \varphi_u(c) \quad \text{for all } c \in C.$$

Here is the complete, self-contained Lean 4 formalization reproducing each of the 5 concrete calculations step by step, using only Mathlib tactics without any `sorry` or `admit`.

Lean 4 formalization: appendix module `BSinMeasurementTheory/Course21241Numerical.lean`.

3.4 21-228 — Discrete Mathematics

Let $x_1, \dots, x_n \in \mathbb{Q}^k$ be a finite sequence of rational vectors.

- (a) Prove that if every permutation π of $\{1, \dots, n\}$ produces a sequence $x_{\pi(1)}, \dots, x_{\pi(n)}$ satisfying some symmetric linear condition L , then L depends only on the multiset $\{x_1, \dots, x_n\}$, not the order.
- (b) Given two sequences $x_1, \dots, x_n$ and $y_1, \dots, y_n$, count (with an exact combinatorial formula, not an estimate) the number of pairs of permutations (π, σ) for which $\sum_i x_{\pi(i)} = \sum_i y_{\sigma(i)}$ termwise, under stated genericity assumptions on the vectors.
- (c) Prove your formula reduces correctly to the $n = 2$ and $n = 3$ cases by direct enumeration.

Requires: permutations/combinations, combinatorial proof technique, the direct toolkit for Scott's Theorems 1.2 and 3.2.

Prove it in English and then in Lean 4, using as much Mathlib as possible. No uses of sorry, admit or axiom are allowed. If you do any numerical calculations for examples to illustrate the proof, reproduce them after the proof in Lean.

Mathematical Analysis and English Proof

Part (a)

Proposition: Let $x_1, \dots, x_n \in \mathbb{Q}^k$ be a sequence of vectors. If every permutation $\pi \in S_n$ produces a sequence $(x_{\pi(1)}, \dots, x_{\pi(n)})$ satisfying a symmetric linear condition L , then L depends only on the underlying multiset $\{x_1, \dots, x_n\}$, not on the order.

Proof:

1. **Symmetry and Multiset Equivalence:** Two sequences $u = (u_1, \dots, u_n)$ and $v = (v_1, \dots, v_n)$ in $(\mathbb{Q}^k)^n$ represent the same multiset if and only if they lie in the same orbit under the natural action of the symmetric group S_n given by

$$\pi \cdot (u_1, \dots, u_n) = (u_{\pi(1)}, \dots, u_{\pi(n)}).$$

A condition L is *symmetric* if for every permutation $\pi \in S_n$ and every sequence $u \in (\mathbb{Q}^k)^n$,

$$L(u_{\pi(1)}, \dots, u_{\pi(n)}) \iff L(u_1, \dots, u_n).$$

Consequently, the truth value (or evaluation) of L is constant on every orbit of the S_n -action. Since the quotient space $(\mathbb{Q}^k)^n / S_n$ is canonically bijective to the set of multisets of size n with elements in $\mathbb{Q}^k$, L factors through this quotient. Therefore, L depends only on the multiset $\{x_1, \dots, x_n\}$.

2. **Structure of Symmetric Linear Forms:** If $L : (\mathbb{Q}^k)^n \rightarrow \mathbb{Q}^m$ is a linear mapping, it can be written in block form as

$$L(u_1, \dots, u_n) = \sum_{i=1}^n A_i u_i$$

for linear maps $A_i : \mathbb{Q}^k \rightarrow \mathbb{Q}^m$. Symmetry under the transposition $(i \ j) \in S_n$ implies that for all vectors $v \in \mathbb{Q}^k$, evaluating L on the tuple $(0, \dots, v, \dots, 0)$ (with v in position i) gives the same result as evaluating on the tuple with v in position j . Thus:

$$A_i v = A_j v \quad \forall v \in \mathbb{Q}^k \implies A_1 = A_2 = \dots = A_n =: A.$$

Hence, every symmetric linear map has the form:

$$L(u_1, \dots, u_n) = A \left(\sum_{i=1}^n u_i \right).$$

Because vector addition in $\mathbb{Q}^k$ is commutative and associative, the sum $\sum_{i=1}^n x_i$ is an invariant of the multiset $\{x_1, \dots, x_n\}$, depending in no way on the order of the terms. ■

Part (b)

Problem: Given sequences $x = (x_1, \dots, x_n)$ and $y = (y_1, \dots, y_n)$ in $\mathbb{Q}^k$, find an exact combinatorial formula for the number of pairs of permutations $(\pi, \sigma) \in S_n \times S_n$ such that $\sum_i x_{\pi(i)} = \sum_i y_{\sigma(i)}$ *termwise*.

1. Genericity Assumptions:

- **Assumption 1 (Pairwise Distinctness):** The vectors within the sequence x are pairwise distinct: $x_i \neq x_j$ for all $i \neq j$ (i.e., the mapping $i \mapsto x_i$ is injective).
- **Assumption 2 (Multiset Compatibility):** The multiset of vectors in y equals the multiset of vectors in x . That is, there exists a permutation $\tau \in S_n$ such that $y_i = x_{\tau(i)}$ for all $i \in \{1, \dots, n\}$. (If the multisets do not match, no matching exists, yielding 0 pairs.)

2. Derivation of the Exact Formula:

- The termwise equality of the sums means that corresponding terms match:

$$x_{\pi(i)} = y_{\sigma(i)} \quad \forall i \in \{1, \dots, n\}.$$

- Substituting $y_j = x_{\tau(j)}$ gives:

$$x_{\pi(i)} = x_{\tau(\sigma(i))} \quad \forall i \in \{1, \dots, n\}.$$

- By **Assumption 1** (injectivity of x), this holds if and only if the indices themselves agree:

$$\pi(i) = \tau(\sigma(i)) \quad \forall i \in \{1, \dots, n\}.$$

- In the symmetric group S_n , this is the identity $\pi = \tau \circ \sigma$, which is uniquely invertible:

$$\sigma = \tau^{-1} \circ \pi.$$

- Hence, for **any** choice of $\pi \in S_n$, there exists **exactly one** permutation $\sigma \in S_n$ satisfying the termwise condition, namely $\sigma = \tau^{-1} \circ \pi$.
- The assignment $\pi \mapsto (\pi, \tau^{-1} \circ \pi)$ is a bijection between S_n and the set of valid pairs.
- Since $|S_n| = n!$, the exact count of pairs (π, σ) is:

$$\boxed{N = n!}$$

(Remark: If the distinctness assumption is relaxed so that the vectors have distinct values with multiplicities $m_1, \dots, m_r$ where $\sum m_j = n$, the formula generalizes to $n! \prod_{j=1}^r m_j!$. Under the generic assumption $m_j = 1$, this evaluates to $n!$.)

Part (c)

Verification by Direct Enumeration:

Case $n = 2$:

- The symmetric group S_2 has $2! = 2$ elements: $e = \text{id} = (1)(2)$ and $\tau_{12} = (1\ 2)$.
- The total number of pairs in $S_2 \times S_2$ is $2 \times 2 = 4$.
- Without loss of generality, let $y = x$ (so $\tau = \text{id}$). The 4 pairs are:

1. $(\text{id}, \text{id}): x_1 = x_1 \text{ and } x_2 = x_2 \implies \text{Valid}$.
2. $(\text{id}, (1\ 2)): x_1 = x_2 \text{ and } x_2 = x_1 \implies \text{Invalid (since } x_1 \neq x_2)$.
3. $((1\ 2), \text{id}): x_2 = x_1 \text{ and } x_1 = x_2 \implies \text{Invalid (since } x_1 \neq x_2)$.
4. $((1\ 2), (1\ 2)): x_2 = x_2 \text{ and } x_1 = x_1 \implies \text{Valid}$.

- Number of valid pairs: 2, which matches $2! = 2$.

Case $n = 3$:

- S_3 contains $3! = 6$ permutations:

$$\{\text{id}, (1\ 2), (1\ 3), (2\ 3), (1\ 2\ 3), (1\ 3\ 2)\}.$$

- The product $S_3 \times S_3$ has $6 \times 6 = 36$ pairs (π, σ) .
- For each permutation $\pi \in S_3$, the condition $x_{\pi(i)} = x_{\sigma(i)}$ for all $i \in \{1, 2, 3\}$ forces $\sigma(i) = \pi(i)$ for all i (since x_1, x_2, x_3 are pairwise distinct).
- Thus $\sigma = \pi$ is the unique solution for each π .
- The valid pairs are precisely the 6 diagonal pairs:

$$\{(\pi, \pi) \mid \pi \in S_3\}.$$

- All remaining $36 - 6 = 30$ pairs fail termwise equality.
- Number of valid pairs: 6, which matches $3! = 6$.

Lean 4 Formalization

The following complete Lean 4 script formalizes the results with no uses of `sorry`, `admit`, or `axiom`.
Lean 4 formalization: `appendix module BSinMeasurementTheory/Course21228.lean`.

4 Phase 2: Analysis, algebras, and formal proof

Completeness, finite Boolean algebras, and Scott's Theorems 1.1–1.3 in Lean.

4.1 21-355 — Principles of Real Analysis I

Let (x_n) be a bounded sequence in $\mathbb{R}$.

- (a) Prove the Bolzano–Weierstrass theorem for this sequence from the completeness axiom (least upper bound property) — do not cite it, derive it.
- (b) Prove that $\limsup x_n$ and $\liminf x_n$ exist and characterize each as the limit of a specific monotone sequence you construct.
- (c) Give a sequence where $\limsup x_n \neq \liminf x_n$, and prove using an ε -argument (not just intuition) that (x_n) has no limit.
- (d) Prove that $[a, b] \subset \mathbb{R}$ is compact directly from the open cover definition — no shortcuts via sequential compactness unless you first prove the two notions equivalent.

Requires: completeness, sequences, compactness — proved from axioms, not quoted.

Prove it in English and then in Lean 4, using as much Mathlib as possible. No uses of sorry, admit or axiom are allowed. If you do any numerical calculations for examples to illustrate the proof, reproduce them after the proof in Lean.

4.1.1 Part 1: English Proofs

(a) Bolzano–Weierstrass Theorem from the Completeness Axiom

Theorem: Every bounded sequence $(x_n)_{n \in \mathbb{N}}$ in $\mathbb{R}$ has a convergent subsequence.

Axiom Used (Completeness / Least Upper Bound Property)

Every non-empty subset of $\mathbb{R}$ that is bounded above has a least upper bound (supremum) in $\mathbb{R}$. Equivalently, every non-empty subset bounded below has an infimum.

Lemma 1 (Monotone Convergence Theorem)

Every bounded, monotonic sequence in $\mathbb{R}$ converges.

Proof:

Let $(a_k)_{k \in \mathbb{N}}$ be non-decreasing and bounded above by M . The range $A = \{a_k : k \in \mathbb{N}\}$ is non-empty and bounded above, so by the completeness axiom, $L = \sup A$ exists.

For any $\varepsilon > 0$, $L - \varepsilon$ is not an upper bound of A , so there exists $K \in \mathbb{N}$ such that $a_K > L - \varepsilon$. Since (a_k) is non-decreasing, for all $k \geq K$,

$$L - \varepsilon < a_K \leq a_k \leq L < L + \varepsilon \implies |a_k - L| < \varepsilon.$$

Thus, $\lim_{k \rightarrow \infty} a_k = L$. The case of a non-increasing sequence bounded below is symmetric by considering $-a_k$. ■

Lemma 2 (Monotone Subsequence / Peak Points)

Every real sequence $(x_n)_{n \in \mathbb{N}}$ has a monotonic subsequence.

Proof:

We say an index $m \in \mathbb{N}$ is a *peak point* of the sequence (x_n) if

$$x_n \leq x_m \quad \text{for all } n > m.$$

We divide into two cases:

1. Case 1: There are infinitely many peak points.

Enumerate the peak points in increasing order: $m_0 < m_1 < m_2 < \dots$

Since m_k is a peak point and $m_{k+1} > m_k$, we have $x_{m_{k+1}} \leq x_{m_k}$.

Thus, $(x_{m_k})_{k \in \mathbb{N}}$ is a non-increasing subsequence.

2. Case 2: There are only finitely many peak points.

Let N be an integer strictly greater than all peak points (or $N = 0$ if there are none).

Choose $n_0 = N$. Since $n_0 \geq N$, n_0 is not a peak point, which means there exists $n_1 > n_0$ with $x_{n_1} > x_{n_0}$.

Similarly, $n_1 > N$ is not a peak point, so there exists $n_2 > n_1$ with $x_{n_2} > x_{n_1}$.

Continuing inductively, we obtain indices $n_0 < n_1 < n_2 < \dots$ such that $x_{n_k} < x_{n_{k+1}}$ for all $k \in \mathbb{N}$.

Thus, $(x_{n_k})_{k \in \mathbb{N}}$ is a strictly increasing subsequence. ■

Derivation of Bolzano–Weierstrass

Let (x_n) be bounded. By Lemma 2, (x_n) has a monotonic subsequence (x_{n_k}) . Because (x_n) is bounded, (x_{n_k}) is also bounded. By Lemma 1 (which follows directly from the completeness axiom), (x_{n_k}) converges. ■

(b) Existence and Characterization of $\limsup x_n$ and $\liminf x_n$

Let $(x_n)_{n \in \mathbb{N}}$ be a bounded sequence: there exist $m, M \in \mathbb{R}$ such that $m \leq x_n \leq M$ for all $n \in \mathbb{N}$.

For each $k \in \mathbb{N}$, consider the tail set $S_k = \{x_n : n \geq k\}$. 1. Each $S_k \neq \emptyset$ (since $x_k \in S_k$) and $S_k \subseteq [m, M]$. 2. By the completeness axiom: - $y_k = \sup S_k = \sup_{n \geq k} x_n$ exists, and $m \leq y_k \leq M$. - $z_k = \inf S_k = \inf_{n \geq k} x_n$ exists, and $m \leq z_k \leq M$.

Monotonicity of the Tail Extremities

Since $S_{k+1} = \{x_n : n \geq k+1\} \subseteq \{x_n : n \geq k\} = S_k$: - Any upper bound of S_k is an upper bound of S_{k+1} , so $\sup S_{k+1} \leq \sup S_k$, meaning $y_{k+1} \leq y_k$. Thus, (y_k) is a non-increasing sequence bounded below by m . - Any lower bound of S_k is a lower bound of S_{k+1} , so $\inf S_{k+1} \geq \inf S_k$, meaning $z_{k+1} \geq z_k$. Thus, (z_k) is a non-decreasing sequence bounded above by M .

Existence of Limits

By Lemma 1 (Monotone Convergence Theorem): - The non-increasing bounded sequence (y_k) converges to its infimum:

$$\lim_{k \rightarrow \infty} y_k = \inf_{k \in \mathbb{N}} y_k = \inf_{k \in \mathbb{N}} \left(\sup_{n \geq k} x_n \right) =: \limsup_{n \rightarrow \infty} x_n.$$

- The non-decreasing bounded sequence (z_k) converges to its supremum:

$$\lim_{k \rightarrow \infty} z_k = \sup_{k \in \mathbb{N}} z_k = \sup_{k \in \mathbb{N}} \left(\inf_{n \geq k} x_n \right) =: \liminf_{n \rightarrow \infty} x_n.$$

Both limits exist as finite real numbers. ■

(c) Example where $\limsup x_n \neq \liminf x_n$ and ε -Proof of Divergence

Consider the sequence $x_n = (-1)^n$ for $n \in \mathbb{N}$.

Computation of $\limsup$ and $\liminf$

For every $k \in \mathbb{N}$, the tail $S_k = \{(-1)^n : n \geq k\}$ contains both $(-1)^{2k} = 1$ and $(-1)^{2k+1} = -1$. Hence:

$$S_k = \{-1, 1\}.$$

Therefore: - $y_k = \sup S_k = 1$ for all $k \implies \limsup_{n \rightarrow \infty} x_n = \lim_{k \rightarrow \infty} 1 = 1$. - $z_k = \inf S_k = -1$ for all $k \implies \liminf_{n \rightarrow \infty} x_n = \lim_{k \rightarrow \infty} (-1) = -1$.

Since $1 \neq -1$, $\limsup x_n \neq \liminf x_n$.

Direct ε -Proof of Divergence

We prove by contradiction that (x_n) does not converge to any $L \in \mathbb{R}$.

Suppose there exists $L \in \mathbb{R}$ such that $\lim_{n \rightarrow \infty} x_n = L$.

Set $\varepsilon = 1 > 0$.

By definition of convergence, there exists $N \in \mathbb{N}$ such that for all $n \geq N$:

$$|x_n - L| < 1.$$

Choose the even index $n = 2N \geq N$:

$$|x_{2N} - L| = |1 - L| < 1.$$

Choose the odd index $n = 2N + 1 \geq N$:

$$|x_{2N+1} - L| = |-1 - L| = |L + 1| < 1.$$

Using the triangle inequality:

$$2 = |1 - (-1)| = |(1 - L) + (L + 1)| \leq |1 - L| + |L + 1| < 1 + 1 = 2,$$

which yields $2 < 2$, a contradiction.

Hence, (x_n) has no limit in $\mathbb{R}$. ■

(d) Heine–Borel: $[a, b]$ is Compact Directly from Open Covers

Theorem: Every open cover of $[a, b]$ has a finite subcover.

Proof:

If $a > b$, $[a, b] = \emptyset$, which is covered by the empty collection. Assume $a \leq b$.

Let $\mathcal{U} = \{U_\alpha\}_{\alpha \in \Lambda}$ be an open cover of $[a, b]$:

$$[a, b] \subseteq \bigcup_{\alpha \in \Lambda} U_\alpha, \quad \text{with each } U_\alpha \text{ open in } \mathbb{R}.$$

Define the set:

$$S = \{x \in [a, b] : [a, x] \text{ is covered by finitely many sets from } \mathcal{U}\}.$$

1. S is non-empty:

Since $a \in [a, b]$, there is some $\alpha_0 \in \Lambda$ such that $a \in U_{\alpha_0}$. Then $[a, a] = \{a\} \subseteq U_{\alpha_0}$, so $[a, a]$ is covered by the single set U_{α_0} . Thus $a \in S$.

2. S is bounded above:

By definition, $S \subseteq [a, b]$, so every $x \in S$ satisfies $x \leq b$. Thus b is an upper bound for S .

3. **Existence of the supremum:**

By the completeness axiom, $c = \sup S$ exists in $\mathbb{R}$, and $a \leq c \leq b$.

4. **Claim:** $c \in S$:

Since $c \in [a, b]$, there exists $\alpha_c \in \Lambda$ such that $c \in U_{\alpha_c}$.

Since U_{α_c} is open, there exists $\varepsilon > 0$ such that $(c - \varepsilon, c + \varepsilon) \subseteq U_{\alpha_c}$.

Because $c = \sup S$, $c - \varepsilon$ is not an upper bound of S , so there exists $x_0 \in S$ with $c - \varepsilon < x_0 \leq c$.

Since $x_0 \in S$, $[a, x_0]$ is covered by a finite subcollection $\{U_1, \dots, U_m\} \subseteq \mathcal{U}$.

Then:

$$[a, c] \subseteq [a, x_0] \cup (c - \varepsilon, c + \varepsilon) \subseteq U_1 \cup \dots \cup U_m \cup U_{\alpha_c}.$$

Hence $[a, c]$ is covered by $\{U_1, \dots, U_m, U_{\alpha_c}\}$, a finite subcollection of $\mathcal{U}$. Therefore, $c \in S$.

5. **Claim:** $c = b$:

Suppose for contradiction that $c < b$.

Using the same open set U_{α_c} containing c , choose $\delta > 0$ such that $[c, c + \delta) \subseteq U_{\alpha_c} \cap [a, b]$.

Let $x_1 = \min(b, c + \delta/2) > c$.

Then $[a, x_1] = [a, c] \cup [c, x_1]$. Since $[a, c]$ is covered by $\{U_1, \dots, U_m, U_{\alpha_c}\}$ and $[c, x_1] \subseteq U_{\alpha_c}$,

$$[a, x_1] \subseteq U_1 \cup \dots \cup U_m \cup U_{\alpha_c}.$$

Thus $x_1 \in S$, which contradicts c being an upper bound of S since $x_1 > c$.

Therefore, $c = b$.

Since $c \in S$ and $c = b$, we have $b \in S$. By definition of S , $[a, b]$ is covered by a finite subcollection of $\mathcal{U}$. ■

4.1.2 Part 2: Formal Proofs in Lean 4

Below is the complete Lean 4 code formalizing each part without sorry, admit, or non-standard axioms.

Lean 4 formalization: `appendix module BSinMeasurementTheory/Course21355.lean`.

4.2 21-373 — Algebraic Structures

Let B be a finite Boolean algebra with atoms $a_1, \dots, a_n$.

- (a) Prove every element of B is a unique join of a subset of the atoms.
- (b) Prove B is isomorphic (as a Boolean algebra) to the power set of $\{a_1, \dots, a_n\}$, and that this identifies B with the set of characteristic functions on $\{1, \dots, n\}$.
- (c) Separately: let G be a finite group acting on B by automorphisms. Prove the fixed-point set B^G is a Boolean subalgebra, and determine its atoms in terms of the orbits of $a_1, \dots, a_n$.

Requires: groups (part c), and — the self-study piece the syllabus flags as not separately covered — atoms/characteristic-function identification for finite Boolean algebras, which Theorem 4.1 needs directly.

Prove it in English and then in Lean 4, using as much Mathlib as possible. No uses of `sorry`, `admit` or `axiom` are allowed. If you do any numerical calculations for examples to illustrate the proof, reproduce them after the proof in Lean.

Here is the complete proof in English, followed by the formalization in Lean 4 with no uses of `sorry`, `admit`, or `axiom`, and a concrete computational example.

Mathematical Proof in English

Let $(B, \vee, \wedge, \neg, 0, 1)$ be a finite Boolean algebra. Recall that an element $a \in B$ is an **atom** if $a > 0$ and there is no element $x \in B$ with $0 < x < a$ (equivalently, for all $x \leq a$, either $x = 0$ or $x = a$). Let $A = \{a_1, \dots, a_n\}$ be the set of all atoms of B .

Part (a): Every element is a unique join of a subset of the atoms

Lemma 1 (Disjointness of distinct atoms): If $a, b \in A$ with $a \neq b$, then $a \wedge b = 0$. *Proof:* Since $a \wedge b \leq a$ and a is an atom, $a \wedge b \in \{0, a\}$. If $a \wedge b = a$, then $a \leq b$. Since b is an atom and $a \neq 0$, this forces $a = b$, contradicting $a \neq b$. Hence $a \wedge b = 0$. ■

Lemma 2 (Atom under a join): For any atom $a \in A$ and any subset $S \subseteq A$,

$$a \leq \bigvee_{s \in S} s \iff a \in S.$$

Proof: - ($\Leftarrow$) If $a \in S$, clearly $a \leq \bigvee_{s \in S} s$. - ($\Rightarrow$) By distributivity of $\wedge$ over $\vee$:

$$a = a \wedge \left(\bigvee_{s \in S} s \right) = \bigvee_{s \in S} (a \wedge s).$$

By Lemma 1, $a \wedge s = 0$ for all $s \in S \setminus \{a\}$. If $a \notin S$, then $a \wedge s = 0$ for every $s \in S$, which gives $a = \bigvee_{s \in S} 0 = 0$, contradicting that a is an atom ($a > 0$). Thus $a \in S$. ■

Existence: For any $x \in B$, define $S_x = \{a \in A \mid a \leq x\}$. Clearly $\bigvee S_x \leq x$. Suppose $\bigvee S_x < x$. In a Boolean algebra, consider the complement $y = x \wedge \neg(\bigvee S_x)$. Since $\bigvee S_x < x$, we have $y > 0$. Because B is finite, the non-empty set $\{z \in B \mid 0 < z \leq y\}$ has a minimal element a^* . Minimality implies a^* is an atom of B , so $a^* \in A$. Since $a^* \leq y \leq x$, we have $a^* \in S_x$. But also $a^* \leq y \leq \neg(\bigvee S_x)$, which implies $a^* \wedge \bigvee S_x = 0$. Since $a^* \in S_x$, $a^* \leq \bigvee S_x$, giving $a^* = a^* \wedge \bigvee S_x = 0$, a contradiction since $a^* > 0$. Hence $x = \bigvee S_x$.

Uniqueness: Suppose $x = \bigvee S = \bigvee T$ for two subsets $S, T \subseteq A$. By Lemma 2, for every atom $a \in A$:

$$a \in S \iff a \leq \bigvee S \iff a \leq \bigvee T \iff a \in T.$$

Hence $S = T$. ■

Part (b): Isomorphism with $\mathcal{P}(\{a_1, \dots, a_n\})$ and identification with $\{0, 1\}^n$

Define $\phi : B \rightarrow \mathcal{P}(A)$ by $\phi(x) = \{a \in A \mid a \leq x\} = S_x$, with inverse $\psi(S) = \bigvee S$. By Part (a), ϕ and ψ are mutually inverse bijections.

We verify that ϕ preserves the Boolean algebra operations: 1. **Lattice operations:** - $\phi(x \wedge y) = \{a \in A \mid a \leq x \wedge y\} = \{a \in A \mid a \leq x \text{ and } a \leq y\} = \phi(x) \cap \phi(y)$. - $\phi(x \vee y) = \{a \in A \mid a \leq x \vee y\}$. By distributivity, for any atom a , $a = (a \wedge x) \vee (a \wedge y)$. Since $a \wedge x \in \{0, a\}$ and $a \wedge y \in \{0, a\}$ and $a \neq 0$, at least one must equal a . Thus $a \leq x \vee y \iff a \leq x \text{ or } a \leq y$, so $\phi(x \vee y) = \phi(x) \cup \phi(y)$. 2. **Bounds and complement:** - $\phi(0) = \{a \in A \mid a \leq 0\} = \emptyset$. - $\phi(1) = \{a \in A \mid a \leq 1\} = A$. - For any atom a , $a \leq \neg x \iff a \wedge x = 0 \iff a \not\leq x$. Thus $\phi(\neg x) = A \setminus \phi(x) = (\phi(x))^c$.

Hence ϕ is an isomorphism of Boolean algebras: $B \cong \mathcal{P}(A)$.

Identification with characteristic functions: The power set $\mathcal{P}(A)$ is naturally isomorphic to $\{0, 1\}^A$ via the characteristic function map $S \mapsto \chi_S$, where $\chi_S(a) = 1$ if $a \in S$ and 0 otherwise. Using the indexing $A = \{a_1, \dots, a_n\}$, this corresponds to:

$$\chi : B \xrightarrow{\sim} \{0, 1\}^n, \quad x \mapsto (c_1, \dots, c_n) \quad \text{where } c_i = \begin{cases} 1 & \text{if } a_i \leq x \\ 0 & \text{if } a_i \not\leq x. \end{cases}$$

The Boolean operations on B map to the pointwise Boolean operations on $\{0, 1\}^n$:

$$(c \vee d)_i = \max(c_i, d_i), \quad (c \wedge d)_i = \min(c_i, d_i), \quad (\neg c)_i = 1 - c_i.$$

Part (c): Action of a finite group G by automorphisms

Let G act on B such that each $g \in G$ is a Boolean algebra automorphism. The fixed-point set is $B^G = \{x \in B \mid \forall g \in G, g \cdot x = x\}$.

1. **B^G is a Boolean subalgebra:** For any $x, y \in B^G$ and $g \in G$:

- $g \cdot 0 = 0 \implies 0 \in B^G$.
- $g \cdot 1 = 1 \implies 1 \in B^G$.
- $g \cdot (x \vee y) = (g \cdot x) \vee (g \cdot y) = x \vee y \implies x \vee y \in B^G$.
- $g \cdot (x \wedge y) = (g \cdot x) \wedge (g \cdot y) = x \wedge y \implies x \wedge y \in B^G$.
- $g \cdot (\neg x) = \neg(g \cdot x) = \neg x \implies \neg x \in B^G$. Hence B^G is closed under all Boolean operations and contains the bounds, so it is a Boolean subalgebra of B .

2. **Action on atoms and orbits:** Since each $g \in G$ is an automorphism, it preserves the order: $x \leq y \iff g \cdot x \leq g \cdot y$. Thus a is an atom if and only if $g \cdot a$ is an atom. Hence G acts as a permutation group on the set of atoms $A = \{a_1, \dots, a_n\}$. This action partitions A into disjoint orbits $\mathcal{O}_1, \dots, \mathcal{O}_k$.

3. **Atoms of B^G :** For each orbit $\mathcal{O}_j$, define $b_j = \bigvee_{a \in \mathcal{O}_j} a$.

- $b_j \in B^G$: For any $g \in G$, $g \cdot b_j = \bigvee_{a \in \mathcal{O}_j} (g \cdot a) = \bigvee_{a' \in \mathcal{O}_j} a' = b_j$, since g permutes $\mathcal{O}_j$.
- $b_j > 0$: Since orbits are non-empty and atoms are non-zero, $b_j > 0$.

- **b_j is an atom of B^G :** Suppose $x \in B^G$ satisfies $x \leq b_j$. Consider the set $S_x = \{a \in A \mid a \leq x\}$. Since $x \leq b_j$, $S_x \subseteq \mathcal{O}_j$. Moreover, since $x \in B^G$, for any $g \in G$ and $a \in A$:

$$a \leq x \iff g \cdot a \leq g \cdot x = x.$$

Hence S_x is a G -invariant subset of $\mathcal{O}_j$. Since $\mathcal{O}_j$ is a single transitive orbit, its only invariant subsets are $\emptyset$ and $\mathcal{O}_j$:

- If $S_x = \emptyset$, then $x = \bigvee \emptyset = 0$.
- If $S_x = \mathcal{O}_j$, then $x = \bigvee \mathcal{O}_j = b_j$. Thus, for any $x \in B^G$ with $x \leq b_j$, either $x = 0$ or $x = b_j$.
- **Exhaustion:** Any $x \in B^G$ has an invariant support S_x , which must be a union of orbits: $S_x = \bigcup_{j \in J} \mathcal{O}_j$. Thus $x = \bigvee_{j \in J} b_j$. Therefore, the atoms of B^G are precisely the orbit joins $b_1, \dots, b_k$. ■

Formalization in Lean 4

Lean 4 formalization: appendix module `BSinMeasurementTheory/Course21373.lean`.

Numerical / Computational Example

To illustrate the theorem: - Let $B = \mathcal{P}(\{0, 1, 2\})$ with $n = 3$ atoms: $a_0 = \{0\}$, $a_1 = \{1\}$, $a_2 = \{2\}$. - Let $G = \mathbb{Z}/2\mathbb{Z} = \{e, \sigma\}$ act by the transposition $(0 \ 1)$ exchanging 0 and 1, while fixing 2. - The action partitions the atoms into two orbits:

$$\mathcal{O}_1 = \{a_0, a_1\} = \{\{0\}, \{1\}\}, \quad \mathcal{O}_2 = \{a_2\} = \{\{2\}\}.$$

- The corresponding orbit joins are:

$$b_1 = a_0 \vee a_1 = \{0, 1\}, \quad b_2 = a_2 = \{2\}.$$

- The fixed-point subalgebra B^G consists of the $2^2 = 4$ sets:

$$B^G = \{\emptyset, \{0, 1\}, \{2\}, \{0, 1, 2\}\}.$$

- The atoms of B^G are $\{0, 1\}$ and $\{2\}$, exactly matching b_1 and b_2 .

We verify this calculation in Lean:

Lean 4 formalization: appendix module `BSinMeasurementTheory/Course21373Numerical.lean`.

4.3 21-321 — Interactive Theorem Proving

Please

- (a) State and prove Scott's Theorem 1.1 (the finite-dimensional separating-hyperplane form) as a formal theorem, using `Mathlib`'s convex-cone and dual-space machinery. Your statement must quantify correctly over an arbitrary finite-dimensional real vector space, not a fixed $\mathbb{R}^n$.

- (b) Prove it without `sorry`, `admit`, or any axiom beyond what `Mathlib` already assumes.
- (c) Write a second, independent formalization of Theorem 1.2 (the combinatorial recasting over rational vectors) and prove the two theorems are equivalent as a third Lean theorem — not just informally equivalent.

Requires: dependent type theory, tactic-mode proof, correct universal quantification, and using existing library machinery rather than reproving convexity from scratch.

Prove it in English and then in Lean 4, using as much `Mathlib` as possible. No uses of `sorry`, `admit` or axiom are allowed. If you do any numerical calculations for examples to illustrate the proof, reproduce them after the proof in Lean.

1. Mathematical Background and Statement of Scott's Theorems

In his seminal paper *Measurement Structures and Linear Inequalities* (1964), Dana Scott established the foundational connection between qualitative measurement structures (such as subjective probability, semiordeers, and additive conjoint measurement) and the geometry of linear inequalities and separating hyperplanes.

****Theorem 1.1 (Finite-Dimensional Separating Hyperplane / Geometric Form)****

Let V be an arbitrary finite-dimensional real vector space, and let $X, Y \subset V$ be finite sets. There exists a linear functional $f \in V^*$ such that:

$$f(x) \geq 0 \quad (\forall x \in X) \quad \text{and} \quad f(y) > 0 \quad (\forall y \in Y)$$

if and only if for every collection of non-negative real multipliers $\lambda : X \rightarrow \mathbb{R}_{\geq 0}$ and $\mu : Y \rightarrow \mathbb{R}_{\geq 0}$ such that $\sum_{y \in Y} \mu(y) > 0$,

$$\sum_{x \in X} \lambda(x)x + \sum_{y \in Y} \mu(y)y \neq 0.$$

Proof Idea. - ($\implies$) **Necessity:** Suppose such an f exists. If $\sum_{x \in X} \lambda(x)x + \sum_{y \in Y} \mu(y)y = 0$ with $\lambda(x) \geq 0, \mu(y) \geq 0$, applying f yields:

$$\sum_{x \in X} \lambda(x)f(x) + \sum_{y \in Y} \mu(y)f(y) = f(0) = 0.$$

Since $\lambda(x)f(x) \geq 0$ for all x and $\mu(y)f(y) \geq 0$ for all y , each term must be zero. But $\sum_{y \in Y} \mu(y) > 0$ means there is some $y_0 \in Y$ with $\mu(y_0) > 0$. Since $f(y_0) > 0$, we have $\mu(y_0)f(y_0) > 0$, a contradiction.

- ($\impliedby$) **Sufficiency:** Consider the convex cone $C = \text{cone}(X) + \text{cone}_+(Y)$. The non-cancellation condition states that $0 \notin \text{conv}(Y) + \text{cone}(X)$, which is a compact/closed convex separation problem. By the Minkowski–Hahn–Banach separating hyperplane theorem on finite-dimensional V , the disjoint closed convex set and closed cone can be strictly separated by a hyperplane $f \in V^*$.

****Theorem 1.2 (Combinatorial / Rational Cancellation Form)****

Let V_0 be a finite-dimensional $\mathbb{Q}$ -vector space (or let X, Y be vectors with rational coordinates). There exists a $\mathbb{Q}$ -linear functional $f_{\mathbb{Q}} \in V_0^*$ such that:

$$f_{\mathbb{Q}}(x) \geq 0 \quad (\forall x \in X) \quad \text{and} \quad f_{\mathbb{Q}}(y) > 0 \quad (\forall y \in Y)$$

if and only if for every collection of natural numbers (or non-negative integers) $n : X \rightarrow \mathbb{N}$ and $m : Y \rightarrow \mathbb{N}$ such that $\sum_{y \in Y} m(y) > 0$,

$$\sum_{x \in X} n(x)x + \sum_{y \in Y} m(y)y \neq 0.$$

This integer-weighted condition is precisely **Scott's Finite Cancellation Axiom**.

****Theorem 1.3 (Equivalence of the Real and Combinatorial Forms)****

The real hyperplane separation condition (Theorem 1.1) and the rational/combinatorial finite cancellation condition (Theorem 1.2) are equivalent: 1. Real solvability implies rational/combinatorial non-cancellation. 2. If integer cancellation were violated, cleared denominators would give a rational (hence real) violation. 3. By the density of $\mathbb{Q}$ in $\mathbb{R}$ and linear programming duality (or Fourier–Motzkin elimination), a finite homogeneous system of linear inequalities with rational coefficients has a real solution if and only if it has a rational solution.

2. Complete Lean 4 Formalization

Below is the complete Lean 4 formalization using Mathlib's linear algebra and dual space machinery. Lean 4 formalization: appendix module `BSinMeasurementTheory/Course21321.lean`.

3. Concrete Numerical Example (Kraft–Pratt–Seidenberg Violation)

To illustrate how Scott's finite cancellation detects non-representability: Let $\Omega = \{\omega_1, \omega_2, \omega_3, \omega_4, \omega_5\}$. Consider the comparative probability relations: 1. $A_1 = \{\omega_1, \omega_2\} \prec B_1 = \{\omega_3, \omega_4, \omega_5\} \implies \mathbf{1}_{B_1} - \mathbf{1}_{A_1} \in Y$ 2. $A_2 = \{\omega_2, \omega_3\} \prec B_2 = \{\omega_1, \omega_4, \omega_5\} \implies \mathbf{1}_{B_2} - \mathbf{1}_{A_2} \in Y$ 3. $A_3 = \{\omega_1, \omega_3\} \prec B_3 = \{\omega_2, \omega_4, \omega_5\} \implies \mathbf{1}_{B_3} - \mathbf{1}_{A_3} \in Y$ 4. $A_4 = \{\omega_4, \omega_5\} \prec B_4 = \{\omega_1, \omega_2, \omega_3\} \implies \mathbf{1}_{B_4} - \mathbf{1}_{A_4} \in Y$

Summing all 4 difference vectors with weights $m = (1, 1, 1, 1)$:

$$\sum_{i=1}^4 (\mathbf{1}_{B_i} - \mathbf{1}_{A_i}) = (1 - 2) + (1 - 2) + (1 - 2) + (3 - 3) = \mathbf{0} \in \mathbb{R}^5$$

Because $\sum m_i = 4 > 0$ and the sum of difference vectors equals $\mathbf{0}$, Scott's Finite Cancellation fails, proving that no additive probability measure representing these relations can exist.

Lean 4 formalization: appendix module `BSinMeasurementTheory/Course21321Kraft.lean`.

5 Phase 3: Duality and set theory

Linear-programming duality and the ultrafilter engine of the infinite case.

5.1 21-292 — Operations Research I

Consider the LP: minimize $c^T x$ subject to $Ax \geq b$, $x \geq 0$, where $A \in \mathbb{Q}^{m \times n}$.

- (a) State and prove LP duality (weak and strong) for this pair.
- (b) Using the simplex method by hand on a concrete 3×3 instance you construct where the feasible cone is exactly the setting of Scott's Theorem 1.1, exhibit the separating hyperplane as the dual optimal solution.
- (c) Prove that Kuhn–Tucker's separating-hyperplane theorem (the one Scott cites without reproving) follows from LP strong duality applied to your construction — i.e., derive Scott's cited result from the simplex-based machinery, not the other way around.

Requires: simplex method, duality, and the actual logical dependency Scott leans on.

Prove it in English and then in Lean 4, using as much Mathlib as possible. No uses of sorry, admit or axiom are allowed. If you do any numerical calculations for examples to illustrate the proof, reproduce them after the proof in Lean.

(a) Linear Programming Duality: State and Proof

Let $A \in \mathbb{Q}^{m \times n}$, $b \in \mathbb{Q}^m$, and $c \in \mathbb{Q}^n$. We consider the primal-dual pair of linear programs in standard inequality form:

$$\text{(Primal LP)} \quad \text{minimize } c^T x \quad \text{subject to } Ax \geq b, \ x \geq 0$$

$$\text{(Dual LP)} \quad \text{maximize } b^T y \quad \text{subject to } A^T y \leq c, \ y \geq 0$$

1. Weak Duality Theorem

Theorem (Weak Duality): If $x \in \mathbb{Q}^n$ is primal feasible ($Ax \geq b$, $x \geq 0$) and $y \in \mathbb{Q}^m$ is dual feasible ($A^T y \leq c$, $y \geq 0$), then:

$$b^T y \leq c^T x$$

Proof: Since $y \geq 0$ and $Ax - b \geq 0$, their inner product is non-negative:

$$y^T (Ax - b) \geq 0 \implies y^T Ax \geq y^T b = b^T y$$

Similarly, since $x \geq 0$ and $c - A^T y \geq 0$:

$$(c - A^T y)^T x \geq 0 \implies c^T x \geq (A^T y)^T x = y^T Ax$$

Combining the two inequalities:

$$b^T y \leq y^T Ax \leq c^T x$$

Equality $b^T y = c^T x$ holds if and only if the complementary slackness conditions hold:

$$y^T (Ax - b) = 0 \quad \text{and} \quad (c - A^T y)^T x = 0$$

2. Strong Duality Theorem

Theorem (Strong Duality): If the primal LP has a finite optimal solution x^* , then the dual LP has an optimal solution y^* such that:

$$c^T x^* = b^T y^*$$

Proof via the Simplex Algorithm: 1. Introduce slack variables $s = Ax - b \geq 0$ to put the primal constraints in equality form: $Ax - s = b$, with $x, s \geq 0$. 2. Under Bland's pivot rule (or lexicographic perturbation), the simplex algorithm is guaranteed to terminate in finitely many steps at an optimal basic feasible solution x^* associated with basis matrix B . 3. At the optimal tableau, the vector of reduced costs is non-negative:

$$\bar{c}^T = c^T - c_B^T B^{-1} A \geq 0$$

4. Define the dual candidate vector $y^* = (B^{-1})^T c_B$. - **Dual Feasibility:** $c - A^T y^* = \bar{c} \geq 0 \implies A^T y^* \leq c$, and the slack components yield $y^* \geq 0$. - **Objective Value Match:** $b^T y^* = (y^*)^T b = c_B^T B^{-1} b = c_B^T x_B^* = c^T x^*$. 5. By Weak Duality, since y^* attains the upper bound $c^T x^*$, y^* is an optimal dual solution.

(b) Concrete 3×3 Instance & Hand Simplex Execution

In the setting of Dana Scott's 1964 paper (*Measurement Structures and Linear Inequalities*), we test whether a point lies in the convex cone generated by column vectors of A or can be separated by a linear functional y^* .

Let us construct a 3×3 instance:

$$A = \begin{pmatrix} 2 & 1 & 0 \\ 1 & 2 & 0 \\ 0 & 0 & 1 \end{pmatrix}, \quad b = \begin{pmatrix} 3 \\ 3 \\ 1 \end{pmatrix}, \quad c = \begin{pmatrix} 1 \\ 1 \\ 2 \end{pmatrix}$$

$$\begin{aligned} \text{Primal: } \min \quad & x_1 + x_2 + 2x_3 \\ \text{s.t. } \quad & 2x_1 + x_2 \geq 3 \\ & x_1 + 2x_2 \geq 3 \\ & x_3 \geq 1, \quad x_1, x_2, x_3 \geq 0 \end{aligned}$$

Dual Formulation:

$$\begin{aligned} \text{Dual: } \max \quad & 3y_1 + 3y_2 + y_3 \\ \text{s.t. } \quad & 2y_1 + y_2 \leq 1 \\ & y_1 + 2y_2 \leq 1 \\ & y_3 \leq 2, \quad y_1, y_2, y_3 \geq 0 \end{aligned}$$

Step-by-Step Simplex Iteration:

- **Phase I/II Basis:** The binding constraints at the primal optimum are $2x_1 + x_2 = 3$, $x_1 + 2x_2 = 3$, and $x_3 = 1$.
- Solving the active linear system gives the primal optimal vertex:

$$x^* = \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix}, \quad c^T x^* = 1(1) + 1(1) + 2(1) = 4$$

- The basis matrix $B = A = \begin{pmatrix} 2 & 1 & 0 \\ 1 & 2 & 0 \\ 0 & 0 & 1 \end{pmatrix}$.

- Inverting B :

$$B^{-1} = \begin{pmatrix} 2/3 & -1/3 & 0 \\ -1/3 & 2/3 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

- The dual optimal solution (separating hyperplane normal) is obtained directly from the simplex multipliers:

$$y^* = (B^{-1})^T c = \begin{pmatrix} 2/3 & -1/3 & 0 \\ -1/3 & 2/3 & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} 1 \\ 1 \\ 2 \end{pmatrix} = \begin{pmatrix} 1/3 \\ 1/3 \\ 2 \end{pmatrix}$$

- Checking dual feasibility and value:

$$A^T y^* = \begin{pmatrix} 2(1/3) + 1/3 \\ 1/3 + 2(1/3) \\ 1(2) \end{pmatrix} = \begin{pmatrix} 1 \\ 1 \\ 2 \end{pmatrix} \leq c \quad (\text{tight})$$

$$b^T y^* = 3(1/3) + 3(1/3) + 1(2) = 1 + 1 + 2 = 4$$

- Primal cost equals dual objective: $c^T x^* = b^T y^* = 4$.

(c) Derivation of Scott's Theorem 1.1 / Kuhn-Tucker Separation from LP Duality

Scott's Theorem 1.1 (Scott, 1964): Let $v_1, \dots, v_m \in \mathbb{Q}^n$. Exactly one of the following holds: 1. There exists $y \in \mathbb{Q}^n$ such that $v_i^T y > 0$ for all $i \in \{1, \dots, m\}$. 2. There exist $\lambda_1, \dots, \lambda_m \geq 0$, not all zero, such that $\sum_{i=1}^m \lambda_i v_i = 0$.

Proof using LP Strong Duality:

Consider the linear program:

$$\min 0^T y \quad \text{subject to } Vy \geq \mathbf{1}$$

where $V \in \mathbb{Q}^{m \times n}$ has rows v_i^T , and $\mathbf{1} = (1, \dots, 1)^T$.

- By scaling, condition (1) is equivalent to the existence of $y \in \mathbb{Q}^n$ satisfying $Vy \geq \mathbf{1}$ (primal feasibility).
- The dual of this LP is:

$$\max \mathbf{1}^T \lambda \quad \text{subject to } V^T \lambda = 0, \lambda \geq 0$$

- **Case A:** If the primal is feasible, then by Weak Duality, the dual objective is bounded from above by $0^T y = 0$. Since $\lambda \geq 0$, $\mathbf{1}^T \lambda \leq 0 \implies \lambda = 0$. Thus no non-trivial non-negative combination can sum to zero.
- **Case B:** If the primal is infeasible, then the dual is either infeasible or unbounded. But $\lambda = 0$ is always dual feasible ($V^T 0 = 0, 0 \geq 0$), so the dual must be unbounded. An unbounded dual implies the existence of a ray $\lambda \geq 0$ with $\mathbf{1}^T \lambda > 0$ (i.e. $\lambda \neq 0$) such that $V^T \lambda = \sum_{i=1}^m \lambda_i v_i = 0$.

This completes the logical reduction of Kuhn-Tucker / Scott Theorem 1.1 directly from LP duality.

Lean 4 Formalization

The complete Lean 4 proof below formalizes: 1. Weak LP Duality in full generality for inner products. 2. Constructive certificates of optimality and hyperplane separation. 3. The exact 3×3 Scott-type cone instance verified without `sorry`, `admit`, or non-standard axioms.

Lean 4 formalization: `appendix module BSinMeasurementTheory/Course21292.lean`.

Numerical Calculations

The numerical values computed during the simplex method and formal verification:

1. **Primal Cost:**

$$\text{dot}(c, x^*) = (1)(1) + (1)(1) + (2)(1) = 1 + 1 + 2 = 4$$

2. **Dual Cost:**

$$\text{dot}(b, y^*) = (3)(1/3) + (3)(1/3) + (1)(2) = 1 + 1 + 2 = 4$$

3. **Primal Slack Vector** ($Ax^* - b$):

$$Ax^* - b = \begin{pmatrix} 2(1) + 1(1) + 0(1) - 3 \\ 1(1) + 2(1) + 0(1) - 3 \\ 0(1) + 0(1) + 1(1) - 1 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix}$$

4. **Dual Slack Vector** ($c - A^T y^*$):

$$c - A^T y^* = \begin{pmatrix} 1 - (2(1/3) + 1(1/3) + 0(2)) \\ 1 - (1(1/3) + 2(1/3) + 0(2)) \\ 2 - (0(1/3) + 0(1/3) + 1(2)) \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix}$$

Lean 4 formalization: `appendix module BSinMeasurementTheory/Course21292Numerical.lean`.

5.2 21-329 — Set Theory

Let B be an infinite Boolean algebra.

- (a) Prove that every proper filter on B extends to an ultrafilter, using Zorn's Lemma explicitly (state the poset, the chains, and verify the hypothesis).
- (b) Prove that the set of ultrafilters on B , given the Stone topology, is nonempty precisely because of (a).
- (c) Prove that if B is countably infinite, the set of ultrafilters on B has cardinality $2^{\aleph_0}$, using a cardinal-arithmetic argument (not just citing the result).

Requires: Zorn's Lemma applied correctly, cardinal arithmetic — the exact tools the unproven Hahn–Banach extension in Scott's closing remark runs on.

Prove it in English and then in Lean 4, using as much Mathlib as possible. No uses of sorry, admit or axiom are allowed. If you do any numerical calculations for examples to illustrate the proof, describe the calculations in English and then code them in Lean 4 after that.

Here is the complete proof in English, followed by a formalization in Lean 4.

****Mathematical Proof****

****(a) Ultrafilter Extension Theorem via Zorn's Lemma****

Let $(B, \wedge, \vee, \neg, 0, 1)$ be a Boolean algebra.

1. Definitions:

- A subset $F \subseteq B$ is a **filter** if:
 1. $1 \in F$,
 2. $\forall x, y \in F, x \wedge y \in F$,
 3. $\forall x \in F, y \in B, x \leq y \implies y \in F$.
- A filter F is **proper** if $0 \notin F$ (equivalently, $F \neq B$).
- An **ultrafilter** is a maximal proper filter with respect to subset inclusion $\subseteq$.

2. The Poset: Let F_0 be a fixed proper filter on B . Define the poset:

$$\mathcal{P} = \{F \subseteq B \mid F \text{ is a proper filter on } B \text{ and } F_0 \subseteq F\}$$

ordered by subset inclusion $\subseteq$.

- $\mathcal{P}$ is non-empty since $F_0 \in \mathcal{P}$.

3. Verification of the Chain Condition (Zorn's Hypothesis): Let $\mathcal{C} \subseteq \mathcal{P}$ be a non-empty chain (a totally ordered family of filters under $\subseteq$). Define:

$$U = \bigcup_{F \in \mathcal{C}} F$$

We verify that $U \in \mathcal{P}$:

- **Contains F_0 :** Since $\mathcal{C} \neq \emptyset$, choose $F \in \mathcal{C}$. Then $F_0 \subseteq F \subseteq U$.
- **Contains 1:** For any $F \in \mathcal{C}$, $1 \in F \subseteq U$.
- **Upward Closed:** If $x \in U$ and $x \leq y$, there is some $F \in \mathcal{C}$ with $x \in F$. Since F is a filter, $y \in F \subseteq U$.
- **Closed under Finite Meets:** If $x, y \in U$, there exist $F_1, F_2 \in \mathcal{C}$ such that $x \in F_1$ and $y \in F_2$. Since $\mathcal{C}$ is a chain, either $F_1 \subseteq F_2$ or $F_2 \subseteq F_1$. Without loss of generality, assume $F_1 \subseteq F_2$. Then $x, y \in F_2$, so $x \wedge y \in F_2 \subseteq U$.

- **Properness** ($0 \notin U$): If $0 \in U$, then $0 \in F$ for some $F \in \mathcal{C}$, contradicting that every $F \in \mathcal{P}$ is proper ($0 \notin F$). Thus $0 \notin U$.

Hence U is a proper filter extending F_0 , so $U \in \mathcal{P}$, and $F \subseteq U$ for all $F \in \mathcal{C}$. Thus, U is an upper bound for $\mathcal{C}$ in $\mathcal{P}$.

4. **Conclusion by Zorn's Lemma:** Every chain in $\mathcal{P}$ has an upper bound in $\mathcal{P}$. By Zorn's Lemma, $\mathcal{P}$ has a maximal element M .

- If M' is any proper filter on B containing M , then $F_0 \subseteq M \subseteq M'$, so $M' \in \mathcal{P}$.
- By the maximality of M in $\mathcal{P}$, we must have $M' = M$. Therefore, M is a maximal proper filter (ultrafilter) on B extending F_0 . ■

****(b) Non-emptiness of the Stone Space****

The Stone space $S(B)$ is the set of all ultrafilters on B equipped with the Stone topology. 1. Since B is an infinite Boolean algebra, $0 \neq 1$. 2. The principal filter generated by 1, $F_0 = \{1\} = \{x \in B \mid 1 \leq x\}$, is a proper filter because $0 \neq 1$ implies $0 \notin F_0$. 3. By part (a), F_0 extends to an ultrafilter U on B . 4. Hence $U \in S(B)$, which proves $S(B) \neq \emptyset$. ■

****(c) Cardinality of Ultrafilters on a Countably Infinite Boolean Algebra****

Let B be a countably infinite Boolean algebra, so $|B| = \aleph_0$.

1. **Upper Bound:** Every ultrafilter is a subset of B , so $S(B) \subseteq \mathcal{P}(B)$.

$$|S(B)| \leq |\mathcal{P}(B)| = 2^{|B|} = 2^{\aleph_0}$$

2. **Lower Bound:** Since B is infinite, we can inductively construct a dyadic tree of non-zero elements $\{a_s\}_{s \in \{0,1\}^{<\omega}} \subseteq B \setminus \{0\}$ such that:

- $a_\emptyset = 1$,
- For each sequence s , $a_{s \smallfrown 0} \wedge a_{s \smallfrown 1} = 0$,
- $a_{s \smallfrown 0} \leq a_s$ and $a_{s \smallfrown 1} \leq a_s$.

For each infinite path $\sigma \in \{0,1\}^{\mathbb{N}}$, the family $\{a_{\sigma|_n} \mid n \in \mathbb{N}\}$ is a descending chain of non-zero elements, hence it has the finite intersection property and generates a proper filter:

$$F_\sigma = \{x \in B \mid \exists n \in \mathbb{N}, a_{\sigma|_n} \leq x\}$$

By part (a), each F_σ extends to an ultrafilter $U_\sigma \in S(B)$.

- **Injectivity:** Let $\sigma \neq \tau \in \{0,1\}^{\mathbb{N}}$. Let n be the minimal index where $\sigma(n) \neq \tau(n)$, and let $s = \sigma|_n = \tau|_n$. Without loss of generality, $\sigma(n) = 0$ and $\tau(n) = 1$. Then $a_{s \smallfrown 0} \in F_\sigma \subseteq U_\sigma$ and $a_{s \smallfrown 1} \in F_\tau \subseteq U_\tau$. Since $a_{s \smallfrown 0} \wedge a_{s \smallfrown 1} = 0$, U_σ and U_τ cannot be equal (otherwise $0 \in U_\sigma$, impossible). Thus $\sigma \mapsto U_\sigma$ is an injection from $\{0,1\}^{\mathbb{N}}$ into $S(B)$.

Therefore:

$$|S(B)| \geq |\{0, 1\}^{\mathbb{N}}| = 2^{\aleph_0}$$

3. Conclusion by Cantor–Bernstein–Schröder:

$$2^{\aleph_0} \leq |S(B)| \leq 2^{\aleph_0} \implies |S(B)| = 2^{\aleph_0}$$

■

****Lean 4 Formal Proof****

The following complete Lean 4 formalization proves the extension theorem via Zorn’s Lemma and the cardinal arithmetic bounds without `sorry`, `admit`, or extra axioms.

Lean 4 formalization: `appendix module BSinMeasurementTheory/Course21329.lean`.

6 Phase 4: Stone spaces and measures

Stone representation and Riesz representation.

6.1 21-651 — General Topology

Let B be a Boolean algebra and let X be its Stone space (the space of ultrafilters on B with the topology generated by $\{\hat{b} : b \in B\}$, $\hat{b} = \{U : b \in U\}$).

- (a) Prove X is compact.
- (b) Prove X is Hausdorff and totally disconnected.
- (c) Prove the sets $\hat{b}$ are exactly the clopen sets of X , and conclude B is isomorphic to the Boolean algebra of clopen subsets of X (Stone’s representation theorem).
- (d) Prove your compactness argument in (a) as an instance of Tychonoff’s theorem applied to 2^B with X realized as a closed subspace — do not just cite Tychonoff, embed and verify closedness.

Requires: compactness, Tychonoff’s theorem, separation axioms — exactly what Kelley’s method needs to represent an infinite Boolean algebra as a Stone space.

Prove it in English and then in Lean 4, using as much Mathlib as possible. No uses of `sorry`, `admit` or axiom are allowed. If you do any numerical calculations for examples to illustrate the proof, reproduce them after the proof in English and then Lean. Every time you give a concrete example in English you must follow it with the equivalent Lean code. We are using Lean v4.34. Please refer to <https://leanprover-community.github.io/mathlib-overview.html> for API.

Here is the complete proof and formalization written to standard **Mathlib** specifications, starting with the full mathematical English exposition followed by the complete, machine-checked Lean 4 code using `import Mathlib` without any `sorry`, `admit`, or custom axioms.

6.1.1 Mathematical Proof

Let $(B, \wedge, \vee, \neg, 0, 1)$ be a Boolean algebra.

Setup and Topological Realization

1. **Ultrafilters as Homomorphisms:** Every ultrafilter U on B corresponds bijectively to a Boolean homomorphism $u : B \rightarrow \mathbf{2}$, where $\mathbf{2} = \{\text{False}, \text{True}\}$ is the two-element Boolean algebra, via $u(b) = \text{True} \iff b \in U$.
2. **The Ambient Space $\mathbf{2}^B$:** Equip $\mathbf{2}$ with the discrete topology. The function space $\mathbf{2}^B = \prod_{b \in B} \mathbf{2}$ is equipped with the product topology (the topology of pointwise convergence).
3. **The Stone Space X :** $X = \text{Stone}(B) = \text{Hom}_{\mathbf{BA}}(B, \mathbf{2}) \subseteq \mathbf{2}^B$, endowed with the subspace topology induced from $\mathbf{2}^B$.

(d) Realization as a Closed Subspace of $\mathbf{2}^B$ & (a) Compactness

1. **Closedness of the Embedding:** A function $f \in \mathbf{2}^B$ is a Boolean homomorphism if and only if it satisfies the pointwise equations:
 - $f(1) = \text{True}$
 - $f(0) = \text{False}$
 - $\forall a, b \in B, f(a \wedge b) = (f(a) \wedge f(b))$
 - $\forall a, b \in B, f(a \vee b) = (f(a) \vee f(b))$
 - $\forall a \in B, f(\neg a) = \neg f(a)$

Since $\mathbf{2}$ is discrete, all logical connectives on $\mathbf{2}$ are continuous. Coordinate evaluation $\pi_b : \mathbf{2}^B \rightarrow \mathbf{2}$ is continuous for every $b \in B$. Therefore, each condition defines an equalizer of two continuous maps, which is a closed set in $\mathbf{2}^B$:

$$X = \pi_1^{-1}(\{\text{True}\}) \cap \pi_0^{-1}(\{\text{False}\}) \cap \bigcap_{a, b \in B} \text{Eq}(\pi_{a \wedge b}, \pi_a \wedge \pi_b) \cap \bigcap_{a, b \in B} \text{Eq}(\pi_{a \vee b}, \pi_a \vee \pi_b) \cap \bigcap_{a \in B} \text{Eq}(\pi_{\neg a}, \neg \circ \pi_a)$$

Being an intersection of closed sets, X is a **closed subspace** of $\mathbf{2}^B$.

2. **Compactness (a):** The two-point discrete space $\mathbf{2}$ is compact and Hausdorff. By **Tychonoff's Theorem**, the product space $\mathbf{2}^B$ is compact. Because X is a closed subspace of the compact space $\mathbf{2}^B$, $X = \text{Stone}(B)$ is **compact**.

(b) Hausdorff and Totally Disconnected

1. **Hausdorff (T_2):** $\mathbf{2}^B$ is a product of T_2 spaces, hence T_2 . Any subspace of a T_2 space is T_2 , so X is Hausdorff.

2. **Totally Disconnected:** The product topology on 2^B has a subbasis of cylinder sets $\pi_b^{-1}(\{i\})$ which are both open and closed. Thus 2^B is zero-dimensional (having a basis of clopen sets). Subspaces of zero-dimensional Hausdorff spaces are zero-dimensional and therefore **totally disconnected**.

(c) Clopen Sets & Stone's Representation Theorem

1. **Clopen Sets:** For each $b \in B$, define the Stone evaluation map:

$$\sigma(b) = \hat{b} = \{u \in X \mid u(b) = \text{True}\} = \pi_b^{-1}(\{\text{True}\}) \cap X$$

Since $\{\text{True}\}$ is open and closed in 2 , $\hat{b}$ is clopen in X . Its complement is:

$$X \setminus \hat{b} = \{u \in X \mid u(b) = \text{False}\} = \{u \in X \mid u(\neg b) = \text{True}\} = \widehat{\neg b}$$

2. **Boolean Isomorphism:** The map $\sigma : B \rightarrow \text{Clopen}(X)$ is a homomorphism preserving $\top, \perp, \wedge, \vee, \neg$:

- $\sigma(0) = \emptyset, \sigma(1) = X$
- $\sigma(a \wedge b) = \sigma(a) \cap \sigma(b)$
- $\sigma(a \vee b) = \sigma(a) \cup \sigma(b)$
- $\sigma(\neg a) = X \setminus \sigma(a)$ By the Ultrafilter Lemma, distinct elements $a \neq b \in B$ are separated by some ultrafilter u , making σ injective. Since the basic clopens $\{\hat{b}\}$ generate the topology and X is compact, every clopen set is a finite union of basic clopens, hence in the image of σ . Thus $B \cong \text{Clopen}(X)$.

Concrete Numerical Example

English: Consider the 4-element Boolean algebra $B = \mathcal{P}(\{0, 1\}) = \{\emptyset, \{0\}, \{1\}, \{0, 1\}\}$. - The ultrafilters are the principal ultrafilters u_0, u_1 :

$$u_0(S) = (0 \in S), \quad u_1(S) = (1 \in S)$$

- The Stone space has 2 points $X = \{u_0, u_1\}$. - Clopen evaluation sets: - $\sigma(\emptyset) = \emptyset$ - $\sigma(\{0\}) = \{u_0\}$ - $\sigma(\{1\}) = \{u_1\}$ - $\sigma(\{0, 1\}) = \{u_0, u_1\} = X$

6.1.2 Lean 4 Formalization

Lean 4 formalization: appendix module `BSinMeasurementTheory/Course21651.lean`.

6.2 21-720 — Measure and Integration

Let X be a compact Hausdorff space and $\Lambda : C(X) \rightarrow \mathbb{R}$ a positive linear functional.

- (a) Prove Λ is bounded, i.e., continuous with respect to the sup norm.
- (b) State and prove the Riesz Representation Theorem for this Λ : construct the regular Borel measure μ such that $\Lambda(f) = \int_X f d\mu$ for all $f \in C(X)$, including proving μ is well-defined and regular.
- (c) Specialize to X a Stone space of a Boolean algebra B : show the resulting measure μ restricts to a finitely additive measure on B itself, via $\mu(b) := \mu(\hat{b})$.

Requires: Lebesgue-style integration theory, signed/regular measures, and Riesz Representation — the exact mechanism Kelley uses to turn a linear functional on an infinite Boolean algebra into a genuine measure.

Prove it in English and then in Lean 4, using as much Mathlib as possible. No uses of sorry, admit or axiom are allowed. If you do any numerical calculations for examples to illustrate the proof, reproduce them after the proof in English and then Lean. Every time you give a concrete example in English you must follow it with the equivalent Lean code. We are using Lean v4.34. Please refer to <https://leanprover-community.github.io/mathlib-overview.html> for API.

Mathematical Exposition in English

Let X be a compact Hausdorff topological space, and let $C(X) = C(X, \mathbb{R})$ denote the Banach space of continuous real-valued functions on X , equipped with the supremum norm:

$$\|f\|_\infty = \sup_{x \in X} |f(x)|.$$

Since X is compact, every continuous function is bounded and achieves its bounds.

A linear functional $\Lambda : C(X) \rightarrow \mathbb{R}$ is **positive** if:

$$f \geq 0 \implies \Lambda(f) \geq 0,$$

where $f \geq 0$ means $f(x) \geq 0$ for all $x \in X$.

Part (a): Boundedness of Λ

Theorem: Λ is continuous (bounded) with operator norm $\|\Lambda\|_{op} = \Lambda(\mathbf{1})$, where $\mathbf{1} \in C(X)$ is the constant function $\mathbf{1}(x) = 1$.

Proof: For any $f \in C(X)$ and every point $x \in X$:

$$-\|f\|_\infty \leq f(x) \leq \|f\|_\infty.$$

In terms of functions in $C(X)$, this implies:

$$\|f\|_\infty \mathbf{1} - f \geq 0 \quad \text{and} \quad \|f\|_\infty \mathbf{1} + f \geq 0.$$

Applying the positivity and linearity of Λ : 1. $\Lambda(\|f\|_\infty \mathbf{1} - f) = \|f\|_\infty \Lambda(\mathbf{1}) - \Lambda(f) \geq 0 \implies \Lambda(f) \leq \Lambda(\mathbf{1})\|f\|_\infty$. 2. $\Lambda(\|f\|_\infty \mathbf{1} + f) = \|f\|_\infty \Lambda(\mathbf{1}) + \Lambda(f) \geq 0 \implies -\Lambda(f) \leq \Lambda(\mathbf{1})\|f\|_\infty$.

Combining these two inequalities gives:

$$|\Lambda(f)| \leq \Lambda(\mathbf{1})\|f\|_\infty.$$

Since $\|\mathbf{1}\|_\infty = 1$, we have $|\Lambda(\mathbf{1})| = \Lambda(\mathbf{1})\|\mathbf{1}\|_\infty$, which establishes:

$$\|\Lambda\|_{op} = \sup_{f \neq 0} \frac{|\Lambda(f)|}{\|f\|_\infty} = \Lambda(\mathbf{1}) < \infty.$$

Hence, Λ is bounded and continuous. ■

Part (b): Riesz Representation Theorem

Theorem (Riesz–Markov–Kakutani): Let X be a compact Hausdorff space and $\Lambda : C(X) \rightarrow \mathbb{R}$ a positive linear functional. There exists a unique regular Borel measure μ on X such that:

$$\Lambda(f) = \int_X f d\mu \quad \text{for all } f \in C(X).$$

Construction and Proof:

1. **Definition on open sets and outer measure:** For any open set $U \subseteq X$, define:

$$\mu(U) := \sup \{ \Lambda(f) : f \in C(X), 0 \leq f \leq \mathbf{1}_U \}.$$

For an arbitrary subset $E \subseteq X$, define the outer measure μ^* :

$$\mu^*(E) := \inf \{ \mu(U) : U \supseteq E, U \text{ is open} \}.$$

2. μ^* is an outer measure:

- $\mu^*(\emptyset) = 0$ since the only continuous function $0 \leq f \leq \mathbf{1}_\emptyset$ is $f = 0$, so $\Lambda(f) = 0$.
- Monotonicity: $E_1 \subseteq E_2 \implies \mu^*(E_1) \leq \mu^*(E_2)$ by definition of the infimum over containing open sets.
- Countable subadditivity: Let $E \subseteq \bigcup_{n=1}^\infty E_n$. For any $\epsilon > 0$, choose open $U_n \supseteq E_n$ such that $\mu(U_n) \leq \mu^*(E_n) + 2^{-n}\epsilon$. For any $f \in C(X)$ with $0 \leq f \leq \mathbf{1}_{\bigcup U_n}$, the compact support $\text{supp}(f)$ is covered by finitely many U_n , say $U_1, \dots, U_m$. Subordinating a partition of unity $h_1 + \dots + h_m = 1$ on $\text{supp}(f)$, we have:

$$\Lambda(f) = \sum_{k=1}^m \Lambda(fh_k) \leq \sum_{k=1}^m \mu(U_k) \leq \sum_{n=1}^\infty \mu^*(E_n) + \epsilon.$$

Taking the supremum over f and letting $\epsilon \rightarrow 0$ gives $\mu^*(E) \leq \sum_{n=1}^\infty \mu^*(E_n)$.

3. **Carathéodory Measurability of Borel sets:** A set $A \subseteq X$ is Carathéodory measurable if for every test set $T \subseteq X$:

$$\mu^*(T) \geq \mu^*(T \cap A) + \mu^*(T \setminus A).$$

By Urysohn's Lemma on compact Hausdorff spaces, every open set U satisfies this condition. Because the collection of measurable sets forms a σ -algebra containing all open sets, it contains all Borel sets $\mathcal{B}(X)$. The restriction $\mu = \mu^*|_{\mathcal{B}(X)}$ is a countably additive Borel measure.

4. Regularity:

- **Outer regularity:** For any Borel set E , $\mu(E) = \inf\{\mu(U) : U \supseteq E, U \text{ open}\}$ holds directly from the definition of μ^* .
- **Inner regularity:** Since X is compact, for any Borel set E , the complement $X \setminus E$ is Borel. Applying outer regularity to $X \setminus E$:

$$\mu(E) = \mu(X) - \mu(X \setminus E) = \sup\{\mu(K) : K \subseteq E, K \text{ compact}\}.$$

Thus, μ is regular.

5. **Integral representation:** For $f \in C(X)$ with $f \geq 0$, subdivide $[0, \|f\|_\infty]$ by $y_k = \frac{k\|f\|_\infty}{N}$ and open level sets $U_k = \{x \in X : f(x) > y_k\}$. Both $\Lambda(f)$ and $\int_X f d\mu$ are squeezed between the lower and upper Darboux sums $\sum_{k=1}^N (y_k - y_{k-1})\mu(U_k)$. As $N \rightarrow \infty$, the difference vanishes, proving $\Lambda(f) = \int_X f d\mu$. Linearity extends the identity to all $f \in C(X)$.

6. **Uniqueness:** If ν is another regular Borel measure representing Λ , then for any open U :

$$\nu(U) = \sup_{K \subseteq U} \nu(K) = \sup_{0 \leq f \leq \mathbf{1}_U} \int_X f d\nu = \sup_{0 \leq f \leq \mathbf{1}_U} \Lambda(f) = \mu(U).$$

Since μ and ν agree on all open sets and are outer regular, $\mu(E) = \nu(E)$ for all $E \in \mathcal{B}(X)$. ■

Part (c): Specialization to the Stone Space of a Boolean Algebra

Let $B = (B, \vee, \wedge, \neg, \perp, \top)$ be a Boolean algebra. 1. The **Stone space** $X = \text{Ult}(B)$ is the compact, Hausdorff, totally disconnected space of ultrafilters on B , with basic open sets:

$$\hat{b} := \{u \in \text{Ult}(B) : b \in u\}, \quad \text{for } b \in B.$$

2. The Stone representation map $b \mapsto \hat{b}$ is an isomorphism of Boolean algebras from B onto the algebra $\text{Clopen}(X)$ of clopen subsets of X , preserving joins, meets, complements, $\perp$, and $\top$:

$$\widehat{a \vee b} = \hat{a} \cup \hat{b}, \quad \widehat{a \wedge b} = \hat{a} \cap \hat{b}, \quad \widehat{\perp} = \emptyset, \quad \widehat{\top} = X.$$

3. Since each $\hat{b}$ is clopen, its characteristic function $\mathbf{1}_{\hat{b}} : X \rightarrow \mathbb{R}$ is continuous: $\mathbf{1}_{\hat{b}} \in C(X)$.

Define the restriction $\mu_B : B \rightarrow \mathbb{R}$ by:

$$\mu_B(b) := \mu(\hat{b}) = \int_X \mathbf{1}_{\hat{b}} d\mu = \Lambda(\mathbf{1}_{\hat{b}}).$$

Finite Additivity on B : - **Non-negativity:** $\mathbf{1}_{\hat{b}} \geq 0 \implies \mu_B(b) = \Lambda(\mathbf{1}_{\hat{b}}) \geq 0$. - **Null element:** $\mu_B(\perp) = \Lambda(\mathbf{1}_{\emptyset}) = \Lambda(0) = 0$. - **Disjoint Additivity:** If $a, b \in B$ with $a \wedge b = \perp$, then $\hat{a} \cap \hat{b} = \widehat{a \wedge b} = \widehat{\perp} = \emptyset$. The indicators satisfy:

$$\mathbf{1}_{\widehat{a \vee b}} = \mathbf{1}_{\hat{a} \cup \hat{b}} = \mathbf{1}_{\hat{a}} + \mathbf{1}_{\hat{b}}.$$

By linearity of Λ :

$$\mu_B(a \vee b) = \Lambda(\mathbf{1}_{\widehat{a \vee b}}) = \Lambda(\mathbf{1}_{\hat{a}} + \mathbf{1}_{\hat{b}}) = \Lambda(\mathbf{1}_{\hat{a}}) + \Lambda(\mathbf{1}_{\hat{b}}) = \mu_B(a) + \mu_B(b).$$

Thus, μ_B is a finitely additive non-negative measure on B . ■

Concrete Numerical Example

Example: Consider the 4-element Boolean algebra $B = \mathcal{P}(\{0, 1\})$. The Stone space is the 2-point discrete space $X = \{0, 1\}$. Continuous functions $C(X)$ are represented as pairs $(f(0), f(1)) \in \mathbb{R}^2$. Define the positive linear functional $\Lambda : C(X) \rightarrow \mathbb{R}$ by:

$$\Lambda(f) = 3f(0) + 2f(1).$$

Calculations: 1. Functional Norm:

$$\Lambda(1) = \Lambda(1, 1) = 3(1) + 2(1) = 5.$$

Thus $\|\Lambda\|_{op} = 5$. 2. **Measure values on B :** - $\mu_B(\emptyset) = \Lambda(0, 0) = 3(0) + 2(0) = 0$ - $\mu_B(\{0\}) = \Lambda(1, 0) = 3(1) + 2(0) = 3$ - $\mu_B(\{1\}) = \Lambda(0, 1) = 3(0) + 2(1) = 2$ - $\mu_B(\{0, 1\}) = \Lambda(1, 1) = 3(1) + 2(1) = 5$.

Finite Additivity Check:

$$\mu_B(\{0\} \cup \{1\}) = \mu_B(\{0\}) + \mu_B(\{1\}) = 3 + 2 = 5 = \mu_B(\{0, 1\}).$$

Lean 4 Formalization

The following complete Lean 4 code formalizes the theorem for (a), the Stone representation measure construction for (c), and the numerical calculations with zero uses of `sorry`, `admit`, or `axiom`.

Lean 4 formalization: `appendix module BSinMeasurementTheory/Course21720.lean`.

7 Phase 5: The missing extension

Hahn–Banach, the step Scott cites and leaves unproved.

7.1 21-640 — Introduction to Functional Analysis

Let B be an infinite Boolean algebra, viewed as a sublattice of $C(X)$ (its Stone space) via $b \mapsto \mathbf{1}_{\hat{b}}$, and suppose φ_0 is a positive linear functional defined only on the (not necessarily dense) linear span of $\{\mathbf{1}_{\hat{b}} : b \in B\}$.

- State the Hahn–Banach theorem (dominated extension form) precisely, including the role of the sublinear dominating functional.
- Prove φ_0 extends to a positive linear functional φ on all of $C(X)$, exhibiting the dominating functional explicitly (the sup-norm works — prove it satisfies sublinearity and dominates φ_0 on the subspace).
- Prove this extension is exactly the missing step in Scott’s Theorem 4.1: combine with your Riesz Representation result from 21-720 to produce a regular Borel measure extending the finitely additive measure on B to a genuine measure on the Boolean algebra of all clopen sets of X .

Requires: Hahn–Banach itself, dual-space reasoning, and assembling it with the prior two courses into the actual argument Scott gestures at and leaves unproved.

Prove it in English and then in Lean 4, using as much Mathlib as possible. No uses of sorry, admit or axiom are allowed. If you do any numerical calculations for examples to illustrate the proof, reproduce them after the proof in English and then Lean. Every time you give a concrete example in English you must follow it with the equivalent Lean code. We are using Lean v4.34. Please refer to <https://leanprover-community.github.io/mathlib-overview.html> for API. Put all of the Lean code in a single section at the end which is self-contained and checkable.

(a) Hahn–Banach Theorem (Dominated Extension Form)

Precise Statement

Let V be a real vector space, and let $p : V \rightarrow \mathbb{R}$ be a **sublinear functional**, which means: 1. **Subadditivity**: $p(x + y) \leq p(x) + p(y)$ for all $x, y \in V$. 2. **Positive Homogeneity**: $p(cx) = cp(x)$ for all $c \geq 0$ and $x \in V$.

Let $M \subseteq V$ be a real linear subspace, and let $\varphi_0 : M \rightarrow \mathbb{R}$ be a linear functional satisfying the dominance condition:

$$\varphi_0(x) \leq p(x) \quad \forall x \in M$$

Then there exists a linear functional $\varphi : V \rightarrow \mathbb{R}$ such that: 1. **Extension**: $\varphi(x) = \varphi_0(x)$ for all $x \in M$. 2. **Global Dominance**: $\varphi(x) \leq p(x)$ for all $x \in V$.

Role of the Sublinear Functional

The sublinear functional p acts as a convex upper envelope controlling growth. In a 1-dimensional extension step (adjoining a single vector $v \notin M$), the condition $\varphi_0(m_1 - m_2) \leq p(m_1 - m_2) \leq p(m_1 + v) + p(-v - m_2)$ guarantees the existence of a real constant $c = \varphi(v)$ satisfying:

$$\sup_{m \in M} (\varphi_0(m) - p(m - v)) \leq c \leq \inf_{m \in M} (p(m + v) - \varphi_0(m))$$

Zorn's Lemma then extends this 1-step extension to arbitrary infinite-dimensional subspaces.

(b) Extension of φ_0 to a Positive Linear Functional on $C(X)$

Let X be the Stone space of the Boolean algebra B . Then X is a compact Hausdorff (totally disconnected) space. The elements of B correspond canonically to the clopen subsets $\hat{b} \subseteq X$, and their indicator functions $\mathbf{1}_{\hat{b}} \in C(X, \mathbb{R})$.

Let $M = \text{span}_{\mathbb{R}}\{\mathbf{1}_{\hat{b}} : b \in B\} \subseteq C(X, \mathbb{R})$, and let $\varphi_0 : M \rightarrow \mathbb{R}$ be a positive linear functional (i.e., $g \in M, g \geq 0 \implies \varphi_0(g) \geq 0$). Note that $\mathbf{1}_X = \mathbf{1}_{\hat{1}} \in M$, and $C := \varphi_0(\mathbf{1}_X) \geq 0$.

Step 1: Definition of the Sublinear Functional

Define $p : C(X, \mathbb{R}) \rightarrow \mathbb{R}$ by:

$$p(f) = C \|f\|_{\infty} = \varphi_0(\mathbf{1}_X) \sup_{x \in X} |f(x)|$$

1. **Positive Homogeneity**: For $c \geq 0$,

$$p(cf) = C \|cf\|_{\infty} = cC \|f\|_{\infty} = cp(f)$$

2. **Subadditivity**: By the triangle inequality for the supremum norm,

$$p(f + g) = C \|f + g\|_{\infty} \leq C(\|f\|_{\infty} + \|g\|_{\infty}) = p(f) + p(g)$$

Thus p is a sublinear functional on $C(X, \mathbb{R})$.

Step 2: Dominance on the Subspace M

Let $g \in M$. Since $-\|g\|_\infty \leq g(x) \leq \|g\|_\infty$ for all $x \in X$, we have:

$$\|g\|_\infty \mathbf{1}_X - g \geq 0$$

Since φ_0 is positive and linear on M :

$$\varphi_0(\|g\|_\infty \mathbf{1}_X - g) \geq 0 \implies \varphi_0(g) \leq \varphi_0(\|g\|_\infty \mathbf{1}_X) = \|g\|_\infty \varphi_0(\mathbf{1}_X) = p(g)$$

Thus $\varphi_0(g) \leq p(g)$ holds for all $g \in M$.

Step 3: Application of Hahn–Banach

By the Hahn–Banach dominated extension theorem (Part a), there exists a linear functional $\varphi : C(X, \mathbb{R}) \rightarrow \mathbb{R}$ such that: 1. $\varphi|_M = \varphi_0$. 2. $\varphi(f) \leq p(f) = C\|f\|_\infty$ for all $f \in C(X, \mathbb{R})$.

Step 4: Positivity of the Extension φ

Let $f \in C(X, \mathbb{R})$ with $f \geq 0$ pointwise. Consider the function $h = \|f\|_\infty \mathbf{1}_X - f$. Since $0 \leq f(x) \leq \|f\|_\infty$, we have $0 \leq h(x) \leq \|f\|_\infty$, which implies $\|h\|_\infty \leq \|f\|_\infty$.

Applying the dominance property to h :

$$\varphi(h) \leq p(h) = C\|h\|_\infty \leq C\|f\|_\infty$$

By linearity of φ :

$$\varphi(h) = \varphi(\|f\|_\infty \mathbf{1}_X - f) = \|f\|_\infty \varphi(\mathbf{1}_X) - \varphi(f) = C\|f\|_\infty - \varphi(f)$$

Combining the inequalities:

$$C\|f\|_\infty - \varphi(f) \leq C\|f\|_\infty \implies -\varphi(f) \leq 0 \implies \varphi(f) \geq 0$$

Therefore, φ is a positive linear functional on all of $C(X, \mathbb{R})$.

(c) Completing Scott's Theorem 4.1 via Riesz Representation

Dana Scott's Theorem 4.1 concerns extending a finitely additive measure μ_0 on a Boolean algebra B to a countably additive regular Borel measure μ on the Stone space $X = \text{Stone}(B)$.

1. **Integration on Simple Functions:** Any $g \in M$ can be written as a linear combination of disjoint indicator functions $g = \sum_{i=1}^n c_i \mathbf{1}_{\hat{b}_i}$ where $\{\hat{b}_i\}$ forms a partition of unity in B . Setting $\varphi_0(g) = \sum_{i=1}^n c_i \mu_0(b_i)$ gives a well-defined positive linear functional on M with $\varphi_0(\mathbf{1}_{\hat{b}}) = \mu_0(b)$ and $\varphi_0(\mathbf{1}_X) = \mu_0(1_B) = 1$.
2. **Dominated Extension:** By part (b), φ_0 extends to a positive linear functional $\varphi \in C(X)^*$.
3. **Riesz–Markov–Kakutani Representation:** X is a compact Hausdorff space. By the Riesz Representation Theorem, for every positive linear functional φ on $C(X)$, there exists a unique regular Borel measure μ on the Borel σ -algebra $\mathcal{B}(X)$ such that:

$$\varphi(f) = \int_X f d\mu \quad \forall f \in C(X)$$

4. **Restriction to Clopen Sets:** For any $b \in B$, the set $\hat{b}$ is clopen (hence Borel), and:

$$\mu(\hat{b}) = \int_X \mathbf{1}_{\hat{b}} d\mu = \varphi(\mathbf{1}_{\hat{b}}) = \varphi_0(\mathbf{1}_{\hat{b}}) = \mu_0(b)$$

This completes the proof of Scott's Theorem 4.1 by producing the countably additive regular Borel measure μ on X extending μ_0 .

Concrete Illustrative Example

Let $B = \mathcal{P}(\{1, 2\})$, so $X = \{1, 2\}$ with the discrete topology. $C(X, \mathbb{R}) \cong \mathbb{R}^2$ with functions represented by $(f(1), f(2))$.

Let $M = \mathbb{R} \cdot (1, 1) \subset \mathbb{R}^2$, and define $\varphi_0 : M \rightarrow \mathbb{R}$ by $\varphi_0(c, c) = c$. Here $\mathbf{1}_X = (1, 1)$ and $\varphi_0(\mathbf{1}_X) = 1$.

For $f = (x, y) \in \mathbb{R}^2$: - $\|f\|_\infty = \max(|x|, |y|)$. - $p(f) = 1 \cdot \max(|x|, |y|)$. - For $(c, c) \in M$, $\varphi_0(c, c) = c \leq |c| = p(c, c)$. - One valid Hahn–Banach extension is $\varphi(x, y) = \frac{1}{3}x + \frac{2}{3}y$. - Extension property: $\varphi(c, c) = \frac{1}{3}c + \frac{2}{3}c = c = \varphi_0(c, c)$. - Dominance: $\frac{1}{3}x + \frac{2}{3}y \leq \frac{1}{3}\max(|x|, |y|) + \frac{2}{3}\max(|x|, |y|) = \max(|x|, |y|) = p(x, y)$. - Positivity: $x \geq 0, y \geq 0 \implies \frac{1}{3}x + \frac{2}{3}y \geq 0$.

Self-Contained Lean 4 Verification

Lean 4 formalization: `appendix module BSinMeasurementTheory/Course21640.lean.`

8 Phase 6: Assembly and consistency check

The infinite extension of Theorem 4.1, and the finite/infinite agreement check.

8.1 21-410/21-599 — Independent Study

Reconstruct the full extension of Scott 1964 Measurement Theory Theorem 4.1 to infinite Boolean algebras end to end:

- Starting only from Kraft–Pratt–Seidenberg (1959) and Kelley (1959), write out the complete proof — Stone space construction, positive functional, Hahn–Banach extension, Riesz representation, restriction back to B — as a single connected argument, citing your own solutions to the 21-329, 21-651, 21-720, and 21-640 problems above at each step rather than re-deriving them.
- Identify precisely where finite additivity is upgraded to countable additivity (or note if it isn't, and Scott's "measure" is only finitely additive) — this is the one gap most expositions of Kelley's method gloss over.
- Formalize the finite-dimensional theorems (1.1–4.1 as stated, no extension) completely in Lean 4, then formalize as much of (a) as is tractable given current Mathlib coverage of Stone duality — and document precisely which lemma, if any, blocks full formalization.

Requires: everything above, assembled — and the honesty to report a genuine obstruction rather than paper over it.

Prove it in English and then in Lean 4, using as much Mathlib as possible. No uses of sorry, admit or axiom are allowed. If you do any numerical calculations for examples to illustrate the proof, reproduce them after the proof in English and then Lean. Every time you give a concrete example in English you must follow it with the equivalent Lean code. We are using Lean v4.34. Self contained not allowed please refer to <https://leanprover-community.github.io/mathlib-overview.html>

Put all the Lean code at the end in a single block of proofs and examples.

Leave no holes: If there is a “missing glue lemma”, then state it and prove it.

(a) Extension of Scott’s Theorem 4.1 to Infinite Boolean Algebras

1. Setup and Scott Cancellation Condition

Let $B = \langle B, \vee, \wedge, \neg, \perp, \top \rangle$ be an arbitrary (possibly infinite) Boolean algebra. A binary relation $\lesssim$ on B represents qualitative probability (with strict relation $a \prec b \iff a \lesssim b \wedge \neg(b \lesssim a)$).

Following Kraft–Pratt–Seidenberg (1959) and Scott (1964), $\lesssim$ satisfies the **finite cancellation condition** if for every integer $n \geq 1$ and all tuples $(a_1, \dots, a_n), (b_1, \dots, b_n) \in B^n$:

$$\sum_{i=1}^n \mathbf{1}_{a_i} = \sum_{i=1}^n \mathbf{1}_{b_i} \implies \left(\bigwedge_{i=1}^{n-1} a_i \lesssim b_i \implies b_n \lesssim a_n \right),$$

subject to non-triviality: $\perp \prec \top$ and $\perp \lesssim a$ for all $a \in B$.

2. Stone Space Construction

By Stone’s Representation Theorem, let:

$$X = \text{Stone}(B) = \{\mathcal{U} \subseteq B \mid \mathcal{U} \text{ is an ultrafilter on } B\}.$$

Endow X with the topology generated by basic clopen sets $\widehat{a} = \{\mathcal{U} \in X \mid a \in \mathcal{U}\}$ for each $a \in B$. - X is a compact, Hausdorff, totally disconnected space. - The map $a \mapsto \widehat{a}$ is a Boolean isomorphism:

$$\widehat{a \vee b} = \widehat{a} \cup \widehat{b}, \quad \widehat{a \wedge b} = \widehat{a} \cap \widehat{b}, \quad \widehat{\neg a} = X \setminus \widehat{a}, \quad \widehat{\perp} = \emptyset, \quad \widehat{\top} = X.$$

- For every $a \in B$, its indicator function $\mathbf{1}_{\widehat{a}} : X \rightarrow \mathbb{R}$ is continuous: $\mathbf{1}_{\widehat{a}} \in C(X, \mathbb{R})$.

3. Vector Space of Step Functions and the Preference Cone

Let $V \subseteq C(X, \mathbb{R})$ be the linear subspace of step functions generated by B :

$$V = \text{span}_{\mathbb{R}} \{\mathbf{1}_{\widehat{a}} \mid a \in B\}.$$

Define two convex cones in V : 1. The pointwise non-negative cone:

$$P = \{f \in V \mid \forall x \in X, f(x) \geq 0\}.$$

2. The qualitative preference cone:

$$K = \text{cone}(\{\mathbf{1}_{\widehat{b}} - \mathbf{1}_{\widehat{a}} \mid a \lesssim b\} \cup P).$$

By the Kraft–Pratt–Seidenberg / Scott finite cancellation condition, no strictly negative function can be formed as a conical combination of preference differences:

$$K \cap (-P^\circ) = \emptyset, \quad \text{where } -P^\circ = \{f \in V \mid \forall x \in X, f(x) < 0\}.$$

4. Hahn–Banach / M. Riesz Positive Extension

Equip $C(X, \mathbb{R})$ with the supremum norm $\|f\|_\infty = \sup_{x \in X} |f(x)|$, and define the sublinear functional $p(f) = \sup_{x \in X} f(x)$.

On the one-dimensional subspace $\mathbb{R}\mathbf{1}_X$, define $L_0(c\mathbf{1}_X) = c$. Because K is disjoint from the open cone $\{f \in C(X, \mathbb{R}) \mid p(f) < 0\}$, the **Hahn–Banach Separation Theorem** (M. Riesz / Kantorovich extension) extends L_0 to a positive linear functional:

$$\Lambda : C(X, \mathbb{R}) \rightarrow \mathbb{R}$$

satisfying: 1. $\Lambda(\mathbf{1}_X) = 1$, 2. $\Lambda(f) \geq 0$ for all $f \geq 0$, 3. $\Lambda(\mathbf{1}_{\hat{b}} - \mathbf{1}_{\hat{a}}) \geq 0$ whenever $a \precsim b$.

5. Restriction back to B

Define the measure $\mu : B \rightarrow [0, 1]$ by:

$$\mu(a) := \Lambda(\mathbf{1}_{\hat{a}}).$$

- **Normalization:** $\mu(\top) = \Lambda(\mathbf{1}_X) = 1$, $\mu(\perp) = \Lambda(\mathbf{0}) = 0$. - **Finite Additivity:** If $a \wedge b = \perp$, then $\hat{a} \cap \hat{b} = \emptyset$, so $\mathbf{1}_{\widehat{a \vee b}} = \mathbf{1}_{\hat{a}} + \mathbf{1}_{\hat{b}}$ pointwise. By linearity of Λ :

$$\mu(a \vee b) = \Lambda(\mathbf{1}_{\widehat{a \vee b}}) = \Lambda(\mathbf{1}_{\hat{a}}) + \Lambda(\mathbf{1}_{\hat{b}}) = \mu(a) + \mu(b).$$

- **Order Agreement:** If $a \precsim b$, then $\mu(b) - \mu(a) = \Lambda(\mathbf{1}_{\hat{b}} - \mathbf{1}_{\hat{a}}) \geq 0 \implies \mu(a) \leq \mu(b)$.

(b) Finite Additivity vs. Countable Additivity

Theorem (Additivity Classification): Scott's representation theorem yields an exact **finitely additive** probability measure on B . It does **not** upgrade to countable (σ -)additivity in general.

The Exact Point of the Obstruction

1. **The Stone Space σ -Algebra Gap:** By the Riesz–Markov–Kakutani Theorem, Λ corresponds to a unique regular Borel probability measure $\tilde{\mu}$ on the Borel σ -algebra $\mathcal{B}(X)$ of $X = \text{Stone}(B)$. However, B embeds only onto $\text{Clopen}(X)$, which is an algebra, not a σ -algebra.
2. **Topological Boundary of Countable Supremums:** For an infinite pairwise disjoint sequence $(a_n)_{n=1}^\infty$ in a complete Boolean algebra B with supremum $a = \bigvee_{n=1}^\infty a_n$:

$$\bigcup_{n=1}^\infty \hat{a}_n \subsetneq \hat{a} = \overline{\bigcup_{n=1}^\infty \hat{a}_n}.$$

The boundary set $N = \hat{a} \setminus \bigcup_{n=1}^\infty \hat{a}_n$ is a closed, nowhere-dense subset of X .

3. **Countable Additivity Criterion:** Countable additivity $\mu(a) = \sum_{n=1}^\infty \mu(a_n)$ holds if and only if $\tilde{\mu}(N) = 0$ for every nowhere-dense set $N \subseteq X$.

4. **Kelley’s Theorem (1959):** A complete Boolean algebra carries a strictly positive, countably additive measure if and only if B satisfies the **Countable Chain Condition (CCC)** and **weak (σ, ∞) -distributivity** (strictly positive Kelley intersection numbers $\kappa(B \setminus \{0\}) > 0$). Scott’s condition controls only finite cancellation, yielding finite additivity.

Concrete Worked Example

English Calculation

Consider the four-atom Boolean algebra $B = \mathcal{P}(\{0, 1, 2, 3\})$ with state weights:

$$w = (w(0), w(1), w(2), w(3)) = (0.1, 0.2, 0.3, 0.4).$$

Let $A = \{0, 1\}$ and $B = \{1, 2, 3\}$. 1. **Normalization:** $\sum_{x=0}^3 w(x) = 0.1 + 0.2 + 0.3 + 0.4 = 1.0$. 2. **Evaluations:**

$$\mu(A) = w(0) + w(1) = 0.1 + 0.2 = 0.3,$$

$$\mu(B) = w(1) + w(2) + w(3) = 0.2 + 0.3 + 0.4 = 0.9.$$

3. Intersection & Union:

$$A \cap B = \{1\} \implies \mu(A \cap B) = w(1) = 0.2,$$

$$A \cup B = \{0, 1, 2, 3\} \implies \mu(A \cup B) = 1.0.$$

4. Inclusion-Exclusion:

$$\mu(A \cup B) + \mu(A \cap B) = 1.0 + 0.2 = 1.2 = 0.3 + 0.9 = \mu(A) + \mu(B).$$

(c) Complete Lean 4 Formalization

Lean 4 formalization: appendix module `BSinMeasurementTheory/Course21410.lean`.

8.2 21-322 — Topics in Formal Mathematics (opportunistic)

Whichever term you take this, it should overlap real analysis or order theory in Lean. As a standing problem regardless of that term’s specific topic:

Formalize, in Lean 4, the statement and proof that a finite Boolean algebra’s atom structure (21-373, part a–b above) is preserved under the Stone-space construction in the degenerate finite case — i.e., prove formally that when B is finite, the general infinite-case machinery from Phase 4–6 reduces exactly to the elementary atoms/characteristic-functions picture from Phase 2, with no discrepancy.

Requires: connecting the “easy” finite self-study material to the full machinery, formally — the check that Phases 2 and 4–6 actually agree at the point where they overlap.

Prove it in English and then in Lean 4, using as much Mathlib as possible. No uses of sorry, admit or axiom are allowed. If you do any numerical calculations for examples to illustrate the proof, reproduce them after the proof in English and then Lean. Every time you give a concrete example in English you must follow it with the equivalent Lean code. We are using Lean v4.34. Self contained not allowed please refer to <https://leanprover-community.github.io/mathlib-overview.html>

Mathematical Exposition in English

1. The Finite-to-Infinite Duality Gap and the Reduction Problem

In Phases 4–6, Dana Scott’s measurement theorem is extended to general Boolean algebras by embedding B into the continuous function space $C(X, \mathbb{R})$ over its Stone spectrum:

$$X = \text{Stone}(B) = \text{Hom}_{\mathbf{BA}}(B, \mathbf{2}),$$

extending the positive functional φ_0 from the simple-function subspace $M = \text{span}\{\mathbf{1}_{\hat{b}} \mid b \in B\}$ to all of $C(X, \mathbb{R})$ via the Hahn–Banach theorem, and representing it as an integral $\Lambda(f) = \int_X f d\mu$ by the Riesz–Markov–Kakutani theorem.

In Phase 2 (21-373), a finite Boolean algebra B is shown to be purely atomic: $B \cong \mathcal{P}(A) \cong \{0, 1\}^A$, where $A = \text{Atoms}(B)$.

The purpose of this theorem is to prove that the two constructions **coincide with zero discrepancy** when B is finite: 1. Every Stone homomorphism $u \in \text{Stone}(B)$ is the principal evaluation at a unique atom $a \in A$. 2. The Stone topological space X is discrete, so every subset is clopen: $\text{Clopen}(X) = \mathcal{P}(X) \cong \mathcal{P}(A)$. 3. Under this identification, the Stone clopen representation $\sigma(b) = \hat{b}$ is identical to the atomic support representation $\phi(b) = \{a \in A \mid a \leq b\}$. 4. The positive linear functional integral $\Lambda(\mathbf{1}_{\hat{b}})$ collapses directly to the finite atom-weight sum $\sum_{a \leq b} w(a)$.

2. Concrete Example in English

Let $\alpha = \{0, 1, 2\}$ with $B = \mathcal{P}(\alpha)$, which has 3 atoms $\{0\}, \{1\}, \{2\}$. 1. **Atoms and Stone Homomorphisms:** The Stone space consists of 3 points $X = \{u_0, u_1, u_2\}$ where $u_i(s) = \text{true} \iff i \in s$. 2. **Clopen Sets:** For $s = \{0, 2\}$, the Stone image is:

$$\hat{s} = \{u_0, u_2\} \subseteq X.$$

Under $u_i \leftrightarrow i$, $\hat{s}$ is identified with $\{0, 2\}$. 3. **Measure Collapse:** Let the weights assigned to the atoms be $w = (0.2, 0.5, 0.3)$. - The Stone measure evaluation is:

$$\Lambda(\mathbf{1}_{\hat{s}}) = \sum_{a \in \alpha} w(a) \cdot \mathbf{1}_{\hat{s}}(u_a) = 0.2(1) + 0.5(0) + 0.3(1) = 0.5.$$

- The discrete atomic sum is:

$$\sum_{a \in s} w(a) = w(0) + w(2) = 0.2 + 0.3 = 0.5.$$

Both values match at 0.5.

Complete Lean 4 Formalization

The code below compiles cleanly in Lean v4.34 with `import Mathlib`. It contains no `sorry`, `admit`, or extra axioms.

Lean 4 formalization: `appendix module BSinMeasurementTheory/Course21322.lean`.

9 References

[Sco64] Dana S. Scott, *Measurement Structures and Linear Inequalities*, Journal of Mathematical Psychology 1 (1964), 233–247.

[KP59] L. G. Kraft, J. D. Pratt, and A. Seidenberg, *Intuitive Probability on Finite Sets*, Annals of Mathematical Statistics 30 (1959), 408–419.

[Kel59] J. L. Kelley, *Measures on Boolean Algebras*, Pacific Journal of Mathematics 9 (1959), 1165–1177.

A Lean module index

Checked by lake build. Complete sources: <https://github.com/catskillsresearch/fabbro>. Each subsection is the corresponding library file.

Role	File
Root importer	<code>BSinMeasurementTheory.lean</code>
21-120-21-122	<code>BSinMeasurementTheory/ Course21120FTC.lean</code>
21-120-21-122	<code>BSinMeasurementTheory/ Course21120Taylor.lean</code>
21-127	<code>BSinMeasurementTheory/Course21127. lean</code>
21-241	<code>BSinMeasurementTheory/Course21241. lean</code>
21-241	<code>BSinMeasurementTheory/ Course21241Numerical.lean</code>
21-228	<code>BSinMeasurementTheory/Course21228. lean</code>
21-355	<code>BSinMeasurementTheory/Course21355. lean</code>
21-373	<code>BSinMeasurementTheory/Course21373. lean</code>
21-373	<code>BSinMeasurementTheory/ Course21373Numerical.lean</code>
21-321	<code>BSinMeasurementTheory/Course21321. lean</code>
21-321	<code>BSinMeasurementTheory/ Course21321Kraft.lean</code>
21-292	<code>BSinMeasurementTheory/Course21292. lean</code>
21-292	<code>BSinMeasurementTheory/ Course21292Numerical.lean</code>
21-329	<code>BSinMeasurementTheory/Course21329. lean</code>

Role	File
21-651	BSinMeasurementTheory/Course21651.lean
21-720	BSinMeasurementTheory/Course21720.lean
21-640	BSinMeasurementTheory/Course21640.lean
21-410-21-599	BSinMeasurementTheory/Course21410.lean
21-322	BSinMeasurementTheory/Course21322.lean

A.1 BSinMeasurementTheory.lean

Lean 4 source

```
import BSinMeasurementTheory.Course21120FTC
import BSinMeasurementTheory.Course21120Taylor
import BSinMeasurementTheory.Course21127
import BSinMeasurementTheory.Course21228
import BSinMeasurementTheory.Course21241
import BSinMeasurementTheory.Course21241Numerical
import BSinMeasurementTheory.Course21292
import BSinMeasurementTheory.Course21292Numerical
import BSinMeasurementTheory.Course21321
import BSinMeasurementTheory.Course21321Kraft
import BSinMeasurementTheory.Course21322
import BSinMeasurementTheory.Course21329
import BSinMeasurementTheory.Course21355
import BSinMeasurementTheory.Course21373
import BSinMeasurementTheory.Course21373Numerical
import BSinMeasurementTheory.Course21410
import BSinMeasurementTheory.Course21640
import BSinMeasurementTheory.Course21651
import BSinMeasurementTheory.Course21720
```

A.2 BSinMeasurementTheory/Course21120FTC.lean

Lean 4 source

```
import Mathlib

open Real intervalIntegral MeasureTheory

namespace Course21120FTC

/-- Definition of  $f(x) = \int_0^x \frac{\sin(t)}{(1+t)^2} dt$  -/
noncomputable def f (x : ℝ) : ℝ := ∫ t in (0 : ℝ)..x, sin (t ^ 2) / (1 + t ^ 2)
```

```

/-- The integrand is continuous everywhere on ?. -/
lemma continuous_integrand : Continuous (fun t : ? => sin (t ^ 2) / (1 + t ^ 2)) := by
  apply Continuous.div
  * exact continuous_sin.comp (continuous_pow 2)
  * exact continuous_const.add (continuous_pow 2)
  * intro t
    have h : 0 <= t ^ 2 := sq_nonneg t
    linarith

/-- Fundamental theorem of calculus derivative statement for f. -/
lemma f_hasDerivAt (x : ?) : HasDerivAt f (sin (x ^ 2) / (1 + x ^ 2)) x :=
  HasStrictDerivAt.hasDerivAt (continuous_integrand.integral_hasStrictDerivAt 0 x)

/-- (a) Part 1: f is differentiable everywhere on ?. -/
theorem part_a_differentiable (x : ?) : DifferentiableAt ? f x :=
  HasDerivAt.differentiableAt (f_hasDerivAt x)

/-- (a) Part 2: f'(x) = sin(x?) / (1 + x?). -/
theorem part_a_deriv (x : ?) : deriv f x = sin (x ^ 2) / (1 + x ^ 2) :=
  HasDerivAt.deriv (f_hasDerivAt x)

end Course21120FTC

```

A.3 BSinMeasurementTheory/Course21120Taylor.lean

Lean 4 source

```

import Mathlib

namespace Course21120Taylor

/-!
#### Taylor Series Estimation of f(0.5)

Integrand Taylor expansion around t = 0:
g(t) = sin(t?) / (1 + t?) = t? - t? + (5/6) t? - (5/6) t? + O(t1?)

Integrating term-by-term from 0 to x:
T?(x) = x?/3 - x?/5 + (5/42) x? - (5/54) x?
-/

/-- Degree-9 Taylor polynomial approximation for f(x) -/
def T9 (x : ?) : ? :=
  x ^ 3 / 3 - x ^ 5 / 5 + 5 * x ^ 7 / 42 - 5 * x ^ 9 / 54

/-! ##### Step 1: Verification of individual terms at x = 1/2 -/

theorem term1_eq : (1 / 2 : ?) ^ 3 / 3 = 1 / 24 := by norm_num
theorem term2_eq : (1 / 2 : ?) ^ 5 / 5 = 1 / 160 := by norm_num
theorem term3_eq : 5 * (1 / 2 : ?) ^ 7 / 42 = 5 / 5376 := by norm_num
theorem term4_eq : 5 * (1 / 2 : ?) ^ 9 / 54 = 5 / 27648 := by norm_num

/-! ##### Step 2: Exact sum of the terms -/

```

```

/-- The exact rational value of  $T(1/2)$  is  $34997 / 967680$  -/
theorem T9_half_eq : T9 (1 / 2) = 34997 / 967680 := by
  dsimp [T9]
  norm_num

/-! ##### Step 3: Four-decimal-place accuracy bounds -/

/--  $T(1/2)$  lies strictly between 0.03615 and 0.03625,
    so it rounds to 0.0362 to 4 decimal places. -/
theorem T9_half_rounds_to_0_0362 :
  (3615 : ?) / 100000 < T9 (1 / 2) /\ T9 (1 / 2) < 3625 / 100000 := by
  rw [T9_half_eq]
  norm_num

/-- Explicit distance to 0.0362 is less than 0.00005 (half of  $10^{-?}$ ) -/
theorem T9_half_close_to_0_0362 :
  T9 (1 / 2) - 362 / 10000 < 5 / 100000 /\
  362 / 10000 - T9 (1 / 2) < 5 / 100000 := by
  rw [T9_half_eq]
  norm_num

/-! ##### Step 4: Verification of the Truncation Error Bound -/

/-- The next non-zero term in the alternating series bounds the tail error:
     $|R(0.5)| \leq (101 / 1320) * (1/2)^{11}$  -/
def truncation_error : ? := (101 / 1320) * (1 / 2) ^ 11

theorem truncation_error_eq : truncation_error = 101 / 2703360 := by
  dsimp [truncation_error]
  norm_num

/-- The truncation error is strictly less than  $10^{-?}$  (0.0001) -/
theorem truncation_error_lt : truncation_error < 1 / 10000 := by
  rw [truncation_error_eq]
  norm_num

/-! ##### Step 5: Evaluations (Inspect in Lean InfoView) -/

-- Float value of each term:
#eval (1.0 / 24.0 : Float)           -- 0.041666666666666664
#eval (1.0 / 160.0 : Float)          -- 0.00625
#eval (5.0 / 5376.0 : Float)         -- 0.000929985119047619
#eval (5.0 / 27648.0 : Float)        -- 0.0001808449074074074

-- Float value of  $T(0.5)$ :
#eval (34997.0 / 967680.0 : Float)   -- 0.03616588128306879

-- Float value of the truncation error:
#eval (101.0 / 2703360.0 : Float)     -- 0.000037360913825757576

end Course21120Taylor

```

A.4 BSinMeasurementTheory/Course21127.lean

Lean 4 source

```
import Mathlib

namespace Course21127

/-!
# Part (a): Equivalence Relation on  $\text{Int} \times (\text{Int} \setminus \{0\})$ 
-/

/-- The domain  $S = \text{Int} \times (\text{Int} \setminus \{0\})$  -/
def S : Type := Int × { z : Int // z != 0 }

/-- The relation  $(a, b) \sim (c, d) \iff a * d = b * c$  -/
def rel (x y : S) : Prop :=
  x.1 * y.2.val = x.2.val * y.1

lemma rel_refl (x : S) : rel x x := by
  dsimp [rel]
  ring

lemma rel_symm {x y : S} (h : rel x y) : rel y x := by
  dsimp [rel] at *
  calc
    y.1 * x.2.val = x.2.val * y.1 := by ring
    _ = x.1 * y.2.val := h.symm
    _ = y.2.val * x.1 := by ring

lemma rel_trans {x y z : S} (h1 : rel x y) (h2 : rel y z) : rel x z := by
  dsimp [rel] at *
  have hd : y.2.val != 0 := y.2.property
  have H : (x.1 * z.2.val) * y.2.val = (x.2.val * z.1) * y.2.val := by
    calc
      (x.1 * z.2.val) * y.2.val = (x.1 * y.2.val) * z.2.val := by ring
      _ = (x.2.val * y.1) * z.2.val := by rw [h1]
      _ = x.2.val * (y.1 * z.2.val) := by ring
      _ = x.2.val * (y.2.val * z.1) := by rw [h2]
      _ = (x.2.val * z.1) * y.2.val := by ring
  have H2 : (x.1 * z.2.val - x.2.val * z.1) * y.2.val = 0 := by
    calc (x.1 * z.2.val - x.2.val * z.1) * y.2.val
      _ = (x.1 * z.2.val) * y.2.val - (x.2.val * z.1) * y.2.val := by ring
      _ = (x.2.val * z.1) * y.2.val - (x.2.val * z.1) * y.2.val := by rw [H]
      _ = 0 := by ring
  cases mul_eq_zero.mp H2 with
  | inl h => exact sub_eq_zero.mp h
  | inr h => exact False.elim (hd h)

/-- (a) `rel` is an equivalence relation on  $S$ . -/
def setoid : Setoid S where
  r := rel
  iseqv := <rel_refl, rel_symm, rel_trans>

instance : Setoid S := setoid
```

```

/-!
# Part (b): Bijection between Quotient and ?
-/

lemma rat_div_eq_of_mul_eq {a b c d : ?} (hb : b != 0) (hd : d != 0) (h : a * d =
  b * c) :
  a / b = c / d := by
  rw [div_eq_div_iff hb hd]
  linarith

lemma rat_mul_eq_of_div_eq {a b c d : ?} (hb : b != 0) (hd : d != 0) (h : a / b =
  c / d) :
  a * d = b * c := by
  have := (div_eq_div_iff hb hd).mp h
  linarith

/-- The explicit map from a pair (a, b) to ?. -/
def toRat (p : S) : ? := (p.1 : ?) / (p.2.val : ?)

/-- The map respects the equivalence relation (well-defined). -/
lemma toRat_wellDefined (x y : S) (h : rel x y) : toRat x = toRat y := by
  dsimp [toRat]
  have hx : ((x.2.val : ?) != 0) := by
    intro hz
    exact x.2.property (by exact_mod_cast hz)
  have hy : ((y.2.val : ?) != 0) := by
    intro hz
    exact y.2.property (by exact_mod_cast hz)
  have h_int : x.1 * y.2.val = x.2.val * y.1 := h
  have h_cast : (x.1 : ?) * (y.2.val : ?) = (x.2.val : ?) * (y.1 : ?) := by
    exact_mod_cast h_int
  exact rat_div_eq_of_mul_eq hx hy h_cast

/-- The induced map on the quotient set. -/
def toRatQuot : Quotient setoid -> ? :=
  Quotient.lift toRat toRat_wellDefined

/-- (b) Part 1: Injectivity -/
theorem toRatQuot_injective : Function.Injective toRatQuot := by
  intro q1 q2
  refine Quotient.inductionOn_2 q1 q2 ?_
  intro x y h
  dsimp [toRatQuot, toRat] at h
  have hx : ((x.2.val : ?) != 0) := by
    intro hz
    exact x.2.property (by exact_mod_cast hz)
  have hy : ((y.2.val : ?) != 0) := by
    intro hz
    exact y.2.property (by exact_mod_cast hz)
  have h_mul := rat_mul_eq_of_div_eq hx hy h
  have h_int : x.1 * y.2.val = x.2.val * y.1 := by
    exact_mod_cast h_mul
  apply Quotient.sound
  exact h_int

/-- (b) Part 2: Surjectivity -/

```

```

theorem toRatQuot_surjective : Function.Surjective toRatQuot := by
  intro q
  have hden : (q.den : Int) != 0 := by
    intro hz
    have : q.den = 0 := by exact_mod_cast hz
    exact q.den_nz this
  use ?(q.num, <(q.den : Int), hden>)?
  change toRat (q.num, <(q.den : Int), hden>) = q
  dsimp [toRat]
  have : ((q.den : Int) : ?) = (q.den : ?) := by push_cast; rfl
  rw [this]
  exact Rat.num_div_den q

/-- (b) Part 3: Bijectivity -/
theorem toRatQuot_bijective : Function.Bijective toRatQuot :=
  <toRatQuot_injective, toRatQuot_surjective>

/-- The induced quotient set is in bijection with ?. -/
noncomputable def quotientEquivRat : Quotient setoid ? ? :=
  Equiv.ofBijective toRatQuot toRatQuot_bijective

/-!
# Part (c): Refutation of the False Claim and the Correct Theorem
-/

/-- A subset S subseteq ? is well-ordered under the usual order if every nonempty
subset has a least element in the set. -/
def IsWellOrderedSet (S : Set ?) : Prop :=
  forall T : Set ?, T subseteq S -> T.Nonempty -> exists m in T, forall x
    in T, m <= x

/-- The false claim stated in the prompt using Set.Ici to avoid untyped binder
problems. -/
def FalseClaim : Prop :=
  forall S : Set ?, S subseteq Set.Ici (0 : ?) -> S.Nonempty ->
    IsWellOrderedSet S ->
    (S = {} \/\ Set.Infinite S)

lemma singleton_zero_well_ordered : IsWellOrderedSet { (0 : ?) } := by
  intro T hT <t, ht>
  have ht0 : t = 0 := hT ht
  subst ht0
  refine <0, ht, ?_>
  intro x hx
  have hx0 : x = 0 := hT hx
  subst hx0
  exact le_rfl

lemma singleton_zero_nonneg : ({ (0 : ?) } : Set ?) subseteq Set.Ici 0 :=
  Set.singleton_subset_iff.mpr (Set.mem_Ici.mpr le_rfl)

lemma singleton_zero_nonempty : ({ (0 : ?) } : Set ?).Nonempty :=
  <0, rfl>

/-- (c) Refutation by contradiction:
The claim implies that {0} must be empty or infinite, both of which are false. -/

```

```

theorem false_claim_is_false : not FalseClaim := by
  intro hclaim
  have h_cases := hclaim {0} singleton_zero_nonneg singleton_zero_nonempty
    singleton_zero_well_ordered
  cases h_cases with
  | inl h_empty =>
    have h0 : (0 : ?) in ({ (0 : ?) } : Set ?) := rfl
    rw [h_empty] at h0
    exact h0
  | inr h_inf =>
    exact h_inf (Set.finite_singleton 0)

/-- Correct positive statement proved by induction:
  Every nonempty finite list of rationals contains a least element. -/
lemma list_has_min : forall (a : ?) (l : List ?), exists m in a :: l, forall x
  in a :: l, m <= x
| a, [] => by
  refine <a, List.mem_cons.mpr (Or.inl rfl), ?_>
  intro x hx
  rcases List.mem_cons.mp hx with hxa | hx_nil
  * rw [hxa]
  * cases hx_nil
| a, b :: l => by
  rcases list_has_min (min a b) l with <m, hm, hle>
  have hm_orig : m in a :: b :: l := by
    rcases List.mem_cons.mp hm with hm_eq | h_in
    * rw [hm_eq]
    cases le_total a b with
    | inl hab =>
      rw [min_eq_left hab]
      exact List.mem_cons.mpr (Or.inl rfl)
    | inr hba =>
      rw [min_eq_right hba]
      exact List.mem_cons.mpr (Or.inr (List.mem_cons.mpr (Or.inl rfl)))
  * exact List.mem_cons.mpr (Or.inr (List.mem_cons.mpr (Or.inr h_in)))
  refine <m, hm_orig, ?_>
  intro x hx
  rcases List.mem_cons.mp hx with hxa | hx1
  * rw [hxa]
  have h_min_in : min a b in (min a b) :: l := List.mem_cons.mpr (Or.inl rfl)
  exact le_trans (hle (min a b) h_min_in) (min_le_left a b)
  * rcases List.mem_cons.mp hx1 with hxb | hx2
  * rw [hxb]
  have h_min_in : min a b in (min a b) :: l := List.mem_cons.mpr (Or.inl rfl)
  exact le_trans (hle (min a b) h_min_in) (min_le_right a b)
  * have hx_in : x in (min a b) :: l := List.mem_cons.mpr (Or.inr hx2)
    exact hle x hx_in

/-!
# Numerical Examples Reproducing Calculations
-/

-- Example 1: (1, 2) ~ (2, 4) since 1 * 4 = 2 * 2 = 4
def pair1 : S := (1, <2, by decide>)
def pair2 : S := (2, <4, by decide>)

```

```

example : rel pair1 pair2 := by
  dsimp [rel, pair1, pair2]

-- Example 2: Both pairs map to 1/2 in ? under toRat
example : toRat pair1 = 1 / 2 := by
  norm_num [toRat, pair1]

example : toRat pair2 = 1 / 2 := by
  norm_num [toRat, pair2]

-- Example 3: Minimal element in the set {1/2, 1/3} is 1/3
example : min (1 / 2 : ?) (1 / 3) = 1 / 3 := by
  norm_num

end Course21127

```

A.5 BSinMeasurementTheory/Course21241.lean

Lean 4 source

```

import Mathlib

open BigOperators
open Finset

namespace Course21241

/-!
# Convex Cones, Dual Bases, and Separation

This file formalizes:
(a) Explicit basis and dual basis constructions for  $V = \text{Fin } 3 \rightarrow \mathbb{R}$ .
(b) Riesz representation under the standard inner product and separation of  $v \notin C$ .
(c) The dual cone  $C^*$  is a convex cone, identified in the dual basis coordinates.
-/

abbrev V := Fin 3 → ℝ

/-- The non-negative orthant cone C in  $V = \mathbb{R}^3$  -/
def C : Set V := {x | forall i : Fin 3, 0 ≤ x i}

/-! ### Part (a): Basis and Dual Basis Construction -/

/-- Standard primal basis vectors of  $V$  -/
def e (i : Fin 3) : V := fun j => if i = j then 1 else 0

/-- Explicit dual basis functionals  $e_{\text{star } i}$  in  $V^*$  -/
def e_star (i : Fin 3) : V → ℝ where
  toFun x := x i
  map_add' _ _ := rfl
  map_smul' _ _ := rfl

```

```

/-- Duality relation between primal and dual bases:  $e_{\text{star } i} (e_j) = \delta_{ij}$  -/
theorem e_star_apply_e (i j : Fin 3) :
  e_star i (e j) = if i = j then (1 : ?) else 0 := by
  dsimp [e_star, e]
  fin_cases i <|> fin_cases j <|> rfl

/-- Every vector  $x$  in  $V$  expands uniquely in the primal basis -/
theorem vector_eq_sum (x : V) :
  x = ? i : Fin 3, x i * e i := by
  ext j
  simp only [Finset.sum_apply, Pi.smul_apply, smul_eq_mul]
  rw [Fin.sum_univ_three]
  fin_cases j <|> { dsimp [e]; ring }

/-- The basis vectors  $\{e_0, e_1, e_2\}$  are linearly independent -/
theorem e_linearIndependent :
  LinearIndependent ? e := by
  rw [linearIndependent_iff']
  intro s g hg i hi
  have h := congr_fun hg i
  simp only [Finset.sum_apply, Pi.smul_apply, smul_eq_mul, e, Pi.zero_apply] at h
  rw [Finset.sum_eq_single_of_mem i hi] at h
  * simp only [ite_true, mul_one] at h
  exact h
  * intro j _ hj
  simp only [hj, ite_false, mul_zero]

/-- Every linear functional  $\phi$  in  $V^*$  expands uniquely in the dual basis -/
theorem dual_eq_sum (phi : V → ?[?] ?) :
  phi = ? i : Fin 3, phi (e i) * e_star i := by
  apply LinearMap.ext
  intro x
  have hx : x = ? i : Fin 3, x i * e i := vector_eq_sum x
  conv_lhs => rw [hx]
  rw [map_sum]
  simp only [LinearMap.sum_apply, LinearMap.smul_apply, smul_eq_mul]
  dsimp [e_star]
  apply Finset.sum_congr rfl
  intro i _
  rw [LinearMap.map_smul, smul_eq_mul, mul_comm]

/-! ### Part (b): Inner Product and Separating Functional -/

/-- Standard Euclidean inner product on  $V$  -/
def dot (u v : V) : ? := ? i : Fin 3, u i * v i

lemma dot_comm (u v : V) : dot u v = dot v u := by
  dsimp [dot]
  apply Finset.sum_congr rfl
  intro i _
  ring

lemma dot_e (u : V) (i : Fin 3) : dot u (e i) = u i := by
  dsimp [dot]
  rw [Fin.sum_univ_three]
  fin_cases i <|> { dsimp [e]; ring }

```

```

/-- Riesz map from V to V* -/
def toDual (u : V) : V → ?[?] ? where
  toFun v := dot u v
  map_add' v w := by
    dsimp [dot]
    rw [← Finset.sum_add_distrib]
    apply Finset.sum_congr rfl
    intro i _
    ring
  map_smul' c v := by
    dsimp [dot]
    rw [Finset.mul_sum]
    apply Finset.sum_congr rfl
    intro i _
    ring

/-- Inverse Riesz map from V* to V -/
def toVec (phi : V → ?[?] ?) : V := fun i => phi (e i)

theorem toDual_toVec (phi : V → ?[?] ?) : toDual (toVec phi) = phi := by
  apply LinearMap.ext
  intro x
  have hx : x = ? i : Fin 3, x i * e i := vector_eq_sum x
  conv_rhs => rw [hx]
  rw [map_sum]
  change (? i : Fin 3, phi (e i) * x i) = ? i : Fin 3, phi (x i * e i)
  apply Finset.sum_congr rfl
  intro i _
  rw [LinearMap.map_smul, smul_eq_mul, mul_comm]

theorem toVec_toDual (u : V) : toVec (toDual u) = u := by
  ext i
  exact dot_e u i

/-- Characterization of functionals nonnegative on C -/
theorem nonneg_on_C_iff (u : V) :
  (forall c in C, 0 ≤ dot u c) <=> forall i : Fin 3, 0 ≤ u i := by
  constructor
  * intro h i
    have hei : e i in C := by
      intro j
      dsimp [e]
      split_ifs <=> linarith
    have h_dot := h (e i) hei
    rw [dot_e] at h_dot
    exact h_dot
  * intro hu c hc
    dsimp [dot]
    apply Finset.sum_nonneg
    intro i _
    exact mul_nonneg (hu i) (hc i)

/-- Existence of a separating vector under the inner product for any v notin C -/
theorem exists_separating_vector (v : V) (hv : v notin C) :
  exists u : V, (forall c in C, 0 ≤ dot u c) /\ dot u v < 0 := by

```

```

change not (forall i : Fin 3, 0 <= v i) at hv
push Not at hv
rcases hv with <i, hi>
use e i
constructor
* rw [nonneg_on_C_iff]
  intro j
  dsimp [e]
  split_ifs <;> linarith
* rw [dot_comm, dot_e]
  exact hi

/-! ### Numerical Separation Example -/

/-- A concrete point v = (-1, 1, 1) notin C -/
def v_example : V := fun i =>
  if i = 0 then -1 else 1

/-- Separating vector u = e 0 = (1, 0, 0) -/
def u_example : V := e 0

theorem v_example_not_mem_C : v_example notin C := by
  intro h
  have h0 := h 0
  dsimp [v_example] at h0
  linarith

theorem dot_u_example_v_example : dot u_example v_example = -1 := by
  dsimp [u_example, dot]
  rw [Fin.sum_univ_three]
  dsimp [e, v_example]
  ring

theorem u_example_separates_v_example : dot u_example v_example < 0 := by
  rw [dot_u_example_v_example]
  linarith

theorem u_example_nonneg_on_C (c : V) (hc : c in C) : 0 <= dot u_example c := by
  dsimp [u_example, dot]
  rw [Fin.sum_univ_three]
  dsimp [e]
  have hc0 := hc 0
  linarith

/-! ### Part (c): The Dual Cone C* and its Dual Basis Representation -/

/-- The dual cone C* in V* -/
def dualCone (C : Set V) : Set (V -> ?[?] ?) :=
  {phi | forall c in C, 0 <= phi c}

theorem dualCone_smul_mem {phi : V -> ?[?] ?} (hphi : phi in dualCone C) {a : ?} (
  ha : 0 <= a) :
  a * phi in dualCone C := by
  intro c hc_mem
  simp only [LinearMap.smul_apply, smul_eq_mul]
  exact mul_nonneg ha (hphi c hc_mem)

```

```

theorem dualCone_add_mem {phi psi : V -> ?[?] ?} (hphi : phi in dualCone C) (hpsi :
  psi in dualCone C) :
  phi + psi in dualCone C := by
  intro c hc_mem
  simp only [LinearMap.add_apply]
  exact add_nonneg (hphi c hc_mem) (hpsi c hc_mem)

theorem dualCone_conic_combination {phi psi : V -> ?[?] ?} (hphi : phi in dualCone
  C) (hpsi : psi in dualCone C)
  {a b : ?} (ha : 0 <= a) (hb : 0 <= b) :
  a * phi + b * psi in dualCone C := by
  intro c hc_mem
  simp only [LinearMap.add_apply, LinearMap.smul_apply, smul_eq_mul]
  exact add_nonneg (mul_nonneg ha (hphi c hc_mem)) (mul_nonneg hb (hpsi c hc_mem))

theorem mem_dualCone_iff (phi : V -> ?[?] ?) :
  phi in dualCone C <-> forall i : Fin 3, 0 <= phi (e i) := by
  constructor
  * intro hphi i
    have hei : e i in C := by
      intro j
      dsimp [e]
      split_ifs <=> linarith
    exact hphi (e i) hei
  * intro hphi c hc
    have hc_sum : c = ? i : Fin 3, c i * e i := vector_eq_sum c
    conv_rhs => rw [hc_sum]
    rw [map_sum]
    simp only [LinearMap.map_smul, smul_eq_mul]
    apply Finset.sum_nonneg
    intro i _
    exact mul_nonneg (hc i) (hphi i)

/-- Dual cone represented as non-negative linear combinations of {e_star i} -/
def dualCone_representation : Set (V -> ?[?] ?) :=
  {phi | exists a : Fin 3 -> ?, (forall i, 0 <= a i) /\ phi = ? i : Fin 3, a i
    * e_star i}

theorem dualCone_eq_representation :
  dualCone C = dualCone_representation := by
  ext phi
  constructor
  * intro hphi
    rw [mem_dualCone_iff] at hphi
    use fun i => phi (e i)
    refine <hphi, ?_>
    exact dual_eq_sum phi
  * rintro <a, ha, rfl>
    rw [mem_dualCone_iff]
    intro i
    simp only [LinearMap.sum_apply, LinearMap.smul_apply, smul_eq_mul]
    have h_sum : (? j : Fin 3, a j * e_star j (e i)) = a i := by
      rw [Fin.sum_univ_three]
      fin_cases i <=> simp [e_star_apply_e]
    rw [h_sum]

```

```

exact ha i

end Course21241

```

A.6 BSinMeasurementTheory/Course21241Numerical.lean

Lean 4 source

```

import Mathlib

open BigOperators
open Finset

namespace Course21241Numerical

abbrev V := Fin 3 -> ?

/-- The convex cone C = {x in ?? : x_1 >= 0, x_2 >= 0, x_3 >= 0} -/
def C : Set V := {x | forall i : Fin 3, 0 <= x i}

/-- Standard basis vector e_i -/
def e (i : Fin 3) : V := fun j => if i = j then 1 else 0

/-- Standard Euclidean inner product on V -/
def dot (u v : V) : ? := ? i : Fin 3, u i * v i

/-- Embedding of V into V* via the inner product -/
def toDual (u : V) : V -> ?[?] ? where
  toFun v := dot u v
  map_add' v w := by
    dsimp [dot]
    rw [← Finset.sum_add_distrib]
    apply Finset.sum_congr rfl
    intro i _
    ring
  map_smul' c v := by
    dsimp [dot]
    rw [Finset.mul_sum]
    apply Finset.sum_congr rfl
    intro i _
    ring

/-!
### Choice of point v notin C:
v = (-1, 1, 1) ? v_0 = -1 < 0 ? v notin C
-/

def v : V := fun i => if i = 0 then -1 else 1

lemma v_zero : v 0 = -1 := rfl
lemma v_one : v 1 = 1 := rfl
lemma v_two : v 2 = 1 := rfl

```

```

/-- Coordinates of v are (-1, 1, 1) -/
theorem v_coords : v 0 = -1 /\ v 1 = 1 /\ v 2 = 1 :=
  <v_zero, v_one, v_two>

/-- v_0 = -1 < 0 -/
theorem v_zero_neg : v 0 < 0 := by
  rw [v_zero]
  norm_num

/-- v notin C because v 0 < 0 -/
theorem v_not_mem_C : v notin C := by
  intro hc
  have h0 : 0 <= v 0 := hc 0
  rw [v_zero] at h0
  norm_num at h0

/-!
### Separating vector u in V:
u = e_0 = (1, 0, 0)
-/

def u : V := e 0

lemma u_zero : u 0 = 1 := rfl
lemma u_one : u 1 = 0 := rfl
lemma u_two : u 2 = 0 := rfl

/-- u is e_0 -/
theorem u_eq_e_zero : u = e 0 := rfl

/-- Coordinates of u are (1, 0, 0) -/
theorem u_coords : u 0 = 1 /\ u 1 = 0 /\ u 2 = 0 :=
  <u_zero, u_one, u_two>

/-!
### Inner product value at v:
<u, v> = ?_{i=0}^2 u_i v_i = (1)(-1) + (0)(1) + (0)(1) = -1 < 0
-/

theorem dot_u_v_sum :
  dot u v = ? i : Fin 3, u i * v i := rfl

/-- <u, v> = (1)(-1) + (0)(1) + (0)(1) -/
theorem dot_u_v_expansion :
  dot u v = (1 : ?) * (-1) + 0 * 1 + 0 * 1 := by
  dsimp [dot]
  rw [Fin.sum_univ_three]
  rw [u_zero, u_one, u_two, v_zero, v_one, v_two]

/-- <u, v> = -1 -/
theorem dot_u_v_eval : dot u v = -1 := by
  rw [dot_u_v_expansion]
  ring

/-- <u, v> < 0 -/
theorem dot_u_v_neg : dot u v < 0 := by

```

```

    rw [dot_u_v_eval]
    norm_num

/-!
### Inner product on arbitrary c in C:
    <u, c> = ?_{i=0}? u_i c_i = (1) c_0 + (0) c_1 + (0) c_2 = c_0 >= 0
-/

theorem dot_u_c_sum (c : V) :
    dot u c = ? i : Fin 3, u i * c i := rfl

/-- <u, c> = (1) c_0 + (0) c_1 + (0) c_2 -/
theorem dot_u_c_expansion (c : V) :
    dot u c = (1 : ?) * c 0 + 0 * c 1 + 0 * c 2 := by
    dsimp [dot]
    rw [Fin.sum_univ_three]
    rw [u_zero, u_one, u_two]

/-- <u, c> = c_0 -/
theorem dot_u_c_eq_coord (c : V) :
    dot u c = c 0 := by
    rw [dot_u_c_expansion]
    ring

/-- <u, c> >= 0 for all c in C because c_0 >= 0 -/
theorem dot_u_c_nonneg (c : V) (hc : c in C) :
    0 <= dot u c := by
    rw [dot_u_c_eq_coord]
    exact hc 0

/-!
### Linear functional separation:
    phi_u(v) = -1 < 0 <= c_0 = phi_u(c) for all c in C
-/

/-- The linear functional phi_u corresponding to u under the standard inner product -/
def phi_u : V -> ?[?] ? := toDual u

/-- phi_u(v) = -1 -/
theorem phi_u_v_eval : phi_u v = -1 := by
    change dot u v = -1
    exact dot_u_v_eval

/-- phi_u(c) = c_0 -/
theorem phi_u_c_eval (c : V) : phi_u c = c 0 := by
    change dot u c = c 0
    exact dot_u_c_eq_coord c

/-- phi_u(v) < 0 -/
theorem phi_u_v_neg : phi_u v < 0 := by
    rw [phi_u_v_eval]
    norm_num

/-- 0 <= phi_u(c) for all c in C -/
theorem phi_u_c_nonneg (c : V) (hc : c in C) : 0 <= phi_u c := by
    rw [phi_u_c_eval]

```

```

exact hc 0

/-- Separation:  $\phi_u(v) < 0 \leq \phi_u(c)$  for all  $c$  in  $C$  -/
theorem phi_u_separates (c : V) (hc : c in C) :
   $\phi_u v < 0 \wedge 0 \leq \phi_u c :=$ 
  <phi_u_v_neg, phi_u_c_nonneg c hc>

/-- Complete separation chain:  $\phi_u(v) = -1 < 0 \leq \phi_u(c) = c_0$  -/
theorem phi_u_separation_chain (c : V) (hc : c in C) :
   $\phi_u v = -1 \wedge \phi_u v < 0 \wedge 0 \leq \phi_u c \wedge \phi_u c = c_0 :=$ 
  <phi_u_v_eval, phi_u_v_neg, phi_u_c_nonneg c hc, phi_u_c_eval c>

end Course21241Numerical

```

A.7 BSinMeasurementTheory/Course21228.lean

Lean 4 source

```

import Mathlib.Data.Fintype.Basic
import Mathlib.Data.Fintype.Perm
import Mathlib.Data.Fintype.Card
import Mathlib.Data.Fintype.Prod
import Mathlib.Algebra.BigOperators.Group.Finset.Basic

open Equiv

namespace Course21228

/-!
### Part (a): Symmetric Conditions and Multiset Invariance
-/

section PartA

variable {n : Nat} {a : Type*}

/-- A condition `L` on length-`n` sequences is symmetric if it is invariant
under any permutation of the sequence. -/
def IsSymmetric (L : (Fin n → a) → Prop) : Prop :=
  forall (x : Fin n → a) (? : Perm (Fin n)), L (x comp ?) <-> L x

/-- If two sequences represent the same multiset (i.e. one is a permutation of the
other),
any symmetric condition `L` yields the same truth value on both. -/
theorem symmetric_depends_only_on_multiset
  (L : (Fin n → a) → Prop) (hL : IsSymmetric L)
  (x y : Fin n → a) (? : Perm (Fin n)) (h : y = x comp ?) :
  L y <-> L x := by
  rw [h]
  exact hL x ?

/-- For any additive commutative monoid, the total sum is invariant under permutations
. -/

```

```

theorem sum_perm_invariant {M : Type*} [AddCommMonoid M]
  (x : Fin n → M) (? : Perm (Fin n)) :
  ? i : Fin n, (x comp ?) i = ? i : Fin n, x i := by
  exact Equiv.sum_comp ? x

end PartA

/-!
### Part (b): Counting Pairs of Permutations Under Genericity
-/

section PartB

variable {n : Nat} {a : Type*}

/-- Two sequences `x` and `y` are equal termwise after permuting by `p.1` and `p.2`. -/
def ValidPair (x y : Fin n → a) (p : Perm (Fin n) × Perm (Fin n)) : Prop :=
  forall i : Fin n, x (p.1 i) = y (p.2 i)

/-- Synthesis of decidability for `ValidPair` when equality on `a` is decidable.
    This enables typeclass resolution for `Fintype { p // ValidPair x y p }`. -/
instance [DecidableEq a] (x y : Fin n → a) : DecidablePred (ValidPair x y) := by
  intro p
  unfold ValidPair
  infer_instance

/-- Bijection between `Perm (Fin n)` and the valid pairs `(?, sigma)`.
    Genericity assumptions:
    1. `hij` : vectors in `x` are pairwise distinct (`Function.Injective x`).
    2. `hy` : `y` is a permutation of `x` via `tau : Perm (Fin n)`. -/
def validPairsEquiv (x y : Fin n → a) (hij : Function.Injective x)
  (tau : Perm (Fin n)) (hy : y = x comp tau) :
  Perm (Fin n) → { p : Perm (Fin n) × Perm (Fin n) // ValidPair x y p } where
toFun ? := <(? , ?.trans tau.symm), by
  intro i
  dsimp [ValidPair]
  rw [hy]
  change x (? i) = x (tau (tau.symm (? i)))
  rw [apply_symm_apply]>
invFun p := p.1.1
left_inv ? := rfl
right_inv := by
  rintro <(? , sigma>, h_valid>
  apply Subtype.ext
  dsimp only
  refine Prod.ext rfl ?_
  apply Equiv.ext
  intro i
  have h : x (? i) = y (sigma i) := h_valid i
  rw [hy] at h
  have h_eq : ? i = tau (sigma i) := hij h
  exact (symm_apply_eq tau).mpr h_eq

/-- The exact number of valid pairs is `n!`. -/
theorem validPairs_card [DecidableEq a] (x y : Fin n → a)
  (hij : Function.Injective x) (tau : Perm (Fin n)) (hy : y = x comp tau) :

```

```

    Fintype.card { p : Perm (Fin n) x Perm (Fin n) // ValidPair x y p } = Nat.
    factorial n := by
    rw [ <- Fintype.card_congr (validPairsEquiv x y hinv tau hy)]
    rw [Fintype.card_perm]
    rw [Fintype.card_fin]

end PartB

/-!
### Part (c): Reduction to n = 2 and n = 3
-/

section PartC

variable {a : Type*}

/-- Exact reduction to n = 2: 2! = 2 pairs. -/
theorem validPairs_card_two [DecidableEq a] (x y : Fin 2 -> a)
  (hinv : Function.Injective x) (tau : Perm (Fin 2)) (hy : y = x comp tau) :
  Fintype.card { p : Perm (Fin 2) x Perm (Fin 2) // ValidPair x y p } = 2 := by
  rw [validPairs_card x y hinv tau hy]
  rfl

/-- Exact reduction to n = 3: 3! = 6 pairs. -/
theorem validPairs_card_three [DecidableEq a] (x y : Fin 3 -> a)
  (hinv : Function.Injective x) (tau : Perm (Fin 3)) (hy : y = x comp tau) :
  Fintype.card { p : Perm (Fin 3) x Perm (Fin 3) // ValidPair x y p } = 6 := by
  rw [validPairs_card x y hinv tau hy]
  rfl

/-- Direct verification of the underlying symmetric group cardinalities. -/
theorem perm_fin_two_card : Fintype.card (Perm (Fin 2)) = 2 := by decide
theorem perm_fin_three_card : Fintype.card (Perm (Fin 3)) = 6 := by decide

end PartC

/-!
### Numerical Evaluation
-/

#eval Nat.factorial 2 -- 2
#eval Nat.factorial 3 -- 6

end Course21228

```

A.8 BSinMeasurementTheory/Course21355.lean

Lean 4 source

```

import Mathlib

open Filter Topology

```

```

namespace Course21355

noncomputable section

/-!
### (a) Bolzano-Weierstrass Theorem
Mathlib derives sequential compactness of closed bounded intervals in ? from the
completeness axiom / Heine-Borel compactness (`IsCompact.tendsto_subseq`).
-/

theorem bolzano_weierstrass (x : Nat → ?) (h_bdd : exists M : ?, forall n, |x n|
  ≤ M) :
  exists (phi : Nat → Nat) (l : ?), StrictMono phi /\ Tendsto (x comp phi)
    atTop (? l) := by
  rcases h_bdd with <M, hM>
  have h_mem : forall n, x n in Set.Icc (-M) M := fun n => abs_le.mp (hM n)
  have h_freq : exists ? n in atTop, x n in Set.Icc (-M) M :=
    Filter.frequently_atTop.2 (fun N => <N, le_refl, h_mem N>)
  rcases isCompact_Icc.tendsto_subseq' h_freq with <l, _, phi, hphi_mono, hphi_lim>
  exact <phi, l, hphi_mono, hphi_lim>

/-!
### (b) Existence and Characterization of limsup and liminf
The tail supremum sequence y_k = bigcup n, x (n + k) is monotone non-increasing,
and the tail infimum sequence z_k = ? n, x (n + k) is monotone non-decreasing.
Both converge to the limsup and liminf via monotone convergence in ?.
-/

def tailSup (x : Nat → ?) (k : Nat) : ? := bigcup n : Nat, x (n + k)
def tailInf (x : Nat → ?) (k : Nat) : ? := ? n : Nat, x (n + k)

theorem tailSup_antitone (x : Nat → ?) (M : ?) (hM : forall n, x n ≤ M) :
  Antitone (tailSup x) := by
  intro j k hjk
  apply ciSup_le
  intro n
  have h_eq : n + k = (n + (k - j)) + j := by omega
  rw [h_eq]
  have h_bdd : BddAbove (Set.range (fun i : Nat => x (i + j))) :=
    <M, fun r <i, hi> => hi |> hM (i + j)>
  exact le_ciSup h_bdd (n + (k - j))

theorem tailInf_monotone (x : Nat → ?) (m : ?) (hm : forall n, m ≤ x n) :
  Monotone (tailInf x) := by
  intro j k hjk
  apply le_ciInf
  intro n
  have h_eq : n + k = (n + (k - j)) + j := by omega
  rw [h_eq]
  have h_bdd : BddBelow (Set.range (fun i : Nat => x (i + j))) :=
    <m, fun r <i, hi> => hi |> hm (i + j)>
  exact ciInf_le h_bdd (n + (k - j))

theorem limsup_liminf_exist (x : Nat → ?) (m M : ?) (h : forall n, m ≤ x n /\
  x n ≤ M) :
  (exists L_sup : ?, Tendsto (tailSup x) atTop (? L_sup)) /\

```

```

    (exists L_inf : ?, Tendsto (tailInf x) atTop (? L_inf)) := by
have h_sup_bdd : BddBelow (Set.range (tailSup x)) := by
  refine <m, fun y <k, hk> => ?_>
  subst hk
  have h1 : m <= x (0 + k) := (h (0 + k)).1
  have hb : BddAbove (Set.range (fun n : Nat => x (n + k))) :=
    <M, fun r <i, hi> => hi |> (h (i + k)).2>
  have h2 : x (0 + k) <= tailSup x k := le_ciSup hb 0
  exact le_trans h1 h2
have h_inf_bdd : BddAbove (Set.range (tailInf x)) := by
  refine <M, fun y <k, hk> => ?_>
  subst hk
  have hb : BddBelow (Set.range (fun n : Nat => x (n + k))) :=
    <m, fun r <i, hi> => hi |> (h (i + k)).1>
  have h1 : tailInf x k <= x (0 + k) := ciInf_le hb 0
  have h2 : x (0 + k) <= M := (h (0 + k)).2
  exact le_trans h1 h2
exact <<? k, tailSup x k, tendsto_atTop_ciInf (tailSup_antitone x M (fun n => (h n)
.2)) h_sup_bdd>,
  <bigcup k, tailInf x k, tendsto_atTop_ciSup (tailInf_monotone x m (fun n =>
(h n).1)) h_inf_bdd>>

/-!
### (c) Divergence of  $x_n = (-1)^n$  via an  $\epsilon$ -argument ( $\epsilon = 1$ )
-/

def altSeq (n : Nat) : ? := (-1 : ?) ^ n

lemma altSeq_even (k : Nat) : altSeq (2 * k) = 1 := by
  dsimp [altSeq]
  rw [pow_mul]
  have : (-1 : ?) ^ 2 = 1 := by norm_num
  rw [this, one_pow]

lemma altSeq_odd (k : Nat) : altSeq (2 * k + 1) = -1 := by
  dsimp [altSeq]
  rw [pow_add, pow_one, pow_mul]
  have : (-1 : ?) ^ 2 = 1 := by norm_num
  rw [this, one_pow, one_mul]

theorem altSeq_no_limit : not exists (L : ?), Tendsto altSeq atTop (? L) := by
  rintro <L, hL>
  rw [Metric.tendsto_atTop] at hL
  have h_eps : (0 : ?) < 1 := by norm_num
  rcases hL 1 h_eps with <N, hN>
  have h_even : dist (altSeq (2 * N)) L < 1 := hN (2 * N) (Nat.le_mul_of_pos_left _ (
    by norm_num))
  have h_odd : dist (altSeq (2 * N + 1)) L < 1 := hN (2 * N + 1) (by omega)
  rw [altSeq_even N, Real.dist_eq] at h_even
  rw [altSeq_odd N, Real.dist_eq] at h_odd
  have h_triangle : (2 : ?) <= |1 - L| + |-1 - L| := by
    have h2 : (2 : ?) = (1 - L) - (-1 - L) := by ring
    rw [h2]
    exact le_trans (le_abs_self _) (abs_sub (1 - L) (-1 - L))
  have h_lt : |1 - L| + |-1 - L| < 2 := by linarith
  linarith

```

```

/-!
### (d) Compactness of [a, b] from Open Covers
-/

theorem compact_Icc (a b : ?) : IsCompact (Set.Icc a b) :=
  isCompact_Icc

/-!
### Numerical Calculations
Computable evaluation of the alternating values:
-/
def altSeqZ (n : Nat) : Int := (-1 : Int) ^ n

#eval altSeqZ 0 -- 1
#eval altSeqZ 1 -- -1
#eval altSeqZ 2 -- 1
#eval altSeqZ 3 -- -1

end

end Course21355

```

A.9 BSinMeasurementTheory/Course21373.lean

Lean 4 source

```

import Mathlib

open Finset
open Classical

namespace Course21373

variable {B : Type*} [BooleanAlgebra B]

/-! ### Definitions -/

/-- An atom in a Boolean algebra is a non-bottom element whose only sub-elements are
    False and itself.
    Named `IsBooleanAtom` to avoid collision with Mathlib's order-theoretic `IsAtom`.
-/
def IsBooleanAtom (a : B) : Prop :=
  a != False /\ forall x, x <= a -> x = False \/ x = a

/-! ### Part (a): Atoms and Unique Join Representation -/

/-- Distinct atoms are disjoint. -/
theorem isAtom_inf_eq_bot {a b : B} (ha : IsBooleanAtom a) (hb : IsBooleanAtom b) (hne
  : a != b) :
  a sqcap b = False := by
  have hle : a sqcap b <= a := inf_le_left
  cases ha.2 (a sqcap b) hle with

```

```

| inl h => exact h
| inr h =>
  have hle2 : a <= b := by
    calc a = a sqcap b := h.symm
    _ <= b := inf_le_right
  cases hb.2 a hle2 with
  | inl hbot => exact (ha.1 hbot).elim
  | inr heq => exact (hne heq).elim

/-- An atom is below a join of two elements iff it is below at least one. -/
theorem isAtom_le_sup_iff {a b c : B} (ha : IsBooleanAtom a) :
  a <= b sqsup c <-> a <= b \/\ a <= c := by
  constructor
  * intro h
    have hdist : a = (a sqcap b) sqsup (a sqcap c) := by
      calc a = a sqcap (b sqsup c) := (inf_eq_left.mpr h).symm
      _ = (a sqcap b) sqsup (a sqcap c) := inf_sup_left a b c
    cases ha.2 (a sqcap b) inf_le_left with
    | inr hab =>
      left
      calc a = a sqcap b := hab.symm
      _ <= b := inf_le_right
    | inl hab =>
      right
      have hac : a = a sqcap c := by
        calc a = (a sqcap b) sqsup (a sqcap c) := hdist
        _ = False sqsup (a sqcap c) := congr_arg (* sqsup (a sqcap c)) hab
        _ = a sqcap c := bot_sup_eq (a sqcap c)
      calc a = a sqcap c := hac
      _ <= c := inf_le_right
  * rintro (h | h)
  * exact le_trans h le_sup_left
  * exact le_trans h le_sup_right

/-- An atom is below a finite join iff it is below some member. -/
theorem isAtom_le_finsup {a : B} (ha : IsBooleanAtom a) (s : Finset B) :
  a <= s.sup id <-> exists b in s, a <= b := by
  induction s using Finset.induction_on with
  | empty =>
    constructor
    * intro h
      rw [Finset.sup_empty] at h
      exact (ha.1 (le_bot_iff.mp h)).elim
    * rintro <b, hb, _>
      cases hb
  | @insert b s _ ih =>
    rw [Finset.sup_insert]
    dsimp only [id]
    rw [isAtom_le_sup_iff ha, ih]
    simp only [Finset.mem_insert]
    constructor
    * rintro (h | <c, hc, hac>)
    * exact <b, Or.inl rfl, h>
    * exact <c, Or.inr hc, hac>
    * rintro <c, rfl | hc, hac>
    * exact Or.inl hac

```

```

      * exact Or.inr <c, hc, hac>

/-- If `s` consists of atoms, an atom `a` <= s.sup id` iff `a` in `s`. -/
theorem isAtom_le_finset_sup_of_atoms {a : B} (ha : IsBooleanAtom a) {s : Finset B}
  (hs : forall b in s, IsBooleanAtom b) :
  a <= s.sup id <-> a in s := by
  rw [isAtom_le_finset_sup ha]
  constructor
  * rintro <b, hb, hab>
    have hb_atom := hs b hb
    cases hb_atom.2 a hab with
    | inl hbot => exact (ha.1 hbot).elim
    | inr heq =>
      rw [heq]
      exact hb
  * intro ha_mem
    exact <a, ha_mem, le_refl>

/-- Uniqueness of representation: two subsets of atoms with equal join must be equal.
-/
theorem finset_sup_inj_of_atoms {s t : Finset B}
  (hs : forall b in s, IsBooleanAtom b) (ht : forall b in t, IsBooleanAtom b)
  (h_eq : s.sup id = t.sup id) : s = t := by
  ext a
  constructor
  * intro ha
    have ha_atom := hs a ha
    have hle : a <= s.sup id := (isAtom_le_finset_sup_of_atoms ha_atom hs).mpr ha
    rw [hle] at hle
    exact (isAtom_le_finset_sup_of_atoms ha_atom ht).mp hle
  * intro ha
    have ha_atom := ht a ha
    have hle : a <= t.sup id := (isAtom_le_finset_sup_of_atoms ha_atom ht).mpr ha
    rw [hle] at hle
    exact (isAtom_le_finset_sup_of_atoms ha_atom hs).mp hle

/-! ### Part (b): Representation Map and Characteristic Functions -/

/-- The canonical map sending an element to the atoms below it. -/
noncomputable def repMap (A : Finset B) (x : B) : Finset B :=
  A.filter (fun a => a <= x)

theorem repMap_inf (A : Finset B) (x y : B) :
  repMap A (x sqcap y) = repMap A x ∩ repMap A y := by
  ext a
  simp only [repMap, Finset.mem_filter, Finset.mem_inter, le_inf_iff]
  tauto

theorem repMap_sup {A : Finset B} (hA : forall a in A, IsBooleanAtom a) (x y : B) :
  repMap A (x sqsup y) = repMap A x ∪ repMap A y := by
  ext a
  simp only [repMap, Finset.mem_filter, Finset.mem_union]
  constructor
  * rintro <haA, hle>
    have ha_atom := hA a haA
    rw [isAtom_le_sup_iff ha_atom] at hle

```

```

    cases hle with
    | inl h => exact Or.inl <haA, h>
    | inr h => exact Or.inr <haA, h>
* rintro (<haA, h> | <haA, h>)
* exact <haA, le_trans h le_sup_left>
* exact <haA, le_trans h le_sup_right>

theorem repMap_bot {A : Finset B} (hA : forall a in A, IsBooleanAtom a) :
  repMap A False = {} := by
  ext a
  simp only [repMap, Finset.mem_filter]
  constructor
* rintro <haA, hle>
  have ha_bot : a = False := le_bot_iff.mp hle
  exact ((hA a haA).1 ha_bot).elim
* intro h
  cases h

/-- `repMap` is the inverse of `Finset.sup id` on subsets of atoms. -/
theorem repMap_sup_id {A : Finset B} (hA : forall a in A, IsBooleanAtom a) {s :
  Finset B} (hs : s ⊆ A) :
  repMap A (s.sup id) = s := by
  ext a
  simp only [repMap, Finset.mem_filter]
  constructor
* rintro <haA, hle>
  have ha_atom := hA a haA
  exact (isAtom_le_finset_sup_of_atoms ha_atom (fun b hb => hA b (hs hb))).mp hle
* intro ha
  have haA := hs ha
  have ha_atom := hA a haA
  refine <haA, ?_>
  exact (isAtom_le_finset_sup_of_atoms ha_atom (fun b hb => hA b (hs hb))).mpr ha

/-- Natural equivalence between subsets of `{0, ..., n-1}` and characteristic functions
  . -/
def finsetToCharFun (n : Nat) : Finset (Fin n) → (Fin n → Bool) where
  toFun s := fun i => decide (i in s)
  invFun f := Finset.filter (fun i => f i = true) Finset.univ
  left_inv s := by
    ext i
    simp
  right_inv f := by
    ext i
    simp

/-! ### Part (c): Group Action, Fixed Subalgebra, and Orbit Atoms -/

variable {G : Type*} [SMul G B]

/-- The fixed-point set of the action. -/
def fixedPoints (G : Type*) (B : Type*) [SMul G B] : Set B :=
  {x : B | forall g : G, g * x = x}

/-- An action of G on B by Boolean algebra automorphisms. -/
class BooleanAction (G : Type*) (B : Type*) [BooleanAlgebra B] [SMul G B] : Prop where

```

```

smul_sup : forall (g : G) (x y : B), g * (x sqsup y) = (g * x) sqsup (g * y)
smul_inf : forall (g : G) (x y : B), g * (x sqcap y) = (g * x) sqcap (g * y)
smul_compl : forall (g : G) (x : B), g * (x?) = (g * x)?
smul_bot : forall (g : G), g * (False : B) = False
smul_top : forall (g : G), g * (top : B) = top

theorem fixed_bot [BooleanAction G B] : (False : B) in fixedPoints G B := fun g =>
  BooleanAction.smul_bot g

theorem fixed_top [BooleanAction G B] : (top : B) in fixedPoints G B := fun g =>
  BooleanAction.smul_top g

theorem fixed_sup [BooleanAction G B] {x y : B}
  (hx : x in fixedPoints G B) (hy : y in fixedPoints G B) :
  x sqsup y in fixedPoints G B := by
  intro g
  rw [BooleanAction.smul_sup, hx g, hy g]

theorem fixed_inf [BooleanAction G B] {x y : B}
  (hx : x in fixedPoints G B) (hy : y in fixedPoints G B) :
  x sqcap y in fixedPoints G B := by
  intro g
  rw [BooleanAction.smul_inf, hx g, hy g]

theorem fixed_compl [BooleanAction G B] {x : B}
  (hx : x in fixedPoints G B) :
  x? in fixedPoints G B := by
  intro g
  rw [BooleanAction.smul_compl, hx g]

/-- An atom of the fixed-point subalgebra. -/
def IsFixedAtom (G : Type*) (B : Type*) [BooleanAlgebra B] [SMul G B] (b : B) : Prop
:=
  b in fixedPoints G B /\ b != False /\ forall x in fixedPoints G B, x <=
    b -> x = False \/ x = b

/-- Distribution of meet over a finite join. -/
theorem inf_finset_sup (x : B) (s : Finset B) :
  x sqcap s.sup id = s.sup (fun a => x sqcap a) := by
  induction s using Finset.induction_on with
  | empty => simp only [Finset.sup_empty, inf_bot_eq]
  | @insert b s _ ih =>
    rw [Finset.sup_insert]
    dsimp only [id]
    rw [inf_sup_left, ih, Finset.sup_insert]

theorem sup_congr_bot {s : Finset B} {f : B -> B} (h : forall a in s, f a = False) :
  s.sup f = False := by
  induction s using Finset.induction_on with
  | empty => simp only [Finset.sup_empty]
  | @insert b s _ ih =>
    rw [Finset.sup_insert]
    have hb : f b = False := h b (Finset.mem_insert_self b s)
    have hs : forall a in s, f a = False := fun a ha => h a (Finset.
      mem_insert_of_mem ha)

```

```

    rw [hb, ih hs, bot_sup_eq]

/-- The join over an orbit of atoms is an atom of the fixed-point subalgebra. -/
theorem orbit_sup_isFixedAtom {O : Finset B}
  (hO_ne : O.Nonempty)
  (hO_atoms : forall a in O, IsBooleanAtom a)
  (hO_fixed : O.sup id in fixedPoints G B)
  (hO_min : forall x in fixedPoints G B, (forall a in O, a <= x) \/\ (
    forall a in O, a sqcap x = False)) :
  IsFixedAtom G B (O.sup id) := by
  refine <hO_fixed, ?_, ?_>
  * -- b != False
    rcases hO_ne with <a, ha>
    have ha_atom := hO_atoms a ha
    intro hbot
    have hle : a <= O.sup id := (isAtom_le_finset_sup_of_atoms ha_atom hO_atoms).mpr
    ha
    rw [hbot] at hle
    exact ha_atom.1 (le_bot_iff.mp hle)
  * -- x <= b -> x = False \/\ x = b
    intro x hx hle
    rcases hO_min x hx with (hall | hnone)
  * right
    have hsup_le : O.sup id <= x := by
      apply Finset.sup_le
      intro a ha
      exact hall a ha
    exact le_antisymm hle hsup_le
  * left
    have hx_eq : x = x sqcap O.sup id := (inf_eq_left.mpr hle).symm
    rw [hx_eq, inf_finset_sup]
    apply sup_congr_bot
    intro a ha
    rw [inf_comm]
    exact hnone a ha

end Course21373

```

A.10 BSinMeasurementTheory/Course21373Numerical.lean

Lean 4 source

```

import Mathlib
namespace Course21373Numerical

/--! ### Numerical / Computational Example -/

def orbit1 : Finset (Fin 3) := {0, 1}
def orbit2 : Finset (Fin 3) := {2}

-- Both orbit joins are disjoint and partition the full set of atoms
#eval (orbit1 I orbit2 == ({} : Finset (Fin 3))) -- true
#eval (orbit1 U orbit2 == {0, 1, 2}) -- true

```

```

-- The 4 fixed points of the action in P({0,1,2})
def fixed_elements : List (Finset (Fin 3)) :=
  [{}, orbit1, orbit2, orbit1 ∪ orbit2]

#eval fixed_elements.length          -- 4

end Course21373Numerical

```

A.11 BSinMeasurementTheory/Course21321.lean

Lean 4 source

```

import Mathlib

open BigOperators

namespace Course21321

namespace Scott1964

/--
## Section 1: Theorem 1.1 (Separating Hyperplane / Real Duality Form)
Quantified universally over an arbitrary real vector space `V`.
-/

variable {V : Type*} [AddCommGroup V] [Module ? V]

/-- A real linear functional strictly separating `Y` from 0 while keeping `X` non-
negative. -/
def RealSolvable (X Y : Finset V) : Prop :=
  exists f : V → ℝ, (forall x in X, 0 ≤ f x) /\ (forall y in Y, 0 <
    f y)

/-- The real non-cancellation condition: no non-trivial non-negative linear
combination sums to 0. -/
def RealNoCancellation (X Y : Finset V) : Prop :=
  forall (c_coeff d_coeff : V → ℝ),
    (forall x in X, 0 ≤ c_coeff x) →
    (forall y in Y, 0 ≤ d_coeff y) →
    (0 < ∑ y in Y, d_coeff y) →
    (∑ x in X, c_coeff x * x) + (∑ y in Y, d_coeff y * y) ≠ 0

/-- Scott's Theorem 1.1: Solvability implies real non-cancellation. -/
theorem theorem_1_1_real_separation [DecidableEq V] {X Y : Finset V}
  (hsol : RealSolvable X Y) : RealNoCancellation X Y := by
  intro c_coeff d_coeff hc hd hd_pos hsum
  rcases hsol with <f, hfX, hfY>
  have f_linear_sum : f (∑ x in X, c_coeff x * x) + (∑ y in Y, d_coeff y * y) =
    0 := by
    rw [hsum, map_zero]
  rw [map_add, map_sum, map_sum] at f_linear_sum

```

```

have f_terms_X : forall x in X, 0 <= c_coeff x * f x := by
  intro x hx
  exact mul_nonneg (hc x hx) (hfX x hx)
have f_terms_Y : forall y in Y, 0 <= d_coeff y * f y := by
  intro y hy
  exact mul_nonneg (hd y hy) (le_of_lt (hfY y hy))
have exists_strict_pos : exists y0 in Y, 0 < d_coeff y0 * f y0 := by
  by_contra! hall
  have hall_zero : forall y in Y, d_coeff y = 0 := by
    intro y hy
    have hle := hall y hy
    have hnonneg := hd y hy
    have hfpos := hfY y hy
    obtain heq | hgt := le_iff_eq_or_lt.mp hnonneg
    * exact heq.symm
    * exfalse
    have : 0 < d_coeff y * f y := mul_pos hgt hfpos
    linarith
  have sum_zero : ? y in Y, d_coeff y = 0 :=
    Finset.sum_eq_zero hall_zero
  linarith
rcases exists_strict_pos with <y0, hy0, hy0_pos>
have sum_Y_pos : 0 < ? y in Y, f (d_coeff y * y) := by
  rw [ <- Finset.add_sum_erase _ _ hy0 ]
  have h_rest : 0 <= ? y in Y.erase y0, f (d_coeff y * y) := by
    apply Finset.sum_nonneg
    intro y hy
    have hyY := Finset.mem_of_mem_erase hy
    rw [map_smul, smul_eq_mul]
    exact f_terms_Y y hyY
  rw [map_smul, smul_eq_mul]
  linarith
have hX_nonneg : 0 <= ? x in X, f (c_coeff x * x) := by
  apply Finset.sum_nonneg
  intro x hx
  rw [map_smul, smul_eq_mul]
  exact f_terms_X x hx
have total_pos : 0 < (? x in X, f (c_coeff x * x)) + (? y in Y, f (d_coeff y * y)) := by
  linarith
linarith

/-!
## Section 2: Theorem 1.2 (Combinatorial / Integer Cancellation Form)
Quantified universally over an arbitrary rational vector space `V_Q`.
-/

variable {V_Q : Type*} [AddCommGroup V_Q] [Module ? V_Q]

/-- Existence of a rational separating linear functional. -/
def RatSolvable (X Y : Finset V_Q) : Prop :=
  exists f : V_Q -> ?[?] ?, (forall x in X, 0 <= f x) /\ (forall y in Y, 0
    < f y)

/-- Scott's Combinatorial Finite Cancellation condition with natural-number
multipliers. -/

```

```

def CombinatorialNoCancellation (X Y : Finset V_Q) : Prop :=
  forall (n m : V_Q → Nat),
    (0 < ? y in Y, m y) →
      (? x in X, (n x : ?) * x) + (? y in Y, (m y : ?) * y) != 0

/-- Scott's Theorem 1.2: Rational solvability implies combinatorial non-cancellation.
-/
theorem theorem_1_2_combinatorial_cancellation [DecidableEq V_Q] {X Y : Finset V_Q}
  (hsol : RatSolvable X Y) : CombinatorialNoCancellation X Y := by
  intro n m hm_pos hsum
  rcases hsol with <f, hfX, hfY>
  have f_sum : f ((? x in X, (n x : ?) * x) + (? y in Y, (m y : ?) * y)) = 0 := by
    rw [hsum, map_zero]
  rw [map_add, map_sum, map_sum] at f_sum
  have f_terms_X : forall x in X, 0 <= (n x : ?) * f x := by
    intro x hx
    exact mul_nonneg (Nat.cast_nonneg (n x)) (hfX x hx)
  have exists_strict_pos : exists y0 in Y, 0 < (m y0 : ?) * f y0 := by
    by_contra! hall
    have hall_zero : forall y in Y, m y = 0 := by
      intro y hy
      have hle := hall y hy
      have hfpos := hfY y hy
      by_contra hm_ne
      have hm_gt : 0 < (m y : ?) := Nat.cast_pos.mpr (Nat.pos_of_ne_zero hm_ne)
      have : 0 < (m y : ?) * f y := mul_pos hm_gt hfpos
      linarith
    have sum_zero : ? y in Y, m y = 0 :=
      Finset.sum_eq_zero hall_zero
    linarith
  rcases exists_strict_pos with <y0, hy0, hy0_pos>
  have sum_Y_pos : 0 < ? y in Y, f ((m y : ?) * y) := by
    rw [← Finset.add_sum_erase _ _ hy0]
  have h_rest : 0 <= ? y in Y.erase y0, f ((m y : ?) * y) := by
    apply Finset.sum_nonneg
    intro y hy
    have hyY := Finset.mem_of_mem_erase hy
    rw [map_smul, smul_eq_mul]
    exact mul_nonneg (Nat.cast_nonneg (m y)) (le_of_lt (hfY y hyY))
  rw [map_smul, smul_eq_mul]
  linarith
  have hX_nonneg : 0 <= ? x in X, f ((n x : ?) * x) := by
    apply Finset.sum_nonneg
    intro x hx
    rw [map_smul, smul_eq_mul]
    exact f_terms_X x hx
  have total_pos : 0 < (? x in X, f ((n x : ?) * x)) + (? y in Y, f ((m y : ?) * y)) := by
    linarith
  linarith

/-!
## Section 3: Theorem 1.3 (Equivalence of Combinatorial and Rational Conditions)
-/

/-- Rational non-cancellation condition over non-negative rational coefficients. -/

```

```

def RatNoCancellation (X Y : Finset V_Q) : Prop :=
  forall (c_coeff d_coeff : V_Q -> ?),
    (forall x in X, 0 <= c_coeff x) ->
    (forall y in Y, 0 <= d_coeff y) ->
    (0 < ? y in Y, d_coeff y) ->
    (? x in X, c_coeff x * x) + (? y in Y, d_coeff y * y) != 0

/-- Equivalence: Rational non-cancellation implies combinatorial non-cancellation. -/
theorem combinatorial_of_rat_no_cancellation {X Y : Finset V_Q}
  (hrat : RatNoCancellation X Y) : CombinatorialNoCancellation X Y := by
  intro n m hm_pos hsum
  have hc : forall x in X, 0 <= (n x : ?) := fun x _ => Nat.cast_nonneg (n x)
  have hd : forall y in Y, 0 <= (m y : ?) := fun y _ => Nat.cast_nonneg (m y)
  have hd_sum : (0 : ?) < ? y in Y, (m y : ?) := by
    have hpos : (0 : ?) < ((? y in Y, m y : Nat) : ?) := Nat.cast_pos.mpr hm_pos
    rwa [Nat.cast_sum] at hpos
  exact hrat (fun x => (n x : ?)) (fun y => (m y : ?)) hc hd hd_sum hsum

end Scott1964

end Course21321

```

A.12 BSinMeasurementTheory/Course21321Kraft.lean

Lean 4 source

```

import Mathlib

open BigOperators

namespace Course21321Kraft

noncomputable section

namespace KraftExample

/-!
### Definition of the Difference Vectors in ??
Let ? = {0, 1, 2, 3, 4}.
We define 4 difference vectors corresponding to comparative probability claims:
  y_1 = (-1, -1, 1, 1, 0)
  y_2 = (-1, 1, -1, 0, 1)
  y_3 = (1, -1, -1, 1, 0)
  y_? = (1, 1, 1, -2, -1)
-/

def y1 : Fin 5 -> ? := ![-1, -1, 1, 1, 0]
def y2 : Fin 5 -> ? := ![-1, 1, -1, 0, 1]
def y3 : Fin 5 -> ? := ![1, -1, -1, 1, 0]
def y4 : Fin 5 -> ? := ![1, 1, 1, -2, -1]

/-!
### Numerical Cancellation Check: ? y_i = 0

```

```

Verified coordinate-by-coordinate using `fin_cases` and `norm_num`.
-/

theorem kraft_sum_zero : y1 + y2 + y3 + y4 = 0 := by
  ext i
  fin_cases i <|> norm_num [y1, y2, y3, y4]

/-!
### Proof that No Linear Functional Can Strictly Separate These Vectors
Applying any linear functional `f` with `f(y_i) > 0` produces a contradiction:
  0 = f(0) = f(y_1 + y_2 + y_3 + y?) = f(y_1) + f(y_2) + f(y_3) + f(y?) > 0
-/

theorem no_strictly_positive_functional (f : (Fin 5 -> ?) -> ?[?] ?)
  (h1 : 0 < f y1) (h2 : 0 < f y2) (h3 : 0 < f y3) (h4 : 0 < f y4) : False := by
  have h_sum : f (y1 + y2 + y3 + y4) = 0 := by
    rw [kraft_sum_zero, map_zero]
  have h_pos : 0 < f (y1 + y2 + y3 + y4) := by
    rw [map_add, map_add, map_add]
    linarith
  linarith

/-!
### Connection to Scott's Solvability Condition
Defining Y = {y_1, y_2, y_3, y?}, we prove `not RealSolvable {} Y`.
-/

def Y : Finset (Fin 5 -> ?) := {y1, y2, y3, y4}

/-- Scott's Real Solvability on ({}, Y) fails. -/
theorem kraft_not_solvable :
  not (exists f : (Fin 5 -> ?) -> ?[?] ?, forall y in Y, 0 < f y) := by
  intro <f, hf>
  have hy1 : y1 in Y := by simp [Y]
  have hy2 : y2 in Y := by simp [Y]
  have hy3 : y3 in Y := by simp [Y]
  have hy4 : y4 in Y := by simp [Y]
  exact no_strictly_positive_functional f (hf y1 hy1) (hf y2 hy2) (hf y3 hy3) (hf y4
    hy4)

end KraftExample

end

end Course21321Kraft

```

A.13 BSinMeasurementTheory/Course21292.lean

Lean 4 source

```

import Mathlib

open scoped BigOperators

```

```

namespace Course21292

namespace LPDuality

-- Vector inner product over Rational numbers for Fin n
def dot {n : Nat} (u v : Fin n → ?) : ? :=
  ? i : Fin n, u i * v i

-- Non-negativity of a vector
def NonNeg {n : Nat} (v : Fin n → ?) : Prop :=
  forall i : Fin n, 0 <= v i

-- Matrix-vector multiplication (A * x)
def matMulVec {m n : Nat} (A : Fin m → Fin n → ?) (x : Fin n → ?) : Fin m →
  ? :=
  fun i => dot (A i) x

-- Transpose matrix-vector multiplication (A? * y)
def matTransposeMulVec {m n : Nat} (A : Fin m → Fin n → ?) (y : Fin m → ?) :
  Fin n → ? :=
  fun j => ? i : Fin m, y i * A i j

/-!
### General Weak Duality Theorem
-/

theorem transpose_dot_assoc {m n : Nat} (A : Fin m → Fin n → ?)
  (y : Fin m → ?) (x : Fin n → ?) :
  dot y (matMulVec A x) = dot (matTransposeMulVec A y) x := by
  dsimp [dot, matMulVec, matTransposeMulVec]
  -- Distribute multiplications inside the sums before commuting
  simp_rw [Finset.mul_sum, Finset.sum_mul]
  rw [Finset.sum_comm]
  apply Finset.sum_congr rfl
  intro j _
  apply Finset.sum_congr rfl
  intro i _
  ring

theorem weak_duality {m n : Nat} (A : Fin m → Fin n → ?)
  (b : Fin m → ?) (c : Fin n → ?)
  (x : Fin n → ?) (y : Fin m → ?)
  (hx_nonneg : NonNeg x)
  (hy_nonneg : NonNeg y)
  (h_primal_feas : forall i, b i <= matMulVec A x i)
  (h_dual_feas : forall j, matTransposeMulVec A y j <= c j) :
  dot b y <= dot c x := by
  have h1 : dot b y <= dot (matMulVec A x) y := by
    dsimp [dot]
    have h_term : forall i : Fin m, b i * y i <= matMulVec A x i * y i := by
      intro i
      have hy : 0 <= y i := hy_nonneg i
      have hfeas : b i <= matMulVec A x i := h_primal_feas i
      nlinarith
    exact Finset.sum_le_sum fun i _ => h_term i

```

```

have h_comm : dot (matMulVec A x) y = dot y (matMulVec A x) := by
  dsimp [dot]
  apply Finset.sum_congr rfl
  intro i _
  ring
rw [h_comm, transpose_dot_assoc] at h1
have h2 : dot (matTransposeMulVec A y) x <= dot c x := by
  dsimp [dot]
  have h_term : forall j : Fin n, matTransposeMulVec A y j * x j <= c j * x j :=
  by
    intro j
    have hx : 0 <= x j := hx_nonneg j
    have hfeas : matTransposeMulVec A y j <= c j := h_dual_feas j
    nlinarith
  exact Finset.sum_le_sum fun j _ => h_term j
exact le_trans h1 h2

/-!
### Concrete 3x3 Instance Verification (Scott 1.1 Cone Setting)
-/

def A_inst : Fin 3 -> Fin 3 -> ?
| 0, 0 => 2 | 0, 1 => 1 | 0, 2 => 0
| 1, 0 => 1 | 1, 1 => 2 | 1, 2 => 0
| 2, 0 => 0 | 2, 1 => 0 | 2, 2 => 1

def b_inst : Fin 3 -> ?
| 0 => 3 | 1 => 3 | 2 => 1

def c_inst : Fin 3 -> ?
| 0 => 1 | 1 => 1 | 2 => 2

def x_star : Fin 3 -> ?
| 0 => 1 | 1 => 1 | 2 => 1

def y_star : Fin 3 -> ?
| 0 => 1/3 | 1 => 1/3 | 2 => 2

-- Primal Cost Evaluation: dot(c, x*) = 4
theorem primal_cost_eval : dot c_inst x_star = 4 := by
  dsimp [dot, c_inst, x_star]
  rw [Fin.sum_univ_three]
  norm_num

-- Dual Cost Evaluation: dot(b, y*) = 4
theorem dual_cost_eval : dot b_inst y_star = 4 := by
  dsimp [dot, b_inst, y_star]
  rw [Fin.sum_univ_three]
  norm_num

-- Primal Feasibility Proof
theorem x_star_feasible :
  NonNeg x_star /\ (forall i, b_inst i <= matMulVec A_inst x_star i) := by
  constructor
  * intro i
  fin_cases i <|> norm_num [x_star, NonNeg]

```

```

* intro i
  fin_cases i <|> {
    dsimp [b_inst, matMulVec, dot, A_inst, x_star]
    rw [Fin.sum_univ_three]
    norm_num
  }

-- Dual Feasibility Proof
theorem y_star_feasible :
  NonNeg y_star /\ (forall j, matTransposeMulVec A_inst y_star j <= c_inst j)
:= by
  constructor
* intro i
  fin_cases i <|> norm_num [y_star, NonNeg]
* intro j
  fin_cases j <|> {
    dsimp [c_inst, matTransposeMulVec, A_inst, y_star]
    rw [Fin.sum_univ_three]
    norm_num
  }

-- Certificate of Zero Duality Gap
theorem strong_duality_certificate :
  dot c_inst x_star = dot b_inst y_star := by
  rw [primal_cost_eval, dual_cost_eval]

end LPDuality

end Course21292

```

A.14 B SinMeasurementTheory/Course21292Numerical.lean

Lean 4 source

```

import Mathlib

open scoped BigOperators

namespace Course21292Numerical

namespace LPDualityCalculations

-- 1. Definitions of Vector/Matrix Operations
def dot {n : Nat} (u v : Fin n → ?) : ? :=
  ? i : Fin n, u i * v i

def matMulVec {m n : Nat} (A : Fin m → Fin n → ?) (x : Fin n → ?) : Fin m →
  ? :=
  fun i => dot (A i) x

def matTransposeMulVec {m n : Nat} (A : Fin m → Fin n → ?) (y : Fin m → ?) :
  Fin n → ? :=
  fun j => ? i : Fin m, y i * A i j

```

```

-- Definition of the Slack Vectors
def primalSlack (A : Fin 3 -> Fin 3 -> ?) (b : Fin 3 -> ?) (x : Fin 3 -> ?) :
  Fin 3 -> ? :=
  fun i => matMulVec A x i - b i

def dualSlack (A : Fin 3 -> Fin 3 -> ?) (c : Fin 3 -> ?) (y : Fin 3 -> ?) :
  Fin 3 -> ? :=
  fun j => c j - matTransposeMulVec A y j

-- 2. Instance Data
def A_inst : Fin 3 -> Fin 3 -> ?
| 0, 0 => 2 | 0, 1 => 1 | 0, 2 => 0
| 1, 0 => 1 | 1, 1 => 2 | 1, 2 => 0
| 2, 0 => 0 | 2, 1 => 0 | 2, 2 => 1

def b_inst : Fin 3 -> ?
| 0 => 3 | 1 => 3 | 2 => 1

def c_inst : Fin 3 -> ?
| 0 => 1 | 1 => 1 | 2 => 2

def x_star : Fin 3 -> ?
| 0 => 1 | 1 => 1 | 2 => 1

def y_star : Fin 3 -> ?
| 0 => 1/3 | 1 => 1/3 | 2 => 2

/-!
### Formal Proofs of the Four Numerical Calculations
-/

-- (1) Primal Cost = 4
theorem primal_cost_eq_four : dot c_inst x_star = 4 := by
  dsimp [dot, c_inst, x_star]
  rw [Fin.sum_univ_three]
  norm_num

-- (2) Dual Cost = 4
theorem dual_cost_eq_four : dot b_inst y_star = 4 := by
  dsimp [dot, b_inst, y_star]
  rw [Fin.sum_univ_three]
  norm_num

-- (3) Primal Slack Vector = 0
theorem primal_slack_is_zero : primalSlack A_inst b_inst x_star = 0 := by
  ext i
  fin_cases i <|> {
    dsimp [primalSlack, matMulVec, dot, A_inst, b_inst, x_star]
    rw [Fin.sum_univ_three]
    norm_num
  }

-- (4) Dual Slack Vector = 0
theorem dual_slack_is_zero : dualSlack A_inst c_inst y_star = 0 := by
  ext j

```

```

fin_cases j <=> {
  dsimp [dualSlack, matTransposeMulVec, A_inst, c_inst, y_star]
  rw [Fin.sum_univ_three]
  norm_num
}

-- Summary: Exact Complementary Slackness & Zero Duality Gap
theorem complementary_slackness_and_optimality :
  dot c_inst x_star = dot b_inst y_star /\
  primalSlack A_inst b_inst x_star = 0 /\
  dualSlack A_inst c_inst y_star = 0 := by
  refine <by rw [primal_cost_eq_four, dual_cost_eq_four],
    primal_slack_is_zero,
    dual_slack_is_zero>

end LPDualityCalculations

end Course21292Numerical

```

A.15 BSinMeasurementTheory/Course21329.lean

Lean 4 source

```

import Mathlib.Order.Zorn
import Mathlib.SetTheory.Cardinal.Basic
import Mathlib.SetTheory.Cardinal.Arithmetic

universe u

namespace Course21329

section FilterExtension

variable {B : Type u} [Lattice B] [BoundedOrder B]

/-- A filter on a bounded lattice `B` is a non-empty, upward-closed set closed under
meets. -/
def IsFilter (F : Set B) : Prop :=
  top in F /\
  (forall ?x y : B?, x in F -> x <= y -> y in F) /\
  (forall ?x y : B?, x in F -> y in F -> x sqcap y in F)

/-- A filter is proper if it does not contain the bottom element `False`. -/
def IsProperFilter (F : Set B) : Prop :=
  IsFilter F /\ False notin F

/-- An ultrafilter is a maximal proper filter under subset inclusion. -/
def IsUltrafilter (U : Set B) : Prop :=
  IsProperFilter U /\ forall ?F : Set B?, IsProperFilter F -> U subseteq F ->
  U = F

/-- Union of a non-empty chain of proper filters is a proper filter. -/
lemma sUnion_chain_isProperFilter (c : Set (Set B))

```

```

(hc_sub : forall F in c, IsProperFilter F)
(hc_chain : IsChain (* subseq *) c)
(hc_ne : c.Nonempty) :
IsProperFilter (?_0 c) := by
refine <<?_, ?_, ?_>, ?_>
-- 1. top in ?_0 c
* rcases hc_ne with <F, hF>
  exact Set.mem_sUnion_of_mem (hc_sub F hF).1.1 hF
-- 2. Upward closed
* intro x y hx hxy
  rcases Set.mem_sUnion.mp hx with <F, hF, hxF>
  exact Set.mem_sUnion_of_mem ((hc_sub F hF).1.2.1 hxF hxy) hF
-- 3. Closed under infima (meets)
* intro x y hx hy
  rcases Set.mem_sUnion.mp hx with <F, hF, hxF>
  rcases Set.mem_sUnion.mp hy with <G, hG, hyG>
  rcases hc_chain.total hF hG with hle | hle
  * exact Set.mem_sUnion_of_mem ((hc_sub G hG).1.2.2 (hle hxF) hyG) hG
  * exact Set.mem_sUnion_of_mem ((hc_sub F hF).1.2.2 hxF (hle hyG)) hF
-- 4. False notin ?_0 c (Properness)
* intro hbot
  rcases Set.mem_sUnion.mp hbot with <F, hF, hbotF>
  exact (hc_sub F hF).2 hbotF

/-- (a) Ultrafilter Extension Theorem:
Every proper filter `F_0` extends to an ultrafilter `U` using Zorn's Lemma. -/
theorem ultrafilter_extension (F_0 : Set B) (hF_0 : IsProperFilter F_0) :
exists U : Set B, IsUltrafilter U /\ F_0 subseq U := by
let S := {F : Set B | IsProperFilter F /\ F_0 subseq F}
have hzorn : forall c subseq S, IsChain (* subseq *) c -> c.Nonempty ->
exists ub in S, forall s in c, s subseq ub := by
intro c hcS hcchain hcne
refine <<?_0 c, <?_, ?_>, fun s hs => Set.subset_sUnion_of_mem hs>
* apply sUnion_chain_isProperFilter c (fun F hF => (hcS hF).1) hcchain hcne
* rcases hcne with <F, hF>
  exact (hcS hF).2.trans (Set.subset_sUnion_of_mem hF)
rcases zorn_subset_nonempty S hzorn F_0 <hF_0, Set.Subset.refl F_0> with <U, hF_0U,
hUmax>
refine <U, <hUmax.1.1, ?_>, hF_0U>
intro G hGprop hUG
have hGS : G in S := <hGprop, hF_0U.trans hUG>
exact Set.Subset.antisymm hUG (hUmax.2 hGS hUG)

/-- The principal filter generated by `top`. -/
def topFilter : Set B := { top }

lemma topFilter_isProperFilter (h_nontriv : (False : B) != top) :
IsProperFilter (topFilter (B := B)) := by
refine <<?_, ?_, ?_>, ?_>
* exact Set.mem_singleton top
* intro x y hx hxy
  rw [topFilter, Set.mem_singleton_iff] at hx |-
  subst hx
  exact top_le_iff.mp hxy
* intro x y hx hy
  rw [topFilter, Set.mem_singleton_iff] at hx hy |-

```

```

    subst hx; subst hy
    rw [inf_idem]
* intro hbot
    rw [topFilter, Set.mem_singleton_iff] at hbot
    exact h_nontriv hbot

/-- (b) If `False != top` (which holds in any infinite Boolean algebra),
the set of ultrafilters is non-empty. -/
theorem stoneSpace_nonempty (h_nontriv : (False : B) != top) :
  exists U : Set B, IsUltrafilter U := by
    rcases ultrafilter_extension (topFilter (B := B)) (topFilter_isProperFilter
      h_nontriv) with <U, hU, _>
    exact <U, hU>

end FilterExtension

section CardinalArithmetic

open Cardinal

/-- (c) Cardinal-arithmetic argument:
If `|B| = ?_0` and `2^?_0 <= |S(B)| <= 2^|B|`, then `|S(B)| = 2^?_0`. -/
theorem ultrafilter_cardinality {S_B : Type*} (card_B : Cardinal)
  (hB : card_B = aleph0)
  (h_upper : #S_B <= 2 ^ card_B)
  (h_lower : 2 ^ aleph0 <= #S_B) :
  #S_B = 2 ^ aleph0 := by
    rw [hB] at h_upper
    exact le_antisymm h_upper h_lower

end CardinalArithmetic

end Course21329

```

A.16 BSinMeasurementTheory/Course21651.lean

Lean 4 source

```

import Mathlib

set_option linter.unusedSectionVars false

open Topology Set

namespace Course21651

namespace StoneRepresentation

variable (B : Type*) [BooleanAlgebra B]

/-!
### Characterization of Boolean Homomorphisms to Bool
-/

```

```

/-- Predicate characterizing Boolean algebra homomorphisms from `B` to `Bool`. -/
def isHom (f : B -> Bool) : Prop :=
  f top = true /\
  f False = false /\
  (forall x y, f (x sqcap y) = (f x && f y)) /\
  (forall x y, f (x sqsup y) = (f x || f y)) /\
  (forall x, f (x?) = (!f x))

/-- The Stone space  $\text{Stone}(B)$  realized as the subtype of homomorphisms
    in  $B \rightarrow \text{Bool}$ . -/
@[reducible]
def StoneSpace : Type _ := { f : B -> Bool // isHom B f }

/-!
### (d) Realization as a Closed Subspace of  $B \rightarrow \text{Bool}$ 
-/

theorem continuous_bool_and (x y : B) :
  Continuous (fun f : B -> Bool => f x && f y) := by
  have : (fun f : B -> Bool => f x && f y) = (fun p : Bool x Bool => p.1 && p.2)
  comp (fun f => (f x, f y)) := rfl
  rw [this]
  exact continuous_of_discreteTopology.comp
    ((@continuous_apply B (fun _ => Bool) _ x).prodMk (@continuous_apply B (fun _ =>
    Bool) _ y))

theorem continuous_bool_or (x y : B) :
  Continuous (fun f : B -> Bool => f x || f y) := by
  have : (fun f : B -> Bool => f x || f y) = (fun p : Bool x Bool => p.1 || p.2)
  comp (fun f => (f x, f y)) := rfl
  rw [this]
  exact continuous_of_discreteTopology.comp
    ((@continuous_apply B (fun _ => Bool) _ x).prodMk (@continuous_apply B (fun _ =>
    Bool) _ y))

theorem continuous_bool_not (x : B) :
  Continuous (fun f : B -> Bool => !f x) := by
  have : (fun f : B -> Bool => !f x) = (! *) comp (fun f => f x) := rfl
  rw [this]
  exact continuous_of_discreteTopology.comp (@continuous_apply B (fun _ => Bool) _ x)

/-- The set of homomorphisms as an explicit intersection of closed equalizer sets in  $B
    \rightarrow \text{Bool}$ . -/
def isHomSet : Set (B -> Bool) :=
  { f | f top = true } ∩
  { f | f False = false } ∩
  { f | f (x sqcap y) = (f x && f y) } ∩
  { f | f (x sqsup y) = (f x || f y) } ∩
  { f | f (x?) = (!f x) }

theorem isHomSet_isClosed : IsClosed (isHomSet B) := by
  dsimp [isHomSet]
  refine IsClosed.inter (IsClosed.inter (IsClosed.inter (IsClosed.inter ?_ ?_) ?_) ?_)
    ?_
  * exact isClosed_eq (@continuous_apply B (fun _ => Bool) _ top) continuous_const

```

```

* exact isClosed_eq (@continuous_apply B (fun _ => Bool) _ False) continuous_const
* refine isClosed_iInter (fun x => isClosed_iInter (fun y => ?_))
  exact isClosed_eq (@continuous_apply B (fun _ => Bool) _ (x sqcap y)) (
    continuous_bool_and B x y)
* refine isClosed_iInter (fun x => isClosed_iInter (fun y => ?_))
  exact isClosed_eq (@continuous_apply B (fun _ => Bool) _ (x sqsup y)) (
    continuous_bool_or B x y)
* refine isClosed_iInter (fun x => ?_)
  exact isClosed_eq (@continuous_apply B (fun _ => Bool) _ (x?)) (
    continuous_bool_not B x)

theorem isHom_iff_mem_isHomSet (f : B -> Bool) :
  isHom B f <-> f in isHomSet B := by
  dsimp [isHom, isHomSet]
  constructor
* intro <ht, hb, hi, hs, hc>
  refine <<<<ht, hb>, ?_>, ?_>, ?_>
  * rw [Set.mem_iInter]; intro x; rw [Set.mem_iInter]; intro y; exact hi x y
  * rw [Set.mem_iInter]; intro x; rw [Set.mem_iInter]; intro y; exact hs x y
  * rw [Set.mem_iInter]; intro x; exact hc x
* intro <<<<ht, hb>, hi>, hs>, hc>
  refine <ht, hb, ?_, ?_, ?_>
  * intro x y; exact (Set.mem_iInter.mp (Set.mem_iInter.mp hi x)) y
  * intro x y; exact (Set.mem_iInter.mp (Set.mem_iInter.mp hs x)) y
  * intro x; exact Set.mem_iInter.mp hc x

/-!
### (a) & (b) Compactness, Hausdorff, and Total Disconnectedness
-/

instance : T2Space (StoneSpace B) := inferInstance

instance : TotallyDisconnectedSpace (StoneSpace B) := inferInstance

instance [Finite B] : CompactSpace (StoneSpace B) := inferInstance

/-!
### (c) Stone Clopen Map and Properties (Stone's Representation Theorem)
-/

/-- The Stone mapping  $\hat{b} = \{u \in X \mid u(b) = \mathrm{true}\}$ . -/
def stoneMap (b : B) : Set (StoneSpace B) :=
  { u | u.1 b = true }

theorem stoneMap_top : stoneMap B top = univ := by
  ext <u, hu>
  change (u top = true) <-> True
  simp [hu.1]

theorem stoneMap_bot : stoneMap B False = {} := by
  ext <u, hu>
  change (u False = true) <-> False
  have h := hu.2.1
  cases hu_bot : u False
  * simp
  * rw [hu_bot] at h; contradiction

```

```

theorem stoneMap_inf (x y : B) :
  stoneMap B (x sqcap y) = stoneMap B x I stoneMap B y := by
  ext <u, hu>
  change (u (x sqcap y) = true) <-> (u x = true /\ u y = true)
  rw [hu.2.2.1 x y, Bool.and_eq_true]

theorem stoneMap_sup (x y : B) :
  stoneMap B (x sqsup y) = stoneMap B x U stoneMap B y := by
  ext <u, hu>
  change (u (x sqsup y) = true) <-> (u x = true \/ u y = true)
  rw [hu.2.2.2.1 x y, Bool.or_eq_true]

theorem stoneMap_compl (x : B) :
  stoneMap B (x?) = (stoneMap B x)? := by
  ext <u, hu>
  change (u (x?) = true) <-> not (u x = true)
  rw [hu.2.2.2.2 x]
  cases u x <|> simp

theorem isClopen_stoneMap (b : B) : IsClopen (stoneMap B b) := by
  have h_cont : Continuous (fun (u : StoneSpace B) => u.1 b) :=
    (@continuous_apply B (fun _ => Bool) _ b).comp continuous_subtype_val
  have h_open : IsOpen (stoneMap B b) := by
    change IsOpen ((fun (u : StoneSpace B) => u.1 b) ^-1' ({true} : Set Bool))
    exact (isOpen_discrete ({true} : Set Bool)).preimage h_cont
  have h_closed : IsClosed (stoneMap B b) := by
    change IsClosed ((fun (u : StoneSpace B) => u.1 b) ^-1' ({true} : Set Bool))
    exact (isClosed_discrete ({true} : Set Bool)).preimage h_cont
  exact <h_closed, h_open>

/-!
### Concrete Numerical Example: B = ?({0, 1})  ~ =  (Fin 2 -> Bool)
-/

abbrev TwoAlgebra := Fin 2 -> Bool

def u0_fun : TwoAlgebra -> Bool := fun s => s 0
def u1_fun : TwoAlgebra -> Bool := fun s => s 1

theorem u0_isHom : isHom TwoAlgebra u0_fun :=
  <rfl, rfl, fun _ _ => rfl, fun _ _ => rfl, fun _ => rfl>

theorem u1_isHom : isHom TwoAlgebra u1_fun :=
  <rfl, rfl, fun _ _ => rfl, fun _ _ => rfl, fun _ => rfl>

def u0 : StoneSpace TwoAlgebra := <u0_fun, u0_isHom>
def u1 : StoneSpace TwoAlgebra := <u1_fun, u1_isHom>

def elem_bot : TwoAlgebra := fun _ => false
def elem_zero : TwoAlgebra := fun i => (i == 0)
def elem_one : TwoAlgebra := fun i => (i == 1)
def elem_top : TwoAlgebra := fun _ => true

-- Verifications of the Stone map on the 4 elements of the Boolean algebra
theorem example_bot : stoneMap TwoAlgebra elem_bot = {} :=

```

```

stoneMap_bot TwoAlgebra

theorem example_top : stoneMap TwoAlgebra elem_top = univ :=
  stoneMap_top TwoAlgebra

theorem example_u0_in_zero : u0 in stoneMap TwoAlgebra elem_zero := rfl

theorem example_u1_not_in_zero : u1 notin stoneMap TwoAlgebra elem_zero := by
  intro h; cases h

theorem example_u1_in_one : u1 in stoneMap TwoAlgebra elem_one := rfl

theorem example_u0_not_in_one : u0 notin stoneMap TwoAlgebra elem_one := by
  intro h; cases h

end StoneRepresentation

end Course21651

```

A.17 BSinMeasurementTheory/Course21720.lean

Lean 4 source

```

import Mathlib.Topology.ContinuousMap.Compact

namespace Course21720

/-!
# Positive Linear Functionals, Riesz Representation, and Stone Duality

This file formalizes:
1. Positivity implies boundedness for linear functionals on  $C(X, \mathbb{R})$  with  $X$  compact.
2. The finite additivity of measures induced on Boolean algebras via Stone duality.
3. Concrete numerical calculations verifying bounds and additivity.
-/

section Theory

variable {X : Type*} [TopologicalSpace X] [CompactSpace X]

/-- A linear functional on  $C(X, \mathbb{R})$  is positive if  $(\forall x, 0 \leq f x) \rightarrow 0 \leq \int f$ . -/
def IsPositiveLinearMap (f : C(X, ℝ) → ℝ) : Prop :=
  forall f : C(X, ℝ), (forall x : X, 0 ≤ f x) → 0 ≤ ∫ f

/-- Part (a): A positive linear functional on  $C(X, \mathbb{R})$  for compact  $X$ 
is bounded by  $\int 1 * f$ . -/
theorem positive_linear_map_bounded (f : C(X, ℝ) → ℝ) (hpos :
  IsPositiveLinearMap f) :
  (f : C(X, ℝ)) : |∫ f| ≤ ∫ 1 * f := by
  have h1 : forall x : X, 0 ≤ (∫ f * (1 : C(X, ℝ)) - f) x := by
    intro x
    rw [ContinuousMap.sub_apply, ContinuousMap.smul_apply, ContinuousMap.one_apply,

```

```

    smul_eq_mul, mul_one]
  have := ContinuousMap.apply_le_norm f x
  linarith
have h2 : forall x : X, 0 <= (?? * (1 : C(X, ?)) + f) x := by
  intro x
  rw [ContinuousMap.add_apply, ContinuousMap.smul_apply, ContinuousMap.one_apply,
    smul_eq_mul, mul_one]
  have := ContinuousMap.neg_norm_le_apply f x
  linarith
have h1 := hpos (?? * (1 : C(X, ?)) - f) h1
have h2 := hpos (?? * (1 : C(X, ?)) + f) h2
rw [map_sub, LinearMap.map_smul, smul_eq_mul, sub_nonneg] at h1
rw [map_add, LinearMap.map_smul, smul_eq_mul, <- sub_le_iff_le_add'] at h2
rw [abs_le]
constructor
* rw [mul_comm]
  linarith
* rw [mul_comm]
  linarith

/-- Part (c): Definition of a finitely additive measure on a Boolean algebra `B`. -/
structure FinitelyAdditiveMeasure (B : Type*) [BooleanAlgebra B] where
  toFun : B -> ?
  nonneg' : forall b, 0 <= toFun b
  bot_zero' : toFun False = 0
  add_disjoint' : forall a b, Disjoint a b -> toFun (a sqsup b) = toFun a + toFun b

/-- Given an assignment of continuous indicator functions `chi : B -> C(X, ?)`
preserving
disjoint sums and the empty set, any positive linear functional induces a finitely
additive measure on `B`. -/
def measureOfStoneRepresentation {B : Type*} [BooleanAlgebra B]
  (chi : B -> C(X, ?))
  (h_bot : chi False = 0)
  (h_disj : forall a b : B, Disjoint a b -> chi (a sqsup b) = chi a + chi b)
  (h_pos_ind : forall b : B, forall x : X, 0 <= chi b x)
  (? : C(X, ?) -> ?[?] ?) (hpos : IsPositiveLinearMap ?) :
  FinitelyAdditiveMeasure B where
toFun b := ? (chi b)
nonneg' b := hpos (chi b) (h_pos_ind b)
bot_zero' := by
  rw [h_bot, map_zero]
add_disjoint' a b h := by
  rw [h_disj a b h, map_add]

end Theory

/-!
### Concrete Numerical Calculations in Lean 4
-/

section ConcreteExample

/-- Linear map instance for the concrete functional `(f) = 3 f(0) + 2 f(1)`. -/
def lambda_linear : (? x ?) -> ?[?] ? where

```

```

toFun f := 3 * f.1 + 2 * f.2
map_add' x y := by
  dsimp
  ring
map_smul' c x := by
  dsimp
  ring

/-- Value of ? on the constant function 1 = (1, 1). -/
theorem concrete_lambda_one : lambda_linear (1, 1) = 5 := by
  dsimp [lambda_linear]
  norm_num

/-- Values on the 4 elements of the Boolean algebra P({0, 1}). -/
theorem concrete_measure_bot : lambda_linear (0, 0) = 0 := by
  dsimp [lambda_linear]
  norm_num

theorem concrete_measure_singleton_0 : lambda_linear (1, 0) = 3 := by
  dsimp [lambda_linear]
  norm_num

theorem concrete_measure_singleton_1 : lambda_linear (0, 1) = 2 := by
  dsimp [lambda_linear]
  norm_num

theorem concrete_measure_top : lambda_linear (1, 1) = 5 := by
  dsimp [lambda_linear]
  norm_num

/-- Verification of finite additivity for disjoint singletons:
  ?({0} ∪ {1}) = ?({0}) + ?({1}) = 3 + 2 = 5. -/
theorem concrete_additivity :
  lambda_linear (1 + 0, 0 + 1) = lambda_linear (1, 0) + lambda_linear (0, 1) := by
  dsimp [lambda_linear]
  norm_num

end ConcreteExample

end Course21720

```

A.18 BSinMeasurementTheory/Course21640.lean

Lean 4 source

```

import Mathlib

namespace Course21640

noncomputable section

/-!
# Dominated Extension, Positivity, and Scott's Theorem Step

```

This file proves the linear-algebraic and order-theoretic steps of the dominated functional extension used to complete Scott's Theorem.

-/

/-- A sublinear functional on a real vector space. -/

```
structure SublinearFunctional (V : Type*) [AddCommGroup V] [Module ? V] where
  toFun : V -> ?
  map_add_le : forall x y : V, toFun (x + y) <= toFun x + toFun y
  map_smul_nonneg : forall (c : ?) (x : V), 0 <= c -> toFun (c * x) = c * toFun x
```

```
instance (V : Type*) [AddCommGroup V] [Module ? V] :
  CoeFun (SublinearFunctional V) (fun _ => V -> ?) where
  coe p := p.toFun
```

/-- Step (b): Given a linear functional `phi` dominated by `p(f) = C \* ?f?` with `0 <= C`,

`phi` is positive on non-negative elements. -/

theorem positive_of_dominated_by_norm

```
{V : Type*} [NormedAddCommGroup V] [NormedSpace ? V]
(phi : V -> ?[?] ?) (one : V) (C : ?) (hC : 0 <= C)
(h_one : phi one = C)
(h_dom : forall f : V, phi f <= C * ?f?)
(f : V) (hf_norm : ??f? * one - f? <= ?f?) :
0 <= phi f := by
have h_le : phi (?f? * one - f) <= C * ??f? * one - f? := h_dom (?f? * one - f)
have h_decomp : phi (?f? * one - f) = ?f? * C - phi f := by
  rw [map_sub, map_smul, h_one, smul_eq_mul]
  rw [h_decomp] at h_le
have h_bound : C * ??f? * one - f? <= C * ?f? :=
  mul_le_mul_of_nonneg_left hf_norm hC
linarith
```

/-!

Concrete Numerical Example

We verify the calculations for $V = \mathbb{R}^2$ with the sup norm (L-infinity).

-/

/-- The sup-norm on $\mathbb{R}^2$. -/

```
def supNorm (v : ? x ?) : ? := max (|v.1|) (|v.2|)
```

/-- The sublinear dominating functional on $\mathbb{R}^2$ associated with weight $C = 1$. -/

```
def p_concrete (v : ? x ?) : ? := supNorm v
```

/-- Subadditivity check for concrete p on $\mathbb{R}^2$. -/

theorem p_concrete_subadditive (u v : ? x ?) :

```
p_concrete (u + v) <= p_concrete u + p_concrete v := by
  dsimp [p_concrete, supNorm]
  apply max_le
  * have h2 : |u.1 + v.1| <= |u.1| + |v.1| := abs_add_le u.1 v.1
    have hu : |u.1| <= max (|u.1|) (|u.2|) := le_max_left _ _
    have hv : |v.1| <= max (|v.1|) (|v.2|) := le_max_left _ _
    linarith
  * have h2 : |u.2 + v.2| <= |u.2| + |v.2| := abs_add_le u.2 v.2
    have hu : |u.2| <= max (|u.1|) (|u.2|) := le_max_right _ _
    have hv : |v.2| <= max (|v.1|) (|v.2|) := le_max_right _ _
```

```

linarith

/-- Positive homogeneity of concrete p. -/
theorem p_concrete_smul (c : ?) (hc : 0 <= c) (v : ? x ?) :
  p_concrete (c * v) = c * p_concrete v := by
  dsimp [p_concrete, supNorm]
  rw [abs_mul, abs_mul, abs_of_nonneg hc]
  exact (mul_max_of_nonneg (|v.1|) (|v.2|) hc).symm

/-- Extension functional phi(x, y) = (1/3) x + (2/3) y. -/
def phi_ext (v : ? x ?) : ? := (1 / 3) * v.1 + (2 / 3) * v.2

/-- Verification that phi extends phi_0 on the diagonal subspace M = span{(1,1)}. -/
theorem phi_ext_extends (c : ?) : phi_ext (c, c) = c := by
  dsimp [phi_ext]
  ring

/-- Verification of dominance for phi: phi(x, y) <= max(|x|, |y|). -/
theorem phi_ext_dominated (x y : ?) : phi_ext (x, y) <= p_concrete (x, y) := by
  dsimp [phi_ext, p_concrete, supNorm]
  have hx : x <= |x| := le_abs_self x
  have hy : y <= |y| := le_abs_self y
  have hx_max : |x| <= max (|x|) (|y|) := le_max_left _ _
  have hy_max : |y| <= max (|x|) (|y|) := le_max_right _ _
  linarith

/-- Positivity verification: if x >= 0 and y >= 0, then phi(x, y) >= 0. -/
theorem phi_ext_positive (x y : ?) (hx : 0 <= x) (hy : 0 <= y) :
  0 <= phi_ext (x, y) := by
  dsimp [phi_ext]
  positivity

end

end Course21640

```

A.19 BSinMeasurementTheory/Course21410.lean

Lean 4 source

```

import Mathlib

open scoped BigOperators Classical

namespace Course21410

namespace ScottMeasurementTheory

/-!
### Finite-Dimensional Step Function Space and Additive Measures
-/

/-- A finitely additive probability measure on subsets of a finite space `a`. -/

```

```

structure FinitelyAdditiveMeasure (a : Type*) [Fintype a] where
  prob : Set a → ?
  prob_nonneg : forall A, 0 ≤ prob A
  prob_univ : prob Set.univ = 1
  prob_empty : prob {} = 0
  prob_disjoint : forall A B, Disjoint A B → prob (A ∪ B) = prob A + prob B

/-- Linear evaluation functional from weight assignments. -/
noncomputable def measureOfWeights {a : Type*} [Fintype a]
  (w : a → ?) (hw_nonneg : forall x, 0 ≤ w x) (hw_sum : ? x, w x = 1) :
    FinitelyAdditiveMeasure a where
  prob A := ? x : a, if x in A then w x else 0
  prob_nonneg A := by
    apply Finset.sum_nonneg
    intro x _
    split_ifs <|> linarith [hw_nonneg x]
  prob_univ := by
    simp [hw_sum]
  prob_empty := by
    simp
  prob_disjoint A B hAB := by
    rw [← Finset.sum_add_distrib]
    apply Finset.sum_congr rfl
    intro x _
    by_cases hA : x in A
    * have hB : x notin B := fun h => Set.disjoint_left.mp hAB hA h
      have hu : x in A ∪ B := Or.inl hA
      split_ifs
      ring
    * by_cases hB : x in B
      * have hu : x in A ∪ B := Or.inr hB
        split_ifs
        ring
      * have hu : x notin A ∪ B := by
          intro h
          rcases h with h1 | h2
          * exact hA h1
          * exact hB h2
        split_ifs
        ring

/-!
### Glue Lemmas: Finite Additivity, Inclusion-Exclusion, and Monotonicity
-/

/-- GLUE LEMMA 1: Inclusion-exclusion for finitely additive measures. -/
theorem measure_union {a : Type*} [Fintype a]
  (? : FinitelyAdditiveMeasure a) (A B : Set a) :
    ?.prob (A ∪ B) + ?.prob (A ∩ B) = ?.prob A + ?.prob B := by
  have h1 : A ∪ B = (A \ B) ∪ B := by
    ext x
    constructor
    * intro h
      rcases h with hA | hB
    * by_cases hB : x in B
      * exact Or.inr hB

```

```

      * exact Or.inl <hA, hB>
      * exact Or.inr hB
    * intro h
      rcases h with <hA, _> | hB
      * exact Or.inl hA
      * exact Or.inr hB
  have h1_disj : Disjoint (A \ B) B := by
    rw [Set.disjoint_left]
    intro x hx
    exact hx.2
  have h2 : A = (A \ B) ∪ (A ∩ B) := by
    ext x
    constructor
    * intro hA
      by_cases hB : x ∈ B
      * exact Or.inr <hA, hB>
      * exact Or.inl <hA, hB>
    * intro h
      rcases h with <hA, _> | <hA, _>
      * exact hA
      * exact hA
  have h2_disj : Disjoint (A \ B) (A ∩ B) := by
    rw [Set.disjoint_left]
    intro x hx hx'
    exact hx.2 hx'.2
  have eq1 : ?.prob (A ∪ B) = ?.prob (A \ B) + ?.prob B := by
    nth_rw 1 [h1]
    exact ?.prob_disjoint (A \ B) B h1_disj
  have eq2 : ?.prob A = ?.prob (A \ B) + ?.prob (A ∩ B) := by
    nth_rw 1 [h2]
    exact ?.prob_disjoint (A \ B) (A ∩ B) h2_disj
  rw [eq1, eq2]
  ring

/-- GLUE LEMMA 2: Monotonicity of finitely additive measures. -/
theorem measure_mono {a : Type*} [Fintype a]
  (w : a → ℝ) (hw_nonneg : forall x, 0 ≤ w x) (hw_sum : ∑ x, w x = 1)
  (A B : Set a) (hAB : A ⊆ B) :
  (measureOfWeights w hw_nonneg hw_sum).prob A ≤ (measureOfWeights w hw_nonneg
    hw_sum).prob B := by
  dsimp [measureOfWeights]
  apply Finset.sum_le_sum
  intro x _
  split_ifs with hA hB
  * exact le_rfl
  * exfalso
    exact hB (hAB hA)
  * exact hw_nonneg x
  * exact le_rfl

/-- GLUE LEMMA 3: Preservation of qualitative preference order by linear functional
restriction. -/
theorem qualitative_agreement {a : Type*} [Fintype a]
  (w : a → ℝ) (hw_nonneg : forall x, 0 ≤ w x) (hw_sum : ∑ x, w x = 1)
  (A B : Set a)

```

```

    (h_diff : 0 <= (? x : a, if x in B then w x else 0) - (? x : a, if x in A
    then w x else 0)) :
    (measureOfWeights w hw_nonneg hw_sum).prob A <= (measureOfWeights w hw_nonneg
    hw_sum).prob B := by
    dsimp [measureOfWeights]
    linarith

/-!
### Concrete Example in Lean 4
-/

def w4 : Fin 4 -> ?
| 0 => 0.1
| 1 => 0.2
| 2 => 0.3
| 3 => 0.4

lemma w4_nonneg : forall x : Fin 4, 0 <= w4 x := by
  intro x
  fin_cases x <|> norm_num [w4]

lemma w4_sum : ? x : Fin 4, w4 x = 1 := by
  rw [Fin.sum_univ_four]
  norm_num [w4]

noncomputable def ?4 : FinitelyAdditiveMeasure (Fin 4) :=
  measureOfWeights w4 w4_nonneg w4_sum

def setA : Set (Fin 4) := {0, 1}
def setB : Set (Fin 4) := {1, 2, 3}

theorem example_probA : ?4.prob setA = 0.3 := by
  dsimp [?4, measureOfWeights, setA]
  rw [Fin.sum_univ_four]
  simp only [Set.mem_insert_iff, Set.mem_singleton_iff, Fin.ext_iff]
  norm_num [w4]

theorem example_probB : ?4.prob setB = 0.9 := by
  dsimp [?4, measureOfWeights, setB]
  rw [Fin.sum_univ_four]
  simp only [Set.mem_insert_iff, Set.mem_singleton_iff, Fin.ext_iff]
  norm_num [w4]

theorem example_prob_inter : ?4.prob (setA ∩ setB) = 0.2 := by
  dsimp [?4, measureOfWeights, setA, setB]
  rw [Fin.sum_univ_four]
  simp only [Set.mem_inter_iff, Set.mem_insert_iff, Set.mem_singleton_iff, Fin.ext_iff]
  ]
  norm_num [w4]

theorem example_prob_union : ?4.prob (setA ∪ setB) = 1.0 := by
  dsimp [?4, measureOfWeights, setA, setB]
  rw [Fin.sum_univ_four]
  simp only [Set.mem_union, Set.mem_insert_iff, Set.mem_singleton_iff, Fin.ext_iff]
  norm_num [w4]

```

```

theorem example_inclusion_exclusion :
  ?4.prob (setA U setB) + ?4.prob (setA I setB) = ?4.prob setA + ?4.prob setB :=
  by
  rw [example_prob_union, example_prob_inter, example_probA, example_probB]
  norm_num

end ScottMeasurementTheory

end Course21410

```

A.20 BSinMeasurementTheory/Course21322.lean

Lean 4 source

```

import Mathlib

open scoped BigOperators Classical

namespace Course21322

namespace FiniteStoneCollapse

set_option linter.unusedSectionVars false

/-!
### Stone Homomorphisms on a Finite Power Set Algebra
-/

variable {a : Type*} [Fintype a] [DecidableEq a]

/-- Predicate characterizing Boolean algebra homomorphisms from `Set a` to `Bool`. -/
def isHom (f : Set a -> Bool) : Prop :=
  f Set.univ = true /\
  f {} = false /\
  (forall x y, f (x I y) = (f x && f y)) /\
  (forall x y, f (x U y) = (f x || f y)) /\
  (forall x, f (x?) = (!f x))

/-- The Stone space  $\mathcal{S}X = \mathrm{Stone}(\mathcal{P}(\alpha))$ . -/
def StoneSpace (a : Type*) [Fintype a] [DecidableEq a] : Type _ :=
  { f : Set a -> Bool // isHom f }

/-- The principal evaluation homomorphism associated with an atom `a : a`. -/
noncomputable def principalHom (a : a) : Set a -> Bool :=
  fun s => decide (a in s)

/-- Verification that `principalHom a` is a genuine Boolean homomorphism. -/
theorem principalHom_isHom (a : a) : isHom (principalHom a) := by
  refine <?_, ?_, ?_, ?_, ?_>
  * dsimp [principalHom]; simp
  * dsimp [principalHom]; simp
  * intro x y; dsimp [principalHom]
  by_cases hx : a in x <;> by_cases hy : a in y <;> simp [hx, hy]

```

```

* intro x y; dsimp [principalHom]
by_cases hx : a in x <;> by_cases hy : a in y <;> simp [hx, hy]
* intro x; dsimp [principalHom]
by_cases hx : a in x <;> simp [hx]

/-- Embedding of atoms into the Stone space. -/
noncomputable def atomToStone (a : a) : StoneSpace a :=
  <principalHom a, principalHom_isHom a>

/-!
### Isomorphism: Atoms ? Stone Ultrafilters
-/

/-- Distinct atoms map to distinct Stone homomorphisms. -/
theorem atomToStone_injective : Function.Injective (atomToStone (a := a)) := by
  intro a b hab
  have hb : b in ({a} : Set a) := by
    have : (atomToStone b).1 ({a} : Set a) = true := by
      rw [← hab]
    simp [atomToStone, principalHom]
  simp [atomToStone, principalHom] using this
  exact (Set.mem_singleton_iff.mp hb).symm

/-- Every Stone homomorphism on `Set a` is the principal evaluation at some atom. -/
theorem stoneToAtom_surjective (u : StoneSpace a) :
  exists a : a, atomToStone a = u := by
  rcases u with <f, hf>
  have h_univ : f Set.univ = true := hf.1
  have h_sum : exists a : a, f {a} = true := by
    by_contra! hall
    have h_zero : forall s : Finset a, f (? a in s, ({a} : Set a)) = false := by
      intro s
      induction s using Finset.induction_on with
      | empty =>
        have : (? a in ({} : Finset a), ({a} : Set a)) = {} := by ext; simp
        rw [this, hf.2.1]
      | @insert x s _ ih =>
        have h_union : (? a in insert x s, ({a} : Set a)) = {x} U (? a in s, ({a} : Set a)) := by
          ext t; simp
        rw [h_union, hf.2.2.2.1 {x} (? a in s, ({a} : Set a))]
        have hx : f {x} = false := Bool.eq_false_iff.mpr (hall x)
        rw [hx, ih]
        rfl
    have h_all : (? a in (Finset.univ : Finset a), ({a} : Set a)) = Set.univ := by
      ext t; simp
    have h_false := h_zero Finset.univ
    rw [h_all] at h_false
    rw [h_false] at h_univ
    contradiction
  rcases h_sum with <a, ha>
  use a
  apply Subtype.ext
  ext s
  dsimp [atomToStone, principalHom]
  by_cases has : a in s

```

```

* have h_le : {a} ⊆ s := Set.singleton_subset_iff.mpr has
have h_inter : {a} ∩ s = {a} := Set.inter_eq_self_of_subset_left h_le
have h_f_inter := hf.2.2.1 {a} s
rw [h_inter, ha] at h_f_inter
have : f s = true := by
  cases hfs : f s
  * rw [hfs, Bool.and_false] at h_f_inter; contradiction
  * rfl
simp [has, this]
* have h_le : {a} ⊆ s? := Set.singleton_subset_iff.mpr has
have h_inter : {a} ∩ s? = {a} := Set.inter_eq_self_of_subset_left h_le
have h_f_inter := hf.2.2.1 {a} (s?)
rw [h_inter, ha] at h_f_inter
have h_compl := hf.2.2.2.2 s
have h_fs_false : f s = false := by
  cases hfs : f s
  * rfl
  * rw [hfs, Bool.not_true] at h_compl
    rw [h_compl, Bool.and_false] at h_f_inter
    contradiction
simp [has, h_fs_false]

/-- Canonical bijection between the atom set and the Stone space of a finite algebra.
-/
noncomputable def atomEquivStone : a ? StoneSpace a :=
  Equiv.ofBijective atomToStone <atomToStone_injective, stoneToAtom_surjective>

/-!
### Identification of Stone Clopens and Atomic Characteristic Sets
-/

/-- The Stone mapping  $\widehat{s} = \{u \in X \mid u(s) = \mathrm{true}\}$ . -/
def stoneMap (s : Set a) : Set (StoneSpace a) :=
  { u | u.1 s = true }

/-- Preservation of representation: an atom `a` belongs to `s` iff its ultrafilter
`atomToStone a` belongs to the Stone clopen `stoneMap s`. -/
theorem atom_mem_iff_stone_mem (s : Set a) (a : a) :
  a in s <=> atomToStone a in stoneMap s := by
  dsimp [atomToStone, principalHom, stoneMap]
  exact decide_eq_true_iff.symm

/-- The preimage of the Stone clopen `stoneMap s` under the atomic isomorphism
is exactly the original subset `s`. -/
theorem stoneMap_preimage_eq (s : Set a) :
  atomToStone ^-1' (stoneMap s) = s := by
  ext a
  rw [Set.mem_preimage]
  exact (atom_mem_iff_stone_mem s a).symm

/-!
### Measure Reduction: Stone Integration to Discrete Weight Sum
-/

/-- Linear evaluation on step functions over the Stone space with weights `w`. -/
noncomputable def stoneMeasure (w : a -> ?) (s : Set a) : ? :=

```

```

    ? a : a, if atomToStone a in stoneMap s then w a else 0

/-- Discrete atomic measure sum from Phase 2 / Phase 3. -/
noncomputable def atomicMeasure (w : a -> ?) (s : Set a) : ? :=
  ? a : a, if a in s then w a else 0

/-- THE COLLAPSE THEOREM: The Stone space measure evaluation reduces identically
    to the finite atomic characteristic sum with ZERO discrepancy. -/
theorem stone_measure_eq_atomic_measure (w : a -> ?) (s : Set a) :
  stoneMeasure w s = atomicMeasure w s := by
  dsimp [stoneMeasure, atomicMeasure]
  apply Finset.sum_congr rfl
  intro a _
  rw [ <- atom_mem_iff_stone_mem ]

/-!
### Concrete Numerical Example: 3-Atom Boolean Algebra
-/

def w3 : Fin 3 -> ?
| 0 => 0.2
| 1 => 0.5
| 2 => 0.3

lemma w3_sum : ? x : Fin 3, w3 x = 1 := by
  rw [Fin.sum_univ_three]
  norm_num [w3]

def testSet : Set (Fin 3) := {0, 2}

/-- Direct calculation of the atomic measure on {0, 2}: 0.2 + 0.3 = 0.5 -/
theorem atomic_test_eval : atomicMeasure w3 testSet = 0.5 := by
  dsimp [atomicMeasure, testSet]
  rw [Fin.sum_univ_three]
  simp only [Set.mem_insert_iff, Set.mem_singleton_iff, Fin.ext_iff]
  norm_num [w3]

/-- Calculation via the Stone representation map on {0, 2}: 0.2 + 0.3 = 0.5 -/
theorem stone_test_eval : stoneMeasure w3 testSet = 0.5 := by
  rw [stone_measure_eq_atomic_measure]
  exact atomic_test_eval

end FiniteStoneCollapse

end Course21322

```

Total: 20 modules, 2723 lines of Lean.