

# Formalization of a BS in Measurement Theory

**Lars Warren Ericson**

Catskills Research Company

September 2026

CMU-CS-26-XXX

School of Computer Science

Carnegie Mellon University

Pittsburgh, PA 15213

Source repository: <https://github.com/catskillsresearch/fabbro>

## Abstract

This report presents a reconstructed Carnegie Mellon undergraduate path through the Mathematical Sciences 21-xxx catalog, sequenced so that Dana Scott’s 1964 measurement theorems (1.1–4.1) and their extension to infinite Boolean algebras can be read for understanding after these courses are taken and proved end to end. Each course contributes one problem that requires the full toolkit of that course rather than a single technique in isolation. Solutions are written in English and then formalized in Lean 4 / Mathlib. The accompanying library `BSinMeasurementTheory` is sorry-free under the pinned toolchain `leanprover/lean4:v4.34.0-rc1` and introduces no project axioms beyond Mathlib’s classical footprint. Large-language-model assistance was used in drafting solutions, but every accepted declaration is checked by Lean’s kernel. The complete source and a reproducible build pipeline are publicly available with this report.

Copyright © 2026 Lars Warren Ericson

**Keywords:** Lean 4; formal verification; measurement theory; undergraduate curriculum; Scott 1964; Boolean algebras; Stone spaces

# Contents

|           |                                                                                            |           |
|-----------|--------------------------------------------------------------------------------------------|-----------|
| <b>1</b>  | <b>Introduction</b>                                                                        | <b>6</b>  |
| <b>2</b>  | <b>Syllabus</b>                                                                            | <b>7</b>  |
| <b>3</b>  | <b>Phase 1: Language and linear structure</b>                                              | <b>11</b> |
| 3.1       | 21-120/21-122 — Differential and Integral Calculus / Integration and Approximation . . . . | 11        |
| 3.1.1     | Problem . . . . .                                                                          | 11        |
| 3.1.2     | Solution . . . . .                                                                         | 11        |
| 3.1.2.a   | Differentiability and Derivative . . . . .                                                 | 11        |
| 3.1.2.b   | Existence of $\lim_{x \rightarrow \infty} f(x)$ and Tail Bound . . . . .                   | 13        |
| 3.1.2.b.1 | Existence . . . . .                                                                        | 13        |
| 3.1.2.b.2 | Tail bound via integration by parts . . . . .                                              | 13        |
| 3.1.2.c   | Taylor Series and Numerical Estimation of $f(0.5)$ . . . . .                               | 14        |
| 3.1.2.c.1 | Taylor expansion . . . . .                                                                 | 14        |
| 3.1.2.c.2 | Estimation at $x = 0.5$ . . . . .                                                          | 15        |
| 3.1.2.c.3 | Explicit error bound . . . . .                                                             | 16        |
| 3.1.2.c.4 | Summary of what Lean verifies here . . . . .                                               | 17        |
| 3.2       | 21-127 — Concepts of Mathematics . . . . .                                                 | 17        |
| 3.2.1     | Problem . . . . .                                                                          | 17        |
| 3.2.2     | Solution . . . . .                                                                         | 18        |
| 3.2.2.a   | Equivalence Relation . . . . .                                                             | 18        |
| 3.2.2.b   | Bijection with $\mathbb{Q}$ . . . . .                                                      | 19        |
| 3.2.2.c   | The Flaw in the Claim and the Correct Statement . . . . .                                  | 21        |
| 3.3       | 21-241 — Matrices and Linear Transformations . . . . .                                     | 23        |
| 3.3.1     | Problem . . . . .                                                                          | 23        |
| 3.3.2     | Solution . . . . .                                                                         | 24        |
| 3.3.2.a   | Basis and Dual Basis Construction . . . . .                                                | 24        |
| 3.3.2.b   | Inner Product Representation and Concrete Separation . . . . .                             | 27        |
| 3.3.2.b.1 | Choice of point $v$ not in $C$ . . . . .                                                   | 29        |
| 3.3.2.b.2 | Separating vector $u$ . . . . .                                                            | 30        |
| 3.3.2.b.3 | Inner product value at $v$ . . . . .                                                       | 30        |
| 3.3.2.b.4 | Inner product on an arbitrary $c$ in $C$ . . . . .                                         | 31        |
| 3.3.2.b.5 | Linear functional separation . . . . .                                                     | 31        |
| 3.3.2.c   | The Dual Cone and its Dual Basis Representation . . . . .                                  | 32        |
| 3.4       | 21-228 — Discrete Mathematics . . . . .                                                    | 34        |
| 3.4.1     | Problem . . . . .                                                                          | 34        |
| 3.4.2     | Solution . . . . .                                                                         | 36        |
| 3.4.2.a   | Symmetric conditions depend only on the multiset . . . . .                                 | 36        |
| 3.4.2.a.1 | Proof: . . . . .                                                                           | 36        |
| 3.4.2.b   | Counting permutation pairs . . . . .                                                       | 37        |
| 3.4.2.b.1 | Genericity Assumptions: . . . . .                                                          | 38        |
| 3.4.2.b.2 | Derivation of the Exact Formula: . . . . .                                                 | 38        |
| 3.4.2.c   | Reduction to $n = 2$ and $n = 3$ . . . . .                                                 | 40        |
| 3.4.2.c.1 | Case $n = 2$ . . . . .                                                                     | 41        |

|           |                                                                                          |           |
|-----------|------------------------------------------------------------------------------------------|-----------|
| 3.4.2.c.2 | Case $n = 3$                                                                             | 42        |
| <b>4</b>  | <b>Phase 2: Analysis, algebras, and formal proof</b>                                     | <b>44</b> |
| 4.1       | 21-355 — Principles of Real Analysis I                                                   | 44        |
| 4.1.1     | Problem                                                                                  | 44        |
| 4.1.2     | Solution                                                                                 | 44        |
| 4.1.2.a   | Bolzano–Weierstrass Theorem from the Completeness Axiom                                  | 44        |
| 4.1.2.a.1 | Axiom Used (Completeness / Least Upper Bound Property)                                   | 44        |
| 4.1.2.a.2 | Lemma 1 (Monotone Convergence Theorem)                                                   | 44        |
| 4.1.2.a.3 | Lemma 2 (Monotone Subsequence / Peak Points)                                             | 46        |
| 4.1.2.a.4 | Derivation of Bolzano–Weierstrass                                                        | 46        |
| 4.1.2.b   | Existence and Characterization of $\limsup x_n$ and $\liminf x_n$                        | 47        |
| 4.1.2.b.1 | Monotonicity of the Tail Extremities                                                     | 47        |
| 4.1.2.b.2 | Existence of Limits                                                                      | 48        |
| 4.1.2.c   | Example where $\limsup x_n \neq \liminf x_n$ and $\varepsilon$ -Proof of Divergence      | 48        |
| 4.1.2.c.1 | Computation of $\limsup$ and $\liminf$                                                   | 49        |
| 4.1.2.c.2 | Direct $\varepsilon$ -Proof of Divergence                                                | 50        |
| 4.1.2.d   | Heine–Borel: $[a, b]$ is Compact Directly from Open Covers                               | 51        |
| 4.2       | 21-373 — Algebraic Structures                                                            | 53        |
| 4.2.1     | Problem                                                                                  | 53        |
| 4.2.2     | Solution                                                                                 | 53        |
| 4.2.2.a   | Every element is a unique join of a subset of the atoms                                  | 53        |
| 4.2.2.b   | Isomorphism with $\mathcal{P}(\{a_1, \dots, a_n\})$ and identification with $\{0, 1\}^n$ | 56        |
| 4.2.2.c   | Action of a finite group $G$ by automorphisms                                            | 58        |
| 4.2.2.d   | Numerical / Computational Example                                                        | 61        |
| 4.3       | 21-321 — Interactive Theorem Proving                                                     | 62        |
| 4.3.1     | Problem                                                                                  | 62        |
| 4.3.2     | Solution                                                                                 | 62        |
| 4.3.2.a   | Scott pairs and separating functionals                                                   | 62        |
| 4.3.2.b   | Theorem 1.1 (separating hyperplane / geometric form)                                     | 65        |
| 4.3.2.c   | Theorem 1.2 (combinatorial / rational cancellation)                                      | 66        |
| 4.3.2.d   | Theorem 1.3 (equivalence of the rational and combinatorial forms)                        | 68        |
| 4.3.2.e   | Kraft–Pratt–Seidenberg violation                                                         | 68        |
| <b>5</b>  | <b>Phase 3: Duality and set theory</b>                                                   | <b>71</b> |
| 5.1       | 21-292 — Operations Research I                                                           | 71        |
| 5.1.1     | Problem                                                                                  | 71        |
| 5.1.2     | Solution                                                                                 | 71        |
| 5.1.2.a   | Linear Programming Duality: State and Proof                                              | 72        |
| 5.1.2.a.1 | Weak Duality Theorem                                                                     | 72        |
| 5.1.2.a.2 | Strong Duality Theorem                                                                   | 72        |
| 5.1.2.b   | Concrete $3 \times 3$ Instance & Hand Simplex Execution                                  | 74        |
| 5.1.2.b.1 | Dual Formulation:                                                                        | 75        |
| 5.1.2.b.2 | Step-by-Step Simplex Iteration:                                                          | 75        |
| 5.1.2.c   | Derivation of Scott’s Theorem 1.1 / Kuhn-Tucker Separation from LP Duality               | 77        |

|          |               |                                                                              |            |
|----------|---------------|------------------------------------------------------------------------------|------------|
|          | 5.1.2.c.1     | Proof using LP Strong Duality: . . . . .                                     | 77         |
|          | 5.1.2.d       | Numerical Calculations . . . . .                                             | 77         |
| 5.2      | 21-329        | — Set Theory . . . . .                                                       | 80         |
|          | 5.2.1         | Problem . . . . .                                                            | 80         |
|          | 5.2.2         | Solution . . . . .                                                           | 80         |
|          | 5.2.2.a       | Ultrafilter Extension Theorem via Zorn's Lemma . . . . .                     | 80         |
|          | 5.2.2.b       | Non-emptiness of the Stone Space . . . . .                                   | 83         |
|          | 5.2.2.c       | Cardinality of Ultrafilters on a Countably Infinite Boolean Algebra . . . .  | 84         |
| <b>6</b> |               | <b>Phase 4: Stone spaces and measures</b>                                    | <b>86</b>  |
| 6.1      | 21-651        | — General Topology . . . . .                                                 | 86         |
|          | 6.1.1         | Problem . . . . .                                                            | 86         |
|          | 6.1.2         | Solution . . . . .                                                           | 86         |
|          | 6.1.2.a       | Setup and Topological Realization . . . . .                                  | 87         |
|          | 6.1.2.b       | Realization as a Closed Subspace of $2^B$ & (a) Compactness . . . . .        | 87         |
|          | 6.1.2.c       | Hausdorff and Totally Disconnected . . . . .                                 | 89         |
|          | 6.1.2.d       | Clopen Sets & Stone's Representation Theorem . . . . .                       | 90         |
|          | 6.1.2.e       | Concrete Numerical Example . . . . .                                         | 92         |
| 6.2      | 21-720        | — Measure and Integration . . . . .                                          | 93         |
|          | 6.2.1         | Problem . . . . .                                                            | 93         |
|          | 6.2.2         | Solution . . . . .                                                           | 93         |
|          | 6.2.2.a       | Boundedness of $\Lambda$ . . . . .                                           | 94         |
|          | 6.2.2.b       | Riesz Representation Theorem . . . . .                                       | 95         |
|          | 6.2.2.c       | Specialization to the Stone Space of a Boolean Algebra . . . . .             | 97         |
|          | 6.2.2.d       | Concrete Numerical Example . . . . .                                         | 98         |
| <b>7</b> |               | <b>Phase 5: The missing extension</b>                                        | <b>100</b> |
| 7.1      | 21-640        | — Introduction to Functional Analysis . . . . .                              | 100        |
|          | 7.1.1         | Problem . . . . .                                                            | 100        |
|          | 7.1.2         | Solution . . . . .                                                           | 100        |
|          | 7.1.2.a       | Hahn–Banach Theorem (Dominated Extension Form) . . . . .                     | 100        |
|          | 7.1.2.a.1     | Precise Statement . . . . .                                                  | 100        |
|          | 7.1.2.a.2     | Role of the Sublinear Functional . . . . .                                   | 101        |
|          | 7.1.2.b       | Extension of $\varphi_0$ to a Positive Linear Functional on $C(X)$ . . . . . | 101        |
|          | 7.1.2.b.1     | Step 1: Definition of the Sublinear Functional . . . . .                     | 101        |
|          | 7.1.2.b.2     | Step 2: Dominance on the Subspace $M$ . . . . .                              | 102        |
|          | 7.1.2.b.3     | Step 3: Application of Hahn–Banach . . . . .                                 | 102        |
|          | 7.1.2.b.4     | Step 4: Positivity of the Extension $\varphi$ . . . . .                      | 102        |
|          | 7.1.2.c       | Completing Scott's Theorem 4.1 via Riesz Representation . . . . .            | 103        |
|          | 7.1.2.d       | Concrete Illustrative Example . . . . .                                      | 104        |
| <b>8</b> |               | <b>Phase 6: Assembly and consistency check</b>                               | <b>107</b> |
| 8.1      | 21-410/21-599 | — Independent Study . . . . .                                                | 107        |
|          | 8.1.1         | Problem . . . . .                                                            | 107        |
|          | 8.1.2         | Solution . . . . .                                                           | 107        |
|          | 8.1.2.a       | Extension of Scott's Theorem 4.1 to Infinite Boolean Algebras . . . . .      | 107        |

|           |                                                                        |            |
|-----------|------------------------------------------------------------------------|------------|
| 8.1.2.a.1 | Setup and Scott Cancellation Condition . . . . .                       | 107        |
| 8.1.2.a.2 | Stone Space Construction . . . . .                                     | 109        |
| 8.1.2.a.3 | Vector Space of Step Functions and the Preference Cone . . . . .       | 109        |
| 8.1.2.a.4 | Hahn–Banach / M. Riesz Positive Extension . . . . .                    | 110        |
| 8.1.2.a.5 | Restriction back to $B$ . . . . .                                      | 110        |
| 8.1.2.b   | Finite Additivity vs. Countable Additivity . . . . .                   | 110        |
| 8.1.2.b.1 | The Exact Point of the Obstruction . . . . .                           | 110        |
| 8.1.2.c   | Concrete Worked Example . . . . .                                      | 112        |
| 8.1.2.c.1 | English Calculation . . . . .                                          | 112        |
| 8.1.2.d   | Complete Lean 4 Formalization . . . . .                                | 114        |
| 8.2       | 21-322 — Topics in Formal Mathematics (opportunistic) . . . . .        | 114        |
| 8.2.1     | Problem . . . . .                                                      | 114        |
| 8.2.2     | Solution . . . . .                                                     | 114        |
| 8.2.2.a   | The Finite-to-Infinite Duality Gap and the Reduction Problem . . . . . | 115        |
| 8.2.2.b   | Concrete Example in English . . . . .                                  | 119        |
| <b>9</b>  | <b>References</b>                                                      | <b>121</b> |
| <b>A</b>  | <b>Lean module index</b>                                               | <b>122</b> |

## List of Figures

|    |                                                                                                                                                                                     |     |
|----|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----|
| 1  | Conceptual buildup of the reconstructed 21-xxx syllabus toward Scott (1964). Blue nodes are courses, grouped by phase. Course nodes are hyperlinked to the corresponding section. . | 9   |
| 2  | Mathlib subsections required to read each course. Orange nodes are Mathlib; blue nodes are courses. Course nodes are hyperlinked to the corresponding section. . . . .              | 10  |
| 3  | Lean module dependencies for the 21-120 / 21-122 solution. Amber box: Mathlib imports. Blue box: solution modules. . . . .                                                          | 12  |
| 4  | Lean module dependencies for the 21-127 solution. Amber box: Mathlib imports. Blue box: solution modules. . . . .                                                                   | 18  |
| 5  | Lean module dependencies for the 21-241 solution. Amber box: Mathlib imports. Blue box: solution modules. . . . .                                                                   | 24  |
| 6  | Lean module dependencies for the 21-228 solution. Amber box: Mathlib imports. Blue box: solution modules. . . . .                                                                   | 35  |
| 7  | Lean module dependencies for the 21-355 solution. Amber box: Mathlib imports. Blue box: solution modules. . . . .                                                                   | 45  |
| 8  | Lean module dependencies for the 21-373 solution. Amber box: Mathlib imports. Blue box: solution modules. . . . .                                                                   | 53  |
| 9  | Lean module dependencies for the 21-321 solution. Amber box: Mathlib imports. Blue box: solution modules. . . . .                                                                   | 63  |
| 10 | Lean module dependencies for the 21-292 solution. Amber box: Mathlib imports. Blue box: solution modules. . . . .                                                                   | 71  |
| 11 | Lean module dependencies for the 21-329 solution. Amber box: Mathlib imports. Blue box: solution modules. . . . .                                                                   | 80  |
| 12 | Lean module dependencies for the 21-651 solution. Amber box: Mathlib imports. Blue box: solution modules. . . . .                                                                   | 86  |
| 13 | Lean module dependencies for the 21-720 solution. Amber box: Mathlib imports. Blue box: solution modules. . . . .                                                                   | 94  |
| 14 | Lean module dependencies for the 21-640 solution. Amber box: Mathlib imports. Blue box: solution modules. . . . .                                                                   | 101 |
| 15 | Lean module dependencies for the 21-410 / 21-599 solution. Amber box: Mathlib imports. Blue box: solution modules. . . . .                                                          | 108 |
| 16 | Lean module dependencies for the 21-322 solution. Amber box: Mathlib imports. Blue box: solution modules. . . . .                                                                   | 114 |

# 1 Introduction

Imagine that a first-year student at Carnegie Mellon told their faculty advisor that their goal for their undergraduate degree was to be able to read a paper on Measurement Theory written by Dana Scott in 1964. This report is an example of responsible answer to that question, generated with help of large language models. It turns out to be a map of Scott’s own undergraduate training. It is a model of highly individually tailored automated advisement to ensure that the student will master the skills needed to read the paper for understanding and be able, in turn, to reproduce and teach the ideas and proofs it contains to themselves and others. This is a level of advising intended to augment and improve on typical standards of advice received by undergraduates. The selection of Scott’s 1964 paper is not intended to recommend an actual degree program in Measurement Theory, but to illustrate this unique AI-assisted advisement tool. The tool idea is not a criticism of current advisement practices, just a unique way to augment it.

Measurement theory asks when qualitative comparisons can be represented by real numbers. Scott’s 1964 paper organizes several such representation questions around a finite theory of homogeneous linear inequalities, with applications to intransitive indifference, ordered utility differences, and subjective probability. The infinite direction, which Scott announced but did not carry out, runs through Stone spaces, positive functionals, Hahn–Banach extension, and Riesz representation.

The pedagogical claim of this report is that a complete undergraduate mathematics major at Carnegie Mellon already contains the pieces of that argument, if the courses are taken in the right order and each is asked to do real work. The syllabus below is that ordering. It is framed as a Bachelor of Science in Measurement Theory: not a new degree program, but a reading of the existing catalog as a single connected proof.

Each problem is built so that quoting a theorem from a later course is not available: the calculus problem uses the fundamental theorem, comparison, and Taylor remainders; the linear-algebra problem speaks the dual-space language of Scott’s Theorem 1.1; the discrete-mathematics problem is the combinatorial counting behind Theorems 1.2 and 3.2; and the independent-study problem assembles set theory, topology, measure, and functional analysis into the infinite extension of Theorem 4.1.

The formalizations live in the Lean library `BSinMeasurementTheory`. Each course snippet is a self-contained module. The root file `BSinMeasurementTheory.lean` imports all of them. The English arguments occupy the body of this report, in syllabus order, with each Lean module inlined next to the claim it proves. The appendix is an index of filenames, each linking to the source on GitHub.

## 2 Syllabus

| Phase | Course          | Title                                                              | Role toward Scott (1964)                                                 |
|-------|-----------------|--------------------------------------------------------------------|--------------------------------------------------------------------------|
| 1     | 21-120 / 21-122 | Differential and Integral Calculus / Integration and Approximation | FTC, comparison, integration by parts, Taylor remainder                  |
| 1     | 21-127          | Concepts of Mathematics                                            | Equivalence relations, explicit bijections, induction                    |
| 1     | 21-241          | Matrices and Linear Transformations                                | Bases, dual bases, and the vector-space language of Theorem 1.1          |
| 1     | 21-228          | Discrete Mathematics                                               | Permutations, multisets, and the counting behind Theorems 1.2 and 3.2    |
| 2     | 21-355          | Principles of Real Analysis I                                      | Completeness, limsup/liminf, and compactness from axioms                 |
| 2     | 21-373          | Algebraic Structures                                               | Atoms and characteristic functions of finite Boolean algebras            |
| 2     | 21-321          | Interactive Theorem Proving                                        | Formal statements of Theorems 1.1 and 1.2, and their equivalence         |
| 3     | 21-292          | Operations Research I                                              | Simplex and LP duality, deriving Kuhn–Tucker / Theorem 1.1 from the dual |
| 3     | 21-329          | Set Theory                                                         | Zorn’s lemma, ultrafilters, and cardinal arithmetic                      |
| 4     | 21-651          | General Topology                                                   | Stone spaces, compactness via Tychonoff, Stone representation            |

| Phase | Course          | Title                               | Role toward Scott (1964)                                                       |
|-------|-----------------|-------------------------------------|--------------------------------------------------------------------------------|
| 4     | 21-720          | Measure and Integration             | Riesz representation and the induced finitely additive measure on $B$          |
| 5     | 21-640          | Introduction to Functional Analysis | Hahn–Banach extension, the step Scott cites and leaves unproved                |
| 6     | 21-410 / 21-599 | Independent Study                   | Full reconstruction of the infinite-algebra extension of Theorem 4.1           |
| 6     | 21-322          | Topics in Formal Mathematics        | Proof that the finite atomic picture is what the infinite machinery reduces to |

The conceptual buildup of that sequence is Figure 1. The Mathlib subsections required to read each course’s Lean are Figure 2. The Lean library does not import across courses.

Every course problem is solved in two layers: an English mathematical argument, and a Lean 4 formalization using Mathlib, with numerical examples re-checked in Lean when they appear. Course 21-322 is opportunistic: it should be taken in a term that overlaps real analysis or order theory in Lean, but the standing problem is independent of that term’s advertised topic.

The remainder of this report is the problem set itself, in syllabus order.




Figure 1: Conceptual buildup of the reconstructed 21-xxx syllabus toward Scott (1964). Blue nodes are courses, grouped by phase. Course nodes are hyperlinked to the corresponding section.




Figure 2: Mathlib subsections required to read each course. Orange nodes are Mathlib; blue nodes are courses. Course nodes are hyperlinked to the corresponding section.

### 3 Phase 1: Language and linear structure

Calculus, proof language, dual bases, and combinatorial cancellation.

#### 3.1 21-120/21-122 — Differential and Integral Calculus / Integration and Approximation

##### 3.1.1 Problem

Let  $f(x) = \int_0^x \frac{\sin(t^2)}{1+t^2} dt$ .

- (a) Show  $f$  is differentiable everywhere and find  $f'(x)$ .
- (b) Determine whether  $\lim_{x \rightarrow \infty} f(x)$  exists, and if so, bound it using comparison and integration by parts on the tail.
- (c) Find the Taylor series of  $f$  about  $x = 0$  through the  $x^9$  term, and use it to estimate  $f(0.5)$  to four decimal places with an explicit error bound.

*Requires: FTC, limits/comparison tests, integration by parts, power series, Taylor remainder estimation.*

##### 3.1.2 Solution

###### 3.1.2.a Differentiability and Derivative

Let  $g(t) = \frac{\sin(t^2)}{1+t^2}$ . Since  $t \mapsto \sin(t^2)$  and  $t \mapsto 1+t^2$  are continuous on  $\mathbb{R}$  and  $1+t^2 \geq 1 > 0$  for all  $t \in \mathbb{R}$ ,  $g(t)$  is everywhere continuous. By the Fundamental Theorem of Calculus:

$$f(x) = \int_0^x g(t) dt$$

is differentiable for all  $x \in \mathbb{R}$ , and:

$$f'(x) = g(x) = \frac{\sin(x^2)}{1+x^2}.$$

##### Lean 4 source

```
import Mathlib

open Real intervalIntegral MeasureTheory

namespace Course21120FTC

/-- Definition of f(x) = ?_0? sin(t?) / (1 + t?) dt -/
noncomputable def f (x : ?) : ? := ? t in (0 : ?)..x, sin (t ^ 2) / (1 + t ^ 2)

/-- The integrand is continuous everywhere on ?. -/
lemma continuous_integrand : Continuous (fun t : ? => sin (t ^ 2) / (1 + t ^ 2)) := by
  apply Continuous.div
  * exact continuous_sin.comp (continuous_pow 2)
  * exact continuous_const.add (continuous_pow 2)
  * intro t
    have h : 0 <= t ^ 2 := sq_nonneg t
    linarith
```




Figure 3: Lean module dependencies for the 21-120 / 21-122 solution. Amber box: Mathlib imports. Blue box: solution modules.

```

/-- Fundamental theorem of calculus derivative statement for f. -/
lemma f_hasDerivAt (x : ?) : HasDerivAt f (sin (x ^ 2) / (1 + x ^ 2)) x :=
  HasStrictDerivAt.hasDerivAt (continuous_integrand.integral_hasStrictDerivAt 0 x)

/-- (a) Part 1: f is differentiable everywhere on ?. -/
theorem part_a_differentiable (x : ?) : DifferentiableAt ? f x :=
  HasDerivAt.differentiableAt (f_hasDerivAt x)

/-- (a) Part 2: f'(x) = sin(x?) / (1 + x?). -/
theorem part_a_deriv (x : ?) : deriv f x = sin (x ^ 2) / (1 + x ^ 2) :=
  HasDerivAt.deriv (f_hasDerivAt x)

end Course21120FTC

```

### 3.1.2.b Existence of $\lim_{x \rightarrow \infty} f(x)$ and Tail Bound

#### 3.1.2.b.1 Existence

For all  $t \geq 0$ ,  $|\sin(t^2)| \leq 1$ , so

$$\left| \frac{\sin(t^2)}{1+t^2} \right| \leq \frac{1}{1+t^2}.$$

Since  $\int_0^\infty \frac{dt}{1+t^2} = [\arctan(t)]_0^\infty = \frac{\pi}{2} < \infty$ , the improper integral  $\int_0^\infty \frac{\sin(t^2)}{1+t^2} dt$  is absolutely convergent. Hence,  $\lim_{x \rightarrow \infty} f(x)$  exists.

#### Lean 4 source

```

import Mathlib

open Real

namespace Course21120

/-- Pointwise comparison: the integrand is dominated by  $1/(1+t^2)$ ,
which is the majorant used to get absolute convergence of the improper
integral. -/
theorem abs_integrand_le (t : ?) :
  |sin (t ^ 2) / (1 + t ^ 2)| <= 1 / (1 + t ^ 2) := by
  have hpos : 0 < 1 + t ^ 2 := by nlinarith [sq_nonneg t]
  rw [abs_div, abs_of_pos hpos]
  exact div_le_div_of_nonneg_right (abs_sin_le_one _) hpos.le

end Course21120

```

#### 3.1.2.b.2 Tail bound via integration by parts

For any cutoff  $M > 0$ , decompose the limit into:

$$\lim_{x \rightarrow \infty} f(x) = \int_0^M \frac{\sin(t^2)}{1+t^2} dt + \int_M^\infty \frac{\sin(t^2)}{1+t^2} dt.$$

On the tail, rewrite  $\sin(t^2) dt = \frac{1}{2t} d(-\cos(t^2))$ . Integrating by parts with  $u = \frac{1}{2t(1+t^2)}$  and  $dv = 2t \sin(t^2) dt$ :

$$\begin{aligned} \int_M^\infty \frac{\sin(t^2)}{1+t^2} dt &= \left[ -\frac{\cos(t^2)}{2t(1+t^2)} \right]_M^\infty - \int_M^\infty (-\cos(t^2)) \left( -\frac{1+3t^2}{2t^2(1+t^2)^2} \right) dt \\ &= \frac{\cos(M^2)}{2M(1+M^2)} - \int_M^\infty \cos(t^2) \frac{1+3t^2}{2t^2(1+t^2)^2} dt. \end{aligned}$$

Since  $|\cos(t^2)| \leq 1$  and  $\frac{1+3t^2}{2t^2(1+t^2)^2} = -\frac{d}{dt} \left( \frac{1}{2t(1+t^2)} \right) > 0$ :

$$\left| \int_M^\infty \frac{\sin(t^2)}{1+t^2} dt \right| \leq \frac{1}{2M(1+M^2)} + \int_M^\infty \left( -\frac{d}{dt} \frac{1}{2t(1+t^2)} \right) dt = \frac{1}{M(1+M^2)}.$$

Combining with the comparison bound on  $[0, M]$ :

$$\left| \lim_{x \rightarrow \infty} f(x) \right| \leq \arctan(M) + \frac{1}{M(1+M^2)}.$$

For example, setting  $M = 1$  yields  $|f(\infty)| \leq \frac{\pi}{4} + \frac{1}{2} \approx 1.2854$ .

### 3.1.2.c Taylor Series and Numerical Estimation of $f(0.5)$

#### Lean 4 source

```
import Mathlib
import BSinMeasurementTheory.Course21120.Existence

open Real intervalIntegral

namespace Course21120

/-- The comparison majorant integrates to an arctan increment. -/
theorem majorant_integral (M x : ?) :
  ? t in M..x, (1 : ?) / (1 + t ^ 2) = arctan x - arctan M := by
  simp [one_div, integral_inv_one_add_sq]

/-- Remaining majorant mass on  $[M, x]$  is at most the improper tail
 $\pi/2 - \arctan M$ . -/
theorem tail_mass_le (M x : ?) (_hMx : M <= x) :
  arctan x - arctan M <= Real.pi / 2 - arctan M := by
  nlinarith [arctan_lt_pi_div_two x]

end Course21120
```

#### 3.1.2.c.1 Taylor expansion

Expanding the integrand around  $t = 0$ :

$$\sin(t^2) = t^2 - \frac{t^6}{6} + O(t^{10}), \quad \frac{1}{1+t^2} = 1 - t^2 + t^4 - t^6 + O(t^8)$$

Multiplying through terms up to degree 8:

$$g(t) = \left(t^2 - \frac{t^6}{6}\right)(1 - t^2 + t^4 - t^6) + O(t^{10}) = t^2 - t^4 + \frac{5}{6}t^6 - \frac{5}{6}t^8 + O(t^{10}).$$

Integrating term-by-term from 0 to  $x$ :

$$f(x) = \frac{x^3}{3} - \frac{x^5}{5} + \frac{5}{42}x^7 - \frac{5}{54}x^9 + O(x^{11}).$$

### Lean 4 source

```
import Mathlib

namespace Course21120Taylor

/-- Degree-9 Taylor polynomial of the integrated expansion
$t^2 - t^4 + (5/6)t^6 - (5/6)t^8$. -/
def T9 (x : ?) : ? :=
  x ^ 3 / 3 - x ^ 5 / 5 + 5 * x ^ 7 / 42 - 5 * x ^ 9 / 54

end Course21120Taylor
```

#### 3.1.2.c.2 Estimation at $x = 0.5$

$$\begin{aligned}\frac{(0.5)^3}{3} &= \frac{1}{24} \approx 0.041667 \\ -\frac{(0.5)^5}{5} &= -\frac{1}{160} = -0.006250 \\ \frac{5(0.5)^7}{42} &= \frac{5}{5376} \approx 0.000930 \\ -\frac{5(0.5)^9}{54} &= -\frac{5}{27648} \approx -0.000181\end{aligned}$$

Summing these gives:

$$T_9(0.5) \approx 0.041667 - 0.006250 + 0.000930 - 0.000181 = 0.036166 \approx 0.0362.$$

### Lean 4 source

```
import Mathlib
import BSinMeasurementTheory.Course21120.TaylorExpansion

namespace Course21120Taylor

theorem term1_eq : (1 / 2 : ?) ^ 3 / 3 = 1 / 24 := by norm_num
theorem term2_eq : (1 / 2 : ?) ^ 5 / 5 = 1 / 160 := by norm_num
theorem term3_eq : 5 * (1 / 2 : ?) ^ 7 / 42 = 5 / 5376 := by norm_num
theorem term4_eq : 5 * (1 / 2 : ?) ^ 9 / 54 = 5 / 27648 := by norm_num

/-- The exact rational value of $T_9(1/2)$ is $34997 / 967680$. -/
```

```

theorem T9_half_eq : T9 (1 / 2) = 34997 / 967680 := by
  dsimp [T9]
  norm_num

/-- $T_9(1/2)$ lies strictly between $0.03615$ and $0.03625$. -/
theorem T9_half_rounds_to_0_0362 :
  (3615 : ?) / 100000 < T9 (1 / 2) /\ T9 (1 / 2) < 3625 / 100000 := by
  rw [T9_half_eq]
  norm_num

/-- Explicit distance to $0.0362$ is less than $0.00005$. -/
theorem T9_half_close_to_0_0362 :
  T9 (1 / 2) - 362 / 10000 < 5 / 100000 /\
  362 / 10000 - T9 (1 / 2) < 5 / 100000 := by
  rw [T9_half_eq]
  norm_num

#eval (1.0 / 24.0 : Float)
#eval (1.0 / 160.0 : Float)
#eval (5.0 / 5376.0 : Float)
#eval (5.0 / 27648.0 : Float)
#eval (34997.0 / 967680.0 : Float)

end Course21120Taylor

```

### 3.1.2.c.3 Explicit error bound

On  $[0, 0.5]$ , the series for the integrand is alternating with terms decreasing monotonically in magnitude. By the alternating series estimation theorem, the truncation error in the integrand is bounded by the magnitude of the next term:

$$R_{\text{integrand}}(t) \leq \left(1 - \frac{1}{6} + \frac{1}{120}\right) t^{10} = \frac{101}{120} t^{10}.$$

Integrating yields:

$$|R_f(0.5)| \leq \int_0^{0.5} \frac{101}{120} t^{10} dt = \frac{101}{1320} (0.5)^{11} = \frac{101}{1320 \times 2048} \approx 3.73 \times 10^{-5} < 0.0001.$$

Thus, to four decimal places:

$$f(0.5) \approx 0.0362 \quad (\text{with error} < 4 \times 10^{-5}).$$

### Lean 4 source

```

import Mathlib
import BSinMeasurementTheory.Course21120.TaylorExpansion

namespace Course21120Taylor

/-- Alternating-series tail bound
$|R(0.5)| \le (101 / 1320) \cdot (1/2)^{11}$. -/
def truncation_error : ? := (101 / 1320) * (1 / 2) ^ 11

```

```

theorem truncation_error_eq : truncation_error = 101 / 2703360 := by
  dsimp [truncation_error]
  norm_num

/-- The truncation error is strictly less than  $10^{-4}$ . -/
theorem truncation_error_lt : truncation_error < 1 / 10000 := by
  rw [truncation_error_eq]
  norm_num

#eval (101.0 / 2703360.0 : Float)

end Course21120Taylor

```

### 3.1.2.c.4 Summary of what Lean verifies here

#### 1. Term-by-term calculation:

- $x^3/3 = 1/24 \approx 0.041667$
- $x^5/5 = 1/160 = 0.006250$
- $5x^7/42 = 5/5376 \approx 0.000930$
- $5x^9/54 = 5/27648 \approx 0.000181$

2. **Exact value:**  $T_9(1/2) = \frac{34997}{967680} \approx 0.0361659$ .

3. **Rounding:** Lean proves  $0.03615 < T_9(1/2) < 0.03625$ , confirming it rounds to 0.0362.

4. **Error bound:** Lean proves the remainder bound  $\frac{101}{2703360} \approx 3.74 \times 10^{-5} < 10^{-4}$ .

## 3.2 21-127 — Concepts of Mathematics

### 3.2.1 Problem

Define a relation  $\sim$  on  $\mathbb{Z} \times (\mathbb{Z} \setminus \{0\})$  by  $(a, b) \sim (c, d) \iff ad = bc$ .

- (a) Prove  $\sim$  is an equivalence relation.
- (b) Prove the induced quotient set is in bijection with  $\mathbb{Q}$ , exhibiting the bijection explicitly and proving it well-defined, injective, and surjective.
- (c) Prove by strong induction that every nonempty subset of  $\mathbb{Q}_{\geq 0}$  that is well-ordered under the usual order is either empty or has infinitely many elements — or find the flaw in that claim and prove the correct statement instead, using contradiction.

*Requires: relations/equivalence classes, functions/bijections, induction, proof by contradiction, and — the real test — recognizing a false claim rather than forcing a proof of it.*




Figure 4: Lean module dependencies for the 21-127 solution. Amber box: Mathlib imports. Blue box: solution modules.

### 3.2.2 Solution

#### 3.2.2.a Equivalence Relation

We define  $S = \mathbb{Z} \times (\mathbb{Z} \setminus \{0\})$  and  $(a, b) \sim (c, d) \iff ad = bc$ : - **Reflexivity**:  $ab = ba$  holds by commutativity of multiplication in  $\mathbb{Z}$ . - **Symmetry**:  $ad = bc \implies cb = da$ , so  $(c, d) \sim (a, b)$ . - **Transitivity**: If  $ad = bc$  and  $cf = de$ , multiplying gives  $(af)d = (ad)f = (bc)f = b(cf) = b(de) = (be)d$ . Since  $d \neq 0$  and  $\mathbb{Z}$  is an integral domain (has no zero divisors),  $(af - be)d = 0 \implies af = be$ .

#### Lean 4 source

```

import Mathlib

namespace Course21127

/-- Integers with a nonzero denominator: the domain of the fraction relation. -/
structure NonzeroDenomInt where
  num : Int
  den : Int
  den_ne_zero : den != 0

/-- The relation (a, b) ~ (c, d) <-> a * d = b * c -/
def Rel (x y : NonzeroDenomInt) : Prop :=
  x.num * y.den = x.den * y.num

lemma rel_refl (x : NonzeroDenomInt) : Rel x x := by
  dsimp [Rel]

```

```

ring

lemma rel_symm {x y : NonzeroDenomInt} (h : Rel x y) : Rel y x := by
  dsimp [Rel] at *
  calc
    y.num * x.den = x.den * y.num := by ring
    _ = x.num * y.den := h.symm
    _ = y.den * x.num := by ring

lemma rel_trans {x y z : NonzeroDenomInt} (h1 : Rel x y) (h2 : Rel y z) : Rel x z :=
  by
  dsimp [Rel] at *
  have hd : y.den != 0 := y.den_ne_zero
  have H : (x.num * z.den) * y.den = (x.den * z.num) * y.den := by
    calc
      (x.num * z.den) * y.den = (x.num * y.den) * z.den := by ring
      _ = (x.den * y.num) * z.den := by rw [h1]
      _ = x.den * (y.num * z.den) := by ring
      _ = x.den * (y.den * z.num) := by rw [h2]
      _ = (x.den * z.num) * y.den := by ring
  have H2 : (x.num * z.den - x.den * z.num) * y.den = 0 := by
    calc
      (x.num * z.den - x.den * z.num) * y.den
      _ = (x.num * z.den) * y.den - (x.den * z.num) * y.den := by ring
      _ = (x.den * z.num) * y.den - (x.den * z.num) * y.den := by rw [H]
      _ = 0 := by ring
  cases mul_eq_zero.mp H2 with
  | inl h => exact sub_eq_zero.mp h
  | inr h => exact False.elim (hd h)

instance nonzeroDenomIntSetoid : Setoid NonzeroDenomInt where
  r := Rel
  iseqv := <rel_refl, rel_symm, rel_trans>

end Course21127

```

### 3.2.2.b Bijection with $\mathbb{Q}$

The canonical map is  $F(a, b) = \frac{a}{b} \in \mathbb{Q}$ : - **Well-defined:** If  $ad = bc$ , then  $\frac{a}{b} = \frac{c}{d}$  in  $\mathbb{Q}$  because clearing denominators yields  $ad - bc = 0$ . - **Injective:** If  $\frac{a}{b} = \frac{c}{d}$  in  $\mathbb{Q}$ , then  $ad = bc$  in  $\mathbb{Z}$ , meaning  $[(a, b)] = [(c, d)]$ . - **Surjective:** Every  $q \in \mathbb{Q}$  has canonical numerator  $q.\text{num} \in \mathbb{Z}$  and denominator  $q.\text{den} \in \mathbb{N}^+$ , with  $q = \frac{q.\text{num}}{q.\text{den}}$ . Since  $q.\text{den} \neq 0$ ,  $F([(q.\text{num}, q.\text{den})]) = q$ .

#### Lean 4 source

```

import Mathlib
import BSinMeasurementTheory.Course21127.NonzeroDenomInt

namespace Course21127

lemma rat_div_eq_of_mul_eq {a b c d : ?} (hb : b != 0) (hd : d != 0) (h : a * d =
  b * c) :
  a / b = c / d := by
  rw [div_eq_div_iff hb hd]
  linarith

```

```

lemma rat_mul_eq_of_div_eq {a b c d : ?} (hb : b != 0) (hd : d != 0) (h : a / b =
  c / d) :
  a * d = b * c := by
  have := (div_eq_div_iff hb hd).mp h
  linarith

/-- The explicit map from a pair (a, b) to Q. -/
def toRat (p : NonzeroDenomInt) : ? := (p.num : ?) / (p.den : ?)

/-- The map respects the equivalence relation (well-defined). -/
lemma toRat_wellDefined (x y : NonzeroDenomInt) (h : Rel x y) : toRat x = toRat y :=
  by
  dsimp [toRat]
  have hx : ((x.den : ?) != 0) := by
    intro hz
    exact x.den_ne_zero (by exact_mod_cast hz)
  have hy : ((y.den : ?) != 0) := by
    intro hz
    exact y.den_ne_zero (by exact_mod_cast hz)
  have h_int : x.num * y.den = x.den * y.num := h
  have h_cast : (x.num : ?) * (y.den : ?) = (x.den : ?) * (y.num : ?) := by
    exact_mod_cast h_int
  exact rat_div_eq_of_mul_eq hx hy h_cast

/-- The induced map on the quotient set. -/
def toRatQuot : Quotient nonzeroDenomIntSetoid -> ? :=
  Quotient.lift toRat toRat_wellDefined

/-- (b) Part 1: Injectivity -/
theorem toRatQuot_injective : Function.Injective toRatQuot := by
  intro q1 q2
  refine Quotient.inductionOn_2 q1 q2 ?_
  intro x y h
  dsimp [toRatQuot, toRat] at h
  have hx : ((x.den : ?) != 0) := by
    intro hz
    exact x.den_ne_zero (by exact_mod_cast hz)
  have hy : ((y.den : ?) != 0) := by
    intro hz
    exact y.den_ne_zero (by exact_mod_cast hz)
  have h_mul := rat_mul_eq_of_div_eq hx hy h
  have h_int : x.num * y.den = x.den * y.num := by
    exact_mod_cast h_mul
  apply Quotient.sound
  exact h_int

/-- (b) Part 2: Surjectivity -/
theorem toRatQuot_surjective : Function.Surjective toRatQuot := by
  intro q
  have hden : (q.den : Int) != 0 := by
    intro hz
    have : q.den = 0 := by exact_mod_cast hz
    exact q.den_nz this
  use Quotient.mk nonzeroDenomIntSetoid <q.num, (q.den : Int), hden>
  change toRat <q.num, (q.den : Int), hden> = q
  dsimp [toRat]

```

```

have : ((q.den : Int) : ?) = (q.den : ?) := by push_cast; rfl
rw [this]
exact Rat.num_div_den q

/-- (b) Part 3: Bijectivity -/
theorem toRatQuot_bijective : Function.Bijective toRatQuot :=
  <toRatQuot_injective, toRatQuot_surjective>

/-- The induced quotient set is in bijection with  $\mathbb{Q}$ . -/
noncomputable def quotientEquivRat : Quotient nonzeroDenomIntSetoid ? ? :=
  Equiv.ofBijective toRatQuot toRatQuot_bijective

end Course21127

```

### 3.2.2.c The Flaw in the Claim and the Correct Statement

The prompt poses the challenge: > “*Prove by strong induction that every nonempty subset of  $\mathbb{Q}_{\geq 0}$  that is well-ordered under the usual order is either empty or has infinitely many elements — or find the flaw in that claim and prove the correct statement instead, using contradiction.*”

**The Flaws in the Claim:** 1. **Syntactic contradiction:** A subset assumed to be *nonempty* cannot be *empty*. 2. **Mathematical falsehood:** The claim asserts that any nonempty well-ordered subset of  $\mathbb{Q}_{\geq 0}$  must be infinite. This is completely false. Any finite nonempty subset of  $\mathbb{Q}_{\geq 0}$  (for instance, the singleton  $\{0\}$ , or  $\{0, 1\}$ , or  $\{1/2\}$ ) is: - A subset of  $\mathbb{Q}_{\geq 0}$ , - Nonempty, - Well-ordered under the standard  $\leq$  order (every nonempty subset has a least element), - Yet strictly **finite**, not infinite. 3. **Why strong induction fails here (the trap):** An attempted proof by induction would try to construct an infinite sequence  $x_0 < x_1 < x_2 < \dots$  by repeatedly taking successors. This fails because a finite well-ordered subset has a **maximal element** with no strictly greater element in the set.

**The Correct Theorems:** 1. **Refutation by contradiction:** The claim is false. Assuming the claim holds for  $S = \{0\}$  implies  $\{0\} = \emptyset$  or  $\{0\}$  is infinite, both of which immediately contradict the properties of singletons. 2. **Positive statement via induction:** Every finite nonempty subset of  $\mathbb{Q}$  (and hence of  $\mathbb{Q}_{\geq 0}$ ) has a least element, and is therefore well-ordered. We formalize this by induction on lists.

#### Lean 4 source

```

import Mathlib

namespace Course21127

/-- A subset of  $\mathbb{Q}$  is well-ordered under the usual order if every nonempty
subset has a least element in the set. -/
class IsWellOrderedSet (S : Set ?) : Prop where
  least : forall T : Set ?, T subseteq S -> T.Nonempty -> exists m in T,
    forall x in T, m <= x

/-- The false claim stated in the prompt using Set.Ici to avoid untyped binder
problems. -/
def FalseClaim : Prop :=
  forall S : Set ?, S subseteq Set.Ici (0 : ?) -> S.Nonempty ->
    IsWellOrderedSet S ->
      (S = {} \/\ Set.Infinite S)

lemma singleton_zero_well_ordered : IsWellOrderedSet {(0 : ?)} where
  least T hT hne := by

```

```

    rcases hne with <t, ht>
    have ht0 : t = 0 := hT ht
    subst ht0
    refine <0, ht, ?_>
    intro x hx
    have hx0 : x = 0 := hT hx
    subst hx0
    exact le_rfl

lemma singleton_zero_nonneg : ({(0 : ?)} : Set ?) ⊆ Set.Ici 0 :=
  Set.singleton_subset_iff.mpr (Set.mem_Ici.mpr le_rfl)

lemma singleton_zero_nonempty : ({(0 : ?)} : Set ?).Nonempty :=
  <0, rfl>

/-- (c) Refutation by contradiction:
    The claim implies that {0} must be empty or infinite, both of which are false. -/
theorem false_claim_is_false : not FalseClaim := by
  intro hclaim
  have h_cases := hclaim {0} singleton_zero_nonneg singleton_zero_nonempty
  singleton_zero_well_ordered
  cases h_cases with
  | inl h_empty =>
    have h0 : (0 : ?) in ({(0 : ?)} : Set ?) := rfl
    rw [h_empty] at h0
    exact h0
  | inr h_inf =>
    exact h_inf (Set.finite_singleton 0)

/-- Correct positive statement proved by induction:
    Every nonempty finite list of rationals contains a least element. -/
lemma list_has_min : forall (a : ?) (l : List ?), exists m in a :: l, forall x
  in a :: l, m <= x
| a, [] => by
  refine <a, List.mem_cons.mpr (Or.inl rfl), ?_>
  intro x hx
  rcases List.mem_cons.mp hx with hxa | hx_nil
  * rw [hxa]
  * cases hx_nil
| a, b :: l => by
  rcases list_has_min (min a b) l with <m, hm, hle>
  have hm_orig : m in a :: b :: l := by
    rcases List.mem_cons.mp hm with hm_eq | h_in
    * rw [hm_eq]
    cases le_total a b with
    | inl hab =>
      rw [min_eq_left hab]
      exact List.mem_cons.mpr (Or.inl rfl)
    | inr hba =>
      rw [min_eq_right hba]
      exact List.mem_cons.mpr (Or.inr (List.mem_cons.mpr (Or.inl rfl)))
  * exact List.mem_cons.mpr (Or.inr (List.mem_cons.mpr (Or.inr h_in)))
  refine <m, hm_orig, ?_>
  intro x hx
  rcases List.mem_cons.mp hx with hxa | hx_l
  * rw [hxa]

```

```

    have h_min_in : min a b in (min a b) :: l := List.mem_cons.mpr (Or.inl rfl)
    exact le_trans (hle (min a b) h_min_in) (min_le_left a b)
  * rcases List.mem_cons.mp hx1 with hxb | hx2
  * rw [hxb]
    have h_min_in : min a b in (min a b) :: l := List.mem_cons.mpr (Or.inl rfl)
    exact le_trans (hle (min a b) h_min_in) (min_le_right a b)
  * have hx_in : x in (min a b) :: l := List.mem_cons.mpr (Or.inr hx2)
    exact hle x hx_in
end Course21127

```

## Lean 4 source

```

import Mathlib
import BSinMeasurementTheory.Course21127.NonzeroDenomInt
import BSinMeasurementTheory.Course21127.ToRat

namespace Course21127

-- Example 1: (1, 2) ~ (2, 4) since 1 * 4 = 2 * 2 = 4
def pair1 : NonzeroDenomInt := <1, 2, by decide>
def pair2 : NonzeroDenomInt := <2, 4, by decide>

example : Rel pair1 pair2 := by
  dsimp [Rel, pair1, pair2]

-- Example 2: Both pairs map to 1/2 in ? under toRat
example : toRat pair1 = 1 / 2 := by
  norm_num [toRat, pair1]

example : toRat pair2 = 1 / 2 := by
  norm_num [toRat, pair2]

-- Example 3: Minimal element in the set {1/2, 1/3} is 1/3
example : min (1 / 2 : ?) (1 / 3) = 1 / 3 := by
  norm_num
end Course21127

```

## 3.3 21-241 — Matrices and Linear Transformations

### 3.3.1 Problem

Let  $V$  be a finite-dimensional real vector space,  $C \subset V$  a convex cone with nonempty interior, and  $v \notin C$ .

- Construct a basis of  $V$  and its dual basis of  $V^*$  explicitly for the case  $V = \mathbb{R}^3$ ,  $C = \{x : x_1, x_2, x_3 \geq 0\}$ .
- Prove that a linear functional  $\varphi \in V^*$  separating  $v$  from  $C$  (i.e.,  $\varphi(v) < 0 \leq \varphi(c)$  for all  $c \in C$ ) corresponds to a specific vector under the standard inner product, and compute it for a concrete  $v \notin C$ .
- Prove that the set of linear functionals nonnegative on all of  $C$  is itself a convex cone in  $V^*$ , and identify its dual basis representation.

*Requires: basis/dual basis construction, linear transformations, the vector-space language* Scott's Theorem 1.1 is stated in — not just matrix computation.

Prove it in English and then in Lean 4, using as much Mathlib as possible. No uses of sorry, admit or axiom are allowed. If you do any numerical calculations for examples to illustrate the proof, reproduce them after the proof in Lean.

### 3.3.2 Solution




Figure 5: Lean module dependencies for the 21-241 solution. Amber box: Mathlib imports. Blue box: solution modules.

#### Geometric Separation and Scott's Theorem 1.1.

Let  $V$  be a finite-dimensional real vector space. A non-empty subset  $C \subseteq V$  is a **convex cone** if: 1. For all  $x, y \in C$ ,  $x + y \in C$  (closed under addition). 2. For all  $x \in C$  and all  $\lambda \geq 0$ ,  $\lambda x \in C$  (closed under non-negative scaling).

Scott's Theorem 1.1 (the geometric Hahn–Banach separation theorem for convex cones) establishes that if  $C \subset V$  is a convex cone with non-empty interior  $\text{int}(C) \neq \emptyset$ , and  $v \in V \setminus C$ , then there exists a non-zero linear functional  $\varphi \in V^*$  strictly separating  $v$  from  $\text{int}(C)$  and non-strictly separating  $v$  from  $C$ :

$$\varphi(v) < 0 \leq \varphi(c), \quad \forall c \in C.$$

The separation level must be 0: if  $\varphi(c) \geq \alpha$  for all  $c \in C$ , then for any  $\lambda > 0$ ,  $\varphi(\lambda c) = \lambda \varphi(c) \geq \alpha \implies \varphi(c) \geq \alpha/\lambda$ . Sending  $\lambda \rightarrow \infty$  forces  $\varphi(c) \geq 0$ . Furthermore, since  $0 \in \overline{C}$ ,  $\alpha \leq \varphi(0) = 0$ .

#### 3.3.2.a Basis and Dual Basis Construction

Let  $V = \mathbb{R}^3$  and  $C = \{x = (x_0, x_1, x_2) \in \mathbb{R}^3 \mid x_0 \geq 0, x_1 \geq 0, x_2 \geq 0\}$  be the non-negative orthant.

1. **Primal Basis of  $V$ :** Define the standard basis vectors  $e_0, e_1, e_2 \in V$  by:

$$e_i(j) = \delta_{ij} = \begin{cases} 1 & \text{if } i = j, \\ 0 & \text{if } i \neq j. \end{cases}$$

Explicitly:

$$e_0 = (1, 0, 0), \quad e_1 = (0, 1, 0), \quad e_2 = (0, 0, 1).$$

- **Spanning:** Any vector  $x \in V$  can be uniquely expressed as:

$$x = \sum_{i=0}^2 x_i e_i.$$

- **Linear Independence:** If  $\sum_{i=0}^2 c_i e_i = 0$ , evaluating the  $j$ -th coordinate gives  $c_j = 0$  for each  $j \in \{0, 1, 2\}$ .

2. **Dual Basis of  $V^*$ :** The algebraic dual space is  $V^* = \text{Hom}_{\mathbb{R}}(V, \mathbb{R})$ . The dual basis  $\{e_0^*, e_1^*, e_2^*\} \subset V^*$  is defined by the coordinate projections:

$$e_i^*(x) := x_i = \text{proj}_i(x).$$

- **Duality relation:**

$$e_i^*(e_j) = e_j(i) = \delta_{ij}.$$

- **Expansion in the dual basis:** For any linear map  $\varphi \in V^*$ , by linearity:

$$\varphi(x) = \varphi\left(\sum_{i=0}^2 x_i e_i\right) = \sum_{i=0}^2 x_i \varphi(e_i) = \sum_{i=0}^2 \varphi(e_i) e_i^*(x),$$

$$\text{meaning } \varphi = \sum_{i=0}^2 \varphi(e_i) e_i^*.$$

#### Lean 4 source

```
import Mathlib

namespace Course21241

abbrev V := Fin 3 -> ?

/-- The non-negative orthant cone C in V = ?? -/
def C : Set V := {x | forall i : Fin 3, 0 <= x i}

end Course21241
```

#### Lean 4 source

```
import Mathlib
import BSinMeasurementTheory.Course21241.Space

open BigOperators
open Finset

namespace Course21241

/-- Standard primal basis vectors of V -/
```

```

def e (i : Fin 3) : V := fun j => if i = j then 1 else 0

/-- Explicit dual basis functionals e_star i in V* -/
def e_star (i : Fin 3) : V -> ?[?] ? where
  toFun x := x i
  map_add' _ _ := rfl
  map_smul' _ _ := rfl

/-- Duality relation between primal and dual bases: e_star i (e j) = ?_{ij} -/
theorem e_star_apply_e (i j : Fin 3) :
  e_star i (e j) = if i = j then (1 : ?) else 0 := by
  dsimp [e_star, e]
  fin_cases i <|> fin_cases j <|> rfl

/-- Every vector x in V expands uniquely in the primal basis -/
theorem vector_eq_sum (x : V) :
  x = ? i : Fin 3, x i * e i := by
  ext j
  simp only [Finset.sum_apply, Pi.smul_apply, smul_eq_mul]
  rw [Fin.sum_univ_three]
  fin_cases j <|> { dsimp [e]; ring }

/-- The basis vectors {e 0, e 1, e 2} are linearly independent -/
theorem e_linearIndependent :
  LinearIndependent ? e := by
  rw [linearIndependent_iff']
  intro s g hg i hi
  have h := congr_fun hg i
  simp only [Finset.sum_apply, Pi.smul_apply, smul_eq_mul, e, Pi.zero_apply] at h
  rw [Finset.sum_eq_single_of_mem i hi] at h
  * simp only [ite_true, mul_one] at h
  exact h
  * intro j _ hj
  simp only [hj, ite_false, mul_zero]

/-- Every linear functional phi in V* expands uniquely in the dual basis -/
theorem dual_eq_sum (phi : V -> ?[?] ?) :
  phi = ? i : Fin 3, phi (e i) * e_star i := by
  apply LinearMap.ext
  intro x
  have hx : x = ? i : Fin 3, x i * e i := vector_eq_sum x
  conv_lhs => rw [hx]
  rw [map_sum]
  simp only [LinearMap.sum_apply, LinearMap.smul_apply, smul_eq_mul]
  dsimp [e_star]
  apply Finset.sum_congr rfl
  intro i _
  rw [LinearMap.map_smul, smul_eq_mul, mul_comm]
end Course21241

```

### 3.3.2.b Inner Product Representation and Concrete Separation

1. **Riesz Isomorphism under the Standard Inner Product:** Equip  $V$  with the standard Euclidean inner product:

$$\langle u, x \rangle = \sum_{i=0}^2 u_i x_i.$$

Every vector  $u \in V$  induces a linear functional  $\varphi_u(x) = \langle u, x \rangle$ . Conversely, every linear functional  $\varphi \in V^*$  uniquely corresponds to the vector:

$$u_\varphi := \sum_{i=0}^2 \varphi(e_i) e_i = (\varphi(e_0), \varphi(e_1), \varphi(e_2)).$$

Indeed:

$$\langle u_\varphi, x \rangle = \sum_{i=0}^2 \varphi(e_i) x_i = \varphi \left( \sum_{i=0}^2 x_i e_i \right) = \varphi(x).$$

2. **Non-negativity on  $C$ :**

$$\forall c \in C, \langle u, c \rangle \geq 0 \iff \forall i \in \{0, 1, 2\}, u_i \geq 0.$$

- ( $\implies$ ) Since  $e_i \in C$ , we have  $u_i = \langle u, e_i \rangle \geq 0$ .
- ( $\impliedby$ ) If  $u_i \geq 0$  for all  $i$  and  $c \in C$  (so  $c_i \geq 0$ ), then  $\langle u, c \rangle = \sum_{i=0}^2 u_i c_i \geq 0$ .

3. **Separating Vector for  $v \notin C$ :** If  $v \notin C$ , then by definition  $\exists k \in \{0, 1, 2\}$  such that  $v_k < 0$ . Choosing  $u = e_k$ , we verify:

- For all  $c \in C$ :  $\langle u, c \rangle = \langle e_k, c \rangle = c_k \geq 0$ .
- At  $v$ :  $\langle u, v \rangle = \langle e_k, v \rangle = v_k < 0$ . Thus  $\varphi(x) = \langle e_k, x \rangle = e_k^*(x)$  is the desired separating functional.

#### Lean 4 source

```
import Mathlib
import BSinMeasurementTheory.Course21241.Basis

open BigOperators
open Finset

namespace Course21241

/-- Standard Euclidean inner product on V -/
def dot (u v : V) : ? := ? i : Fin 3, u i * v i

lemma dot_comm (u v : V) : dot u v = dot v u := by
  dsimp [dot]
  apply Finset.sum_congr rfl
  intro i _
  ring
```

```

lemma dot_e (u : V) (i : Fin 3) : dot u (e i) = u i := by
  dsimp [dot]
  rw [Fin.sum_univ_three]
  fin_cases i <|> { dsimp [e]; ring }

/-- Riesz map from V to V* -/
def toDual (u : V) : V → ?[?] ? where
  toFun v := dot u v
  map_add' v w := by
    dsimp [dot]
    rw [← Finset.sum_add_distrib]
    apply Finset.sum_congr rfl
    intro i _
    ring
  map_smul' c v := by
    dsimp [dot]
    rw [Finset.mul_sum]
    apply Finset.sum_congr rfl
    intro i _
    ring

/-- Inverse Riesz map from V* to V -/
def toVec (phi : V → ?[?] ?) : V := fun i => phi (e i)

theorem toDual_toVec (phi : V → ?[?] ?) : toDual (toVec phi) = phi := by
  apply LinearMap.ext
  intro x
  have hx : x = ? i : Fin 3, x i * e i := vector_eq_sum x
  conv_rhs => rw [hx]
  rw [map_sum]
  change (? i : Fin 3, phi (e i) * x i) = ? i : Fin 3, phi (x i * e i)
  apply Finset.sum_congr rfl
  intro i _
  rw [LinearMap.map_smul, smul_eq_mul, mul_comm]

theorem toVec_toDual (u : V) : toVec (toDual u) = u := by
  ext i
  exact dot_e u i

/-- Characterization of functionals nonnegative on C -/
theorem nonneg_on_C_iff (u : V) :
  (forall c in C, 0 ≤ dot u c) <-> forall i : Fin 3, 0 ≤ u i := by
  constructor
  * intro h i
  have hei : e i in C := by
    intro j
    dsimp [e]
    split_ifs <|> linarith
  have h_dot := h (e i) hei
  rw [dot_e] at h_dot
  exact h_dot
  * intro hu c hc
  dsimp [dot]
  apply Finset.sum_nonneg
  intro i _
  exact mul_nonneg (hu i) (hc i)

```

```

/-- Existence of a separating vector under the inner product for any  $v \notin C$  -/
theorem exists_separating_vector (v : V) (hv : v ∉ C) :
  exists u : V, (forall c in C, 0 ≤ dot u c) /\ dot u v < 0 := by
  change not (forall i : Fin 3, 0 ≤ v i) at hv
  push Not at hv
  rcases hv with <i, hi>
  use e i
  constructor
  * rw [nonneg_on_C_iff]
    intro j
    dsimp [e]
    split_ifs <;> linarith
  * rw [dot_comm, dot_e]
    exact hi
end Course21241

```

#### 4. Concrete Numerical Example: Let $v = (-1, 1, 1)$ .

- $v \notin C$  because  $v_0 = -1 < 0$ .
- Choose  $u = e_0 = (1, 0, 0)$ .
- Separation check:

$$\langle u, v \rangle = 1 \cdot (-1) + 0 \cdot 1 + 0 \cdot 1 = -1 < 0.$$

- Cone non-negativity: for any  $c = (c_0, c_1, c_2) \in C$ ,

$$\langle u, c \rangle = c_0 \geq 0.$$

##### 3.3.2.b.1 Choice of point $v$ not in $C$

###### Lean 4 source

```

import Mathlib
import BSinMeasurementTheory.Course21241.Space

namespace Course21241

def v : V := fun i => if i = 0 then -1 else 1

lemma v_zero : v 0 = -1 := rfl
lemma v_one : v 1 = 1 := rfl
lemma v_two : v 2 = 1 := rfl

theorem v_coords : v 0 = -1 /\ v 1 = 1 /\ v 2 = 1 :=
  <v_zero, v_one, v_two>

theorem v_zero_neg : v 0 < 0 := by
  rw [v_zero]
  norm_num

theorem v_not_mem_C : v ∉ C := by
  intro hc

```

```

have h0 : 0 <= v 0 := hc 0
rw [v_zero] at h0
norm_num at h0

end Course21241

```

### 3.3.2.b.2 Separating vector $u$

#### Lean 4 source

```

import Mathlib
import BSinMeasurementTheory.Course21241.Basis

namespace Course21241

def u : V := e 0

lemma u_zero : u 0 = 1 := rfl
lemma u_one : u 1 = 0 := rfl
lemma u_two : u 2 = 0 := rfl

theorem u_eq_e_zero : u = e 0 := rfl

theorem u_coords : u 0 = 1 /\ u 1 = 0 /\ u 2 = 0 :=
  <u_zero, u_one, u_two>

end Course21241

```

### 3.3.2.b.3 Inner product value at $v$

#### Lean 4 source

```

import Mathlib
import BSinMeasurementTheory.Course21241.PointV
import BSinMeasurementTheory.Course21241.SeparatingU
import BSinMeasurementTheory.Course21241.Separation

open BigOperators
open Finset

namespace Course21241

theorem dot_u_v_sum :
  dot u v = ? i : Fin 3, u i * v i := rfl

theorem dot_u_v_expansion :
  dot u v = (1 : ?) * (-1) + 0 * 1 + 0 * 1 := by
  dsimp [dot]
  rw [Fin.sum_univ_three]
  rw [u_zero, u_one, u_two, v_zero, v_one, v_two]

theorem dot_u_v_eval : dot u v = -1 := by
  rw [dot_u_v_expansion]
  ring

```

```

theorem dot_u_v_neg : dot u v < 0 := by
  rw [dot_u_v_eval]
  norm_num

end Course21241

```

### 3.3.2.b.4 Inner product on an arbitrary $c$ in $C$

#### Lean 4 source

```

import Mathlib
import BSinMeasurementTheory.Course21241.SeparatingU
import BSinMeasurementTheory.Course21241.Separation

open BigOperators
open Finset

namespace Course21241

theorem dot_u_c_sum (c : V) :
  dot u c = ? i : Fin 3, u i * c i := rfl

theorem dot_u_c_expansion (c : V) :
  dot u c = (1 : ?) * c 0 + 0 * c 1 + 0 * c 2 := by
  dsimp [dot]
  rw [Fin.sum_univ_three]
  rw [u_zero, u_one, u_two]

theorem dot_u_c_eq_coord (c : V) :
  dot u c = c 0 := by
  rw [dot_u_c_expansion]
  ring

theorem dot_u_c_nonneg (c : V) (hc : c in C) :
  0 <= dot u c := by
  rw [dot_u_c_eq_coord]
  exact hc 0

end Course21241

```

### 3.3.2.b.5 Linear functional separation

#### Lean 4 source

```

import Mathlib
import BSinMeasurementTheory.Course21241.DotAtV
import BSinMeasurementTheory.Course21241.DotOnC

namespace Course21241

def phi_u : V -> ?[?] ? := toDual u

theorem phi_u_v_eval : phi_u v = -1 := by
  change dot u v = -1
  exact dot_u_v_eval

```

```

theorem phi_u_c_eval (c : V) : phi_u c = c 0 := by
  change dot u c = c 0
  exact dot_u_c_eq_coord c

theorem phi_u_v_neg : phi_u v < 0 := by
  rw [phi_u_v_eval]
  norm_num

theorem phi_u_c_nonneg (c : V) (hc : c in C) : 0 <= phi_u c := by
  rw [phi_u_c_eval]
  exact hc 0

theorem phi_u_separates (c : V) (hc : c in C) :
  phi_u v < 0 /\ 0 <= phi_u c :=
  <phi_u_v_neg, phi_u_c_nonneg c hc>

theorem phi_u_separation_chain (c : V) (hc : c in C) :
  phi_u v = -1 /\ phi_u v < 0 /\ 0 <= phi_u c /\ phi_u c = c 0 :=
  <phi_u_v_eval, phi_u_v_neg, phi_u_c_nonneg c hc, phi_u_c_eval c>

end Course21241

```

### 3.3.2.c The Dual Cone and its Dual Basis Representation

1. **The Dual Cone is a Convex Cone:** The dual cone of  $C$  is defined by:

$$C^* := \{\varphi \in V^* \mid \forall c \in C, \varphi(c) \geq 0\}.$$

- **Closed under non-negative scaling:** Let  $\varphi \in C^*$  and  $\lambda \geq 0$ . For any  $c \in C$ :

$$(\lambda\varphi)(c) = \lambda \cdot \varphi(c) \geq 0 \quad (\text{product of non-negative reals}).$$

- **Closed under addition:** Let  $\varphi, \psi \in C^*$ . For any  $c \in C$ :

$$(\varphi + \psi)(c) = \varphi(c) + \psi(c) \geq 0 + 0 = 0.$$

- **Conic combinations:** If  $\varphi, \psi \in C^*$  and  $a, b \geq 0$ , then  $a\varphi + b\psi \in C^*$ . Thus  $C^*$  is a convex cone in  $V^*$ .

2. **Dual Basis Representation:** Using the dual basis expansion from Part (a), any  $\varphi \in V^*$  is  $\varphi = \sum_{i=0}^2 a_i e_i^*$ , where  $a_i = \varphi(e_i)$ .

- If  $\varphi \in C^*$ , then since  $e_i \in C$ ,  $a_i = \varphi(e_i) \geq 0$  for all  $i$ .
- Conversely, if  $a_i \geq 0$  for all  $i$ , then for any  $c \in C$ :

$$\varphi(c) = \sum_{i=0}^2 a_i e_i^*(c) = \sum_{i=0}^2 a_i c_i \geq 0.$$

Therefore:

$$C^* = \left\{ \sum_{i=0}^2 a_i e_i^* \mid a_0, a_1, a_2 \geq 0 \right\}.$$

Under the dual basis  $\{e_0^*, e_1^*, e_2^*\}$ , the dual cone  $C^*$  is canonically identified with the non-negative orthant  $\mathbb{R}_{\geq 0}^3$ .

### Lean 4 source

```
import Mathlib
import BSinMeasurementTheory.Course21241.Basis

open BigOperators
open Finset

namespace Course21241

/-- The dual cone C* in V* -/
def dualCone (C : Set V) : Set (V → ℝ) :=
  {phi | forall c in C, 0 ≤ phi c}

theorem dualCone_smul_mem {phi : V → ℝ} (hphi : phi in dualCone C) {a : ℝ} (
  ha : 0 ≤ a) :
  a * phi in dualCone C := by
  intro c hc_mem
  simp only [LinearMap.smul_apply, smul_eq_mul]
  exact mul_nonneg ha (hphi c hc_mem)

theorem dualCone_add_mem {phi psi : V → ℝ} (hphi : phi in dualCone C) (hpsi :
  psi in dualCone C) :
  phi + psi in dualCone C := by
  intro c hc_mem
  simp only [LinearMap.add_apply]
  exact add_nonneg (hphi c hc_mem) (hpsi c hc_mem)

theorem dualCone_conic_combination {phi psi : V → ℝ} (hphi : phi in dualCone
  C) (hpsi : psi in dualCone C)
  {a b : ℝ} (ha : 0 ≤ a) (hb : 0 ≤ b) :
  a * phi + b * psi in dualCone C := by
  intro c hc_mem
  simp only [LinearMap.add_apply, LinearMap.smul_apply, smul_eq_mul]
  exact add_nonneg (mul_nonneg ha (hphi c hc_mem)) (mul_nonneg hb (hpsi c hc_mem))

theorem mem_dualCone_iff (phi : V → ℝ) :
  phi in dualCone C ↔ forall i : Fin 3, 0 ≤ phi (e i) := by
  constructor
  * intro hphi i
  have hei : e i in C := by
    intro j
    dsimp [e]
    split_ifs <=> linarith
  exact hphi (e i) hei
  * intro hphi c hc
  have hc_sum : c = ∑ i : Fin 3, c i * e i := vector_eq_sum c
  conv_rhs => rw [hc_sum]
  rw [map_sum]
  simp only [LinearMap.map_smul, smul_eq_mul]
  apply Finset.sum_nonneg
  intro i _
```

```

exact mul_nonneg (hc i) (hphi i)

/-- Dual cone represented as non-negative linear combinations of {e_star i} -/
def dualCone_representation : Set (V -> ?[?] ?) :=
  {phi | exists a : Fin 3 -> ?, (forall i, 0 <= a i) /\ phi = ? i : Fin 3, a i
    * e_star i}

theorem dualCone_eq_representation :
  dualCone C = dualCone_representation := by
  ext phi
  constructor
  * intro hphi
    rw [mem_dualCone_iff] at hphi
    use fun i => phi (e i)
    refine <hphi, ?_>
    exact dual_eq_sum phi
  * rintro <a, ha, rfl>
    rw [mem_dualCone_iff]
    intro i
    simp only [LinearMap.sum_apply, LinearMap.smul_apply, smul_eq_mul]
    have h_sum : (? j : Fin 3, a j * e_star j (e i)) = a i := by
      rw [Fin.sum_univ_three]
      fin_cases i <;> simp [e_star_apply_e]
    rw [h_sum]
    exact ha i

end Course21241

```

### 3.4 21-228 — Discrete Mathematics

#### 3.4.1 Problem

Let  $x_1, \dots, x_n \in \mathbb{Q}^k$  be a finite sequence of rational vectors.

- Prove that if every permutation  $\pi$  of  $\{1, \dots, n\}$  produces a sequence  $x_{\pi(1)}, \dots, x_{\pi(n)}$  satisfying some symmetric linear condition  $L$ , then  $L$  depends only on the multiset  $\{x_1, \dots, x_n\}$ , not the order.
- Given two sequences  $x_1, \dots, x_n$  and  $y_1, \dots, y_n$ , count (with an exact combinatorial formula, not an estimate) the number of pairs of permutations  $(\pi, \sigma)$  for which  $\sum_i x_{\pi(i)} = \sum_i y_{\sigma(i)}$  termwise, under stated genericity assumptions on the vectors.
- Prove your formula reduces correctly to the  $n = 2$  and  $n = 3$  cases by direct enumeration.

*Requires: permutations/combinations, combinatorial proof technique, the direct toolkit for Scott's Theorems 1.2 and 3.2.*

Prove it in English and then in Lean 4, using as much Mathlib as possible. No uses of sorry, admit or axiom are allowed. If you do any numerical calculations for examples to illustrate the proof, reproduce them after the proof in Lean.




Figure 6: Lean module dependencies for the 21-228 solution. Amber box: Mathlib imports. Blue box: solution modules.

### 3.4.2 Solution

#### 3.4.2.a Symmetric conditions depend only on the multiset

**Proposition:** Let  $x_1, \dots, x_n \in \mathbb{Q}^k$  be a sequence of vectors. If every permutation  $\pi \in S_n$  produces a sequence  $(x_{\pi(1)}, \dots, x_{\pi(n)})$  satisfying a symmetric linear condition  $L$ , then  $L$  depends only on the underlying multiset  $\{x_1, \dots, x_n\}$ , not on the order.

##### 3.4.2.a.1 Proof:

1. **Symmetry and Multiset Equivalence:** Two sequences  $u = (u_1, \dots, u_n)$  and  $v = (v_1, \dots, v_n)$  in  $(\mathbb{Q}^k)^n$  represent the same multiset if and only if they lie in the same orbit under the natural action of the symmetric group  $S_n$  given by

$$\pi \cdot (u_1, \dots, u_n) = (u_{\pi(1)}, \dots, u_{\pi(n)}).$$

A condition  $L$  is *symmetric* if for every permutation  $\pi \in S_n$  and every sequence  $u \in (\mathbb{Q}^k)^n$ ,

$$L(u_{\pi(1)}, \dots, u_{\pi(n)}) \iff L(u_1, \dots, u_n).$$

Consequently, the truth value (or evaluation) of  $L$  is constant on every orbit of the  $S_n$ -action. Since the quotient space  $(\mathbb{Q}^k)^n / S_n$  is canonically bijective to the set of multisets of size  $n$  with elements in  $\mathbb{Q}^k$ ,  $L$  factors through this quotient. Therefore,  $L$  depends only on the multiset  $\{x_1, \dots, x_n\}$ .

2. **Structure of Symmetric Linear Forms:** If  $L : (\mathbb{Q}^k)^n \rightarrow \mathbb{Q}^m$  is a linear mapping, it can be written in block form as

$$L(u_1, \dots, u_n) = \sum_{i=1}^n A_i u_i$$

for linear maps  $A_i : \mathbb{Q}^k \rightarrow \mathbb{Q}^m$ . Symmetry under the transposition  $(i \ j) \in S_n$  implies that for all vectors  $v \in \mathbb{Q}^k$ , evaluating  $L$  on the tuple  $(0, \dots, v, \dots, 0)$  (with  $v$  in position  $i$ ) gives the same result as evaluating on the tuple with  $v$  in position  $j$ . Thus:

$$A_i v = A_j v \quad \forall v \in \mathbb{Q}^k \implies A_1 = A_2 = \dots = A_n =: A.$$

Hence, every symmetric linear map has the form:

$$L(u_1, \dots, u_n) = A \left( \sum_{i=1}^n u_i \right).$$

Because vector addition in  $\mathbb{Q}^k$  is commutative and associative, the sum  $\sum_{i=1}^n x_i$  is an invariant of the multiset  $\{x_1, \dots, x_n\}$ , depending in no way on the order of the terms. ■

#### Lean 4 source

```
import Mathlib.Data.Fintype.Perm
import Mathlib.Algebra.BigOperators.Group.Finset.Basic

open Equiv

namespace Course21228
```

```

variable {n : Nat} {a : Type*}

/-- A condition `L` on length-`n` sequences is symmetric if it is invariant
    under any permutation of the sequence. -/
class IsSymmetric (L : (Fin n -> a) -> Prop) : Prop where
  perm_iff : forall (x : Fin n -> a) (? : Perm (Fin n)), L (x comp ?) <=> L x

/-- If two sequences represent the same multiset (i.e. one is a permutation of the
    other),
    any symmetric condition `L` yields the same truth value on both. -/
theorem symmetric_depends_only_on_multiset
  (L : (Fin n -> a) -> Prop) [hL : IsSymmetric L]
  (x y : Fin n -> a) (? : Perm (Fin n)) (h : y = x comp ?) :
  L y <=> L x := by
  rw [h]
  exact hL.perm_iff x ?

/-- For any additive commutative monoid, the total sum is invariant under permutations
    . -/
theorem sum_perm_invariant {M : Type*} [AddCommMonoid M]
  (x : Fin n -> M) (? : Perm (Fin n)) :
  ? i : Fin n, (x comp ?) i = ? i : Fin n, x i := by
  exact Equiv.sum_comp ? x

end Course21228

```

### 3.4.2.b Counting permutation pairs

**Problem:** Given sequences  $x = (x_1, \dots, x_n)$  and  $y = (y_1, \dots, y_n)$  in  $\mathbb{Q}^k$ , find an exact combinatorial formula for the number of pairs of permutations  $(\pi, \sigma) \in S_n \times S_n$  such that  $\sum_i x_{\pi(i)} = \sum_i y_{\sigma(i)}$  *termwise*.

#### Lean 4 source

```

import Mathlib.Data.Fintype.Perm
import Mathlib.Algebra.BigOperators.Group.Finset.Basic

open Equiv

namespace Course21228

variable {n : Nat} {a : Type*}

/-- A condition `L` on length-`n` sequences is symmetric if it is invariant
    under any permutation of the sequence. -/
class IsSymmetric (L : (Fin n -> a) -> Prop) : Prop where
  perm_iff : forall (x : Fin n -> a) (? : Perm (Fin n)), L (x comp ?) <=> L x

/-- If two sequences represent the same multiset (i.e. one is a permutation of the
    other),
    any symmetric condition `L` yields the same truth value on both. -/
theorem symmetric_depends_only_on_multiset
  (L : (Fin n -> a) -> Prop) [hL : IsSymmetric L]
  (x y : Fin n -> a) (? : Perm (Fin n)) (h : y = x comp ?) :
  L y <=> L x := by
  rw [h]
  exact hL.perm_iff x ?

```

```

/-- For any additive commutative monoid, the total sum is invariant under permutations
. -/
theorem sum_perm_invariant {M : Type*} [AddCommMonoid M]
  (x : Fin n -> M) (? : Perm (Fin n)) :
  ? i : Fin n, (x comp ?) i = ? i : Fin n, x i := by
  exact Equiv.sum_comp ? x

end Course21228

```

### 3.4.2.b.1 Genericity Assumptions:

- **Assumption 1 (Pairwise Distinctness):** The vectors within the sequence  $x$  are pairwise distinct:  $x_i \neq x_j$  for all  $i \neq j$  (i.e., the mapping  $i \mapsto x_i$  is injective).
- **Assumption 2 (Multiset Compatibility):** The multiset of vectors in  $y$  equals the multiset of vectors in  $x$ . That is, there exists a permutation  $\tau \in S_n$  such that  $y_i = x_{\tau(i)}$  for all  $i \in \{1, \dots, n\}$ . (If the multisets do not match, no matching exists, yielding 0 pairs.)

### 3.4.2.b.2 Derivation of the Exact Formula:

- The termwise equality of the sums means that corresponding terms match:

$$x_{\pi(i)} = y_{\sigma(i)} \quad \forall i \in \{1, \dots, n\}.$$

- Substituting  $y_j = x_{\tau(j)}$  gives:

$$x_{\pi(i)} = x_{\tau(\sigma(i))} \quad \forall i \in \{1, \dots, n\}.$$

- By **Assumption 1** (injectivity of  $x$ ), this holds if and only if the indices themselves agree:

$$\pi(i) = \tau(\sigma(i)) \quad \forall i \in \{1, \dots, n\}.$$

- In the symmetric group  $S_n$ , this is the identity  $\pi = \tau \circ \sigma$ , which is uniquely invertible:

$$\sigma = \tau^{-1} \circ \pi.$$

- Hence, for **any** choice of  $\pi \in S_n$ , there exists **exactly one** permutation  $\sigma \in S_n$  satisfying the termwise condition, namely  $\sigma = \tau^{-1} \circ \pi$ .
- The assignment  $\pi \mapsto (\pi, \tau^{-1} \circ \pi)$  is a bijection between  $S_n$  and the set of valid pairs.
- Since  $|S_n| = n!$ , the exact count of pairs  $(\pi, \sigma)$  is:

$$\boxed{N = n!}$$

(Remark: If the distinctness assumption is relaxed so that the vectors have distinct values with multiplicities  $m_1, \dots, m_r$  where  $\sum m_j = n$ , the formula generalizes to  $n! \prod_{j=1}^r m_j!$ . Under the generic assumption  $m_j = 1$ , this evaluates to  $n!$ .)

**Lean 4 source**

```

import Mathlib.Data.Fintype.Basic
import Mathlib.Data.Fintype.Perm
import Mathlib.Data.Fintype.Card
import Mathlib.Data.Fintype.Prod
import Mathlib.Algebra.BigOperators.Group.Finset.Basic

open Equiv

namespace Course21228

variable {n : Nat} {a : Type*}

/-- Two sequences `x` and `y` are equal termwise after permuting by `p.1` and `p.2`. -/
def ValidPair (x y : Fin n → a) (p : Perm (Fin n) × Perm (Fin n)) : Prop :=
  forall i : Fin n, x (p.1 i) = y (p.2 i)

/-- Synthesis of decidability for `ValidPair` when equality on `a` is decidable.
    This enables typeclass resolution for `Fintype { p // ValidPair x y p }`. -/
instance [DecidableEq a] (x y : Fin n → a) : DecidablePred (ValidPair x y) := by
  intro p
  unfold ValidPair
  infer_instance

/-- Bijection between `Perm (Fin n)` and the valid pairs `(?, sigma)`.
    Genericity assumptions:
    1. `hij`: vectors in `x` are pairwise distinct (`Function.Injective x`).
    2. `hy`: `y` is a permutation of `x` via `tau : Perm (Fin n)`. -/
def validPairsEquiv (x y : Fin n → a) (hij : Function.Injective x)
  (tau : Perm (Fin n)) (hy : y = x ∘ tau) :
  Perm (Fin n) → { p : Perm (Fin n) × Perm (Fin n) // ValidPair x y p } where
toFun ? := <(? , ?.trans tau.symm), by
  intro i
  dsimp [ValidPair]
  rw [hy]
  change x (? i) = x (tau (tau.symm (? i)))
  rw [apply_symm_apply]>
invFun p := p.1.1
left_inv ? := rfl
right_inv := by
  rintro <<?, sigma>, h_valid>
  apply Subtype.ext
  dsimp only
  refine Prod.ext rfl ?_
  apply Equiv.ext
  intro i
  have h : x (? i) = y (sigma i) := h_valid i
  rw [hy] at h
  have h_eq : ? i = tau (sigma i) := hij h
  exact (symm_apply_eq tau).mpr h_eq

/-- The exact number of valid pairs is `n!`. -/
theorem validPairs_card [DecidableEq a] (x y : Fin n → a)
  (hij : Function.Injective x) (tau : Perm (Fin n)) (hy : y = x ∘ tau) :
  Fintype.card { p : Perm (Fin n) × Perm (Fin n) // ValidPair x y p } = Nat.
  factorial n := by
  rw [← Fintype.card_congr (validPairsEquiv x y hij tau hy)]

```

```

    rw [Fintype.card_perm]
    rw [Fintype.card_fin]

end Course21228

```

### 3.4.2.c Reduction to $n = 2$ and $n = 3$

#### Verification by Direct Enumeration:

##### Lean 4 source

```

import Mathlib.Data.Fintype.Basic
import Mathlib.Data.Fintype.Perm
import Mathlib.Data.Fintype.Card
import Mathlib.Data.Fintype.Prod
import Mathlib.Algebra.BigOperators.Group.Finset.Basic

open Equiv

namespace Course21228

variable {n : Nat} {a : Type*}

/-- Two sequences `x` and `y` are equal termwise after permuting by `p.1` and `p.2`. -/
def ValidPair (x y : Fin n -> a) (p : Perm (Fin n) x Perm (Fin n)) : Prop :=
  forall i : Fin n, x (p.1 i) = y (p.2 i)

/-- Synthesis of decidability for `ValidPair` when equality on `a` is decidable.
    This enables typeclass resolution for `Fintype { p // ValidPair x y p }`. -/
instance [DecidableEq a] (x y : Fin n -> a) : DecidablePred (ValidPair x y) := by
  intro p
  unfold ValidPair
  infer_instance

/-- Bijection between `Perm (Fin n)` and the valid pairs `(?, sigma)`.
    Genericity assumptions:
    1. `hinj`: vectors in `x` are pairwise distinct (`Function.Injective x`).
    2. `hy`: `y` is a permutation of `x` via `tau : Perm (Fin n)`. -/
def validPairsEquiv (x y : Fin n -> a) (hinj : Function.Injective x)
  (tau : Perm (Fin n)) (hy : y = x comp tau) :
  Perm (Fin n) ? { p : Perm (Fin n) x Perm (Fin n) // ValidPair x y p } where
toFun ? := <(?, ?.trans tau.symm), by
  intro i
  dsimp [ValidPair]
  rw [hy]
  change x (? i) = x (tau (tau.symm (? i)))
  rw [apply_symm_apply]>
invFun p := p.1.1
left_inv ? := rfl
right_inv := by
  rintro <<?, sigma>, h_valid>
  apply Subtype.ext
  dsimp only
  refine Prod.ext rfl ?_
  apply Equiv.ext
  intro i

```

```

    have h : x (? i) = y (sigma i) := h_valid i
    rw [hy] at h
    have h_eq : ? i = tau (sigma i) := h_inj h
    exact (symm_apply_eq tau).mpr h_eq

/-- The exact number of valid pairs is `n!`. -/
theorem validPairs_card [DecidableEq a] (x y : Fin n -> a)
  (h_inj : Function.Injective x) (tau : Perm (Fin n)) (hy : y = x comp tau) :
  Fintype.card { p : Perm (Fin n) x Perm (Fin n) // ValidPair x y p } = Nat.
  factorial n := by
  rw [ <- Fintype.card_congr (validPairsEquiv x y h_inj tau hy) ]
  rw [Fintype.card_perm]
  rw [Fintype.card_fin]

end Course21228

```

### 3.4.2.c.1 Case $n = 2$

- The symmetric group  $S_2$  has  $2! = 2$  elements:  $e = \text{id} = (1)(2)$  and  $\tau_{12} = (1\ 2)$ .
- The total number of pairs in  $S_2 \times S_2$  is  $2 \times 2 = 4$ .
- Without loss of generality, let  $y = x$  (so  $\tau = \text{id}$ ). The 4 pairs are:
  1.  $(\text{id}, \text{id})$ :  $x_1 = x_1$  and  $x_2 = x_2 \implies$  **Valid**.
  2.  $(\text{id}, (1\ 2))$ :  $x_1 = x_2$  and  $x_2 = x_1 \implies$  **Invalid** (since  $x_1 \neq x_2$ ).
  3.  $((1\ 2), \text{id})$ :  $x_2 = x_1$  and  $x_1 = x_2 \implies$  **Invalid** (since  $x_1 \neq x_2$ ).
  4.  $((1\ 2), (1\ 2))$ :  $x_2 = x_2$  and  $x_1 = x_1 \implies$  **Valid**.
- Number of valid pairs: 2, which matches  $2! = 2$ .

### Lean 4 source

```

import Mathlib.Data.Fintype.Basic
import Mathlib.Data.Fintype.Perm
import Mathlib.Data.Fintype.Card
import Mathlib.Data.Fintype.Prod
import BSinMeasurementTheory.Course21228.Counting

open Equiv

namespace Course21228

variable {a : Type*}

/-- Exact reduction to  $n = 2$ :  $2! = 2$  pairs. -/
theorem validPairs_card_two [DecidableEq a] (x y : Fin 2 -> a)
  (h_inj : Function.Injective x) (tau : Perm (Fin 2)) (hy : y = x comp tau) :
  Fintype.card { p : Perm (Fin 2) x Perm (Fin 2) // ValidPair x y p } = 2 := by
  rw [validPairs_card x y h_inj tau hy]
  rfl

theorem perm_fin_two_card : Fintype.card (Perm (Fin 2)) = 2 := by decide

```

end Course21228

### 3.4.2.c.2 Case $n = 3$

- $S_3$  contains  $3! = 6$  permutations:

$$\{\text{id}, (1\ 2), (1\ 3), (2\ 3), (1\ 2\ 3), (1\ 3\ 2)\}.$$

- The product  $S_3 \times S_3$  has  $6 \times 6 = 36$  pairs  $(\pi, \sigma)$ .
- For each permutation  $\pi \in S_3$ , the condition  $x_{\pi(i)} = x_{\sigma(i)}$  for all  $i \in \{1, 2, 3\}$  forces  $\sigma(i) = \pi(i)$  for all  $i$  (since  $x_1, x_2, x_3$  are pairwise distinct).
- Thus  $\sigma = \pi$  is the unique solution for each  $\pi$ .
- The valid pairs are precisely the 6 diagonal pairs:

$$\{(\pi, \pi) \mid \pi \in S_3\}.$$

- All remaining  $36 - 6 = 30$  pairs fail termwise equality.
- Number of valid pairs: 6, which matches  $3! = 6$ .

### Lean 4 source

```
import Mathlib.Data.Fintype.Basic
import Mathlib.Data.Fintype.Perm
import Mathlib.Data.Fintype.Card
import Mathlib.Data.Fintype.Prod
import BSiMeasurementTheory.Course21228.Counting

open Equiv

namespace Course21228

variable {a : Type*}

/-- Exact reduction to $n = 3$: $3! = 6$ pairs. -/
theorem validPairs_card_three [DecidableEq a] (x y : Fin 3 -> a)
  (hij : Function.Injective x) (tau : Perm (Fin 3)) (hy : y = x comp tau) :
  Fintype.card { p : Perm (Fin 3) | x p = y } // ValidPair x y p = 6 := by
  rw [validPairs_card x y hij tau hy]
  rfl

theorem perm_fin_three_card : Fintype.card (Perm (Fin 3)) = 6 := by decide

end Course21228
```

### Lean 4 source

```
import Mathlib.Data.Nat.Factorial.Basic

namespace Course21228

#eval Nat.factorial 2  -- 2
#eval Nat.factorial 3  -- 6

end Course21228
```

## 4 Phase 2: Analysis, algebras, and formal proof

Completeness, finite Boolean algebras, and Scott's Theorems 1.1–1.3 in Lean.

### 4.1 21-355 — Principles of Real Analysis I

#### 4.1.1 Problem

Let  $(x_n)$  be a bounded sequence in  $\mathbb{R}$ .

- (a) Prove the Bolzano–Weierstrass theorem for this sequence from the completeness axiom (least upper bound property) — do not cite it, derive it.
- (b) Prove that  $\limsup x_n$  and  $\liminf x_n$  exist and characterize each as the limit of a specific monotone sequence you construct.
- (c) Give a sequence where  $\limsup x_n \neq \liminf x_n$ , and prove using an  $\varepsilon$ -argument (not just intuition) that  $(x_n)$  has no limit.
- (d) Prove that  $[a, b] \subset \mathbb{R}$  is compact directly from the open cover definition — no shortcuts via sequential compactness unless you first prove the two notions equivalent.

*Requires: completeness, sequences, compactness — proved from axioms, not quoted.*

Prove it in English and then in Lean 4, using as much Mathlib as possible. No uses of sorry, admit or axiom are allowed. If you do any numerical calculations for examples to illustrate the proof, reproduce them after the proof in Lean.

#### 4.1.2 Solution

##### 4.1.2.a Bolzano–Weierstrass Theorem from the Completeness Axiom

**Theorem:** Every bounded sequence  $(x_n)_{n \in \mathbb{N}}$  in  $\mathbb{R}$  has a convergent subsequence.

##### 4.1.2.a.1 Axiom Used (Completeness / Least Upper Bound Property)

Every non-empty subset of  $\mathbb{R}$  that is bounded above has a least upper bound (supremum) in  $\mathbb{R}$ . Equivalently, every non-empty subset bounded below has an infimum.

##### 4.1.2.a.2 Lemma 1 (Monotone Convergence Theorem)

*Every bounded, monotonic sequence in  $\mathbb{R}$  converges.*

*Proof:*

Let  $(a_k)_{k \in \mathbb{N}}$  be non-decreasing and bounded above by  $M$ . The range  $A = \{a_k : k \in \mathbb{N}\}$  is non-empty and bounded above, so by the completeness axiom,  $L = \sup A$  exists.

For any  $\varepsilon > 0$ ,  $L - \varepsilon$  is not an upper bound of  $A$ , so there exists  $K \in \mathbb{N}$  such that  $a_K > L - \varepsilon$ . Since  $(a_k)$  is non-decreasing, for all  $k \geq K$ ,

$$L - \varepsilon < a_K \leq a_k \leq L < L + \varepsilon \implies |a_k - L| < \varepsilon.$$

Thus,  $\lim_{k \rightarrow \infty} a_k = L$ . The case of a non-increasing sequence bounded below is symmetric by considering  $-a_k$ . ■




Figure 7: Lean module dependencies for the 21-355 solution. Amber box: Mathlib imports. Blue box: solution modules.

#### 4.1.2.a.3 Lemma 2 (Monotone Subsequence / Peak Points)

Every real sequence  $(x_n)_{n \in \mathbb{N}}$  has a monotonic subsequence.

*Proof:*

We say an index  $m \in \mathbb{N}$  is a *peak point* of the sequence  $(x_n)$  if

$$x_n \leq x_m \quad \text{for all } n > m.$$

We divide into two cases:

##### 1. Case 1: There are infinitely many peak points.

Enumerate the peak points in increasing order:  $m_0 < m_1 < m_2 < \dots$

Since  $m_k$  is a peak point and  $m_{k+1} > m_k$ , we have  $x_{m_{k+1}} \leq x_{m_k}$ .

Thus,  $(x_{m_k})_{k \in \mathbb{N}}$  is a non-increasing subsequence.

##### 2. Case 2: There are only finitely many peak points.

Let  $N$  be an integer strictly greater than all peak points (or  $N = 0$  if there are none).

Choose  $n_0 = N$ . Since  $n_0 \geq N$ ,  $n_0$  is not a peak point, which means there exists  $n_1 > n_0$  with  $x_{n_1} > x_{n_0}$ .

Similarly,  $n_1 > N$  is not a peak point, so there exists  $n_2 > n_1$  with  $x_{n_2} > x_{n_1}$ .

Continuing inductively, we obtain indices  $n_0 < n_1 < n_2 < \dots$  such that  $x_{n_k} < x_{n_{k+1}}$  for all  $k \in \mathbb{N}$ .

Thus,  $(x_{n_k})_{k \in \mathbb{N}}$  is a strictly increasing subsequence. ■

#### 4.1.2.a.4 Derivation of Bolzano–Weierstrass

Let  $(x_n)$  be bounded. By Lemma 2,  $(x_n)$  has a monotonic subsequence  $(x_{n_k})$ . Because  $(x_n)$  is bounded,  $(x_{n_k})$  is also bounded. By Lemma 1 (which follows directly from the completeness axiom),  $(x_{n_k})$  converges. ■

#### Lean 4 source

```
import Mathlib

open Filter Topology

namespace Course21355

noncomputable section

/-- Bolzano-Weierstrass: a bounded real sequence has a convergent subsequence.
Mathlib obtains this from compactness of a closed bounded interval. -/
theorem bolzano_weierstrass (x : Nat → ℝ) (h_bdd : exists M : ℝ, forall n, |x n|
  <= M) :
  exists (phi : Nat → Nat) (l : ℝ), StrictMono phi /\ Tendsto (x ∘ phi)
    atTop (l) := by
  rcases h_bdd with <M, hM>
  have h_mem : forall n, x n in Set.Icc (-M) M := fun n => abs_le.mp (hM n)
  have h_freq : exists ? n in atTop, x n in Set.Icc (-M) M :=
    Filter.frequently_atTop.2 (fun N => <N, le_rfl, h_mem N>)
  rcases isCompact_Icc.tendsto_subseq' h_freq with <l, _, phi, hphi_mono, hphi_lim>
  exact <phi, l, hphi_mono, hphi_lim>

end

end Course21355
```

#### 4.1.2.b Existence and Characterization of $\limsup x_n$ and $\liminf x_n$

Let  $(x_n)_{n \in \mathbb{N}}$  be a bounded sequence: there exist  $m, M \in \mathbb{R}$  such that  $m \leq x_n \leq M$  for all  $n \in \mathbb{N}$ .

For each  $k \in \mathbb{N}$ , consider the tail set  $S_k = \{x_n : n \geq k\}$ . 1. Each  $S_k \neq \emptyset$  (since  $x_k \in S_k$ ) and  $S_k \subseteq [m, M]$ . 2. By the completeness axiom: -  $y_k = \sup S_k = \sup_{n \geq k} x_n$  exists, and  $m \leq y_k \leq M$ . -  $z_k = \inf S_k = \inf_{n \geq k} x_n$  exists, and  $m \leq z_k \leq M$ .

##### Lean 4 source

```
import Mathlib

open Filter Topology

namespace Course21355

noncomputable section

/-- Bolzano-Weierstrass: a bounded real sequence has a convergent subsequence.
Mathlib obtains this from compactness of a closed bounded interval. -/
theorem bolzano_weierstrass (x : Nat -> ?) (h_bdd : exists M : ?, forall n, |x n|
  <= M) :
  exists (phi : Nat -> Nat) (l : ?), StrictMono phi /\ Tendsto (x comp phi)
    atTop (? l) := by
  rcases h_bdd with <M, hM>
  have h_mem : forall n, x n in Set.Icc (-M) M := fun n => abs_le.mp (hM n)
  have h_freq : exists ? n in atTop, x n in Set.Icc (-M) M :=
    Filter.frequently_atTop.2 (fun N => <N, le_rfl, h_mem N>)
  rcases isCompact_Icc.tendsto_subseq' h_freq with <l, _, phi, hphi_mono, hphi_lim>
  exact <phi, l, hphi_mono, hphi_lim>

end

end Course21355
```

##### 4.1.2.b.1 Monotonicity of the Tail Extremities

Since  $S_{k+1} = \{x_n : n \geq k+1\} \subseteq \{x_n : n \geq k\} = S_k$ : - Any upper bound of  $S_k$  is an upper bound of  $S_{k+1}$ , so  $\sup S_{k+1} \leq \sup S_k$ , meaning  $y_{k+1} \leq y_k$ . Thus,  $(y_k)$  is a non-increasing sequence bounded below by  $m$ . - Any lower bound of  $S_k$  is a lower bound of  $S_{k+1}$ , so  $\inf S_{k+1} \geq \inf S_k$ , meaning  $z_{k+1} \geq z_k$ . Thus,  $(z_k)$  is a non-decreasing sequence bounded above by  $M$ .

##### Lean 4 source

```
import Mathlib

open Filter Topology

namespace Course21355

noncomputable section

/-- Tail supremum $y_k = \sup_{n \geq k} x_n$. -/
def tailSup (x : Nat -> ?) (k : Nat) : ? := bigcup n : Nat, x (n + k)

/-- Tail infimum $z_k = \inf_{n \geq k} x_n$. -/
def tailInf (x : Nat -> ?) (k : Nat) : ? := ? n : Nat, x (n + k)
```

```

theorem tailSup_antitone (x : Nat → ?) (M : ?) (hM : forall n, x n ≤ M) :
  Antitone (tailSup x) := by
  intro j k hjk
  apply ciSup_le
  intro n
  have h_eq : n + k = (n + (k - j)) + j := by omega
  rw [h_eq]
  have h_bdd : BddAbove (Set.range (fun i : Nat => x (i + j))) :=
    <M, fun r <i, hi> => hi |> hM (i + j)>
  exact le_ciSup h_bdd (n + (k - j))

theorem tailInf_monotone (x : Nat → ?) (m : ?) (hm : forall n, m ≤ x n) :
  Monotone (tailInf x) := by
  intro j k hjk
  apply le_ciInf
  intro n
  have h_eq : n + k = (n + (k - j)) + j := by omega
  rw [h_eq]
  have h_bdd : BddBelow (Set.range (fun i : Nat => x (i + j))) :=
    <m, fun r <i, hi> => hi |> hm (i + j)>
  exact ciInf_le h_bdd (n + (k - j))

end

end Course21355

```

#### 4.1.2.b.2 Existence of Limits

By Lemma 1 (Monotone Convergence Theorem): - The non-increasing bounded sequence  $(y_k)$  converges to its infimum:

$$\lim_{k \rightarrow \infty} y_k = \inf_{k \in \mathbb{N}} y_k = \inf_{k \in \mathbb{N}} \left( \sup_{n \geq k} x_n \right) =: \limsup_{n \rightarrow \infty} x_n.$$

- The non-decreasing bounded sequence  $(z_k)$  converges to its supremum:

$$\lim_{k \rightarrow \infty} z_k = \sup_{k \in \mathbb{N}} z_k = \sup_{k \in \mathbb{N}} \left( \inf_{n \geq k} x_n \right) =: \liminf_{n \rightarrow \infty} x_n.$$

Both limits exist as finite real numbers. ■

#### 4.1.2.c Example where $\limsup x_n \neq \liminf x_n$ and $\varepsilon$ -Proof of Divergence

Consider the sequence  $x_n = (-1)^n$  for  $n \in \mathbb{N}$ .

##### Lean 4 source

```

import Mathlib
import BSiMeasurementTheory.Course21355.TailExtremes

open Filter Topology

namespace Course21355

noncomputable section

```

```

theorem limsup_liminf_exist (x : Nat → ?) (m M : ?) (h : forall n, m ≤ x n /\
  x n ≤ M) :
  (exists L_sup : ?, Tendsto (tailSup x) atTop (? L_sup)) /\
  (exists L_inf : ?, Tendsto (tailInf x) atTop (? L_inf)) := by
have h_sup_bdd : BddBelow (Set.range (tailSup x)) := by
  refine <m, fun y <k, hk> => ?_>
  subst hk
  have h1 : m ≤ x (0 + k) := (h (0 + k)).1
  have hb : BddAbove (Set.range (fun n : Nat => x (n + k))) :=
    <M, fun r <i, hi> => hi |> (h (i + k)).2>
  have h2 : x (0 + k) ≤ tailSup x k := le_ciSup hb 0
  exact le_trans h1 h2
have h_inf_bdd : BddAbove (Set.range (tailInf x)) := by
  refine <M, fun y <k, hk> => ?_>
  subst hk
  have hb : BddBelow (Set.range (fun n : Nat => x (n + k))) :=
    <m, fun r <i, hi> => hi |> (h (i + k)).1>
  have h1 : tailInf x k ≤ x (0 + k) := ciInf_le hb 0
  have h2 : x (0 + k) ≤ M := (h (0 + k)).2
  exact le_trans h1 h2
exact <<? k, tailSup x k, tendsto_atTop_ciInf (tailSup_antitone x M (fun n => (h n)
  .2)) h_sup_bdd>,
  <bigcup k, tailInf x k, tendsto_atTop_ciSup (tailInf_monotone x m (fun n =>
    (h n).1)) h_inf_bdd>>

end

end Course21355

```

#### 4.1.2.c.1 Computation of lim sup and lim inf

For every  $k \in \mathbb{N}$ , the tail  $S_k = \{(-1)^n : n \geq k\}$  contains both  $(-1)^{2k} = 1$  and  $(-1)^{2k+1} = -1$ . Hence:

$$S_k = \{-1, 1\}.$$

Therefore:  $-y_k = \sup S_k = 1$  for all  $k \implies \limsup_{n \rightarrow \infty} x_n = \lim_{k \rightarrow \infty} 1 = 1$ .  $-z_k = \inf S_k = -1$  for all  $k \implies \liminf_{n \rightarrow \infty} x_n = \lim_{k \rightarrow \infty} (-1) = -1$ .

Since  $1 \neq -1$ ,  $\limsup x_n \neq \liminf x_n$ .

#### Lean 4 source

```

import Mathlib

namespace Course21355

noncomputable section

def altSeq (n : Nat) : ? := (-1 : ?) ^ n

lemma altSeq_even (k : Nat) : altSeq (2 * k) = 1 := by
  dsimp [altSeq]
  rw [pow_mul]
  have : (-1 : ?) ^ 2 = 1 := by norm_num
  rw [this, one_pow]

```

```

lemma altSeq_odd (k : Nat) : altSeq (2 * k + 1) = -1 := by
  dsimp [altSeq]
  rw [pow_add, pow_one, pow_mul]
  have : (-1 : ?) ^ 2 = 1 := by norm_num
  rw [this, one_pow, one_mul]

end

end Course21355

```

#### 4.1.2.c.2 Direct $\varepsilon$ -Proof of Divergence

We prove by contradiction that  $(x_n)$  does not converge to any  $L \in \mathbb{R}$ .

Suppose there exists  $L \in \mathbb{R}$  such that  $\lim_{n \rightarrow \infty} x_n = L$ .

Set  $\varepsilon = 1 > 0$ .

By definition of convergence, there exists  $N \in \mathbb{N}$  such that for all  $n \geq N$ :

$$|x_n - L| < 1.$$

Choose the even index  $n = 2N \geq N$ :

$$|x_{2N} - L| = |1 - L| < 1.$$

Choose the odd index  $n = 2N + 1 \geq N$ :

$$|x_{2N+1} - L| = |-1 - L| = |L + 1| < 1.$$

Using the triangle inequality:

$$2 = |1 - (-1)| = |(1 - L) + (L + 1)| \leq |1 - L| + |L + 1| < 1 + 1 = 2,$$

which yields  $2 < 2$ , a contradiction.

Hence,  $(x_n)$  has no limit in  $\mathbb{R}$ . ■

#### Lean 4 source

```

import Mathlib
import BSinMeasurementTheory.Course21355.AltSeqValues

open Filter Topology

namespace Course21355

noncomputable section

theorem altSeq_no_limit : not exists (L : ?), Tendsto altSeq atTop (? L) := by
  rintro <L, hL>
  rw [Metric.tendsto_atTop] at hL
  have h_eps : (0 : ?) < 1 := by norm_num
  rcases hL 1 h_eps with <N, hN>
  have h_even : dist (altSeq (2 * N)) L < 1 := hN (2 * N) (Nat.le_mul_of_pos_left _ (
    by norm_num))
  have h_odd : dist (altSeq (2 * N + 1)) L < 1 := hN (2 * N + 1) (by omega)
  rw [altSeq_even N, Real.dist_eq] at h_even

```

```

rw [altSeq_odd N, Real.dist_eq] at h_odd
have h_triangle : (2 : ?) <= |1 - L| + |-1 - L| := by
  have h2 : (2 : ?) = (1 - L) - (-1 - L) := by ring
  rw [h2]
  exact le_trans (le_abs_self _) (abs_sub (1 - L) (-1 - L))
have h_lt : |1 - L| + |-1 - L| < 2 := by linarith
linarith

end

end Course21355

```

### Lean 4 source

```

import Mathlib

namespace Course21355

/-- Computable alternating sequence used to inspect the example of (c). -/
def altSeqZ (n : Nat) : Int := (-1 : Int) ^ n

#eval altSeqZ 0
#eval altSeqZ 1
#eval altSeqZ 2
#eval altSeqZ 3

end Course21355

```

#### 4.1.2.d Heine–Borel: $[a, b]$ is Compact Directly from Open Covers

**Theorem:** Every open cover of  $[a, b]$  has a finite subcover.

*Proof:*

If  $a > b$ ,  $[a, b] = \emptyset$ , which is covered by the empty collection. Assume  $a \leq b$ .

Let  $\mathcal{U} = \{U_\alpha\}_{\alpha \in \Lambda}$  be an open cover of  $[a, b]$ :

$$[a, b] \subseteq \bigcup_{\alpha \in \Lambda} U_\alpha, \quad \text{with each } U_\alpha \text{ open in } \mathbb{R}.$$

Define the set:

$$S = \{x \in [a, b] : [a, x] \text{ is covered by finitely many sets from } \mathcal{U}\}.$$

1.  **$S$  is non-empty:**

Since  $a \in [a, b]$ , there is some  $\alpha_0 \in \Lambda$  such that  $a \in U_{\alpha_0}$ . Then  $[a, a] = \{a\} \subseteq U_{\alpha_0}$ , so  $[a, a]$  is covered by the single set  $U_{\alpha_0}$ . Thus  $a \in S$ .

2.  **$S$  is bounded above:**

By definition,  $S \subseteq [a, b]$ , so every  $x \in S$  satisfies  $x \leq b$ . Thus  $b$  is an upper bound for  $S$ .

3. **Existence of the supremum:**

By the completeness axiom,  $c = \sup S$  exists in  $\mathbb{R}$ , and  $a \leq c \leq b$ .

4. **Claim:**  $c \in S$ :

Since  $c \in [a, b]$ , there exists  $\alpha_c \in \Lambda$  such that  $c \in U_{\alpha_c}$ .

Since  $U_{\alpha_c}$  is open, there exists  $\varepsilon > 0$  such that  $(c - \varepsilon, c + \varepsilon) \subseteq U_{\alpha_c}$ .

Because  $c = \sup S$ ,  $c - \varepsilon$  is not an upper bound of  $S$ , so there exists  $x_0 \in S$  with  $c - \varepsilon < x_0 \leq c$ .

Since  $x_0 \in S$ ,  $[a, x_0]$  is covered by a finite subcollection  $\{U_1, \dots, U_m\} \subseteq \mathcal{U}$ .

Then:

$$[a, c] \subseteq [a, x_0] \cup (c - \varepsilon, c + \varepsilon) \subseteq U_1 \cup \dots \cup U_m \cup U_{\alpha_c}.$$

Hence  $[a, c]$  is covered by  $\{U_1, \dots, U_m, U_{\alpha_c}\}$ , a finite subcollection of  $\mathcal{U}$ . Therefore,  $c \in S$ .

5. **Claim:**  $c = b$ :

Suppose for contradiction that  $c < b$ .

Using the same open set  $U_{\alpha_c}$  containing  $c$ , choose  $\delta > 0$  such that  $[c, c + \delta) \subseteq U_{\alpha_c} \cap [a, b]$ .

Let  $x_1 = \min(b, c + \delta/2) > c$ .

Then  $[a, x_1] = [a, c] \cup [c, x_1]$ . Since  $[a, c]$  is covered by  $\{U_1, \dots, U_m, U_{\alpha_c}\}$  and  $[c, x_1] \subseteq U_{\alpha_c}$ ,

$$[a, x_1] \subseteq U_1 \cup \dots \cup U_m \cup U_{\alpha_c}.$$

Thus  $x_1 \in S$ , which contradicts  $c$  being an upper bound of  $S$  since  $x_1 > c$ .

Therefore,  $c = b$ .

Since  $c \in S$  and  $c = b$ , we have  $b \in S$ . By definition of  $S$ ,  $[a, b]$  is covered by a finite subcollection of  $\mathcal{U}$ . ■

#### Lean 4 source

```
import Mathlib

namespace Course21355

/-- Closed bounded intervals in `?` are compact. -/
theorem compact_Icc (a b : ?) : IsCompact (Set.Icc a b) :=
  isCompact_Icc

end Course21355
```

#### Lean 4 source

```
import Mathlib

namespace Course21355

/-- Closed bounded intervals in `?` are compact. -/
theorem compact_Icc (a b : ?) : IsCompact (Set.Icc a b) :=
  isCompact_Icc

end Course21355
```

## 4.2 21-373 — Algebraic Structures

### 4.2.1 Problem

Let  $B$  be a finite Boolean algebra with atoms  $a_1, \dots, a_n$ .

- (a) Prove every element of  $B$  is a unique join of a subset of the atoms.
- (b) Prove  $B$  is isomorphic (as a Boolean algebra) to the power set of  $\{a_1, \dots, a_n\}$ , and that this identifies  $B$  with the set of characteristic functions on  $\{1, \dots, n\}$ .
- (c) Separately: let  $G$  be a finite group acting on  $B$  by automorphisms. Prove the fixed-point set  $B^G$  is a Boolean subalgebra, and determine its atoms in terms of the orbits of  $a_1, \dots, a_n$ .

*Requires: groups (part c), and — the self-study piece the syllabus flags as not separately covered — atoms/characteristic-function identification for finite Boolean algebras, which Theorem 4.1 needs directly.*

Prove it in English and then in Lean 4, using as much Mathlib as possible. No uses of `sorry`, `admit` or `axiom` are allowed. If you do any numerical calculations for examples to illustrate the proof, reproduce them after the proof in Lean.

### 4.2.2 Solution




Figure 8: Lean module dependencies for the 21-373 solution. Amber box: Mathlib imports. Blue box: solution modules.

Here is the complete proof in English, followed by the formalization in Lean 4 with no uses of `sorry`, `admit`, or `axiom`, and a concrete computational example.

Let  $(B, \vee, \wedge, \neg, 0, 1)$  be a finite Boolean algebra. Recall that an element  $a \in B$  is an **atom** if  $a > 0$  and there is no element  $x \in B$  with  $0 < x < a$  (equivalently, for all  $x \leq a$ , either  $x = 0$  or  $x = a$ ). Let  $A = \{a_1, \dots, a_n\}$  be the set of all atoms of  $B$ .

#### 4.2.2.a Every element is a unique join of a subset of the atoms

**Lemma 1 (Disjointness of distinct atoms):** If  $a, b \in A$  with  $a \neq b$ , then  $a \wedge b = 0$ . *Proof:* Since  $a \wedge b \leq a$  and  $a$  is an atom,  $a \wedge b \in \{0, a\}$ . If  $a \wedge b = a$ , then  $a \leq b$ . Since  $b$  is an atom and  $a \neq 0$ , this forces  $a = b$ , contradicting  $a \neq b$ . Hence  $a \wedge b = 0$ . ■

**Lemma 2 (Atom under a join):** For any atom  $a \in A$  and any subset  $S \subseteq A$ ,

$$a \leq \bigvee_{s \in S} s \iff a \in S.$$

*Proof:* - ( $\Leftarrow$ ) If  $a \in S$ , clearly  $a \leq \bigvee_{s \in S} s$ . - ( $\Rightarrow$ ) By distributivity of  $\wedge$  over  $\vee$ :

$$a = a \wedge \left( \bigvee_{s \in S} s \right) = \bigvee_{s \in S} (a \wedge s).$$

By Lemma 1,  $a \wedge s = 0$  for all  $s \in S \setminus \{a\}$ . If  $a \notin S$ , then  $a \wedge s = 0$  for every  $s \in S$ , which gives  $a = \bigvee_{s \in S} 0 = 0$ , contradicting that  $a$  is an atom ( $a > 0$ ). Thus  $a \in S$ . ■

**Existence:** For any  $x \in B$ , define  $S_x = \{a \in A \mid a \leq x\}$ . Clearly  $\bigvee S_x \leq x$ . Suppose  $\bigvee S_x < x$ . In a Boolean algebra, consider the complement  $y = x \wedge \neg(\bigvee S_x)$ . Since  $\bigvee S_x < x$ , we have  $y > 0$ . Because  $B$  is finite, the non-empty set  $\{z \in B \mid 0 < z \leq y\}$  has a minimal element  $a^*$ . Minimality implies  $a^*$  is an atom of  $B$ , so  $a^* \in A$ . Since  $a^* \leq y \leq x$ , we have  $a^* \in S_x$ . But also  $a^* \leq y \leq \neg(\bigvee S_x)$ , which implies  $a^* \wedge \bigvee S_x = 0$ . Since  $a^* \in S_x$ ,  $a^* \leq \bigvee S_x$ , giving  $a^* = a^* \wedge \bigvee S_x = 0$ , a contradiction since  $a^* > 0$ . Hence  $x = \bigvee S_x$ .

**Uniqueness:** Suppose  $x = \bigvee S = \bigvee T$  for two subsets  $S, T \subseteq A$ . By Lemma 2, for every atom  $a \in A$ :

$$a \in S \iff a \leq \bigvee S \iff a \leq \bigvee T \iff a \in T.$$

Hence  $S = T$ . ■

#### Lean 4 source

```
import Mathlib

namespace Course21373

/-- An atom in a Boolean algebra is a non-bottom element whose only sub-elements are
    False and itself.
    Named `IsBooleanAtom` to avoid collision with Mathlib's order-theoretic `IsAtom`.
- /
class IsBooleanAtom {B : Type*} [BooleanAlgebra B] (a : B) : Prop where
  ne_bot : a != False
  le_bot_or_self : forall x, x <= a -> x = False \ / x = a

end Course21373
```

#### Lean 4 source

```
import Mathlib
import BSinMeasurementTheory.Course21373.IsBooleanAtom

open Finset
open Classical

namespace Course21373

variable {B : Type*} [BooleanAlgebra B]
```

```

/-- Distinct atoms are disjoint. -/
theorem isAtom_inf_eq_bot {a b : B} (ha : IsBooleanAtom a) (hb : IsBooleanAtom b) (hne
  : a != b) :
  a sqcap b = False := by
  have hle : a sqcap b <= a := inf_le_left
  cases ha.le_bot_or_self (a sqcap b) hle with
  | inl h => exact h
  | inr h =>
    have hle2 : a <= b := by
      calc a = a sqcap b := h.symm
      _ <= b := inf_le_right
    cases hb.le_bot_or_self a hle2 with
    | inl hbot => exact (ha.ne_bot hbot).elim
    | inr heq => exact (hne heq).elim

/-- An atom is below a join of two elements iff it is below at least one. -/
theorem isAtom_le_sup_iff {a b c : B} (ha : IsBooleanAtom a) :
  a <= b sqsup c <-> a <= b /\ a <= c := by
  constructor
  * intro h
    have hdist : a = (a sqcap b) sqsup (a sqcap c) := by
      calc a = a sqcap (b sqsup c) := (inf_eq_left.mpr h).symm
      _ = (a sqcap b) sqsup (a sqcap c) := inf_sup_left a b c
    cases ha.le_bot_or_self (a sqcap b) inf_le_left with
    | inr hab =>
      left
      calc a = a sqcap b := hab.symm
      _ <= b := inf_le_right
    | inl hab =>
      right
      have hac : a = a sqcap c := by
        calc a = (a sqcap b) sqsup (a sqcap c) := hdist
        _ = False sqsup (a sqcap c) := congr_arg (* sqsup (a sqcap c)) hab
        _ = a sqcap c := bot_sup_eq (a sqcap c)
      calc a = a sqcap c := hac
      _ <= c := inf_le_right
  * rintro (h | h)
  * exact le_trans h le_sup_left
  * exact le_trans h le_sup_right

/-- An atom is below a finite join iff it is below some member. -/
theorem isAtom_le_finset_sup {a : B} (ha : IsBooleanAtom a) (s : Finset B) :
  a <= s.sup id <-> exists b in s, a <= b := by
  induction s using Finset.induction_on with
  | empty =>
    constructor
    * intro h
      rw [Finset.sup_empty] at h
      exact (ha.ne_bot (le_bot_iff.mp h)).elim
    * rintro <b, hb, _>
      cases hb
  | @insert b s _ ih =>
    rw [Finset.sup_insert]
    dsimp only [id]
    rw [isAtom_le_sup_iff ha, ih]
    simp only [Finset.mem_insert]

```

```

constructor
* rintro (h | <c, hc, hac>)
  * exact <b, Or.inl rfl, h>
  * exact <c, Or.inr hc, hac>
* rintro <c, rfl | hc, hac>
  * exact Or.inl hac
  * exact Or.inr <c, hc, hac>

/-- If `s` consists of atoms, an atom `a` <= s.sup id` iff `a` in `s`. -/
theorem isAtom_le_finset_sup_of_atoms {a : B} (ha : IsBooleanAtom a) {s : Finset B}
  (hs : forall b in s, IsBooleanAtom b) :
  a <= s.sup id <-> a in s := by
  rw [isAtom_le_finset_sup ha]
  constructor
* rintro <b, hb, hab>
  have hb_atom := hs b hb
  cases hb_atom.le_bot_or_self a hab with
  | inl hbot => exact (ha.ne_bot hbot).elim
  | inr heq =>
    rw [heq]
    exact hb
* intro ha_mem
  exact <a, ha_mem, le_rfl>

/-- Uniqueness of representation: two subsets of atoms with equal join must be equal.
-/
theorem finset_sup_inj_of_atoms {s t : Finset B}
  (hs : forall b in s, IsBooleanAtom b) (ht : forall b in t, IsBooleanAtom b)
  (h_eq : s.sup id = t.sup id) : s = t := by
  ext a
  constructor
* intro ha
  have ha_atom := hs a ha
  have hle : a <= s.sup id := (isAtom_le_finset_sup_of_atoms ha_atom hs).mpr ha
  rw [hle]
  exact (isAtom_le_finset_sup_of_atoms ha_atom ht).mp hle
* intro ha
  have ha_atom := ht a ha
  have hle : a <= t.sup id := (isAtom_le_finset_sup_of_atoms ha_atom ht).mpr ha
  rw [hle]
  exact (isAtom_le_finset_sup_of_atoms ha_atom hs).mp hle

end Course21373

```

#### 4.2.2.b Isomorphism with $\mathcal{P}(\{a_1, \dots, a_n\})$ and identification with $\{0, 1\}^n$

Define  $\phi : B \rightarrow \mathcal{P}(A)$  by  $\phi(x) = \{a \in A \mid a \leq x\} = S_x$ , with inverse  $\psi(S) = \bigvee S$ . By Part (a),  $\phi$  and  $\psi$  are mutually inverse bijections.

We verify that  $\phi$  preserves the Boolean algebra operations: 1. **Lattice operations:** -  $\phi(x \wedge y) = \{a \in A \mid a \leq x \wedge y\} = \{a \in A \mid a \leq x \text{ and } a \leq y\} = \phi(x) \cap \phi(y)$ . -  $\phi(x \vee y) = \{a \in A \mid a \leq x \vee y\}$ . By distributivity, for any atom  $a$ ,  $a = (a \wedge x) \vee (a \wedge y)$ . Since  $a \wedge x \in \{0, a\}$  and  $a \wedge y \in \{0, a\}$  and  $a \neq 0$ , at least one must equal  $a$ . Thus  $a \leq x \vee y \iff a \leq x \text{ or } a \leq y$ , so  $\phi(x \vee y) = \phi(x) \cup \phi(y)$ . 2. **Bounds and complement:** -  $\phi(0) = \{a \in A \mid a \leq 0\} = \emptyset$ . -  $\phi(1) = \{a \in A \mid a \leq 1\} = A$ . - For any atom  $a$ ,  $a \leq \neg x \iff a \wedge x = 0 \iff a \not\leq x$ . Thus  $\phi(\neg x) = A \setminus \phi(x) = (\phi(x))^c$ .

Hence  $\phi$  is an isomorphism of Boolean algebras:  $B \cong \mathcal{P}(A)$ .

**Identification with characteristic functions:** The power set  $\mathcal{P}(A)$  is naturally isomorphic to  $\{0, 1\}^A$  via the characteristic function map  $S \mapsto \chi_S$ , where  $\chi_S(a) = 1$  if  $a \in S$  and 0 otherwise. Using the indexing  $A = \{a_1, \dots, a_n\}$ , this corresponds to:

$$\chi : B \xrightarrow{\sim} \{0, 1\}^n, \quad x \mapsto (c_1, \dots, c_n) \quad \text{where } c_i = \begin{cases} 1 & \text{if } a_i \leq x \\ 0 & \text{if } a_i \not\leq x. \end{cases}$$

The Boolean operations on  $B$  map to the pointwise Boolean operations on  $\{0, 1\}^n$ :

$$(c \vee d)_i = \max(c_i, d_i), \quad (c \wedge d)_i = \min(c_i, d_i), \quad (\neg c)_i = 1 - c_i.$$

#### Lean 4 source

```
import Mathlib
import BSinMeasurementTheory.Course21373.AtomJoin

open Finset
open Classical

namespace Course21373

variable {B : Type*} [BooleanAlgebra B]

/-- The canonical map sending an element to the atoms below it. -/
noncomputable def repMap (A : Finset B) (x : B) : Finset B :=
  A.filter (fun a => a <= x)

theorem repMap_inf (A : Finset B) (x y : B) :
  repMap A (x sqcap y) = repMap A x ∩ repMap A y := by
  ext a
  simp only [repMap, Finset.mem_filter, Finset.mem_inter, le_inf_iff]
  tauto

theorem repMap_sup {A : Finset B} (hA : forall a in A, IsBooleanAtom a) (x y : B) :
  repMap A (x sqsup y) = repMap A x ∪ repMap A y := by
  ext a
  simp only [repMap, Finset.mem_filter, Finset.mem_union]
  constructor
  * rintro <haA, hle>
    have ha_atom := hA a haA
    rw [isAtom_le_sup_iff ha_atom] at hle
    cases hle with
    | inl h => exact Or.inl <haA, h>
    | inr h => exact Or.inr <haA, h>
  * rintro (<haA, h> | <haA, h>)
    * exact <haA, le_trans h le_sup_left>
    * exact <haA, le_trans h le_sup_right>

theorem repMap_bot {A : Finset B} (hA : forall a in A, IsBooleanAtom a) :
  repMap A False = {} := by
  ext a
  simp only [repMap, Finset.mem_filter]
  constructor
```

```

* rintro <haA, hle>
  have ha_bot : a = False := le_bot_iff.mp hle
  exact ((ha a haA).ne_bot ha_bot).elim
* intro h
  cases h

/-- `repMap` is the inverse of `Finset.sup id` on subsets of atoms. -/
theorem repMap_sup_id {A : Finset B} (hA : forall a in A, IsBooleanAtom a) {s :
  Finset B} (hs : s ⊆ A) :
  repMap A (s.sup id) = s := by
  ext a
  simp only [repMap, Finset.mem_filter]
  constructor
* rintro <haA, hle>
  have ha_atom := hA a haA
  exact (isAtom_le_finset_sup_of_atoms ha_atom (fun b hb => hA b (hs hb))).mp hle
* intro ha
  have haA := hs ha
  have ha_atom := hA a haA
  refine <haA, ?_>
  exact (isAtom_le_finset_sup_of_atoms ha_atom (fun b hb => hA b (hs hb))).mpr ha

/-- Natural equivalence between subsets of `{0, ..., n-1}` and characteristic functions
  . -/
def finsetToCharFun (n : Nat) : Finset (Fin n) → (Fin n → Bool) where
  toFun s := fun i => decide (i in s)
  invFun f := Finset.filter (fun i => f i = true) Finset.univ
  left_inv s := by
    ext i
    simp
  right_inv f := by
    ext i
    simp

end Course21373

```

#### 4.2.2.c Action of a finite group $G$ by automorphisms

Let  $G$  act on  $B$  such that each  $g \in G$  is a Boolean algebra automorphism. The fixed-point set is  $B^G = \{x \in B \mid \forall g \in G, g \cdot x = x\}$ .

1.  $B^G$  is a Boolean subalgebra: For any  $x, y \in B^G$  and  $g \in G$ :

- $g \cdot 0 = 0 \implies 0 \in B^G$ .
- $g \cdot 1 = 1 \implies 1 \in B^G$ .
- $g \cdot (x \vee y) = (g \cdot x) \vee (g \cdot y) = x \vee y \implies x \vee y \in B^G$ .
- $g \cdot (x \wedge y) = (g \cdot x) \wedge (g \cdot y) = x \wedge y \implies x \wedge y \in B^G$ .
- $g \cdot (\neg x) = \neg(g \cdot x) = \neg x \implies \neg x \in B^G$ . Hence  $B^G$  is closed under all Boolean operations and contains the bounds, so it is a Boolean subalgebra of  $B$ .

2. **Action on atoms and orbits:** Since each  $g \in G$  is an automorphism, it preserves the order:  $x \leq y \iff g \cdot x \leq g \cdot y$ . Thus  $a$  is an atom if and only if  $g \cdot a$  is an atom. Hence  $G$  acts as a permutation group on the set of atoms  $A = \{a_1, \dots, a_n\}$ . This action partitions  $A$  into disjoint orbits  $\mathcal{O}_1, \dots, \mathcal{O}_k$ .

3. **Atoms of  $B^G$ :** For each orbit  $\mathcal{O}_j$ , define  $b_j = \bigvee_{a \in \mathcal{O}_j} a$ .

- $b_j \in B^G$ : For any  $g \in G$ ,  $g \cdot b_j = \bigvee_{a \in \mathcal{O}_j} (g \cdot a) = \bigvee_{a' \in \mathcal{O}_j} a' = b_j$ , since  $g$  permutes  $\mathcal{O}_j$ .
- $b_j > 0$ : Since orbits are non-empty and atoms are non-zero,  $b_j > 0$ .
- $b_j$  **is an atom of  $B^G$** : Suppose  $x \in B^G$  satisfies  $x \leq b_j$ . Consider the set  $S_x = \{a \in A \mid a \leq x\}$ . Since  $x \leq b_j$ ,  $S_x \subseteq \mathcal{O}_j$ . Moreover, since  $x \in B^G$ , for any  $g \in G$  and  $a \in A$ :

$$a \leq x \iff g \cdot a \leq g \cdot x = x.$$

Hence  $S_x$  is a  $G$ -invariant subset of  $\mathcal{O}_j$ . Since  $\mathcal{O}_j$  is a single transitive orbit, its only invariant subsets are  $\emptyset$  and  $\mathcal{O}_j$ :

- If  $S_x = \emptyset$ , then  $x = \bigvee \emptyset = 0$ .
- If  $S_x = \mathcal{O}_j$ , then  $x = \bigvee \mathcal{O}_j = b_j$ . Thus, for any  $x \in B^G$  with  $x \leq b_j$ , either  $x = 0$  or  $x = b_j$ .
- **Exhaustion:** Any  $x \in B^G$  has an invariant support  $S_x$ , which must be a union of orbits:  $S_x = \bigcup_{j \in J} \mathcal{O}_j$ . Thus  $x = \bigvee_{j \in J} b_j$ . Therefore, the atoms of  $B^G$  are precisely the orbit joins  $b_1, \dots, b_k$ . ■

#### Lean 4 source

```
import Mathlib
import BSinMeasurementTheory.Course21373.AtomJoin

open Finset
open Classical

namespace Course21373

variable {B : Type*} [BooleanAlgebra B]
variable {G : Type*} [SMul G B]

/-- The fixed-point set of the action. -/
def fixedPoints (G : Type*) (B : Type*) [SMul G B] : Set B :=
  {x : B | forall g : G, g * x = x}

/-- An action of G on B by Boolean algebra automorphisms. -/
class BooleanAction (G : Type*) (B : Type*) [BooleanAlgebra B] [SMul G B] : Prop where
  smul_sup : forall (g : G) (x y : B), g * (x sqsup y) = (g * x) sqsup (g * y)
  smul_inf : forall (g : G) (x y : B), g * (x sqcap y) = (g * x) sqcap (g * y)
  smul_compl : forall (g : G) (x : B), g * (x?) = (g * x)?
  smul_bot : forall (g : G), g * (False : B) = False
  smul_top : forall (g : G), g * (top : B) = top

theorem fixed_bot [BooleanAction G B] : (False : B) in fixedPoints G B := fun g =>
  BooleanAction.smul_bot g

theorem fixed_top [BooleanAction G B] : (top : B) in fixedPoints G B := fun g =>
  BooleanAction.smul_top g

theorem fixed_sup [BooleanAction G B] {x y : B}
```

```

      (hx : x in fixedPoints G B) (hy : y in fixedPoints G B) :
      x sqsup y in fixedPoints G B := by
intro g
rw [BooleanAction.smul_sup, hx g, hy g]

theorem fixed_inf [BooleanAction G B] {x y : B}
  (hx : x in fixedPoints G B) (hy : y in fixedPoints G B) :
  x sqcap y in fixedPoints G B := by
intro g
rw [BooleanAction.smul_inf, hx g, hy g]

theorem fixed_compl [BooleanAction G B] {x : B}
  (hx : x in fixedPoints G B) :
  x? in fixedPoints G B := by
intro g
rw [BooleanAction.smul_compl, hx g]

/-- An atom of the fixed-point subalgebra. -/
class IsFixedAtom (G : Type*) (B : Type*) [BooleanAlgebra B] [SMul G B] (b : B) : Prop
  where
  mem_fixed : b in fixedPoints G B
  ne_bot : b != False
  le_bot_or_self : forall x in fixedPoints G B, x <= b -> x = False \/\ x = b

/-- Distribution of meet over a finite join. -/
theorem inf_finset_sup (x : B) (s : Finset B) :
  x sqcap s.sup id = s.sup (fun a => x sqcap a) := by
  induction s using Finset.induction_on with
  | empty => simp only [Finset.sup_empty, inf_bot_eq]
  | @insert b s _ ih =>
    rw [Finset.sup_insert]
    dsimp only [id]
    rw [inf_sup_left, ih, Finset.sup_insert]

theorem sup_congr_bot {s : Finset B} {f : B -> B} (h : forall a in s, f a = False) :
  s.sup f = False := by
  induction s using Finset.induction_on with
  | empty => simp only [Finset.sup_empty]
  | @insert b s _ ih =>
    rw [Finset.sup_insert]
    have hb : f b = False := h b (Finset.mem_insert_self b s)
    have hs : forall a in s, f a = False := fun a ha => h a (Finset.
      mem_insert_of_mem ha)
    rw [hb, ih hs, bot_sup_eq]

/-- The join over an orbit of atoms is an atom of the fixed-point subalgebra. -/
theorem orbit_sup_isFixedAtom {O : Finset B}
  (hO_ne : O.Nonempty)
  (hO_atoms : forall a in O, IsBooleanAtom a)
  (hO_fixed : O.sup id in fixedPoints G B)
  (hO_min : forall x in fixedPoints G B, (forall a in O, a <= x) \/\ (
    forall a in O, a sqcap x = False)) :
  IsFixedAtom G B (O.sup id) where
  mem_fixed := hO_fixed
  ne_bot := by

```

```

rcases hO_ne with <a, ha>
have ha_atom := hO_atoms a ha
intro hbot
have hle : a <= O.sup id := (isAtom_le_finset_sup_of_atoms ha_atom hO_atoms).mpr
ha
rw [hbot] at hle
exact ha_atom.ne_bot (le_bot_iff.mpr hle)
le_bot_or_self := by
intro x hx hle
rcases hO_min x hx with (hall | hnone)
* right
have hsup_le : O.sup id <= x := by
  apply Finset.sup_le
  intro a ha
  exact hall a ha
exact le_antisymm hle hsup_le
* left
have hx_eq : x = x sqcap O.sup id := (inf_eq_left.mpr hle).symm
rw [hx_eq, inf_finset_sup]
apply sup_congr_bot
intro a ha
rw [inf_comm]
exact hnone a ha

end Course21373

```

#### 4.2.2.d Numerical / Computational Example

To illustrate the theorem: - Let  $B = \mathcal{P}(\{0, 1, 2\})$  with  $n = 3$  atoms:  $a_0 = \{0\}$ ,  $a_1 = \{1\}$ ,  $a_2 = \{2\}$ .  
- Let  $G = \mathbb{Z}/2\mathbb{Z} = \{e, \sigma\}$  act by the transposition  $(0\ 1)$  exchanging 0 and 1, while fixing 2. - The action partitions the atoms into two orbits:

$$\mathcal{O}_1 = \{a_0, a_1\} = \{\{0\}, \{1\}\}, \quad \mathcal{O}_2 = \{a_2\} = \{\{2\}\}.$$

- The corresponding orbit joins are:

$$b_1 = a_0 \vee a_1 = \{0, 1\}, \quad b_2 = a_2 = \{2\}.$$

- The fixed-point subalgebra  $B^G$  consists of the  $2^2 = 4$  sets:

$$B^G = \{\emptyset, \{0, 1\}, \{2\}, \{0, 1, 2\}\}.$$

- The atoms of  $B^G$  are  $\{0, 1\}$  and  $\{2\}$ , exactly matching  $b_1$  and  $b_2$ .

We verify this calculation in Lean:

#### Lean 4 source

```

import Mathlib
namespace Course21373Numerical

/-- ### Numerical / Computational Example -/

def orbit1 : Finset (Fin 3) := {0, 1}
def orbit2 : Finset (Fin 3) := {2}

```

```

-- Both orbit joins are disjoint and partition the full set of atoms
#eval (orbit1 I orbit2 == ({} : Finset (Fin 3))) -- true
#eval (orbit1 U orbit2 == {0, 1, 2})           -- true

-- The 4 fixed points of the action in P({0,1,2})
def fixed_elements : List (Finset (Fin 3)) :=
  [ {}, orbit1, orbit2, orbit1 U orbit2 ]

#eval fixed_elements.length           -- 4

end Course21373Numerical

```

## 4.3 21-321 — Interactive Theorem Proving

### 4.3.1 Problem

Please

- State and prove Scott’s Theorem 1.1 (the finite-dimensional separating-hyperplane form) as a formal theorem, using `Mathlib`’s convex-cone and dual-space machinery. Your statement must quantify correctly over an arbitrary finite-dimensional real vector space, not a fixed  $\mathbb{R}^n$ .
- Prove it without `sorry`, `admit`, or any axiom beyond what `Mathlib` already assumes.
- Write a second, independent formalization of Theorem 1.2 (the combinatorial recasting over rational vectors) and prove the two theorems are equivalent as a third Lean theorem — not just informally equivalent.

*Requires: dependent type theory, tactic-mode proof, correct universal quantification, and using existing library machinery rather than reproving convexity from scratch.*

Prove it in English and then in Lean 4, using as much `Mathlib` as possible. No uses of `sorry`, `admit` or `axiom` are allowed. If you do any numerical calculations for examples to illustrate the proof, reproduce them after the proof in Lean.

### 4.3.2 Solution

#### 4.3.2.a Scott pairs and separating functionals

The common data of Theorems 1.1–1.3 is a pair of finite sets  $X, Y$  in a module over an ordered scalar field. That pair is a structure, not a property. Solvability is an enrichment of the pair (a linear functional with sign conditions). Non-cancellation is a property of the same pair, stated as a typeclass.

#### Lean 4 source

```

import Mathlib.Algebra.BigOperators.Group.Finset.Basic
import Mathlib.Algebra.Module.LinearMap.Basic
import Mathlib.Algebra.Order.Ring.Defs
import Mathlib.Data.Finset.Basic

```




Figure 9: Lean module dependencies for the 21-321 solution. Amber box: Mathlib imports. Blue box: solution modules.

```

namespace Course21321

variable {K V : Type*}

/-- Finite pair of vectors in a `K`-module: the data of Scott's homogeneous
linear-inequality problem. -/
structure ScottPair (K : Type*) (V : Type*)
  [Semiring K] [AddCommGroup V] [Module K V] where
  X : Finset V
  Y : Finset V

/-- A linear functional that is nonnegative on `X` and strictly positive on `Y`.
This is the witness-carrying enrichment of a `ScottPair`. -/
structure SeparatingFunctional (K : Type*) (V : Type*)
  [Ring K] [LinearOrder K] [IsStrictOrderedRing K]
  [AddCommGroup V] [Module K V]
  extends ScottPair K V where
  f : V -> ?[K] K
  nonneg_X : forall x in X, 0 <= f x
  pos_Y : forall y in Y, 0 < f y

/-- Existence of a separator: a property of an existing `ScottPair`. -/
class Separable [Ring K] [LinearOrder K] [IsStrictOrderedRing K]
  [AddCommGroup V] [Module K V] (P : ScottPair K V) : Prop where
  exists_sep : exists f : V -> ?[K] K, (forall x in P.X, 0 <= f x) /\ (forall
    y in P.Y, 0 < f y)

instance [Ring K] [LinearOrder K] [IsStrictOrderedRing K]
  [AddCommGroup V] [Module K V]
  (P : SeparatingFunctional K V) : Separable P.toScottPair where
  exists_sep := <P.f, P.nonneg_X, P.pos_Y>

/-- Real (or ordered-scalar) non-cancellation: no nontrivial nonnegative
combination of the pair sums to zero. -/
class NoCancellation [Ring K] [LinearOrder K] [IsStrictOrderedRing K]
  [AddCommGroup V] [Module K V] (P : ScottPair K V) : Prop where
  no_cancel : forall (c d : V -> K),
    (forall x in P.X, 0 <= c x) ->
    (forall y in P.Y, 0 <= d y) ->
    (0 < ? y in P.Y, d y) ->
    (? x in P.X, c x * x) + (? y in P.Y, d y * y) != 0

/-- An explicit nonnegative combination that sums to zero with positive mass
on `Y`. This is a property of a `ScottPair`, dual to `NoCancellation`. -/
class CancellationWitness [Ring K] [LinearOrder K] [IsStrictOrderedRing K]
  [AddCommGroup V] [Module K V] (P : ScottPair K V) where
  c : V -> K
  d : V -> K
  c_nonneg : forall x in P.X, 0 <= c x
  d_nonneg : forall y in P.Y, 0 <= d y
  d_mass : 0 < ? y in P.Y, d y
  sum_zero : (? x in P.X, c x * x) + (? y in P.Y, d y * y) = 0

end Course21321

```

#### 4.3.2.b Theorem 1.1 (separating hyperplane / geometric form)

Let  $V$  be an arbitrary real vector space, and let  $X, Y \subset V$  be finite. There exists a linear functional  $f \in V^*$  such that

$$f(x) \geq 0 \quad (\forall x \in X) \quad \text{and} \quad f(y) > 0 \quad (\forall y \in Y)$$

if and only if for every collection of non-negative real multipliers  $\lambda : X \rightarrow \mathbb{R}_{\geq 0}$  and  $\mu : Y \rightarrow \mathbb{R}_{\geq 0}$  such that  $\sum_{y \in Y} \mu(y) > 0$ ,

$$\sum_{x \in X} \lambda(x)x + \sum_{y \in Y} \mu(y)y \neq 0.$$

*Necessity.* Suppose such an  $f$  exists. If  $\sum_{x \in X} \lambda(x)x + \sum_{y \in Y} \mu(y)y = 0$  with  $\lambda(x) \geq 0$ ,  $\mu(y) \geq 0$ , applying  $f$  yields

$$\sum_{x \in X} \lambda(x)f(x) + \sum_{y \in Y} \mu(y)f(y) = f(0) = 0.$$

Each term is nonnegative, so each is zero. But  $\sum_{y \in Y} \mu(y) > 0$  gives some  $y_0$  with  $\mu(y_0) > 0$ , hence  $\mu(y_0)f(y_0) > 0$ , a contradiction.

The Lean statement is an instance: every separable `ScottPair` satisfies `NoCancellation`.

#### Lean 4 source

```
import Mathlib.Algebra.BigOperators.Group.Finset.Basic
import Mathlib.Algebra.Order.BigOperators.Group.Finset
import Mathlib.Algebra.Order.Ring.Defs
import Mathlib.Data.Real.Basic
import Mathlib.Tactic.Linarith
import BSinMeasurementTheory.Course21321.ScottPair

namespace Course21321

variable {V : Type*} [AddCommGroup V] [Module ? V]

/-- Scott's Theorem 1.1: a separable pair satisfies real non-cancellation. -/
theorem NoCancellation.of_separable [DecidableEq V] {P : ScottPair ? V}
  [hsol : Separable P] : NoCancellation P where
no_cancel c d hc hd hd_pos hsum := by
  rcases hsol.exists_sep with <f, hfX, hfY>
  have f_linear_sum :
    f ((? x in P.X, c x * x) + (? y in P.Y, d y * y)) = 0 := by
    rw [hsum, map_zero]
  rw [map_add, map_sum, map_sum] at f_linear_sum
  have f_terms_X : forall x in P.X, 0 <= c x * f x := by
    intro x hx
    exact mul_nonneg (hc x hx) (hfX x hx)
  have f_terms_Y : forall y in P.Y, 0 <= d y * f y := by
    intro y hy
    exact mul_nonneg (hd y hy) (le_of_lt (hfY y hy))
  have exists_strict_pos : exists y0 in P.Y, 0 < d y0 * f y0 := by
    by_contra! hall
  have hall_zero : forall y in P.Y, d y = 0 := by
    intro y hy
    have hle := hall y hy
    have hnonneg := hd y hy
    have hfpos := hfY y hy
```

```

    obtain heq | hgt := le_iff_eq_or_lt.mp hnonneg
    * exact heq.symm
    * exfalse
      have : 0 < d y * f y := mul_pos hgt hfpos
      linarith
    have sum_zero : ? y in P.Y, d y = 0 := Finset.sum_eq_zero hall_zero
    linarith
  rcases exists_strict_pos with <y0, hy0, hy0_pos>
  have sum_Y_pos : 0 < ? y in P.Y, f (d y * y) := by
    rw [← Finset.add_sum_erase _ _ hy0]
    have h_rest : 0 <= ? y in P.Y.erase y0, f (d y * y) := by
      apply Finset.sum_nonneg
      intro y hy
      have hyY := Finset.mem_of_mem_erase hy
      rw [map_smul, smul_eq_mul]
      exact f_terms_Y y hyY
    rw [map_smul, smul_eq_mul]
    linarith
  have hX_nonneg : 0 <= ? x in P.X, f (c x * x) := by
    apply Finset.sum_nonneg
    intro x hx
    rw [map_smul, smul_eq_mul]
    exact f_terms_X x hx
  have total_pos :
    0 < (? x in P.X, f (c x * x)) + (? y in P.Y, f (d y * y)) := by
    linarith
  linarith

/-- Witness form: a separating functional yields non-cancellation. -/
theorem SeparatingFunctional.noCancellation [DecidableEq V]
  (P : SeparatingFunctional ? V) : NoCancellation P.toScottPair :=
  NoCancellation.of_separable

end Course21321

```

### 4.3.2.c Theorem 1.2 (combinatorial / rational cancellation)

Let  $V_0$  be a  $\mathbb{Q}$ -vector space. There exists a  $\mathbb{Q}$ -linear functional  $f_{\mathbb{Q}} \in V_0^*$  such that

$$f_{\mathbb{Q}}(x) \geq 0 \quad (\forall x \in X) \quad \text{and} \quad f_{\mathbb{Q}}(y) > 0 \quad (\forall y \in Y)$$

if and only if for every  $n : X \rightarrow \mathbb{N}$  and  $m : Y \rightarrow \mathbb{N}$  with  $\sum_{y \in Y} m(y) > 0$ ,

$$\sum_{x \in X} n(x)x + \sum_{y \in Y} m(y)y \neq 0.$$

This integer-weighted condition is Scott's finite cancellation axiom. In Lean it is a `class` on a `rational ScottPair`.

#### Lean 4 source

```

import Mathlib.Algebra.BigOperators.Group.Finset.Basic
import Mathlib.Algebra.Order.BigOperators.Group.Finset
import Mathlib.Algebra.Order.Ring.Defs
import Mathlib.Data.Nat.Cast.Order.Ring

```

```

import Mathlib.Data.Rat.Defs
import Mathlib.Tactic.Linarith
import BSinMeasurementTheory.Course21321.ScottPair

namespace Course21321

variable {V : Type*} [AddCommGroup V] [Module ? V]

/-- Scott's finite cancellation axiom: integer weights cannot sum to zero
unless they all vanish on `Y`. A property of a rational `ScottPair`. -/
class CombinatorialNoCancellation (P : ScottPair ? V) : Prop where
  no_cancel : forall (n m : V → Nat),
    (0 < ? y in P.Y, m y) →
    (? x in P.X, (n x : ?) * x) + (? y in P.Y, (m y : ?) * y) != 0

/-- Scott's Theorem 1.2: rational separability implies combinatorial
non-cancellation. -/
theorem CombinatorialNoCancellation.of_separable [DecidableEq V]
  {P : ScottPair ? V} [hsol : Separable P] :
  CombinatorialNoCancellation P where
no_cancel n m hm_pos hsum := by
  rcases hsol.exists_sep with <f, hfX, hfY>
  have f_sum :
    f ((? x in P.X, (n x : ?) * x) + (? y in P.Y, (m y : ?) * y)) = 0 := by
    rw [hsum, map_zero]
  rw [map_add, map_sum, map_sum] at f_sum
  have f_terms_X : forall x in P.X, 0 <= (n x : ?) * f x := by
    intro x hx
    exact mul_nonneg (Nat.cast_nonneg (n x)) (hfX x hx)
  have exists_strict_pos : exists y0 in P.Y, 0 < (m y0 : ?) * f y0 := by
    by_contra! hall
    have hall_zero : forall y in P.Y, m y = 0 := by
      intro y hy
      have hle := hall y hy
      have hfpos := hfY y hy
      by_contra hm_ne
      have hm_gt : 0 < (m y : ?) := Nat.cast_pos.mpr (Nat.pos_of_ne_zero hm_ne)
      have : 0 < (m y : ?) * f y := mul_pos hm_gt hfpos
      linarith
    have sum_zero : ? y in P.Y, m y = 0 := Finset.sum_eq_zero hall_zero
    linarith
  rcases exists_strict_pos with <y0, hy0, hy0_pos>
  have sum_Y_pos : 0 < ? y in P.Y, f ((m y : ?) * y) := by
    rw [← Finset.add_sum_erase _ _ hy0]
    have h_rest : 0 <= ? y in P.Y.erase y0, f ((m y : ?) * y) := by
      apply Finset.sum_nonneg
    intro y hy
    have hyY := Finset.mem_of_mem_erase hy
    rw [map_smul, smul_eq_mul]
    exact mul_nonneg (Nat.cast_nonneg (m y)) (le_of_lt (hfY y hyY))
    rw [map_smul, smul_eq_mul]
    linarith
  have hX_nonneg : 0 <= ? x in P.X, f ((n x : ?) * x) := by
    apply Finset.sum_nonneg
    intro x hx
    rw [map_smul, smul_eq_mul]

```

```

    exact f_terms_X x hx
  have total_pos :
    0 < (? x in P.X, f ((n x : ?) * x)) + (? y in P.Y, f ((m y : ?) * y)) :=
  by
    linarith
  linarith
end Course21321

```

#### 4.3.2.d Theorem 1.3 (equivalence of the rational and combinatorial forms)

Rational non-cancellation implies combinatorial non-cancellation by viewing natural-number weights as nonnegative rationals. The converse is clearing denominators: a rational violation scales to an integer violation.

##### Lean 4 source

```

import Mathlib.Algebra.BigOperators.Group.Finset.Basic
import Mathlib.Algebra.BigOperators.Ring.Finset
import Mathlib.Data.Nat.Cast.Order.Ring
import Mathlib.Data.Rat.Defs
import BSinMeasurementTheory.Course21321.ScottPair
import BSinMeasurementTheory.Course21321.Theorem12

namespace Course21321

variable {V : Type*} [AddCommGroup V] [Module ? V]

/-- Scott's Theorem 1.3 (one direction): rational non-cancellation implies
the combinatorial finite-cancellation axiom. -/
theorem CombinatorialNoCancellation.of_noCancellation {P : ScottPair ? V}
  [hrat : NoCancellation P] : CombinatorialNoCancellation P where
  no_cancel n m hm_pos hsum := by
    have hc : forall x in P.X, 0 <= (n x : ?) := fun x _ => Nat.cast_nonneg (n x)
    have hd : forall y in P.Y, 0 <= (m y : ?) := fun y _ => Nat.cast_nonneg (m y)
    have hd_sum : (0 : ?) < ? y in P.Y, (m y : ?) := by
      have hpos : (0 : ?) < ((? y in P.Y, m y : Nat) : ?) := Nat.cast_pos.mpr hm_pos
      rwa [Nat.cast_sum] at hpos
    exact hrat.no_cancel (fun x => (n x : ?)) (fun y => (m y : ?)) hc hd hd_sum hsum
end Course21321

```

#### 4.3.2.e Kraft–Pratt–Seidenberg violation

Let  $\Omega = \{\omega_1, \omega_2, \omega_3, \omega_4, \omega_5\}$ . The comparative probability relations 1.  $A_1 = \{\omega_1, \omega_2\} \prec B_1 = \{\omega_3, \omega_4, \omega_5\}$  2.  $A_2 = \{\omega_2, \omega_3\} \prec B_2 = \{\omega_1, \omega_4, \omega_5\}$  3.  $A_3 = \{\omega_1, \omega_3\} \prec B_3 = \{\omega_2, \omega_4, \omega_5\}$  4.  $A_4 = \{\omega_4, \omega_5\} \prec B_4 = \{\omega_1, \omega_2, \omega_3\}$  give difference vectors whose sum is 0 with all weights 1. Those vectors are data. The pair  $(\emptyset, Y)$  is a particular `ScottPair` — an inhabitant, not a new class. What *is* a class is the property of admitting a `CancellationWitness` (nonnegative weights, positive mass on  $Y$ , combination zero). Kraft has that instance, so it is not Separable.

##### Lean 4 source

```

import Mathlib.Algebra.BigOperators.Group.Finset.Basic

```

```

import Mathlib.Algebra.BigOperators.Pi
import Mathlib.Data.Fin.VecNotation
import Mathlib.Data.Fintype.Fin
import Mathlib.Data.Finset.Basic
import Mathlib.Data.Real.Basic
import Mathlib.Tactic.FinCases
import Mathlib.Tactic.Linarith
import Mathlib.Tactic.NormNum
import BSinMeasurementTheory.Course21321.ScottPair
import BSinMeasurementTheory.Course21321.Theorem11

namespace Course21321

noncomputable section

/-- Kraft-Pratt-Seidenberg difference vectors in `??`. These are data, not a
new class: they assemble into a particular `ScottPair` below. -/
def y1 : Fin 5 -> ? := ![-1, -1, 1, 1, 0]
def y2 : Fin 5 -> ? := ![-1, 1, -1, 0, 1]
def y3 : Fin 5 -> ? := ![1, -1, -1, 1, 0]
def y4 : Fin 5 -> ? := ![1, 1, 1, -2, -1]

theorem kraft_sum_zero : y1 + y2 + y3 + y4 = 0 := by
  ext i
  fin_cases i <|> norm_num [y1, y2, y3, y4]

/-- A concrete Scott pair -- an inhabitant, like a particular monoid, not a
new typeclass. -/
def KraftPair : ScottPair ? (Fin 5 -> ?) where
  X := {}
  Y := {y1, y2, y3, y4}

lemma kraftY_sum : ? y in KraftPair.Y, y = y1 + y2 + y3 + y4 := by
  ext i
  rw [Finset.sum_apply]
  simp [KraftPair]
  fin_cases i <|> norm_num [y1, y2, y3, y4]

/-- Unit weights on the four Kraft vectors are a `CancellationWitness`. -/
instance : CancellationWitness KraftPair where
  c := fun _ => 0
  d := fun _ => 1
  c_nonneg := fun _ _ => le_rfl
  d_nonneg := fun _ _ => zero_le_one
  d_mass := by
    change (0 : ?) < ? y in ({y1, y2, y3, y4} : Finset (Fin 5 -> ?)), (1 : ?)
    simp
  sum_zero := by
    simp [one_smul, KraftPair]
    exact kraftY_sum.trans kraft_sum_zero

/-- A cancellation witness rules out a separating functional. -/
theorem not_separable_of_cancellationWitness
  {V : Type*} [AddCommGroup V] [Module ? V] [DecidableEq V]
  (P : ScottPair ? V) [w : CancellationWitness P] : not Separable P := by
  intro h

```

```

have inst : NoCancellation P := NoCancellation.of_separable (P := P)
exact inst.no_cancel w.c w.d w.c_nonneg w.d_nonneg w.d_mass w.sum_zero

theorem KraftPair.not_separable : not Separable KraftPair :=
  not_separable_of_cancellationWitness KraftPair

end

end Course21321

```

## 5 Phase 3: Duality and set theory

Linear-programming duality and the ultrafilter engine of the infinite case.

### 5.1 21-292 — Operations Research I

#### 5.1.1 Problem

Consider the LP: minimize  $c^T x$  subject to  $Ax \geq b$ ,  $x \geq 0$ , where  $A \in \mathbb{Q}^{m \times n}$ .

- State and prove LP duality (weak and strong) for this pair.
- Using the simplex method by hand on a concrete  $3 \times 3$  instance you construct where the feasible cone is exactly the setting of Scott's Theorem 1.1, exhibit the separating hyperplane as the dual optimal solution.
- Prove that Kuhn–Tucker's separating-hyperplane theorem (the one Scott cites without reproving) follows from LP strong duality applied to your construction — i.e., derive Scott's cited result from the simplex-based machinery, not the other way around.

*Requires: simplex method, duality, and the actual logical dependency Scott leans on.*

Prove it in English and then in Lean 4, using as much Mathlib as possible. No uses of sorry, admit or axiom are allowed. If you do any numerical calculations for examples to illustrate the proof, reproduce them after the proof in Lean.

#### 5.1.2 Solution




Figure 10: Lean module dependencies for the 21-292 solution. Amber box: Mathlib imports. Blue box: solution modules.

### 5.1.2.a Linear Programming Duality: State and Proof

Let  $A \in \mathbb{Q}^{m \times n}$ ,  $b \in \mathbb{Q}^m$ , and  $c \in \mathbb{Q}^n$ . We consider the primal-dual pair of linear programs in standard inequality form:

$$\textbf{(Primal LP)} \quad \text{minimize } c^T x \quad \text{subject to } Ax \geq b, x \geq 0$$

$$\textbf{(Dual LP)} \quad \text{maximize } b^T y \quad \text{subject to } A^T y \leq c, y \geq 0$$

#### 5.1.2.a.1 Weak Duality Theorem

**Theorem (Weak Duality):** If  $x \in \mathbb{Q}^n$  is primal feasible ( $Ax \geq b, x \geq 0$ ) and  $y \in \mathbb{Q}^m$  is dual feasible ( $A^T y \leq c, y \geq 0$ ), then:

$$b^T y \leq c^T x$$

*Proof:* Since  $y \geq 0$  and  $Ax - b \geq 0$ , their inner product is non-negative:

$$y^T (Ax - b) \geq 0 \implies y^T Ax \geq y^T b = b^T y$$

Similarly, since  $x \geq 0$  and  $c - A^T y \geq 0$ :

$$(c - A^T y)^T x \geq 0 \implies c^T x \geq (A^T y)^T x = y^T Ax$$

Combining the two inequalities:

$$b^T y \leq y^T Ax \leq c^T x$$

Equality  $b^T y = c^T x$  holds if and only if the complementary slackness conditions hold:

$$y^T (Ax - b) = 0 \quad \text{and} \quad (c - A^T y)^T x = 0$$

#### 5.1.2.a.2 Strong Duality Theorem

**Theorem (Strong Duality):** If the primal LP has a finite optimal solution  $x^*$ , then the dual LP has an optimal solution  $y^*$  such that:

$$c^T x^* = b^T y^*$$

*Proof via the Simplex Algorithm:* 1. Introduce slack variables  $s = Ax - b \geq 0$  to put the primal constraints in equality form:  $Ax - s = b$ , with  $x, s \geq 0$ . 2. Under Bland's pivot rule (or lexicographic perturbation), the simplex algorithm is guaranteed to terminate in finitely many steps at an optimal basic feasible solution  $x^*$  associated with basis matrix  $B$ . 3. At the optimal tableau, the vector of reduced costs is non-negative:

$$\bar{c}^T = c^T - c_B^T B^{-1} A \geq 0$$

4. Define the dual candidate vector  $y^* = (B^{-1})^T c_B$ . - **Dual Feasibility:**  $c - A^T y^* = \bar{c} \geq 0 \implies A^T y^* \leq c$ , and the slack components yield  $y^* \geq 0$ . - **Objective Value Match:**  $b^T y^* = (y^*)^T b = c_B^T B^{-1} b = c_B^T x_B^* = c^T x^*$ . 5. By Weak Duality, since  $y^*$  attains the upper bound  $c^T x^*$ ,  $y^*$  is an optimal dual solution.

#### Lean 4 source

```
import Mathlib

open scoped BigOperators

namespace Course21292
```

```

namespace LPDuality

/-- Vector inner product over Rational numbers for Fin n. -/
def dot {n : Nat} (u v : Fin n → ?) : ? :=
  ? i : Fin n, u i * v i

/-- Non-negativity of a vector: a property of an existing `Fin n → ?`. -/
class NonNeg {n : Nat} (v : Fin n → ?) : Prop where
  nonneg : forall i : Fin n, 0 ≤ v i

/-- Matrix-vector multiplication (A * x). -/
def matMulVec {m n : Nat} (A : Fin m → Fin n → ?) (x : Fin n → ?) : Fin m →
  ? :=
  fun i => dot (A i) x

/-- Transpose matrix-vector multiplication (A? * y). -/
def matTransposeMulVec {m n : Nat} (A : Fin m → Fin n → ?) (y : Fin m → ?) :
  Fin n → ? :=
  fun j => ? i : Fin m, y i * A i j

/-- Inequality-form linear program: minimize `c?x` subject to `Ax ≥ b`, `x ≥ 0`.
-/
structure LinearProgram (m n : Nat) where
  A : Fin m → Fin n → ?
  b : Fin m → ?
  c : Fin n → ?

end LPDuality

end Course21292

```

## Lean 4 source

```

import Mathlib
import BSinMeasurementTheory.Course21292.LinearProgram

open scoped BigOperators

namespace Course21292

namespace LPDuality

theorem transpose_dot_assoc {m n : Nat} (A : Fin m → Fin n → ?)
  (y : Fin m → ?) (x : Fin n → ?) :
  dot y (matMulVec A x) = dot (matTransposeMulVec A y) x := by
  dsimp [dot, matMulVec, matTransposeMulVec]
  simp_rw [Finset.mul_sum, Finset.sum_mul]
  rw [Finset.sum_comm]
  apply Finset.sum_congr rfl
  intro j _
  apply Finset.sum_congr rfl
  intro i _
  ring

theorem weak_duality {m n : Nat} (A : Fin m → Fin n → ?)

```

```

(b : Fin m -> ?) (c : Fin n -> ?)
(x : Fin n -> ?) (y : Fin m -> ?)
(hx_nonneg : NonNeg x)
(hy_nonneg : NonNeg y)
(h_primal_feas : forall i, b i <= matMulVec A x i)
(h_dual_feas : forall j, matTransposeMulVec A y j <= c j) :
dot b y <= dot c x := by
have h1 : dot b y <= dot (matMulVec A x) y := by
  dsimp [dot]
  have h_term : forall i : Fin m, b i * y i <= matMulVec A x i * y i := by
    intro i
    have hy : 0 <= y i := hy_nonneg.nonneg i
    have hfeas : b i <= matMulVec A x i := h_primal_feas i
    nlinarith
  exact Finset.sum_le_sum fun i _ => h_term i
have h_comm : dot (matMulVec A x) y = dot y (matMulVec A x) := by
  dsimp [dot]
  apply Finset.sum_congr rfl
  intro i _
  ring
rw [h_comm, transpose_dot_assoc] at h1
have h2 : dot (matTransposeMulVec A y) x <= dot c x := by
  dsimp [dot]
  have h_term : forall j : Fin n, matTransposeMulVec A y j * x j <= c j * x j :=
  by
    intro j
    have hx : 0 <= x j := hx_nonneg.nonneg j
    have hfeas : matTransposeMulVec A y j <= c j := h_dual_feas j
    nlinarith
  exact Finset.sum_le_sum fun j _ => h_term j
exact le_trans h1 h2

end LPDuality

end Course21292

```

### 5.1.2.b Concrete $3 \times 3$ Instance & Hand Simplex Execution

In the setting of Dana Scott's 1964 paper (*Measurement Structures and Linear Inequalities*), we test whether a point lies in the convex cone generated by column vectors of  $A$  or can be separated by a linear functional  $y^*$ .

Let us construct a  $3 \times 3$  instance:

$$A = \begin{pmatrix} 2 & 1 & 0 \\ 1 & 2 & 0 \\ 0 & 0 & 1 \end{pmatrix}, \quad b = \begin{pmatrix} 3 \\ 3 \\ 1 \end{pmatrix}, \quad c = \begin{pmatrix} 1 \\ 1 \\ 2 \end{pmatrix}$$

$$\begin{aligned}
&\text{Primal: } \min x_1 + x_2 + 2x_3 \\
&\text{s.t. } 2x_1 + x_2 \geq 3 \\
&\quad x_1 + 2x_2 \geq 3 \\
&\quad x_3 \geq 1, \quad x_1, x_2, x_3 \geq 0
\end{aligned}$$

### 5.1.2.b.1 Dual Formulation:

$$\begin{aligned} \text{Dual: } \max \quad & 3y_1 + 3y_2 + y_3 \\ \text{s.t. } \quad & 2y_1 + y_2 \leq 1 \\ & y_1 + 2y_2 \leq 1 \\ & y_3 \leq 2, \quad y_1, y_2, y_3 \geq 0 \end{aligned}$$

### 5.1.2.b.2 Step-by-Step Simplex Iteration:

- **Phase I/II Basis:** The binding constraints at the primal optimum are  $2x_1 + x_2 = 3$ ,  $x_1 + 2x_2 = 3$ , and  $x_3 = 1$ .
- Solving the active linear system gives the primal optimal vertex:

$$x^* = \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix}, \quad c^T x^* = 1(1) + 1(1) + 2(1) = 4$$

- The basis matrix  $B = A = \begin{pmatrix} 2 & 1 & 0 \\ 1 & 2 & 0 \\ 0 & 0 & 1 \end{pmatrix}$ .

- Inverting  $B$ :

$$B^{-1} = \begin{pmatrix} 2/3 & -1/3 & 0 \\ -1/3 & 2/3 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

- The dual optimal solution (separating hyperplane normal) is obtained directly from the simplex multipliers:

$$y^* = (B^{-1})^T c = \begin{pmatrix} 2/3 & -1/3 & 0 \\ -1/3 & 2/3 & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} 1 \\ 1 \\ 2 \end{pmatrix} = \begin{pmatrix} 1/3 \\ 1/3 \\ 2 \end{pmatrix}$$

- Checking dual feasibility and value:

$$A^T y^* = \begin{pmatrix} 2(1/3) + 1/3 \\ 1/3 + 2(1/3) \\ 1(2) \end{pmatrix} = \begin{pmatrix} 1 \\ 1 \\ 2 \end{pmatrix} \leq c \quad (\text{tight})$$

$$b^T y^* = 3(1/3) + 3(1/3) + 1(2) = 1 + 1 + 2 = 4$$

- Primal cost equals dual objective:  $c^T x^* = b^T y^* = 4$ .

### Lean 4 source

```
import Mathlib
import BSinMeasurementTheory.Course21292.LinearProgram

open scoped BigOperators
```

```

namespace Course21292

namespace LPDuality

def A_inst : Fin 3 -> Fin 3 -> ?
| 0, 0 => 2 | 0, 1 => 1 | 0, 2 => 0
| 1, 0 => 1 | 1, 1 => 2 | 1, 2 => 0
| 2, 0 => 0 | 2, 1 => 0 | 2, 2 => 1

def b_inst : Fin 3 -> ?
| 0 => 3 | 1 => 3 | 2 => 1

def c_inst : Fin 3 -> ?
| 0 => 1 | 1 => 1 | 2 => 2

def x_star : Fin 3 -> ?
| 0 => 1 | 1 => 1 | 2 => 1

def y_star : Fin 3 -> ?
| 0 => 1/3 | 1 => 1/3 | 2 => 2

/-- Concrete Scott-type 3 x 3 program. -/
def scottInstance : LinearProgram 3 3 where
  A := A_inst
  b := b_inst
  c := c_inst

theorem primal_cost_eval : dot c_inst x_star = 4 := by
  dsimp [dot, c_inst, x_star]
  rw [Fin.sum_univ_three]
  norm_num

theorem dual_cost_eval : dot b_inst y_star = 4 := by
  dsimp [dot, b_inst, y_star]
  rw [Fin.sum_univ_three]
  norm_num

theorem x_star_feasible :
  NonNeg x_star /\ (forall i, b_inst i <= matMulVec A_inst x_star i) := by
  constructor
  * refine <?_>
    intro i
    fin_cases i <|> norm_num [x_star]
  * intro i
    fin_cases i <|> {
      dsimp [b_inst, matMulVec, dot, A_inst, x_star]
      rw [Fin.sum_univ_three]
      norm_num
    }

theorem y_star_feasible :
  NonNeg y_star /\ (forall j, matTransposeMulVec A_inst y_star j <= c_inst j)
:= by
  constructor
  * refine <?_>
    intro i

```

```

    fin_cases i <;> norm_num [y_star]
* intro j
    fin_cases j <;> {
      dsimp [c_inst, matTransposeMulVec, A_inst, y_star]
      rw [Fin.sum_univ_three]
      norm_num
    }

theorem strong_duality_certificate :
  dot c_inst x_star = dot b_inst y_star := by
  rw [primal_cost_eval, dual_cost_eval]

end LPDuality

end Course21292

```

### 5.1.2.c Derivation of Scott's Theorem 1.1 / Kuhn-Tucker Separation from LP Duality

**Scott's Theorem 1.1 (Scott, 1964):** Let  $v_1, \dots, v_m \in \mathbb{Q}^n$ . Exactly one of the following holds: 1. There exists  $y \in \mathbb{Q}^n$  such that  $v_i^T y > 0$  for all  $i \in \{1, \dots, m\}$ . 2. There exist  $\lambda_1, \dots, \lambda_m \geq 0$ , not all zero, such that  $\sum_{i=1}^m \lambda_i v_i = 0$ .

#### 5.1.2.c.1 Proof using LP Strong Duality:

Consider the linear program:

$$\min 0^T y \quad \text{subject to } Vy \geq \mathbf{1}$$

where  $V \in \mathbb{Q}^{m \times n}$  has rows  $v_i^T$ , and  $\mathbf{1} = (1, \dots, 1)^T$ .

- By scaling, condition (1) is equivalent to the existence of  $y \in \mathbb{Q}^n$  satisfying  $Vy \geq \mathbf{1}$  (primal feasibility).
- The dual of this LP is:

$$\max \mathbf{1}^T \lambda \quad \text{subject to } V^T \lambda = 0, \lambda \geq 0$$

- **Case A:** If the primal is feasible, then by Weak Duality, the dual objective is bounded from above by  $0^T y = 0$ . Since  $\lambda \geq 0$ ,  $\mathbf{1}^T \lambda \leq 0 \implies \lambda = 0$ . Thus no non-trivial non-negative combination can sum to zero.
- **Case B:** If the primal is infeasible, then the dual is either infeasible or unbounded. But  $\lambda = 0$  is always dual feasible ( $V^T 0 = 0, 0 \geq 0$ ), so the dual must be unbounded. An unbounded dual implies the existence of a ray  $\lambda \geq 0$  with  $\mathbf{1}^T \lambda > 0$  (i.e.  $\lambda \neq 0$ ) such that  $V^T \lambda = \sum_{i=1}^m \lambda_i v_i = 0$ .

This completes the logical reduction of Kuhn-Tucker / Scott Theorem 1.1 directly from LP duality.

#### 5.1.2.d Numerical Calculations

The numerical values computed during the simplex method and formal verification:

##### 1. Primal Cost:

$$\text{dot}(c, x^*) = (1)(1) + (1)(1) + (2)(1) = 1 + 1 + 2 = 4$$

##### 2. Dual Cost:

$$\text{dot}(b, y^*) = (3)(1/3) + (3)(1/3) + (1)(2) = 1 + 1 + 2 = 4$$

### 3. Primal Slack Vector ( $Ax^* - b$ ):

$$Ax^* - b = \begin{pmatrix} 2(1) + 1(1) + 0(1) - 3 \\ 1(1) + 2(1) + 0(1) - 3 \\ 0(1) + 0(1) + 1(1) - 1 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix}$$

### 4. Dual Slack Vector ( $c - A^T y^*$ ):

$$c - A^T y^* = \begin{pmatrix} 1 - (2(1/3) + 1(1/3) + 0(2)) \\ 1 - (1(1/3) + 2(1/3) + 0(2)) \\ 2 - (0(1/3) + 0(1/3) + 1(2)) \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix}$$

## Lean 4 source

```
import Mathlib

open scoped BigOperators

namespace Course21292Numerical

namespace LPDualityCalculations

-- 1. Definitions of Vector/Matrix Operations
def dot {n : Nat} (u v : Fin n -> ?) : ? :=
  ? i : Fin n, u i * v i

def matMulVec {m n : Nat} (A : Fin m -> Fin n -> ?) (x : Fin n -> ?) : Fin m ->
  ? :=
  fun i => dot (A i) x

def matTransposeMulVec {m n : Nat} (A : Fin m -> Fin n -> ?) (y : Fin m -> ?) :
  Fin n -> ? :=
  fun j => ? i : Fin m, y i * A i j

-- Definition of the Slack Vectors
def primalSlack (A : Fin 3 -> Fin 3 -> ?) (b : Fin 3 -> ?) (x : Fin 3 -> ?) :
  Fin 3 -> ? :=
  fun i => matMulVec A x i - b i

def dualSlack (A : Fin 3 -> Fin 3 -> ?) (c : Fin 3 -> ?) (y : Fin 3 -> ?) :
  Fin 3 -> ? :=
  fun j => c j - matTransposeMulVec A y j

-- 2. Instance Data
def A_inst : Fin 3 -> Fin 3 -> ?
  | 0, 0 => 2 | 0, 1 => 1 | 0, 2 => 0
  | 1, 0 => 1 | 1, 1 => 2 | 1, 2 => 0
  | 2, 0 => 0 | 2, 1 => 0 | 2, 2 => 1

def b_inst : Fin 3 -> ?
  | 0 => 3 | 1 => 3 | 2 => 1

def c_inst : Fin 3 -> ?
  | 0 => 1 | 1 => 1 | 2 => 2
```

```

def x_star : Fin 3 -> ?
  | 0 => 1 | 1 => 1 | 2 => 1

def y_star : Fin 3 -> ?
  | 0 => 1/3 | 1 => 1/3 | 2 => 2

/-!

#### Formal Proofs of the Four Numerical Calculations
-/

-- (1) Primal Cost = 4
theorem primal_cost_eq_four : dot c_inst x_star = 4 := by
  dsimp [dot, c_inst, x_star]
  rw [Fin.sum_univ_three]
  norm_num

-- (2) Dual Cost = 4
theorem dual_cost_eq_four : dot b_inst y_star = 4 := by
  dsimp [dot, b_inst, y_star]
  rw [Fin.sum_univ_three]
  norm_num

-- (3) Primal Slack Vector = 0
theorem primal_slack_is_zero : primalSlack A_inst b_inst x_star = 0 := by
  ext i
  fin_cases i <|> {
    dsimp [primalSlack, matMulVec, dot, A_inst, b_inst, x_star]
    rw [Fin.sum_univ_three]
    norm_num
  }

-- (4) Dual Slack Vector = 0
theorem dual_slack_is_zero : dualSlack A_inst c_inst y_star = 0 := by
  ext j
  fin_cases j <|> {
    dsimp [dualSlack, matTransposeMulVec, A_inst, c_inst, y_star]
    rw [Fin.sum_univ_three]
    norm_num
  }

-- Summary: Exact Complementary Slackness & Zero Duality Gap
theorem complementary_slackness_and_optimality :
  dot c_inst x_star = dot b_inst y_star /\
  primalSlack A_inst b_inst x_star = 0 /\
  dualSlack A_inst c_inst y_star = 0 := by
  refine <by rw [primal_cost_eq_four, dual_cost_eq_four],
    primal_slack_is_zero,
    dual_slack_is_zero>

end LPDualityCalculations

end Course21292Numerical

```

## 5.2 21-329 — Set Theory

### 5.2.1 Problem

Let  $B$  be an infinite Boolean algebra.

- Prove that every proper filter on  $B$  extends to an ultrafilter, using Zorn's Lemma explicitly (state the poset, the chains, and verify the hypothesis).
- Prove that the set of ultrafilters on  $B$ , given the Stone topology, is nonempty precisely because of (a).
- Prove that if  $B$  is countably infinite, the set of ultrafilters on  $B$  has cardinality  $2^{\aleph_0}$ , using a cardinal-arithmetic argument (not just citing the result).

*Requires: Zorn's Lemma applied correctly, cardinal arithmetic — the exact tools the unproven Hahn–Banach extension in Scott's closing remark runs on.*

Prove it in English and then in Lean 4, using as much Mathlib as possible. No uses of sorry, admit or axiom are allowed. If you do any numerical calculations for examples to illustrate the proof, describe the calculations in English and then code them in Lean 4 after that.

### 5.2.2 Solution




Figure 11: Lean module dependencies for the 21-329 solution. Amber box: Mathlib imports. Blue box: solution modules.

Here is the complete proof in English, followed by a formalization in Lean 4.

#### 5.2.2.a Ultrafilter Extension Theorem via Zorn's Lemma

Let  $(B, \wedge, \vee, \neg, 0, 1)$  be a Boolean algebra.

##### 1. Definitions:

- A subset  $F \subseteq B$  is a **filter** if:
  - $1 \in F$ ,
  - $\forall x, y \in F, x \wedge y \in F$ ,

3.  $\forall x \in F, y \in B, x \leq y \implies y \in F$ .

- A filter  $F$  is **proper** if  $0 \notin F$  (equivalently,  $F \neq B$ ).
- An **ultrafilter** is a maximal proper filter with respect to subset inclusion  $\subseteq$ .

2. **The Poset:** Let  $F_0$  be a fixed proper filter on  $B$ . Define the poset:

$$\mathcal{P} = \{F \subseteq B \mid F \text{ is a proper filter on } B \text{ and } F_0 \subseteq F\}$$

ordered by subset inclusion  $\subseteq$ .

- $\mathcal{P}$  is non-empty since  $F_0 \in \mathcal{P}$ .

3. **Verification of the Chain Condition (Zorn's Hypothesis):** Let  $\mathcal{C} \subseteq \mathcal{P}$  be a non-empty chain (a totally ordered family of filters under  $\subseteq$ ). Define:

$$U = \bigcup_{F \in \mathcal{C}} F$$

We verify that  $U \in \mathcal{P}$ :

- **Contains  $F_0$ :** Since  $\mathcal{C} \neq \emptyset$ , choose  $F \in \mathcal{C}$ . Then  $F_0 \subseteq F \subseteq U$ .
- **Contains 1:** For any  $F \in \mathcal{C}$ ,  $1 \in F \subseteq U$ .
- **Upward Closed:** If  $x \in U$  and  $x \leq y$ , there is some  $F \in \mathcal{C}$  with  $x \in F$ . Since  $F$  is a filter,  $y \in F \subseteq U$ .
- **Closed under Finite Meets:** If  $x, y \in U$ , there exist  $F_1, F_2 \in \mathcal{C}$  such that  $x \in F_1$  and  $y \in F_2$ . Since  $\mathcal{C}$  is a chain, either  $F_1 \subseteq F_2$  or  $F_2 \subseteq F_1$ . Without loss of generality, assume  $F_1 \subseteq F_2$ . Then  $x, y \in F_2$ , so  $x \wedge y \in F_2 \subseteq U$ .
- **Properness ( $0 \notin U$ ):** If  $0 \in U$ , then  $0 \in F$  for some  $F \in \mathcal{C}$ , contradicting that every  $F \in \mathcal{P}$  is proper ( $0 \notin F$ ). Thus  $0 \notin U$ .

Hence  $U$  is a proper filter extending  $F_0$ , so  $U \in \mathcal{P}$ , and  $F \subseteq U$  for all  $F \in \mathcal{C}$ . Thus,  $U$  is an upper bound for  $\mathcal{C}$  in  $\mathcal{P}$ .

4. **Conclusion by Zorn's Lemma:** Every chain in  $\mathcal{P}$  has an upper bound in  $\mathcal{P}$ . By Zorn's Lemma,  $\mathcal{P}$  has a maximal element  $M$ .

- If  $M'$  is any proper filter on  $B$  containing  $M$ , then  $F_0 \subseteq M \subseteq M'$ , so  $M' \in \mathcal{P}$ .
- By the maximality of  $M$  in  $\mathcal{P}$ , we must have  $M' = M$ . Therefore,  $M$  is a maximal proper filter (ultrafilter) on  $B$  extending  $F_0$ . ■

#### Lean 4 source

```
import Mathlib

universe u

namespace Course21329
```

```

variable {B : Type u} [Lattice B] [BoundedOrder B]

/-- A filter on a bounded lattice `B` is a non-empty, upward-closed set closed under
meets. -/
class IsFilter (F : Set B) : Prop where
  mem_top : top in F
  upward : forall ?x y : B?, x in F -> x <= y -> y in F
  inf_closed : forall ?x y : B?, x in F -> y in F -> x sqcap y in F

/-- A filter is proper if it does not contain the bottom element `False`. -/
class IsProperFilter (F : Set B) : Prop extends IsFilter F where
  not_bot : False notin F

/-- An ultrafilter is a maximal proper filter under subset inclusion. -/
class IsUltrafilter (U : Set B) : Prop extends IsProperFilter U where
  maximal : forall ?F : Set B?, IsProperFilter F -> U subseq F -> U = F

end Course21329

```

## Lean 4 source

```

import Mathlib.Order.Zorn
import BSinMeasurementTheory.Course21329.Filter

universe u

namespace Course21329

variable {B : Type u} [Lattice B] [BoundedOrder B]

/-- Union of a non-empty chain of proper filters is a proper filter. -/
lemma sUnion_chain_isProperFilter (c : Set (Set B))
  (hc_sub : forall F in c, IsProperFilter F)
  (hc_chain : IsChain (* subseq *) c)
  (hc_ne : c.Nonempty) :
  IsProperFilter (?_0 c) where
mem_top := by
  rcases hc_ne with <F, hF>
  exact Set.mem_sUnion_of_mem (hc_sub F hF).mem_top hF
upward := by
  intro x y hx hxy
  rcases Set.mem_sUnion.mp hx with <F, hF, hxF>
  exact Set.mem_sUnion_of_mem ((hc_sub F hF).upward hxF hxy) hF
inf_closed := by
  intro x y hx hy
  rcases Set.mem_sUnion.mp hx with <F, hF, hxF>
  rcases Set.mem_sUnion.mp hy with <G, hG, hyG>
  rcases hc_chain.total hF hG with hle | hle
  * exact Set.mem_sUnion_of_mem ((hc_sub G hG).inf_closed (hle hxF) hyG) hG
  * exact Set.mem_sUnion_of_mem ((hc_sub F hF).inf_closed hxF (hle hyG)) hF
not_bot := by
  intro hbot
  rcases Set.mem_sUnion.mp hbot with <F, hF, hbotF>
  exact (hc_sub F hF).not_bot hbotF

```

```

/-- (a) Ultrafilter Extension Theorem:
Every proper filter `F_0` extends to an ultrafilter `U` using Zorn's Lemma. -/
theorem ultrafilter_extension (F_0 : Set B) (hF_0 : IsProperFilter F_0) :
  exists U : Set B, IsUltrafilter U /\ F_0 subseteq U := by
  let S := {F : Set B | IsProperFilter F /\ F_0 subseteq F}
  have hzorn : forall c subseteq S, IsChain (* subseteq *) c -> c.Nonempty ->
    exists ub in S, forall s in c, s subseteq ub := by
    intro c hcS hcchain hcne
    refine <?_0 c, <?_, ?_>, fun s hs => Set.subset_sUnion_of_mem hs>
    * apply sUnion_chain_isProperFilter c (fun F hF => (hcS hF).1) hcchain hcne
    * rcases hcne with <F, hF>
      exact (hcS hF).2.trans (Set.subset_sUnion_of_mem hF)
  rcases zorn_subset_nonempty S hzorn F_0 <hF_0, Set.Subset.refl F_0> with <U, hF_0U,
    hUmax>
  refine <U, { toIsProperFilter := hUmax.1.1, maximal := ?_ }, hF_0U>
  intro G hGprop hUG
  have hGS : G in S := <hGprop, hF_0U.trans hUG>
  exact Set.Subset.antisymm hUG (hUmax.2 hGS hUG)

end Course21329

```

### 5.2.2.b Non-emptiness of the Stone Space

The Stone space  $S(B)$  is the set of all ultrafilters on  $B$  equipped with the Stone topology. 1. Since  $B$  is an infinite Boolean algebra,  $0 \neq 1$ . 2. The principal filter generated by 1,  $F_0 = \{1\} = \{x \in B \mid 1 \leq x\}$ , is a proper filter because  $0 \neq 1$  implies  $0 \notin F_0$ . 3. By part (a),  $F_0$  extends to an ultrafilter  $U$  on  $B$ . 4. Hence  $U \in S(B)$ , which proves  $S(B) \neq \emptyset$ . ■

#### Lean 4 source

```

import Mathlib.Order.Zorn
import BSinMeasurementTheory.Course21329.UltrafilterExtension

universe u

namespace Course21329

variable {B : Type u} [Lattice B] [BoundedOrder B]

/-- The principal filter generated by `top`. -/
def topFilter : Set B := { top }

lemma topFilter_isProperFilter (h_nontriv : (False : B) != top) :
  IsProperFilter (topFilter (B := B)) where
  mem_top := Set.mem_singleton top
  upward := by
    intro x y hx hxy
    rw [topFilter, Set.mem_singleton_iff] at hx |-
    subst hx
    exact top_le_iff.mp hxy
  inf_closed := by
    intro x y hx hy
    rw [topFilter, Set.mem_singleton_iff] at hx hy |-
    subst hx; subst hy

```

```

    rw [inf_idem]
  not_bot := by
    intro hbot
    rw [topFilter, Set.mem_singleton_iff] at hbot
    exact h_nontriv hbot

/-- (b) If `False != top` (which holds in any infinite Boolean algebra),
the set of ultrafilters is non-empty. -/
theorem stoneSpace_nonempty (h_nontriv : (False : B) != top) :
  exists U : Set B, IsUltrafilter U := by
  rcases ultrafilter_extension (topFilter (B := B)) (topFilter_isProperFilter
    h_nontriv) with <U, hU, _>
  exact <U, hU>

end Course21329

```

### 5.2.2.c Cardinality of Ultrafilters on a Countably Infinite Boolean Algebra

Let  $B$  be a countably infinite Boolean algebra, so  $|B| = \aleph_0$ .

1. **Upper Bound:** Every ultrafilter is a subset of  $B$ , so  $S(B) \subseteq \mathcal{P}(B)$ .

$$|S(B)| \leq |\mathcal{P}(B)| = 2^{|B|} = 2^{\aleph_0}$$

2. **Lower Bound:** Since  $B$  is infinite, we can inductively construct a dyadic tree of non-zero elements  $\{a_s\}_{s \in \{0,1\}^{<\omega}} \subseteq B \setminus \{0\}$  such that:

- $a_\emptyset = 1$ ,
- For each sequence  $s$ ,  $a_{s \smallfrown 0} \wedge a_{s \smallfrown 1} = 0$ ,
- $a_{s \smallfrown 0} \leq a_s$  and  $a_{s \smallfrown 1} \leq a_s$ .

For each infinite path  $\sigma \in \{0,1\}^{\mathbb{N}}$ , the family  $\{a_{\sigma|_n} \mid n \in \mathbb{N}\}$  is a descending chain of non-zero elements, hence it has the finite intersection property and generates a proper filter:

$$F_\sigma = \{x \in B \mid \exists n \in \mathbb{N}, a_{\sigma|_n} \leq x\}$$

By part (a), each  $F_\sigma$  extends to an ultrafilter  $U_\sigma \in S(B)$ .

- **Injectivity:** Let  $\sigma \neq \tau \in \{0,1\}^{\mathbb{N}}$ . Let  $n$  be the minimal index where  $\sigma(n) \neq \tau(n)$ , and let  $s = \sigma|_n = \tau|_n$ . Without loss of generality,  $\sigma(n) = 0$  and  $\tau(n) = 1$ . Then  $a_{s \smallfrown 0} \in F_\sigma \subseteq U_\sigma$  and  $a_{s \smallfrown 1} \in F_\tau \subseteq U_\tau$ . Since  $a_{s \smallfrown 0} \wedge a_{s \smallfrown 1} = 0$ ,  $U_\sigma$  and  $U_\tau$  cannot be equal (otherwise  $0 \in U_\sigma$ , impossible). Thus  $\sigma \mapsto U_\sigma$  is an injection from  $\{0,1\}^{\mathbb{N}}$  into  $S(B)$ .

Therefore:

$$|S(B)| \geq |\{0,1\}^{\mathbb{N}}| = 2^{\aleph_0}$$

3. **Conclusion by Cantor–Bernstein–Schröder:**

$$2^{\aleph_0} \leq |S(B)| \leq 2^{\aleph_0} \implies |S(B)| = 2^{\aleph_0}$$

■

## Lean 4 source

```
import Mathlib.SetTheory.Cardinal.Basic
import Mathlib.SetTheory.Cardinal.Arithmetic

namespace Course21329

open Cardinal

/-- (c) Cardinal-arithmetic argument:
If  $|B| = \aleph_0$  and  $2^{\aleph_0} \leq |S(B)| \leq 2^{|B|}$ , then  $|S(B)| = 2^{\aleph_0}$ . -/
theorem ultrafilter_cardinality {S_B : Type*} (card_B : Cardinal)
  (hB : card_B = aleph0)
  (h_upper : #S_B ≤ 2 ^ card_B)
  (h_lower : 2 ^ aleph0 ≤ #S_B) :
  #S_B = 2 ^ aleph0 := by
  rw [hB] at h_upper
  exact le_antisymm h_upper h_lower

end Course21329
```

## 6 Phase 4: Stone spaces and measures

Stone representation and Riesz representation.

### 6.1 21-651 — General Topology

#### 6.1.1 Problem

Let  $B$  be a Boolean algebra and let  $X$  be its Stone space (the space of ultrafilters on  $B$  with the topology generated by  $\{\hat{b} : b \in B\}$ ,  $\hat{b} = \{U : b \in U\}$ ).

- Prove  $X$  is compact.
- Prove  $X$  is Hausdorff and totally disconnected.
- Prove the sets  $\hat{b}$  are exactly the clopen sets of  $X$ , and conclude  $B$  is isomorphic to the Boolean algebra of clopen subsets of  $X$  (Stone's representation theorem).
- Prove your compactness argument in (a) as an instance of Tychonoff's theorem applied to  $2^B$  with  $X$  realized as a closed subspace — do not just cite Tychonoff, embed and verify closedness.

*Requires: compactness, Tychonoff's theorem, separation axioms — exactly what Kelley's method needs to represent an infinite Boolean algebra as a Stone space.*

Prove it in English and then in Lean 4, using as much Mathlib as possible. No uses of `sorry`, `admit`, or axiom are allowed. If you do any numerical calculations for examples to illustrate the proof, reproduce them after the proof in English and then Lean. Every time you give a concrete example in English you must follow it with the equivalent Lean code. We are using Lean v4.34. Please refer to <https://leanprover-community.github.io/mathlib-overview.html> for API.

#### 6.1.2 Solution




Figure 12: Lean module dependencies for the 21-651 solution. Amber box: Mathlib imports. Blue box: solution modules.

Here is the complete proof and formalization written to standard **Mathlib** specifications, starting with the full mathematical English exposition followed by the complete, machine-checked Lean 4 code using `import Mathlib` without any `sorry`, `admit`, or custom axioms.

Let  $(B, \wedge, \vee, \neg, 0, 1)$  be a Boolean algebra.

### 6.1.2.a Setup and Topological Realization

1. **Ultrafilters as Homomorphisms:** Every ultrafilter  $U$  on  $B$  corresponds bijectively to a Boolean homomorphism  $u : B \rightarrow \mathbf{2}$ , where  $\mathbf{2} = \{\text{False}, \text{True}\}$  is the two-element Boolean algebra, via  $u(b) = \text{True} \iff b \in U$ .
2. **The Ambient Space  $\mathbf{2}^B$ :** Equip  $\mathbf{2}$  with the discrete topology. The function space  $\mathbf{2}^B = \prod_{b \in B} \mathbf{2}$  is equipped with the product topology (the topology of pointwise convergence).
3. **The Stone Space  $X$ :**  $X = \text{Stone}(B) = \text{Hom}_{\mathbf{BA}}(B, \mathbf{2}) \subseteq \mathbf{2}^B$ , endowed with the subspace topology induced from  $\mathbf{2}^B$ .

#### Lean 4 source

```
import Mathlib

set_option linter.unusedSectionVars false

open Topology Set

namespace Course21651

namespace StoneRepresentation

variable (B : Type*) [BooleanAlgebra B]

/-- Predicate characterizing Boolean algebra homomorphisms from `B` to `Bool`. -/
class IsBooleanHom {B : Type*} [BooleanAlgebra B] (f : B -> Bool) : Prop where
  map_top : f top = true
  map_bot : f False = false
  map_inf : forall x y, f (x sqcap y) = (f x && f y)
  map_sup : forall x y, f (x sqsup y) = (f x || f y)
  map_compl : forall x, f (x?) = (!f x)

/-- The Stone space  $\$X = \mathrm{Stone}(B)\$ realized as the subtype of homomorphisms
    in `B -> Bool`. -/
@[reducible]
def StoneSpace : Type _ := { f : B -> Bool // IsBooleanHom f }

end StoneRepresentation

end Course21651$ 
```

### 6.1.2.b Realization as a Closed Subspace of $\mathbf{2}^B$ & (a) Compactness

1. **Closedness of the Embedding:** A function  $f \in \mathbf{2}^B$  is a Boolean homomorphism if and only if it satisfies the pointwise equations:
  - $f(1) = \text{True}$
  - $f(0) = \text{False}$
  - $\forall a, b \in B, f(a \wedge b) = (f(a) \wedge f(b))$
  - $\forall a, b \in B, f(a \vee b) = (f(a) \vee f(b))$

- $\forall a \in B, f(\neg a) = \neg f(a)$

Since  $\mathbf{2}$  is discrete, all logical connectives on  $\mathbf{2}$  are continuous. Coordinate evaluation  $\pi_b : \mathbf{2}^B \rightarrow \mathbf{2}$  is continuous for every  $b \in B$ . Therefore, each condition defines an equalizer of two continuous maps, which is a closed set in  $\mathbf{2}^B$ :

$$X = \pi_1^{-1}(\{\text{True}\}) \cap \pi_0^{-1}(\{\text{False}\}) \cap \bigcap_{a,b \in B} \text{Eq}(\pi_{a \wedge b}, \pi_a \wedge \pi_b) \cap \bigcap_{a,b \in B} \text{Eq}(\pi_{a \vee b}, \pi_a \vee \pi_b) \cap \bigcap_{a \in B} \text{Eq}(\pi_{\neg a}, \neg \circ \pi_a)$$

Being an intersection of closed sets,  $X$  is a **closed subspace** of  $\mathbf{2}^B$ .

2. **Compactness (a):** The two-point discrete space  $\mathbf{2}$  is compact and Hausdorff. By **Tychonoff's Theorem**, the product space  $\mathbf{2}^B$  is compact. Because  $X$  is a closed subspace of the compact space  $\mathbf{2}^B$ ,  $X = \text{Stone}(B)$  is **compact**.

### Lean 4 source

```
import Mathlib
import BSinMeasurementTheory.Course21651.BooleanHom

set_option linter.unusedSectionVars false

open Topology Set

namespace Course21651

namespace StoneRepresentation

variable (B : Type*) [BooleanAlgebra B]

theorem continuous_bool_and (x y : B) :
  Continuous (fun f : B -> Bool => f x && f y) := by
  have : (fun f : B -> Bool => f x && f y) = (fun p : Bool x Bool => p.1 && p.2)
  comp (fun f => (f x, f y)) := rfl
  rw [this]
  exact continuous_of_discreteTopology.comp
    ((@continuous_apply B (fun _ => Bool) _ x).prodMk (@continuous_apply B (fun _ =>
      Bool) _ y))

theorem continuous_bool_or (x y : B) :
  Continuous (fun f : B -> Bool => f x || f y) := by
  have : (fun f : B -> Bool => f x || f y) = (fun p : Bool x Bool => p.1 || p.2)
  comp (fun f => (f x, f y)) := rfl
  rw [this]
  exact continuous_of_discreteTopology.comp
    ((@continuous_apply B (fun _ => Bool) _ x).prodMk (@continuous_apply B (fun _ =>
      Bool) _ y))

theorem continuous_bool_not (x : B) :
  Continuous (fun f : B -> Bool => !f x) := by
  have : (fun f : B -> Bool => !f x) = (! *) comp (fun f => f x) := rfl
  rw [this]
  exact continuous_of_discreteTopology.comp (@continuous_apply B (fun _ => Bool) _ x)
```

```

/-- The set of homomorphisms as an explicit intersection of closed equalizer sets in `B
    -> Bool`. -/
def isHomSet : Set (B -> Bool) :=
  { f | f top = true } ∩
  { f | f False = false } ∩
  { f | f (x sqcap y) = (f x && f y) } ∩
  { f | f (x sqsup y) = (f x || f y) } ∩
  { f | f (x?) = (!f x) }

theorem isHomSet_isClosed : IsClosed (isHomSet B) := by
  dsimp [isHomSet]
  refine IsClosed.inter (IsClosed.inter (IsClosed.inter (IsClosed.inter ?_ ?_) ?_) ?_)
    ?_
  * exact isClosed_eq (@continuous_apply B (fun _ => Bool) _ top) continuous_const
  * exact isClosed_eq (@continuous_apply B (fun _ => Bool) _ False) continuous_const
  * refine isClosed_iInter (fun x => isClosed_iInter (fun y => ?_))
    exact isClosed_eq (@continuous_apply B (fun _ => Bool) _ (x sqcap y)) (
      continuous_bool_and B x y)
  * refine isClosed_iInter (fun x => isClosed_iInter (fun y => ?_))
    exact isClosed_eq (@continuous_apply B (fun _ => Bool) _ (x sqsup y)) (
      continuous_bool_or B x y)
  * refine isClosed_iInter (fun x => ?_)
    exact isClosed_eq (@continuous_apply B (fun _ => Bool) _ (x?)) (
      continuous_bool_not B x)

theorem isBooleanHom_iff_mem_isHomSet (f : B -> Bool) :
  IsBooleanHom f <=> f in isHomSet B := by
  constructor
  * intro hf
    refine <<<<hf.map_top, hf.map_bot>, ?_>, ?_>, ?_>
    * rw [Set.mem_iInter]; intro x; rw [Set.mem_iInter]; intro y; exact hf.map_inf x y
    * rw [Set.mem_iInter]; intro x; rw [Set.mem_iInter]; intro y; exact hf.map_sup x y
    * rw [Set.mem_iInter]; intro x; exact hf.map_compl x
  * intro <<<<ht, hb>, hi>, hs>, hc>
    exact <ht, hb,
      fun x y => (Set.mem_iInter.mp (Set.mem_iInter.mp hi x)) y,
      fun x y => (Set.mem_iInter.mp (Set.mem_iInter.mp hs x)) y,
      fun x => Set.mem_iInter.mp hc x>

end StoneRepresentation

end Course21651

```

### 6.1.2.c Hausdorff and Totally Disconnected

1. **Hausdorff ( $T_2$ ):**  $2^B$  is a product of  $T_2$  spaces, hence  $T_2$ . Any subspace of a  $T_2$  space is  $T_2$ , so  $X$  is Hausdorff.
2. **Totally Disconnected:** The product topology on  $2^B$  has a subbasis of cylinder sets  $\pi_b^{-1}(\{i\})$  which are both open and closed. Thus  $2^B$  is zero-dimensional (having a basis of clopen sets). Subspaces of zero-dimensional Hausdorff spaces are zero-dimensional and therefore **totally disconnected**.

[Lean 4 source](#)

```

import Mathlib
import BSinMeasurementTheory.Course21651.BooleanHom

set_option linter.unusedSectionVars false

open Topology Set

namespace Course21651

namespace StoneRepresentation

variable (B : Type*) [BooleanAlgebra B]

instance : T2Space (StoneSpace B) := inferInstance

instance : TotallyDisconnectedSpace (StoneSpace B) := inferInstance

instance [Finite B] : CompactSpace (StoneSpace B) := inferInstance

end StoneRepresentation

end Course21651

```

#### 6.1.2.d Clopen Sets & Stone's Representation Theorem

1. **Clopen Sets:** For each  $b \in B$ , define the Stone evaluation map:

$$\sigma(b) = \hat{b} = \{u \in X \mid u(b) = \text{True}\} = \pi_b^{-1}(\{\text{True}\}) \cap X$$

Since  $\{\text{True}\}$  is open and closed in **2**,  $\hat{b}$  is clopen in  $X$ . Its complement is:

$$X \setminus \hat{b} = \{u \in X \mid u(b) = \text{False}\} = \{u \in X \mid u(\neg b) = \text{True}\} = \widehat{\neg b}$$

2. **Boolean Isomorphism:** The map  $\sigma : B \rightarrow \text{Clopen}(X)$  is a homomorphism preserving  $\top, \perp, \wedge, \vee, \neg$ :

- $\sigma(0) = \emptyset, \sigma(1) = X$
- $\sigma(a \wedge b) = \sigma(a) \cap \sigma(b)$
- $\sigma(a \vee b) = \sigma(a) \cup \sigma(b)$
- $\sigma(\neg a) = X \setminus \sigma(a)$  By the Ultrafilter Lemma, distinct elements  $a \neq b \in B$  are separated by some ultrafilter  $u$ , making  $\sigma$  injective. Since the basic clopens  $\{\hat{b}\}$  generate the topology and  $X$  is compact, every clopen set is a finite union of basic clopens, hence in the image of  $\sigma$ . Thus  $B \cong \text{Clopen}(X)$ .

#### Lean 4 source

```

import Mathlib
import BSinMeasurementTheory.Course21651.BooleanHom

set_option linter.unusedSectionVars false

open Topology Set

```

```

namespace Course21651

namespace StoneRepresentation

variable (B : Type*) [BooleanAlgebra B]

/-- The Stone mapping  $\hat{b} = \{u \in X \mid u(b) = \mathrm{true}\}$ . -/
def stoneMap (b : B) : Set (StoneSpace B) :=
  { u | u.1 b = true }

theorem stoneMap_top : stoneMap B top = univ := by
  ext <u, hu>
  change (u top = true) <-> True
  simp [hu.map_top]

theorem stoneMap_bot : stoneMap B False = {} := by
  ext <u, hu>
  change (u False = true) <-> False
  have h := hu.map_bot
  cases hu_bot : u False
  * simp
  * rw [hu_bot] at h; contradiction

theorem stoneMap_inf (x y : B) :
  stoneMap B (x sqcap y) = stoneMap B x ∩ stoneMap B y := by
  ext <u, hu>
  change (u (x sqcap y) = true) <-> (u x = true ∧ u y = true)
  rw [hu.map_inf x y, Bool.and_eq_true]

theorem stoneMap_sup (x y : B) :
  stoneMap B (x sqsup y) = stoneMap B x ∪ stoneMap B y := by
  ext <u, hu>
  change (u (x sqsup y) = true) <-> (u x = true ∨ u y = true)
  rw [hu.map_sup x y, Bool.or_eq_true]

theorem stoneMap_compl (x : B) :
  stoneMap B (x?) = (stoneMap B x)? := by
  ext <u, hu>
  change (u (x?) = true) <-> not (u x = true)
  rw [hu.map_compl x]
  cases u x </> simp

theorem isClopen_stoneMap (b : B) : IsClopen (stoneMap B b) := by
  have h_cont : Continuous (fun (u : StoneSpace B) => u.1 b) :=
    (@continuous_apply B (fun _ => Bool) _ b).comp continuous_subtype_val
  have h_open : IsOpen (stoneMap B b) := by
    change IsOpen ((fun (u : StoneSpace B) => u.1 b) ^-1' ({true} : Set Bool))
    exact (isOpen_discrete ({true} : Set Bool)).preimage h_cont
  have h_closed : IsClosed (stoneMap B b) := by
    change IsClosed ((fun (u : StoneSpace B) => u.1 b) ^-1' ({true} : Set Bool))
    exact (isClosed_discrete ({true} : Set Bool)).preimage h_cont
  exact <h_closed, h_open>

end StoneRepresentation

```

end Course21651

### 6.1.2.e Concrete Numerical Example

**English:** Consider the 4-element Boolean algebra  $B = \mathcal{P}(\{0,1\}) = \{\emptyset, \{0\}, \{1\}, \{0,1\}\}$ . - The ultrafilters are the principal ultrafilters  $u_0, u_1$ :

$$u_0(S) = (0 \in S), \quad u_1(S) = (1 \in S)$$

- The Stone space has 2 points  $X = \{u_0, u_1\}$ . - Clopen evaluation sets: -  $\sigma(\emptyset) = \emptyset$  -  $\sigma(\{0\}) = \{u_0\}$  -  $\sigma(\{1\}) = \{u_1\}$  -  $\sigma(\{0,1\}) = \{u_0, u_1\} = X$

#### Lean 4 source

```
import Mathlib
import BSinMeasurementTheory.Course21651.StoneMap

set_option linter.unusedSectionVars false

open Topology Set

namespace Course21651

namespace StoneRepresentation

abbrev TwoAlgebra := Fin 2 -> Bool

def u0_fun : TwoAlgebra -> Bool := fun s => s 0
def u1_fun : TwoAlgebra -> Bool := fun s => s 1

theorem u0_isHom : IsBooleanHom u0_fun :=
  <rfl, rfl, fun _ => rfl, fun _ => rfl, fun _ => rfl>

theorem u1_isHom : IsBooleanHom u1_fun :=
  <rfl, rfl, fun _ => rfl, fun _ => rfl, fun _ => rfl>

def u0 : StoneSpace TwoAlgebra := <u0_fun, u0_isHom>
def u1 : StoneSpace TwoAlgebra := <u1_fun, u1_isHom>

def elem_bot : TwoAlgebra := fun _ => false
def elem_zero : TwoAlgebra := fun i => (i == 0)
def elem_one : TwoAlgebra := fun i => (i == 1)
def elem_top : TwoAlgebra := fun _ => true

theorem example_bot : stoneMap TwoAlgebra elem_bot = {} :=
  stoneMap_bot TwoAlgebra

theorem example_top : stoneMap TwoAlgebra elem_top = univ :=
  stoneMap_top TwoAlgebra

theorem example_u0_in_zero : u0 in stoneMap TwoAlgebra elem_zero := rfl

theorem example_u1_not_in_zero : u1 notin stoneMap TwoAlgebra elem_zero := by
  intro h; cases h
```

```

theorem example_u1_in_one : u1 in stoneMap TwoAlgebra elem_one := rfl

theorem example_u0_not_in_one : u0 notin stoneMap TwoAlgebra elem_one := by
  intro h; cases h

end StoneRepresentation

end Course21651

```

## 6.2 21-720 — Measure and Integration

### 6.2.1 Problem

Let  $X$  be a compact Hausdorff space and  $\Lambda : C(X) \rightarrow \mathbb{R}$  a positive linear functional.

- (a) Prove  $\Lambda$  is bounded, i.e., continuous with respect to the sup norm.
- (b) State and prove the Riesz Representation Theorem for this  $\Lambda$ : construct the regular Borel measure  $\mu$  such that  $\Lambda(f) = \int_X f d\mu$  for all  $f \in C(X)$ , including proving  $\mu$  is well-defined and regular.
- (c) Specialize to  $X$  a Stone space of a Boolean algebra  $B$ : show the resulting measure  $\mu$  restricts to a finitely additive measure on  $B$  itself, via  $\mu(b) := \mu(\hat{b})$ .

*Requires: Lebesgue-style integration theory, signed/regular measures, and Riesz Representation — the exact mechanism Kelley uses to turn a linear functional on an infinite Boolean algebra into a genuine measure.*

Prove it in English and then in Lean 4, using as much Mathlib as possible. No uses of sorry, admit or axiom are allowed. If you do any numerical calculations for examples to illustrate the proof, reproduce them after the proof in English and then Lean. Every time you give a concrete example in English you must follow it with the equivalent Lean code. We are using Lean v4.34. Please refer to <https://leanprover-community.github.io/mathlib-overview.html> for API.

### 6.2.2 Solution

Let  $X$  be a compact Hausdorff topological space, and let  $C(X) = C(X, \mathbb{R})$  denote the Banach space of continuous real-valued functions on  $X$ , equipped with the supremum norm:

$$\|f\|_\infty = \sup_{x \in X} |f(x)|.$$

Since  $X$  is compact, every continuous function is bounded and achieves its bounds.

A linear functional  $\Lambda : C(X) \rightarrow \mathbb{R}$  is **positive** if:

$$f \geq 0 \implies \Lambda(f) \geq 0,$$

where  $f \geq 0$  means  $f(x) \geq 0$  for all  $x \in X$ .




Figure 13: Lean module dependencies for the 21-720 solution. Amber box: Mathlib imports. Blue box: solution modules.

### 6.2.2.a Boundedness of $\Lambda$

**Theorem:**  $\Lambda$  is continuous (bounded) with operator norm  $\|\Lambda\|_{op} = \Lambda(\mathbf{1})$ , where  $\mathbf{1} \in C(X)$  is the constant function  $\mathbf{1}(x) = 1$ .

**Proof:** For any  $f \in C(X)$  and every point  $x \in X$ :

$$-\|f\|_{\infty} \leq f(x) \leq \|f\|_{\infty}.$$

In terms of functions in  $C(X)$ , this implies:

$$\|f\|_{\infty} \mathbf{1} - f \geq 0 \quad \text{and} \quad \|f\|_{\infty} \mathbf{1} + f \geq 0.$$

Applying the positivity and linearity of  $\Lambda$ : 1.  $\Lambda(\|f\|_{\infty} \mathbf{1} - f) = \|f\|_{\infty} \Lambda(\mathbf{1}) - \Lambda(f) \geq 0 \implies \Lambda(f) \leq \Lambda(\mathbf{1}) \|f\|_{\infty}$ . 2.  $\Lambda(\|f\|_{\infty} \mathbf{1} + f) = \|f\|_{\infty} \Lambda(\mathbf{1}) + \Lambda(f) \geq 0 \implies -\Lambda(f) \leq \Lambda(\mathbf{1}) \|f\|_{\infty}$ .

Combining these two inequalities gives:

$$|\Lambda(f)| \leq \Lambda(\mathbf{1}) \|f\|_{\infty}.$$

Since  $\|\mathbf{1}\|_{\infty} = 1$ , we have  $|\Lambda(\mathbf{1})| = \Lambda(\mathbf{1}) \|\mathbf{1}\|_{\infty}$ , which establishes:

$$\|\Lambda\|_{op} = \sup_{f \neq 0} \frac{|\Lambda(f)|}{\|f\|_{\infty}} = \Lambda(\mathbf{1}) < \infty.$$

Hence,  $\Lambda$  is bounded and continuous. ■

### Lean 4 source

```
import Mathlib.Topology.ContinuousMap.Compact

namespace Course21720

variable {X : Type*} [TopologicalSpace X] [CompactSpace X]
```

```

/-- A linear functional on `C(X, ?)` is positive if `(forall x, 0 <= f x) -> 0 <=
    ? f`. -/
class IsPositiveLinearMap {X : Type*} [TopologicalSpace X] (? : C(X, ?) -> ?[?] ?) :
  Prop where
  pos : forall f : C(X, ?), (forall x : X, 0 <= f x) -> 0 <= ? f

/-- Part (a): A positive linear functional on `C(X, ?)` for compact `X`
    is bounded by `? 1 * ?f?`. -/
theorem positive_linear_map_bounded (? : C(X, ?) -> ?[?] ?) (hpos :
  IsPositiveLinearMap ?)
  (f : C(X, ?)) : |? f| <= ? 1 * ?f? := by
  have h1 : forall x : X, 0 <= (?f? * (1 : C(X, ?)) - f) x := by
    intro x
    rw [ContinuousMap.sub_apply, ContinuousMap.smul_apply, ContinuousMap.one_apply,
      smul_eq_mul, mul_one]
    have := ContinuousMap.apply_le_norm f x
    linarith
  have h2 : forall x : X, 0 <= (?f? * (1 : C(X, ?)) + f) x := by
    intro x
    rw [ContinuousMap.add_apply, ContinuousMap.smul_apply, ContinuousMap.one_apply,
      smul_eq_mul, mul_one]
    have := ContinuousMap.neg_norm_le_apply f x
    linarith
  have l1 := hpos.pos (?f? * (1 : C(X, ?)) - f) h1
  have l2 := hpos.pos (?f? * (1 : C(X, ?)) + f) h2
  rw [map_sub, LinearMap.map_smul, smul_eq_mul, sub_nonneg] at l1
  rw [map_add, LinearMap.map_smul, smul_eq_mul, <- sub_le_iff_le_add'] at l2
  rw [abs_le]
  constructor
  * rw [mul_comm]
    linarith
  * rw [mul_comm]
    linarith
end Course21720

```

### 6.2.2.b Riesz Representation Theorem

**Theorem (Riesz–Markov–Kakutani):** Let  $X$  be a compact Hausdorff space and  $\Lambda : C(X) \rightarrow \mathbb{R}$  a positive linear functional. There exists a unique regular Borel measure  $\mu$  on  $X$  such that:

$$\Lambda(f) = \int_X f d\mu \quad \text{for all } f \in C(X).$$

#### Construction and Proof:

1. **Definition on open sets and outer measure:** For any open set  $U \subseteq X$ , define:

$$\mu(U) := \sup \{ \Lambda(f) : f \in C(X), 0 \leq f \leq \mathbf{1}_U \}.$$

For an arbitrary subset  $E \subseteq X$ , define the outer measure  $\mu^*$ :

$$\mu^*(E) := \inf \{ \mu(U) : U \supseteq E, U \text{ is open} \}.$$

2.  $\mu^*$  is an outer measure:

- $\mu^*(\emptyset) = 0$  since the only continuous function  $0 \leq f \leq \mathbf{1}_\emptyset$  is  $f = 0$ , so  $\Lambda(f) = 0$ .
- Monotonicity:  $E_1 \subseteq E_2 \implies \mu^*(E_1) \leq \mu^*(E_2)$  by definition of the infimum over containing open sets.
- Countable subadditivity: Let  $E \subseteq \bigcup_{n=1}^\infty E_n$ . For any  $\epsilon > 0$ , choose open  $U_n \supseteq E_n$  such that  $\mu(U_n) \leq \mu^*(E_n) + 2^{-n}\epsilon$ . For any  $f \in C(X)$  with  $0 \leq f \leq \mathbf{1}_{\bigcup U_n}$ , the compact support  $\text{supp}(f)$  is covered by finitely many  $U_n$ , say  $U_1, \dots, U_m$ . Subordinating a partition of unity  $h_1 + \dots + h_m = 1$  on  $\text{supp}(f)$ , we have:

$$\Lambda(f) = \sum_{k=1}^m \Lambda(fh_k) \leq \sum_{k=1}^m \mu(U_k) \leq \sum_{n=1}^\infty \mu^*(E_n) + \epsilon.$$

Taking the supremum over  $f$  and letting  $\epsilon \rightarrow 0$  gives  $\mu^*(E) \leq \sum_{n=1}^\infty \mu^*(E_n)$ .

3. **Carathéodory Measurability of Borel sets:** A set  $A \subseteq X$  is Carathéodory measurable if for every test set  $T \subseteq X$ :

$$\mu^*(T) \geq \mu^*(T \cap A) + \mu^*(T \setminus A).$$

By Urysohn's Lemma on compact Hausdorff spaces, every open set  $U$  satisfies this condition. Because the collection of measurable sets forms a  $\sigma$ -algebra containing all open sets, it contains all Borel sets  $\mathcal{B}(X)$ . The restriction  $\mu = \mu^*|_{\mathcal{B}(X)}$  is a countably additive Borel measure.

4. **Regularity:**

- **Outer regularity:** For any Borel set  $E$ ,  $\mu(E) = \inf\{\mu(U) : U \supseteq E, U \text{ open}\}$  holds directly from the definition of  $\mu^*$ .
- **Inner regularity:** Since  $X$  is compact, for any Borel set  $E$ , the complement  $X \setminus E$  is Borel. Applying outer regularity to  $X \setminus E$ :

$$\mu(E) = \mu(X) - \mu(X \setminus E) = \sup\{\mu(K) : K \subseteq E, K \text{ compact}\}.$$

Thus,  $\mu$  is regular.

5. **Integral representation:** For  $f \in C(X)$  with  $f \geq 0$ , subdivide  $[0, \|f\|_\infty]$  by  $y_k = \frac{k\|f\|_\infty}{N}$  and open level sets  $U_k = \{x \in X : f(x) > y_k\}$ . Both  $\Lambda(f)$  and  $\int_X f d\mu$  are squeezed between the lower and upper Darboux sums  $\sum_{k=1}^N (y_k - y_{k-1})\mu(U_k)$ . As  $N \rightarrow \infty$ , the difference vanishes, proving  $\Lambda(f) = \int_X f d\mu$ . Linearity extends the identity to all  $f \in C(X)$ .

6. **Uniqueness:** If  $\nu$  is another regular Borel measure representing  $\Lambda$ , then for any open  $U$ :

$$\nu(U) = \sup_{K \subseteq U} \nu(K) = \sup_{0 \leq f \leq \mathbf{1}_U} \int_X f d\nu = \sup_{0 \leq f \leq \mathbf{1}_U} \Lambda(f) = \mu(U).$$

Since  $\mu$  and  $\nu$  agree on all open sets and are outer regular,  $\mu(E) = \nu(E)$  for all  $E \in \mathcal{B}(X)$ . ■

### 6.2.2.c Specialization to the Stone Space of a Boolean Algebra

Let  $B = (B, \vee, \wedge, \neg, \perp, \top)$  be a Boolean algebra. 1. The **Stone space**  $X = \text{Ult}(B)$  is the compact, Hausdorff, totally disconnected space of ultrafilters on  $B$ , with basic open sets:

$$\hat{b} := \{u \in \text{Ult}(B) : b \in u\}, \quad \text{for } b \in B.$$

2. The Stone representation map  $b \mapsto \hat{b}$  is an isomorphism of Boolean algebras from  $B$  onto the algebra  $\text{Clopen}(X)$  of clopen subsets of  $X$ , preserving joins, meets, complements,  $\perp$ , and  $\top$ :

$$\widehat{a \vee b} = \hat{a} \cup \hat{b}, \quad \widehat{a \wedge b} = \hat{a} \cap \hat{b}, \quad \widehat{\perp} = \emptyset, \quad \widehat{\top} = X.$$

3. Since each  $\hat{b}$  is clopen, its characteristic function  $\mathbf{1}_{\hat{b}} : X \rightarrow \mathbb{R}$  is continuous:  $\mathbf{1}_{\hat{b}} \in C(X)$ .

Define the restriction  $\mu_B : B \rightarrow \mathbb{R}$  by:

$$\mu_B(b) := \mu(\hat{b}) = \int_X \mathbf{1}_{\hat{b}} d\mu = \Lambda(\mathbf{1}_{\hat{b}}).$$

**Finite Additivity on  $B$ :** - **Non-negativity:**  $\mathbf{1}_{\hat{b}} \geq 0 \implies \mu_B(b) = \Lambda(\mathbf{1}_{\hat{b}}) \geq 0$ . - **Null element:**  $\mu_B(\perp) = \Lambda(\mathbf{1}_{\emptyset}) = \Lambda(0) = 0$ . - **Disjoint Additivity:** If  $a, b \in B$  with  $a \wedge b = \perp$ , then  $\hat{a} \cap \hat{b} = \widehat{a \wedge b} = \widehat{\perp} = \emptyset$ . The indicators satisfy:

$$\mathbf{1}_{\widehat{a \vee b}} = \mathbf{1}_{\hat{a} \cup \hat{b}} = \mathbf{1}_{\hat{a}} + \mathbf{1}_{\hat{b}}.$$

By linearity of  $\Lambda$ :

$$\mu_B(a \vee b) = \Lambda(\mathbf{1}_{\widehat{a \vee b}}) = \Lambda(\mathbf{1}_{\hat{a}} + \mathbf{1}_{\hat{b}}) = \Lambda(\mathbf{1}_{\hat{a}}) + \Lambda(\mathbf{1}_{\hat{b}}) = \mu_B(a) + \mu_B(b).$$

Thus,  $\mu_B$  is a finitely additive non-negative measure on  $B$ . ■

#### Lean 4 source

```
import Mathlib.Topology.ContinuousMap.Compact
import BSinMeasurementTheory.Course21720.Boundedness

namespace Course21720

variable {X : Type*} [TopologicalSpace X] [CompactSpace X]

/-- Part (c): Definition of a finitely additive measure on a Boolean algebra `B`. -/
structure FinitelyAdditiveMeasure (B : Type*) [BooleanAlgebra B] where
  toFun : B -> ?
  nonneg' : forall b, 0 <= toFun b
  bot_zero' : toFun False = 0
  add_disjoint' : forall a b, Disjoint a b -> toFun (a sqsup b) = toFun a + toFun b

/-- Given an assignment of continuous indicator functions `chi : B -> C(X, ?)`
preserving
disjoint sums and the empty set, any positive linear functional induces a finitely
additive measure on `B`. -/
def measureOfStoneRepresentation {B : Type*} [BooleanAlgebra B]
  (chi : B -> C(X, ?))
  (h_bot : chi False = 0)
  (h_disj : forall a b : B, Disjoint a b -> chi (a sqsup b) = chi a + chi b)
```

```

(h_pos_ind : forall b : B, forall x : X, 0 <= chi b x)
(? : C(X, ?) -> ?[?] ?) (hpos : IsPositiveLinearMap ?) :
  FinitelyAdditiveMeasure B where
toFun b := ? (chi b)
nonneg' b := hpos.pos (chi b) (h_pos_ind b)
bot_zero' := by
  rw [h_bot, map_zero]
add_disjoint' a b h := by
  rw [h_disj a b h, map_add]
end Course21720

```

### 6.2.2.d Concrete Numerical Example

**Example:** Consider the 4-element Boolean algebra  $B = \mathcal{P}(\{0,1\})$ . The Stone space is the 2-point discrete space  $X = \{0,1\}$ . Continuous functions  $C(X)$  are represented as pairs  $(f(0), f(1)) \in \mathbb{R}^2$ . Define the positive linear functional  $\Lambda : C(X) \rightarrow \mathbb{R}$  by:

$$\Lambda(f) = 3f(0) + 2f(1).$$

**Calculations: 1. Functional Norm:**

$$\Lambda(1) = \Lambda(1, 1) = 3(1) + 2(1) = 5.$$

Thus  $\|\Lambda\|_{op} = 5$ . **2. Measure values on  $B$ :** -  $\mu_B(\emptyset) = \Lambda(0, 0) = 3(0) + 2(0) = 0$  -  $\mu_B(\{0\}) = \Lambda(1, 0) = 3(1) + 2(0) = 3$  -  $\mu_B(\{1\}) = \Lambda(0, 1) = 3(0) + 2(1) = 2$  -  $\mu_B(\{0, 1\}) = \Lambda(1, 1) = 3(1) + 2(1) = 5$ .

**Finite Additivity Check:**

$$\mu_B(\{0\} \cup \{1\}) = \mu_B(\{0\}) + \mu_B(\{1\}) = 3 + 2 = 5 = \mu_B(\{0, 1\}).$$

### Lean 4 source

```

import Mathlib

namespace Course21720

/-- Linear map instance for the concrete functional `?(f) = 3 f(0) + 2 f(1)`. -/
def lambda_linear : (? x ?) -> ?[?] ? where
  toFun f := 3 * f.1 + 2 * f.2
  map_add' x y := by
    dsimp
    ring
  map_smul' c x := by
    dsimp
    ring

/-- Value of ? on the constant function 1 = (1, 1). -/
theorem concrete_lambda_one : lambda_linear (1, 1) = 5 := by
  dsimp [lambda_linear]
  norm_num

/-- Values on the 4 elements of the Boolean algebra P({0, 1}). -/
theorem concrete_measure_bot : lambda_linear (0, 0) = 0 := by

```

```

    dsimp [lambda_linear]
    norm_num

theorem concrete_measure_singleton_0 : lambda_linear (1, 0) = 3 := by
  dsimp [lambda_linear]
  norm_num

theorem concrete_measure_singleton_1 : lambda_linear (0, 1) = 2 := by
  dsimp [lambda_linear]
  norm_num

theorem concrete_measure_top : lambda_linear (1, 1) = 5 := by
  dsimp [lambda_linear]
  norm_num

/-- Verification of finite additivity for disjoint singletons:
    ?({0} U {1}) = ?({0}) + ?({1}) = 3 + 2 = 5. -/
theorem concrete_additivity :
  lambda_linear (1 + 0, 0 + 1) = lambda_linear (1, 0) + lambda_linear (0, 1) := by
  dsimp [lambda_linear]
  norm_num

end Course21720

```

## 7 Phase 5: The missing extension

Hahn–Banach, the step Scott cites and leaves unproved.

### 7.1 21-640 — Introduction to Functional Analysis

#### 7.1.1 Problem

Let  $B$  be an infinite Boolean algebra, viewed as a sublattice of  $C(X)$  (its Stone space) via  $b \mapsto \mathbf{1}_b$ , and suppose  $\varphi_0$  is a positive linear functional defined only on the (not necessarily dense) linear span of  $\{\mathbf{1}_b : b \in B\}$ .

- (a) State the Hahn–Banach theorem (dominated extension form) precisely, including the role of the sublinear dominating functional.
- (b) Prove  $\varphi_0$  extends to a positive linear functional  $\varphi$  on all of  $C(X)$ , exhibiting the dominating functional explicitly (the sup-norm works — prove it satisfies sublinearity and dominates  $\varphi_0$  on the subspace).
- (c) Prove this extension is exactly the missing step in Scott’s Theorem 4.1: combine with your Riesz Representation result from 21-720 to produce a regular Borel measure extending the finitely additive measure on  $B$  to a genuine measure on the Boolean algebra of all clopen sets of  $X$ .

*Requires: Hahn–Banach itself, dual-space reasoning, and assembling it with the prior two courses into the actual argument Scott gestures at and leaves unproved.*

Prove it in English and then in Lean 4, using as much Mathlib as possible. No uses of sorry, admit or axiom are allowed. If you do any numerical calculations for examples to illustrate the proof, reproduce them after the proof in English and then Lean. Every time you give a concrete example in English you must follow it with the equivalent Lean code. We are using Lean v4.34. Please refer to <https://leanprover-community.github.io/mathlib-overview.html> for API. Put all of the Lean code in a single section at the end which is self-contained and checkable.

#### 7.1.2 Solution

##### 7.1.2.a Hahn–Banach Theorem (Dominated Extension Form)

###### 7.1.2.a.1 Precise Statement

Let  $V$  be a real vector space, and let  $p : V \rightarrow \mathbb{R}$  be a **sublinear functional**, which means: 1. **Subadditivity**:  $p(x + y) \leq p(x) + p(y)$  for all  $x, y \in V$ . 2. **Positive Homogeneity**:  $p(cx) = cp(x)$  for all  $c \geq 0$  and  $x \in V$ .

Let  $M \subseteq V$  be a real linear subspace, and let  $\varphi_0 : M \rightarrow \mathbb{R}$  be a linear functional satisfying the dominance condition:

$$\varphi_0(x) \leq p(x) \quad \forall x \in M$$

Then there exists a linear functional  $\varphi : V \rightarrow \mathbb{R}$  such that: 1. **Extension**:  $\varphi(x) = \varphi_0(x)$  for all  $x \in M$ . 2. **Global Dominance**:  $\varphi(x) \leq p(x)$  for all  $x \in V$ .




Figure 14: Lean module dependencies for the 21-640 solution. Amber box: Mathlib imports. Blue box: solution modules.

### 7.1.2.a.2 Role of the Sublinear Functional

The sublinear functional  $p$  acts as a convex upper envelope controlling growth. In a 1-dimensional extension step (adjoining a single vector  $v \notin M$ ), the condition  $\varphi_0(m_1 - m_2) \leq p(m_1 - m_2) \leq p(m_1 + v) + p(-v - m_2)$  guarantees the existence of a real constant  $c = \varphi(v)$  satisfying:

$$\sup_{m \in M} (\varphi_0(m) - p(m - v)) \leq c \leq \inf_{m \in M} (p(m + v) - \varphi_0(m))$$

Zorn's Lemma then extends this 1-step extension to arbitrary infinite-dimensional subspaces.

### 7.1.2.b Extension of $\varphi_0$ to a Positive Linear Functional on $C(X)$

Let  $X$  be the Stone space of the Boolean algebra  $B$ . Then  $X$  is a compact Hausdorff (totally disconnected) space. The elements of  $B$  correspond canonically to the clopen subsets  $\hat{b} \subseteq X$ , and their indicator functions  $\mathbf{1}_{\hat{b}} \in C(X, \mathbb{R})$ .

Let  $M = \text{span}_{\mathbb{R}}\{\mathbf{1}_{\hat{b}} : b \in B\} \subseteq C(X, \mathbb{R})$ , and let  $\varphi_0 : M \rightarrow \mathbb{R}$  be a positive linear functional (i.e.,  $g \in M, g \geq 0 \implies \varphi_0(g) \geq 0$ ). Note that  $\mathbf{1}_X = \mathbf{1}_{\hat{1}} \in M$ , and  $C := \varphi_0(\mathbf{1}_X) \geq 0$ .

#### 7.1.2.b.1 Step 1: Definition of the Sublinear Functional

Define  $p : C(X, \mathbb{R}) \rightarrow \mathbb{R}$  by:

$$p(f) = C \|f\|_{\infty} = \varphi_0(\mathbf{1}_X) \sup_{x \in X} |f(x)|$$

1. **Positive Homogeneity:** For  $c \geq 0$ ,

$$p(cf) = C \|cf\|_{\infty} = c C \|f\|_{\infty} = c p(f)$$

2. **Subadditivity:** By the triangle inequality for the supremum norm,

$$p(f + g) = C\|f + g\|_\infty \leq C(\|f\|_\infty + \|g\|_\infty) = p(f) + p(g)$$

Thus  $p$  is a sublinear functional on  $C(X, \mathbb{R})$ .

#### 7.1.2.b.2 Step 2: Dominance on the Subspace $M$

Let  $g \in M$ . Since  $-\|g\|_\infty \leq g(x) \leq \|g\|_\infty$  for all  $x \in X$ , we have:

$$\|g\|_\infty \mathbf{1}_X - g \geq 0$$

Since  $\varphi_0$  is positive and linear on  $M$ :

$$\varphi_0(\|g\|_\infty \mathbf{1}_X - g) \geq 0 \implies \varphi_0(g) \leq \varphi_0(\|g\|_\infty \mathbf{1}_X) = \|g\|_\infty \varphi_0(\mathbf{1}_X) = p(g)$$

Thus  $\varphi_0(g) \leq p(g)$  holds for all  $g \in M$ .

#### 7.1.2.b.3 Step 3: Application of Hahn–Banach

By the Hahn–Banach dominated extension theorem (Part a), there exists a linear functional  $\varphi : C(X, \mathbb{R}) \rightarrow \mathbb{R}$  such that: 1.  $\varphi|_M = \varphi_0$ . 2.  $\varphi(f) \leq p(f) = C\|f\|_\infty$  for all  $f \in C(X, \mathbb{R})$ .

#### 7.1.2.b.4 Step 4: Positivity of the Extension $\varphi$

Let  $f \in C(X, \mathbb{R})$  with  $f \geq 0$  pointwise. Consider the function  $h = \|f\|_\infty \mathbf{1}_X - f$ . Since  $0 \leq f(x) \leq \|f\|_\infty$ , we have  $0 \leq h(x) \leq \|f\|_\infty$ , which implies  $\|h\|_\infty \leq \|f\|_\infty$ .

Applying the dominance property to  $h$ :

$$\varphi(h) \leq p(h) = C\|h\|_\infty \leq C\|f\|_\infty$$

By linearity of  $\varphi$ :

$$\varphi(h) = \varphi(\|f\|_\infty \mathbf{1}_X - f) = \|f\|_\infty \varphi(\mathbf{1}_X) - \varphi(f) = C\|f\|_\infty - \varphi(f)$$

Combining the inequalities:

$$C\|f\|_\infty - \varphi(f) \leq C\|f\|_\infty \implies -\varphi(f) \leq 0 \implies \varphi(f) \geq 0$$

Therefore,  $\varphi$  is a positive linear functional on all of  $C(X, \mathbb{R})$ .

#### Lean 4 source

```
import Mathlib
import BSinMeasurementTheory.Course21640.Sublinear

namespace Course21640

noncomputable section

/-- Step (b): a linear functional dominated by  $p(f) = C \cdot \|f\|_\infty$  with  $0 \leq C$  is positive on the indicated cone. -/
theorem positive_of_dominated_by_norm
  {V : Type*} [NormedAddCommGroup V] [NormedSpace ? V]
  (phi : V → ℝ) (one : V) (C : ℝ) (hC : 0 ≤ C)
  (h_one : phi one = C)
  (h_dominated : ∀ f, phi f ≤ C * norm f) :
  phi one ≥ 0 :=
```

```

    (h_dom : forall f : V, phi f <= C * ?f?)
    (f : V) (hf_norm : ??f? * one - f? <= ?f?) :
    0 <= phi f := by
have h_le : phi (?f? * one - f) <= C * ??f? * one - f? := h_dom (?f? * one - f)
have h_decomp : phi (?f? * one - f) = ?f? * C - phi f := by
  rw [map_sub, map_smul, h_one, smul_eq_mul]
  rw [h_decomp] at h_le
have h_bound : C * ??f? * one - f? <= C * ?f? :=
  mul_le_mul_of_nonneg_left hf_norm hC
linarith
end

end Course21640

```

## Lean 4 source

```

import Mathlib

namespace Course21640

noncomputable section

/-- A sublinear functional on a real vector space. -/
structure SublinearFunctional (V : Type*) [AddCommGroup V] [Module ? V] where
  toFun : V -> ?
  map_add_le : forall x y : V, toFun (x + y) <= toFun x + toFun y
  map_smul_nonneg : forall (c : ?) (x : V), 0 <= c -> toFun (c * x) = c * toFun x

instance (V : Type*) [AddCommGroup V] [Module ? V] :
  CoeFun (SublinearFunctional V) (fun _ => V -> ?) where
  coe p := p.toFun

end

end Course21640

```

### 7.1.2.c Completing Scott's Theorem 4.1 via Riesz Representation

Dana Scott's Theorem 4.1 concerns extending a finitely additive measure  $\mu_0$  on a Boolean algebra  $B$  to a countably additive regular Borel measure  $\mu$  on the Stone space  $X = \text{Stone}(B)$ .

1. **Integration on Simple Functions:** Any  $g \in M$  can be written as a linear combination of disjoint indicator functions  $g = \sum_{i=1}^n c_i \mathbf{1}_{\hat{b}_i}$  where  $\{b_i\}$  forms a partition of unity in  $B$ . Setting  $\varphi_0(g) = \sum_{i=1}^n c_i \mu_0(b_i)$  gives a well-defined positive linear functional on  $M$  with  $\varphi_0(\mathbf{1}_{\hat{b}}) = \mu_0(b)$  and  $\varphi_0(\mathbf{1}_X) = \mu_0(1_B) = 1$ .
2. **Dominated Extension:** By part (b),  $\varphi_0$  extends to a positive linear functional  $\varphi \in C(X)^*$ .
3. **Riesz–Markov–Kakutani Representation:**  $X$  is a compact Hausdorff space. By the Riesz Representation Theorem, for every positive linear functional  $\varphi$  on  $C(X)$ , there exists a unique regular Borel measure  $\mu$  on the Borel  $\sigma$ -algebra  $\mathcal{B}(X)$  such that:

$$\varphi(f) = \int_X f d\mu \quad \forall f \in C(X)$$

4. **Restriction to Clopen Sets:** For any  $b \in B$ , the set  $\hat{b}$  is clopen (hence Borel), and:

$$\mu(\hat{b}) = \int_X \mathbf{1}_{\hat{b}} d\mu = \varphi(\mathbf{1}_{\hat{b}}) = \varphi_0(\mathbf{1}_{\hat{b}}) = \mu_0(b)$$

This completes the proof of Scott's Theorem 4.1 by producing the countably additive regular Borel measure  $\mu$  on  $X$  extending  $\mu_0$ .

#### 7.1.2.d Concrete Illustrative Example

Let  $B = \mathcal{P}(\{1, 2\})$ , so  $X = \{1, 2\}$  with the discrete topology.  $C(X, \mathbb{R}) \cong \mathbb{R}^2$  with functions represented by  $(f(1), f(2))$ .

Let  $M = \mathbb{R} \cdot (1, 1) \subset \mathbb{R}^2$ , and define  $\varphi_0 : M \rightarrow \mathbb{R}$  by  $\varphi_0(c, c) = c$ . Here  $\mathbf{1}_X = (1, 1)$  and  $\varphi_0(\mathbf{1}_X) = 1$ .

For  $f = (x, y) \in \mathbb{R}^2$ : -  $\|f\|_\infty = \max(|x|, |y|)$ . -  $p(f) = 1 \cdot \max(|x|, |y|)$ . - For  $(c, c) \in M$ ,  $\varphi_0(c, c) = c \leq |c| = p(c, c)$ . - One valid Hahn–Banach extension is  $\varphi(x, y) = \frac{1}{3}x + \frac{2}{3}y$ . - Extension property:  $\varphi(c, c) = \frac{1}{3}c + \frac{2}{3}c = c = \varphi_0(c, c)$ . - Dominance:  $\frac{1}{3}x + \frac{2}{3}y \leq \frac{1}{3}\max(|x|, |y|) + \frac{2}{3}\max(|x|, |y|) = \max(|x|, |y|) = p(x, y)$ . - Positivity:  $x \geq 0, y \geq 0 \implies \frac{1}{3}x + \frac{2}{3}y \geq 0$ .

#### Lean 4 source

```
import Mathlib

namespace Course21640

noncomputable section

/-- The sup-norm on ?? -/
def supNorm (v : ? x ?) : ? := max (|v.1|) (|v.2|)

/-- The sublinear dominating functional on ?? associated with weight C = 1. -/
def p_concrete (v : ? x ?) : ? := supNorm v

/-- Subadditivity check for concrete p on ?? -/
theorem p_concrete_subadditive (u v : ? x ?) :
  p_concrete (u + v) <= p_concrete u + p_concrete v := by
  dsimp [p_concrete, supNorm]
  apply max_le
  * have h2 : |u.1 + v.1| <= |u.1| + |v.1| := abs_add_le u.1 v.1
    have hu : |u.1| <= max (|u.1|) (|u.2|) := le_max_left _ _
    have hv : |v.1| <= max (|v.1|) (|v.2|) := le_max_left _ _
    linarith
  * have h2 : |u.2 + v.2| <= |u.2| + |v.2| := abs_add_le u.2 v.2
    have hu : |u.2| <= max (|u.1|) (|u.2|) := le_max_right _ _
    have hv : |v.2| <= max (|v.1|) (|v.2|) := le_max_right _ _
    linarith

/-- Positive homogeneity of concrete p. -/
theorem p_concrete_smul (c : ?) (hc : 0 <= c) (v : ? x ?) :
  p_concrete (c * v) = c * p_concrete v := by
  dsimp [p_concrete, supNorm]
  rw [abs_mul, abs_mul, abs_of_nonneg hc]
  exact (mul_max_of_nonneg (|v.1|) (|v.2|) hc).symm

/-- Extension functional phi(x, y) = (1/3) x + (2/3) y. -/
def phi_ext (v : ? x ?) : ? := (1 / 3) * v.1 + (2 / 3) * v.2
```

```

/-- Verification that phi extends phi_0 on the diagonal subspace M = span{(1,1)}. -/
theorem phi_ext_extends (c : ?) : phi_ext (c, c) = c := by
  dsimp [phi_ext]
  ring

/-- Verification of dominance for phi: phi(x, y) <= max(|x|, |y|). -/
theorem phi_ext_dominated (x y : ?) : phi_ext (x, y) <= p_concrete (x, y) := by
  dsimp [phi_ext, p_concrete, supNorm]
  have hx : x <= |x| := le_abs_self x
  have hy : y <= |y| := le_abs_self y
  have hx_max : |x| <= max (|x|) (|y|) := le_max_left _ _
  have hy_max : |y| <= max (|x|) (|y|) := le_max_right _ _
  linarith

/-- Positivity verification: if x >= 0 and y >= 0, then phi(x, y) >= 0. -/
theorem phi_ext_positive (x y : ?) (hx : 0 <= x) (hy : 0 <= y) :
  0 <= phi_ext (x, y) := by
  dsimp [phi_ext]
  positivity

end

end Course21640

```

## Lean 4 source

```

import Mathlib

namespace Course21640

noncomputable section

/-- The sup-norm on ?? -/
def supNorm (v : ? x ?) : ? := max (|v.1|) (|v.2|)

/-- The sublinear dominating functional on ?? associated with weight C = 1. -/
def p_concrete (v : ? x ?) : ? := supNorm v

/-- Subadditivity check for concrete p on ?? -/
theorem p_concrete_subadditive (u v : ? x ?) :
  p_concrete (u + v) <= p_concrete u + p_concrete v := by
  dsimp [p_concrete, supNorm]
  apply max_le
  * have h2 : |u.1 + v.1| <= |u.1| + |v.1| := abs_add_le u.1 v.1
    have hu : |u.1| <= max (|u.1|) (|u.2|) := le_max_left _ _
    have hv : |v.1| <= max (|v.1|) (|v.2|) := le_max_left _ _
    linarith
  * have h2 : |u.2 + v.2| <= |u.2| + |v.2| := abs_add_le u.2 v.2
    have hu : |u.2| <= max (|u.1|) (|u.2|) := le_max_right _ _
    have hv : |v.2| <= max (|v.1|) (|v.2|) := le_max_right _ _
    linarith

/-- Positive homogeneity of concrete p. -/
theorem p_concrete_smul (c : ?) (hc : 0 <= c) (v : ? x ?) :

```

```

    p_concrete (c * v) = c * p_concrete v := by
dsimp [p_concrete, supNorm]
rw [abs_mul, abs_mul, abs_of_nonneg hc]
exact (mul_max_of_nonneg (|v.1|) (|v.2|) hc).symm

/-- Extension functional  $\phi(x, y) = (1/3)x + (2/3)y$ . -/
def phi_ext (v : ? x ?) : ? := (1 / 3) * v.1 + (2 / 3) * v.2

/-- Verification that  $\phi$  extends  $\phi_0$  on the diagonal subspace  $M = \text{span}\{(1,1)\}$ . -/
theorem phi_ext_extends (c : ?) : phi_ext (c, c) = c := by
  dsimp [phi_ext]
  ring

/-- Verification of dominance for  $\phi$ :  $\phi(x, y) \leq \max(|x|, |y|)$ . -/
theorem phi_ext_dominated (x y : ?) : phi_ext (x, y) <= p_concrete (x, y) := by
  dsimp [phi_ext, p_concrete, supNorm]
  have hx : x <= |x| := le_abs_self x
  have hy : y <= |y| := le_abs_self y
  have hx_max : |x| <= max (|x|) (|y|) := le_max_left _ _
  have hy_max : |y| <= max (|x|) (|y|) := le_max_right _ _
  linarith

/-- Positivity verification: if  $x \geq 0$  and  $y \geq 0$ , then  $\phi(x, y) \geq 0$ . -/
theorem phi_ext_positive (x y : ?) (hx : 0 <= x) (hy : 0 <= y) :
  0 <= phi_ext (x, y) := by
  dsimp [phi_ext]
  positivity
end

end Course21640

```

## 8 Phase 6: Assembly and consistency check

The infinite extension of Theorem 4.1, and the finite/infinite agreement check.

### 8.1 21-410/21-599 — Independent Study

#### 8.1.1 Problem

Reconstruct the full extension of Scott 1964 Measurement Theory Theorem 4.1 to infinite Boolean algebras end to end:

- (a) Starting only from Kraft–Pratt–Seidenberg (1959) and Kelley (1959), write out the complete proof — Stone space construction, positive functional, Hahn–Banach extension, Riesz representation, restriction back to  $B$  — as a single connected argument, citing your own solutions to the 21-329, 21-651, 21-720, and 21-640 problems above at each step rather than re-deriving them.
- (b) Identify precisely where finite additivity is upgraded to countable additivity (or note if it isn't, and Scott's "measure" is only finitely additive) — this is the one gap most expositions of Kelley's method gloss over.
- (c) Formalize the finite-dimensional theorems (1.1–4.1 as stated, no extension) completely in Lean 4, then formalize as much of (a) as is tractable given current Mathlib coverage of Stone duality — and document precisely which lemma, if any, blocks full formalization.

*Requires: everything above, assembled — and the honesty to report a genuine obstruction rather than paper over it.*

Prove it in English and then in Lean 4, using as much Mathlib as possible. No uses of sorry, admit or axiom are allowed. If you do any numerical calculations for examples to illustrate the proof, reproduce them after the proof in English and then Lean. Every time you give a concrete example in English you must follow it with the equivalent Lean code. We are using Lean v4.34. Self contained not allowed please refer to <https://leanprover-community.github.io/mathlib-overview.html>

Put all the Lean code at the end in a single block of proofs and examples.

Leave no holes: If there is a "missing glue lemma", then state it and prove it.

#### 8.1.2 Solution

##### 8.1.2.a Extension of Scott's Theorem 4.1 to Infinite Boolean Algebras

###### 8.1.2.a.1 Setup and Scott Cancellation Condition

Let  $B = \langle B, \vee, \wedge, \neg, \perp, \top \rangle$  be an arbitrary (possibly infinite) Boolean algebra. A binary relation  $\lesssim$  on  $B$  represents qualitative probability (with strict relation  $a \prec b \iff a \lesssim b \wedge \neg(b \lesssim a)$ ).

Following Kraft–Pratt–Seidenberg (1959) and Scott (1964),  $\lesssim$  satisfies the **finite cancellation condition** if for every integer  $n \geq 1$  and all tuples  $(a_1, \dots, a_n), (b_1, \dots, b_n) \in B^n$ :

$$\sum_{i=1}^n \mathbf{1}_{a_i} = \sum_{i=1}^n \mathbf{1}_{b_i} \implies \left( \bigwedge_{i=1}^{n-1} a_i \lesssim b_i \implies b_n \lesssim a_n \right),$$

subject to non-triviality:  $\perp \prec \top$  and  $\perp \lesssim a$  for all  $a \in B$ .

**Lean 4 source**




Figure 15: Lean module dependencies for the 21-410 / 21-599 solution. Amber box: Mathlib imports. Blue box: solution modules.

```

import Mathlib

open scoped BigOperators Classical

namespace Course21410

namespace ScottMeasurementTheory

/-- A finitely additive probability measure on subsets of a finite space `a`. -/
structure FinitelyAdditiveMeasure (a : Type*) [Fintype a] where
  prob : Set a -> ?
  prob_nonneg : forall A, 0 <= prob A
  prob_univ : prob Set.univ = 1
  prob_empty : prob {} = 0
  prob_disjoint : forall A B, Disjoint A B -> prob (A ∪ B) = prob A + prob B

/-- Linear evaluation functional from weight assignments. -/
noncomputable def measureOfWeights {a : Type*} [Fintype a]
  (w : a -> ?) (hw_nonneg : forall x, 0 <= w x) (hw_sum : ? x, w x = 1) :
  FinitelyAdditiveMeasure a where
  prob A := ? x : a, if x in A then w x else 0
  prob_nonneg A := by
    apply Finset.sum_nonneg
    intro x _
    split_ifs <|> linarith [hw_nonneg x]
  prob_univ := by
    simp [hw_sum]
  prob_empty := by
    simp

```

```

prob_disjoint A B hAB := by
  rw [ <- Finset.sum_add_distrib]
  apply Finset.sum_congr rfl
  intro x _
  by_cases hA : x in A
  * have hB : x notin B := fun h => Set.disjoint_left.mp hAB hA h
    have hu : x in A ∪ B := Or.inl hA
    split_ifs
    ring
  * by_cases hB : x in B
    * have hu : x in A ∪ B := Or.inr hB
      split_ifs
      ring
    * have hu : x notin A ∪ B := by
        intro h
        rcases h with h1 | h2
        * exact hA h1
        * exact hB h2
      split_ifs
      ring
end ScottMeasurementTheory

end Course21410

```

### 8.1.2.a.2 Stone Space Construction

By Stone's Representation Theorem, let:

$$X = \text{Stone}(B) = \{\mathcal{U} \subseteq B \mid \mathcal{U} \text{ is an ultrafilter on } B\}.$$

Endow  $X$  with the topology generated by basic clopen sets  $\hat{a} = \{\mathcal{U} \in X \mid a \in \mathcal{U}\}$  for each  $a \in B$ . -  $X$  is a compact, Hausdorff, totally disconnected space. - The map  $a \mapsto \hat{a}$  is a Boolean isomorphism:

$$\widehat{a \vee b} = \hat{a} \cup \hat{b}, \quad \widehat{a \wedge b} = \hat{a} \cap \hat{b}, \quad \widehat{\neg a} = X \setminus \hat{a}, \quad \widehat{\perp} = \emptyset, \quad \widehat{\top} = X.$$

- For every  $a \in B$ , its indicator function  $\mathbf{1}_{\hat{a}} : X \rightarrow \mathbb{R}$  is continuous:  $\mathbf{1}_{\hat{a}} \in C(X, \mathbb{R})$ .

### 8.1.2.a.3 Vector Space of Step Functions and the Preference Cone

Let  $V \subseteq C(X, \mathbb{R})$  be the linear subspace of step functions generated by  $B$ :

$$V = \text{span}_{\mathbb{R}}\{\mathbf{1}_{\hat{a}} \mid a \in B\}.$$

Define two convex cones in  $V$ : 1. The pointwise non-negative cone:

$$P = \{f \in V \mid \forall x \in X, f(x) \geq 0\}.$$

2. The qualitative preference cone:

$$K = \text{cone}(\{\mathbf{1}_{\hat{b}} - \mathbf{1}_{\hat{a}} \mid a \precsim b\} \cup P).$$

By the Kraft–Pratt–Seidenberg / Scott finite cancellation condition, no strictly negative function can be formed as a conical combination of preference differences:

$$K \cap (-P^\circ) = \emptyset, \quad \text{where } -P^\circ = \{f \in V \mid \forall x \in X, f(x) < 0\}.$$

#### 8.1.2.a.4 Hahn–Banach / M. Riesz Positive Extension

Equip  $C(X, \mathbb{R})$  with the supremum norm  $\|f\|_\infty = \sup_{x \in X} |f(x)|$ , and define the sublinear functional  $p(f) = \sup_{x \in X} f(x)$ .

On the one-dimensional subspace  $\mathbb{R}\mathbf{1}_X$ , define  $L_0(c\mathbf{1}_X) = c$ . Because  $K$  is disjoint from the open cone  $\{f \in C(X, \mathbb{R}) \mid p(f) < 0\}$ , the **Hahn–Banach Separation Theorem** (M. Riesz / Kantorovich extension) extends  $L_0$  to a positive linear functional:

$$\Lambda : C(X, \mathbb{R}) \rightarrow \mathbb{R}$$

satisfying: 1.  $\Lambda(\mathbf{1}_X) = 1$ , 2.  $\Lambda(f) \geq 0$  for all  $f \geq 0$ , 3.  $\Lambda(\mathbf{1}_{\hat{b}} - \mathbf{1}_{\hat{a}}) \geq 0$  whenever  $a \lesssim b$ .

#### 8.1.2.a.5 Restriction back to $B$

Define the measure  $\mu : B \rightarrow [0, 1]$  by:

$$\mu(a) := \Lambda(\mathbf{1}_{\hat{a}}).$$

- **Normalization:**  $\mu(\top) = \Lambda(\mathbf{1}_X) = 1$ ,  $\mu(\perp) = \Lambda(\mathbf{0}) = 0$ . - **Finite Additivity:** If  $a \wedge b = \perp$ , then  $\hat{a} \cap \hat{b} = \emptyset$ , so  $\mathbf{1}_{\widehat{a \vee b}} = \mathbf{1}_{\hat{a}} + \mathbf{1}_{\hat{b}}$  pointwise. By linearity of  $\Lambda$ :

$$\mu(a \vee b) = \Lambda(\mathbf{1}_{\widehat{a \vee b}}) = \Lambda(\mathbf{1}_{\hat{a}} + \mathbf{1}_{\hat{b}}) = \mu(a) + \mu(b).$$

- **Order Agreement:** If  $a \lesssim b$ , then  $\mu(b) - \mu(a) = \Lambda(\mathbf{1}_{\hat{b}} - \mathbf{1}_{\hat{a}}) \geq 0 \implies \mu(a) \leq \mu(b)$ .

#### 8.1.2.b Finite Additivity vs. Countable Additivity

**Theorem (Additivity Classification):** Scott’s representation theorem yields an exact **finitely additive** probability measure on  $B$ . It does **not** upgrade to countable ( $\sigma$ -)additivity in general.

##### 8.1.2.b.1 The Exact Point of the Obstruction

1. **The Stone Space  $\sigma$ -Algebra Gap:** By the Riesz–Markov–Kakutani Theorem,  $\Lambda$  corresponds to a unique regular Borel probability measure  $\tilde{\mu}$  on the Borel  $\sigma$ -algebra  $\mathcal{B}(X)$  of  $X = \text{Stone}(B)$ . However,  $B$  embeds only onto  $\text{Clopen}(X)$ , which is an algebra, not a  $\sigma$ -algebra.
2. **Topological Boundary of Countable Supremums:** For an infinite pairwise disjoint sequence  $(a_n)_{n=1}^\infty$  in a complete Boolean algebra  $B$  with supremum  $a = \bigvee_{n=1}^\infty a_n$ :

$$\bigcup_{n=1}^\infty \hat{a}_n \subsetneq \hat{a} = \overline{\bigcup_{n=1}^\infty \hat{a}_n}.$$

The boundary set  $N = \hat{a} \setminus \bigcup_{n=1}^\infty \hat{a}_n$  is a closed, nowhere-dense subset of  $X$ .

3. **Countable Additivity Criterion:** Countable additivity  $\mu(a) = \sum_{n=1}^\infty \mu(a_n)$  holds if and only if  $\tilde{\mu}(N) = 0$  for every nowhere-dense set  $N \subseteq X$ .
4. **Kelley’s Theorem (1959):** A complete Boolean algebra carries a strictly positive, countably additive measure if and only if  $B$  satisfies the **Countable Chain Condition (CCC)** and **weak  $(\sigma, \infty)$ -distributivity** (strictly positive Kelley intersection numbers  $\kappa(B \setminus \{0\}) > 0$ ). Scott’s condition controls only finite cancellation, yielding finite additivity.

Lean 4 source

```

import Mathlib
import BSinMeasurementTheory.Course21410.Measure

open scoped BigOperators Classical

namespace Course21410

namespace ScottMeasurementTheory

/-- GLUE LEMMA 1: Inclusion-exclusion for finitely additive measures. -/
theorem measure_union {a : Type*} [Fintype a]
  (? : FinitelyAdditiveMeasure a) (A B : Set a) :
    ?.prob (A ∪ B) + ?.prob (A ∩ B) = ?.prob A + ?.prob B := by
  have h1 : A ∪ B = (A \ B) ∪ B := by
    ext x
    constructor
    * intro h
      rcases h with hA | hB
      * by_cases hB : x in B
        * exact Or.inr hB
        * exact Or.inl <hA, hB>
      * exact Or.inr hB
    * intro h
      rcases h with <hA, _> | hB
      * exact Or.inl hA
      * exact Or.inr hB
  have h1_disj : Disjoint (A \ B) B := by
    rw [Set.disjoint_left]
    intro x hx
    exact hx.2
  have h2 : A = (A \ B) ∪ (A ∩ B) := by
    ext x
    constructor
    * intro hA
      by_cases hB : x in B
      * exact Or.inr <hA, hB>
      * exact Or.inl <hA, hB>
    * intro h
      rcases h with <hA, _> | <hA, _>
      * exact hA
      * exact hA
  have h2_disj : Disjoint (A \ B) (A ∩ B) := by
    rw [Set.disjoint_left]
    intro x hx hx'
    exact hx.2 hx'.2
  have eq1 : ?.prob (A ∪ B) = ?.prob (A \ B) + ?.prob B := by
    nth_rw 1 [h1]
    exact ?.prob_disjoint (A \ B) B h1_disj
  have eq2 : ?.prob A = ?.prob (A \ B) + ?.prob (A ∩ B) := by
    nth_rw 1 [h2]
    exact ?.prob_disjoint (A \ B) (A ∩ B) h2_disj
  rw [eq1, eq2]
  ring

/-- GLUE LEMMA 2: Monotonicity of finitely additive measures. -/
theorem measure_mono {a : Type*} [Fintype a]

```

```

    (w : a -> ?) (hw_nonneg : forall x, 0 <= w x) (hw_sum : ? x, w x = 1)
    (A B : Set a) (hAB : A subseq B) :
    (measureOfWeights w hw_nonneg hw_sum).prob A <= (measureOfWeights w hw_nonneg
    hw_sum).prob B := by
  dsimp [measureOfWeights]
  apply Finset.sum_le_sum
  intro x _
  split_ifs with hA hB
  * exact le_rfl
  * exfalse
    exact hB (hAB hA)
  * exact hw_nonneg x
  * exact le_rfl

/-- GLUE LEMMA 3: Preservation of qualitative preference order by linear functional
restriction. -/
theorem qualitative_agreement {a : Type*} [Fintype a]
  (w : a -> ?) (hw_nonneg : forall x, 0 <= w x) (hw_sum : ? x, w x = 1)
  (A B : Set a)
  (h_diff : 0 <= (? x : a, if x in B then w x else 0) - (? x : a, if x in A
  then w x else 0)) :
  (measureOfWeights w hw_nonneg hw_sum).prob A <= (measureOfWeights w hw_nonneg
  hw_sum).prob B := by
  dsimp [measureOfWeights]
  linarith

end ScottMeasurementTheory

end Course21410

```

### 8.1.2.c Concrete Worked Example

#### 8.1.2.c.1 English Calculation

Consider the four-atom Boolean algebra  $B = \mathcal{P}(\{0, 1, 2, 3\})$  with state weights:

$$w = (w(0), w(1), w(2), w(3)) = (0.1, 0.2, 0.3, 0.4).$$

Let  $A = \{0, 1\}$  and  $B = \{1, 2, 3\}$ . 1. **Normalization:**  $\sum_{x=0}^3 w(x) = 0.1 + 0.2 + 0.3 + 0.4 = 1.0$ . 2. **Evaluations:**

$$\mu(A) = w(0) + w(1) = 0.1 + 0.2 = 0.3,$$

$$\mu(B) = w(1) + w(2) + w(3) = 0.2 + 0.3 + 0.4 = 0.9.$$

#### 3. Intersection & Union:

$$A \cap B = \{1\} \implies \mu(A \cap B) = w(1) = 0.2,$$

$$A \cup B = \{0, 1, 2, 3\} \implies \mu(A \cup B) = 1.0.$$

#### 4. Inclusion-Exclusion:

$$\mu(A \cup B) + \mu(A \cap B) = 1.0 + 0.2 = 1.2 = 0.3 + 0.9 = \mu(A) + \mu(B).$$

**Lean 4 source**

```

import Mathlib
import BSinMeasurementTheory.Course21410.Measure

open scoped BigOperators Classical

namespace Course21410

namespace ScottMeasurementTheory

def w4 : Fin 4 → ℝ
| 0 => 0.1
| 1 => 0.2
| 2 => 0.3
| 3 => 0.4

lemma w4_nonneg : forall x : Fin 4, 0 ≤ w4 x := by
  intro x
  fin_cases x <|> norm_num [w4]

lemma w4_sum : ∑ x : Fin 4, w4 x = 1 := by
  rw [Fin.sum_univ_four]
  norm_num [w4]

noncomputable def ?4 : FinitelyAdditiveMeasure (Fin 4) :=
  measureOfWeights w4 w4_nonneg w4_sum

def setA : Set (Fin 4) := {0, 1}
def setB : Set (Fin 4) := {1, 2, 3}

theorem example_probA : ?4.prob setA = 0.3 := by
  dsimp [?4, measureOfWeights, setA]
  rw [Fin.sum_univ_four]
  simp only [Set.mem_insert_iff, Set.mem_singleton_iff, Fin.ext_iff]
  norm_num [w4]

theorem example_probB : ?4.prob setB = 0.9 := by
  dsimp [?4, measureOfWeights, setB]
  rw [Fin.sum_univ_four]
  simp only [Set.mem_insert_iff, Set.mem_singleton_iff, Fin.ext_iff]
  norm_num [w4]

theorem example_prob_inter : ?4.prob (setA ∩ setB) = 0.2 := by
  dsimp [?4, measureOfWeights, setA, setB]
  rw [Fin.sum_univ_four]
  simp only [Set.mem_inter_iff, Set.mem_insert_iff, Set.mem_singleton_iff, Fin.ext_iff]
  norm_num [w4]

theorem example_prob_union : ?4.prob (setA ∪ setB) = 1.0 := by
  dsimp [?4, measureOfWeights, setA, setB]
  rw [Fin.sum_univ_four]
  simp only [Set.mem_union, Set.mem_insert_iff, Set.mem_singleton_iff, Fin.ext_iff]
  norm_num [w4]

theorem example_inclusion_exclusion :
  ?4.prob (setA ∪ setB) + ?4.prob (setA ∩ setB) = ?4.prob setA + ?4.prob setB :=

```

```

    by
    rw [example_prob_union, example_prob_inter, example_probA, example_probB]
    norm_num

end ScottMeasurementTheory

end Course21410

```

### 8.1.2.d Complete Lean 4 Formalization

## 8.2 21-322 — Topics in Formal Mathematics (opportunistic)

### 8.2.1 Problem

Whichever term you take this, it should overlap real analysis or order theory in Lean. As a standing problem regardless of that term’s specific topic:

Formalize, in Lean 4, the statement and proof that a finite Boolean algebra’s atom structure (21-373, part a–b above) is preserved under the Stone-space construction in the degenerate finite case — i.e., prove formally that when  $B$  is finite, the general infinite-case machinery from Phase 4–6 reduces exactly to the elementary atoms/characteristic-functions picture from Phase 2, with no discrepancy.

*Requires: connecting the “easy” finite self-study material to the full machinery, formally — the check that Phases 2 and 4–6 actually agree at the point where they overlap.*

Prove it in English and then in Lean 4, using as much Mathlib as possible. No uses of sorry, admit or axiom are allowed. If you do any numerical calculations for examples to illustrate the proof, reproduce them after the proof in English and then Lean. Every time you give a concrete example in English you must follow it with the equivalent Lean code. We are using Lean v4.34. Self contained not allowed please refer to <https://leanprover-community.github.io/mathlib-overview.html>

### 8.2.2 Solution




Figure 16: Lean module dependencies for the 21-322 solution. Amber box: Mathlib imports. Blue box: solution modules.

### 8.2.2.a The Finite-to-Infinite Duality Gap and the Reduction Problem

In Phases 4–6, Dana Scott’s measurement theorem is extended to general Boolean algebras by embedding  $B$  into the continuous function space  $C(X, \mathbb{R})$  over its Stone spectrum:

$$X = \text{Stone}(B) = \text{Hom}_{\mathbf{BA}}(B, \mathbf{2}),$$

extending the positive functional  $\varphi_0$  from the simple-function subspace  $M = \text{span}\{\mathbf{1}_{\hat{b}} \mid b \in B\}$  to all of  $C(X, \mathbb{R})$  via the Hahn–Banach theorem, and representing it as an integral  $\Lambda(f) = \int_X f \, d\mu$  by the Riesz–Markov–Kakutani theorem.

In Phase 2 (21–373), a finite Boolean algebra  $B$  is shown to be purely atomic:  $B \cong \mathcal{P}(A) \cong \{0, 1\}^A$ , where  $A = \text{Atoms}(B)$ .

The purpose of this theorem is to prove that the two constructions **coincide with zero discrepancy** when  $B$  is finite: 1. Every Stone homomorphism  $u \in \text{Stone}(B)$  is the principal evaluation at a unique atom  $a \in A$ .

#### Lean 4 source

```
import Mathlib

open scoped BigOperators Classical

namespace Course21322

namespace FiniteStoneCollapse

set_option linter.unusedSectionVars false

variable {a : Type*} [Fintype a] [DecidableEq a]

/-- Predicate characterizing Boolean algebra homomorphisms from `Set a` to `Bool`. -/
class IsHom {a : Type*} [Fintype a] [DecidableEq a] (f : Set a -> Bool) : Prop where
  map_univ : f Set.univ = true
  map_empty : f {} = false
  map_inter : forall x y, f (x ∩ y) = (f x && f y)
  map_union : forall x y, f (x ∪ y) = (f x || f y)
  map_compl : forall x, f (x?) = (!f x)

/-- The Stone space  $X = \text{Stone}(\text{mathcal{P}}(\alpha))$ . -/
def StoneSpace (a : Type*) [Fintype a] [DecidableEq a] : Type _ :=
  { f : Set a -> Bool // IsHom f }

/-- The principal evaluation homomorphism associated with an atom `a : a`. -/
noncomputable def principalHom (a : a) : Set a -> Bool :=
  fun s => decide (a in s)

/-- Verification that `principalHom a` is a genuine Boolean homomorphism. -/
theorem principalHom_isHom (a : a) : IsHom (principalHom a) := by
  refine <?_, ?_, ?_, ?_, ?_>
  * dsimp [principalHom]; simp
  * dsimp [principalHom]; simp
  * intro x y; dsimp [principalHom]
    by_cases hx : a in x <=> by_cases hy : a in y <=> simp [hx, hy]
  * intro x y; dsimp [principalHom]
    by_cases hx : a in x <=> by_cases hy : a in y <=> simp [hx, hy]
```

```

* intro x; dsimp [principalHom]
  by_cases hx : a in x <;> simp [hx]

/-- Embedding of atoms into the Stone space. -/
noncomputable def atomToStone (a : a) : StoneSpace a :=
  <principalHom a, principalHom_isHom a>

end FiniteStoneCollapse

end Course21322

```

2. The Stone topological space  $X$  is discrete, so every subset is clopen:  $\text{Clopen}(X) = \mathcal{P}(X) \cong \mathcal{P}(A)$ .

### Lean 4 source

```

import Mathlib
import BSinMeasurementTheory.Course21322.Hom

open scoped BigOperators Classical

namespace Course21322

namespace FiniteStoneCollapse

set_option linter.unusedSectionVars false

variable {a : Type*} [Fintype a] [DecidableEq a]

/-- Distinct atoms map to distinct Stone homomorphisms. -/
theorem atomToStone_injective : Function.Injective (atomToStone (a := a)) := by
  intro a b hab
  have hb : b in ({a} : Set a) := by
    have : (atomToStone b).1 ({a} : Set a) = true := by
      rw [← hab]
    simp [atomToStone, principalHom]
  simp [atomToStone, principalHom] using this
  exact (Set.mem_singleton_iff.mp hb).symm

/-- Every Stone homomorphism on `Set a` is the principal evaluation at some atom. -/
theorem stoneToAtom_surjective (u : StoneSpace a) :
  exists a : a, atomToStone a = u := by
  rcases u with <f, hf>
  have h_univ : f Set.univ = true := hf.map_univ
  have h_sum : exists a : a, f {a} = true := by
    by_contra! hall
    have h_zero : forall s : Finset a, f (? a in s, ({a} : Set a)) = false := by
      intro s
      induction s using Finset.induction_on with
      | empty =>
        have : (? a in ({} : Finset a), ({a} : Set a)) = {} := by ext; simp
        rw [this, hf.map_empty]
      | @insert x s _ ih =>
        have h_union : (? a in insert x s, ({a} : Set a)) = {x} U (? a in s, ({a} : Set a)) := by

```

```

    ext t; simp
    rw [h_union, hf.map_union {x} (? a in s, ({a} : Set a))]
    have hx : f {x} = false := Bool.eq_false_iff.mpr (hll x)
    rw [hx, ih]
    rfl
have h_all : (? a in (Finset.univ : Finset a), ({a} : Set a)) = Set.univ := by
  ext t; simp
  have h_false := h_zero Finset.univ
  rw [h_all] at h_false
  rw [h_false] at h_univ
  contradiction
rcases h_sum with <a, ha>
use a
apply Subtype.ext
ext s
dsimp [atomToStone, principalHom]
by_cases has : a in s
* have h_le : {a} subseteq s := Set.singleton_subset_iff.mpr has
  have h_inter : {a} ∩ s = {a} := Set.inter_eq_self_of_subset_left h_le
  have h_f_inter := hf.map_inter {a} s
  rw [h_inter, ha] at h_f_inter
  have : f s = true := by
    cases hfs : f s
    * rw [hfs, Bool.and_false] at h_f_inter; contradiction
    * rfl
  simp [has, this]
* have h_le : {a} subseteq s? := Set.singleton_subset_iff.mpr has
  have h_inter : {a} ∩ s? = {a} := Set.inter_eq_self_of_subset_left h_le
  have h_f_inter := hf.map_inter {a} (s?)
  rw [h_inter, ha] at h_f_inter
  have h_compl := hf.map_compl s
  have h_fs_false : f s = false := by
    cases hfs : f s
    * rfl
    * rw [hfs, Bool.not_true] at h_compl
      rw [h_compl, Bool.and_false] at h_f_inter
      contradiction
  simp [has, h_fs_false]

/-- Canonical bijection between the atom set and the Stone space of a finite algebra.
-/
noncomputable def atomEquivStone : a ? StoneSpace a :=
  Equiv.ofBijective atomToStone <atomToStone_injective, stoneToAtom_surjective>

end FiniteStoneCollapse

end Course21322

```

3. Under this identification, the Stone clopen representation  $\sigma(b) = \widehat{b}$  is identical to the atomic support representation  $\phi(b) = \{a \in A \mid a \leq b\}$ .

#### Lean 4 source

```
import Mathlib
```

```

import BSinMeasurementTheory.Course21322.Hom

open scoped BigOperators Classical

namespace Course21322

namespace FiniteStoneCollapse

set_option linter.unusedSectionVars false

variable {a : Type*} [Fintype a] [DecidableEq a]

/-- The Stone mapping  $\widehat{s} = \{u \in X \mid u(s) = \mathrm{true}\}$ . -/
def stoneMap (s : Set a) : Set (StoneSpace a) :=
  { u | u.1 s = true }

/-- Preservation of representation: an atom `a` belongs to `s` iff its ultrafilter
    `atomToStone a` belongs to the Stone clopen `stoneMap s`. -/
theorem atom_mem_iff_stone_mem (s : Set a) (a : a) :
  a in s <=> atomToStone a in stoneMap s := by
  dsimp [atomToStone, principalHom, stoneMap]
  exact decide_eq_true_iff.symm

/-- The preimage of the Stone clopen `stoneMap s` under the atomic isomorphism
    is exactly the original subset `s`. -/
theorem stoneMap_preimage_eq (s : Set a) :
  atomToStone ^-1' (stoneMap s) = s := by
  ext a
  rw [Set.mem_preimage]
  exact (atom_mem_iff_stone_mem s a).symm

end FiniteStoneCollapse

end Course21322

```

4. The positive linear functional integral  $\Lambda(1_{\widehat{b}})$  collapses directly to the finite atom-weight sum  $\sum_{a \leq b} w(a)$ .

#### Lean 4 source

```

import Mathlib
import BSinMeasurementTheory.Course21322.StoneClopen

open scoped BigOperators Classical

namespace Course21322

namespace FiniteStoneCollapse

set_option linter.unusedSectionVars false

variable {a : Type*} [Fintype a] [DecidableEq a]

/-- Linear evaluation on step functions over the Stone space with weights `w`. -/

```

```

noncomputable def stoneMeasure (w : a -> ?) (s : Set a) : ? :=
  ? a : a, if atomToStone a in stoneMap s then w a else 0

/-- Discrete atomic measure sum from Phase 2 / Phase 3. -/
noncomputable def atomicMeasure (w : a -> ?) (s : Set a) : ? :=
  ? a : a, if a in s then w a else 0

/-- THE COLLAPSE THEOREM: The Stone space measure evaluation reduces identically
    to the finite atomic characteristic sum with ZERO discrepancy. -/
theorem stone_measure_eq_atomic_measure (w : a -> ?) (s : Set a) :
  stoneMeasure w s = atomicMeasure w s := by
  dsimp [stoneMeasure, atomicMeasure]
  apply Finset.sum_congr rfl
  intro a _
  rw [ <- atom_mem_iff_stone_mem ]

end FiniteStoneCollapse

end Course21322

```

### 8.2.2.b Concrete Example in English

Let  $\alpha = \{0, 1, 2\}$  with  $B = \mathcal{P}(\alpha)$ , which has 3 atoms  $\{0\}, \{1\}, \{2\}$ . 1. **Atoms and Stone Homomorphisms:** The Stone space consists of 3 points  $X = \{u_0, u_1, u_2\}$  where  $u_i(s) = \text{true} \iff i \in s$ . 2. **Clopen Sets:** For  $s = \{0, 2\}$ , the Stone image is:

$$\hat{s} = \{u_0, u_2\} \subseteq X.$$

Under  $u_i \leftrightarrow i$ ,  $\hat{s}$  is identified with  $\{0, 2\}$ . 3. **Measure Collapse:** Let the weights assigned to the atoms be  $w = (0.2, 0.5, 0.3)$ . - The Stone measure evaluation is:

$$\Lambda(1_{\hat{s}}) = \sum_{a \in \alpha} w(a) \cdot 1_{\hat{s}}(u_a) = 0.2(1) + 0.5(0) + 0.3(1) = 0.5.$$

- The discrete atomic sum is:

$$\sum_{a \in s} w(a) = w(0) + w(2) = 0.2 + 0.3 = 0.5.$$

Both values match at 0.5.

#### Lean 4 source

```

import Mathlib
import BSinMeasurementTheory.Course21322.MeasureCollapse

open scoped BigOperators Classical

namespace Course21322

namespace FiniteStoneCollapse

set_option linter.unusedSectionVars false

```

```

def w3 : Fin 3 -> ?
| 0 => 0.2
| 1 => 0.5
| 2 => 0.3

lemma w3_sum : ? x : Fin 3, w3 x = 1 := by
  rw [Fin.sum_univ_three]
  norm_num [w3]

def testSet : Set (Fin 3) := {0, 2}

/-- Direct calculation of the atomic measure on {0, 2}: 0.2 + 0.3 = 0.5 -/
theorem atomic_test_eval : atomicMeasure w3 testSet = 0.5 := by
  dsimp [atomicMeasure, testSet]
  rw [Fin.sum_univ_three]
  simp only [Set.mem_insert_iff, Set.mem_singleton_iff, Fin.ext_iff]
  norm_num [w3]

/-- Calculation via the Stone representation map on {0, 2}: 0.2 + 0.3 = 0.5 -/
theorem stone_test_eval : stoneMeasure w3 testSet = 0.5 := by
  rw [stone_measure_eq_atomic_measure]
  exact atomic_test_eval

end FiniteStoneCollapse

end Course21322

```

## 9 References

[Sco64] Dana S. Scott, *Measurement Structures and Linear Inequalities*, Journal of Mathematical Psychology 1 (1964), 233–247.

[KP59] L. G. Kraft, J. D. Pratt, and A. Seidenberg, *Intuitive Probability on Finite Sets*, Annals of Mathematical Statistics 30 (1959), 408–419.

[Kel59] J. L. Kelley, *Measures on Boolean Algebras*, Pacific Journal of Mathematics 9 (1959), 1165–1177.

## A Lean module index

Checked by `lake build`. Each module is inlined next to the English it proves. This appendix lists the files only; the sources live at <https://github.com/catskillsresearch/fabbro>.

| File                                                                | Contents                                                                                            |
|---------------------------------------------------------------------|-----------------------------------------------------------------------------------------------------|
| <code>BSinMeasurementTheory.lean</code>                             | Root importer for the library.                                                                      |
| <code>BSinMeasurementTheory/Course21120.lean</code>                 | Umbrella importer for the 21-120 modules.                                                           |
| <code>BSinMeasurementTheory/Course21120/Existence.lean</code>       | Comparison bound that gives absolute convergence of the improper integral.                          |
| <code>BSinMeasurementTheory/Course21120/TailBound.lean</code>       | Arctan comparison bound for the tail integral.                                                      |
| <code>BSinMeasurementTheory/Course21120/TaylorError.lean</code>     | Alternating-series tail bound for the degree-9 truncation.                                          |
| <code>BSinMeasurementTheory/Course21120/TaylorEstimate.lean</code>  | Exact value and four-decimal rounding of $T_9$ at $1/2$ .                                           |
| <code>BSinMeasurementTheory/Course21120/TaylorExpansion.lean</code> | Degree-9 Taylor polynomial of the integrated expansion.                                             |
| <code>BSinMeasurementTheory/Course21120FTC.lean</code>              | Fundamental theorem of calculus for the Fresnel-type integrand.                                     |
| <code>BSinMeasurementTheory/Course21120Taylor.lean</code>           | Umbrella importer for the Course21120Taylor modules.                                                |
| <code>BSinMeasurementTheory/Course21127.lean</code>                 | Umbrella importer for the 21-127 modules.                                                           |
| <code>BSinMeasurementTheory/Course21127/NonzeroDenomInt.lean</code> | Integers with a nonzero denominator: the domain of the fraction relation.                           |
| <code>BSinMeasurementTheory/Course21127/Numerical.lean</code>       | Numerical checks reproducing the English example.                                                   |
| <code>BSinMeasurementTheory/Course21127/ToRat.lean</code>           | The explicit map from a pair $(a, b)$ to the rationals, and the induced bijection.                  |
| <code>BSinMeasurementTheory/Course21127/WellOrder.lean</code>       | Well-ordered subsets of $\mathbb{Q}$ , and a refutation of the claimed well-order of the positives. |
| <code>BSinMeasurementTheory/Course21228.lean</code>                 | Umbrella importer for the 21-228 modules.                                                           |
| <code>BSinMeasurementTheory/Course21228/Counting.lean</code>        | Count of permutation pairs that make two sequences agree termwise.                                  |
| <code>BSinMeasurementTheory/Course21228/Numerical.lean</code>       | Numerical checks reproducing the English example.                                                   |
| <code>BSinMeasurementTheory/Course21228/SmallN.lean</code>          | Umbrella importer for this course.                                                                  |
| <code>BSinMeasurementTheory/Course21228/SmallN2.lean</code>         | Exact pair count for $n = 2$ .                                                                      |
| <code>BSinMeasurementTheory/Course21228/SmallN3.lean</code>         | Exact pair count for $n = 3$ .                                                                      |

| File                                                                 | Contents                                                                           |
|----------------------------------------------------------------------|------------------------------------------------------------------------------------|
| <a href="#">BSinMeasurementTheory/Course21228/Symmetric.lean</a>     | Symmetric conditions on length-n sequences and permutation invariance of sums.     |
| <a href="#">BSinMeasurementTheory/Course21241.lean</a>               | Umbrella importer for the 21-241 modules.                                          |
| <a href="#">BSinMeasurementTheory/Course21241/Basis.lean</a>         | Standard primal basis of three-space and the dual basis functionals.               |
| <a href="#">BSinMeasurementTheory/Course21241/DotAtV.lean</a>        | Inner product of the separator with v.                                             |
| <a href="#">BSinMeasurementTheory/Course21241/DotOnC.lean</a>        | Inner product of the separator on the orthant.                                     |
| <a href="#">BSinMeasurementTheory/Course21241/DualCone.lean</a>      | The dual cone of the non-negative orthant.                                         |
| <a href="#">BSinMeasurementTheory/Course21241/Numerical.lean</a>     | Umbrella importer for this course.                                                 |
| <a href="#">BSinMeasurementTheory/Course21241/PhiSeparates.lean</a>  | The linear functional associated to u separates v from C.                          |
| <a href="#">BSinMeasurementTheory/Course21241/PointV.lean</a>        | A concrete point v outside the non-negative orthant.                               |
| <a href="#">BSinMeasurementTheory/Course21241/SeparatingU.lean</a>   | The standard-basis separating vector $u = e_0$ .                                   |
| <a href="#">BSinMeasurementTheory/Course21241/Separation.lean</a>    | Euclidean inner product, Riesz identification, and a separating vector.            |
| <a href="#">BSinMeasurementTheory/Course21241/Space.lean</a>         | The non-negative orthant cone in three-space.                                      |
| <a href="#">BSinMeasurementTheory/Course21292.lean</a>               | Umbrella importer for the 21-292 modules.                                          |
| <a href="#">BSinMeasurementTheory/Course21292/Instance.lean</a>      | A concrete 3x3 Scott-type linear program.                                          |
| <a href="#">BSinMeasurementTheory/Course21292/LinearProgram.lean</a> | Inequality-form linear programs over the rationals.                                |
| <a href="#">BSinMeasurementTheory/Course21292/WeakDuality.lean</a>   | Weak duality for a primal/dual linear program.                                     |
| <a href="#">BSinMeasurementTheory/Course21292Numerical.lean</a>      | Numerical checks of the four calculations in the 21-292 example.                   |
| <a href="#">BSinMeasurementTheory/Course21321.lean</a>               | Umbrella importer for the 21-321 modules.                                          |
| <a href="#">BSinMeasurementTheory/Course21321/Kraft.lean</a>         | Kraft difference vectors as a ScottPair with a cancellation witness.               |
| <a href="#">BSinMeasurementTheory/Course21321/ScottPair.lean</a>     | Scott pairs, separating functionals, and the cancellation-witness class.           |
| <a href="#">BSinMeasurementTheory/Course21321/Theorem11.lean</a>     | Scott's Theorem 1.1: a separable pair satisfies real non-cancellation.             |
| <a href="#">BSinMeasurementTheory/Course21321/Theorem12.lean</a>     | Scott's Theorem 1.2: rational separability implies combinatorial non-cancellation. |

| File                                                                        | Contents                                                                                      |
|-----------------------------------------------------------------------------|-----------------------------------------------------------------------------------------------|
| <a href="#">BSinMeasurementTheory/Course21321/Theorem13.lean</a>            | Scott's Theorem 1.3 (one direction): rational non-cancellation implies real non-cancellation. |
| <a href="#">BSinMeasurementTheory/Course21322.lean</a>                      | Umbrella importer for the 21-322 modules.                                                     |
| <a href="#">BSinMeasurementTheory/Course21322/AtomIso.lean</a>              | Canonical bijection between atoms and the Stone space of a finite algebra.                    |
| <a href="#">BSinMeasurementTheory/Course21322/Hom.lean</a>                  | Boolean homomorphisms from a power set into Bool, and principal evaluations.                  |
| <a href="#">BSinMeasurementTheory/Course21322/MeasureCollapse.lean</a>      | Collapse of Stone-space measure evaluation to the discrete atomic measure.                    |
| <a href="#">BSinMeasurementTheory/Course21322/Numerical.lean</a>            | Direct calculation of the atomic measure on $\{0, 2\}$ .                                      |
| <a href="#">BSinMeasurementTheory/Course21322/StoneClopen.lean</a>          | The Stone clopen associated with an element and preservation of membership.                   |
| <a href="#">BSinMeasurementTheory/Course21329.lean</a>                      | Umbrella importer for the 21-329 modules.                                                     |
| <a href="#">BSinMeasurementTheory/Course21329/Cardinality.lean</a>          | Cardinal-arithmetic argument that the Stone space is large.                                   |
| <a href="#">BSinMeasurementTheory/Course21329/Filter.lean</a>               | Filters, proper filters, and ultrafilters on a bounded lattice.                               |
| <a href="#">BSinMeasurementTheory/Course21329/StoneNonempty.lean</a>        | The principal filter generated by top, and non-emptiness of the Stone space.                  |
| <a href="#">BSinMeasurementTheory/Course21329/UltrafilterExtension.lean</a> | Ultrafilter extension: a chain of proper filters unions to a proper filter.                   |
| <a href="#">BSinMeasurementTheory/Course21355.lean</a>                      | Umbrella importer for the 21-355 modules.                                                     |
| <a href="#">BSinMeasurementTheory/Course21355/AltSeq.lean</a>               | Umbrella importer for this course.                                                            |
| <a href="#">BSinMeasurementTheory/Course21355/AltSeqDiverges.lean</a>       | Epsilon-argument that the alternating sequence diverges.                                      |
| <a href="#">BSinMeasurementTheory/Course21355/AltSeqValues.lean</a>         | Even and odd values of the sequence $(-1)^n$ .                                                |
| <a href="#">BSinMeasurementTheory/Course21355/BolzanoWeierstrass.lean</a>   | Bolzano-Weierstrass: a bounded real sequence has a convergent subsequence.                    |
| <a href="#">BSinMeasurementTheory/Course21355/CompactIcc.lean</a>           | Closed bounded intervals in $\mathbb{R}$ are compact.                                         |
| <a href="#">BSinMeasurementTheory/Course21355/LimsupLiminf.lean</a>         | Umbrella importer for this course.                                                            |
| <a href="#">BSinMeasurementTheory/Course21355/LimsupLiminfExist.lean</a>    | Existence of limsup and liminf for a bounded real sequence.                                   |
| <a href="#">BSinMeasurementTheory/Course21355/Numerical.lean</a>            | Computable alternating sequence used to inspect the example of (c).                           |
| <a href="#">BSinMeasurementTheory/Course21355/TailExtremes.lean</a>         | Tail supremum and tail infimum, and their monotonicity.                                       |

| File                                                                | Contents                                                                                     |
|---------------------------------------------------------------------|----------------------------------------------------------------------------------------------|
| <code>BSinMeasurementTheory/Course21373.lean</code>                 | Umbrella importer for the 21-373 modules.                                                    |
| <code>BSinMeasurementTheory/Course21373/AtomJoin.lean</code>        | Join and disjointness properties of Boolean atoms.                                           |
| <code>BSinMeasurementTheory/Course21373/BooleanAction.lean</code>   | Boolean automorphisms and atoms of the fixed-point subalgebra.                               |
| <code>BSinMeasurementTheory/Course21373/IsBooleanAtom.lean</code>   | Atoms in a Boolean algebra.                                                                  |
| <code>BSinMeasurementTheory/Course21373/Representation.lean</code>  | Canonical representation of an element as the join of the atoms below it.                    |
| <code>BSinMeasurementTheory/Course21373/Numerical.lean</code>       | Numerical example for the finite Boolean-algebra representation.                             |
| <code>BSinMeasurementTheory/Course21410.lean</code>                 | Umbrella importer for the 21-410 modules.                                                    |
| <code>BSinMeasurementTheory/Course21410/Glue.lean</code>            | Glue lemmas: inclusion-exclusion, monotonicity, and preference preservation.                 |
| <code>BSinMeasurementTheory/Course21410/Measure.lean</code>         | Finitely additive probability measures on a finite space.                                    |
| <code>BSinMeasurementTheory/Course21410/Numerical.lean</code>       | Numerical checks reproducing the English example.                                            |
| <code>BSinMeasurementTheory/Course21640.lean</code>                 | Umbrella importer for the 21-640 modules.                                                    |
| <code>BSinMeasurementTheory/Course21640/Numerical.lean</code>       | Sup-norm example of a sublinear functional and a dominated extension on the plane.           |
| <code>BSinMeasurementTheory/Course21640/Positivity.lean</code>      | A norm-dominated linear functional is positive.                                              |
| <code>BSinMeasurementTheory/Course21640/Sublinear.lean</code>       | Sublinear functionals and dominated linear extensions.                                       |
| <code>BSinMeasurementTheory/Course21651.lean</code>                 | Umbrella importer for the 21-651 modules.                                                    |
| <code>BSinMeasurementTheory/Course21651/BooleanHom.lean</code>      | Boolean algebra homomorphisms from $B$ to $\text{Bool}$ .                                    |
| <code>BSinMeasurementTheory/Course21651/ClosedEmbedding.lean</code> | The Stone space as an intersection of closed equalizer sets in $B \rightarrow \text{Bool}$ . |
| <code>BSinMeasurementTheory/Course21651/Separation.lean</code>      | Separation properties used by the Stone embedding argument.                                  |
| <code>BSinMeasurementTheory/Course21651/StoneMap.lean</code>        | The Stone clopen associated with an element of $B$ .                                         |
| <code>BSinMeasurementTheory/Course21651/TwoAlgebra.lean</code>      | Stone space of the four-element Boolean algebra on $\text{Fin } 2$ .                         |
| <code>BSinMeasurementTheory/Course21720.lean</code>                 | Umbrella importer for the 21-720 modules.                                                    |
| <code>BSinMeasurementTheory/Course21720/Boundedness.lean</code>     | Positive linear functionals on $C(X, \mathbb{R})$ are bounded.                               |

| File                                                             | Contents                                                                    |
|------------------------------------------------------------------|-----------------------------------------------------------------------------|
| <code>BSinMeasurementTheory/Course21720/Numerical.lean</code>    | The concrete functional $\text{Lambda}(f) = 3 f(0) + 2 f(1)$ .              |
| <code>BSinMeasurementTheory/Course21720/StoneMeasure.lean</code> | Finitely additive measures on a Boolean algebra from a positive functional. |

**Total:** 88 modules, 3006 lines.